

UNIFORMLY RECTIFIABLE METRIC SPACES SATISFY THE WEAK CONSTANT DENSITY CONDITION

JARED KRANDEL

CONTENTS

1.	Introduction	1
2.	Preliminaries	4
3.	Oscillation of means of L^2 functions	6
4.	Bi-Lipschitz images satisfy the WCD	11
5.	Uniformly rectifiable metric spaces satisfy the WCD	14
	References	19

ABSTRACT. The weak constant density condition is a quantitative regularity property originally investigated by David and Semmes in their foundational work on uniform rectifiability. Roughly speaking, a space satisfies this condition if in *most* balls, the space supports a measure with *nearly constant* density in a neighborhood of scales and locations. In this paper, we prove that uniformly rectifiable metric spaces satisfy the weak constant density condition, an extension of a theorem of David and Semmes. In order to do this, we first prove a strengthened version of a separate theorem of David and Semmes which weakly controls the oscillation of the means of L^∞ functions over normed balls.

1. INTRODUCTION

The relation between the rectifiability properties and density properties of sets and measures has long been a topic of interest in geometric measure theory. For a metric space X , we say that $E \subseteq X$ with $\mathcal{H}^n(E) < \infty$ is n -rectifiable if E can be covered \mathcal{H}^n almost everywhere by a countable union of Lipschitz images of subsets of \mathbb{R}^n . In Euclidean space, n -rectifiable sets give a natural generalization of n -dimensional C^1 manifolds. One of many interesting characterizations of n -rectifiability in Euclidean spaces involves the Hausdorff density, which measures how the \mathcal{H}^n measure of small balls around a point compare to the \mathcal{H}^n measure of balls of equal radius in \mathbb{R}^n .

Theorem 1.1. *Let $E \subseteq \mathbb{R}^d$ be \mathcal{H}^n measurable with $\mathcal{H}^n(E) < \infty$. E is n -rectifiable if and only if for \mathcal{H}^n -a.e. $x \in E$,*

$$(1.1) \quad \lim_{r \rightarrow 0} \frac{\mathcal{H}^n(E \cap B(x, r))}{(2r)^n} = 1.$$

The backward direction was proven by Besicovitch for $n = 1$, $d = 2$ [Bes28],[Bes38], Marstrand for $n = 2$, $d = 3$ [Mar61] and Mattila for general $n \leq d$ [Mat75]. Later, Preiss showed that any measure in \mathbb{R}^d whose n -dimensional density merely exists and is positive and finite \mathcal{H}^n -a.e. is n -rectifiable, generalizing this result significantly [Pre87].

The forward direction of Theorem 1.1 follows more readily from the almost everywhere differentiability of Lipschitz maps, but it was not until Kircheim gave a notion of metric differentiability for maps from \mathbb{R}^n into X that Theorem 1.1 received the following one-sided metric space analog.

Theorem 1.2 (See [Kir94] Theorem 9). *Let $E \subseteq X$ be n -rectifiable. Then (1.1) holds at \mathcal{H}^n -a.e. $x \in E$.*

It follows from work of Preiss and Tis̄er [PT92] that the converse of Theorem 1.2 holds when $n = 1$, but it remains an interesting and difficult open question whether the converse holds for general n .

The weak constant density condition (WCD) is one condition meant to provide an analog of (1.1) in the world of *uniform rectifiability* pioneered by David and Semmes in their foundational works [DS91] and [DS93].

Definition 1.3 (uniform n -rectifiability). We say that a set $E \subseteq \mathbb{R}^d$ is *uniformly n -rectifiable* if there exists a constant $C_0 > 0$ such that E is Ahlfors (C_0, n) -regular, i.e., for all $x \in E$ and $0 < r < \text{diam}(E)$,

$$(1.2) \quad C_0^{-1}r^n \leq \mathcal{H}^n(B(x, r) \cap E) \leq C_0r^n,$$

and E has *Big Pieces of Lipschitz images of \mathbb{R}^n* (BPLI), i.e., there exist constants $L, \theta > 0$ such that for all $x \in E$ and $0 < r < \text{diam}(E)$, there exists an L -Lipschitz map $f : B(0, r) \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^d$ such that

$$(1.3) \quad \mathcal{H}^n(B(x, r) \cap E \cap f(B(0, r))) \geq \theta r^n.$$

One can think of this as a stronger form of n -rectifiability in which one requires a uniform percentage of the measure of each ball to be covered by a Lipschitz image. David and Semmes introduced the WCD as a way of quantifying (1.1) by requiring that in almost every *ball*, there exists a measure supported on the set with *nearly constant* density nearby.

Definition 1.4 (Weak constant density condition, Carleson sets and measures). Let $E \subseteq \mathbb{R}^d$ be Ahlfors n -regular, let $C_0, \epsilon_0 > 0$, and define

$$(1.4) \quad \mathcal{G}_{\text{cd}}(C_0, \epsilon_0) = \left\{ (x, r) \in E \times \mathbb{R}^+ \left| \begin{array}{l} \exists \text{ Ahlfors } (C_0, n)\text{-regular } \mu, \text{ spt}(\mu) = E, \\ \forall y \in B(x, r), 0 < t \leq r, \\ |\mu(E \cap B(y, t)) - t^n| \leq \epsilon_0 r^n \end{array} \right. \right\},$$

$$(1.5) \quad \mathcal{B}_{\text{cd}}(C_0, \epsilon_0) = E \times \mathbb{R}^+ \setminus \mathcal{G}_{\text{cd}}(C_0, \epsilon_0).$$

We say that E satisfies the weak constant density condition if there exists $C_0 > 0$ such that for all $\epsilon_0 > 0$, $\mathcal{B}_{\text{cd}}(C_0, \epsilon_0)$ is a Carleson set. That is, there exists a constant $C_1 > 0$ such that for all $z \in E$ and $0 < r < \text{diam}(E)$,

$$\int_{B(z, r) \cap E} \int_0^r \chi_{\mathcal{B}_{\text{cd}}(C_0, \epsilon_0)}(x, t) d\mathcal{H}^n(x) \frac{dt}{t} \leq C_1 r^n.$$

If this holds, we say that $\chi_{\mathcal{B}_{\text{cd}}(C_0, \epsilon_0)} d\mathcal{H}^n(x) \frac{dt}{t}$ is a *Carleson measure* and say that $\mathcal{B}_{\text{cd}}(C_0, \epsilon_0)$ is C_1 -*Carleson*.

For related quantitative conditions involving densities, see [CGLT16], [AH22], and [TT15]. The work of David, Semmes, and Tolsa combine to prove the following Theorem:

Theorem 1.5. *Let $E \subseteq \mathbb{R}^d$ be Ahlfors n -regular. Then E is uniformly n -rectifiable if and only if E satisfies the WCD.*

David and Semmes proved the forward implication in Chapter 6 of [DS91] using a characterization of uniform rectifiability (condition C2 of [DS91]) more closely related to the boundedness of singular integral operators. We will say more about this when we discuss our result.

They proved the reverse implication only in the case $n = 1, 2$, and $d - 1$. Their proof uses the fact that if a measure is very close to having constant density in a large neighborhood of scales and locations, then its support is well-approximated by the support of an n -uniform measure, a measure μ for which there exists $c > 0$ such that $\mu(B(x, r)) = cr^n$ for all $x \in \text{spt}(\mu)$ and $r > 0$. Because uniform measures in Euclidean space are completely classified for $n = 1, 2$ (they are all multiples of Hausdorff measure on a plane) and for $n = d - 1$ (they are Hausdorff measure on products of planes and light cones [KP87]), David and Semmes are able to show that a WCD set is very close to flat on most balls which are good for the WCD. The absence of a classification for uniform measures in intermediate dimensions prevented a direct adaptation of their arguments. However, Tolsa completed the proof of the reverse direction in Theorem 1.5 in [Tol15] by replacing elements of David and Semmes's argument specific to their examples of uniform measures with general flatness properties of uniform measures derived by Preiss [Pre87] in addition to new arguments using the Riesz transform.

In general, classifying uniform measures is a difficult open problem, but see [Nim22] for an interesting family of examples. For further studies of uniform measures in Euclidean spaces see [Pre87] (and [DL08] for a more gentle presentation of Preiss), [KP02], [Nim17], and [Nim19]. For research into uniform measures in the Heisenberg group see [CMT20] and [Mer22] and for a related result in ℓ_∞^3 , see [Lor03].

Just as Theorem 1.5 provides a quantitative analog of Theorem 1.1, one might expect a quantitative analog of Kirchheim's result, Theorem 1.2, to hold for *uniformly rectifiable metric spaces*, i.e., metric spaces which are Ahlfors n -regular and have big pieces of Lipschitz images of subsets of \mathbb{R}^n . In this paper, we provide such an analog by proving the following theorem.

Theorem A. *Uniformly n -rectifiable metric spaces satisfy the WCD.*

Our proof is made possible by the recent work of Bate, Hyde, and Schul [BHS23] which adapted a substantial portion of the theory of uniformly rectifiable subsets of Euclidean spaces to metric spaces. While our argument uses this theory, it does not follow David and Semmes's original proof. Roughly speaking, David and Semmes proved the Euclidean version of Theorem A by showing that for most balls centered on a uniformly n -rectifiable set E , there exists an n -plane P such that the pushforward of Hausdorff measure for E onto P must be very close to symmetric. They do this by showing that the non-symmetric balls contribute substantially to the value of a Carleson measure defined using integrals over E of a family of smooth odd functions designed to detect asymmetry.

We prove Theorem A by first proving WCD estimates for bi-Lipschitz images. Then, using the fact that uniformly rectifiable spaces have very big pieces of bi-Lipschitz images in Banach spaces proven by Bate, Hyde, and Schul [BHS23], we

adapt the bi-Lipschitz image arguments to the general case. To handle the bi-Lipschitz image case, we first prove Lemma 3.6, a form of quantitative Lebesgue differentiation theorem for L^2 functions similar to theorems considered by David and Semmes (See [DS93] Lemma IV.2.2.14 and Corollary IV.2.2.19, and see Remark 3.7 for a discussion of the difference with our result), although our proof proceeds by contradiction, a method which differs significantly from their proofs. We apply this lemma to the Jacobian of the metric derivative of our bi-Lipschitz function $f : \mathbb{R}^n \rightarrow \Sigma$ to control the variation of its averages over neighborhoods of scales and locations and receive control over the variation of the Hausdorff measure of Σ using the area formula. To the knowledge of the author, this gives a new proof of the WCD even in the Euclidean case.

We note here that the naive converse of Theorem 1.5 is false: There exist Ahlfors regular metric spaces which satisfy the WCD, yet are not uniformly rectifiable. Indeed, the metric space $(X, d) = (\mathbb{R}, |\cdot|_{\text{Euc}}^{1/2})$ is in fact 2-uniform: $\mathcal{H}^2(B(x, r)) = 2r^2$ for all $x \in \mathbb{R}$ and $r \geq 0$, hence X satisfies the weak constant density condition, yet X is purely 2-unrectifiable (notice that the Hausdorff 2-density is everywhere $1/2$ so that this space does not give a counterexample to the potential converse of Theorem 1.2). Some different examples of this failure even in the case $n = 1$ are given by Bate [Bat23]. He proves that every 1-uniform metric measure space is either \mathbb{R} , a particular union of disjoint circles of radius d , or a purely unrectifiable “limit” of the circle spaces. These last two spaces are examples of 1-uniform spaces which are not uniformly rectifiable.

Analyzing connectedness plays a special role in the proof because any 1-uniform connected component must be locally isometric to \mathbb{R} , implying any connected 1-uniform space is itself \mathbb{R} . From these examples, it seems reasonable to think that some connectedness and topological conditions are necessary hypotheses for any type of converse to hold. It also follows from work of Schul [Sch07], [Sch09], and Fassler and Violo [FV23] (see also [Hah05]) that any Ahlfors 1-regular connected subset of a metric space is uniformly 1-rectifiable, although perhaps adding some form of weaker hypothesis could provide an interesting converse to our result in the one-dimensional case using Bate’s classification.

2. PRELIMINARIES

Whenever we write $A \lesssim B$, we mean that there exists some constant C independent of A and B such that $A \leq CB$. If we write $A \lesssim_{a,b,c} B$ for some constants a, b, c , then we mean that the implicit constant C mentioned above is allowed to depend on a, b, c . We will sometimes write $A \succ_{a,b,c} B$ to mean that both $A \lesssim_{a,b,c} B$ and $B \lesssim_{a,b,c} A$ hold. We use the notation $f : E \twoheadrightarrow F$ to mean f is a surjective map from E to F .

Let (X, d) be a metric space. For any subset $F \subseteq X$, integer $n \geq 0$, and constant $0 < \delta \leq \infty$, we define

$$\mathcal{H}_\delta^n(F) = \inf \left\{ \sum \text{diam}(E_i)^d : F \subseteq \bigcup E_i, \text{diam}(E_i) < \delta \right\}$$

where $\text{diam}(E) = \sup_{x,y \in E} d(x,y)$. The Hausdorff n -measure of F is defined as

$$\mathcal{H}^n(F) = \lim_{\delta \rightarrow 0} \mathcal{H}_\delta^n(F).$$

Occasionally, we will specify a subset $\Sigma \subseteq X$ and write $\mathcal{H}_\Sigma^n = \mathcal{H}^n|_\Sigma$. For any \mathcal{H}^n measurable $A \subseteq X$ and measurable $f : A \subseteq X \rightarrow \mathbb{R}$, we define

$$\int_A f = \frac{1}{\mathcal{H}^n(A)} \int_A f(x) d\mathcal{H}^n(x).$$

We let $\mathcal{D}(\mathbb{R}^n)$ denote the family of dyadic cubes in \mathbb{R}^n . For $Q \in \mathcal{D}(\mathbb{R}^n)$, we let $\ell(Q)$ denote the side length of Q . If $R \in \mathcal{D}(\mathbb{R}^n)$ and $k \in \mathbb{Z}$, we define

$$\begin{aligned} \mathcal{D}(R) &= \{ Q \in \mathcal{D}(\mathbb{R}^n) \mid Q \subseteq R \}, \\ \mathcal{D}_k(R) &= \{ Q \in \mathcal{D}(R) \mid \ell(Q) = 2^{-k}\ell(R) \}. \end{aligned}$$

We will also need a version of ‘‘cubes’’ associated to a metric space. David [Dav88] introduced this idea first, and it was later generalized by [Chr90] and [HM12]. The following formulation draws most from the latter two.

Theorem 2.1 (Christ-David cubes). *Let X be a doubling metric space. Let X_k be a nested sequence of maximal ρ^k -nets for X where $\rho < 1/1000$ and let $c_0 = 1/500$. For each $k \in \mathbb{Z}$ there is a collection \mathcal{D}_k of ‘‘cubes,’’ which are Borel subsets of X such that the following hold.*

- (i) $X = \bigcup_{Q \in \mathcal{D}_k} Q$.
- (ii) If $Q, Q' \in \mathcal{D} = \bigcup \mathcal{D}_k$ and $Q \cap Q' \neq \emptyset$, then $Q \subseteq Q'$ or $Q' \subseteq Q$.
- (iii) For $Q \in \mathcal{D}$, let $k(Q)$ be the unique integer so that $Q \in \mathcal{D}_k$ and set $\ell(Q) = 5\rho^{k(Q)}$. Then there is $x_Q \in X_k$ so that

$$B(x_Q, c_0\ell(Q)) \subseteq Q \subseteq B(x_Q, \ell(Q))$$

and

$$X_k = \{x_Q : Q \in \mathcal{D}_k\}.$$

- (iv) If X is Ahlfors n -regular, then there exists $C \geq 1$ such that

$$\mathcal{H}^n(\{x \in Q \mid d(x, X \setminus Q) \leq \eta\rho^k\}) \lesssim \eta^{1/C} \ell(Q)^n$$

for all $Q \in \mathcal{D}$ and $\eta > 0$.

In addition, we define

$$B_Q = B(x_Q, \ell(Q)).$$

In analogy to the dyadic cube notation, for any $R \in \mathcal{D}$ and $k \in \mathbb{Z}$ we also write

$$\begin{aligned} \mathcal{D}(R) &= \{ Q \in \mathcal{D} \mid Q \subseteq R \}, \\ \mathcal{D}_k(R) &= \{ Q \in \mathcal{D}(R) \mid \ell(Q) = \rho^{-k}\ell(R) \}. \end{aligned}$$

We will actually prove a form of the WCD adapted to Christ-David cubes. The following two lemmas will allow us to show that the cube WCD in Definition 4.1 implies the WCD from Definition 1.4. Recall from Theorem 2.1 that $c_0 = \frac{1}{500}$.

Lemma 2.2. *Let X be a doubling metric space with doubling constant C_d . There exists $N(C_d) \in \mathbb{N}$ such that the following holds: There exist N Christ-David systems of cubes $\{\mathcal{D}_i\}_{i=1}^N$ for X such that for any $x \in X$, $0 < t < \text{diam}(X)$, there exists $i \in \{1, \dots, N\}$ and $Q \in \mathcal{D}_i$ with $\ell(Q) \leq \frac{5}{\rho c_0}t$ such that $x \in \frac{c_0}{4}B_Q$ and $t < \frac{c_0}{4}\ell(Q)$.*

Proof. Fix $\rho < \frac{1}{1000}$. For each k , let \tilde{X}_k be a maximal $c_0\rho^k$ -net for X . We now iteratively construct maximal ρ^k -nets $X_k^1, X_k^2, \dots, X_k^N, \dots$ in the following way. Let X_k^1 be a completion of a maximal ρ^k -separated subset of \tilde{X}_k to a maximal ρ^k -net for X . Given X_k^i for any $i > 0$, construct X_k^{i+1} by completing a maximal ρ^k -separated

subset of $Y_k^i := \tilde{X}_k \setminus (X_k^1 \cup X_k^2 \cup \dots \cup X_k^i)$ to a maximal ρ^k -net for X . We claim that this process terminates in $N(C_d)$ steps, giving for each $k \in \mathbb{Z}$ a collection of maximal ρ^k -nets X_k^1, \dots, X_k^N . Indeed, let B be a ball of radius $2\rho^k$. By doubling, there exists $N(C_d) < \infty$ such that $\#(B \cap \tilde{X}_k) \leq N(C_d)$. Suppose that $\frac{1}{2}B \cap Y_k^j \neq \emptyset$ for some $j > 0$. Then, because X_k^{j+1} is maximal, there exists some $x \in B \cap Y_k^j$ such that $x \in X_k^{j+1}$. Therefore, $\#(B \cap Y_k^{j+1}) < \#(B \cap Y_k^j)$ whenever $\frac{1}{2}B \cap Y_k^j \neq \emptyset$. This means $\frac{1}{2}B \cap Y_k^{N+1} = \emptyset$ for any such B , implying $Y_k^{N+1} = \emptyset$ and $\tilde{X}_k \subseteq \cup_{i=1}^N X_k^i$ as desired.

We now show that the lemma follows from this. Recall that Theorem 2.1 takes as input a collection $\{X_k\}_{k \in \mathbb{Z}}$ of maximal ρ^k -nets for X and outputs a system of cubes \mathcal{D} such that every $x_k^\alpha \in X_k$ is the ‘‘center’’ of a cube $Q_k^\alpha \in \mathcal{D}$ with $B_X(x_k^\alpha, c_0 5\rho^k) = c_0 B_{Q_k^\alpha} \subseteq Q_k^\alpha$. We apply Theorem 2.1 to the collection $\{X_k^i\}_{k \in \mathbb{Z}}$ for every $1 \leq i \leq N$ and receive a Christ-David system \mathcal{D}_i such that each point $\tilde{x}_k \in \tilde{X}_k$ is the center of some $Q \in \mathcal{D}_i$ for some i . So, let $x \in X$, $0 < t < \text{diam}(X)$, and let $k \in \mathbb{Z}$ such that $c_0 \rho^{k-1} \leq t < c_0 \rho^k$. Because \tilde{X}_k is a maximal $c_0 \rho^k$ -net for X , there exists $\tilde{x}_k \in \tilde{X}_k$ such that $d(x, \tilde{x}_k) < c_0 \rho^k$. Because $\tilde{X}_k \subseteq \cup_{i=1}^N X_k^i$, there then exists $1 \leq i \leq N$ and $Q \in \mathcal{D}_i$ such that $\tilde{x}_k = x_Q$ so that $x \in B(x_Q, c_0 \rho^k) = \frac{1}{5}B(x_Q, c_0 \ell(Q)) = \frac{c_0}{5}B_Q$. Similarly, $\frac{\rho^{c_0}}{5}\ell(Q) \leq t < \frac{c_0}{5}\ell(Q)$. ■

Lemma 2.3. *Let X be Ahlfors (C_0, n) -regular. If X satisfies the cube WCD of Definition 4.1, then X satisfies the WCD.*

Proof. We first note that if an $a_Q > 0$ as in (4.1) exists, then $(2C_0)^{-1} \leq a_Q \leq 2C_0$ by Ahlfors regularity. Therefore, whenever $|\mathcal{H}^n(X \cap B(y, r)) - a_Q r^n| \leq \epsilon_0 \ell(Q)$, we have

$$|(a_Q)^{-1} \mathcal{H}^n(X \cap B(y, r)) - r^n| = (a_Q)^{-1} |\mathcal{H}^n(X \cap B(y, r)) - a_Q r^n| \leq (2C_0)^{-1} \epsilon_0 \ell(Q).$$

This means that one can replace \mathcal{H}^n with a multiple of \mathcal{H}^n and $a_Q r^n$ with r^n in the definition of the cube WCD at the cost of increasing ϵ_0 . Therefore, it suffices to show that the complement of

$$\mathcal{G}_0(C_0, \epsilon_0) = \left\{ (x, t) \in X \times \mathbb{R}^+ \mid \begin{array}{l} \exists a_{(x,t)} > 0, \text{ such that } \forall y \in B(x, t), 0 < r \leq t, \\ |\mathcal{H}^n(X \cap B(y, r)) - a_{(x,t)} r^n| \leq \epsilon_0 t^n \end{array} \right\}$$

is a Carleson set. In order to show this, we apply Lemma 2.2 to X and receive a finite number of Christ-David systems $\{\mathcal{D}_i\}_{i=1}^N$ with N depending only on n and C_0 such that for any $x \in X$, $0 < t < \text{diam}(X)$, there exists $i \in \{1, \dots, N\}$ and $Q \in \mathcal{D}_i$ with $\ell(Q) \lesssim t$ such that $x \in \frac{c_0}{4}B_Q$ and $t < \frac{c_0}{4}\ell(Q)$. It follows that if $Q \in \mathcal{G}_{\text{cd}}(C_0, \epsilon_0)$, then $(x, t) \in \mathcal{G}_0(C_0, C(n)\epsilon_0)$ for any $x \in \frac{c_0}{4}B_Q$ and $\frac{\rho^2 c_0}{4}\ell(Q) \leq t < \frac{c_0}{4}\ell(Q)$ by choosing $a_{(x,t)} = a_Q$. Therefore, if X satisfies the cube WCD then \mathcal{D}_i has a Carleson packing condition for its bad set, implying a Carleson condition for the bad balls of the WCD with a larger choice of ϵ_0 and with larger Carleson constant. ■

3. OSCILLATION OF MEANS OF L^2 FUNCTIONS

In this section, we review necessary facts about wavelets and prove Lemma 3.6, one of our main tools for the proof of Theorem A.

Definition 3.1. We follow the presentation of [Tol12]. Given $h : \mathbb{R}^n \rightarrow \mathbb{R}$ and $Q \in \mathcal{D}(\mathbb{R}^n)$, define

$$\Delta_Q h(x) = \begin{cases} \int_P h(z) dz - \int_Q h(z) dz & \text{if } x \in P, \text{ where } P \text{ is a child of } Q, \\ 0 & \text{otherwise} \end{cases}$$

If $h \in L^2(\mathbb{R}^n)$, then

$$h = \sum_{Q \in \mathcal{D}(\mathbb{R}^n)} \Delta_Q h \quad \text{and} \quad h \chi_Q = \int_Q h + \sum_{R \subseteq Q} \Delta_R h$$

where the sums converge in L^2 and $\langle \Delta_Q h, \Delta_{Q'} h \rangle_{L^2} = 0$ when $Q \neq Q'$ so that $\|h\|_2 = \sum_{Q \in \mathcal{D}} \|\Delta_Q h\|_2^2$. One can view $\Delta_Q h$ as a projection of h onto the subspace of L^2 formed by the Haar wavelets h_Q^ϵ , $\epsilon \in \{0, 1\}^n \setminus \{(0, 0, \dots, 0)\}$ associated to Q .

We now use the wavelet-like decomposition of h to define coefficients $\Delta_k^h(Q)$ which, roughly speaking, measure the variation in means of h from Q through to its k -th generation descendants.

Definition 3.2. For any $k \in \mathbb{N}$, define

$$(3.1) \quad \Delta_k^h(Q)^2 = \sum_{j=0}^k \sum_{R \in \mathcal{D}_j(Q)} \|\Delta_R h\|_2^2.$$

Remark 3.3 (Properties of Δ_k^h). Notice that if $h \in L^\infty(\mathbb{R}^n)$, then Δ_k^h has a form of geometric lemma since

$$\sum_{Q \subseteq Q_0} \Delta_k^h(Q)^2 = \sum_{Q \subseteq Q_0} \sum_{j=0}^k \sum_{R \in \mathcal{D}_j(Q)} \|\Delta_R h\|_2^2 \lesssim_{k,n} \sum_{R \subseteq Q_0} \|\Delta_R h\|_2^2 \lesssim_{\|h\|_\infty} \ell(Q_0)^n.$$

This gives the Carleson condition

$$(3.2) \quad \sum_{\substack{Q \subseteq Q_0 \\ \Delta_k^h(Q) > \delta \ell(Q)^{n/2}}} \ell(Q)^n \lesssim_\delta \sum_{Q \subseteq Q_0} \Delta_k^h(Q)^2 \lesssim_{k,n,\|h\|_\infty} \ell(Q_0)^n.$$

Δ_k^h also scales appropriately in the following manner: Let $Q, \tilde{Q} \in \mathcal{D}(\mathbb{R}^n)$ and let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be the affine map sending \tilde{Q} onto Q by

$$(3.3) \quad T(x) = x_Q + \left(\frac{x - x_{\tilde{Q}}}{\ell(\tilde{Q})} \right) \ell(Q)$$

where $x_{\tilde{Q}}$ is the center of \tilde{Q} . Let $\tilde{h} \in L^2(\tilde{Q})$ and set $h = \tilde{h} \circ T^{-1}$. Notice that

$$\|h\|_{L^2(Q)}^2 = \int_{T(\tilde{Q})} (\tilde{h} \circ T^{-1})^2 = \int_{\tilde{Q}} \tilde{h}^2 \frac{\ell(Q)^n}{\ell(\tilde{Q})^n} = \frac{\ell(Q)^n}{\ell(\tilde{Q})^n} \|\tilde{h}\|_2^2.$$

Similarly, notice that if $V \subseteq Q$ and $\tilde{V} \subseteq \tilde{Q}$ with $T(\tilde{V}) = V$, then

$$\begin{aligned} \|\Delta_V h\|_2^2 &= \int_V (\Delta_V h(x))^2 dx = \int_{T(\tilde{V})} (\Delta_{\tilde{V}} \tilde{h}(T^{-1}(x)))^2 dx \\ &= \left(\frac{\ell(Q)}{\ell(\tilde{Q})} \right)^n \int_{\tilde{V}} (\Delta_{\tilde{V}} \tilde{h})^2 dx = \left(\frac{\ell(Q)}{\ell(\tilde{Q})} \right)^n \|\Delta_{\tilde{V}} \tilde{h}\|_2^2 \end{aligned}$$

which gives $\Delta_k^h(Q)^2 = \left(\frac{\ell(Q)}{\ell(\tilde{Q})} \right)^n \Delta_k^{\tilde{h}}(\tilde{Q})^2$.

Definition 3.4 (normed balls). Given $L > 0$, we define the set of norms on \mathbb{R}^n which are L -bi-Lipschitz to the Euclidean norm by

$$\mathcal{N}_L = \{\|\cdot\| : L^{-1}\|x\| \leq |x| \leq L\|x\|\}.$$

Given a dyadic cube $Q \in \mathcal{D}(\mathbb{R}^n)$ and $L > 0$, we define a collection of L -bi-Lipschitz normed balls inside Q by

$$\mathcal{B}_L(Q) = \{B_{\|\cdot\|}(x, r) \subseteq Q : \|\cdot\| \in \mathcal{N}_L, r \geq L^{-1}\ell(Q)\}.$$

The following lemma gives a form of compactness result for $\mathcal{B}_L(Q)$.

Lemma 3.5. *Fix $Q \in \mathcal{D}(\mathbb{R}^n)$ and $L > 0$. Let $B_j = B_{\|\cdot\|_j} \in \mathcal{B}_L(Q)$ for $j \in \mathbb{N}$. There exists a subsequence of $B_j = B_{\|\cdot\|_j}(x_j, r_j)$ and a normed ball $B = B_{\|\cdot\|}(x, r) \in \mathcal{B}_L(Q)$ for which the following holds: For every $\eta > 0$, there exists $j_0 \in \mathbb{N}$ such that for all $j \geq j_0$,*

$$(3.4) \quad B_{\|\cdot\|}(x, (1 - 5\eta)r) \subseteq B_j \subseteq B_{\|\cdot\|}(x, (1 + 5\eta)r).$$

Proof. Because $\|\cdot\|_j \in \mathcal{N}_L$ for all j , the functions $f_j : B(0, 1) \rightarrow \mathbb{R}$ given by $f_j(x) = \|x\|_j$ are an equicontinuous, uniformly bounded family of continuous functions. Therefore, they subconverge uniformly to some limit function $f : B(0, 1) \rightarrow \mathbb{R}$. It is straightforward to show this function gives a norm $\|\cdot\|$ when extended homogeneously to \mathbb{R}^n . By passing to further subsequences, we can assume that $x_j \rightarrow x \in Q$ and $r_j \rightarrow r$ with $L^{-1}\ell(Q) \leq r$. We set $B = B_{\|\cdot\|}(x, r)$ and fix $\eta > 0$ as in the statement of the lemma. By the convergences assumed, we can take j_0 large enough such that for $j \geq j_0$, we have $\|x_j - x\| \leq \eta \min\{r, r_j\}$, $|r_j - r| \leq \eta \min\{r, r_j\}$, and

$$(1 - \eta)\|x\| \leq \|x\|_j \leq (1 + \eta)\|x\|, \text{ for all } x \in \mathbb{R}^n,$$

We now aim to prove (3.4). Let $y \in B_j$. Then

$$\begin{aligned} \|y - x\| &\leq \|y - x_j\| + \|x_j - x\| \leq (1 + 2\eta)\|y - x_j\|_j + \eta r \\ &\leq (1 + 2\eta)(1 + \eta)r + \eta r \leq (1 + 5\eta)r. \end{aligned}$$

so that $y \in B_{\|\cdot\|}(x, (1 + 5\eta)r)$. On the other hand, if $z \in B_{\|\cdot\|}(x, (1 - 5\eta)r)$, then

$$\begin{aligned} \|z - x_j\|_j &\leq \|z - x\|_j + \|x - x_j\|_j \leq (1 + 2\eta)\|z - x\| + (1 + 2\eta)\|x - x_j\| \\ &\leq (1 + 2\eta)(1 - 5\eta)r_j + (1 + 2\eta)\eta r_j \leq r_j \end{aligned}$$

showing $z \in B_j$. ■

The following lemma shows that in order to control the means of h over balls in $\mathcal{B}_L(Q)$, it is sufficient for $\Delta_k^h(Q)$ to be sufficiently small.

Lemma 3.6. *For all $\epsilon, M, L > 0$, $n \in \mathbb{N}$, there exist $k(\epsilon, M, L, n) \in \mathbb{N}$ and $\delta(\epsilon, M, L, n) > 0$ such that the following holds: Suppose $h \in L^2(\mathbb{R}^n)$ and $Q \in \mathcal{D}(\mathbb{R}^n)$ are such that*

- (i) $h \geq 0$ a.e.,
- (ii) $\|h\|_{L^2(Q)}^2 \leq M\ell(Q)^n$,
- (iii) $\Delta_k^h(Q)^2 \leq \delta\ell(Q)^n$.

Then, for any normed ball $B \in \mathcal{B}_L(Q)$, we have

$$(3.5) \quad \left| \int_B h - \int_Q h \right| \leq \epsilon.$$

Proof. Suppose the conclusion of the lemma is false. Then there exist ϵ, M, L, n and a sequence of maps $\tilde{h}_j \in L^2(\mathbb{R}^n)$ and cubes $\tilde{Q}_j \in \mathcal{D}(\mathbb{R}^n)$ with $\tilde{h}_j \geq 0$, $\|\tilde{h}_j\|_2 \leq M\ell(\tilde{Q}_j)^n$, and normed balls $\tilde{B}_j \in \mathcal{B}_L(\tilde{Q}_j)$ so that (3.5) does not hold for $\tilde{h}_j, \tilde{Q}_j, \tilde{B}_j$, yet $\Delta_j^{h_j}(\tilde{Q}_j) \leq \frac{1}{j}\ell(\tilde{Q}_j)^{n/2}$. For any $j \in \mathbb{N}$, let T_j be the affine transformation sending \tilde{Q}_j onto $Q = [0, 1]^n$ as in (3.3) and define $h_j : Q \rightarrow \mathbb{R}$ as in Remark 3.3 by

$$h_j = \tilde{h}_j \circ T_j^{-1}.$$

It follows from Remark 3.3 that $\|h_j\|_{L^2(Q)}^2 \leq M$ and $\Delta_j^{h_j}(Q)^2 \leq \frac{1}{j}$. We also define $B_j = T_j(\tilde{B}_j)$ to be the appropriately translated and scaled copy of \tilde{B}_j .

By the weak compactness of bounded closed balls in L^2 , there exists some $h \in L^2(Q)$ such that $h_j \rightharpoonup h$ in L^2 for some subsequence of h_j . By further refining subsequences and using Lemma 3.5, we can further assume the subsequence is chosen so that a limiting normed ball $B = B_{\|\cdot\|}(x, r) \in \mathcal{B}_L(Q)$ as in the lemma's conclusion exists. Let $c = \int_Q h$ and let $c_j = \int_Q h_j$.

We will first show that $h = c$ by showing that $\Delta_V h = 0$ for all $V \subseteq Q$. By weak convergence we have

$$(3.6) \quad c_j = \int_Q h_j \rightarrow \int_Q h = c.$$

Write $h_j = c_j + \sum_{R \subseteq Q} \Delta_R h_j$ and $h = c + \sum_{R \subseteq Q} \Delta_R h$. Fix $V \subseteq Q$ and observe

$$\begin{aligned} \int_Q h_j \Delta_V h &= \int_Q \left(c_j + \sum_{R \subseteq Q} \Delta_R h_j \right) \Delta_V h = c_j \int_Q \Delta_V h + \sum_{R \subseteq Q} \langle \Delta_R h_j, \Delta_V h \rangle \\ &= \langle \Delta_V h_j, \Delta_V h \rangle. \end{aligned}$$

where the final equality follows since $\langle \Delta_R f_1, \Delta_V f_2 \rangle = 0$ whenever $f_1, f_2 \in L^2$ and $R \neq V$. Similarly, we have

$$\int_Q h \Delta_V h = \langle h, \Delta_V h \rangle = \langle \Delta_V h, \Delta_V h \rangle = \|\Delta_V h\|_2^2$$

Using weak convergence again, we get

$$\langle \Delta_V h_j, \Delta_V h \rangle = \int_Q h_j \Delta_V h \rightarrow \int_Q h \Delta_V h = \|\Delta_V h\|_2^2.$$

Using Cauchy-Schwarz, we can now conclude that $\|\Delta_V h\|_2 \leq \lim_j \|\Delta_V h_j\|_2$. We claim that $\|\Delta_V h\|_2 = 0$. Indeed, if j is sufficiently large, then both $V \in \mathcal{D}_{j'}(Q)$ for some $j' \leq j$ and $\|\Delta_V h\|_2 \leq 2\|\Delta_V h_j\|_2$. This means

$$\|\Delta_V h\|_2^2 \leq 4\|\Delta_V h_j\|_2^2 \leq 4 \sum_{k=0}^j \sum_{R \in \mathcal{D}_k(Q)} \|\Delta_R h_j\|_2^2 = 4\Delta_j^{h_j}(Q) \leq \frac{4}{j}.$$

for all large j . This shows that $\Delta_V h = 0$ for all $V \subseteq Q$, hence $h = c$ as desired.

We will now show how this leads to a contradiction. Let $\eta > 0$ and choose j large enough so that

$$B_1 := B_{\|\cdot\|}(x, (1 - 5\eta)r) \subseteq B_j \subseteq B_{\|\cdot\|}(x, (1 + 5\eta)r) =: B_2.$$

Using the fact that $h_j \geq 0$ (this is the first use of this hypothesis),

$$(3.7) \quad \int_{B_1} h_j \leq \int_{B_j} h_j \leq \int_{B_2} h_j$$

so that

$$\frac{\mathcal{L}^n(B_1)}{\mathcal{L}^n(B_j)} \int_{B_1} h_j - \int_Q h_j \leq \int_{B_j} h_j - \int_Q h_j \leq \frac{\mathcal{L}^n(B_2)}{\mathcal{L}^n(B_j)} \int_{B_2} h_j - \int_Q h_j.$$

Because $\mathcal{L}^n(B_1) \leq \mathcal{L}^n(B_j) \leq \mathcal{L}^n(B_2)$ and $\frac{\mathcal{L}^n(B_2)}{\mathcal{L}^n(B_1)} \leq (1 + c'(n)\eta)$, we can assume without loss of generality that

$$\limsup_j \left| \int_{B_j} h_j - \int_Q h_j \right| \leq \limsup_j \left| \frac{\mathcal{L}^n(B_2)}{\mathcal{L}^n(B_1)} \int_{B_2} h_j - \int_Q h_j \right| \leq |(1 + c'\eta)c - c| \lesssim_n \eta c.$$

Since this holds for all $\eta > 0$, we get $\limsup_j \left| \int_{B_j} h_j - \int_Q h_j \right| = 0$. On the other hand, by hypothesis

$$\left| \int_{B_j} h_j - \int_Q h_j \right| > \epsilon$$

for all j , giving a contradiction. \blacksquare

Remark 3.7. Suppose that we only want to conclude (3.5) with normed balls $B \in \mathcal{B}_L(Q)$ replaced by $Q' \in \mathcal{D}_j(Q)$ for $j \leq k \in \mathbb{N}$. The following stronger condition holds even without the positivity assumption for h : Let $\alpha : \mathcal{D}(\mathbb{R}^n) \rightarrow \mathcal{D}(\mathbb{R}^n)$ where $\alpha(Q) \in \mathcal{D}_j(Q)$. There exists $C_0(k, n) > 0$ such that

$$(3.8) \quad \sum_{Q \subseteq R} \left| \int_{\alpha(Q)} h - \int_Q h \right|^2 \ell(Q)^n \leq C_0 \|h\|_2^2.$$

The proof is straightforward: because $\alpha(Q) \in \mathcal{D}_j(Q)$, there is a chain of at most $k + 1$ cubes $\alpha(Q) = Q_j \subseteq Q_{j-1} \subseteq \dots \subseteq Q_0 = Q$ such that Q_{j+1} is a child of Q_j . Therefore, we can use the triangle inequality to write

$$\left| \int_{\alpha(Q)} h - \int_Q h \right|^2 \ell(Q)^n \lesssim_{k,n} \sum_{i=1}^j \left| \int_{Q_i} h - \int_{Q_{i-1}} h \right|^2 \ell(Q_{i-1})^n \leq \sum_{i=1}^j \|\Delta_{Q_i} h\|_2^2.$$

Because each cube $Q' \subseteq R$ can appear in at most $N(n, k) < \infty$ chains of the above type, this gives

$$\left| \int_{\alpha(Q)} h - \int_Q h \right|^2 \ell(Q)^n \lesssim_{n,k} \sum_{Q \subseteq R} \sum_{i=1}^{j(Q)} \|\Delta_{Q_i} h\|_2^2 \lesssim_{n,k} \sum_{Q \subseteq R} \|\Delta_R h\|_2^2 = \|h\|_2^2.$$

The reader should also see [DS93] Lemma IV.2.2.14 for a version of this statement where $\alpha(Q)$ is only required to be “ N -close” to Q rather than contained in Q . The main difference in Lemma 3.6 comes from averaging over normed balls rather than dyadic cubes.

Corollary 3.8 (cf. [DS93] Corollary IV.2.2.19). *Let $L, \epsilon, M > 0$ and let $h \in L^\infty(\mathbb{R}^n)$ with $\|h\|_\infty \leq M$. Let*

$$\mathcal{G}_h = \left\{ Q \in \mathcal{D}(\mathbb{R}^n) \mid \left| \int_B h - \int_Q h \right| \leq \epsilon \text{ for all } B \in \mathcal{B}_L(Q) \right\}.$$

Then $\mathcal{B}_h = \mathcal{D}(\mathbb{R}^n) \setminus \mathcal{G}_h$ is $C(M, L, n, \epsilon)$ -Carleson.

Proof. Let $\tilde{h} = h + \|h\|_\infty$ so that $0 \leq \tilde{h} \leq 2M$ a.e. Choose $k, \delta > 0$ such that the conclusion of Lemma 3.6 holds with our given $\epsilon, L, (2M)^2$. Let

$$\mathcal{B}_{\tilde{h}} = \left\{ Q \in \mathcal{D}(\mathbb{R}^n) \mid \exists B \in \mathcal{B}_L(Q), \left| \int_B \tilde{h} - \int_Q \tilde{h} \right| > \epsilon \right\}$$

and fix $R \in \mathcal{D}$. By Lemma 3.6, $Q \in \mathcal{B}_{\tilde{h}}$ implies $\Delta_k^{\tilde{h}}(Q) > \delta \ell(Q)^{n/2}$ so that by (3.2).

$$\sum_{\substack{Q \subseteq R \\ Q \in \mathcal{B}_{\tilde{h}}}} \ell(Q)^n \leq \sum_{\substack{Q \subseteq R \\ \Delta_k^{\tilde{h}}(Q) > \delta \ell(Q)^{n/2}}} \ell(Q)^n \lesssim_{\delta, k, n, M} \ell(R)^n.$$

The result follows since $\int_B \tilde{h} - \int_Q \tilde{h} = \int_B h - \int_Q h$ so that $\mathcal{B}_h = \mathcal{B}_{\tilde{h}}$. \blacksquare

4. BI-LIPSCHITZ IMAGES SATISFY THE WCD

In this section, we use the tools from Section 3 to prove that metric spaces which are bi-Lipschitz images of Euclidean spaces satisfy the WCD. In this section and the next, we will use the following version of the WCD adapted to Christ-David cubes using only \mathcal{H}^n .

Definition 4.1 (Cube weak constant density condition). Let (X, d) be an Ahlfors n -regular metric space, \mathcal{D} be a system of Christ-David cubes for X , and let $C_0, \epsilon_0 > 0$. Define

$$(4.1) \quad \mathcal{G}_{\text{cd}}(C_0, \epsilon_0) = \left\{ Q \in \mathcal{D} \mid \begin{array}{l} \exists a_Q > 0, \text{ such that } \forall y \in \frac{\epsilon_0}{2} B_Q, 0 < r \leq \frac{\epsilon_0}{2} \ell(Q), \\ |\mathcal{H}^n(B(y, r)) - a_Q r^n| \leq \epsilon_0 \ell(Q)^n \end{array} \right\},$$

$$(4.2) \quad \mathcal{B}_{\text{cd}}(C_0, \epsilon_0) = \mathcal{D} \setminus \mathcal{G}_{\text{cd}}(C_0, \epsilon_0).$$

We say that X satisfies the cube WCD if there exists $C_0 > 0$ such that for all choices of system \mathcal{D} and $\epsilon_0 > 0$, $\mathcal{B}_{\text{cd}}(C_0, \epsilon_0)$ is Carleson.

See Lemma 2.3 for a proof that this version of the WCD implies the version given in Definition 1.4. We will also need to review some of the theory of rectifiability in metric spaces.

Definition 4.2 (metric derivatives, jacobians). Let $f : \mathbb{R}^n \rightarrow \Sigma$ be L -Lipschitz. We say a seminorm on \mathbb{R}^n $|Df|(x)$ is a *metric derivative* of f at x if

$$\lim_{y, z \rightarrow x} \frac{d(f(y), f(z)) - |Df|(x)(y - z)}{|y - x| + |z - x|} = 0.$$

Given a seminorm s on \mathbb{R}^n , define $\mathcal{J}(s)$, the *jacobian* of s , by

$$\mathcal{J}(s) = \alpha(n)n \left(\int_{\mathbb{S}^{n-1}} (s(x))^{-n} d\mathcal{H}^{n-1}(x) \right)^{-1}.$$

Kirchheim used these ideas to prove the following metric analogs of Rademacher's theorem and the area formula for Lipschitz maps from \mathbb{R}^n into metric spaces.

Theorem 4.3 (cf. [Kir94] Theorem 2, Corollary 8). *Let $f : \mathbb{R}^n \rightarrow \Sigma$ be L -Lipschitz and let $\mathcal{J}_f(x) = \mathcal{J}(|Df|(x))$. A metric derivative for f exists at \mathcal{L}^n almost every $x \in \mathbb{R}^n$. In addition, for any Lebesgue integrable function $g : \mathbb{R}^n \rightarrow \mathbb{R}$,*

$$\int_{\mathbb{R}^n} g(x) \mathcal{J}_f(x) d\mathcal{L}^n(x) = \int_{\Sigma} \left(\sum_{x \in f^{-1}(y)} g(x) \right) d\mathcal{H}^n(y).$$

In their work on Lipschitz analogs of Sard’s Theorem, Azzam and Schul developed the following quantitative measure of how far a function f is from being given by a seminorm.

Definition 4.4. Let $f : \mathbb{R}^n \rightarrow X$ and $Q \in \mathcal{D}(\mathbb{R}^n)$. Define

$$\text{md}_f(Q) = \frac{1}{\ell(Q)} \inf_{\|\cdot\|} \sup_{x, y \in Q} \left| d(f(x), f(y)) - \|x - y\| \right|$$

Norms which are close to the infimum in the definition of $\text{md}_f(Q)$ can be thought of as “coarse” metric derivatives for f inside Q (note that they are biased towards approximating points whose distances are on the scale of Q). Azzam and Schul proved the following metric quantitative differentiation theorem involving these coefficients.

Theorem 4.5 ([AS14] Theorem 1.1). *Let $f : \mathbb{R}^n \rightarrow X$ be an L -Lipschitz function. Let $\delta > 0$. Then for each $R \in \mathcal{D}(\mathbb{R}^n)$,*

$$\sum_{\substack{Q \in \mathcal{D}(R) \\ \text{md}_f(3Q) > \delta L}} \ell(Q)^n \leq C(\delta, n) \ell(R)^n.$$

Finally, we will need to extend the standard system of dyadic cubes.

Definition 4.6 (one-third trick lattices). The following family of dyadic systems were introduced by Okikiolu [Oki92]. For any $e \in \{0, 1\}^n$ and cube $Q_0 \in \mathcal{D}(\mathbb{R}^n)$, define the shifted dyadic lattice

$$\begin{aligned} \mathcal{D}_j^e(Q_0) &= \left\{ Q + \frac{\ell(Q)}{3} e \mid Q \in \mathcal{D}_j(Q_0) \right\}, \\ \mathcal{D}^e(Q_0) &= \bigcup_{j \geq 0} \mathcal{D}_j^e(\mathbb{R}^n) \end{aligned}$$

and set

$$\tilde{\mathcal{D}}(Q_0) = \bigcup_{e \in \{0, 1\}^n} \mathcal{D}^e(Q_0).$$

$\tilde{\mathcal{D}}(Q_0)$ has the following property: For any $x \in Q_0$ and $j \geq 0$, there exists $Q \in \tilde{\mathcal{D}}(Q_0)$ such that $x \in \frac{2}{3}Q$ (See [Ler03] Proposition 3.2).

We now begin setting up the proof of the WCD for bi-Lipschitz images. We use the following good family of dyadic cubes from our collection of dyadic trees $\tilde{\mathcal{D}}$ to do analysis in the domain of our bi-Lipschitz maps.

Definition 4.7 (L -good I_Q). Let $f : [0, 1]^n \rightarrow \Sigma$ be L -bi-Lipschitz. Fix $Q \in \mathcal{D}(\Sigma)$. We call a dyadic cube $I_Q \in \tilde{\mathcal{D}}$ L -good for Q if the following hold:

- (i) $\ell(I_Q) \asymp_L \ell(Q)$,
- (ii) $3B_Q \subseteq f(I_Q)$,

where the implicit constant in 4.7(i) is independent of Q and I_Q .

Using the special property of the one-third trick lattices and the definition of bi-Lipschitz maps, the following lemma is standard.

Lemma 4.8. *Let $f : [0, 1]^n \rightarrow \Sigma$ be L -bi-Lipschitz. For each $Q \in \mathcal{D}(\Sigma)$ with $\ell(Q) \lesssim_L 1$, there exists an L -good $I_Q \in \tilde{\mathcal{D}}$.*

For $k \in \mathbb{N}$, $\delta > 0$, define

$$(4.3) \quad \mathcal{G}_\Sigma(k, \delta) = \left\{ Q \in \mathcal{D} \mid \begin{array}{l} \exists L\text{-good } I_Q \in \tilde{\mathcal{D}} \text{ with} \\ \Delta_k^{\mathcal{J}_f}(I_Q) \leq \delta \ell(I_Q)^{n/2}, \text{ md}_f(I_Q) \leq \delta \end{array} \right\},$$

$$(4.4) \quad \mathcal{B}_\Sigma(k, \delta) = \mathcal{D} \setminus \mathcal{G}_\Sigma(k, \delta).$$

The strategy of the proof is to first show that $\mathcal{G}_\Sigma \subseteq \mathcal{G}_{\text{cd}}$ and then show that \mathcal{B}_Σ is Carleson, and therefore \mathcal{B}_{cd} is Carleson since $\mathcal{B}_{\text{cd}} \subseteq \mathcal{B}_\Sigma$.

Lemma 4.9. *Let $f : [0, 1]^n \rightarrow \Sigma$ be L -bi-Lipschitz and let $\epsilon > 0$. There exist $k(\epsilon, L, n)$, $\delta(\epsilon, L, n) > 0$ such that the following holds: For any $Q \in \mathcal{G}_\Sigma(k, \delta)$ there exists a constant $c_Q \asymp_{L, n} 1$ such that for any normed ball $B \in \mathcal{B}_L(I_Q)$,*

$$|\mathcal{H}^n(f(B)) - c_Q \mathcal{L}^n(B)| \leq \epsilon \mathcal{L}^n(B)$$

Proof. Let $I_Q \in \tilde{\mathcal{D}}$ for Q be as in (4.3), let $\epsilon > 0$, and assume k, δ are small enough to satisfy the hypotheses of Lemma 3.6 with respect to $0 \leq \mathcal{J}_f \in L^\infty$ and ϵ . Then $\Delta_k^{\mathcal{J}_f}(I_Q) \leq \delta$ implies that

$$\left| \int_B \mathcal{J}_f - \int_Q \mathcal{J}_f \right| \leq \epsilon.$$

for any normed ball $B \in \mathcal{B}_L(I_Q)$. By setting $c_Q = \int_Q \mathcal{J}_f$ and using the area formula, we get the desired inequality. \blacksquare

Lemma 4.10. *Let $f : [0, 1]^n \rightarrow \Sigma$ be L -bi-Lipschitz, let C_0 be a regularity constant for Σ , and let $\epsilon_0 > 0$. There exist $k(\epsilon_0, L, n)$, $\delta(\epsilon_0, L, n) > 0$ such that $\mathcal{G}_\Sigma(k, \delta) \subseteq \mathcal{G}_{\text{cd}}(2C_0, \epsilon_0)$. In fact, for any $Q \in \mathcal{G}_\Sigma(k, \delta)$, there exists a constant $(2C_0)^{-1} \leq a_Q \leq 2C_0$ such that for any $y \in B_Q$, $0 < r \leq \ell(Q)$, we have*

$$(4.5) \quad |\mathcal{H}^n(B(y, r)) - a_Q r^n| \leq \epsilon_0 \ell(Q)^n.$$

That is, the condition on cubes in $\mathcal{G}_{\text{cd}}(2C_0, \epsilon_0)$ is attained with a multiple of \mathcal{H}^n .

Proof. First, we note that if a constant a_Q such as in (4.5) exists, it must satisfy $(2C_0)^{-1} \leq a_Q \leq 2C_0$ for small enough ϵ_0 because Σ is (C_0, n) -regular. Let I_Q be as in (4.3) and let $\epsilon > 0$. By Lemma 4.9, we can choose k large enough and $\delta > 0$ small enough so there exists $c_Q \asymp_{L, n} 1$ such that for any $B \in \mathcal{B}_{2L}(I_Q)$

$$(4.6) \quad |\mathcal{H}^n(f(B)) - c_Q \mathcal{L}^n(B)| \leq \epsilon \mathcal{L}^n(B).$$

In addition, the fact that $\text{md}_f(I_Q) \leq \delta$ implies that there exists a norm $\|\cdot\|_Q$ such that

$$(4.7) \quad \sup_{x, y \in I_Q} |d(f(x), f(y)) - \|x - y\|_Q| \leq \delta \ell(I_Q).$$

Let $c_{\|\cdot\|_Q} \asymp_{L, n} 1$ be such that $\mathcal{L}^n(B_{\|\cdot\|_Q}(0, r)) = c_{\|\cdot\|_Q} r^n$. We set

$$a_Q = c_Q c_{\|\cdot\|_Q}$$

and begin the proof of (4.5).

Let $y_0 = f^{-1}(y)$. We claim that there exists a constant $c_1(n, L) > 0$ such that

$$(4.8) \quad B_1 := B_{\|\cdot\|_Q}(y_0, (1 - c_1\delta)r) \subseteq f^{-1}(B(y, r)) \subseteq B_{\|\cdot\|_Q}(y_0, (1 + c_1\delta)r) =: B_2.$$

For the first inclusion, let $x_0 \in B_1$. By (4.7),

$$d(f(x_0), f(y_0)) \leq \|x_0 - y_0\|_Q + 3\delta\ell(I_Q) \leq (1 - c_1\delta)r + C(L, n)\delta r < r$$

where the final inequality holds if c_1 is large enough. Similarly, let $z_0 \in f^{-1}(B(y, r)) \subseteq I_Q$. Then

$$\|z_0 - y_0\|_Q \leq d(f(z_0), f(y_0)) + \delta\ell(3I_Q) \leq r + C(L, n)\delta r \leq (1 + c_1\delta)r$$

with the same restriction on δ as above. This finishes the proof of (4.8). Because $3B_Q \subseteq f(I_Q)$, we immediately have that $B_1, B_2 \subseteq I_Q$ for small enough δ so that $B_1, B_2 \in \mathcal{B}_{2L}(I_Q)$. Using (4.6), this implies the existence of a constant $c_2(n, L)$ so that

$$\mathcal{H}^n(B(y, r)) \leq \mathcal{H}^n(f(B_2)) \leq (c_Q + \epsilon)c_{\|\cdot\|_Q}(1 + c_1\delta)^n r^n \leq a_Q r^n + c_2(\epsilon + \delta)\ell(Q)^n.$$

A similar computation using $\mathcal{H}^n(f(B_1))$ gives a similar lower bound for $\mathcal{H}^n(B(y, r))$. This shows that

$$|\mathcal{H}^n_\Sigma(B(y, r)) - a_Q r^n| \leq c_2(\epsilon + \delta)\ell(Q)^n$$

By choosing ϵ small enough, then k large enough and δ small enough, we get the conclusion of the lemma. \blacksquare

Lemma 4.11. *Let $f : [0, 1]^n \rightarrow \Sigma$ be L -bi-Lipschitz. For any $k, \delta > 0$, $\mathcal{B}_\Sigma(k, \delta)$ is $C(k, \delta, n, L)$ -Carleson.*

Proof. Let $R \in \mathcal{D}$. By Lemma 4.8, Remark 3.3, and Theorem 4.5 we have

$$\begin{aligned} \sum_{\substack{Q \in \mathcal{B}(k, \delta) \\ Q \subseteq R}} \ell(Q)^n &\lesssim \sum_{\substack{Q \subseteq R \\ \Delta_k^{\mathcal{J}^f}(I_Q) > \delta\ell(I_Q)^{n/2}}} \ell(Q)^n + \sum_{\substack{Q \subseteq R \\ \text{md}_f(I_Q) > \delta}} \ell(Q)^n + \sum_{\substack{Q \subseteq R \\ \ell(Q) > C(L)}} \ell(Q)^n \\ &\lesssim_{L, n} \sum_{\substack{I_Q \subseteq I_R \\ \Delta_k^{\mathcal{J}^f}(I_Q) > \delta\ell(I_Q)^{n/2}}} \ell(I_Q)^n + \sum_{\substack{I_Q \subseteq I_R \\ \text{md}_f(I_Q) > \delta}} \ell(I_Q)^n + C(L, n)\ell(R)^n \\ &\lesssim_{k, \delta, L, n} \ell(I_R)^n + \ell(R)^n \lesssim_{L, n} \ell(R)^n. \end{aligned} \quad \blacksquare$$

Theorem 4.12. *The WCD holds for any bi-Lipschitz image of $[0, 1]^n$.*

Proof. Let f be L -bi-Lipschitz $f : [0, 1]^n \rightarrow \Sigma$. Choose $C_0(L, n)$ such that Σ is Ahlfors (C_0, n) -regular and let $\epsilon_0 > 0$. Choose k large enough and δ small enough with respect to ϵ_0, L, n so that $\mathcal{G}_\Sigma(k, \delta) \subseteq \mathcal{G}_{\text{cd}}(2C_0, \epsilon_0)$. That is, the conclusion of Lemma 4.10 holds. Then $\mathcal{B}_{\text{cd}}(\epsilon_0, 2C_0) \subseteq \mathcal{B}_\Sigma(k, \delta)$. Lemma 4.11 implies that $\mathcal{B}_\Sigma(k, \delta)$ is $C(\epsilon_0, L, n)$ -Carleson, implying $\mathcal{B}_{\text{cd}}(2C_0, \epsilon_0)$ is also $C(\epsilon_0, L, n)$ -Carleson which says exactly that Σ satisfies the WCD. \blacksquare

5. UNIFORMLY RECTIFIABLE METRIC SPACES SATISFY THE WCD

The goal of this section is to prove Theorem A: uniformly n -rectifiable metric spaces satisfy the WCD. We will prove this via approximating by bi-Lipschitz images. That is, we will use the fact that uniformly rectifiable metric spaces have very big pieces of bi-Lipschitz images to transfer our bi-Lipschitz image estimates to the uniformly rectifiable case. The primary tool for this argument is the following abstract analog of the John-Nirenberg-Stromberg theorem.

Lemma 5.1 ([BHS23] Lemma 4.2.8, [DS93] Lemma IV.1.12). *Let X be an Ahlfors n -regular metric space and \mathcal{D} a system of Christ-David cubes for X . Let $\alpha : \mathcal{D} \rightarrow [0, \infty)$ be given and suppose there are $N, \eta > 0$ such that*

$$(5.1) \quad \mathcal{H}^n \left(\left\{ x \in R \mid \sum_{\substack{Q \subseteq R \\ x \in Q}} \alpha(Q) \leq N \right\} \right) \geq \eta \ell(R)^n$$

for all $R \in \mathcal{D}$. Then,

$$\sum_{Q \subseteq R} \alpha(Q) \ell(Q)^n \lesssim_{N, \eta} \ell(R)^n$$

for all $R \in \mathcal{D}$.

For our application, we will take $\alpha(Q) = \chi_{\mathcal{B}_{\text{cd}}(C_0, \epsilon_0)}(Q)$ where $\mathcal{B}_{\text{cd}}(C_0, \epsilon_0)$ is as in (4.2). If we can show that (5.1) holds for this choice of α , then we will conclude

$$(5.2) \quad \sum_{Q \subseteq R} \chi_{\mathcal{B}_{\text{cd}}(C_0, \epsilon_0)}(Q) \ell(Q)^n = \sum_{\substack{Q \subseteq R \\ Q \in \mathcal{B}_{\text{cd}}(C_0, \epsilon_0)}} \ell(Q)^n \lesssim_{N, \eta} \ell(R)^n$$

which is exactly the desired Carleson packing condition for $\mathcal{B}_{\text{cd}}(C_0, \epsilon_0)$. We will need the following result from Bate, Hyde, and Schul's paper which states that uniformly rectifiable metric spaces have very big pieces of bi-Lipschitz images.

Theorem 5.2 (cf. [BHS23] Theorem B, Proposition 9.0.2). *Let $\epsilon > 0$ and let X be uniformly n -rectifiable. There is an $L \geq 1$ depending only on ϵ, n , the Ahlfors regularity constant for X , and the BPLI constants for X such that for each $x \in X$ and $r > 0$ there exists $F \subseteq B(x, r)$, satisfying $\mathcal{H}_X^n(B(x, r) \setminus F) \leq \epsilon r^d$ and an L -bi-Lipschitz map $g : F \rightarrow \mathbb{R}^n$.*

Remark 5.3. If we embed X isometrically into ℓ_∞ , then we can take the map $g^{-1} : g(F) \rightarrow F \subseteq \ell_\infty$ above and extend it to an $L'(L, n)$ -bi-Lipschitz map $f : \mathbb{R}^n \rightarrow \ell_\infty$ satisfying the same conclusions with respect to the isometric embedding of X . (See [BHS23] Lemma 4.3.2 for a proof.)

We now begin setting up the proof of Theorem A. Fix a uniformly n -rectifiable metric space X with regularity constant C_0 and a system of Christ-David cubes \mathcal{D} for X . Let $\epsilon_0 > 0$ and $R \in \mathcal{D}(X)$. By applying Theorem 5.2 to the ball $3B_R$, we get an L -bi-Lipschitz map $f : \mathbb{R}^n \rightarrow \Sigma \subseteq \ell_\infty$ such that

$$(5.3) \quad \mathcal{H}_X^n(3B_R \setminus \Sigma) \leq \frac{\epsilon_0}{4} \ell(R)^n.$$

We will only need to use f near where it parameterizes $3B_R$, so it suffices to consider $f|_{I_R}$ where I_R is L -good for R (See Definition 4.7). We can assume without loss of generality that $I_R = [0, 1]^n$ so that the results of the previous section for bi-Lipschitz images of $[0, 1]^n$ apply to f .

Because Σ has such large intersection with $3B_R$, we can use the following lemma to find a substantial subset $\tilde{R} \subseteq R$ such that for every $x \in \tilde{R}$, every cube $Q \subseteq R$ with $x \in Q$ has very large intersection with Σ .

Lemma 5.4. *Let X be a doubling metric space with a system of Christ-David cubes \mathcal{D} . Let $\epsilon > 0$, $F \subseteq X$ measurable, and let $R \in \mathcal{D}$ be such that $\mathcal{H}^n(R \setminus F) \leq$*

$\epsilon \mathcal{H}^n(R)$. Define

$$(5.4) \quad \tilde{R} = \left\{ x \in R \mid \begin{array}{l} \text{For all } Q \in \mathcal{D} \text{ such that } x \in Q \subseteq R, \\ \mathcal{H}^n(Q \cap F) \geq (1 - 2\epsilon) \mathcal{H}^n(Q) \end{array} \right\}.$$

We have $\mathcal{H}^n(\tilde{R}) \geq \epsilon \mathcal{H}^n(R)$.

Proof. This proof is essentially contained in the proof of Lemma IV.2.2.38 in [DS93], but we need to be precise about the constant ϵ . If $x \in R \setminus \tilde{R}$, then x is contained in some cube Q such that $\mathcal{H}^n(Q \cap F) \leq (1 - 2\epsilon) \mathcal{H}^n(Q)$. Let $\{Q_i\}_i$ be a maximal disjoint family of such cubes so that $R \setminus \tilde{R} = \bigcup_i Q_i$. Then

$$(5.5) \quad \begin{aligned} \mathcal{H}^n((R \setminus \tilde{R}) \cap F) &= \sum_i \mathcal{H}^n(Q_i \cap F) \leq (1 - 2\epsilon) \sum_i \mathcal{H}^n(Q_i) \\ &\leq (1 - 2\epsilon) \mathcal{H}^n(R \setminus \tilde{R}) \leq (1 - 2\epsilon) \mathcal{H}^n(R). \end{aligned}$$

On the other hand,

$$(5.6) \quad \begin{aligned} \mathcal{H}^n((R \setminus \tilde{R}) \cap F) &= \mathcal{H}^n((R \cap F) \setminus \tilde{R}) \geq \mathcal{H}^n(R \cap F) - \mathcal{H}^n(\tilde{R}) \\ &\geq (1 - \epsilon) \mathcal{H}^n(R) - \mathcal{H}^n(\tilde{R}). \end{aligned}$$

Combining (5.5) and (5.6) and rearranging gives

$$\mathcal{H}^n(\tilde{R}) \geq (1 - \epsilon) \mathcal{H}^n(R) - (1 - 2\epsilon) \mathcal{H}^n(R) = \epsilon \mathcal{H}^n(R). \quad \blacksquare$$

While this lemma allows us to control the measure of the part of X outside of Σ , we will also use separate control of the maximal distance of points in $Q \in \mathcal{D}(R)$ from Σ as measured by the following quantity.

Definition 5.5. Let (Z, d) be a metric space and suppose $X, Y \subseteq Z$. For $x \in X$ and $0 < r < \text{diam}(X)$, define

$$I_{X,Y}(x, r) = \frac{1}{r} \sup_{\substack{y \in X \cap B(x, r) \\ \text{dist}(y, Y) \leq r}} \text{dist}(y, Y)$$

The following lemma gives Carleson control over cubes where $I_{X,Y}$ is large.

Lemma 5.6 ([BHS23] Lemma 4.2.6). *Let (Z, d) be a metric space with $X, Y \subseteq Z$ Ahlfors (C_0, n) -regular subsets and \mathcal{D} a system of Christ-David cubes for X . For any $\delta > 0$, the set $\{Q \in \mathcal{D} \mid I_{X,Y}(3B_Q) > \delta\}$ is $C(C_0, n, \delta)$ -Carleson.*

We can now define the good family of descendants of R we want to consider. Let $E = f^{-1}(X)$. For any $k \in \mathbb{N}$ and $\delta > 0$, consider the following three conditions applicable to $Q \in \mathcal{D}(R) \subseteq \mathcal{D}(X)$:

- (i) $\mathcal{H}^n(Q \setminus \Sigma) \leq \frac{\epsilon_0}{2} \mathcal{H}^n(Q)$,
- (ii) $I_{X,\Sigma}(3B_Q) \leq \delta$,
- (iii) $\exists L$ -good $I_Q \in \tilde{\mathcal{D}}(\mathbb{R}^n)$ for which the following hold:
 - (a) $\Delta_k^{\mathcal{J}^{f \cdot X^E}}(I_Q) \leq \delta \ell(I_Q)^{n/2}$,
 - (b) $\text{md}_f(I_Q) \leq \delta$.

Define

$$\begin{aligned} \mathcal{G}_R(k, \delta) &= \{ Q \in \mathcal{D}(R) \mid Q \text{ satisfies (i), (ii), and (iii)} \}, \\ \mathcal{B}_R(k, \delta) &= \mathcal{D} \setminus \mathcal{G}_R(k, \delta). \end{aligned}$$

We first show that $\mathcal{G}_R(k, \delta)$ cubes are good for the WCD for X . The reader should compare the following lemma with Lemma 4.10.

Lemma 5.7. *Let X be uniformly n -rectifiable with regularity constant C_0 , let $\epsilon_0 > 0$, and let $R \in \mathcal{D}(X)$. Let $f : \mathbb{R}^n \rightarrow \Sigma$ be L -bi-Lipschitz and satisfy (5.3). There exist $k, \delta > 0$ dependent on C_0, ϵ_0, n, L such that $\mathcal{G}_R(k, \delta) \subseteq \mathcal{G}_{cd}(2C_0, \epsilon_0)$.*

Proof. Let $Q \in \mathcal{G}_R(k, \delta)$, $y \in \frac{\epsilon_0}{2}B_Q$, $0 < r \leq \frac{\epsilon_0}{2}\ell(Q)$ and let I_Q be the cube guaranteed from condition (iii). Notice that $B(y, r) \subseteq c_0B_Q$. By condition (ii), there exists $y_0 \in \Sigma$ satisfying $|y - y_0| \leq 3\delta\ell(Q)$. Let $r_{\pm} = r \pm 3\delta\ell(Q)$ so that

$$B_{\Sigma}(y_0, r_-) \cap X \subseteq B_X(y, r) \cap \Sigma \subseteq B_{\Sigma}(y_0, r_+) \cap X.$$

Since $\text{md}_f(I_Q) \leq \delta$, the proof of Lemma 4.10, specifically of (3.4), shows that there exists a norm $\|\cdot\|_Q$ and $c_1(n, L) > 0$ such that the balls

$$\begin{aligned} B_1 &= B_{\|\cdot\|_Q}(f^{-1}(y_0), (1 - c_1\delta)r_-), \\ B_2 &= B_{\|\cdot\|_Q}(f^{-1}(y_0), (1 + c_1\delta)r_+) \end{aligned}$$

satisfy $B_1, B_2 \in \mathcal{B}_{2L}(I_Q)$ and

$$(5.7) \quad B_1 \cap E \subseteq f^{-1}(B_X(y, r) \cap \Sigma) \subseteq B_2 \cap E.$$

Let $\epsilon > 0$. By taking δ small enough and k large enough so that the hypotheses of Lemma 3.6 are satisfied for $\mathcal{J}_{f\chi_E}$, the fact that $\Delta_k^{\mathcal{J}_{f\chi_E}}(I_Q) \leq \delta\ell(I_Q)^{n/2}$ gives

$$\left| \int_B \mathcal{J}_{f\chi_E} - \int_Q \mathcal{J}_{f\chi_E} \right| \leq \epsilon$$

for any normed ball $B \in \mathcal{B}_{2L}(I_Q)$. Set $c_Q = \int_Q \mathcal{J}_{f\chi_E}$. After rearranging and applying the area formula, this becomes

$$(5.8) \quad |\mathcal{H}^n(f(B \cap E)) - c_Q \mathcal{L}^n(B)| \leq \epsilon \mathcal{L}^n(B).$$

Let $\mathcal{L}^n(B_{\|\cdot\|_Q}(0, r)) = c_{\|\cdot\|_Q} r^n$ and set $a_Q = c_Q c_{\|\cdot\|_Q}$. Combining (5.7) and (5.8) gives

$$\begin{aligned} \mathcal{H}^n(B_X(y, r) \cap \Sigma) &\leq \mathcal{H}^n(f(B_2 \cap E)) \leq (c_Q + \epsilon) \mathcal{L}^n(B_2) = a_Q(1 + c_Q^{-1}\epsilon)(1 - c_1\delta)^n r_+^n \\ &= a_Q(1 + c_Q^{-1}\epsilon)(1 - c_1\delta)^n (r + 3\delta\ell(Q))^n \\ &\leq a_Q r^n + C(n, L)(\epsilon + \delta)\ell(Q)^n. \end{aligned}$$

A similar argument using B_1 gives a similar lower bound so that

$$|\mathcal{H}^n(B_X(y, r) \cap \Sigma) - a_Q r^n| \leq C(n, L)(\epsilon + \delta)\ell(Q)^n.$$

Finally, using (i), we have

$$\begin{aligned} |\mathcal{H}^n(B_X(y, r)) - a_Q r^n| &\leq \mathcal{H}^n(B_X(y, r) \setminus \Sigma) + |\mathcal{H}^n(B_X(y, r) \cap \Sigma) - a_Q r^n| \\ &\leq \mathcal{H}^n(c_0B_Q \cap X \setminus \Sigma) + C(n, L)(\epsilon + \delta)\ell(Q)^n \\ &\leq \frac{\epsilon_0}{2}\ell(Q)^n + C(n, L)(\epsilon + \delta)\ell(Q)^n \leq \epsilon_0\ell(Q)^n \end{aligned}$$

where the final inequality follows by first fixing ϵ sufficiently small in terms of ϵ_0, L, n then δ small and k large in terms of $\epsilon, \epsilon_0, n, L$. \blacksquare

We now show that $\mathcal{B}_R(k, \delta)$ is not too big. The reader should compare this lemma with Lemma 4.11.

Lemma 5.8. *Let X be uniformly n -rectifiable with regularity constant C_0 , let $\epsilon_0, \delta, k > 0$, and let $R \in \mathcal{D}(X)$. Let $f : \mathbb{R}^n \rightarrow \Sigma$ be L -bi-Lipschitz and satisfy (5.3). There exist constants $N, \eta > 0$ dependent on $k, \delta, \epsilon_0, n, L$ such that*

$$(5.9) \quad \mathcal{H}^n \left(\left\{ x \in R \mid \sum_{\substack{Q \subseteq R \\ x \in Q}} \chi_{\mathcal{B}_R(k, \delta)} \leq N \right\} \right) \geq \eta \ell(R)^n.$$

Proof. Define

$$\begin{aligned} \mathcal{B}_1 &= \{ Q \subseteq R \mid I_{X, \Sigma}(3B_Q) > \delta \}, \\ \mathcal{B}_2 &= \{ Q \subseteq R \mid \text{there is no } I_Q \text{ satisfying (iii)} \}. \end{aligned}$$

Lemma 5.6 shows that \mathcal{B}_1 is $C(\delta, C_0, n)$ -Carleson and Lemma 4.11 shows that \mathcal{B}_2 is $C(k, \delta, n, L)$ -Carleson. Let \tilde{R} be as in (5.4). By Chebyshev's inequality,

$$\begin{aligned} \mathcal{H}^n \left(\left\{ x \in \tilde{R} \mid \sum_{\substack{Q \subseteq R \\ x \in Q}} \chi_{\mathcal{B}_R(k, \delta)}(Q) > N \right\} \right) &\leq \frac{1}{N} \int_{\tilde{R}} \sum_{\substack{Q \subseteq R \\ x \in Q}} \chi_{\mathcal{B}_R(k, \delta)}(Q) \\ &\lesssim \frac{1}{N} \int_{\tilde{R}} \sum_{\substack{Q \subseteq R \\ x \in Q}} \chi_{\mathcal{B}_1}(Q) + \chi_{\mathcal{B}_2}(Q) \\ &\leq \frac{1}{N} \sum_{\substack{Q \subseteq R \\ Q \in \mathcal{B}_1 \cup \mathcal{B}_2}} \ell(Q)^n \lesssim \frac{1}{N} \ell(R)^n. \end{aligned}$$

The result follows by taking N sufficiently large since the left hand side of (5.9) is bounded below by

$$\mathcal{H}^n(\tilde{R}) - \frac{C}{N} \ell(R)^n \geq \epsilon \mathcal{H}^n(R) - \frac{C}{N} \ell(R)^n \gtrsim_\epsilon \ell(R)^n. \quad \blacksquare$$

We finally observe that these pieces combine to prove Theorem A:

Proof of Theorem A. Choose $R \in \mathcal{D}(X)$ and apply Theorem 5.2 to get an L -bi-Lipschitz map $f : \mathbb{R}^n \rightarrow \Sigma \subseteq \ell_\infty$ satisfying (5.3). Fix k large enough and δ small enough in terms of C_0, ϵ_0, n, L so that $\mathcal{G}_R(k, \delta) \subseteq \mathcal{G}_{\text{cd}}(2C_0, \epsilon_0)$. That is, the conclusion of Lemma 5.7 holds. Recall the definition of \tilde{R} from (5.4) and define

$$\tilde{\mathcal{D}}(R) = \{ Q \in \mathcal{D}(R) : \exists x \in \tilde{R} \cap Q \}.$$

We have $\mathcal{B}_{\text{cd}}(2C_0, \epsilon_0) \cap \tilde{\mathcal{D}}(R) \subseteq \mathcal{B}_R(k, \delta)$ so that $\chi_{\mathcal{B}_{\text{cd}}(2C_0, \epsilon_0)}(Q) \leq \chi_{\mathcal{B}_R(k, \delta)}(Q)$ for all $Q \in \tilde{\mathcal{D}}(R)$. Lemma 5.8 gives the existence of N, η independent of R so that

$$\mathcal{H}^n \left(\left\{ x \in R \mid \sum_{\substack{Q \subseteq R \\ x \in Q}} \chi_{\mathcal{B}_{\text{cd}}(2C_0, \epsilon_0)} \leq N \right\} \right) \geq \eta \ell(R)^n.$$

By Lemma 5.1 this implies $\mathcal{B}_{\text{cd}}(2C_0, \epsilon_0)$ is Carleson, implying X satisfies the WCD. \blacksquare

REFERENCES

- [AH22] Jonas Azzam and Matthew Hyde. The weak lower density condition and uniform rectifiability. *Ann. Fenn. Math.*, 47(2):791–819, 2022.
- [AS14] Jonas Azzam and Raanan Schul. A quantitative metric differentiation theorem. *Proc. Amer. Math. Soc.*, 142(4):1351–1357, 2014.
- [Bat23] David Bate. On 1-regular and 1-uniform metric measure spaces. Youtube, 2023. <https://www.youtube.com/watch?v=9zXjfKeWGv4&t=2096s>.
- [Bes28] A.S. Besicovitch. On the fundamental geometrical properties of linearly measurable plane sets of points. *Mathematische Annalen*, 98:422–464, 1928.
- [Bes38] A.S. Besicovitch. On the fundamental geometrical properties of linearly measurable plane sets of points (ii). *Mathematische Annalen*, 115:296–329, 1938.
- [BHS23] David Bate, Matthew Hyde, and Raanan Schul. Uniformly rectifiable metric spaces: Lipschitz images, bi-lateral weak geometric lemma and corona decompositions. arXiv preprint, arXiv:2306.12933, 2023.
- [CGLT16] Vasileios Chousionis, John Garnett, Triet Le, and Xavier Tolsa. Square functions and uniform rectifiability. *Trans. Amer. Math. Soc.*, 368(9):6063–6102, 2016.
- [Chr90] M. Christ. A $T(b)$ theorem with remarks on analytic capacity and the Cauchy integral. *Colloq. Math.*, 60/61(2):601–628, 1990.
- [CMT20] Vasilis Chousionis, Valentino Magnani, and Jeremy T. Tyson. On uniform measures in the Heisenberg group. *Adv. Math.*, 363:106980, 42, 2020.
- [Dav88] Guy David. Morceaux de graphes lipschitziens et intégrales singulières sur une surface. *Rev. Mat. Iberoamericana*, 4(1):73–114, 1988.
- [DL08] Camillo De Lellis. *Rectifiable sets, densities and tangent measures*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008.
- [DS91] Guy David and Stephen Semmes. Singular integrals and rectifiable sets in \mathbf{R}^n : Beyond Lipschitz graphs. *Astérisque*, (193):152, 1991.
- [DS93] Guy David and Stephen Semmes. *Analysis of and on uniformly rectifiable sets*, volume 38 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1993.
- [FV23] Katrin Fassler and Ivan Yuri Violo. On various carleson-type geometric lemmas and uniform rectifiability in metric spaces. arXiv preprint, arXiv:2310.10519, 2023.
- [Hah05] Immo Hahlomaa. Menger curvature and Lipschitz parametrizations in metric spaces. *Fund. Math.*, 185(2):143–169, 2005.
- [HM12] T. Hytönen and H. Martikainen. Non-homogeneous Tb theorem and random dyadic cubes on metric measure spaces. *J. Geom. Anal.*, 22(4):1071–1107, 2012.
- [Kir94] Bernd Kirchheim. Rectifiable metric spaces: local structure and regularity of the Hausdorff measure. *Proc. Amer. Math. Soc.*, 121(1):113–123, 1994.
- [KP87] Oldřich Kowalski and David Preiss. Besicovitch-type properties of measures and submanifolds. *J. Reine Angew. Math.*, 379:115–151, 1987.
- [KP02] Bernd Kirchheim and David Preiss. Uniformly distributed measures in Euclidean spaces. *Math. Scand.*, 90(1):152–160, 2002.
- [Ler03] Gilad Lerman. Quantifying curvelike structures of measures by using l_2 Jones quantities. *Comm. Pure Appl. Math.*, 56(9):1294–1365, 2003.
- [Lor03] Andrew Lorent. Rectifiability of measures with locally uniform cube density. *Proc. London Math. Soc. (3)*, 86(1):153–249, 2003.
- [Mar61] J. M. Marstrand. Hausdorff two-dimensional measure in 3-space. *Proc. London Math. Soc. (3)*, 11:91–108, 1961.
- [Mat75] Pertti Mattila. Hausdorff m regular and rectifiable sets in n -space. *Trans. Amer. Math. Soc.*, 205:263–274, 1975.
- [Mer22] Andrea Merlo. Geometry of 1-codimensional measures in Heisenberg groups. *Invent. Math.*, 227(1):27–148, 2022.
- [Nim17] A. Dali Nimer. A sharp bound on the Hausdorff dimension of the singular set of a uniform measure. *Calc. Var. Partial Differential Equations*, 56(4):Paper No. 111, 31, 2017.
- [Nim19] A. Dali Nimer. Uniformly distributed measures have big pieces of Lipschitz graphs locally. *Ann. Acad. Sci. Fenn. Math.*, 44(1):389–405, 2019.

- [Nim22] A. Dali Nimer. Conical 3-uniform measures: a family of new examples and characterizations. *J. Differential Geom.*, 121(1):57–99, 2022.
- [Oki92] Kate Okikiolu. Characterization of subsets of rectifiable curves in \mathbf{R}^n . *Journal of the London Mathematical Society*, s2-46(2):336–348, 1992.
- [Pre87] David Preiss. Geometry of measures in \mathbf{R}^n : distribution, rectifiability, and densities. *Ann. of Math. (2)*, 125(3):537–643, 1987.
- [PT92] David Preiss and Jaroslav Tišer. On Besicovitch’s $\frac{1}{2}$ -problem. *J. London Math. Soc. (2)*, 45(2):279–287, 1992.
- [Sch07] Raanan Schul. Ahlfors-regular curves in metric spaces. *Ann. Acad. Sci. Fenn. Math.*, 32(2):437–460, 2007.
- [Sch09] Raanan Schul. Bi-Lipschitz decomposition of Lipschitz functions into a metric space. *Rev. Mat. Iberoam.*, 25(2):521–531, 2009.
- [Tol12] Xavier Tolsa. Mass transport and uniform rectifiability. *Geom. Funct. Anal.*, 22(2):478–527, 2012.
- [Tol15] Xavier Tolsa. Uniform measures and uniform rectifiability. *J. Lond. Math. Soc. (2)*, 92(1):1–18, 2015.
- [TT15] Xavier Tolsa and Tatiana Toro. Rectifiability via a square function and Preiss’ theorem. *Int. Math. Res. Not. IMRN*, (13):4638–4662, 2015.

DEPARTMENT OF MATHEMATICS, STONY BROOK UNIVERSITY, STONY BROOK, NY 11794-3651

E-mail address: `jared.krandel@stonybrook.edu`

Jared Krandel was partially supported by the National Science Foundation under Grants No. DMS-1763973.