

Blender-producing mechanisms and a dichotomy for local dynamics for heterodimensional cycles

Dongchen Li

Dedicated to Professor Dmitry Turaev on his 60th birthday

Abstract. Blenders are special hyperbolic sets used to produce various robust dynamical phenomena which appear fragile at first glance. We prove for C^r diffeomorphisms ($r = 2, \dots, \infty, \omega$) that blenders naturally exist (without perturbation) near non-degenerate heterodimensional cycles of coindex-1, and the existence is determined by arithmetic properties of moduli of topological conjugacy for diffeomorphisms with heterodimensional cycles. In particular, we obtain a C^r -generic dichotomy for dynamics in any small neighborhood U of a non-degenerate heterodimensional cycle: either there exist infinitely many blenders accumulating on the cycle, forming robust heterodimensional dynamics in most cases, or there are no orbits other than those constituting the cycle lying entirely in U .

Keywords. blender, heterodimensional cycle, nonhyperbolic dynamics, chaotic dynamics.

AMS subject classification. 37C05, 37C29, 37C75, 37D25.

Contents

1	Introduction	2
1.1	Blenders and moduli of topological conjugacy	3
1.2	Simple hyperbolic dynamics	6
1.3	Robust heterodimensional dynamics arisen from perturbations	7
1.4	A dichotomy for local dynamics near heterodimensional cycles	9
1.5	Plan of the proofs	9
2	Results for different types of heterodimensional cycles	10
3	First return maps	13
3.1	Non-degeneracy conditions	13
3.2	Saddle-focus case	16
3.3	Double-focus case	17
4	Blenders near saddle-focus heterodimensional cycles	19
4.1	A criterion for the existence of blenders	19
4.2	Existence of blenders when at least one of θ and $\frac{\omega}{2\pi}$ is irrational	21
4.3	Homoclinic relations in the saddle-focus case	24
4.4	Proofs of Theorem 1 and the first case in Theorem 3	25
5	Blenders near double-focus heterodimensional cycles	26
5.1	Existence of blenders when at least one of $\theta, \frac{\omega_1}{2\pi}, \frac{\omega_2}{2\pi}$ is irrational	26

5.2	Homoclinic relations in the double-focus case	27
5.3	Proofs of Theorem 2 and the second case in Theorem 3	28
5.4	Proof of Proposition 5.1	29
6	Robust heterodimensional dynamics in generic one-parameter unfolding families.	
	Proof of Theorem 4	35
6.1	Unfolding saddle-focus heterodimensional cycles	35
6.2	Unfolding double-focus heterodimensional cycles	37
7	Purely rational moduli. Proof of Theorem B	38
7.1	Simple hyperbolic dynamics in the saddle-focus case	38
7.2	Simple hyperbolic dynamics in the double-focus case	42
	Acknowledgments	42
	References	43

1 Introduction

A blender is a hyperbolic set such that its projection to a certain “central” subspace has a non-empty interior, and this property persists for any C^1 -close system. An immediate consequence is that non-transverse intersections with the stable or unstable set can be unremovable by C^1 -small perturbations. This opens the way for constructing various C^1 -robust dynamical phenomena, including those which appear fragile at first glance. As the first such example, a large class of diffeomorphisms exhibiting the robust nonhyperbolic transitivity was constructed by Bonatti and Díaz in their discovery paper [11] of blenders. Subsequently, various non-trivial results involving blenders have been obtained: the persistence of heterodimensional cycles [13, 15, 23], abundance of C^1 -robust homoclinic tangencies [4, 14, 22], typicality of Newhouse phenomenon in the space of families of diffeomorphisms [7], robustly fast growth of number of periodic orbits [1, 2, 8], robust existence of nonhyperbolic ergodic measures [10], C^1 density of stable ergodicity [3, 30], robust transitivity in Hamiltonian dynamics [27], persistence of homoclinics to saddle-center periodic points [24], and robust bifurcations in complex dynamics [9, 20, 34], among others.

Essentially, there are two strategies for obtaining a blender. One can either explicitly construct blenders from (relatively) specific systems, see e.g. [7, 8, 27], or identify conditions under which blenders emerge near particular dynamical objects. In the latter case, blenders are mostly found near heterodimensional cycles, except that, in [3, 26], blenders arise from horseshoes with sufficiently large fractal dimension, while in [24] blenders are built near one-dimensional whiskered tori with homoclinics. In this paper, we investigate the creation of blenders near heterodimensional cycles. The first major contribution in this direction was made by Díaz in [17] where he implicitly constructed a blender. Later, Bonatti and Díaz developed [13] a general theory in the C^1 topology, establishing that blenders can appear after a C^1 -small perturbation which unfolds a heterodimensional cycle of coindex 1. A conservative version of this result was also obtained in [29]. More recently, together with Turaev, the author showed in [23] that, for every C^r diffeomorphisms ($r = 2, \dots, \infty, \omega$) having a coindex-1 heterodimensional cycle, there exist C^r -small perturbations which give rise to blenders. The method employed there is different from [13] and, in particular, we identified certain classes of heterodimensional cycles which give rise to blenders without being destroyed.

The present paper serves as a continuation of the blender-related investigation initiated in [23].

The main goal is to provide a comprehensive description (Theorem A) of the relation between the natural (i.e., non-perturbative) existence of blenders and arithmetic properties of moduli of topological conjugacy of the C^r diffeomorphisms¹ having heterodimensional cycles of coindex 1. The result characterizes the blender-producing mechanisms for coindex-1 heterodimensional dynamics in full generality. As a consequential result, we establish a dichotomy (Corollary A and Theorem B) for the local dynamics near non-degenerate heterodimensional cycles, depending on the existence/non-existence of rational dependencies between the moduli: one has either rich dynamics, that is, the cycle is a limit of infinitely many blenders, which in most cases form robust heterodimensional dynamics; or simple hyperbolic dynamics, that is, there are only finitely many locally maximal and compact hyperbolic sets in any sufficiently small neighborhood of the cycle and no heterodimensional connections (except for the cycle itself). Moreover, such dichotomy can extend to nearby systems (Corollary B). We also obtain several perturbative results (Theorem C) which can be used as a standard tool for creating blender-involved robust heterodimensional dynamics in the C^r topology, giving possibilities to improve the regularity of many C^1 results based on the Bonatti-Díaz construction in [13].

1.1 Blenders and moduli of topological conjugacy

There are many definitions of blenders, see e.g. [6, 10–12, 14, 27]. The blenders found in this paper are the same as in [23], which we call *standard*. They carry more dynamical meaning than the original geometrically defined blenders [11–13]. Roughly speaking, a standard blender is a zero-dimensional partially hyperbolic set obtained from a Markov partition with finite (but arbitrarily large) number of elements, such that the iterated function system which arises from the partition satisfies the so-called covering property (see Proposition 4.2). It is a generalization of the *blender-horseshoe* defined in [14], whose corresponding Markov partition had exactly two elements. We refer interested readers to [23, Appendix], where a detailed construction of standard blenders is given. Here for stating our results, it suffices to use a version of the geometric definition [12, Definition 6.11].

In the rest of the paper, let \mathcal{M} be a manifold of dimension at least three and let $Diff^r(\mathcal{M})$ be the space of C^r diffeomorphisms of \mathcal{M} .

Definition 1.1 (Blenders). A basic (i.e., compact, transitive, locally maximal) hyperbolic set Λ of a diffeomorphism $f \in Diff^1(\mathcal{M})$ is called a center-stable (cs) blender if there exists a C^1 -open set \mathcal{D} of d^{ss} -dimensional discs (embedded copies of $\mathbb{R}^{d^{ss}}$) with $d^{ss} = \dim(W^s(\Lambda)) - 1$, such that for every system g which is C^1 -close to f , for the hyperbolic continuation Λ_g of the basic set Λ , the set $W^u(\Lambda_g)$ intersects every element from \mathcal{D} ; a center-unstable (cu) blender is a cs-blender of f^{-1} .

Recall that the index of a transitive hyperbolic set is the rank of its unstable bundle. We denote it by $\text{ind}(\cdot)$. Following [23], we use

Definition 1.2 (Heterodimensional dynamics). We say that a diffeomorphism $f \in Diff^1(\mathcal{M})$ has *heterodimensional dynamics* if there are two basic hyperbolic sets Λ_1 and Λ_2 such that they have different indices and their invariant manifolds intersect cyclically, i.e., $W^u(\Lambda_1) \cap W^s(\Lambda_2) \neq \emptyset$ and $W^u(\Lambda_2) \cap W^s(\Lambda_1) \neq \emptyset$. The quantity $|\text{ind}(\Lambda_1) - \text{ind}(\Lambda_2)|$ is called the coindex of the heterodimensional dynamics.

¹ The results in this paper can be translated without effort to the setting of flows on manifolds of dimension four or higher, see Remark 3.5.

In literature, the above heterodimensional dynamics are called ‘heterodimensional cycles’. However, we reserve the word ‘cycle’ for the simplest version of such dynamics, which is the precise object studied in the paper.

Definition 1.3 (Heterodimensional cycles). A *heterodimensional cycle* of f is a closed invariant set consisting of four orbits: two hyperbolic periodic orbits L_1 and L_2 with $\text{ind}(L_1) \neq \text{ind}(L_2)$, and two heteroclinic orbits, one from $W^u(L_1) \cap W^s(L_2)$ and the other from $W^u(L_2) \cap W^s(L_1)$.

See Figure 1 for an illustration. We will only study codimension-1 heterodimensional cycles, i.e., those with $|\text{ind}(L_1) - \text{ind}(L_2)| = 1$, so for brevity we omit the word ‘codimension-1’ in the remaining paper. Note that we lose no generality by considering only cycles associated with periodic orbits. Indeed, since hyperbolic periodic orbits are dense in a hyperbolic basic set and their stable/unstable invariant manifolds approximate the stable/unstable leaves of the hyperbolic set (see e.g. [21, Theorem 6.4.15]), one can always obtain heterodimensional cycles from heterodimensional dynamics by an arbitrarily C^r -small perturbation.




Figure 1: [23, Figure 1]. A heterodimensional cycle for a three-dimensional diffeomorphism. It consists of two fixed points L_1 and L_2 , a fragile heteroclinic orbit Γ^0 (blue dots) in the non-transverse intersection $W^u(L_1) \cap W^s(L_2)$, and a robust heteroclinic orbit Γ^1 (green dots) in the transverse intersection $W^s(L_1) \cap W^u(L_2)$ (red curves).

We further consider heterodimensional cycles which fulfill certain non-degeneracy conditions defined in Section 3.1. All these conditions can be achieved by an arbitrarily C^r -small perturbation and are C^r -open in the set of diffeomorphisms having a heterodimensional cycle.

The dynamics near a non-degenerate heterodimensional cycle is largely determined by the central multipliers² of the two involved periodic orbits L_1 and L_2 . Let us assume $\text{ind}(L_1) + 1 = \text{ind}(L_2)$, and denote $d_1 := \text{ind}(L_1)$ and $d_2 := \text{ind}(L_2)$. When the multipliers of L_1 are in general position, there is either one real multiplier, say λ , or a pair of conjugate complex multipliers, say λ and λ^* , closest to the unit circle from inside, which we call *center-stable*. Similarly, for L_2 , there is either one real multiplier, say γ , or a pair of conjugate complex multipliers, say γ and γ^* , closest to the unit circle from outside, which we call *center-unstable*. See Section 2 for a detailed discussion. It is well-known [28, 33] that the ratio

$$\theta := -\frac{\ln |\lambda|}{\ln |\gamma|}$$

² The multipliers of a periodic orbit L are the eigenvalues of the derivative of the period map $f^{\text{per}(L)}$ evaluated at any point of L .

and the arguments

$$\omega_1 := \text{Arg}(\lambda) \in [0, \pi] \quad \text{and} \quad \omega_2 := \text{Arg}(\gamma) \in [0, \pi], \quad (1)$$

are *moduli*, i.e., invariants of topological conjugacy, for diffeomorphisms having non-degenerate heterodimensional cycles. Note that, the arguments are also defined for real λ and γ , in which case one has $\omega_1 \in \{0, \pi\}$ and, respectively, $\omega_2 \in \{0, \pi\}$. We prove

Theorem A. *Let $f \in \text{Diff}^r(\mathcal{M})$, with $r = 2, \dots, \infty, \omega$, have a non-degenerate heterodimensional cycle, and let U be any small neighborhood of the cycle.*

- *If $\omega_1/2\pi$ and $\omega_2/2\pi$ are both irrational, then there exist both a cs-blender and a cu-blender in U . The blenders have different indices and their stable and unstable sets intersect cyclically and C^1 -robustly;*
- *If exactly one of $\omega_1/2\pi$ and $\omega_2/2\pi$ is irrational, then there exists at least one blender³ in U ;*
- *If θ is irrational with at least one of λ and γ being nonreal, and $\omega_1/2\pi$ and $\omega_2/2\pi$ do not belong to the set*

$$\left\{ \frac{p}{q} \mid p, q \in \mathbb{N} \text{ are coprime with } 3 \leq q \leq 8 \right\},$$

then there exists at least one blender in U .

In each of the above cases, the blender has index d_1 if it is center-stable, and index d_2 if it is center-unstable.

By definition, if $\omega_1/2\pi$ is irrational, the multiplier λ must be nonreal, and, if it is rational, λ can be either real or nonreal; similarly for $\omega_2/2\pi$ and γ . Hence, the result concerns all the possibilities of the modulus values when at least one of λ and γ is nonreal, except for the possibility that all $\omega_1/2\pi, \omega_2/2\pi$ and θ are rational, considered in Theorem B below. Theorem A is our main result and it reflects the essential relation between values of the moduli and the existence of blenders. However, it is too general for practical use in the sense that not much information of the blenders is provided. In fact, this theorem follows from several much finer results collected in Theorems 1 and 2 in Section 2, where the homoclinic relations between the found blenders and periodic orbits involved in the cycle are established. We stress that Theorem A has a non-perturbative nature. This is the main difference from the Bonatti-Díaz construction in [13], where the cycle is destroyed to create blenders.

The case where both λ and γ are real is fully studied in [23], where such cycles are further classified to types I and II, according to whether an orientation-like property is satisfied by the central dynamics along the non-transverse heteroclinic orbit, see [23, Section 2.5].

Theorem ([23, Theorem 1]). *For any non-degenerate type-I cycle and any neighborhood U of it, if θ is irrational, then there exists a blender in U .*

In summary, we obtain

Corollary A. *For any non-degenerate heterodimensional cycle which is not type-II, it is accumulated by infinitely many blenders if*

- *either $\frac{\omega_i}{2\pi}$ is irrational for some $i \in \{1, 2\}$,*

³ This blender, and also the one in the third item, can be center-stable or center-unstable. This depends on the f , and will be made precise in Propositions 4.3 and 5.1

- or θ is irrational, and $\frac{\omega_i}{2\pi} = \frac{p_i}{q_i}$ for $i = 1, 2$, where $p_i, q_i \in \mathbb{Z}$ are coprime with $q_i > 8$.

Note that being type II particularly means that both central multipliers are real. This result significantly strengthened [23, Theorem D], where more strict arithmetic conditions (e.g. rational independence among the moduli and 1) are imposed in the cases where at least one of λ and γ is nonreal.

We remark that the “superposition” of a properly chosen finite collection of the blenders in Corollary A carries dynamical properties which normally do not hold for a single blender. In a forthcoming paper [25], we show that these blenders can be used to produce uncountably many C^1 -robust homoclinic tangencies, and they can also be embedded into a family to form a parablender in the sense of [7], giving rise to robust coexistence of infinitely many sinks for generic families of diffeomorphisms.

1.2 Simple hyperbolic dynamics

In the last remaining case where θ , $\omega_1/2\pi$ and $\omega_2/2\pi$ are all rational (including the possibilities of λ and/or γ being real), there are generally no blenders near the heterodimensional cycle. To state the results precisely, let us introduce some necessary notions.

Denote the heterodimensional cycle of f by Γ , and the two involved heteroclinic orbits by $\Gamma^0 \subset W^u(L_1) \cap W^s(L_2)$ and $\Gamma^1 \subset W^u(L_2) \cap W^s(L_1)$, where we assumed that $\text{ind}(L_1) + 1 = \text{ind}(L_2)$. Recall that the hyperbolic periodic orbits admit unique continuations for all diffeomorphisms C^1 -close to f . Thus, for any small neighborhood U of Γ , we can find a codimension-1 subspace $\mathcal{H}_f \subset \text{Diff}^r(\mathcal{M})$ such that each $g \in \mathcal{H}_f$ has a cycle $\Gamma_g \subset U$, involving the continuations of L_1 and L_2 , and heteroclinic orbits close to Γ^0 and Γ^1 . Since θ depends continuously on diffeomorphisms, inside \mathcal{H}_f there is a codimension-2 subspace $\mathcal{H}_{f,\theta} = \mathcal{H}_f \cap \{\theta = \text{const}\}$, where all diffeomorphisms have the same θ values as f . Similarly, one can define two codimension-3 subspaces $\mathcal{H}_{f,\theta,\omega_1}$ and $\mathcal{H}_{f,\theta,\omega_2}$ and one codimension-4 subspace $\mathcal{H}_{f,\theta,\omega_1,\omega_2}$ by intersecting $\mathcal{H}_{f,\theta}$ with $\{\omega_1 = \text{const}\}$, $\{\omega_2 = \text{const}\}$ and both of them, respectively.

We define the splitting parameter μ for the heterodimensional cycle as the continuous functional of a neighborhood $\mathcal{U} \subset \text{Diff}^r(\mathcal{M})$ of f which measures the distance between $W^u(L_1)$ and $W^s(L_2)$ in certain neighborhood (see Remark 3.1). Then, $\mu(g) = 0$ for $g \in \mathcal{U}$ corresponds to the codimension-1 subspace \mathcal{H}_f . For a one-parameter family $\{f_\varepsilon\}$ of C^r diffeomorphisms with $f_0 = f$, we call it a *generic unfolding family* (when there is one parameter) or a *rank-1 unfolding family* (when there is more than one parameter) if

$$\text{rank} \left. \frac{d\mu(\varepsilon)}{d\varepsilon} \right|_{\varepsilon=0} = 1. \quad (2)$$

Let \mathcal{N} be the set of orbits which lie entirely in U .

Definition 1.4 (Simple hyperbolic dynamics). A generic one-parameter unfolding family $\{f_\varepsilon\}$ is said to have *simple hyperbolic dynamics* if

- at $\varepsilon = 0$, \mathcal{N} is the union of L_1 , L_2 , Γ^0 , and the orbits of transverse intersection of $W^u(L_2)$ with $W^s(L_1)$ near Γ^1 ;
- at all sufficiently small $\varepsilon \neq 0$, \mathcal{N} is comprised

- either by L_2 , an index- d_1 uniformly-hyperbolic compact set Λ_1 containing L_1 , and transverse heteroclinic orbits in $W^u(L_2) \cap W^s(\Lambda_1)$,
- or by L_1 , an index- d_2 uniformly-hyperbolic compact set Λ_2 containing L_2 , and transverse heteroclinic orbits in $W^u(\Lambda_2) \cap W^s(L_1)$.

In particular, there exist no blenders at $\varepsilon = 0$ and no heterodimensional dynamics at $\varepsilon \neq 0$.

Theorem B. *Let $f \in \text{Diff}^r(\mathcal{M})$, with $r = 2, \dots, \infty, \omega$, have a heterodimensional cycle with all θ , $\omega_1/2\pi$ and $\omega_2/2\pi$ being rational. There exists a set $\mathcal{H}'_f \subset \mathcal{H}_f$ such that any generic one-parameter unfolding family $\{g_\varepsilon\}$ with $g_0 \in \mathcal{H}'_f$ has only simple hyperbolic dynamics defined above. Moreover, the set \mathcal{H}'_f satisfies that*

- if λ and γ are both real, then \mathcal{H}'_f is open and dense in $\mathcal{H}_{f,\theta}$,
- if λ is nonreal and γ is real, then \mathcal{H}'_f is open and dense in $\mathcal{H}_{f,\theta,\omega_1}$,
- if λ is real and γ is nonreal, then \mathcal{H}'_f is open and dense in $\mathcal{H}_{f,\theta,\omega_2}$, and
- if λ and γ are both nonreal, then \mathcal{H}'_f is open and dense in $\mathcal{H}_{f,\theta,\omega_1,\omega_2}$.

The proof is given in Section 7. The result implies that the blenders which exists for the moduli values in Corollary A depart from the heterodimensional cycle immediately once we change the moduli to rational values within \mathcal{H}_f .

1.3 Robust heterodimensional dynamics arisen from perturbations

A direct consequence of the existence of blenders near a heterodimensional cycle is the emergence of robust heterodimensional dynamics, defined as

Definition 1.5 (Robust heterodimensional dynamics). We say that a diffeomorphism f exhibits C^1 -robust heterodimensional dynamics if it has heterodimensional dynamics involving two hyperbolic basic sets Λ_1 and Λ_2 , where at least one of them is non-trivial, and there exists a C^1 -neighborhood \mathcal{U} of f such that every $g \in \mathcal{U}$ has heterodimensional dynamics involving the continuations of Λ_1 and Λ_2 .

It is proven that robust heterodimensional dynamics can always be obtained by an arbitrarily small perturbation of a heterodimensional cycle, see [13, 15] for the C^1 topology and [23] for the C^r topology. In this paper we obtained a parametric version of such result, which improves [23, Theorem B] in terms of the class of families allowed. Specifically, we consider the case where at least one of λ and γ is nonreal.

Let f have a heterodimensional cycle involving L_1 and L_2 with $\text{ind}(L_1) + 1 = \text{ind}(L_2)$. We further classify the unfolding families with more than one parameter. We already defined rank-1 unfolding families by (2). When there are at least two parameters, a family $\{f_\varepsilon\}$ is called a *rank-2 unfolding family* if

- $\text{rank} \frac{d(\mu(\varepsilon), \omega_i(\varepsilon))}{d\varepsilon} \Big|_{\varepsilon=0} = 2$, $\frac{\omega_i(0)}{2\pi} \notin \left\{0, \frac{1}{2}\right\}$, $\frac{\omega_{3-i}(\varepsilon)}{2\pi} \not\equiv \frac{p}{q}$ for $3 \leq q \leq 8$,
where $i = 1, 2$, or
- $\text{rank} \frac{d(\mu(\varepsilon), \theta(\varepsilon))}{d\varepsilon} \Big|_{\varepsilon=0} = 2$, $\left(\frac{\omega_1(0)}{2\pi}, \frac{\omega_2(0)}{2\pi}\right) \notin \left\{0, \frac{1}{2}\right\}^2$, $\frac{\omega_{1,2}(\varepsilon)}{2\pi} \not\equiv \frac{p}{q}$ for $3 \leq q \leq 8$.

Here p, q are coprime integers, and ' \neq ' means not equal in a small neighborhood of $\varepsilon = 0$. Recall that we take $\omega_{1,2} \in [0, \pi]$ (see (1)). Hence, the multipliers λ and γ are real if and only if $\omega_1/2\pi \in \{0, 1/2\}$ and $\omega_2/2\pi \in \{0, 1/2\}$, respectively. If the family has least three parameters, we call it a *rank-3 unfolding family* if

- $\text{rank} \frac{d(\mu(\varepsilon), \omega_i(\varepsilon), \theta(\varepsilon))}{d\varepsilon} \Big|_{\varepsilon=0} = 3, \quad \frac{\omega_i(0)}{2\pi} \notin \left\{0, \frac{1}{2}\right\}, \quad \frac{\omega_{3-i}(\varepsilon)}{2\pi} \neq \frac{p}{q} \quad \text{for } 3 \leq |q| \leq 8,$
where $i = 1, 2$, or
- $\text{rank} \frac{d(\mu(\varepsilon), \omega_1(\varepsilon), \omega_2(\varepsilon))}{d\varepsilon} \Big|_{\varepsilon=0} = 3, \quad \left(\frac{\omega_1(0)}{2\pi}, \frac{\omega_2(0)}{2\pi}\right) \notin \left\{0, \frac{1}{2}\right\}^2.$

Note that the rank-2 and rank-3 unfolding families are generic in the space of families with at least two and, respectively, three parameters.

Recall that two hyperbolic periodic points P and Q are *homoclinically related* if they have the same index and their invariant manifolds intersect cyclically and transversely: $W^s(P) \pitchfork W^u(Q) \neq \emptyset$ and $W^u(P) \pitchfork W^s(Q) \neq \emptyset$. Two hyperbolic basic sets are homoclinically related if they have two periodic orbits which are homoclinically related.

Theorem C. *Let $f \in \text{Diff}^r(\mathcal{M})$, with $r = 3^4, \dots, \infty, \omega$, have a non-degenerate heterodimensional cycle such that at least one of λ and γ is nonreal. Let $\{f_\varepsilon\}$ be either a rank- i ($i = 2, 3$) unfolding family, or a rank-1 unfolding family with f satisfying the condition in Corollary A. Given any neighborhood U of the cycle, there exist open regions in the ε -space accumulating on $\varepsilon = 0$ such that each corresponding diffeomorphism f_ε has robust heterodimensional dynamics in U , which involve two non-trivial hyperbolic basic sets Λ_1 of index d_1 and Λ_2 of index d_2 . The sets Λ_1 and Λ_2 are homoclinically related to the continuations of L_1 and L_2 , respectively. Moreover, we have that*

- *if the family is rank-1 or -2, then either Λ_1 is a cs-blender or Λ_2 is a cu-blender or both;*
- *if the family is rank-3, then Λ_1 is a cs-blender and Λ_2 is a cu-blender.*

The theorem follows from Theorem 4 in Section 2, which concerns one-parameter unfoldings of heterodimensional cycles with different arithmetic properties of the moduli. It should be noted that not for all cases the cycle needs to be unfolded. For example, in the second case of rank-3 unfolding families, the result is achieved by keeping $\mu = 0$ and adjusting ω_i so that $\omega_i/2\pi$ are both irrational. If we do not require the homoclinic relations between Λ_i and the continuations of L_i , then the cycle can be preserved also in the first case of rank-3 unfolding families. This fact can be inferred from Theorem 2.

The bifurcations of cycles with real λ and γ are studied in [23]. For completeness, we summarize the result as

Theorem 1.6 ([23, Theorem 2 and Corollary 2]). *Let f has a non-degenerate heterodimensional cycle with both real central multipliers. Suppose that*

- *when the cycle is type-I, $\{f_\varepsilon\}$ is rank-2 or rank-1 with irrational $\theta(0)$; and*
- *when the cycle is type-II, $\{f_\varepsilon\}$ is rank-2 .*

⁴ Here we require one more regularity, because this guarantees a smooth dependence on the parameters of the coordinates used in the proof. For $r = 2$, the result is still true with a minor adjustment to the arguments, see the discussion above [23, Corollary 2] for details.

Given any neighborhood U of the cycle, there exist open regions in the ε -space accumulating on $\varepsilon = 0$ such that each corresponding diffeomorphism f_ε has robust heterodimensional dynamics in U , which involve two non-trivial hyperbolic basic sets Λ_1 of index d_1 and Λ_2 of index d_2 .

When both λ and γ are real, it is shown in [23] that the coexisting of a cs-blender and a cu-blender cannot be achieved by local C^r -small bifurcations.

1.4 A dichotomy for local dynamics near heterodimensional cycles

In closing the introduction, we present a result on the complexity of the dynamics near a heterodimensional cycle. Note from Theorem B that by taking the union of $\mathcal{H}'_f \subset \mathcal{H}_f$ over different rational values of the moduli, we obtain a dense subset $\mathcal{S}_f \subset \mathcal{H}_f$ where each diffeomorphism g has simple hyperbolic dynamics defined in Section 1.1. Since non-degenerate cycles are open and dense in \mathcal{H}_f (see Section 3.1), the above observation along with Corollary A, Theorem C and Theorem 1.6 leads to

Corollary B. *Let $f \in \text{Diff}^r(\mathcal{M})$, with $r = 3, \dots, \infty, \omega$, have a heterodimensional cycle Γ . There exists a dense subset $\mathcal{S} \in \mathcal{H}_f$, and, when the cycle is not type-I, also exists a dense subset $\mathcal{C} \in \mathcal{H}_f \setminus \mathcal{S}$, such that*

- any close arc transverse to \mathcal{H}_f at $g \in \mathcal{S}$ has simple hyperbolic dynamics near the cycle Γ_g , and
- for $g \in \mathcal{C}$, the cycle Γ_g is a limit of blenders, and any close arc transverse to \mathcal{H}_f at g intersects a sequence of open sets accumulating on g , which correspond to diffeomorphisms having robust heterodimensional dynamics involving blenders.

1.5 Plan of the proofs

In Section 2, to facilitate the discussion of the problem, we first define two types of heterodimensional cycles, *saddle-focus* and *double-focus*, depending on whether only one or both central multipliers are nonreal. Then, according to the types and finer arithmetic properties of $\theta, \omega_1/2\pi, \omega_2/2\pi$ (e.g. rational independence), we split Theorem A into slightly more technical parts, Theorems 1–3. After that, based on these theorem, we state a bifurcation result, Theorem 4, which implies Theorem C.

In Section 3, we describe the non-degeneracy conditions and introduce the first return maps associated with heterodimensional cycles. We will present the cross-form formulas for these maps which are obtained in [23].

We prove the non-perturbative results Theorems 1–3 in Sections 4 and 5 for saddle-focus, and, respectively, double-focus cycles. A key step is the use of a criterion for the existence of blenders, which is adapted to the cross-form formulas, such that the problem of finding blenders is reduced to finding sequences $\{(k_n, m_n)\} \subset \mathbb{N}^2$ such that the set

$$\left\{ m_n - k_n\theta, k_n \frac{\omega_1}{2\pi}, m_n \frac{\omega_2}{2\pi} \right\} \subset \mathbb{R} \times S^1 \times S^1$$

has suitable accumulation points. The criterion is introduced in Section 4.1.

In Section 6, we prove the perturbative result, Theorem 4. This is mainly about how the invariant

manifolds of the periodic orbits move when the cycle is unfolded. The arguments are based on combinations of various results in [23].

Finally, we prove Theorem B in Section 7. The result for the case where both central multipliers are real is a direct consequence of [23, Theorem 6]. We extend this theorem for saddle-focus and double-focus cycles.

2 Results for different types of heterodimensional cycles

In this section, we state several detailed results considering more subcases of the arithmetic properties of the moduli. These results imply Theorems A and C. We distinguish three main types of heterodimensional cycles, defined as follows. Let $f \in \text{Diff}^r(\mathcal{M})$ ($r = 2, \dots, \infty$) have a heterodimensional cycle involving two hyperbolic periodic orbits L_1 and L_2 satisfying $\text{ind}(L_1) + 1 = \text{ind}(L_2)$. Denoting $d = \dim(\mathcal{M})$ and $d_i = \text{ind}(L_i)$ ($i = 1, 2$), we can order the multipliers of L_i as

$$|\lambda_{i,d-d_i}| \leq \dots \leq |\lambda_{i,2}| \leq |\lambda_{i,1}| \leq 1 \leq |\gamma_{i,1}| \leq |\gamma_{i,2}| \leq \dots \leq |\gamma_{i,d_i}|. \quad (3)$$

Up to an arbitrarily C^r -small perturbation, we can assume that the multipliers are in general position. So, the center-stable multiplier(s) either is $\lambda_{1,1}$ being simple and real, or are $\lambda_{1,1}$ and $\lambda_{1,2} = \lambda_{1,1}^*$ being a pair of simple complex conjugate numbers; and the center-unstable multiplier(s) either is $\gamma_{2,1}$ being simple and real or are $\gamma_{2,1}$ and $\gamma_{2,2} = \gamma_{2,1}^*$ being a pair of simple complex conjugate numbers. We say that the heterodimensional cycle is of

- saddle type, if $\lambda_{1,1} =: \lambda$ and $\gamma_{2,1} =: \gamma$ are real and satisfy $|\lambda_{1,2}| < |\lambda_{1,1}|$ and $|\gamma_{2,1}| < |\gamma_{2,2}|$;
- saddle-focus type, if either

$$\lambda_{1,1} = \lambda_{1,2}^* = \lambda e^{i\omega}, \quad \omega \in (0, \pi), \quad \text{and} \quad \gamma_{2,1} =: \gamma \quad \text{is real,}$$

where $\lambda > |\lambda_{1,3}|$ and $|\gamma_{2,1}| < |\gamma_{2,2}|$, or

$$\gamma_{2,1} = \gamma_{2,2}^* = \gamma e^{i\omega}, \quad \omega \in (0, \pi), \quad \text{and} \quad \lambda_{1,1} =: \lambda \quad \text{is real,}$$

where $\gamma < |\gamma_{2,3}|$ and $|\lambda_{1,2}| < |\lambda_{1,1}|$;

- double-focus type, if

$$\lambda_{1,1} = \lambda_{1,2}^* = \lambda e^{i\omega_1}, \quad \omega_1 \in (0, \pi), \quad \text{and} \quad \gamma_{2,1} = \gamma_{2,2}^* = \gamma e^{i\omega_2}, \quad \omega_2 \in (0, \pi),$$

where $\lambda > |\lambda_{1,3}|$ and $0 < \gamma < |\gamma_{2,3}|$.

As the second case for saddle-focus type can be reduced to the first one by considering f^{-1} , we always assume that for saddle-focus cycles $\lambda_{1,1}$ is nonreal and $\gamma_{2,1}$ is real.

As mentioned, in this paper we focus on saddle-focus and double-focus heterodimensional cycles (while the saddle ones have been comprehensively studied in [23]). To formulate our results in a compact way, we use a version of the notion of activation introduced in [13].

Definition 2.1 (Activation). A cs-blender Λ of index i is said to be *activated* by a hyperbolic set H if $\text{ind}(H) = i + 1$, $W^u(H)$ intersects $W^s(\Lambda)$ transversely, and $W^s(H)$ contains a disc in the open set

\mathcal{D} in Definition 1.1 (with $d^{ss} = d - i - 1$) so that the non-transverse intersection $W^s(H) \cap W^u(\Lambda)$ is C^1 -robustly non-empty.

Similarly, a cu-blender Λ of index i is *activated* by a hyperbolic set H if $\text{ind}(H) = i - 1$, $W^s(H) \pitchfork W^u(\Lambda) \neq \emptyset$, and $W^u(H)$ contains a disc in an open set \mathcal{D} of embeddings of $(i - 1)$ -dimensional discs, each of which intersects C^1 -robustly the set $W^s(\Lambda)$.

Let us start with saddle-focus heterodimensional cycles. As mentioned, we always assume that the center-stable multipliers are a pair of complex conjugate numbers and the center-unstable multiplier is real. Throughout the remaining paper, we use the notation p/q to refer to rational numbers with p, q being coprime integers; and, in all cases, a cs-blender always has index d_1 and a cu-blender always has index d_2 .

Theorem 1. *Let $f \in \text{Diff}^r(\mathcal{M})$ have a non-degenerate saddle-focus heterodimensional cycle, and let U be any small neighborhood of the cycle.*

- (1) *If $\omega/2\pi$ is irrational, then there exists a cu-blender in U , homoclinically related to L_2 .*
- (2) *If $\omega/2\pi$ is irrational, and $\theta, \omega/2\pi, 1$ are rationally independent (so θ is also irrational), then there exists a cs-blender in U , in addition to the cu-blender of case (1) which is homoclinically related to L_2 . The two blenders activate each other, and in particular L_2 activates the cs-blender.*
- (3) *If $\omega/2\pi = p/q$ with $q \geq 9$ and θ is irrational, then there exists a blender in U .*

Remark 2.2. The type of the blender in case (3) depends on certain coefficients of the first return map, see case 1 of Proposition 4.3.

The theorem is proved in Section 4. Case (2) here, and also case (2) in Theorem 2 below, follow from [23, Theorem 7], but we reprove them with a more general approach. Let us explain briefly how the lower bound for q appears in case (3). The existence of blenders is determined by the central dynamics of the first return maps $T_{k,m}$ along a heterodimensional cycle, where k and m represent the number of iterations spent near the two periodic orbits L_1 and L_2 , respectively. When the center-stable multiplier of L_1 is nonreal, the rotation brought by it leads to the emergence of trigonometric terms like $\sin(k\omega + \eta)$ in the formula for the one-dimensional central part of the first return map. To prove the existence of blenders, we need certain inequalities involving these trigonometric terms to be satisfied (see (22)), and in most cases we only need $\sin(k\omega + \eta)$ to take sufficiently many different values for different k . This is ensured by a lower bound on q . See Section 3.2 for details.

For double-focus heterodimensional cycles, we have

Theorem 2. *Let $f \in \text{Diff}^r(\mathcal{M})$ have a non-degenerate double-focus heterodimensional cycle, and let U be any small neighborhood of the cycle.*

- (1) *Suppose that at least one of $\omega_i/2\pi$ ($i = 1, 2$) is irrational.*
 - (i) *If $\omega_1/2\pi$ is irrational and $\omega_2/2\pi = p/q$ with $q \geq 8$, then there exists a cu-blender in U , homoclinically related to L_2 .*
 - (ii) *If $\omega_2/2\pi$ is irrational and $\omega_1/2\pi = p/q$ with $q \geq 8$, then there exists a cs-blender in U , homoclinically related to L_1 .*
 - (iii) *If both $\omega_1/2\pi$ and $\omega_2/2\pi$ are irrational, then in U there exist simultaneously a cs-blender, homoclinically related to L_1 , and a cu-blender, homoclinically related to L_2 ; the two blenders activate each other, and in particular L_1 activates the cu-blender and L_2 activates the cs-blender.*

(2) Suppose that θ and exactly one of $\omega_i/2\pi$ are irrational.

- (i) If $\theta, \omega_1/2\pi, 1$ are rationally independent and $\omega_2/2\pi = p/q$ with $q \geq 8$, then there exists in U a cs-blender, in addition to the cu-blender of (1.i) which is homoclinically related to L_2 .
- (ii) If $\theta^{-1}, \omega_2/2\pi, 1$ are rationally independent and $\omega_1/2\pi = p/q$ with $q \geq 8$, then there exists a cu-blender in U , in addition to the cs-blender of (1.ii) which is homoclinically related to L_1 .

In each of the two cases, the two blenders activate each other, and in particular L_2 activates the cs-blender in (2.i) and L_1 activates the cu-blender in (2.ii).

(3) If θ is irrational, and $\omega_1/2\pi = p_1/q_1$ and $\omega_2/2\pi = p_2/q_2$ such that either $q_1 \geq 9$ and $q_2 \geq 8$ or $q_2 \geq 9$ and $q_1 \geq 8$, then there always exists a blender in U .

In all cases, a cs-blender always has index d_1 and a cu-blender always has index d_2 .

Remark 2.3. The type of the blender in case (3) depends on certain coefficients of the first return map, see case 1 of Proposition 5.1.

This theorem is proved in Section 5, which along with Theorem 1 implies Theorem A. The reason for the lower bounds of q is the same as for the saddle-focus cycles, where the related inequalities are given in (32).

The next result concerns the possibilities for the simultaneous appearance of a cs-blender and a cu-blender in cases (3) of Theorems 1 and 2. In fact, such coexistence is ensured when q is larger than a number which depends on the diffeomorphism and is locally constant. Recall that, for a diffeomorphism g having a non-degenerate cycle Γ , we denote by \mathcal{H}_g the codimension-1 subspace in $\text{Diff}^r(\mathcal{M})$ where every $f \in \mathcal{H}_g$ has a heterodimensional cycle Γ_f near Γ . Also note that being non-degenerate is C^r -open in \mathcal{H}_g , see Section 3.1.

Theorem 3. *Let g have a non-degenerate heterodimensional cycle of saddle-focus or double-focus type Γ . For any sufficiently small neighborhood \mathcal{U} of g in \mathcal{H}_g , there exists a number $N(\mathcal{U})$ such that a diffeomorphism $f \in \mathcal{U}$ has a pair of mutually activating cs-blender and cu-blender in any neighborhood of Γ_f if*

- in the saddle-focus case, $\theta(f)$ is irrational and $\omega(f)/2\pi = p/q$ with $q \geq N(\mathcal{U})$, or
- in the double-focus case, $\theta(f)$ is irrational and $\omega_i(f)/2\pi = p_i/q_i$ with either $q_1 \geq N(\mathcal{U})$ and $q_2 \geq 8$ or $q_2 \geq N(\mathcal{U})$ and $q_1 \geq 8$.

We prove this theorem in Sections 4.4 and 5.3. In light of this result, by taking $q \rightarrow \infty$, it is intuitive to expect coexisting blenders near saddle-focus cycles when both θ and $\omega/2\pi$ are irrational. It is true if $\theta, \omega/2\pi$ and 1 are rationally independent, as in case (1) of Theorem 1. However, in the absence of such rational independence, we cannot prove coexisting blenders, see case 2 of Proposition 4.3. Similar comments apply to double-focus cycles, see case 2.2 of Proposition 5.1.

Let us now present some perturbative results based on the above theorems. The results show that, by splitting the heterodimensional cycle, one builds homoclinic relations between the involved periodic orbits and the blenders found above, and hence obtains robust heterodimensional dynamics. Recall that a parameter family $\{f_\varepsilon\}$ is of rank-1 if $d\mu/d\varepsilon \neq 0$ at $\varepsilon = 0$, see (2). Let us denote the cs-blenders and cu-blenders in Theorems 1–3 by Λ^{cs} and Λ^{cu} , respectively. We use the same notations $L_{1,2}, \Lambda^{cs}, \Lambda^{cu}$ for their continuations at $\varepsilon \neq 0$.

Theorem 4. Consider any rank-1 unfolding family $\{f_\varepsilon\}$ of C^r ($r = 3, \dots, \infty, \omega$) diffeomorphisms with $f_0 = f$ from either of Theorems 1–3. Let U be any neighborhood of the cycle of f . There exists a sequence $\{\varepsilon_j\}$ accumulating on $\varepsilon = 0$ such that for each f_{ε_j} ,

- in Theorem 1.(1) and Theorem 2.(1.i), Λ^{cu} is homoclinically related to L_2 , and is activated by a non-trivial hyperbolic basic set in U which is homoclinically related to L_1 ;
- in Theorem 2.(1.ii), Λ^{cs} is homoclinically related to L_1 , and is activated by a non-trivial hyperbolic basic set in U which is homoclinically related to L_2 ;
- in Theorem 1.(2), Theorem 2.(2) and Theorem 3, Λ^{cs} and Λ^{cu} activate each other, and are homoclinically related to L_1 and L_2 , respectively;
- in Theorem 1.(3) and Theorem 2.(3), if the blender is center-stable, it is homoclinically related to L_1 , and is activated by a non-trivial hyperbolic basic set in U which is homoclinically related to L_2 ; if the blender is center-unstable, then it is homoclinically related to L_2 , and is activated by a non-trivial hyperbolic basic set in U which is homoclinically related to L_1 .

Moreover, in all cases, the periodic orbits, blenders and hyperbolic sets involved in the robust heterodimensional dynamics, as well as the heteroclinic orbits constituting the intersections involved in the activations and homoclinic relations all lie in U .

This theorem is proved in Section 6. Theorem C follows immediately from the above results, by using the additional parameters to change $\theta, \omega, \omega_{1,2}$ to fulfill the hypotheses required in corresponding cases. The last part of Theorem 4 shows that all the objects related to the obtained robust heterodimensional dynamics are localized arbitrarily close to the original cycle. This fact is important to the (local) stabilization theory of heterodimensional cycles [5, 15, 23].

3 First return maps

In this section, we introduce the formulas of the first return maps of non-degenerate saddle-focus and double-focus heterodimensional cycles, which are obtained in [23]. For that, we first define the required non-degeneracy conditions.

3.1 Non-degeneracy conditions

Let $f \in \text{Diff}^r(\mathcal{M})$ with $r = 2, \dots, \infty, \omega$ have a heterodimensional cycle involving two hyperbolic periodic orbits L_1 and L_2 . Assume $d_1 := \text{ind}(L_1)$ and $d_2 := \text{ind}(L_2)$ satisfy $d_1 + 1 = d_2$. Let Γ^0 be a heteroclinic orbit in $W^u(L_1) \cap W^s(L_2)$, and Γ^1 be a heteroclinic orbit in $W^s(L_1) \cap W^u(L_2)$. We denote the heterodimensional cycle as the closed invariant set $\Gamma = L_1 \cup L_2 \cup \Gamma^0 \cup \Gamma^1$. The orbit Γ^0 from the non-transverse intersection is called a *fragile heteroclinic orbit* since it can be destroyed by a small perturbation, while Γ^1 from the transverse intersection is called a *robust heteroclinic orbit* as it persists in small perturbations, see Figure 1. Following [23], we impose below non-degeneracy conditions C1–C4 on the heterodimensional cycle.

Take two points O_1 and O_2 from the two periodic orbits L_1 and L_2 . Let us first collect some preliminary facts on the local dynamics of hyperbolic periodic points (see e.g. [32]). Let $\text{per}(O_i)$

($i = 1, 2$) be the periods of O_i and denote $F_i := f^{per(O_i)}$. We call F_i the *local maps*. Let U_{0i} be some sufficiently small neighborhoods of O_i . In U_{01} , the map F_1 has a $(d - d_1)$ -dimensional local stable invariant manifold $W_{loc}^s(O_1)$ and a d_1 -dimensional local unstable manifold $W_{loc}^u(O_1)$. On $W_{loc}^s(O_1)$ there exists a strong-stable C^r -smooth foliation \mathcal{F}^{ss} , which particularly includes the $(d - d_1 - 2)$ -dimensional strong-stable invariant manifold $W_{loc}^{ss}(O_1)$. There also exists a $(d_1 + 2)$ -dimensional extended-unstable invariant manifold $W_{loc}^{uE}(O_1)$ tangent at O_1 to the eigenspace corresponding to the two conjugate nonreal center-stable multipliers and the unstable multipliers $\gamma_{1,1}, \dots, \gamma_{1,d_1}$ (recall that the multipliers are introduced in (3) and in the saddle-focus case we always assume that L_1 has nonreal center-stable multipliers). The manifold $W_{loc}^{uE}(O_1)$ is not uniquely defined; however, any two such manifolds contain $W_{loc}^u(O_1)$ and are tangent to each other at points of $W_{loc}^u(O_1)$.

Similarly, there exists a strong-unstable C^r -smooth foliation \mathcal{F}^{uu} on the d_2 -dimensional local unstable invariant manifold $W_{loc}^u(O_2)$. Denote by d^{cu} the number of center-unstable multipliers of O_2 , so $d^{cu} = 1$ if the cycle is of saddle-focus type, or $d^{cu} = 2$ if it is of double-focus type. The manifold $W_{loc}^u(O_2)$ includes the $(d_2 - d^{cu})$ -dimensional strong-unstable invariant manifold $W_{loc}^{uu}(O_2)$. Here we also have a $(d - d_2 + d^{cu})$ -dimensional extended-stable invariant manifold $W_{loc}^{sE}(O_2)$, corresponding to the center-unstable multiplier(s) and the stable multipliers $\lambda_{2,1}, \dots, \lambda_{2,d-d_2}$. Any two such manifolds contain $W_{loc}^s(O_2)$ and are tangent to each other at the points of $W_{loc}^s(O_2)$.

Next, using the heteroclinic orbits Γ^0 and Γ^1 , we define *transition maps* connecting neighborhoods of O_1 and O_2 . Take in Γ^0 a pair of points $M_1^- \in W_{loc}^u(O_1)$ and $M_2^+ \in W_{loc}^s(O_2)$ so that $f^{n_1}(M_1^-) = M_2^+$ for some integer $n_1 > 0$, and in Γ^1 a pair of points $M_2^- \in W_{loc}^u(O_2)$ and $M_1^+ \in W_{loc}^s(O_1)$ so that $f^{n_2}(M_2^-) = M_1^+$ for some integer $n_2 > 0$. The transition maps are then defined as $F_{12} := f^{n_1}$, taking a small neighborhood of M_1^- to a small neighborhood of M_2^+ , and $F_{21} := f^{n_2}$, taking a small neighborhood of M_2^- to a small neighborhood of M_1^+ .

Remark 3.1. The splitting parameter introduced in Section 1.1 can be defined to measure the signed distance between $F_{12}(W_{loc}^u(O_1))$ and $W_{loc}^s(O_2)$ in a small neighborhood of M_2^+ .

Figure 2 provides an illustration in the straightened coordinates. We introduce the following non-degeneracy conditions:

- C1:** $F_{12}^{-1}(W_{loc}^{sE}(O_2)) \cap W_{loc}^u(O_1) \neq \emptyset$ at the point M_1^- and $F_{12}(W_{loc}^{uE}(O_1)) \cap W_{loc}^s(O_2) \neq \emptyset$ at M_2^+ ;
- C2:** the leaf of \mathcal{F}^{uu} at the point M_2^- is not tangent to $F_{21}^{-1}(W_{loc}^s(O_1))$ and the leaf of \mathcal{F}^{ss} at the point M_1^+ is not tangent to $F_{21}(W_{loc}^u(O_2))$, and
- C3:** $\Gamma^1 \cap (W_{loc}^{ss}(O_1) \cup W_{loc}^{uu}(O_2)) = \emptyset$, i.e., the points M_1^+ and M_2^- lie outside of $W_{loc}^{ss}(O_1)$ and $W_{loc}^{uu}(O_2)$, respectively.

The first two conditions are about the non-degeneracy of the non-transverse and transverse intersections. Condition **C2** is C^r -open, and condition **C1** is C^r -open in the \mathcal{H}_f , the class of diffeomorphisms with the heterodimensional cycle defined above Theorem B. One can always make f fulfill conditions **C1** and **C2** by an arbitrarily small C^r -perturbation; more specifically, in the smooth case one adds a local perturbation to f and in the analytic case one follows the scheme used in [16, 18]. Condition **C3** is also C^r -open, and, if it is not satisfied, then, without perturbing f , we can always find another robust heteroclinic orbit which satisfies **C3** and is arbitrarily close to Γ^1 . To see this, note that condition **C2** shows that the curve $\ell_1 = W_{loc}^s(O_1) \cap F_{21}(W_{loc}^u(O_2))$ is not tangent to $W_{loc}^{ss}(O_1) \cup F_{21}(W_{loc}^{uu}(O_2))$. Hence, there is always another point in ℓ_1 curve lying outside of that $W_{loc}^{ss}(O_1)$ and $W_{loc}^{uu}(O_2)$. We then take this orbit of this point as the new robust heteroclinic orbit in our cycle.




Figure 2: A heterodimensional cycle satisfying **C1–C3** in the straightened coordinates. Here $W_{loc}^u(O_1)$ is the vertical y -axis, $W_{loc}^s(O_1)$ is the horizontal (x, z) -plane, $W_{loc}^s(O_2)$ is the vertical v -axis, $W_{loc}^u(O_2)$ is the horizontal (u, w) -plane, ℓ^{ss} is a strong-stable leaf through M_1^+ , ℓ^{uu} is a strong-unstable leaf through M_2^- . The two red curves are $\ell_1 = W_{loc}^s(O_1) \cap F_{21}(W_{loc}^u(O_2))$ and $\ell_2 = F_{21}^{-1}\ell_1$.

There is one more non-degeneracy condition. To state it precisely, let us introduce C^r -coordinates $(x, y, z) \in \mathbb{R}^2 \times \mathbb{R}^{d_1} \times \mathbb{R}^{d-d_1-2}$ in U_{O_1} such that the local invariant manifolds as well as the leaves of the foliation \mathcal{F}^{ss} are straightened (see Figure 2). So we have

$$W_{loc}^s(O_1) = \{y = 0\}, \quad W_{loc}^u(O_1) = \{x = 0, z = 0\}, \quad W_{loc}^{ss}(O_1) = \{x = 0, y = 0\}.$$

These coordinates can be chosen such that the extended-unstable manifold $W_{loc}^{uE}(O_1)$ is tangent to $\{z = 0\}$ when $x = 0$ and $z = 0$.

Similarly, we introduce C^r -coordinates $(u, v, w) \in \mathbb{R}^{d^{cu}} \times \mathbb{R}^{d-d_2} \times \mathbb{R}^{d_2-d^{cu}}$ in U_{O_2} such that the local invariant manifolds as well as the leaves of the foliation \mathcal{F}^{uu} are straightened, so we have

$$W_{loc}^s(O_2) = \{u = 0, w = 0\}, \quad W_{loc}^u(O_2) = \{v = 0\}, \quad W_{loc}^{uu}(O_2) = \{u = 0, v = 0\}.$$

These coordinates can be chosen such that the extended-sable manifold $W_{loc}^{sE}(O_2)$ is tangent to $\{w = 0\}$ when $u = 0$ and $v = 0$.

With [19, Lemmas 5 and 6], the coordinates can be chosen to further satisfy that the restrictions

$$F_1|_{W_{loc}^s(O_1)} : (x, z) \mapsto (\bar{x}, \bar{z}) \quad \text{and} \quad F_2|_{W_{loc}^u(O_2)} : (u, w) \mapsto (\bar{u}, \bar{w})$$

are linear in the center-stable and, respectively, center-unstable coordinates:

$$\begin{aligned} \text{Saddle-focus case:} \quad & \bar{x} = \lambda \begin{pmatrix} \cos \omega & \sin \omega \\ -\sin \omega & \cos \omega \end{pmatrix} x \quad \text{and} \quad \bar{u} = \gamma u; \\ \text{Double-focus case:} \quad & \bar{x} = \lambda \begin{pmatrix} \cos \omega_1 & \sin \omega_1 \\ -\sin \omega_1 & \cos \omega_1 \end{pmatrix} x \quad \text{and} \quad \bar{u} = \gamma \begin{pmatrix} \cos \omega_2 & \sin \omega_2 \\ -\sin \omega_2 & \cos \omega_2 \end{pmatrix} u. \end{aligned} \tag{4}$$

Recall that for saddle-focus cycles we omit the case where the center-stable multiplier is real but the center-unstable multipliers are nonreal, since it can be reduced to the above case by considering f^{-1} . In these coordinates, we have $M_1^+ = (x^+, 0, z^+)$ and $M_2^- = (u^-, 0, w^-)$, and hence condition **C3** reads

$x^+ \neq 0$ and $u^- \neq 0$.

Recall that the transition map F_{21} takes a small neighborhood of M_2^- to a small neighborhood of M_1^+ . By counting dimension, the two intersections $\ell_1 = W_{loc}^s(O_1) \cap F_{21}(W_{loc}^u(O_2))$ and $\ell_2 = F_{21}^{-1}\ell_1$ are two one-dimensional curves. Denote by $(dx, dz)_{M_1^+}$ the tangent vectors to ℓ_1 at the point M_1^+ , and by $(du, dw)_{M_2^-}$ the tangent vectors to ℓ_2 at the point M_2^- .

C4: when the center-stable multipliers are a pair of complex conjugate numbers (so $x \in \mathbb{R}^2$), the vector $dx_{M_1^+}$ is not parallel to the vector x^+ . When the center-unstable multipliers are a pair of complex conjugate numbers (so $u \in \mathbb{R}^2$), the vector $du_{M_2^-}$ is not parallel to the vector u^- .

This condition is independent of the choice of the coordinates which straighten the local invariant manifolds and foliations, and linearize the central dynamics as in (4). To see this, one notices that any transformation between two sets of such coordinates leaves the x - and u -plane invariant and restricts to a linear map on them. Condition **C4** is also independent of the choice of the points M_1^+ and M_2^- from Γ^1 , since replacing them by their iterates contribute the same factor to $dx_{M_1^+}$ and x^+ , and, respectively, to $du_{M_2^-}$ and u^- .

A saddle-focus or double-focus heterodimensional cycle is called *non-degenerate* if it satisfies conditions C1–C4, and any such cycle can be made non-degenerate by an arbitrarily C^r -small perturbation.

Remark 3.2. For heterodimensional cycles of saddle type, i.e., both center-stable and center-unstable multipliers are real, the first three non-degeneracy conditions are the same. The fourth condition is different. In this case, the central dynamics on $W_{loc}^s(O_1)$ and $W_{loc}^u(O_2)$ are given by $\bar{x} = \lambda x$, $\bar{u} = \gamma u$ with $x \in \mathbb{R}$ and $u \in \mathbb{R}$. By condition **C2**, the restriction $F_{21}|_{\ell_2} : \ell_2 \rightarrow \ell_1, (u, v, w) \mapsto (x, v, z)$ has the following non-zero derivative:

$$b^* = \left. \frac{\partial x}{\partial u} \right|_{M_2^-}.$$

The fourth condition is $|b^*u^-/x^+| \neq 1$ (which is condition C4.1 in [23, Section 2.3]).

3.2 Saddle-focus case

The first return map of a saddle-focus heterodimensional cycle is defined as the composition $T_{k,m} := F_{21} \circ F_2^m \circ F_{12} \circ F_1^k$, which, for suitable choices of (k, m) , takes points in a small neighborhood of M_1^+ back to the neighborhood. It is shown in [23, Sections 6.1.1–6.1.3] that there exist coordinates $(X, Y, Z) \in \mathbb{R} \times \mathbb{R}^{d_1} \times \mathbb{R}^{d-d_1-1}$ near M_1^+ such that the first return map can be expressed in a cross-form which manifests the partial hyperbolicity for a collection of pairs (k, m) . In what follows we introduce this formula.

Recall that we denote the pair of complex conjugate center-stable multipliers by $\lambda e^{\pm i\omega}$ ($0 < \omega < \pi$) and the real center-unstable multiplier by γ . The coordinates (X, Y, Z) are obtained from those in (4), and a small neighborhood of M_1^+ in the new coordinates is given by the cube

$$\Pi = [-\delta, \delta] \times [-\delta, \delta]^{d_1} \times [-\delta, \delta]^{d-d_1-1}, \quad (5)$$

where $d_1 = \text{ind}(O_1)$. It is proven in [23] that there exist coefficients $A, B, b, \eta_{1,2}$ (which are expressed in terms of the derivatives $DF_{12}|_{M_1^-}$ and $DF_{21}|_{M_2^+}$) and a positive number $C(\delta)$ depending on δ such

that, for all sufficiently large k and m satisfying

$$|\lambda^k \gamma^m B \sin(k\omega + \eta_2) - bu^-| < C(\delta), \quad (6)$$

(where u^- is the u -coordinate of M_2^- given in (4)), one has that $(\bar{X}, \bar{Y}, \bar{Z}) := T_{k,m}(X, Y, Z) \in \Pi$ for a point $(X, Y, Z) \in \Pi$ if and only if ⁵

$$\begin{aligned} \bar{X} &= A_{k,m}X + B_{k,m} + \phi_1(X, \bar{Y}, Z), \\ Y &= \phi_3(X, \bar{Y}, Z), \quad \bar{Z} = \phi_4(X, \bar{Y}, Z), \end{aligned} \quad (7)$$

where the functions ϕ satisfy

$$\phi_1 = o(\delta) + o(1)_{k,m \rightarrow \infty}, \quad \frac{\partial \phi_1}{\partial (X, \bar{Y}, Z)} = o(1)_{\delta \rightarrow 0} + o(1)_{k,m \rightarrow \infty}, \quad \|\phi_{2,3}\|_{C^1} = o(1)_{k,m \rightarrow \infty},$$

and

$$A_{k,m} = \lambda^k \gamma^m A \sin(k\omega + \eta_1), \quad B_{k,m} = \lambda^k \gamma^m B \sin(k\omega + \eta_2) - bu^-. \quad (8)$$

The non-degeneracy conditions ensure (see [23, Section 6.1.3]) that $A > 0, B > 0, b \neq 0, u^- \neq 0$ and

$$\tan \eta_1 \neq \tan \eta_2. \quad (9)$$

The partial hyperbolicity of $T_{k,m}$ follows immediately from a computation of the invariant cone fields using (7). Denote by $(\Delta X, \Delta Y, \Delta Z)$ the vectors in the tangent spaces.

Lemma 3.3 ([23, Lemma 6.1]). *Given any $K > 0$, one can choose δ sufficiently small such that, for all sufficiently large k, m for which $\sin(k\omega + \eta_1)$ and $\sin(k\omega + \eta_2)$ are bounded away from zero, the cone fields*

$$\begin{aligned} \mathcal{C}^{cu} &= \{(\Delta X, \Delta Y, \Delta Z) : \|\Delta Z\| \leq K(|\Delta X| + \|\Delta Y\|)\}, \\ \mathcal{C}^{uu} &= \{(\Delta X, \Delta Y, \Delta Z) : \max\{|\Delta X|, \|\Delta Z\|\} \leq K\|\Delta Y\|\}, \end{aligned} \quad (10)$$

which are defined on Π , are forward-invariant in the sense that if a point $M \in \Pi$ has its image $\bar{M} = T_{k,m}(M)$ in Π , then the cone at M is mapped strictly into the cone at \bar{M} by $DT_{k,m}$; the cone fields

$$\begin{aligned} \mathcal{C}^{cs} &= \{(\Delta X, \Delta Y, \Delta Z) : \|\Delta Y\| \leq K(|\Delta X| + \|\Delta Z\|)\}, \\ \mathcal{C}^{ss} &= \{(\Delta X, \Delta Y, \Delta Z) : \max\{|\Delta X|, \|\Delta Y\|\} \leq K\|\Delta Z\|\}, \end{aligned} \quad (11)$$

which are defined on Π , are backward-invariant in the sense that if a point $\bar{M} \in \Pi$ has its preimage $M = T_{k,m}^{-1}(\bar{M})$ in Π , then the cone at \bar{M} is mapped strictly into the cone at M by $DT_{k,m}^{-1}$. Moreover, vectors in \mathcal{C}^{uu} and, if $|A_{k,m}| > 1$, also vectors in \mathcal{C}^{cu} are expanded by $DT_{k,m}$; vectors in \mathcal{C}^{ss} and, if $|A_{k,m}| < 1$, also vectors in \mathcal{C}^{cs} are contracted by $DT_{k,m}$.

3.3 Double-focus case

In this case, we consider the first return map $T_{k,m} := F_2^m \circ F_{12} \circ F_1^k \circ F_{21}$ near M_2^- . A cross-form formula for $T_{k,m}$ is obtained in [23, Sections 6.4.1–6.4.3] with coordinates $(U, V, W) \in \mathbb{R} \times \mathbb{R}^{d-d_2} \times \mathbb{R}^{d_2-1}$

⁵ Equation (7) is (6.18) in [23] with dropping the subscript of X_1 and grouping the coordinates $(X_2, Z) =: Z$.

in a small neighborhood M_2^- , which we describe below. For simplicity, many coefficients here will carry the same notations as those in the saddle-focus case, but they are different coefficients. This will not cause ambiguity as we never discuss the two cases at the same time in the proofs.

Recall that, for a double-focus heterodimensional cycle, we denote the center-stable multipliers by $\lambda e^{\pm i\omega_1}$ ($0 < \omega_1 < \pi$) and the center-unstable multipliers by $\gamma e^{\pm i\omega_2}$ ($0 < \omega_2 < \pi$). The coordinates (U, V, W) are obtained from those in (4), and a small neighborhood of M_2^- in the new coordinates is given by the cube

$$\Pi = [-\delta, \delta] \times [-\delta, \delta]^{d-d_2} \times [-\delta, \delta]^{d_2-1}, \quad (12)$$

where $d_2 = \text{ind}(O_2)$. It is proven that there exist coefficients $A^*, B^*, b, \eta_{1,2,3}$ (which are expressed in terms derivatives $DF_{12}|_{M_1^-}$ and $DF_{21}|_{M_2^+}$) and a positive number $c(\delta) = o(1)_{\delta \rightarrow 0}$ depending on δ such that, for all sufficiently large k and m with m satisfying

$$\min\{|\cos m\omega_2|, |\cos(m\omega_2 + \eta_3)|, |\cos(m\omega_2 + \eta_3) + b \sin(m\omega_2 + \eta_3)|\} > c(\delta), \quad (13)$$

one has that $(\bar{U}, \bar{V}, \bar{W}) := F_{k,m}(U, V, W) \in \Pi$ for a point $(U, V, W) \in \Pi$ if and only if ⁶

$$\begin{aligned} \bar{U} &= A_{k,m}U + B_{k,m} + \phi_1(U, V, \bar{W}), \\ \bar{V} &= \phi_2(U, V, \bar{W}), \quad W = \phi_3(U, V, \bar{W}), \end{aligned} \quad (14)$$

where the functions ϕ satisfy

$$\phi_1 = o(\delta) + o(1)_{k,m \rightarrow \infty}, \quad \frac{\partial \phi_1}{\partial(U, V, \bar{W})} = o(1)_{\delta \rightarrow 0} + o(1)_{k,m \rightarrow \infty}, \quad \|\phi_{2,3}\|_{C^1} = o(\delta) + o(1)_{k,m \rightarrow \infty},$$

and

$$A_{k,m} = \lambda^k \gamma^m A(m\omega_2) \sin(k\omega_1 + \eta_1), \quad B_{k,m} = \lambda^k \gamma^m B(m\omega_2) \sin(k\omega_1 + \eta_2) - \xi(m\omega_2) \quad (15)$$

with

$$\begin{aligned} A(m\omega_2) &= \frac{A^*}{\cos(m\omega_2 + \eta_3) + b \sin(m\omega_2 + \eta_3)}, \quad B(m\omega_2) = \frac{B^*}{\cos(m\omega_2 + \eta_3) + b \sin(m\omega_2 + \eta_3)}, \\ \xi(m\omega_2) &= \frac{\cos(m\omega_2 + \eta_3)u_1^- + \sin(m\omega_2 + \eta_3)u_2^-}{\cos(m\omega_2 + \eta_3) + b \sin(m\omega_2 + \eta_3)}. \end{aligned} \quad (16)$$

Here $u^- = (u_1^-, u_2^-)$ are the u -coordinates of M_2^- in the coordinates (4). The non-degeneracy conditions ensure that $A^* > 0, B^* > 0, b \neq 0, u^- \neq 0$ and $\tan \eta_1 \neq \tan \eta_2$, see [23, Section 6.4.3].

One sees that formula (14) assumes the same form as (7), with replacing U, V, W by X, Z, Y , respectively. By a completely parallel computation, one finds the following counterpart to Lemma 3.3, which characterizes the partial hyperbolicity of $T_{k,m}$.

Lemma 3.4. *Given any $K > 0$, one can choose δ sufficiently small, such that for all sufficiently large k, m satisfying (13) and (32), the cone fields*

$$\begin{aligned} \mathcal{C}^{cu} &= \{(\Delta U, \Delta V, \Delta W) : \|\Delta V\| \leq K(|\Delta U| + \|\Delta W\|)\}, \\ \mathcal{C}^{uu} &= \{(\Delta U, \Delta V, \Delta W) : \max\{|\Delta U|, \|\Delta V\|\} \leq K\|\Delta W\|\}, \end{aligned} \quad (17)$$

⁶ Equation (14) is (6.44) in [23] with dropping the subscript of U_1 , grouping the coordinates $(U_2, W) =: W$, and dividing $(1 + b_{41} \tan(m\omega + \eta_3))$ to both sides of (6.44).

which are defined on Π , are forward-invariant in the sense that if a point $M \in \Pi$ has its image $\bar{M} = T_{k,m}(M)$ in Π , then the cone at M is mapped into the cone at \bar{M} by $DT_{k,m}$; the cone fields

$$\begin{aligned} \mathcal{C}^{cs} &= \{(\Delta U, \Delta V, \Delta W) : \|\Delta W\| \leq K(|\Delta U| + \|\Delta V\|)\}, \\ \mathcal{C}^{ss} &= \{(\Delta U, \Delta V, \Delta W) : \max\{|\Delta U|, \|\Delta W\|\} \leq K\|\Delta V\|\}, \end{aligned} \quad (18)$$

which are defined on Π , are backward-invariant in the sense that if a point $\bar{M} \in \Pi$ has its preimage $M = T_{k,m}^{-1}(\bar{M})$ in Π , then the cone at \bar{M} is mapped into the cone at M by $DT_{k,m}^{-1}$. Moreover, vectors in \mathcal{C}^{uu} and, if $|A_{k,m}| > 1$, also vectors in \mathcal{C}^{cu} are expanded by $DT_{k,m}$; vectors in \mathcal{C}^{ss} and, if $|A_{k,m}| < 1$, also vectors in \mathcal{C}^{cs} are contracted by $DT_{k,m}$.

Remark 3.5. The results in this paper are solely based on the analysis of the first return maps, so they can be transferred easily to the continuous-time case, i.e., when f is a flow on any manifold of dimension at least four. Indeed, one just needs to define F_i for each $i = 1, 2$ as the Poincaré map of a local transverse cross-section at some point O_i of the periodic orbit, and define F_{12} by the trajectories of the flow from a neighborhood of M_1^- to a neighborhood of M_2^+ , and similarly for F_{21} .

4 Blenders near saddle-focus heterodimensional cycles

In this section, we prove Theorem 1 and the first case in Theorem 3. Both theorems can be divided into two parts, where one concerns the existence of blenders and the other concerns the homoclinic relations. We will obtain results for these two parts separately, and then prove the theorems. For the blender part, we use a criterion which summarizes the construction in [23, Section 4.2].

4.1 A criterion for the existence of blenders

Let us first introduce the notion of *activating pair* to further characterize the embedding set \mathcal{D} in Definition 1.1. Take a cube

$$Q = \{(x_1, \dots, x_d) \mid x_i \in I_i\} \subset \mathbb{R}^d,$$

where I_i are closed intervals in \mathbb{R} . A k -dimensional disc S is said to *cross* Q if the intersection $S \cap Q$ is given by $(x_{i_{k+1}}, \dots, x_{i_d}) = s(x_{i_1}, \dots, x_{i_k})$, where (i_1, \dots, i_d) is some permutation of $(1, \dots, d)$, and s is a smooth function defined on $I_{i_1} \times \dots \times I_{i_k}$. It crosses Q *properly* with respect to a cone field \mathcal{C} defined on Q if the tangent spaces of $S \cap Q$ lie in \mathcal{C} , i.e., the tangent spaces of S lie in \mathcal{C} .

For an invariant cone field \mathcal{C} , we use $\dim(\mathcal{C})$ to denote the largest dimension of linear subspaces of the tangent spaces contained in \mathcal{C} .

Definition 4.1 (Activating pair). A pair (Q, \mathcal{C}) consisting of a cube and a cone field is called an activating pair for a cs-blender Λ if $\dim(\mathcal{C}) = \dim W^s(\Lambda) - 1$ and if any surface S crossing Q properly with respect to \mathcal{C} intersects $W^u(\Lambda)$. The pair (Q, \mathcal{C}) is an activating pair for a cu-blender Λ if $\dim(\mathcal{C}) = \dim W^u(\Lambda) - 1$ and any surface S crossing Q properly with respect to \mathcal{C} intersects $W^s(\Lambda)$. The cube Q is called an activating domain.

For example, if (Π, \mathcal{C}^{ss}) , with Π in (5) and \mathcal{C}^{ss} in (11), is an activating pair of some cs-blender, then all curves crossing Π properly with respect to \mathcal{C}^{ss} belong to the embedding set \mathcal{D} in Definition 1.1.

Each such curve is the graph of some smooth function $(X, Y) = s(Z)$, which is defined on $[-\delta, \delta]^{d-d_1-1}$ and whose tangent spaces lie in \mathcal{C}^{ss} .

Let $\delta > 0$ and take a cube

$$Q = [-\delta, \delta] \times [-\delta, \delta]^{d_Y} \times [-\delta, \delta]^{d_Z}$$

with $1 + d_Y + d_Z = d$. Let $\{T_n\}_{n \in \mathbb{N}}$ be a sequence of maps defined on pair wise disjoint subsets $\{\sigma_n\}$ of Q such that $T_i(\sigma_i) \cap T_j(\sigma_j) \cap Q = \emptyset$ for $i \neq j$. Define $T : \bigcup_n \sigma_n \rightarrow \mathbb{R}^d$ by $T(X, Y, Z) = T_n(X, Y, Z)$ if $(X, Y, Z) \in \sigma_n$. Suppose for every $n \in \mathbb{N}$ and $(X, Y, Z) \in \Pi$, one has that $(\bar{X}, \bar{Y}, \bar{Z}) = T_n(X, Y, Z)$ if and only if

$$\begin{aligned} \bar{X} &= A_n X + B_n + \phi_1(X, \bar{Y}, Z; n), \\ Y &= \phi_2(X, \bar{Y}, Z; n), \\ \bar{Z} &= \phi_3(X, \bar{Y}, Z; n), \end{aligned} \tag{19}$$

where the constants A_n and B_n , and the C^1 functions ϕ defined on Q are allowed to depend on δ , and we have the estimates

$$\phi_1 = o(\delta) + o(1)_{n \rightarrow \infty}, \quad \frac{\partial \phi_1}{\partial (X, \bar{Y}, Z)} = o(1)_{\delta \rightarrow 0} + o(1)_{n \rightarrow \infty}, \quad \|\phi_{2,3}\|_{C^1} = o(1)_{\delta \rightarrow 0} + o(1)_{n \rightarrow \infty}.$$

Proposition 4.2 ([23, Proposition 4.12]). *Suppose there exists $\delta_0 > 0$ such that the constants A_n and B_n satisfy that*

1. *for each $\delta \in (0, \delta_0)$, the set $\{B_n\}$ is dense in an interval I which contains zero and is independent of δ ;*
2. *for all $\delta \in (0, \delta_0)$ and all $n \in \mathbb{N}$,*
 - 2.1. *either $0 < C_1 < |A_{k_n, m_n}| < C_2 < 1$ with $C_{1,2}$ independent of n ,*
 - 2.2. *or $1 < C_3 < |A_{k_n, m_n}| < C_4 < \infty$ with $C_{3,4}$ independent of n .*

Then, there exist $c \in (0, 1)$ and

$$Q' = [-c\delta, c\delta] \times [-\delta, \delta]^{d_Y} \times [-\delta, \delta]^{d_Z}, \tag{20}$$

such that for every $\delta \in (0, \delta_0)$ one can find a finite set $\mathcal{N} \subset \mathbb{N}$ with arbitrarily large elements such that the restriction of T to $\bigcup_{n \in \mathcal{N}} \sigma_n \cap Q'$ has a blender with an activating pair (Q', \mathcal{C}) , where the cone field \mathcal{C} can be taken independent of δ . Moreover,

- *if condition 2.1 holds, Λ is a cs-blender of index d_Y and cones in \mathcal{C} are around Z -coordinates;*
- *if condition 2.2 holds, Λ is a cu-blender of index $(d_Y + 1)$ and cones in \mathcal{C} are around Y -coordinates.*

In our case, the maps T_n will be the return maps T_{k_n, m_n} for some sequence of pairs satisfying conditions 1 and 2. For completeness we sketch the proof of this proposition.

Sketch of proof. Let us first prove the proposition when condition 2.1 is satisfied. The map $T_n^\times : (X, \bar{Y}, Z) \mapsto (\bar{X}, Y, \bar{Z})$ given by (19) is a contraction. Take a finite set $\mathcal{N} \subset \mathbb{N}$ such that T_n^\times takes Q into itself for all $n \in \mathcal{N}$. Then, by a fixed point lemma for a direct product of metric spaces (see e.g. Theorem 6.2 in [31]), there is a one-to-one correspondence between the set $\Sigma := \{0, \dots, \text{Card } \mathcal{N} - 1\}^{\mathbb{Z}}$ of two-sided sequences of $\text{Card } \mathcal{N}$ symbols and the set $\hat{\Lambda}$ of points whose orbits under the iterated

function system $\{T_n\}_{n \in \mathcal{N}}$ never leave Q . This shows that $\hat{\Lambda}$ is a compact, transitive, locally maximal invariant set of T . Evidently from formula (19), $\hat{\Lambda}$ is also hyperbolic.

Let $(\Delta X, \Delta Y, \Delta Z)$ denote vectors in tangent spaces. Given any constant $K > 0$, it is easy to check that, by taking δ sufficiently small and n sufficiently large, the cone field defined by

$$\mathcal{C}^{ss} = \{(\Delta X, \Delta Y, \Delta Z) : |\Delta X| + \|\Delta Y\| < K\|Z\|\}, \quad (21)$$

is backward invariant in the sense that for any $(X, Y, Z) \in \Pi$ if $T_n(X, Y, Z) = (\bar{X}, \bar{Y}, \bar{Z}) \in \Pi$ for some $n \in \mathcal{N}$, then the cone at $(\bar{X}, \bar{Y}, \bar{Z})$ is mapped inside the one at (X, Y, Z) by DT_n^{-1} .

By assumption 1, there exists $c \in (0, 1)$ such that B_n is dense in $[-c\delta, c\delta]$. Let Q' be defined as in (20). Consider the open set \mathcal{S} of discs S of the form $(X, Y) = s(Z)$ for some smooth function s , which crosses Q' properly with respect to \mathcal{C}^{ss} . We will prove that the set $\Lambda = \hat{\Lambda} \cap Q'$ is a cs-blender, which amounts to showing that $S \cap W^u(\Lambda) \neq \emptyset$ for every $S \in \mathcal{S}$. By construction, the local unstable manifold of Λ is the set of points in Q whose backward orbits under $\{T_n\}_{n \in \mathcal{N}}$ stay in Q' . Thus, it suffices to prove the so-called *covering property* that for any $S \in \mathcal{S}$ there exists $n \in \mathcal{N}$ such that the intersection $T_n^{-1}(S) \cap Q'$ contains a piece in \mathcal{S} . Indeed, this gives a sequence of positive integers $\{n_i \in \mathcal{N}\}$ and a sequence of nested compact sets $\{\hat{S}_i \subset S\}$ such that each preimage $T_{n_i}(\hat{S}_i)$ is a disc in \mathcal{S} , which particularly implies that $\bigcap \hat{S}_i$ contains a point whose backward orbit under $\{T_n\}_{n \in \mathcal{N}}$ lies in Q .

Let us prove the covering property. Due to the backward invariance of \mathcal{C}^{ss} , one only needs to check that $T_n^{-1}(S) \cap Q$ has a connected component crossing Q . Also observe that by the strong contraction in Z and strong expansion in Y , it suffices to show that, for any $S \in \mathcal{S}$, there exists $n \in \mathcal{N}$ such that the X -coordinates of points of $T_n^{-1}(S)$ lie in $I := [-c\delta, c\delta]$. This is equivalent to prove that

- for $n \in \mathcal{N}$, the union of the images of the interior of I under the affine maps $g_n : \bar{X} = A_n X + B_n$ covers a neighborhood of I , and that
- the overlaps between $g_n(I)$ are large enough so that there exists some $n^* \in \mathcal{N}$ such that the X -coordinates of points of S lie completely in $g_{n^*}(I)$.

To find a set \mathcal{N} satisfying the above conditions, we note that the center of $g_n(I)$ is B_n and the length of it is greater than some constant $L_1 > 0$. Since $\{B_n\}$ is dense in I , it follows that there exist finitely many n such that the union of the $L_1/2$ -neighborhoods of B_n covers a neighborhood of I . The overlap requirement can be satisfied by taking K in (21) sufficiently small so that the total variation of X -coordinates of S is sufficiently small. The covering property is proven.

Finally, one sees that the blender found above can be arbitrarily close to the origin by taking the integers in \mathcal{N} sufficiently large. This completes the proof when condition 2.1 holds.

When condition 3.1 holds, define $\hat{T}_n := T_n^{-1}|_{T_n(\sigma_n) \cap Q}$. The maps \hat{T}_n assume the same form as (19) but with new A_n whose absolute values are uniformly bounded away from 0 and 1. Then, apply the above arguments to $\{\hat{T}_n\}$. \square

4.2 Existence of blenders when at least one of θ and $\frac{\omega}{2\pi}$ is irrational

Recall that for a saddle-focus heterodimensional cycle, we denote the complex center-stable multipliers by $\lambda e^{\pm i\omega}$, where $\lambda > 0$ and $0 < \omega < \pi$, and the real center-unstable multiplier by γ . Let $A > 0, B > 0, b, u^-, \eta_{1,2}$ be the coefficients in (8). We introduce a condition on the modulus ω .

A1: there exist infinitely many $k > 0$ such that

$$bu^- \sin(k\omega + \eta_2) > 0, \quad \sin(k\omega + \eta_1) \neq 0, \quad \alpha_k := \frac{bu^- A \sin(k\omega + \eta_1)}{\sin(k\omega + \eta_2)} \neq \pm 1. \quad (22)$$

Observe that condition **A1** is automatically satisfied if $\omega/2\pi$ is irrational, or if $\omega/2\pi = p/q$ with $q \geq 9$. To see the latter, note first that the last inequality in (22) can be rewritten as

$$|bu^- \sin(k\omega + \eta_1)| \neq |\sin(k\omega + \eta_1) \cos(\eta_2 - \eta_1) + \sin(k\omega + \eta_1) \sin(\eta_2 - \eta_1)|,$$

or

$$|(bu^- - \cos(\eta_2 - \eta_1)) \tan(k\omega + \eta_1)| \neq |\sin(\eta_2 - \eta_1)|.$$

It follows that to achieve simultaneously this inequality and the second one in (22) it is enough to have at least 4 different values of $k\omega \bmod 2\pi$ in $[0, 1/2)$ or in $[1/2, 0)$. It is easy to check that, when $\omega/2\pi = p/q$, there are at least $\lfloor (q-1)/2 \rfloor$ different values of $k\omega \bmod 2\pi$ both in $[0, 1/2)$ and in $[1/2, 0)$, where $\lfloor a \rfloor$ means the largest integer which is smaller than or equal to a . Choosing the values from the suitable interval also gives $bu^- \sin(k\omega + \eta_2) > 0$. Thus, when $\lfloor (q-1)/2 \rfloor \geq 4$ or $q \geq 9$, there exists some $r \in \{0, \dots, q-1\}$ such that all $k = r + iq$ with $i = 0, 1, 2, \dots$ satisfy the inequalities in (22).

Proposition 4.3. *Let δ be the size of Π defined in (5), and \mathcal{C}^{ss} and \mathcal{C}^{uu} be given by Lemma 3.3 with a sufficiently small K . Let a non-degenerate saddle-focus heterodimensional cycle satisfy condition **A1** and U be any neighborhood of the cycle.*

1. *If $\theta = \ln \lambda / \ln |\gamma|$ is irrational and $\omega/2\pi = p/q$, then, for every sufficiently large k satisfying condition **A1**, there exists a cs-blender in U when $|\alpha_k| < 1$ (defined in (22)) or a cu-blender in U when $|\alpha_k| > 1$.*
2. *If $\omega/2\pi$ is irrational, and $\theta, \omega/2\pi$ and 1 are not rationally independent, then there exists a cu-blender in U .*
3. *If $\theta, \omega/2\pi$ and 1 are rationally independent, then there exist simultaneously a cs-blender and a cu-blender in U .*

In each of the above cases, the cs-blender has index d_1 and an activating pair (Π', \mathcal{C}^{ss}) , while the cu-blender has index $(d_2 = d_1 + 1)$ and an activating pair (Π', \mathcal{C}^{uu}) , where the cube Π' is defined as

$$\Pi' = [-c\delta, c\delta] \times [-\delta, \delta]^{d_1} \times [-\delta, \delta]^{d-d_1-1} \subset \Pi, \quad (23)$$

for some $c \in (0, 1)$.

Proof. We prove the proposition by finding in each case a sequence of pairs $\{(k_n, m_n)\}$ such that the sequence of return maps $\{T_{k_n, m_n}\}$ satisfies the conditions of Proposition 4.2. Let $A > 0, B > 0, b, u^-, \eta_{1,2}$ be the coefficients in (7).

(Case 1: θ is irrational and $\omega/2\pi$ is rational.) Take $\omega/2\pi = p/q$. Since condition **A1** is satisfied, there exists $k = r \in \{0, \dots, q-1\}$ satisfying (22). We particularly have

$$\sin\left(\frac{2\pi r p}{q} + \eta_2\right) B b u^- =: R B b u^- > 0. \quad (24)$$

Since θ is irrational, the set

$$\{m - (kq + r)\theta\}_{k \in \mathbb{N}, m \in 2\mathbb{N}}$$

is dense in \mathbb{R} , and, hence, $\{\gamma^{m-(kq+r)\theta}\}_{k \in \mathbb{N}, m \in 2\mathbb{N}}$ is dense in $[0, \infty]$. This along with (24) implies that one can find a sequence $\{(k'_n q + r, m'_n)\}$ such that $\{\gamma^{m'_n - (k'_n q + r)\theta}\}$ is dense in a small neighborhood of bu^-/BR . Thus, the sequence

$$\{|\gamma|^{m'_n - (k'_n q + r)\theta} B \sin((k'_n q + r)\omega + \eta_2) - bu^-\} \quad (25)$$

is dense in a small neighborhood Δ of zero.

Recall $\theta = -\ln \lambda / \ln |\gamma|$ (note that $\lambda > 0$). When the integer m is even, we can write

$$\lambda^k \gamma^m = |\gamma|^{m - \theta k}. \quad (26)$$

Now denote $k_n = k'_n q + r$ and $m_n = m'_n$. The preceding paragraph shows that $\sin(k_n \omega + \eta_2)$ is bounded away from zero. Also, since $\lambda > 0$ and m_n are even, relation (26) implies that $\lambda^{k_n} \gamma^{m_n} = \gamma^{m_n - k_n \theta}$. It follows that by taking Δ sufficiently small, the condition (6) is satisfied with $(k, m) = (k_n, m_n)$. Hence, the first return maps $T_n := T_{k_n, m_n}$ take the form (7). By comparing (8) with (25) we find that the sequence (25) is exactly the constant term B_{k_n, m_n} in (7). Thus, setting $X = X_1$ and $Z^{new} = (X_2, Z)$, we find that the sequence $\{T_n\}$ satisfies condition 1 of Proposition 4.2.

Next, let us check condition 2 of Proposition 4.2. To find the values of $|A_{k_n, m_n}|$, we use (8) to write

$$A_{k_n, m_n} = \frac{(B_{k_n, m_n} + bu^-) A \sin(k_n \omega + \eta_1)}{\sin(k_n \omega + \eta_2)}. \quad (27)$$

On the one hand, by taking Δ sufficiently small the contribution of B_{k_n, m_n} to A_{k_n, m_n} can be ignored. On the other hand, by the choice of k_n one has $\sin(k_n \omega + \eta_1) = \sin(r\omega + \eta_1)$ and $\sin(k_n \omega + \eta_2) = \sin(r\omega + \eta_2)$. Thus, we have

$$A_{k_n, m_n} \sim \frac{bu^- A \sin(r\omega + \eta_1)}{\sin(r\omega + \eta_2)} =: \alpha_r, \quad (28)$$

Since assumption (22) ensures $|\alpha_r| \neq 1$, this gives condition 2.1 of Proposition 4.2 if $|\alpha_r| < 1$, or condition 2.2 if $|\alpha_r| > 1$. Therefore, using Proposition 4.2 we obtain in the former case a cs-blender with an activating pair (Π', \mathcal{C}_1) for some cone field \mathcal{C}_1 , and in the latter case a cu-blender with an activating pair (Π', \mathcal{C}_2) for some cone field \mathcal{C}_2 . According to the proof of Proposition 4.2, these cone fields \mathcal{C}_1 and \mathcal{C}_2 are just \mathcal{C}^{ss} and \mathcal{C}^{uu} in Lemma 3.3, respectively.

(Case 2: $\omega/2\pi$ is irrational, and $\theta, \omega/2\pi$ and 1 are rationally dependent.) Since $\omega/2\pi$ is irrational, the rational dependence allows us to write

$$\theta = \frac{p}{q} \frac{\omega}{2\pi} + \frac{p'}{q'}.$$

Consider the function

$$h(s) = |\gamma|^{-\frac{p}{q}s} B \sin(2\pi s + \eta_2) - bu^-,$$

which is defined on \mathbb{R} . Let $s = s_0$ be a zero of h , i.e., $h(s_0) = 0$. It is obvious that s_0 can be taken with arbitrarily large absolute value. Since the set

$$\left\{ qq'k \frac{\omega}{2\pi} - qq'l \right\}_{k, l \in 2\mathbb{N}}$$

is dense in \mathbb{R} , one can find a sequence $\{qq'k'_n\omega/2\pi - qq'l_n\}$ dense in a small neighborhood of s_0 such that, by continuity of $h(s)$, the sequence

$$\{|\gamma|^{m_0 - \frac{p}{q}(qq'k'_n\frac{\omega}{2\pi} - qq'l_n)} B \sin(qq'k'_n\omega + \eta_2) - bu^-\} \quad (29)$$

is dense in a small neighborhood Δ of zero.

By (26) and (A1), for even m we have

$$\lambda^k \gamma^m = |\gamma|^{m - k\frac{p}{q}\frac{\omega}{2\pi} - k\frac{p'}{q'}}.$$

Taking $k_n = qq'k'_n$ and $m_n = p'qk'_n + pq'l_n$, one readily finds that B_{k_n, m_n} in (7) equals (29). Arguing as in case (1), we obtain condition 1 of Proposition 4.2 with $\{T_n := T_{k_n, m_n}\}$.

It is immediate from (29) that $\sin(k_n\omega + \eta_2)$ can be arbitrarily close to zero by taking $|s_0|$ sufficiently large. Since B_{k_n, m_n} is negligible by taking Δ small, it follows from (9) and (27) that $|A_{k_n, m_n}| > 1$ for all n , and, particularly, they are bounded away from 1 and infinity. So, applying Proposition 4.2 with condition 2.2 gives a cu-blender. The activating pair is obtained as in case 1.

(Case 3: $\theta, \omega/2\pi$ and 1 are rationally independent.) The result is an easy consequence of the denseness of the set

$$\left\{ \left(k\theta - m, k\frac{\omega}{2\pi} - l \right) \right\}_{k \in \mathbb{N}, m \in 2\mathbb{N}, l \in \mathbb{Z}}$$

in \mathbb{R}^2 . Indeed, one just needs to choose a sequence $\{(k_n, m_n)\}$ satisfying that $\sin(k_n\omega + \eta_2)$ is dense in a small interval Δ' and $\lambda^{k_n} \gamma^{m_n} = |\gamma|^{m_n - k_n\theta} \rightarrow t$ for some constant t such that

$$\{B_{k_n, m_n} = |\gamma|^{m_n - k_n\theta} B \sin(k_n\omega + \eta_2) - bu^-\}$$

is dense in a small neighborhood Δ of zero. The latter fact gives condition 1 of Proposition 4.2. The former one implies that, with sufficiently small Δ , the values $|A_{k_n, m_n}|$ in (27) can lie between zero and 1 by taking Δ' close to zero, which leads to a cs-blender, or lie between 1 and infinity by taking Δ' such that $\sin(k_n\omega + \eta_2)$ is close to zero, which leads to a cu-blender. The activating pair is obtained as in case 1. \square

4.3 Homoclinic relations in the saddle-focus case

We first make some observations on the blenders obtained in Section 4.2. It follows from the proof of Proposition 4.3 that if a cs-blender Λ_1 corresponds to a finite set \mathcal{P}_1 of pairs (k, m) such that $0 < C_1 < |A_{k, m}| < C_2 < 1$ for $(k, m) \in \mathcal{P}_1$, where $A_{k, m}$ is as in (7) and $C_{1,2}$ are some constants. Similarly, a cu-blender Λ_2 corresponds to a finite set \mathcal{P}_2 of pairs (k, m) such that $1 < C_3 < |A_{k, m}| < C_4 < \infty$ for $(k, m) \in \mathcal{P}_2$, where $C_{3,4}$ are some constants. It follows from the sketch proof of Proposition 4.2 that the sets Λ_i ($i = 1, 2$) consist of points whose orbits under $T_{k, m}$ with $(k, m) \in \mathcal{P}_i$ never leave the cube Π defined in (5) and the activating domain is Π' defined in (23). These sets are hyperbolic basic sets equipped with the cone fields in Lemma 3.3.

For each $i = 1, 2$ and a point $M_i \in \Lambda_i \cap \Pi'$, define its local invariant manifolds $W_{loc}^s(M_i)$ and $W_{loc}^u(M_i)$ as the connected components through M_i of $W_{loc}^s(\Lambda_i) \cap \Pi$ and, respectively, of $W_{loc}^u(\Lambda_i) \cap \Pi$. One readily finds from Lemma 3.3 that those manifolds are given by smooth functions of the following forms:

- $W_{loc}^s(M_1) : Y = \psi_{M_1}^s(X, Z)$ defined on $[-\delta, \delta] \times [-\delta, \delta]^{d-d_1-1}$;
- $W_{loc}^u(M_1) : (X, Z) = \psi_{M_1}^u(Y)$ defined on $[-\delta, \delta]^{d_1}$, and, in particular, $W_{loc}^u(M_1)$ crosses Π' properly with respect to \mathcal{C}^{uu} defined in (10);
- $W_{loc}^s(M_2) : (X, Y) = \psi_{M_2}^s(Z)$ defined on $[-\delta, \delta]^{d-d_1-1}$, and, in particular, $W_{loc}^s(M_2)$ crosses Π' properly with respect to \mathcal{C}^{ss} defined in (11);
- $W_{loc}^u(M_2) : Z = \psi_{M_2}^u(X, Y)$ defined on $[-\delta, \delta] \times [-\delta, \delta]^{d_1}$.

An immediate observation is that when a cs-blender and a cu-blender found in Proposition 4.3 coexist (i.e., in case 3), they activate each other in the sense of Definition 2.1.

We next collect some relevant results in [23] about the invariant manifolds of the periodic points O_1 and O_2 . It is shown in equation (3.23) of [23] that

$$W_{loc}^s(O_1) = \{Y = 0\} \quad \text{and} \quad F_{21}(W_{loc}^u(O_2)) = \{Z = 0\}. \quad (30)$$

Comparing these equations with those for $W_{loc}^u(M_1)$ and $W_{loc}^s(M_2)$, we obtain that

$$W^u(O_2) \pitchfork W^s(\Lambda_2) \neq \emptyset \quad \text{and} \quad W^s(O_1) \pitchfork W^u(\Lambda_1) \neq \emptyset. \quad (31)$$

One also has the following

Lemma 4.4 ([23, Lemma 6.5]). *If $\omega/2\pi$ is irrational, then there exist infinitely many k such that $F_1^{-k} \circ F_{12}^{-1}(W_{loc}^s(O_2))$ is a disc of the form*

$$(X, Y) = s(Z),$$

which crosses Π' defined by (23) properly with respect to \mathcal{C}^{ss} defined by (11).

Comparing the above two formulas with those for $W_{loc}^s(M_2)$ and $W_{loc}^u(M_2)$, one finds that O_2 is homoclinically related to the cu-blender found in Proposition 4.3 when $\omega/2\pi$ is irrational. In this case, it follows from the lambda-lemma that O_2 also activates the cs-blender if the cu-blender does. In summary, we obtain

Lemma 4.5. *The periodic points $O_{1,2}$ and the blenders found in Proposition 4.3 have the following homoclinic relations:*

- *in case 2, the cu-blender is homoclinically related to O_2 , and*
- *in case 3, the cu-blender is homoclinically related to O_2 ; the two blenders activate each other, and in particular O_2 activates the cs-blender.*

4.4 Proofs of Theorem 1 and the first case in Theorem 3

Theorem 1 follows immediately from Proposition 4.3 and Lemma 4.5.

Proof of the first case in Theorem 3. Since being non-degenerate is an open and dense property in \mathcal{H}_f , we take the neighborhood \mathcal{U} of g sufficiently small such that Γ_f is non-degenerate for all $f \in \mathcal{U}$. Take any $f \in \mathcal{U}$ with irrational $\theta(f)$ and rational $\omega(f)/2\pi = p/q$. According to the proof of Proposition

4.2, to obtain both a cs-blender and a cu-blender for f , it suffices to find $r_1, r_2 \in \{0, \dots, q-1\}$ such that $0 < |\alpha_{r_1}| < 1$ and $1 < |\alpha_{r_2}| < \infty$ (see (28)). Obviously, there exists N_f depending on $A, b, u^-, \eta_{1,2}$ such that if $q > N_f$ then one can always find such r_1 and r_2 . Since the cycle is non-degenerate for all $f \in (U)$, the involved coefficients are well defined and depend continuously on f . Thus, coexisting blenders appear whenever $q > N(\mathcal{U}) = \sup_{N \in \mathcal{U}} q_f$. The homoclinic relation follows from the formulas for the invariant manifolds of points of the blenders listed in Section 4.3. \square

5 Blenders near double-focus heterodimensional cycles

In this section, we prove Theorem 2 and the second case in Theorem 3, in the same fashion as for saddle-focus cycles. We first state the blender results and homoclinic relation results separately, and then prove the Theorems.

5.1 Existence of blenders when at least one of $\theta, \frac{\omega_1}{2\pi}, \frac{\omega_2}{2\pi}$ is irrational

Recall that for a double-focus heterodimensional cycle, we denote the center-stable multipliers by $\lambda e^{\pm i\omega_1}$ and the center-unstable multipliers by $\gamma e^{\pm i\omega_2}$, where $\lambda > 0, \gamma > 0$ and $0 < \omega_{1,2} < \pi$. Let $A^* > 0, B^* > 0, b, u^-, \eta_{1,2,3}$ be the coefficients in (15) and (16). We first introduce a condition on the moduli ω_1 and ω_2 .

A2: there exist infinitely many pairs (k, m) satisfying (13) and the inequalities

$$\begin{aligned} & (\cos(m\omega_2 + \eta_3)u_1^- + \sin(m\omega_2 + \eta_3)u_2^-) \sin(k\omega_1 + \eta_2) > 0, \quad \sin(k\omega_1 + \eta_1) \neq 0, \\ \alpha_{k,m} := & \frac{A^*(\cos(m\omega_2 + \eta_3)u_1^- + \sin(m\omega_2 + \eta_3)u_2^-)}{(\cos(m\omega_2 + \eta_3) + b \sin(m\omega_2 + \eta_3))^2} \cdot \frac{\sin(k\omega_1 + \eta_1)}{\sin(k\omega_1 + \eta_2)} \neq \pm 1. \end{aligned} \quad (32)$$

Obviously, condition **A2** is satisfied when both $\omega_1/2\pi$ and $\omega_2/2\pi$ are irrational. This condition also holds if $\omega_1/2\pi$ is irrational and $\omega_2/2\pi = p/q$ with $q \geq 8$, and if $\omega_1/2\pi = p_1/q_1$ with $q_1 \geq 9$ and $\omega_2/2\pi = p_2/q_2$ with $q_2 \geq 8$. Indeed, when $q_2 \geq 8$, there are at least 6 different values of $m\omega_2 \bmod 2\pi$ in $[0, 2\pi)$, so at least one value will satisfy (13) and $\cos(m\omega_2 + \eta_3)u_1^- + \sin(m\omega_2 + \eta_3) \neq 0$. Taking this value fixes the factors involving m in (32). Then, arguing as in the saddle-focus case for condition **A1**, we find that, when ω_1 is irrational or $q_1 \geq 9$, there is at least one value of $k\omega_1 \bmod 2\pi$ validating all inequalities in (32).

For the sake of clarity in proving our results, we classify the various possibilities of the arithmetic relations among $\theta, \omega_1/2\pi, \omega_2/2\pi$ into 7 cases.

Proposition 5.1. *Let δ be the size of Π defined in (12), and \mathcal{C}^{ss} and \mathcal{C}^{uu} be given by Lemma 3.4 with a sufficiently small K . Let a non-degenerate double-focus heterodimensional cycle satisfy condition **A2** and U be any neighborhood of the cycle.*

1. *If θ is irrational, and $\omega_1/2\pi, \omega_2/2\pi$ are rational, then, for every sufficiently large pair (k, m) satisfying condition **A2**, there exists a cs-blender in U if $|\alpha_{k,m}| < 1$ (defined in (32)), or a cu-blender in U if $|\alpha_{k,m}| > 1$.*

- 2.1. *If $\omega_1/2\pi$ is irrational, $\omega_2/2\pi$ is rational, and $\theta, \omega_1/2\pi, 1$ are rationally independent, then there exist simultaneously a cs-blender and a cu-blender in U .*

2.2. If $\omega_1/2\pi$ is irrational, $\omega_2/2\pi$ is rational, and $\theta, \omega_1/2\pi, 1$ are rationally dependent, then there exists a cu-blender in U .

3.1. If $\omega_1/2\pi, \omega_2/2\pi$ are irrational, and $\theta, \omega_1/2\pi, \theta\omega_2/2\pi, 1$ are rationally independent, then there exist simultaneously a cs-blender and a cu-blender in U .

3.2. If $\omega_1/2\pi, \omega_2/2\pi$ are irrational, $\theta, \omega_1/2\pi, \theta\omega_2/2\pi, 1$ are rationally dependent, but $\theta, \omega_1/2\pi, 1$ are rationally independent, then there exist simultaneously a cs-blender and a cu-blender in U .

3.3.1. If $\omega_1/2\pi, \omega_2/2\pi$ are irrational, $\theta, \omega_1/2\pi, 1$ are rationally dependent, and $\omega_1/2\pi, \theta\omega_2/2\pi, 1$ are rationally dependent, then there exists a cu-blender in U .

3.3.2. If $\omega_1/2\pi, \omega_2/2\pi$ are irrational, $\theta, \omega_1/2\pi, 1$ are rationally dependent, and $\omega_1/2\pi, \theta\omega_2/2\pi, 1$ are rationally independent, then there exists a cu-blender in U .

In each of the above cases, the cs-blender has index d_1 and an activating pair (Π', \mathcal{C}^{ss}) , while the cu-blender has index $(d_2 = d_1 + 1)$ and an activating pair (Π', \mathcal{C}^{uu}) , where the cube Π' is defined as

$$\Pi' = [-c\delta, c\delta] \times [-\delta, \delta]^{d-d_2} \times [-\delta, \delta]^{d_2-1} \subset \Pi, \quad (33)$$

for some $c \in (0, 1)$.

We postpone the proof of this proposition to Section 5.3, after we prove the theorems.

5.2 Homoclinic relations in the double-focus case

Let us find out the homoclinic relations among the found blenders in Proposition 5.1 and the periodic points O_1 and O_2 . Denote by Λ_1 and Λ_2 the cs- and cu-blenders. Since the proof of Proposition 5.1 (see Section 5.4) is also based on the use of Proposition 4.2, the structures of Λ_i ($i = 1, 2$) are completely the same as those in the saddle-focus case, i.e., they consist of points whose orbits under the corresponding sets of first return maps never leave the cube Π defined in (12), and their activating domain is Π' defined in (33). (See the discussion at the beginning of Section 4.3). One concludes from Lemma 3.4 that for any points $M_1 \in \Lambda_1 \cap \Pi'$ and $M_2 \in \Lambda_2 \cap \Pi'$, their local invariant manifolds are given by smooth functions of the following forms:

- $W_{loc}^s(M_1) : W = \psi_{M_1}^s(U, V)$ defined on $[-\delta, \delta] \times [-\delta, \delta]^{d-d_2}$ with c in (33);
- $W_{loc}^u(M_1) : (U, V) = \psi_{M_1}^u(W)$ defined on $[-\delta, \delta]^{d_2-1}$, and, in particular, $W_{loc}^u(M_1)$ crosses Π properly with respect to \mathcal{C}^{uu} defined in (17);
- $W_{loc}^s(M_2) : (U, W) = \psi_{M_2}^s(V)$ defined on $[-\delta, \delta]^{d-d_2}$, and, in particular, $W_{loc}^s(M_2)$ crosses Π properly with respect to \mathcal{C}^{ss} defined in (18);
- $W_{loc}^u(M_2) : V = \psi_{M_2}^u(U, W)$ defined on $[-\delta, \delta] \times [-\delta, \delta]^{d_2-1}$.

Obviously, whenever two blenders in Proposition 5.1 coexist (i.e., cases 2.1, 3.1 and 3.2), they activate each other in the sense of Definition 2.1.

In what follows we collect some results in [23] on the invariant manifolds of the periodic points O_1 and O_2 . First, by equations (6.42) and (6.36) in [23] (see the discussion at the end of [23, Section

6.4]), one has

$$W_{loc}^u(O_2) = \{V = 0\} \quad \text{and} \quad F_{21}^{-1}(W_{loc}^s(O_1)) \cap \Pi' = \{W = \psi(U, V)\},$$

where $\psi = O(U^2 + \delta V^2)$ is some smooth function defined on $[-c\delta, c\delta] \times [-\delta, \delta]^{d-d_2}$. Comparing these equations with those for $W_{loc}^s(M_2)$ and $W_{loc}^u(M_1)$, we obtain that

$$W^u(O_2) \pitchfork W^s(\Lambda_2) \neq \emptyset \quad \text{and} \quad W^s(O_1) \pitchfork W^u(\Lambda_1) \neq \emptyset. \quad (34)$$

For other heteroclinic intersections, we use the following

Lemma 5.2 ([23, Lemma 6.8]). *If $\omega_2/2\pi$ is irrational, then there exist infinitely many m such that $F_2^m \circ F_{12}(W_{loc}^u(O_1)) \cap \Pi'$ is a disc of the form $(U, V) = s(W)$, crossing Π' properly with respect to C^{uu} defined by (17).*

Lemma 5.3 ([23, Lemma 6.9]). *If $\omega_1/2\pi$ is irrational, then there exist infinitely many k such that $F_{21}^{-1} \circ F_1^{-k} \circ F_{12}^{-1}(W_{loc}^s(O_2))$ is a disc of the form $(U, W) = s_2(V)$, crossing Π' defined by (33) properly with respect to C^{ss} defined by (18).*

Thus, if $\omega_2/2\pi$ is irrational, then $W^u(O_1) \pitchfork W^s(\Lambda_1) \neq \emptyset$, which along with (34) implies that O_1 is homoclinically related to the cs-blender Λ_1 ; if $\omega_1/2\pi$ is irrational, then $W^s(O_2) \pitchfork W^u(\Lambda_2) \neq \emptyset$, which along with (34) implies that hence O_2 is homoclinically related to the cu-blender Λ_2 . In summary, we obtain

Lemma 5.4. *The periodic points $O_{1,2}$ and the blenders found in Proposition 5.1 have the following homoclinic relations:*

- *in case 2.1, the cu-blender is homoclinically related to O_2 ; the two blenders activate each other, and, in particular, O_2 activates the cs-blender,*
- *in case 2.2, the cu-blender is homoclinically related to O_2 ,*
- *in cases 3.1 and 3.2, the cs-blender is homoclinically related to O_1 and the cu-blender is homoclinically related to O_2 ; the two blenders activate each other, and, in particular, O_1 activates the cu-blender and O_2 activates the cs-blender, and*
- *in cases 3.3.1 and 3.3.2, the cu-blender is homoclinically related to O_2 and activated by O_1 .*

5.3 Proofs of Theorem 2 and the second case in Theorem 3

Proof of Theorem 2. Cases 2.1 and 2.2 in Proposition 5.1 together with Lemma 5.4 cover case (1.i), and, by considering f^{-1} , also case (1.ii). Since the rational independence of $\theta, \omega_1/2\pi, 1$ implies the irrationality of θ , case (2.i) follows from case 2.1 in Proposition 5.1 and Lemma 5.4. Using f^{-1} , we also obtain case (2.ii). Case (3) follows immediately from case 1 in Proposition 5.1.

What remains is case (1.iii), whose hypothesis is covered by those of cases 3.1, 3.2, 3.3.1 and 3.3.2 in Proposition 5.1. We are done if either of the hypotheses in cases 3.1 and 3.2 holds. Now suppose we are under the hypothesis of case 3.3.1. With Lemma 5.4, this gives us a cu-blender homoclinically related to O_2 and activated by O_1 . Considering f^{-1} leads to one of cases 3.1, 3.2, 3.3.1 and 3.3.2. Since the roles of O_1 and O_2 interchange for f^{-1} , it follows from Proposition 5.1 and Lemma 5.4 that

we always have a cu-blender of f^{-1} , homoclinically related to O_1 and activated by O_2 . This blender is a cs-blender of f . Using the lambda-lemma, we see that this cs-blender and the previous cu-blender activate each other, giving the desired result for case (1.iii). The proof with the hypothesis of case 3.3.1 is the same. \square

Proof of the second case in Theorem 3. The proof is essentially the same as in the saddle-focus case. The only difference is that the quantity α_k there is replaced by $\alpha_{k,m}$ defined in (32), which depends on the coefficients $A^*, b, u_1^-, u_2^-, \eta_{1,2,3}$ in (15). Take the neighborhood \mathcal{U} of g sufficiently small such that Γ_f is non-degenerate for all $f \in \mathcal{U}$. Take any $f \in \mathcal{U}$ with rational $\omega_1(f)/2\pi$ and $\omega_2(f)/2\pi$, and denote their denominators by q_1 and q_2 , respectively. Since the coefficients depend on f continuously, there exists a lower bound $N_1(\mathcal{U})$ such that if $q_1 \geq N_1(\mathcal{U})$ and $q_2 \geq 8$, then one can find $r_1, r'_1 \in \{0, \dots, q_1 - 1\}$ and $r_2 \in \{0, \dots, q_2 - 1\}$ satisfying that $|\alpha_{r_1, r_2}| < 1$ and $|\alpha_{r'_1, r_2}| > 1$. Note that here taking $q_2 \geq 8$ is only for the validation of condition A2, and after fixing r_2 one only needs to change r_1 adjust the values of $|\alpha_{r_1, r_2}|$. Applying the same arguments to f^{-1} gives a lower bound $N_2(\mathcal{U})$ for q_2 . Setting $N(\mathcal{U}) = \max\{N_1(\mathcal{U}), N_2(\mathcal{U})\}$ proves the theorem. \square

5.4 Proof of Proposition 5.1

We first prove an auxiliary result (Lemma 5.5 below). Define two functions

$$g : \mathbb{R} \times S^1 \times S^1 \rightarrow \mathbb{R}, \quad g(t, s, w) = \gamma^t A(2\pi w) \sin(2\pi s + \eta_1), \quad (35)$$

$$h : \mathbb{R} \times S^1 \times S^1 \rightarrow \mathbb{R}, \quad h(t, s, w) = \gamma^t B(2\pi w) \sin(2\pi s + \eta_2) - \xi(2\pi w), \quad (36)$$

where $S^1 = [0, 1]$ with the endpoints identified, and A, B, ξ are given in (16). We consider two classes of lines and one class of planes in \mathbb{R}^3 :

$$\text{(proper horizontal line)} \quad \{s = s^*, w = w^*\}, \quad (37)$$

where s^* and w^* are constants satisfying

$$0 < \xi(2\pi w^*) B(2\pi w^*) \sin(2\pi s^* + \eta_2) < \infty, \quad \frac{\xi(2\pi w^*) A(2\pi w^*) \sin(2\pi s^* + \eta_1)}{\sin(2\pi s^* + \eta_2)} \notin \{0, \pm 1, \infty\}; \quad (38)$$

$$\text{(proper tilted line)} \quad \{t = a_1 s + b, w = a_2 s\} \quad \text{or} \quad \{t = a_1 s + b, w = w^*\}, \quad (39)$$

where $a_1^2 + b^2 \neq 0, a_2 \neq 0$, and w^* satisfies $A(2\pi w^*) B(2\pi w^*) \xi(2\pi w^*) \notin \{0, \infty\}$;

$$\text{(proper plane)} \quad \{w = at + bs\} \quad \text{or} \quad \{w = w^*\}, \quad (40)$$

where $a^2 + b^2 \neq 0$, and w^* satisfies $A(2\pi w^*) B(2\pi w^*) \xi(2\pi w^*) \notin \{0, \infty\}$.

Lemma 5.5. *For a sequence $E = \{(t_n, s_n, w_n)\}$ of points in \mathbb{R}^3 ,*

1. *if the closure $cl(E)$ contains a proper horizontal line, then there exists a subsequence for which $\{h(t_n, s_n, w_n)\}$ is dense near zero and $\{|g(t_n, s_n, w_n)|\}$ belongs to some closed interval either in $(0, 1)$ or in $(1, \infty)$, depending on s^* and w^* in (37) and coefficients in g and h ;*
2. *if the closure $cl(E)$ contains a proper tilted line, then there exists a subsequence for which $\{h(t_n, s_n, w_n)\}$ is dense near zero and $\{|g(t_n, s_n, w_n)|\}$ belongs to some closed interval in $(1, \infty)$;*

3. if the closure $cl(E)$ contains a proper plane, then there exist two subsequences such that $\{h(t_n, s_n, w_n)\}$ is dense near zero for both of them and $\{|g(t_n, s_n, w_n)|\}$ belongs to some closed interval in $(0, 1)$ for one of them and in $(1, \infty)$ for the other.

Proof. To simplify the proof, we consider g and h as functions $\mathbb{R}^3 \rightarrow \mathbb{R}$, which makes no difference by the formulas of g and h . Also note that if a sequence approximates a proper line or a proper plane in $\mathbb{R} \times S^1 \times S^1$, then its lift in \mathbb{R}^3 also approximates the lift of the proper line or proper plane.

(Case 1.) Let the proper horizontal line be given by (37). Its image by h is parametrized as

$$h_1(t) := h(t, s^*, w^*) = \gamma^t B(2\pi w^*) \sin(2\pi s^* + \eta_2) - \xi(2\pi w^*).$$

It follows from the first inequality in (38) that this image covers a neighborhood of zero. Since $cl(E)$ contains the proper horizontal line, by continuity of h_1 , we obtain a subsequence for which $\{h(t_n, s_n, w_n)\}$ is dense near zero. Let us denote the subsequence by the same notations. By (35) and (36), we have

$$\begin{aligned} |g(t_n, s_n, w_n)| &= \left| \frac{(h(t_n, s_n, w_n) + \xi(2\pi w_n))A(2\pi w_n) \sin(2\pi s_n + \eta_1)}{B(2\pi w_n) \sin(2\pi s_n + \eta_2)} \right| \\ &\rightarrow \left| \frac{\xi(2\pi w^*)A(2\pi w^*) \sin(2\pi s^* + \eta_1)}{B(2\pi w^*) \sin(2\pi s^* + \eta_2)} \right|. \end{aligned} \quad (41)$$

This gives the desired result on $|g(t_n, s_n, w_n)|$ due to the second inequality in (38).

(Case 2.) Let the proper tilted line be given by (39). By (16), one can find a small interval Δ_i arbitrarily close to $s = (-\eta_2/2\pi) + i$ with $i \in \mathbb{Z}$ such that

$$0 < \xi(2\pi w)B(2\pi w) \sin(2\pi s + \eta_2) < \infty, \quad \frac{\xi(2\pi w)A(2\pi w) \sin(2\pi s + \eta_1)}{\sin(2\pi s + \eta_2)} \notin \{0, \pm 1, \infty\}, \quad (42)$$

for all $s \in \Delta_i$ with $w = a_2 s$ or $w = w^*$. Since $\xi(2\pi w)$ is bounded, taking n sufficiently large and Δ_i sufficiently close to $s = (-\eta_2/2\pi) + i$ ensures that the images of the functions

$$h_{21}(s) := h(a_1 s + b, s, a_2 w) = \gamma^{a_1 s + b} B(2\pi a_2 s) \sin(2\pi s + \eta_2) - \xi(2\pi a_2 s), \quad s \in \Delta_i$$

obtained with $w = a_2 s$ in (39), and

$$h_{22}(s) := h(a_1 s + b, s, w^*) = \gamma^{a_1 s + b} B(2\pi w^*) \sin(2\pi s + \eta_2) - \xi(2\pi w^*), \quad s \in \Delta_i$$

obtained with $w = w^*$ in (39) both cover a small neighborhood of zero. Now, take a subsequence of E approximating the line segment obtained by intersecting the tilted line with $\{(t, w, w) \mid s \in \Delta_i\}$. Obviously, for this subsequence $\{h(t_n, s_n, w_n)\}$ is dense near zero. Moreover, when Δ_i is taking sufficiently close, the value of $\sin(2\pi s + \eta_2)$ is arbitrarily close to zero, which together with (41) implies that $\{|g(t_n, s_n, w_n)|\}$ lie in some closed interval in $(1, \infty)$.

(Case 3.) We now take a small interval Δ_i arbitrarily close to $s = (-\eta_2/2\pi) + i$ and a small interval Δ'_i arbitrarily close to $s = -\eta_1/2\pi + i$. Define two functions

$$h_{31}(t, s) := h(t, s, at + bs) = \gamma^t B(2\pi(at + bs)) \sin(2\pi s + \eta_2) - \xi(at + bs),$$

in the first case of (40), and

$$h_{32}(t, s) := h(t, s, w^*) = \gamma^t B(2\pi w^*) \sin(2\pi s + \eta_2) - \xi(w^*)$$

in the second case of (40). Obviously, one can find t^* such that, for all $s \in \Delta_i$ and either $w = at^* + bs$ or $w = w^*$, the inequalities in (42) are satisfied and $h_{31}(\Delta_i)$ or $h_{32}(\Delta_i)$, depending on whether $w = at^* + bs$ or $w = w^*$, covers a small neighborhood of zero. Then, consider the subsequence approximating the intersection of the proper plane with $\{t = t^*, s \in \Delta_i\}$. Proceeding as in case 2 shows that $\{h(t_n, s_n, w_n)\}$ is dense near zero and $\{|g(t_n, s_n, w_n)|\}$ lie in some closed interval in $(1, \infty)$. Considering a similar number t^* for $s \in \Delta'_i$ gives another required subsequence. \square

Proof of Proposition 5.1. Denote

$$t_n = m_n - k_n \theta, \quad s_n = k_n \frac{\omega_1}{2\pi} \bmod 1, \quad w_n = m_n \frac{\omega_2}{2\pi} \bmod 1. \quad (43)$$

Recall that in the double-focus case we have $\lambda > 0, \gamma > 0$, so $\lambda^k \gamma^m = \gamma^{m-k\theta}$. Comparing $A_{k,m}$ and $B_{k,m}$ in (15) with functions g and h given by (35) and (36), yields

$$A_{k_n, m_n} = g(t_n, s_n, w_n) \quad \text{and} \quad B_{k_n, m_n} = h(t_n, s_n, w_n).$$

It follows that, for any sequence $\{(t_n, s_n, m_n)\}$ falling in one of the cases of Lemma 5.5, Proposition 4.2 is applicable to the system $\{T_n := T_{k_n, m_n}\}$ (passing to the subsequence in the lemma) given by (14), with the correspondence $U \rightarrow X, W \rightarrow Y, V \rightarrow Z$. More specifically, we have that

- if $\{(t_n, s_n, m_n)\}$ approximates a proper horizontal line, then there exists a blender whose type depends on the coefficients in (14);
- if $\{(t_n, s_n, m_n)\}$ approximates a proper tilted line, then there exists a cu-blender; and
- if $\{(t_n, s_n, m_n)\}$ approximates a proper plane, then there exist simultaneously a cs-blender and a cu-blender.

Moreover, according to the proof of Proposition 4.2, the obtained blender has an activating pair (Π', \mathcal{C}^{ss}) if it is center-unstable or (Π', \mathcal{C}^{uu}) if it is center-stable, where Π is defined in (33) and \mathcal{C}^{ss} and \mathcal{C}^{uu} are given by Lemma 3.4.

We thus prove the Proposition 5.1 by showing that, for each of the cases in the statement of the proposition, one can find a sequence $\{(k_n, m_n)\}$ such that $\{(t_n, s_n, w_n)\}$ given by (43) approximates a proper line or a proper plane. Recall that $\omega_{1,2} \in (0, \pi)$ and by p/q we always mean that p and q are positive coprime integers.

(1) θ is irrational, $\omega_1/2\pi, \omega_2/2\pi$ are rational. Denote

$$\frac{\omega_1}{2\pi} = \frac{p_1}{q_1} \quad \text{and} \quad \frac{\omega_2}{2\pi} = \frac{p_2}{q_2}.$$

Since condition A2 is satisfied, there exist integers $r_1 \in \{0, \dots, q_1 - 1\}$ and $r_2 \in \{0, \dots, q_2 - 1\}$ such that inequalities in (38) are satisfied with $s^* = r_1 \omega_1 / 2\pi$ and $w^* = r_2 \omega_2 / 2\pi$. Then,

$$\left\{ s = r_1 \frac{\omega_1}{2\pi} \bmod 1, w = r_2 \frac{\omega_2}{2\pi} \bmod 1 \right\}$$

is a proper horizontal line of the form (37). The irrationality of θ implies that we can take a sequence

$\{(k'_n, m'_n)\}$ of pairs of positive integers such that the set

$$\{q_2 m'_n + r_2 - (q_1 k'_n + r_1)\theta\}$$

is dense in \mathbb{R} . Setting $k_n = q_1 k'_n + r_1$ and $m_n = q_2 m'_n + r_2$ in (43) gives a sequence whose closure contains above proper horizontal line. So, we get a cs-blender if $|\alpha_{r_1, r_2}| < 1$, or a cu-blender if $|\alpha_{r_1, r_2}| > 1$.

(2.1) $\omega_1/2\pi$ is irrational, $\omega_2/2\pi$ is rational, and $\theta, \omega_1/2\pi, 1$ are rationally independent. Let p_2, q_2, r_2 be as in the preceding case. By the rational independence among $\theta, \omega_1/2\pi, 1$, one can find a sequence $\{(k'_n, m'_n, i'_n) \in \mathbb{N}^3\}$ such that the set

$$\left\{ \left(q_2 m'_n + r_2 - k'_n \theta, k'_n \frac{\omega_1}{2\pi} - i'_n \right) \right\}$$

is dense in \mathbb{R}^2 . Setting $k_n = k'_n$ and $m_n = q_2 m'_n + r_2$ in (43) gives a sequence whose closure contains the proper plane $\{w = r_2 \omega_2 / 2\pi\}$. So, we get simultaneously a cs-blender and a cu-blender.

Before proceed to other cases, we make the following

Claim. To prove that there exists $\{(t_n, s_n, w_n)\}$ given by (43) approximating a proper line or a proper plane, it suffices to show that each point of this proper line or proper plane is accumulated by points of the set

$$\left\{ \left(m_n - k_n \theta, k_n \frac{\omega_1}{2\pi} \bmod 1, m_n \frac{\omega_2}{2\pi} \bmod 1 \right) \right\} \quad (44)$$

for some sequence $\{(k_n, m_n)\}$.

Indeed, the assumption particularly implies that, for each rational point in the proper line or proper plane, we have such a sequence $\{(k_n, m_n)\}$, and hence the union of these countably many sequences gives a new sequence which defines the desired $\{(t_n, s_n, w_n)\}$ by (43).

(2.2) $\omega_1/2\pi$ is irrational, $\omega_2/2\pi$ is rational, and $\theta, \omega_1/2\pi, 1$ are rationally dependent. Denote

$$\theta = \frac{p_1 \omega_1}{q_1 2\pi} + \frac{p_2}{q_2} \quad \text{and} \quad \frac{\omega_2}{2\pi} = \frac{p_3}{q_3}. \quad (45)$$

With r_2 as in case 1, the set $\{t = -sp_1/q_1 + r_2, w = r_2 p_3 / q_3 \bmod 1\}$ is a proper tilted line. Let us show that each point of this line can be approximated by the set (44).

By the irrationality of $\omega_1/2\pi$, the set

$$\left\{ q_2 q_3 k \frac{\omega_1}{2\pi} - q_1 q_3 i \right\}_{k, i \in \mathbb{N}^2}$$

is dense in \mathbb{R} . So, for any $s \in [0, 1]$, one can find a sequence $\{(k'_n, i'_n)\}$ such that

$$q_2 q_3 k'_n \frac{\omega_1}{2\pi} - q_1 q_3 i'_n \rightarrow s.$$

Denote $k_n = q_2 q_3 k'_n$ and $m_n = r_2 + p_2 q_3 k'_n + p_1 q_3 i'_n$. The relation (45) then implies

$$m_n - k_n \theta = (r_2 + p_2 q_3 k'_n + p_1 q_3 i'_n) - \frac{p_1}{q_1} k_n \frac{\omega_1}{2\pi} - p_2 q_3 k'_n$$

$$\begin{aligned}
&= r_2 - \frac{p_1}{q_1} \left(k_n \frac{\omega_1}{2\pi} - q_1 q_3 i'_n \right) \\
&\rightarrow r_2 - s \frac{p_1}{q_1},
\end{aligned}$$

$$\begin{aligned}
m_n \frac{\omega_2}{2\pi} \pmod{1} &= (r_2 + p_2 q_3 k'_n + p_1 q_3 i'_n) \frac{p_3}{q_3} \pmod{1} \\
&= \frac{r_2 p_3}{q_3} \pmod{1},
\end{aligned}$$

which give the desired approximation. By the claim, we have a cu-blender.

(3.1) $\omega_1/2\pi, \omega_2/2\pi$ are irrational, and $\theta, \omega_1/2\pi, \theta\omega_2/2\pi, 1$ are rationally independent. The rational independence implies that, for any point (t, s, w) , there exists $\{(k_n, m_n, i_n, \ell_n)\}$ such that

$$m_n - k_n \theta \rightarrow t, \quad k_n \frac{\omega_1}{2\pi} - i_n \rightarrow s, \quad k_n \theta \frac{\omega_2}{2\pi} - \ell_n \rightarrow w - t \frac{\omega_2}{2\pi}. \quad (46)$$

Combining the first and third relations, yields

$$(m_n - t) \frac{\omega_2}{2\pi} - \ell_n \rightarrow w - t \frac{\omega_2}{2\pi},$$

which implies

$$m_n - k_n \theta \rightarrow t, \quad k_n \frac{\omega_1}{2\pi} - i_n \rightarrow s, \quad m_n \frac{\omega_2}{2\pi} - \ell_n \rightarrow w, \quad (47)$$

Therefore, any point of any proper plane can be approximated by a sequence of the form (44). By the claim, we get simultaneously a cs-blender and a cu-blender.

(3.2) $\omega_1/2\pi, \omega_2/2\pi$ are irrational, $\theta, \omega_1/2\pi, \theta\omega_2/2\pi, 1$ are rationally dependent, but $\theta, \omega_1/2\pi, 1$ are rationally independent. Denote

$$\theta \frac{\omega_2}{2\pi} = \frac{p_1}{q_1} \theta + \frac{p_2}{q_2} \frac{\omega_1}{2\pi} + \frac{p_3}{q_3}. \quad (48)$$

Since $\theta, \omega_1/2\pi, 1$ are rationally independent, the set

$$\left\{ \left(q_1 m - q_3 k \theta, q_3 k \frac{\omega_1}{2\pi} - q_2 i \right) \right\}_{k, m, i \in \mathbb{N}}$$

is dense in \mathbb{R}^2 . Hence, for any point (t, s) there exists a sequence (k'_n, m'_n, i'_n) such that

$$q_1 m'_n - q_3 k'_n \theta \rightarrow t \quad \text{and} \quad q_3 k'_n \frac{\omega_1}{2\pi} - q_2 i'_n \pmod{1} \rightarrow s.$$

Denote $k_n = q_3 k'_n$ and $m_n = q_1 m'_n$. The relation (48) shows that

$$\begin{aligned}
m_n \frac{\omega_2}{2\pi} \pmod{1} &= k_n \left(\frac{p_1}{q_1} \theta + \frac{p_2}{q_2} \frac{\omega_1}{2\pi} + \frac{p_3}{q_3} \right) - \frac{\omega_2}{2\pi} t \pmod{1} \\
&\rightarrow \left(\frac{p_1}{q_1} - \frac{\omega_2}{2\pi} \right) t + \frac{p_2}{q_2} s \pmod{1} \\
&=: at + bs \pmod{1}.
\end{aligned}$$

Thus, any point of the proper plane $\{w = at + bs \pmod{1}\}$ can be approximated by a sequence of the form 44. By the claim, we get simultaneously a cs-blender and a cu-blender.

(3.3.1) $\omega_1/2\pi, \omega_2/2\pi$ are irrational, $\theta, \omega_1/2\pi, 1$ are rationally dependent, and $\omega_1/2\pi, \theta\omega_2/2\pi, 1$ are

rationally dependent. Denote

$$\theta = \frac{p_1}{q_1} \frac{\omega_1}{2\pi} + \frac{p_2}{q_2} \quad \text{and} \quad \theta \frac{\omega_2}{2\pi} = \frac{p_3}{q_3} \frac{\omega_1}{2\pi} + \frac{p_4}{q_4}. \quad (49)$$

The irrationality of $\omega_1/2\pi$ implies that the set

$$\left\{ q_2 q_4 k \frac{\omega_1}{2\pi} - q_1 q_3 i \right\}_{k,i \in \mathbb{N}}$$

is dense in \mathbb{R} . For any $s \in [0, 1]$ and any sequence (k'_n, i'_n) such that

$$q_2 q_4 k'_n \frac{\omega_1}{2\pi} - q_1 q_3 i'_n \rightarrow s,$$

denote $k_n = q_2 q_4 k'_n$ and $m_n = p_2 q_4 k'_n + p_1 q_3 i'_n$. The relations in (49) show that

$$\begin{aligned} m_n - k_n \theta &= p_2 q_4 k'_n + p_1 q_3 i'_n - q_2 q_4 k'_n \frac{p_1}{q_1} \left(\frac{\omega_1}{2\pi} + \frac{p_2}{q_2} \right) \\ &= -\frac{p_1}{q_1} \left(q_2 q_4 k'_n \frac{\omega_1}{2\pi} - q_1 q_3 i'_n \right) \\ &\rightarrow -\frac{p_1}{q_1} s, \\ k_n \theta \frac{\omega_2}{2\pi} \pmod{1} &= \frac{p_3}{q_3} \left(q_2 q_4 k'_n \frac{\omega_1}{2\pi} - q_1 q_3 i'_n \right) + p_3 q_1 i'_n + p_4 q_2 k'_n \pmod{1} \\ &\rightarrow \frac{p_3}{q_3} s \pmod{1}. \end{aligned}$$

Observing the relation between (46) and (47), one finds

$$m_n \frac{\omega_2}{2\pi} \rightarrow \left(\frac{p_3}{q_3} - \frac{\omega_2 p_1}{2\pi q_1} \right) s =: as.$$

Thus, any point of the proper tilted line $\{t = -p_1 s/q_1, w = as\}$ can be approximated by a sequence of the form (44). The claim then gives a cu-blender.

(3.3.2) $\omega_1/2\pi, \omega_2/2\pi$ are irrational, $\theta, \omega_1/2\pi, 1$ are rationally dependent, and $\omega_1/2\pi, \theta\omega_2/2\pi, 1$ are rationally independent. We still denote θ as in (49). For any $s \in [0, 1]$ and any sequence (k'_n, i'_n) such that

$$q_2 k'_n \frac{\omega_1}{2\pi} - q_1 i'_n \rightarrow s,$$

denote $k_n = q_2 k'_n$ and $m_n = p_2 k'_n + p_1 i'_n$. The first relation in (49) gives

$$m_n - k_n \theta = p_2 k'_n + p_1 i'_n - q_2 k'_n \left(\frac{p_1}{q_1} \frac{\omega_1}{2\pi} + \frac{p_2}{q_2} \right) \rightarrow -\frac{p_1}{q_1} s.$$

It then follows from the rational independence among $\omega_1/2\pi, \theta\omega_2/2\pi, 1$ and the correspondence relation between (46) and (47) that any point of any proper tilted line in the plane $\{t = -p_1 s/q_1\}$ can be approximated by a sequence of the form (44). The claim then gives a cu-blender. \square

6 Robust heterodimensional dynamics in generic one-parameter unfolding families. Proof of Theorem 4

Finding robust heterodimensional dynamics is essentially equivalent to connecting the invariant manifolds of the blenders and periodic orbits O_i ($i = 1, 2$). As shown in Sections 4.3 and 5.2, the structure of the blenders obtained in this paper is quite clear in the small neighborhood Π of M_1^+ (for saddle-focus cycles) and of M_2^- (for double-focus cycles), with explicit formulas for their local invariant manifolds. The homoclinic relations or activations between blenders and the points O_i depend on how the local invariant manifolds of O_i behave along the dynamics of the cycle. By construction, the intersections of $W^s(O_1)$ with the unstable manifolds of the blenders and the intersections of $W^u(O_2)$ with the stable manifolds of the blenders are automatic (see (31) and (34)). The other intersections depend on the positions of the iterates

$$F_{21} \circ F_2^m \circ F_{12}(W_{loc}^u(O_1)) \quad \text{and} \quad F_1^{-k} \circ F_{12}^{-1}(W_{loc}^s(O_2))$$

for saddle-focus cycles, and

$$F_2^m \circ F_{12}(W_{loc}^u(O_1)) \quad \text{and} \quad F_{21}^{-1} \circ F_1^{-k} \circ F_{12}^{-1}(W_{loc}^s(O_2))$$

for double-focus cycles. Basically, one gets the intersection of $W^u(O_1)$ with the stable manifolds of the blenders if the first iterate in each case crosses Π' , and gets the intersection of $W^s(O_2)$ with the unstable manifolds of the blenders if the second iterate in each case crosses Π' . According to Lemma 4.4, the second iterate in the saddle-focus case always hits Π' if $\omega/2\pi$ is irrational. Similarly, in the double-focus case, Lemmas 5.2 and 5.3 show that the two iterates cross Π' if $\omega_2/2\pi$ and, respectively, $\omega_1/2\pi$ is irrational. Thus, the proof of Theorem 3 is essentially about verifying the intersections of these iterates with Π' when $\omega/2\pi$ and $\omega_i/2\pi$ are rational and the cycle is unfolded.

Recall that μ is the splitting parameter of a non-degenerate heterodimensional cycle, see Remark 3.1. A one-parameter unfolding family $\{f_\varepsilon\}$ with $f_0 = f$ having a non-degenerate cycle satisfies $d\mu(0)/d\varepsilon \neq 0$ (see (2)). Since it suffices to prove the results for one-parameter families, we simply use μ as a new parameter, replacing ε .

6.1 Unfolding saddle-focus heterodimensional cycles

Let us first deal with saddle-focus cycles, where the center-stable multipliers are nonreal and the center-unstable one is real, denoted by $\lambda e^{i\omega}$ and γ , respectively. In the coordinates (4), we have $M_1^- = (0, y^-, 0)$ and $M_2^+ = (0, v^+, 0)$ for some constants $y^- \neq 0$ and $v^+ \neq 0$. The parameter μ enters the Taylor expansion of the transition map F_{12} as

$$\begin{aligned} u &= \mu + a_{11}x + a_{12}(y - y^-) + a_{13}z + \dots \\ v - v^+ &= a_{21}x + a_{22}(y - y^-) + a_{23}z + \dots \\ w &= a_{31}x + a_{32}(y - y^-) + a_{33}z + \dots, \end{aligned}$$

where all coefficients y^-, v^+, a_{ij} depend on μ . For brevity, we denote below the continuations of hyperbolic objects (e.g. $O_{1,2}$ and the blenders) by the same letters and drop the term ‘continuation’.

Remark 6.1. In [23], the parameter is denoted by $\hat{\mu}$, which is written in the cross-form of F_{12} . When

$r \geq 3$ (which is our case), it is the same as our μ defined above. See [23, Section 6.1.2]

Let the coefficient $A, B, u^-, b, \eta_{1,2}$ be as in (8). The cube Π' with constant c is defined in (23), and the cone fields \mathcal{C}^{ss} and \mathcal{C}^{uu} are given by Lemma 3.3. We have the following

Lemma 6.2 ([23, Lemma 6.4]⁷). *Fix any sufficiently small δ . For every sufficiently large m , if*

$$|\mu - \gamma^{-m} u^-| < \frac{1}{2} |\gamma^{-m} b^{-1}| q \delta, \quad (50)$$

then $F_{21} \circ F_2 \circ F_{12}(W_{loc}^u(O_1))$ is a disc of the form $(X, Z) = s(Y)$ for some smooth function s , which crosses Π' properly with respect to \mathcal{C}^{uu} .

When $\omega/2\pi$ is rational, the following result can be obtained by a minor modification of the proof of [23, Lemma 6.5].

Lemma 6.3. *Suppose $\omega/2\pi = p/q$. Fix a sufficiently small δ and any $r \in \{0, \dots, q-1\}$. For every $k = r + iq$ with sufficiently large i , if*

$$|\mu + \lambda^k b^{-1} B \sin(2\pi r + \eta_2)| < \frac{1}{2} |\lambda^k A \sin(2\pi r + \eta_1)| q \delta, \quad (51)$$

then $F_1^{-k} \circ F_{12}^{-1}(W_{loc}^s(O_2))$ is a disc of the form $(X, Y) = s(Z)$ for some smooth function s , which crosses Π' defined by (23) properly with respect to \mathcal{C}^{ss} defined by (11).

Proof. We need a formula for $F_1^{-k} \circ F_{12}^{-1}(W_{loc}^s(O_2))$. Such formula at $\mu = 0$ is given by (6.30) of [23], and a parametric version with non-zero μ can be obtained by setting $\hat{\mu}$ (which is the μ here by Remark 6.1) non-zero in (6.8) and then applying the coordinate transformation (6.11). In the end, one finds that $F_1^{-k} \circ F_{12}^{-1}(W_{loc}^s(O_2))$ is given by

$$\begin{aligned} X &= \frac{-b\lambda^{-k}\mu - B \sin(2\pi r + \eta_2)}{A \sin(2\pi r + \eta_1)} + O(\delta^{\frac{3}{2}}) + o(1)_{k \rightarrow \infty}, \\ Y &= o(\hat{\gamma}^{-k}), \end{aligned}$$

where $\hat{\gamma} > |\gamma|$ is some constant and the right-hand sides are functions of Z defined on $[-\delta, \delta]^{d-d_1-1}$. It immediately follows that, for all sufficiently small δ and sufficiently large k , if μ satisfies (51), then

$$|X| < q\delta, \quad \|Y\| < \delta, \quad \frac{\partial(X, Y)}{\partial Z} = O(\delta^{\frac{3}{2}}) + o(1)_{k \rightarrow \infty},$$

where the first two inequalities mean that $F_1^{-k} \circ F_{12}^{-1}(W_{loc}^s(O_2))$ crosses Π' and the last one implies that the crossing is proper with respect to \mathcal{C}^{ss} . \square

Proof of Theorem 4 (the saddle-focus case). The results in Theorem 4 for the saddle-focus case are those concerning Theorems 1 and 3. By (2), we only need to consider the family $\{f_\mu\}$ with $f_0 = f$.

Since blenders persist under C^1 -small perturbations, the homoclinic relations and activations obtained in Theorem 1 are preserved for all small μ . Hence, to prove the result concerning Theorem 1.(1), it suffices to show that O_1 activates the cu-blender Λ^{cu} and is homoclinically related to some non-trivial hyperbolic basic set. Then, by the lambda-lemma this set also activates Λ^{cu} . Since $W_{loc}^s(O_1) =$

⁷ As mentioned, the coordinates in [23] are (X_1, X_2, Y, Z) , and in this paper we dropped the subscript of X_1 and grouped X_2, Z as a new Z coordinate. Moreover, the coefficient b_{11} there is denoted by b in this paper.

$\{Y = 0\}$ by (30), Lemma 6.2 implies that there exists a sequence $\mu_j \rightarrow 0$ such that $F_{21} \circ F_2 \circ F_{12}(W_{loc}^u(O_1)) \cap W_{loc}^s(O_1) \neq \emptyset$, which gives a hyperbolic basic set homoclinically related to O_1 . To see that O_1 activates Λ^{cu} , we note first that $W^s(O_1) \cap W^u(\Lambda^{cu}) \neq \emptyset$ by (30) and the formula for $W_{loc}^u(M_2)$ in Section 4.3. Then, since Λ^{cu} is obtained from Proposition 4.3 and hence has an activating pair (Π', \mathcal{C}^{uu}) , the proper crossing part of Lemma 6.2 shows that O_1 indeed activates Λ^{cu} .

To prove the result concerning Theorem 1.(2), one only needs to create a homoclinic relation between the cs-blender Λ^{cs} and O_1 by changing μ . The intersection $W^s(O_1) \cap W^u(\Lambda^{cs}) \neq \emptyset$ is automatic at $\mu = 0$ by (30) and the formula for $W_{loc}^u(M_1)$ in Section 4.3. With the formula for $W_{loc}^s(M_1)$ in the same section, one obtains by Lemma 6.2 a sequence $\mu_j \rightarrow 0$ for which $W^u(O_1) \cap W^s(\Lambda^{cs}) \neq \emptyset$.

Before proceeding to the rational ω cases, let us make an observation. By similar arguments to the above cases, we obtain that, when Lemma 6.3 holds, O_2 is homoclinically related to the cu-blender in Proposition 4.3, if exists; it activates the cs-blender, if exists; and, it is always homoclinically related to a non-trivial hyperbolic basic set. We thus proved

Claim. *For every sufficiently small μ satisfying both conditions (50) and (51), each of O_1 and O_2 is homoclinically related to a non-trivial hyperbolic basic set; and, moreover, whenever there is a cs-blender from Proposition 4.3, it is homoclinically related to O_1 and activated by O_2 , and whenever there is a cu-blender from Proposition 4.3, it is homoclinically related to O_2 and activated by O_1 .*

Then, the results concerning Theorem 3 and Theorem 1.(3) follow immediately from

Claim 6.4. *There exists a sequence $\mu_j \rightarrow 0$ with each μ_j satisfying both conditions (50) and (51).*

This completes the proof of Theorem 4 for saddle-focus cycles. Let us prove the above claim. Note that (50) and (51) give two intervals I_m^u and, respectively, I_k^s of μ values such that the inequality is satisfied if μ lies in corresponding interval. It then suffices to find a sequence $\{k_n, m_n\}$ with $k_n, m_n \rightarrow \infty$ such that $I_{m_n}^u \cap I_{k_n}^s \neq \emptyset$. The centers of these two intervals are given by

$$c_m^u = \gamma^{-m} u^- \quad \text{and} \quad c_k^s = -\lambda^k b^{-1} B \sin(k\omega + \eta_2),$$

and the radius of I_m^u is $|\gamma^{-m} b^{-1}| q \delta / 2$. Now set $k_n = k'_n q + r$ and $m_n = m'_n$ as in (25). It follows from (26) that

$$\frac{|c_{m_n}^u - c_{k_n}^s|}{|\gamma^{-m_n} b^{-1}| q \delta / 2} = 2 \cdot \frac{|\lambda^{k_n} \gamma^{m_n} b^{-1} B \sin(k_n \omega + \eta_2) - u^-|}{b^{-1} \delta} \rightarrow 0 \quad \text{as} \quad n \rightarrow \infty,$$

which shows that $I_{m_n}^u \cap I_{k_n}^s \neq \emptyset$ for all sufficiently large n . □

6.2 Unfolding double-focus heterodimensional cycles

Here we have the following counterparts to Lemmas 6.2 and 6.2, and Claim 6.4:

- If ω_2/π is rational, then there exists a sequence of intervals I_m^u such that, for all sufficiently large m , if $\mu \in I_m^u$ then $F_2^m \circ F_{12}(W_{loc}^u(O_1)) \cap \Pi'$ is a disc of the form $(U, V) = s(W)$, crossing Π' in (33) properly with respect to \mathcal{C}^{uu} in Lemma 3.4.
- If ω_1/π is rational, then there exists a sequence of intervals I_k^s such that, for all sufficiently large k , if $\mu \in I_k^s$ then $F_{21}^{-1} \circ F_1^{-k} \circ F_{12}^{-1}(W_{loc}^s(O_2)) \cap \Pi'$ is a disc of the form $(U, W) = s_2(V)$, crossing Π' in (33) properly with respect to \mathcal{C}^{ss} in Lemma 3.4.

- If both ω_1/π and ω_2/π are rational, then there exists $\{k_n, m_n\}$ such that $I_{m_n}^u \cap I_{k_n}^s \neq \emptyset$ for all large n .

The first two items are perturbative versions of Lemma 5.2 and Lemma 5.3, respectively. They can be obtained by modifying the proofs in [23, Section 6.4.5] in the same way as we did to obtain Lemma 6.3, the perturbative version of Lemma 4.4. The third item can also be proved in the same way as Claim 6.4. We hence omit their proofs. Theorem 4 for double-focus cycles then follows from the routine checks as in Section 6.1, with using the above results and the lemmas in Section 5.2. \square

7 Purely rational moduli. Proof of Theorem B

Since the set of non-degenerate cycles is open and dense in \mathcal{H}_f (see Section 3.1), we can always assume that the cycle in Theorem B is non-degenerate. The first item of Theorem B concerns the saddle case, i.e., when both the center-stable multiplier λ and center-unstable multiplier γ are real. In this case, the result follows immediately from

Theorem 7.1 ([23, Theorem 6]). *Let $f \in \text{Diff}^r(\mathcal{M})$ with $r = 2, \dots, \infty, \omega$ have a non-degenerate (see Remark 3.2) saddle heterodimensional cycle with rational θ , and let U be any neighborhood of the cycle. There exists a nowhere dense subset $E \subset \mathbb{R}^2$ such that, if*

$$\left(|ab|, \left| \frac{u^-}{ax^+} \right| \right) \notin E,$$

then any generic one-parameter unfolding family $\{f_\varepsilon\}$ with $f_0 = f$ has simple hyperbolic dynamics as in Definition 1.4.

Here a, b are coefficients in the transition maps, and x^+ and u^- are coordinate components of O_1 and, respectively, O_2 in linearizing coordinates similar to (4), see [23, Section 3] for details.

We extend the above theorem to the saddle-focus and double-focus cases. For that, we need to investigate the set of orbits which lies entirely in a small neighborhood U of a heterodimensional cycle. Recall that a cycle is the union $\Gamma = L_1 \cup L_2 \cup \Gamma^0 \cup \Gamma^1$, where $L_{1,2}$ are the orbits of the periodic points $O_{1,2}$ and Γ^0 and Γ^1 are the two heteroclinic orbits in $W^u(L_1) \cap W^s(L_2)$ and $W^u(L_2) \cap W^s(L_1)$, respectively. An orbit which lies entirely in U , except for those in the invariant manifolds of $L_{1,2}$, must intersect the neighborhood Π of M^+ infinitely many times for both forward and backward iterations. In particular, an intersection of Π with any orbit gives a pair of two points M and \bar{M} in this orbit satisfying $\bar{M} = T_{k,m}(M)$. In what follows we show that (Lemma 7.2), for any sequence of pairs $\{(k_i, m_i)\}$ corresponding to intersection points of Π with any set of orbits, the assumption of rational moduli imposes a strong relation on the pairs, which thus leads to simple hyperbolic dynamics.

7.1 Simple hyperbolic dynamics in the saddle-focus case

Let us first deal with the saddle-focus cycles, where the center-stable and center-unstable multipliers are $\lambda e^{\pm i\omega}$ and γ , and we assume that θ and $\omega/2\pi$ are rational.

Consider a generic unfolding family $\{f_\mu\}$. To study the dynamics, we need a perturbative version of formula (7). This can be obtained by first applying the coordinate transformation (6.11) in [23]

to equation (6.10) there with keeping the $\hat{\mu}$ term, then making the notation change $b_{11} \rightarrow b, X_1 \rightarrow X, (X_2, Z) \rightarrow Z$, and finally using the Implicit Function Theorem in an obvious way to express the nonlinearities as functions of (X, \bar{Y}, Z) . In the end, we obtain that any two points $(X, Y, Z), (\bar{X}, \bar{Y}, \bar{Z}) \in \Pi^8$ (the domain defined in (5)) satisfy $(\bar{X}, \bar{Y}, \bar{Z}) = T_{k,m}(X, Y, Z)$ for some pair (k, m) if and only if

$$\begin{aligned}\bar{X} &= b\gamma^m \mu + A_{k,m}X + B_{k,m} + \phi_1(X, \bar{Y}, Z), \\ Y &= \phi_3(X, \bar{Y}, Z), \quad \bar{Z} = \phi_4(X, \bar{Y}, Z),\end{aligned}$$

where the coefficients and the functions are the same as in (7), but with smooth dependence on μ . Here we also use a finer estimate for ϕ_1 which is given by equation (6.12) in [23]:

$$\phi_1 = \gamma^m(o(\lambda^k) + o(\gamma^{-m})).$$

Let $\omega/2\pi = p/q$. There exist two finite sets $E_{\eta_i} \subset \mathbb{R}$ ($i = 1, 2$) such that if $\eta_i \notin E_{\eta_i}$ then

$$\sin(2\pi r + \eta_1) \neq 0 \quad \text{and} \quad \sin(2\pi r + \eta_2) \neq 0. \quad (52)$$

For these $\eta_{1,2} \notin E_{\eta}$, we use (8) to rewrite the above formula as

$$\begin{aligned}\bar{X} &= b\gamma^m \mu + a_r b \lambda^k \gamma^m X + a_r b x_r^+ \lambda^k \gamma^m - b u^- + \phi_1(X, \bar{Y}, Z), \\ Y &= \phi_3(X, \bar{Y}, Z), \quad \bar{Z} = \phi_4(X, \bar{Y}, Z),\end{aligned} \quad (53)$$

where

$$a_r = b^{-1} A \sin(2\pi r + \eta_1), \quad x_r^+ = \frac{B \sin(2\pi r + \eta_2)}{A \sin(2\pi r + \eta_1)}, \quad r \in \{0, \dots, q-1\}. \quad (54)$$

Lemma 7.2. *Let $\theta = p'/q'$ and $\omega/2\pi = p/q$. Assume that for all $r_1, r_2 \in \{0, \dots, q-1\}$ we have*

$$\left| \frac{u^-}{a_{r_1} x_{r_1}^+} \right| \notin \bigcup_{i=0}^{q-1} cl \left\{ |\gamma|^{\frac{s}{q'}} \cdot \frac{|a_{r_2} x_{r_2}^+ (a_{r_1} x_{r_1}^+)^{-1}| - \lambda^\ell}{1 - \gamma^{-n}} \right\}_{s \in \mathbb{Z}, \ell \in \mathbb{N}, n \in \mathbb{N}} =: E_{r_1, r_2}. \quad (55)$$

Consider any (finite or infinite) sequence of different pairs $\{(k_i, m_i)\}$ such that there exist some sequence of points $\{M_i \in \Pi\}$ and some small μ for which $T_{k_i, m_i}(M_i) \in \Pi$ with Π defined in (5). Then, for all sufficiently small δ , either all m_i are equal to each other, $m_i = m$, and $\lambda^{k_i} = O(\delta)\gamma^{-m}$ for all i , or all k_i are equal to each other, $k_i = k$, and $\gamma^{-m_i} = O(\delta)\lambda^k$ for all i .

Proof. Let us first consider the special case where the sequence $\{(k_i, m_i)\}$ has two pairs. By assumption the system

$$\bar{M}_1 = T_{k_1, m_1}(M_1), \quad \bar{M}_2 = T_{k_2, m_2}(M_2) \quad (56)$$

have at least one solution $(k_1, m_1, M_1, \bar{M}_1, k_2, m_2, M_2, \bar{M}_2)$ with $k_i, m_i \in \mathbb{N}$ and $M_i, \bar{M}_i \in \Pi$. This system imposes two equations involving the X -equation in (53), which are

$$\begin{aligned}b\gamma^m \mu + a_{r_1} b \lambda^{k_1} \gamma^{m_1} X_1 + a_{r_1} b x_{r_1}^+ \lambda^{k_1} \gamma^{m_1} - b u^- + \gamma^m(o(\lambda^k) + o(\gamma^{-m})) &= \bar{X}_1, \\ b\gamma^m \mu + a_{r_2} b \lambda^{k_2} \gamma^{m_2} X_2 + a_{r_2} b x_{r_2}^+ \lambda^{k_2} \gamma^{m_2} - b u^- + \gamma^m(o(\lambda^k) + o(\gamma^{-m})) &= \bar{X}_2,\end{aligned}$$

for some $r_1, r_2 \in \{0, \dots, q-1\}$. Since $X_i, \bar{X}_i \in [-\delta, \delta]$ by assumption, the new system has solutions

⁸ One notices that here we do not have condition (6). Indeed, this condition is equivalent to condition (6.17) in [23], which is to ensure that the image $T_{k,m}(M)$ of a point $M \in \Pi$ lies also in Π , and this fact then allows one to use the Implicit Function Theorem to deduce formula (6.18) (i.e., formula (7) in the present paper) from (6.12). Since we now directly assume $\bar{M} = (\bar{X}, \bar{Y}, \bar{Z}) \in \Pi$, condition (6) is no longer needed.

if and only if there exist $X_i = K_i\delta$ and $\bar{X}_i = C_i\delta$ for some $K_i, C_i \in [-1, 1]$ (where δ is the size of Π defined in (5)) such that

$$\begin{aligned}\mu + a_{r_1}\lambda^{k_1}(x_{r_1}^+ + K_1\delta) - \gamma^{-m_1}u^- &= \gamma^{-m_1}b^{-1}C_1\delta, \\ \mu + a_{r_2}\lambda^{k_2}(x_{r_2}^+ + K_2\delta) - \gamma^{-m_2}u^- &= \gamma^{-m_2}b^{-1}C_2\delta.\end{aligned}$$

Subtracting the second equation for the first one yields

$$\lambda^{k_1} - \lambda^{k_2} \frac{a_{r_2}(x_{r_2}^+ + K_2\delta)}{a_{r_1}(x_{r_1}^+ + K_1\delta)} = \gamma^{-m_1} \frac{C_1\delta + bu^-}{a_{r_1}b(x_{r_1}^+ + K_1\delta)} - \gamma^{-m_2} \frac{C_2\delta + bu^-}{a_{r_1}b(x_{r_1}^+ + K_1\delta)}. \quad (57)$$

From this point, the proof is the same as that of [23, Lemma 5.7]. Indeed, comparing (57) with (5.36) in [23], we see that the only difference is that $a_{r_1} = a_{r_2} = a$ and $x_{r_1}^+ = x_{r_2}^+ = x^+$ in [23]. Modifying the proof there in the obvious way, one finds that equation (57) has no solutions satisfying $(k_1, m_1) \neq (k_2, m_2)$ unless

$$\left| \frac{u^-}{a_{r_1}x_{r_1}^+} \right| \notin cl \left\{ |\gamma|^{\frac{s}{q'}} \cdot \frac{|a_{r_2}x_{r_2}^+(a_{r_1}x_{r_1}^+)^{-1}| - \lambda^\ell}{1 - \gamma^{-n}} \right\}_{s \in \mathbb{Z}, \ell \in \mathbb{N}, n \in \mathbb{N}}.$$

Proceeding as in [23] with considering all possible pairs (r_1, r_2) , we obtain the desired conclusion on k_i and m_i for an arbitrary two-element sequence $\{(k_1, m_1), (k_2, m_2)\}$.

To finish the proof of the lemma, let us consider a sequence $\{(k_i, m_i)\}$ with more than two elements. Apply the above result to the first two elements, we have $m_1 = m_2 = m$ or $k_1 = k_2 = k$. We assume the former case and the proof in the latter case is the same. Then, we have

$$\lambda^{k_1} = O(\delta)\gamma^{-m_1}.$$

We claim that $m_i = m$ for all i . If this was not true, then there would be some i such that $m_i \neq m = m_1$. Applying the two-element result to $\{(k_1, m_1), (k_i, m_i)\}$, yields $k_1 = k_i$ and $\gamma^{-m_1} = O(\delta)\lambda^{k_1}$, contradicting the above inequality. Thus, all m_i are the same, and the relation $\lambda^{k_i} = O(\delta)\gamma^{-m}$ is obtained by applying the two-element result to $\{(k_i, m), (k_j, m)\}$ for every (i, j) pair. \square

Recall that E_{η_i} ($i = 1, 2$) are the set of η_i values for which (52) is satisfied, and E_{r_1, r_2} are the sets in (55).

Theorem 7.3. *Let f have a non-degenerate saddle-focus heterodimensional cycle with both $\theta = p'/q'$ and $\omega/2\pi = p/q$ being rational. If*

$$\eta_i \notin E_{\eta_i}, \quad \left| \frac{u^-}{a_{r_1}x_{r_1}^+} \right| \notin E_{r_1, r_2}, \quad |a_r b| \notin \{|\gamma|^{\frac{s}{q'}}\}_{s \in \mathbb{Z}} =: E'_r,$$

with $i \in \{1, 2\}$ and $r_1, r_2, r \in \{0, \dots, q-1\}$, then any generic one-parameter unfolding family $\{f_\varepsilon\}$ with $f_0 = f$ has simple hyperbolic dynamics as in Definition 1.4.

Since each of the sets E_{η_i} , E_{r_1, r_2} and E'_r is nowhere dense in \mathbb{R} , this theorem covers the second and third items of Theorem B. The proof is essentially a word-by-word repetition of the proof of [23, Theorem 6], with formula (53) in place of [23, equation (3.20)] and Lemma 7.2 in place of [23, Lemma 5.7]. For completeness, we sketch the proof below.

Sketch of proof. First note that Lemma 7.2 holds when k_i or m_i is infinite, with taking $\lambda^\infty = 0$ and $\gamma^{-\infty} = 0$. The case $k_i = \infty$ corresponds to $M_i \in W_{loc}^s(O_1)$ and $\bar{M}_i \in F_{21} \circ F_2^{m_i} \circ F_{12}(W_{loc}^u(O_1))$. The case $m_i = \infty$ corresponds to $M_i \in F_1^{-k_i} \circ F_{12}^{-1}(W_{loc}^s(O_2))$ and $\bar{M}_i \in F_{21}(W_{loc}^u(O_2))$.

The case $k_1 = m_1 = \infty$ corresponds to the existence of the fragile heteroclinic intersection of $F_{12}(W_{loc}^u(O_1))$ with $W_{loc}^s(O_2)$, where one can take $M_1 \in W_{loc}^s(O_1)$ and $\bar{M}_1 \in T_{21}(W_{loc}^u(O_2))$. This happens at $\mu = 0$. Suppose there is another pair (k_2, m_2) such that $\bar{M}_2 = T_{k_2, m_2}(M_2)$. Then, one can easily check that applying Lemma 7.2 to the sequence $\{(k_1, m_1), (k_2, m_2)\}$ forces $k_2 = m_2 = \infty$, which implies the dynamics described in as in Definition 1.4 for $\mu = 0$.

For $\mu \neq 0$, consider the set \mathcal{L} of orbits which lie entirely in U and different from $L_{1,2}$ and from the heteroclinic orbits in the intersection of $F_{21}(W_{loc}^u(O_2))$ with $W_{loc}^s(O_1)$. Any orbit $L \in \mathcal{L}$ gives a sequence $\{M_i\}$ of intersection points with Π , satisfying $M_{i+1} = T_{k_i, m_i}(M_i)$ for some (k_i, m_i) . Note that the case $k_i = m_i = \infty$ is excluded since $F_{12}(W_{loc}^u(O_1)) \cap W_{loc}^s(O_2) = \emptyset$ at $\mu \neq 0$. Depending on $\{M_i\}$, the sequence $\{(k_i, m_i)\}$ can be infinite or finite. We modify the sequence $\{(k_i, m_i)\}$ as follows such that it is always infinite.

- If $\{M_i\}$ is infinite, then the sequence $\{(k_i, m_i)\}$ is infinite with finite k_i, m_i , and we keep it;
- If $\{M_i\}$ is finite from the left, then the leftmost point M_{i_l} lies either in $F_{21}(W_{loc}^u(O_2))$ or in $F_{21} \circ F_2^m \circ F_{12}(W_{loc}^u(O_1))$ for some m . But the former is impossible since $L \in \mathcal{L}$. In this case, we extend $\{(k_i, m_i)\}$ to an infinite sequence by adding $(k_{i_l-j}, m_{i_l-j}) = (\infty, m)$ for all $j \geq 1$;
- If $\{M_i\}$ is finite from the right, then the rightmost point M_{i_r} lies either in $W_{loc}^s(O_1)$ or in $F_1^{-k} \circ F_{12}^{-1}(W_{loc}^s(O_2))$ for some k . But the former is impossible since $L \in \mathcal{L}$. In this case, we extend $\{(k_i, m_i)\}$ by adding $(k_{i_r+j}, m_{i_r+j}) = (k, \infty)$ for all $j \geq 0$.

Lemma 7.2 then implies three possibilities:

1. there exist positive integers k and m such that $k_i = k$ and $m_i = m$ for all i and for all $L \in \mathcal{L}$. In this case, one achieves the desired result by using the fact that the map (53) is hyperbolic when $|a_r b \lambda^k \gamma^m|$ to be bounded away from 0 and 1, which is equivalent to require $|a_r b| \notin \{|\gamma|^{\frac{s}{q}}\}$, and hence the maps $T_{k,m}$ are hyperbolic with contraction in (X, Z) and expansion in Y when $0 < |a_r b \lambda^k \gamma^m| < 1$, and with contraction in Z and expansion in (X, Y) when $1 < |a_r b \lambda^k \gamma^m|$.
2. there exists positive integer m such that $m_i = m$ for all i and for all $L \in \mathcal{L}$, while some k_i are different. The desired result follows from the fact that $|a_r b \lambda^k \gamma^m| < 1$ by the relation $\lambda^{k_i} = O(\delta) \gamma^{-m}$ given by Lemma 7.2, and hence the maps $T_{k_i, m}$ are hyperbolic with contraction in (X, Z) and expansion in Y ;
3. there exists positive integer k such that $k_i = k$ for all i and for all $L \in \mathcal{L}$, while some m_i are different. The desired result follows from the fact that $|a_r b \lambda^k \gamma^m| > 1$ by the relation $\gamma^{-m_i} = O(\delta) \lambda^k$, and hence the maps $T_{k_i, m}$ are hyperbolic with contraction in Z and expansion in (X, Y) .

□

7.2 Simple hyperbolic dynamics in the double-focus case

The arguments for double-focus cycles are completely parallel. In this case, the central multipliers are $\lambda e^{\pm i\omega_1}$ and $\gamma e^{\pm i\omega_2}$, and

$$\theta = \frac{p'}{q'}, \quad \frac{\omega_1}{2\pi} = \frac{p_1}{q_1}, \quad \frac{\omega_2}{2\pi} = \frac{p_2}{q_2}$$

are rational. First note that there is a set E_{η_3} such that if $\eta_3 \in E_{\eta_3}$, then the condition (13) is satisfied. For these η_3 values, one can obtain a perturbative version of formula (14) by performing the calculation in the proof of [23, Lemma 6.6] with adding the parameter μ to the right hand side of the first equation in (6.22) there. We obtain that, for point $(U, V, W) \in \Pi$ (defined in (12)), one has

$$\begin{aligned} \bar{U} &= \ell_{r_2} \lambda^k \mu + A_{k,m} U + B_{k,m} + \phi_1(U, V, \bar{W}), \\ \bar{V} &= \phi_2(U, V, \bar{W}), \quad W = \phi_3(U, V, \bar{W}), \end{aligned} \tag{58}$$

where $A_{k,m}, B_{k,m}$ and the functions ϕ are as in (14) with smooth dependence on μ , and

$$\ell_{r_2} = \frac{2}{\sqrt{1 + \ell^2 \cos(2\pi r_2 + \eta_3)}}, \quad r_2 \in \{0, \dots, q_2 - 1\}.$$

with ℓ ⁹ being certain coefficient in DF_{12} . Denoting

$$a_{r_1, r_2} = A(2\pi r_2) \sin(2\pi r_1 + \eta_1), \quad b_{r_1, r_2} = B(2\pi r_2) \sin(2\pi r_1 + \eta_2), \quad c_{r_2} = \xi(2\pi r_2),$$

with $r_i \in \{0, \dots, q_i\}$ ($i = 1, 2$), one rewrites (58) as

$$\begin{aligned} \bar{U} &= \ell_{r_2} \lambda^k \mu + a_{r_1, r_2} \lambda^k \gamma^m U + b_{r_1, r_2} \lambda^k \gamma^m + c_{r_2} + \phi_1(U, V, \bar{W}), \\ \bar{V} &= \phi_2(U, V, \bar{W}), \quad W = \phi_3(U, V, \bar{W}). \end{aligned}$$

Arguing as in the saddle-focus case, one easily obtains a counterpart to Lemma 7.2 and hence the result covering the last item of Theorem B:

Theorem 7.4. *Let f have a non-degenerate double-focus heterodimensional cycle with $\theta = p'/q', \omega_1/2\pi = p_1/q_1, \omega_2/2\pi = p_2/q_2$ being rational. There exist nowhere dense subsets of \mathbb{R} : $E_{\eta_i}, E_{r_1, r_2, r'_1, r'_2}, E'_{r_1, r_2}$ with $i \in \{1, 2, 3\}, r_1, r'_1 \in \{0, \dots, q_1 - 1\}$ and $r_2, r'_2 \in \{0, \dots, q_2 - 1\}$ such that if*

$$\eta_i \notin E_{\eta_i}, \quad \left| \frac{1}{b_{r_1, r_2}} \right| \notin E_{r_1, r_2, r'_1, r'_2}, \quad |a_{r_1, r_2}| \notin E'_{r_1, r_2},$$

then any generic one-parameter unfolding family $\{f_\varepsilon\}$ with $f_0 = f$ has simple hyperbolic dynamics as in Definition 1.4.

Acknowledgment

I am grateful to Professor Dmitry Turaev for his invaluable guidance during my career and for bringing me the interesting topics on heterodimensional cycles. I also thank Xiaolong Li for useful discussions that improve the presentation of the paper. This work was supported by the Leverhulme Trust grant RPG 2021-072.

⁹ This is the coefficient 'a₁₄' in equation (6.22) of [23]

References

- [1] M. Asaoka, K. Shinohara and D. V. Turaev, Degenerate behavior in non-hyperbolic semigroup actions on the interval: fast growth of periodic points and universal dynamics, *Mathematische Annalen*, **368:3-4** (2017), 1277–1309. DOI: [10.1007/s00208-016-1468-0](https://doi.org/10.1007/s00208-016-1468-0)
- [2] M. Asaoka, K. Shinohara and D. V. Turaev, Fast growth of the number of periodic points arising from heterodimensional connections, *Compositio Mathematica*, **157:9** (2021), 1899–1963. DOI: [10.1112/S0010437X21007405](https://doi.org/10.1112/S0010437X21007405)
- [3] A. Avila, S. Crovisier, A. Wilkinson, C^1 density of stable ergodicity, *Advances in Mathematics*, **379** (2021), 107496. DOI: [10.1016/j.aim.2020.107496](https://doi.org/10.1016/j.aim.2020.107496)
- [4] P. Barrientos, L. Díaz and S. Pérez, Homoclinic tangencies leading to robust heterodimensional cycles, *Math. Z.*, **302** (2022), 519–558. DOI: [10.1007/s00209-022-03065-w](https://doi.org/10.1007/s00209-022-03065-w)
- [5] P. Barrientos, L. Díaz and D. Li, Nondominated cycles of codimension one, *in preparation*.
- [6] P. Barrientos, Y. Ki and A. Raibekas, Symbolic blender-horseshoes and applications, *Nonlinearity*, **27** (2014), 2805.
- [7] P. Berger, Generic family with robustly infinitely many sinks, *Inventiones mathematicae*, **205:1** (2016), 121–172. DOI: [10.1007/s00222-015-0632-6](https://doi.org/10.1007/s00222-015-0632-6)
- [8] P. Berger, Generic family displaying robustly a fast growth of the number of periodic points, *Acta Math.*, **227:2** (2021), 205–262. DOI: [10.4310/ACTA.2021.v227.n2.a1](https://doi.org/10.4310/ACTA.2021.v227.n2.a1)
- [9] S. Biebler, Lattès maps and the interior of the bifurcation locus, *J. Mod. Dyn.*, **15** (2019), 95–130. DOI: [10.3934/jmd.2019014](https://doi.org/10.3934/jmd.2019014)
- [10] J. Bochi, C. Bonatti and L. J. Díaz, Robust criterion for the existence of nonhyperbolic ergodic measures, *Comm. Math. Phys.*, **344** (2016), 751–795. DOI: [10.1007/s00220-016-2644-5](https://doi.org/10.1007/s00220-016-2644-5)
- [11] C. Bonatti and L. J. Díaz, Persistent transitive diffeomorphisms, *Annals of Mathematics*, **143:2** (1996), 357–396.
- [12] C. Bonatti, L. J. Díaz and M. Viana, *Dynamics Beyond Uniform Hyperbolicity*, Springer, Berlin, Heidelberg, New York, 2000.
- [13] C. Bonatti and L. J. Díaz, Robust heterodimensional cycles and C^1 -generic dynamics, *Journal of the Institute of Mathematics of Jussieu*, **7:3** (2008), 469–525.
- [14] C. Bonatti and L. J. Díaz, Abundance of C^1 -robust homoclinic tangencies, *Trans. Amer. Math. Soc.*, **364** (2012), 5111–5148.
- [15] C. Bonatti, L. J. Díaz and S. Kiriki, Stabilization of heterodimensional cycles, *Nonlinearity*, **25:4** (2012), 931–960.
- [16] H. W. Broer and F. M. Tangerman, From a differentiable to a real analytic perturbation theory, applications to the Kupka Smale theorems, *Ergodic Theory and Dynamical Systems*, **6(3)** (1986), 345–362.
- [17] L. J. Díaz, Robust nonhyperbolic dynamics and heterodimensional cycles, *Ergodic Theory and Dynamical Systems*, **15** (1995), 291–315.

- [18] S. V. Gonchenko, D. V. Turaev and L. P. Shilnikov, Homoclinic tangencies of arbitrarily high orders in conservative and dissipative two-dimensional maps, *Nonlinearity*, **20** (2007), 241–275.
- [19] S. V. Gonchenko, L. P. Shilnikov and D. V. Turaev, On dynamical properties of multidimensional diffeomorphisms from Newhouse regions, *Nonlinearity*, **21** (2008), 923–972.
- [20] R. Dujardin, Non density of stability for holomorphic mappings on \mathbb{C}^2 , *J. Éc. polytech. Math.*, **4** (2017), 813–843. [DOI: 10.5802/jep.57](#)
- [21] A. Katok and B. Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems (Encyclopedia of Mathematics and its Applications)*, Cambridge University Press, Cambridge, 1995.
- [22] D. Li, X. Li, K. Shinohara and D. V. Turaev, Robust heterodimensional cycles in two-parameter unfolding of homoclinic tangencies, *arXiv:2203.14075*.
- [23] D. Li and D. V. Turaev, Persistence of heterodimensional cycles, *arXiv:2105.03739*, to appear in *Inventiones mathematicae*.
- [24] D. Li and D. V. Turaev, Symplectic blenders and persistence of homoclinics to saddle-center periodic orbits, *in preparation*.
- [25] D. Li and D. V. Turaev, Typicality of robust coexistence of infinitely many sinks for families of diffeomorphisms, *in preparation*.
- [26] C. Moreira and W. Silva, On the geometry of horseshoes in higher dimensions, *arXiv:1210.2623*.
- [27] M. Nassiri and E. Pujals, Robust transitivity in Hamiltonian dynamics, *Annales scientifiques de l'École Normale Supérieure*, **45:2** (2012), 191–239.
- [28] S. E. Newhouse, J. Palis and F. Takens, Bifurcations and stability of families of diffeomorphisms, *Publications Mathématiques de l'IHÉS*, **57** (1983), 5–71.
- [29] F. Rodriguez-Hertz, M. A. Rodriguez-Hertz, A. Tahzibi and R. Ures, Creation of blenders in the conservative setting, *Nonlinearity*, **23** (2011), 201–223. [DOI: 10.1088/0951-7715/23/2/001](#)
- [30] F. Rodriguez-Hertz, M. A. Rodriguez-Hertz, A. Tahzibi and R. Ures, New criteria for ergodicity and nonuniform hyperbolicity, *Duke Math. J.*, **160:3** (2011), 599–629. [DOI: 10.1215/00127094-1444314](#)
- [31] L. P. Shilnikov, On a Poincaré-Birkhoff problem, *Math. USSR Sbornik*, **3** (1967), 91–102.
- [32] L. P. Shilnikov, A. L. Shilnikov, D. V. Turaev and L. O. Chua, *Methods Of Qualitative Theory In Nonlinear Dynamics (Part I)*, 2nd World Sci.-Singapore, New Jersey, London, Hong Kong, 2001.
- [33] S. J. van Strien, *One parameter families of vector fields*, PhD Thesis, Utrecht, 1982.
- [34] J. Taflin, Blenders near polynomial product maps of \mathbb{C}^2 , *J. Eur. Math. Soc. (JEMS)*, **23** (2017), 3555–3589.

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, UNITED KINGDOM
 E-mail address: dongchenli416@gmail.com