PHYLOGENETIC DIVERSITY INDICES FROM AN AFFINE AND PROJECTIVE VIEWPOINT

V. MOULTON, A. SPILLNER, AND K. WICKE

ABSTRACT. Phylogenetic diversity indices are commonly used to rank the elements in a collection of species or populations for conservation purposes. The derivation of these indices is typically based on some quantitative description of the evolutionary history of the species in question, which is often given in terms of a phylogenetic tree. Both rooted and unrooted phylogenetic trees can be employed, and there are close connections between the indices that are derived in these two different ways. In this paper, we introduce more general phylogenetic diversity indices that can be derived from collections of subsets (clusters) and collections of bipartitions (splits) of the given set of species. Such indices could be useful, for example, in case there is some uncertainty in the topology of the tree being used to derive a phylogenetic diversity index. As well as characterizing some of the indices that we introduce in terms of their special properties, we provide a link between cluster-based and split-based phylogenetic diversity indices that uses a discrete analogue of the classical link between affine and projective geometry. This provides a unified framework for many of the various phylogenetic diversity indices used in the literature based on rooted and unrooted phylogenetic trees, generalizations and new proofs for previous results concerning treebased indices, and a way to define some new phylogenetic diversity indices that naturally arise as affine or projective variants of each other.

1. INTRODUCTION

Evolutionary isolation metrics or phylogenetic diversity indices provide quantitative measures of biodiversity and are increasingly popular tools to prioritize species for conservation (Isaac et al., 2007; Redding et al., 2008, 2014; Redding and Mooers, 2006; Tucker et al., 2016; Vane-Wright et al., 1991). These indices quantify the importance of a species to overall biodiversity by assessing its unique and shared evolutionary history as indicated by its placement in an underlying phylogeny. Preserving phylogenetic diversity and the "Tree of Life" has become an integral component of conservation considerations (see, e.g., the "Phylogenetic Diversity Task Force"¹ initiated by the IUCN). Indeed, conservation initiatives like the EDGE of Existence programme² (Gumbs et al., 2023; Isaac et al., 2007) incorporate phylogenetic diversity indices in their identification of species that are both evolutionary distinct and globally endangered.

Date: March 26, 2024.

School of Computing Sciences, University of East Anglia, UK.

Merseburg University of Applied Sciences, Germany.

Department of Mathematical Sciences, New Jersey Institute of Technology, USA.

¹https://www.pdtf.org/

²https://www.edgeofexistence.org/



FIGURE 1. (a) A rooted phylogenetic tree on the set $X = \{a, b, c, d, e\}$ of species. The root vertex is *r* and all edges are weighted. The table gives the value $FP_r(x)$ of the fair proportion index on this rooted tree for each $x \in X$. (b) The unrooted phylogenetic tree with weighted edges on the same set *X* of species obtained by suppressing the root of the tree in (a). The table gives the value $FP_u(x)$ of the fair proportion index on this unrooted tree for each $x \in X$.

Moreover, the "guide to phylogenetic metrics for conservation, community ecology and macroecology" by Tucker et al. (2016) has been cited more than 700 times since its publication, thus demonstrating an even more widespread interest and application of phylogenetic tools, and in particular different phylogenetic diversity indices, within conservation settings.

Mathematically, with a multitude of phylogenetic diversity indices at hand, there is now an increasing interest in understanding how the different indices relate to each other. Much of the previous work in this direction has focused on comparing and analyzing different indices derived from rooted phylogenetic trees (Bordewich and Semple, 2023; Manson, 2022; Manson and Steel, 2023; Wicke and Steel, 2019). Phylogenetic diversity indices have also been defined for unrooted trees (Haake et al., 2008; Wicke and Steel, 2019), and an exploration of the relationship between indices derived via rooted and unrooted phylogenetic trees is presented by Wicke and Steel (2019).

As one might expect, phylogenetic diversity indices for rooted and unrooted trees are closely related. To illustrate this, consider the much studied *fair proportion index* (Isaac et al., 2007; Redding, 2003). For the rooted phylogenetic tree with edge weights in Figure 1(a), the value $FP_r(x)$ of the rooted fair proportion index for a species $x \in X$ is computed by adding, over all edges that are contained in the path from the root *r* to the leaf labeled by *x*, the weight of the edge divided by the total number of species for which the path from the root to the leaf labeled by that species also contains that edge. For example, for species *e* there are three edges in the path from *r* to *e* and we obtain

(1)
$$FP_r(e) = \frac{3}{3} + \frac{2}{2} + \frac{4}{1} = 6.$$

In (Wicke and Steel, 2019) the fair proportion index has also been defined for unrooted phylogenetic trees. Consider the unrooted phylogenetic tree with edge weights in Figure 1(b). The removal of an edge breaks the tree into two subtrees. The value $FP_u(x)$ of the unrooted fair proportion index for a species $x \in X$ is one half of the value obtained by adding, over all edges in the unrooted tree, the weight of the edge divided by the number of species that lie in the same subtree as x after removal of the edge. For example, for species a we obtain

(2)
$$FP_u(a) = \frac{1}{2} \cdot \left(\frac{3}{1} + \frac{2}{4} + \frac{7}{2} + \frac{6}{4} + \frac{2}{3} + \frac{1}{4} + \frac{4}{4}\right) = \frac{125}{24}.$$

While, at first glance, there is only some similarity in the way the values of the fair proportion index for rooted and unrooted phylogenetic trees are computed from the edge weights, there is a deeper connection. For example, as can be seen in Figure 1, $\sum_{x \in X} FP_r(x) = \sum_{x \in X} FP_u(x) = 25$, which is the total weight of the edges of the phylogenetic tree from which the values are computed.

To better understand this type of observation, in this paper we consider indices from the viewpoint of affine and projective clustering. This way of thinking about clustering has its origins in (Dress, 1997), and since then has become a useful tool in phylogenetic combinatorics (see, e.g., Dress 2012, Ch. 9 and Kleinman et al. 2013). More specifically, in this paper we extend the study of phylogenetic diversity indices into the more general setting of collections of *clusters* (subsets of a set) and collections of *splits* (bipartitions of a set). These settings correspond to affine and projective viewpoints of clustering, respectively (see Section 5). Considering collections of clusters and splits in general can be beneficial since it allows for the representation of data that is not tree-like or where it is difficult to determine the correct topology for a phylogenetic tree. Indeed, phylogenetic diversity indices have already been introduced for split systems (see, e.g., Abhari et al. 2023).

To illustrate this way of thinking, as hinted above, collections of clusters naturally arise when computing the rooted fair proportion index. In particular, clusters arise from rooted phylogenetic trees by taking, for each edge, the subset of species for which the path from the root to that species contains the edge (e.g. in in Figure 1(a) the edge with weight 3 next to the root gives rise to the cluster $\{c, d, e\}$). Thus, the sum used to compute the fair proportion index of e in Equation (1) is just the sum of the values $\frac{\omega(C)}{|C|}$ taken over all clusters C that contain e, where $\omega(C)$ is the weight of the edge giving rise to cluster C and |C| denotes the number of species in C. Similarly, we can interpret Equation (2) in terms of splits, using the fact that splits arise from unrooted phylogenetic trees by taking, for each edge, the split obtained by removing the edge and considering the subsets of species in the two resulting subtrees (e.g. in Figure 1(b) the edge with weight 7 gives rise to the split $\{\{a,b\}, \{c,d,e\}\}$). Then the sum used to compute the unrooted fair proportion index of *a* in Equation (2) is just the sum of the values $\frac{\lambda(S)}{2|A|}$ taken over all splits S coming from the tree, where $\lambda(S)$ is the weight of the edge giving rise to S and A is the part in S that contains a. Clearly, the sums used to compute FP_r and FP_u can be generalized to any collection of weighted clusters or splits, respectively, and, as we shall see, the more general phylogenetic diversity indices that result in this way have similar properties to their tree-based counterparts.

Thinking about phylogenetic diversity indices in an affine and projective way, leads us to two key questions that we will consider in this paper:

(i) How do properties of tree-based phylogenetic diversity indices extend to indices defined via collections of clusters and splits?

(ii) How can the relationships between collections of clusters and collections of splits be exploited to relate cluster- and split-based phylogenetic diversity indices?

In this contribution, we give answers to both of these questions, introducing the concept of phylogenetic diversity indices based on collections of clusters and splits, and giving characterizations for some of these indices in terms of their special properties. We also present a general framework to systematically relate cluster- and split-based phylogenetic diversity indices via a process that is commonly used in phylogenetic combinatorics. This provides concise proofs for generalizations of previous results for trees as well as ways to define new indices. We illustrate our new concepts and results by focusing on a few well-known tree-based phylogenetic diversity indices including the fair proportion index, the Shapely value (Haake et al., 2008; Shapley, 1953), and the equal splits index (Redding and Mooers, 2006).

The rest of this paper is structured as follows. In Section 2 we formally define cluster-based phylogenetic diversity indices and present some key properties that such indices may have. Then, in Section 3, we present a characterization of the general cluster-based fair proportion index. In Section 4 we present a characterization of the Shapley value and use its relationship to the fair proportion index to describe the first building block of our framework. In Section 5 we then give the complete framework, and illustrate some of its applications in Section 6 using the fair proportion index and the equal splits index as examples. We conclude in Section 7 discussing some potential interesting directions for future work.

2. Cluster-based indices

Let X be a non-empty finite set. We denote the power set of X by $\mathscr{P}(X)$. We call a non-empty subset $C \subseteq X$ a *cluster* on X and call a non-empty collection $\mathscr{C} \subseteq \mathscr{P}(X) \setminus \{\emptyset\}$ a *cluster system* on X. In this section we introduce the concept of a phylogenetic diversity index on a cluster system, and illustrate some basic properties of these indices by considering a generalization of the fair proportion index for rooted trees that we introduced in the introduction.

To motivate the definition of a phylogenetic diversity index on a cluster system, we briefly look again at rooted phylogenetic trees. Fixing a rooted phylogenetic tree \mathscr{T} on a set X of species, a phylogenetic diversity index Φ on \mathscr{T} assigns, to each weighting ω of the edges in \mathscr{T} , a vector $\Phi(\omega) \in \mathbb{R}^X$. To give an example, let Φ be the fair proportion index on the rooted phylogenetic tree in Figure 1(a). Then, for the weighting ω of its edges given in Figure 1(a), we can write

(3)
$$\Phi(\omega) = (5,4,7,3,6),$$

or, in more detail, $(\Phi(\omega))(a) = 5$, $(\Phi(\omega))(b) = 4$,..., $(\Phi(\omega))(e) = 6$.

As described in the introduction, each edge in a rooted phylogenetic tree on X is associated with a cluster on X. In Figure 2(a) the clusters associated with the edges of the rooted phylogenetic tree in Figure 1(a) are given, where each cluster is weighted by the length of the corresponding edge. Note that this cluster system \mathscr{C} has a special property, namely it is a *hierarchy*, that is, $C \cap C' \in \{\emptyset, C, C'\}$ holds for all $C, C' \in \mathscr{C}$. In particular, as we see in this example, hierarchies are essentially those cluster systems that can be represented by a rooted phylogenetic tree on X (see, e.g., Semple and Steel 2003, Thm. 3.5.2 for a more precise statement of this fact using the concept of a rooted X-tree).

(a)			(b)							
()	С	$\omega(C)$			a	b	С	d	е	
	$\{a\}$	3		$\{a\}$	1	0	0	0	0	
	$\{b\}$	2		$\{b\}$	0	1	0	0	0	
	$\{c\}$	6		$\{c\}$	0	0	1	0	0	
	$\{d\}$	1		$\{d\}$	0	0	0	1	0	
	$\{e\}$	4		$\{e\}$	0	0	0	0	1	
	$\{a,b\}$	4		$\{a,b\}$	$\frac{1}{2}$	$\frac{1}{2}$	0	0	0	
	$\{d,e\}$	2		$\{d,e\}$	0	0	0	$\frac{1}{2}$	$\frac{1}{2}$	
	$\{c,d,e\}$	3		$\{c,d,e\}$	0	0	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$	

FIGURE 2. (a) The weighted clusters on X corresponding to the edges of the rooted phylogenetic tree in Figure 1(a). (b) The matrix Γ from Equation (5) for the fair proportion index on \mathscr{C} , where \mathscr{C} is the cluster system consisting of the clusters given in (a).

Bearing these facts in mind, for an arbitrary cluster system \mathscr{C} on X, we consider the space $\mathbb{L}(\mathscr{C})$ consisting of all weightings $\omega : \mathscr{C} \to \mathbb{R}$. We then define a *phylogenetic diversity index* on \mathscr{C} to be a map $\Phi : \mathbb{L}(\mathscr{C}) \to \mathbb{R}^X$. For example, following the intuitive description in the introduction, we define the *fair-proportion index* on a cluster system \mathscr{C} on X by putting, for each $\omega \in \mathbb{L}(\mathscr{C})$ and all $x \in X$,

(4)
$$(FP(\boldsymbol{\omega}))(x) = \sum_{C \in \mathscr{C}: \ x \in C} \frac{\boldsymbol{\omega}(C)}{|C|}.$$

It can then be checked that (4) applied to the weighted cluster system in Figure 2(a) yields precisely the vector we saw in (3).

We now introduce three key properties of cluster-based indices which generalize properties of tree-based indices described in the literature. We will illustrate these properties for the fair proportion index and, as we shall see, these properties are also shared by some of the other phylogenetic diversity indices that we consider later on.

Let \mathscr{C} be a cluster system on X. A phylogenetic diversity index Φ on \mathscr{C} is *additive* if

(A)
$$\Phi(\omega_1 + \omega_2) = \Phi(\omega_1) + \Phi(\omega_2)$$
 for all $\omega_1, \omega_2 \in \mathbb{L}(\mathscr{C})$,

and Φ is *homogeneous* if

(H)
$$\Phi(a \cdot \omega) = a \cdot \Phi(\omega)$$
 for all $\omega \in \mathbb{L}(\mathscr{C})$ and all $a \in \mathbb{R}$.

Properties (A) and (H) together mean that Φ is a *linear map*, in which case we call Φ *linear*. Linearity is a natural property to require even on a rooted phylogenetic tree \mathscr{T} , where it implies, for example, that, applying the phylogenetic diversity index to a weighting obtained by taking the average over several different edge weightings of \mathscr{T} amounts to averaging the phylogenetic diversity index. Note that every linear phylogenetic diversity index Φ on \mathscr{C} corresponds to a $|\mathscr{C}| \times |X|$ -matrix $\Gamma = \Gamma_{\Phi} = (\gamma_{(C,x)})$ such that

(5)
$$(\Phi(\omega))(x) = \sum_{C \in \mathscr{C}} \omega(C) \cdot \gamma_{(C,x)}$$

for all $\omega \in \mathbb{L}(\mathscr{C})$ and all $x \in X$. For example, the matrix Γ corresponding to the fair proportion index on the cluster system in Figure 2(a) is given in Figure 2(b). Finally, we call a phylogenetic diversity index Φ on \mathscr{C} complete if

(C)
$$\sum_{x \in X} (\Phi(\omega))(x) = \sum_{C \in \mathscr{C}} \omega(C)$$
 holds for all $\omega \in \mathbb{L}(\mathscr{C})$.

For tree-based phylogenetic diversity indices, completeness is often required as part of their definition (see, e.g., Bordewich and Semple 2023; Wicke and Steel 2019). For example, we have seen in the introduction for the fair proportion index on a rooted phylogenetic tree that $\sum_{x \in X} FP_r(x)$ equals the total weight of the edges in the tree. Property (C) expresses this fact in terms of clusters. Note that if a linear phylogenetic diversity index Φ is complete, then we have $\sum_{x \in X} \gamma_{(C,x)} = 1$ for all $C \in \mathscr{C}$ (cf. Wicke and Steel 2019, Eq. (2)), where $\Gamma = (\gamma_{(C,x)})$ is the matrix from Equation (5).

We conclude this section by showing that the fair proportion index satisfies all three of the above properties.

Lemma 2.1. The fair proportion index is a complete, linear phylogenetic diversity index on \mathscr{C} for any cluster system \mathscr{C} on X.

Proof: As we have seen in the example in Figure 2, the fair proportion index can be described by a matrix $\Gamma = (\gamma_{(C,x)})$ where the row associated with a cluster $C \in \mathscr{C}$ contains |C| entries equal to $\frac{1}{|C|}$ and |X| - |C| entries equal to 0.

3. A CHARACTERIZATION OF THE FAIR PROPORTION INDEX

In general, it is of interest to characterize phylogenetic diversity indices in terms of their key properties, as this can help to understand better how they are related to one another. In this section, as an illustration for cluster-based indices, we shall present a characterization of the fair proportion index. This generalizes the characterization of the fair proportion index on rooted phylogenetic trees given by Manson and Steel (2023, Thm. 6).

Our characterization will require three properties. The first two properties concern linear phylogenetic diversity indices Φ on a cluster system \mathscr{C} on X, and are given in terms of the matrix corresponding to Φ . For all $C \in \mathscr{C}$, let ch(C) denote the set of those $C' \in \mathscr{C}$ with $C' \subsetneq C$ such that there is no $C'' \in \mathscr{C}$ with $C' \subsetneq C'' \subsetneq C$. We say that Φ satisfies the *neutrality condition* if

(NC) the entries of the matrix Γ_{Φ} in Equation (5) are such that $\gamma_{(C,x)} = \gamma_{(C,y)}$ holds for all $C \in \mathscr{C}$ with $ch(C) = \emptyset$ and all $x, y \in C$.

A property similar to (NC) was introduced by Manson and Steel (2023) for rooted X-trees. In addition, we say that Φ is a *descendant diversity index* if

(DD) Φ is complete, all entries of the matrix Γ_{Φ} in Equation (5) are non-negative and, for all $C \in \mathscr{C}, \gamma_{(C,x)} = 0$ if $x \notin C$.

Property (DD) was introduced by Bordewich and Semple (2023) for the special case where the cluster system \mathscr{C} is a hierarchy (using the equivalent description of hierarchies in terms rooted *X*-trees).



FIGURE 3. Collapsing the edge with weight 6 in the rooted phylogenetic tree \mathscr{T} on $X = \{a, b, c, d\}$ yields the rooted phylogenetic tree \mathscr{T}^* on *X*.

The third property is a bit more complicated, and thus we first motivate it using rooted trees as in (Manson and Steel, 2023). Let \mathscr{T} be a rooted phylogenetic tree on X with edge weights and let \mathscr{T}^* be the rooted phylogenetic tree on X obtained by collapsing one of the edges of \mathscr{T} . This is illustrated in Figure 3. In addition, let Φ and Φ^* be phylogenetic diversity indices on \mathscr{T} and \mathscr{T}^* , respectively. Both Φ and Φ^* yield a vector in \mathbb{R}^X for all weightings of the edges of \mathscr{T} and \mathscr{T}^* , respectively. The topology of the rooted phylogenetic trees, however, may have an impact on how the weights of the edges are used to compute these vectors by Φ and Φ^* , respectively. Therefore, since the topologies of \mathscr{T} and \mathscr{T}^* differ, the vector in \mathbb{R}^X that we obtain by Φ^* for \mathscr{T}^* will usually not coincide with the vector that we obtain in the limit, as the weight of the edge in \mathscr{T} tends to 0, by Φ (keeping the weights of all other edges in \mathscr{T} in constant).

With this in mind, let \mathscr{C} be a cluster system on X and let $C \in \mathscr{C}$ such that $\mathscr{C}^* = \mathscr{C} \setminus \{C\}$ is non-empty. A phylogenetic diversity index Φ on \mathscr{C} is *downward continuous* with respect to a phylogenetic diversity index Φ^* on \mathscr{C}^* if

(DC) for all $\omega \in \mathbb{L}(\mathscr{C})$ we have

(6)
$$\lim_{\omega(C)\to 0} \Phi(\omega) = \Phi^*(\omega^*),$$

where $\omega^* \in \mathbb{L}(\mathscr{C}^*)$ is the weighting with $\omega^*(D) = \omega(D)$ for all $D \in \mathscr{C}^*$.

With the properties (NC), (DD) and (DC) in hand, we now present our characterization of the fair proportion index.

Theorem 3.1. Suppose we have, for each cluster system \mathscr{C} on X, a phylogenetic diversity index $\Phi_{\mathscr{C}}$ on \mathscr{C} . Then the following are equivalent:

- (i) For all cluster systems \mathscr{C} on X, $\Phi_{\mathscr{C}}$ is the fair proportion index on \mathscr{C} .
- (ii) For all cluster systems \mathscr{C} on X, $\Phi_{\mathscr{C}}$ is a descendant diversity index that satisfies the neutrality condition and is downward continuous with respect to $\Phi_{\mathscr{C}\setminus\{C\}}$ for all $C \in \mathscr{C}$ such that $\mathscr{C}\setminus\{C\} \neq \emptyset$.

Proof: We first show that (i) implies (ii). Consider a cluster system \mathscr{C} on X and put $\Phi = \Phi_{\mathscr{C}}$. By assumption, Φ is the fair proportion index on \mathscr{C} . Thus, in view of Lemma 2.1, Φ is linear and complete. Moreover, as illustrated by the example in Figure 2(b), it follows immediately from the definition of the fair proportion index in (4) that Φ is a descendant diversity index and satisfies the neutrality condition.

It remains to establish downward continuity. Consider a cluster $C \in \mathscr{C}$ and assume that $\mathscr{C}^* = \mathscr{C} \setminus \{C\} \neq \emptyset$. Put $\Phi^* = \Phi_{\mathscr{C}^*}$. Let $\Gamma = \Gamma_{\Phi}$ and $\Gamma^* = \Gamma_{\Phi^*}$ be the matrices whose entries satisfy Equation (5) for Φ and Φ^* , respectively. By assumption, Φ is the fair proportion index on \mathscr{C} and Φ^* is the fair proportion index on \mathscr{C}^* Therefore, it follows again from the definition of the fair proportion index in (4) that deleting the row corresponding to the cluster *C* from the matrix Γ yields the matrix Γ^* . But this immediately implies that Equation (6) holds for all $\omega \in \mathbb{L}(\mathscr{C})$, as required.

Next we show that (ii) implies (i). Let \mathscr{C} be a cluster system on *X*. By assumption, $\Phi = \Phi_{\mathscr{C}}$ is a descendant diversity index and, therefore, linear. Let $\Gamma = \Gamma_{\Phi}$ be the matrix whose entries satisfy Equation (5) for Φ . In view of the definition of the fair proportion index in (4), it suffices to show that the entries of Γ satisfy

$$Y_{(C,x)} = \begin{cases} \frac{1}{|C|} & \text{for } x \in C\\ 0 & \text{for } x \notin C \end{cases}$$

for all $C \in \mathscr{C}$ and all $x \in X$. We use induction on $|\mathscr{C}|$ to show this.

To establish the base case of the induction, assume $|\mathscr{C}| = 1$. Consider $C \in \mathscr{C}$ and $x \in X$. In view of $|\mathscr{C}| = 1$ we have $ch(C) = \emptyset$. Thus, in view of the assumption that Φ is a descendant diversity index and satisfies the neutrality condition, we have $\gamma_{(C,x)} = \frac{1}{|C|}$ for all $x \in C$ and $\gamma_{(C,x)} = 0$ for all $x \in X \setminus C$, as required.

Next assume $|\mathscr{C}| \geq 2$. Consider $C \in \mathscr{C}$ and put $\mathscr{C}^* = \mathscr{C} \setminus \{\mathscr{C}\}$. By the assumption that Φ is downward continuous with respect to $\Phi^* = \Phi_{\mathscr{C}^*}$, the matrix $\Gamma^* = \Gamma_{\Phi^*}$ whose entries satisfy Equation (5) for Φ^* is obtained by deleting the row corresponding to cluster *C* from Γ . Thus, by induction, we have

$$\gamma_{(D,x)} = \begin{cases} \frac{1}{|D|} & \text{for } x \in D\\ 0 & \text{for } x \notin D \end{cases}$$

for all $D \in \mathscr{C} \setminus \{C\}$ and all $x \in X$. Since this holds for all $C \in \mathscr{C}$, this finishes the inductive proof.

4. The Shapely value

The Shapely value is a well-known phylogenetic diversity index that can be computed using rooted phylogenetic trees and that has its origins in game theory. Interestingly, to understand a generalization of this index in the cluster setting, it is necessary to consider mappings on slightly more general spaces than those used in the definition of cluster-based phylogenetic diversity indices in Section 2. In this section, we shall explain this, and then give a characterization of a cluster-based version of the Shapely value.

Let *X* be a finite set. A *game* is a map $g : \mathscr{P}(X) \to \mathbb{R}$. The elements of *X* are referred to as the *players* in this context and the value g(C) for some $C \in \mathscr{P}(X)$ can be interpreted as the gain when the players in *C* form a coalition. The aim of the game is then to quantify, for each player $x \in X$, the value $v(x) \in \mathbb{R}$ of the player with respect to a given game (see, e.g., Branzei et al. 2008 for a more detailed exposition of these concepts). Formally speaking, we are thus interested in

maps v from $\mathbb{R}^{\mathscr{P}(X)}$ to \mathbb{R}^X , and the *Shapley value* is one specific such map v given by

(7)
$$(v(g))(x) = \frac{1}{|X|!} \cdot \sum_{M \in \mathscr{P}(X): \ x \in M} \left[(|M| - 1)! \cdot (|X| - |M|)! \cdot (g(M) - g(M \setminus \{x\})) \right]$$

This map was originally proposed by Shapley (1953).

In a biological context, the players of Shapley's game are species and from a rooted phylogenetic tree \mathscr{T} on X with edge weights we obtain a game g by setting g(M) = PD(M) for each $M \in \mathscr{P}(X)$, where PD(M) is the *phylogenetic diversity* of M. The value PD(M) is defined as the total weight of those edges in T that lie on a path from the root to some species in M (Faith, 1992). For example, for the rooted phylogenetic tree in Figure 1(a) we obtain

 $PD(\{a, b, d\}) = 3 + 2 + 4 + 3 + 2 + 1 = 15.$

We now explain a way to generalize these considerations to cluster systems \mathscr{C} on *X*. First we need to define the phylogenetic diversity of a subset of *X* relative to a weighted cluster system. Let $\omega \in \mathbb{L}(\mathscr{C})$. Then the *phylogenetic diversity* of a subset *M* of *X* with respect to ω is defined as

(8)
$$PD(M) = PD_{\omega}(M) = \sum_{C \in \mathscr{C}: \ M \cap C \neq \emptyset} \omega(C).$$

To further explore properties of the Shapley value in the context of our work, it will be convenient to consider the set

 $\mathbb{PD}(\mathscr{C}) = \{g \in \mathbb{R}^{\mathscr{P}(X)} : \text{ there exists } \omega \in \mathbb{L}(\mathscr{C}) \text{ with } g = PD_{\omega}\},\$

that is, the set of games in $\mathbb{R}^{\mathscr{P}(X)}$ for which there is some $\omega \in \mathbb{L}(\mathscr{C})$ which gives rise to this game. The next technical lemma summarizes the key structural properties of this set.

Lemma 4.1. Let \mathscr{C} be a cluster system on X. Then $\mathbb{PD}(\mathscr{C})$ is a linear subspace of $\mathbb{R}^{\mathscr{P}(X)}$ that has dimension $|\mathscr{C}|$.

Proof: For $C \in \mathscr{P}(X)$, let $g_C : \mathscr{P}(X) \to \mathbb{R}$ be the game defined by putting

(9)
$$g_C(M) = \begin{cases} 1 & \text{if } C \cap M \neq \emptyset \\ 0 & \text{if } C \cap M = \emptyset. \end{cases}$$

In view of (8) and (9), $\mathbb{PD}(\mathscr{C})$ is the linear span of the games g_C for $C \in \mathscr{C}$:

$$PD_{\omega}(M) = \sum_{C \in \mathscr{C}: \ M \cap C \neq \emptyset} \omega(C) = \sum_{C \in \mathscr{C}} \omega(C) \cdot g_C(M)$$

Thus, it suffices to show that the games g_C , $C \in \mathscr{C}$, are linearly independent. To see this, consider the square matrix A whose rows and columns are each in one-to-one correspondence with the elements of $\mathscr{P}(X) \setminus \{\emptyset\}$. For all $C, M \in \mathscr{P}(X) \setminus \{\emptyset\}$ the entry of A in the row corresponding to C and the column corresponding to M is 1 if $C \cap M \neq \emptyset$ and is 0 otherwise. A is the so-called *intersection matrix* of $\mathscr{P}(X) \setminus \{\emptyset\}$ and it is known that A has full rank (see, e.g., Jukna 2011, p. 216). Thus, in particular, the rows corresponding to $C \in \mathscr{C}$ are linearly independent.

Now, as explained above, for a cluster system \mathscr{C} on *X*, we restrict in (7) to games g = PD in $\mathbb{PD}(\mathscr{C})$. More specifically, we define the Shapley value *relative to the cluster system* \mathscr{C} as the map $SV : \mathbb{PD}(\mathscr{C}) \to \mathbb{R}^X$ obtained by putting

(10)
$$(SV(PD))(x) = \frac{1}{|X|!} \cdot \sum_{M \in \mathscr{P}(X): \ x \in M} \left[(|M| - 1)! \cdot (|X| - |M|)! \cdot (PD(M) - PD(M \setminus \{x\})) \right]$$

for all $PD \in \mathbb{PD}(\mathscr{C})$ and all $x \in X$. Note that, in view of Lemma 4.1, $\mathbb{PD}(\mathscr{C})$ may be a proper subspace of $\mathbb{R}^{\mathscr{P}(X)}$ (i.e. the set of all games). As we will see below, any characterization of the Shapley value relative to a cluster system must take this into account (see also Dubey 1975 for a more general discussion of this aspect).

The sharp-eyed reader will have noticed that the Shapley value relative to a cluster system \mathscr{C} is *not* a phylogenetic diversity index on \mathscr{C} , as the latter is defined as a map from $\mathbb{L}(\mathscr{C})$ to \mathbb{R}^X . However, we can resolve this issue by slightly generalizing our cluster-based definition of phylogenetic diversity indices. Let \mathbb{L} be a linear subspace of $\mathbb{R}^{\mathscr{P}(X)}$. Then we define a phylogenetic diversity index on \mathbb{L} to be a map $\Phi : \mathbb{L} \to \mathbb{R}^X$. This encompasses then the Shapley value as a phylogenetic diversity index on $\mathbb{L} = \mathbb{PD}(\mathscr{C})$ for all cluster systems \mathscr{C} on X. Moreover, viewing $\mathbb{L}(\mathscr{C})$ as the linear subspace

$$\mathbb{L} = \{ \boldsymbol{\omega} \in \mathbb{R}^{\mathscr{P}(X)} : \boldsymbol{\omega}(C) = 0 \text{ for all } C \notin \mathscr{C} \},\$$

it also encompasses phylogenetic diversity indices on \mathscr{C} as defined in Section 2. In fact, we can say even more about these relationships, which we will return to in the next section.

For the remainder of this section, we focus on giving a characterization of the Shapley value relative to a cluster system. This will involve the following two properties. We say that a phylogenetic diversity index Φ on a linear subspace \mathbb{L} of $\mathbb{R}^{\mathscr{P}(X)}$ satisfies *Pareto efficiency* if

(PE) $\sum_{x \in X} (\Phi(\omega))(x) = \omega(X)$ for all $\omega \in \mathbb{L}$.

Remark 4.2. The properties of completeness and Pareto efficiency are tightly linked. Let \mathscr{C} be a cluster system on X and note that $\sum_{C \in \mathscr{C}} \omega(C) = PD_{\omega}(X)$ holds for all $\omega \in \mathbb{L}(\mathscr{C})$. Therefore, every complete phylogenetic diversity index Φ on $\mathbb{L}(\mathscr{C})$ corresponds to a phylogenetic diversity index Φ' on $\mathbb{PD}(\mathscr{C})$ that satisfies Pareto efficiency, where Φ' is obtained such that the diagram in Figure 4 commutes, that is,

(11) $\Phi'(PD_{\omega}) = \Phi(\omega)$

for all $\omega \in \mathbb{L}(\mathscr{C})$.

We say that a phylogenetic diversity index Φ on a linear subspace \mathbb{L} of $\mathbb{R}^{\mathscr{P}(X)}$ satisfies *group proportionality* (cf. Haake et al. 2008) if

(GP)
$$(\Phi(a \cdot g_C))(x) = \begin{cases} \frac{a}{|C|} & \text{if } x \in C \\ 0 & \text{if } x \notin C, \end{cases}$$
 for all $C \in \mathscr{P}(X) \setminus \{\emptyset\}$ and all $a \in \mathbb{R}$

with g_C the game as defined in (9). Note that a similar characterization to that given in the following theorem was established by Wicke and Steel (2019, Thm. 7) for the special case of cluster systems that form a hierarchy.



FIGURE 4. This diagram depicts the relationship between a phylogenetic diversity index Φ on $\mathbb{L}(\mathscr{C})$ for a cluster system \mathscr{C} on X and a phylogenetic diversity index Φ' on $\mathbb{PD}(\mathscr{C})$ as described by Equation (11).

Theorem 4.3. Let \mathscr{C} be a cluster system on X. The Shapley value is the unique phylogenetic diversity index on $\mathbb{PD}(\mathscr{C})$ that is additive and satisfies Pareto efficiency and group proportionality.

Proof: Assume that Φ' is the Shapley value on $\mathbb{PD}(\mathscr{C})$. It is known (see, e.g., Aumann 1994) that Φ' satisfies Pareto efficiency for all $\omega \in \mathbb{R}^{\mathscr{P}(X)}$ and is additive for all $\omega_1, \omega_2 \in \mathbb{R}^{\mathscr{P}(X)}$. Thus, these two properties hold, in particular, for all $\omega, \omega_1, \omega_2 \in \mathbb{PD}(\mathscr{C}) \subseteq \mathbb{R}^{\mathscr{P}(X)}$.

To establish that Φ' also satisfies group proportionality, consider $x \in X$, $C \in \mathscr{C}$ and $a \in \mathbb{R}$. We calculate the value $(\Phi'(a \cdot g_C))(x)$ using Formula (10) (similar calculations are used in the proofs of Haake et al. 2008, Thm. 4 and Coronado et al. 2018, Thm. 1):

If $x \notin C$ we have $g_C(M \cup \{x\}) - g_C(M) = 0$ for all $M \in \mathscr{P}(X)$, implying $\Phi'(a \cdot g_C))(x) = 0$, as required. So assume that $x \in C$, put c = |C| and put n = |X|. Then, in view of the fact that only $M \in \mathscr{P}(X)$ with $M \cap C = \emptyset$ contribute to $(\Phi'(a \cdot g_C))(x)$, we have

$$(\Phi'(a \cdot g_C))(x) = \frac{a}{n!} \cdot \sum_{m=0}^{n-c} m! \cdot (n-m-1)! \cdot \binom{n-c}{m} = \frac{a \cdot (n-c)! \cdot (c-1)!}{n!} \cdot \sum_{j=c-1}^{n-1} \binom{j}{c-1} = a \cdot \frac{(n-c)! \cdot (c-1)!}{n!} \cdot \binom{n}{c} = \frac{a}{c},$$

as required, where we used the formula for the sum along a diagonal in Pascal's triangle to obtain the first equality in the second line.

Uniqueness now follows from the fact that, in view of the proof of Lemma 4.1, $\mathbb{PD}(\mathscr{C})$ is the linear span of $\{g_C : C \in \mathscr{C}\}$.

Interestingly, as shown by Fuchs and Jin (2015), the vector in \mathbb{R}^X that results from computing the Shapley value on the game *PD* obtained from an edge-weighted rooted phylogenetic tree always coincides with the vector that we obtain by computing the fair proportion index on the rooted phylogenetic tree. In fact, this is a particular instance of (11). The following Corollary of Theorem 4.3 makes this more precise.

Corollary 4.4. Let \mathscr{C} be a cluster system on X, Φ be the fair proportion index on $\mathbb{L}(\mathscr{C})$, and Φ' be the Shapley value on $\mathbb{PD}(\mathscr{C})$. Then

$$\Phi(\omega) = \Phi'(\sum_{C \in \mathscr{C}} \omega(C) \cdot g_C) = \Phi'(PD_{\omega})$$

holds for all $\omega \in \mathbb{L}(\mathscr{C})$ *.*

Proof: This follows immediately from the definition of the fair proportion index together with the fact that, by Theorem 4.3, the Shapley value is additive and satisfies group proportionality.

It is remarked in the discussion by Coronado et al. (2018) that Corollary 4.4 can also be derived using arguments based on so-called phylogenetic networks (for more on the connection between such networks and diversity indices see Section 7). Moreover, the fact that the Shapley value on $\mathbb{PD}(\mathscr{C})$ satisfies Pareto efficiency means that it apportions the phylogenetic diversity of *X* among the elements of *X*. In view of Corollary 4.4 this then also holds for the fair proportion index on $\mathbb{L}(\mathscr{C})$ and, in view of Remark 4.2, this corresponds to the fact that the fair proportion index is complete, as can be seen in the example in Figure 2(a):

$$PD_{\omega}(X) = \sum_{C \in \mathscr{C}} \omega(C) = 25 = \sum_{x \in X} (FP(\omega))(x).$$

5. AN AFFINE AND PROJECTIVE FRAMEWORK FOR PHYLOGENETIC DIVERSITY INDICES

As mentioned in the introduction, the notion of phylogenetic diversity indices has also been considered on unrooted phylogenetic trees (Haake et al., 2008; Wicke and Steel, 2019) and, just as rooted phylogenetic trees can be encoded by a collection of clusters, unrooted phylogenetic trees on a set X of species can be encoded by a collection \mathscr{S} of bipartitions, or splits, of X (see, e.g., Steel 2016, Ch. 2). In the area of phylogenetic combinatorics, the interplay between collections of clusters and collections of splits has been studied in terms of affine and projective models of clustering, respectively, in analogy with the interplay between affine and projective geometry in classical geometry (Dress 2012, p. 207; see also Dress 1997). One of the key ideas that we will exploit from this theory is that we can map a collection \mathscr{S} of splits of X in a natural way to a cluster system $\mathscr{C}(\mathscr{S})$ on X (defined in (13) below) and, in this way, derive split-based indices from cluster-based indices. In this section, we will make this more precise, and illustrate the resulting framework using the fair proportion index and the Shapely value as examples.

First, we formally define the concepts mentioned above. A *split S* of *X* is a bipartition of *X* into two non-empty subsets *A* and *B*, that is, $A \cup B = X$ and $A \cap B = \emptyset$. We denote such a split as an unordered pair A|B = B|A. A *split system* \mathscr{S} on *X* is a non-empty set of splits of *X*. By $\mathscr{S}(X)$ we denote the set of all splits of *X* and, for a split system $\mathscr{S} \subseteq \mathscr{S}(X)$, we denote by $\mathbb{L}(\mathscr{S})$ the set of all weightings $\lambda : \mathscr{S}(X) \to \mathbb{R}$ with $\lambda(S) = 0$ for all $S \in \mathscr{S}(X) \setminus \mathscr{S}$. In addition, we denote by $\mathbb{PD}(\mathscr{S})$ the set of all weightings $PD : \mathscr{P}(X) \to \mathbb{R}$ that can be written as

(12)
$$PD(M) = PD_{\lambda}(M) = \sum_{A|B \in \mathscr{S}: A \cap M \neq \emptyset, B \cap M \neq \emptyset} \lambda(A|B)$$

for some $\lambda \in \mathbb{L}(\mathscr{S})$. The value $PD_{\lambda}(M)$ is usually called the *phylogenetic diversity* of *M* with respect to the weighting λ of the splits in \mathscr{S} (see, e.g., Spillner et al. 2008).



FIGURE 5. A diagram of the various maps we consider to study relationships between phylogenetic diversity indices. The left part of the diagram we have already seen in Figure 4. In analogy to this, the right part of the diagram depicts phylogenetic diversity indices Ψ and Ψ' on $\mathbb{L}(\mathscr{S})$ and $\mathbb{PD}(\mathscr{S})$, respectively, where \mathscr{S} is a split system on X. Finally τ associates with each weighting λ of the splits in \mathscr{S} a weighting $\omega = \tau(\lambda)$ of the clusters in a cluster system $\mathscr{C} = \mathscr{C}(\mathscr{S})$ that arises from \mathscr{S} by (13).

Figure 5 gives an overview of the various spaces we shall consider and the maps between them. In addition to the maps already introduced in in Figure 4 in Section 4, we also consider, for split systems \mathscr{S} on X, maps τ from $\mathbb{L}(\mathscr{S})$ to $\mathbb{L}(\mathscr{C})$ where \mathscr{C} is the cluster system

(13)
$$\mathscr{C}(\mathscr{S}) = \bigcup_{S \in \mathscr{S}} S$$

on X mentioned above. In particular, we are interested in maps τ for which various parts of the diagram in Figure 5 commute.

As an illustration of this setup, we now revisit the relationship between the fair proportion index and the Shapely value. Let \mathscr{S} be a split system on X. Then the *Shapley value* on $\mathbb{PD}(\mathscr{S})$ is defined as in (10). Equivalently, as shown by Haake et al. (2008) for trees and by Volkmann et al. (2014) for split systems in general, the Shapley value on $\mathbb{PD}(\mathscr{S})$ can also be computed as

(14)
$$(SV(PD_{\lambda}))(x) = \sum_{A|B \in \mathscr{S}: x \in A} \frac{|B|}{|X| \cdot |A|} \cdot \lambda(A|B).$$

for all $\lambda \in \mathbb{L}(\mathscr{S})$ and all $x \in X$.

Now consider the map $\tau : \mathbb{L}(\mathscr{S}) \to \mathbb{L}(\mathscr{C}(\mathscr{S}))$ defined by putting, for $\lambda \in \mathbb{L}(\mathscr{S})$,

(15)
$$(\tau(\lambda))(A) = \frac{|B|}{|X|} \cdot \lambda(A|B) \text{ and } (\tau(\lambda))(B) = \frac{|A|}{|X|} \cdot \lambda(A|B)$$

for all $A, B \in \mathscr{C}(\mathscr{S})$ such that A|B is a split in \mathscr{S} . For example, consider the split system \mathscr{S} with weighting λ in Figure 6(a). Using Formula (14), we obtain $SV(a) = \frac{11}{2}$ in this example and we also have $FP(a) = \frac{11}{2}$ for the fair proportion index as defined in (4) applied to the cluster system $\mathscr{C}(\mathscr{S})$ with weighting $\omega = \tau(\lambda)$. We conclude this section by showing that this is not a coincidence.

(a)	S	$\lambda(S)$	$^{(b)}$ C	(\mathcal{O})
-	5	A(D)		$\omega(C)$
	$\{a,b\} \{c,d,e\}$	10	$\{c\}$	16
	$\{a,c\} \{b,d,e\}$	5	$\{a,b\}$	6
	$\{c\} \{a,b,d,e\}$	20	$\{a, c\}$	3
			$\{c, d, e\}$	4
			$\{b, d, e\}$	2
			$\{a, b, d, e\}$	4

FIGURE 6. (a) A split system \mathscr{S} on $X = \{a, b, c, d, e\}$ with weighting λ . (b) The associated cluster system $\mathscr{C}(\mathscr{S})$ on X as defined in (13) and the weighting $\omega = \tau(\lambda)$ as defined in (15).

Theorem 5.1. Let \mathscr{S} be a split system on X, Φ be the fair proportion index on $\mathbb{L}(\mathscr{C}(\mathscr{S}))$ and Ψ' be the Shapley value on $\mathbb{PD}(\mathscr{S})$. If τ is as defined in (15), then

(16)
$$\Phi(\tau(\lambda)) = \Psi'(PD_{\lambda})$$

for all $\lambda \in \mathbb{L}(\mathscr{S})$.

Proof: Let $\lambda \in \mathbb{L}(\mathscr{S})$ and put $\omega = \tau(\lambda)$. Since the maps Φ , τ , Ψ' and *PD* are all linear, it suffices to show Equation (16) for the case that one element of \mathscr{S} , say S = A|B has weight 1 (i.e. $\lambda(A|B) = 1$), whereas $\lambda(S') = 0$ for all $S' \neq S$. Then we have $\omega(A) = |B|/|X|$, $\omega(B) = |A|/|X|$, and $\omega(C) = 0$ for all $C \in \mathscr{C}(\mathscr{S})$ with $C \neq A, B$. Now let $x \in X$, and assume without loss of generality that $x \in A$. Then,

$$(\Phi(\boldsymbol{\omega}))(x) = \sum_{C \in \mathscr{C}(\mathscr{S}): \ x \in C} \frac{\boldsymbol{\omega}(C)}{|C|} = \frac{\boldsymbol{\omega}(A)}{|A|} = \frac{\frac{|B|}{|X|}}{|A|} = \frac{|B|}{|X| \cdot |A|}.$$

ומו

On the other hand, in view of (14) we have

$$(\Psi'(PD_{\lambda}))(x) = \sum_{A'|B' \in \mathscr{S}: \ x \in A'} \frac{|B'|}{|X| \cdot |A'|} \lambda(S) = \frac{|B|}{|X| \cdot |A|}$$

as well. This completes the proof.

6. COMPLETE DIVERSITY INDICES

In this section we shall consider Figure 5 once again, considering an alternative definition for the map τ that can be used to translate, for any split system \mathscr{S} on X, the property of completeness from a cluster-based index Φ on $\mathbb{L}(\mathscr{C}(\mathscr{S}))$ to an associated split-based index $\Psi = \Psi_{\tau}(\Phi)$ on $\mathbb{L}(\mathscr{S})$. In particular, we will see that this immediately implies the completeness of the fair proportion index on unrooted phylogenetic trees that was established by Wicke and Steel (2019) (for example, see Figure 1(b) in the introduction). In addition, we illustrate the application of these considerations to a generalization of the so-called equal splits index that appears in (Wicke and Steel, 2019).

We begin by proving a result concerning completeness. Let \mathscr{S} be a split system on X. A phylogenetic diversity index Ψ on $\mathbb{L}(\mathscr{S})$ is *complete* if

(C') $\sum_{x \in X} (\Psi(\lambda))(x) = \sum_{S \in \mathscr{S}} \lambda(S)$ holds for all $\lambda \in \mathbb{L}(\mathscr{S})$. Define the map $\tau : \mathbb{L}(\mathscr{S}) \to \mathbb{L}(\mathscr{C}(\mathscr{S}))$ by putting

(17)
$$(\tau(\lambda))(C) = \frac{1}{2} \cdot \lambda(C|(X-C))$$

for all $C \in \mathscr{C}(\mathscr{S})$. Then, for a phylogenetic diversity index Φ on $\mathbb{L}(\mathscr{C}(\mathscr{S}))$, we define the phylogenetic diversity index $\Psi = \Psi_{\tau}(\Phi)$ on $\mathbb{L}(\mathscr{S})$ by putting $\Psi(\lambda) = \Phi(\tau(\lambda))$ for all $\lambda \in \mathbb{L}(\mathscr{S})$.

Theorem 6.1. Let \mathscr{S} be a split system on X and Φ a complete linear phylogenetic diversity index on $\mathbb{L}(\mathscr{C}(\mathscr{S}))$. If τ is as defined in (17), then $\Psi_{\tau}(\Phi)$ is a complete linear phylogenetic diversity index on $\mathbb{L}(\mathscr{S})$.

Proof: Let Φ be a complete linear phylogenetic diversity index on $\mathbb{L}(\mathscr{C}(\mathscr{S}))$. We first show that Ψ_{τ} is linear. Let $\lambda_1, \lambda_2 \in \mathbb{L}(\mathscr{S})$ and $a \in \mathbb{R}$. Then, noting that τ is linear, we have

$$\begin{aligned} (\Psi_{\tau}(\Phi))(a \cdot \lambda_1 + \lambda_2) &= \Phi(\tau(a \cdot \lambda_1 + \lambda_2)) = \Phi(a \cdot \tau(\lambda_1) + \tau(\lambda_2)) \\ &= a \cdot \Phi(\tau(\lambda_1)) + \Phi(\tau(\lambda_2)) = a \cdot (\Psi_{\tau}(\Phi))(\lambda_1) + (\Psi_{\tau}(\Phi))(\lambda_2), \end{aligned}$$

as required.

It remains to show that Ψ_{τ} is complete. Let $\lambda \in \mathbb{L}(\mathscr{S})$. Then we have

$$\sum_{x \in X} ((\Psi_{\tau}(\Phi))(\lambda))(x) = \sum_{x \in X} (\Phi(\tau(\lambda)))(x) = \sum_{C \in \mathscr{C}(\mathscr{S})} (\tau(\lambda))(C)$$
$$= \sum_{C \in \mathscr{C}(\mathscr{S})} \frac{1}{2} \cdot \lambda(C|X-C) = \sum_{S \in \mathscr{S}} \lambda(S),$$

as required.

The following Corollary 6.2 includes, as a special case, the completeness of the fair proportion index on unrooted phylogenetic trees that was established by Wicke and Steel (2019, Thm. 10). To see this, it suffices to consider, for an unrooted phylogenetic tree on X, the split system consisting of those splits of X that can be obtained by removing an edge from the tree.

Corollary 6.2. Let \mathscr{S} be a split system on X and Φ be the fair proportion index on $\mathbb{L}(\mathscr{C}(\mathscr{S}))$. If τ is as defined in (17), then $\Psi_{\tau}(\Phi)$ is a complete linear phylogenetic diversity index on $\mathbb{L}(\mathscr{S})$ and we have

(18)
$$((\Psi_{\tau}(\Phi))(\lambda))(x) = \sum_{A|B \in \mathscr{S}: x \in A} \frac{\lambda(S)}{2 \cdot |A|}$$

for all $\lambda \in \mathbb{L}(\mathscr{S})$ and all $x \in X$.

Proof: In view of Lemma 2.1, Theorem 6.1 implies that $\Psi_{\tau}(\Phi)$ is a complete linear phylogenetic diversity index on $\mathbb{L}(\mathscr{S})$. Moreover, (18) follows from (4), (13), and (17).

(a)								(b)	
()	С	$\boldsymbol{\omega}(C)$	$m_a(C)$	$m_b(C)$	$m_c(C)$	$m_d(C)$	$m_e(C)$	(°) x	ES(x)
	$\{a\}$	2	1	0	0	0	0	a	4
	$\{b\}$	3	0	1	0	0	0	b	5
	$\{c\}$	1	0	0	1	0	0	С	7
	$\{d\}$	4	0	0	0	1	0	d	8
	$\{a,b,c\}$	5	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$	0	0	е	1
	$\{c,d\}$	7	0	0	$\frac{1}{2}$	$\frac{1}{2}$	0		
	X	3	$\frac{1}{9}$	$\frac{1}{9}$	$\frac{5}{18}$	$\frac{1}{6}$	$\frac{1}{3}$		

FIGURE 7. (a) A cluster system \mathscr{C} on $X = \{a, b, c, d, e\}$ with weighting ω and the quantities $m_x(C)$ as defined for all $x \in X$ and $C \in \mathscr{C}$ in (19). (b) The equal splits index ES(x) for all $x \in X$ obtained from \mathscr{C} and ω by (20).

We now turn our attention to a generalization of the *equal splits* index, a phylogenetic diversity index that was introduced in the setting of rooted phylogenetic trees by Redding and Mooers (2006). We first define our generalization for cluster systems \mathscr{C} on X. For all $C \in \mathscr{C}$, let cl(C) denote the set of those $x \in C$ that are not contained in any cluster $C' \in ch(C)$. Then put

(19)
$$m_{x}(C) = \begin{cases} 0 & \text{if } x \notin C, \\ \frac{1}{|ch(C)| + |cl(C)|} & \text{if } x \in cl(C) \\ \sum_{C' \in ch(C)} \frac{m_{x}(C')}{|ch(C)| + |cl(C)|} & \text{otherwise,} \end{cases}$$

for all $x \in X$ and all $C \in \mathscr{C}$. Note that $m_x(C) = 1/|C|$ if $x \in C$ and $ch(C) = \emptyset$ (as in this case |ch(C)| = 0 and |cl(C)| = |C|). The equal splits index is defined by putting

(20)
$$(ES(\boldsymbol{\omega}))(x) = \sum_{C \in \mathscr{C}} m_x(C) \cdot \boldsymbol{\omega}(C)$$

for all $\omega \in \mathbb{L}(\mathscr{C})$ and all $x \in X$. As an example, consider the cluster system \mathscr{C} with weighting ω in Figure 7(a). For the cluster C = X we have $ch(X) = \{\{a, b, c\}, \{c, d\}\}$ and $cl(X) = \{e\}$, which yields, by (19), $m_e(X) = \frac{1}{3}$. The resulting values of the equal splits index are given in Figure 7(b).

The equal splits index on $\mathbb{L}(\mathscr{C})$ is linear with the corresponding $|\mathscr{C}| \times |X|$ -matrix Γ in Equation (5) having the entries $\gamma_{(C,x)} = m_x(C)$. Moreover, as can be seen in the example in Figure 7(a), the sum of the entries in each row of Γ equals 1. The next theorem establishes that this is always the case.

Theorem 6.3. For all cluster systems \mathscr{C} on X the equal splits index is a complete linear phylogenetic diversity index on $\mathbb{L}(\mathscr{C})$.

Proof: Let \mathscr{C} be a cluster system on *X*. We already noted above that the equal splits index is linear. Thus, it remains to establish that the equal splits index is complete. More specifically, it suffices to show that $\sum_{x \in X} m_x(C) = 1$ for all $C \in \mathscr{C}$. We show this by induction on |ch(C)|. For the base case |ch(C)| = 0 we have |cl(C)| = |C| and

$$\sum_{x\in X} m_x(C) = \sum_{x\in C} m_x(C) + \sum_{x\in X\setminus C} m_x(C) = |C| \cdot \frac{1}{|C|} + 0 = 1,$$

as required. Next assume |ch(C)| > 0. Then we have

$$\begin{split} \sum_{x \in X} m_x(C) &= \sum_{x \in X \setminus C} m_x(C) + \sum_{x \in cl(C)} m_x(C) + \sum_{x \in C \setminus cl(C)} m_x(C) \\ &= 0 + \frac{|cl(C)|}{|ch(C)| + |cl(C)|} + \sum_{x \in C \setminus cl(C)} \sum_{C' \in ch(C)} \frac{m_x(C')}{|ch(C)| + |cl(C)|} \\ &= \frac{|cl(C)|}{|ch(C)| + |cl(C)|} + \sum_{C' \in ch(C)} \sum_{x \in C \setminus cl(C)} \frac{m_x(C')}{|ch(C)| + |cl(C)|} \\ &= \frac{|cl(C)|}{|ch(C)| + |cl(C)|} + \sum_{C' \in ch(C)} \sum_{x \in X} \frac{m_x(C')}{|ch(C)| + |cl(C)|} \\ &= \frac{|cl(C)|}{|ch(C)| + |cl(C)|} + |ch(C)| \cdot \frac{1}{|ch(C)| + |cl(C)|} \\ &= 1, \end{split}$$

where the equality in the fourth line holds in view of the fact that $m_x(C') = 0$ for all $x \in X \setminus (C \setminus cl(C))$ and for all $C' \in ch(C)$, and the equality in the fifth line holds by induction.

Our final result in this section, which is an immediate consequence of Theorem 6.1 and Theorem 6.3, summarizes how we obtain, via the map τ defined in (17), a complete linear split-based phylogenetic diversity index from the cluster-based equal splits index.

Corollary 6.4. Let \mathscr{S} be a split system on X and Φ be the equal splits index on $\mathbb{L}(\mathscr{C}(\mathscr{S}))$. If τ is as defined in (17), then $\Psi_{\tau}(\Phi)$ is a complete linear phylogenetic diversity index on $\mathbb{L}(\mathscr{S})$.

As a concrete example of the last corollary, we present a computation of the phylogenetic diversity index $\Psi_{\tau}(\Phi)$. We consider a data set comprising 32 populations of spotted owls (*Strix occidentalis*) that was analyzed by Volkmann et al. (2014). In Table S1 of (Volkmann et al., 2014) pairwise genetic distances between these populations are given. Here, for illustration purposes, we select a subset $X = \{a, f, h, m, r, s\}$ of six of these populations.

We first compute an unrooted phylogenetic network from the pairwise distances between them using the implementation of NeighborNet (Bryant and Moulton, 2004) in SplitsTree (Huson and Bryant, 2005) (the same methodology was applied by Volkmann et al. 2014 to all 32 populations). The resulting phylogenetic network is shown in Figure 8. Note that each band of parallel edges in this network corresponds to a split of the six populations, so that the network represents a weighted split system \mathscr{S} on X that consists of 10 splits of X. In addition, the length of the edges in the band corresponding to a split S gives the weight $\lambda(S)$ of the split. For example, the band of three parallel horizontal edges in the center of the network in Figure 8 corresponds to the split $S = \{a, f, r\} | \{h, m, s\}$ which has weight $\lambda(S) = 0.133$. The total weight of all splits in \mathscr{S} is 1.980 (all weights rounded to three decimal places).

From the split system \mathscr{S} on X with weighting λ we compute the cluster system $\mathscr{C} = \mathscr{C}(\mathscr{S})$ on X with weighting $\omega = \tau(\lambda)$ (in Figure 10 in the Appendix we present the Hasse diagram for the 20 clusters in \mathscr{C} , where the weight $\omega(C)$ is given below each cluster C in the diagram).



FIGURE 8. A phylogenetic network visualizing a weighted split system \mathscr{S} on the set $X = \{a, f, h, m, r, s\}$ of owl populations described in the text. The elements in X represent sampling locations of spotted owls in western North America (a = Aguascalientes, Mexico; f = San Fransisco Peaks, AZ; h = Huachuca Mountains, AZ; m = Marin County, CA; r = Capitol Reef National Park, UT; s = San Bernardino Mountains, CA).

x	$\Psi(x)$	SV(x)
a	0.302	0.276
f	0.185	0.119
h	0.322	0.378
т	0.403	0.464
r	0.348	0.233
S	0.420	0.513

FIGURE 9. The values of the phylogenetic diversity indices $\Psi = \Psi_{\tau}(\Phi)$ and SV for six owl populations as described in the main text.

Finally, we compute the matrix $\Gamma = \Gamma_{\Phi}$ for the equal splits index Φ on $\mathbb{L}(\mathscr{C})$ (see Figure 11 in the Appendix) from which we obtain the values of the phylogenetic diversity index $\Psi = \Psi_{\tau}(\Phi)$ given in Figure 9. For comparison purposes, we also compute the Shapley value index, *SV*, based on the NeighborNet³ in Figure 8 using SplitsTreeCE (Huson and Bryant, 2005).

As can be checked in Figure 9, the sum of the values of the index Ψ yields the total weight 1.980 of all splits in \mathscr{S} , as it should be for a complete phylogenetic diversity index. The ranking of the six populations given by Φ and SV is similar (but not identical): populations s and m are ranked at the top and population f is ranked at the bottom by both indices. In future work, it

³In fact this is the Shapely value of the underlying split system as derived by Volkmann et al. (2014).

could be interesting to further investigate the differences in rankings obtained from these and other split-based diversity indices.

7. CONCLUSION

In this paper, we have presented a framework for phylogenetic diversity indices defined on linear spaces coming from weighted cluster and split systems. Using some examples of popular tree-based phylogenetic diversity indices from the literature, we have shown that this framework can yield generalizations of these indices for cluster and split systems as well as allowing us to gain a better understanding of their interrelationships.

Note that in our framework presented in Figure 5, by associating to any split system \mathscr{S} on X the cluster system $\mathscr{C}(\mathscr{S})$ on X and then considering maps τ , we have focused on deriving splitbased indices from cluster-based indices. In the affine and projective clustering approach, however, there are also ways to associate to any cluster system \mathscr{C} on X a split system $\mathscr{S}(\mathscr{C})$ on X(see, e.g., Dress 2012, Sec. 9.3). Thus, it could be interesting to investigate how this fact might be used to derive cluster-based indices from split-based indices,

In our results, we have considered cluster and split systems in general, special examples of which include hierarchical cluster systems and compatible split systems which correspond to phylogenetic trees. There are, however, various other special classes of cluster and split systems that could be interesting to investigate within our framework. For example, in (Volkmann et al., 2014) two specific phylogenetic diversity indices for so-called *circular split systems* were computed and, in the light of the recent results by Abhari et al. (2023), it would be interesting to look at phylogenetic diversity indices for such split systems more generally. This possibility is illustrated in the example of six owl populations presented at the end of Section 6 since the split systems underlying NeighborNets are always circular (Bryant and Moulton, 2004). It would also be interesting to consider diversity indices coming from *weak hierarchies*, a special type of cluster system introduced by Bandelt and Dress (1989). The advantage of considering such specialized cluster and split systems is that they can be efficiently computed from biological data, making them potentially more useful for applications.

In the literature, various approaches have been proposed to generalize tree-based phylogenetic diversity indices using phylogenetic networks, a graph-theoretical generalization of phylogenetic trees (Coronado et al., 2018; Volkmann et al., 2014; Wicke and Fischer, 2018). Such networks are essentially directed, acyclic, graphs with a single root and whose set of leaves corresponds to some collection of species. The fair proportion index, for example, is generalized in terms of such networks by Coronado et al. (2018). In general, phylogenetic networks give rise to cluster systems (see, e.g., Steel 2016, Sec. 10.3.4) by, for example, taking the set of leaves that lie below a vertex or an edge in the network (just as with rooted phylogenetic trees). Thus, it could be interesting to explore how phylogenetic diversity indices defined in terms of phylogenetic networks, such as, for example, those considered by Wicke and Fischer (2018), fit into our cluster based framework. This could also be interesting to investigate for *split networks* such as the one presented in Figure 8, which are a certain type of undirected phylogenetic network (see, e.g., Dress 2012, Sec. 4.4).

Finally, in another direction, it could be interesting to apply our framework to establish properties and generalizations of other tree-based phylogenetic diversity indices that we did not consider in this paper. Indeed, as we have demonstrated, sometimes expressing indices in terms of clusters or splits can lead to more concise proofs for showing that they have certain properties. For example, it would be interesting to extend the Pauplin index considered by Wicke and Steel (2019) for unrooted phylogenetic trees to more general split systems and study its properties, as well as to consider some of the other related questions asked in (Wicke and Steel, 2019, Sec. 6) within our new framework.

Data availability. No data was generated.

Acknowledgment. All authors thank Schloss Dagstuhl – Leibniz Centre for Informatics – for hosting the Seminar 19443 *Algorithms and Complexity in Phylogenetics* in October 2019, where this work was initiated.

REFERENCES

- Abhari, N., Colijn, C., Mooers, A., and Tupper, P. (2023). Capturing diversity: split systems and circular approximations for conservation. *Journal of Theoretical Biology*.
- Aumann, R. J. (1994). The Shapley value. In *Game-Theoretic Methods in General Equilibrium Analysis*, pages 61–66. Springer.
- Bandelt, H.-J. and Dress, A. W. (1989). Weak hierarchies associated with similarity measures—an additive clustering technique. *Bulletin of Mathematical Biology*, 51(1):133–166.
- Bordewich, M. and Semple, C. (2023). Quantifying the difference between phylogenetic diversity and diversity indices. arXiv:2304.10725.
- Branzei, R., Dimitrov, D., and Tijs, S. (2008). *Models in cooperative game theory*, volume 556. Springer Science & Business Media.
- Bryant, D. and Moulton, V. (2004). Neighbor-net: an agglomerative method for the construction of phylogenetic networks. *Molecular Biology and Evolution*, 21(2):255–265.
- Coronado, T. M., Riera, G., and Rosselló, F. (2018). The fair proportion is a Shapley value on phylogenetic networks too. In *Enjoying Natural Computing*, pages 77–87. Springer.
- Dress, A. (1997). Towards a theory of holistic clustering. In *Mathematical Hierarchies and Biology*, pages 271–289. American Mathematical Society.
- Dress, A. (2012). Basic phylogenetic combinatorics. Cambridge University Press.
- Dubey, P. (1975). On the uniqueness of the Shapley value. *International Journal of Game Theory*, 4:131–139.
- Faith, D. P. (1992). Conservation evaluation and phylogenetic diversity. *Biological Conservation*, 61(1):1–10.
- Fuchs, M. and Jin, E. Y. (2015). Equality of Shapley value and fair proportion index in phylogenetic trees. *Journal of Mathematical Biology*, 71(5):1133–1147.
- Gumbs, R., Gray, C. L., Böhm, M., Burfield, I. J., Couchman, O. R., Faith, D. P., Forest, F., Hoffmann, M., Isaac, N. J. B., Jetz, W., Mace, G. M., Mooers, A. O., Safi, K., Scott, O., Steel, M., Tucker, C. M., Pearse, W. D., Owen, N. R., and Rosindell, J. (2023). The EDGE2 protocol:

Advancing the prioritisation of evolutionarily distinct and globally endangered species for practical conservation action. *PLOS Biology*, 21(2):e3001991.

- Haake, C.-J., Kashiwada, A., and Su, F. E. (2008). The Shapley value of phylogenetic trees. *Journal of Mathematical Biology*, 56(4):479–497.
- Huson, D. H. and Bryant, D. (2005). Application of phylogenetic networks in evolutionary studies. *Molecular Biology and Evolution*, 23(2):254–267.
- Isaac, N. J., Turvey, S. T., Collen, B., Waterman, C., and Baillie, J. E. (2007). Mammals on the EDGE: Conservation priorities based on threat and phylogeny. *PLoS ONE*, 2(3):e296.
- Jukna, S. (2011). Extremal combinatorics: with applications in computer science. Springer.
- Kleinman, A., Harel, M., and Pachter, L. (2013). Affine and projective tree metric theorems. *Annals of Combinatorics*, 17:205–228.
- Manson, K. (2022). Many phylogenetic diversity indices are not robust to extinctions. bioRxiv:2022.06.28.498028.
- Manson, K. and Steel, M. (2023). Spaces of phylogenetic diversity indices: combinatorial and geometric properties. *Bulletin of Mathematical Biology*, 85(78).
- Redding, D. W. (2003). Incorporating genetic distinctness and reserve occupancy into a conservation priorisation approach. Master's thesis, University Of East Anglia, Norwich, UK.
- Redding, D. W., Hartmann, K., Mimoto, A., Bokal, D., DeVos, M., and Mooers, A. (2008). Evolutionarily distinctive species often capture more phylogenetic diversity than expected. *Journal of Theoretical Biology*, 251(4):606–615.
- Redding, D. W., Mazel, F., and Mooers, A. Ø. (2014). Measuring evolutionary isolation for conservation. *PLoS ONE*, 9(12):e113490.
- Redding, D. W. and Mooers, A. Ø. (2006). Incorporating evolutionary measures into conservation prioritization. *Conservation Biology*, 20(6):1670–1678.
- Semple, C. and Steel, M. (2003). Phylogenetics. Oxford University Press.
- Shapley, L. S. (1953). A value for *n*-person games. *Contributions to the Theory of Games*, 2(28):307–317.
- Spillner, A., Nguyen, B. T., and Moulton, V. (2008). Computing phylogenetic diversity for split systems. *IEEE/ACM Transactions on Computational Biology and Bioinformatics*, 5(2):235–244.
- Steel, M. (2016). *Phylogeny: Discrete and random processes in evolution*. SIAM, Philadelphia PA.
- Tucker, C. M., Cadotte, M. W., Carvalho, S. B., Davies, T. J., Ferrier, S., Fritz, S. A., Grenyer, R., Helmus, M. R., Jin, L. S., Mooers, A. O., Pavoine, S., Purschke, O., Redding, D. W., Rosauer, D. F., Winter, M., and Mazel, F. (2016). A guide to phylogenetic metrics for conservation, community ecology and macroecology. *Biological Reviews*, 92(2):698–715.
- Vane-Wright, R., Humphries, C., and Williams, P. (1991). What to protect?—Systematics and the agony of choice. *Biological Conservation*, 55(3):235–254.
- Volkmann, L., Martyn, I., Moulton, V., Spillner, A., and Mooers, A. O. (2014). Prioritizing populations for conservation using phylogenetic networks. *PLoS One*, 9(2).
- Wicke, K. and Fischer, M. (2018). Phylogenetic diversity and biodiversity indices on phylogenetic networks. *Mathematical Biosciences*, 298:80–90.

Wicke, K. and Steel, M. (2019). Combinatorial properties of phylogenetic diversity indices. *Journal of Mathematical Biology*, pages 1–29.

APPENDIX



FIGURE 10. The Hasse diagram for the clusters in the cluster system $\mathscr{C}(\mathscr{S})$ on $X = \{a, f, h, m, r, s\}$ computed from the split system \mathscr{S} on X represented by the phylogenetic network in Figure 8. The number below each cluster is the weight of the cluster obtained by the map τ from the weights of the splits in \mathscr{S} .

	a	f	h	m	r	s
$\{a\}$	1	0	0	0	0	0
$\{h\}$	0	0	1	0	0	0
$\{m\}$	0	0	0	1	0	0
$\{r\}$	0	0	0	0	1	0
$\{s\}$	0	0	0	0	0	1
$\{a, f\}$	$\frac{1}{2}$	$\frac{1}{2}$	0	0	0	0
$\{a,m\}$	$\frac{\overline{1}}{2}$	Ō	0	$\frac{1}{2}$	0	0
$\{m,s\}$	0	0	0	$\frac{1}{2}$	0	$\frac{1}{2}$
$\{a, f, r\}$	$\frac{1}{4}$	$\frac{1}{4}$	0	0	$\frac{1}{2}$	0
$\{a, m, s\}$	$\frac{1}{4}$	0	0	$\frac{1}{2}$	0	$\frac{1}{4}$
$\{f, h, r\}$	0	$\frac{1}{3}$	$\frac{1}{3}$	0	$\frac{1}{3}$	0
$\{h,m,s\}$	0	0	$\frac{1}{2}$	$\frac{1}{4}$	0	$\frac{1}{4}$
$\{a, f, h, r\}$	$\frac{1}{8}$	$\frac{7}{24}$	$\frac{1}{6}$	0	$\frac{5}{12}$	0
$\{f,h,r,s\}$	0	$\frac{1}{6}$	$\frac{1}{6}$	0	$\frac{1}{6}$	$\frac{1}{2}$
$\{h,m,r,s\}$	0	0	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{2}$	$\frac{1}{8}$
$\{a,f,h,m,r\}$	$\frac{5}{16}$	$\frac{7}{48}$	$\frac{1}{12}$	$\frac{1}{4}$	$\frac{5}{24}$	0
$\{a,f,h,m,s\}$	$\frac{1}{4}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{4}$	0	$\frac{1}{6}$
$\{a,f,h,r,s\}$	$\frac{1}{16}$	$\frac{11}{48}$	$\frac{1}{6}$	0	$\frac{7}{24}$	$\frac{1}{4}$
$\{a,f,m,r,s\}$	$\frac{1}{4}$	$\frac{1}{8}$	Õ	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{8}$
$\{f, h, m, r, s\}$	0	$\frac{1}{12}$	$\frac{5}{24}$	$\frac{1}{16}$	$\frac{1}{3}$	$\frac{5}{16}$

FIGURE 11. The matrix Γ for the equal splits index on $\mathbb{L}(\mathscr{C})$ for the cluster system \mathscr{C} in Figure 10.