Near-Optimal differentially private low-rank trace regression with guaranteed private initialization

Mengyue, Zha *

(March 26, 2024)

Abstract

We study differentially private (DP) estimation of a rank-r matrix $M \in \mathbb{R}^{d_1 \times d_2}$ under the trace regression model with Gaussian measurement matrices. Theoretically, the sensitivity of non-private spectral initialization is precisely characterized, and the differentialprivacy-constrained minimax lower bound for estimating M under the Schatten-q norm is established. Methodologically, the paper introduces a computationally efficient algorithm for DP-initialization with a sample size of $n \geq \tilde{O}(r^2(d_1 \vee d_2))$. Under certain regularity conditions, the DP-initialization falls within a local ball surrounding M. We also propose a differentially private algorithm for estimating M based on Riemannian optimization (DP-RGrad), which achieves a near-optimal convergence rate with the DP-initialization and sample size of $n \geq \tilde{O}(r(d_1 + d_2))$. Finally, the paper discusses the non-trivial gap between the minimax lower bound and the upper bound of low-rank matrix estimation under the trace regression model. It is shown that the estimator given by DP-RGrad attains the optimal convergence rate in a weaker notion of differential privacy. Our powerful technique for analyzing the sensitivity of initialization requires no eigengap condition between r non-zero singular values.

1 Introduction

The trace regression model (Rohde and Tsybakov, 2011; Koltchinskii et al., 2011), as an extension of the standard regression model, has been widely applied in various fields such as ma-

^{*}Department of Mathematics, Hong Kong University of Science and Technology, mzha@connect.ust.hk. Mengyue, Zha's research was supported by Hong Kong PhD Fellowship Scheme.

trix completion, compressed sensing, and multi-task learning (Negahban and Wainwright, 2011; Koltchinskii et al., 2011; Hamidi and Bayati, 2022). Previous studies have proposed both convex and non-convex approaches for optimal estimation procedures for the model. However, the increasing demand for privacy protection has added new complexities to this extensively studied problem. Differential privacy (DP) (Dwork et al., 2006), a framework for protecting individual privacy, has been widely adopted in industrial and governmental applications (Erlingsson et al., 2014; Apple Differential Privacy Team, 2017; Abowd et al., 2020). This paper aims to develop a near-optimal differentially private method for low-rank matrix estimation under the trace regression model.

Trace regression model Let $M \in \mathbb{R}^{d_1 \times d_2}$ be an unknown rank-*r* matrix and $X_i \in \mathbb{R}^{d_1 \times d_2}$ be the measurement matrix for $i = 1, \dots, n$. Suppose the noisy observation y_i satisfies

$$y_i = \langle X_i, M \rangle + \xi_i \quad \text{for} \quad i = 1, \cdots, n,$$
 (1)

where the model noise $\xi_i \stackrel{i.i.d.}{\sim} \mathcal{N}(0, \sigma_{\xi}^2)$ and the inner product between X_i and M in Euclidean space is given by $\langle X_i, M \rangle := \operatorname{Tr}(X_i^{\top}M)$. The goal of the present paper is to estimate the unknown rank-r matrix $M \in \mathbb{R}^{d_1 \times d_2}$ under trace regression model defined by (1), subject to differential privacy, based on n independent observations $Z := \{(X_i, y_i)\}_{i=1}^n$.

Our approaches are built upon the Gaussian mechanism Dwork et al. (2006). The main difficulty in applying the Gaussian mechanism is sharply characterizing sensitivities of statistics whose privacy is under protection. Listed below are definitions of sensitivity, differential privacy (DP), and Gaussian mechanism. Interested readers may refer to Dwork et al. (2006, 2014); Vadhan (2017) for proofs and other details. Let $\|\cdot\|$ denotes the spectral norm and $\|\cdot\|_F$ denotes the Frobenius norm.

Sensitivity The sensitivity of a function f that maps a dataset Z into $\mathbb{R}^{d_1 \times d_2}$ is defined by $\Delta_f := \sup_{\text{neighbouring}(Z,Z')} \|f(Z) - f(Z')\|_{\text{F}}$, where and the supremum is taken over all neighbouring datasets Z and Z' that differ by at most one observation.

Differential privacy Let $\varepsilon > 0$ and $0 < \delta < 1$, then we say the randomized algorithm A is (ε, δ) -differentially private if $\mathbb{P}(A(Z) \in \mathcal{Q}) \leq e^{\varepsilon} \mathbb{P}(A(Z') \in \mathcal{Q}) + \delta$ for all neighbouring data sets Z, Z' and all subset $\mathcal{Q} \subset \mathbb{R}^{d_1 \times d_2}$.

Gaussian mechanism The randomized algorithm defined by A(Z) = f(Z) + E is (ε, δ) -DP where $E \in \mathbb{R}^{d_1 \times d_2}$ has i.i.d. $N(0, 2\Delta_f^2 \varepsilon^{-2} \log(1.25/\delta))$ entries.

RIP of Gaussian measurement matrices The sensitivity of any statistic involving $\{X_i\}_{i \in [n]}$ depends on the properties of the measurement matrices. Besides, it has been previously established since Candes and Tao (2005) that the restricted isometry property (RIP) on measurement matrices is crucial to the recovery of the unknown matrix M. Hence, assumptions on $\{X_i\}_{i \in [n]}$ are necessary and the present paper considers $\{X_i\}_{i \in [n]}$ with Gaussian design.

Assumption 1 (Gaussian design). The vectorization of measurement matrices X_1, \ldots, X_n are independent Gaussian $\operatorname{vec}(X_i) \sim \mathcal{N}(\mathbf{0}, \Lambda_i)$ where Λ_i 's are known, symmetric and positive definite. There exist absolute constants $C_l, C_u > 0$ such that $C_l \leq \lambda_{\min}(\Lambda_i) \leq \lambda_{\max}(\Lambda_i) \leq C_u$.

The following Lemma 1 shows that under Assumption 1, the measurement matrices $\{X_i\}_{i=1}^n$ satisfy the restricted isometry property (RIP) with high probability, see the proof in E.1.

Lemma 1. Under the Assumption 1, for any $B \in \mathbb{R}^{d_1 \times d_2}$ of rank r, there exist constants $c_1, c_2, c_3 > 0$ and $c_5 > c_4 > 0$ such that if $n \ge c_1 r(d_1 + d_2)$, with probability at least $1 - c_2 \exp(-c_3 r(d_1 + d_2))$, we have $c_4 \sqrt{C_u C_l} \|B\|_F^2 \le \frac{1}{n} \sum_{i=1}^n \langle X_i, B \rangle^2 \le c_5 \sqrt{C_u C_l} \|B\|_F^2$.

Notations Suppose M is of rank-r and its singular value decomposition is of the form $M = U\Sigma V^{\top} \in \mathbb{R}^{d_1 \times d_2}$ where $U \in \mathbb{O}_{d_1,r}$, $V \in \mathbb{O}_{d_2,r}$ and $\Sigma = \text{diag}\{\sigma_1 \cdots \sigma_r\}$ with $\sigma_1 \geq \cdots \geq \sigma_r$. Here, $\mathbb{O}_{d,r}$ denotes the set of $d \times r$ matrices satisfying $H^{\top}H = I_r$. Let $\kappa := \sigma_1/\sigma_r$ be the condition number and $\kappa_{\xi} := \sigma_{\xi}/\sigma_r$ be the signal-to-noise ratio. Let \widetilde{O} stand for the typical big-O notation up to logarithmic factors and $\widetilde{O}_p(\cdot)$ stand for \widetilde{O} holds with high probability.

1.1 Main results

The paper presents several key results related to differentially private low-rank matrix estimation. Firstly, we propose a private initialization \widetilde{M}_0 (as detailed in Algorithm 1). Secondly, we establish the privacy-constrained minimax lower bound under the general Shatten-q norm (as detailed in Theorem 2). Finally, we introduce a private estimator \widetilde{M}_{l^*} (as detailed in Algorithm 2) that achieves the near-optimal convergence rate under the Frobenius norm. The sensitivity analysis of \widetilde{M}_0 heavily relies on a spectral representation formula for asymmetric matrices (See Lemma 2). We prove in Corollary 1 that the private initialization \widetilde{M}_0 satisfies $\|\widetilde{M}_0 - M\|_F \leq \sqrt{2r} \|\widetilde{M}_0 - M\| \leq c_0 \sigma_r$, with high probability (w.h.p.), for a small constant $0 < c_0 < 1$, provided that $n \geq \widetilde{O}\left((\kappa^4 r^2 + \kappa^2 \kappa_{\xi}^2 r)(d_1 \vee d_2)\right)$.

Theorem 2 establishes the DP-constrained minimax risk of estimating the rank-r matrix $M \in \mathbb{R}^{d_1 \times d_2}$ under model (1) and general Schatten-q norm. Specifically, the minimax risk under Frobenius norm is in the order of $\sigma_{\xi} \sqrt{\frac{r(d_1 \vee d_2)}{n}} + \sigma_{\xi} \frac{r(d_1 \vee d_2)}{n\varepsilon}$.

Finally, we show in Theorem 3 that with a sample size of $n \geq \widetilde{O}\left(\left(\kappa_{\xi}^2 \vee \kappa_{\xi}\right) r(d_1 \vee d_2)\right)$ and any initialization satisfying (2), Algorithm 2 achieves geometric convergence rate. The private estimator \widetilde{M}_{l^*} attains the near-optimal convergence rate

$$\left\|\widetilde{M}_{l^*} - M\right\|_F \le \widetilde{O}_p\left(\sigma_{\xi}\sqrt{\frac{r(d_1+d_2)}{n}} + (\sigma_{\xi}+\sigma_r)\frac{r(d_1+d_2)}{n\varepsilon}\right)$$

1.2 Motivations and related works

The trace regression model has been extensively researched, resulting in well-known optimal procedures and theoretical properties. Both convex (Rohde and Tsybakov, 2011; Koltchinskii et al., 2011; Candes and Plan, 2011; Negahban and Wainwright, 2011) and non-convex methods (Burer and Montei 2003; Chen and Wainwright, 2015; Zheng and Lafferty, 2016; Wei et al., 2016) have achieved the optimal convergence rate of the order $\sigma_{\xi} \sqrt{\frac{r(d_1 \vee d_2)}{n}}$ without the constraint from differential privacy. However, the DP-constrained minimax rate of low-rank matrix estimation under the trace regression model is still unknown. (Near) Optimal DP-algorithms have been developed for statistical problems such as learning Gaussians Kamath et al. (2019); Kuditipudi et al. (2023); Brown et al. (2023) or heavy-tailed distributions Kamath et al. (2020), (sparse or generalized) linear regression Wang (2018); Cai et al. (2021, 2023), and PCA Blum et al. (2005); Dwork et al. (2014); Chaudhuri et al. (2012); Liu et al. (2022). Previous works on DP-regression Cai et al. (2021, 2023) assume that all measurements have bounded ℓ_2 norm. This assumption presents a significant limitation to studying the role of measurements play in the estimation error. Additionally, by treating measurements as a fixed vector or matrix, the statistical properties of measurements are disregarded. As a result, the opportunity for optimal statistical analysis subject to privacy concerns is inevitably lost. Recently, (McSherry and Mironov, 2009; Liu et al., 2015; Jain et al., 2018; Chien et al., 2021; Wang et al., 2023) propose gradient-descent-based algorithms for DP low-rank matrix completion. These algorithms have attained near-optimal sample complexity. However, the problem of sample-efficient, differentially private initialization remains under-explored. Additionally, it is unknown how to establish the minimax risk of lowrank matrix estimation with the constraints of differential privacy, especially when the matrix is asymmetric.

1.3 Organization

Section 2 proposes a DP-initialization algorithm and presents its privacy and utility guarantees. In Section 3, we establish a DP-constrained minimax lower bound (5) for estimating the rank-r matrix M under the trace regression model. Section 4 presents the DP-estimator based on nonconvex optimization and derives the upper bound of the DP-estimator's error, as stated in (7). We discuss the score attack argument and the non-trivial gap between the upper bound of (7) and the DP-constrained minimax lower bound (5) in Section 5. Proofs are given in Appendix A to F.

2 DP-initialization

Section 2.1 presents an (ε, δ) -DP initialization \widetilde{M}_0 , as stated in Algorithm 1. In Section 2.2, we introduce a spectral representation formula (See Lemma 2) that is crucial to sensitivity analysis on the initialization. With the help of the spectral representation formula, the privacy and utility guarantees of the DP-initialization \widetilde{M}_0 are given in Section 2.3.

2.1 Algorithm for DP-initialization

We begin with $\widehat{L} = n^{-1} \sum_{i=1}^{n} \max \left(\Lambda_i^{-1} \operatorname{vec} (X_i) \right) y_i$, which is unbiased for M. Suppose that the leading-r left and right singular vectors of \widehat{L} is given by the columns of $\widehat{U} \in \mathbb{O}_{d_1,r}$ and $\widehat{V} \in \mathbb{O}_{d_2,r}$, respectively. Then, $\widehat{M}_0 := \operatorname{SVD}_r(\widehat{L})$ is a non-private estimator for M. Let $\widehat{\Sigma} := \widehat{U}^{\top} \widehat{L} \widehat{V}$, then we have

$$\widehat{M}_0 = \mathrm{SVD}_r(\widehat{L}) = \widehat{U}\widehat{U}^\top \widehat{L}\widehat{V}\widehat{V}^\top = \widehat{U}\widehat{\Sigma}\widehat{V}^\top$$

It is reasonable to think about privatizing \widehat{U} , \widehat{V} , and $\widehat{\Sigma}$, separately. We first privatize the empirical spectral projector $\widehat{U}\widehat{U}^{\top}$ and $\widehat{V}\widehat{V}^{\top}$ by Gaussian mechanism. Thanks to postprocessing property Dwork et al. (2006), we obtain $\widetilde{U} \in \mathbb{O}^{d_1,r}$ and $\widetilde{V} \in \mathbb{O}^{d_2,r}$ whose columns are differentially private and orthogonal. Secondly, we privatize the $r \times r$ matrix $\widetilde{U}^{\top}\widehat{L}\widetilde{V}$ by Gaussian mechanism and obtain $\widetilde{\Sigma} \in \mathbb{R}^{r \times r}$ which is a private surrogate for $\widehat{\Sigma} = \widehat{U}^{\top}\widehat{L}\widehat{V}$. Finally, we take $\widetilde{M}_0 = \widetilde{U}\widetilde{\Sigma}\widetilde{V}^{\top}$ as the DP-initialization. We display the pseudo-code of the proposed DP-initialization in Algorithm 1. The privacy of \widetilde{M}_0 is guaranteed by the composition property Dwork et al. (2006).

Algorithm 1 Differentially private initialization for trace regression

Input: the data set $\{(X_i, y_i)\}_{i=1}^n$; the covariance matrices $\{\Lambda_i\}_{i=1}^n$; sensitivity $\Delta^{(1)}$, $\Delta^{(2)} > 0$; rank r; nuisance variance σ_{ξ}^2 ; privacy budget $\varepsilon > 0$ and $0 < \delta < 1$.

Output: (ε, δ) -differentially private initialization \widetilde{M}_0 .

Compute the unbiased sample estimator \hat{L} and its top-*r* left and right singular vectors:

$$\widehat{L} \leftarrow n^{-1} \sum_{i=1}^{n} \max\left(\Lambda_i^{-1} \operatorname{vec}\left(X_i\right)\right) y_i \quad \text{and} \quad \widehat{M}_0 = \widehat{U}\widehat{\Sigma}\widehat{V} \leftarrow \operatorname{SVD}_r(\widehat{L})$$

Compute $(\varepsilon/3, \delta/3)$ -differentially private singular subspaces by adding artificial Gaussian noise:

$$\widetilde{U} \leftarrow \text{SVD}_r \left(\widehat{U} \widehat{U}^\top + E_U \right) \text{ with } (E_U)_{ij} = (E_U)_{ji} \overset{\text{i.i.d.}}{\sim} N \left(0, \frac{18\Delta^{(1)^2}}{\varepsilon^2} \log(3/\delta) \right), \forall 1 \le i \le j \le d_1$$
$$\widetilde{V} \leftarrow \text{SVD}_r \left(\widehat{V} \widehat{V}^\top + E_V \right) \text{ with } (E_V)_{ij} = (E_U)_{ji} \overset{\text{i.i.d.}}{\sim} N \left(0, \frac{18\Delta^{(1)^2}}{\varepsilon^2} \log(3/\delta) \right), \forall 1 \le i \le j \le d_2$$

Compute $(\varepsilon/3, \delta/3)$ -differentially private estimates of singular values up to rotations:

$$\widetilde{\Sigma} \leftarrow \widetilde{U}^{\top} \widehat{L} \widetilde{V} + E_{\Sigma} \operatorname{with} (E_{\Sigma})_{ij} = (E_{\Sigma})_{ji} \overset{i.i.d.}{\sim} N\left(0, \frac{18\Delta^{(2)^2}}{\varepsilon^2} \log(3/\delta)\right), \forall 1 \le i \le j \le n$$

Compute (ε, δ) -differentially private initialization: $\widetilde{M}_0 \leftarrow \widetilde{U}\widetilde{\Sigma}\widetilde{V}^\top$ Return: \widetilde{M}_0

To this end, we define the sensitivities of $\Delta^{(1)}$ and $\Delta^{(2)}$ appear in Algorithm 1. Let

$$\widehat{L}^{(i)} := n^{-1} \sum_{j \neq i}^{n} \max\left(\Lambda_{j}^{-1} \operatorname{vec}\left(X_{j}\right)\right) Y_{j} + n^{-1} \operatorname{mat}\left(\Lambda_{i}^{-1} \operatorname{vec}\left(X_{i}'\right)\right) y_{i}'$$

where (X'_i, y'_i) is an i.i.d. copy of (X_i, y_i) . Then, the estimator $\widehat{L}^{(i)}$ differs with \widehat{L} only by the *i*-th pair of observations. Suppose the top-*r* left and right singular vectors of $\widehat{L}^{(i)}$ are given by $U^{(i)}$ and $V^{(i)\top}$, respectively. The sensitivity of $\widehat{U}\widehat{U}^{\top}$ is defined by

$$\Delta_{\widehat{U}\widehat{U}^{\top}} = \sup_{\text{neighbouring}(Z,Z')} \left\| \widehat{U}(Z)\widehat{U}(Z)^{\top} - \widehat{U}(Z')\widehat{U}(Z')^{\top} \right\|_{F} = \max_{i \in [n]} \left\| \widehat{U}\widehat{U}^{\top} - \widehat{U}^{(i)}\widehat{U}^{(i)\top} \right\|_{F},$$

and the sensitivity $\Delta_{\widehat{V}\widehat{V}^{\top}}$ of $\widehat{V}\widehat{V}^{\top}$ is defined similarly. We refer to $\Delta^{(1)} \triangleq \Delta_{\widehat{U}\widehat{U}^{\top}} \vee \Delta_{\widehat{V}\widehat{V}^{\top}}$ as the sensitivity of singular subspaces and define the sensitivity

$$\Delta^{(2)} \triangleq \Delta_{\widetilde{U}^{\top}\widehat{L}\widetilde{V}} = \sup_{\text{neighbouring}(Z,Z')} \left\| \widetilde{U}\widehat{L}(Z)\widetilde{V}^{\top} - \widetilde{U}\widehat{L}(Z')\widetilde{V}^{\top} \right\|_{F} = \max_{i \in [n]} \left\| \widetilde{U}^{\top} \left(\widehat{L} - \widehat{L}^{(i)}\right)\widetilde{V} \right\|_{F}.$$

As privatizing $\widehat{\Sigma} = \widehat{U}^{\top} \widehat{L} \widehat{V}$ by Gaussian mechanism, the scale of artificial noise avoids growing with an unnecessary $\sqrt{d_1} \vee \sqrt{d_2}$ but rather growing with a smaller quantity \sqrt{r} . This benefit motivates us to privatize \widehat{U} , \widehat{V} and $\widehat{\Sigma}$, separately. However, it is technically challenging to characterize $\Delta^{(1)} = \Delta_{\widehat{U}\widehat{U}^{\top}} \vee \Delta_{\widehat{V}\widehat{V}^{\top}}$ due to the non-linear dependence of $\widehat{U}\widehat{U}^{\top}$ and $\widehat{V}\widehat{V}^{\top}$ on the dataset $Z = \{(X_i, y_i)\}_{i=1}^n$. To address this challenge, we introduce an explicit spectral representation formula (See Lemma 2) to obtain a sharp upper bound on the sensitivity of the singular subspaces.

2.2 Spetral representation formula

This section introduces a spectral representation formula for asymmetric matrices (See Lemma 2). To begin with, we quickly explain the standard *symmetric dilation* trick (See e.g., Section 2.1.17 in Tropp et al. (2015)) and define auxiliary operators used in Lemma 2.

Symmetric dilation and auxiliary operators For any $M \in \mathbb{R}^{d_1 \times d_2}$, the symmetric dialation M_* of M is a $(d_1 + d_2) \times (d_1 + d_2)$ matrix defined by $M_* := \begin{pmatrix} 0 & M \\ M^\top & 0 \end{pmatrix}$. It is easy to check that $M_* = M_*^\top$ and $||M_*|| = ||M||$. Further, if we assume that M is of rank r and has the form of SVD $M = U\Sigma V^\top \in \mathbb{R}^{d_1 \times d_2}$, then M_* is of rank-2r and has eigendecomposition of the form

$$\frac{1}{\sqrt{2}} \begin{pmatrix} U & U \\ V & -V \end{pmatrix} \cdot \begin{pmatrix} \Sigma & 0 \\ 0 & -\Sigma \end{pmatrix} \cdot \frac{1}{\sqrt{2}} \begin{pmatrix} U & U \\ V & -V \end{pmatrix}^{\top} := U_{M^*} \Sigma_{M^*} U_{M^*}^{\top}.$$

The 2r eigenvectors of M_* is given by the columns of $U_{M^*} \in \mathbb{O}_{(d_1+d_2),2r}$. For integer $t \ge 1$, we define operators

$$Q^{-t} := U_{M^*} \Sigma_{M^*}^{-t} U_{M^*}^{\top}$$
 and $Q^{-0} := Q^{\perp} \triangleq (U_{M^*})_{\perp} (U_{M^*})_{\perp}^{\top} = I_{d_1+d_2} - U_{M^*} U_{M^*}^{\top}$.

Lemma 2 (Spectral representation formula). Let $M \in \mathbb{R}^{d_1 \times d_2}$ be any rank-r matrix with singular values $\sigma_1 \geq \cdots \geq \sigma_r > 0$ and $\widehat{L} = M + \Delta \in \mathbb{R}^{d_1 \times d_2}$ be a perturbation of M where $\Delta \in \mathbb{R}^{d_1 \times d_2}$ is the deviation matrix. Suppose the top-r left and right singular vectors of \widehat{L} and M, are given by the columns of \widehat{U} , \widehat{V} and U, V, respectively. Suppose that $2\|\Delta\| \leq \sigma_r$, then

$$\begin{pmatrix} \widehat{U}\widehat{U}^{\top} - UU^{\top} & 0\\ 0 & \widehat{V}\widehat{V}^{\top} - VV^{\top} \end{pmatrix} = \sum_{k \ge 1} \mathcal{S}_{M_*,k}(\Delta_*).$$

Here, Δ_* is the symmetric dilation of $\Delta := \widehat{L} - M$ and the k-th order term $\mathcal{S}_{M_*,k}(\Delta_*)$ is a summation of $\binom{2k}{k}$ terms defined by $\mathcal{S}_{M_*,k}(\Delta_*) = \sum_{\mathbf{s}:s_1+\ldots+s_{k+1}=k} (-1)^{1+\tau(\mathbf{s})} \cdot Q^{-s_1} \Delta_* Q^{-s_2} \ldots \Delta_* Q^{-s_{k+1}}$, where $\mathbf{s} = (s_1, \ldots, s_{k+1})$ contains non-negative indices and $\tau(\mathbf{s}) = \sum_{j=1}^{k+1} \mathbb{I}(s_j > 0)$.

In Lemma 2, the spectral projectors $\widehat{U}\widehat{U}^{\top}$ and $\widehat{V}\widehat{V}^{\top}$ of the matrix $\widehat{L} = M + \Delta$, is explicitly represented in terms of the symmetric dilation of Δ , with the help of auxiliary operators Q^{-0} and Q^{-t} for integer $t \geq 1$. The proof of Lemma 2 is deferred to Appendix E.2. Note that Lemma 2 accommodates a diverging condition number and requires no eigengap condition between rnon-zero singular values. In the proof of Theorem 1, we shall see that $\widehat{V}\widehat{V}^{\top} - \widehat{V}^{(i)}\widehat{V}^{(i)\top}$ and $\widehat{U}\widehat{U}^{\top} - \widehat{U}^{(i)}\widehat{U}^{(i)\top}$ are mainly contributed by the 1-st order approximation $\mathcal{S}_{M_{*},1}(\Delta_{*}) - \mathcal{S}_{M_{*},1}(\Delta_{*}^{(i)})$ where $\Delta_{*}^{(i)}$ is the symmetric dilation of $\Delta^{(i)} := \widehat{L}^{(i)} - M$.

2.3 Privacy and utility guarantees of the initialization

In this section, we study the privacy and utility guarantees of the initialization \widetilde{M}_0 . Theorem 1 characterizes the sensitivities $\Delta^{(1)}$ and $\Delta^{(2)}$ needed to guarantee an (ε, δ) -DP \widetilde{M}_0 , and present the upper bounds of $\|\widetilde{M}_0 - M\|$ and $\|\widetilde{M}_0 - M\|_F$. The proof of Theorem 1 is provided in Appendix A.

Theorem 1 (Privacy and utility guarantees of the initialization \widetilde{M}_0). Consider i.i.d. observations $Z = \{z_1, \dots, z_n\}$ drawn from the trace regression model stated in (1) where $z_i := (X_i, y_i)$ for $i = 1, \dots, n$. Let the true rank-r regression coefficients matrix be $M \in \mathbb{R}^{d_1 \times d_2}$. Suppose that $\{X_i\}_{i \in [n]}$ satisfy the Assumption 1. Under the mild condition $n \geq \frac{\log^2 n}{(d_1 \vee d_2) \log(d_1 + d_2)}$, there exists absolute constants $C_1, C_2, C_3 > 0$ such that

$$n \ge n_0 \triangleq C_1 C_l^{-1} r \left(\frac{\sigma_{\xi} + \sqrt{C_u r} \sigma_1}{\sigma_r} \right)^2 (d_1 \lor d_2) \log(d_1 + d_2);$$

if Algorithm 1 takes in sensitivities at least $\Delta^{(1)} = C_2 \left(\frac{\sqrt{C_l^{-1}} \left(\sqrt{C_u r} \sigma_1 + \sigma_{\xi} \right)}{\sigma_r} \right) \frac{\sqrt{r}}{n} \log n$ and $\Delta^{(2)} = C_2 \sqrt{C_l^{-1}} \left(\sqrt{C_u r} \sigma_1 + \sigma_{\xi} \right) \frac{\sqrt{r}}{n} \log n$, then Algorithm 1 is guaranteed to be (ε, δ) -DP. Moreover, the

output \widetilde{M}_0 of Algorithm 1 satisfies

r

$$\begin{split} \|\widetilde{M}_{0} - M\| \bigvee \left(\|\widetilde{M}_{0} - M\|_{F} / \sqrt{2r} \right) \\ \leq \underbrace{C_{3} \sqrt{C_{l}^{-1}} \left(\sqrt{C_{u} r} \sigma_{1} + \sigma_{\xi} \right) \frac{\sigma_{1}}{\sigma_{r}} \sqrt{\frac{(d_{1} \vee d_{2}) \log(d_{1} + d_{2})}{n}}_{e_{1}}}_{e_{1}} \\ + \underbrace{C_{3} \sqrt{C_{l}^{-1}} \left(\sqrt{C_{u} r} \sigma_{1} + \sigma_{\xi} \right) \left(\frac{\sigma_{1}}{\sigma_{r}} \frac{\sqrt{r(d_{1} \vee d_{2})}}{n\varepsilon} + \frac{r}{n\varepsilon} \right) \log n \log^{\frac{1}{2}}(\frac{3.75}{\delta})}_{e_{2}}}_{e_{2}} \end{split}$$

with probability at least $1 - (d_1 + d_2)^{-10} - n^{-9} - \exp(-d_1) - \exp(-d_2) - 10^{-20r}$.

In Theorem 1, the sample size condition $n \ge n_0$ ensures that the spectral norm of perturbations is small enough, i.e., $\|\Delta\| + \max_{i \in [n]} \|\Delta^{(i)}\| \le \sigma_r/2$, to apply Lemma 2 and obtain a sharp characterization on $\Delta^{(1)} \triangleq \Delta_{\widehat{U}\widehat{U}^{\top}} \lor \Delta_{\widehat{V}\widehat{V}^{\top}}$. Theorem 1 also provides an upper bound on the sensitivity of *pseudo singular values*, which is of the order $\Delta^{(2)} \triangleq \Delta_{\widetilde{U}^{\top}\widehat{L}\widetilde{V}} \asymp \sigma_1\Delta^{(1)}$. Based on these results, Algorithm 1 outputs an (ε, δ) -DP initialization \widetilde{M}_0 under the sample size condition

$$n \ge \widetilde{O}\left((\kappa^2 r^2 + \kappa_{\xi} r)(d_1 \lor d_2)\right),$$

with an upper bound on the error $\|\widetilde{M}_0 - M\|$ consisting of two terms. The first term e_1 accounts for the statistical error of \widehat{M}_0 and is greater than the the optimal rate $\sigma_{\xi}\sqrt{\frac{d_1\vee d_2}{n}}$ (Koltchinskii, 2011). The second term e_2 can be further decomposed into the cost of privacy on the singular subspaces which is of the order $\widetilde{O}_p\left(\frac{\sigma_1}{\sigma_r}\left(\sigma_1\sqrt{r}+\sigma_{\xi}\right)\frac{\sqrt{(d_1\vee d_2)}}{n\varepsilon}\right)$, and the cost of privacy arises from privatizing the singular values by Gaussian mechanism which is of the order $\widetilde{O}_p\left((\sigma_1\sqrt{r}+\sigma_{\xi})r/(n\varepsilon)\right)$.

Next, Corollary 1 gives the sample size required by a DP-initialization \widetilde{M}_0 that falls within a local ball of M. The proof of Corollary 1 is deferred to Appendix A.5.

Corollary 1. Under the conditions stated in Theorem 1, as the sample size is sufficiently large

$$n \geq C_1 \max\left\{\underbrace{C_l^{-1} \left(\frac{\sqrt{C_u r} \sigma_1 + \sigma_{\xi}}{\sigma_r}\right)^2 \kappa^2 r(d_1 \vee d_2) \log(d_1 + d_2)}_{n_1}, \underbrace{\sqrt{C_l^{-1} \left(\frac{\sqrt{C_u r} \sigma_1 + \sigma_{\xi}}{\sigma_r}\right) \left(\kappa r \sqrt{d_1 \vee d_2} + r^{\frac{3}{2}}\right) \log n \frac{\log^{\frac{1}{2}}(\frac{3.75}{\delta})}{\varepsilon}}_{n_2}}_{n_2}\right\},$$

for some absolute constant $c_2 > 0$, then we have, for some small constant $0 < c_0 < 1$,

$$\|\widetilde{M}_0 - M\|_F \le \sqrt{2r} \|\widetilde{M}_0 - M\| \le c_0 \sigma_r.$$
⁽²⁾

In Corollary 1, the error due to $||M|| \cdot \left(\left\| \widehat{V}\widehat{V}^{\top} - VV^{\top} \right\| + \left\| \widehat{U}\widehat{U}^{\top} - UU^{\top} \right\| \right)$ dominates the statistical error $||\widehat{L} - M||$ and the sample size n_1 is required to control these two terms; the sample size n_2 controls the error due to privatizing the singular subspaces and singular values. According to Corollary 1, as the sample size $n \ge \widetilde{O}\left((\kappa^4 r^2 + \kappa^2 \kappa_{\xi}^2 r)(d_1 \lor d_2)\right)$, the (ε, δ) -DP \widetilde{M}_0 is guaranteed to fall into a local ball surrounding M, as stated in (2). The condition (2) is a pre-requisite for Algorithm 2 to converge geometrically, as discussed in Theorem 3.

3 Minimax lower bounds

This section applies DP-Fano's lemma (See Lemma 3) to establish the DP-constrained minimax lower bound of estimating the matrix $M \in \mathbb{M}_r := \{M \in \mathbb{R}^{d_1 \times d_2} : \operatorname{rank}(M) = r\}$ under trace regression model

$$f_M(y_i|X_i) = \frac{1}{\sqrt{2\pi\sigma_{\xi}}} \exp\left(\frac{-\left(y_i - \langle X_i, M \rangle\right)^2}{2\sigma^2}\right); X_i \sim \mathcal{N}\left(\mathbf{0}, \Lambda_i\right).$$
(3)

Suppose we observe an i.i.d. sample $\{(X_i, y_i), (X'_i, y'_i)\}_{i \in [n]}$ of size 2n drawn from (3). Then, we have

$$\bar{y}_i := y_i + y'_i = \left\langle \left(\begin{array}{cc} 0 & M \\ M^\top & 0 \end{array} \right), \left(\begin{array}{cc} 0 & X_i \\ X'^\top & 0 \end{array} \right) \right\rangle + \xi_i + \xi'_i,$$

where the underlying matrix M_* . Let $f(X_i, X'_i)$ be the joint distribution of X_i and X'_i ; $f_{M_*}(\bar{y}_i | X_i, X'_i)$ be the conditional distribution of \bar{y}_i given X_i, X'_i ; and denote the joint distribution of \bar{y}_i and X_i, X'_i as $f_{M_*}(\bar{y}_i, X_i, X'_i)$. It is clear that $f_{M_*}(\bar{y}_i | X_i, X'_i)$ is given by the distribution of

$$\mathcal{N}\left(\left\langle \left(\begin{array}{cc} 0 & M \\ M^{\top} & 0 \end{array}\right), \left(\begin{array}{cc} 0 & X_i \\ X_i^{\prime \top} & 0 \end{array}\right)\right\rangle, 2\sigma_{\xi}^2\right).$$

Let \otimes represent the tensor product of marginal laws. For a given matrix $\Sigma = \text{diag}\{\sigma_1, \dots, \sigma_r\}$ where $C\sigma \geq \sigma_1 \dots \geq \sigma_r \geq c\sigma$ for some constants $\sigma > 0$ and C > c > 0, we consider the family of normal distribution under trace regression model:

$$\mathcal{P}_{\Sigma} := \Big\{ \bigotimes_{i=1}^{n} f_{M_{*}}(\bar{y}_{i}, X_{i}, X_{i}') : M_{*} = (U\Sigma V^{\top})_{*}, U \in \mathbb{O}_{d_{1}, r}, V \in \mathbb{O}_{d_{2}, r} \Big\}.$$

By definition, each distribution $P_{M_*} \in \mathcal{P}_{\Sigma}$ is indexed by $U_{M_*} = \frac{1}{\sqrt{2}} \begin{pmatrix} U & U \\ V & -V \end{pmatrix} \in \mathbb{O}_{d_1+d_2,2r}$ whose columns are the *r* eigenvectors of M_* . Next, we employ DP-Fano's lemma to derive the minimax lower bound of estimating M_* by a sample drawn from \mathcal{P}_{Σ} . Let $\mathrm{KL}(\cdot \| \cdot)$ and $\mathrm{TV}(\cdot, \cdot)$ denote the Kullback-Leibler divergence and total variation distance between two distributions.

Lemma 3 (DP-Fano's lemma, Acharya et al. (2021)). Let $\mathcal{P} := \{P : P = \mu^{(1)} \times \cdots \times \mu^{(n)}\}$ be a family of product measures indexed by a parameter from a pseudo-metric space (Θ, ρ) . Denote $\theta(P) \in \Theta$ the parameter associated with the distribution P. Let $\mathcal{Q} = \{P_1, \cdots, P_N\} \subset \mathcal{P}$ contain N probability measures and there exist constants $\rho_0, l_0, t_0 > 0$ such that for all $i \neq i' \in [N]$, $\rho(\theta(P_i), \theta(P_{i'})) \ge \rho_0$, $\operatorname{KL}(P_i || P_{i'}) \le l_0$, $\sum_{k \in [n]} \operatorname{TV}(\mu_i^{(k)}, \mu_{i'}^{(k)}) \le t_0$, where $P_i = \mu_i^{(1)} \times \cdots \times \mu_i^{(n)}$ and $P_{i'} = \mu_{i'}^{(1)} \times \cdots \times \mu_{i'}^{(n)}$. Suppose $\delta \lesssim e^{-n}$, then

$$\inf_{A \in \mathcal{A}_{\varepsilon,\delta}(\mathcal{P})} \sup_{P \in \mathcal{P}} \mathbb{E}_A \ \rho(A, \theta(P)) \ge \max\left\{\frac{\rho_0}{2} \left(1 - \frac{l_0 + \log 2}{\log N}\right), \frac{\rho_0}{4} \left(1 \bigwedge \frac{N - 1}{\exp\left(4\varepsilon t_0\right)}\right)\right\}, \quad (4)$$

where the infimum is taken over all the (ε, δ) -DP randomized algorithm defined by $\mathcal{A}_{\varepsilon,\delta}(\mathcal{P}) := \{A : Z \mapsto \Theta \text{ and } A \text{ is } (\varepsilon, \delta)\text{-differentially private for all } Z \sim P \in \mathcal{P} \}$.

To apply Fano's lemma, we need to construct a large subset with well-separated elements for $\mathbb{O}_{d_1+d_2,2r}$. By Lemma 6, there exists a subset $\mathcal{S}_q^{(d_1+d_2)} \subset \mathbb{O}_{d_1+d_2,2r}$ with cardinality $\left|\mathcal{S}_q^{(d_1+d_2)}\right| \geq 2^{2r(d_1+d_2-2r)}$ such that for any $H \neq H' \in \mathcal{S}_q^{(d_1+d_2)}$,

$$||HH^{\top} - H'H'^{\top}||_q \gtrsim \tau \varepsilon_0 (2r)^{1/q} \text{ and } ||HH^{\top} - H'H'^{\top}||_F \lesssim 2\sqrt{r}\varepsilon_0,$$

for some small constants $\tau, \varepsilon_0 > 0$, where $\|\cdot\|_q$ denotes the Schatten-q norm. We then consider the family of distributions

$$\mathcal{P}_{\sigma} = \left\{ \bigotimes_{i=1}^{n} f_{M_*}(\bar{y}_i, X_i, X_i') : M_* = \sigma H H^{\top}, H \in \mathcal{S}_q^{(d_1+d_2)} \right\} \subset \mathcal{P}_{\Sigma},$$

whose cardinality $N := |\mathcal{P}_{\sigma}| \geq 2^{2r(d_1+d_2-2r)}$. Let $M_* = \sigma H H^{\top}$ and $M'_* = \sigma H' H'^{\top}$. As shown in Appendix B, for any $H \neq H' \in \mathcal{S}_q^{(d_1+d_2)}$, we have

$$\operatorname{KL}\left(\bigotimes_{i=1}^{n} f_{M_{*}}(\bar{y}_{i}, X_{i}, X_{i}') \|\bigotimes_{i=1}^{n} f_{M_{*}'}(\bar{y}_{i}, X_{i}, X_{i}')\right) \lesssim \frac{n}{\sigma_{\xi}^{2}} C_{u} \sigma^{2} r \varepsilon_{0}^{2},$$

and $\sum_{k \in [n]} \text{TV}\left(f_{M_*}(\bar{y}_i, X_i, X'_i), f_{M'_*}(\bar{y}_i, X_i, X'_i)\right) \lesssim n \frac{\sqrt{C_u \sigma}}{\sigma_{\xi}} \sqrt{r} \varepsilon_0$. To this end, we obtain Theorem 2 by applying Lemma 3 with the bounded KL divergence and TV distance, together with the facts that $\mathcal{P}_{\sigma} \subset \mathcal{P}_{\Sigma}$. The proof of Theorem 2 is deferred to Appendix B.

Theorem 2. Consider a sample of size n drawn from the distribution $P_{M^*} \in \mathcal{P}_{\Sigma}$, then for any $\delta \leq e^{-n}$ and any $q \in [1, \infty]$, there exists a constant c > 0

$$\inf_{\widetilde{M}} \sup_{P \in \mathcal{P}_{\Sigma}} \mathbb{E} \left\| \widetilde{M} - M \right\|_{q} \ge c \frac{\sigma_{\xi}}{\sqrt{C_{u}}} \left(r^{1/q} \sqrt{\frac{d_{1} \vee d_{2}}{n}} + r^{\frac{1}{2} + \frac{1}{q}} \frac{d_{1} \vee d_{2}}{n\varepsilon} \right) \bigwedge r^{1/q} \sigma_{q}$$

where the infimum is taken over all possible (ε, δ) -DP algorithms. It suffices to choose $q = 1, 2, \infty$ to obtain the bounds in the nuclear norm, Frobenius norm, and spectral norm, respectively. For example, when q = 2, there exists a constant c > 0

$$\inf_{\widetilde{M}} \sup_{P \in \mathcal{P}_{\Sigma}} \mathbb{E} \left\| \widetilde{M} - M \right\|_{F} \ge c \left(\underbrace{\frac{\sigma_{\xi}}{\sqrt{C_{u}}} \sqrt{\frac{r(d_{1} \vee d_{2})}{n}}}_{l_{1}} + \underbrace{\frac{\sigma_{\xi}}{\sqrt{C_{u}}} \frac{r(d_{1} \vee d_{2})}{n\varepsilon}}_{l_{2}} \right) \bigwedge r^{1/2} \sigma.$$
(5)

In Theorem 2, the lower bound (5) consists of two terms, the statistical error l_1 and the cost of privacy l_2 . The next section proposes a DP-estimator that attains the minimax lower bound (5), up to an additional factor σ_r and some logarithmic factors. As a supplement to DP-Fano's Lemma which works for $\delta \leq e^{-n}$, we also try the score attack argument, which is valid for a wider range of $\delta \leq n^{1+\gamma}$ where $\gamma > 0$ is a constant. Theorem 5 presents the DP-constrained lower bound established by the score attack argument. The content and proof of Theorem 5 are deferred to Appendix D. We also point out that it is trivial to derive the minimax lower bound of the case $d_1 = d_2 = d$ based on DP-Fano's Lemma since there is no need to apply the trick of symmetrization.

4 Upper bounds with differential privacy

In this section, we present Algorithm 2, DP-RGrad, and show that DP-RGrad attains the nearoptimal convergence rate for differentially privately estimating low-rank matrices under the trace regression model. Our approach is based on privatizing the Riemannian gradient descent (RGrad) by the Gaussian mechanism. Interested readers may refer to Vandereycken (2013); Edelman et al. (1998); Adler et al. (2002); Absil et al. (2008) for the basics of RGrad. Let the estimate we obtain after *l* iterations be the rank-*r* matrix $M_l \in \mathbb{R}^{d_1 \times d_2}$ whose SVD has the form $M_l = U_l \Sigma_l V_l^{\top}$. It is well-known in Absil et al. (2008); Vandereycken (2013) that the tangent space of \mathbb{M}_r at M_l is given by $\mathbb{T}_l := \{Z \in \mathbb{R}^{d_1 \times d_2} : Z = U_l R^{\top} + L V_l^{\top}, R \in \mathbb{R}^{d_2 \times r}, L \in \mathbb{R}^{d_1 \times r}\}$. The projection of the gradient G_l onto \mathbb{T}_l is $\mathcal{P}_{\mathbb{T}_l}(G_l) = U_l U_l^{\top} G_l + G_l V_l V_l^{\top} - U_l U_l^{\top} G_l V_l V_l^{\top}$, which is of rank at most 2r. Let the noisy gradient descent on the tangent space be $M_l - \eta_l \mathcal{P}_{\mathbb{T}_l}(G_l) + \mathcal{P}_{\mathbb{T}_l} N_l$ where $N_l \in \mathbb{R}^{d_1 \times d_2}$ is the Gaussian noise matrix. Then, we retract it back to \mathbb{M}_r and obtain

$$M_{l+1} = \text{SVD}_r \left(M_l - \eta_l \mathcal{P}_{\mathbb{T}_l} \left(G_l \right) + \mathcal{P}_{\mathbb{T}_l} N_l \right).$$
(6)

We update the estimate as defined in (6) for $1 = 0, \dots, l^* - 1$ where l^* is the total number of iterations. Thanks to the composition property and Gaussian mechanism, we only need to ensure

that each iteration is $(\varepsilon/l^*, \delta/l^*)$ -DP. For trace regression model defined in (1), empirical mean squared loss is defined as $\mathcal{L}_n(M_l; Z) := \frac{1}{2n} \sum_{i=1}^n (\langle X_i, M_l \rangle - y_i)^2$ and the empirical Euclidean gradient is $G_l := \nabla \mathcal{L}_n(M_l; Z) = \frac{1}{n} \sum_{i=1}^n (\langle X_i, M_l \rangle - y_i) X_i$. The sensitivity of the *l*-th iteration is $\Delta_l := \max_{\text{neighbouring}(Z,Z')} \|M_l - \eta \mathcal{P}_{\mathbb{T}_l}(G_l(Z)) - [M_l - \eta \mathcal{P}_{\mathbb{T}_l}(G_l(Z'))]\|_F$.

Algorithm 2 DP-RGrad for trace regression

Input: the loss function \mathcal{L} ; the data set $\{(X_i, y_i)\}_{i=1}^n$; sensitivities $\{\Delta_l\}_{l \in [l^*]}$; DP-initialization \widetilde{M}_0 ; rank r; nuisance variance σ_{ξ}^2 ; privacy budget $\varepsilon > 0, \delta \in (0, 1)$.

Output: (ε, δ) -differentially private estimate M_{l^*} for trace regression. Initialization: $M_0 \leftarrow \widetilde{M}_0$.

for $l + 1 \in [l^*]$ do

$$M_{l+1} \leftarrow \text{SVD}_r \left(M_l - \eta_l \mathcal{P}_{\mathbb{T}_l} \left(G_l \right) + \mathcal{P}_{\mathbb{T}_l} N_l \right)$$

where G_l is the empirical Euclidean gradient

$$G_l := \nabla \mathcal{L}_n(M_l; Z) = \frac{1}{n} \sum_{i=1}^n \left(\langle X_i, M_l \rangle - y_i \right) X_i,$$

and $N_l \in \mathbb{R}^{d_1 \times d_2}$ has entries i.i.d. to

$$\mathcal{N}\left(0, \frac{2\Delta_l^2 l^{*2}}{\varepsilon^2} \log\left(\frac{1.25l^*}{\delta}\right)\right).$$

Return: $\widetilde{M}_{l^*} \longleftarrow M_{l^*}$

Theorem 3 establishes the error bound of the estimator \widetilde{M}_{l^*} given by Algorithm 2. The proof of Theorem 3 is deferred to Appendix C.

Theorem 3. Consider i.i.d. observations $Z = \{z_1, \dots, z_n\}$ drawn from the trace regression model stated in (1) where the true low-rank regression coefficients matrix being $M \in \mathbb{M}_r$. Here, $z_i := (X_i, y_i)$ for $i = 1, \dots, n$ and we assume that $\{X_i\}_{i \in [n]}$ satisfy the Assumption 1. Under the Assumption 1 and the condition that $(d_1 + d_2) > \log n$, suppose that Algorithm 2 takes in an (ε, δ) -DP initialization such that for some small constant $0 < c_0 < 1$, $\|\widetilde{M}_0 - M\|_F \le \sqrt{2r}\|\widetilde{M}_0 - M\| \le c_0\sigma_r$, and the sensitivities Δ_l take the value

$$\Delta_l = C_3 \frac{\eta}{n} (\sigma_{\xi} + \sigma_r \sqrt{C_u}) \sqrt{C_u r (d_1 + d_2) \log n}, \quad \text{for all} \quad l = 1, \cdots, l^*,$$

for some absolute constant $C_3 > 0$, then we have, Algorithm 2 is $(2\varepsilon, 2\delta)$ -differentially private.

Moreover, as the sample size

$$n \ge c_4 \max\left\{\underbrace{\underbrace{c_1 r(d_1 + d_2)}_{n_3}, \underbrace{\eta^2 \kappa_{\xi}^2 C_u r(d_1 + d_2) \log(d_1 + d_2)}_{n_4}}_{\eta \sqrt{C_u} \left(\kappa_{\xi} + \sqrt{C_u}\right) r(d_1 + d_2) \log^{3/2}(n) \frac{\log^{1/2} \left(\frac{1.25 \log(n)}{\delta}\right)}{\varepsilon}\right\},$$

for some small constant $0 < c_4 < 1$, number of iteration $l^* = O(\log n)$, and the step size $0 < \eta < 1$, we have the output \widetilde{M}_{l^*} of Algorithm 2 satisfies

$$\left\|\widetilde{M}_{l^{*}} - M\right\|_{F} \leq \underbrace{C_{4}\sigma_{\xi}\sqrt{C_{u}}\sqrt{\frac{r(d_{1}+d_{2})}{n}}\log^{1/2}(d_{1}+d_{2})}_{u_{1}} + \underbrace{C_{4}\sqrt{C_{u}}(\sigma_{\xi}+\sigma_{r}\sqrt{C_{u}})\frac{r(d_{1}+d_{2})}{n\varepsilon}\log^{3/2}n\log^{1/2}\left(\frac{1.25\log(n)}{\delta}\right)}_{u_{2}}.$$

with probability at least

$$1 - \tilde{c}_2 \exp\left(-\tilde{c}_3 r(d_1 + d_2)\right) - (d_1 + d_2)^{-10} - n^{-9} - e^{-d_1} - e^{-d_2} - 10^{-20r} - \left((d_1 + d_2)^{-10} + \exp\left(-(d_1 + d_2)\right) + n^{-9} + \exp\left(-10C_u(d_1 + d_2)\right)n^{-9}\right) \log n.$$

According to Theorem 3, the upper bound of $\|\widetilde{M}_{l^*} - M\|_F$ can be decomposed into the the statistical error u_1 and the cost of privacy u_2 . The term u_1 matches the the optimal rate $l_1 \sim \sigma_{\xi} \sqrt{\frac{r(d_1 \vee d_2)}{n}}$, only up to logarithmic factors. However, the term u_2 differs from the theoretical lower bound of the cost of privacy $l_2 \sim \sigma_{\xi} \frac{r(d_1+d_2)}{n\varepsilon}$, by a non-trivial factor σ_r , apart from logarithmic factors. In conclusion, Theorem 3 shows that as the sample size $n \gtrsim \widetilde{O}\left(\left(\kappa_{\xi}^2 \vee \kappa_{\xi}\right) r(d_1 \vee d_2)\right)$, the estimator \widetilde{M}_{l^*} given by Algorithm 2 attains the near-optimal convergence rate

$$\widetilde{O}_p\left(\sigma_{\xi}\sqrt{\frac{r(d_1+d_2)}{n}} + (\sigma_{\xi}+\sigma_r)\frac{r(d_1+d_2)}{n\varepsilon}\right).$$
(7)

The sample size requirement of Theorem 3 has the following explanations. The sample size n_3 is required to guarantee that the RIP condition stated in Lemma 1 occurs with high probability. The sample size n_4 is necessary to control the statistical error contributed by $\sum_{i \in [n]} \xi_i X_i$ in each iteration where ξ_i is the model noise. The sample size n_5 arises from controlling the cost of privacy due to $\mathcal{P}_{\mathbb{T}_l} N_l$ in each iteration.

5 Discussion

In this section, we discuss the non-trivial gap σ_r between $u_2 \sim (\sigma_{\xi} + \sigma_r) \frac{r(d_1+d_2)}{n\varepsilon}$ and $l_2 \sim \sigma_{\xi} \frac{r(d_1 \vee d_2)}{n\varepsilon}$. Note that l_2 is free of σ_r while u_2 contains the factor σ_r arising from sensitivities

$$\Delta_l \asymp \frac{\eta}{n} (\sigma_{\xi} + \sigma_r \sqrt{C_u}) \sqrt{C_u r (d_1 + d_2) \log n} \quad \text{for} \quad l = 1, \cdots, l^*$$

The quantity $\sigma_r \sqrt{C_u}$ of Δ_l arises from $\|\langle M_l - M, X_i \rangle\|_{\psi_2} \leq \sqrt{C_u} \|M_l - M\|_F$, as elaborated in (S.12). Here, $\|\cdot\|_{\psi_2}$ denotes the sub-Gaussian norm. Therefore, we cannot get rid of the factor σ_r once the measurement matrices $\{X_i\}_{i\in[n]}$ are subject to differential privacy. In many real applications, however, the measurement matrices $\{X_i\}_{i\in[n]}$ are fixed with deterministic designs. People publish $\{X_i\}_{i\in[n]}$ to the public with little concern on the privacy of $\{X_i\}_{i\in[n]}$. Although the exposure of $\{X_i\}_{i\in[n]}$ alone will not reveal any information on M, the privacy of M suffers from leakage when the public has access to the joint observations $\{(X_i, y_i)\}_{i\in[n]}$. We, therefore, introduce the following notion of privacy for neighboring datasets sharing the same measurement matrix.

Definition 1 (weak (ε, δ) -differential privacy). The algorithm A that maps Z into $\mathbb{R}^{d_1 \times d_2}$ is weak (ε, δ) -differentially private over the dataset Z if $\mathbb{P}(A(Z) \in \mathcal{Q}) \leq e^{\varepsilon} \mathbb{P}(A(Z') \in \mathcal{Q}) + \delta$, for all neighbouring data set Z, Z' sharing the same measurement X and all subset $\mathcal{Q} \subset \mathbb{R}^{d_1 \times d_2}$.

In Theorem 7, Appendix F, we show that as $n \gtrsim \widetilde{O}\left(\left(\kappa_{\xi}^2 \lor \kappa_{\xi}\right) r(d_1 \lor d_2)\right)$, the estimator \widetilde{M}_{l^*} given by Algorithm 2 attains the optimal convergence rate $\widetilde{O}_p\left(\sigma_{\xi}\sqrt{\frac{r(d_1+d_2)}{n}} + \sigma_{\xi}\frac{r(d_1+d_2)}{n\varepsilon}\right)$ in the sense of weak differential privacy. The analogs of Theorem 1, Corollary 1 and Theorem 3 in the sense of weak differential privacy can be found as Theorem 6, Corollary 2 and Theorem 7 in Appendix F. It is interesting to explore in future work whether the score attack argument or DP-Fano's Lemma can be generalized to include the non-trivial factor σ_r .

Acknowledgement

The author would like to express sincere gratitude to Hong Kong PhD Fellowship Scheme and Hong Kong RGC GRF Grant 16301622 for providing financial support for this research. The author also wishes to acknowledge the invaluable guidance provided by Prof. Dong, Xia throughout the research process. Additionally, the author would like to extend heartfelt thanks to Mr. Zetao, Fei for his constructive criticism during the paper revision. Their contributions have been instrumental in the successful completion of this research.

A Proof of Theorem 1

The proof of Theorem 1 consists of four parts. In Part A.1, we list several existing results that are useful in the proofs later. In Part A.2, Lemma 2 works as the main technique to derive the sensitivity $\Delta^{(1)}$. Part A.3 derives the sensitivity $\Delta^{(2)}$. Part A.4 establishes the upper bounds of $\|\widetilde{M}_0 - M\|$ and $\|\widetilde{M}_0 - M\|_F$ based on the $\Delta^{(1)}$ and $\Delta^{(2)}$.

A.1 Part 1

The following Theorem 4 proposed by Proposition 2, Koltchinskii (2011) will be frequently used to establish the upper bound of the spectral norm of a summation of independent random matrices.

Theorem 4 (Bernstein's inequality, Koltchinskii (2011)). Let B_1, \dots, B_n be independent $d_1 \times d_2$ matrices such that for some $\alpha \ge 1$ and all $i \in [n]$

$$\mathbb{E}B_i = 0, \quad \|\Lambda_{\max}(B_i)\|_{\Psi_{\alpha}} =: K < +\infty.$$

Let

$$S^{2} := \max\left\{\Lambda_{\max}\left(\sum_{i=1}^{n} \mathbb{E}B_{i}B_{i}^{\top}\right)/n, \Lambda_{\max}\left(\sum_{i=1}^{n} \mathbb{E}B_{i}^{\top}B_{i}\right)/n\right\}.$$

Then, for some constant C > 0 and for all t > 0,

$$\mathbb{P}\left(\left\|\frac{1}{n}\sum_{i=1}^{n}B_{i}\right\| \geq CS\sqrt{\frac{t+\log(d_{1}+d_{2})}{n}} + CK\log^{1/\alpha}\left(\frac{K}{S}\right)\frac{t+\log(d_{1}+d_{2})}{n}\right) \leq \exp(-t).$$

Theorem 4 applies to bound the spectral norm of $\Delta := \hat{L} - M$. The existing result for the case of heavy-tailed noise can be found in Theorem 6, Shen et al. (2023). Adapting the existing result to the case of Gaussian noise, we have that for some absolute constant $C_0 > 0$,

$$\|\Delta\| = \|\widehat{L} - M\| \le C_0 \sqrt{C_l^{-1}} \left(\sigma_{\xi} + \sqrt{C_u} \sqrt{r} \sigma_1\right) \sqrt{\frac{(d_1 \lor d_2) \log(d_1 + d_2)}{n}},$$
(S.1)

with probability at least $1 - (d_1 + d_2)^{-10}$. The following Lemma originated from Lemma 18, Shen et al. (2023), is useful to analyze the matrix permutation due to singular value decomposition.

Lemma 4 (Matrix Permutation, Shen et al. (2023)). Let $M \in \mathbb{R}^{d_1 \times d_2}$ be a rank-r matrix with singular value decomposition of the form $M = U\Sigma V^{\top}$ where $\Sigma = \text{diag} \{\sigma_1, \sigma_2, \cdots, \sigma_r\}$ with

 $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_r > 0$. For any $\widehat{M} \in \mathbb{R}^{d \times d}$ satisfying $\|\widehat{M} - M\|_{\mathrm{F}} < \sigma_r/8$, then

$$\left\| \operatorname{SVD}_{r}(\widehat{M}) - M \right\| \leq \|\widehat{M} - M\| + 40 \frac{\|\widehat{M} - M\|^{2}}{\sigma_{r}},$$
$$\left\| \operatorname{SVD}_{r}(\widehat{M}) - M \right\|_{F} \leq \|\widehat{M} - M\|_{F} + 40 \frac{\|\widehat{M} - M\|\|\widehat{M} - M\|_{F}}{\sigma_{r}}$$

and

$$\left\|\widehat{U}\widehat{U}^{\top} - UU^{\top}\right\| \le \frac{8}{\sigma_r} \|\widehat{M} - M\|, \quad \left\|\widehat{V}\widehat{V}^{\top} - VV^{\top}\right\| \le \frac{8}{\sigma_r} \|\widehat{M} - M\|,$$

where the leading r left singular vectors of \widehat{M} are given by the columns of $\widehat{U} \in \mathbb{R}^{d_1 \times r}$ and the leading r right singular vectors of \widehat{M} are given by the columns of $\widehat{V} \in \mathbb{R}^{d_2 \times r}$.

A.2 Part 2

The second part aims to derive the sensivitity

$$\Delta^{(1)} := \max_{i \in [n]} \left(\|\widehat{U}\widehat{U}^{\top} - \widehat{U}^{(i)}\widehat{U}^{(i)\top}\|_F \lor \|\widehat{V}\widehat{V}^{\top} - \widehat{V}^{(i)}\widehat{V}^{(i)\top}\|_F \right).$$

Before moving on, we present Lemma 5, which provides conclusions on Δ and $\Delta^{(i)}$, frequently used in the proof later. The proof of Lemma 5 can be found in Appendix E.4.

Lemma 5. Under model (1), Assumption 1, and the condition $n \ge \frac{\log^2 n}{(d_1 \lor d_2) \log(d_1 + d_2)}$, there exists some absolute constant $C_0, C_1 > 0$ such that the event

$$\mathcal{E}_{*} := \left\{ \max_{i \in [n]} \left\| \Delta - \Delta^{(i)} \right\| \le C_{0} \cdot n^{-1} \sqrt{C_{l}^{-1}} \left(\sqrt{C_{u} r} \sigma_{1} + \sigma_{\xi} \right) \log n \right\}$$
$$\bigcap \left\{ \left\| \Delta \right\| + \max_{i \in [n]} \left\| \Delta^{(i)} \right\| \le C_{0} \sqrt{C_{l}^{-1}} \left(\sigma_{\xi} + \sqrt{C_{u}} \sqrt{r} \sigma_{1} \right) \sqrt{\frac{(d_{1} \lor d_{2}) \log(d_{1} + d_{2})}{n}} \right\},$$

holds with probability at least $1 - (d_1 + d_2)^{-10} - n^{-9}$. Conditioned on the event \mathcal{E}_* , as the sample size satisfies

$$n \ge C_1 C_l^{-1} r \left(\frac{\sigma_{\xi} + \sqrt{C_u r} \sigma_1}{\sigma_r}\right)^2 (d_1 \lor d_2) \log(d_1 + d_2), \tag{S.2}$$

we have

$$\|\Delta_*\| \vee \max_{i \in [n]} \|\Delta_*^{(i)}\| = \|\Delta\| \vee \max_{i \in [n]} \|\Delta^{(i)}\| \le \frac{\sigma_r}{8\sqrt{2r}} < \frac{\sigma_r}{5+\delta} < \frac{\sigma_r}{2},$$
 (S.3)

and

$$|\Delta_*\|_F \vee \max_{i \in [n]} \|\Delta_*^{(i)}\|_F = \max_{i \in [n]} \|\Delta\|_F \vee \|\Delta^{(i)}\|_F \le \frac{\sigma_r}{8}$$

for some constant $\delta > 0$, where $\Delta_*^{(i)}$ is the symmetric dilation of $\Delta^{(i)} := L^{(i)} - M$.

The following analysis is proceeded under the sample size condition (S.2) and is mainly conditioned on the event \mathcal{E}_* which happens with probability at least $1 - (d_1 + d_2)^{-10} - n^{-9}$.

Step 1: expansion. Conditioned on \mathcal{E}_* , we are able to apply Lemma 2 to Δ_* and $\Delta_*^{(i)}$ and get

$$\begin{pmatrix} \widehat{U}\widehat{U}^{\top} - \widehat{U}^{(i)}\widehat{U}^{(i)\top} & 0\\ 0 & \widehat{V}\widehat{V}^{\top} - \widehat{V}^{(i)}\widehat{V}^{(i)\top} \end{pmatrix} = \sum_{k\geq 1} \mathcal{S}_{M_*,k}(\Delta_*) - \sum_{k\geq 1} \mathcal{S}_{M_*,k}(\Delta_*^{(i)}).$$

Our goal is to bound $\Delta^{(1)}$ which satisfies

$$\Delta^{(1)} \leq \max_{i \in [n]} \left(\| \widehat{U}\widehat{U}^{\top} - \widehat{U}^{(i)}\widehat{U}^{(i)\top} \|_{F} + \| \widehat{V}\widehat{V}^{\top} - \widehat{V}^{(i)}\widehat{V}^{(i)\top} \|_{F} \right)$$

$$\leq \max_{i \in [n]} \left(\left\| \mathcal{S}_{M_{*},1}(\Delta_{*}) - \mathcal{S}_{M_{*},1}(\Delta_{*}^{(i)}) \right\|_{F} + \left\| \sum_{k \geq 2} \mathcal{S}_{M_{*},k}(\Delta_{*}) - \sum_{k \geq 2} \mathcal{S}_{M_{*},k}(\Delta_{*}^{(i)}) \right\|_{F} \right).$$

Step 2: bounding the first order term. By the definition of $\mathcal{S}_{M_*,1}(\Delta_*)$ and $\mathcal{S}_{M_*,1}(\Delta_*^{(i)})$,

$$\max_{i \in [n]} \| \mathcal{S}_{M_{*},1}(\Delta_{*}) - \mathcal{S}_{M_{*},1}(\Delta_{*}^{(i)}) \| = \max_{i \in [n]} \| Q^{-1}(\Delta - \Delta^{(i)})^{\top} Q_{\perp} + Q_{\perp}(\Delta - \Delta^{(i)})^{\top} Q^{-1} \|$$

$$\leq \frac{2\sqrt{r}}{\sigma_{r}} \max_{i \in [n]} \| \Delta - \Delta^{(i)} \| \leq C_{4} \frac{\sqrt{r}}{n\sigma_{r}} \sqrt{C_{l}^{-1}} \left(\sqrt{C_{u}r}\sigma_{1} + \sigma_{\xi} \right) \log n,$$
(S.4)

conditioned on the event \mathcal{E}_* , for some absolute constant $C_4 > 0$.

Step 3: bounding the higher order terms. Let I_k be the index set for terms in $\mathcal{S}_{M_*,k}$

$$I_{k} = \left\{ \mathbf{s} : \mathbf{s} = (s_{1}, \dots, s_{k+1}), \sum_{m=1}^{k+1} s_{m} = k, s_{m} \ge 0 \quad \forall m \in [k+1] \right\},\$$

with the cardinality $|I_k| = \binom{2k}{k}$. We define

$$\mathcal{T}_{M_{*},k,\mathbf{s},l}\left(\Delta_{*}-\Delta_{*}^{(i)}\right) := Q^{-s_{1}}\Delta_{*}^{(i)}Q^{-s_{2}}\cdots Q^{-s_{l}}\left(\Delta_{*}-\Delta_{*}^{(i)}\right)Q^{s_{l+1}}\cdots Q^{-s_{k}}\Delta_{*}Q^{s_{k+1}},$$

for $k \geq 2, \mathbf{s} = (s_1, \cdots, s_{k+1}) \in I_k$ and $l \in [k]$. Since $|I_k| = \binom{2k}{k}$, the higher order terms $\max_{i \in [n]} \left\| \sum_{k \geq 2} \mathcal{S}_{M_*,k}(\Delta_*) - \sum_{k \geq 2} \mathcal{S}_{M_*,k}\left(\Delta_*^{(i)}\right) \right\|_{\mathrm{F}}$

$$= \max_{i \in [n]} \left\| \sum_{k \ge 2} \sum_{\mathbf{s} \in I_k} \sum_{l \in [k]} \mathcal{T}_{M_*,k,\mathbf{s},l} \left(\Delta_* - \Delta_*^{(i)} \right) \right\|_{\mathcal{F}} \le \max_{i \in [n]} \sum_{k \ge 2} \binom{2k}{k} \sum_{l \in [k]} \left\| \mathcal{T}_{M_*,k,\mathbf{s},l} \left(\Delta_* - \Delta_*^{(i)} \right) \right\|_{\mathcal{F}}.$$
 (S.5)

It is sufficient to find a upper bound of $\left\| \mathcal{T}_{M_*,k,\mathbf{s},l} \left(\Delta_* - \Delta_*^{(i)} \right) \right\|_{\mathrm{F}}$. Denote

$$D_{\max} := C_1 \sqrt{C_l^{-1}} \left(\sigma_{\xi} + \sqrt{C_u} \sqrt{r} \sigma_1 \right) \sqrt{\frac{(d_1 \vee d_2) \log(d_1 + d_2)}{n}},$$

which appeared in the event \mathcal{E}_* as an upper bound of $\|\Delta\| + \max_{i \in [n]} \|\Delta^{(i)}\|$.

Conditioned on \mathcal{E}_* , for all $i \in [n], k \ge 2, \mathbf{s} \in I_k$ and $l \in [k]$,

$$\left\|\mathcal{T}_{M_{*},k,\mathbf{s},l}\left(\Delta-\Delta^{(i)}\right)\right\|_{\mathrm{F}} \leq \sqrt{2r} \left\|\mathcal{T}_{M_{*},k,\mathbf{s},l}\left(\Delta-\Delta^{(i)}\right)\right\| \leq \frac{\sqrt{2r}}{\sigma_{r}} \left\|\Delta_{*}-\Delta^{(i)}_{*}\right\| \left(\frac{D_{\max}}{\sigma_{r}}\right)^{k-1},$$

where the first inequality is because $\mathcal{T}_{M_*,k,\mathbf{s},l}\left(\Delta_* - \Delta_*^{(i)}\right)$ is of rank at most 2r. Let a be a function defined by $a(k) = \binom{2k}{k}k$, then a(2) = 12 and $\frac{a(k+1)}{a(k)} \leq 5$ for all integer $k \geq 2$,

$$\max_{i \in [n]} \sum_{k \ge 2} {\binom{2k}{k}} \sum_{l \in [k]} \left\| \mathcal{T}_{M_*,k,\mathbf{s},l} \left(\Delta_* - \Delta_*^{(i)} \right) \right\|_{\mathrm{F}} \\
\leq \max_{i \in [n]} {\binom{4}{2}} \sum_{k \ge 0} 5^k \left(\frac{\left\| D_{\max} \right\|}{\sigma_r} \right)^k \frac{\sqrt{2r}}{\sigma_r} \left\| \Delta_* - \Delta_*^{(i)} \right\| \left(\frac{D_{\max}}{\sigma_r} \right) \\
\leq \max_{i \in [n]} a(2) \frac{\sqrt{2r}}{\sigma_r} \left\| \Delta_* - \Delta_*^{(i)} \right\| \sum_{k \ge 0} \left(\frac{5}{5+\delta} \right)^k \left(\frac{D_{\max}}{\sigma_r} \right) \\
\leq \max_{i \in [n]} a(2) \left(\frac{5+\delta}{\delta} \right) \left(\frac{D_{\max}}{\sigma_r} \right) \frac{\sqrt{2r}}{\sigma_r} \left\| \Delta_* - \Delta_*^{(i)} \right\|,$$
(S.6)

where the last step is due to (S.3), which is guaranteed by the sample size condition (S.2) together with the event \mathcal{E}_* . Combining (S.5) and (S.6), since $\left\|\Delta_* - \Delta_*^{(i)}\right\| = \left\|\Delta - \Delta^{(i)}\right\|$, conditioned on the event \mathcal{E}_* , we have

$$\max_{i \in [n]} \left\| \sum_{k \ge 2} \mathcal{S}_{M_*,k}(\Delta_*) - \sum_{k \ge 2} \mathcal{S}_{M_*,k}\left(\Delta_*^{(i)}\right) \right\|_{\mathrm{F}} \le C_4\left(\frac{12}{\delta}\right) \frac{\sqrt{2r}}{n} \sqrt{C_l^{-1}} \left(\frac{\sqrt{C_u r} \,\sigma_1 + \sigma_\xi}{\sigma_r}\right) \log n,$$

for some absolute constant $C_3 > 0$. In conclusion, conditioned on \mathcal{E}_* , as the sample size $n \geq C_1 C_l^{-1} r \left(\frac{\sigma_{\xi} + \sqrt{C_u r} \sigma_1}{\sigma_r}\right)^2 (d_1 \vee d_2) \log(d_1 + d_2)$, for some absolute constant $C_1 > 0$, we have $\Delta^{(1)} \leq C_4 \frac{\sqrt{r}}{n} \sqrt{C_l^{-1}} \left(\frac{\sqrt{C_u r} \sigma_1 + \sigma_{\xi}}{\sigma_r}\right) \log n$, for some absolute constant C_4 .

Let $E_U \in \mathbb{R}^{d_1 \times d_1}$ be a symmetric matrix where the entries $(E_U)_{ij}$ i.i.d. to $\mathcal{N}(0, \frac{18\Delta^{(1)^2}}{\varepsilon^2}\log(\frac{3.75}{\delta}))$ for $1 \leq i \leq j \leq d_1$. Then, conditioned on the event \mathcal{E}_* and (S.2), for some absolute constant $\widetilde{C}_4 > 0$, $||E_U|| \leq \widetilde{C}_4 \frac{\sqrt{rd_1}}{n\varepsilon} \sqrt{C_l^{-1}} \left(\frac{\sqrt{C_u r} \sigma_1 + \sigma_{\xi}}{\sigma_r}\right) \log n \log^{\frac{1}{2}}(\frac{3.75}{\delta})$, with probability at least $1 - e^{-d_1}$. Moreover, by Gaussian mechanism, $\widehat{U}\widehat{U}^{\top} + E_U$ is $(\varepsilon/3, \delta/3)$ -DP and thus \widetilde{U} is also $(\varepsilon/3, \delta/3)$ -DP thanks to the post-processing property of differential privacy. Furthermore, by Davis-Kahan's Theorem, for some absolute constant $\tilde{c_0}>0$

$$\begin{aligned} \left\| \widetilde{U}\widetilde{U}^{\top} - UU^{\top} \right\| &\leq 1 \wedge \left(\left\| \widehat{U}\widehat{U}^{\top} - UU^{\top} \right\| + \left\| E_{U} \right\| \right) \stackrel{(a)}{\leq} 1 \wedge \left(\left(\frac{8}{\sigma_{r}} \| \widehat{L} - M \| \wedge 1 \right) + \| E_{U} \| \right) \\ &\leq 1 \wedge \widetilde{c_{0}} \left(\sqrt{C_{l}^{-1}} \left(\frac{\sqrt{C_{u}r} \sigma_{1} + \sigma_{\xi}}{\sigma_{r}} \right) \sqrt{\frac{(d_{1} \vee d_{2}) \log(d_{1} + d_{2})}{n}} \\ &+ \sqrt{C_{l}^{-1}} \left(\frac{\sqrt{C_{u}r} \sigma_{1} + \sigma_{\xi}}{\sigma_{r}} \right) \frac{\sqrt{rd_{1}}}{n\varepsilon} \log n \log^{\frac{1}{2}}(\frac{3.75}{\delta}) \right), \end{aligned}$$
(S.7)

where we apply Lemma 4 in step (a).

Let $E_V \in \mathbb{R}^{d_2 \times d_2}$ be a symmetric matrix with $(E_V)_{ij}$ i.i.d. to $\mathcal{N}(0, \frac{18\Delta^{(1)^2}}{\varepsilon^2}\log(\frac{3.75}{\delta}))$ for $1 \leq i \leq j \leq d_2$, then for some absolute constant $\widetilde{C}_4 > 0$, conditioned on the event \mathcal{E}_* and (S.2), we have $||E_V|| \leq \widetilde{C}_4 \frac{\sqrt{rd_2}}{n\varepsilon} \sqrt{C_l^{-1}} \left(\frac{\sqrt{C_u r} \sigma_1 + \sigma_{\xi}}{\sigma_r}\right) \log n \log^{\frac{1}{2}}(\frac{3.75}{\delta})$, with probability at least $1 - e^{-d_2}$. Moreover, by Gaussian mechanism, $\widehat{V}\widehat{V}^{\top} + E_U$ is $(\varepsilon/3, \delta/3)$ -DP and \widetilde{V} is also $(\varepsilon/3, \delta/3)$ -DP thanks to the post-processing property of differential privacy. Furthermore, by Davis-Kahan's Theorem, for some absolute constant $\widetilde{c}_0 > 0$

$$\left\| \widetilde{V}\widetilde{V}^{\top} - VV^{\top} \right\| \leq 1 \wedge \widetilde{c_0} \left(\sqrt{C_l^{-1}} \left(\frac{\sqrt{C_u r} \sigma_1 + \sigma_{\xi}}{\sigma_r} \right) \sqrt{\frac{(d_1 \vee d_2) \log(d_1 + d_2)}{n}} + \sqrt{C_l^{-1}} \left(\frac{\sqrt{C_u r} \sigma_1 + \sigma_{\xi}}{\sigma_r} \right) \frac{\sqrt{rd_2}}{n\varepsilon} \log n \log^{\frac{1}{2}}(\frac{3.75}{\delta}) \right).$$
(S.8)

A.3 Part 3

Given the $(\varepsilon/3, \delta/3)$ -DP singular vectors $\widetilde{U} \in \mathbb{O}^{d_1 \times r}$ and $\widetilde{V} \in \mathbb{O}^{d_2 \times r}$ obtained in Part A.2, we derive the sensitivity $\Delta^{(2)} := \max_{i \in [n]} \left\| \widetilde{U}^{\top} \left(\widehat{L} - \widehat{L}^{(i)} \right) \widetilde{V} \right\|_F$. Conditioned on the event \mathcal{E}_* ,

$$\Delta^{(2)} := \max_{i \in [n]} \left\| \widetilde{U}^{\top} \left(\widehat{L} - \widehat{L}^{(i)} \right) \widetilde{V} \right\|_{F} \le \max_{i \in [n]} \sqrt{r} \left\| \widehat{L} - \widehat{L}^{(i)} \right\|$$
$$= \max_{i \in [n]} \sqrt{r} \left\| \widehat{L} - M + M - \widehat{L}^{(i)} \right\| = \max_{i \in [n]} \sqrt{r} \left\| \Delta - \Delta^{(i)} \right\|$$
$$\le C_{3} \cdot n^{-1} \sqrt{C_{l}^{-1}} \sqrt{r} \left(\sqrt{C_{u} r} \sigma_{1} + \sigma_{\xi} \right) \log n.$$

Let E_{Σ} be a $r \times r$ matrix with entries i.i.d. to $\mathcal{N}(0, 18\Delta^{(2)^2}\log(\frac{3.75}{\delta})/\varepsilon^2)$, then

$$\|E_{\Sigma}\| \le \widetilde{C}_4 \cdot \frac{r}{n\varepsilon} \sqrt{C_l^{-1}} \left(\sqrt{C_u r} \sigma_1 + \sigma_{\xi} \right) \log n \log^{\frac{1}{2}} \left(\frac{3.75}{\delta} \right), \tag{S.9}$$

for some absolute constant $\widetilde{C}_4 > 0$ with probability at least 10^{-20r} . Moreover, by Gaussian mechanism, $\widetilde{\Sigma} = \widetilde{U}^{\top} \widehat{L} \widetilde{V} + E_{\Sigma}$ is $(\varepsilon/3, \delta/3)$ -differentially private. Thanks to the composition property of differential privacy, the output of Algorithm 1 $\widetilde{M}_0 = \widetilde{U} \widetilde{\Sigma} \widetilde{V}^{\top}$, is (ε, δ) -differentially private.

A.4 Part 4

In this part, we derive the upper bound of $\|\widetilde{M}_0 - M\|$. Note that

$$\|\widetilde{M}_0 - M\| = \left\|\widetilde{U}\widetilde{\Sigma}\widetilde{V}^\top - U\Sigma V^\top\right\| = \left\|\widetilde{U}(\widetilde{U}^\top\widehat{L}\widetilde{V} + E_{\Sigma})\widetilde{V}^\top - UU^\top MVV^\top\right\|,$$

is a $d_1 \times d_2$ matrix of rank at most 2r. Since

$$\begin{split} \left\| \widetilde{U} (\widetilde{U}^{\top} \widehat{L} \widetilde{V} + E_{\Sigma}) \widetilde{V}^{\top} - UU^{\top} M V V^{\top} \right\| &\leq \left\| \widetilde{U} \widetilde{U}^{\top} \widehat{L} \widetilde{V} \widetilde{V}^{\top} - UU^{\top} M V V^{\top} \right\| + \left\| \widetilde{U} E_{\Sigma} \widetilde{V}^{\top} \right\| \\ &\leq \left\| \left(\widetilde{U} \widetilde{U}^{\top} - UU^{\top} \right) \widehat{L} \widetilde{V} \widetilde{V}^{\top} \right\| + \left\| UU^{\top} \left(\widehat{L} - M \right) \widetilde{V} \widetilde{V}^{\top} \right\| + \left\| UU^{\top} M \left(\widetilde{V} \widetilde{V}^{\top} - VV^{\top} \right) \right\| + \left\| \widetilde{U} E_{\Sigma} \widetilde{V}^{\top} \right\| \\ &\leq \left\| \widetilde{U} \widetilde{U}^{\top} - UU^{\top} \right\| \left\| \widehat{L} \right\| + \left\| \widehat{L} - M \right\| + \left\| M \right\| \left\| \widetilde{V} \widetilde{V}^{\top} - VV^{\top} \right\| + \left\| E_{\Sigma} \right\| \\ &\leq \left\| \widetilde{U} \widetilde{U}^{\top} - UU^{\top} \right\| \left\| \widehat{L} - M \right\| + \left\| \widetilde{U} \widetilde{U}^{\top} - UU^{\top} \right\| \left\| M \right\| + \left\| \widehat{L} - M \right\| + \left\| M \right\| \left\| \widetilde{V} \widetilde{V}^{\top} - VV^{\top} \right\| + \left\| E_{\Sigma} \right\| \\ &\leq 2 \left\| \widehat{L} - M \right\| + \left\| M \right\| \left(\left\| \widetilde{V} \widetilde{V}^{\top} - VV^{\top} \right\| + \left\| \widetilde{U} \widetilde{U}^{\top} - UU^{\top} \right\| \right) + \left\| E_{\Sigma} \right\| , \end{split}$$

$$(S.10)$$

it is sufficient to plug in the upper bound (S.1) of $\|\widehat{L} - M\|$, $\|M\| = \sigma_1$, as well as the upper bounds (S.7) of $\|\widetilde{U}\widetilde{U}^{\top} - UU^{\top}\|$, (S.8) of $\|\widetilde{V}\widetilde{V}^{\top} - VV^{\top}\|$ and (S.9) of $\|E_{\Sigma}\|$. In conclusion, conditioned on the event \mathcal{E}_* , as the sample size

$$n \ge C_1 C_l^{-1} r \left(\frac{\sqrt{C_u r} \sigma_1 + \sigma_\xi}{\sigma_r}\right)^2 (d_1 \lor d_2) \log(d_1 + d_2),$$

for some absolute constant $C_1 > 0$, we have

$$\begin{split} \|\widetilde{M}_{0} - M\| &\leq C_{5}\sqrt{C_{l}^{-1}}\left(\sqrt{C_{u}r}\sigma_{1} + \sigma_{\xi}\right)\frac{\sigma_{1}}{\sigma_{r}}\sqrt{\frac{(d_{1} \vee d_{2})\log(d_{1} + d_{2})}{n}} \\ &+ C_{5}\sqrt{C_{l}^{-1}}\left(\sqrt{C_{u}r}\sigma_{1} + \sigma_{\xi}\right)\frac{\sigma_{1}}{\sigma_{r}}\frac{\sqrt{r(d_{1} \vee d_{2})}}{n\varepsilon}\log n\log^{\frac{1}{2}}(\frac{3.75}{\delta}) \\ &+ C_{5}\sqrt{C_{l}^{-1}}\left(\sqrt{C_{u}r}\sigma_{1} + \sigma_{\xi}\right)\frac{r}{n\varepsilon}\log n\log^{\frac{1}{2}}(\frac{3.75}{\delta}), \end{split}$$

for some absolute constant $C_5 > 0$ with probability at least $1 - \exp(-d_1) - \exp(-d_2) - 10^{-20r}$.

A.5 Proof of Corollary 1

The proof of Corollary 1 is obtained by setting the upper bound of $\|\widetilde{M}_0 - M\|_F \leq \sqrt{2r} \|\widetilde{M}_0 - M\|$ given by Theorem 1 smaller than the order of σ_r .

B Proof of Theorem 2

We first present some preliminary results on the KL-divergence and total variation distance between Gaussian distributions. Let $\mathcal{N}(\mu_1, \Sigma_1)$ and $\mathcal{N}(\mu_2, \Sigma_2)$ be two *p*-dimensional multivariate Gaussians, then

$$\operatorname{KL}\left(\mathcal{N}\left(\mu_{1},\Sigma_{1}\right)\left\|\mathcal{N}\left(\mu_{2},\Sigma_{2}\right)\right)\right.$$
$$=\frac{1}{2}\left(\operatorname{Tr}\left(\Sigma_{2}^{-1}\Sigma_{1}-I_{p}\right)+\left(\mu_{2}-\mu_{1}\right)^{\top}\Sigma_{2}^{-1}\left(\mu_{2}-\mu_{1}\right)+\log\left(\frac{\operatorname{det}\Sigma_{2}}{\operatorname{det}\Sigma_{1}}\right)\right)$$

Let $M_* = \sigma H H^{\top}$ and $M'_* = \sigma H' H'^{\top}$ where $H \neq H' \in \mathcal{S}_q^{(d_1+d_2)}$, then

$$\begin{split} & \operatorname{KL}\left(f_{M_*}(\bar{y}_i, X_i, X'_i) \| f_{M'_*}(\bar{y}_i, X_i, X'_i)\right) \\ &= \mathbb{E}_{f_{M_*}(\bar{y}_i, X_i, X'_i)} \left[\log \frac{f_{M_*}(\bar{y}_i, X_i, X'_i)}{f_{M'_*}(\bar{y}_i, X_i, X'_i)} \right] \\ &= \mathbb{E}_{X_i, X'_i} \mathbb{E}_{f_{M_*}(\bar{y}_i | X_i, X'_i)} \left[-\frac{\left(\frac{\bar{y}_i - \left\langle \begin{pmatrix} 0 & M \\ M^\top & 0 \end{pmatrix}, \begin{pmatrix} 0 & X_i \\ X'_i^\top & 0 \end{pmatrix} \right\rangle \right)^2}{4\sigma_{\xi}^2} \right] \\ &+ \mathbb{E}_{X_i, X'_i} \mathbb{E}_{f_{M_*}(\bar{y}_i | X_i, X'_i)} \left[-\frac{\left(\frac{\bar{y}_i - \left\langle \begin{pmatrix} 0 & M' \\ M'^\top & 0 \end{pmatrix}, \begin{pmatrix} 0 & X_i \\ X'_i^\top & 0 \end{pmatrix} \right\rangle \right)^2}{4\sigma_{\xi}^2} \right] \\ &= \mathbb{E}_{X_i, X'_i} \left[\frac{\left\langle M_* - M'_*, \begin{pmatrix} 0 & X_i \\ X'_i^\top & 0 \end{pmatrix} \right\rangle \cdot \left\langle M_* - M'_*, \begin{pmatrix} 0 & X_i \\ X'_i^\top & 0 \end{pmatrix} \right\rangle}{4\sigma_{\xi}^2} \right] \\ &\lesssim \frac{1}{\sigma_{\xi}^2} C_u \| M_* - M'_* \|_F^2 \lesssim \frac{1}{\sigma_{\xi}^2} C_u \sigma^2 r \varepsilon_0^2, \end{split}$$

and further by Pinsker's inequality, we have

TV
$$\left(f_{M_*}(\bar{y}_i, X_i, X'_i), f_{M'_*}(\bar{y}_i, X_i, X'_i)\right) \lesssim \frac{\sqrt{C_u}\sigma}{\sigma_{\xi}} \sqrt{r}\varepsilon_0.$$

For any probability measures $P_{M_*} \neq P_{M'_*} \in \mathcal{P}_{\sigma}$, we have

$$\operatorname{KL}\left(\bigotimes_{i=1}^{n} f_{M_{*}}(\bar{y}_{i}, X_{i}, X_{i}') \|\bigotimes_{i=1}^{n} f_{M_{*}'}(\bar{y}_{i}, X_{i}, X_{i}')\right) \lesssim \frac{n}{\sigma_{\xi}^{2}} C_{u} \sigma^{2} r \varepsilon_{0}^{2}.$$

and

$$\sum_{k \in [n]} \operatorname{TV}\left(f_{M_*}(\bar{y}_i, X_i, X_i'), f_{M'_*}(\bar{y}_i, (X_i^{\perp})_*)\right) \lesssim n \frac{\sqrt{C_u}\sigma}{\sigma_{\xi}} \sqrt{r} \varepsilon_0,$$

The next Lemma 6 states that there exists a sufficiently large subsect of $\mathbb{O}_{d_1+d_2,2r}$ such that the elements in the subsets are well separated.

Lemma 6. For any $r \leq d$, there exists a subset $S_q^{(d)} \subset \mathbb{O}_{d,r}$ with cardinality $\left|S_q^{(d)}\right| \geq 2^{r(d-r)}$ such that for any $U_i \neq U_j \in S_q^{(d)}$,

$$\|U_i U_i^{\top} - U_j U_j^{\top}\|_q \ge \sqrt{\varepsilon_0^2 (1 - \varepsilon_0^2)} \|V_i - V_j\|_q \gtrsim \sqrt{\varepsilon_0^2 (1 - \varepsilon_0^2)} \|V_i V_i^{\top} - V_j V_j^{\top}\|_q \gtrsim \sqrt{\varepsilon_0^2 (1 - \varepsilon_0^2)} r^{1/q}$$

where $\|\cdot\|_q$ denotes the Schatten-q norm and, meanwhile,

$$\|U_i U_i^{\top} - U_j U_j^{\top}\|_{\mathbf{F}} \lesssim \|U_i - U_j\|_{\mathbf{F}} \le \varepsilon_0 \|V_i - V_j\|_{\mathbf{F}} \le \sqrt{2r}\varepsilon_0.$$

By Lemma 6, there exists a subset $S_q^{(d_1+d_2)} \subset \mathbb{O}_{d_1+d_2,2r}$ with cardinality $\left|S_q^{(d_1+d_2)}\right| \ge 2^{2r(d_1+d_2-2r)}$ such that for any $H \neq H' \in S_q^{(d_1+d_2)}$,

$$\|HH^{\top} - H'H'^{\top}\|_q \gtrsim \sqrt{\varepsilon_0^2 (1 - \varepsilon_0^2)} (2r)^{1/q},$$

and meanwhile,

$$\|HH^{\top} - H'H'^{\top}\|_{\mathrm{F}} \lesssim 2\sqrt{r}\varepsilon_0.$$

To invoke Lemma 3, we define the metric $\rho : \mathbb{O}_{d_1+d_2,2r} \times \mathbb{O}_{d_1+d_2,2r} \mapsto \mathbb{R}^+$ as $\rho(H, H') :=$ $||HH^\top - H'H'^\top||_q$ for any $q \in [1, \infty]$ and take $\rho_0 \asymp \tau \varepsilon_0 r^{1/q}$,

$$l_0 = c_0 \frac{n}{\sigma_{\xi}^2} C_u \sigma^2 r \varepsilon_0^2$$
 and $t_0 = c_0 n \frac{\sqrt{C_u} \sigma}{\sigma_{\xi}} \sqrt{r} \varepsilon_0$,

for some small absolute constant $c_0, \tau > 0$. Then, by Lemma 3, for any (ε, δ) -DP estimator \widetilde{H} ,

$$\sup_{P \in \mathcal{P}_{\sigma}} \mathbb{E} \| \widetilde{H} \widetilde{H}^{\top} - H H^{\top} \|_{q} \\ \geqslant \max \left\{ \frac{\tau \varepsilon_{0} r^{1/q}}{2} \left(1 - \frac{c_{0} \frac{n}{\sigma_{\xi}^{2}} C_{u} \sigma^{2} r \varepsilon_{0}^{2} + \log 2}{\log N} \right), \frac{\tau \varepsilon_{0} r^{1/q}}{4} \left(1 \wedge \frac{N - 1}{\exp \left(4 \varepsilon c_{0} n \frac{\sqrt{C_{u} \sigma}}{\sigma_{\xi}} \sqrt{r} \varepsilon_{0} \right)} \right) \right\}.$$

Recall that $N \ge 2^{2r(d_1+d_2)/2}$ if $d_1 + d_2 \ge 4r$. We can take

$$\varepsilon_0 \simeq \frac{\sigma_{\xi}}{\sqrt{C_u}\sigma} \left(\sqrt{\frac{d_1+d_2}{n}} + r^{\frac{1}{2}} \frac{d_1+d_2}{n\varepsilon} \right),$$

to get

$$\sup_{P\in\mathcal{P}_{\sigma}} \mathbb{E}\left\|\widetilde{H}\widetilde{H}^{\top} - HH^{\top}\right\|_{q} \gtrsim \frac{\sigma_{\xi}}{\sqrt{C_{u}\sigma}} \left(r^{1/q}\sqrt{\frac{d_{1}+d_{2}}{n}} + r^{\frac{1}{2}+\frac{1}{q}}\frac{d_{1}+d_{2}}{n\varepsilon}\right) \bigwedge r^{1/q},$$

where the last term is due to a trivial upper of $\|\widetilde{H}\widetilde{H}^{\top} - HH^{\top}\|_q \leq (4r)^{1/q}$. Since σ is already known, it suffices to estimate HH^{\top} differentially privately by the estimator $\widetilde{H}\widetilde{H}^{\top}$, and an estimator $(\widetilde{M})_*$ for the matrix M_* is given by $\sigma \widetilde{H}\widetilde{H}^{\top}$. Therefore,

$$\sup_{P \in \mathcal{P}_{\sigma}} \mathbb{E} \left\| \widetilde{M}_{*} - M_{*} \right\|_{q} \geq \sigma \cdot \sup_{P \in \mathcal{P}_{\sigma}} \mathbb{E} \left\| \widetilde{H} \widetilde{H}^{\top} - H H^{\top} \right\|_{q}.$$

Due to $\mathcal{P}_{\sigma} \subset \mathcal{P}_{\Sigma}$, we have

$$\sup_{P \in \mathcal{P}_{\Sigma}} \mathbb{E} \|\widetilde{M}_{*} - M_{*}\|_{q} \geq \sup_{P \in \mathcal{P}_{\sigma}} \mathbb{E} \|\widetilde{M}_{*} - M_{*}\|_{q} \gtrsim \frac{\sigma_{\xi}}{\sqrt{C_{u}}} \left(r^{1/q} \sqrt{\frac{d_{1} + d_{2}}{n}} + r^{\frac{1}{2} + \frac{1}{q}} \frac{d_{1} + d_{2}}{n\varepsilon} \right) \bigwedge r^{1/q} \sigma.$$

There is a one-to-one mapping between \widetilde{M} and $(\widetilde{M})_*$. Let $\widetilde{M} - M = U_{\Delta} \Sigma_{\Delta} V_{\Delta}^{\top}$, then $\|\widetilde{M} - M\|_q^q = \|\Sigma_{\Delta}\|_q^q$. Note that

$$(\widetilde{M})_* - M_* = \begin{pmatrix} 0 & \widetilde{M} - M \\ \widetilde{M}^\top - M^\top & 0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} U_\Delta & U_\Delta \\ V_\Delta & -V_\Delta \end{pmatrix} \begin{pmatrix} \Sigma_\Delta & 0 \\ 0 & \Sigma_\Delta \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} U_\Delta & U_\Delta \\ V_\Delta & -V_\Delta \end{pmatrix}^\top$$

and

$$\left\| (\widetilde{M})_* - M_* \right\|_q^q = \left\| \left(\begin{array}{cc} \Sigma_A & 0\\ 0 & \Sigma_a \end{array} \right) \right\|_q^q = 2 \left\| \Sigma_\Delta \right\|_q^q = 2 \left\| \widetilde{M} - M \right\|_q^q.$$

Therefore, $\left\| (\widetilde{M})_* - M * \right\|_q = 2^{1/q} \|\widetilde{M} - M\|_q$ and

$$\inf_{\widetilde{M}} \sup_{P \in \mathcal{P}_{\Sigma}} \mathbb{E} \| \widetilde{M} - M \|_q \gtrsim \frac{\sigma_{\xi}}{\sqrt{C_u}} \left(r^{1/q} \sqrt{\frac{d_1 + d_2}{n}} + r^{\frac{1}{2} + \frac{1}{q}} \frac{d_1 + d_2}{n\varepsilon} \right) \bigwedge r^{1/q} \sigma, \tag{S.11}$$

where we use the fact $d_1 + d_2 \lesssim d_1 \lor d_2$ and infimum is taken over all possible (ε, δ) -DP algorithms.

C Proof of Theorem 3

In Appendix C, we aim to prove Theorem 3. The proof is composed of three Parts. In Part C.1, we characterize the sensitivity Δ_l for iterations $l = 1, \dots, l^*$ and bound $\|\mathcal{P}_{\mathbb{T}_l}N_l\|_F$. In Part C.2, take mathematical induction to prove that if the RIP-condition holds and both Δ_l and $\|M_l - M\|_F$ are bounded with high probability, then we also have Δ_{l+1} and $\|M_{l+1} - M\|_F$ are bounded with high probability. In Part C.3, we choose an appropriate l^* as the total number of iterations and give the convergence rate of $\|\widetilde{M}_{l^*} - M\|_F$.

C.1 Part 1

In Part C.1, we focus on upper bounding $\|\mathcal{P}_{\mathbb{T}_l}N_l\|_F$. The first step is to characterize the sensitivity of the *l*-th iteration for $l \in [l^*]$. Let

$$G_l^{(i)} := \frac{1}{n} \sum_{j \neq i} \left(\langle X_j, M_l \rangle - y_j \right) X_j + \frac{1}{n} \left[\langle X'_i, M_l \rangle - y'_i \right] X'_i,$$

which is the gradient of *l*-th iteration obtained by the dataset differes with the original one only by the *i*-th pair of observation. The sensitivity of the gradient descent on the tagent space \mathbb{T}_l is

$$\Delta_{l} := \max_{\substack{\text{neighbouring}(Z,Z')}} \|M_{l} - \eta \mathcal{P}_{\mathbb{T}_{l}}(G_{l}(Z)) - [M_{l} - \eta \mathcal{P}_{\mathbb{T}_{l}}(G_{l}(Z'))]\|_{F}$$
$$= \max_{i \in [n]} \|M_{l} - \eta \mathcal{P}_{\mathbb{T}_{l}}(G_{l}) - \left[M_{l} - \eta \mathcal{P}_{\mathbb{T}_{l}}(G_{l}^{(i)})\right]\|_{F}.$$

By the definition of G_l and $G_l^{(i)}$,

$$\Delta_{l} \leq \frac{\eta}{n} \max_{i \in [n]} \left[\left\| \mathcal{P}_{\mathbb{T}_{l}} \left(\left\langle X_{i}, M_{l} \right\rangle - y_{i} \right) X_{i} \right\|_{F} + \left\| \mathcal{P}_{\mathbb{T}_{l}} \left(\left\langle X_{i}', M_{l} \right\rangle - y_{i}' \right) X_{i}' \right\|_{F} \right],$$

where for all $i \in [n]$ and $l + 1 \in [l^*]$

$$\begin{aligned} \left\| \mathcal{P}_{\mathbb{T}_{l}}\left(\left\langle X_{i}, M_{l} \right\rangle - y_{i} \right) X_{i} \right\|_{F} &\leq \left| \left\langle X_{i}, M_{l} - M \right\rangle - \xi_{i} \right| \left\| \mathcal{P}_{\mathbb{T}_{l}} X_{i} \right\|_{F} \\ &\leq \left(\left| \xi_{i} \right| + \left| \left\langle X_{i}, M_{l} - M \right\rangle \right| \right) \sqrt{2r} \left\| X_{i} \right\|. \end{aligned}$$
(S.12)

Here, the last inequality uses the fact that for any $B \in \mathbb{R}^{d_1 \times d_2}$, the matrix $\mathcal{P}_{\mathbb{T}_l} B$ is of rank at most 2r. Since both ξ_i and $\langle M_l - M, X_i \rangle$ are sub-Gaussians with $\|\xi_i\|_{\psi_2} = \sigma_{\xi}$ and $\|\langle M_l - M, X_i \rangle\|_{\psi_2} \leq \|M_l - M\|_F \sqrt{C_u}$, we turn to Lemma 7 to upper bound $|\xi_i| + |\langle X_i, M_l - M \rangle|$.

Lemma 7 (Vershynin (2018)). For any sub-Gaussian random variable $B \in \mathbb{R}$,

$$\mathbb{P}(|B| \ge t) \le 2 \exp\left(-ct^2/\|B\|_{\psi_2}\right), \quad \forall t \ge 0$$

According to the tail probability of sub-Gaussian random variable stated in Lemma 7, we have with probability at least $1 - n^{-10}$,

$$|\xi_i| + |\langle X_i, M_l - M \rangle| \le C_1 (\sigma_{\xi} + ||M_l - M||_F \sqrt{C_u}) \log^{1/2} n,$$
(S.13)

for some absolute constant $C_1 > 0$. Shen et al. (2023) offeres the following result on $||X_i||$.

Lemma 8 (Shen et al. (2023)). Suppose the vectorization of $X \in \mathbb{R}^{d_1 \times d_2}$ follows mean zero multivariate Gaussian distribution $N(\mathbf{0}, \Lambda)$ where $\Lambda \in \mathbb{R}^{d_1 d_2 \times d_1 d_2}$ satisfies $\lambda_{\max}(\Lambda) \leq C_u$. Then, for some constant c > 0

$$\mathbb{P}\left(\|X\| \ge t + c\sqrt{C_u\left(d_1 + d_2\right)}\right) \le \exp\left(-t^2\right).$$

It implies $|||X|||_{\psi_2} \leq c_1 \sqrt{C_u(d_1+d_2)}$ and $|||X|||_{\psi_1} \leq c_2 \sqrt{C_u(d_1+d_2)}$ for some constants $c_1, c_2 > 0.$

Thus, for some absolute constant $C_2 > 0$, with probability at least $1 - \exp(-10C_u(d_1 + d_2)) n^{-10}$

$$||X_i|| \le C_2 \sqrt{C_u (d_1 + d_2) + \log n}.$$
(S.14)

Combining (S.12), (S.13) and (S.14) and taking maximum over n, for some constant $C_3 > 0$, we have the event

$$\mathcal{E}'_{\Delta_l} := \left\{ \Delta_l \le C_3 \frac{\eta}{n} (\sigma_{\xi} + \|M_l - M\|_F \sqrt{C_u}) \sqrt{C_u r (d_1 + d_2 + \log n) \log n} \right\},$$
(S.15)

happens with probability at least $1 - n^{-9} - \exp(-10C_u(d_1 + d_2)) n^{-9}$. In the event \mathcal{E}'_{Δ_l} stated in (S.15), the sensitivity Δ_l still relies on $||M_l - M||_F$. To get an upper bound irrelevant with l, we take condition on the event

$$\mathcal{E}_l = \left\{ \|M_l - M\|_F \le c_0 \sigma_r \right\},\,$$

and obtain that for some absolute constant $\widetilde{C}_3 > 0$, the event

$$\mathcal{E}_{\Delta_l} := \left\{ \Delta_l \le \widetilde{C}_3 \frac{\eta}{n} (\sigma_{\xi} + \sigma_r \sqrt{C_u}) \sqrt{C_u r (d_1 + d_2 + \log n) \log n} \right\},\tag{S.16}$$

happens with the probability $\mathbb{P}(\mathcal{E}_{\Delta_l}) \geq 1 - n^{-9} - \exp(-10C_u(d_1 + d_2)) n^{-9}$.

In the l+1-th iteration of Algorithm 2, the matrix M_l and operator $\mathcal{P}_{\mathbb{T}_l}$ are known. Moreover, the rank r approximation SVD_r is irrelevant with the data set $Z = \{(X_i, y_i)\}_{i=1}^n$. Thanks to the post-processing property and composition property of differential privacy, we only need to guarantee that $M_l - \eta_l \mathcal{P}_{\mathbb{T}_l}(G_l)$ is $(\varepsilon/l^*, \delta/l^*)$ -DP where the gradient

$$G_l = \frac{1}{n} \sum_{i=1}^n \left(\langle X_i, M_l \rangle - y_i \right) X_i,$$

is the only component depends on the data set Z. Let N_l be a $d_1 \times d_2$ matrix with entries i.i.d. to normal distribution with varicance $\frac{l^{*2}\Delta_l^2}{\varepsilon^2} \log\left(\frac{1.25l^*}{\delta}\right)$. Under the condition that $d_1 + d_2 \gtrsim \log n$ and conditioned on the event $\mathcal{E}_{\Delta_l} \cap \mathcal{E}_l$, we have for some constant $C_4 > 0$

$$\|\mathcal{P}_{\mathbb{T}_l}N_l\|_F \le C_4 \eta l^* \frac{r(d_1+d_2)}{n\varepsilon} (\sigma_{\xi} + \sigma_r \sqrt{C_u}) \sqrt{C_u \log n} \log^{1/2} \left(\frac{1.25l^*}{\delta}\right),$$

with probability at least $1 - \exp(-(d_1 + d_2))$.

C.2 Part 2

In Part C.2, we take mathematical induction to prove that when the events

$$\mathcal{E}_l^* := \mathcal{E}_{\mathrm{RIP}} \cap \mathcal{E}_l \cap \mathcal{E}_{\Delta_l},$$

occurs with high probability, the event \mathcal{E}_{l+1}^* occurs with high probability as well. Here, \mathcal{E}_{RIP} is defined as the event where the RIP condition of $\{X_i\}_{i \in [n]}$ holds, See (S.17). Step 1: \mathcal{E}_0^* is true with high probability.

We first consider the RIP condition. According to Lemma 1, for any $B \in \mathbb{R}^{d_1 \times d_2}$ of rank r, there exist constants $c_1, c_2, c_3 > 0$ and $0 < c_4 < c_5$ such that when $n \ge c_1 r(d_1 + d_2)$, with probability at least $1 - c_2 \exp(-c_3 r(d_1 + d_2))$,

$$(1 - R_r) \|B\|_{\mathrm{F}}^2 \le \frac{1}{n} \sum_{i=1}^n \langle X_i, B \rangle^2 \le (1 + R_r) \|B\|_{\mathrm{F}}^2,$$

where $R_r := (1 - c_4 \sqrt{C_u C_l}) \vee (c_5 \sqrt{C_u C_l} - 1)$. The values of R_{2r} , R_{3r} and R_{4r} are defined similarly. Therefore, under the condition that $n \ge \tilde{c}_1 r(d_1 + d_2)$, for some constants $\tilde{c}_1, \tilde{c}_2, \tilde{c}_3, \tilde{c}_4, \tilde{c}_5 > 0$,

$$\mathcal{E}_{\text{RIP}} := \left\{ R_r \lor R_{2r} \lor R_{3r} \lor R_{4r} \le \left(1 - \widetilde{c}_4 \sqrt{C_u C_l} \right) \lor \left(\widetilde{c}_5 \sqrt{C_u C_l} - 1 \right) \right\}, \quad (S.17)$$

happens with probability at least $1 - \tilde{c}_2 \exp(-\tilde{c}_3 r(d_1 + d_2))$.

As for \mathcal{E}_0 , we refer to Corollary 1, which shows that as the sample size

$$n \ge \widetilde{O}\left((\kappa^4 r^2 + \kappa^2 \kappa_{\xi}^2 r)(d_1 \lor d_2)\right),\,$$

the event \mathcal{E}_0 happens with probability at least $1-(d_1+d_2)^{-10}-n^{-9}-\exp(-d_1)-\exp(-d_2)-10^{-20r}$. Conditioned on \mathcal{E}_0 , plugging l = 0 to the event \mathcal{E}'_{Δ_l} defined in (S.15), we have the event \mathcal{E}_{Δ_l} defined in (S.16) happens with probability at least $1-n^{-9}-\exp(-10C_u(d_1+d_2))n^{-9}$. To this end, we have

$$\mathbb{P}\left(\mathcal{E}_{0}^{*}\right) = \mathbb{P}\left(\mathcal{E}_{\mathrm{RIP}} \cap \mathcal{E}_{0} \cap \mathcal{E}_{\Delta_{0}}\right) \ge 1 - \tilde{c}_{2} \exp\left(-\tilde{c}_{3}r(d_{1}+d_{2})\right)$$
$$- (d_{1}+d_{2})^{-10} - n^{-9} - e^{-d_{1}} - e^{-d_{2}} - 10^{-20r}$$
$$- n^{-9} - \exp\left(-10C_{u}(d_{1}+d_{2})\right)n^{-9}.$$

Step 2: induction. The following analysis is conditioned on the event \mathcal{E}_l^* . Let $\mathcal{X} : \mathbb{R}^{d_1 \times d_2} \to \mathbb{R}^n$ be an operator defined by $\mathcal{X}(B) = (\langle X_1, B \rangle, \cdots, \langle X_n, B \rangle)^\top \in \mathbb{R}^n$, for all $B \in \mathbb{R}^{d_1 \times d_2}$. It is easy to check that the adjoint operator of \mathcal{X} is $\mathcal{X}^* : \mathbb{R}^n \to \mathbb{R}^{d_1 \times d_2}$ which is defined by $\mathcal{X}^*(b) :=$ $\sum_{i=1}^{n} b_i X_i, \text{ for all } b \in \mathbb{R}^n. \text{ Therefore, } \mathcal{X}^* \mathcal{X}(M_l) = \sum_{i=1}^{n} \langle X_i, M_l \rangle X_i \text{ and } \mathcal{X}^*(\xi) = \sum_{i=1}^{n} \xi_i X_i,$ where $\xi := (\xi_1, \cdots, \xi_n) \in \mathbb{R}^n$ and accordingly,

$$P_{\mathbb{T}_l}G_l = \frac{1}{n} \left[\mathcal{P}_{\mathbb{T}_l} \mathcal{X}^* \mathcal{X} \left(M_l - M \right) - \mathcal{P}_{\mathbb{T}_l} \mathcal{X}(\xi) \right].$$

Our first goal is to upper bound

$$\begin{split} \|M_{l} - M - \eta \mathcal{P}_{\mathbb{T}_{l}}(G_{l})\|_{F} &= \|M_{l} - M - \frac{\eta}{n} \mathcal{P}_{\mathbb{T}_{l}} \mathcal{X}^{*} \mathcal{X} (M_{l} - M) - \mathcal{P}_{\mathbb{T}_{l}} \mathcal{X}(\xi)\|_{F} \\ &\leq \left(1 - \frac{\eta}{n}\right) \|M_{l} - M\|_{F} + \frac{\eta}{n} \|(\mathcal{I} - \mathcal{P}_{\mathbb{T}_{l}} \mathcal{X}^{*} \mathcal{X} \mathcal{P}_{\mathbb{T}_{l}}) (M_{l} - M)\|_{F} \\ &\xrightarrow{D_{1}} \\ &+ \underbrace{\frac{\eta}{n} \left\|\mathcal{P}_{\mathbb{T}_{l}} \mathcal{X}^{*} \mathcal{X} \mathcal{P}_{\mathbb{T}_{l}^{\perp}} (M_{l} - M)\right\|_{F}}_{D_{2}} + \underbrace{\frac{\eta}{n} \left\|\mathcal{P}_{\mathbb{T}_{l}} \mathcal{X}^{*} (\xi)\right\|_{F}}_{D_{3}}. \end{split}$$
(S.18)

Lemma 9 characterizes the operators $\mathcal{P}_{\mathbb{T}_l} - \mathcal{P}_{\mathbb{T}_l} \mathcal{X}^* \mathcal{X} \mathcal{P}_{\mathbb{T}_l}$ and $\mathcal{P}_{T_l} \mathcal{X}^* \mathcal{X} \mathcal{P}_{\mathbb{T}_l^{\perp}}$, which are critical to upper bound D_1 and D_2 in (S.18).

Lemma 9 (Wei et al. (2016), Luo and Zhang (2022)). Suppose the event \mathcal{E}_{RIP} happens, then the following conclusions hold

1. $\|\mathcal{P}_{\mathbb{T}_l} - \mathcal{P}_{\mathbb{T}_l}\mathcal{X}^*\mathcal{X}\mathcal{P}_{\mathbb{T}_l}\| \leq R_{2r}.$

2.
$$\|\mathcal{P}_{T_l}\mathcal{X}^*\mathcal{X}\mathcal{P}_{\mathbb{T}_l^{\perp}}(M_l-M)\|_F = R_{4r} \|\mathcal{P}_{\mathbb{T}_l^{\perp}}(M_l-M)\|_F$$

where $\|\mathcal{P}_{\mathbb{T}_{l}^{\perp}}(M_{l}-M)\|_{F} \leq \frac{1}{\sigma_{r}}\|M_{l}-M\|\|M_{l}-M\|_{F}$ according to Wei et al. (2016).

According to Lemma 9, conditioned on the event \mathcal{E}_l^*

$$D_{1} = \frac{\eta}{n} \left\| \mathcal{I} - \mathcal{P}_{\mathbb{T}_{l}} \mathcal{X}^{*} \mathcal{X} \mathcal{P}_{\mathbb{T}_{l}} \left(M_{l} - M \right) \right\|_{F}$$

$$\leq \frac{\eta}{n} \left[\left\| \mathcal{P}_{\mathbb{T}_{l}} - \mathcal{P}_{\mathbb{T}_{l}} \mathcal{X}^{*} \mathcal{X} \mathcal{P}_{\mathbb{T}_{l}} \left(M_{l} - M \right) \right\|_{F} + \left\| \mathcal{P}_{\mathbb{T}_{l}^{\perp}} \left(M_{l} - M \right) \right\|_{F} \right] \leq \frac{\eta}{n} \left(R_{2r} + c_{0} \right) \left\| M_{l} - M \right\|_{F},$$

and $D_2 = \frac{\eta}{n} \| \mathcal{P}_{T_l} \mathcal{X}^* \mathcal{X} \mathcal{P}_{\mathbb{T}_l} (M_l - M) \|_F \leq \frac{\eta}{n} R_{4r} c_0 \| M_l - M \|_F$. To this end, the only term unknown in (S.18) is

$$D_3 = \left\| \mathcal{P}_{\mathbb{T}_l} \mathcal{X}^* \left(\xi \right) \right\|_F = \left\| \mathcal{P}_{\mathbb{T}_l} \sum_{i=1}^n \xi_i X_i \right\|_F \le \sqrt{2r} \left\| \sum_{i=1}^n \xi_i X_i \right\|,$$

where $\|\sum_{i=1}^{n} \xi_i X_i\|$ is the spectral norm of a summation of n i.i.d. mean zero sub-exponential random matrices. We upper bound $\|\sum_{i=1}^{n} \xi_i X_i\|$ by Theorem 4. Let $B_i := \xi_i X_i$ for all $i = 1, \dots, n$, then

$$K := \max_{i \in [n]} \|\|B_i\|\|_{\psi_1} = \max_{i \in [n]} \|\|\xi_i X_i\|\|_{\psi_1} \le \max_{i \in [n]} \|\xi_i\|_{\psi_2} \|\|X_i\|\|_{\psi_2} \le \sqrt{C_u (d_1 + d_2)\sigma_{\xi}^2}.$$

Since $\mathbb{E}\xi_i^4 \leq 3\sigma_{\xi}^4$ and $\mathbb{E}||X_i||^4 \leq 3C_u(d_1+d_2)^2$,

$$\|\mathbb{E}B_i B_i^{\top}\| = \mathbb{E}\xi_i^2 \mathbb{E}\|X_i\|^2 \le \frac{C_u(d_1 + d_2)}{2\sigma_{\xi}^2} \mathbb{E}\xi_i^4 + \frac{\sigma_{\xi}^2}{2C_u(d_1 + d_2)} \mathbb{E}\|X_i\|^4 \le 3\sigma_{\xi}^2 C_u(d_1 + d_2),$$

where the first inequality uses the fact $ab \leq \frac{a^2}{2c} + \frac{cb^2}{2}$. Similarly, $\|\mathbb{E}B_iB_i^{\top}\| \leq 3\sigma_{\xi}^2C_u(d_1+d_2)$.

Therefore,

$$S^{2} := \|\mathbb{E}B_{i}B_{i}^{\top}\| \vee \|\mathbb{E}B_{i}^{\top}B_{i}\| \leq 3\sigma_{\xi}^{2}C_{u}(d_{1}+d_{2}).$$

Applying Thorem 4 with $\alpha = 1$, $K = c_1 \sqrt{C_u (d_1 + d_2) \sigma_{\xi}^2}$ and $S = \sqrt{3\sigma_{\xi}^2 C_u (d_1 + d_2)}$, we have

$$\mathbb{P}\left(\frac{1}{n}\left\|\sum_{i=1}^{n}\xi_{i}X_{i}\right\| \geq C_{5}\sqrt{C_{u}(d_{1}+d_{2})\sigma_{\xi}^{2}}\sqrt{\frac{\log(d_{1}+d_{2})}{n}}\right) \leq (d_{1}+d_{2})^{-10}$$

In conclusion, with probability at least $1 - (d_1 + d_2)^{-10}$

$$D_3 \le C_1 \sqrt{r} \left\| \sum_{i=1}^n \xi_i X_i \right\|_F \le C_1 \sqrt{C_u \sigma_{\xi}^2 n r (d_1 + d_2) \log(d_1 + d_2)},$$

for some constant $C_1 > 0$.

Conditioned on \mathcal{E}_l^* , we plug D_1 , D_2 and D_3 into (S.18) and obtain that for some small constant $0 < c_0 < 1$ and absolute constant $C_2 > 0$

$$\begin{split} \|M_{l} - M - \eta \mathcal{P}_{\mathbb{T}_{l}}(G_{l}) + \mathcal{P}_{\mathbb{T}_{l}}N_{l}\|_{F} &\leq \|M_{l} - M - \eta \mathcal{P}_{\mathbb{T}_{l}}(G_{l})\|_{F} + \|\mathcal{P}_{\mathbb{T}_{l}}N_{l}\|_{F} \\ &\leq (1 - \rho_{0}) \|M_{l} - M\|_{F} + C_{2}\eta\sigma_{\xi}\sqrt{C_{u}}\sqrt{\frac{r(d_{1} + d_{2})}{n}}\log^{1/2}(d_{1} + d_{2}) \\ &+ C_{2}\eta l^{*}\sqrt{C_{u}}(\sigma_{\xi} + \sigma_{r}\sqrt{C_{u}})\frac{r(d_{1} + d_{2})}{n\varepsilon}\log^{1/2}n\log^{1/2}\left(\frac{1.25l^{*}}{\delta}\right), \end{split}$$

with probability at least $1 - (d_1 + d_2)^{-10} - \exp(-(d_1 + d_2))$ where we define

$$\rho_0 := \frac{\eta}{n} \left(1 - R_{2r} - c_0 - R_{4r} c_0 \right).$$

Suppose that $c_0 \leq \frac{1}{R_{2r}(1+R_{4r})} \wedge \frac{1}{8}$, the step size $\eta \leq n$ being a small constant, then we have $0 \leq \rho_0 < 1$. Further, as for some absolute constant $C_3 > 0$, the sample size satisfies

$$n \ge C_3 \max\left\{ \eta^2 \left(\frac{\sigma_{\xi}}{\sigma_r}\right)^2 C_u r(d_1 + d_2) \log(d_1 + d_2), \\ \eta l^* \sqrt{C_u} \left(\frac{\sigma_{\xi} + \sigma_r \sqrt{C_u}}{\sigma_r}\right) \frac{r(d_1 + d_2)}{\varepsilon} \log^{1/2} n \log^{1/2} \left(\frac{1.25l^*}{\delta}\right) \right\},$$

we have for some small constant $0 < \rho_1 < 1$,

$$\|M_l - M - \eta \mathcal{P}_{\mathbb{T}_l}(G_l) + \mathcal{P}_{\mathbb{T}_l}N_l\|_F \le (1 - \rho_1) \|M_l - M\|_F \le (1 - \rho_1)c_0\sigma_r.$$

Applying Lemma 4, we obtain that under the condition $40c_0 < \frac{\rho_1}{2}$,

$$\|M_{l+1} - M\|_{F} \leq (1 + 40c_{0}) \|M_{l} - M - \eta \mathcal{P}_{\mathbb{T}_{l}}(G_{l}) + \mathcal{P}_{\mathbb{T}_{l}}N_{l}\|_{F}$$

$$\leq (1 + 40c_{0}) (1 - \rho_{1})\|M_{l} - M\|_{F}$$

$$\leq \left(1 - \frac{\rho_{1}}{2}\right)\|M_{l} - M\|_{F} \leq \left(1 - \frac{\rho_{1}}{2}\right)c_{0}\sigma_{r}.$$
 (S.19)

In summary, conditioned on the event \mathcal{E}_l^* , the event

$$\mathcal{E}_{l+1} := \{ \|M_{l+1} - M\|_F \le c_0 \sigma_r \} \,,$$

occurs with probability at least $1 - (d_1 + d_2)^{-10} - \exp(-(d_1 + d_2))$. Besides, according to \mathcal{E}'_{Δ_l} defined in (S.15), the event

$$\mathcal{E}_{\Delta_{l+1}} := \left\{ \Delta_{l+1} \le \widetilde{C}_3 \frac{\eta}{n} \left(\sigma_{\xi} + \sigma_r \sqrt{C_u} \right) \sqrt{C_u r \left(d_1 + d_2 + \log n \right) \log n} \right\},\,$$

occurs with probability $1 - n^{-9} - \exp(-10C_u(d_1 + d_2))n^{-9}$. Therefore, conditioned on \mathcal{E}_l^* , the event \mathcal{E}_{l+1}^* happens with probability at least $1 - (d_1 + d_2)^{-10} - \exp(-(d_1 + d_2)) - n^{-9} - \exp(-10C_u(d_1 + d_2))n^{-9}$.

To this end, we has finished the induction and conclude Part C.2 by

$$\mathbb{P}\left(\bigcap_{i=0}^{l} \mathcal{E}_{i}^{*}\right) \geq 1 - \widetilde{c}_{2} \exp\left(-\widetilde{c}_{3}r(d_{1}+d_{2})\right) - (d_{1}+d_{2})^{-10} - n^{-9} - e^{-d_{1}} - e^{-d_{2}} - 10^{-20r} - l\left((d_{1}+d_{2})^{-10} + \exp\left(-(d_{1}+d_{2})\right)\right) - (l+1)\left(n^{-9} + \exp\left(-10C_{u}(d_{1}+d_{2})\right)n^{-9}\right).$$

C.3 Part 3

In Part C.3, we derive the convergence rate of $||M_{l^*} - M||_F$ and choose an appropriate value for l^* . Conditioned on the event $\bigcap_{i=0}^{l^*-1} \mathcal{E}_i^*$, according to (S.19), with probability at least $1 - (d_1 + d_2)^{-10} - \exp(-(d_1 + d_2))$

$$\begin{split} \left\|\widetilde{M}_{l^*} - M\right\|_F &= \|M_{l^*} - M\|_F \\ &\leq (1 - \rho_0)^{l^*} \|M_0 - M\|_F + \left(\sum_{l=0}^{l^*-1} (1 - \rho_0)^{l^*-l-1}\right) C_2 \eta \sigma_\xi \sqrt{C_u} \sqrt{\frac{r(d_1 + d_2)}{n}} \log^{1/2} (d_1 + d_2) \\ &+ \left(\sum_{l=0}^{l^*-1} (1 - \rho_0)^{l^*-l-1}\right) C_2 \eta l^* \sqrt{C_u} (\sigma_\xi + \sigma_r \sqrt{C_u}) \frac{r(d_1 + d_2)}{n\varepsilon} \log^{1/2} n \log^{1/2} \left(\frac{1.25l^*}{\delta}\right). \end{split}$$

Let $\|M_0 - M^*\|_F = c_0^*$ and $l^* := \log(c_0^* n) / \rho_0$, then we have $(1 - \rho_0)^{l^*} \|M_0 - M\|_F \asymp \frac{1}{n}$, indicating that there is little reason to run the algorithm further than $O(\log n)$ iterations.

In conclusion,

$$\begin{split} \left\| \widetilde{M}_{l^*} - M \right\|_F &\leq C_3 \sigma_{\xi} \sqrt{C_u} \sqrt{\frac{r(d_1 + d_2)}{n}} \log^{1/2} (d_1 + d_2) \\ &+ C_3 \sqrt{C_u} (\sigma_{\xi} + \sigma_r \sqrt{C_u}) \frac{r(d_1 + d_2)}{n\varepsilon} \log^{1/2} n \log^{3/2} \left(\frac{1.25l^*}{\delta} \right). \end{split}$$

with probability at least

$$1 - \tilde{c}_2 \exp\left(-\tilde{c}_3 r(d_1 + d_2)\right) - (d_1 + d_2)^{-10} - n^{-9} - e^{-d_1} - e^{-d_2} - 10^{-20r} - \left((d_1 + d_2)^{-10} + \exp\left(-(d_1 + d_2)\right) + n^{-9} + \exp\left(-10C_u(d_1 + d_2)\right)n^{-9}\right) \log n$$

D The lower bound derived by score attack argument

Let $\mathbb{M}_r := \{M \in \mathbb{R}^{d_1 \times d_2} : \operatorname{rank}(M) = r\}$. This section establishes the minimax lower bound of differentially privately estimating the matrix $M \in \bigcup_{k=1}^r \mathbb{M}_k$, within the trace regression model based on an alternative approach, score attack argument Cai et al. (2023).

The score attack argument involves designing a test statistic and establishing the lower bound of the statistic with the help of a prior distribution of the parameters to estimate. It is unclear, however, how to construct a prior distribution for the low-rank matrix M such that the prior complies with the parameter space \mathbb{M}_r and the *score attack* is easy to compute at the same time. Compared to DP-fano's Lemma (See Lemma 3) which requires $\delta \leq e^{-n}$, the score attack argument is valid for a wider range of $\delta \leq n^{1+\gamma}$ where $\gamma > 0$ is a constant. We first define some necessary notations for the elaboration of score attack argument. For any matrix $B, C \in \mathbb{R}^{d_1 \times d_2}$, we denote $\operatorname{supp}(B) := \{(i, j) \in [d_1] \times [d_2] : B_{ij} \neq 0\}$ as the support of B and the matrix C restricted on $\operatorname{supp} B$ is $[C]_{\operatorname{supp}(B)} = \sum_{i=1}^{d_1} \sum_{j=1}^{d_2} C_{ij} e_i e_j^{\top} \mathbb{I}(B_{ij} \neq 0)$ where e_i is the *i*-th canonical basis in \mathbb{R}^{d_1} and e_j is the *j*-th canonical basis in \mathbb{R}^{d_2} .

To apply score attack argument, we relax the problem to deriving minimax lower bounds over $\mathbb{M}_{r,d_1} := \{M \in \mathbb{R}^{d_1 \times d_2} : \operatorname{supp}(M) \subset [d_1] \times [r]\} \subset \bigcup_{k=1}^r \mathbb{M}_k$. The benefit is that there exists a trivial prior of $M \in \mathbb{M}_{r,d_1}$ such that $M_{ij} \stackrel{i.i.d.}{\sim} \mathcal{N}(0,1)$ for $(i,j) \in [d_1] \times [r]$ and $M_{ij} = 0$ otherwise. Similarly, we may consider establish minimax lower bound over $\mathbb{M}_{r,d_2} := \{M \in \mathbb{R}^{d_1 \times d_2} : \operatorname{supp}(M) \subset [r] \times [d_2]\} \subset \bigcup_{k=1}^r \mathbb{M}_k$. For any $M \in \mathbb{M}_{r,d_2}$, there is a trivial prior as well. Let A be a randomized algorithm mapping a dataset Z to a $d_1 \times d_2$ matrix. We define the DP-constrained minimax risk over $\bigcup_{k=1}^r \mathbb{M}_k$ as

$$\operatorname{risk}(\bigcup_{k=1}^{r} \mathbb{M}_{k}) := \inf_{A} \sup_{M \in \mathbb{M}_{r}} \mathbb{E} \|A(Z) - M\|_{F}^{2},$$

where A is taken over all (ϵ, δ) -DP algorithms. Similarly, we define risk (\mathbb{M}_{r,d_1}) and risk (\mathbb{M}_{r,d_2}) . Since $\mathbb{M}_{r,d_1} \subset \bigcup_{k=1}^r \mathbb{M}_k$ and $\mathbb{M}_{r,d_2} \subset \bigcup_{k=1}^r \mathbb{M}_k$, we have

$$\operatorname{risk}(\bigcup_{k=1}^{r} \mathbb{M}_{k}) \ge \operatorname{risk}(\mathbb{M}_{r,d_{1}}) \bigvee \operatorname{risk}(\mathbb{M}_{r,d_{2}}),$$
(S.20)

which indicates that the lower bound of $\operatorname{risk}(\bigcup_{k=1}^{r} \mathbb{M}_{k})$ will be an immediate result once we successfully lower bound $\operatorname{risk}(\mathbb{M}_{r,d_{1}})$ and $\operatorname{risk}(\mathbb{M}_{r,d_{2}})$.

Next, we construct *score attacks* to derive the lower bounds of $risk(\mathbb{M}_{r,d_1})$ and $risk(\mathbb{M}_{r,d_2})$. Let z = (X, y) be the pair of a measurement matrix and its corresponding response variable, drawn independently from (3). The score function is defined by

$$S_M(z) := \nabla_M \log f_M(z) = \nabla_M \log f_M(y|X),$$

and the score attack is defined by

$$\mathcal{A}_{M}^{(1)}(z, A(Z)) := \left\langle [A(Z) - M]_{[d_{1}] \times [r]}, S_{M}(z) \right\rangle,$$

where A is an (ε, δ) -DP algorithm to estimate $M \in \mathbb{M}_r$; z = (X, y) is a piece of datum that we want to test whether it belongs to $Z = \{(X_i, y_i)\}_{i=1}^n$; the quantity $[A(Z) - M]_{[d_1] \times [r]}$ is obtained by restricting $A(Z) - M \in \mathbb{R}^{d_1 \times d_2}$ to the index set $[d_1] \times [r]$. Under some regularity conditions, the score attack $\mathcal{A}_M^{(1)}(z, A(Z))$ will lead to the lower bound of risk (\mathbb{M}_r, d_1) . Similarly, we derive the lower bound of risk (\mathbb{M}_r, d_2) with the help of the attack

$$\mathcal{A}_M^{(2)}(z, A(Z)) := \left\langle \left[A(Z) - M \right]_{[r] \times [d_2]}, S_M(z) \right\rangle.$$

Finally, Theorem 5 establishes the lower bound for estimating the low-rank matrix M. The proof of Theorem 5.

Theorem 5. Consider i.i.d. observations $Z = \{z_1, \dots, z_n\}$ drawn from the trace regression model defined in (1), where $z_i := (X_i, y_i)$ for $i = 1, \dots, n$. We assume that $\{X_i\}_{i \in [n]}$ satisfy the Assumption 1, $r(d_1 \vee d_2) \leq n\varepsilon$, $0 < \varepsilon < 1$ and $\delta \leq n^{-(1+\gamma)}$ for some $\gamma > 0$, then

$$\operatorname{risk}(\bigcup_{k=1}^{r} \mathbb{M}_{k}) = \inf_{A} \sup_{M \in \bigcup_{k=1}^{r} \mathbb{M}_{k}} \mathbb{E} \|A(Z) - M\|_{F}^{2} \gtrsim \underbrace{\sigma_{\xi}^{2} \frac{r(d_{1} \vee d_{2})}{n}}_{a_{1}} + \underbrace{\sigma_{\xi}^{2} \frac{r^{2}(d_{1} \vee d_{2})^{2}}{n^{2}\varepsilon^{2}}}_{a_{2}}.$$
 (S.21)

By Theorem 5, the lower bound of $\operatorname{risk}(\mathbb{M}_r)$ consists of two terms where the first term a_1 accounts for the statistical error and the second term a_2 is the cost of privacy. The proof for a_1 can be found in Rohde and Tsybakov (2011) and the *cost of privacy* is deduced in the following proof.

Proof of Theorem 5. We now start proving Theorem 5 by score attack argument. Throughout the proof, we assume that $Z = \{z_1, \dots, z_n\}$ is an i.i.d. sample drawn from f_M and Z'_i is a neighbouring data set of Z obtained by replacing z_i with an independent copy $z'_i \sim f_M$. Besides, we mainly focus on the case $M \in \mathbb{M}_{r,d_1}$ and states the result for the case $M \in \mathbb{M}_{r,d_2}$ in Remark 1. Let

$$\mathcal{A}_{M}^{(1)}(z, A(Z)) := \left\langle \left[A(Z) - M\right]_{[d_{1}] \times [r]}, S_{M}(z) \right\rangle.$$

We derive the lower bound of $\operatorname{risk}(\mathbb{M}_{r,d_1}) := \inf_A \sup_{M \in \mathbb{M}_{r,d_1}} \mathbb{E} \|A(Z) - M\|_F^2$, in three steps. For ease of notation, we define

$$A'_i := \mathcal{A}_M(z_i, A(Z'_i))$$
 and $A_i := \mathcal{A}_M(z_i, A(Z))$.

Step 1: bounding the summation. The following Lemma 10 bounds $\mathbb{E} |A'_i|$; Lemma 11 develops the upper bound of $\sum_{i \in [n]} \mathbb{E} A_i$ based on $\mathbb{E} |A'_i|$ discussed in Lemma 10 and a tunning parameter T. The proof of Lemma 10 and 11 can be found in Appendix E.7 and E.8.

Lemma 10. For $i \in [n]$, we have $\mathbb{E}A'_i = 0$ and $\mathbb{E}|A'_i| \le \sqrt{\mathbb{E}||A(Z) - M||_F^2} \sqrt{\frac{C_u}{\sigma_{\xi}^2}}$.

Lemma 11. Let A be an (ε, δ) -DP algorithm with $0 < \varepsilon < 1$ and $\delta \ge 0$, under model (1), by choosing $T = \sqrt{2/\sigma_{\xi}^2} r d_1 \sqrt{\log(\frac{1}{\delta})}$, we have

$$\sum_{i \in [n]} \mathbb{E}A_i \le 2n\varepsilon \sqrt{\mathbb{E}} \|A(Z) - M\|_F^2 \sqrt{C_u/\sigma_\xi^2} + 4\sqrt{2\delta r d_1} \sqrt{\log(1/\delta)/\sigma_\xi^2}.$$
 (S.22)

Step 2: lower bounding the summation. Under some regularity conditions, the following Lemma 12 characterize the quantity $\sum_{i \in [n]} A_i$ as a summation of functions of M. Lemma 13 lower bounds the summation of functions by assigning an appropriate prior distribution π to M. The proof of Lemma 12 and 13 can be found in Appendix E.8.

Lemma 12. If for every $(i, j) \in [d_1] \times [r]$, $\log f_M(Z)$ is continuously differentiable with respect to M_{ij} and $\left|\frac{\partial}{\partial M_{ij}}\log f_M(Z)\right| < h_{ij}(Z)$ such that $\mathbb{E}|h_{ij}(Z)A(M)_{ij}| < \infty$, we have

$$\sum_{i \in [n]} \mathbb{E} \mathcal{A}_M^1(z_i, A(Z)) = \sum_{(i,j) \in [d_1] \times [r]} \frac{\partial}{\partial M_{ij}} \mathbb{E} A(Z)_{ij}.$$

Lemma 12 has its general form stated in Theorem 2.1, Cai et al. (2023). Let g_{ij} be a function defined by $g_{ij}(M) := (\mathbb{E}_{Z|M}A(Z))_{ij}$ for all $(i, j) \in [d_1] \times [r]$, then

$$\sum_{(i,j)\in[d_1]\times[r]}\frac{\partial}{\partial M_{ij}}\mathbb{E}A(Z)_{ij} = \mathbb{E}_{\pi}\left(\sum_{(i,j)\in[d_1]\times[r]}\frac{\partial}{\partial M_{ij}}g_{ij}\right).$$

Lemma 13 lower bounds this quantity by assigning the prior distribution π to M such that $M_{ij} \sim \mathcal{N}(0,1)$ for all $(i,j) \in [d_1] \times [r]$ and otherwise, $M_{ij} = 0$.

Lemma 13. Let $M \in \mathbb{M}_{r,d_1}$ be distributed according to a density π whose marginal densities are $\{\pi_{ij}\}$ for $i = 1, \dots, d_1$ and $j = 1, \dots, d_2$ such that $\pi_{ij} \sim \mathcal{N}(0, 1)$ for all $(i, j) \in [d_1] \times [r]$, and otherwise, π_{ij} be the density function such that $\mathbb{P}(M_{ij} = 0) = 1$. Then,

$$\mathbb{E}_{\pi}\left(\sum_{(i,j)\in[d_1]\times[r]}\frac{\partial}{\partial M_{ij}}g_{ij}\right) \geq \sum_{(i,j)\in[d_1]\times[r]}\mathbb{E}_{\pi_{ij}}M_{ij}^2 - \sqrt{C}\sqrt{\mathbb{E}_{\pi_{ij}}M_{ij}^2} = rd_1 - \sqrt{Crd_1}\gtrsim rd_1.$$

Combining Lemma 12 and 13, we obtain

$$\sum_{i \in [n]} \mathbb{E}A_i = \sum_{i \in [n]} \mathbb{E}\mathcal{A}_M(z_i, A(Z)) \gtrsim rd_1.$$
(S.23)

Step 3: combining the upper and lower bounds. Combining the lower bound (S.23) of $\sum_{i \in [n]} \mathbb{E}A_i$ and the upper bound (S.22) of $\sum_{i \in [n]} \mathbb{E}A_i$, we have

$$2n\varepsilon\sqrt{\mathbb{E}_{\pi}\mathbb{E}_{Z|M}}\|A(Z) - M\|_{F}^{2}\sqrt{C_{u}/\sigma_{\xi}^{2}} \gtrsim rd_{1} - 4\sqrt{2}\delta rd_{1}\sqrt{\log(1/\delta)/\sigma_{\xi}^{2}}.$$

Under the assumption that $\delta < n^{-(1+\gamma)}$ for some $\gamma > 0$, we have $rd_1 - 4\sqrt{2}\delta rd_1\sqrt{\log(1/\delta)/\sigma_{\xi}^2} \gtrsim rd_1$, and therefore $\mathbb{E}_{\pi}\mathbb{E}_{Z|M} \|A(Z) - M\|_F^2 \gtrsim \sigma_{\xi}^2 \cdot \frac{r^2d_1^2}{n^2\varepsilon^2}$. Since the sup-risk is greater than the Bayesian risk,

$$\sup_{M \in \mathbb{M}_{r,d_1}} \mathbb{E}_{Z|M} \left\| A(Z) - M \right\|_F^2 \gtrsim \sigma_{\xi}^2 \cdot \frac{r^2 d_1^2}{n^2 \varepsilon^2}$$

Furthermore, due to $\mathbb{M}_{r,d_1} \subset \bigcup_{k=1}^r \mathbb{M}_k$, we have $\sup_{M \in \bigcup_{k=1}^r \mathbb{M}_k} \mathbb{E}_{Z|M} \|A(Z) - M\|_F^2 \gtrsim \sigma_{\xi}^2 \cdot \frac{r^2 d_1^2}{n^2 \varepsilon^2}$ and

$$\inf_{A} \sup_{M \in \bigcup_{k=1}^{r} \mathbb{M}_{k}} \mathbb{E}_{Z|M} \|A(Z) - M\|_{F}^{2} \gtrsim \sigma_{\xi}^{2} \cdot \frac{r^{2}d_{1}^{2}}{n^{2}\varepsilon^{2}}$$

where A is an (ε, δ) -DP algorithm that satisfies $\mathbb{E}_{Z|M} \|A(Z) - M\|_F^2 \lesssim 1$. This conclusion extends to all differentially private A if we assume that $rd_1 \lesssim n\varepsilon$ such that $\frac{r^2d_1^2}{n^2\varepsilon^2} \lesssim 1$.

Remark 1. Lemma 11 and 13 are also applicable to the case where the parameter space is M_{r,d_2} . For $M \in \mathbb{M}_{r,d_2}$, Lemma 11 implies that

$$\sum_{i \in [n]} \mathbb{E}A_i \le 2n\varepsilon\sqrt{\mathbb{E}\|A(Z) - M\|_F^2}\sqrt{C_u/\sigma_\xi^2} + 4\sqrt{2}\delta r d_2\sqrt{\log(1/\delta)/\sigma_\xi^2};$$

and Lemma 13 results in $\mathbb{E}\left(\sum_{(i,j)\in [d_1]\times [r]}\frac{\partial}{\partial M_{ij}}g_{ij}\right)\gtrsim rd_2$. Therefore, as $\delta < n^{-(1+\gamma)}$ for some $\gamma > 0$, the minimax lower bound

$$\inf_{A} \sup_{M \in \bigcup_{k=1}^{r} \mathbb{M}_{k}} \mathbb{E}_{Z|M} \|A(Z) - M\|_{F}^{2} \gtrsim \sigma_{\xi}^{2} \cdot \frac{r^{2} d_{2}^{2}}{n^{2} \varepsilon^{2}},$$

where A is an (ε, δ) -DP algorithm that satisfies $\mathbb{E}_{Z|M} \|A(Z) - M\|_F^2 \lesssim 1$. Similarly, this conclusion extends to all differentially private A if we assume that $rd_2 \lesssim n\varepsilon$ such that $\frac{r^2d_2^2}{n^2\varepsilon^2} \lesssim 1$.

E Proofs of Technical Lemmas

E.1 Proof of Lemma 1

Lemma 1 is a consequence of Proposition 10.4, Vershynin (2015) and Lemma 1, Chen et al. (2019) by setting $c_l^2 = C_l$ and $c_u^2 = C_u$ and $\tau^2 \approx \frac{\sqrt{C_u}}{\sqrt{C_l}}$. See the definiton of c_l^2 and c_u^2 in Lemma 1, Chen et al. (2019).

E.2 Proof of Lemma 2

The proof of Lemma 2 involves applying the symmetric dilation trick to Theorem 1, Xia (2021).

Lemma 14 (Theorem 1, Xia (2021)). Let $B \in d \times d$ be a rank-r symmetric matrix with eigendecomposition of the form $B = \Theta \Lambda \Theta^{\top}$ where $\Theta \in \mathbb{O}_{d,r}$ and the diagonal matrix $\Lambda = \{\lambda_1, \dots, \lambda_r\}$ has the eigenvalues of B arranging in the non-increasing order. Let $\widehat{B} = B + \Delta_B$ be another $d \times d$ symmetric matrix and leading r eigen vector of \widehat{B} is given by $\widehat{\Theta} \in \mathbb{O}_{r,d}$. Then,

$$\widehat{\Theta}\widehat{\Theta}^{\top} - \Theta\Theta^{\top} = \sum_{k \ge 1} \mathcal{S}_{B,k}(\Delta_B),$$

where the k-th order term $\mathcal{S}_{M_*,k}(\Delta)$ is a summation of $\binom{2k}{k}$ terms defined by

$$\mathcal{S}_{B,k}(\Delta_B) = \sum_{\mathbf{s}:s_1 + \ldots + s_{k+1} = k} (-1)^{1+\tau(\mathbf{s})} \cdot Q^{-s_1} \Delta_B Q^{-s_2} \ldots \Delta_B Q^{-s_{k+1}}$$

where $\mathbf{s} = (s_1, \ldots, s_{k+1})$ contains non-negative indices and $\tau(\mathbf{s}) = \sum_{j=1}^{k+1} \mathbb{I}(s_j > 0)$.

Lemma 14 provides an explicit representation formula for the spectral projector $\widehat{\Theta}\widehat{\Theta}^{\top}$ given that *B* is symmetric and of rank-*r*. Since we are interested in the asymmetric rank-*r* matrix $M = U\Sigma V^{\top} \in \mathbb{R}^{d_1 \times d_2} \in \mathbb{R}^{d_1 \times d_2}$, we apply the symmetric dilation trick to *M* and obtain the rank-2*r* symmetric matrix M_* has eigendecomposition of the form

$$M^* = U_{M^*} \Sigma_{M^*} U_{M^*}^{\top} = \frac{1}{\sqrt{2}} \begin{pmatrix} U & U \\ V & -V \end{pmatrix} \begin{pmatrix} \Sigma & 0 \\ 0 & -\Sigma \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} U & U \\ V & -V \end{pmatrix}^{\top}.$$

The proof is finished by applying Lemma 14 with $B = M_*$, $\hat{B} = M_* + \Delta_*$, $d = d_1 + d_2$, the rank be 2r and

$$\Theta = \frac{1}{\sqrt{2}} \begin{pmatrix} U & U \\ V & -V \end{pmatrix} \quad \text{and} \quad \widehat{\Theta} = \frac{1}{\sqrt{2}} \begin{pmatrix} \widehat{U} & \widehat{U} \\ \widehat{V} & -\widehat{V} \end{pmatrix}.$$

E.3 Proof of Lemma 3 and 4

See the proof of Lemma 3 in Acharya et al. (2021) and Cai et al. (2024). See the proof of 4 in Shen et al. (2023).

E.4 Proof of Lemma 5

$$\max_{i \in [n]} \left\| \Delta - \Delta^{(i)} \right\| \leq \frac{1}{n} \left\| \operatorname{mat}(\Lambda_i^{-1} \operatorname{vec}(X_i)) \left(\langle X_i, M \rangle + \xi_i \right) \right\| \\ + \max_{i \in [n]} \frac{1}{n} \left\| \operatorname{mat}((\Lambda_i')^{-1} \operatorname{vec}(X_i')) \left(\langle X_i', M \rangle + \xi_i \right) \right\|,$$

where $\|\xi_i\|_{\psi_2} = \sigma_{\xi}$ and $\langle X_i, M \rangle \sim N\left(0, \operatorname{vec}\left(M\right)^\top \Lambda_i \operatorname{vec}\left(M\right)\right), \quad \Lambda_i^{-1} \operatorname{vec}\left(X_i\right) \sim N\left(0, \Lambda_i^{-1}\right).$

$$\begin{aligned} \left\| \left\| \xi_{i} + \langle X_{i}, M \rangle \operatorname{mat} \left(\Lambda_{i}^{-1} \operatorname{vec} \left(X_{i} \right) \right) - M \right\| \right\|_{\Psi_{1}} \\ &\leq \left\| \xi_{i} + \langle X_{i}, M \rangle \right\|_{\Psi_{2}} \left\| \left\| \operatorname{mat} \left(\Lambda_{i}^{-1} \operatorname{vec} \left(X_{i} \right) \right) \right\| \right\|_{\Psi_{2}} \leq c_{0} \sqrt{C_{l}^{-1}} \left(\sigma_{\xi} + \sqrt{C_{u}} \sqrt{r} \sigma_{1} \right), \end{aligned}$$

for some absolute constant $c_0 > 0$. Therefore, for some absolute constant $C_3 > 0$, with probability at least $1 - n^{-10}$,

$$\left\| \operatorname{mat}(\Lambda_i^{-1}\operatorname{vec}(X_i))\left(\langle X_i, M \rangle + \xi_i\right) \right\| \le C_3 \sqrt{C_l^{-1}} \left(\sigma_{\xi} + \sqrt{C_u} \sqrt{r} \sigma_1\right) \log n.$$

Taking maximum over n, with probability at least $1 - n^{-9}$

$$\max_{i \in [n]} \left\| \Delta - \Delta^{(i)} \right\| \le C_3 \cdot n^{-1} \sqrt{C_l^{-1}} \left(\sqrt{C_u r} \sigma_1 + \sigma_\xi \right) \log n.$$

In (S.1), we have already shown that for some absolute constant $C_1 > 0$,

$$\|\Delta\| = \|\widehat{L} - M\| \le C_1 \sqrt{C_l^{-1}} \left(\sigma_{\xi} + \sqrt{C_u} \sqrt{r} \sigma_1\right) \sqrt{\frac{(d_1 \vee d_2) \log(d_1 + d_2)}{n}},\tag{S.24}$$

with probability at least $1 - (d_1 + d_2)^{-10}$. Note that for all $i \in [n]$, $\|\Delta^{(i)}\| = \|\Delta - (\Delta - \Delta^{(i)})\| \le \|\Delta\| + \|\Delta - \Delta^{(i)}\|$, and thus

$$\|\Delta\| + \max_{i \in [n]} \|\Delta^{(i)}\| \le 2\|\Delta\| + \max_{i \in [n]} \|\Delta - \Delta^{(i)}\|.$$

As long as the sample size $n \ge \frac{\log^2 n}{(d_1 \lor d_2) \log(d_1 + d_2)}$, there exists an absolute constant $C_0 > 0$ such that

$$\|\Delta\| + \max_{i \in [n]} \|\Delta^{(i)}\| \le C_0 \sqrt{C_l^{-1}} \left(\sigma_{\xi} + \sqrt{C_u} \sqrt{r} \sigma_1\right) \sqrt{\frac{(d_1 \lor d_2) \log(d_1 + d_2)}{n}}$$

with probability at least $1 - (d_1 + d_2)^{-10} - n^{-9}$.

E.5 Proof of Lemma 6

By (Pajor, 1998, Proposition 8) and (Koltchinskii and Xia, 2015, Lemma 5), for any $q \in [1, \infty]$, there exists an absolute constant c' > 0 and a subset $S_q^{(d-r)} \subset \mathbb{O}_{d-r,r}$ such that for any $V_i \neq V_j \in S_q^{(d-r)}$, $\|V_i V_i^\top - V_j V_j^\top\|_q \ge c' r^{1/q}$, and the cardinality of $S_q^{(d-r)}$ is at least $2^{r(d-r)}$. Here, $\|\cdot\|_q$ denotes the Schatten-q norm of a matrix. In particular, spectral norm is Schatten- ∞ norm, Frobenius norm is Schatten-2 norm, and nuclear norm is Schatten-1 norm. Let $\varepsilon_0 > 0$ be a small number to be decided later. Now, for each $V \in S_q^{(d-r)}$, we define

$$U = \left(\begin{array}{c} \sqrt{1 - \varepsilon_0^2} I_r \\ \sqrt{\varepsilon_0^2} V \end{array}\right)$$

such that $U \in \mathbb{R}^{d \times r}$ and $U^{\top}U = I_r$. This means that, for any $V \in \mathcal{S}_q^{(d-r)}$, we can construct a $U \in \mathbb{O}_{d,r}$. This defines a subset $\mathcal{S}_q^{(d)} \subset \mathbb{O}_{d,r}$ with $\operatorname{Card}(\mathcal{S}_q^{(d)}) \geq 2^{r(d-r)}$ such that for any $U_i \neq U_j \in \mathcal{S}_q^{(d)}$,

$$\|U_i U_i^\top - U_j U_j^\top\|_q \ge \sqrt{\varepsilon_0^2 (1 - \varepsilon_0^2)} \|V_i - V_j\|_q \gtrsim \sqrt{\varepsilon_0^2 (1 - \varepsilon_0^2)} \|V_i V_i^\top - V_j V_j^\top\|_q \gtrsim \sqrt{\varepsilon_0^2 (1 - \varepsilon_0^2)} r^{1/q}$$

and, meanwhile,

$$\|U_i U_i^{\top} - U_j U_j^{\top}\|_{\mathbf{F}} \lesssim \|U_i - U_j\|_{\mathbf{F}} \le \varepsilon_0 \|V_i - V_j\|_{\mathbf{F}} \le \sqrt{2r\varepsilon_0}$$

E.6 Proof of Lemma 7, 8 and 9

See the proof of Lemma 7 in Vershynin (2018), Lemma 8 in Shen et al. (2023) and Lemma 9 in Wei et al. (2016) and Luo and Zhang (2022).

E.7 Proof of Lemma 10

Since Z'_i is independent of z_i and $\mathbb{E}\mathcal{S}_M(z_i) = \mathbb{E}\nabla_M f_M(y_i|X_i) = 0$,

$$\mathbb{E}\mathcal{A}_{M}^{1}(z_{i},A(Z_{i}')) = \mathbb{E}\left\langle \left[A(Z_{i}')-M\right]_{[d_{1}]\times[r]},\mathcal{S}_{M}(z_{i})\right\rangle = \left\langle \mathbb{E}\left[A(Z_{i}')-M\right]_{[d_{1}]\times[r]},\mathbb{E}\mathcal{S}_{M}(z_{i})\right\rangle = 0,$$

As for $\mathbb{E}A_i = \mathbb{E}\mathcal{A}^1_M(z_i, A(Z))$, we apply Jensen's inequality and have

$$\mathbb{E} \left| \mathcal{A}_{M}^{1}(z_{i}, A(Z_{i}')) \right| \leq \sqrt{\mathbb{E} \left| \mathcal{A}_{M}^{1}(z_{i}, A(Z_{i}')) \right|^{2}} \\
\leq \sqrt{\mathbb{E} \operatorname{vec} \left(\left[A(Z_{i}') - M \right]_{[d_{1}] \times [r]} \right)^{\top} \operatorname{vec}(\mathcal{S}_{M}(z_{i})) \operatorname{vec}(\mathcal{S}_{M}(z_{i}))^{\top} \operatorname{vec} \left(\left[A(Z_{i}') - M \right]_{[d_{1}] \times [r]} \right)} \\
\leq \sqrt{\left\| \mathbb{E} \operatorname{vec}(\mathcal{S}_{M}(z_{i})) \operatorname{vec}(\mathcal{S}_{M}(z_{i}))^{\top} \right\|} \cdot \sqrt{\mathbb{E} \left\| \left[A(Z_{i}') - M \right]_{[d_{1}] \times [r]} \right\|_{F}^{2}}, \quad (S.25)$$

where the second line is due to $\langle B, C \rangle = \operatorname{vec}(B)^{\top} \operatorname{vec}(C)$ and the last inequality is because Z'_i is independent of z_i . By the definition of $\mathcal{S}_M(z_i) = \frac{1}{\sigma_{\xi}^2}(y_i - \langle X_i, M \rangle)X_i$ and the independence between ξ_i and X_i ,

$$\left\|\mathbb{E}\operatorname{vec}(\mathcal{S}_{M}(z_{i}))\operatorname{vec}(\mathcal{S}_{M}(z_{i}))^{\top}\right\| = \left\|\mathbb{E}\left(\frac{y_{i} - \langle X_{i}, M \rangle}{\sigma_{\xi}^{2}}\right)^{2} \mathbb{E}\operatorname{vec}\left(X_{i}\right)\operatorname{vec}\left(X_{i}\right)^{\top}\right\| = \frac{1}{\sigma_{\xi}^{2}}\left\|\Lambda_{i}\right\| \leq \frac{C_{u}}{\sigma_{\xi}^{2}}.$$

Plugging the upper bound of $\|\mathbb{E}\operatorname{vec}(\mathcal{S}_M(z_i))\operatorname{vec}(\mathcal{S}_M(z_i))^{\top}\|$ into (S.25),

$$\mathbb{E}\left|\mathcal{A}_{M}^{1}(z_{i}, A(Z_{i}'))\right| \leq \sqrt{\mathbb{E}\left\|A(Z) - M\right\|_{F}^{2}} \sqrt{\frac{C_{u}}{\sigma_{\xi}^{2}}},$$

E.8 Proof of Lemma 11, Proof of Lemma 12, Lemma 13

Lemma 11 is a trivial consequence by setting $T = \sqrt{2/\sigma_{\xi}^2} r d_1 \sqrt{\log(\frac{1}{\delta})}$ to Proposition 2.1, Cai et al. (2023). Lemma 12 is a trivial consequence of Theorem 2.1, Cai et al. (2023) along with the definition of $\mathcal{A}_M^1(z_i, A(Z))$. Lemma 13 is a trivial consequence of Proposition 2.2, Cai et al. (2023) by taking $M_{ij} \sim \mathcal{N}(0, 1)$ for $(i, j) \in [d_1] \times [r]$ and $M_{ij} = 0$ otherwise.

E.9 Proof of Lemma 14

See the proof of Lemma 14 in Xia (2021).

F Weak Differential privacy

This section proposes a weaker definition than differential privacy such that the sensitivities are free of $\{X_i\}_{i \in [n]}$.

Definition 2 (weak (ε, δ) -differential privacy). Let Z be a given data set and Z' be a weak neighbouring data set of Z, i.e., Z and Z' differs by at most one pair of observations $z \in Z$ and $z' \in Z'$ sharing the same measurement X. The algorithm A that maps Z into $\mathbb{R}^{d_1 \times d_2}$ is weak (ε, δ) -differentially private over the dataset Z if

$$\mathbb{P}(A(Z) \in \mathcal{Q}) \le e^{\varepsilon} \mathbb{P}(A(Z') \in \mathcal{Q}) + \delta,$$
(S.26)

for all weak neighbouring data set Z, Z' and all subset $\mathcal{Q} \subset \mathbb{R}^{d_1 \times d_2}$.

Compared to the standard (ε, δ) -DP, weak (ε, δ) -differential privacy is a less powerful constraint. Definition 2 only requires the algorithm A to preserve the property (S.26) over weak neighbouring datasets, i.e., datasets that differs by at most one pair of observations sharing the same measurement X. As we consider a pair of observations z = (X, y) and z' = (X, y') under the model (1), where $y = \langle X, M \rangle + \xi$ and $y' = \langle X, M \rangle + \xi'$, the difference $y - y' = \xi - \xi'$ is free of the measurement X.

Next, we list the Theorem 6, Corollary 2 and Theorem 7 as the analogues of Theorem 1, Corollary 1 and Theorem 3. All proofs for this section are deferred to the end of this section.

Theorem 6 (Weak DP and utility guarantees of the initialization \widetilde{M}_0). Consider i.i.d. observations $Z = \{z_1, \dots, z_n\}$ drawn from the trace regression model stated in (1) where $z_i := (X_i, y_i)$ for $i = 1, \dots, n$. Let the true low-rank regression coefficients matrix being $M \in \mathbb{M}_r$. Suppose that $\{X_i\}_{i \in [n]}$ satisfy the Assumption 1. Under the mild condition $n \geq \frac{\sigma_{\xi}}{\sigma_{\xi} + \sqrt{C_u r \sigma_1}}$, there exists absolute constants $C_1, C_2, C_3 > 0$ such that as the sample size $n \geq n_0$, the sensitivity for leading r left and right singular vectors takes the value

$$\Delta_{weak}^{(1)} := \max_{i \in [n]} \left(\| \widehat{U}\widehat{U}^{\top} - \widehat{U}^{(i)}\widehat{U}^{(i)\top} \|_F \vee \| \widehat{V}\widehat{V}^{\top} - \widehat{V}^{(i)}\widehat{V}^{(i)\top} \|_F \right) = C_2 \sqrt{C_l^{-1}} \frac{\sigma_{\xi}}{\sigma_r} \frac{\sqrt{r}}{n} \log n;$$

the sensitivity for the r singular values takes the value

$$\Delta_{weak}^{(2)} := \max_{i \in [n]} \left\| \widetilde{U}^{\top} \left(\widehat{L} - \widehat{L}^{(i)} \right) \widetilde{V} \right\|_F = C_2 \sqrt{C_l^{-1}} \sigma_{\xi} \frac{\sqrt{r}}{n} \log n,$$

and Algorithm 1 is weak (ε, δ) -differentially private. Moreover,

$$\|\widetilde{M}_{0}^{weak} - M\| \bigvee \left(\|\widetilde{M}_{0}^{weak} - M\|_{F} / \sqrt{2r} \right) \leq e_{1} + \underbrace{C_{3}\sqrt{C_{l}^{-1}}\sigma_{\xi} \left(\frac{\sigma_{1}}{\sigma_{r}} \frac{\sqrt{r(d_{1} \vee d_{2})}}{n\varepsilon} + \frac{r}{n\varepsilon} \right) \log n \log^{\frac{1}{2}}(\frac{3.75}{\delta})}_{e_{2}^{weak}}$$

with probability at least $1 - (d_1 + d_2)^{-10} - n^{-9} - \exp(-d_1) - \exp(-d_2) - 10^{-20r}$.

Theorem 6 requires the same sample size condition $n \ge n_0$ as Theorem 1, however, the sensitivities $\Delta_{weak}^{(1)}$ and $\Delta_{weak}^{(2)}$ derived under weak DP, differs with their DP counterpart $\Delta^{(1)}$ and $\Delta^{(2)}$ by the factor $\sqrt{C_u r \sigma_1}$. This leads to a smaller cost of privacy e_2^{weak} than the cost of privacy e_2 we obtained under stronger standard DP-constraints, as presented in Theorem 1. **Corollary 2.** Under the conditions stated in Theorem 6, as the sample size is sufficiently large such that for some absolute constant $c_2 > 0$,

$$n \ge C_1 \max\left\{n_1, \underbrace{\sqrt{C_l^{-1}}\left(\frac{\sigma_{\xi}}{\sigma_r}\right)\left(\kappa r \sqrt{d_1 \vee d_2} + r^{\frac{3}{2}}\right) \log n \frac{\log^{\frac{1}{2}}\left(\frac{3.75}{\delta}\right)}{\varepsilon}}_{n_2^{weak}}\right\},$$

we have for some small constant $0 < c_0 < 1$, $\|\widetilde{M}_0 - M\|_F \leq \sqrt{2r} \|\widetilde{M}_0 - M\| \leq c_0 \sigma_r$.

Compared with Corollary 1, Corollary 2 requires smaller sample size as $n_2^{weak} \leq n_2$.

Theorem 7. Consider i.i.d. observations $Z = \{z_1, \dots, z_n\}$ drawn from the trace regression model stated in (1) where the true low-rank regression coefficients matrix being $M \in \mathbb{M}_r$. Here, $z_i := (X_i, y_i)$ for $i = 1, \dots, n$ and we assume that $\{X_i\}_{i \in [n]}$ satisfy the Assumption 1 and $(d_1 + d_2) > \log n$. Suppose the weak (ε, δ) -DP initialization satisfies 2, then Algorithm 2 is weak $(2\varepsilon, 2\delta)$ -differentially private with the sensitivities

$$\Delta_l = C_3 \frac{\eta}{n} \sigma_{\xi} \sqrt{C_u r (d_1 + d_2) \log n},$$

for some absolute constant $C_3 > 0$. Moreover, as the sample size

$$n \ge c_4 \max\left\{n_3, n_4, \underbrace{\eta \sqrt{C_u} \kappa_{\xi} r(d_1 + d_2) \log^{3/2}(n) \frac{\log^{1/2} \left(\frac{1.25 \log(n)}{\delta}\right)}{\varepsilon}}_{n_5^{weak}}\right\},$$

for some small constant $0 < c_4 < 1$, number of iteration $l^* = O(\log n)$, and the step size $0 < \eta < 1$, we have the output of Algorithm 2 satisfies

$$\left\|\widetilde{M}_{l^*} - M\right\|_F \le u_1 + \underbrace{C_4 \sqrt{C_u} \sigma_\xi \frac{r(d_1 + d_2)}{n\varepsilon} \log^{3/2} n \log^{1/2} \left(\frac{1.25 \log(n)}{\delta}\right)}_{u_0^{weak}}.$$

with probability at least

$$1 - \tilde{c}_2 \exp\left(-\tilde{c}_3 r(d_1 + d_2)\right) - (d_1 + d_2)^{-10} - n^{-9} - e^{-d_1} - e^{-d_2} - 10^{-20r} - \left((d_1 + d_2)^{-10} + \exp\left(-(d_1 + d_2)\right) + n^{-9} + \exp\left(-10C_u(d_1 + d_2)\right)n^{-9}\right) \log n^{-9}$$

Theorem 7 shows that as the sample size $n \gtrsim \widetilde{O}\left(\left(\kappa_{\xi}^2 \lor \kappa_{\xi}\right) r(d_1 \lor d_2)\right)$, the estimator \widetilde{M}_{l^*} given by Algorithm 2 attains the optimal convergence rate $\widetilde{O}_p\left(\sigma_{\xi}\sqrt{\frac{r(d_1+d_2)}{n}} + \sigma_{\xi}\frac{r(d_1+d_2)}{n\varepsilon}\right)$, in the sense of weak differential privacy.

The proofs of Theorem 6, 7 and Corollary 2 will be a trivial concequence of replacing the first part of Lemma 5 by the following Lemma 15

Lemma 15. Under model (1), Assumption 1, and the condition $n \ge \frac{\sigma_{\xi}}{\sigma_{\xi} + \sqrt{C_u r \sigma_1}}$, there exists some absolute constant $C_0, C_1 > 0$ such that the event

$$\mathcal{E}_{*} := \left\{ \max_{i \in [n]} \left\| \Delta - \Delta^{(i)} \right\| \le C_{0} \cdot n^{-1} \sqrt{C_{l}^{-1}} \sigma_{\xi} \log n \right\}$$
$$\bigcap \left\{ \left\| \Delta \right\| + \max_{i \in [n]} \left\| \Delta^{(i)} \right\| \le C_{0} \sqrt{C_{l}^{-1}} \left(\sigma_{\xi} + \sqrt{C_{u}} \sqrt{r} \sigma_{1} \right) \sqrt{\frac{(d_{1} \lor d_{2}) \log(d_{1} + d_{2})}{n}} \right\},$$

holds with probability at least $1 - (d_1 + d_2)^{-10} - n^{-9}$.

Proof of Lemma 15. We only need to focus on $\max_{i \in [n]} \|\Delta - \Delta^{(i)}\|$ since the rest of the proof is the same as Lemma 5.

$$\max_{i \in [n]} \left\| \Delta - \Delta^{(i)} \right\| \le \max_{i \in [n]} \frac{1}{n} \left\| \operatorname{mat}(\Lambda_i^{-1} \operatorname{vec}(X_i)) \left(\xi_i - \xi_i' \right) \right\|,$$

where $\|\xi_i\|_{\psi_2} = \|\xi'_i\|_{\psi_2} = \sigma_{\xi}$ and $\Lambda_i^{-1} \operatorname{vec}(X_i) \sim N(0, \Lambda_i^{-1})$ for all $i = 1, \dots, n$. Therefore, for some absolute constant $c_0 > 0$,

$$\left\| \left\| \operatorname{mat}(\Lambda_i^{-1}\operatorname{vec}(X_i))\left(\xi_i - \xi_i'\right) \right\| \right\|_{\Psi_1} \le c_0 \sqrt{C_l^{-1}} \sigma_{\xi}.$$

We complete the proof by appying tail bound for sub-exponential random variable and taking a maximum over n.

References

- Abowd, J. M., I. M. Rodriguez, W. N. Sexton, P. E. Singer, and L. Vilhuber (2020). The modernization of statistical disclosure limitation at the us census bureau. US Census Bureau.
- Absil, P.-A., R. Mahony, and R. Sepulchre (2008). Optimization Algorithms on Matrix Manifolds. Princeton, NJ: Princeton University Press.
- Acharya, J., Z. Sun, and H. Zhang (2021). Differentially private assouad, fano, and le cam. In Algorithmic Learning Theory, pp. 48–78. PMLR.
- Adler, R. L., J. Dedieu, J. Y. Margulies, M. Martens, and M. Shub (2002, 07). Newton's method on Riemannian manifolds and a geometric model for the human spine. *IMA Journal* of Numerical Analysis 22(3), 359–390.

Apple Differential Privacy Team (2017). Learning with privacy at scale.

- Blum, A., C. Dwork, F. McSherry, and K. Nissim (2005, 06). Practical privacy: The sulq framework. Proceedings of the ACM SIGACT-SIGMOD-SIGART Symposium on Principles of Database Systems, 128–138.
- Brown, G., S. Hopkins, and A. Smith (2023, 12–15 Jul). Fast, sample-efficient, affine-invariant private mean and covariance estimation for subgaussian distributions. In G. Neu and L. Rosasco (Eds.), Proceedings of Thirty Sixth Conference on Learning Theory, Volume 195 of Proceedings of Machine Learning Research, pp. 5578–5579. PMLR.
- Burer, S. and R. D. Monteiro (2003). A nonlinear programming algorithm for solving semidefinite programs via low-rank factorization. *Mathematical Programming* 95(2), 329–357.
- Cai, T. T., Y. Wang, and L. Zhang (2021). The cost of privacy: Optimal rates of convergence for parameter estimation with differential privacy. *The Annals of Statistics* 49, 2825–2850.
- Cai, T. T., Y. Wang, and L. Zhang (2023). Score attack: A lower bound technique for optimal differentially private learning. arXiv preprint arXiv:2303.07152.
- Cai, T. T., D. Xia, and M. Zha (2024). Optimal differentially private pca and estimation for spiked covariance matrices. arXiv preprint arXiv:2401.03820.
- Candes, E. J. and Y. Plan (2011). Tight oracle inequalities for low-rank matrix recovery from a minimal number of noisy random measurements. *IEEE Transactions on Information The*ory 57(4), 2342–2359.
- Candes, E. J. and T. Tao (2005). Decoding by linear programming. *IEEE transactions on information theory* 51(12), 4203–4215.
- Chaudhuri, K., A. Sarwate, and K. Sinha (2012). Near-optimal differentially private principal components. *Advances in Neural Information Processing Systems* 25.
- Chen, H., G. Raskutti, and M. Yuan (2019). Non-convex projected gradient descent for generalized low-rank tensor regression. *The Journal of Machine Learning Research* 20(1), 172–208.
- Chen, Y. and M. J. Wainwright (2015). Fast low-rank estimation by projected gradient descent: General statistical and algorithmic guarantees.

- Chien, S., P. Jain, W. Krichene, S. Rendle, S. Song, A. Thakurta, and L. Zhang (2021). Private alternating least squares: Practical private matrix completion with tighter rates. pp. 1877– 1887.
- Dwork, C., F. McSherry, K. Nissim, and A. Smith (2006). Calibrating noise to sensitivity in private data analysis. *Theory of Cryptography Conference*, 265–284.
- Dwork, C., A. Roth, et al. (2014). The algorithmic foundations of differential privacy. Foundations and Trends® in Theoretical Computer Science 9(3-4), 211-407.
- Dwork, C., K. Talwar, A. Thakurta, and L. Zhang (2014). Analyze gauss: optimal bounds for privacy-preserving principal component analysis. *Proceedings of the forty-sixth annual ACM* symposium on Theory of computing, 11–20.
- Edelman, A., T. A. Arias, and S. T. Smith (1998). The geometry of algorithms with orthogonality constraints. SIAM journal on Matrix Analysis and Applications 20(2), 303–353.
- Erlingsson, U., V. Pihur, and A. Korolova (2014). Rappor: Randomized aggregatable privacypreserving ordinal response. In *Proceedings of the 2014 ACM SIGSAC Conference on Computer and Communications Security*, CCS '14, New York, NY, USA, pp. 1054–1067. Association for Computing Machinery.
- Hamidi, N. and M. Bayati (2022). On low-rank trace regression under general sampling distribution. Journal of Machine Learning Research 23(321), 1–49.
- Jain, P., O. D. Thakkar, and A. Thakurta (2018). Differentially private matrix completion revisited. pp. 2215–2224.
- Kamath, G., J. Li, V. Singhal, and J. Ullman (2019). Privately learning high-dimensional distributions. In *Conference on Learning Theory*, pp. 1853–1902. PMLR.
- Kamath, G., V. Singhal, and J. Ullman (2020, 09–12 Jul). Private mean estimation of heavytailed distributions. In J. Abernethy and S. Agarwal (Eds.), Proceedings of Thirty Third Conference on Learning Theory, Volume 125 of Proceedings of Machine Learning Research, pp. 2204–2235. PMLR.
- Koltchinskii, V. (2011). Von neumann entropy penalization and low-rank matrix estimation.

- Koltchinskii, V., K. Lounici, and A. B. Tsybakov (2011). Nuclear-norm penalization and optimal rates for noisy low-rank matrix completion.
- Koltchinskii, V. and D. Xia (2015). Optimal estimation of low rank density matrices. J. Mach. Learn. Res. 16(53), 1757–1792.
- Kuditipudi, R., J. Duchi, and S. Haque (2023, 12–15 Jul). A pretty fast algorithm for adaptive private mean estimation. In G. Neu and L. Rosasco (Eds.), *Proceedings of Thirty Sixth Conference on Learning Theory*, Volume 195 of *Proceedings of Machine Learning Research*, pp. 2511–2551. PMLR.
- Liu, X., W. Kong, P. Jain, and S. Oh (2022). Dp-pca: Statistically optimal and differentially private pca. Advances in Neural Information Processing Systems 35, 29929–29943.
- Liu, Z., Y.-X. Wang, and A. Smola (2015). Fast differentially private matrix factorization. In Proceedings of the 9th ACM Conference on Recommender Systems, pp. 171–178.
- Luo, Y. and A. R. Zhang (2022). Tensor-on-tensor regression: Riemannian optimization, over-parameterization, statistical-computational gap, and their interplay. *arXiv preprint* arXiv:2206.08756.
- McSherry, F. and I. Mironov (2009). Differentially private recommender systems: Building privacy into the netflix prize contenders. In *Proceedings of the 15th ACM SIGKDD international* conference on Knowledge discovery and data mining, pp. 627–636.
- Negahban, S. and M. J. Wainwright (2011). Estimation of (near) low-rank matrices with noise and high-dimensional scaling.
- Pajor, A. (1998). Metric entropy of the grassmann manifold. Convex Geometric Analysis 34 (181-188), 0942–46013.
- Rohde, A. and A. B. Tsybakov (2011). Estimation of high-dimensional low-rank matrices.
- Shen, Y., J. Li, J.-F. Cai, and D. Xia (2023). Computationally efficient and statistically optimal robust high-dimensional linear regression. arXiv preprint arXiv:2305.06199.
- Tropp, J. A. et al. (2015). An introduction to matrix concentration inequalities. Foundations and Trends (R) in Machine Learning 8(1-2), 1–230.

- Vadhan, S. (2017). The complexity of differential privacy. Tutorials on the Foundations of Cryptography: Dedicated to Oded Goldreich, 347–450.
- Vandereycken, B. (2013). Low-rank matrix completion by riemannian optimization. SIAM Journal on Optimization 23(2), 1214–1236.
- Vershynin, R. (2015). Estimation in high dimensions: a geometric perspective. In Sampling Theory, a Renaissance: Compressive Sensing and Other Developments, pp. 3–66. Springer.
- Vershynin, R. (2018). High-dimensional probability: An introduction with applications in data science, Volume 47. Cambridge university press.
- Wang, L., B. Zhao, and M. Kolar (2023, 25–27 Apr). Differentially private matrix completion through low-rank matrix factorization. 206, 5731–5748.
- Wang, Y.-X. (2018). Revisiting differentially private linear regression: optimal and adaptive prediction and estimation in unbounded domain. In UAI 2018.
- Wei, K., J.-F. Cai, T. F. Chan, and S. Leung (2016). Guarantees of riemannian optimization for low rank matrix recovery. SIAM Journal on Matrix Analysis and Applications 37(3), 1198–1222.
- Xia, D. (2021). Normal approximation and confidence region of singular subspaces. *Electronic Journal of Statistics* 15(2), 3798–3851.
- Zheng, Q. and J. Lafferty (2016). A convergent gradient descent algorithm for rank minimization and semidefinite programming from random linear measurements.