

# Log-rank test with coarsened exact matching <sup>\*</sup>

Tomoya Baba<sup>1,2</sup> and Nakahiro Yoshida<sup>1,2,3</sup>

<sup>1</sup>Graduate School of Mathematical Sciences, University of Tokyo <sup>†</sup>

<sup>2</sup>Japan Science and Technology Agency CREST

<sup>3</sup>The Institute of Statistical Mathematics

March 28, 2024

*Summary* It is of special importance in the clinical trial to compare survival times between the treatment group and the control group. Propensity score methods with a logistic regression model are often used to reduce the effects of confounders. However, the modeling of complex structures between the covariates, the treatment assignment and the survival time is difficult. In this paper, we consider coarsened exact matching (CEM), which does not need any parametric models, and we propose the weighted log-rank statistic based on CEM. We derive asymptotic properties of the weighted log-rank statistic, such as the weak convergence to a Gaussian process in Skorokhod space, in particular the asymptotic normality, under the null hypothesis and the consistency of the log-rank test. Simulation studies show that the log-rank statistic based on CEM is more robust than the log-rank statistic based on the propensity score.

*Keywords and phrases* Log-rank test, matching, central limit theorem, causal inference.

## 1 Introduction

In a clinical trial, suppose that each individual in a given collection  $\mathbb{G}$  of individuals is assigned to one of groups  $\mathbb{G}^z$  ( $z \in \{0, 1\}$ ): the treatment group  $\mathbb{G}^1$  and the control group  $\mathbb{G}^0$ . In this paper, we consider a comparison between survival times of the individuals  $\mathbb{G}^1$  and  $\mathbb{G}^0$ , and assess the treatment effect through the survival data. The survival time of the individual  $i$  is denoted by  $T^i$ . For a random variable  $U^i: \Omega \rightarrow (0, \infty)$  denoting the censoring time for  $i \in \mathbb{G}$ , we set

$$\tilde{T}^i = T^i \wedge U^i, \quad \delta^i = 1_{\{T^i \leq U^i\}}$$

for  $i \in \mathbb{G}$ . Moreover, it is supposed that each individual  $i \in \mathbb{G}$  has a covariate  $X^i$ , that is a random variable taking values in  $\mathbb{R}^d$ . Then,  $\tilde{T}^i$ ,  $\delta^i$  and  $X^i$  ( $i \in \mathbb{G}$ ) are observable in the trial.

Let  $Z^i$  denote the assignment of the individual  $i \in \mathbb{G}$ , that is,  $Z^i = z$  if  $i \in \mathbb{G}^z$ . The variables  $Z^i$  are in general random. If the sequence  $(Z^i)_{i \in \mathbb{G}}$  is generated by a Bernoulli trial, then the clinical

---

<sup>\*</sup>This work was in part supported by Japan Science and Technology Agency CREST JPMJCR2115; the research project with Kitasato University; Japan Society for the Promotion of Science Grants-in-Aid for Scientific Research No. 23H03354 (Scientific Research); and by a Cooperative Research Program of the Institute of Statistical Mathematics.

<sup>†</sup>Graduate School of Mathematical Sciences, University of Tokyo: 3-8-1 Komaba, Meguro-ku, Tokyo 153-8914, Japan. e-mail: nakahiro@ms.u-tokyo.ac.jp

trial is randomly controlled. In practice, however, quite possibly the distribution of the assignments may depend on the covariates the individuals have. In such a situation, it is necessary to adjust the distributions of the covariates between  $\mathbb{G}^1$  and  $\mathbb{G}^0$  to correctly compare the outcomes observed in the trial. Based on the observations  $(X^i, \tilde{T}^i, \delta^i)$ ,  $i \in \mathbb{G}$ , we test the hypothesis that there is no effect of the treatment. If the distributions of the covariates are the same for  $\mathbb{G}^1$  and  $\mathbb{G}^0$ , one can use an ordinary log-rank test as a test statistic. However, it does not make sense if the distributions are different, since the so assessed “treatment effect” may be biased by the difference between the distributions of covariates. It is often the case that patients of better conditions are apt to be selected than others in an invasive treatment, and this selection bias would affect the assessment of the treatment effect. If a perfectly randomized clinical trial is impossible ethically or for any other reason, then it is necessary to adjust the covariates’ distribution to correctly measure the treatment effects.

Propensity score methods, such as matching and inverse probability of treatment weighting (IPTW), are widely used to estimate treatment effects from observational data, including time-to-event data: see Austin (2014). There is some research on causal inference in survival analysis. For example, Austin (2013) considers the Cox regression model and conducts a Monte Carlo simulation to compare the performance of propensity score methods to estimate the hazard ratio. Xie and Liu (2005) proposes a weighted Kaplan-Meier estimator and a log-rank test statistic with IPTW. The asymptotic theory of the Cox regression model with IPTW is developed by Chen and Tsiatis (2001) and Guo et al (2021). Propensity score methods need an estimation of the unknown propensity score, and logistic regression is commonly used. On the other hand, there are nonparametric matching methods. Although these methods have the advantage of not requiring any models for data, there is little mathematical research except for Abadie and Imbens (2012).

In this paper, we propose a weighted log-rank test combined a matching method, and study asymptotic properties of the test. This paper is finally aiming at proving asymptotic behavior of the test statistic when  $\#\mathbb{G}^0 \gg \#\mathbb{G}^1$ . Such a problem occurs in the studies of rare diseases and in advanced medicine, where the number of treated patients is much smaller than ones cured by ordinary treatments. For example, as mentioned below, the new therapy with heart sheets is the case.

This work was inspired by the joint research with Osaka University on the statistical analysis of the autologous stem cell-sheet transplantation therapy for treating cardiomyopathy, that was introduced by Professor Yoshiki Sawa and his group at the medical school of Osaka University. The authors appreciate Professor Shigeru Miyagawa (Osaka University, Graduate School of Medicine) and his research group members. Many thanks go to Professor Nobuhiko Okada (Kitasato University) with his group for their great supports for the joint research project, and also to Professor Masahiro Takeuchi (University of Tokyo), Dr. Takumi Suzuki (University of Tokyo) and Dr. Madoka Takeuchi (University of Tokyo) for their valuable comments and helps.

## 2 Weighted log-rank statistic with CEM

Coarsened exact matching (CEM) is a statistical method that classifies all individuals into matched and unmatched individuals: see Iacus et al (2011, 2019). The algorithm of CEM is as follows. The groups  $\mathbb{G}^z$  is given as  $\mathbb{G}^z = \{i \in \mathbb{G}; Z_i = z\}$  for every  $z \in \{1, 0\}$ . They are exclusive sets of individuals that are in general random. Set a partition  $(a)_{a \in \mathbb{A}}$  of a measurable set  $\mathcal{X}$  of  $\mathbb{R}^d$ . Suppose that  $\mathbb{A}$  is a countable set and each element  $a \in \mathbb{A}$  is measurable. Input the covariates  $X^i$  ( $i \in \mathbb{G}$ ) and define  $\mathcal{P}$  by

$$\mathcal{P} = \left\{ (i, j) \in \mathbb{G}^1 \times \mathbb{G}^0; (X^i, X^j) \in \bigsqcup_{a \in \mathbb{A}} (a \times a) \right\}.$$

Let  $G^z = \varpi_z \mathcal{P}$  for  $z \in \{1, 0\}$ , where  $\varpi_z$  is the projection  $\mathbb{G}^1 \times \mathbb{G}^0 \rightarrow \mathbb{G}^z$ .

For  $i \in \mathbb{G}$ , the processes  $(N_t^i)_{t \in \mathbb{R}_+}$  and  $(Y_t^i)_{t \in \mathbb{R}_+}$  are defined by

$$N_t^i = 1_{\{\tilde{T}^i \leq t, \delta^i = 1\}} \quad \text{and} \quad Y_t^i = 1_{\{\tilde{T}^i \geq t\}},$$

respectively. Write  $a^i$  for  $a \in \mathbb{A}$  when  $X^i \in a$ . We define the weights process  $(w_t^i)_{t \in \mathbb{R}_+}$  for  $i \in \mathbb{G}$  as follows:

$$w_t^i = \begin{cases} 1 & \text{if } i \in G^1 \\ \left( \sum_{j \in \mathbb{G}^0} 1_{\{X^j \in a^i\}} Y_t^j \right)^- \sum_{i' \in \mathbb{G}^1} 1_{\{X^{i'} \in a^i\}} Y_t^{i'} & \text{if } i \in G^0 \\ 0 & \text{if } i \notin G^1 \sqcup G^0, \end{cases} \quad (2.1)$$

where  $x^- = 1_{\{x \neq 0\}} x^{-1}$  for  $x \in \mathbb{R}$ . Remark that in (2.1),

$$\left( \sum_{j \in \mathbb{G}^0} 1_{\{X^j \in a^i\}} Y_t^j \right)^- \sum_{i' \in \mathbb{G}^1} 1_{\{X^{i'} \in a^i\}} Y_t^{i'} = \left( \sum_{j \in G^0} 1_{\{X^j \in a^i\}} Y_t^j \right)^- \sum_{i' \in G^1} 1_{\{X^{i'} \in a^i\}} Y_t^{i'}$$

if  $i \in G^0$ . Let

$$\mathcal{N}_t^i = \int_0^t w_s^i dN_s^i, \quad \bar{\mathcal{N}}_t^z = \sum_{i \in G^z} \mathcal{N}_t^i, \quad \mathcal{Y}_t^i = w_t^i Y_t^i, \quad \bar{\mathcal{Y}}_t^z = \sum_{i \in G^z} \mathcal{Y}_t^i.$$

Remark that  $(\mathcal{N}_t^i)_{t \in \mathbb{R}_+}$  and  $(\bar{\mathcal{N}}_t^z)_{t \in \mathbb{R}_+}$  are no longer counting processes.

To assess the treatment effect, we consider the weighted log-rank statistic

$$\mathcal{W}_t = \int_0^t K_s^n \{ (\bar{\mathcal{Y}}_s^1)^- d\bar{\mathcal{N}}_s^1 - (\bar{\mathcal{Y}}_s^0)^- d\bar{\mathcal{N}}_s^0 \} \quad (t \in \mathbb{R}_+), \quad (2.2)$$

where

$$K_s^n = \{ (\bar{\mathcal{Y}}_0^1 + \bar{\mathcal{Y}}_0^0) (\bar{\mathcal{Y}}_0^1 \bar{\mathcal{Y}}_0^0)^- \}^{1/2} (\bar{\mathcal{Y}}_s^1 + \bar{\mathcal{Y}}_s^0)^- \bar{\mathcal{Y}}_s^1 \bar{\mathcal{Y}}_s^0 W_s^n \quad (s \in \mathbb{R}_+) \quad (2.3)$$

and  $W_s^n$  is a locally bounded process predictable with respect to the filtration  $\mathbb{F}$  specified below. When  $W_s^n = 1$ , the weighted log-rank statistic  $\mathcal{W}_t$  becomes the usual log-rank statistic.

### 3 Main results

Suppose that the random elements  $T^i$ ,  $X^i$  and  $U^i$  are defined on a complete probability space  $(\Omega, \mathcal{F}, P)$ . The completeness is not restrictive since completion of a probability space is always possible. Denote by  $\mathbb{N}$  the collection of all  $P$ -null sets. To prove the main result, we need some results from the martingale theory. We begin with computing the compensator of the process  $N^i$ . Define the process  $(\dot{N}_t^i)_{t \in \mathbb{R}_+}$  by  $\dot{N}_t^i = 1_{\{\tilde{T}^i \leq t, \delta^i = 0\}}$  ( $t \in \mathbb{R}_+$ ), and the filtration  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$  by


$$\mathcal{F}_t = \sigma[X^i, Z^i, N_s^i, \dot{N}_s^i \mid i \in \mathbb{G}, s \leq t] \vee \sigma[\mathbb{N}].$$

Then the filtration  $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$  is right-continuous:  $\mathcal{F}_t = \bigcap_{\epsilon > 0} \mathcal{F}_{t+\epsilon}$ . See Brémaud (1981).

For  $i \in \mathbb{G}$ , denote by

$$(P^{T^i}(B \mid X^i = x, Z^i = z); B \in \mathbb{B}[\mathbb{R}_+], x \in \mathbb{R}^d, z \in \{0, 1\})$$

Table 1: The causal diagram of the random variables.



a regular conditional probability of  $T^i$  given  $(X^i, Z^i) = (x, z)$ . Let

$$\Lambda^i(t \mid X^i = x, Z^i = z) = \int_0^t \frac{P^{T^i}(ds \mid X^i = x, Z^i = z)}{P^{T^i}([s, \infty) \mid X^i = x, Z^i = z)} \quad (3.1)$$

and

$$\Lambda^i(t \mid X^i, Z^i) = \Lambda^i(t \mid X^i = x, Z^i = z) \Big|_{x=X^i, z=Z^i}.$$

In what follows, we assume the following condition.

- [C] (i) The random elements  $(T^i, U^i)$  ( $i \in \mathbb{G}$ ) are  $\mathcal{F}_0$ -conditionally independent.  
(ii) For each  $i \in \mathbb{G}$ ,  $T^i$  and  $U^i$  are  $\mathcal{F}_0$ -conditionally independent.  
(iii) For each  $i \in \mathbb{G}$ ,  $T^i$  and  $\sigma[X^j, Z^j \mid j \neq i]$  are  $\sigma[X^i, Z^i]$ -conditionally independent.  
(iv) In  $i \in \mathbb{G}$ , the conditional law  $\mathcal{L}\{T^i \mid X^i = x, Z^i = z\}$  is identical.

Table 1 shows the dependency of  $i$ -th random variables. For random variables  $X$  and  $Y$ , the arrow  $X \rightarrow Y$  denotes that the value of  $X$  affects the value of  $Y$ .

Under Condition [C] (iv), we can choose a common version  $\Lambda(t \mid x, z)$  of  $\Lambda^i(t \mid X^i = x, Z^i = z)$  such that

$$\Lambda^i(t \mid X^i = x, Z^i = z) = \Lambda(t \mid x, z) \quad (t \in \mathbb{R}_+, x \in \mathbb{R}^d, z \in \{0, 1\})$$

for all  $i \in \mathbb{G}$ . We consider the following conditions.

- [A1] There exist a  $\sigma$ -finite measure  $\rho$  on  $([0, \tau], \mathbb{B}([0, \tau]))$  and a constant  $L$  such that  $\rho$  has no atom,  $\rho([0, \tau]) < \infty$ ,  $\Lambda(\cdot \cap [0, \tau] \mid x, z)$  is dominated by  $\rho$ , and

$$|p(s \mid x, z) - p(s \mid y, z)| \leq L|x - y|$$

for all  $x, y \in \mathcal{X}$  and  $z \in \{0, 1\}$ , for a version of the Radon-Nikodym derivative  $p(\cdot \mid x, z) = d\Lambda(\cdot \cap [0, \tau] \mid x, z)/d\rho$ . Moreover,

$$\|p\|_\infty := \sup_{s \in [0, \tau], x \in \mathcal{X}, z \in \{0, 1\}} p(s \mid x, z) < \infty. \quad (3.2)$$

Usually,  $\rho$  is the Lebesgue measure on  $[0, \tau]$  or some measure equivalent to it. However, logically we may consider a measure like the Cantor distribution. The continuity of the  $\Lambda([0, \cdot], \mid x, z)$  can be relaxed but this condition is adopted for simplicity of the analysis.

We regard a parameter  $n \in \mathbb{N}$  as the driver of the asymptotic theory as  $n \rightarrow \infty$ . In what follows,  $\mathbb{G}^z$ ,  $G^z$  ( $z \in \{0, 1\}$ ) and  $\mathbb{A}$  for a partition of  $\mathcal{X}$  will depend on  $n$ . Let  $n_z = \#G^z$  for  $z \in \{0, 1\}$ . The

number  $n_z$  is generally random and satisfies  $n_z \leq \#\mathbb{G}^z$ . The number  $n$  is not necessarily equal to  $\bar{n} = n_0 + n_1$ . For example,  $n = \#\mathbb{G}$  individuals enter the trial, and the number of individuals randomly or deterministically assigned to the treatment/control group after matching is  $n_z$ . The number  $\bar{n}$  can be much less than  $n$  due to matching. In this example, it is also possible to assume that the parameter  $n$  only specifies  $n_z$  as a function  $n_z(n)$  of  $n$ , without corresponding to any number in reality.

As suggested in Introduction, we will consider the situation where  $\#\mathbb{G}^0 \gg \#\mathbb{G}^1$ . For example, it is the case where the number of observations is limited for the new therapy whose effect we want to know, while one can use big registry data obtained in the traditional treatments. In such a situation, the conditions become asymmetric between the two groups, as set as follows.

[A2] There exists a (bounded) measurable function  $q = (q_s)_{s \in [0, \tau]}$  on  $[0, \tau]$  such that

$$\sup_{s \in [0, \tau]} \left| n_1^- \sum_{i \in G^1} Y_s^i p(s | X^i, Z^i) - q_s \right| \xrightarrow{p} 0$$

as  $n \rightarrow \infty$ .

[A3] As  $n \rightarrow \infty$ ,

$$P \left[ \inf_{i \in G^1} \sum_{j \in G^0} 1_{\{X^j \in a^i\}} Y_\tau^j = 0 \right] \rightarrow 0.$$

[A4]  $\sup_{t \in [0, \tau], i \in G^0} w_t^i \xrightarrow{p} 0$  as  $n \rightarrow \infty$ .

[A5]  $n_1^- \xrightarrow{p} 0$  and  $n_1^{1/2} d_n \xrightarrow{p} 0$  as  $n \rightarrow \infty$ , for  $d_n = \sup_{a \in \mathbb{A}} \text{diam}(a)$ .

[A6] There exists a bounded measurable function  $W^\infty = (W_s^\infty)_{s \in [0, \tau]}$  on  $[0, \tau]$  such that

$$\sup_{s \in [0, \tau]} |W_s^n - W_s^\infty| \xrightarrow{p} 0$$

as  $n \rightarrow \infty$ .

By constructing the log-rank statistic, we want to test the hypothesis

[H<sub>0</sub>]  $\Lambda(t | x, 1) = \Lambda(t | x, 0)$  ( $t \in [0, \tau]$ ,  $x \in \mathcal{X}$ ).

Before we consider asymptotic properties of the weighted log-rank statistic as  $n \rightarrow \infty$ , we should clarify the dependency of variables on  $n$ . The random elements  $(X^i, Z^i, T^i, U^i)$  ( $i \in \mathbb{G}$ ) may depend on  $n$ . The sets  $\mathbb{G}$ ,  $\mathbb{G}^z$  ( $z = 0, 1$ ) and  $\mathbb{A}$  depend on  $n$  as we have mentioned before. On the other hand,  $\mathcal{X}$ ,  $\Lambda^i(t | X^i = x, Z^i = z)$  and  $p(s | x, z)$  do not depend on  $n$ .

The following central limit theorem enables us to make a critical region of the test. The D-space on  $[0, \tau]$  equipped with the Skorokhod topology is denoted by  $D([0, \tau])$ .

**Theorem 3.1.** *Suppose that conditions [C], [A1]-[A6] and [H<sub>0</sub>] are fulfilled. Then  $\mathcal{W} \xrightarrow{d} \mathbb{W}$  in  $D([0, \tau])$  as  $n \rightarrow \infty$ , where  $\mathbb{W} = (\mathbb{W}_t)_{t \in [0, \tau]}$  is a centered Gaussian process with independent increments and  $\text{Var}[\mathbb{W}_t] = \frac{1}{2} \int_0^t (W_s^\infty)^2 q_s \rho(ds)$  for  $t \in [0, \tau]$ .*

A proof of Theorem 3.1 is in Section 7. Given a consistent estimator  $\mathcal{V}_\tau$  of the asymptotic variance  $\text{Var}[\mathbb{W}_\tau]$  of  $\mathbb{W}_\tau$ , based on  $\mathcal{W}$ , the critical region of the test of  $[H_0]$  is, for example,

$$\mathcal{C} = \{\mathcal{V}_\tau^{-1/2}\mathbb{W}_\tau \geq z_\alpha\},$$

where  $z_\alpha$  is the upper  $\alpha$ -point of the standard normal distribution. The shape of the critical region is chosen depending on the alternative hypothesis. We can construct a consistent estimator  $\mathcal{V}_\tau$  as follows.

**Proposition 3.2.** *We define the estimator of  $\text{Var}[\mathbb{W}_\tau]$  in Theorem 3.1 by setting*

$$\mathcal{V}_\tau = \frac{1}{n_1} \sum_{i \in G^1} \frac{1}{2} \int_0^\tau (W_s^n)^2 dN_s^i.$$

*Under conditions [C], [A1], [A2], [A5] and [A6], we have  $\mathcal{V}_\tau \xrightarrow{p} \text{Var}[\mathbb{W}_\tau]$ .*

Proof of Proposition 3.2 is given in section 8.

We consider the consistency of the log-rank statistic  $\mathcal{V}_\tau^{-1/2}\mathbb{W}_\tau$  from the convergence  $\mathbb{W}_\tau \xrightarrow{d} \mathbb{W}$  obtained in Theorem 3.1, where  $\mathcal{V}_\tau$  is a consistent estimator of the variance  $\text{Var}[\mathbb{W}_\tau]$ . Suppose that [A1] and [A6] hold. We consider the following alternative hypothesis.

[H<sub>1</sub>] The function  $p(t | x, z)$  in [A1] satisfies

$$p(t | x, 1) \leq p(t | x, 0) \quad (t \in [0, \tau], x \in \mathcal{X}).$$

Moreover, there exists  $B \in \mathcal{B}([0, \tau])$  such that  $\rho(B) > 0$  and

$$\sup_{t \in B, x \in \mathcal{X}} W_t^\infty \{p(t | x, 1) - p(t | x, 0)\} < 0$$

for the process  $W^\infty$  in [A6].

We consider the following condition to show the consistency of the log-rank statistic.

[A7] There exists a positive number  $\tilde{q}_\tau$  such that

$$n_1^- \sum_{i \in G^1} Y_\tau^i \xrightarrow{p} \tilde{q}_\tau$$

as  $n \rightarrow \infty$ .

**Theorem 3.3.** *Suppose that conditions [A1]-[A7] and [H<sub>1</sub>] hold. Let  $\mathcal{V}_\tau$  be a consistent estimator of  $\text{Var}[\mathbb{W}_\tau]$  in Theorem 3.1. Then, we have*

$$P(\mathcal{V}_\tau^{-1/2}\mathbb{W}_\tau < -R) \rightarrow 1 \quad (n \rightarrow \infty) \tag{3.3}$$

for all  $R > 0$ .

See Section 9 for the proof of Theorem 3.3.

## 4 Sufficient conditions for the assumptions of the main theorems

When  $(X^i, Z^i, T^i, U^i)$  ( $i \in \mathbb{G}$ ) is an i.i.d. sequence, conditions [A1]-[A7] can be simplified. Let  $\#\mathbb{G} = n$ . Suppose that the sequence of random elements  $\{(X^i, Z^i, T^i, U^i)\}_{i \in \mathbb{G}}$  is i.i.d. for each  $n \in \mathbb{N}$  and the conditional distribution  $\mathcal{L}\{(X^1, T^1, U^1) \mid Z^1 = z\}$  does not depend on  $n$  for each  $z \in \{0, 1\}$ . Suppose that  $\mathcal{X} \subset \mathbb{R}^d$  is compact. We consider the following conditions.

[B1] Constants  $\beta, \theta \in (0, 1)$  and  $\bar{d} \in \{1, \dots, d\}$  satisfy  $\beta < 2\theta < \bar{d}^{-1}$ .

[B2]  $-\infty < \liminf_{n \rightarrow \infty} n^{-\beta+1} P(Z^1 = 1) \leq \limsup_{n \rightarrow \infty} n^{-\beta+1} P(Z^1 = 1) < \infty$ .

[B3] The conditional distribution  $\mathcal{L}\{X^1 \mid Z^1 = z\}$  has a density  $f(\cdot \mid z)$  with respect to a  $\sigma$ -finite measure  $\nu$  on  $\mathbb{R}^d$ . It holds that

$$\begin{aligned} \operatorname{ess\,sup}_{x \in \mathcal{X}} f(x \mid 1) &< \infty, \\ \operatorname{ess\,inf}_{x \in \mathcal{X}} (f(x \mid 0) E[Y_\tau^1 \mid X^1 = x, Z^1 = 0]) &> 0, \end{aligned}$$

where  $\operatorname{ess\,sup}_{x \in \mathcal{X}}$  and  $\operatorname{ess\,inf}_{x \in \mathcal{X}}$  are with respect to the measure  $\nu$ .

[B4]  $P(\tilde{T}^1 = s, X^1 \in \mathcal{X} \mid Z^1 = 1) = 0$  for all  $s \in [0, \tau]$ , and  $P(\tilde{T}^1 \geq \tau, X^1 \in \mathcal{X} \mid Z^1 = 1) > 0$ .

[B5]  $d_n = O(n^{-\theta})$  as  $n \rightarrow \infty$  and

$$0 < \liminf_{n \rightarrow \infty} n^{\bar{d}\theta} \inf_{a \in \mathbb{A}} \nu(a) \leq \limsup_{n \rightarrow \infty} n^{\bar{d}\theta} \sup_{a \in \mathbb{A}} \nu(a) < \infty.$$

[A1'] Condition [A1] holds and the function  $p(\cdot \mid x, 1)$  in [A1] is continuous on  $[0, \tau]$  for all  $x \in \mathcal{X}$ .

**Remark 4.1.** (i) We may regard  $\bar{d}$  as the number of the continuous covariates. Suppose that the covariates  $X^i$  take values in  $\mathbb{R}^{\bar{d}} \times \mathbb{Z}^{d-\bar{d}}$  and [B3] holds for  $\nu = m_L \times m_C$ , where  $m_L$  is the Lebesgue measure on  $\mathbb{R}^{\bar{d}}$  and  $m_C$  is the counting measure on  $\mathbb{Z}^{d-\bar{d}}$ . Consider the set of strata of CEM  $\mathbb{A}$  satisfying  $d_n = O(n^{-\theta})$ , for example,

$$\mathbb{A} = \left\{ \prod_{\ell=1}^{\bar{d}} (k_\ell n^{-\theta}, k_{\ell+1} n^{-\theta}) \mid k_1, \dots, k_d \in \mathbb{Z} \right\}.$$

Then,  $\inf_{a \in \mathbb{A}} \nu(a) = \sup_{a \in \mathbb{A}} \nu(a) = n^{-\bar{d}\theta}$  for large enough  $n \in \mathbb{N}$  and [B5] are fulfilled.

(ii) Under [A1], in [B4] the condition that  $P(\tilde{T}^1 = s \mid Z^1 = 1) = 0$  for all  $s \in [0, \tau]$  can be replaced by the condition that  $P(U^1 = s \mid Z^1 = 1) = 0$  for all  $s \in [0, \tau]$ .

For each  $n \in \mathbb{N}$ , let  $V^i = (X^i, Z^i, T^i, U^i)$  ( $i \in \mathbb{G}$ ). We have

$$E[g(V^i) \mid X^i \in \mathcal{X}, Z^i = 1] = \frac{E[g(V^i) 1_{\{X^i \in \mathcal{X}\}} \mid Z^i = 1]}{P(X^i \in \mathcal{X} \mid Z^i = 1)}$$

for a nonnegative measurable function  $g: \mathbb{R}^d \times \{0, 1\} \times (0, \infty) \times (0, \infty) \rightarrow \mathbb{R}$ . Note that  $P(X^i \in \mathcal{X} \mid Z^i = 1) > 0$  under [B4]. We define

$$g_s = E[Y_s^1 p(s \mid X^1, Z^1) \mid X^1 \in \mathcal{X}, Z^1 = 1] \quad (s \in [0, \tau]).$$

Consider a centered Gaussian process  $\mathbb{W} = (\mathbb{W}_t)_{t \in [0, \tau]}$  with independent increments and  $\text{Var}[\mathbb{W}_t] = \frac{1}{2} \int_0^t (W_s^\infty)^2 q_s \rho(ds)$ . We define the estimator  $\mathcal{V}_\tau$  of  $\text{Var}[\mathbb{W}_\tau]$  as follows.

$$\mathcal{V}_\tau = \frac{1}{2} \int_0^\tau (W_s^\infty)^2 dN_s^i.$$

**Theorem 4.2.** *Suppose that conditions [C], [A1'], [A6] and [B1]-[B5] are satisfied. Then, the following properties (1)-(3) hold.*

- (1) *If  $[H_0]$  is satisfied, then  $\mathcal{W} \rightarrow^d \mathbb{W}$  in  $D([0, \tau])$  as  $n \rightarrow \infty$ .*
- (2)  *$\mathcal{V}_\tau \rightarrow^p \text{Var}[\mathbb{W}_\tau]$  ( $n \rightarrow \infty$ ).*
- (3) *If  $[H_1]$  is satisfied, then  $P((\mathcal{V}_\tau^{1/2})^{-1} \mathcal{W}_\tau < -R) \rightarrow 1$  as  $n \rightarrow \infty$  for all  $R > 0$ .*

See section 10 for the proof of Theorem 4.2.

## 5 Causal inference

In the context of the causal inference, we consider a random field  $\mathbb{T}^i : \Omega \times \{0, 1\} \rightarrow (0, \infty)$  for  $i \in \mathbb{G} = \mathbb{G}^1 \sqcup \mathbb{G}^0$  on a sample space  $\Omega$ , to describe the potential outcome of the individual  $i \in \mathbb{G}$ , i.e., the random field  $\mathbb{T}^i$  expresses counterfactual outcomes. When causal inference in question is survival analysis aiming at discovery of the treatment effect on survival time,  $\mathbb{T}^i$  denotes the potential survival time of the individual  $i$ , and  $T^i = \mathbb{T}(Z^i)$ . When  $i \in \mathbb{G}^z$ ,  $\mathbb{T}^i(z)$  is observed, however, the counterfactual value of  $\mathbb{T}^i(1-z)$  is never observed from the trial. In the causal inferential framework, one is interested in comparing, in some measure, the two conditional laws  $\mathcal{L}\{\mathbb{T}^i(z) \mid X^i = x, Z^i = z\}$  and  $\mathcal{L}\{\mathbb{T}^i(1-z) \mid X^i = x, Z^i = z\}$ . Though  $\mathbb{T}^i(1-z)$  is counterfactual when  $Z^i = z$ ,  $\mathcal{L}\{\mathbb{T}^i(1-z) \mid X^i = x, Z^i = z\}$  is estimable if the condition that

$$[\mathbf{N}] \quad \mathcal{L}\{\mathbb{T}^i \mid X^i = x, Z^i = z\} = \mathcal{L}\{\mathbb{T}^i \mid X^i = x\} \text{ for } x \in \mathcal{X} \text{ and } z \in \{0, 1\}.$$


is met. Indeed, under [N],  $\mathcal{L}\{\mathbb{T}^i(1-z) \mid X^i = x, Z^i = z\} = \mathcal{L}\{\mathbb{T}^i(1-z) \mid X^i = x, Z^i = 1-z\} = \mathcal{L}\{T^i \mid X^i = x, Z^i = 1-z\}$ . Therefore, under the setting [N], one can consider a test about equality between the effects  $\mathbb{T}^i(1)$  and  $\mathbb{T}^i(0)$  given  $x$ . It should be remarked that this test is of different nature from a test like  $[H_0]$ . To clarify the idea, consider a simple example such that  $\mathbb{T}^i(z) = z + |X^i| + \delta^i$ , where  $\delta^i$  is a random variable having dependency on  $Z^i$ . Condition [N] does not hold for this  $\mathbb{T}^i$ , so the comparison between  $\mathbb{T}^i(1)$  and  $\mathbb{T}^i(0)$  in the above sense is impossible. However, like the test of  $[H_0]$ , the comparison between  $\mathcal{L}\{\mathbb{T}^i(1) \mid X^i = x, Z^i = 1\}$  and  $\mathcal{L}\{\mathbb{T}^i(0) \mid X^i = x, Z^i = 0\}$  is still possible. This is a comparison of distributions that is taking the effects of the unobserved confounder  $\delta^i$  into account. In other words, if the property [N] holds, then our test gives a causal inferential test of equality between  $\mathcal{L}\{\mathbb{T}^i(1) \mid X^i = x, Z^i = z\}$  and  $\mathcal{L}\{\mathbb{T}^i(0) \mid X^i = x, Z^i = z\}$  in some measure.

## 6 Simulation Studies

We conducted simulations to compare the performance of CEM and IPTW used in the log-rank test and to confirm the consistency of the log-rank statistics with CEM. In the first experiment, we iterated to generate data under the null hypothesis and draw two histograms of the log-rank statistics using CEM and IPTW. While we suppose a parametric model and estimate the propensity score when calculating IPTW, we do not need any models when calculating the log-rank statistic with CEM.



Table 2: The causal diagram of the random variables with potential outcomes.



Thus, we considered the two cases, with and without misspecification, in estimating the propensity score to calculate IPTW. In the second experiment, we generated data under the alternative hypothesis and calculated the log-rank statistics with CEM. We drew three histograms of the log-rank statistics based on data of different sample sizes to observe the consistency.

The detailed definition of the simulation procedures is as follows. We generate i.i.d. samples  $\{(X^i, Z^i, \tilde{T}^i, U^i)\}_{i=1}^n$ , where  $n = 10000$  in the first experiment and  $n = 3000, 10000$  and  $30000$  in the second experiment. Let  $X^i = (X_1^i, X_2^i)$  ( $i = 1, \dots, n$ ) be  $\mathbb{R} \times \{0, 1\}$ -valued random variables. Suppose that  $X_1^i$  and  $X_2^i$  are independent,  $X_1^i$  follows the standard normal distribution  $N(0, 1)$  and  $X_2^i$  satisfies  $P(X_2^i = 1) = P(X_2^i = 0) = 1/2$  for each  $i = 1, \dots, n$ . We consider the following propensity score models for generating  $\{Z^i\}_{i=1}^n$ .

**Model (1)** a logistic regression model without interaction

$$P(Z^i = 1 \mid X^i) = \frac{\exp(\alpha_1 + X_1^i + X_2^i)}{1 + \exp(\alpha_1 + X_1^i + X_2^i)}.$$

**Model (2)** a logistic regression model with interaction

$$P(Z^i = 1 \mid X^i) = \frac{\exp(\alpha_2 + \eta^i X_1^i)}{1 + \exp(\alpha_2 + \eta^i X_1^i)}, \quad \eta^i = \begin{cases} 1 & (\text{if } X_2^i = 1) \\ -1 & (\text{if } X_2^i = 0) \end{cases}.$$

We define  $\alpha_1$  and  $\alpha_2$  so that the size of the treatment group  $\#\{i \in \{1, \dots, n\} \mid Z^i = 1\}$  is nearly equal to  $\sqrt{n}$ . Let  $\alpha_1 = -5.7$  and  $\alpha_2 = -6.4$  in the first experiment and  $\alpha_2 = -5.7, -6.4, -7.0$  for  $n = 3000, 10000, 30000$  respectively in the second experiment. We generate the potential outcomes  $\mathbb{T}^i(z)$  ( $z = 0, 1, i = 1, \dots, n$ ) by using the Cox hazards model. For  $x = (x_1, x_2) \in \mathbb{R} \times \{0, 1\}$  and  $z \in \{0, 1\}$ , let  $\Lambda(t \mid x, z)$  ( $t \geq 0$ ) denote the cumulative hazard function of the conditional distribution of  $\mathbb{T}^i(z)$  given  $X^i = x$ . In other words, let

$$\Lambda(t \mid x, z) = \int_0^t \frac{P^{\mathbb{T}^i(z)}(ds \mid X^i = x)}{P^{\mathbb{T}^i(z)}([s, \infty) \mid X^i = x)}.$$

If the condition  $[N]$  holds, this function equals the function  $\Lambda(t \mid x, z)$  in Equation (3.1). When we generate  $\{\mathbb{T}^i(z)\}_{i=1}^n$  ( $z = 0, 1$ ) occurring under the null hypothesis and alternative hypothesis, we calculate the conditional distribution of  $\{\mathbb{T}^i(z)\}$  given  $X^i = x$  and  $Z^i = z$  from the Cox hazard model

$$\Lambda(t \mid x, z) = \begin{cases} \int_0^t h_0(s) \exp(0.2x_1 + x_2) ds & (\text{under the null hypothesis}) \\ \int_0^t h_0(s) \exp(-z + 0.2x_1 + x_2) ds & (\text{under the alternative hypothesis}) \end{cases}$$

where  $h_0(s)$  is the constant function  $h_0(s) = e^{-1.5}$ . Suppose that  $U^i$  ( $i = 1, \dots, n$ ) follows the uniform distribution  $U(0, 10)$ .

Let  $\tau = 8$ . From the observations  $\{(X^i, Z^i, \tilde{T}^i, \delta^i)\}_{i=1}^n$ , we calculate the log-rank statistics  $\mathcal{W}_\tau$  and the estimator of their asymptotic variances  $\mathcal{V}_\tau$  using CEM and IPTW. Whether using CEM and IPTW, the definition of  $\mathcal{W}_\tau$  is given in (2.2) and (2.3). Let  $W^n = 1$ . The differences between the definition of the log-rank statistics with CEM and IPTW are the weights process  $w_s^i$  and the estimator of the asymptotic variance  $\mathcal{V}_\tau$ . First, we consider the case with CEM proposed in this paper. Let  $\mathcal{X} = (-5, 5] \times \{0, 1\}$ . We define the strata of the CEM as follows.

$$\mathbb{A} = \left\{ \left( \frac{k}{\lfloor n^{3/8} \rfloor}, \frac{k+1}{\lfloor n^{3/8} \rfloor} \right) \times \{z\} \mid k \in \mathbb{Z}, -5 \lfloor n^{3/8} \rfloor \leq k < 5 \lfloor n^{3/8} \rfloor, z = 0, 1 \right\}.$$

The process  $w_s^i$  for the log-rank statistic with CEM is defined as (2.1) and the estimator of the asymptotic variance  $\mathcal{V}_\tau$  is given in Proposition 3.2. Next, we define  $w_s^i$  and  $\mathcal{V}_\tau$  for the log-rank statistic with IPTW. This statistic is introduced in Xie and Liu (2005). We calculate  $\hat{w}^i$  as follows and let  $w_s^i = \hat{w}^i$  for all  $s \geq 0$ .

$$\begin{aligned} \hat{w}^i &= Z^i / \hat{p}^i + (1 - Z^i) / (1 - \hat{p}^i), \\ \hat{p}^i &= \frac{\exp(\hat{\alpha}_0 + \hat{\alpha}_1 X_1^i + \hat{\alpha}_2 X_2^i)}{1 + \exp(\hat{\alpha}_0 + \hat{\alpha}_1 X_1^i + \hat{\alpha}_2 X_2^i)}, \end{aligned}$$

where  $(\hat{\alpha}_0, \hat{\alpha}_1, \hat{\alpha}_2)$  is the maximum likelihood estimator of the parameter  $(\alpha_0, \alpha_1, \alpha_2)$  in the logistic regression model

$$P(Z^i = 1 \mid X^i) = \frac{\exp(\alpha_0 + \alpha_1 X_1^i + \alpha_2 X_2^i)}{1 + \exp(\alpha_0 + \alpha_1 X_1^i + \alpha_2 X_2^i)}.$$

Let

$$\begin{aligned} \bar{N}_s &= \sum_{i \in \mathbb{G}} N_s^i, \quad \bar{Y}_s = \sum_{i \in \mathbb{G}} Y_s^i, \\ U_s &= \{\bar{\mathcal{Y}}_s^0(\bar{\mathcal{Y}}_s^1 + \bar{\mathcal{Y}}_s^0)^{-}\}^2 \sum_{i \in \mathbb{G}^1} (w^i)^2 Y_s^i + \{\bar{\mathcal{Y}}_s^1(\bar{\mathcal{Y}}_s^1 + \bar{\mathcal{Y}}_s^0)^{-}\}^2 \sum_{i \in \mathbb{G}^0} (w^i)^2 Y_s^i \end{aligned}$$

for  $s \geq 0$  and  $i \in \mathbb{G}$  and we define

$$\mathcal{V}_\tau = (\bar{\mathcal{Y}}_0^1 + \bar{\mathcal{Y}}_0^0)^{-} \bar{\mathcal{Y}}_0^1 \bar{\mathcal{Y}}_0^0 \int_0^\tau U_s \{\bar{Y}_s(\bar{Y}_s - 1)\}^{-} (\bar{Y}_s - \Delta \bar{N}_s) d\bar{N}_s.$$

Figures 1-4 show the result of the first experiment. From Figure 1 and Figure 2, we can see that if we generate data that follows Model (1), then the performance of the log-rank statistics with CEM and IPTW is almost the same. Moreover, the distribution of  $\mathcal{V}_\tau^{-1/2} / \mathcal{W}_\tau$  converges to the standard normal distribution. On the other hand, Figure 3 and 4 show that if we generate data that follows Model (2), then only the log-rank statistic with CEM works, as well as if data follows Model (1). Figure 5 shows the result of the second experiment. As the sample size increases, the histogram moves away from the origin. This result demonstrates the consistency of the log-rank test statistic with CEM. We can conclude that the log-rank test with CEM performs well compared to the log-rank test with IPTW if the sample size of the control group is large enough, even if the relationship between the covariates and the treatment assignment is unknown.




Figure 1: Histogram and Q-Q plot of  $\nu_\tau^{-1/2} \mathcal{W}_\tau$  with IPTW. Data is generated under the null hypothesis by using Model (1).




Figure 2: Histogram and Q-Q plot of  $\nu_\tau^{-1/2} \mathcal{W}_\tau$  with CEM. Data is generated under the null hypothesis by using Model (1).




Figure 3: Histogram and Q-Q plot of  $\nu_\tau^{-1/2} \mathcal{W}_\tau$  with IPTW. Data is generated under the null hypothesis by using Model (2).




Figure 4: Histogram and Q-Q plot of  $\mathcal{V}_\tau^{-1/2}\mathcal{W}_\tau$  with CEM. Data is generated under the null hypothesis by using Model (2).




Figure 5: Histograms of  $\mathcal{V}_\tau^{-1/2}\mathcal{W}_\tau$  with CEM. Data is generated under the alternative hypothesis by using Model (2). The orange line represents the 5% quantile of the standard normal distribution.

## 7 Proof of Theorem 3.1

For  $i \in \mathbb{G}$ , define the process  $A^i = (A_t^i)_{t \in [0, \tau]}$  by

$$A_t^i = \int_0^t Y_s^i \Lambda^i(ds | X^i, Z^i).$$

**Proposition 7.1.** *Suppose that (3.2) of [A1] is satisfied. Then  $A^i$  is a (unique) compensator of  $N^i$  on  $[0, \tau]$  with respect to  $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, \tau]}$ .*

Let  $M_t^i = N_t^i - A_t^i$  for  $t \in [0, \tau]$  and  $i \in \mathbb{G}$ .

**Proposition 7.2.** *Suppose that [C] and (3.2) of [A1] are satisfied. Then  $M^i$  is a square-integrable  $(\mathcal{F}_t)_{t \in [0, \tau]}$ -martingale for each  $i \in \mathbb{G}$ , and the predictable quadratic covariation process between  $M^i$  and  $M^{i'}$  is  $\langle M^i, M^{i'} \rangle = 1_{\{i=i'\}} A^i$  for  $i, i' \in \mathbb{G}$ .*

See appendix for the proof of Proposition 7.1 and 7.2. Let

$$\mathcal{M}_t^z = \sum_{i \in G^z} \int_0^t K_s^n (\bar{\mathcal{Y}}_s^z)^- w_s^i dM_s^i,$$

and

$$\mathcal{E}_t^z = \int_0^t K_s^n (\bar{\mathcal{Y}}_s^z)^- \sum_{i \in G^z} w_s^i Y_s^i \Lambda(ds | X^i, z).$$

Obviously,

$$\mathcal{E}_t^z = \int_0^t K_s^n (\bar{\mathcal{Y}}_s^z)^- \sum_{i \in G^z} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} w_s^i Y_s^i \Lambda(ds | X^i, z).$$

Then, it holds that

$$\mathcal{W}_t = \mathcal{M}_t^1 - \mathcal{M}_t^0 + \mathcal{E}_t^1 - \mathcal{E}_t^0. \quad (7.1)$$

We will prove  $\mathcal{M}^1$  converges in distribution to Gaussian process in  $D[0, \tau]$  and  $\mathcal{M}^0$  and  $\mathcal{E}^1 - \mathcal{E}^0$  converges to 0 in probability under conditions [C], [A1]-[A6] and  $[H_0]$ .

For  $z \in \{0, 1\}$  and  $i \in \mathbb{G}$ , define the process  $H^{z,i} = (H_s^{z,i})_{s \in \mathbb{R}_+}$  by

$$H_s^{z,i} = K_s^n (\bar{\mathcal{Y}}_s^z)^- w_s^i Y_s^i \quad (s \in \mathbb{R}_+) \quad (7.2)$$

for  $z \in \{0, 1\}$  and  $i \in \mathbb{G}$ . Define  $\dot{\mathcal{E}}_t^z$  and  $\ddot{\mathcal{E}}_t^z$  by

$$\dot{\mathcal{E}}_t^z = \int_0^t \sum_{i \in G^z} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} H_s^{z,i} \Lambda(ds | X_a^n, z) \quad (t \in \mathbb{R}_+)$$

and

$$\ddot{\mathcal{E}}_t^z = \int_0^t \sum_{i \in G^z} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} H_s^{z,i} \{ \Lambda(ds | X^i, z) - \Lambda(ds | X_a^n, z) \} \quad (t \in \mathbb{R}_+),$$

respectively, where  $X_a^n$  is a random variable satisfying  $X_a^n \in a$ . We have  $\mathcal{E}_t^z = \dot{\mathcal{E}}_t^z + \ddot{\mathcal{E}}_t^z$ .

Define the event  $\Omega^n$  by

$$\Omega^n = \left\{ \inf_{i \in G^1} \sum_{j \in G^0} 1_{\{X^j \in a^i\}} Y_\tau^j > 0 \right\}. \quad (7.3)$$

Condition [A3] is equivalent to  $P(\Omega^n) \rightarrow 1$  as  $n \rightarrow \infty$ .

**Lemma 7.3.** (a) On the event  $\Omega^n$ ,

$$\bar{\mathcal{Y}}_s^0 = \bar{\mathcal{Y}}_s^1 \quad (s \in [0, \tau]). \quad (7.4)$$

In particular,  $\bar{\mathcal{Y}}_0^0 = \bar{\mathcal{Y}}_0^1 = n_1$  on  $\Omega^n$ .

(b) On the event  $\Omega^n$ , the functions  $s \mapsto \bar{\mathcal{Y}}_s^z$  are non increasing for  $z \in \{0, 1\}$ .

(c) On the event  $\Omega^n$ ,

$$K_s^n = 2^{-1/2}(\bar{\mathcal{Y}}_0^1)^{-1/2}\bar{\mathcal{Y}}_s^1 W_s^n \quad (s \in [0, \tau]).$$

*Proof.* Since

$$1_{\{X^{i'} \in a^i\}} = 1_{\{a^i = a^{i'}\}} = 1_{\{X^i \in a^{i'}\}}$$

for  $i, i' \in \mathbb{G}$  and

$$\sum_{j \in G^0} 1_{\{X^j \in a^i\}} Y_\tau^j > 0 \quad (\forall \omega \in \Omega^n, i \in G^1),$$

it follows that

$$\begin{aligned} \bar{\mathcal{Y}}_s^0 1_{\Omega^n} &= \sum_{i \in G^0} w_s^i Y_s^i 1_{\Omega^n} \\ &= \sum_{i \in G^0} \left( \sum_{j \in G^0} 1_{\{X^j \in a^i\}} Y_s^j \right)^- \sum_{i' \in G^1} 1_{\{X^{i'} \in a^i\}} Y_s^{i'} Y_s^i 1_{\Omega^n} \\ &= \sum_{i' \in G^1} \sum_{i \in G^0} \left( \sum_{j \in G^0} 1_{\{X^j \in a^i\}} Y_s^j \right)^- 1_{\{a^i = a^{i'}\}} Y_s^{i'} Y_s^i 1_{\Omega^n} \\ &= \sum_{i' \in G^1} \left( \sum_{j \in G^0} 1_{\{X^j \in a^{i'}\}} Y_s^j \right)^- \sum_{i \in G^0} 1_{\{X^i \in a^{i'}\}} Y_s^i Y_s^{i'} 1_{\Omega^n} \\ &= \sum_{i' \in G^1} Y_s^{i'} 1_{\Omega^n} \\ &= \bar{\mathcal{Y}}_s^1 1_{\Omega^n} \end{aligned}$$

for all  $s \in [0, \tau]$ . Thus, we proved (a). We also have, on  $\Omega^n$ ,  $\bar{\mathcal{Y}}_s^0 = \bar{\mathcal{Y}}_s^1 = \sum_{i \in G^1} Y_s^i$ , and the last function is nonincreasing, which shows (b). Since  $\bar{\mathcal{Y}}_s^0 = \bar{\mathcal{Y}}_s^1$  for  $s \in [0, \tau]$  on  $\Omega^n$  by (a),

$$\begin{aligned} K_s^n &= \{(\bar{\mathcal{Y}}_0^1 + \bar{\mathcal{Y}}_0^0)(\bar{\mathcal{Y}}_0^1 \bar{\mathcal{Y}}_0^0)^-\}^{1/2} (\bar{\mathcal{Y}}_s^1 + \bar{\mathcal{Y}}_s^0)^- \bar{\mathcal{Y}}_s^1 \bar{\mathcal{Y}}_s^0 W_s^n \\ &= 2^{-1/2} (\bar{\mathcal{Y}}_0^1)^{-1/2} \bar{\mathcal{Y}}_s^1 W_s^n \quad \text{on } \Omega^n, \end{aligned}$$

that is (c). □

**Lemma 7.4.** Under  $[H_0]$ ,  $\hat{\mathcal{E}}_t^1 1_{\Omega^n} = \hat{\mathcal{E}}_t^0 1_{\Omega^n}$  for all  $t \in [0, \tau]$  a.s.

*Proof.* Fix any  $s \in [0, \tau]$  and Borel set  $B \in \mathbb{B}([0, \tau])$ . Since

$$1_{\{X^{i'} \in a^i\}} = 1_{\{a^i = a^{i'}\}} = 1_{\{X^i \in a^{i'}\}}$$

for  $i, i' \in \mathbb{G}$ , we can calculate as follows on  $\Omega^n$ .

$$\begin{aligned}
& \sum_{i \in G^0} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} w_s^i Y_s^i \Lambda(B \mid X_a^n, 0) \\
&= \sum_{i \in G^0} \left( \sum_{j \in G^0} 1_{\{X^j \in a^i\}} Y_s^j \right)^{-} \sum_{i' \in G^1} 1_{\{X^{i'} \in a^i\}} Y_s^{i'} Y_s^i \Lambda(B \mid X_{a^i}^n, 0) \\
&= \sum_{i \in G^0} \sum_{i' \in G^1} \left( \sum_{j \in G^0} 1_{\{X^j \in a^i\}} Y_s^j \right)^{-} 1_{\{a^{i'} = a^i\}} Y_s^{i'} Y_s^i \Lambda(B \mid X_{a^i}^n, 0) \\
&= \sum_{i' \in G^1} \left( \sum_{j \in G^0} 1_{\{X^j \in a^{i'}\}} Y_s^j \right)^{-} \sum_{i \in G^0} 1_{\{X^i \in a^{i'}\}} Y_s^i Y_s^{i'} \Lambda(B \mid X_{a^{i'}}^n, 0) \\
&= \sum_{i' \in G^1} Y_s^{i'} \Lambda(B \mid X_{a^{i'}}^n, 0) \\
&= \sum_{i' \in G^1} \sum_{a \in \mathbb{A}} 1_{\{X^{i'} \in a\}} w_s^{i'} Y_s^{i'} \Lambda(B \mid X_a^n, 0). \tag{7.5}
\end{aligned}$$

We use  $[H_0]$ , (7.4) of Lemma 7.3, and (7.5), to show

$$\begin{aligned}
& 1_{\Omega^n} \sum_{i \in G^0} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} H_s^{0,i} \Lambda(B \mid X_a^n, 0) \\
&= 1_{\Omega^n} K_s^n (\overline{\mathcal{Y}}_s^0)^{-} \sum_{i \in G^0} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} w_s^i Y_s^i \Lambda(B \mid X_a^n, 1) \\
&= 1_{\Omega^n} K_s^n (\overline{\mathcal{Y}}_s^1)^{-} \sum_{i \in G^1} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} w_s^i Y_s^i \Lambda(B \mid X_a^n, 1).
\end{aligned}$$

Consequently, we proved the assertion of the lemma.  $\square$

**Lemma 7.5.** *If conditions [A1], [A3], [A5], and [A6] are satisfied, then*

$$\sup_{t \in [0, \tau]} |\check{\mathcal{E}}_t^z| = O_p(\sqrt{n_1} d_n) = o_p(1)$$

as  $n \rightarrow \infty$ .

*Proof.* Let  $z \in \{0, 1\}$  and  $t \in [0, \tau]$ . From Lemma 7.3 (c),

$$\begin{aligned}
|H_s^{z,i}| &= |K_s^n (\overline{\mathcal{Y}}_s^z)^{-} w_s^i Y_s^i| \\
&= 2^{-1/2} (\overline{\mathcal{Y}}_0^1)^{-1/2} |W_s^n| w_s^i Y_s^i \quad \text{on } \Omega^n
\end{aligned}$$

for  $s \in [0, t]$  and  $i \in \mathbb{G}$ . Then, on  $\Omega^n$ ,

$$\begin{aligned}
|\tilde{\mathcal{E}}_t^z| &= \left| \int_0^t \sum_{i \in G^z} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} H_s^{z,i} \{p(s | X^i, z) - p(s | X_a^n, z)\} \rho(ds) \right| \\
&\leq \int_0^t \sum_{i \in G^z} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} |H_s^{z,i}| \rho(ds) L d_n \quad (\because [A1]) \\
&\leq L d_n \int_0^t \sum_{i \in G^z} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} 2^{-1/2} (\bar{\mathcal{Y}}_0^1)^{-1/2} |W_s^n| w_s^i Y_s^i \rho(ds) \\
&\leq L d_n \sup_{r \in [0, \tau]} |W_r^n| \int_0^t (\bar{\mathcal{Y}}_0^1)^{-1/2} \sum_{i \in G^z} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} w_s^i Y_s^i \rho(ds) \\
&\leq L d_n \sup_{r \in [0, \tau]} |W_r^n| \int_0^\tau (\bar{\mathcal{Y}}_0^1)^{-1/2} \bar{\mathcal{Y}}_s^1 \rho(ds) \\
&\leq n_1^{1/2} d_n L \sup_{r \in [0, \tau]} |W_r^n| \rho([0, \tau]),
\end{aligned}$$

since  $(\bar{\mathcal{Y}}_0^1)^{-1/2} \bar{\mathcal{Y}}_s^1 \leq (\bar{\mathcal{Y}}_0^1)^{1/2} \leq n_1$ . Remark that  $s \mapsto \bar{\mathcal{Y}}_s^1$  is nonincreasing according to Lemma 7.3 (b). Consequently, we obtain the desired estimate since  $P(\Omega^n) \rightarrow 1$  by [A3] and  $\sup_{r \in [0, \tau]} |W_r^n| = O_p(1)$  by [A6].  $\square$

Next, we calculate  $\langle \mathcal{M}^z \rangle$  to prove asymptotic normality of  $\mathcal{M}^z$  by applying Theorem 11.3.

**Lemma 7.6.** *Suppose that [A1], [A2], [A4] and [A6] are fulfilled. Then*

$$\langle \mathcal{M}^0 \rangle_t \xrightarrow{p} 0 \quad (7.6)$$

and

$$\langle \mathcal{M}^1 \rangle_t \xrightarrow{p} \frac{1}{2} \int_{[0, t]} (W_s^\infty)^2 q_s \rho(ds) \quad (7.7)$$

as  $n \rightarrow \infty$  for every  $t \in [0, \tau]$ .

*Proof.* Let  $t \in [0, \tau]$ . Since [A3] implies  $P(\Omega^n) \rightarrow 1$  as  $n \rightarrow \infty$ , it suffices to show that  $\langle \mathcal{M}^0 \rangle_t 1_{\Omega^n} \rightarrow 0$  and  $\langle \mathcal{M}^1 \rangle_t 1_{\Omega^n} \rightarrow \frac{1}{2} \int_{[0, t]} (W_s^\infty)^2 q_s \rho(ds)$ . On  $\Omega^n$ , we have

$$\begin{aligned}
\langle \mathcal{M}^1 \rangle_t &= \sum_{i \in G^1} \int_0^t \{K_s^n (\bar{\mathcal{Y}}_s^1)^{-w_s^i}\}^2 d\langle M^i \rangle_s \\
&= \sum_{i \in G^1} \int_0^t \{K_s^n (\bar{\mathcal{Y}}_s^1)^{-w_s^i}\}^2 Y_s^i \Lambda^i(ds | X^i, Z^i) \\
&= \sum_{i \in G^1} \int_0^t 2^{-1} n_1^- (W_s^n)^2 Y_s^i \Lambda^i(ds | X^i, Z^i) \quad (\because \text{Lemma 7.3}) \\
&= 2^{-1} \int_0^t (W_s^n)^2 n_1^- \sum_{i \in G^1} Y_s^i p(s | X^i, Z^i) \rho(ds).
\end{aligned}$$



Then, the convergence (7.7) follows from [A6] and [A2]. Similarly, On  $\Omega^n$ ,

$$\begin{aligned}
\langle \mathcal{M}^0 \rangle_t &= \sum_{i \in G^0} \int_0^t \{K_s^n(\bar{\mathcal{Y}}_s^1)^- w_s^i\}^2 d\langle M^i \rangle_s \\
&= \sum_{i \in G^0} \int_0^t \{K_s^n(\bar{\mathcal{Y}}_s^1)^- w_s^i\}^2 Y_s^i \Lambda^i(ds | X^i, Z^i) \\
&= n_1^- \sum_{i \in G^0} \int_0^t 2^{-1} (W_s^n)^2 Y_s^i (w_s^i)^2 \Lambda^i(ds | X^i, Z^i) \quad (\because \text{Lemma 7.3}) \\
&\leq \sup_{r \in [0, \tau]} (W_r^n)^2 \cdot \sup_{r \in [0, \tau], j \in G^0} w_r^j \cdot n_1^- \sum_{i \in G^0} \int_0^t 2^{-1} Y_s^i w_s^i \Lambda^i(ds | X^i, Z^i) \\
&= \sup_{r \in [0, \tau]} (W_r^n)^2 \cdot \sup_{r \in [0, \tau], j \in G^0} w_r^j \cdot n_1^- \sum_{i \in G^0} \int_0^t 2^{-1} Y_s^i w_s^i p(s | X^i, Z^i) \rho(ds) \\
&\leq \sup_{r \in [0, \tau]} (W_r^n)^2 \cdot \sup_{r \in [0, \tau], j \in G^0} w_r^j \cdot n_1^- \int_0^t 2^{-1} \sum_{i \in G^0} Y_s^i w_s^i \|p\|_\infty \rho(ds) \\
&= \sup_{r \in [0, \tau]} (W_r^n)^2 \cdot \sup_{r \in [0, \tau], j \in G^0} w_r^j \cdot n_1^- \int_0^t 2^{-1} \bar{\mathcal{Y}}_s^1 \|p\|_\infty \rho(ds) \\
&\leq \sup_{r \in [0, \tau]} (W_r^n)^2 \cdot \sup_{r \in [0, \tau], i \in G^0} w_r^i \cdot 2^{-1} \rho([0, \tau]) \|p\|_\infty.
\end{aligned}$$

By [A1], [A4] and [A6], we obtain (7.7).  $\square$

**Lemma 7.7.** *Suppose that [C] and (3.2) of [A1] are satisfied. Then*

$$P(\Delta N_t^i = \Delta N_t^j \geq 1 \text{ for some } t \in [0, \tau]) = 0 \quad (7.8)$$

for  $i \neq j$ .

*Proof.* The process  $[M^i, M^j] - \langle M^i, M^j \rangle \in \mathcal{L}$ , the set of local martingales with the initial value 0. Suppose that  $i \neq j$ . Then  $R := [M^i, M^j] \in \mathcal{L}$  by Proposition 7.2. On the other hand, the local martingale  $R$  is purely discontinuous because it has finite variation, i.e., the continuous martingale part of  $R$  is null. Therefore, by Theorem 4.52 in Jacod and Shiryaev (2013),  $R = \sum_{s \leq \cdot} \Delta M_s^i \Delta M_s^j = \sum_{s \leq \cdot} \Delta N_s^i \Delta N_s^j$ , hence  $0 \leq R \leq 1$ , in particular,  $R$  is a martingale. Thus, for any  $t \in \mathbb{R}_+$ ,  $0 = E[R_0] = E[R_t] = E[\sum_{s \leq t} \Delta N_s^i \Delta N_s^j]$ , which proves (7.8).  $\square$

*Proof of Theorem 3.1.* We obtain the convergence  $\sup_{t \in [0, \tau]} |\mathcal{M}_t^0| \xrightarrow{p} 0$  as  $n \rightarrow \infty$  by Lengart's inequality and (7.6) of Lemma 7.6. Let  $\epsilon > 0$ . Set  $\mathcal{A}_t = \sum_{s \leq t} |\Delta \mathcal{M}_s^1|^2 1_{\{|\Delta \mathcal{M}_s^1| > \epsilon\}}$  for  $t \in [0, \tau]$ . We will show that  $\mathcal{A}_\tau^p \xrightarrow{p} 0$  as  $n \rightarrow \infty$ , where  $\mathcal{A}^p$  is the compensator of  $\mathcal{A}$ . Let  $t \in [0, \tau]$ . Since  $A^i$  is continuous, we have

$$\begin{aligned}
\mathcal{A}_t &= \sum_{s \leq t} \left( \sum_{i \in G^1} K_s^n(\bar{\mathcal{Y}}_s^1)^- \Delta M_s^i \right)^2 1_{\{|\sum_{j \in G^1} K_s^n(\bar{\mathcal{Y}}_s^1)^- \Delta M_s^j| > \epsilon\}} \\
&= \sum_{s \leq t} \left( \sum_{i \in G^1} K_s^n(\bar{\mathcal{Y}}_s^1)^- \Delta N_s^i \right)^2 1_{\{|\sum_{j \in G^1} K_s^n(\bar{\mathcal{Y}}_s^1)^- \Delta N_s^j| > \epsilon\}}.
\end{aligned}$$

By Lemma 7.7,

$$\begin{aligned}
& \sum_{s \leq t} \left( \sum_{i \in G^1} K_s^n (\bar{\mathcal{Y}}_s^1)^- \Delta N_s^i \right)^2 1_{\{|\sum_{j \in G^1} K_s^n (\bar{\mathcal{Y}}_s^1)^- \Delta N_s^j| > \epsilon\}} \\
&= \sum_{s \leq t} \sum_{i \in G^1} \left( K_s^n (\bar{\mathcal{Y}}_s^1)^- \Delta N_s^i \right)^2 1_{\{|\sum_{j \in G^1} K_s^n (\bar{\mathcal{Y}}_s^1)^- \Delta N_s^j| > \epsilon\}} \\
&= \sum_{s \leq t} \sum_{i \in G^1} \left( K_s^n (\bar{\mathcal{Y}}_s^1)^- \Delta N_s^i \right)^2 1_{\{|K_s^n (\bar{\mathcal{Y}}_s^1)^- \Delta N_s^i| > \epsilon\}} \\
&= \sum_{s \leq t} \sum_{i \in G^1} \left( K_s^n (\bar{\mathcal{Y}}_s^1)^- \right)^2 1_{\{|K_s^n (\bar{\mathcal{Y}}_s^1)^-| > \epsilon\}} \Delta N_s^i \\
&= \sum_{i \in G^1} \int_0^t \left( K_s^n (\bar{\mathcal{Y}}_s^1)^- \right)^2 1_{\{|K_s^n (\bar{\mathcal{Y}}_s^1)^-| > \epsilon\}} dN_s^i.
\end{aligned}$$

Therefore, it holds that

$$\mathcal{A}_t^p = \sum_{i \in G^1} \int_0^t \left( K_s^n (\bar{\mathcal{Y}}_s^1)^- \right)^2 1_{\{|K_s^n (\bar{\mathcal{Y}}_s^1)^-| > \epsilon\}} dA_s^i.$$

By Lemma 7.3 (c), we have

$$\begin{aligned}
\mathcal{A}_\tau^p 1_{\Omega^n} &\leq \sum_{i \in G^1} \int_0^\tau (2n_1)^- (W_s^n)^2 1_{\{(2n_1)^- (W_s^n)^2 > \epsilon^2\}} dA_s^i 1_{\Omega^n} \\
&\leq \sum_{i \in G^1} \int_0^\tau (4n_1^2 \epsilon^2)^- (W_s^n)^4 1_{\{(2n_1)^- (W_s^n)^2 > \epsilon^2\}} dA_s^i 1_{\Omega^n} \\
&\leq \sup_{s \in [0, \tau]} (W_s^n)^4 \cdot (4n_1^2 \epsilon^2)^- \sum_{i \in G^1} A_\tau^i.
\end{aligned}$$

Conditions [A5] and [A6] imply that  $\sup_{s \in [0, \tau]} (W_s^n)^4 \cdot n_1^- \rightarrow^p 0$ . Moreover,  $\{n_1^- \sum_{i \in G^1} A_\tau^i\}_{n \in \mathbb{N}}$  is tight since

$$E \left[ n_1^- \sum_{i \in G^1} A_\tau^i \right] = E \left[ n_1^- \sum_{i \in G^1} E[A_\tau^i | \mathcal{F}_0] \right] = E \left[ n_1^- \sum_{i \in G^1} E[N_\tau^i | \mathcal{F}_0] \right] \leq 1.$$

Thus, we have  $\mathcal{A}_\tau^p 1_{\Omega^n} \rightarrow^p 0$ . By [A3], we obtain  $\mathcal{A}_\tau^p \rightarrow^p 0$ . From this convergence together with (7.7) of Lemma 7.6, we apply Rebolledo's martingale central limit theorem (Theorem 11.3) to obtain the convergence  $\mathcal{M}^1 \rightarrow^d \mathbb{W}$ . Since the sequences  $\mathcal{M}^0$  and  $\mathcal{E}^1 - \mathcal{E}^0$  are C-tight by Lemmas 7.4 and 7.5, the decomposition (7.1) derives the convergence  $\mathcal{W} \rightarrow^d \mathbb{W}$  in  $D([0, \tau])$ .  $\square$

## 8 Proof of Proposition 3.2

Since  $W_s^\infty$  is bounded, [A2] implies

$$\left| (2n_1)^- \sum_{i \in G^1} \int_0^\tau (W_s^\infty)^2 dA_s^i - \text{Var}[\mathbb{W}_\tau] \right|$$

$$\begin{aligned}
&\leq \frac{1}{2} \int_0^\tau (W_s^\infty)^2 \left| n_1^- \sum_{i \in G^1} Y_s^i p(s | X^i, Z^i) - q_s \right| \rho(ds) \\
&\leq \frac{1}{2} \sup_{s \in [0, \tau]} (W_s^\infty)^2 \cdot \sup_{s \in [0, \tau]} \left| n_1^- \sum_{i \in G^1} Y_s^i p(s | X^i, Z^i) - q_s \right| \cdot \rho([0, \tau]) \\
&\xrightarrow{p} 0.
\end{aligned} \tag{8.1}$$

We have

$$\begin{aligned}
&\left| (2n_1)^- \sum_{i \in G^1} \int_0^\tau (W_s^n)^2 dA_s^i - (2n_1)^- \sum_{i \in G^1} \int_0^\tau (W_s^\infty)^2 dA_s^i \right| \\
&\leq \sup_{s \in [0, \tau]} |(W_s^n)^2 - (W_s^\infty)^2| \cdot (2n_1)^- \sum_{i \in G^1} A_\tau^i \\
&\xrightarrow{p} 0
\end{aligned} \tag{8.2}$$

where we used [A6] and the estimate  $A_\tau^i \leq \|p\|_\infty \rho([0, \tau])$ . Furthermore, using the Lenglart inequality to the locally square-integrable martingale, we see

$$\mathcal{V}_\tau - (2n_1)^- \sum_{i \in G^1} \int_0^\tau (W_s^n)^2 dA_s^i = (2n_1)^- \sum_{i \in G^1} \int_0^\tau (W_s^n)^2 dM_s^i \xrightarrow{p} 0 \tag{8.3}$$

since

$$(n_1^-)^2 \sum_{i \in G^1} \int_0^\tau (W_s^n)^4 dN_s^i \leq n_1^- \sup_{s \in [0, \tau]} |W_s^n|^4 \xrightarrow{p} 0.$$

It follows from equations (8.1), (8.2) and (8.3) that  $\mathcal{V}_\tau \xrightarrow{p} \text{Var}[W_\tau]$ .  $\square$

## 9 Proof of Theorem 3.3

We have shown that

$$\begin{aligned}
\dot{\mathcal{E}}_\tau^1 &= \int_0^\tau \sum_{i \in G^1} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} H_s^{1,i} \Lambda(ds | X_a^n, 1) \\
&= \int_0^\tau \sum_{i \in G^1} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} H_s^{1,i} \Lambda(ds | X^i, 1) + o_p(1)
\end{aligned} \tag{9.1}$$

in Lemma 7.5. By (7.5) of Lemma 7.4 and the proof of Lemma 7.5, it holds that

$$\begin{aligned}
\dot{\mathcal{E}}_t^0 &= \int_0^\tau \sum_{i \in G^0} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} H_s^{0,i} \Lambda(ds | X_a^n, 0) \\
&= \int_0^\tau \sum_{i \in G^1} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} H_s^{1,i} \Lambda(ds | X_a^n, 0) + o_p(1) \\
&= \int_0^\tau \sum_{i \in G^1} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} H_s^{1,i} \Lambda(ds | X^i, 0) + o_p(1).
\end{aligned} \tag{9.2}$$

Equations (9.1), (9.2) and Lemma 7.3 implies

$$\begin{aligned}
\dot{\mathcal{E}}_\tau^1 - \dot{\mathcal{E}}_\tau^0 &= \int_0^\tau \sum_{i \in G^1} \sum_{a \in \mathbb{A}} 1_{\{X^i \in a\}} H_s^{1,i} \{\Lambda(ds | X^i, 1) - \Lambda(ds | X^i, 0)\} + o_p(1) \\
&= \int_0^\tau \sum_{i \in G^1} K_s^n (\bar{\mathcal{Y}}_s^1)^{-} Y_s^i \{\Lambda(ds | X^i, 1) - \Lambda(ds | X^i, 0)\} + o_p(1) \\
&= \int_0^\tau W_s^n 2^{-1/2} n_1^{-1/2} \sum_{i \in G^1} Y_s^i \{p(s | X^i, 1) - p(s | X^i, 0)\} \rho(ds) + o_p(1).
\end{aligned}$$

Therefore, by [A7] and [H1], we have

$$\begin{aligned}
(n_1^-)^{1/2} (\dot{\mathcal{E}}_\tau^1 - \dot{\mathcal{E}}_\tau^0) &= \int_0^\tau W_s^n 2^{-1/2} n_1^- \sum_{i \in G^1} Y_s^i \{p(s | X^i, 1) - p(s | X^i, 0)\} \rho(ds) 1_{\Omega^n} + o_p(1) \\
&\leq \sup_{s \in B, x \in \mathcal{X}} W_s^n \{p(s | x, 1) - p(s | x, 0)\} \cdot n_1^- \sum_{i \in G^1} Y_\tau^i \cdot \rho(B) + o_p(1) \\
&\xrightarrow{P} \sup_{s \in B, x \in \mathcal{X}} W_s^\infty \{p(s | x, 1) - p(s | x, 0)\} \cdot \tilde{q}_\tau \cdot \rho(B) \\
&< 0.
\end{aligned}$$

Since  $\ddot{\mathcal{E}}_\tau^z = o_p(1)$  ( $z = 0, 1$ ) by Lemma 7.5 and  $\mathcal{M}_\tau^z = O_p(1)$  ( $z = 0, 1$ ) by the proof of Theorem 3.1, it holds that  $P(\mathcal{W}_\tau < -R) \rightarrow 1$  as  $n \rightarrow \infty$  for all  $R > 0$ .  $\square$

## 10 Proof of Theorem 4.2

By Theorem 3.1, Proposition 3.2 and Theorem 3.3, if [C] and [A1]-[A7] are fulfilled, then (1)-(3) hold. Thus, it is enough to show that [A2]-[A5] and [A7].

First, we show [A3] and [A4] from [B1]-[B3] and [B5]. For all  $a \in \mathbb{A}$ , we have

$$\begin{aligned}
E \left[ \sum_{i \in G^0} 1_{\{X^i \in a\}} Y_\tau^i \right] &= nE[1_{\{X^1 \in a, Z^1=0\}} Y_\tau^1] \\
&= n \int_a E[Y_\tau^1 | X^1 = x, Z^1 = 0] P^{X^1}(dx | Z^1 = 0) P(Z^1 = 0) \\
&\geq nP(Z^1 = 0) \cdot \operatorname{ess\,inf}_{x \in \mathcal{X}} (f(x | 0) E[Y_\tau^1 | X^1 = x, Z^1 = 0]) \cdot \inf_{a \in \mathbb{A}} \nu(a).
\end{aligned}$$

By [B2], [B3] and [B5], there exists  $C > 0$  such that

$$\inf_{a \in \mathbb{A}} E \left[ \sum_{i \in G^0} 1_{\{X^i \in a\}} Y_\tau^i \right] \geq Cn^{1-\bar{d}\theta} \tag{10.1}$$

for all  $n \in \mathbb{N}$ . In the same way, we have

$$\begin{aligned}
E \left[ \sum_{i \in \mathbb{G}^1} 1_{\{X^i \in a\}} \right] &= nP(X^1 \in a, Z^1 = 1) \\
&= nP(X^1 \in a \mid Z^1 = 1)P(Z^1 = 1) \\
&= nP(Z^1 = 1) \int_a f(x \mid 1) d\nu(dx) \\
&\leq nP(Z^1 = 1) \cdot \operatorname{ess\,sup}_{x \in \mathcal{X}} f(x \mid 1) \cdot \sup_{a \in \mathbb{A}} \nu(a).
\end{aligned}$$

Thus, from [B2], [B3] and [B5], there exists a constant  $C' > 0$  such that

$$\sup_{a \in \mathbb{A}} E \left[ \sum_{i \in \mathbb{G}^1} 1_{\{X^i \in a\}} \right] \leq C' n^{\beta - \bar{d}\theta} \quad (10.2)$$

for all  $n \in \mathbb{N}$ . By [B1], it holds that  $1 - \bar{d}\theta - \beta/2 > (1 - \beta)/2 > 0$ . Therefore, we can take  $\delta \in (0, 1 - \bar{d}\theta - \beta/2)$ . We define the events  $\Omega^{n,0}$  and  $\Omega^{n,1}$  as follows.

$$\begin{aligned}
\Omega^{n,0} &= \left\{ \inf_{a \in \mathbb{A}} \sum_{i \in \mathbb{G}^0} 1_{\{X^i \in a\}} Y_\tau^i > \frac{C}{2} n^{1 - \bar{d}\theta} \right\}, \\
\Omega^{n,1} &= \left\{ \sup_{a \in \mathbb{A}} \sum_{i \in \mathbb{G}^1} 1_{\{X^i \in a\}} < 2C' n^{1 - \bar{d}\theta - \delta} \right\}.
\end{aligned}$$

Let  $\Omega^n$  be the event defined in (7.3). Since

$$\Omega^{n,0} \subset \left\{ \inf_{a \in \mathbb{A}} \sum_{i \in \mathbb{G}^0} 1_{\{X^i \in a\}} Y_\tau^i > 0 \right\} \subset \left\{ \inf_{j \in \mathbb{G}^1} \sum_{i \in \mathbb{G}^0} 1_{\{X^i \in a_j\}} Y_\tau^i > 0 \right\} = \Omega^n,$$

we can obtain [A3] if we show that  $P(\Omega^{n,0}) \rightarrow 1$ . Moreover, if we show that  $P(\Omega^{n,0} \cap \Omega^{n,1}) \rightarrow 1$ , the following estimate yields [A4].

$$\begin{aligned}
\sup_{t \in [0, \tau], i \in \mathbb{G}^0} w_t^i &\leq \sup_{a \in \mathbb{A}} \left( \sum_{i \in \mathbb{G}^1} 1_{\{X^i \in a\}} \right) \left( \sum_{i' \in \mathbb{G}^0} 1_{\{X^{i'} \in a\}} Y_\tau^{i'} \right)^{-1} \\
&\leq \frac{4C'}{C} n^{-\delta} \quad \text{on } \Omega^{n,0} \cap \Omega^{n,1}.
\end{aligned}$$

By (10.1) and [B1], we have

$$\begin{aligned}
P((\Omega^{n,0})^c) &= P \left( \inf_{a \in \mathbb{A}} \sum_{i \in \mathbb{G}^0} 1_{\{X^i \in a\}} Y_\tau^i \leq \frac{C}{2} n^{1 - \bar{d}\theta} \right) \\
&\leq P \left( \sup_{a \in \mathbb{A}} \left| \sum_{i \in \mathbb{G}^0} 1_{\{X^i \in a\}} Y_\tau^i - E \left[ \sum_{i \in \mathbb{G}^0} 1_{\{X^i \in a\}} Y_\tau^i \right] \right| \geq \frac{C}{2} n^{1 - \bar{d}\theta} \right)
\end{aligned}$$

$$\begin{aligned}
&\leq \sum_{a \in \mathbb{A}} P \left( \left| \sum_{i \in \mathbb{G}^0} 1_{\{X^i \in a\}} Y_\tau^i - E \left[ \sum_{i \in \mathbb{G}^0} 1_{\{X^i \in a\}} Y_\tau^i \right] \right| \geq \frac{C}{2} n^{1-\bar{d}\theta} \right) \\
&\leq \frac{4}{C^2} n^{-2+2\bar{d}\theta} \sum_{a \in \mathbb{A}} \text{Var} \left[ \sum_{i \in \mathbb{G}^0} 1_{\{X^i \in a\}} Y_\tau^i \right] \\
&\leq \frac{4}{C^2} n^{-1+2\bar{d}\theta} \sum_{a \in \mathbb{A}} E[1_{\{X^1 \in a, Z^1=0\}} Y_\tau^1] \\
&\leq \frac{4}{C^2} n^{-1+2\bar{d}\theta} E[1_{\{Z^1=0\}} Y_\tau^1] \\
&\rightarrow 0 \quad (n \rightarrow \infty).
\end{aligned}$$

In the same way, from (10.2) and [B1], we have

$$\begin{aligned}
P((\Omega^{n,1})^c) &= P \left( \sup_{a \in \mathbb{A}} \sum_{i \in \mathbb{G}^1} 1_{\{X^i \in a\}} \geq 2C' n^{1-\bar{d}\theta-\delta} \right) \\
&\leq P \left( \sup_{a \in \mathbb{A}} \left| \sum_{i \in \mathbb{G}^1} 1_{\{X^i \in a\}} - E \left[ \sum_{i \in \mathbb{G}^1} 1_{\{X^i \in a\}} \right] \right| \geq C' n^{1-\bar{d}\theta-\delta} \right) \\
&\leq \sum_{a \in \mathbb{A}} P \left( \left| \sum_{i \in \mathbb{G}^1} 1_{\{X^i \in a\}} - E \left[ \sum_{i \in \mathbb{G}^1} 1_{\{X^i \in a\}} \right] \right| \geq C' n^{1-\bar{d}\theta-\delta} \right) \\
&\leq \sum_{a \in \mathbb{A}} C'^{-2} n^{-2+2\bar{d}\theta+2\delta} \text{Var} \left[ \sum_{i \in \mathbb{G}^1} 1_{\{X^i \in a\}} \right] \\
&= \sum_{a \in \mathbb{A}} C'^{-2} n^{-1+2\bar{d}\theta+2\delta} \text{Var}[1_{X^1 \in a, Z^1=1}] \\
&\leq \sum_{a \in \mathbb{A}} C'^{-2} n^{-1+2\bar{d}\theta+2\delta} P(X^1 \in a, Z^1 = 1) \\
&= C^{-2} n^{-1+2\bar{d}\theta+2\delta} P(Z^1 = 1) \\
&= O(n^{-2+2\bar{d}\theta+2\delta+\beta}) \\
&\rightarrow 0.
\end{aligned}$$

Therefore, we obtain [A3] and [A4].

We recall that  $V^i = (X^i, Z^i, T^i, U^i)$  ( $i \in \mathbb{G}$ ). Let  $g: \mathbb{R}^d \times \{0, 1\} \times (0, \infty) \times (0, \infty) \rightarrow \mathbb{R}$  be a measurable function satisfying  $E[|g(V^1)|^2 | Z^1 = 1] < \infty$ . Note that  $E[|g(V^1)|^2 | Z^1 = 1]$  does not depend on  $n$  since the conditional distribution  $\mathcal{L}\{V^1 | Z^1 = 1\}$  does not depend on  $n$ . Suppose that [B1]-[B5] hold. In what follows, we consider large enough  $n \in \mathbb{N}$  such that  $P(Z^1 = 1) > 0$ . We prove the weak law of large numbers

$$\frac{1}{nP(Z^1 = 1)} \sum_{i \in \mathbb{G}^1} g(V^i) \xrightarrow{p} E[g(V^1) 1_{\{X^1 \in \mathcal{X}\}} | Z^1 = 1] \quad (10.3)$$

and

$$n_1^- \sum_{i \in \mathbb{G}^1} g(V^i) \xrightarrow{p} E[g(V^1) | X^1 \in \mathcal{X}, Z^1 = 1]. \quad (10.4)$$

On the event  $\Omega^{n,0}$ , for all  $a \in \mathbb{A}$ , there exists at least one individual  $i \in \mathbb{G}^0$  such that  $X^i \in a$ . Therefore, we have

$$\begin{aligned} G^1 &= \{i \in \mathbb{G}^1 \mid X^i \in a \text{ and } X^j \in a \text{ for some } j \in \mathbb{G}^0 \text{ and } a \in \mathbb{A}\} \\ &= \{i \in \mathbb{G}^1 \mid X^i \in \mathcal{X}\} \quad \text{on } \Omega^{n,0}. \end{aligned} \quad (10.5)$$

We have shown that if [B1]-[B3] and [B5] holds, then  $P(\Omega^{n,0}) \rightarrow 1$ . From  $P(\Omega^{n,0}) \rightarrow 1$  and (10.5), it follows that

$$\begin{aligned} \frac{1}{nP(Z^1 = 1)} \sum_{i \in \mathbb{G}^1} g(V^i) &= \frac{1}{nP(Z^1 = 1)} \sum_{i \in \mathbb{G}^1} g(V^i) 1_{\Omega^{n,0}} + o_p(1) \\ &= \frac{1}{nP(Z^1 = 1)} \sum_{i \in \mathbb{G}^1} g(V^i) 1_{\{X^i \in \mathcal{X}\}} 1_{\Omega^{n,0}} + o_p(1) \\ &= \frac{1}{nP(Z^1 = 1)} \sum_{i \in \mathbb{G}^1} g(V^i) 1_{\{X^i \in \mathcal{X}\}} + o_p(1). \end{aligned}$$

Thus, we need to show that

$$\frac{1}{nP(Z^1 = 1)} \sum_{i \in \mathbb{G}^1} g(V^i) 1_{\{X^i \in \mathcal{X}\}} \xrightarrow{p} E[g(V^1) 1_{\{X^1 \in \mathcal{X}\}} \mid Z^1 = 1]. \quad (10.6)$$

for (10.3). Note that the strong law of large numbers may not hold in (10.6) since the distribution of  $(X^i, Z^i, T^i, U^i)$  depends on  $n$ . We have

$$\begin{aligned} E \left[ \frac{1}{nP(Z^1 = 1)} \sum_{i \in \mathbb{G}^1} g(V^i) 1_{\{X^i \in \mathcal{X}\}} \right] &= \frac{1}{nP(Z^1 = 1)} E \left[ \sum_{i \in \mathbb{G}} g(V^i) 1_{\{X^i \in \mathcal{X}, Z^i = 1\}} \right] \\ &= \frac{1}{P(Z^1 = 1)} E[g(V^1) 1_{\{X^1 \in \mathcal{X}, Z^1 = 1\}}] \\ &= E[g(V^1) 1_{\{X^1 \in \mathcal{X}\}} \mid Z^1 = 1] \end{aligned}$$

and

$$\begin{aligned} \text{Var} \left[ \frac{1}{nP(Z^1 = 1)} \sum_{i \in \mathbb{G}^1} g(V^i) 1_{\{X^i \in \mathcal{X}\}} \right] &= \frac{1}{n^2 P(Z^1 = 1)^2} \text{Var} \left[ \sum_{i \in \mathbb{G}} g(V^i) 1_{\{X^i \in \mathcal{X}, Z^i = 1\}} \right] \\ &= \frac{1}{nP(Z^1 = 1)^2} \text{Var}[g(V^1) 1_{\{X^1 \in \mathcal{X}, Z^1 = 1\}}] \\ &\leq \frac{1}{nP(Z^1 = 1)^2} E[(g(V^1))^2 1_{\{X^1 \in \mathcal{X}, Z^1 = 1\}}] \\ &= \frac{1}{nP(Z^1 = 1)} E[(g(V^1))^2 1_{\{X^1 \in \mathcal{X}\}} \mid Z^1 = 1] \\ &= O_p(n^{-\beta}) \end{aligned}$$

by [B2]. Therefore, we have

$$\frac{1}{nP(Z^1 = 1)} \sum_{i \in \mathbb{G}^1} g(V^i) 1_{\{X^i \in \mathcal{X}\}} \rightarrow E[g(V^1) 1_{\{X^1 \in \mathcal{X}\}} \mid Z^1 = 1] \quad \text{in } L^2.$$

Thus, we have (10.6) and we obtain (10.3). By applying (10.3) to  $g = 1$ , we have

$$\frac{n_1}{nP(Z^1 = 1)} \rightarrow^p P(X^1 \in \mathcal{X} \mid Z^1 = 1) > 0, \quad (10.7)$$

where  $P(X^1 \in \mathcal{X} \mid Z^1 = 1) > 0$  is derived from [B4]. It follows from (10.3) and (10.7) that

$$n_1^- \sum_{i \in G^1} g(V^i) \rightarrow^p \frac{E[g(V^1)1_{\{X^1 \in \mathcal{X}\}} \mid Z^1 = 1]}{P(X^1 \in \mathcal{X} \mid Z^1 = 1)}$$

and (10.4) is proved.

We derive [A5] from [B1]-[B5]. By (10.7) and [B2], it holds that  $n_1 = O_p(n^\beta)$  and  $n_1^- = O_p(n^{-\beta})$ . Therefore, by [B5] and [B1], we have  $n_1^- = o_p(1)$  and  $n_1 d_n^2 = O_p(n^{\beta-2\theta}) = o_p(1)$ , which implies [A5].

Condition [A7] follows from [B1]-[B5]. Indeed, when  $g(x, z, t, u) = 1_{\{t \wedge u \geq \tau\}}$  ( $x \in \mathbb{R}^d, z \in \{0, 1\}, t, u \in (0, \infty)$ ), the weak law of large number (10.4) gives

$$n_1^- \sum_{i \in G^1} Y_\tau^i \rightarrow^p \tilde{q}_\tau := E[Y_\tau^1 \mid X^1 \in \mathcal{X}, Z^1 = 1].$$

By [B4], we have

$$\tilde{q}_\tau = \frac{P(\tilde{T}^1 \geq \tau, X^1 \in \mathcal{X} \mid Z^1 = 1)}{P(X \in \mathcal{X} \mid Z^1 = 1)} > 0.$$

Finally, we show [A2] under [A1'] and [B1]-[B5]. The sequence of càdlàg processes  $(\mathbb{V}_s^n := n_1^- \sum_{i \in G^1} Y_{s+}^i p(s \mid X^i, Z^i))_{n \in \mathbb{N}}$  is tight in  $D([0, \tau])$ . The proof is as follows.

For every  $\alpha \in D([0, \tau])$  and  $\delta > 0$ , we define

$$w(\alpha, \delta) = \sup_{s, t \in [0, \tau], |s-t| \leq \delta} |\alpha_s - \alpha_t|.$$

Condition [A1'] implies

$$\sup_{n \in \mathbb{N}, s \in [0, \tau]} |\mathbb{V}_s^n| \leq \|p\|_\infty = \sup_{s \in [0, \tau], x \in \mathcal{X}, z \in \{0, 1\}} p(s \mid x, z) < \infty.$$

Hence, it is enough to show that

$$\lim_{\delta \downarrow 0} \limsup_{n \rightarrow \infty} P(w(\mathbb{V}^n, \delta) \geq \epsilon) = 0 \quad \text{for all } \epsilon > 0 \quad (10.8)$$

to obtain the (C-)tightness of  $(\mathbb{V}^n)$ . Let  $\delta > 0$  and  $n \in \mathbb{N}$ . If  $s, t \in [0, \tau]$  satisfy  $s \leq t$  and  $t - s \leq \delta$ , then

$$\begin{aligned} |\mathbb{V}_s^n - \mathbb{V}_t^n| &\leq \left| n_1^- \sum_{i \in G^1} Y_{s+}^i p(s \mid X^i, Z^i) - n_1^- \sum_{i \in G^1} Y_{t+}^i p(s \mid X^i, Z^i) \right| \\ &\quad + \left| n_1^- \sum_{i \in G^1} Y_{t+}^i p(s \mid X^i, Z^i) - n_1^- \sum_{i \in G^1} Y_{t+}^i p(t \mid X^i, Z^i) \right| \\ &\leq n_1^- \sum_{i \in G^1} (Y_{s+}^i - Y_{t+}^i) \|p\|_\infty + n_1^- \sum_{i \in G^1} |(p(s \mid X^i, Z^i) - p(t \mid X^i, Z^i))| \\ &\leq \|p\|_\infty \left( n_1^- \sum_{i \in G^1} Y_{s+}^i - n_1^- \sum_{i \in G^1} Y_{t+}^i \right) + \sup_{x \in \mathcal{X}} |p(s \mid x, 1) - p(t \mid x, 1)| \\ &\leq \|p\|_\infty w \left( n_1^- \sum_{i \in G^1} Y_{+}^i, \delta \right) + \sup_{s, t \in [0, \tau], |s-t| \leq \delta, x \in \mathcal{X}} |p(s \mid x, 1) - p(t \mid x, 1)|, \end{aligned}$$



where  $Y_+^i$  denotes the process  $(Y_{s+}^i)_{s \in [0, \tau]}$ . By taking the supremum over  $s$  and  $t$ , we have

$$w(\mathbb{V}^n, \delta) \leq \|p\|_\infty w \left( n_1^- \sum_{i \in G^1} Y_+^i, \delta \right) + \sup_{s, t \in [0, \tau], |s-t| \leq \delta, x \in \mathcal{X}} |p(s | x, 1) - p(t | x, 1)|.$$

Therefore, by Proposition 3.26 in Chapter 6 of Jacod and Shiryaev (2013), we need to show that

$$\lim_{\delta \downarrow 0} \limsup_{n \rightarrow \infty} P \left( w \left( n_1^- \sum_{i \in G^1} Y_+^i, \delta \right) \geq \epsilon \right) = 0 \quad \text{for all } \epsilon > 0 \quad (10.9)$$

and

$$\lim_{\delta \downarrow 0} \sup_{s, t \in [0, \tau], |s-t| \leq \delta, x \in \mathcal{X}} |p(s | x, 1) - p(t | x, 1)| = 0. \quad (10.10)$$

to obtain (10.8).

By [A1'], there exists  $L > 0$  such that for all  $s, t \in [0, \tau]$  and  $x, y \in \mathcal{X}$

$$\begin{aligned} |p(s | x, 1) - p(t | y, 1)| &\leq |p(s | x, 1) - p(t | x, 1)| + |p(t | x, 1) - p(t | y, 1)| \\ &\leq |p(s | x, 1) - p(t | x, 1)| + L|x - y|. \end{aligned}$$

From this and [A1'], the function  $(s, x) \mapsto p(s | x, 1)$  is continuous on  $[0, \tau] \times \mathcal{X}$ . Since  $\mathcal{X}$  is compact,  $p(s | x, 1)$  is uniformly continuous on  $[0, \tau] \times \mathcal{X}$ . Therefore, (10.10) holds.

For each  $s \in [0, \tau]$ , applying (10.4) to  $g(x, z, t, u) = 1_{\{t \wedge u > s\}}$  yields

$$n_1^- \sum_{i \in G^1} Y_{s+}^i \xrightarrow{p} \tilde{q}_s := E[Y_{s+}^1 | X^1 \in \mathcal{X}, Z^1 = 1]. \quad (10.11)$$

The function  $\tilde{q}$  is continuous on  $[0, \infty)$  and it holds that  $\tilde{q}_s = E[Y_s^1 | X^1 \in \mathcal{X}, Z^1 = 1]$  for all  $s \in [0, \tau]$ . Indeed, by the dominated convergence theorem, we have

$$\begin{aligned} \tilde{q}_{s+} &= \lim_{h \downarrow 0} E[Y_{(s+h)+}^1 | X^1 \in \mathcal{X}, Z^1 = 1] \\ &= \lim_{h \downarrow 0} E[1_{\{\tilde{T}^1 > s+h\}} | X^1 \in \mathcal{X}, Z^1 = 1] \\ &= E[1_{\{\tilde{T}^1 > s\}} | X^1 \in \mathcal{X}, Z^1 = 1] \\ &= E[Y_{s+}^1 | X^1 \in \mathcal{X}, Z^1 = 1] \\ &= \tilde{q}_s \end{aligned}$$

for all  $s \in [0, \tau]$ . Thus,  $\tilde{q}$  is right-continuous on  $[0, \infty)$ . By the same way, for all  $s \in (0, \tau]$ , we have

$$\begin{aligned} \tilde{q}_{s-} &= \lim_{h \downarrow 0} E[Y_{(s-h)+}^1 | X^1 \in \mathcal{X}, Z^1 = 1] \\ &= \lim_{h \downarrow 0} E[1_{\{\tilde{T}^1 > s-h\}} | X^1 \in \mathcal{X}, Z^1 = 1] \\ &= E[1_{\{\tilde{T}^1 \geq s\}} | X^1 \in \mathcal{X}, Z^1 = 1] \\ &= E[Y_s^1 | X^1 \in \mathcal{X}, Z^1 = 1]. \end{aligned}$$

Therefore, for all  $s \in [0, \tau]$ , it follows from [B4] that

$$\Delta \tilde{q}_s = -E[1_{\{\tilde{T}^1 = s\}} | X^1 \in \mathcal{X}, Z^1 = 1] = 0$$

and hence

$$\tilde{q}_s = \tilde{q}_{s-} = E[Y_s^1 \mid X^1 \in \mathcal{X}, Z^1 = 1].$$

Since  $n_1^- \sum_{i \in G^1} Y_+^i$  is decreasing and  $\tilde{q}$  is continuous, it follows from (10.11) and Theorem 3.37 (a) in Chapter 6 of Jacod and Shiryaev (2013) that  $n_1^- \sum_{i \in G^1} Y_+^i \rightarrow^d \tilde{q}$  in  $D([0, \tau])$ . In particular,  $n_1^- \sum_{i \in G^1} Y_+^i$  is C-tight and hence (10.9) holds. The proof of the (C-)tightness of  $(\mathbb{V}^n)$  is complete.

For all  $s \in [0, \tau]$ , by (10.4) with  $g(x, z, t, u) = 1_{\{t \wedge u > s\}} p(s \mid x, z)$ , we have

$$\mathbb{V}_s^n \rightarrow^p q_s = E[Y_{s+}^1 p(s \mid X^1, Z^1) \mid X^1 \in \mathcal{X}, Z^1 = 1].$$

We can show that  $q$  is continuous on  $[0, \tau]$  and  $q_s = E[Y_s^1 p(s \mid X^1, Z^1) \mid X^1 \in \mathcal{X}, Z^1 = 1]$  in the same way as for  $\tilde{q}$ . Since  $(\mathbb{V}^n - q)$  is tight,  $\mathbb{V}^n - q \rightarrow^d 0$  in  $D([0, \tau])$  holds. By the continuous mapping theorem, we obtain

$$\sup_{s \in [0, \tau]} |\mathbb{V}_s^n - q_s| \rightarrow^p 0.$$

From this and

$$\left| n_1^- \sum_{i \in G^1} Y_s^i p(s \mid X^i, Z^i) - q_s \right| = |\mathbb{V}_{s-}^n - q_{s-}| \leq \sup_{t \in [0, \tau]} |\mathbb{V}_t^n - q_t|$$

for all  $s \in (0, \tau]$ , it holds that

$$\sup_{s \in (0, \tau]} \left| n_1^- \sum_{i \in G^1} Y_s^i p(s \mid X^i, Z^i) - q_s \right| \leq \sup_{t \in [0, \tau]} |\mathbb{V}_t^n - q_t| \rightarrow^p 0.$$

Moreover, we have

$$n_1^- \sum_{i \in G^1} Y_0^i p(0 \mid X^i, Z^i) = \mathbb{V}_0^n \rightarrow^p q_0.$$

Therefore, [A2] holds. □

## 11 Appendix

In this appendix, we prove Proposition 7.1 and 7.2. We recall the following results, for convenience of reference. For a probability space  $(\Omega, \mathcal{F}, P)$  and sub  $\sigma$ -fields  $\mathcal{A}, \mathcal{B}$  and  $\mathcal{H}$  of  $\mathcal{F}$ , we write  $\mathcal{A} \perp\!\!\!\perp_{\mathcal{H}} \mathcal{B}$  if  $\mathcal{A}$  and  $\mathcal{B}$  are  $\mathcal{H}$ -conditionally independent, that is,  $P(A \cap B \mid \mathcal{H}) = P(A \mid \mathcal{H})P(B \mid \mathcal{H})$  a.s. for all  $A \in \mathcal{A}$  and  $B \in \mathcal{B}$ .

**Lemma 11.1.** *Let  $(\Omega, \mathcal{F}, P)$  be a probability space.*

(a) *Let  $V: \Omega \rightarrow \mathbb{R}$  be an integrable random variable and let  $\mathcal{G}, \mathcal{H}$  be sub  $\sigma$ -fields of  $\mathcal{F}$ . If  $\sigma[V] \perp\!\!\!\perp_{\mathcal{H}} \mathcal{G}$ , then  $E[V \mid \mathcal{G} \vee \mathcal{H}] = E[V \mid \mathcal{H}]$ .*

(b) *Suppose that  $\mathcal{V} \perp\!\!\!\perp_{\mathcal{H}} \mathcal{G}$  for sub  $\sigma$ -fields  $\widehat{\mathcal{G}}, \mathcal{G}, \mathcal{V}, \mathcal{H}$  of  $\mathcal{F}$  such that  $\widehat{\mathcal{G}} \subset \mathcal{G}$ . Then  $\mathcal{V} \perp\!\!\!\perp_{\widehat{\mathcal{G}} \vee \mathcal{H}} \mathcal{G}$ .*

*Proof.* (a) is obvious. Indeed, for all bounded  $\mathcal{G}$ -measurable random variable  $G$  and bounded  $\mathcal{H}$ -measurable random variable  $H$ , it holds that

$$E[VGH] = E[E[VG \mid \mathcal{H}]H] = E[E[V \mid \mathcal{H}]E[G \mid \mathcal{H}]H] = E[E[V \mid \mathcal{H}]GH]$$

and we obtain (a).

We will show (b). Let  $V, G, \widehat{G}, H$  be bounded random variables measurable with respect to  $\mathcal{V}, \mathcal{G}, \widehat{\mathcal{G}}, \mathcal{H}$ , respectively. Then

$$\begin{aligned}
E[VG\widehat{G}H] &= E[E[V|\mathcal{H}]E[G\widehat{G}|\mathcal{H}]H] \quad (\because \mathcal{V} \perp_{\mathcal{H}} \mathcal{G}, \widehat{\mathcal{G}} \subset \mathcal{G}) \\
&= E[E[V|\mathcal{H}]G\widehat{G}H] \\
&= E[E[V|\widehat{\mathcal{G}} \vee \mathcal{H}]G\widehat{G}H] \quad (\because (a)) \\
&= E[E[V|\widehat{\mathcal{G}} \vee \mathcal{H}]E[G|\widehat{\mathcal{G}} \vee \mathcal{H}]\widehat{G}H]
\end{aligned}$$

and hence  $E[VG|\widehat{\mathcal{G}} \vee \mathcal{H}] = E[V|\widehat{\mathcal{G}} \vee \mathcal{H}]E[G|\widehat{\mathcal{G}} \vee \mathcal{H}]$  a.s. This proves (b).  $\square$

**Lemma 11.2.** *We consider the same random variables as section 3 and suppose that condition [C] holds. Let  $i \in \mathbb{G}$ .*

(a)  $P^{T^i}(\cdot|\mathcal{F}_0) = P^{T^i}(\cdot|X^i, Z^i)$  a.s. for a regular conditional distribution.

(b)  $T^i \perp_{\mathcal{F}_0} U^i$ .

(c) For any  $t \geq 0$ , we define sub  $\sigma$ -fields  $\mathcal{G}_t^i$  and  $\widehat{\mathcal{G}}_t^i$  of  $\mathcal{F}$  by setting  $\mathcal{G}_t^i = \sigma[N_\theta^i, \dot{N}_\theta^i | \theta \leq t]$  and  $\widehat{\mathcal{G}}_t^i = \sigma[N_\theta^j, \dot{N}_\theta^j | \theta \leq t, j \neq i]$ . Let  $0 \leq s \leq t \leq \tau$ . Then, it holds that  $\widehat{\mathcal{G}}_t^i \perp_{\mathcal{G}_s^i \vee \mathcal{F}_0} \mathcal{G}_t^i$ .

*Proof.* (a) follows from [C] (iii) and Lemma 11.1 (a). Condition (b) is the same as [C] (ii). Let  $\mathcal{V} = \widehat{\mathcal{G}}_t^i$ ,  $\mathcal{G} = \mathcal{G}_t^i$ ,  $\widehat{\mathcal{G}} = \mathcal{G}_s^i$  and  $\mathcal{H} = \mathcal{F}_0$  respectively in Lemma 11.1 (b), we obtain (c).  $\square$

*Proof of Proposition 7.1.* Suppose that  $F \in \mathcal{F}_0$ ,  $0 \leq s < t \leq \tau$  and  $p_k, q_\ell \in [0, s]$  for  $k \in \{1, \dots, K\}$  and  $\ell \in \{1, \dots, L\}$ . Let  $\bar{p} = \max_{1 \leq k \leq K} p_k$  and  $\bar{q} = \max_{1 \leq \ell \leq L} q_\ell$ .

$$\begin{aligned}
&E \left[ (N_t^i - N_s^i) \prod_{k=1}^K 1_{\{N_{p_k}^i = 0\}} \prod_{\ell=1}^L 1_{\{\dot{N}_{q_\ell}^i = 0\}} 1_F \right] \tag{11.1} \\
&= E \left[ 1_{\{s < T^i \leq t, T^i \leq U^i\}} 1_{\{T^i \wedge U^i > \bar{p}\} \cup \{T^i > U^i\}} 1_{\{T^i \wedge U^i > \bar{q}\} \cup \{T^i \leq U^i\}} 1_F \right] \\
&= E \left[ 1_{\{s < T^i \leq t, T^i \leq U^i\}} 1_F \right] \\
&= E \left[ E[1_{\{s < T^i \leq t, T^i \leq U^i\}} | \mathcal{F}_0] 1_F \right] \\
&= E \left[ \int_s^t E[1_{\{\theta \leq U^i\}} | \mathcal{F}_0] P^{T^i}(d\theta | \mathcal{F}_0) 1_F \right] \quad (\because \text{Lemma 11.2 (b)}) \\
&= E \left[ \int_s^t P(U^i \geq \theta | \mathcal{F}_0) P^{T^i}(d\theta | \mathcal{F}_0) 1_F \right].
\end{aligned}$$

On the other hand,

$$\begin{aligned}
& E \left[ (A_t^i - A_s^i) \prod_{k=1}^K 1_{\{N_{p_k}=0\}} \prod_{\ell=1}^L 1_{\{N_{q_\ell}=0\}} 1_F \right] \tag{11.2} \\
&= E \left[ \int_s^t 1_{\{T^i \wedge U^i \geq \theta\}} \frac{P^{T^i}(d\theta|\mathcal{F}_0)}{P^{T^i}([\theta, \infty)|\mathcal{F}_0)} 1_{\{T^i \wedge U^i > \bar{p}\} \cup \{T^i > U^i\}} 1_{\{T^i \wedge U^i > \bar{q}\} \cup \{T^i \leq U^i\}} 1_F \right] \\
&= E \left[ \int_s^t 1_{\{T^i \wedge U^i \geq \theta\}} \frac{P^{T^i}(d\theta|\mathcal{F}_0)}{P^{T^i}([\theta, \infty)|\mathcal{F}_0)} 1_F \right] \\
&= E \left[ \int_{\mathbb{R}_+^2} \int_s^t 1_{\{\alpha \wedge \beta \geq \theta\}} \frac{P^{T^i}(d\theta|\mathcal{F}_0)}{P^{T^i}([\theta, \infty)|\mathcal{F}_0)} P^{(T^i, U^i)}(d\alpha d\beta|\mathcal{F}_0) 1_F \right] \\
&= E \left[ \int_s^t P(T^i \wedge U^i \geq \theta|\mathcal{F}_0) \frac{P^{T^i}(d\theta|\mathcal{F}_0)}{P^{T^i}([\theta, \infty)|\mathcal{F}_0)} 1_F \right] \quad (\because \text{Fubini's theorem}) \\
&= E \left[ \int_s^t P(T^i \geq \theta|\mathcal{F}_0) P(U^i \geq \theta|\mathcal{F}_0) \frac{P^{T^i}(d\theta|\mathcal{F}_0)}{P^{T^i}([\theta, \infty)|\mathcal{F}_0)} 1_F \right] \quad (\because \text{Lemma 11.2 (b)}) \\
&= E \left[ \int_s^t P(U^i \geq \theta|\mathcal{F}_0) P^{T^i}(d\theta|\mathcal{F}_0) 1_F \right].
\end{aligned}$$

Therefore, the two expressions (11.1) and (11.2) coincide. If  $K = L = 0$  and  $F = \Omega$ , we obtain  $E[A_t^i - A_s^i] = E[N_t^i - N_s^i] < \infty$ . As a matter of fact, more generally,

$$E \left[ (N_t^i - N_s^i) \prod_{k=1}^K 1_{\{N_{p_k}=\epsilon_k\}} \prod_{\ell=1}^L 1_{\{N_{q_\ell}=\eta_\ell\}} 1_F \right] = E \left[ (A_t^i - A_s^i) \prod_{k=1}^K 1_{\{N_{p_k}=\epsilon_k\}} \prod_{\ell=1}^L 1_{\{N_{q_\ell}=\eta_\ell\}} 1_F \right]$$

for any  $\epsilon_1, \dots, \epsilon_K, \eta_1, \dots, \eta_L \in \{0, 1\}$ . Therefore,

$$E[N_t^i - N_s^i | \mathcal{G}_s^i \vee \mathcal{F}_0] = E[A_t^i - A_s^i | \mathcal{G}_s^i \vee \mathcal{F}_0] \quad a.s.$$

Consequently, Lemma 11.2 (c) and Lemma 11.1 (a) implies

$$E[N_t^i - N_s^i | \mathcal{F}_s] = E[A_t^i - A_s^i | \mathcal{F}_s] \quad a.s.$$

By definition,  $(A_t^i)_{t \in [0, \tau]}$  is càdlàg  $\mathbb{F}$ -predictable, and hence the proof is completed.  $\square$

*Proof of Proposition 7.2.* Since  $M^i$  has bounded variation and  $\Delta A^i = 0$ , we have

$$[M^i]_t = \sum_{s \leq t} (\Delta M_s^i)^2 = \sum_{s \leq t} (\Delta N_s^i)^2 = N_t^i \tag{11.3}$$

Thus,  $\langle M^i \rangle = A^i$  since  $[M^i] - \langle M^i \rangle$  is a local martingale. The local martingale  $M^i = (M_t^i)_{t \in [0, \tau]}$  is square-integrable since  $\sup_{t \in [0, \tau]} E[(M_t^i)^2] \leq E[\langle M^i \rangle_\tau] = E[A_\tau^i] < \infty$ .

Let  $i, i' \in \mathbb{G}$  be distinct and  $0 \leq s \leq t \leq \tau$ . By Lemma 11.2 (c) and Lemma 11.1 (a), it holds that

$$\begin{aligned}
E[M_t^i M_t^{i'} | \mathcal{F}_s] &= E[E[M_t^i | \widehat{\mathcal{G}}_t^i \vee \mathcal{G}_s^i \vee \mathcal{F}_0] M_t^{i'} | \mathcal{F}_s] \\
&= E[E[M_t^i | \mathcal{G}_s^i \vee \mathcal{F}_0] M_t^{i'} | \mathcal{F}_s] \\
&= E[E[M_t^i | \widehat{\mathcal{G}}_s^i \vee \mathcal{G}_s^i \vee \mathcal{F}_0] M_t^{i'} | \mathcal{F}_s] \\
&= E[E[M_t^i | \mathcal{F}_s] M_t^{i'} | \mathcal{F}_s] \\
&= M_s^i M_s^{i'}.
\end{aligned}$$

This gives  $\langle M^i, M^{i'} \rangle_t = 1_{\{i=i'\}} \int_0^t (1 - \Delta A_s^i) dA_s^i = 1_{\{i=i'\}} A_t^i$ .  $\square$

Finally, we recall Rebolledo's martingale central limit theorem. See Helland (1982) for the proof of this theorem.

**Theorem 11.3.** *For each  $n \in \mathbb{N}$ , let  $M^n$  be a locally square integrable martingale defined on a filtered probability space  $(\Omega^n, \mathcal{F}^n, (\mathcal{F}_t^n)_{t \in \mathbb{R}_+}, P^n)$ . Suppose that the filtration  $(\mathcal{F}_t^n)_{t \in \mathbb{R}_+}$  satisfies the usual condition. Let  $\alpha: \mathbb{R}_+ \rightarrow \mathbb{R}$  be a continuous function and  $(W_t)$  be a process with independent increments,  $E[W_t] = 0$  and  $E[W_t W_s] = \alpha(s \wedge t)$ . For  $\epsilon > 0$ , we define the adapted increasing process  $\sigma^\epsilon[M^n]$  by setting*

$$\sigma^\epsilon[M^n]_t = \sum_{s \leq t} |\Delta M_s^n|^2 1_{\{|\Delta M_s^n| > \epsilon\}} \quad (t \in \mathbb{R}_+).$$

*Note that  $\sigma^\epsilon[M^n]$  is locally integrable since  $\sigma^\epsilon[M^n]_t \leq \sigma^\epsilon[M^n]_{t-} + |\Delta M_t^n|^2$  and  $M^n$  is a locally square integrable martingale. In particular, we can consider the compensator of the process  $\sigma^\epsilon[M^n]$ . Suppose that the following conditions hold.*

(i)  $M_0^n \rightarrow^p 0$ .

(ii)  $\langle M^n \rangle_t \rightarrow^p \alpha(t)$  for all  $t \in [0, \tau]$ .

(iii)  $\sigma^\epsilon[M^n]_\tau^p \rightarrow^p 0$ .

Then, we have

$$M^n \rightarrow^d W \text{ on } D[0, \tau]$$

where  $D[0, \tau]$  is the space of all càdlàg function on  $[0, \tau]$  equipped with Skorohod topology.

## Conflict of Interest

The authors declare no conflicts of interest associated with this paper.

## References

- Abadie A, Imbens GW (2012) A martingale representation for matching estimators. *Journal of the American Statistical Association* 107(498):833–843
- Austin PC (2013) The performance of different propensity score methods for estimating marginal hazard ratios. *Statistics in medicine* 32(16):2837–2849
- Austin PC (2014) The use of propensity score methods with survival or time-to-event outcomes: reporting measures of effect similar to those used in randomized experiments. *Statistics in medicine* 33(7):1242–1258
- Billingsley P (2013) *Convergence of probability measures*. John Wiley & Sons
- Brémaud P (1981) *Point processes and queues*. Springer
- Chen PY, Tsiatis AA (2001) Causal inference on the difference of the restricted mean lifetime between two groups. *Biometrics* 57(4):1030–1038
- Fleming TR, Harrington DP (2013) *Counting processes and survival analysis*, vol 625. John Wiley & Sons

- Guo YM, Zhou J, Sun LQ (2021) Adjusted log-rank test with double inverse weighting under dependent censoring. *Acta Mathematica Sinica, English Series* 37(10):1573–1585
- Helland IS (1982) Central limit theorems for martingales with discrete or continuous time. *Scandinavian Journal of Statistics* pp 79–94
- Iacus SM, King G, Porro G (2011) Multivariate matching methods that are monotonic imbalance bounding. *Journal of the American Statistical Association* 106(493):345–361
- Iacus SM, King G, Porro G (2019) A theory of statistical inference for matching methods in causal research. *Political Analysis* 27(1):46–68
- Jacod J, Shiryaev A (2013) *Limit theorems for stochastic processes*, vol 288. Springer Science & Business Media
- Xie J, Liu C (2005) Adjusted kaplan–meier estimator and log-rank test with inverse probability of treatment weighting for survival data. *Statistics in medicine* 24(20):3089–3110