A short proof of the Dvoretzky–Kiefer–Wolfowitz–Massart inequality

Henry W J Reeve University of Bristol henry.reeve@bristol.ac.uk

Abstract

The Dvoretzky–Kiefer–Wolfowitz–Massart inequality gives a sub-Gaussian tail bound on the supremum norm distance between the empirical distribution function of a random sample and its population counterpart. We provide a short proof of a result that improves the existing bound in two respects. First, our one-sided bound holds without any restrictions on the failure probability, thereby verifying a conjecture of Birnbaum and McCarty (1958). Second, it is local in the sense that it holds uniformly over sub-intervals of the real line with an error rate that adapts to the behaviour of the population distribution function on the interval. ¹

Let X_1, \ldots, X_n be independent random variables with distribution function $F : \mathbb{R} \to [0, 1]$. Let \hat{F}_n denote the empirical distribution function,

$$\hat{F}_n(t) := \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{\{X_i \le t\}}$$

for $t \in \mathbb{R}$. Understanding the deviation of the empirical distribution function from its population counterpart is a longstanding goal within probability theory with a multitude of applications to statistics. The Glivenko–Cantelli theorem (Glivenko, 1933; Cantelli, 1933) gives almost sure convergence of the empirical distribution function to the population distribution function with respect to the supremum norm, initiating a line of work towards understanding the rate of convergence. Classical results of Bernstein (1924), and subsequent work by Bennett (1962) and Hoeffding (1963), may be viewed as giving finite-sample pointwise guarantees for specific value of $t \in \mathbb{R}$. Dvoretzky et al. (1956) proved a sub-Gaussian bound on the supremum norm distance between the empirical and population distribution functions. Subsequent work of (Devroye and Wise, 1979; Hu, 1985; Shorack and Wellner, 1986) led to improvements in the bound. Motivated by an asymptotic analysis due to Smirnov (1944), and further numerical computations, Birnbaum and McCarty (1958) gave a conjecture on the optimal leading constant. A breakthrough result of Massart (1990) achieved the optimal constant, settling the conjecture of Birnbaum and McCarty (1958), subject to a mild constraint on the failure probability. It follows from Smirnov (1944) that the leading term

 $^{^{1}}$ I would like to express my gratitude to Richard J. Samworth for encouragement and useful feedback which substantially improved the quality of the exposition.

in Massart (1990) cannot be improved. We recommend Dudley (2014) for an exposition of the proof of Massart (1990), and its connections to Bretagnolle and Massart (1989). Maillard (2021) derived an integral expression for the probability that the empirical distribution function exceeds a given distance from their population counterpart, uniformly over a given interval. Recently, Bartl and Mendelson (2023) and Blanchard and Voráček (2023) give more interpretable local bounds.

Let's first introduce some notation. Given $a, b \in [0, 1]$ we let

$$\operatorname{kl}(a||b) := a \log\left(\frac{a}{b}\right) + (1-a) \log\left(\frac{1-a}{1-b}\right),$$

where we adopt the convention that $0 \log 0 = 0 \log(0/0) = 0$ and $z \log(z/0) = \infty$ for z > 0. Thus, kl(a||b) denotes the Kullback-Leibler divergence between two Bernoulli random variables with respective success probabilities a and b. For each $p \in (0, e^{-\varepsilon}]$ there is a unique value of $\eta \in [0, 1-p]$ with $kl(p+\eta||p) = \varepsilon$ (Lemma 5) and we denote this unique value by $\omega(p,\varepsilon)$. We extend ω to $[0,1] \times [0,\infty)$ by letting $\omega(p,\varepsilon) := 1-p$ if $p \in (e^{-\varepsilon}, 1]$; we also let $\omega(p,\varepsilon) := 0$ whenever p = 0, so that $p \mapsto \omega(p,\varepsilon)$ is continuous (Lemma 6). Given $\delta \in (0, 1)$ we let $\varepsilon(n, \delta) := \log(1/\delta)/n$.

Our main result is the following.

Theorem 1. Given any interval $\mathcal{I} \subseteq \mathbb{R}$ and $\delta \in (0, 1)$ we have

$$\mathbb{P}\left\{\sup_{t\in\mathcal{I}}\left\{\hat{F}_n(t)-F(t)\right\}>\sup_{t\in\mathcal{I}}\omega\left(F(t),\varepsilon(n,\delta)\right)\right\}\leq\delta.$$

In order to give a more interpretable bound we write $\sigma(p) := \sqrt{p(1-p)}$ for $p \in [0,1]$ and introduce the function $\varphi_0 : [0,1] \times (0,\infty) \to (0,1]$ defined by

$$\varphi_0(p,\varepsilon) := \sqrt{4\sigma^2(p)\rho_\varepsilon^2 + \sigma^2(\rho_\varepsilon)} + \sigma(\rho_\varepsilon)(1-2p)$$

$$= \sqrt{4\sigma^2(p)\rho_\varepsilon^2 + \sigma^2(\rho_\varepsilon)} + \sigma(\rho_\varepsilon)\operatorname{sign}(1-2p)\sqrt{1-4\sigma^2(p)},$$
(1)

where $\rho_{\varepsilon} := 9/(9+2\varepsilon) \in (0,1)$. Note that $\varphi_0(p,\varepsilon) \leq 1$ for all $p \in [0,1]$. Indeed, the map $p \mapsto \varphi_0(p,\varepsilon)$ is concave with a unique maximum of $\sup_{p \in [0,1]} \varphi_0(p,\varepsilon) = 1$ attained when $1 - 2p = \sigma(\rho_{\varepsilon})/\rho_{\varepsilon}$. Moreover, we have $\lim_{\varepsilon \searrow 0} \rho_{\varepsilon} = 1$, so $\lim_{\varepsilon \searrow 0} \sigma(\rho_{\varepsilon}) = 0$, and hence $\lim_{\varepsilon \searrow 0} \varphi_0(p,\varepsilon) = 2\sigma(p)$ for all $p \in [0,1]$. Given an interval $\mathcal{I} \subseteq \mathbb{R}$, let $p_{\min} := \inf_{t \in \mathcal{I}} F(t)$ and $p_{\max} := \sup_{t \in \mathcal{I}} F(t)$, and set

$$\varphi(\mathcal{I},\varepsilon) := \begin{cases} \varphi_0(p_{\max},\varepsilon) & \text{if } 2p_{\max} < 1 - \sigma(\rho_{\varepsilon})/\rho_{\varepsilon} \\ 1 & \text{if } 2p_{\min} \le 1 - \sigma(\rho_{\varepsilon})/\rho_{\varepsilon} \le 2p_{\max} \\ \varphi_0(p_{\min},\varepsilon) & \text{if } 2p_{\min} > 1 - \sigma(\rho_{\varepsilon})/\rho_{\varepsilon}. \end{cases}$$

Our first corollary is designed to highlight the role played by variance in the local bound as well as the mild asymmetry in deviations around one half.

Corollary 1. Given any interval $\mathcal{I} \subseteq \mathbb{R}$ and $\delta \in (0, 1)$ we have

$$\mathbb{P}\left\{\sup_{t\in\mathcal{I}}\left\{\hat{F}_n(t)-F(t)\right\}>\varphi\left(\mathcal{I},\varepsilon(n,\delta)\right)\sqrt{\frac{\varepsilon(n,\delta)}{2}}\right\}\leq\delta.$$

Our next corollary is a refinement of the celebrated Dvoretzky–Kiefer–Wolfowitz–Massart inequality (Dvoretzky et al., 1956; Massart, 1990).

Corollary 2. Given any $\xi \ge 0$ and $\zeta := 2\xi^2/3$ we have

$$\mathbb{P}\left\{\sup_{t\in\mathbb{R}}\sqrt{n}\left\{\hat{F}_n(t) - F(t)\right\} > \xi\right\} \le \exp\left(-2\xi^2 - \frac{\zeta^2}{n}\left\{1 + \frac{4\zeta}{5n} + \frac{425\zeta^2}{525n^2}\right\}\right).$$

In comparison, Massart (1990, Theorem 1) gives a slightly larger bound of $\exp(-2\xi^2)$ for $\xi \geq \min\{\sqrt{\log(2)/2}, 1.0841n^{-1/6}\}$. Massart (1990) remarks that whilst this captures the regime of interest for statistical purposes, the proof therein does not extend to smaller values of ξ . Corollary 1 verifies the conjecture of Birnbaum and McCarty (1958) for all values of $\xi \geq 0$ and Corollary 2 proves a slightly stronger result.

Our final corollary is a variance-adaptive bound for the entire real line, at the expense of an extra logarithmic factor.

Corollary 3. Suppose that $\delta \in (0, 1)$, $\beta > 1$ and let $\varepsilon_{\beta}(n, \delta) := \log(2\lceil \log_{\beta}(n) \rceil / \delta) / n$. Then,

$$\mathbb{P}\left\{\hat{F}_n(t) - F(t) > \left(\beta\varphi_0(F(t),\varepsilon_\beta(n,\delta))\sqrt{\varepsilon_\beta(n,\delta)/2}\right) \vee \left(\frac{1}{n}\right) \text{ for some } t \in \mathbb{R}\right\} \le \delta.$$

Corollary 3 is a slight improvement on recent results from Bartl and Mendelson (2023, Theorem 1.2) and Blanchard and Voráček (2023, Theorem 5) where similar bounds are given but the universal constants are left unspecified. We remark that Corollary 3 can be restated as a simultaneous lower-confidence bound by taking

$$\mathcal{U}(q,\varepsilon) := \frac{3\beta \left((2q-1)(3\beta-1)\varepsilon + \sqrt{\epsilon \{18\sigma^2(q) + \varepsilon(3\beta-1)^2\}} \right)}{9 + 2\varepsilon(3\beta-1)^2}.$$

Indeed, Corollary 3 may be restated as follows

$$\mathbb{P}\left\{F(t) > \hat{F}_n(t) - \mathcal{U}(\hat{F}_n(t), \varepsilon_\beta(n, \delta)) \lor \left(\frac{1}{n}\right) \text{ for all } t \in \mathbb{R}\right\} \ge 1 - \delta.$$

The proof of Theorem 1 is given in Section 1, with the proof a technical lemma and the corrollaries given in Section 2.

1 Proof of Theorem 1

To begin the proof we first introduce a reverse filtration $(\mathcal{G}_t)_{t\in\mathbb{R}}$ (a collection of sigma algebras such that $\mathcal{G}_{t_1} \subseteq \mathcal{G}_{t_0}$ whenever $t_0 \leq t_1$). For each $t \in \mathbb{R}$ we let \mathcal{G}_t denote the sigma-algebra by all events of the form $\{X_i \leq s\}$ with $s \geq t$ and $i \in [n]$ are measurable.

Given $\lambda \geq 0$ let $(M_{\lambda}(t))_{t \geq t_0}$ denote the non-negative stochastic process defined by

$$M_{\lambda}(t) := \frac{1}{(1+\lambda)^n} \left(1 + \frac{\lambda}{F(t)}\right)^{n\tilde{F}_n(t)}.$$

Lemma 1. Suppose $t_0 \in \mathbb{R}$ satisfies $F(t_0) > 0$. Then $(M_{\lambda}(t))_{t \geq t_0}$ is a reverse martingale with respect to $(\mathcal{G}_t)_{t \geq t_0}$. Moreover, $\mathbb{E}\{M_{\lambda}(t)\} = 1$ for all $t \geq t_0$.

Proof. Take $s_0 \in [t_0, \infty)$ and $s_1 \in (s_0, \infty)$. Observe that, conditionally on \mathcal{G}_{s_1} , the random variable $W := n\hat{F}_n(s_0)$ has the distribution of a Binomial random variable with $m := n\hat{F}_n(s_1)$ trials and success probability $p := F(s_0)/F(s_1)$. Thus, for each $\lambda \geq 0$ we have

$$\mathbb{E}\left\{\left(1+\frac{\lambda}{F(s_0)}\right)^{n\hat{F}_n(s_0)} \mid \mathcal{G}_{s_1}\right\} = \mathbb{E}\left\{\left(1+\frac{\lambda}{pF(s_1)}\right)^W \mid \mathcal{G}_{s_1}\right\}$$
$$= \left\{p\left(1+\frac{\lambda}{pF(s_1)}\right) + (1-p)\right\}^m = \left(1+\frac{\lambda}{F(s_1)}\right)^{n\hat{F}_n(s_1)}$$

which entails the reverse martingale property. Applying the dominated convergence theorem with the bound $(1 + \lambda/F(s_0))^n$ yields that for all $t \ge t_0$,

$$\mathbb{E}\{M_{\lambda}(t)\} = \lim_{s \nearrow \infty} \mathbb{E}\{M_{\lambda}(s)\} = \mathbb{E}\left\{\lim_{s \nearrow \infty} M_{\lambda}(s)\right\} = 1.$$

Next, for each $\varepsilon > 0$, we define a function $\Delta \equiv \Delta_{\varepsilon} : (0,1] \times (0,\infty) \to [0,\infty)$ by

$$\Delta(r,\lambda) \equiv \Delta_{\varepsilon}(r,\lambda) := \frac{\log(1+\lambda) + \varepsilon}{\log(1+\lambda/r)} - r.$$

Note that the inequality $(1 + \lambda/r)^r \leq (1 + \lambda)$ for $r \in [0, 1]$ ensures that Δ is non-negative. Given $\lambda > 0$ and $\delta \in (0, 1)$ we define

$$\mathcal{E}(\lambda,\delta) := \bigcap_{t \in \mathbb{R}: F(t) > 0} \left\{ \hat{F}_n(t) - F(t) \le \Delta_{\varepsilon(n,\delta)}(F(t),\lambda) \right\}.$$

Lemma 2. For all $\lambda > 0$ and $\delta \in (0, 1)$ we have $\mathbb{P}\{\mathcal{E}(\lambda, \delta)\} \ge 1 - \delta$.

Proof. Given $k \in \mathbb{N}$ we let $\mathbb{T}_k := \{t \in [-2^k, 2^k] : t = 2^{-k} \lfloor 2^k t \rfloor$ and $F(t) > 0\}$. By Lemma 1 the process $\{M_\lambda(t)\}_{t \in \mathbb{T}_k}$ is a reverse martingale with $\mathbb{E}(M_\lambda(t)) = 1$ for all $k \in \mathbb{N}$ and $t \in \mathbb{T}_k$. Hence, by Doob's maximal inequality (Dudley, 2018, Theorem 10.4.2) we have $\mathbb{P}\{\mathcal{E}_k(\lambda, \delta)\} \ge 1 - \delta$ where

$$\mathcal{E}_k(\lambda,\delta) := \left\{ \max_{t \in \mathbb{T}_k} M_\lambda(t) < 1/\delta \right\} = \bigcap_{t \in \mathbb{T}_k} \left\{ \hat{F}_n(t) < F(t) + \Delta_{\varepsilon(n,\delta)} (F(t),\lambda) \right\},\$$

where the equality between the events follows by taking logarithms. In addition, since $t \mapsto M_{\lambda}(t)$ is almost surely right-continuous and $\bigcup_{k \in \mathbb{N}} \mathbb{T}_k$ is dense in $\{t \in \mathbb{R} : F(t) > 0\}$ we have $\mathcal{E}(\lambda, \delta) = \bigcap_{k \in \mathbb{N}} \mathcal{E}_k(\lambda, \delta)$. Thus, since the events $\{\mathcal{E}_k(\lambda, \delta)\}_{k \in \mathbb{N}}$ form a decreasing sequence we have $\mathbb{P}\{\mathcal{E}(\lambda, \delta)\} \ge 1 - \delta$.

Lemma 3. Given any non-empty closed interval $\mathcal{P} \subseteq (0, e^{-\varepsilon})$ there exists $(p_{\star}, \lambda_{\star}) \in \mathcal{P} \times (0, \infty)$ with $\Delta(p_{\star}, \lambda_{\star}) = \inf_{\lambda \in (0, \infty)} \Delta(p_{\star}, \lambda) = \sup_{p \in \mathcal{P}} \Delta(p, \lambda_{\star}) = \omega(p_{\star}, \varepsilon)$.

The proof of Lemma 3 is somewhat technical and is given in Section 2.

Proof of Theorem 1. Choose $\zeta \in (0,1)$ and suppose that the closed interval $\mathcal{P}_{\zeta} := \operatorname{cl}(\{F(t) : t \in \mathcal{I}\}) \cap [\zeta, e^{-\varepsilon} - \zeta]$ is non-empty. By applying Lemma 3 with $\varepsilon = \varepsilon(n, \delta)$ we obtain $(p_{\star}, \lambda_{\star}) \in \mathcal{P} \times (0, \infty)$ so that $\Delta(p_{\star}, \lambda_{\star}) = \sup_{p \in \mathcal{P}_{\zeta}} \Delta(p, \lambda_{\star}) = \omega(p_{\star}, \varepsilon) > 0$. Moreover, by Lemma 2 we have $\mathbb{P}\{\mathcal{E}(\lambda_{\star}, \delta)\} \geq 1 - \delta$. On the event $\mathcal{E}(\lambda_{\star}, \delta)$, for each $t \in \mathcal{I}$ with $F(t) \in [\zeta, e^{-\varepsilon} - \zeta]$ we have

$$\hat{F}_n(t) - F(t) < \Delta(F(t), \lambda) \le \sup_{p \in \mathcal{P}_{\zeta}} \Delta(p, \lambda_\star) = \Delta(p_\star, \lambda_\star) = \omega(p_\star, \varepsilon)$$
$$\le \sup_{p \in \operatorname{cl}(\{F(t): t \in \mathcal{I}\})} \omega(p, \varepsilon) = \sup_{p \in \{F(t): t \in \mathcal{I}\}} \omega(p, \varepsilon) = \sup_{t \in \mathcal{I}} \omega(F(t), \varepsilon),$$

where we applied the continuity of $p \mapsto \omega(p, \varepsilon)$ (Lemma 6) for the penulitmate equality. Note that the bound holds vacuously if $\mathcal{P}_{\zeta} = \emptyset$. Moreover, if $t \in \mathcal{I}$ satisfies $F(t) \in \{0\} \cup [e^{-\varepsilon}, 1]$ then we have $\hat{F}_n(t) - F(t) \leq \omega(F(t), \varepsilon) \leq \sup_{t \in \mathcal{I}} \omega(F(t), \varepsilon)$ almost surely (Lemma 7). Hence,

$$\mathbb{P}\left\{\hat{F}_n(t) - F(t) > \sup_{t \in \mathcal{I}} \omega(F(t), \varepsilon(n, \delta)) \text{ for some } t \in \mathcal{I} \text{ with } F(t) \notin (0, \zeta) \cup (e^{-\varepsilon} - \zeta)\right\} \le \delta.$$

Since this bound holds for all sufficiently small ζ , the conclusion of the theorem follows by continuity of probability.

2 Proofs of technical lemmas

Throughout we shall consider $[-\infty, \infty] := \mathbb{R} \cup \{-\infty, \infty\}$ with the usual topology, generated by the topological basis consisting of all intervals of the form (a, b), $(a, \infty]$ or $[-\infty, b)$ where $a, b \in \mathbb{R}$. We shall use the following elementary form of the implicit function theorem (Lemma 4). Given a Borel set $\mathcal{Y} \subseteq [-\infty, \infty]$ we shall say that a function $g : \mathcal{Y} \to [-\infty, \infty]$ is sign-monotonic if

$$(g(y_a), g(y_b)) \in ([-\infty, 0) \times [-\infty, \infty]) \cup ([-\infty, \infty] \times (0, \infty]),$$
(2)

for all $y_a, y_b \in \mathcal{Y}$ with $y_a < y_b$. Note that a sufficient condition for $g : \mathcal{Y} \to [-\infty, \infty]$ to be sign-monotonic is that g is both non-decreasing and $|\{y \in \mathcal{Y} : g(y) = 0\}| \leq 1$.

Lemma 4. Let $\mathcal{X} \subseteq \mathbb{R}^d$ be a Borel set, $\mathcal{Y} \subseteq [-\infty, \infty]$ is a non-empty interval which is bounded from below and let $\Gamma : \mathcal{X} \times \mathcal{Y} \to [-\infty, \infty]$ be a function. Suppose that for each $x \in \mathcal{X}$ the function $y \mapsto \Gamma(x, y)$ is sign-monotonic. Suppose further that for each $y \in \mathcal{Y}$, the function $x \mapsto \Gamma(x, y)$ is continuous and define a function $\gamma : \mathcal{X} \to cl(\mathcal{Y})$ by

$$\gamma(x) := \begin{cases} \sup \mathcal{Y} & \text{if } \Gamma(x, y) < 0 \text{ for all } y \in \mathcal{Y} \\ \inf\{y \in \mathcal{Y} : \Gamma(x, y) \ge 0\} & \text{otherwise.} \end{cases}$$

Then γ is continuous and $\Gamma(x, y)(y - \gamma(x)) > 0$ for all $x \in \mathcal{X}$ and $y \in \mathcal{Y} \setminus \{\gamma(x)\}$.

Whilst we expect that Lemma 4 is known, we include a proof in Section 4 for the convenience of the reader.

Lemma 5. For each $p \in (0,1)$, the function $\eta \mapsto \text{kl}(p+\eta||p)$ is strictly increasing and continuous bijection from [0, 1-p] to $[0, \log(1/p)]$. Hence, for $p \in (0, e^{-\varepsilon}]$, there is a unique value of $\eta \in [0, 1-p]$ with $\text{kl}(p+\eta||p) = \varepsilon$ and if $p \in (e^{-\varepsilon}, 1]$ then $\text{kl}(p+\eta||p) < \varepsilon$ for all $\eta \in [0, 1-p]$.

Proof. Given $p \in (0, 1)$ and $\eta_0 \in [0, 1 - p)$, we have $p + \eta_0 < 1$ so the map $\eta \mapsto \text{kl}(p + \eta || p)$ is differentiable at η_0 , with derivative

$$\log\left(\frac{\eta_0+p}{p}\right) + \log\left(\frac{1-p}{1-p-\eta_0}\right) > 0.$$

Hence, $\eta \mapsto \operatorname{kl}(p + \eta || p)$ is also strictly increasing and continuous function on [0, 1 - p). Moreover, since $p \in (0, 1)$ we have $\lim_{a \nearrow 1} \operatorname{kl}(a || p) = \log(1/p) = \operatorname{kl}(1 || p)$ so that $\eta \mapsto \operatorname{kl}(p + \eta || p)$ is strictly increasing and continuous bijection from [0, 1 - p] to $[0, \log(1/p)]$. Finally, note that $\varepsilon \in [0, \log(1/p)] = \{\operatorname{kl}(p + \eta || p) : \eta \in [0, 1 - p]\}$ if and only if $p \in (0, e^{-\varepsilon})$.

Lemma 6. Given $\varepsilon > 0$ the function $p \mapsto \omega(p, \varepsilon)$ is continuous.

Proof. Let's define a function $\Gamma_{\rm kl}: [0,1]^2 \to [0,\infty]$ by

$$\Gamma_{\rm kl}(x,y) := \begin{cases} -\varepsilon & \text{if } y \le x \\ {\rm kl}(y||x) - \varepsilon & \text{otherwise.} \end{cases}$$

For each $x \in (0, 1)$, it follows from Lemma 5 that $y \mapsto \Gamma_{kl}(x, y)$ is sign-monotonic. Furthermore, if $x \in \{0, 1\}$ then $y \mapsto \Gamma_{kl}(x, y)$ is a non-decreasing and piecewise constant with $\{y \in [0, 1] : \Gamma_{kl}(x, y) = 0\} = \emptyset$. As such, $y \mapsto \Gamma_{kl}(x, y)$ is sign-monotonic for all $x \in [0, 1]$. Note also that for each $y \in [0, 1]$, the map $x \mapsto kl(y||x)$ is continuous; and consequently so is $x \mapsto \Gamma_{kl}(x, y)$. Thus, by Lemma 4 the function $\gamma_{kl} : [0, 1] \to [0, 1]$ by

$$\gamma_{\rm kl}(x) := \begin{cases} 1 & \text{if } {\rm kl}(y||x) < \varepsilon \text{ for all } y \in [0,1] \\ {\rm inf}\{y \in [0,1] : {\rm kl}(y||x) \ge \varepsilon\} & \text{otherwise.} \end{cases}$$

is also continuous. Finally note that for all $p \in [0, 1]$ we have $\omega(p, \varepsilon) = \gamma_{kl}(p) - p$, so that $p \mapsto \omega(p, \varepsilon)$ is also continuous.

Proof of Lemma 3. Observe that Δ is twice differentiable on $(0,1] \times (0,\infty)$ with

$$\partial_r \Delta(r,\lambda) = \frac{\lambda \left\{ \varepsilon + \log\left(1+\lambda\right) \right\}}{r\left(\lambda+r\right)\log^2\left(1+\lambda/r\right)} - 1,$$

$$\partial_{r,r}\Delta(r,\lambda) = \frac{\lambda^2 \left\{ \varepsilon + \log\left(1+\lambda\right) \right\} \left\{ 2 - \left(1+2r/\lambda\right)\log\left(1+\lambda/r\right) \right\}}{r^2 \left(r+\lambda\right)^2 \log^3\left(1+\lambda/r\right)}.$$

By applying the inequality $\log(1+z) > 2z/(2+z)$ for z > 0 we deduce $\partial_{r,r}\Delta(r,\lambda) < 0$ for r > 0, so $r \mapsto \Delta(r,\lambda)$ is strictly concave. Now consider the function $\Gamma_0: (0,\infty) \times \mathcal{P} \to [-\infty,\infty]$ defined by $\Gamma_0(\lambda,r) := -\partial_r \Delta(r,\lambda)$. For each $\lambda \in (0,\infty)$, the strict concavity of $r \mapsto \Delta(r,\lambda)$ implies that $r \mapsto \Gamma_0(\lambda,r)$ is strictly increasing and as such $|\{r \in \mathcal{P} : \Gamma_0(\lambda,r) = 0\}| \leq 1$. Moreover, for each $r \in \mathcal{P}$ the function $\lambda \mapsto \Gamma_0(\lambda,r)$ is continuous. Hence, by Lemma 4 there exists a continuous function $\overline{p} : (0, \infty) \to \mathcal{P}$ such that $(\overline{p}(\lambda) - r)\partial_r \Delta(r, \lambda) < 0$ for all $\lambda \in (0, \infty)$ and $p \in \mathcal{P} \setminus \{\overline{p}(\lambda)\}$. Hence, we have $\Delta(r, \lambda) \leq \Delta(\overline{p}(\lambda), \lambda)$ for all $(r, \lambda) \in \mathcal{P} \times (0, \infty)$. Next, we define a function $\Gamma_1 : \mathcal{P} \times (0, \infty) \to \mathbb{R}$ by

 $\Gamma_1(p,\lambda) := (p+\lambda)\log\left(1+\lambda/p\right) - (1+\lambda)\left\{\varepsilon + \log\left(\lambda+1\right)\right\},\,$

so that $\partial_{\lambda}\Delta(p,\lambda) = \Gamma_1(p,\lambda)/\{(1+\lambda)(p+\lambda)\log^2(1+\lambda/p)\}$. Note that for each $\lambda \in (0,\infty)$ the map $p \mapsto \Gamma_1(p,\lambda)$ is continuous on \mathcal{P} with

$$\partial_{\lambda}\Gamma_{1}(p,\lambda) = \log\left(1 + \lambda/p\right) - \{\varepsilon + \log\left(1 + \lambda\right)\},\$$

so $\Gamma_1(p,\lambda) = 0$ implies $\partial_{\lambda}\Gamma_1(p,\lambda) > 0$. Thus, $\lambda \mapsto \Gamma_1(p,\lambda)$ is sign-monotonic for all $p \in \mathcal{P}$. Consequently, there exists a continuous function $\underline{\lambda} : \mathcal{P} \to [0,\infty]$ such that $\Gamma_1(p,\lambda)(\underline{\lambda}(p) - \lambda) > 0$, and consequently, $\partial_{\lambda}\Delta(p,\lambda)(\underline{\lambda}(p)-\lambda) > 0$ for all $p \in \mathcal{P}$ and $\lambda \in (0,\infty) \setminus \{\underline{\lambda}(p)\}$. Note also that for each $p \in \mathcal{P} \subseteq (0, e^{-\varepsilon})$ we have $\lim_{\lambda \searrow 0} \Gamma_0(p,\lambda) = -\varepsilon < 0$ and $\lim_{\lambda \nearrow \infty} \Gamma(p,\lambda) = \infty$ so by by the continuity of $\lambda \mapsto \Gamma(p,\lambda)$ there must be at least one root in $(0,\infty)$, which must coincide with $\underline{\lambda}(p)$ i.e. $\Gamma(p,\underline{\lambda}(p)) = 0$. Hence, $\underline{\lambda}(p) \in (0,\infty)$ and $\Delta(p,\underline{\lambda}(p)) \leq \Delta(p,\lambda)$ for all $\lambda \in (0,\infty)$.

Next, we define a function $\Phi : \mathcal{P} \to \mathbb{R}$ by $\Phi(\rho) = \overline{p}(\underline{\lambda}(\rho)) - \rho$ and choose $p_{\min}, p_{\max} \in (0, e^{-\varepsilon})$ so that $\mathcal{P} = [p_{\min}, p_{\max}]$. Notice that Φ is continuous with $\Phi(p_{\min}) \ge 0$ and $\Phi(p_{\max}) \le 0$ so by the intermediate value theorem there must exist some $p_{\star} \in \mathcal{P}$ with $\Phi(p_{\star}) = 0$. Letting $\lambda_{\star} := \underline{\lambda}(p_{\star})$ we obtain a pair $(p_{\star}, \lambda_{\star}) \in \mathcal{P} \times (0, \infty)$ satisfying $\Delta(p_{\star}, \lambda_{\star}) = \inf_{\lambda \in (0,\infty)} \Delta(p_{\star}, \lambda) = \sup_{p \in \mathcal{P}} \Delta(p, \lambda_{\star})$.

In addition, given any $p \in (0, e^{-\varepsilon})$ we have $\Gamma_1(p, \underline{\lambda}(p)) = 0$ which yields $p + \Delta\{p, \underline{\lambda}(p)\} = (p + \underline{\lambda}(p))/(1 + \underline{\lambda}(p)) \in (p, 1)$ and $\operatorname{kl}(p + \Delta\{p, \underline{\lambda}(p)\} ||p) = \varepsilon$ upon rearranging. Hence, $\omega(p, \underline{\lambda}(p)) = \Delta(p, \varepsilon) > 0$. Since $\lambda_{\star} = \underline{\lambda}(p_{\star})$ this completes the proof of the lemma. \Box

For the next lemma we define an event \mathcal{E}_0 by

$$\mathcal{E}_0 := \bigcap_{t \in \mathbb{R}: F(t) \in \{0\} \cup [e^{-\varepsilon}, 1]} \left\{ \hat{F}_n(t) - F(t) \le \omega(F(t), \varepsilon) \right\}.$$

Lemma 7. We have $\mathbb{P}(\mathcal{E}_0) = 1$.

Proof. Let's introduce $J_0 := \{t \in \mathbb{R} : F(t) = 0\}$ and $\tilde{\mathcal{E}}_0 := \{\sup_{t \in \mathbb{R}} \hat{F}_n(t) \leq 1\} \cap \bigcap_{i=1}^n \{X_i \notin J_0\}$, so that $\mathbb{P}(\tilde{\mathcal{E}}_0) = 1$. Now suppose the event $\tilde{\mathcal{E}}_0$ holds. If F(t) = 0 then $(0,t] \subseteq J_0$ so that $\hat{F}_n(t) = 0$, and so $\hat{F}_n(t) - F(t) = 0 = \omega(F(t), \varepsilon)$. On the other hand, if $F(t) \geq e^{-\varepsilon}$ then $\hat{F}_n(t) - F(t) \leq 1 - F(t) = \omega(F(t), \varepsilon)$, where we have used the definition of $\omega(p,\varepsilon)$ for $\log(1/p) < \varepsilon$ and the continuity of $p \mapsto \omega(p,\varepsilon)$ at $p = e^{-\varepsilon}$ (Lemma 6).

Given $(p,\eta) \in (0,1)$ and $\eta \in [0,1-p]$ we let

$$h_m(p,\eta) := \frac{\eta^2}{2(p+\eta/3)(1-p-\eta/3)}$$

Lemma 8. For all $p \in (0, 1)$ and $\eta \in [0, 1 - p]$ we have $h_m(p, \eta) \leq kl(p + \eta || p)$.

Proof. Given $z \in (0, 1]$ let's write $f_z : (0, z] \to \mathbb{R}$ for the function

$$f_z(\eta) := \mathrm{kl}(z||z-\eta) - h_m(z-\eta,\eta).$$

Observe that $f_z(0) = 0$ and for $\eta < z \le 1$ we have

$$\partial_{\eta} f_z(\eta) = \frac{\eta^3 \left(16\eta^2 - 42\eta z + 21\eta + 27z^2 - 27z + 9\right)}{(3z - 2\eta)^2 (3 + 2\eta - 3z)^2 (z - \eta)(1 + \eta - z)} > 0$$

Hence, $kl(p + \eta || p) = h_m(p, \eta) + f_{p+\eta}(\eta) \ge h_m(p, \eta)$, which establishes the first part of the lemma.

3 Proofs of the corollaries

Proof of Corollary 1. Let $p_{\min} := \inf_{t \in \mathcal{I}} F(t)$, $p_{\max} := \sup_{t \in \mathcal{I}} F(t)$ and take $\varepsilon = \varepsilon(n, \delta)$. Given $t \in \mathcal{I}$ we have $p = F(t) \in [p_{\min}, p_{\max}]$ we have $\operatorname{kl}(p + \eta | p) < \varepsilon$ for any $\eta \in (0, \omega(p, \varepsilon))$. Thus, by Lemma 8 we have $h_m(p, \eta) < \varepsilon$. By rearranging this bound with $\rho_{\varepsilon} = 9/(9 + 2\varepsilon)$, so $\sigma^2(\rho_{\varepsilon})/\rho_{\varepsilon}^2 = 2\varepsilon/9$ we deduce that

$$\eta < \frac{3(\sqrt{\varepsilon\{\varepsilon+18p(1-p)\}}+\varepsilon\{1-2p\})}{2\varepsilon+9} = \rho_{\varepsilon} \left\{ \sqrt{4p(1-p)+\frac{2\varepsilon}{9}} + (1-2p)\sqrt{\frac{2\varepsilon}{9}} \right\} \sqrt{\frac{\varepsilon}{2}} \\ = \left\{ \sqrt{4\sigma^{2}(p)\rho_{\varepsilon}^{2}+\sigma^{2}(\rho_{\varepsilon})} + (1-2p)\sigma(\rho_{\varepsilon}) \right\} \sqrt{\frac{\varepsilon}{2}} = \varphi_{0}(p,\varepsilon)\sqrt{\frac{\varepsilon}{2}}.$$

Since this result for all $t \in \mathcal{I}$ and $\eta \in (0, \omega(p, \varepsilon))$ we deduce that

$$\sup_{t \in \mathcal{I}} \omega(F(t), \varepsilon) \le \sup_{p \in [p_{\min}, p_{\max}]} \varphi_0(p, \varepsilon) \sqrt{\frac{\varepsilon}{2}}.$$

Moreover, the function $p \mapsto \varphi_0(p, \varepsilon)$ is unimodal, strictly increasing on $[0, 1/2 - \sqrt{\varepsilon/3}]$ and strictly decreasing on $[1/2 - \sqrt{\varepsilon/3}, 1]$. Thus, we have

$$\varphi(\mathcal{I},\varepsilon) = \sup_{p \in [p_{\min}, p_{\max}]} \varphi_0(p,\varepsilon) \ge \sup_{t \in \mathcal{I}} \omega(F(t),\varepsilon) \sqrt{\frac{2}{\varepsilon}}$$

and the conclusion of Corollary 1 follows from Theorem 1.

Proof of Corollary 2. Given $\xi \ge 0$, $\zeta := 2\xi^2/3$, $\eta = \xi/\sqrt{n}$ and $p \in [0, 1 - \eta]$ it follows from Fedotov et al. (2003, Theorem 7) that

$$\mathrm{kl}(p+\eta||p) \ge 2\eta^2 + \frac{4\eta^4}{9} + \frac{32\eta^6}{135} + \frac{7072\eta^8}{42525} > n\left(2\xi^2 + \frac{\zeta^2}{n}\left\{1 + \frac{4\zeta}{5n} + \frac{425\zeta^2}{525n^2}\right\}\right) =:\log(1/\delta).$$

Thus, $\omega(p, \varepsilon(n, \delta)) \leq \eta$ for all $p \in [0, 1]$. By Theorem 1 we deduce that

$$\mathbb{P}\left\{\sup_{t\in\mathcal{I}}\left\{\hat{F}_n(t)-F(t)\right\}>\eta\right\}\leq\delta.$$

Rearranging yields the conclusion of Corollary 2.

Proof of Corollary 3. By computing first and second derivatives we note that the map $p \mapsto \varphi_0(p,\varepsilon)\sqrt{\varepsilon/2}$ is concave with a unique maximum attained at $p_\star := \{1 - \sigma(\rho_\varepsilon)/\rho_\varepsilon\}/2$. Hence, we may construct a sequence of closed intervals $\{A_r\}_{r=1}^{\lceil \log_\beta(n) \rceil - 1}$ and $\{B_r\}_{r=1}^{\lceil \log_\beta(n) \rceil - 1}$ such that $A_r \subseteq [0, p_\star], B_r \subseteq [p_\star, 1]$ and

$$\inf_{p \in A_r} \varphi_0 \{ F(t), \varepsilon_\beta(n, \delta) \} = \inf_{p \in B_r} \varphi_0 \{ F(t), \varepsilon_\beta(n, \delta) \} = \sqrt{\frac{2\beta^{-2(r+1)}}{\varepsilon_\beta(n, \delta)}},$$
$$\sup_{p \in A_r} \varphi_0 \{ F(t), \varepsilon_\beta(n, \delta) \} = \sup_{p \in B_r} \varphi_0 \{ F(t), \varepsilon_\beta(n, \delta) \} = \sqrt{\frac{2\beta^{-2r}}{\varepsilon_\beta(n, \delta)}}.$$

By Corollary 1, for each $r = 1, \ldots, \lceil \log_{\beta}(n) \rceil - 1$ we have

$$\begin{split} & \mathbb{P}\Big\{\hat{F}_{n}(t) - F(t) > \beta\varphi_{0}\big\{F(t), \varepsilon_{\beta}(n,\delta)\big\}\sqrt{\frac{\varepsilon_{\beta}(n,\delta)}{2}} \text{ for some } t \in \mathbb{R} \text{ with } F(t) \in A_{r}\Big\} \\ & \leq \mathbb{P}\Big\{\sup_{t \in \mathbb{R} : F(t) \in A_{r}}\big\{\hat{F}_{n}(t) - F(t)\big\} > \inf_{p \in A_{r}}\beta\varphi_{0}\big\{p, \varepsilon_{\beta}(n,\delta)\big\}\sqrt{\frac{\varepsilon_{\beta}(n,\delta)}{2}}\Big\} \\ & \leq \mathbb{P}\Big\{\sup_{t \in \mathbb{R} : F(t) \in A_{r}}\big\{\hat{F}_{n}(t) - F(t)\big\} > \sup_{p \in A_{r}}\varphi_{0}\big\{p, \varepsilon_{\beta}(n,\delta)\big\}\sqrt{\frac{\varepsilon_{\beta}(n,\delta)}{2}}\Big\} \\ & = \mathbb{P}\Big\{\sup_{t \in \mathbb{R} : F(t) \in A_{r}}\big\{\hat{F}_{n}(t) - F(t)\big\} > \varphi\big\{A_{r}, \varepsilon_{\beta}(n,\delta)\big\}\sqrt{\frac{\varepsilon_{\beta}(n,\delta)}{2}}\Big\} \le \frac{\delta}{2\lceil\log_{\beta}(n)\rceil}, \end{split}$$

and similarly with B_r in place of A_r . We also let $A_{\lceil \log_\beta(n) \rceil} := [0, \inf A_{\lceil \log_\beta(n) \rceil-1}]$ and $B_{\lceil \log_\beta(n) \rceil} := [\sup B_{\lceil \log_\beta(n) \rceil-1}, 1]$ so that

$$\sup_{p \in A_{\lceil \log_{\beta}(n) \rceil} \cup B_{\lceil \log_{\beta}(n) \rceil}} \varphi_0 \{ F(t), \varepsilon_{\beta}(n, \delta) \} \le \sqrt{\frac{2\beta^{-2\lceil \log_{\beta}(n) \rceil}}{\varepsilon_{\beta}(n, \delta)}} \le \frac{1}{n} \sqrt{\frac{2}{\varepsilon_{\beta}(n, \delta)}}$$

Thus, applying Corollary 1 once more yields

$$\mathbb{P}\left\{\hat{F}_{n}(t) - F(t) > \frac{1}{n} \text{ for some } t \in \mathbb{R} \text{ with } F(t) \in A_{\lceil \log_{\beta}(n) \rceil}\right\}$$
$$\leq \mathbb{P}\left\{\sup_{t \in \mathbb{R}: F(t) \in A_{\lceil \log_{\beta}(n) \rceil}} \left\{\hat{F}_{n}(t) - F(t)\right\} > \varphi\left\{A_{\lceil \log_{\beta}(n) \rceil}, \varepsilon_{\beta}(n, \delta)\right\} \sqrt{\frac{\varepsilon_{\beta}(n, \delta)}{2}}\right\} \leq \frac{\delta}{2\lceil \log_{\beta}(n) \rceil}$$

and similarly with $B_{\lceil \log_{\beta}(n) \rceil}$ in place of $A_{\lceil \log_{\beta}(n) \rceil}$. Finally, letting C_0 denote the (possibly empty) closed open interval $C_0 := [0, 1] \setminus \left\{ \bigcup_{r=1}^{\lceil \log_{\beta}(n) \rceil} (A_r \cup B_r) \right\}$ we have

$$\mathbb{P}\left\{\hat{F}_{n}(t) - F(t) > \beta\varphi_{0}\left\{F(t), \varepsilon_{\beta}(n, \delta)\right\} \sqrt{\frac{\varepsilon_{\beta}(n, \delta)}{2}} \text{ for some } t \in \mathbb{R} \text{ with } F(t) \in C_{0}\right\}$$
$$= \mathbb{P}\left\{\sup_{t \in \mathbb{R} : F(t) \in C_{0}}\left\{\hat{F}_{n}(t) - F(t)\right\} > \inf_{p \in C_{0}}\beta\varphi_{0}\left\{p, \varepsilon_{\beta}(n, \delta)\right\} \sqrt{\frac{\varepsilon_{\beta}(n, \delta)}{2}}\right\}$$
$$= \mathbb{P}\left\{\sup_{t \in \mathbb{R} : F(t) \in C_{0}}\left\{\hat{F}_{n}(t) - F(t)\right\} > 1\right\} = 0.$$

Hence, the conclusion of Corollary 3 follows by a union bound.

4 Proof of the topological implicit function theorem

Proof of Lemma 4. First let's show that $\Gamma(x_0, y_0)(y_0 - \gamma(x_0)) > 0$ for all $x_0 \in \mathcal{X}$ and $y_0 \in \mathcal{Y} \setminus \{\gamma(x)\}$. If $y_0 \in \mathcal{Y}$ and $y_0 < \gamma(x)$ then $\Gamma(x_0, y_0) < 0$ and so $\Gamma(x_0, y_0)(y_0 - \gamma(x_0)) > 0$. On the other hand, if $y_0 \in \mathcal{Y}$ and $y_0 > \gamma(x_0)$ then $\gamma(x_0) = \inf\{y \in \mathcal{Y} : \Gamma(x_0, y) \ge 0\} \in cl(\mathcal{Y})$. Since \mathcal{Y} is an interval we can choose $y_1 \in (\gamma(x_0), y_0) \subseteq \mathcal{Y}$, so that $\Gamma(x_0, y_1) \ge 0$ and hence $\Gamma(x_0, y_0) > 0$ since $y \mapsto \Gamma(x_0, y)$ is sign-monotonic.

We begin by showing that γ is upper semi-continuous. Fix some $x_0 \in \mathcal{X}$ and $\epsilon > 0$. If $\gamma(x_0) = \sup \mathcal{Y}$, then $\gamma(x_1) \leq \gamma(x_0)$ for all $x_1 \in \mathcal{X}$. Suppose then that $\gamma(x_0) = \inf\{y \in \mathcal{Y} : \Gamma(x_0, y) < 0\} < \sup \mathcal{Y}$ and we can choose $y_0 \in (\gamma(x_0), \gamma(x_0) + \epsilon) \cap \mathcal{Y}$. Since $y_0 > \gamma(x_0)$ we have $\Gamma(x_0, y_0) > 0$. By the continuity of $x \mapsto \Gamma(x, y_0)$ we deduce that there exists an open set $U \subseteq \mathbb{R}^d$ with $x_0 \in U \cap \mathcal{X}$ and for all $x_1 \in U \cap \mathcal{X}$ we have $\Gamma(x_1, y_0) > 0$. As such, we have $\gamma(x_1) \leq y_0 < \gamma(x_0) + \epsilon$. Thus, γ is upper semi-continuous.

Next, we show that γ is lower semi-continuous. Again, we begin by choosing $x_0 \in \mathcal{X}$ and $\epsilon > 0$. If $\gamma(x_0) = \inf \mathcal{Y}$, then $\gamma(x_1) \geq \gamma(x_0)$ for all $x_1 \in \mathcal{X}$. Suppose that $\gamma(x_0) > \inf \mathcal{Y}$ so that we can choose $y_0 \in (\gamma(x_0) - \epsilon, \gamma(x_0)) \cap (\inf \mathcal{Y}, \gamma(x_0)) \subseteq (\inf \mathcal{Y}, \sup \mathcal{Y}) \subseteq \mathcal{Y}$, where we have used the fact that \mathcal{Y} is an interval. Since $y_0 < \gamma(x_0)$ we have $\Gamma(x_0, y_0) < 0$. Hence, by the continuity of $x \mapsto \Gamma(x, y_0)$, there exists an open set $U \subseteq \mathbb{R}^d$ with $x_0 \in U \cap \mathcal{X}$ and for all $x_1 \in U \cap \mathcal{X}$ we have $\Gamma(x_1, y_0) < 0$. Thus, applying the sign-monotonicity of $y \mapsto \Gamma(x_0, y)$, we have $\gamma(x_0) - \epsilon < y_0 \leq \inf\{y \in \mathcal{Y} : \Gamma(x_1, y) \geq 0\} \leq \sup \mathcal{Y}$. Hence, $\gamma(x_1) > \gamma(x_0) - \epsilon$, and we have shown that γ is also lower semi-continuous.

References

- Bartl, D. and Mendelson, S. (2023). On a variance dependent Dvoretzky-Kiefer-Wolfowitz inequality. arXiv preprint arXiv:2308.04757.
- Bennett, G. (1962). Probability inequalities for the sum of independent random variables. Journal of the American Statistical Association, 57(297):33–45.
- Bernstein, S. (1924). On a modification of Chebyshev's inequality and of the error formula of laplace. Ann. Sci. Inst. Sav. Ukraine, Sect. Math, 1(4):38–49.
- Birnbaum, Z. and McCarty, R. (1958). A distribution-free upper confidence bound for $\Pr\{Y < X\}$, based on independent samples of X and Y. The Annals of Mathematical Statistics, pages 558–562.
- Blanchard, M. and Voráček, V. (2023). Tight bounds for local Glivenko–Cantelli. arXiv preprint arXiv:2308.01896.
- Bretagnolle, J. and Massart, P. (1989). Hungarian Constructions from the Nonasymptotic Viewpoint. *The Annals of Probability*, 17(1):239 256.
- Cantelli, F. P. (1933). Sulla determinazione empirica delle leggi di probabilita. *Giorn. Ist. Ital. Attuari*, 4(421-424).
- Devroye, L. P. and Wise, G. L. (1979). On the recovery of discrete probability densities from imperfect measurements. *Journal of the Franklin Institute*, 307(1):1–20.

- Dudley, R. M. (2014). Uniform Central Limit Theorems, volume 142. Cambridge University press.
- Dudley, R. M. (2018). Real Analysis and Probability. CRC Press, Cambridge.
- Dvoretzky, A., Kiefer, J., and Wolfowitz, J. (1956). Asymptotic minimax character of the sample distribution function and of the classical multinomial estimator. *The Annals of Mathematical Statistics*, 27(3):642–669.
- Fedotov, A. A., Harremoës, P., and Topsoe, F. (2003). Refinements of Pinsker's inequality. *IEEE Transactions on Information Theory*, 49(6):1491–1498.
- Glivenko, V. (1933). Sulla determinazione empirica delle leggi di probabilita. *Gion. Ist. Ital. Attauri.*, 4:92–99.
- Hoeffding, W. (1963). Probability inequalities for sums of bounded random variables. *Journal* of the American Statistical Association, 58(301):13–30.
- Hu, I. (1985). A Uniform Bound for the Tail Probability of Kolmogorov-Smirnov Statistics. The Annals of Statistics, 13(2):821 – 826.
- Maillard, O.-A. (2021). Local Dvoretzky–Kiefer–Wolfowitz confidence bands. Mathematical Methods of Statistics, 30(1):16–46.
- Massart, P. (1990). The tight constant in the Dvoretzky-Kiefer-Wolfowitz Inequality. *The* Annals of Probability, 18(3):1269 – 1283.
- Shorack, G. R. and Wellner, J. A. (1986). *Empirical Processes with Applications to Statistics*. SIAM.
- Smirnov, N. V. (1944). Approximate laws of distribution of random variables from empirical data. Uspekhi Matematicheskikh Nauk, (10):179–206.