

PROBABILISTIC BIVARIATE BELL POLYNOMIALS

TAEKYUN KIM AND DAE SAN KIM

ABSTRACT. Let Y be a random variable whose moment generating function exists in some neighborhood of the origin. We consider the probabilistic bivariate Bell polynomials associated with Y and the probabilistic bivariate r -Bell polynomials associated with Y . For those polynomials, we derive the recurrence relations corresponding to the ones found by Zheng and Li for the bivariate Bell polynomials and the bivariate r -Bell polynomials.

1. INTRODUCTION

The Stirling number of the second kind $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ enumerates the number of ways to partition a set of n objects into k nonempty subsets (see (1)). More generally, for any $r \in \mathbb{N}$, the r -Stirling number of the second kind $\left\{ \begin{smallmatrix} n+r \\ k+r \end{smallmatrix} \right\}_r$ counts the number of ways to partition a set of $n+r$ objects into $k+r$ nonempty subsets such that the first r elements are in distinct subsets (see (2)).

Spivey discovered a remarkable recurrence relation for the Bell numbers (see (4)). This relation was extended to the Bell polynomials by Gould and Quaintance (see (5)). The corresponding ones for the bivariate Bell polynomials $\phi_n(x, y) = \sum_{k=0}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} (x)_k y^k$ and the bivariate r -Bell polynomials $\phi_{n,r}(x, y) = \sum_{k=0}^n \left\{ \begin{smallmatrix} n+r \\ k+r \end{smallmatrix} \right\}_r (x)_k y^k$ were found by Zheng and Li (see (9), (10)).

Assume that Y is a random variable whose moment generating function exists in some neighbourhood of the origin. Let $(Y_j)_{j \geq 1}$ be a sequence of mutually independent copies of the random variable Y , and let $S_k = Y_1 + Y_2 + \cdots + Y_k$, ($k \geq 1$), with $S_0 = 0$. In this paper, we introduce bivariate extensions of the probabilistic Bell polynomials associated Y and the probabilistic r -Bell polynomials associated with Y , namely the probabilistic bivariate Bell polynomials associated with Y , $\phi_n^Y(x, y)$ (see (16)) and the probabilistic bivariate r -Bell polynomials associated with Y , $\phi_{n,r}^Y(x, y)$ (see (20)). Then we derive the recurrence relations corresponding to the ones found by Zheng and Li for the probabilistic bivariate Bell polynomials associated with Y (see Theorem 2.2)

$$\begin{aligned} & \phi_{m+n}^Y(x, y) \\ &= \sum_{k=0}^n \sum_{j=0}^m \binom{m}{j} \phi_j^Y(x-k, y) (x)_k y^k \frac{1}{k!} \sum_{l_1 + \cdots + l_k = n} \binom{n}{l_1, \dots, l_k} E \left[S_k^{m-j} \prod_{i=1}^k Y_i^{l_i} \right], \end{aligned}$$

2010 *Mathematics Subject Classification.* 11B73; 11B83.

Key words and phrases. probabilistic bivariate Bell polynomials; probabilistic bivariate r -Bell polynomials.

and also for the probabilistic bivariate r -Bell polynomials associated with Y (see Theorems 2.4 and 2.5)

$$\begin{aligned} \phi_{m+n,r}^Y(x,y) &= \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} \phi_{i,r}^Y(x-k,y) \frac{(x)_k y^k}{k!} \sum_{j=k}^m \sum_{l_1+\dots+l_k=j} \binom{m}{j} \binom{j}{l_1, \dots, l_k} r^{m-j} E \left[S_k^{n-i} \prod_{i=1}^k Y_i^{l_i} \right] \\ &= \sum_{i=0}^n \sum_{k=0}^m \frac{1}{k!} \binom{n}{i} (x)_k y^k \phi_i^Y(x-k,y) \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} E \left[(S_k+r)^{n-i} \prod_{i=1}^k Y_i^{l_i} \right] r^{m-j}. \end{aligned}$$

These are done by using the simple formula for Taylor series (see (11)). For the rest of this section, we recall the facts that are needed throughout this paper.

For $n \geq 0$, the Stirling numbers of the second kind are given by

$$(1) \quad x^n = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} (x)_k, \quad (\text{see [1-29]}),$$

where $(x)_0 = 1$, $(x)_k = x(x-1)\cdots(x-k+1)$, $(k \geq 1)$.

The r -Stirling number of the second kind are defined as

$$(2) \quad (x+r)^n = \sum_{k=0}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r (x)_k, \quad (n \geq 0), \quad (\text{see [8, 18-20]}).$$

The Bell polynomials are given by

$$(3) \quad \phi_n(x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k, \quad (n \geq 0), \quad (\text{see [2, 10, 17, 25, 28]}).$$

When $x = 1$, $\phi_n = \phi_n(1)$ are called the Bell numbers.

Spivey found the following recurrence relation for ϕ_n :

$$(4) \quad \phi_{l+n} = \sum_{k=0}^l \sum_{i=0}^n k^{n-i} \binom{n}{i} \left\{ \begin{matrix} l \\ k \end{matrix} \right\} \phi_i, \quad (\text{see [28]}).$$

Gould and Quaintance discovered the following extension of the relation for ϕ_n in (4) to that for $\phi_n(x)$:

$$(5) \quad \phi_{l+n}(x) = \sum_{k=0}^l \sum_{i=0}^n k^{n-i} \binom{n}{i} \left\{ \begin{matrix} l \\ k \end{matrix} \right\} \phi_i(x) x^k, \quad (\text{see [12, 28]}).$$

For $r \in \mathbb{N}$, the r -Bell polynomials are defined as

$$(6) \quad \phi_{n,r}(x) = \sum_{k=0}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r x^k, \quad (n \geq 0), \quad (\text{see [18, 20]}).$$

It is known that the bivariate Bell polynomials are given by

$$(7) \quad \phi_n(x,y) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} (x)_k y^k, \quad (\text{see [30]}).$$

For $r \in \mathbb{N}$, the bivariate r -Bell polynomials are defined as

$$(8) \quad \phi_{n,r}(x,y) = \sum_{k=0}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r (x)_k y^k, \quad (\text{see [30]}).$$

Note that

$$\lim_{y \rightarrow 0} \phi_n \left(\frac{x}{y}, y \right) = \phi_n(x), \quad \lim_{y \rightarrow 0} \phi_{n,r} \left(\frac{x}{y}, y \right) = \phi_{n,r}(x), \quad (n \geq 0).$$

From (7) and (8), we note that

$$(9) \quad \phi_{m+n}(x, y) = \sum_{k=0}^m \sum_{i=0}^n k^{n-i} \binom{n}{i} \left\{ \begin{matrix} m \\ k \end{matrix} \right\} \phi_i(x-k, y) (x)_k y^k,$$

$$(10) \quad \phi_{m+n,r}(x, y) = \sum_{k=0}^m \sum_{i=0}^n k^{n-i} \binom{n}{i} \left\{ \begin{matrix} m+r \\ k+r \end{matrix} \right\}_r \phi_{i,r}(x-k, y) (x)_k y^k, \quad (\text{see [30]}),$$

where m, n are nonnegative integers.

We observe that

$$(11) \quad \begin{aligned} f(x+t) &= \sum_{n=0}^{\infty} \frac{f^{(n)}(x)}{n!} (x+t-x)^n = \sum_{n=0}^{\infty} \frac{f^{(n)}(x)}{n!} t^n \\ &= \sum_{n=0}^{\infty} \frac{t^n D_x^n}{n!} f(x) = e^{tD_x} f(x), \end{aligned}$$

where $D_x = \frac{d}{dx}$.

Assume that Y is a random variable such that the moment generating function of Y ,

$$E[e^{tY}] = \sum_{n=0}^{\infty} \frac{t^n}{n!} E[Y^n], \quad (|t| < r),$$

exists for some $r > 0$, (see [3,16,17]). Let $(Y_j)_{j \geq 1}$ be a sequence of mutually independent copies of the random variable Y , and let $S_k = Y_1 + Y_2 + \cdots + Y_k$, ($k \geq 1$), with $S_0 = 0$.

We recall that the probabilistic Stirling numbers of the second kind associated with Y are given by

$$(12) \quad \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_Y = \frac{1}{k!} \sum_{i=0}^k \binom{k}{i} (-1)^{k-i} E[S_i^n], \quad (0 \leq k \leq n), \quad (\text{see [3, 16, 17]}).$$

The probabilistic Bell polynomials associated with Y are defined as

$$(13) \quad \phi_n^Y(x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_Y x^k, \quad (n \geq 0), \quad (\text{see [16, 17, 27]}).$$

For $r \in \mathbb{N}$, the probabilistic r -Stirling numbers of the second kind associated with Y are given by

$$(14) \quad \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} = \frac{1}{k!} \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} E[(S_j+r)^n], \quad (n \geq k \geq 0), \quad (\text{see [16, 17]}).$$

When $Y = 1$, we have

$$\left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} = \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r, \quad (n \geq k \geq 0).$$

The probabilistic r -Bell polynomials associated with Y are defined as

$$(15) \quad \phi_{n,r}^Y(x) = \sum_{k=0}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} x^k, \quad (n \geq 0), \quad (\text{see [16, 17]}).$$

When $Y = 1$, we have $\phi_{n,r}^Y(x) = \phi_{n,r}(x)$, ($n \geq 0$).

2. PROBABILISTIC BIVARIATE BELL POLYNOMIALS

Let $(Y_j)_{j \geq 1}$ be a sequence of mutually independent copies of the random variable Y , and let

$$S_0 = 0, \quad S_k = Y_1 + Y_2 + \cdots + Y_k, \quad (k \geq 1).$$

Now, we define the *probabilistic bivariate Bell polynomials associated with Y* by

$$(16) \quad \left(1 + y(E[e^{Yt}] - 1)\right)^x = \sum_{n=0}^{\infty} \phi_n^Y(x, y) \frac{t^n}{n!}.$$

From (16), we note that

$$(17) \quad \begin{aligned} \sum_{n=0}^{\infty} \phi_n^Y(x, y) \frac{t^n}{n!} &= \sum_{k=0}^{\infty} (x)_k y^k \frac{1}{k!} (E[e^{Yt}] - 1)^k \\ &= \sum_{k=0}^{\infty} (x)_k y^k \sum_{n=k}^{\infty} \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_Y \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_Y (x)_k y^k \right) \frac{t^n}{n!}. \end{aligned}$$

Therefore, by comparing the coefficients on both sides of (17), we obtain the following theorem.

Theorem 2.1. *For $n \geq 0$, we have*

$$\phi_n^Y(x, y) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_Y (x)_k y^k.$$

When $Y = 1$, $\phi_n^Y(x, y) = \phi_n(x, y)$. As $\lim_{y \rightarrow 0} \phi_n^Y(x, y) = \phi_n^Y(x)$, the probabilistic bivariate Bell polynomials $\phi_n^Y(x, y)$ are also a bivariate extension of the probabilistic Bell polynomials $\phi_n^Y(x)$.

From (16), we note that

$$(18) \quad \begin{aligned} e^{sD_t} \left(1 + y(E[e^{Yt}] - 1)\right)^x &= \sum_{n=0}^{\infty} \frac{s^n}{n!} D_t^n \sum_{m=0}^{\infty} \phi_m^Y(x, y) \frac{t^m}{m!} \\ &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \phi_{m+n}^Y(x, y) \frac{s^n}{n!} \frac{t^m}{m!}. \end{aligned}$$

On the other hand, by (11), we get

$$\begin{aligned}
 (19) \quad & e^{sDt} \left(1 + y(E[e^{Yt}] - 1)\right)^x = \left(1 + y(E[e^{(s+t)Y}] - 1)\right)^x \\
 & = \left(1 + y(E[e^{Yt}] - 1) + yE[e^{Yt}(e^{sY} - 1)]\right)^x \\
 & = \sum_{k=0}^{\infty} \binom{x}{k} \left(1 + y(E[e^{Yt}] - 1)\right)^{x-k} y^k \left(E[e^{Yt}(e^{sY} - 1)]\right)^k \\
 & = \sum_{k=0}^{\infty} \frac{(x)_k}{k!} y^k \sum_{j=0}^{\infty} \phi_j^Y(x-k, y) \frac{t^j}{j!} E\left[e^{sY_1}(e^{sY_2} - 1) \cdots (e^{sY_k} - 1)\right] \\
 & = \sum_{k=0}^{\infty} \frac{(x)_k}{k!} y^k \sum_{n=k}^{\infty} \sum_{l_1+l_2+\dots+l_k=n} \binom{n}{l_1, l_2, \dots, l_k} E\left[e^{sY_1^{l_1} Y_2^{l_2} \cdots Y_k^{l_k}}\right] \frac{s^n}{n!} \sum_{j=0}^{\infty} \phi_j^Y(x-k, y) \frac{t^j}{j!} \\
 & = \sum_{n=0}^{\infty} \sum_{k=0}^n (x)_k y^k \frac{1}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} \frac{s^n}{n!} \sum_{l=0}^{\infty} E\left[S_k^l \prod_{i=1}^k Y_i^{l_i}\right] \frac{t^l}{l!} \sum_{j=0}^{\infty} \phi_j^Y(x-k, y) \frac{t^j}{j!} \\
 & = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \sum_{k=0}^n \sum_{j=0}^m \binom{m}{j} \phi_j^Y(x-k, y) (x)_k y^k \frac{1}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} E\left[S_k^{m-j} \prod_{i=1}^k Y_i^{l_i}\right] \frac{s^n}{n!} \frac{t^m}{m!},
 \end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers.

Therefore, by (18) and (19), we obtain the following theorem.

Theorem 2.2. For $m, n \geq 0$, we have

$$\begin{aligned}
 & \phi_{m+n}^Y(x, y) \\
 & = \sum_{k=0}^n \sum_{j=0}^m \binom{m}{j} \phi_j^Y(x-k, y) (x)_k y^k \frac{1}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} E\left[S_k^{m-j} \prod_{i=1}^k Y_i^{l_i}\right],
 \end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers.

When $Y = 1$, the expression in Theorem 2.2 reduces to

$$\begin{aligned}
 \phi_{m+n}(x, y) & = \sum_{k=0}^n \sum_{j=0}^m \binom{m}{j} \phi_j(x-k, y) (x)_k y^k \frac{1}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} k^{m-j} \\
 & = \sum_{k=0}^n \sum_{j=0}^m \binom{m}{j} \phi_j(x-k, y) (x)_k y^k \left\{ \begin{matrix} n \\ k \end{matrix} \right\} k^{m-j}.
 \end{aligned}$$

For $r \in \mathbb{N}$, we define the *probabilistic bivariate r -Bell polynomials associated with Y* by

$$(20) \quad \left(1 + y(E[e^{Yt}] - 1)\right)^x e^{rt} = \sum_{n=0}^{\infty} \phi_{n,r}^Y(x, y) \frac{t^n}{n!}.$$

From (14), we note that

$$\begin{aligned}
 (21) \quad \sum_{n=0}^{\infty} \phi_{n,r}^Y(x, y) \frac{t^n}{n!} & = \left(1 + y(E[e^{Yt}] - 1)\right)^x e^{rt} = \sum_{k=0}^{\infty} (x)_k \frac{y^k}{k!} (E[e^{Yt}] - 1)^k e^{rt} \\
 & = \sum_{k=0}^{\infty} y^k (x)_k \sum_{n=k}^{\infty} \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} \frac{t^n}{n!} = \sum_{n=0}^{\infty} \sum_{k=0}^n y^k (x)_k \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} \frac{t^n}{n!}.
 \end{aligned}$$

Therefore, by comparing the coefficients on both sides of (21), we obtain the following theorem.

Theorem 2.3. *For $n \geq 0$, we have*

$$\phi_{n,r}^Y(x,y) = \sum_{k=0}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} (x)_k y^k.$$

When $Y = 1$, we obtain

$$\phi_{n,r}(x,y) = \sum_{k=0}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r (x)_k y^k.$$

In addition, we note that

$$\lim_{y \rightarrow 0} \phi_{n,r}^Y\left(\frac{x}{y}, y\right) = \sum_{k=0}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} x^k = \phi_{n,r}^Y(x).$$

By using (11), we have

(22)

$$\begin{aligned} e^{tD_s} \left(1 + y(E[e^{sY}] - 1)\right)^x e^{rs} &= \left(1 + y(E[e^{(s+t)Y}] - 1)\right)^x e^{(s+t)r} \\ &= \sum_{k=0}^{\infty} \binom{x}{k} \left(1 + y(E[e^{tY}] - 1)\right)^{x-k} e^{rt} y^k \left(E[e^{tY}(e^{sY} - 1)]\right)^k e^{sr} \\ &= \sum_{k=0}^{\infty} \frac{(x)_k y^k}{k!} \sum_{i=0}^{\infty} \phi_{i,r}^Y(x-k, y) \frac{t^i}{i!} \sum_{j=k, l_1+\dots+l_k=j}^{\infty} \binom{j}{l_1, \dots, l_k} E\left[e^{S_{kt} \prod_{i=1}^k Y_i^{l_i}}\right] \frac{s^j}{j!} e^{sr} \\ &= \sum_{k=0}^{\infty} \frac{(x)_k y^k}{k!} \sum_{m=k}^{\infty} \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} E\left[e^{S_{kt} \prod_{i=1}^k Y_i^{l_i}}\right] r^{m-j} \frac{s^m}{m!} \sum_{i=0}^{\infty} \phi_{i,r}^Y(x-k, y) \frac{t^i}{i!} \\ &= \sum_{m=0}^{\infty} \sum_{l=0}^{\infty} \sum_{k=0}^m \frac{(x)_k y^k}{k!} \sum_{j=k, l_1+\dots+l_k=j}^m \binom{m}{j} \binom{j}{l_1, \dots, l_k} r^{m-j} \frac{s^m}{m!} E\left[S_k^l \prod_{i=1}^k Y_i^{l_i}\right] \frac{t^l}{l!} \sum_{i=0}^{\infty} \phi_{i,r}^Y(x-k, y) \frac{t^i}{i!} \\ &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} \phi_{i,r}^Y(x-k, y) \frac{(x)_k y^k}{k!} \\ &\quad \times \sum_{j=k, l_1+\dots+l_k=j}^m \binom{m}{j} \binom{j}{l_1, \dots, l_k} r^{m-j} E\left[S_k^{n-i} \prod_{i=1}^k Y_i^{l_i}\right] \frac{t^n s^m}{n! m!}, \end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers.

On the other hand, by (20), we get

$$\begin{aligned} (23) \quad e^{tD_s} \left(1 + y(E[e^{sY}] - 1)\right)^x e^{rs} &= \sum_{n=0}^{\infty} \frac{t^n}{n!} D_s^n \sum_{m=0}^{\infty} \phi_{m,r}^Y(x, y) \frac{s^m}{m!} \\ &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \phi_{m+n,r}^Y(x, y) \frac{t^n s^m}{n! m!}. \end{aligned}$$

Therefore, by (22) and (23), we obtain the following theorem.

Theorem 2.4. *For $m, n \geq 0$, we have*

$$\begin{aligned} &\phi_{m+n,r}^Y(x, y) \\ &= \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} \phi_{i,r}^Y(x-k, y) \frac{(x)_k y^k}{k!} \sum_{j=k, l_1+\dots+l_k=j}^m \binom{m}{j} \binom{j}{l_1, \dots, l_k} r^{m-j} E\left[S_k^{n-i} \prod_{i=1}^k Y_i^{l_i}\right], \end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers.

When $Y = 1$, the expression in Theorem 2.4 boils down to

$$\begin{aligned}\phi_{m+n,r}(x,y) &= \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} \phi_{i,r}(x-k,y) \frac{(x)_k y^k}{k!} \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} r^{m-j} k^{n-i} \\ &= \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} \phi_{i,r}(x-k,y) (x)_k y^k k^{n-i} \left\{ \begin{matrix} m+r \\ k+r \end{matrix} \right\}_r.\end{aligned}$$

Now, we observe that

$$\begin{aligned}(24) \quad e^{tD_s} \left(1 + y(E[e^{sY}] - 1) \right)^x e^{rs} &= \left(1 + y(E[e^{(s+t)Y}] - 1) \right)^x e^{r(s+t)} \\ &= \sum_{k=0}^{\infty} \frac{(x)_k y^k}{k!} \left(1 + y(E[e^{tY}] - 1) \right)^{x-k} E \left[e^{(S_k+r)t} \prod_{i=1}^k (e^{sY_i} - 1) \right] e^{sr} \\ &= \sum_{k=0}^{\infty} \frac{(x)_k y^k}{k!} \sum_{i=0}^{\infty} \phi_i^Y(x-k,y) \frac{t^i}{i!} \sum_{j=k}^{\infty} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} E \left[e^{(S_k+r)t} \prod_{i=1}^k Y_i^{l_i} \right] \frac{s^j}{j!} e^{sr} \\ &= \sum_{k=0}^{\infty} \frac{(x)_k y^k}{k!} \sum_{m=k}^{\infty} \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} E \left[e^{(S_k+r)t} \prod_{i=1}^k Y_i^{l_i} \right] r^{m-j} \frac{s^m}{m!} \sum_{i=0}^{\infty} \phi_i^Y(x-k,y) \frac{t^i}{i!} \\ &= \sum_{m=0}^{\infty} \sum_{k=0}^m \frac{(x)_k y^k}{k!} \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} E \left[e^{(S_k+r)t} \prod_{i=1}^k Y_i^{l_i} \right] r^{m-j} \frac{s^m}{m!} \sum_{i=0}^{\infty} \phi_i^Y(x-k,y) \frac{t^i}{i!} \\ &= \sum_{m=0}^{\infty} \sum_{l=0}^{\infty} \sum_{k=0}^m \frac{(x)_k y^k}{k!} \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} r^{m-j} \frac{s^m}{m!} \\ &\quad \times E \left[(S_k+r)^l \prod_{i=1}^k Y_i^{l_i} \right] \frac{t^l}{l!} \sum_{i=0}^{\infty} \phi_i^Y(x-k,y) \frac{t^i}{i!} \\ &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} \frac{(x)_k y^k}{k!} \phi_i^Y(x-k,y) \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} \\ &\quad \times E \left[(S_k+r)^{n-i} \prod_{i=1}^k Y_i^{l_i} \right] r^{m-j} \frac{s^m}{m!} \frac{t^n}{n!},\end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers.

Therefore, by (23) and (24), we obtain the following theorem.

Theorem 2.5. For $m, n \geq 0$, we have

$$\begin{aligned}\phi_{m+n,r}^Y(x,y) &= \sum_{i=0}^n \sum_{k=0}^m \frac{1}{k!} \binom{n}{i} (x)_k y^k \phi_i^Y(x-k,y) \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} E \left[(S_k+r)^{n-i} \prod_{i=1}^k Y_i^{l_i} \right] r^{m-j},\end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers.

When $Y = 1$, the expression in Theorem 2.5 reduces to

$$\begin{aligned} & \phi_{m+n,r}(x,y) \\ &= \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} (x)_k y^k \phi_i(x-k,y) \frac{1}{k!} \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} r^{m-j} (r+k)^{n-i} \\ &= \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} (x)_k y^k \phi_i(x-k,y) \left\{ \begin{matrix} m+r \\ k+r \end{matrix} \right\}_r (r+k)^{n-i}. \end{aligned}$$

From Theorem 2.4, we note that

(25)

$$\begin{aligned} \phi_{m+n,r}^Y(x) &= \lim_{y \rightarrow 0} \phi_{m+n}^Y\left(\frac{x}{y}, y\right) \\ &= \lim_{y \rightarrow 0} \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} \phi_{i,r}^Y\left(\frac{x}{y} - k, y\right) \left(\frac{x}{y}\right)_k y^k \frac{1}{k!} \\ &\quad \times \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} r^{m-j} E\left[S_k^{n-i} \prod_{i=1}^k Y_i^{l_i}\right] \\ &= \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} \phi_{i,r}^Y(x) x^k \frac{1}{k!} \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} r^{m-j} E\left[S_k^{n-i} \prod_{i=1}^k Y_i^{l_i}\right], \end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers.

Therefore, by (25), we obtain the following theorem.

Theorem 2.6. For $m, n \geq 0$, we have

$$\begin{aligned} & \phi_{m+n,r}^Y(x) \\ &= \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} \phi_{i,r}^Y(x) x^k \frac{1}{k!} \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} r^{m-j} E\left[S_k^{n-i} \prod_{i=1}^k Y_i^{l_i}\right], \end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers.

From Theorem 2.5, we get

(26)

$$\begin{aligned} \phi_{m+n,r}^Y(x) &= \lim_{y \rightarrow 0} \phi_{m+n,r}^Y\left(\frac{x}{y}, y\right) \\ &= \lim_{y \rightarrow 0} \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} \left(\frac{x}{y}\right)_k y^k \phi_i^Y\left(\frac{x}{y} - k, y\right) \frac{1}{k!} \sum_{j=k}^m \binom{m}{j} \\ &\quad \times \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} E\left[(S_k+r)^{n-i} \prod_{i=1}^k Y_i^{l_i}\right] r^{m-j} \\ &= \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} x^k \phi_i^Y(x) \frac{1}{k!} \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} E\left[(S_k+r)^{n-i} \prod_{i=1}^k Y_i^{l_i}\right] r^{m-j}, \end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers.

Therefore, by (26), we obtain the following theorem.

Theorem 2.7. For $m, n \geq 0$, we have

$$\phi_{m+n,r}^Y(x) = \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} x^k \phi_i^Y(x) \frac{1}{k!} \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} \binom{j}{l_1, \dots, l_k} E \left[(S_k+r)^{n-i} \prod_{i=1}^k Y_i^{l_i} \right] r^{m-j},$$

where l_1, l_2, \dots, l_k are positive integers.

When $Y = 1$, the expression in Theorem 2.7 boils down to

$$\begin{aligned} \phi_{m+n}(x) &= \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} x^k \phi_i(x) \frac{1}{k!} \sum_{j=k}^m \binom{m}{j} \sum_{l_1+\dots+l_k=j} r^{m-j} (k+r)^{n-i} \\ &= \sum_{i=0}^n \sum_{k=0}^m \binom{n}{i} x^k \phi_i(x) \left\{ \begin{matrix} m+r \\ k+r \end{matrix} \right\}_r (k+r)^{n-i}. \end{aligned}$$

3. CONCLUSION

Let Y be a random variable such that the moment generating function of Y exists in a neighborhood of the origin. We introduced probabilistic extensions of the bivariate Bell polynomials and the bivariate r -Bell polynomials, namely the probabilistic bivariate Bell polynomials associated with Y and the probabilistic bivariate r -Bell polynomials associated with Y . Zheng and Li found recurrence relations for the bivariate Bell polynomials and the bivariate r -Bell polynomials. In this paper, by utilizing the simple fact (11) we obtained recurrence relations for the probabilistic bivariate Bell polynomials associated with Y and the probabilistic bivariate r -Bell polynomials associated with Y . These relations boil down to the relations found by Zheng and Li when $Y = 1$.

As one of our future projects, we would like to continue to study probabilistic extensions of many special polynomials and numbers and to find their applications to physics, science and engineering as well as to mathematics.

4. DECLARATIONS

The authors declare that they have no conflicts of interest to report regarding the present study.

REFERENCES

- [1] Abbas, M.; Bouroubi, S. *On new identities for Bell's polynomials*, Discrete Math. **293** (2005), no. 1-3, 5-10.
- [2] Abramowitz, M.; Stegun, I.A. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Dover, New York, 1992.
- [3] Adell, J. A. *Probabilistic Stirling numbers of the second kind and applications*, J. Theoret. Probab. **35** (2022), no. 1, 636-652.
- [4] Belbachir, H.; Bousbaa, I. E. *Translated Whitney and r -Whitney numbers: a combinatorial approach*, J. Integer Seq. **16** (2013), no. 8, Article 13.8.6, 7 pp.
- [5] Boubellouta, K.; Boussayoud, A.; Araci, S.; Kerada, M. *Some theorems on generating functions and their applications*, Adv. Stud. Contemp. Math. (Kyungshang) **30** (2020), no. 3, 307-324.
- [6] Broder, A. Z. *The r -Stirling numbers*, Discrete Math. **49** (1984), no. 3, 241-259.
- [7] Carlitz, L. *Some remarks on the Bell numbers*, Fibonacci Quart. **18** (1980), no. 1, 66-73.
- [8] Carlitz, L. *Weighted Stirling numbers of the first and second kind II*, Fibonacci Quart. **18** (1980), no. 3, 242-257.
- [9] Carlitz, L. *Weighted Stirling numbers of the first and second kind I*, Fibonacci Quart. **18** (1980), no. 2, 147-162.
- [10] Comtet, L. *Advanced combinatorics. The art of finite and infinite expansions*, Revised and enlarged edition, D. Reidel Publishing Co., Dordrecht, 1974.

- [11] Gould, H. W.; Quaintance, J. *Implications of Spivey's Bell number formula*, J. Integer Seq. **11** (2008), no. 3, Article 08.3.7, 6 pp.
- [12] Graham, R. L.; Knuth, D. E.; Patashnik, O. *Concrete mathematics. A foundation for computer science*, Second edition, Addison-Wesley Publishing Company, Reading, MA, 1994.
- [13] Kilar, N.; Simsek, Y. *Combinatorial sums involving Fubini type numbers and other special numbers and polynomials: approach trigonometric functions and p-adic integrals*, Adv. Stud. Contemp. Math. (Kyungshang) **31** (2021), no. 1, 75-87.
- [14] Kim, D. S.; Kim, T. *Normal ordering associated with λ -Whitney numbers of the first kind in λ -shift algebra*. Russ. J. Math. Phys. **30** (2023), no. 3, 310-319.
- [15] Kim, T.; Kim, D. S. *Representation by degenerate Frobenius-Euler polynomials*, Georgian Math. J. **29** (2022), no. 5, 741-754.
- [16] Kim, T.; Kim, D. S. *Probabilistic degenerate Bell polynomials associated with random variables*, Russ. J. Math. Phys. **30** (2023), no. 4, 528-542.
- [17] Kim, T.; Kim, D. S.; Kwon, J. *Probabilistic degenerate Stirling polynomials of the second kind and their applications*, Math. Comput. Model. Dyn. Syst. **30** (2024), no. 1, 16-30.
- [18] Kim, T.; Kim, D. S. *Some identities on degenerate r-Stirling numbers via boson operators*, Russ. J. Math. Phys. **29** (2022), no. 4, 508-517.
- [19] Kim, T.; Kim, D. S. *Degenerate Whitney numbers of first and second kind of Dowling lattices*, Russ. J. Math. Phys. **29** (2022), no. 3, 358-377.
- [20] Kim, T.; Kim, D. S. *Degenerate r-Whitney numbers and degenerate r-Dowling polynomials via boson operators*, Adv. in Appl. Math. **140** (2022), Paper No. 102394, 21 pp.
- [21] Kim, T.; Kim, D. S. *Combinatorial identities involving degenerate harmonic and hyperharmonic numbers*, Adv. Appl. Math. **148** (2023), Paper No. 102535, 15 pp.
- [22] Kim, T.; Kim, D. S. *On some degenerate differential and degenerate difference operators*, Russ. J. Math. Phys. **29** (2022), no. 1, 37-46.
- [23] Riordan, J. *An introduction to combinatorial analysis*, Dover Publications, Inc., Mineola, NY, 2002.
- [24] Riordan, J. *Combinatorial identities*, John Wiley & Sons, Inc., New York-London-Sydney, 1968.
- [25] Roman, S. *The umbral calculus*, Pure and Applied Mathematics 111, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1984.
- [26] Ross, S. M. *Introduction to probability models*, Thirteenth edition, Academic Press, London, 2024.
- [27] Soni, R.; Vellaisamy, P.; Pathak, A. K. *A probabilistic generalization of the Bell polynomials*, J. Anal. (2023). <https://doi.org/10.1007/s41478-023-00642-y>
- [28] Spivey, M. Z. *A generalized recurrence for Bell numbers*. J. Integer Seq. **11** (2008), no. 2, Article 08.2.5, 3 pp.
- [29] Ta, B. Q. *Probabilistic approach to Appell polynomials*, Expo. Math. **33** (2015), no. 3, 269-294.
- [30] Zheng, Y.; Li, N. N. *Bivariate extension of Bell polynomials*, J. Integer Seq. **22** (2019), no. 8, Art. 19.8.8, 8 pp.

DEPARTMENT OF MATHEMATICS, KWANGWOON UNIVERSITY, SEOUL 139-701, REPUBLIC OF KOREA

Email address: `tkkim@kw.ac.kr`

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 121-742, REPUBLIC OF KOREA

Email address: `dskim@sogang.ac.kr`