# COMBINATORIAL BASES IN QUANTUM TOROIDAL $\mathfrak{gl}_2$ MODULES

M. JIMBO AND E. MUKHIN

ABSTRACT. We show that many tame modules of the quantum toroidal  $\mathfrak{gl}_2$  algebra can be explicitly constructed in a purely combinatorial way using the theory of *q*-characters. The examples include families of evaluation modules obtained from analytic continuation and automorphism twists of Verma modules of the quantum affine  $\mathfrak{gl}_2$  algebra. The combinatorial bases in the modules are labeled by colored plane partitions with various properties.

#### 1. INTRODUCTION

The problem of finding a good basis in a representation is a central problem in representation theory. There are various approaches depending on the definition of the word "good". The theory of canonical [L90] and crystal bases [K90] in representations of quantum algebras and quantum affine algebras is compatible with the change  $q \rightarrow q^{-1}$  in an appropriate sense. The Gelfand-Tsetlin bases in symmetric groups, see e.g. [OV05], and in Lie algebras, see [GT50], respect the branching rules. The bases of Bethe vectors in various tensor products are eigenbases for a family of commuting Hamiltonians, [B31].

In this note, we discuss bases which we call combinatorial. A quantum affine algebra at level zero has a family of commutative Cartan currents. If a module is tame, that is if the Cartan currents are diagonalizable and each eigenspace is one-dimensional, we get (up to normalization) a combinatorial basis of eigenvectors.

In combinatorial bases, the action of the quantum affine algebra can be written explicitly (up to some constants depending on the normalization of the basis) since the matrix coefficients of generating series are essentially delta functions.

If a module is tame, the combinatorial basis (that is the set of eigenvalues of Cartan currents or the q-character) can be constructed recursively by the algorithm described in [FM1]. Then one needs to solve the recursion and fix the normalization constants – and this is one of the subjects of this article.

In type A, the combinatorial basis in all known cases is labeled by colored plane partitions with various boundary conditions. The creation and annihilation operators of a given color add or delete a box of the same color with explicit coefficients. We develop a general scheme of fixing the normalization constants, see (3.6), (3.7), and Theorem 3.4. Our method is based on viewing plane partitions as "tensor products" of boxes in a lexicographic order.

In general, the problem of determining the coefficients is not solved, see, for example, Section 5.3 of [FJM2].

Sometimes, one can construct a combinatorial basis in a module over a quantum algebra even if the level of the module is not zero or if the eigenvalues of Cartan currents are not distinct.

It happens if one manages to upgrade the action of the quantum algebra to a tame level zero module over an affinization of the quantum algebra. We list known examples.

- Any finite-dimensional irreducible  $U_q\mathfrak{gl}_n$  module can be considered as an evaluation module over  $U_q \widehat{\mathfrak{gl}}_n$ . The evaluation modules are tame and therefore one gets a combinatorial basis depending on a parameter. The *q*-character is known, Lemma 4.7 in [FM2], which gives a parameterization of the combinatorial basis by semi-standard Young tableaux. A semi-standard Young tableau can be interpreted as a plane partition over a Young diagram of height at most *n* with strictly increasing height along columns.
- The modules of minimal models of deformed affine W-algebras of type  $\mathfrak{gl}_n$ . In this case, the combinatorial answer is the cylindrical partitions, see [FFJMM]. It includes the deformed Virasoro algebra (multiplied by an additional Heisenberg current) when n = 2.
- Integrable irreducible  $U_q \widehat{\mathfrak{gl}}_n$ -modules of highest weight  $(0, \ldots, 0, q, 0, \ldots, 0)$  can be extended to a module over the quantum toroidal algebra of type  $\mathfrak{gl}_n$ . This extension depends on a parameter  $q_3$  of the toroidal algebra and it is an evaluation module only for  $q_3^n = q^2$ . The combinatorial basis is parameterized by colored partitions. The corresponding modules over toroidal algebras are known as Fock modules, [FJMM1].
- The supersymmetric analog of the previous item is also known, see [BrM]. The combinatorial basis is parameterized by colored super partitions.

We extend this list for the case of  $\mathfrak{gl}_2$  as follows.

- The  $U_q \widehat{\mathfrak{gl}}_2$  Verma modules. The corresponding evaluation module over the quantum toroidal algebra is tame. We compute the corresponding basis and matrix coefficients. The combinatorial set is the set of pairs of vertical partitions, see Figure 7.
- The relaxed  $U_q \mathfrak{gl}_2$  Verma modules. These are modules of generic level induced from a "one line" representation of  $U_q \mathfrak{gl}_2$ . The corresponding weight diagram is given in Figure 1. The evaluation module over the quantum toroidal algebra is tame and we compute the corresponding basis and matrix coefficients. The combinatorial set is the set of pairs of vertical partitions with an additional single tower over a box, see Figure 8.
- Slanted slope m relaxed  $U_q \widehat{\mathfrak{gl}}_2$  Verma modules,  $m \in \mathbb{Z}$ . These modules are obtained from the relaxed  $U_q \widehat{\mathfrak{gl}}_2$  Verma modules via a twist by an automorphism. The corresponding weight diagram is given in Figure 2. The evaluation module over the quantum toroidal algebra is tame and we compute the corresponding basis and matrix coefficients. The combinatorial set is the set of pairs of vertical partitions with an additional tower over a staircase consisting of 2|m| + 1 boxes, see Figure 10.

One may wonder how rich the class of modules with combinatorial bases is. In other words, how many tame modules do we have? Tame modules for  $U_q \widehat{\mathfrak{gl}}_n$  have been classified in [NT] and they correspond to (unions of) skew Young diagrams. Tame modules for  $U_q \widehat{\mathfrak{o}}_{2n+1}$  (type B) are also known, [BM], the corresponding combinatorial set is given by certain non-intersecting

#### COMBINATORIAL MODULES

paths. From the examples above, it is clear that for quantum toroidal algebras the set of tame modules is quite large and interesting and deserves further study.

Many constructions discussed in the paper can be applied to the case of  $\mathfrak{gl}_n$ , n > 2. We hope to address combinatorial bases of  $\mathfrak{gl}_n$  type in a separate publication.

We also expect, that one can define an affine crystal structure on the combinatorial sets corresponding to slanted relaxed Verma modules.

Our initial motivation was the study of integrable systems whose Hamiltonians are transfer matrices corresponding to quantum affine and quantum toroidal algebras. For highest weight modules, there is a powerful Bethe ansatz method to study the spectrum of the model. The relaxed  $U_q \widehat{\mathfrak{gl}}_2$  Verma modules form the most natural class of non-highest weight modules of generic level for the quantum affine algebra, see [FST]. One can consider them as analytic continuation of Verma modules and therefore one could try to understand the integrable system using continuation of the Bethe ansatz.

Combinatorially, the slanted relaxed Verma modules look as a natural generalization, however, it is not clear how to attack the corresponding integrable system (except for the cases of m = 0, 1).

The paper is constructed as follows. In Section 2 we recall the quantum algebra  $U_q\mathfrak{gl}_2$  and the quantum affine algebra  $U_q \mathfrak{gl}_2$  and give a few simple properties. In particular, in Section 2.3 we introduce the slanted relaxed Verma modules obtained from Verma modules by analytic continuation and automorphism twists. In Section 3, we discuss the quantum toroidal algebra  $\mathcal{E}$  associated to  $\mathfrak{gl}_2$ . In particular, in Section 3.3 we describe a way to construct combinatorial  $\mathcal{E}$ -modules, resulting in Theorem 3.4. In Section 4 we discuss various combinatorial  $\mathcal{E}$ -modules.

# 2. The quantum affine algebra associated to $\mathfrak{gl}_2$

Fix  $q, d \in \mathbb{C}^{\times}$  and set  $q_1 = q^{-1}d, q_2 = q^2, q_3 = q^{-1}d^{-1}$ . Then  $q_1q_2q_3 = 1$ . Fix  $\log q, \log d \in \mathbb{C}$ , so that  $q = e^{\log q}, d = e^{\log d}$ .

We assume that q, d are generic: for rational numbers a, b, the equality  $q^a d^b = 1$  holds if and only if a = b = 0.

We use the standard notation  $[A, B]_p = AB - pBA$  and  $[r] = (q^r - q^{-r})/(q - q^{-1})$ .

2.1. Quantum algebra  $U_q \mathfrak{gl}_2$ . Let  $\mathbb{C}P = \bigoplus_{i \in \mathbb{Z}/2\mathbb{Z}} \mathbb{C}\varepsilon_i$  be a 2-dimensional vector space with the chosen basis and a non-degenerate bilinear form such that  $(\varepsilon_i, \varepsilon_j) = \delta_{i,j}$ . Call the lattice  $P = \bigoplus_{i \in \mathbb{Z}/2\mathbb{Z}} \mathbb{Z}\varepsilon_i$  the weight lattice. Let  $\alpha = \varepsilon_1 - \varepsilon_2$ .

The quantum  $\mathfrak{gl}_2$  algebra  $U_q\mathfrak{gl}_2$  has generators  $e_1, f_1, q^h, h \in P$ , with the defining relations

$$q^{h}q^{h'} = q^{h+h'}, \qquad q^{0} = 1, \qquad q^{h}e_{1} = q^{(h,\alpha)}e_{1}q^{h}, \qquad q^{h}f_{1} = q^{-(h,\alpha)}f_{1}q^{h}, \quad [e_{1},f_{1}] = \frac{K - K^{-1}}{q - q^{-1}},$$

where  $K = q^{\alpha}$ .

The quantum  $\mathfrak{sl}_2$  algebra  $U_q\mathfrak{sl}_2$  is the subalgebra of  $U_q\mathfrak{gl}_2$  generated by  $e_1, f_1, K^{\pm 1}$ . The element  $\mathbf{t} = q^{\varepsilon_1 + \varepsilon_2} \in U_q\mathfrak{gl}_2$  is central and split.

The Casimir element  $\mathcal{C} = qK + q^{-1}K^{-1} + (q - q^{-1})^2 f_1 e_1$  is central in  $U_q \mathfrak{gl}_2$ .

2.2. Quantum affine algebra  $U_q \widehat{\mathfrak{gl}}_2$ . The quantum affine algebra  $U_q \widehat{\mathfrak{gl}}_2$  in the Drinfeld new realization is defined by generators  $x_k^{\pm}$ ,  $h_{j,r}$ ,  $q^h$ ,  $C^{\pm 1}$ , where  $j = 1, 2, k \in \mathbb{Z}, r \in \mathbb{Z} \setminus \{0\}, h \in P$ , with the defining relations

$$\begin{split} C^{\pm 1} &\text{ are central, } \quad CC^{-1} = 1, \quad q^h q^{h'} = q^{h+h'}, \quad q^0 = 1, \quad [q^h, h_{j,r}] = 0, \\ q^h x^{\pm}(z) q^{-h} &= q^{\pm (h,\alpha)} x^{\pm}(z), \\ [h_{i,r}, h_{j,s}] &= \delta_{r+s,0} \,\bar{a}_{i,j}(r) \, \frac{C^r - C^{-r}}{q - q^{-1}}, \\ [h_{i,r}, x^{\pm}(z)] &= \pm \,\bar{a}_{i,j}(r) \, C^{-(r\pm |r|)/2} z^r x^{\pm}(z), \\ [x^+(z), x^-(w)] &= \frac{1}{q - q^{-1}} \left( \delta\left(C\frac{w}{z}\right) \phi^+(w) - \delta\left(C\frac{z}{w}\right) \phi^-(z) \right), \\ (z - q^{\pm 2}w) x^{\pm}(z) x^{\pm}(w) + (w - q^{\pm 2}z) x^{\pm}(w) x^{\pm}(z) = 0. \end{split}$$

Here we set  $x^{\pm}(z) = \sum_{k \in \mathbb{Z}} x_k^{\pm} z^{-k}, \ \phi^{\pm}(z) = K^{\pm 1} \exp\left(\pm (q - q^{-1}) \sum_{r>0} h_{1,\pm r} z^{\mp r}\right)$ , where  $K = q^{\alpha}$ and

$$\bar{a}_{i,j}(r) = \frac{|r|}{r} \left( (q^r + q^{-r}) \delta_{i,j} - 2(1 - \delta_{i,j}) \right).$$

The element  $\mathbf{t} = q^{\varepsilon_1 + \varepsilon_2}$  is central and split.

The subalgebra of  $U_q \widehat{\mathfrak{gl}}_2$  generated by  $x_0^{\pm}$ ,  $q^h$ ,  $h \in P$ , is isomorphic to  $U_q \mathfrak{gl}_2$ . The subalgebra of  $U_q \widehat{\mathfrak{gl}}_2$  generated by  $x_k^{\pm}$ ,  $h_r$ ,  $K^{\pm 1}$ ,  $k \in \mathbb{Z}$ ,  $r \in \mathbb{Z} \setminus \{0\}$ , and  $C^{\pm 1}$ , is the quantum affine  $\mathfrak{sl}_2$  algebra  $U_q \widehat{\mathfrak{sl}}_2$ .

Algebra  $U_q \widehat{\mathfrak{gl}}_2$  contains a Heisenberg subalgebra commuting with  $U_q \widehat{\mathfrak{sl}}_2$ , generated by the elements  $Z_r, r \in \mathbb{Z} \setminus \{0\}$ , given by

$$Z_r = \frac{2}{q^r - q^{-r}} h_{1,r} + \frac{q^r + q^{-r}}{q^r - q^{-r}} h_{2,r}, \quad [Z_r, Z_s] = -\delta_{r+s,0} [2r] \frac{1}{r} \frac{C^r - C^{-r}}{q - q^{-1}}.$$

For  $\kappa \in \mathbb{C}^{\times}$  we denote  $U_{q,\kappa}\widehat{\mathfrak{gl}}_2$  the quotient of  $U_q\widehat{\mathfrak{gl}}_2$  by the relation  $C = \kappa$ .

The algebra  $U_{q,\kappa}\widehat{\mathfrak{gl}}_2$  is  $\mathbb{Z}^2$  graded:

(2.1) 
$$\deg x_k^{\pm} = (\pm 1, k), \qquad \deg h_{i,k} = (0, k), \qquad \deg q^h = (0, 0).$$

We call the first component of the degree the weight and the second component the homogeneous degree.

We denote  $\widetilde{U}_{q,\kappa}\widehat{\mathfrak{gl}}_2$  the completion of the algebra  $U_{q,\kappa}\widehat{\mathfrak{gl}}_2$  with respect to the homogeneous degree in the negative direction.

We have several automorphisms of  $U_q \widehat{\mathfrak{gl}}_2$ .

For  $a \in \mathbb{C}^{\times}$ , the shift of spectral parameter automorphism  $\tau_a : U_q \widehat{\mathfrak{gl}}_2 \to U_q \widehat{\mathfrak{gl}}_2$  maps

(2.2) 
$$x^{\pm}(z) \mapsto x^{\pm}(z/a), \qquad \phi^{\pm}(z) \mapsto \phi^{\pm}(z/a), \qquad q^h \mapsto q^h, \qquad C \to C.$$

The automorphism  $\tau_a$  preserves the degree.

The automorphism  $\omega : U_q \widehat{\mathfrak{gl}}_2 \to U_q \widehat{\mathfrak{gl}}_2$  maps

$$x^{\pm}(z) \mapsto z^{\pm 1} x^{\pm}(z), \qquad \phi^{\pm}(z) \mapsto \phi^{\pm}(z), \qquad q^h \mapsto q^h, \qquad C \to C$$

The automorphism  $\omega$  preserves the weight but not the homogeneous degree.

## 2.3. Examples of representations. We start with representations of $U_a \mathfrak{gl}_2$ .

Let L be a  $\mathbb{C}$ -vector space with basis  $\{v_i\}_{i\in\mathbb{Z}}$ . For  $a, b, c \in \mathbb{C}$ , let

$$f_1 v_i = v_{i+1}, \qquad K v_0 = q^a v_0, \qquad \mathbf{t} \, v_0 = q^b v_0, \qquad e_1 v_0 = c v_{-1}$$

where  $i \in \mathbb{Z}$ . Then for generic a, b, c, this data is uniquely extended to a  $U_q \mathfrak{gl}_2$ -module structure on L. Then one can analytically continue this structure to all values of a, b, c. We call the resulting module L(a, b, c).

Given a, b, for generic c, the module L(a, b, c) is irreducible. If c = 0, then L has a submodule isomorphic to  $U_q\mathfrak{gl}_2$  Verma module with highest weight  $q^a$ . Following [FST], we call L(a, b, c) a relaxed Verma module.

Note that in L(a, b, c), the Casimir element acts by the constant  $\mathcal{C} = q^{a+1} + q^{-a-1} + (q-q^{-1})^2 c$ . Thus, L(a, b, c) is uniquely determined by the values of central elements  $t, \mathcal{C}$  and by the weight of  $v_0$ .

The two modules L(a, b, c) and L(a', b', c') are isomorphic if and only if values of central elements t, C are equal, and  $a - a' \in 2\mathbb{Z}$ .

Let  $U^+ \subset U_q \widehat{\mathfrak{gl}}_2$  be the subalgebra generated by  $q^h, x_k^{\pm}, h_{j,k+1}$  with  $k \ge 0, j = 0, 1, h \in P$ , and C. Let  $\kappa \in \mathbb{C}^{\times}$ . We extend the action of  $U_q \mathfrak{gl}_2$  on L(a, b, c) to  $U^+$  action by setting  $C = \kappa$ and

$$x_k^{\pm} = h_{j,k} = 0, \qquad k > 0.$$

Set

$$\hat{L}(a, b, c; \kappa) = \operatorname{Ind}_{U^+}^{U_q \mathfrak{gl}_2} L(a, b, c).$$

Given a, b, for generic  $c, \kappa$ , the  $U_q \widehat{\mathfrak{gl}}_2$ -module  $\hat{L}(a, b, c; \kappa)$  is irreducible. If c = 0, then it has a submodule isomorphic to  $U_q \widehat{\mathfrak{gl}}_2$  Verma module with highest weight  $(\kappa q^{-a}, q^a)$ . Following [FST], we call  $\hat{L}(a, b, c; \kappa)$  a relaxed  $U_q \widehat{\mathfrak{gl}}_2$  Verma module.



FIGURE 1. Grading in a relaxed  $U_q \widehat{\mathfrak{gl}}_2$  Verma module.

Set deg  $v_0 = (0,0)$ . Then the  $\mathbb{Z}^2$  grading on  $U_q \widehat{\mathfrak{gl}}_2$  induces a  $\mathbb{Z}^2$  grading on  $\hat{L}(a, b, c; \kappa)$ . We note that a vector of weight a + 2k has the first component of degree in  $\hat{L}(a, b, c; \kappa)$  equal to k. The grade diagram of  $\hat{L}(a, b, c; \kappa)$  is given on Figure 1. The dimension  $d_s$  of the space of degree (k, -s) is independent of k and is given by

$$\sum_{s=0}^{\infty} d_s t^s = \frac{1}{\prod_{i=1}^{\infty} (1-t^i)^4} = 1 + 4t + 14t^2 + 40t^3 + 105t^4 + 252t^5 + \dots$$

Let  $\hat{L}_m(a, b, c; \kappa)$  be the module  $\hat{L}(a, b, c; \kappa)$  twisted by automorphism  $\omega^m$ . We call the module  $\hat{L}_m(a, b, c; \kappa)$  the slanted relaxed Verma module of slope m. That is,  $\hat{L}_m(a, b, c; \kappa) = \hat{L}(a, b, c; \kappa)$  as a vector space and for  $v \in \hat{L}_m(a, b, c; \kappa)$ ,  $g \in U_q \widehat{\mathfrak{gl}}_2$ , we have  $g \cdot v = \omega^m(g)v$ .

The automorphism  $\omega$  changes the homogeneous degree. Thus, grade diagram of  $\hat{L}_m(a, b, c; \kappa)$  is obtained from that of  $\hat{L}(a, b, c; \kappa)$  by moving vectors of weight 2s down by by sm. The grade diagram of  $\hat{L}_m(a, b, c; \kappa)$  with m = 2 is given on Figure 2. The dimension of the space down s from the dashed line is still  $d_s$ .



FIGURE 2. Grading in a slope 2 relaxed  $U_q \widehat{\mathfrak{gl}}_2$  Verma module.

# 3. Quantum toroidal $\mathfrak{gl}_2$ algebra

3.1. The perpendicular realization  $\mathcal{E}^{\perp}$ . For  $i, j \in \mathbb{Z}/2\mathbb{Z}$  and  $r \neq 0$ , we set

$$g_{i,j}(z,w) = \begin{cases} z - q_2 w & (i \equiv j), \\ (z - q_1 w)(z - q_3 w) & (i \neq j). \end{cases}$$

The quantum toroidal algebra of type  $\mathfrak{gl}_2$ , which we denote by  $\mathcal{E}^{\perp}$ , is a unital associative algebra generated by  $E_{i,k}^{\perp}, F_{i,k}^{\perp}, H_{i,r}^{\perp}$  and elements  $q^h$ ,  $C^{\pm 1}$ , where  $i \in \mathbb{Z}/2\mathbb{Z}, k \in \mathbb{Z}, r \in \mathbb{Z} \setminus \{0\}$ ,  $h \in P$ . We set  $K_1^{\perp} = K = q^{\alpha}, K_0^{\perp} = K^{-1} = q^{-\alpha}$ .

We give the defining relations in terms of generating series

$$E_i^{\perp}(z) = \sum_{k \in \mathbb{Z}} E_{i,k}^{\perp} z^{-k}, \quad F_i^{\perp}(z) = \sum_{k \in \mathbb{Z}} F_{i,k}^{\perp} z^{-k},$$
  
$$K_i^{\pm,\perp}(z) = (K_i^{\perp})^{\pm 1} \bar{K}_i^{\pm,\perp}(z), \quad \bar{K}_i^{\pm,\perp}(z) = \exp\left(\pm (q - q^{-1}) \sum_{r>0} H_{i,\pm r}^{\perp} z^{\mp r}\right).$$

The relations are as follows.

$$\begin{split} C \text{ is central,} \quad & CC^{-1} = 1, \quad q^h q^{h'} = q^{h+h'}, \quad q^0 = 1, \\ q^h E_i^{\perp}(z) q^{-h} = q^{(h,\alpha_i)} E_i^{\perp}(z), \quad q^h F_i^{\perp}(z) q^{-h} = q^{-(h,\alpha_i)} F_i^{\perp}(z), \quad q^h K_i^{\pm,\perp}(z) = K_i^{\pm,\perp}(z) q^h, \\ & K_i^{\pm,\perp}(z) K_j^{\pm,\perp}(w) = K_j^{\pm,\perp}(w) K_i^{\pm,\perp}(z), \\ & \frac{g_{i,j}(C^{-1}z,w)}{g_{i,j}(Cz,w)} K_i^{-,\perp}(z) K_j^{\pm,\perp}(w) = \frac{g_{j,i}(w, C^{-1}z)}{g_{j,i}(w, Cz)} K_j^{\pm,\perp}(w) K_i^{\pm,\perp}(z), \\ & (-1)^{i+j} g_{i,j}(z,w) K_i^{\pm,\perp}(C^{-(1\pm1)/2}z) E_j^{\perp}(w) + g_{j,i}(w, z) E_j^{\perp}(w) K_i^{\pm,\perp}(C^{-(1\pm1)/2}z) = 0, \\ & (-1)^{i+j} g_{j,i}(w,z) K_i^{\pm,\perp}(C^{-(1\pm1)/2}z) F_j^{\perp}(w) + g_{i,j}(z,w) F_j^{\perp}(w) K_i^{\pm,\perp}(C^{-(1\pm1)/2}z) = 0, \\ & [E_i^{\perp}(z), F_j^{\perp}(w)] = \frac{\delta_{i,j}}{q - q^{-1}} (\delta(C\frac{w}{z}) K_i^{\pm,\perp}(w) - \delta(C\frac{z}{w}) K_i^{-,\perp}(z)), \\ & (-1)^{i+j} g_{i,j}(x,w) E_i^{\perp}(z) E_j^{\perp}(w) + g_{i,j}(z,w) F_j^{\perp}(w) E_i^{\perp}(z) = 0, \\ & (-1)^{i+j} g_{j,i}(w,z) F_i^{\perp}(z) F_j^{\perp}(w) + g_{i,j}(z,w) F_j^{\perp}(w) F_i^{\perp}(z) = 0, \\ & (-1)^{i+j} g_{j,i}(w,z) F_i^{\perp}(z) F_j^{\perp}(w) + g_{i,j}(z,w) F_j^{\perp}(w) F_i^{\perp}(z) = 0, \\ & Sym_{z_1, z_2, z_3} [E_i^{\perp}(z_1), [E_i^{\perp}(z_2), [E_i^{\perp}(z_3), E_{i+1}^{\perp}(w)]_q 2]]_{q^{-2}} = 0, \\ & Sym_{z_1, z_2, z_3} [F_i^{\perp}(z_1), [F_i^{\perp}(z_2), [F_i^{\perp}(z_3), F_{i+1}^{\perp}(w)]_q 2]]_{q^{-2}} = 0. \end{split}$$

Here we use  $\delta(z) = \sum_{k \in \mathbb{Z}} z^k$  and

Sym 
$$f(x_1, ..., x_N) = \frac{1}{N!} \sum_{\pi \in \mathfrak{S}_N} f(x_{\pi(1)}, ..., x_{\pi(N)})$$

The algebra  $\mathcal{E}^{\perp}$  considered here is obtained from  $\mathcal{E}_2$  in [FJMM2] by setting the second central element  $K_0^{\perp}K_1^{\perp}$  to 1, dropping the scaling elements D,  $D^{\perp}$  and adding the split central element  $\mathbf{t} = q^{\varepsilon_1 + \varepsilon_2}$ .

We note that the algebra  $\mathcal{E}^{\perp}$  is invariant under the exchange of  $q_1$  and  $q_3$ .

Algebra  $\mathcal{E}^{\perp}$  is  $(\mathbb{Z}^2 \times \mathbb{Z})$ -graded by

$$\deg E_{i,k}^{\perp} = (1_i, k), \quad \deg F_{i,k}^{\perp} = (-1_i, k), \qquad \deg H_{i,r}^{\perp} = (0, 0, r), \qquad \deg C = \deg q^h = 0,$$

where  $1_0 = (1, 0)$  and  $1_1 = (0, 1)$ .

For i = 0, 1, there exist automorphisms  $\omega_i : \mathcal{E}^{\perp} \to \mathcal{E}^{\perp}$  sending

$$E_j^{\perp}(z) \mapsto z^{\delta_{ij}} E_j^{\perp}(z), \qquad F_j^{\perp}(z) \mapsto z^{-\delta_{ij}} F_j^{\perp}(z), \qquad K_i^{\pm,\perp}(z) \mapsto K_i^{\pm,\perp}(z),$$

and  $q^h \mapsto q^h$ ,  $C \mapsto C$ . Note that  $\omega_0 \omega_1 = \omega_1 \omega_0$ .

We have an embedding  $v^{\perp}$ :  $U_q \widehat{\mathfrak{sl}}_2 \to \mathcal{E}^{\perp}$  such that

$$x^+(z) \mapsto E_1^{\perp}(d^{-1}z), \quad x^-(z) \mapsto F_1^{\perp}(d^{-1}z), \quad \phi^{\pm}(z) \mapsto K_1^{\pm,\perp}(d^{-1}z),$$

and  $C \mapsto C$ . We call the image of  $v^{\perp}$  the vertical  $U_q \widehat{\mathfrak{sl}}_2$ -subalgebra and denote it by  $U_q^{v,\perp} \widehat{\mathfrak{sl}}_2 \subset \mathcal{E}^{\perp}$ .

The subalgebra of  $\mathcal{E}^{\perp}$  generated by  $E_1^{\perp}(z)$ ,  $F_1^{\perp}(z)$ ,  $K_0^{\pm,\perp}(z)$ ,  $K_1^{\pm,\perp}(z)$ ,  $q^h$  and  $C^{\pm 1}$  is isomorphic to  $U_q \widehat{\mathfrak{gl}}_2$ . We call this subalgebra the vertical  $U_q \widehat{\mathfrak{gl}}_2$ -subalgebra and denote it by  $U_q^{v,\perp} \widehat{\mathfrak{gl}}_2 \subset \mathcal{E}^{\perp}$ .

**Proposition 3.1** ([M2],[FJM1]). Let  $u \in \mathbb{C}^{\times}$ . Let  $C = q_3$ . There exists a surjective homomorphism of algebras:  $ev_u : \mathcal{E}^{\perp} \to \widetilde{U}_{q,q_3}\widehat{\mathfrak{gl}}_2$ , such that  $ev_u \circ v^{\perp} = id$  and  $\deg(ev_u(E_{0,k}^{\perp})) = (-1,k)$ ,  $\deg(ev_u(F_{0,k}^{\perp})) = (1,k)$ .

Note that if L is an  $\widetilde{U}_{q,q_3}\widehat{\mathfrak{gl}}_2$ -module and  $L_m$  is the module obtained by twisting L by the automorphism  $\omega^m$ , then the evaluation  $\mathcal{E}^{\perp}$ -module of  $L_m$  is obtained from the evaluation  $\mathcal{E}^{\perp}$ -module of L by twisting by the automorphism  $\omega_1^m \omega_0^{-m}$ .

We call the subalgebra of  $\mathcal{E}^{\perp}$  generated by  $E_{i,0}^{\perp}, F_{i,0}^{\perp}, i = 1, 2$ , the horizontal  $\widehat{\mathfrak{sl}}_2$  subalgebra and denote it by  $U_q^{h,\perp}\widehat{\mathfrak{sl}}_2$ . The horizontal  $\widehat{\mathfrak{sl}}_2$  subalgebra is isomorphic to the quotient of  $U_q\widehat{\mathfrak{sl}}_2$ by the relation C = 1.

3.2. Parallel realization of  $\mathcal{E}^{\perp}$ . Algebra  $\mathcal{E}^{\perp}$  has an alternative presentation. Let  $\mathcal{E}$  be the unital associative algebra generated by coefficients of the generating series

$$E_i(z) = \sum_{k \in \mathbb{Z}} E_{i,k} z^{-k}, \quad F_i(z) = \sum_{k \in \mathbb{Z}} F_{i,k} z^{-k}, \quad K_i^{\pm}(z) = K_i^{\pm 1} \exp\left(\pm (q - q^{-1}) \sum_{r>0} H_{i,\pm r} z^{\mp r}\right),$$

where  $i \in \mathbb{Z}/2\mathbb{Z}$ , and a split central element t, which satisfy the following relations.

$$\begin{aligned} K_i^{\pm}(z)K_j^{\pm}(w) &= K_j^{\pm}(w)K_i^{\pm}(z), \qquad K_i^{\pm}(z)K_j^{\mp}(w) = K_j^{\mp}(w)K_i^{\pm}(z), \\ (-1)^{i+j}g_{i,j}(z,w)K_i^{\pm}(z)E_j(w) + g_{j,i}(w,z)E_j(w)K_i^{\pm}(z) = 0, \\ (-1)^{i+j}g_{j,i}(w,z)K_i^{\pm}(z)F_j(w) + g_{i,j}(z,w)F_j(w)K_i^{\pm}(z) = 0, \end{aligned}$$

COMBINATORIAL MODULES

$$\begin{split} [E_i(z),F_j(w)] &= \frac{\delta_{i,j}}{q-q^{-1}}\delta(z/w)(K_i^+(w)-K_i^-(z)),\\ (-1)^{i+j}g_{i,j}(z,w)E_i(z)E_j(w)+g_{j,i}(w,z)E_j(w)E_i(z)=0,\\ (-1)^{i+j}g_{j,i}(w,z)F_i(z)F_j(w)+g_{i,j}(z,w)F_j(w)F_i(z)=0,\\ \mathrm{Sym}_{z_1,z_2,z_3}[E_i(z_1),[E_i(z_2),[E_i(z_3),E_{i+1}(w)]_{q^2}]]_{q^{-2}}=0,\\ \mathrm{Sym}_{z_1,z_2,z_3}[F_i(z_1),[F_i(z_2),[F_i(z_3),F_{i+1}(w)]_{q^2}]]_{q^{-2}}=0. \end{split}$$

There exists a Drinfeld coproduct  $\Delta$  in  $\mathcal{E}$  given by

(3.1)  

$$\Delta(K_i^{\pm}(z)) = K_i^{\pm}(z) \otimes K_i^{\pm}(z),$$

$$\Delta(E_i(z)) = E_i(z) \otimes 1 + K_i^{-}(z) \otimes E_i(z),$$

$$\Delta(F_i(z)) = F_i(z) \otimes K_i^{+}(z) + 1 \otimes F_i^{\pm}(z)$$

and  $\Delta(t) = t \otimes t$ .

There exists a map  $h: U_q \widehat{\mathfrak{sl}}_2 \to \mathcal{E}$  such that

$$x^+(z) \mapsto E_1(d^{-1}z), \quad x^-(z) \mapsto F_1(d^{-1}z), \quad \phi^{\pm}(z) \mapsto K_1^{\pm}(d^{-1}z), \qquad C \mapsto 1.$$

We denote the image of h by  $U_q^h \widehat{\mathfrak{sl}}_2 \subset \mathcal{E}$ . The algebra  $U_q^h \widehat{\mathfrak{sl}}_2$  is isomorphic to the quantum affine algebra  $U_{q,1} \widehat{\mathfrak{sl}}_2$  with C = 1.

We denote the subalgebra of  $\mathcal{E}$  generated by  $E_{i,0}, F_{i,0}, i = 1, 2$ , by  $U_q^v \widehat{\mathfrak{sl}}_2$ . Define a  $\mathbb{Z}^3$  grading in  $\mathcal{E}$  by

$$deg E_{1,k} = (-k, -k + 1, 0), deg E_{0,k} = (-k, -k - 1, 1), deg F_{1,k} = (-k, -k - 1, 0), deg F_{0,k} = (-k, -k + 1, -1), deg H_{1,r} = (-r, -r, 0), deg H_{0,r} = (-r, -r, 0),$$

and deg t = (0, 0, 0).

It is easy to check it is indeed a grading. The definition of this grading is chosen to make the isomorphism in Proposition 3.2 below graded, cf. Lemma 2.4 in [FJMM2].

The algebras  $\mathcal{E}^{\perp}$  and  $\mathcal{E}$  are, in fact, the same.

**Proposition 3.2** ([M1]). There exists a  $\mathbb{Z}^3$ -graded isomorphism of algebras  $\theta$ :  $\mathcal{E}^{\perp} \to \mathcal{E}$  such that  $\theta(C) = K_0 K_1$ ,  $\theta(U_q^{v,\perp} \widehat{\mathfrak{sl}_2}) = U_q^v \widehat{\mathfrak{sl}_2}$ ,  $\theta(U_q^{h,\perp} \widehat{\mathfrak{sl}_2}) = U_q^h \widehat{\mathfrak{sl}_2}$ , and  $\theta(q^h) = q^h$ ,  $h \in P$ .

The commutative currents  $K_i(z)$  are called Cartan currents.

For  $a \in \mathbb{C}^{\times}$ , the shift of spectral parameter automorphism  $\hat{\tau}_a : \mathcal{E} \to \mathcal{E}$  maps

(3.3) 
$$E_i(z) \mapsto E_i(z/a), \qquad F_i(z) \mapsto F_i(z/a), \qquad K_i^{\pm}(z) \mapsto K_i^{\pm}(z/a),$$

and  $\mathbf{t} \to \mathbf{t}$ . The automorphism  $\hat{\tau}_a$  preserves the degree. Note that the automorphism  $\hat{\tau}_a$  preserves subalgebras  $U_q^h \widehat{\mathfrak{sl}_2}$  and  $U_q^v \widehat{\mathfrak{sl}_2}$ . The restriction of  $\hat{\tau}_a$  to  $U_q^h \widehat{\mathfrak{sl}_2}$  is the automorphism  $\tau_a$ , see (2.2), and the restriction of  $\hat{\tau}_a$  to  $U_q^h \widehat{\mathfrak{sl}_2}$  is the identity map.

Let V be an  $\mathcal{E}$ -module.

For  $\kappa \in \mathbb{C}^{\times}$ , we say V has level  $\kappa$  if the central element  $K_0K_1$  acts in V as the scalar  $\kappa$ .

A vector  $v \in V$  is called a vector of  $\ell$ -weight  $\phi(z) = (\phi^0(z), \phi^1(z))$  if  $K_i^{\pm}(z) v = \phi^i(z) v$ , i = 0, 1.

A module V is called a highest  $\ell$ -weight module of highest  $\ell$ -weight  $\phi(z)$  if it is generated by a vector v of  $\ell$ -weight  $\phi(z)$  satisfying  $E_i(z) v = 0, i = 0, 1$ .

**Proposition 3.3.** [FJM1] Let  $L_{a,b}$  be the irreducible highest weight  $U_q \widehat{\mathfrak{gl}}_2$ -module of highest weight a, b, that is generated by a non-zero vector v such that  $K_0 v = av$ ,  $K_1 v = bv$ ,  $E_{i,0} v = 0$ , i = 0, 1. Set  $q_3 = ab$ . Then the corresponding evaluation  $\mathcal{E}$ -module has highest  $\ell$ -weight

(3.4) 
$$\phi(z) = \left(\frac{az-u}{z-au}, \ \frac{bz-u}{z-bu}\right)$$

Here the parameter u depends on the choice of the evaluation map and can be made arbitrary by twisting by automorphism (3.3).

In all modules considered in this paper, the series  $\phi_i(z)$  will be rational functions regular at zero and infinity, satisfying  $\phi_i(0)\phi_i(\infty) = 1$ , cf. Theorem 2.3 in [FJMM1]. We call such rational functions balanced. The set of all pairs of rational functions  $\phi(z)$  with balanced coordinates is naturally a group under coordinate-wise multiplication.

A module V is called tame if V has a basis of  $\ell$ -weight vectors with distinct  $\ell$ -weights.

3.3. An explicit construction of tame  $\mathcal{E}$ -modules. If a module over quantum affine algebra is tame then the matrix coefficients of  $E_i(z)$  and  $F_i(z)$  are delta functions multiplied by some constants. We give an abstract construction which under some assumption defines the action in a completely combinatorial way.

A box  $\square$  is a set which has two elements: color of the box  $c(\square) \in \{0,1\}$  and coordinate of the box  $(x(\square), y(\square), z(\square)) \in \mathbb{C}^3$ . The position of the box is then given by  $p(\square) = q_1^{-x(\square)}q_2^{-y(\square)}q_3^{-z(\square)}$ . We assume that the set of all boxes is totally ordered.

We do not allow boxes with the same positions and different colors.

If all three coordinates of two boxes differ by integers then we assume that the ordering is lexicographic according to y > z > x. Namely, two such boxes are ordered as  $\Box_1 > \Box_2$  if and only if  $y(\Box_1) > y(\Box_2)$  or if  $y(\Box_1) = y(\Box_2)$  and  $z(\Box_1) > z(\Box_2)$ , or if  $y(\Box_1) = y(\Box_2)$  and  $z(\Box_1) = z(\Box_2)$  and  $x(\Box_1) > x(\Box_2)$ . One can take any ordering satisfying this assumption.

If the x and z coordinates of two boxes differ by integers then we assume  $c(\Box_1) - c(\Box_2) \equiv x(\Box_1) - x(\Box_2) + z(\Box_1) - z(\Box_2)$  modulo 2. In other words, colors alternate in x- and z-directions and do not change in y-direction.

The set of all boxes is divided into two disjoint subsets of positive and negative boxes.

A state  $\lambda = \lambda^+ \sqcup \lambda^-$  is a disjoint union of two finite sets of boxes without repetitions such that all boxes in  $\lambda^+$  are positive, and all boxes in  $\lambda^-$  are negative.

The state  $\emptyset$  is called the reference state. We think of a state  $\lambda$  as having positive boxes in  $\lambda^+$  added to the reference state and negative boxes in  $\lambda^-$  removed from the reference state. So, one can think that the reference state consists of all negative boxes.

Let S be a collection of states including the reference state, and let  $\Psi(z) = (\Psi^0(z), \Psi^1(z))$ be a pair of balanced rational functions. This is our data for constructing an  $\mathcal{E}$ -module under the assumption to be stated below.

For  $\lambda \in S$ , a box  $\Box$  is called convex if either  $\Box \in \lambda^+$  and the removal of this box produces a state,  $((\lambda^+ \setminus \Box) \sqcup \lambda^-) \in S$ , or  $\Box \notin \lambda^-$  is a negative box and the addition of this box to  $\lambda^$ produces a state,  $(\lambda^+ \sqcup (\lambda^- \sqcup \Box)) \in S$ . In both cases we say that the new state is obtained from  $\lambda$  by removing the box  $\Box$  and we denote it by  $\lambda \setminus \Box$ . We denote the set of all convex boxes of color *i* for a state  $\lambda$  by  $CV_i(\lambda)$ .

For  $\lambda \in S$ , a box  $\Box$  is called concave if either  $\Box \notin \lambda^+$  is a positive box and the addition of this box to  $\lambda^+$  produces a state,  $((\lambda^+ \sqcup \Box) \sqcup \lambda^-) \in S$ , or  $\Box \in \lambda^-$  and the removal of this box produces a state,  $(\lambda^+ \sqcup (\lambda^- \setminus \Box)) \in S$ . In both cases we say that the new state is obtained from  $\lambda$  by adding the box  $\Box$  and we denote it by  $\lambda \sqcup \Box$ . We denote the set of all concave boxes of color *i* for a state  $\lambda$  by  $CC_i(\lambda)$ .

A set S of states is connected if any non-empty state  $\lambda$  can be connected to the reference state by a sequence of removal of boxes.

Next we define the  $\ell$ -weight of a state. For  $a \in \mathbb{C}^{\times}$ , introduce the notation for pairs of balanced rational functions

$$0_a = (\frac{qz-a}{z-aq}, 1), \qquad 1_a = (1, \frac{qz-a}{z-aq}).$$

These are quantum affine analogs of  $\widehat{\mathfrak{sl}}_2$  fundamental weights. The quantum affine analogs of  $\widehat{\mathfrak{sl}}_2$  roots are given by

$$A_{0,a}(z) = \left(-\frac{g_{0,0}(a,z)}{g_{0,0}(z,a)}, \frac{g_{0,1}(a,z)}{g_{1,0}(z,a)}\right) = \left(\frac{q_2z-a}{z-q_2a}, \frac{(q_1z-a)(q_3z-a)}{(z-q_1a)(z-q_3a)}\right) = 0_{q^{-1}a}0_{qa}1_{q_1qa}^{-1}1_{q_3qa}$$

$$A_{1,a}(z) = \left(\frac{g_{1,0}(a,z)}{g_{0,1}(z,a)}, -\frac{g_{1,1}(a,z)}{g_{1,1}(z,a)}\right) = \left(\frac{(q_1z-a)(q_3z-a)}{(z-q_1a)(z-q_3a)}, \frac{q_2z-a}{z-q_2a}\right) = 1_{q^{-1}a}1_{qa}0_{q_1qa}^{-1}0_{q_3qa}^{-1}.$$

Note the complete symmetries between  $0 \leftrightarrow 1$  and between  $q_1 \leftrightarrow q_3$ . Note the identities

$$A_{i,a}(b) = A_{i,1}(b/a),$$
  $A_{i,a}(b) = A_{i,b}^{-1}(a),$   $A_{0,a}(a) = (-1,1),$   $A_{1,a}(a) = (1,-1).$ 

We define the reference state to have  $\ell$ -weight  $\Psi(z)$ , that is  $\phi_{\emptyset}(z) = \Psi(z)$ , and a box  $\Box$  to have  $\ell$ -weight  $A_{c(\Box),p(\Box)}^{-1}(z)$ .

We define the  $\ell$ -weight of a state  $\lambda \in S$  by

$$\phi_{\lambda}(z) = (\phi_{\lambda}^{0}(z), \phi_{\lambda}^{1}(z)) = \Psi(z) \prod_{\square \in \lambda^{+}} A_{c(\square), p(\square)}^{-1}(z) \prod_{\square \in \lambda^{-}} A_{c(\square), p(\square)}(z).$$

Given a box  $\Box$  we break the  $\ell$ -weight of a state into a product collecting boxes before and after  $\Box$ :

$$\phi_{\lambda,<\Box}(z) = \Psi(z) \prod_{\Box_1 \in \lambda^+, \Box_1 < \Box} A_{c(\Box_1),p(\Box_1)}^{-1}(z) \prod_{\Box_1 \in \lambda^-, \Box_1 < \Box} A_{c(\Box_1),p(\Box_1)}(z),$$
  
$$\phi_{\lambda,>\Box}(z) = \prod_{\Box_1 \in \lambda^+, \Box_1 > \Box} A_{c(\Box_1),p(\Box_1)}^{-1}(z) \prod_{\Box_1 \in \lambda^-, \Box_1 > \Box} A_{c(\Box_1),p(\Box_1)}(z).$$

We have

$$\phi_{\lambda}(z) = \begin{cases} \phi_{\lambda,<\square}(z) \phi_{\lambda,>\square}(z), & \text{if } \square \notin \lambda, \\ \phi_{\lambda,<\square}(z) \phi_{\lambda,>\square}(z) A_{c(\square),p(\square)}^{-1}(z), & \text{if } \square \in \lambda^+, \\ \phi_{\lambda,<\square}(z) \phi_{\lambda,>\square}(z) A_{c(\square),p(\square)}(z), & \text{if } \square \in \lambda^-. \end{cases}$$

We make the following assumptions:

- (A1) The  $\ell$ -weights of all states are distinct.
- (A2) For each  $\lambda \in S$  the the components  $\phi_{\lambda}^{i}(z)$  of  $\ell$ -weight  $\phi(\lambda)$  are balanced rational functions with only simple poles.
- (A3) The poles of  $\phi_{\lambda}^{i}(z)$  are in a bijection with the set  $CC_{i}(\lambda) \sqcup CV_{i}(\lambda)$  of concave and convex boxes of color *i*.
- (A4) A pole z = a of  $\phi_{\lambda}^{i}(z)$  corresponds to a convex or concave box  $\Box$  such that  $c(\Box) = i$ and  $p(\Box) = a$  and such that  $\phi_{\lambda, \supset \Box}(z)$  is regular at  $z = a = p(\Box)$  if  $\Box$  is positive and  $\phi_{\lambda, <\Box}(z)$  is regular at  $z = a = p(\Box)$  if  $\Box$  is negative.

In particular, the assumptions (A2), (A3) imply that for each state  $\lambda \in S$ , the sets of convex and concave boxes  $CV_i(\lambda)$ ,  $CC_i(\lambda)$  are finite. By (A2) and (A4), the positions of all these boxes are distinct.

Using assumptions (A3), (A4), one can also deduce our previous assumption on coloring but we do not pursue this point.

Now, we define the  $\mathcal{E}$ -module  $V(S, \Psi)$ .

Let  $V = \bigoplus_{\lambda \in S} \mathbb{C} |\lambda\rangle$  be the  $\mathbb{C}$ -vector space with a fixed basis labeled by states in S. Set

(3.5)  

$$K_{i}^{\pm}(z)|\lambda\rangle = \phi_{\lambda}^{i}(z)|\lambda\rangle,$$

$$F_{i}(z)|\lambda\rangle = \sum_{\Box \in CC_{i}(\lambda)} a_{\lambda,\Box}^{i} \,\delta(z/p(\Box)) \,|\lambda \sqcup \Box\rangle,$$

$$E_{i}(z)|\lambda\rangle = \sum_{\Box \in CV_{i}(\lambda)} b_{\lambda,\Box}^{i} \,\delta(z/p(\Box)) \,|\lambda \setminus \Box\rangle,$$

where the coefficients are given by

(3.6) 
$$a_{\lambda,\Box}^{i} = \begin{cases} \phi_{\lambda,\supset\Box}^{i}(p(\Box)), & \text{if } \Box \text{ is positive,} \\ \frac{-1}{q-q^{-1}} \operatorname{res}_{z=p(\Box)} \phi_{\lambda,\supset\Box}^{i}(z) \frac{dz}{z}, & \text{if } \Box \text{ is negative,} \end{cases}$$

(3.7) 
$$b_{\lambda,\Box}^{i} = \begin{cases} \frac{1}{q-q^{-1}} \operatorname{res}_{z=p(\Box)} \phi_{\lambda,<\Box}^{i}(z) \frac{dz}{z}, & \text{if } \Box \text{ is positive,} \\ \phi_{\lambda,<\Box}^{i}(p(\Box)), & \text{if } \Box \text{ is negative} \end{cases}$$

The element t can be set to an arbitrary non-zero number. For brevity, we set t = 1.

We remark that the statement that in tame modules the matrix coefficients of  $F_i(z), E_i(z)$ must be delta functions at poles of  $\ell$ -weights is an easy general fact. However, in general it is not clear how to fix the coefficients, see Section 5.3 in [FJM2]. Here we regard our module as some kind of tensor product, therefore the choice of coefficients is dictated by the coproduct (3.1). Now we check that it works.

**Theorem 3.4.** Let S be a connected set of states and  $\Psi(z)$  a pair of balanced functions such that the assumptions (A1)–(A4) hold. Then formulas (3.5) make  $V = \bigoplus_{\lambda \in S} \mathbb{C} |\lambda\rangle$  into a tame irreducible  $\mathcal{E}$ -module, provided the Serre relations hold on the reference vector  $|\emptyset\rangle$ .

*Proof.* All the relations except for Serre relation are checked directly.

For the *EF* relation notice that for a rational function  $\phi(z) = \prod_{s=1}^{k} (z - b_s)/(z - a_s)$ , the difference of expansions at  $x = \infty$  and at x = 0 gives  $\sum_{s=1}^{k} \delta(z/a_s) \operatorname{res}_{z=a_s} \phi(z) dz/z$ . At the same time we have  $(q - q^{-1})a_{\lambda,\Box}^i b_{\lambda,\Box}^i = \pm \operatorname{res}_{z=p(\Box)} \phi_{\lambda}^i(z) dz/z$  with the appropriate sign.

Then we have a module of the analog of  $\mathcal{E}$  without Serre relations. This module is irreducible, since S is connected and since all states are separated by the  $\ell$ -weights (eigenvalues of Cartan currents).

Then it is well known that the left hand side of the Serre relations are in the center of such algebra. Therefore it is sufficient to check the Serre relation on one vector.  $\Box$ 

We denote by  $V(S, \Psi)$  the  $\mathcal{E}$ -module V constructed by the use of Theorem 3.4 and call it a combinatorial module.

Define the degree of a box of color 0 to be (1,0) and the degree of a box of color 1 to be (0,1). The degree of a collection of boxes is the sum of degrees of the boxes. For a state  $\lambda = \lambda^+ \sqcup \lambda^- \in S$ , define the two component degree of  $\lambda$  by

$$\deg(\lambda) = \deg(\lambda^+) - \deg(\lambda^-) = (\deg_0(\lambda), \deg_1(\lambda)).$$

Note that the degree is normalized by the condition that the degree of the reference state is (0,0).

Define the character of a combinatorial module  $V(S, \Psi)$  as a generating series of degrees of the states:

$$\chi_{V(S,\Psi)}(z_0, z_1) = \sum_{\lambda \in S} z_0^{\deg_0(\lambda)} z_1^{\deg_1(\lambda)}$$

The results of this section, including Theorem 3.4, also hold for quantum toroidal algebras of type  $\mathfrak{gl}_n$  with arbitrary *n* with obvious changes in the coloring.

### 4. Examples of combinatorial *E*-modules.

It seems that Theorem 3.4 can be applied to many combinatorial sets S. We have no classification of all possibilities. Here we give a number of examples starting with known ones previously constructed by different methods and then proceed to new ones. In each case we describe the set S and the initial weight  $\Psi(z)$  and then check the assumptions of Theorem 3.4.

4.1. Vector representations. We define a combinatorial module  $U_0$  associated with the data  $S(U_0), \Psi(U_0)$  given as follows.

Let  $\Psi(U_0) = 0_{q^{-1}} 1_{qq_1}^{-1}$ . Let the set of boxes consist of boxes with coordinates (i, 0, 0), and color  $i \mod 2, i \in \mathbb{Z}$ . The box with coordinates (i, 0, 0) is negative if and only if i is negative.

The set of states is  $S(U_0) = \mathbb{Z}$ . For  $k \ge 0$ , the state k consists of k positive boxes with coordinates  $(0,0,0), (1,0,0), \ldots, (k-1,0,0)$ . For k < 0, the state k consists of |k| negative boxes with coordinates  $(-1,0,0), (-2,0,0), \ldots, (k,0,0)$ . The state 0 is the reference state.

**Proposition 4.1.** The space  $U_0 = V(S(U_0), \Psi(U_0))$  with action (3.5) is an irreducible tame  $\mathcal{E}$ -module of level 1 called the vector module of color 0.

*Proof.* Indeed, by induction on k, clearly the  $\ell$ -weight of the state k is

(4.1) 
$$\phi_k = \begin{cases} 0_{q^{-1}q_1^{-k}} 1_{qq_1^{-k+1}}^{-1}, & \text{if } k \text{ is even,} \\ 0_{qq_1^{-k+1}}^{-1} 1_{q^{-1}q_1^{-k}}, & \text{if } k \text{ is odd.} \end{cases}$$

At the same time there is a convex box (k-1,0,0) and a concave box (k,0,0). We also note that for k > 0,  $\phi_{k,>(k-1,0,0)}(z) = \phi_{-k,<(-k+1,0,0)}(z) = 1$ . Thus, the assumptions (A1)–(A4) clearly hold.

The Serre relations trivially hold on the reference state.

Thus the proposition follows from Theorem 3.4.

One can think about the reference state as the union of all negative boxes with coordinates  $(j, 0, 0), j \in \mathbb{Z}_{\leq 0}$ . Then for all  $k \in \mathbb{Z}$  the state k is the set consisting of all boxes  $(j, 0, 0), j \in \mathbb{Z}_{\leq k}$ . Thus adding a negative box to a state is actually removing a box from this infinite chain of boxes.

The vector representation  $U_1$  of color 1 is obtained by interchanging colors. The module  $U_1$  is isomorphic to  $U_0$  shifted by  $q_1$ . The representation  $U_0$  shifted by  $q_1^2$  is isomorphic to  $U_0$ .

Another vector representation could be obtained by exchanging  $q_1$  and  $q_3$ .

More vector representations can be obtained by "shifts" – by twisting with the shift of spectral parameter automorphism  $\tau_a$ .

Clearly, the characters of the vector representations are given by

$$\chi_{U_i}(z_0, z_1) = (1 + z_i) \sum_{k \in \mathbb{Z}} (z_0 z_1)^k = (1 + z_i) \, \delta(z_0 z_1).$$

The module  $U_i$  is an irreducible module of vertical  $U_q \widehat{\mathfrak{gl}}_2$ . It is a quantization of the natural  $\mathfrak{gl}_2[t, t^{-1}]$ -module  $\mathbb{C}^2[t, t^{-1}]$ .

# 4.2. Fock modules. Next we define the Fock modules $F_i$ as combinatorial $\mathcal{E}$ modules.

Let  $\Psi(F_0) = 0_{q^{-1}}$ .

Let the set of boxes consist of boxes with coordinates (i, 0, j), and color  $i + j \mod 2$ , where  $i, j \in \mathbb{Z}_{\geq 0}$ . All the boxes are positive, we have no negative boxes. The ordering of boxes is lexicographic.

A partition  $\lambda$  is a non-increasing sequence of non-negative numbers  $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_s \geq 0$ called parts. We do not distinguish between partitions which differ by zero parts. We identify a partition with a set of boxes using the Young diagram. A partition  $\lambda$  corresponds to the set of boxes with coordinates  $(i, 0, j), j = 0, \ldots, s - 1, i = 0, \ldots, \lambda_{j-1} - 1$ .

Let the set of states  $S(F_0)$  consist of all partitions. Let the empty partition be the reference state.

14

An example of a state is pictured on Figure 3. (Recall that  $q_1, q_2, q_3$  correspond to coordinates x, y, z respectively and the position of the box is  $p(\Box) = q_1^{-x(\Box)}q_2^{-y(\Box)}q_3^{-z(\Box)}$ .)



FIGURE 3. A state in the Fock module.

This state corresponds to the partition  $\lambda = (4, 2, 1)$ . We color the boxes by white and black colors, corresponding to colors 0 and 1. Thus colors alternate in both x- and z-directions.

There are 4 concave boxes: (4, 0, 0) (white), (2, 0, 1), (1, 0, 2), (0, 0, 3) (black) and 3 convex boxes: (3, 0, 0) (black), (1, 0, 1), (0, 0, 2) (white). Accordingly, the  $\ell$ -weight of this state is

$$\phi_{(4,2,1)} = 0_{q^{-1}q_1^{-4}} 0_{qq_1^{-1}q_3^{-1}}^{-1} 0_{qq_3^{-2}}^{-1} 1_{qq_1^{-3}}^{-1} 1_{q^{-1}q_1^{-2}q_3^{-1}} 1_{q^{-1}q_1^{-1}q_3^{-2}} 1_{q^{-1}q_3^{-3}}$$

Informally, one can think that this state is a tensor product of three vectors corresponding to rows: state 4 from  $U_0$ , state 2 from  $U_1$  shifted by  $q_3^{-1}$ , and state 1 from  $U_0$  shifted by  $q_3^{-2}$ .

**Proposition 4.2.** The space  $F_0 = V(S(F_0), \Psi(F_0))$  with action (3.5) is an irreducible tame highest  $\ell$ -weight module of highest  $\ell$ -weight  $\Psi(F_0)$ . In particular it has level q.

Proof. The poles of the  $\ell$ -weight of a partition correspond to the convex and concave boxes. Indeed, the  $\ell$ -weight of row *i* is a shift of  $\phi_{\lambda_i} \Psi(U_0)^{-1}$  by  $q_3^{-2i}$  if *i* is even and similarly with the change of colors if *i* is odd, see (4.1). Then  $\Psi(F_0)$  multiplied by shifts of  $\Psi(U_0)^{-1}$  and  $\Psi(U_1)^{-1}$  produces only a pole in the convex box in the first column of  $\lambda$ .

Clearly, a partition cannot have two convex boxes with the same position. A partition cannot have two concave boxes with the same position either. Therefore all  $\ell$ -weights are distinct. Thus the assumptions (A1)–(A4) hold.

The Serre relations trivially hold on the reference state as all terms act by zero.

Thus the proposition follows from Theorem 3.4.

The module  $F_0$  described in Proposition 4.2 is called the Fock module of color 0. The Fock module  $F_0$  is the unique irreducible highest  $\ell$ -weight  $\mathcal{E}$ -module with highest  $\ell$ -weight  $\Psi(F_0) = 0_{q^{-1}} = (\frac{qz-q^{-1}}{z-1}, 1).$ 

The Fock module  $F_1$  of color 1 is obtained from  $F_0$  by changing colors. The Fock module  $F_0$  is the unique irreducible highest  $\ell$ -weight  $\mathcal{E}$ -module with highest  $\ell$ -weight  $\Psi(F_1) = \mathbb{1}_{q^{-1}}$ .

The  $\mathcal{E}$  Fock modules in parallel realization were constructed in [FJMM1] using the inductive limit of tensor products of vector representations, an analog of the semi-infinite wedge construction.

The character of the Fock space does not factorize, but we have the usual partition formula when  $z_0 = z_1 = z$ .

$$\chi_{F_i}(z, z) = \frac{1}{\prod_{i=1}^{\infty} (1 - z^i)}$$

The Fock module  $F_0$  restricted to the vertical  $U_q^v \widehat{\mathfrak{gl}}_2 \subset \mathcal{E}$  subalgebra is the irreducible basic module of level q. It is generated by a singular vector v such that  $K_0v = qv$  and  $K_1v = v$ . If  $q = q_3$ , then  $F_0$  is also an evaluation module and  $\Psi(F_0)$  has the form (3.4) with a = q and b = u = 1.

# 4.3. Macmahon modules. We now define the Macmahon module $M_0^{\kappa}$ with $\kappa \in \mathbb{C}^{\times}$ .

Let  $\kappa$  be generic. It means  $\kappa \notin q_1^{\mathbb{Z}} q_2^{\mathbb{Z}}$ . Let

$$\Psi(M_0^{\kappa}) = \left(\frac{z\kappa - \kappa^{-1}}{z - 1}, 1\right).$$

Let the set of boxes consist of boxes with coordinates (i, j, k), and color  $i + k \mod 2$ , where  $i, j, k \in \mathbb{Z}_{\geq 0}$ . All the boxes are positive, we have no negative boxes. The ordering of boxes is lexicographic.

For partitions  $\lambda, \mu$  we say  $\mu \subset \lambda$  if for all  $j, \lambda_j \geq \mu_j$ . A plane partition  $\lambda$  is a set of partitions  $\lambda^{(1)} \supset \lambda^{(2)} \supset \cdots \supset \lambda^{(s)}$  called layers. We do not distinguish between plane partitions which differ by zero layers.

The set of states  $S(M_0^{\kappa})$  is the set of all plane partitions. The reference state is the empty plane partition.

The Young diagram of a plane partition  $\lambda$  is a union of Young diagrams of layers  $\lambda^{(j)}$  placed on height j-1. The boxes corresponding to  $\lambda_i^{(j)}$  have coordinates (0, j-1, i-1), (1, j-1, i-1)1), ...,  $(\lambda_i^{(j)} - 1, j - 1, i - 1).$ 

Note that the colors alternate along  $q_1$  and  $q_3$  axes and repeat along  $q_2$  axes. The starting box with coordinates (0, 0, 0) has color 0.

An example of a state in  $M_0^{\kappa}$  is given in Figure 4. In the figure,  $\lambda^{(1)} = (4, 2, 1), \lambda^{(2)} = (3, 1),$ and  $\lambda^{(3)} = (1)$ .



FIGURE 4. A state in the Macmahon module.

**Proposition 4.3.** The space  $M_0^{\kappa} = V(S(M_0^{\kappa}), \Psi(M_0^{\kappa}))$  with action (3.5) is an irreducible tame highest  $\ell$ -weight  $\mathcal{E}$ -module of highest  $\ell$ -weight  $\Psi(M_0^{\kappa})$ . In particular, the level of  $M_0^{\kappa}$  is  $\kappa$ .

*Proof.* The proof is similar to the proof of Proposition 4.2.

We call the module  $M_0^{\kappa}$  constructed in Proposition 4.3 the Macmahon module of color 0.

The Macmahon module  $M_1^{\kappa}$  of color 1 is the irreducible highest  $\ell$ -weight  $\mathcal{E}$ -module with highest  $\ell$ -weight  $\Psi(M_1^{\kappa}) = (1, \frac{z\kappa - \kappa^{-1}}{z-1})$ . It is obtained from  $M_0^{\kappa}$  by exchanging colors.

The Macmahon modules are building blocks for highest  $\ell$ -weight  $\mathcal{E}$ -modules. The Macmahon module was constructed explicitly in [FJMM1] using inductive limit construction of tensor products of Fock modules.

We note the combinatorial difference between the Fock module  $F_0$  and Macmahon module  $M_0^{\kappa}$ . Both modules are generated from the empty reference state which has a pole z = 1 in the zero component of the  $\ell$ -weight. After adding a white box the  $\ell$ -weight is multiplied by  $A_{0,1}^{-1}$ . This creates two poles at  $z = q_1^{-1}$  and  $z = q_3^{-1}$  in the first component of the  $\ell$ -weight. The difference is in the zero component. In the case of the Fock module, the zero  $z = q^{-1}$  of  $\Psi(F_0) = 0_{q^{-1}}$  cancels the pole of  $A_0^{-1}$  and no more poles are created. While in the Macmahon case such a cancellation does not happen and therefore, one obtains an additional concave box with coordinates (0, 1, 0). In addition, since the zero of  $\Psi(M_0^{\kappa})$  is not used, it can be arbitrary and therefore the Macmahon module depends on a parameter  $\kappa$ .

For special values of  $\kappa$  the Macmahon module  $M_0^{\kappa}$  may develop a submodule due to the factor  $(\kappa p(\Box) - \kappa^{-1})$  in the coefficients  $b_{\lambda,\Box}$  in the action of  $E_0(z)$ . To take advantage of this construction, choose a white box  $\Box$ ,  $c(\Box) = 0$  with coordinates (i, j, k) such that  $\min\{i, j, k\} < 1$  and specialize  $\kappa^2 = p(\Box)^{-1} = q_1^i q_2^j q_3^k$ . Then the action of  $E_0(z)$  cannot remove the chosen box due to vanishing of the coefficient  $b_{\lambda,\Box}$ . As the result we obtain an irreducible tame  $\mathcal{E}$ -module  $\overline{M}_0^{\kappa}$  whose basis is given by all plane partitions which do not contain the chosen white box, see [FJMM1].

The following two cases are of special importance since their restrictions to the horizontal  $U_{a}\widehat{\mathfrak{gl}}_{2}$  are irreducible.

Let  $\kappa = q$ . Then the prohibited box has coordinates (0, 1, 0) and the corresponding module has a basis consisting of colored plane partitions with one layer. Moreover,  $\Psi(M_0^q) = \Psi(F_0)$ . So the irreducible quotient  $\overline{M}_0^q$  is the Fock module  $F_0$ .

We show a state in the Fock space  $F_0$  corresponding to partition  $\lambda = (4, 2, 1)$  in Figure 5. The prohibited box is shown dashed.

Let  $\kappa = q_3$ . Then the prohibited box has coordinates (0, 0, 2). We denote the corresponding irreducible  $\mathcal{E}$ -module  $\bar{M}_0^{q_3}$  by  $G_0$ . One can say that this module has a basis labeled by plane partitions with two layers placed vertically. Importantly, the coloring is different since vertically the coloring does not change.

We show a state in the module  $G_0$  corresponding to the partition  $\boldsymbol{\lambda} = ((4,3), (2,1), (1), (1))$ in Figure 6. The same picture can be seen as two layer partition  $\boldsymbol{\mu} = ((4,2,1,1), (2,1,1))$ placed vertically. The prohibited box is shown dashed.



FIGURE 5. The prohibited box and the Fock module.

Note that we have a similar module corresponding to the choice  $\kappa = q_1$  (the prohibited box has coordinates (2, 0, 0)). Also, if one starts from  $M_1$ , the same choices of  $\kappa$  produces two similar modules. The difference in pictures is the color of the initial box with coordinates (0, 0, 0).



FIGURE 6. A state in the  $G_0$  module.

The character of the Macmahon modules is given by, see Theorem 7.6 in [BM],

$$\chi_{M_0^{\kappa}}(z_0, z_1) = \frac{1}{\prod_{i=1}^{\infty} (1 - (z_0 z_1)^i)^{2i} (1 - z_0 (z_0 z_1)^{i-1})^i (1 - z_1 (z_0 z_1)^{i-1})^{i-1}}$$

In particular, we have the famous Macmahon formula for plane partitions,

$$\chi_{M_0^{\kappa}}(z,z) = \frac{1}{\prod_{i=1}^{\infty} (1-z^i)^i}$$

The Macmahon module  $M_0^{\kappa}$  restricted to the vertical  $U_q^v \widehat{\mathfrak{gl}}_2 \subset \mathcal{E}$  subalgebra is not irreducible. The reference state  $|\emptyset\rangle$  is a highest  $\ell$ -weight vector of weight  $(\kappa, 1)$  and therefore the irreducible  $U_q^v \widehat{\mathfrak{gl}}_2$ -subquotient containing the reference state is a parabolic Verma module. When  $\kappa = q_3$ , the module  $G_0$  is the evaluation parabolic Verma module. In particular, the weight  $\Psi(M_0^{q_3})$  has form (3.4) with  $a = q_3$ , b = u = 1. Note that  $q_3$  is a generic number with respect to the parameter q of  $U_q^v \widehat{\mathfrak{gl}}_2$ .

## 4.4. Evaluation Verma modules. Let $\mu$ be a non-integer complex number. Let

$$\Psi(G_0^{\mu}) = \left(q_3 q^{-\mu} \frac{z - q_3^{-2}}{z - q_2^{-\mu}}, \ q^{\mu} \frac{z - q_3^{-1} q_2^{-\mu}}{z - q_3^{-1}}\right).$$

Let the set of boxes consist of boxes with positions (i, k, 1), color  $i + 1 \mod 2$ , and of boxes with positions  $(i, k + \mu, 0)$ , color  $i \mod 2$ . Here  $i, k \in \mathbb{Z}_{\geq 0}$ . All the boxes are positive, we have no negative boxes. The ordering of boxes with the same z-coordinate is lexicographic. The boxes with z-coordinate 1 are larger than the boxes with z-coordinate 0.

In fact, the last condition actually could be opposite.

The set of states  $S(G_0^{\mu})$  is the set of pairs of partitions positioned vertically. The reference state is the pair of empty partitions.

A state in  $S(G_0^{\mu})$  is shown on Figure 7. Here we read the two partitions vertically: (4, 2, 1, 1) and (2, 1, 1). Since the position of the first partition is shifted by  $(0, \mu, 0)$  we picture it on a pedestal of height  $\mu$  which is infinite in direction of  $q_1$  and one box thick (shaded).



FIGURE 7. A state in the Verma module  $G_0^{\mu}$ .

**Proposition 4.4.** The space  $G_0^{\mu} = V(S(G_0^{\mu}), \Psi(G_0^{\mu}))$  with action (3.5) is an irreducible tame highest  $\ell$ -weight  $\mathcal{E}$ -module of highest  $\ell$ -weight  $\Psi(G_0^{\mu})$ . In particular, the level of  $G_0^{\mu}$  is  $q_3$ .

*Proof.* The proof is similar the proof of Proposition 4.2.

Alternatively, one can construct the module  $G_0^{\mu}$  by starting with the  $G_0$  module. Consider states in  $G_0$  which contain all the boxes in the pedestal (with positions (i, j, 0) where  $i = 0, 1, \ldots, \mu - 1$ ,  $j = 0, 1, \ldots, \lambda - 1$  with (large) positive integers  $\lambda, \mu$ ) and observe that the

matrix coefficients stabilize as  $\lambda \to \infty$  and can be continued with respect to  $\mu$  to generic values. We do not give further details.

The character of the module  $G_0^{\mu}$  is a product of those of vertical partitions. Unlike the case of the Fock modules, the character of the vertical partitions is readily computed similarly to the non-graded case. One simply reads the partition along the columns. Set  $t = z_0 z_1$ . Thus, the character of one vertical partition whose top box has color *i* is

$$\bar{\chi}_i(z_0, z_1) = \frac{1}{\prod_{j=1}^{\infty} (1 - t^j)(1 - z_i t^{j-1})}$$

and

$$\chi_{G_0^{\mu}}(z_0, z_1) = \bar{\chi}_0(z_0, z_1)\bar{\chi}_1(z_0, z_1) = \frac{1}{\prod_{i=1}^{\infty} (1 - t^i)^2 (1 - z_0 t^{i-1})(1 - z_1 t^{i-1})}$$

The module  $G_0^{\mu}$  restricted to the vertical  $U_q^{\nu}\mathfrak{gl}_2 \subset \mathcal{E}$  subalgebra is the Verma module of weight  $(q_3q^{-\mu}, q^{\mu})$ . Moreover,  $G_0^{\mu}$  is the evaluation module obtained from the Verma module, and  $\Psi(G_0^{\mu})$  is given by (3.4) with  $a = q_3q^{-\mu}, b = q^{\mu}$ , and  $u = q^{-\mu}q_3^{-1}$ .

The module  $G_1^{\mu}$  is obtained from  $G_0^{\mu}$  by interchanging colors. We have the equality  $\Psi(G_1^{\mu})(z) = \Psi(G_0^{\beta-\mu})(q^{-\beta+2\mu}z)$ , where  $\beta$  is defined by  $q_3 = q^{\beta}$ . Thus, up to a twist by the shift automorphism  $\hat{\tau}_a$ , the module  $G_1^{\mu}$  is isomorphic to  $G_0^{\beta-\mu}$ .

4.5. Evaluation relaxed Verma modules. Let  $\mu, \nu$  be complex numbers. We assume that  $\mu, \nu$  and  $\mu + \nu$  are not integers. Let

$$\Psi(G_0^{\mu,\nu}) = \left(q_3 q^{-\mu-2\nu} \frac{(z-q_3^{-2})(z-q_2^{-\mu+1})}{(z-q_2^{-\mu-\nu})(z-q_2^{-\mu-\nu+1})}, \ q^{\mu+2\nu} \frac{(z-q_3^{-1}q_2^{-\mu-\nu})(z-q_1^{-1}q_2^{-\mu-\nu})}{(z-q_3^{-1})(z-q_1^{-1}q_2^{-\mu})}\right).$$

Let the set of positive boxes consist of boxes with coordinates (i, k, 1), color  $i + 1 \mod 2$ , of boxes with coordinates  $(i + 1, k + \mu, 0)$ , color  $i + 1 \mod 2$  and of boxes with positions  $(0, \mu + \nu + k, 0)$ , color 0. Here  $i, k \in \mathbb{Z}_{\geq 0}$ . In addition, we have negative boxes with coordinates  $(0, \mu + \nu - k, 0)$ , color 0,  $k \in \mathbb{Z}_{>0}$ .

Thus we have three families of boxes. Each family is ordered lexicographically. In addition, we declare that boxes from the last group are larger than boxes in the second group which are in their turn larger than boxes in the first group. (The last assignment can be changed arbitrarily.)

The set of states  $S(G_0^{\mu,\nu})$  is the set of pairs of partitions positioned vertically and an integer k. The integer k indicates the number of boxes put on the box with coordinates  $(0, \mu + \nu, 0)$  (negative k means removing |k| boxes). The reference state is the pair of empty partitions and integer 0.

A state in  $S(G_0^{\mu,\nu})$  is shown on Figure 8. Here we read the two partitions vertically: (2,2,1) and (2,1) and k=2.

Since the position of the boxes above (0, 0, 0) are shifted by  $(1, \mu + \nu, 1)$  we picture it on a tower of height  $\nu$  which is placed on the previous pedestal (shaded). Note that for negative k, |k| boxes are taken off that tower.

Alternatively, one can start with the  $G_0^{\mu}$  module. Then consider states which contain the boxes in the tower above (0,0,0) (with positions  $(0, j + \mu, 0)$  where  $j = 0, 1, \ldots, \nu - 1$  with

(large) positive integers  $\nu$ ) and observes that the matrix coefficients can be continued with respect to  $\nu$  to generic values. We do not give further details.

For that reason we picture the tower over the first box on the pedestal. Though, since  $\nu$  is generic, we could place it over any prohibited white box.



FIGURE 8. A state in the evaluation relaxed Verma module  $G_0^{\mu,\nu}$ .

**Proposition 4.5.** The space  $G_0^{\mu,\nu} = V(S(G_0^{\mu,\nu}), \Psi(G_0^{\mu,\nu}))$  with action (3.5) is an irreducible tame  $\mathcal{E}$ -module of level  $q_3$ .

*Proof.* The proof follows the proof of Proposition 4.2.

The character of  $G_0^{\mu,\nu}$  is easy to compute. We set  $t = z_0 z_1$ .

$$\chi_{G_0^{\mu,\nu}}(z_0, z_1) = \bar{\chi}_1^2(z_0, z_1) \sum_{i \in \mathbb{Z}} z_0^i = \frac{1}{\prod_{j=1}^\infty (1 - t^j)^2 (1 - z_1 t^{j-1})^2} \sum_{i \in \mathbb{Z}} z_0^i$$
$$= \frac{1}{\prod_{j=1}^\infty (1 - t^j)^2 (1 - z_0^{-1} t^j)^2} \sum_{i \in \mathbb{Z}} z_0^i = \frac{1}{\prod_{j=1}^\infty (1 - t^j)^4} \sum_{i \in \mathbb{Z}} z_0^i = \frac{1}{\prod_{j=1}^\infty (1 - t^j)^4} \,\delta(z_0).$$

Here we used the identity  $\delta(z)f(z) = 1$  which holds for any power series f(z) in  $z^{-1}$ .

The character of  $G_1^{\mu,\nu}$  coincides with the character of  $\hat{L}(a, b, c; \kappa)$ , cf. Figure 1. Note that by (3.2), the degree of a black box on Figure 1 is (-1, 0) while the degree of a white box is (1, -1).

The module  $G_1^{\mu,\nu}$  restricted to the vertical  $U_q^{\nu}\widehat{\mathfrak{gl}}_2 \subset \mathcal{E}$  subalgebra is the irreducible relaxed Verma module  $\hat{L}(a, b, c; \kappa)$  with  $q^a = q_3 q^{-\mu-2\nu}$ ,  $q^b = 1$ ,  $c = (q^{\nu} - q^{-\nu})(q^{a+1+\nu} - q^{-a-1-\nu})/(q - q^{-1})^2$  and  $\kappa = q_3$ , so that  $\mathcal{C} = q_3 q^{-\mu+1} + q_3^{-1} q^{\mu-1}$ . Moreover,  $G_1^{\mu,\nu}$  is the evaluation module obtained from  $\hat{L}(a, b, c; \kappa)$ .

Similarly, the module  $G_0^{\mu,\nu}$  is the evaluation module obtained from slope one relaxed module  $\hat{L}_1(a, b, c; \kappa)$ .

4.6. Evaluation slanted relaxed Verma modules. Let  $\mu, \nu$  be complex numbers. We assume that  $\mu, \nu$  and  $\mu + \nu$  are not integers. Let  $m \ge 1$ .

Let

$$\begin{split} \Psi(G_{0,m}^{\mu,\nu}) &= (\Psi^0(G_{0,m}^{\mu,\nu}), \Psi^1(G_{0,m}^{\mu,\nu})), \\ \Psi^0(G_{0,m}^{\mu,\nu}) &= q^{-\mu-2\nu} q_3^{-m+1} \frac{(z-q_3^{-2})(z-q_1^{-1}q_3^{-1}q_2^{-\mu}) \prod_{i=1}^{m-1} (z-q_2^{-\mu-\nu}q_1^i q_3^{-i})}{\prod_{i=0}^r (z-q_2^{-\mu-\nu+1}q_1^i q_3^{-i})} \\ \Psi^1(G_{0,m}^{\mu,\nu}) &= q^{\mu+2\nu} q_3^m \frac{\prod_{i=0}^{m+1} (z-q_1^{i-1}q_3^{-i}q_2^{-\mu-\nu})}{(z-q_1^{-1}q_2^{-\mu}) \prod_{i=0}^{m-1} (z-q_1^{i+1}q_3^{-i}q_2^{-\mu-\nu})}. \end{split}$$

It is important to note

$$\Psi(G_{0,m}^{\mu,\nu})\prod_{i=0}^{m}A_{0,q_{1}^{i}q_{3}^{-i}q_{2}^{-\mu-\nu}}^{-1}\prod_{i=0}^{m-1}A_{1,q_{1}^{i+1}q_{3}^{-i}q_{2}^{-\mu-\nu}}^{-1}=\Psi(G_{0,m}^{\mu,\nu+1}).$$

Here we take a product of  $\ell$ -weights of m black boxes and m+1 white boxes forming a staircase as shown on Figure 9 (with m = 4) with y-coordinate (height)  $\mu + \nu$ .



FIGURE 9. A top view of the reference state in the slope 4 slanted relaxed Verma module  $G_{0,4}^{\mu,\nu}$ .

Let the set of positive boxes consist of boxes with coordinates (i, k, 1), color  $i + 1 \mod 2$ , of boxes with coordinates  $(i + 1, k + \mu, 0)$ , color  $i + 1 \mod 2$ , of boxes with coordinates  $(-s, \mu + \nu + k, s)$ , color 0, with  $s = 0, \ldots, m$ , and of boxes with coordinates  $(-1 - s, \mu + \nu + k, s)$ , color 1, with  $s = 0, \ldots, m - 1$ . Here  $i, k \in \mathbb{Z}_{\geq 0}$ . In addition, we have negative boxes with boxes with coordinates  $(-s, \mu + \nu - k, s)$ , color 0, where  $s = 0, \ldots, m$ , and with coordinates  $(-1 - s, \mu + \nu - k, s)$ , color 1, with  $s = 0, \ldots, m - 1$ ,  $k \in \mathbb{Z}_{\geq 0}$ .

In other words, we can have boxes vertically in the first two rows shown by dashed lines in Figure 9 and boxes above the staircase or below the staircase. The boxes below the staircase are negative. The boxes on the first row are lifted by  $\mu$ , the ones on the staircase by  $\mu + \nu$ .

Thus we have three families of boxes: on the bottom, on the pedestal and at the tower. Each family is ordered lexicographically. In addition, we declare that boxes from the last group are

larger than boxes in the second group which are in their turn larger than boxes in the first group. (The last assignment can be changed arbitrarily.)



FIGURE 10. A state in the slope 1 slanted relaxed Verma module  $G_{0.1}^{\mu,\nu}$ .

The set of states  $S(G_{0,m}^{\mu,\nu})$  is the set of pairs of partitions positioned vertically and integers  $a_0, \ldots, a_m, b_0, \ldots, b_{m-1}$  such that

$$b_s \ge a_s, \quad b_s \ge a_{s+1}, \qquad s = 0, \dots, m-1.$$

The integer  $a_s$  indicates the number of the white boxes put on the box with coordinates  $(s, \mu + \nu, -s)$  (negative  $a_s$  means removing  $|a_s|$  boxes). The integer  $b_s$  indicates the number of the black boxes put on the box with coordinates  $(s-1, \mu + \nu, -s)$  (negative  $b_s$  means removing  $|b_s|$  boxes).

The reference state is the pair of empty partitions and integer 0.

An example of a state in  $S(G_{0,1}^{\mu,\nu})$  is given in Figure 10. In the picture we have two vertical partitions (2, 2, 1) and (2, 1) and on the tower we have  $a_0 = 2$ ,  $a_1 = 1$ ,  $b_1 = 2$ .

We also show a reference state of  $S(G_{0,2}^{\mu,\nu})$  in Figure 11. Note that one can add boxes to the white and black colored places. In addition one can add negative boxes which amounts to taking off boxes from the staircase tower.

**Proposition 4.6.** The space  $G_{0,m}^{\mu,\nu} = V(S(G_{0,m}^{\mu,\nu}), \Psi(G_{0,m}^{\mu,\nu}))$  with action (3.5) is an irreducible tame  $\mathcal{E}$ -module of level  $q_3$ . Moreover, for any  $k \in \mathbb{Z}$ , we have  $G_{0,m}^{\mu,\nu} \simeq G_{0,m}^{\mu,\nu+k}$ .

*Proof.* The proof is similar to the proof of Proposition 4.2.

Now we compute the character of  $G_{0,m}^{\mu,\nu}$ . We recall our notation  $t = z_0 z_1$  and  $\delta(z) = \sum_{i \in \mathbb{Z}} z^i$ .



FIGURE 11. The reference state in the slope 2 slanted relaxed Verma module  $G_{0,2}^{\mu,\nu}$ .

**Proposition 4.7.** We have

$$\chi_{G_{0,m}^{\mu,\nu}}(z_0, z_1) = \frac{1}{\prod_{j=1}^{\infty} (1-t^j)^4} \,\delta(z_0 t^m)$$

Proof. We have

$$\chi_{G_{0,m}^{\mu,\nu}}(z_0, z_1) = \bar{\chi}_1^2(z_0, z_1)\chi_m(z_0, z_1) = \frac{\chi_m(z_0, z_1)}{\prod_{i=1}^{\infty}(1 - t^i)^2(1 - z_1t^{i-1})^2} = p(t)\sum_{i \in \mathbb{Z}}(z_0t^m)^i,$$

where  $\chi_r(z_0, z_1)$  is the contribution of the tower and p(t) is some series. The last equality takes place since we have  $G_{0,m}^{\mu,\nu} \simeq G_{0,m}^{\mu,\nu+1}$  and

$$\chi_{G_{0,m}^{\mu,\nu+1}}(z_0, z_1) = \chi_{G_{0,m}^{\mu,\nu}}(z_0, z_1)t^m z_0 = \chi_{G_{0,m}^{\mu,\nu}}(z_0, z_1).$$

To compute p(t) we write the character as a series in  $t, z_0$  and compute the coefficient of  $z_0^0$ .

The state in the tower is characterized by integer numbers  $b_0, \ldots, b_{m-1}, a_0, \ldots, a_m$  satisfying  $b_i \ge a_i, a_{i+1}$  which tell the amount of boxes on the tower. Let  $c_i = b_i - a_i$  and  $d_i = b_i - a_{i+1}$ ,  $i = 1, \ldots, m$ . Then  $b_i$  and  $a_i$  are uniquely determined by  $a_0, c_i$  and  $d_i$ . Moreover,  $c_i \ge 0$  and  $d_i \ge 0$ .

Note that if  $a_j < 0$  for some j, then

$$\sum_{i=0}^{m-1} b_i - \sum_{i=0}^{m-1} a_i = \sum_{i=0}^{j-1} c_i - a_j + \sum_{i=j}^{m-1} d_i > 0.$$

So the contribution of this state is  $z_0^i z_1^j = t^j z_0^{i-j}$  with j > i. Since all monomials in  $\bar{\chi}_1(z_0, z_1)$  have the form  $z_0^i z_1^j$  with  $i \leq j$ , such a state does not contribute to the coefficient of  $(z_0)^0$ . Thus we can assume  $a_i \geq 0$  and therefore  $b_i \geq 0$  for all i.

Let the shape  $R_i$  consist of *i* horizontal dominoes aligned to form a staircase, see Figure 12. Adding  $R_i$  to the tower aligning the white end with the box with coordinates (0, x, 0) changes  $d_i$  to  $d_i + 1$  keeping  $a_0, c_j$ , and  $d_k, k \neq i$  intact.

Similarly, let shape  $\bar{R}_i$  consist of *i* vertical dominoes aligned to form a staircase, see Figure 12. Adding  $\bar{R}_i$  to the tower aligning the white end with the box with coordinates (-m, x, m) changes  $c_{m-i}$  to  $c_{m-i} + 1$  keeping  $a_0, d_j$ , and  $c_k, k \neq m - i$  intact.



FIGURE 12. The shapes  $R_i$  and  $R_i$ .

Therefore a state corresponding  $a_i$  and  $b_i$  can be obtained adding to the reference state  $a_0$  full staircases,  $c_i$  shapes  $R_i$  and  $d_i$  shapes  $\bar{R}_i$ . Thus the character of all states with  $a_0 > 0$  is

$$A(z_0,t) = \frac{\bar{\chi}_1^2(z_0,z_1)}{(1-z_0t^m)\prod_{j=1}^m(1-t^j)^2} = \frac{1}{(1-z_0t^m)\prod_{j=1}^m(1-t^j)^2\prod_{i=1}^\infty(1-t^i)^2(1-z_0^{-1}t^i)^2}$$

Let |t| < 1. Then the coefficient of  $z_0$  can be computed as an integral:

$$p(t) = \frac{1}{2\pi i} \int_{|z_0|=1} A(z_0, t) \frac{dz_0}{z_0} = - \operatorname{res}_{z_0=t^{-m}} \frac{A(z_0, t)}{z_0} = \frac{1}{\prod_{i=1}^{\infty} (1 - t^i)^4}.$$

Note that the character of  $G_{1,m}^{\mu,\nu}$  coincides with the character of  $\hat{L}_{-m}(a,b,c;\kappa)$ , cf. Figure 11.

The module  $G_{1,m}^{\mu,\nu}$  restricted to the vertical  $U_q^v \widehat{\mathfrak{gl}}_2 \subset \mathcal{E}$  subalgebra is the irreducible relaxed Verma module  $\hat{L}_{-m}(a, b, c; \kappa)$  with  $q^a = q_3^{m+1}q^{-\mu-2\nu}$ ,  $q^b = 1$ ,  $\kappa = q_3$ , and suitable c. Moreover,  $G_{1,m}^{\mu,\nu}$  is the evaluation module obtained from  $\hat{L}_{-m}(a, b, c; \kappa)$ .

Similarly, the module  $G_{0,m}^{\mu,\nu}$  is the evaluation module obtained from slope m+1 relaxed module  $\hat{L}_{m+1}(a, b, c; \kappa)$ .

Acknowledgments. MJ is partially supported by JSPS KAKENHI Grant Numbers JP19K03509, JP20K03568. EM is partially supported by Simons Foundation grant number #709444.

EM thanks Rikkyo University, where a part of this work was done, for hospitality.

We thank B. Feigin for interesting discussions.

#### References

- [B31] H. Bethe, Zur Theorie der Metalle, Z. Physik **71** (1931), 205–226
- [BM] L. Bezerra and E. Mukhin, Representations of quantum toroidal superalgebras and plane s-partitions, arXiv:2104.05841, 1-42
- [BrM] M. Brito and E. Mukhin, Representations of quantum affine algebras of type  $B_N$ , Trans. Amer. Math. Soc., **369** (2017), no. 4, 2775-2806
- [FFJMM] B. Feigin, E. Feigin, M. Jimbo, T. Miwa, and E. Mukhin, Quantum continuous  $\mathfrak{gl}_{\infty}$  tensor products of Fock modules and  $W_n$ -characters, Kyoto J. Math., **51** (2011), no. 2, 365–392
- [FJMM1] B. Feigin, M. Jimbo, T. Miwa, and E. Mukhin, Representations of quantum toroidal  $\mathfrak{gl}_n$ , J. Algebra, **380** (2013), 78–108
- $\label{eq:generalized} \begin{array}{l} [\mathrm{FJMM2}] \ \mathrm{B.\ Feigin,\ M.\ Jimbo,\ T.\ Miwa,\ and\ E.\ Mukhin,\ Branching\ rules\ for\ quantum\ toroidal\ \mathfrak{gl}_n,\ \mathrm{Adv.\ Math.}, \\ \mathbf{300}\ (2016),\ 229-274 \end{array}$
- [FJM1] B. Feigin, M. Jimbo, and E. Mukhin, Evaluation modules for quantum toroidal gl<sub>n</sub> algebras, Progr. Math., 337 (2021), 393–425
- [FJM2] B. Feigin, M. Jimbo, and E. Mukhin, Combinatorics of vertex operators and deformed W-algebra of type D(2, 1; α), Adv. Math., 403 (2022), Paper No. 108331, 54 pp
- [FM1] E. Frenkel and E. Mukhin, Combinatorics of q-characters of finite-dimensional representations of quantum affine algebras, Comm. Math. Phys., 216 (2001), no. 1, 23–57
- [FM2] E. Frenkel and E. Mukhin, The Hopf algebra Rep  $U_q \hat{\mathfrak{gl}}_{\infty}$ , Selecta Math., 8 (2002), 537-635
- [FST] B. Feigin, A. Semikhatov, and I. Tipunin, Equivalence between chain categories of representations of affine  $\mathfrak{sl}(2)$  and N = 2 superconformal algebras, J. Math. Phys. **39** (1998), no. 7, 3865–3905
- [GT50] I. Gelfand and M. Tsetlin, Finite-dimensional representations of the group of unimodular matrices, Dokl. Akad. Nauk SSSR 71 (1950), 825–828 (Russian). English transl. in: I. M. Gelfand, "Collected papers". Vol II, Berlin: Springer-Verlag 1988, 653–656
- [K90] M. Kashiwara, Crystalizing the q-analogue of universal enveloping algebras, Comm. Math. Phys., 133, no. 2 (1990), 249–260
- [L90] G. Lusztig, Canonical bases arising from quantized enveloping algebras, J. of AMS, 3, (1990), no. 2, 447–498
- [M1] K. Miki, Toroidal braid group action and an automorphism of toroidal algebra  $U_q(\mathfrak{sl}_{n+1,tor})$   $(n \ge 2)$ , Lett. Math. Phys., 47 (1999), no. 4, 365-378
- [M2] K. Miki, Toroidal and level 0  $U'_q \widehat{sl_{n+1}}$  actions on  $U_q \widehat{gl_{n+1}}$  modules, J. Math. Phys., **40** (1999), no. 6, 3191-3210
- [NT] M. Nazarov and V. Tarasov, Yangians and Gelfand-Zetlin bases, Publ. RIMS, Kyoto Univ. 30 (1994) 459–478
- [OV05] A. Vershik and A. Okounkov, A New Approach to the Representation Theory of the Symmetric Groups. II, J. Math. Sciences., 131 (2005), 5471–5494

MJ: PROFESSOR EMERITUS, RIKKYO UNIVERSITY, TOSHIMA-KU, TOKYO 171-8501, JAPAN *Email address*: jimbomm@rikkyo.ac.jp

EM: DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY INDIANAPOLIS, 402 N. BLACKFORD ST., LD 270, INDIANAPOLIS, IN 46202, USA

Email address: emukhin@iu.edu