

# THE CUBIC NONLINEAR SCHRÖDINGER EQUATION WITH ROUGH POTENTIAL

NORBERT J. MAUSER, YIFEI WU, AND XIAOFEI ZHAO

ABSTRACT. We consider the cubic nonlinear Schrödinger equation with a spatially rough potential, a key equation in the mathematical setup for nonlinear Anderson localization. Our study comprises two main parts: new optimal results on the well-posedness analysis on the PDE level, and subsequently a new efficient numerical method, its convergence analysis and simulations that illustrate our analytical results. In the analysis part, our results focus on understanding how the regularity of the solution is influenced by the regularity of the potential, where we provide quantitative and explicit characterizations. Ill-posedness results are also established to demonstrate the sharpness of the obtained regularity characterizations and to indicate the minimum regularity required from the potential for the NLS to be solvable. Building upon the obtained regularity results, we design an appropriate numerical discretization for the model and establish its convergence with an optimal error bound. The numerical experiments in the end not only verify the theoretical regularity results, but also confirm the established convergence rate of the proposed scheme. Additionally, a comparison with other existing schemes is conducted to demonstrate the better accuracy of our new scheme in the case of a rough potential.

## CONTENTS

1. Introduction	2
1.1. Well-posedness theory	3
1.2. Ill-posedness theory	4
1.3. Numerical counterpart	5
2. Preliminaries	6
2.1. Basic notations	6
2.2. Fourier transform	7
2.3. Some operators	8
2.4. Some tool lemmas	8
3. Proof for well-posedness theory	8
3.1. Proof of Theorem 1.1	9
3.2. Proof of Theorem 1.2	19
4. Proof for ill-posedness theory	21
4.1. Proof of Theorem 1.3	22
4.2. Proof of Theorem 1.4	25
4.3. Proof of Theorem 1.5	25
5. Numerical method and convergence analysis	29
5.1. Construction of numerical scheme	29
5.2. Convergence analysis	30
6. Numerical experiments	45
6.1. On regularity of solution	46

2010 *Mathematics Subject Classification.* 65M12, 65M15, 35Q55.

*Key words and phrases.* Nonlinear Schrödinger equation, rough potential, well-posedness, ill-posedness, numerical solution, convergence.

6.2. On accuracy of scheme	47
6.3. Accuracy comparison	49
7. Conclusion	52
Acknowledgements	52
References	52

## 1. INTRODUCTION

We deal with the following cubic nonlinear Schrödinger equation (NLS) with a (“rough”) spatial potential

$$\begin{cases} i\partial_t u(t, x) + \partial_{xx} u(t, x) + \xi(x)u(t, x) = \lambda|u(t, x)|^2 u(t, x), & \text{for } x \in \mathbb{T} \text{ and } t > 0, \\ u(0, x) = u_0(x), & \text{for } x \in \mathbb{T}. \end{cases} \quad (1.1)$$

Here  $u(t, x) : \mathbb{R}^+ \times \mathbb{T} \rightarrow \mathbb{C}$  is the complex-valued unknown “wave function”. We choose the setup on the one-dimensional torus  $\mathbb{T} = (-\pi, \pi)$ , which is an appropriate setting for efficient numerical methods.  $\xi(x) : \mathbb{T} \rightarrow \mathbb{R}$  is a given real-valued potential that can “rough” in terms of regularity.  $u_0(x)$  is a given initial wave function, and  $\lambda \in \mathbb{R}$  a given parameter. Here  $\lambda < 0$  resp-  $\lambda > 0$  corresponds to the focusing resp. defocusing nonlinear self-interaction cases.

The solution of this NLS satisfies the following mass and energy conservation laws:

$$\frac{1}{2\pi} \int_{\mathbb{T}} |u(t, x)|^2 dx = \frac{1}{2\pi} \int_{\mathbb{T}} |u_0(x)|^2 dx, \quad \text{for } t > 0, \quad (1.2a)$$

$$\int_{\mathbb{T}} \left[ |\partial_x u(t, x)|^2 - \xi(x)|u(t, x)|^2 + \frac{\lambda}{2}|u(t, x)|^4 \right] dx = \int_{\mathbb{T}} \left[ |\partial_x u_0|^2 - \xi|u_0|^2 + \frac{\lambda}{2}|u_0|^4 \right] dx. \quad (1.2b)$$

When the potential function  $\xi(x)$  is considered as a rough/random potential, (1.1) is often called the disordered NLS, as it arises in the context of the so-called Anderson localization: In 1958, Philip W. Anderson discovered for the linear and lattice Schrödinger model that the waves in the solution will be localized when the potential  $\xi$  is spatially random/rough enough [1]. Such phenomenon found important applications, e.g., for semiconductors. Note that the NLS (1.1) is the continuous version of a discrete/discretized “lattice” model. Nonlinearities in the Schrödinger equation pose many deep questions for physicists and mathematicians. E.g., in the defocusing NLS case where the interaction tends to push the wave to the far field, there is a long-standing debate about whether the localization dominates in the end or the spreading dominates. There is a wide range of works to address this problem from different aspects, e.g., [8, 20, 21, 28, 34, 37, 41]. Numerical simulations were done to suggest answers on the nonlinear lattice model, see e.g., [20, 21, 34].

However, the mathematical and numerical studies of (1.1) are far from done. With this article we contribute new results to both in a setup on the torus in one space dimension for (1.1) that serves as a valid truncation of localized wave dynamics up to a finite time.

For applications of the nonlinear localization model (1.1) in physics, we refer to [11, 15, 33, 36]. On the mathematical level, let us briefly review some theoretical results on (1.1) related to our work. Without the potential function, i.e.,  $\xi \equiv 0$ , it is known that (1.1) is globally well-posed in  $H^s$  for  $s \geq 0$ ; see e.g., [3, 39]. With  $\xi(x)$  an  $L^\infty$ -potential, Cazenave [10] showed the global well-posedness of (1.1) for small initial data in  $H^1$ . For potentials that are stochastic in time but rather regular in space, we refer to [9] for the well-posedness theory of the model. For less regular spatial random potentials, [12, 23, 24] studied the

spectrum of the linear part of the operator in (1.1), i.e.,  $-\partial_x^2 + \xi$ , and the results revealed the mechanism of localization in the continuous level. Rodnianski and Schlag [35] analyzed the decaying property of the solution for the linear Schrödinger model. For a spatial white noise potential, the work of [13, 14] shows that (1.1) has a solution almost surely in  $H^1$  provided the smallness of the initial data. For a deterministic rough potential or one precise sample of a spatial noise, the sharp regularity of the solution of (1.1) is not clearly known from the existing results.

Although the numerical methods for Schrödinger models have been extensively developed in the literature, see e.g., [2, 6, 18, 27], the existing studies mostly concern a smooth setup and do not work for the NLS equations with rough potentials. A clear understanding of the regularities in (1.1) is a basis for optimal numerical methods and convergence analysis, that allow for reliable simulations of the nonlinear localization or delocalization phenomenon of waves in the model (1.1). A rough potential  $\xi$  is felt in the solution  $u$  of (1.1) and will certainly bring serious numerical difficulties due to the simultaneous low-regularity in  $\xi$  and  $u$ . Indeed, the low-regularity not just affects the spatial discretization accuracy but also affects temporal discretization error, because the truncation error of an approximation usually involves derivatives of the functions. Popular traditional schemes for NLS models like the finite difference methods or operator/time splitting methods or exponential integrators can suffer from severe loss of accuracy when the solution is not smooth enough [7, 17, 32]. In numerical experiments below, we will show their poor performance on approximating (1.1) in case of rough potential.

In this work, we first address the well-posedness of the NLS equation (1.1). Although in the original spirit of Anderson, the potential is generated according to some random distribution, the localization phenomenon as such is not a mere stochastic behaviour. It occurs for any rough enough potential  $\xi$ . Therefore, we consider the deterministic case and focus on the effect of the roughness. In particular, we aim to understand how the regularity of the potential affects the regularity of the solution of (1.1). We then continue with the numerical analysis and propose accurate discretizations for (1.1). Hence, in the second part of the work, we are going to derive an efficient tailored low-regularity integrator for solving (1.1) based on the established regularity results from the first part, and then we address that under a rough potential with certain regularity, what the numerical method could offer for accuracy. The accurate computational results will in turn verify the theoretical regularity results of the solution.

The main theoretical results of this paper are given in the following theorems.

**1.1. Well-posedness theory.** Firstly, let us introduce the following space  $\hat{b}^{s,p}$  to characterize the regularity of a function  $f(x)$  on  $\mathbb{T}$  (particularly for the potential function  $\xi(x)$ ) based on its Fourier coefficients:

$$\hat{b}^{s,p} = \hat{b}^{s,p}(\mathbb{T}) \triangleq \{f(x) : |\hat{f}_0| + \||k|^s \hat{f}_k\|_{l^p} < +\infty\}, \quad f(x) = \sum_{k \in \mathbb{Z}} \hat{f}_k e^{ikx}, \quad x \in \mathbb{T}, \quad (1.3)$$

where  $s \geq 0$  describes the differentiability and  $1 \leq p \leq \infty$  gives the integrability. When  $p = 2$ , it is the usual Sobolev space  $\hat{b}^{s,2} = H^s$ . For simplicity, we denote  $\hat{b}^{0,p}$  by  $\hat{l}^p$ . A critical characteristic index  $\gamma_p$  is defined as

$$\gamma_p \triangleq \frac{3}{2} + \frac{1}{p}. \quad (1.4)$$

Before stating the results, we recall the following definition for the well-posedness.

**Definition 1.1** (Well-posedness). The well-posedness of a time dependent PDE can be defined as follows: Denote by  $C_t(I; X_0)$  the space of continuous functions from the time interval  $I$  to the topological space  $X_0$ . We say that the Cauchy problem is locally well-posed in  $C_t(I; X_0)$  if the following properties hold:

- (1) There is unconditional uniqueness in  $C_t(I; X_0)$  for the problem.
- (2) For every  $u_0 \in X_0$ , there exists a strong solution defined on a maximal time interval  $I = [0, T_{\max})$ , with  $T_{\max} \in (0, +\infty]$ .
- (3) The solution map  $u_0 \mapsto u[u_0]$  is continuous from  $X_0$  to  $X_0$ .

**Remark 1.1.** *The well-posedness defined above is stronger than the common definition, which can be regarded as the unconditional well-posedness.*

Now we state our first well-posedness result for the NLS equation (1.1).

**Theorem 1.1 (Well-posedness for  $\hat{b}^{s,p}$ -potential).** *Let  $\xi \in \hat{b}^{s,p}$  for  $s \geq 0, 2 < p \leq \infty$ , so  $\gamma_p \in (\frac{3}{2}, 2)$ . Then, (1.1) is locally well-posed in  $H^{s+\gamma_p-}(\mathbb{T})$  (notation  $\gamma_p-$  explained in Section 2.1 (iii), (iv)).*

**Remark 1.2.** *In Theorem 1.1, the regularity of the solution is essentially determined by the regularity of the potential. In order to focus on the effect from the potential, we will not address the impact from rough initial data in this work. On the other hand, requiring more regularity from the initial data will neither improve the well-posedness result. We will explain this point in Remark 1.4 and also in Section 6.1 by numerical experiments.*

**Remark 1.3.** *Note that  $L^1(\mathbb{T}) \hookrightarrow \hat{l}^\infty$ . Therefore, Theorem 1.1 with  $s = 0, p = \infty$  covers particularly the case of  $\delta$ -potential, i.e.,  $\xi(x) = \delta(x)$  in (1.1). Note that the well-posedness of NLS under a delta potential is known only in  $H^1$  in the literature [19, 22]. Now Theorem 1.1 states that the regularity  $H^{\frac{3}{2}-}$  can be attained, and this will be shown to be sharp.*

Let  $s \geq 0, 1 \leq r \leq \infty$ . Denote

$$H^{s,r} = H^{s,r}(\mathbb{T}) \triangleq \{f(x) : \|f\|_{L^r(\mathbb{T})} + \|(-\partial_x^2)^{\frac{s}{2}} f\|_{L^r(\mathbb{T})} < +\infty\}.$$

Note that by Young's inequality, we have that  $\hat{b}^{s,p} \hookrightarrow H^{s,p'}(\mathbb{T})$  where  $1/p' + 1/p = 1$ . If we replace  $\hat{b}^{s,p}(\mathbb{T})$  for  $s \geq 0, 2 \leq p < \infty$  by a slightly stronger space  $H^{s,p'}(\mathbb{T})$ , then we can cover the endpoint regularity.

**Theorem 1.2 (Well-posedness for  $H^{s,p'}$ -potential).** *Let  $\xi \in H^{s,p'}(\mathbb{T})$  for  $s \geq 0, 2 \leq p < \infty$  and so  $\gamma_p \in (\frac{3}{2}, 2]$ . Then, (1.1) is locally well-posed in  $H^{s+\gamma_p}(\mathbb{T})$ .*

**1.2. Ill-posedness theory.** When one of the conditions in Definition 1.1 is violated, the Cauchy problem (1.1) is said to be ill-posed in space  $X_0$ . In this work, we refer to the violation of the third condition (around zero solution), which is especially relevant to valid numerical approximations. More precisely, the definition is the following.

**Definition 1.2** (Ill-posedness). Let  $\epsilon > 0$  and denote

$$B(\epsilon) \triangleq \{u_0 \in \mathcal{S} : \|u_0\|_{X_0} \leq \epsilon\}, \quad \text{where } \mathcal{S} \text{ is the Schwartz space.} \quad (1.5)$$

By ‘‘ill-posedness’’ of (1.1) in  $X_0$  we mean that for arbitrarily small  $\epsilon > 0$ , there exists a  $u_0 \in B(\epsilon)$  such that the solution map  $u_0 \mapsto u[u_0]$  is discontinuous from  $X_0$  to  $C([0, 1]; X_0)$ .

The following result shows that for general  $\hat{b}^{s,p}$ -potential  $\xi(x)$ , we can only expect the solution to be in  $H^{s+\gamma_p}(\mathbb{T})$  at most. This hence implies the sharpness of Theorem 1.1.

**Theorem 1.3 (Ill-posedness for  $\hat{b}^{s,p}$ -potential).** *For  $s \geq 0, 2 < p \leq \infty$  and any  $\gamma \geq \gamma_p$ , there exists some  $\xi \in \hat{b}^{s,p}$  such that (1.1) is ill-posed in  $H^{s+\gamma}(\mathbb{T})$ .*

**Remark 1.4.** Note that the case  $s = 0, p = \infty$  is not covered in [Theorem 1.2](#), and indeed here [Theorem 1.3](#) states that the endpoint regularity does not hold for  $\xi \in \hat{l}^\infty$ . As a matter of fact, for the case of  $\xi \in \hat{l}^\infty$ , the counterexample we choose in the proof of [Theorem 1.3](#) later is exactly  $\xi(x) = \delta(x)$ . Thus, for the  $\delta$ -potential of [\(1.1\)](#), the regularity  $C_t^0([0, T]; H^{\frac{3}{2}-}(\mathbb{T}))$  is sharp. Moreover, in the proof we set a smooth initial data

$$u_0(x) = \epsilon(1 + 2 \cos(2x)), \quad \forall \epsilon > 0.$$

This in general means that the smoothness of initial function will not make things better.

For potentials that belong to the usual Sobolev space  $H^s$ , i.e.,  $p = p' = 1/2$  in [Theorem 1.2](#), the following result states that  $H^{s+2}$  is the most regularity that we can expect in general for the solution.

**Theorem 1.4 (Ill-posedness for  $H^s$ -potential).** For  $s \geq 0$  and any  $\gamma > 2$ , there exists some  $\xi \in H^s$  such that [\(1.1\)](#) is ill-posed in  $H^{s+\gamma}(\mathbb{T})$ .

Last but not least, we show by the following theorem that for the NLS equation [\(1.1\)](#) to be well-posed, the potential  $\xi \in \hat{l}^\infty$  is the lowest regularity requirement.

**Theorem 1.5 (Minimum regularity for potential).** For any  $s < 0$  and  $1 \leq p \leq \infty$ , there exists some  $\xi \in \hat{b}^{s,p} \setminus l^\infty$  such that for any  $\gamma \in \mathbb{R}$ , [\(1.1\)](#) is ill-posed in  $H^\gamma(\mathbb{T})$ .

**1.3. Numerical counterpart.** We are particularly interested in understanding the numerical issues for solving the NLS equation [\(1.1\)](#) for the rough potential case  $\xi \in \hat{b}^{s,p}$ . Traditional numerical methods could be completely inaccurate due to the roughness, and so the computational results are not reliable. Hence, the first goal would be a correct and accurate scheme that can work under the roughest case. Here we propose a simple and effective exponential type integration scheme which will be derived later in the spirit of a low-regularity integrator (LRI) [\[32\]](#).

Let  $\tau = \Delta t > 0$  be the time step with the time grid points  $t_n = n\tau$  for  $n = 0, 1, \dots$ , and denote  $u^n(x) \approx u(t_n, x)$  for the numerical solution. Our LRI scheme reads as follows: start with the exact initial value  $u^0(x) = u_0(x)$  and then update as

$$u^{n+1}(x) = i\tau \mathcal{D}_\tau [\xi(x) \mathcal{D}_\tau [u^n(x)]] + \mathcal{N}_\tau \left[ e^{i\tau \partial_x^2} u^n(x) \right], \quad n = 0, 1, \dots, \quad x \in \mathbb{T}, \quad (1.6)$$

where the two operators  $\mathcal{D}_\tau$  and  $\mathcal{N}_\tau$  are defined by

$$\mathcal{D}_\tau[f] \triangleq \left( e^{i\tau \partial_x^2} - 1 \right) (i\tau \partial_x^2)^{-1} f, \quad \mathcal{N}_\tau[f] \triangleq e^{-i\tau \lambda |P_{\leq N} f|^2} f, \quad (1.7)$$

with  $P_{\leq N} f \triangleq \sum_{|k| \leq N} \hat{f}_k e^{ikx}$  “cutting off high modes”, and with  $N = \tau^{-\frac{1}{2} + \varepsilon_0}$  for any  $0 < \varepsilon_0 \leq \frac{1}{8(s+\gamma_p)}$ . Here  $\hat{f}_k$  denotes the Fourier coefficient of  $f = f(x)$  for  $x \in \mathbb{T}$ , and  $P_{\leq N}$  is referred as the filter operator in [\[25, 26\]](#). The derivation of the scheme [\(1.6\)](#) will be given in [Section 5](#). Clearly, [\(1.6\)](#) is explicit in time, and thanks to the period boundary conditions that we chose, it can be efficiently implemented under Fourier spectral discretization [\[40\]](#) using FFT (fast Fourier transforms). In practice, when the number of spatial grid points which is also the number of total frequencies is greater than or equal to  $2N$ , the filter  $P_{\leq N}$  becomes trivial for the implementation. Moreover, such a filter primarily serves for theoretical results, and in practice the scheme performs very similarly without it.

For the semi-discretized (in time) LRI scheme [\(1.6\)](#), the following theorem gives its error estimate up to a fixed time of computation.

**Theorem 1.6 (Error estimate for the numerical method).** Let  $\xi \in \hat{b}^{s,p}$  with  $s \geq 0, 2 \leq p \leq +\infty$  in [\(1.1\)](#). For any  $u_0 \in H^{s+2}(\mathbb{T})$  and any  $0 < T < T_{\max}$ , there exist

constants  $C > 0, \tau_0 > 0$  such that for  $0 < \tau \leq \tau_0$ , the numerical solution given by (1.6) satisfies

$$\max_{0 \leq t_n \leq T} \|u(t_n) - u^n\|_{L^2} \leq C\tau^{\min\{s + \frac{1}{p} + \frac{1}{4} -, 1\}}, \quad (1.8)$$

where  $\tau_0$  and  $C$  depend only on  $T, \lambda, \|\xi\|_{\hat{b}^{s,p}}$  and  $\|u_0\|_{H^{s+2}}$ .

As illustrated in the above result, for the “worst” potential case  $\xi \in \hat{l}^\infty$ , the scheme (1.6) has  $\mathcal{O}(\tau^{\frac{1}{4}-})$  accuracy which is indeed quite low as a convergence rate but at least can guarantee the convergence and correctness of the computational results. The convergence order (1.8) will be verified by numerical experiments. In addition, numerical comparisons will highlight that the accuracy of (1.6) is indeed better than the existing schemes for (1.1). To achieve a higher order accurate scheme would be an extremely challenging future task. With the reliable numerical results provided by the LRI scheme (1.6), we will be able to verify the theoretical results given in Section 1.1. The predicted regularity in the theorems can be exactly observed in our numerical results (see Figure 1-Figure 3).

The main difficulty and innovation of the mathematical analysis in the work is summarized here. For the well-posedness and ill-posedness theorems, the low regularity of the considered  $\xi$ , for instance for  $\xi \in L^1$ , makes the usual Hölder type estimate for the product  $\xi u$  no longer available. Delicate frequency analysis has to be developed here. For the numerical analysis, the roughness limits the measure of error under the  $L^2$ -norm (1.8), where the algebraic property is missing. The established  $L^2$ -estimate here essentially benefits from the design of the approximation part  $\mathcal{D}_\tau[\xi \mathcal{D}_\tau[u^n]]$  in the scheme (1.6), which has a crucial inner product structure.

The rest of our paper is organized as follows. In Section 2, we introduce some preliminaries for the following analysis. In Section 3, we prove the well-posedness results for the NLS equation (1.1). In Section 4, we prove the ill-posedness results for (1.1). The numerical scheme and convergence analysis are given in Section 5. Numerical experiments are done in Section 6 and conclusions are drawn in Section 7.

## 2. PRELIMINARIES

As a preparation for subsequent PDE and numerical analysis, we give some notations and lemmas that will be frequently used in the rest of the paper.

**2.1. Basic notations.** We adopt the following widely used notations from harmonic analysis and partial differential equations [3, 39]:

- (i) For a function  $f(t, x)$  which depends on  $t$  and  $x$ , we simply denote  $f(t) = f(t, \cdot)$ .
- (ii) Denote  $\langle k \rangle = (1 + |k|^2)^{\frac{1}{2}}$  for  $k \in \mathbb{Z}$ .
- (iii) The notation  $a+$  stands for  $a + \epsilon$  with an arbitrary small  $\epsilon > 0$ , and  $a-$  stands for  $a - \epsilon$ .
- (iv) Denote  $\gamma_p = \frac{3}{2} + \frac{1}{p}$  for some given  $p \geq 1$ .
- (v) For a sequence  $\{a_N\}_{N \in \mathbb{N}}$ , denote the discrete norm as  $\|a_N\|_{l_N^p} = (\sum_{N=1}^{\infty} |a_N|^p)^{\frac{1}{p}}$ , and sometimes we will omit the subscript  $N$  in  $l_N^p$  for brevity.
- (vi) Denote by  $C$  a generic positive constant which may have different values at different occurrences, possibly depending on the norms of the solution and  $T$  but independent of the step size  $\tau$  and time level  $n$  in numerical analysis.
- (vii) Denote by  $A \lesssim B$  or  $B \gtrsim A$  the statement “ $A \leq CB$  for some constant  $C > 0$ ”.

(viii) Denote by  $A \sim B$  the statement “ $C^{-1}B \leq A \leq CB$  for some constant  $C > 0$ ”. Namely,  $A \sim B$  is equivalently to  $A \lesssim B \lesssim A$ . Moreover, denote by  $A \ll B$  or  $B \gg A$  the statement  $A \leq C^{-1}B$  for some sufficiently large constant  $C$ .

With the notations above, we often decompose a subset  $E \subset \mathbb{Z}^2 = \{(k_1, k_2) : k_1, k_2 \in \mathbb{Z}\}$  into two parts, i.e.,  $E = E_1 \cup E_2$ , with

$$E_1 = \{(k_1, k_2) \in E : |k_1| \ll |k_2|\} \quad \text{and} \quad E_2 = \{(k_1, k_2) \in E : |k_1| \gtrsim |k_2|\}.$$

This means that we consider the decomposition with

$$E_1 = \{(k_1, k_2) \in E : |k_1| < c|k_2|\} \quad \text{and} \quad E_2 = \{(k_1, k_2) \in E : |k_1| \geq c|k_2|\},$$

where  $c > 0$  is some sufficiently small constant (independent of  $\tau$  and  $n$ ) which can satisfy the requirement in our analysis.

**2.2. Fourier transform.** The inner product and the norm of  $L^2(\mathbb{T})$  are defined by

$$\langle f, g \rangle \triangleq \operatorname{Re} \int_{\mathbb{T}} f(x) \overline{g(x)} dx \quad \text{and} \quad \|f\|_{L^2(\mathbb{T})} \triangleq \sqrt{\langle f, f \rangle}.$$

The Fourier transform of a function  $f \in L^2(\mathbb{T})$  is defined by

$$\mathcal{F}_k[f] \triangleq \frac{1}{2\pi} \int_{\mathbb{T}} e^{-ikx} f(x) dx, \quad k \in \mathbb{Z}.$$

For the simplicity of notation, we also denote  $\hat{f}_k = \mathcal{F}_k[f]$  and  $f = \mathcal{F}_k^{-1}[\hat{f}_k]$ . The following standard properties of the Fourier transform are well known:

(Fourier series expansion)	$f(x) = \sum_{k \in \mathbb{Z}} \hat{f}_k e^{ikx};$
(Plancherel's identity)	$\ f\ _{L^2(\mathbb{T})} = \sqrt{2\pi} \left( \sum_{k \in \mathbb{Z}}  \hat{f}_k ^2 \right)^{\frac{1}{2}};$
(Parseval's identity)	$\langle f, g \rangle = 2\pi \operatorname{Re} \sum_{k \in \mathbb{Z}} \hat{f}_k \overline{\hat{g}_k};$
(Conversion of products to convolutions)	$\mathcal{F}_k[fg] = \sum_{k_1+k_2=k} \hat{f}_{k_1} \hat{g}_{k_2}.$

The Sobolev space  $H^s(\mathbb{T})$  with some  $s \in \mathbb{R}$ , consists of generalized functions  $f = \sum_{k \in \mathbb{Z}} \hat{f}_k e^{ikx}$  such that  $\|f\|_{H^s} < \infty$ , where

$$\|f\|_{H^s} \triangleq \sqrt{2\pi} \left( \sum_{k \in \mathbb{Z}} \langle k \rangle^{2s} |\hat{f}_k|^2 \right)^{\frac{1}{2}}.$$

The operator  $J^s = (1 - \partial_x^2)^{\frac{s}{2}} : H^{s_0}(\mathbb{T}) \rightarrow H^{s_0-s}(\mathbb{T})$  with  $s_0, s \in \mathbb{R}$ , is defined as

$$J^s f \triangleq \sum_{k \in \mathbb{Z}} \langle k \rangle^s \hat{f}_k e^{ikx}, \quad \forall f \in H^{s_0}(\mathbb{T}),$$

and so we have  $\|f\|_{H^s(\mathbb{T})} = \|J^s f\|_{L^2(\mathbb{T})}$ .

We often use the abbreviations  $H^s(\mathbb{T}) = H^s$ ,  $L^p(\mathbb{T}) = L^p$  and  $l_k^p = l_{k \in \mathbb{Z}}^p$ . Moreover, we use the similar abbreviations for the spacetime norms, like  $L_t^q H_x^s(I \times \mathbb{T}) = L_t^q H_x^s(I)$ .

**2.3. Some operators.** Some frequently used operators are defined as follows.

- (i) Projection: for any real number  $N \geq 0$ , the Littlewood–Paley projections  $P_{\leq N} : H^s(\mathbb{T}) \rightarrow H^s(\mathbb{T})$  and  $P_{>N} : H^s(\mathbb{T}) \rightarrow H^s(\mathbb{T})$  are defined as

$$P_{\leq N} f \triangleq \mathcal{F}_k^{-1}(1_{|k| \leq N} \mathcal{F}_k[f]) = \sum_{|k| \leq N} \hat{f}_k e^{ikx}, \quad P_{>N} f \triangleq \mathcal{F}_k^{-1}(1_{|k| > N} \mathcal{F}_k[f]) = \sum_{|k| > N} \hat{f}_k e^{ikx}.$$

- (ii) The inversion of differentiation:  $\partial_x^{-1} : H^s(\mathbb{T}) \rightarrow H^{s+1}(\mathbb{T})$  is defined by

$$\mathcal{F}_k[\partial_x^{-1} f] \triangleq \begin{cases} (ik)^{-1} \hat{f}_k, & \text{for } k \neq 0, \\ 0, & \text{for } k = 0. \end{cases}$$

- (iii) Average: define the *average* of a time-dependent function  $f(t)$  in the interval  $[0, \tau]$  by

$$\mathcal{M}_\tau(f) \triangleq \frac{1}{\tau} \int_0^\tau f(t) dt. \quad (2.1)$$

**2.4. Some tool lemmas.**

**Lemma 2.1** ([30]). *Let  $\alpha, \beta \in \mathbb{R}$ . If  $\alpha, \beta \neq 0$  and  $s \in [0, \tau]$ , then*

$$|\mathcal{M}_\tau(e^{is(\alpha+\beta)}) - \mathcal{M}_\tau(e^{is\alpha})\mathcal{M}_\tau(e^{is\beta})| \lesssim \min \left\{ \left| \frac{\alpha}{\beta} \right|, \left| \frac{\beta}{\alpha} \right|, \tau|\alpha|, \tau|\beta| \right\}.$$

*If  $\alpha + \beta \neq 0$ , then*

$$|\mathcal{M}_\tau(e^{is(\alpha+\beta)}) - \mathcal{M}_\tau(e^{is\alpha})\mathcal{M}_\tau(e^{is\beta})| \lesssim \tau^{-1}|\alpha + \beta|^{-1}.$$

**Lemma 2.2** (Schur's test). *For any  $a > 0$ , let sequences  $\{a_N\}, \{b_N\} \in l_{N \in \mathbb{Z}^N}^2$ , then we have*

$$\sum_{N_1 \leq N} \left( \frac{N_1}{N} \right)^a a_N b_{N_1} \lesssim \|a_N\|_{l_N^2} \|b_N\|_{l_N^2}.$$

**Lemma 2.3** ([5]). *Consider a quantitatively well posed abstract equation in spaces  $D$  and  $S$ ,*

$$u = L(f) + N_k(u, \dots, u),$$

*which means for all  $f \in D$ ,  $u_1, \dots, u_k \in S$  and some constant  $C > 0$ ,*

$$\|L(f)\|_S \leq C\|f\|_D, \quad \|N_k(u_1, \dots, u_k)\|_S \leq C\|u_1\|_S \dots \|u_k\|_S.$$

*Here  $(D, \|\cdot\|_D)$  is a Banach space of initial data and  $(S, \|\cdot\|_S)$  is a Banach space of spacetime functions. Define*

$$A_1(f) \triangleq L(f), \quad A_n(f) \triangleq \sum_{n_1, \dots, n_k \geq 1, n_1 + \dots + n_k = n} N_k(A_{n_1}(f), \dots, A_{n_k}(f)), \quad n > 1.$$

*Then for some  $C_1 > 0$ , all  $f, g \in D$  and all  $n \geq 1$ ,*

$$\|A_n(f) - A_n(g)\|_S \leq C_1^n (\|f\|_D + \|g\|_D)^{n-1} \|f - g\|_D.$$

### 3. PROOF FOR WELL-POSEDNESS THEORY

In this section, we shall give the proofs for the two main well-posedness results, i.e., [Theorem 1.1](#) and [Theorem 1.2](#). Without loss of generality, we assume  $\lambda = -1$  in the NLS model [\(1.1\)](#) for simplicity of notations.

**3.1. Proof of Theorem 1.1.** Let  $T$  be a fixed time with  $0 < T < +\infty$  which will be determined later. By Duhamel's formula, we have that for  $0 \leq t \leq T$ ,

$$u(t) = e^{i(t-t_0)\partial_x^2} u(t_0) + i \int_{t_0}^t e^{i(t-\rho)\partial_x^2} \left( \xi u(\rho) + |u(\rho)|^2 u(\rho) \right) d\rho.$$

Denote  $v(t) = e^{-it\partial_x^2} u(t)$  and set  $t_0 = 0$ , then it reduces to

$$v(t) = u_0 + i \int_0^t e^{-i\rho\partial_x^2} \left( \xi u(\rho) + |u(\rho)|^2 u(\rho) \right) d\rho. \quad (3.1)$$

We denote the operator  $\Phi$  by

$$\Phi(v) \triangleq u_0 + i \int_0^t e^{-i\rho\partial_x^2} \left( \xi u(\rho) + |u(\rho)|^2 u(\rho) \right) d\rho, \quad (3.2)$$

and

$$R_0 \triangleq \max\{2\|u_0\|_{H_x^{s+\gamma}}, 1\}.$$

Then we aim to show that for any  $v \in L_t^\infty H_x^{s+\gamma}((0, T) \times \mathbb{T})$  with

$$\|v\|_{L_t^\infty H_x^{s+\gamma}} \leq R_0, \quad (3.3)$$

it holds that

$$\|\Phi(v)\|_{L_t^\infty H_x^{s+\gamma}} \leq R_0. \quad (3.4)$$

Here and below, we use the abbreviation for the norms  $L_t^\infty H_x^{s+\gamma} = L_t^\infty H_x^{s+\gamma}([0, T] \times \mathbb{T})$ . Moreover, for any  $v_1, v_2 \in L_t^\infty H_x^{s+\gamma}((0, T) \times \mathbb{T})$  with

$$\|v_j\|_{L_t^\infty H_x^{s+\gamma}} \leq R_0, \quad j = 1, 2,$$

it holds that

$$\|\Phi(v_1) - \Phi(v_2)\|_{L_t^\infty H_x^{s+\gamma}} \leq \theta \|v_1 - v_2\|_{L_t^\infty H_x^{s+\gamma}}, \quad \text{for some } \theta \in (0, 1). \quad (3.5)$$

The proof of (3.5) is similar as the proof of (3.4), we only consider the latter. Then it can be reduced to the following lemma.

**Lemma 3.1.** *Let  $s \geq 0, 2 < p \leq +\infty, \gamma = \gamma_p -$  and denote  $\varepsilon_0 = \min\{\frac{1}{2}(\gamma_p - \gamma), \frac{1}{8}\}$ . Then for any  $v$  satisfying (3.3), and any  $N_0 > 0$ , there exists a function  $C_0(N_0) > 0$  which is dependent on  $\|\xi\|_{\dot{b}^{s,p}}$  such that*

$$\|\Phi(v)\|_{L_t^\infty H_x^{s+\gamma}} \leq \frac{1}{2}R_0 + \left[ C_0(N_0)T + N_0^{-\frac{\varepsilon_0}{2}} \right] (R_0 + R_0^3).$$

The proof of Lemma 3.1 will be split into several parts. Firstly, we consider the term

$$\Phi^1(u) \triangleq i \int_0^t e^{-i\rho\partial_x^2} \left( \xi u(\rho) \right) d\rho,$$

and we have the following estimate for it.

**Lemma 3.2.** *Under the assumption of Lemma 3.1,*

$$\|\Phi^1(v)\|_{L_t^\infty H_x^{s+\gamma}} \leq \left[ C_0(N_0)T + N_0^{-\frac{\varepsilon_0}{2}} \right] (R_0 + R_0^3).$$

*Proof.* Taking Fourier transform for  $\Phi^1(v)$ , we have that for any  $k \in \mathbb{Z}$ , its Fourier coefficients denoted by  $I_k$ , equal to

$$I_k \triangleq i \int_0^t \sum_{k_1+k_2=k} e^{i\rho\phi(k,k_2)} \hat{\xi}_{k_1} \hat{v}_{k_2}(\rho) d\rho.$$

Here we denote the phase functions  $\phi$  by

$$\phi(k, k_j) \triangleq |k|^2 - |k_j|^2.$$

Then

$$\left\| i \int_0^t e^{-i\rho\partial_x^2} (\xi u(\rho)) d\rho \right\|_{H^{s+\gamma}} = \|\langle k \rangle^{s+\gamma} I_k\|_{l_k^2}.$$

Note that  $s + \gamma > \frac{1}{2}$ , then by Cauchy-Schwartz's inequality, we have

$$\begin{aligned} \|\langle k \rangle^{s+\gamma} I_k\|_{l_k^2(\{|k| \leq N_0\})} &\lesssim N_0^{s+\gamma} \|\hat{\xi}_k\|_{l_k^\infty} \int_0^t \sum_{k_2} |v_{k_2}| d\rho \\ &\lesssim t N_0^{s+\gamma} \|\hat{\xi}_k\|_{l_k^\infty} \|v\|_{L_t^\infty H_x^{s+\gamma}}. \end{aligned} \quad (3.6)$$

Hence, we only consider the piece  $\|\langle k \rangle^{s+\gamma} I_k\|_{l_k^2(\{|k| > N_0\})}$ . Now we split  $I$  into two parts as

$$I_k = I_{1,k} + I_{2,k},$$

where

$$I_{1,k} \triangleq i \int_0^t \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2)=0}} \hat{\xi}_{k_1} \hat{v}_{k_2}(\rho) d\rho; \quad I_{2,k} \triangleq i \int_0^t \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2) \neq 0}} e^{i\rho\phi(k,k_2)} \hat{\xi}_{k_1} \hat{v}_{k_2}(\rho) d\rho.$$

We first estimate  $I_{1,k}$ . Note that  $\phi(k, k_2) = 0$  is equivalent to  $k = k_2$  or  $k = -k_2$ , and so

$$I_{1,k} = i \int_0^t \hat{\xi}_0 \hat{v}_k(\rho) d\rho + i \int_0^t \hat{\xi}_{2k} \hat{v}_{-k}(\rho) d\rho.$$

Therefore,

$$\begin{aligned} \|\langle k \rangle^{s+\gamma} I_{1,k}\|_{l_k^2} &\lesssim \int_0^t \|\langle k \rangle^{s+\gamma} \hat{\xi}_0 \hat{v}_k(\rho)\|_{l_k^2} d\rho + \int_0^t \|\langle k \rangle^{s+\gamma} \hat{\xi}_{2k} \hat{v}_{-k}(\rho)\|_{l_k^2} d\rho \\ &\lesssim t \|\hat{\xi}_k\|_{l_k^\infty} \|\langle k \rangle^{s+\gamma} \hat{v}_k\|_{L_t^\infty l_k^2} \lesssim t \|\hat{\xi}_k\|_{l_k^\infty} \|v\|_{L_t^\infty H_x^{s+\gamma}}. \end{aligned}$$

Now we start to estimate  $I_{2,k}$ . By integration-by-parts and noting that

$$\partial_t v(t) = e^{-it\partial_x^2} (\xi u + |u|^2 u),$$

we have

$$\begin{aligned} I_{2,k} &= i \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2) \neq 0}} e^{i\rho\phi(k,k_2)} \frac{\hat{\xi}_{k_1} \hat{v}_{k_2}(\rho)}{i\phi(k, k_2)} \Big|_0^t - i \int_0^t \sum_{\substack{k_1+k_2+k_3=k \\ \phi(k,k_2+k_3) \neq 0}} \frac{e^{i\rho\phi(k,k_3)}}{i\phi(k, k_2+k_3)} \hat{\xi}_{k_1} \hat{\xi}_{k_2} \hat{v}_{k_3}(\rho) d\rho \\ &\quad - i \int_0^t \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2) \neq 0}} e^{i\rho k^2} \frac{1}{i\phi(k, k_2)} \hat{\xi}_{k_1} (\widehat{|u|^2 u})_{k_2} d\rho \\ &\triangleq I_{21,k} + I_{22,k} + I_{23,k}. \end{aligned}$$

3.1.1. *Estimates on  $I_{21,k}$ .* For this term, by using the dual and Parseval's identity, we obtain

$$\begin{aligned}
\| \langle k \rangle^{s+\gamma} I_{21,k} \|_{l^2(|k| \geq N_0)} &= \sup_{h: \|h\|_{l^2} = 1} \langle \langle k \rangle^{s+\gamma} I_{21,k}, h_k \rangle \\
&= 2\pi \sup_{h: \|h\|_{l^2} = 1} \operatorname{Re} \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2) \neq 0, |k| \geq N_0}} \langle k \rangle^{s+\gamma} e^{\rho\phi(k,k_2)} \frac{\hat{\xi}_{k_1} \hat{v}_{k_2}(\rho)}{i\phi(k,k_2)} \Big|_0^t \overline{h_k} \\
&\leq 2\pi \sup_{h: \|h\|_{l^2} = 1} \sup_{t \in [0, T]} \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2) \neq 0, |k| \geq N_0, |k_1| \lesssim |k_2|}} \langle k \rangle^{s+\gamma} \frac{|\hat{\xi}_{k_1}| |\hat{v}_{k_2}(\rho)|}{|\phi(k,k_2)|} |h_k| \\
&\quad + 2\pi \sup_{h: \|h\|_{l^2} = 1} \sup_{t \in [0, T]} \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2) \neq 0, |k| \geq N_0, |k_1| \gg |k_2|}} \langle k \rangle^{s+\gamma} \frac{|\hat{\xi}_{k_1}| |\hat{v}_{k_2}(\rho)|}{|\phi(k,k_2)|} |h_k| \\
&\triangleq I_{211} + I_{212}. \tag{3.7}
\end{aligned}$$

For short, we omit  $\sup_{h: \|h\|_{l^2} = 1} \sup_{t \in [0, T]}$  in the front, and turn to check the estimate hold for any  $h$  such that  $\|h\|_{l^2} = 1$  and any  $t \in [0, T]$  in the proof.

• **Estimates on  $I_{211}$ .** Note that  $|\phi(k, k_2)| = (k - k_2)(k + k_2)$ , and  $\phi \neq 0$  implies  $|k \pm k_2| \geq 1$ . Thus,

$$|\phi(k, k_2)| = |k - k_2| |k + k_2| \geq |k| \geq N_0.$$

Then, combining with the estimate

$$|\phi(k, k_2)| = |k_1(k_1 + 2k_2)| \sim \langle k_1 \rangle \langle k_1 + 2k_2 \rangle,$$

we have

$$\begin{aligned}
I_{211} &\lesssim \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2) \neq 0, |k| \geq N_0, |k_1| \lesssim |k_2|}} \langle k_2 \rangle^{s+\gamma} \frac{1}{|\phi(k, k_2)|} |\hat{\xi}_{k_1}| |\hat{v}_{k_2}| |h_k| \\
&\lesssim N_0^{-\varepsilon_0} \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2) \neq 0, |k| \geq N_0, |k_1| \lesssim |k_2|}} \langle k_2 \rangle^{s+\gamma} \frac{1}{|\phi(k, k_2)|^{1-\varepsilon_0}} |\hat{\xi}_{k_1}| |\hat{v}_{k_2}| |h_k| \\
&\lesssim N_0^{-\varepsilon_0} \sum_{k_2} \sum_{k_1} \frac{1}{\langle k_1 \rangle^{1-\varepsilon_0}} \frac{1}{\langle k_1 + 2k_2 \rangle^{1-\varepsilon_0}} |\hat{\xi}_{k_1}| \langle k_2 \rangle^{s+\gamma} |\hat{v}_{k_2}| |h_{k_1+k_2}| \\
&\lesssim N_0^{-\varepsilon_0} \|\hat{\xi}_k\|_{l_k^\infty} \sum_{k_2} \sum_{k_1} \left( \frac{1}{\langle k_1 \rangle^{2-2\varepsilon_0}} + \frac{1}{\langle k_1 + 2k_2 \rangle^{2-2\varepsilon_0}} \right) \langle k_2 \rangle^{s+\gamma} |\hat{v}_{k_2}| |h_{k_1+k_2}| \\
&\lesssim N_0^{-\varepsilon_0} \|\hat{\xi}_k\|_{l_k^\infty} \sum_{k_1} \frac{1}{\langle k_1 \rangle^{2-2\varepsilon_0}} \sum_{k_2} \langle k_2 \rangle^{s+\gamma} |\hat{v}_{k_2}| |h_{k_1+k_2}| \\
&\quad + N_0^{-\varepsilon_0} \|\hat{\xi}_k\|_{l_k^\infty} \sum_{\tilde{k}_1} \frac{1}{\langle \tilde{k}_1 \rangle^{2-2\varepsilon_0}} \sum_{k_2} \langle k_2 \rangle^{s+\gamma} |\hat{v}_{k_2}| |h_{\tilde{k}_1-k_2}|,
\end{aligned}$$

where in the last inequality, we have used the change of the variable  $\tilde{k}_1 = k_1 + 2k_2$ . By the Cauchy-Schwartz inequality, we obtain that

$$|I_{211}| \lesssim N_0^{-\varepsilon_0} \|\hat{\xi}_k\|_{l_k^\infty} \|v\|_{H^{s+\gamma}} \|h_k\|_{l_k^2} \lesssim N_0^{-\varepsilon_0} \|\xi\|_{\hat{b}_{s,p}} \|v\|_{H^{s+\gamma}}. \tag{3.8}$$

• **Estimates on  $I_{212}$ .** When  $|k_1| \gg |k_2|$ , we have

$$|\phi(k, k_2)| \gtrsim |k_1|^2.$$

Then by the Cauchy-Schwartz inequality, we find

$$\begin{aligned} I_{212} &\lesssim \sum_{\substack{k_1+k_2=k \\ \phi(k, k_2) \neq 0, |k| \geq N_0, |k_1| \gg |k_2|}} \langle k_1 \rangle^{\gamma+s-2} |\hat{\xi}_{k_1}| |\hat{v}_{k_2}| |h_{k_1+k_2}| \\ &\lesssim \sum_{k_2} \left( \sum_{k_1} (\langle k_1 \rangle^s |\hat{\xi}_{k_1}|)^p \right)^{\frac{1}{p}} \left( \sum_{k_1: |k_1| \gtrsim N_0} (\langle k_1 \rangle^{\gamma-2} |h_{k_1+k_2}|)^{p'} \right)^{\frac{1}{p'}} |\hat{v}_{k_2}|. \end{aligned}$$

Then by further using the Cauchy-Schwartz inequality, we get

$$I_{212} \lesssim \|\xi\|_{\hat{b}^{s,p}} \|v\|_{H^{s+\gamma}} \|h\|_{L^2} \|\langle k_1 \rangle^{\gamma-2}\|_{L^r(|k_1| \gtrsim N_0)} \sum_{k_2} |\hat{v}_{k_2}|,$$

where  $r$  satisfies that  $\frac{1}{r} = \frac{1}{2} - \frac{1}{p}$  and so  $(\gamma - 2)r < -1$ . Moreover,  $r + s > \frac{1}{2}$  and so we get

$$I_{212} \lesssim N_0^{-\varepsilon_0} \|\xi\|_{\hat{b}^{s,p}} \|v\|_{L_t^\infty H_x^{s+\gamma}}. \quad (3.9)$$

Collecting the estimates (3.7)-(3.9), we obtain

$$\|\langle k \rangle^{s+\gamma} I_{21,k}\|_{L^2(|k| \geq N_0)} \lesssim N_0^{-\varepsilon_0} \|\xi\|_{\hat{b}^{s,p}} \|v\|_{H^{s+\gamma}}.$$

3.1.2. *Estimates on  $I_{22,k}$ .* For this term, we further split it into the following two parts as

$$\begin{aligned} I_{22,k} &= -i \int_0^t \sum_{\substack{k_1+k_2+k_3=k \\ \phi(k, k_2+k_3) \neq 0, \phi(k, k_3)=0}} \frac{e^{i\rho\phi(k, k_3)}}{i\phi(k, k_2+k_3)} \hat{\xi}_{k_1} \hat{\xi}_{k_2} \hat{v}_{k_3}(\rho) d\rho \\ &\quad - i \int_0^t \sum_{\substack{k_1+k_2+k_3=k \\ \phi(k, k_2+k_3) \neq 0, \phi(k, k_3) \neq 0}} \frac{e^{i\rho\phi(k, k_3)}}{i\phi(k, k_2+k_3)} \hat{\xi}_{k_1} \hat{\xi}_{k_2} \hat{v}_{k_3}(\rho) d\rho \\ &\triangleq I_{221,k} + I_{222,k}. \end{aligned}$$

• **Estimates on  $I_{221,k}$ .** Note that  $\phi(k, k_3) = 0$  implies that  $k = k_3$  or  $k = -k_3$ . Therefore,

$$\begin{aligned} I_{221,k} &= -i \int_0^t \sum_{k_1: \phi(k, -k_1+k) \neq 0} \frac{1}{i\phi(k, -k_1+k)} \hat{\xi}_{k_1} \hat{\xi}_{-k_1} \hat{v}_k(\rho) d\rho \\ &\quad - i \int_0^t \sum_{\substack{k_1+k_2=2k \\ \phi(k, k_2-k) \neq 0}} \frac{1}{i\phi(k, k_2-k)} \hat{\xi}_{k_1} \hat{\xi}_{k_2} \hat{v}_{-k}(\rho) d\rho \\ &\triangleq I_{2211,k} + I_{2212,k}. \end{aligned}$$

1) On  $I_{2211,k}$ . Since  $\phi(k, -k_1+k) = k_1(2k - k_1)$ , it can be rewritten as

$$I_{2211,k} = -i \int_0^t \sum_{\substack{k_1 \neq 0 \\ 2k - k_1 \neq 0}} \frac{1}{ik_1(2k - k_1)} \hat{\xi}_{k_1} \hat{\xi}_{-k_1} \hat{v}_k(\rho) d\rho.$$

Therefore, we have that

$$\begin{aligned}
 \|\langle k \rangle^{s+\gamma} I_{2211,k}\|_{l_k^2} &= \left\| \langle k \rangle^{s+\gamma} \int_0^t \sum_{\substack{k_1 \neq 0 \\ 2k-k_1 \neq 0}} \frac{1}{|k_1||2k-k_1|} |\hat{\xi}_{k_1}| |\hat{\xi}_{-k_1}| |\hat{v}_k(\rho)| d\rho \right\|_{l_k^2} \\
 &\lesssim \|\hat{\xi}_k\|_{l_k^\infty}^2 \left\| \langle k \rangle^{s+\gamma} \int_0^t \sum_{\substack{k_1 \neq 0 \\ 2k-k_1 \neq 0}} \left( \frac{1}{|k_1|^2} + \frac{1}{|2k-k_1|^2} \right) |\hat{v}_k(\rho)| d\rho \right\|_{l_k^2} \\
 &\lesssim \|\hat{\xi}_k\|_{l_k^\infty}^2 \int_0^t \|\langle k \rangle^{s+\gamma} |\hat{v}_k(\rho)|\|_{l_k^2} d\rho \lesssim t \|\hat{\xi}_k\|_{l_k^\infty}^2 \|v\|_{L_t^\infty H_x^{s+\gamma}}.
 \end{aligned}$$

2) On  $I_{2212,k}$ . Since  $\phi(k, k_2 - k) = k_1 k_2$ , it can be rewritten as

$$I_{2212,k} = -i \int_0^t \sum_{\substack{k_1+k_2=2k \\ k_1 \neq 0, k_2 \neq 0}} \frac{1}{ik_1 k_2} \hat{\xi}_{k_1} \hat{\xi}_{k_2} \hat{v}_{-k}(\rho) d\rho.$$

Therefore, we have that

$$\begin{aligned}
 \|\langle k \rangle^{s+\gamma} I_{2212,k}\|_{l_k^2} &= \left\| \langle k \rangle^{s+\gamma} \int_0^t \sum_{\substack{k_1+k_2=2k \\ k_1 \neq 0, k_2 \neq 0}} \frac{1}{|k_1||k_2|} \hat{\xi}_{k_1} \hat{\xi}_{k_2} \hat{v}_{-k}(\rho) d\rho \right\|_{l_k^2} \\
 &\lesssim \|\hat{\xi}_k\|_{l_k^\infty}^2 \left\| \langle k \rangle^{s+\gamma} \int_0^t \sum_{\substack{k_1 \neq 0 \\ 2k-k_1 \neq 0}} \left( \frac{1}{|k_1|^2} + \frac{1}{|2k-k_1|^2} \right) |\hat{v}_{-k}(\rho)| d\rho \right\|_{l_k^2} \\
 &\lesssim t \|\hat{\xi}_k\|_{l_k^\infty}^2 \|v\|_{L_t^\infty H_x^{s+\gamma}}.
 \end{aligned}$$

• **Estimates on  $I_{222,k}$ .** By integration-by-parts, we obtain that

$$\begin{aligned}
 &I_{222,k} \\
 = &-i \sum_{\substack{k_1+k_2+k_3=k \\ \phi(k, k_2+k_3) \neq 0, \phi(k, k_3) \neq 0}} \frac{e^{i\rho\phi(k, k_3)}}{i\phi(k, k_2+k_3) \cdot i\phi(k, k_3)} \hat{\xi}_{k_1} \hat{\xi}_{k_2} \hat{v}_{k_3}(\rho) \Big|_0^t \\
 &+ i \int_0^t \sum_{\substack{k_1+k_2+k_3+k_4=k \\ \phi(k, k_2+k_3+k_4) \neq 0, \phi(k, k_3+k_4) \neq 0}} \frac{e^{i\rho\phi(k, k_4)}}{i\phi(k, k_2+k_3+k_4) \cdot i\phi(k, k_3+k_4)} \hat{\xi}_{k_1} \hat{\xi}_{k_2} \hat{\xi}_{k_3} \hat{v}_{k_4}(\rho) d\rho \\
 &+ i \int_0^t \sum_{\substack{k_1+k_2+k_3=k \\ \phi(k, k_2+k_3) \neq 0, \phi(k, k_3) \neq 0}} \frac{e^{i\rho k^2}}{i\phi(k, k_2+k_3) \cdot i\phi(k, k_3)} \hat{\xi}_{k_1} \hat{\xi}_{k_2} (\widehat{|u|^2 u})_{k_3} d\rho \\
 \triangleq &I_{2221,k} + I_{2222,k} + I_{2223,k}.
 \end{aligned}$$

1) On  $I_{2221,k}$ . Note that

$$\phi(k, k_2+k_3) = k_1(2k-k_1); \quad \phi(k, k_3) = (k-k_3)(k+k_3).$$

Moreover, we use the formula that when  $\phi(k, j) \neq 0$ ,

$$\frac{1}{\phi(k, j)} = \frac{1}{2k} \left( \frac{1}{k-j} + \frac{1}{k+j} \right)$$

and obtain that

$$\frac{1}{\phi(k, k_3)} = \frac{1}{2k} \left[ \frac{1}{k - k_3} + \frac{1}{k + k_3} \right], \quad \frac{1}{\phi(k, k_2 + k_3)} = \frac{1}{2k} \left[ \frac{1}{k - (k_2 + k_3)} + \frac{1}{k + k_2 + k_3} \right].$$

Then by the Cauchy-Schwartz inequality, we obtain that when  $\phi(k, k_2 + k_3) \neq 0$ ,

$$\begin{aligned} \left| \frac{1}{\phi(k, k_2 + k_3)} \right| &= \frac{1}{|k_1|^{\frac{1}{2}\varepsilon_0} |2k - k_1|^{\frac{1}{2}\varepsilon_0} |k|^{1-\frac{1}{2}\varepsilon_0}} \left( \frac{1}{|k_1|^{1-\frac{1}{2}\varepsilon_0}} + \frac{1}{|2k - k_1|^{1-\frac{1}{2}\varepsilon_0}} \right) \\ &\lesssim \langle k \rangle^{-1+\frac{1}{2}\varepsilon_0} \left( \frac{1}{\langle k_1 \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle 2k - k_1 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right); \end{aligned} \quad (3.10)$$

and similarly,

$$\left| \frac{1}{\phi(k, k_3)} \right| \lesssim \langle k \rangle^{-1+\frac{1}{2}\varepsilon_0} \left( \frac{1}{\langle k - k_3 \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right). \quad (3.11)$$

Together with (3.10) and (3.11), and using  $k_1 = k - k_2 - k_3$  it gives that

$$\begin{aligned} \frac{1}{|\phi(k, k_2 + k_3)| |\phi(k, k_3)|} &\lesssim \langle k \rangle^{-2+\varepsilon_0} \left[ \frac{1}{\langle k - k_3 \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right] \\ &\quad \cdot \left[ \frac{1}{\langle k - (k_2 + k_3) \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_2 + k_3 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right]. \end{aligned} \quad (3.12)$$

Now we are ready to estimate  $I_{2221, k}$ . By duality, we have that

$$\begin{aligned} \left\| \langle k \rangle^{s+\gamma} |I_{2221, k}| \right\|_{l_k^2(|k| \geq N_0)} &= \sup_{h: \|h\|_{l^2} = 1} \langle \langle k \rangle^{s+\gamma} |I_{2221, k}|, h_k \rangle \\ &= \sup_{h: \|h\|_{l^2} = 1} \sup_{t \in [0, T]} \sum_{\substack{k_1+k_2+k_3=k \\ \phi(k, k_2+k_3) \neq 0, \phi(k, k_3) \neq 0}} \frac{\langle k \rangle^{s+\gamma}}{|\phi(k, k_2 + k_3)| |\phi(k, k_3)|} \\ &\quad \cdot |\hat{\xi}_{k_1}| |\hat{\xi}_{k_2}| |\hat{v}_{k_3}| |h_k|. \end{aligned}$$

As before, we omit  $\sup_{h: \|h\|_{l^2} = 1} \sup_{t \in [0, T]}$  in the front. Then by (3.12), the trivial inequality  $\langle k \rangle^s \leq$

$\langle k_1 \rangle^s \langle k_2 \rangle^s \langle k_3 \rangle^s$ , and the Cauchy-Schwartz inequality, we obtain

$$\begin{aligned} &\left\| \langle k \rangle^{s+\gamma} |I_{2221, k}| \right\|_{l_k^2(|k| \geq N_0)} \\ &\lesssim \sum_{\substack{k_1+k_2+k_3=k, |k| \geq N_0 \\ \phi(k, k_2+k_3) \neq 0, \phi(k, k_3) \neq 0}} \frac{\langle k \rangle^\gamma}{|\phi(k, k_2 + k_3)| |\phi(k, k_3)|} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle k_2 \rangle^s |\hat{\xi}_{k_2}| \langle k_3 \rangle^s |\hat{v}_{k_3}| |h_k| \\ &\lesssim \sum_{\substack{k_1+k_2+k_3=k, |k| \geq N_0 \\ \phi(k, k_2+k_3) \neq 0, \phi(k, k_3) \neq 0}} \langle k \rangle^{-2+\varepsilon_0+\gamma} \left[ \frac{1}{\langle k - k_3 \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right] \\ &\quad \cdot \left[ \frac{1}{\langle k - (k_2 + k_3) \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_2 + k_3 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right] \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle k_2 \rangle^s |\hat{\xi}_{k_2}| \langle k_3 \rangle^s |\hat{v}_{k_3}| |h_k|. \end{aligned}$$

Denote

$$\zeta_{k_3, k}^1 = \langle k \rangle^{-2+\varepsilon_0+\gamma} \left[ \frac{1}{\langle k - k_3 \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right] |h_k|$$

and

$$\zeta_{k_3, k}^2 = \sum_{k_2} \left[ \frac{1}{\langle k - (k_2 + k_3) \rangle^{1 + \frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_2 + k_3 \rangle^{1 + \frac{1}{2}\varepsilon_0}} \right] \langle k - k_2 - k_3 \rangle^s |\hat{\xi}_{k-k_2-k_3}| \langle k_2 \rangle^s |\hat{\xi}_{k_2}|.$$

Then we have that

$$\begin{aligned} \left\| \langle k \rangle^{s+\gamma} |I_{2221, k}| \right\|_{l_k^2(|k| \geq N_0)} &\lesssim \sum_{\substack{k, k_3 \\ |k| \geq N_0}} \zeta_{k_3, k}^1 \zeta_{k_3, k}^2 \langle k_3 \rangle^s |\hat{v}_{k_3}| \\ &\lesssim \sum_{k_3} \langle k_3 \rangle^s |\hat{v}_{k_3}| \cdot \left( \sum_{k: |k| \geq N_0} (\zeta_{k_3, k}^1)^{p'} \right)^{\frac{1}{p'}} \cdot \left( \sum_{k: |k| \geq N_0} (\zeta_{k_3, k}^2)^p \right)^{\frac{1}{p}}. \end{aligned}$$

For convenience, we further denote

$$J_{k_3} = \left\| \zeta_{k_3, k}^1 \right\|_{l_k^{p'}(|k| \geq N_0)}, \quad \tilde{J}_{k_3} = \left\| \zeta_{k_3, k}^2 \right\|_{l_k^p(|k| \geq N_0)}.$$

Then it gives that

$$\left\| \langle k \rangle^{s+\gamma} |I_{2221, k}| \right\|_{l_k^2(|k| \geq N_0)} \lesssim \sum_{k_3} \langle k_3 \rangle^s |\hat{v}_{k_3}| J_{k_3} \tilde{J}_{k_3}, \quad (3.13)$$

Therefore, it reduces to estimate  $J_{k_3}$  and  $\tilde{J}_{k_3}$ .

For the term  $J_{k_3}$ ,

$$J_{k_3} \lesssim \|h_k\|_{l_k^2} \left\| \langle k \rangle^{-2+\varepsilon_0+\gamma} \left[ \frac{1}{\langle k - k_3 \rangle^{1 + \frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3 \rangle^{1 + \frac{1}{2}\varepsilon_0}} \right] \right\|_{l_k^r(|k| \geq N_0)} \lesssim N_0^{-\varepsilon_0} \|h\|_{l^2}, \quad (3.14)$$

where we have used the relationship that  $r$  satisfies that  $\frac{1}{r} = \frac{1}{2} - \frac{1}{p}$  and  $(\gamma - 2)r < -1$ .

For the term  $\tilde{J}_{k_3}$ , we change the variable and write

$$\begin{aligned} \tilde{J}_{k_3} &= \left\| \sum_{k_2} \left( \frac{1}{\langle k - (k_2 + k_3) \rangle^{1 + \frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_2 + k_3 \rangle^{1 + \frac{1}{2}\varepsilon_0}} \right) \langle k - k_2 - k_3 \rangle^s |\hat{\xi}_{k-k_2-k_3}| \langle k_2 \rangle^s |\hat{\xi}_{k_2}| \right\|_{l_k^p} \\ &\lesssim \left\| \sum_{k_1} \frac{1}{\langle k_1 \rangle^{1 + \frac{1}{2}\varepsilon_0}} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle k - k_1 - k_3 \rangle^s |\hat{\xi}_{k-k_1-k_3}| \right\|_{l_k^p} \\ &\quad + \left\| \sum_{\tilde{k}_1} \frac{1}{\langle \tilde{k}_1 \rangle^{1 + \frac{1}{2}\varepsilon_0}} \langle 2k - \tilde{k}_1 \rangle^s |\hat{\xi}_{2k-\tilde{k}_1}| \langle \tilde{k}_1 - k - k_3 \rangle^s |\hat{\xi}_{\tilde{k}_1-k-k_3}| \right\|_{l_k^p} \\ &\lesssim \|\langle k \rangle^s \hat{\xi}_k\|_{l_k^p} \|\langle k \rangle^s \hat{\xi}_k\|_{l_k^\infty} \lesssim \|\xi\|_{\dot{b}^{s,p}}^2. \end{aligned} \quad (3.15)$$

Hence, by the estimates (3.13)-(3.15), we have

$$\left\| \langle k \rangle^{s+\gamma} |I_{2221, k}| \right\|_{l_k^2(|k| \geq N_0)} \lesssim N_0^{-\varepsilon_0} \|h\|_{l^2} \|\xi\|_{\dot{b}^{s,p}}^2 \sum_{k_3} \langle k_3 \rangle^s |\hat{v}_{k_3}| \lesssim N_0^{-\varepsilon_0} \|h\|_{l^2} \|\xi\|_{\dot{b}^{s,p}}^2 \|v\|_{H^{s+\gamma}}.$$

2) On  $I_{2222, k}$ . Similarly as (3.12), we have that

$$\begin{aligned} \frac{1}{|\phi(k, k_2 + k_3 + k_4)| |\phi(k, k_3 + k_4)|} &\lesssim \langle k \rangle^{-2+\varepsilon_0} \left[ \frac{1}{\langle k_1 \rangle^{1 + \frac{1}{2}\varepsilon_0}} + \frac{1}{\langle 2k - k_1 \rangle^{1 + \frac{1}{2}\varepsilon_0}} \right] \\ &\quad \cdot \left[ \frac{1}{\langle k - (k_3 + k_4) \rangle^{1 + \frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3 + k_4 \rangle^{1 + \frac{1}{2}\varepsilon_0}} \right]. \end{aligned} \quad (3.16)$$

Then by the duality, (3.16) and Cauchy-Schwartz's inequality, we have that (again, from the second line we omit  $\sup_{h:\|h\|_{l^2}=1}$  in the front)

$$\begin{aligned}
& \left\| \langle k \rangle^{s+\gamma} |I_{2222,k}| \right\|_{l_k^2(|k| \geq N_0)} = \sup_{h:\|h\|_{l^2}=1} \langle \langle k \rangle^{s+\gamma} |I_{2222,k}|, h_k \rangle \\
& \lesssim \sum_{\substack{k_1+k_2+k_3+k_4=k \\ \phi(k, k_2+k_3+k_4) \neq 0, \phi(k, k_3+k_4) \neq 0}} \int_0^t \frac{\langle k \rangle^{s+\gamma} |\hat{\xi}_{k_1}| |\hat{\xi}_{k_2}| |\hat{\xi}_{k_3}| |\hat{v}_{k_4}(\rho)| |h_k|}{\phi(k, k_2+k_3+k_4) \cdot \phi(k, k_3+k_4)} d\rho \\
& \lesssim \sum_{\substack{k_1+k_2+k_3+k_4=k \\ \phi(k, k_2+k_3+k_4) \neq 0, \phi(k, k_3+k_4) \neq 0}} \int_0^t \langle k \rangle^{-2+\varepsilon_0+\gamma} \left[ \frac{1}{\langle k - (k_3+k_4) \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3+k_4 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right] \\
& \quad \cdot \left[ \frac{1}{\langle k_1 \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle 2k - k_1 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right] \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle k_2 \rangle^s |\hat{\xi}_{k_2}| \langle k_3 \rangle^s |\hat{\xi}_{k_3}| \langle k_4 \rangle^s |\hat{v}_{k_4}| |h_k| d\rho,
\end{aligned}$$

where in the last step we have used the inequality  $\langle k \rangle^s \leq \langle k_1 \rangle^s \langle k_2 \rangle^s \langle k_3 \rangle^s$ . Then we further get that

$$\begin{aligned}
& \left\| \langle k \rangle^{s+\gamma} |I_{2222,k}| \right\|_{l_k^2(|k| \geq N_0)} \\
& \lesssim \int_0^t \sum_{k_4} \sum_k \sum_{k_3} \sum_{k_1} \left[ \frac{1}{\langle k_1 \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{|2k - k_1|^{1+\frac{1}{2}\varepsilon_0}} \right] \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle k - k_1 - k_3 - k_4 \rangle^s |\hat{\xi}_{k-k_1-k_3-k_4}| \\
& \quad \cdot \left[ \frac{1}{\langle k - (k_3+k_4) \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3+k_4 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right] \langle k_3 \rangle^s |\hat{\xi}_{k_3}| \langle k_4 \rangle^s |\hat{v}_{k_4}| |h_k| \langle k \rangle^{-2+\varepsilon_0+\gamma} d\rho \\
& \lesssim \left\| \langle k \rangle^s |\hat{\xi}_k| \right\|_{l_k^\infty}^2 \int_0^t \sum_{k_4} \sum_k \sum_{k_3} \left[ \frac{1}{\langle k - (k_3+k_4) \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3+k_4 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right] \\
& \quad \cdot \langle k_3 \rangle^s |\hat{\xi}_{k_3}| \langle k_4 \rangle^s |\hat{v}_{k_4}| |h_k| \langle k \rangle^{-2+\varepsilon_0+\gamma} d\rho.
\end{aligned}$$

We denote

$$M_k = \sum_{k_3} \left[ \frac{1}{\langle k - (k_3+k_4) \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3+k_4 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right] \langle k_3 \rangle^s |\hat{\xi}_{k_3}|,$$

then we have that

$$\left\| \langle k \rangle^{s+\gamma} |I_{2222,k}| \right\|_{l_k^2(|k| \geq N_0)} \lesssim \left\| \langle k \rangle^s |\hat{\xi}_k| \right\|_{l_k^\infty}^2 \int_0^t \sum_{k_4} \sum_k M_k \langle k_4 \rangle^s |\hat{v}_{k_4}| |h_k| \langle k \rangle^{-2+\varepsilon_0+\gamma} d\rho. \quad (3.17)$$

Now, we estimate the term  $M_k$  as follows

$$\begin{aligned} M_k &\lesssim \left\| \left[ \frac{1}{\langle k - (k_3 + k_4) \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3 + k_4 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right]^{\frac{1}{p'}} \right\|_{l_{k_3}^{p'}} \\ &\cdot \left\| \left[ \frac{1}{\langle k - (k_3 + k_4) \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3 + k_4 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right]^{\frac{1}{p}} \langle k_3 \rangle^s |\hat{\xi}_{k_3}| \right\|_{l_{k_3}^p} \\ &\lesssim \left\| \left[ \frac{1}{\langle k - (k_3 + k_4) \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3 + k_4 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right]^{\frac{1}{p}} \langle k_3 \rangle^s |\hat{\xi}_{k_3}| \right\|_{l_{k_3}^p}. \end{aligned}$$

This gives that

$$\begin{aligned} \|M_k\|_{l_k^p} &\lesssim \left\| \left[ \frac{1}{\langle k - (k_3 + k_4) \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3 + k_4 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right]^{\frac{1}{p}} \langle k_3 \rangle^s |\hat{\xi}_{k_3}| \right\|_{l_{k,k_3}^p} \\ &\lesssim \left\| \left[ \frac{1}{\langle k - (k_3 + k_4) \rangle^{1+\frac{1}{2}\varepsilon_0}} + \frac{1}{\langle k + k_3 + k_4 \rangle^{1+\frac{1}{2}\varepsilon_0}} \right]^{\frac{1}{p}} \langle k_3 \rangle^s |\hat{\xi}_{k_3}| \right\|_{l_k^1} \\ &\lesssim \|\xi\|_{\hat{b}^{s,p}}. \end{aligned} \tag{3.18}$$

Insert the estimate (3.18) into (3.17), we have

$$\begin{aligned} \left\| \langle k \rangle^{s+\gamma} |I_{2222,k}| \right\|_{l_k^2(|k| \geq N_0)} &\lesssim \|\langle k \rangle^s |\hat{\xi}_k|\|_{l_k^\infty}^2 \int_0^t \sum_{k_4} \|M_k\|_{l_k^p} \|\langle k \rangle^{-2+\varepsilon_0+\gamma}\|_{l_k^r} \|h\|_{l^2} \langle k_4 \rangle^s |\hat{v}_{k_4}| d\rho \\ &\lesssim \|\xi\|_{\hat{b}^{s,\infty}}^2 \|\xi\|_{\hat{b}^{s,p}} \int_0^t \sum_{k_4} \langle k_4 \rangle^s |\hat{v}_{k_4}| d\rho \lesssim t \|\xi\|_{\hat{b}^{s,p}}^3 \|v\|_{L_t^\infty H_x^{s+\gamma}}, \end{aligned}$$

where in the last step we have used that  $\gamma > \frac{1}{2}$ .

3) On  $I_{2223,k}$ . Similarly as treating the term  $I_{2222,k}$ , we can obtain

$$\left\| \langle k \rangle^{s+\gamma} |I_{2223,k}| \right\|_{l_k^2(|k| \geq N_0)} \lesssim t \|\xi\|_{\hat{b}^{s,p}}^2 \| |u|^2 u \|_{L_t^\infty H_x^{s+\gamma}} \lesssim t \|\xi\|_{\hat{b}^{s,p}}^2 \|u\|_{L_t^\infty H_x^{s+\gamma}}^3.$$

Hence, we obtain the estimate on  $I_{22,k}$  as follows

$$\begin{aligned} \left\| \langle k \rangle^{s+\gamma} |I_{22,k}| \right\|_{l_k^2(|k| \geq N_0)} &\lesssim t \|\hat{\xi}_k\|_{l_k^\infty}^2 \|v\|_{L_t^\infty H_x^{s+\gamma}}^2 + N_0^{-\varepsilon_0} \|\xi\|_{\hat{b}^{s,p}}^2 \|v\|_{L_t^\infty H_x^{s+\gamma}} \\ &\quad + t \|\xi\|_{\hat{b}^{s,p}}^3 \|v\|_{L_t^\infty H_x^{s+\gamma}} + t \|\xi\|_{\hat{b}^{s,p}}^2 \|u\|_{L_t^\infty H_x^{s+\gamma}}^3. \end{aligned}$$

This completes the estimates of  $I_{222}$  and then finishes **Section 3.1.2**.

3.1.3. *Estimates on  $I_{23,k}$ .* The estimation of this term is very similar as the term  $I_{21}$ , so we almost repeat the estimates of the latter. Firstly, we split it into the following two parts:

$$\begin{aligned} I_{23,k} &= -i \int_0^t \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2) \neq 0, |k_1| \lesssim |k_2|}} e^{i\rho k^2} \frac{1}{i\phi(k,k_2)} \hat{\xi}_{k_1} \widehat{(|u|^2 u)}_{k_2} d\rho \\ &\quad - i \int_0^t \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2) \neq 0, |k_1| \gg |k_2|}} e^{i\rho k^2} \frac{1}{i\phi(k,k_2)} \hat{\xi}_{k_1} \widehat{(|u|^2 u)}_{k_2} d\rho \\ &\triangleq I_{231,k} + I_{232,k}. \end{aligned}$$

• **Estimates on  $I_{231,k}$ .** Since

$$\frac{1}{|\phi(k,k_2)|} \leq \frac{1}{2\langle k_1 \rangle^2} + \frac{1}{2\langle k_1 + 2k_2 \rangle^2}.$$

Hence, we have that

$$\begin{aligned} \langle k \rangle^{s+\gamma} |I_{231,k}| &\lesssim \int_0^t \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2) \neq 0, |k_1| \lesssim |k_2|}} \langle k \rangle^{s+\gamma} \frac{1}{|\phi(k,k_2)|} |\hat{\xi}_{k_1}| \widehat{(|u|^2 u)}_{k_2} d\rho \\ &\lesssim \int_0^t \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2) \neq 0, |k_1| \lesssim |k_2|}} \langle k_2 \rangle^{s+\gamma} \left[ \frac{1}{\langle k_1 \rangle^2} + \frac{1}{\langle k_1 + 2k_2 \rangle^2} \right] |\hat{\xi}_{k_1}| \widehat{(|u|^2 u)}_{k_2} d\rho. \end{aligned}$$

Then treating similarly as  $I_{211,k}$  and by Kato-Ponce's inequality, we have that

$$\|\langle k \rangle^\gamma I_{231,k}\|_{l_k^2} \lesssim \int_0^t \|\hat{\xi}_k\|_{l_k^\infty} \| |u|^2 u \|_{L_t^\infty H_x^{s+\gamma}} d\rho \lesssim t \|\hat{\xi}_k\|_{l_k^\infty} \|u\|_{L_t^\infty H_x^{s+\gamma}}^3.$$

• **Estimates on  $I_{232,k}$ .** Under the restriction in the summation, it gives that

$$|\phi(k,k_2)| \gtrsim \langle k_1 \rangle^2.$$

Then we have that

$$\langle k \rangle^{s+\gamma} |I_{232,k}| \lesssim \int_0^t \sum_{\substack{k_1+k_2=k \\ \phi(k,k_2) \neq 0, |k_1| \gg |k_2|}} \langle k_1 \rangle^{\gamma+s-2} |\hat{\xi}_{k_1}| \widehat{(|u|^2 u)}_{k_2} d\rho.$$

Again, treating similarly as  $I_{211}$  and by Kato-Ponce's inequality, we have that

$$\|\langle k \rangle^{s+\gamma} I_{232,k}\|_{l_k^2} \lesssim \int_0^t \|\xi\|_{\hat{b}^{s,p}} \| |u|^2 u \|_{L_t^\infty H_x^{s+\gamma}} d\rho \lesssim t \|\xi\|_{\hat{b}^{s,p}} \|u\|_{L_t^\infty H_x^{s+\gamma}}^3.$$

Combining with the estimates on  $I_{231}$  and  $I_{232}$ , we obtain that

$$\|\langle k \rangle^{s+\gamma} I_{23,k}\|_{l_k^2} \lesssim t \|\xi\|_{\hat{b}^{s,p}} \|u\|_{L_t^\infty H_x^{s+\gamma}}^3.$$

Collecting the estimates on  $I_1$  and  $I_{21} - I_{23}$  above, we have that there exists a constant  $C > 0$  which depends on  $\|\xi\|_{\hat{b}^{s,p}}$  such that

$$\|\langle k \rangle^{s+\gamma} I_k\|_{L_t^\infty l_k^2([0,T] \times \{|k| > N_0\})} \leq C (T + N_0^{-\frac{\varepsilon_0}{2}}) (\|v\|_{L_t^\infty H_x^{s+\gamma}} + \|v\|_{L_t^\infty H_x^{s+\gamma}}^3).$$

This together with (3.6) give that

$$\begin{aligned} \|\Phi^1(u)\|_{L_t^\infty H_x^{s+\gamma}([0,T])} &= \|\langle k \rangle^{s+\gamma} I_k\|_{L_t^\infty l_k^2} \\ &\leq C \left( T N_0^{s+\gamma} + N_0^{-\frac{\varepsilon_0}{2}} \right) (\|v\|_{L_t^\infty H_x^{s+\gamma}} + \|v\|_{L_t^\infty H_x^{s+\gamma}}^3). \end{aligned}$$

This finishes the proof of [Lemma 3.2](#).  $\square$

*Proof of Lemma 3.1.* By Kato-Ponce's inequality, we have

$$\left\| i \int_0^t e^{-it' \partial_x^2} (|u(t')|^2 u(t')) dt' \right\|_{L_t^\infty H_x^{s+\gamma}([0, T])} \leq CT \|u\|_{L_t^\infty H_x^{s+\gamma}}^3.$$

This combining with [\(3.2\)](#) and [Lemma 3.2](#) yields that under the assumption [\(3.3\)](#),

$$\begin{aligned} \|\Phi(v)\|_{L_t^\infty H_x^{s+\gamma}([0, T])} &\leq \|u^0\|_{H_x^{s+\gamma}} + C(TN_0^{s+\gamma} + N_0^{-\frac{\varepsilon_0}{2}})(R_0 + R_0^3) \\ &\leq \frac{1}{2}R_0 + C(TN_0^{s+\gamma} + N_0^{-\frac{\varepsilon_0}{2}})(R_0 + R_0^3). \end{aligned}$$

This proves the lemma.  $\square$

Now we choosing  $N_0$  and  $T$  in [Lemma 3.1](#) such that

$$N_0^{\frac{\varepsilon_0}{2}} = 8(1 + R_0^2); \quad T = [8C_0(N_0)(1 + R_0^2)]^{-1},$$

then we obtain [\(3.4\)](#) and [\(3.5\)](#) with  $\theta = \frac{3}{4}$ , and thus complete the proof of [Theorem 1.1](#).  $\square$

**3.2. Proof of Theorem 1.2.** In the following, we denote  $T$  (using the same notation) to be a positive constant which is smaller than the lifespan obtained in [Theorem 1.1](#).

We denote  $\varphi = \partial_t u$ , then in order to prove that  $u \in C_t^0([0, T]; H^{s+\frac{3}{2}+\frac{1}{p}}(\mathbb{T}))$ , it is sufficient to prove that  $\varphi \in C_t^0([0, T]; H^{s-\frac{1}{2}+\frac{1}{p}}(\mathbb{T}))$ . Indeed, note that

$$\partial_{xx} u = -i\varphi - \xi u + |u|^2 u.$$

This gives that

$$\|u\|_{H^{s+\frac{3}{2}+\frac{1}{p}}} \lesssim \|\varphi\|_{H^{s-\frac{1}{2}+\frac{1}{p}}} + \|\xi u\|_{H^{s-\frac{1}{2}+\frac{1}{p}}} + \||u|^2 u\|_{H^{s-\frac{1}{2}+\frac{1}{p}}}.$$

Note that  $H^{s,p'} \hookrightarrow \hat{b}^{s,p}$ , then from the proof of [Theorem 1.1](#), we have that

$$\|u\|_{L_t^\infty H_x^{s+\frac{3}{2}+\frac{1}{p}-}([0, T])} \leq 2\|u_0\|_{H^{s+2}}. \quad (3.19)$$

By Sobolev embedding  $L^{p'} \hookrightarrow H^{-\frac{1}{2}+\frac{1}{p}}$  and [\(3.19\)](#), we further have that

$$\begin{aligned} \|u\|_{H^{s+\frac{3}{2}+\frac{1}{p}}} &\lesssim \|\varphi\|_{H^{s-\frac{1}{2}+\frac{1}{p}}} + \|J^s(\xi u)\|_{L^{p'}} + \|J^s(|u|^2 u)\|_{L^{p'}} \\ &\lesssim \|\varphi\|_{H^{s-\frac{1}{2}+\frac{1}{p}}} + \|J^s \xi\|_{L^{p'}} \cdot \|u\|_{L^\infty} \\ &\quad + \|\xi\|_{L^{p'}} \cdot \|J^s u\|_{L^\infty} + \|J^s u\|_{L^{p'}} \|u\|_{L^\infty}^2 \\ &\lesssim \|\varphi\|_{H^{s-\frac{1}{2}+\frac{1}{p}}} + \|J^s \xi\|_{L^{p'}} \cdot \|u\|_{H^{s+\frac{3}{2}+\frac{1}{p}-}} \\ &\quad + \|\xi\|_{L^{p'}} \cdot \|u\|_{H^{s+\frac{3}{2}+\frac{1}{p}-}} + \|u\|_{H^{s+\frac{3}{2}+\frac{1}{p}-}}^3 \\ &\lesssim \|\varphi\|_{H^{s-\frac{1}{2}+\frac{1}{p}}} + \|\xi\|_{H^{s,p'}} \cdot \|u_0\|_{H^{s+2}} + \|u_0\|_{H^{s+2}}^3. \end{aligned} \quad (3.20)$$

This proves the claim that  $\varphi \in C_t^0([0, T]; H^{s-\frac{1}{2}+\frac{1}{p}}(\mathbb{T}))$  implies  $u \in C_t^0([0, T]; H^{s+\frac{3}{2}+\frac{1}{p}}(\mathbb{T}))$ .

Moreover, from [\(1.1\)](#),  $\varphi$  satisfies the following equation

$$\begin{cases} i\partial_t \varphi + \partial_{xx} \varphi + \xi \varphi = \mathcal{O}(u^2 \varphi), & \text{for } x \in \mathbb{T} \text{ and } t \in (0, T], \\ \varphi(0, x) = i(\partial_{xx} u_0 + \xi u_0 - |u_0|^2 u_0) \triangleq \varphi_0, & \text{for } x \in \mathbb{T}. \end{cases} \quad (3.21)$$

Here we denote  $\mathcal{O}(u^2\varphi) = 2|u|^2\varphi + u^2\bar{\varphi}$ . From Duhamel's formula, we have

$$\varphi(t) = e^{it\partial_x^2}\varphi_0 + i \int_0^t e^{i(t-\rho)\partial_x^2} \left( \xi\varphi(\rho) + \mathcal{O}(u^2\varphi(\rho)) \right) d\rho. \quad (3.22)$$

Accordingly, we denote that

$$\Psi(\varphi) = e^{it\partial_x^2}\varphi_0 + i \int_0^t e^{i(t-\rho)\partial_x^2} \left( \xi\varphi(\rho) + \mathcal{O}(u^2\varphi(\rho)) \right) d\rho.$$

In this proof, we denote  $a = \frac{1}{2} - \frac{1}{p}$  for short. Firstly, similarly as (3.20), we have

$$\|\varphi_0\|_{H^{-a+s}} \lesssim \|u_0\|_{H^{s-a+2}} + \|\xi\|_{H^{s,p'}} \cdot \|u_0\|_{H^{s+2}} + \|u_0\|_{H^{s+2}}^3. \quad (3.23)$$

Hence, we have  $\varphi_0 \in H^{-a+s}$ . Similar as the proof of [Theorem 1.1](#), the proof of [Theorem 1.2](#) is reduced to the following lemma.

**Lemma 3.3.** *There exist a constant  $\varepsilon_0$  and a positive function  $C(\cdot)$  such that for any  $N_0 > 0$ , any  $t \in [0, T]$ ,*

$$\|\Psi(\varphi)\|_{H^{-a+s}} \leq \|\varphi_0\|_{H^{-a+s}} + [tC(N_0) + N_0^{-\varepsilon_0}] \cdot (\|\varphi\|_{L_t^\infty H_x^{-a+s}} + \|\varphi\|_{L_t^\infty H_x^{-a+s}}^3). \quad (3.24)$$

*Proof.* We split it into the following two parts.

• **Estimate on**  $\left\| \int_0^t e^{i(t-\rho)\partial_x^2} [\xi\varphi(\rho)] d\rho \right\|_{H^{-a+\rho}}$ . Denote  $w = e^{-it\partial_x^2}\varphi$ , then it reduces to

$$I \triangleq \int_0^t e^{-i\rho\partial_x^2} [\xi \cdot e^{i\rho\partial_x^2} w(\rho)] d\rho. \quad (3.25)$$

Since  $H^{s,p'} \hookrightarrow \hat{b}^{s,p}$ , arguing similarly as the proof of [Lemma 3.2](#), we have that (for which the details are omitted here)

$$\begin{aligned} \|I\|_{H^{-a+s}} &\lesssim [tC(N_0) + N^{-\varepsilon_0}] \cdot (\|w\|_{L_t^\infty H_x^{-a+s}} + \|w\|_{L_t^\infty H_x^{-a+s}}^3) \\ &= [tC(N_0) + N^{-\varepsilon_0}] \cdot (\|\varphi\|_{L_t^\infty H_x^{-a+s}} + \|\varphi\|_{L_t^\infty H_x^{-a+s}}^3). \end{aligned} \quad (3.26)$$

• **Estimate on**  $\left\| \int_0^t e^{i(t-\rho)\partial_x^2} [\mathcal{O}(u^2\varphi)] d\rho \right\|_{H^{-a+\rho}}$ . If  $-a + s \geq 0$ , then by Kato-Ponce's Inequality and (3.19),

$$\begin{aligned} \left\| \int_0^t e^{i(t-\rho)\partial_x^2} [\mathcal{O}(u^2\varphi)] d\rho \right\|_{H^{-a+s}} &\lesssim \int_0^t \|J^{-a+s}(u^2\varphi)\|_{L^2} d\rho \\ &\lesssim \|\varphi\|_{L_t^\infty H_x^{-a+s}} \|u\|_{L_t^\infty H_x^{2-a+s}}^2 \\ &\lesssim \|u_0\|_{H^{s+2}}^2 \|\varphi\|_{L_t^\infty H_x^{-a+s}}. \end{aligned} \quad (3.27)$$

If  $-a + s < 0$ , using dual argument, we note that

$$\|J^{-a+s}(u^2\varphi)\|_{L^2} = \sup_{h: \|h\|_{l^2}=1} \langle J^{-a+s}(u^2\varphi), h \rangle. \quad (3.28)$$

By Hölder's inequality, we have

$$\begin{aligned} &\langle J^{-a+s}(u^2\varphi), h \rangle \\ &= \sum_N \langle P_N J^{-a+s}(u^2\varphi), P_N h \rangle \lesssim \sum_N N^{-a+s} \|P_N(u^2\varphi)\|_{L^2} \cdot \|P_N h\|_{L^2} \\ &\lesssim \sum_{N \gtrsim M} N^{-a+s} \|P_N(u^2 P_M \varphi)\|_{L^2} \cdot \|P_N h\|_{L^2} + \sum_{N \ll M} N^{-a+s} \|P_N(u^2 P_M \varphi)\|_{L^2} \cdot \|P_N h\|_{L^2} \\ &\triangleq S_1 + S_2. \end{aligned} \quad (3.29)$$

For  $S_1$ , by Schur's test in [Lemma 2.2](#) and [\(3.19\)](#), we have

$$\begin{aligned}
S_1 &= \sum_{N \gtrsim M} N^{-a+s} \|P_N(u^2 P_M \varphi)\|_{L^2} \cdot \|P_N h\|_{L^2} \\
&\lesssim \sum_{N \gtrsim M} N^{-a+s} \|u\|_{H^{\frac{1}{2}+}}^2 \|P_M \varphi\|_{L^2} \cdot \|P_N h\|_{L^2} \\
&\lesssim \|u_0\|_{H^{s+2}}^2 \sum_{N \gtrsim M} \left(\frac{M}{N}\right)^{-a+s} \|P_M \varphi\|_{H^{-a+s}} \|P_N h\|_{L^2} \\
&\lesssim \|u_0\|_{H^{s+2}}^2 \|\varphi\|_{H^{-a+s}} \|h\|_{L^2}.
\end{aligned} \tag{3.30}$$

For  $S_2$ , note that when  $N \ll M$ ,  $P_N(u^2 P_M \varphi) = P_N(u P_{>\frac{M}{2}} u P_M \varphi)$ . Then using Hölder's inequality and [\(3.19\)](#), we have

$$\begin{aligned}
S_2 &= \sum_{N \ll M} N^{-a+s} \|P_N(u^2 P_M \varphi)\|_{L^2} \cdot \|P_N h\|_{L^2} \\
&\lesssim \sum_{N \ll M} N^{-a+s} \|u\|_{L^\infty} \|P_{>\frac{M}{2}} u\|_{L^\infty} \|P_M \varphi\|_{L^2} \|P_N h\|_{L^2} \\
&\lesssim \sum_{N \ll M} N^{-a+s} \|u\|_{H^{\frac{1}{2}+}} \|P_{>\frac{M}{2}} u\|_{H^{\frac{1}{2}+}} \|P_M \varphi\|_{L^2} \|P_N h\|_{L^2} \\
&\lesssim \sum_{N \ll M} M^{-2(s+\frac{1}{p})-\frac{1}{2}+} N^{-a+s} \|u\|_{H^{\frac{1}{2}+}} \|P_{>\frac{M}{2}} u\|_{H^{s+\frac{1}{p}+\frac{3}{2}+}} \|P_M \varphi\|_{H^{-a+s}} \|P_N h\|_{L^2} \\
&\lesssim \sum_{N \ll M} M^{-2(s+\frac{1}{p})-\frac{1}{2}+} N^{-a+s} \|u_0\|_{H^{s+2}}^2 \|\varphi\|_{H^{-a+s}} \|h\|_{L^2}.
\end{aligned}$$

Note that  $-2(s+\frac{1}{p})-\frac{1}{2} < 0$ ,  $-a+s < 0$ , by Cauchy-Schwartz's inequality, we obtain that

$$S_2 \lesssim \|u_0\|_{H^{s+2}}^2 \|\varphi\|_{H^{-a+s}} \|h\|_{L^2}. \tag{3.31}$$

Inserting [\(3.30\)](#) and [\(3.31\)](#) into [\(3.29\)](#), and then by [\(3.28\)](#) we have that

$$\|J^{-a+s}(u^2 \varphi)\|_{L^2} \lesssim \|u_0\|_{H^{s+2}}^2 \|\varphi\|_{H^{-a+s}}.$$

This implies that

$$\begin{aligned}
\left\| \int_0^t e^{-i(t-\rho)\partial_x^2} [\mathcal{O}(u^2 \varphi)] d\rho \right\|_{H^{-a+s}} &\lesssim \int_0^t \left\| e^{-i(t-\rho)\partial_x^2} [\mathcal{O}(u^2 \varphi)] \right\|_{H^{-a+s}} d\rho \\
&\lesssim t \|u_0\|_{H^{s+2}}^2 \|\varphi\|_{L_t^\infty H_x^{-a+s}}.
\end{aligned}$$

This together with [\(3.27\)](#) and [\(3.26\)](#), yields [\(3.24\)](#).  $\square$

#### 4. PROOF FOR ILL-POSEDNESS THEORY

In this section, we are going to prove the three ill-posedness results, i.e., [Theorem 1.3](#), [Theorem 1.4](#) and [Theorem 1.5](#). Again, we assume  $\lambda = -1$  in [\(1.1\)](#) for simplicity. In order to apply the tool [Lemma 2.3](#), we consider the following.

Let  $f = f(x)$  be a time-independent function. Denote

$$A_2(f) \triangleq i \int_0^t e^{-i\rho\partial_x^2} \left( \xi e^{i\rho\partial_x^2} f + |e^{i\rho\partial_x^2} f|^2 e^{i\rho\partial_x^2} f \right) d\rho.$$

Taking Fourier transform, we have that for any  $k \in \mathbb{Z}$ ,

$$\begin{aligned} (\widehat{A_2(f)})_k &= i \int_0^t \left( \sum_{k_1+k_2=k} e^{i\rho(k^2-k_2^2)} \hat{\xi}_{k_1} \hat{f}_{k_2} + \sum_{k_1+k_2+k_3=k} e^{i\rho(k^2+k_1^2-k_2^2-k_3^2)} (\widehat{f})_{k_1} \hat{f}_{k_2} \hat{f}_{k_3} \right) d\rho \\ &= i \sum_{k_1+k_2=k} \int_0^t e^{i\rho(k^2-k_2^2)} d\rho \hat{\xi}_{k_1} \hat{f}_{k_2} \\ &\quad + i \sum_{k_1+k_2+k_3=k} \int_0^t e^{i\rho(k^2+k_1^2-k_2^2-k_3^2)} d\rho (\widehat{f})_{k_1} \hat{f}_{k_2} \hat{f}_{k_3}. \end{aligned}$$

The first term is equal to

$$i \sum_{\substack{k_1+k_2=k \\ k^2-k_2^2=0}} \int_0^t e^{i\rho(k^2-k_2^2)} d\rho \hat{\xi}_{k_1} \hat{f}_{k_2} + i \sum_{\substack{k_1+k_2=k \\ k^2-k_2^2 \neq 0}} \int_0^t e^{i\rho(k^2-k_2^2)} d\rho \hat{\xi}_{k_1} \hat{f}_{k_2}.$$

Note that the set

$$\begin{aligned} &\{(k_1, k_2) : k_1 + k_2 = k, k^2 - k_2^2 = 0\} \\ &= \{(k_1, k_2) : k_1 = 0, k_2 = k\} \cup \{(k_1, k_2) : k_1 = 2k, k_2 = -k\} \setminus \{(k_1, k_2) : k_1 = k_2 = 0\}. \end{aligned}$$

We further get that

$$\begin{aligned} &i \sum_{k_1+k_2=k} \int_0^t e^{i\rho(k^2-k_2^2)} d\rho \hat{\xi}_{k_1} \hat{f}_{k_2} \\ &= it\hat{\xi}_0 \hat{f}_k + it\hat{\xi}_{2k} \hat{f}_{-k} - it\hat{\xi}_0 \hat{f}_0 + i \sum_{\substack{k_1+k_2=k \\ k^2-k_2^2 \neq 0}} \frac{1}{i(k^2-k_2^2)} \left( e^{it(k^2-k_2^2)} - 1 \right) \hat{\xi}_{k_1} \hat{f}_{k_2}. \end{aligned}$$

This implies that

$$\begin{aligned} (\widehat{A_2(f)})_k &= it\hat{\xi}_0 \hat{f}_k + it\hat{\xi}_{2k} \hat{f}_{-k} - it\hat{\xi}_0 \hat{f}_0 \\ &\quad + i \sum_{\substack{k_1+k_2=k \\ k^2-k_2^2 \neq 0}} \frac{1}{i(k^2-k_2^2)} \left( e^{it(k^2-k_2^2)} - 1 \right) \hat{\xi}_{k_1} \hat{f}_{k_2} \\ &\quad + i \sum_{k_1+k_2+k_3=k} \int_0^t e^{i\rho(k^2+k_1^2-k_2^2-k_3^2)} d\rho (\widehat{f})_{k_1} \hat{f}_{k_2} \hat{f}_{k_3}. \end{aligned} \tag{4.1}$$

#### 4.1. Proof of **Theorem 1.3**.

• **Case 1:**  $p = \infty$ . We only need to show the ill-posedness in  $H^{s+\frac{3}{2}}$ . To do this, we define the initial data

$$u_0 \triangleq \epsilon. \tag{4.2}$$

Then  $\|u_0\|_{H^2} = \epsilon$ . Moreover, we define

$$\xi \triangleq 1 + \sum_{k \in \mathbb{Z} \setminus \{0\}} |k|^{-s} e^{ikx}. \tag{4.3}$$

Then  $\|\xi\|_{\dot{H}^s, \infty} = 1$ . In particular,  $\xi = 2\pi\delta$  when  $s = 0$ .

Then by (4.1), in which the first three terms vanish, we have that

$$(\widehat{A_2(u_0)})_k = i \sum_{k_1+k_2=k} \frac{1}{i(k^2 - k_2^2)} \left( e^{it(k^2 - k_2^2)} - 1 \right) \hat{\xi}_{k_1}(\widehat{u_0})_{k_2} \quad (4.4a)$$

$$+ i \sum_{k_1+k_2+k_3=k} \int_0^t e^{i\rho(k^2+k_1^2-k_2^2-k_3^2)} d\rho (\widehat{u_0})_{k_1} (\widehat{u_0})_{k_2} (\widehat{u_0})_{k_3}. \quad (4.4b)$$

For (4.4a), according to the definition of  $u_0$  and  $\xi$  we have

$$(4.4a) = \epsilon \frac{1}{k^{2+s}} \left( e^{itk^2} - 1 \right). \quad (4.5)$$

Now we denote the set

$$\mathbb{K} \triangleq \{M_0(2j+1) : j \in \mathbb{Z}^+, 1 \leq j \leq M_1\},$$

where  $M_0, M_1 \geq 10$  are some large constants determined later. Then for  $k \in \mathbb{K}$ , we have

$$k^2 = M_0^2(4j^2 + 4j + 1).$$

Taking

$$t \triangleq \frac{\pi}{2} M_0^{-2}, \quad (4.6)$$

then  $tk^2 = 2(j^2 + j)\pi + \frac{\pi}{2}$ . This yields that  $e^{itk^2} = i$ . This further gives (4.5) that for any  $k \in \mathbb{K}$ ,

$$(4.4a) = \epsilon(i-1) \frac{1}{k^{2+s}}. \quad (4.7)$$

Now we consider the estimate on (4.4a). By (4.7), it reads

$$\begin{aligned} \|(1 + |k|^2)^{\frac{1}{2}(s+\frac{3}{2})} (4.4a)\|_{l_k^2}^2 &\geq \|(1 + |k|^2)^{\frac{1}{2}(s+\frac{3}{2})} (4.4a)\|_{l_k^2(\mathbb{K})}^2 \\ &= \epsilon^2 \sum_{k \in \mathbb{K}} (1 + |k|^2)^{s+\frac{3}{2}} \left| (i-1) \frac{1}{k^{2+s}} \right|^2 \\ &= 2\epsilon^2 \sum_{k \in \mathbb{K}} (1 + |k|^2)^{s+\frac{3}{2}} k^{-(4+2s)}. \end{aligned}$$

Note that in  $\mathbb{K}$ ,

$$(1 + |k|^2)^{s+\frac{3}{2}} k^{-(4+2s)} \geq |k|^{-1}.$$

Therefore, we get that

$$\|(1 + |k|^2)^{\frac{1}{2}(s+\frac{3}{2})} (4.4a)\|_{l_k^2}^2 \geq 2\epsilon^2 \sum_{k \in \mathbb{K}} |k|^{-1} = 2\epsilon^2 M_0^{-1} \sum_{j=1}^{M_1} (2j+1)^{-1} \geq 2\epsilon^2 M_0^{-1} \ln(2M_1+1).$$

For (4.4b), by Cauchy-Schwartz's inequality, we have

$$\begin{aligned} &\|(1 + |k|^2)^{\frac{1}{2}(s+\frac{3}{2})} (4.4b)\|_{l_k^2}^2 \\ &= \sum_k (1 + |k|^2)^{s+\frac{3}{2}} \left| \sum_{k_1+k_2+k_3=k} \int_0^t e^{i\rho(k^2+k_1^2-k_2^2-k_3^2)} d\rho (\widehat{u_0})_{k_1} (\widehat{u_0})_{k_2} (\widehat{u_0})_{k_3} \right|^2 \\ &= t^2 \epsilon^6. \end{aligned} \quad (4.8)$$

Together with the estimates on (4.4a) and (4.4b) above, we obtain

$$\begin{aligned} \|A_2(u_0)\|_{H^{s+\frac{3}{2}}} &= \left\| (1 + |k|^2)^{\frac{1}{2}(s+\frac{3}{2})} (\widehat{A_2(u_0)})_k \right\|_{l_k^2} \\ &\geq \sqrt{2}\epsilon M_0^{-\frac{1}{2}} \sqrt{\ln(2M_1 + 1)} - t\epsilon^3 \\ &= \sqrt{2}\epsilon M_0^{-\frac{1}{2}} \sqrt{\ln(2M_1 + 1)} - \frac{\pi}{2} M_0^{-2} \epsilon^3 \rightarrow +\infty, \quad \text{when } M_1 \rightarrow +\infty. \end{aligned}$$

Then by Lemma 2.3, we obtain the ill-posedness in  $H^{s+\frac{3}{2}}$ , and thus finish the proof of Theorem 1.3 in the case of  $p = \infty$ .

• **Case 2:**  $2 < p < \infty$ . As before, we define the initial data

$$u_0 \triangleq \epsilon.$$

Now, we define

$$\xi \triangleq 1 + \sum_{k \in \mathbb{Z} \setminus \{0\}} \frac{1}{|k|^{s+\frac{1}{p}} (\ln |k|)^\alpha} e^{ikx}, \quad (4.9)$$

where  $\alpha > \frac{1}{p}$  will be decided later. Then

$$\|\xi\|_{\dot{b}^{s,p}} \lesssim 1.$$

Therefore, the same estimates as (4.7) give that

$$\begin{aligned} \|(1 + |k|^2)^{\frac{1}{2}(s+\frac{3}{2}+\frac{1}{p})} (4.4a)\|_{l_k^2}^2 &\geq \|(1 + |k|^2)^{\frac{1}{2}(s+\frac{3}{2}+\frac{1}{p})} (4.4a)\|_{l_k^2(\mathbb{K})}^2 \\ &= 2\epsilon^2 \sum_{k \in \mathbb{K}} (1 + |k|^2)^{s+\frac{3}{2}+\frac{1}{p}} k^{-(4+2s+\frac{2}{p})} (\ln |k|)^{-2\alpha} \\ &\geq 2\epsilon^2 \sum_{k=1}^{M_1} k^{-1} (\ln |k|)^{-2\alpha}. \end{aligned}$$

This together with (4.8) yields that

$$\begin{aligned} \|A_2(u_0)\|_{H^{s+\frac{3}{2}+\frac{1}{p}}} &= \left\| (1 + |k|^2)^{\frac{1}{2}(s+\frac{3}{2}+\frac{1}{p})} (\widehat{A_2(u_0)})_k \right\|_{l_k^2} \\ &\geq \sqrt{2}\epsilon \left( \sum_{k=1}^{M_1} k^{-1} (\ln k)^{-2\alpha} \right)^{\frac{1}{2}} - \frac{\pi}{2} M_0^{-2} \epsilon^3. \end{aligned}$$

Choosing  $\frac{1}{p} < \alpha < \frac{1}{2}$ , then

$$\sum_{k=1}^{M_1} k^{-1} (\ln k)^{-2\alpha} \rightarrow +\infty, \quad \text{when } M_1 \rightarrow +\infty.$$

Then the estimates above give that

$$\|A_2(u_0)\|_{H^{s+\frac{3}{2}+\frac{1}{p}}} \rightarrow +\infty, \quad \text{when } M_1 \rightarrow +\infty.$$

The proof is done by applying Lemma 2.3.  $\square$

4.2. **Proof of Theorem 1.4.** As in the proof of [Theorem 1.3](#), we define the initial data

$$u_0 \triangleq \epsilon,$$

and

$$\xi \triangleq 1 + \sum_{k \in \mathbb{Z} \setminus \{0\}} \frac{1}{|k|^{s+\beta}} e^{ikx}, \quad (4.10)$$

where  $\beta > \frac{1}{2}$  will be decided later. Then  $\|\xi\|_{H^s} \lesssim 1$ . Given  $\gamma > 2$ , arguing similarly as above, we have that

$$\begin{aligned} \|(1 + |k|^2)^{\frac{1}{2}(s+\gamma)}(4.4a)\|_{l_k^2}^2 &\geq \|(1 + |k|^2)^{\frac{1}{2}(s+\gamma)}(4.4a)\|_{l_k^2(\mathbb{K})}^2 \\ &= 2\epsilon^2 \sum_{k \in \mathbb{K}} (1 + |k|^2)^{s+\gamma} |k|^{-(4+2s+2\beta)} \geq 2\epsilon^2 \sum_{k=1}^{M_1} |k|^{2(\gamma-2)-2\beta}. \end{aligned}$$

Choosing  $\beta$  such that  $2\beta - 1 < 2(\gamma - 2)$ , then

$$\|(1 + |k|^2)^{\frac{1}{2}(s+\gamma)}(4.4a)\|_{l_k^2}^2 \geq 2\epsilon^2 M_1^{2(\gamma-2)-2\beta+1}.$$

This together with [\(4.8\)](#) yields that

$$\begin{aligned} \|A_2(u_0)\|_{H^{s+\frac{3}{2}+\frac{1}{p}}} &= \left\| (1 + |k|^2)^{\frac{1}{2}(s+\frac{3}{2}+\frac{1}{p})} (\widehat{A_2(u_0)})_k \right\|_{l_k^2} \\ &\geq \sqrt{2}\epsilon M_1^{\gamma-2-\beta+\frac{1}{2}} - \frac{\pi}{2} M_0^{-2}\epsilon^3 \rightarrow +\infty, \quad \text{when } M_1 \rightarrow +\infty. \end{aligned}$$

The proof is done by applying [Lemma 2.3](#).

4.3. **Proof of Theorem 1.5.** Note that the ill-posedness in  $H^\gamma$  for  $\gamma \geq \frac{3}{2}$  has been essentially presented in [Theorem 1.3](#), so here we only need to further consider  $\gamma < \frac{3}{2}$ .

Define the initial data

$$u_0 \triangleq \epsilon \sum_{k:|k-N|\leq 10} N^{-\gamma} e^{ikx}, \quad (4.11)$$

then

$$\|u_0\|_{H^\gamma} \sim \epsilon.$$

Define the potential

$$\xi \triangleq \sum_{k:|k|>10} \ln(|k|) e^{ikx}, \quad (4.12)$$

then for any  $\epsilon > 0$ ,

$$\|\hat{\xi}_k\|_{\hat{b}^{-\epsilon, \infty}} < +\infty, \quad \text{and} \quad \|\hat{\xi}_k\|_{l_k^\infty} = +\infty.$$

By [\(4.1\)](#), we have that

$$\begin{aligned} (\widehat{A_2(u_0)})_k &= it\hat{\xi}_0(\widehat{u_0})_k + it\hat{\xi}_{2k}(\widehat{u_0})_{-k} - it\hat{\xi}_0(\widehat{u_0})_0 \\ &\quad + i \sum_{\substack{k_1+k_2=k \\ k^2-k_2^2 \neq 0}} \frac{1}{i(k^2-k_2^2)} \left( e^{it(k^2-k_2^2)} - 1 \right) \hat{\xi}_{k_1}(\widehat{u_0})_{k_2} \\ &\quad + i \sum_{k_1+k_2+k_3=k} \int_0^t e^{i\rho(k^2+k_1^2-k_2^2-k_3^2)} d\rho (\widehat{u_0})_{k_1} (\widehat{u_0})_{k_2} (\widehat{u_0})_{k_3}. \end{aligned}$$

According to the definition of  $\xi$  in (4.12), we have  $\hat{\xi}_0 = 0$  and so

$$\widehat{(A_2(u_0))}_k = it\hat{\xi}_{2k}\widehat{(u_0)}_{-k} \quad (4.13a)$$

$$+ i \sum_{\substack{k_1+k_2=k \\ k^2-k_2^2 \neq 0}} \frac{1}{i(k^2-k_2^2)} \left( e^{it(k^2-k_2^2)} - 1 \right) \hat{\xi}_{k_1}\widehat{(u_0)}_{k_2} \quad (4.13b)$$

$$+ i \sum_{k_1+k_2+k_3=k} \int_0^t e^{i\rho(k^2+k_1^2-k_2^2-k_3^2)} d\rho \widehat{(u_0)}_{k_1}\widehat{(u_0)}_{k_2}\widehat{(u_0)}_{k_3}. \quad (4.13c)$$

• **Lower and upper bounds on (4.13a).** By (4.11) and (4.12), we have that

$$\widehat{u_{0k}} = N^{-\gamma}, \text{ for } |k - N| \leq 10; \quad \widehat{u_{0k}} = 0, \text{ for } |k - N| > 10;$$

and

$$\hat{\xi}_k = \ln(|k|), \text{ for } |k| > 10.$$

This gives

$$\begin{aligned} \|\langle k \rangle^\gamma (4.13a)\|_{l_k^2}^2 &= t^2 \sum_k \langle k \rangle^{2\gamma} |\hat{\xi}_{2k}|^2 |\widehat{(u_0)}_{-k}|^2 \\ &= t^2 \sum_{k: |k-N| \leq 10} \langle k \rangle^{2\gamma} \ln(|k|)^2 N^{-2\gamma}. \end{aligned}$$

By choosing  $N$  large enough, we further get that

$$\frac{1}{2}t^2(\ln N)^2 \leq \|\langle k \rangle^\gamma (4.13a)\|_{l_k^2}^2 \leq 20t^2(\ln N)^2. \quad (4.14)$$

• **Upper bound on (4.13b).** We split (4.13b) into the following two parts:

$$(4.13b) = i \sum_{\substack{k_1+k_2=k \\ k^2-k_2^2 \neq 0, |k_2| \geq |k|}} \frac{1}{i(k^2-k_2^2)} \left( e^{it(k^2-k_2^2)} - 1 \right) \hat{\xi}_{k_1}\widehat{(u_0)}_{k_2} \quad (4.15a)$$

$$+ i \sum_{\substack{k_1+k_2=k \\ k^2-k_2^2 \neq 0, |k_2| \ll |k|}} \frac{1}{i(k^2-k_2^2)} \left( e^{it(k^2-k_2^2)} - 1 \right) \hat{\xi}_{k_1}\widehat{(u_0)}_{k_2}. \quad (4.15b)$$

For (4.15a), we first note that if  $k^2 - k_2^2 \neq 0$ , then

$$|k^2 - k_2^2| \geq |k|. \quad (4.16)$$

Indeed,  $|k^2 - k_2^2| = |k + k_2||k - k_2|$ . If  $k \cdot k_2 \geq 0$ , then  $|k + k_2| \geq |k|$  and  $|k - k_2| \geq 1$ , thus we have (4.16). If  $k \cdot k_2 < 0$ , then  $|k - k_2| \geq |k|$  and  $|k + k_2| \geq 1$ , thus we also have (4.16). Moreover, (4.16) combining with the trivial bound

$$|k^2 - k_2^2| = |k + k_2||k - k_2| \geq |k - k_2|,$$

give

$$|k^2 - k_2^2| \geq |k|^{\frac{3}{4}} |k - k_2|^{\frac{1}{4}}. \quad (4.17)$$

Therefore, by (4.11), (4.12) and (4.17), we have that

$$\begin{aligned}
\langle k \rangle^\gamma |(4.15a)| &= \left| \langle k \rangle^\gamma \sum_{\substack{k_1+k_2=k \\ k^2-k_2^2 \neq 0, |k_2| \gtrsim |k| \\ |k_2-N| \leq 10}} \frac{1}{i(k^2-k_2^2)} \left( e^{it(k^2-k_2^2)} - 1 \right) \ln(|k_1|) N^{-\gamma} \right| \\
&\leq \left| \sum_{\substack{k_1+k_2=k \\ k^2-k_2^2 \neq 0, |k_2| \gtrsim |k| \\ |k_2-N| \leq 10}} \frac{1}{i(k^2-k_2^2)} \left( e^{it(k^2-k_2^2)} - 1 \right) \ln(|k_1|) \right| \\
&\leq |k|^{-\frac{3}{4}} \sum_{k_2: |k_2-N| \leq 10} \ln(|k-k_2|) |k-k_2|^{-\frac{1}{4}} \lesssim |k|^{-\frac{3}{4}}.
\end{aligned}$$

This implies

$$\|\langle k \rangle^\gamma (4.15a)\|_{l_k^2} \lesssim \| |k|^{-\frac{3}{4}} \|_{l_k^2} \lesssim 1. \quad (4.18)$$

For (4.15b), under the restriction in the summation, we have  $|k^2-k_2^2| \gtrsim |k|^2$  and  $|k| \sim |k_1|$ . Therefore,

$$\begin{aligned}
\langle k \rangle^\gamma |(4.15b)| &= \left| \langle k \rangle^\gamma \sum_{\substack{k_1+k_2=k \\ k^2-k_2^2 \neq 0, |k_2| \ll |k| \\ |k_2-N| \leq 10}} \frac{1}{i(k^2-k_2^2)} \left( e^{it(k^2-k_2^2)} - 1 \right) \ln(|k_1|) N^{-\gamma} \right| \\
&\lesssim N^{-\gamma} |k|^{\gamma-2} \ln(|k|) \sum_{k_2: |k_2-N| \leq 10} 1 \lesssim N^{-\gamma} |k|^{\gamma-2} \ln(|k|).
\end{aligned}$$

Since  $\gamma < \frac{3}{2}$ , it yields

$$\|\langle k \rangle^\gamma (4.15b)\|_{l_k^2} \lesssim N^{-\gamma} \| |k|^{\gamma-2} \ln(|k|) \|_{l_k^2 \{k \neq 0\}} \lesssim N^{-\gamma}. \quad (4.19)$$

Together with the estimates on (4.18) and (4.19), we obtain

$$\|\langle k \rangle^\gamma (4.13b)\|_{l_k^2} \lesssim \max\{1, N^{-\gamma}\}. \quad (4.20)$$

• **Lower and upper bounds on (4.13c).** By the definition (4.11), we have

$$(4.13c) = iN^{-3\gamma} \sum_{\substack{k_1+k_2+k_3=k \\ |k_1+N| \leq 10, |k_j-N| \leq 10, j=2,3}} \int_0^t e^{i\rho(k^2+k_1^2-k_2^2-k_3^2)} d\rho.$$

Note that in the restriction:  $k_1+k_2+k_3=k$ ,  $|k_1+N| \leq 10$ ,  $|k_j-N| \leq 10$ ,  $j=2,3$ , we have

$$|k^2+k_1^2-k_2^2-k_3^2| = |k_1+k_2||k_1+k_3| \leq 100.$$

Setting  $t \in (0, 10^{-3})$ , then for any  $\rho \in (0, t)$ ,

$$\left| e^{i\rho(k^2+k_1^2-k_2^2-k_3^2)} - 1 \right| \leq \frac{1}{2}.$$

This gives that

$$\frac{1}{2} t N^{-3\gamma} \sum_{\substack{k_1+k_2+k_3=k \\ |k_1+N| \leq 10, |k_j-N| \leq 10, j=2,3}} 1 \leq |(4.13c)| \leq 2t N^{-3\gamma} \sum_{\substack{k_1+k_2+k_3=k \\ |k_1+N| \leq 10, |k_j-N| \leq 10, j=2,3}} 1.$$

For the lower bound, we use the embedding  $l^2 \hookrightarrow l^\infty$  and obtain that

$$\begin{aligned} \|\langle k \rangle^\gamma (4.13c)\|_{l_k^2} &\geq \|\langle k \rangle^\gamma (4.13c)\|_{l_k^\infty} \\ &\geq \frac{1}{2} t N^{-3\gamma} \left\| \langle k \rangle^\gamma \sum_{\substack{k_1+k_2+k_3=k \\ |k_1+N|\leq 10, |k_j-N|\leq 10, j=2,3}} 1 \right\|_{l_k^\infty}. \end{aligned}$$

Note that

$$\left\{ k : k = k_1 + k_2 + k_3, |k_1 + N| \leq 10, |k_j - N| \leq 10, j = 2, 3 \right\} \subset \left\{ k : |k - N| \leq 30 \right\}.$$

Choosing  $N$  large enough, we further have

$$\begin{aligned} \|\langle k \rangle^\gamma (4.13c)\|_{l_k^2} &\geq \frac{1}{4} t N^{-2\gamma} \left\| \sum_{\substack{k_1+k_2+k_3=k \\ |k_1+N|\leq 10, |k_j-N|\leq 10, j=2,3}} 1 \right\|_{l_k^\infty} \\ &\geq 5tN^{-2\gamma}. \end{aligned} \tag{4.21}$$

For the upper bound, we use the embedding  $l^1 \hookrightarrow l^2$  and obtain that

$$\begin{aligned} \|\langle k \rangle^\gamma (4.13c)\|_{l_k^2} &\leq \|\langle k \rangle^\gamma (4.13c)\|_{l_k^1} \\ &\leq 2tN^{-3\gamma} \left\| \langle k \rangle^\gamma \sum_{\substack{k_1+k_2+k_3=k \\ |k_1+N|\leq 10, |k_j-N|\leq 10, j=2,3}} 1 \right\|_{l_k^2(\{|k-N|\leq 30\})} \\ &\leq 4tN^{-2\gamma} \sum_{\substack{k_1, k_2, k_3 \\ |k_1+N|\leq 10, |k_j-N|\leq 10, j=2,3}} 1 \\ &\leq 120tN^{-2\gamma}. \end{aligned} \tag{4.22}$$

Together with (4.21) and (4.22), we obtain

$$5tN^{-2\gamma} \leq \|\langle k \rangle^\gamma (4.13c)\|_{l_k^2} \leq 120tN^{-2\gamma}. \tag{4.23}$$

Now we collect the estimates in (4.14), (4.20) and (4.23), and choose  $N$  large enough to obtain that for  $\gamma \geq 0$ ,

$$\|A_2(u_0)\|_{H^\gamma} = \left\| \langle k \rangle^\gamma (\widehat{A_2(u_0)})_k \right\|_{l_k^2} \geq \frac{\sqrt{2}}{2} t \ln N - C(tN^{-2\gamma} + 1) \geq \frac{1}{2} t \ln N - C;$$

for  $\gamma < 0$ ,

$$\|A_2(u_0)\|_{H^\gamma} = \left\| \langle k \rangle^\gamma (\widehat{A_2(u_0)})_k \right\|_{l_k^2} \geq 5tN^{-2\gamma} - C(t \ln N + N^{-\gamma}) \geq tN^{-2\gamma} - N^{-\gamma}.$$

By choosing  $t = 10^{-3}$ , for either  $\gamma \geq 0$  or  $\gamma < 0$ , we can have

$$\|A_2(u_0)\|_{H^\gamma} \rightarrow +\infty, \quad \text{when } N \rightarrow +\infty.$$

The proof is done by again applying [Lemma 2.3](#).  $\square$

## 5. NUMERICAL METHOD AND CONVERGENCE ANALYSIS

In this section, we shall first give the derivation of the presented LRI scheme (1.6), and then we shall prove its convergence result, i.e., [Theorem 1.6](#).

**5.1. Construction of numerical scheme.** By the Duhamel formula of (1.1) and the variable  $v(t) = e^{-it\partial_x^2}u(t)$ , we have that for any  $n \geq 0$ ,

$$v(t_{n+1}) = v(t_n) + i \int_{t_n}^{t_{n+1}} e^{-i\rho\partial_x^2} \left( \xi u(\rho) - \lambda |u(\rho)|^2 u(\rho) \right) d\rho. \quad (5.1)$$

Then we write

$$v(t_{n+1}) = i \int_{t_n}^{t_{n+1}} e^{-i\rho\partial_x^2} \left( \xi e^{i\rho\partial_x^2} v(t_n) \right) d\rho \quad (5.2)$$

$$+ v(t_n) - i\lambda \int_{t_n}^{t_{n+1}} e^{-i\rho\partial_x^2} \left( |u(\rho)|^2 u(\rho) \right) d\rho \quad (5.3)$$

$$+ R_1^n,$$

where

$$R_1^n \triangleq i \int_{t_n}^{t_{n+1}} e^{-i\rho\partial_x^2} \left[ \xi e^{i\rho\partial_x^2} (v(\rho) - v(t_n)) \right] d\rho \quad n \geq 0.$$

We first consider the integral term (5.2) involving the potential. By taking the Fourier transform, we have

$$\begin{aligned} \widehat{(5.2)}_k &= i \int_{t_n}^{t_{n+1}} \sum_{k_1+k_2=k} e^{i\rho(k^2-k_2^2)} \hat{\xi}_{k_1} \hat{v}_{k_2}(t_n) d\rho \\ &= i\tau e^{it_n k^2} \sum_{k_1+k_2=k} \mathcal{M}_\tau \left( e^{i\rho(k^2-k_2^2)} \right) \hat{\xi}_{k_1} e^{-it_n k_2^2} \hat{v}_{k_2}(t_n), \end{aligned}$$

where  $\mathcal{M}_\tau$  is the average operator given in (2.1). Then we consider the approximation  $\mathcal{M}_\tau \left( e^{i\rho(k^2-k_2^2)} \right) \approx \mathcal{M}_\tau \left( e^{i\rho k^2} \right) \mathcal{M}_\tau \left( e^{-i\rho k_2^2} \right)$ , and note that

$$\begin{aligned} &\sum_k e^{ikx} e^{it_n k^2} \sum_{k_1+k_2=k} \mathcal{M}_\tau \left( e^{i\rho k^2} \right) \mathcal{M}_\tau \left( e^{-i\rho k_2^2} \right) \hat{\xi}_{k_1} e^{-it_n k_2^2} \hat{v}_{k_2}(t_n) \\ &= e^{-it_n \partial_x^2} \mathcal{D}_{-\tau} \left[ \xi \mathcal{D}_\tau [e^{it_n \partial_x^2} v(t_n)] \right] = e^{-it_n \partial_x^2} \mathcal{D}_{-\tau} \left[ \xi \mathcal{D}_\tau [u(t_n)] \right]. \end{aligned}$$

So we can write the integral term (5.2) as

$$(5.2) = i\tau e^{-it_n \partial_x^2} \mathcal{D}_{-\tau} \left[ \xi \mathcal{D}_\tau [u(t_n)] \right] + R_2^n, \quad (5.4)$$

where  $R_2^n$  is the truncation term defined by

$$\widehat{(R_2^n)}_k \triangleq i\tau e^{it_n k^2} \sum_{k_1+k_2=k} \left[ \mathcal{M}_\tau \left( e^{i\rho(k^2-k_2^2)} \right) - \mathcal{M}_\tau \left( e^{i\rho k^2} \right) \mathcal{M}_\tau \left( e^{-i\rho k_2^2} \right) \right] \hat{\xi}_{k_1} e^{-it_n k_2^2} \hat{v}_{k_2}(t_n),$$

for  $n \geq 0$ . For the term (5.3), we simply use the Lie-Trotter splitting method and we denote the remainder as

$$R_3^n \triangleq (5.3) - e^{-it_{n+1} \partial_x^2} \mathcal{N}_\tau \left[ e^{i\tau \partial_x^2} u(t_n) \right], \quad n \geq 0. \quad (5.5)$$

Plugging (5.4), (5.5) back to (5.2) and (5.3) without the remainder terms  $R_2^n, R_3^n$ , and denoting  $v^n \approx v(t_n)$  yield:

$$v^{n+1} = i\tau e^{-it_n \partial_x^2} \mathcal{D}_{-\tau} \left[ \xi \mathcal{D}_\tau [e^{it_n \partial_x^2} v^n] \right] + e^{-it_{n+1} \partial_x^2} \mathcal{N}_\tau \left[ e^{i\tau \partial_x^2} e^{it_n \partial_x^2} v^n \right]. \quad (5.6)$$

Therefore, by letting  $u(t_n) \approx u^n = e^{it_n \partial_x^2} v^n$  with  $u^0 = v^0 = u_0$  and noting  $e^{i\tau \partial_x^2} \mathcal{D}_{-\tau} = \mathcal{D}_\tau$ , we obtain our LRI scheme presented in (1.6):

$$u^{n+1} = i\tau \mathcal{D}_\tau \left[ \xi \mathcal{D}_\tau [u^n] \right] + \mathcal{N}_\tau \left[ e^{i\tau \partial_x^2} u^n \right], \quad n \geq 0.$$

Note that the filter  $P_{\leq N}$  is an additional approximation applied in  $\mathcal{N}_\tau$  to cut off the high frequencies. We remark that such approximation is mainly used for the technical analysis of the convergence under  $L^2$ -norm [25, 26, 31]. It will not essentially affect the computational results in practice, and it could be omitted which will be addressed in a future work.

**5.2. Convergence analysis.** In the rest of this section, we assume  $\lambda = -1$  in (1.1) for simplicity of notations, and we aim to prove the error estimate given in Theorem 1.6.

Denote the part of the scheme (5.6) involving the integration of the potential as

$$\Phi(f) \triangleq f + i\tau e^{-it_n \partial_x^2} \mathcal{D}_{-\tau} \left[ \xi \mathcal{D}_\tau [e^{it_n \partial_x^2} f] \right],$$

and denote the splitting part involving the nonlinearity as

$$\Psi(f) \triangleq e^{-it_{n+1} \partial_x^2} \left( \mathcal{N}_\tau \left[ e^{i\tau \partial_x^2} e^{it_n \partial_x^2} f \right] - e^{i\tau \partial_x^2} e^{it_n \partial_x^2} f \right).$$

Then (5.6) reads

$$v^{n+1} = \Phi(v^n) + \Psi(v^n).$$

Denote the error function as

$$h^n \triangleq v^n - v(t_n), \quad n \geq 0,$$

so  $h^0 \equiv 0$  and we have

$$h^{n+1} = \Phi(v^n) - \Phi(v(t_n)) + \Psi(v^n) - \Psi(v(t_n)) + R_1^n + R_2^n + R_3^n.$$

We shall first analyze to give the stability result of the scheme (5.6) by working on the  $\Phi$  part and the  $\Psi$  part in a sequel. Afterwards, we shall estimate the local truncation errors  $R_1^n, R_2^n, R_3^n$ .

**Lemma 5.1** (Stability of potential part). *Let  $\xi \in \hat{L}^\infty$ , then for any  $f \in H^{\frac{1}{2}+}(\mathbb{T})$  and  $h = f - g \in L^2(\mathbb{T})$ ,*

$$\|\Phi(f) - \Phi(g)\|_{L^2} \leq (1 + C\tau) \|h\|_{L^2}, \quad (5.7)$$

where the constant  $C > 0$  depends only on  $\|\xi\|_{\hat{L}^\infty}$ .

*Proof.* Denote  $\tilde{h} = e^{it_n \partial_x^2} h$ , then  $\|\tilde{h}\|_{L^2} = \|h\|_{L^2}$ . Note that

$$\Phi(f) - \Phi(g) = e^{-it_n \partial_x^2} \left( \tilde{h} + i\tau \mathcal{D}_{-\tau} \left[ \xi \mathcal{D}_\tau \tilde{h} \right] \right).$$

Therefore,

$$\begin{aligned} \|\Phi(f) - \Phi(g)\|_{L^2}^2 &= \left\| \tilde{h} + i\tau \mathcal{D}_{-\tau} \left[ \xi \mathcal{D}_\tau \tilde{h} \right] \right\|_{L^2}^2 \\ &= \|h\|_{L^2}^2 + 2 \left\langle \tilde{h}, i\tau \mathcal{D}_{-\tau} \left[ \xi \mathcal{D}_\tau \tilde{h} \right] \right\rangle + \tau^2 \left\| \mathcal{D}_\tau \left[ \xi \mathcal{D}_\tau \tilde{h} \right] \right\|_{L^2}^2. \end{aligned}$$

where in the last step we have used the relationship:  $e^{i\tau \partial_x^2} \mathcal{D}_{-\tau} = \mathcal{D}_\tau$ .

A key observation (for real-valued  $\xi$ ) is that

$$\left\langle \tilde{h}, i\tau \mathcal{D}_{-\tau} \left[ \xi \mathcal{D}_\tau \tilde{h} \right] \right\rangle = \left\langle \mathcal{D}_\tau \tilde{h}, i\tau \xi \mathcal{D}_\tau \tilde{h} \right\rangle = \tau \operatorname{Im} \int_{\mathbb{T}} \xi |\mathcal{D}_\tau \tilde{h}|^2 dx = 0, \quad (5.8)$$

where in the second step we have used the relationship  $\mathcal{D}_{-\tau} = \overline{\mathcal{D}_\tau}$ .

Moreover, we claim that

$$\tau \left\| \mathcal{D}_\tau \left[ \xi \mathcal{D}_\tau \tilde{h} \right] \right\|_{L^2} \lesssim \sqrt{\tau} \|\xi\|_{\hat{L}^\infty} \|h\|_{L^2}. \quad (5.9)$$

To prove this claim, we take the Fourier transform and write

$$\mathcal{F}_k \left[ \mathcal{D}_\tau (\xi \mathcal{D}_\tau \tilde{h}) \right] = \sum_{k_1+k_2=k} \mathcal{M}_\tau(e^{isk^2}) \mathcal{M}_\tau(e^{-isk_2^2}) \hat{\xi}_{k_1} \hat{h}_{k_2}.$$

Now by duality and Plancherel's identity we have

$$\left\| \mathcal{D}_\tau \left[ \xi \mathcal{D}_\tau \tilde{h} \right] \right\|_{L^2} = \sup_{w: \|w\|_{l^2}=1} \left\langle \sum_{k_1+k_2=k} \mathcal{M}_\tau(e^{isk^2}) \mathcal{M}_\tau(e^{-isk_2^2}) \hat{\xi}_{k_1} \hat{h}_{k_2}, w \right\rangle.$$

As before, we omit  $\sup_{w: \|w\|_{l^2}=1}$  in the front. Then we split it into the following three cases and further write

$$\begin{aligned} \left\| \mathcal{D}_\tau \left[ \xi \mathcal{D}_\tau \tilde{h} \right] \right\|_{L^2} &= \sum_{\substack{k_1+k_2=k \\ |k| \leq N_0, |k_2| \leq N_0}} \mathcal{M}_\tau(e^{isk^2}) \mathcal{M}_\tau(e^{-isk_2^2}) \hat{\xi}_{k_1} \hat{h}_{k_2} w_k \\ &\quad + \sum_{\substack{k_1+k_2=k \\ |k| \leq N_0, |k_2| \geq N_0}} \mathcal{M}_\tau(e^{isk^2}) \mathcal{M}_\tau(e^{-isk_2^2}) \hat{\xi}_{k_1} \hat{h}_{k_2} w_k \\ &\quad + \sum_{\substack{k_1+k_2=k \\ |k| \geq N_0, |k_2| \geq N_0}} \mathcal{M}_\tau(e^{isk^2}) \mathcal{M}_\tau(e^{-isk_2^2}) \hat{\xi}_{k_1} \hat{h}_{k_2} w_k, \end{aligned}$$

where  $N_0 > 0$  will be determined later. Note that

$$|\mathcal{M}_\tau(e^{isk^2})| \lesssim \min\{1, \tau^{-1}|k|^{-2}\},$$

therefore,

$$\begin{aligned} \left\| \mathcal{D}_\tau \left[ \xi \mathcal{D}_\tau \tilde{h} \right] \right\|_{L^2} &\lesssim \|\xi\|_{\hat{L}^\infty} \left( \sum_{\substack{k, k_2 \\ |k| \leq N_0, |k_2| \leq N_0}} |\hat{h}_{k_2}| |w_k| + \sum_{\substack{k, k_2 \\ |k| \leq N_0, |k_2| \geq N_0}} \tau^{-1} |k_2|^{-2} |\hat{h}_{k_2}| |w_k| \right. \\ &\quad \left. + \sum_{\substack{k, k_2 \\ |k| \geq N_0, |k_2| \geq N_0}} \tau^{-2} |k|^{-2} |k_2|^{-2} |\hat{h}_{k_2}| \right) |w_k|. \end{aligned}$$

This gives that

$$\begin{aligned} \left\| \mathcal{D}_\tau \left[ \xi \mathcal{D}_\tau \tilde{h} \right] \right\|_{L^2} &\lesssim \|\xi\|_{\hat{L}^\infty} \left( N_0 + \tau^{-1} N_0^{-1} + \tau^{-2} N_0^{-3} \right) \|\hat{h}_k\|_{l_k^2} \|w_k\|_{l_k^2} \\ &\lesssim \|\xi\|_{\hat{L}^\infty} \left( N_0 + \tau^{-1} N_0^{-1} + \tau^{-2} N_0^{-3} \right) \|h\|_{L^2}. \end{aligned}$$

Choosing  $N_0 = \tau^{-\frac{1}{2}}$ , we obtain (5.9).

Together with (5.8) and (5.9), we have

$$\|\Phi(f) - \Phi(g)\|_{L^2}^2 \leq (1 + C\tau) \|h\|_{L^2}^2,$$

and thus

$$\|\Phi(f) - \Phi(g)\|_{L^2} \leq (1 + C\tau) \|h\|_{L^2}.$$

This finishes the proof of the lemma.  $\square$

**Lemma 5.2** (Stability of splitting part). *Let  $f \in L^2(\mathbb{T})$  and  $g \in H^{\frac{1}{2}+}(\mathbb{T})$ , then*

$$\|\Psi(f) - \Psi(g)\|_{L^2} \leq C\tau (1 + N\|f - g\|_{L^2}^2) \|f - g\|_{L^2}, \quad (5.10)$$

for some constant  $C > 0$  that depends on  $\|g\|_{H^{\frac{1}{2}+}}$ .

*Proof.* For simplicity, we denote

$$\tilde{f} = e^{i\tau\partial_x^2} e^{it_n\partial_x^2} f, \quad \tilde{g} = e^{i\tau\partial_x^2} e^{it_n\partial_x^2} g, \quad \tilde{h} = e^{i\tau\partial_x^2} e^{it_n\partial_x^2} h, \quad h = f - g.$$

Then

$$\Psi(f) - \Psi(g) = e^{-it_{n+1}\partial_x^2} \left( \mathcal{N}_\tau[\tilde{f}] - \mathcal{N}_\tau[\tilde{g}] - \tilde{h} \right).$$

We rewrite

$$\begin{aligned} \mathcal{N}_\tau[\tilde{f}] - \mathcal{N}_\tau[\tilde{g}] &= e^{i\tau|P_{\leq N}\tilde{f}|^2} \tilde{f} - e^{i\tau|P_{\leq N}\tilde{g}|^2} \tilde{g} \\ &= e^{i\tau|P_{\leq N}\tilde{f}|^2} \tilde{h} + \left[ e^{i\tau|P_{\leq N}\tilde{f}|^2} - e^{i\tau|P_{\leq N}\tilde{g}|^2} \right] \tilde{g}. \end{aligned}$$

Therefore,

$$\Psi(f) - \Psi(g) = e^{-it_{n+1}\partial_x^2} \left[ \left( e^{i\tau|P_{\leq N}\tilde{f}|^2} - 1 \right) \tilde{h} + \left( e^{i\tau|P_{\leq N}\tilde{f}|^2} - e^{i\tau|P_{\leq N}\tilde{g}|^2} \right) \tilde{g} \right].$$

Then by the Sobolev and Bernstein inequalities, we have

$$\begin{aligned} \|\Psi(f) - \Psi(g)\|_{L^2} &\lesssim \left\| \left[ e^{i\tau|P_{\leq N}\tilde{f}|^2} - 1 \right] \tilde{h} \right\|_{L^2} + \left\| \left[ e^{i\tau|P_{\leq N}\tilde{f}|^2} - e^{i\tau|P_{\leq N}\tilde{g}|^2} \right] \tilde{g} \right\|_{L^2} \\ &\lesssim \left\| e^{i\tau|P_{\leq N}\tilde{f}|^2} - 1 \right\|_{L^\infty} \|\tilde{h}\|_{L^2} + \left\| e^{i\tau|P_{\leq N}\tilde{f}|^2} - e^{i\tau|P_{\leq N}\tilde{g}|^2} \right\|_{L^2} \|\tilde{g}\|_{L^\infty}. \end{aligned}$$

Note that

$$\left\| e^{i\tau|P_{\leq N}\tilde{f}|^2} - 1 \right\|_{L^\infty} \lesssim \left\| i\tau|P_{\leq N}\tilde{f}|^2 \right\|_{L^\infty} = \tau \|P_{\leq N}\tilde{f}\|_{L^\infty}^2 \lesssim \tau N \|h\|_{L^2}^2 + \tau \|g\|_{H^{\frac{1}{2}+}}^2.$$

Similarly, we have

$$\begin{aligned} \left\| e^{i\tau|P_{\leq N}\tilde{f}|^2} - e^{i\tau|P_{\leq N}\tilde{g}|^2} \right\|_{L^2} &\lesssim \tau \left\| |P_{\leq N}\tilde{f}|^2 - |P_{\leq N}\tilde{g}|^2 \right\|_{L^2} \\ &\lesssim \tau \|P_{\leq N}\tilde{h}\|_{L^2} (\|P_{\leq N}\tilde{f}\|_{L^\infty} + \|P_{\leq N}\tilde{g}\|_{L^\infty}) \\ &\lesssim \tau N^{\frac{1}{2}} \|h\|_{L^2}^2 + \tau \|g\|_{H^{\frac{1}{2}+}} \|h\|_{L^2}. \end{aligned}$$

Applying these two estimates, we further have

$$\begin{aligned} \|\Psi(f) - \Psi(g)\|_{L^2} &\leq C\tau \|h\|_{L^2} + C\tau N^{\frac{1}{2}} \|h\|_{L^2}^2 + C\tau N \|h\|_{L^2}^3 \\ &\leq C\tau \|h\|_{L^2} + C\tau N \|h\|_{L^2}^3, \end{aligned}$$

where the constant  $C > 0$  depends on  $\|g\|_{H^{\frac{1}{2}+}}$ . Hence, we obtain the desired assertion.  $\square$

Now, we move on to estimating the local truncation errors  $R_1^n, R_2^n, R_3^n$  one by one. To estimate  $R_1^n$ , we need the following result.

**Lemma 5.3.** *Let  $\xi \in \hat{b}^{s,p}$  with the conditions of [Theorem 1.6](#) satisfied, so  $u \in L_t^\infty H_x^{s+\gamma_p-}([0, T] \times \mathbb{T})$  with  $\gamma_p = \frac{3}{2} + \frac{1}{p}$ . Then, for any  $0 \leq \beta < s + \frac{1}{p} + \frac{3}{2}$  we have*

$$\|v(t) - v(t_n)\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} \leq C\tau \max \left\{ \tau^{-\frac{1}{2}(\beta - s - \frac{1}{p}) - \frac{1}{4}-}, 1 \right\}, \quad (5.11)$$

where the constant  $C > 0$  depends only on  $\|\xi\|_{\hat{b}^{s,p}}$  and  $\|u\|_{L_t^\infty H_x^{s+\gamma_p-}}$ .

*Proof.* By  $\|v(t) - v(t_n)\|_{H^{s+\gamma_p-}} \leq 2\|v\|_{L_t^\infty H_x^{s+\gamma_p-}}$ , we can deduce that for any  $\beta < s + \gamma_p$ ,

$$\begin{aligned} \|v(t) - v(t_n)\|_{H^\beta} &= \|P_{\leq N_0}(v(t) - v(t_n))\|_{H^\beta} + \|P_{> N_0}(v(t) - v(t_n))\|_{H^\beta} \\ &\lesssim \|P_{\leq N_0}(v(t) - v(t_n))\|_{H^\beta} + N_0^{-s-\gamma_p+\beta+} \|v\|_{L_t^\infty H_x^{s+\gamma_p-}}, \end{aligned} \quad (5.12)$$

where  $N_0 > 0$  will be determined later. Hence, in the following we only need to estimate  $\|P_{\leq N_0}(v(t) - v(t_n))\|_{H^\beta}$ .

By (3.1), we have that

$$\begin{aligned} v(t) - v(t_n) &= i \int_{t_n}^t e^{-i\rho\partial_x^2} \left[ \xi e^{i\rho\partial_x^2} v(\rho) \right] d\rho + i \int_{t_n}^t e^{-i\rho\partial_x^2} |u(\rho)|^2 u(\rho) d\rho \\ &\triangleq I_1 + I_2. \end{aligned}$$

Since  $s + \gamma_p > \frac{1}{2}$ , we can directly have

$$\|I_2\|_{L_t^\infty H_x^{s+\gamma_p-}([t_n, t_{n+1}])} \lesssim \tau \|u\|_{H^{s+\gamma_p-}}^3. \quad (5.13)$$

Then it remains to estimate  $P_{\leq N_0}I_1$ . For  $I_1$ , we take the Fourier transform to have

$$\begin{aligned} (\widehat{I_1})_k &= i \int_{t_n}^t \sum_{k=k_1+k_2} e^{i\rho(k^2-k_2^2)} \hat{\xi}_{k_1} \hat{v}_{k_2}(\rho) d\rho \\ &= i \int_{t_n}^t \sum_{\substack{k_1+k_2=k \\ |k_1| \lesssim |k_2|}} e^{i\rho(k^2-k_2^2)} \hat{\xi}_{k_1} \hat{v}_{k_2}(\rho) d\rho + i \int_{t_n}^t \sum_{\substack{k_1+k_2=k \\ |k_1| \gg |k_2|}} e^{i\rho(k^2-k_2^2)} \hat{\xi}_{k_1} \hat{v}_{k_2}(\rho) d\rho \\ &\triangleq (\widehat{I_{11}})_k + (\widehat{I_{12}})_k. \end{aligned}$$

We shall estimate  $P_{\leq N_0}I_{11}$  and  $P_{\leq N_0}I_{12}$  in a sequel.

For  $P_{\leq N_0}I_{11}$ , by duality and Plancherel's identity we have

$$\begin{aligned} &\|P_{\leq N_0}I_{11}\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} \\ &\lesssim \mathcal{T} \left\| \sup_{h: \|h\|_{l^2}=1} \sum_{\substack{|k_1| \lesssim |k_2| \\ |k_1+k_2| \leq N_0}} |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(t)| |h_{k_1+k_2}| \cdot \langle k_1+k_2 \rangle^\beta \right\|_{L_t^\infty([t_n, t_{n+1}])} \\ &\lesssim \mathcal{T} \left\| \sup_{h: \|h\|_{l^2}=1} \sum_{\substack{|k_1| \lesssim |k_2| \\ |k_1+k_2| \leq N_0}} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle k_2 \rangle^{s+\gamma_p-} |\hat{v}_{k_2}(t)| |h_{k_1+k_2}| \langle k_1 \rangle^{-s} \right. \\ &\quad \left. \cdot \langle k_1+k_2 \rangle^\beta \langle k_2 \rangle^{-s-\gamma_p+} \right\|_{L_t^\infty([t_n, t_{n+1}])} \\ &\lesssim \mathcal{T} \left\| \sup_{h: \|h\|_{l^2}=1} \sum_{\substack{|k_1| \lesssim |k_2| \\ |k_1+k_2| \leq N_0}} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle k_2 \rangle^{s+\gamma_p-} |\hat{v}_{k_2}(t)| |h_{k_1+k_2}| \langle k_1 \rangle^{-s} \right. \\ &\quad \left. \cdot \min\{N_0^\beta, \langle k_2 \rangle^\beta\} \langle k_2 \rangle^{-s-\gamma_p+} \right\|_{L_t^\infty([t_n, t_{n+1}])}. \end{aligned}$$

Denote  $\frac{1}{r} = \frac{1}{2} - \frac{1}{p}$ .

If  $s \leq \frac{1}{r}$ , then our strategy is to first sum up  $k_1$  and then sum up  $k_2$ , and by Cauchy-Schwartz's inequality, we further have

$$\begin{aligned}
& \left\| P_{\leq N_0} I_{11} \right\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} \\
& \lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \left\| \sum_{k_2} \left\| \langle k_1 \rangle^{-s} \right\|_{l^r(\{k_1: |k_1| \lesssim |k_2|\})} \min \left\{ N_0^\beta, \langle k_2 \rangle^\beta \right\} \langle k_2 \rangle^{-s-\gamma_p+} \right. \\
& \quad \left. \cdot \langle k_2 \rangle^{s+\gamma_p-} |\hat{v}_{k_2}(t)| \right\|_{L_t^\infty([t_n, t_{n+1}])} \\
& \lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \left\| \sum_{k_2} \langle k_2 \rangle^{-2s-\gamma_p+\frac{1}{r}+} \min \left\{ N_0^\beta, \langle k_2 \rangle^\beta \right\} \langle k_2 \rangle^{s+\gamma_p-} |\hat{v}_{k_2}(t)| \right\|_{L_t^\infty([t_n, t_{n+1}])} \\
& \lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \|v\|_{L_t^\infty H_x^{s+\gamma_p-}} \left\| \langle k_2 \rangle^{-2s-\gamma_p+\frac{1}{r}+} \min \left\{ N_0^\beta, \langle k_2 \rangle^\beta \right\} \right\|_{l_{k_2}^2}.
\end{aligned}$$

Noticing that

$$\left\| \langle k_2 \rangle^{-2s-\gamma_p+\frac{1}{r}+} \min \left\{ N_0^\beta, \langle k_2 \rangle^\beta \right\} \right\|_{l_{k_2}^2} \lesssim \max \left\{ N_0^{-2s-\frac{2}{p}+\beta-\frac{1}{2}+}, 1 \right\},$$

we have

$$\left\| P_{\leq N_0} I_{11} \right\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} \lesssim \tau \max \left\{ N_0^{-2s-\frac{2}{p}+\beta-\frac{1}{2}+}, 1 \right\} \|\xi\|_{\dot{b}^{s,p}} \|v\|_{L_t^\infty H_x^{s+\gamma_p-}}. \quad (5.14)$$

If  $s > \frac{1}{r}$ , then our strategy is to first sum up  $k_2$  and then sum up  $k_1$ , and by Cauchy-Schwartz's inequality, we have

$$\begin{aligned}
& \left\| P_{\leq N_0} I_{11} \right\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} \\
& \lesssim \tau \|v\|_{L_t^\infty H_x^{s+\gamma_p-}} \sum_{k_1} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle k_1 \rangle^{-s} \left\| \min \left\{ N_0^\beta, \langle k_2 \rangle^\beta \right\} \langle k_2 \rangle^{-s-\gamma_p+} \right\|_{l_{\{k_2: |k_1| \lesssim |k_2|\}}^\infty}.
\end{aligned}$$

Furthermore if  $s + \frac{1}{p} < 1$ , we can apply the following estimate:

$$\left\| \min \left\{ N_0^\beta, \langle k_2 \rangle^\beta \right\} \langle k_2 \rangle^{-s-\gamma_p+} \right\|_{l_{\{k_2: |k_1| \lesssim |k_2|\}}^\infty} \lesssim 1,$$

where we have used the relation:  $\beta < s + \gamma_p$ , and then we can obtain

$$\begin{aligned}
\left\| P_{\leq N_0} I_{11} \right\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} & \lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \|v\|_{L_t^\infty H_x^{s+\gamma_p-}} \left\| \langle k_1 \rangle^{-s} \right\|_{l_{k_1}^{p'}} \\
& \lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \|v\|_{L_t^\infty H_x^{s+\gamma_p-}}.
\end{aligned}$$

Otherwise for  $s + \frac{1}{p} \geq 1$ , we have

$$\left\| \min \left\{ N_0^\beta, \langle k_2 \rangle^\beta \right\} \langle k_2 \rangle^{-s-\gamma_p+} \right\|_{l_{\{k_2: |k_1| \lesssim |k_2|\}}^\infty} \lesssim \langle k_1 \rangle^{s+\frac{1}{p'}-1-} \max \left\{ N_0^{-2s-\frac{2}{p}+\beta-\frac{1}{2}+}, 1 \right\},$$

and we can find

$$\begin{aligned}
& \left\| P_{\leq N_0} I_{11} \right\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} \\
& \lesssim \tau \max \left\{ N_0^{-2s-\frac{2}{p}+\beta-\frac{1}{2}+}, 1 \right\} \|\xi\|_{\dot{b}^{s,p}} \|v\|_{L_t^\infty H_x^{s+\gamma_p-}} \left\| \langle k_1 \rangle^{-1-} \right\|_{l_{k_1}^{p'}} \\
& \lesssim \tau \max \left\{ N_0^{-2s-\frac{2}{p}+\beta-\frac{1}{2}+}, 1 \right\} \|\xi\|_{\dot{b}^{s,p}} \|v\|_{L_t^\infty H_x^{s+\gamma_p-}}.
\end{aligned}$$

Therefore, we obtain that for  $s > \frac{1}{r}$ ,

$$\|P_{\leq N_0} I_{11}\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} \lesssim \tau \max \left\{ N_0^{-2s - \frac{2}{p} + \beta - \frac{1}{2} +}, 1 \right\} \|\xi\|_{\dot{b}^{s,p}} \|v\|_{L_t^\infty H_x^{s+\gamma p^-}}.$$

This combining with (5.14) gives that for either  $s \leq \frac{1}{r}$  or  $s > \frac{1}{r}$ ,

$$\|P_{\leq N_0} I_{11}\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} \lesssim \tau \max \left\{ N_0^{-2s - \frac{2}{p} + \beta - \frac{1}{2} +}, 1 \right\} \|\xi\|_{\dot{b}^{s,p}} \|v\|_{L_t^\infty H_x^{s+\gamma p^-}}. \quad (5.15)$$

For  $P_{\leq N_0} I_{12}$ , we have

$$\begin{aligned} & \|P_{\leq N_0} I_{12}\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} \\ & \lesssim \tau \left\| \sup_{h: \|h\|_{l^2} = 1} \sum_{\substack{|k_1| \gg |k_2| \\ |k_1 + k_2| \leq N_0}} |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(t)| |h_{k_1+k_2}| \langle k_1 + k_2 \rangle^\beta \right\|_{L_t^\infty([t_n, t_{n+1}])} \\ & \lesssim \tau \left\| \sup_{h: \|h\|_{l^2} = 1} \sum_{\substack{|k_1| \gg |k_2| \\ |k_1 + k_2| \leq N_0}} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(t)| |h_{k_1+k_2}| \cdot \langle k_1 \rangle^{\beta-s} \right\|_{L_t^\infty([t_n, t_{n+1}])} \\ & \leq \tau \|\xi\|_{\dot{b}^{s,p}} \left\| \sum_{k_2} \|\langle k_1 \rangle^{\beta-s}\|_{l_{\{k_1: |k_1| \lesssim N_0\}}^r} |\hat{v}_{k_2}(t)| \right\|_{L_t^\infty([t_n, t_{n+1}])}. \end{aligned} \quad (5.16)$$

Noticing

$$\|\langle k_1 \rangle^{\beta-s}\|_{l_{\{k_1: |k_1| \lesssim N_0\}}^r} \lesssim \begin{cases} N_0^{\beta-s+1/r+}, & \beta - s + 1/r \geq 0, \\ 1, & \beta - s + 1/r < 0, \end{cases}$$

(5.16) further gives

$$\begin{aligned} \|P_{\leq N_0} I_{12}\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} & \lesssim \tau \max \{ N_0^{\beta-s+\frac{1}{r}+}, 1 \} \|\xi\|_{\dot{b}^{s,p}} \left\| \sum_{k_2} |\hat{v}_{k_2}(t)| \right\|_{L_t^\infty([t_n, t_{n+1}])} \\ & \lesssim \tau \max \left\{ N_0^{-s-\frac{1}{p}+\beta+\frac{1}{2}+}, 1 \right\} \|\xi\|_{\dot{b}^{s,p}} \|v\|_{L_t^\infty H_x^{s+\gamma p^-}}, \end{aligned} \quad (5.17)$$

where we have used  $s + \gamma p > \frac{1}{2}$  in the last step.

Now (5.13) together with (5.15) and (5.17) lead us to

$$\|P_{\leq N_0}(v(t) - v(t_n))\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} \leq C \tau \max \left\{ N_0^{-2s - \frac{2}{p} + \beta - \frac{1}{2} +}, N_0^{-s - \frac{1}{p} + \beta + \frac{1}{2} +}, 1 \right\}.$$

Noting that  $-2s - \frac{2}{p} + \beta - \frac{1}{2} < -s - \frac{1}{p} + \beta + \frac{1}{2}$ , we have

$$\|P_{\leq N_0}(v(t) - v(t_n))\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} \leq C \tau \max \left\{ N_0^{-s - \frac{1}{p} + \beta + \frac{1}{2} +}, 1 \right\}.$$

This combining with (5.12) give

$$\|v(t) - v(t_n)\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} \leq C \left( \tau \max \left\{ N_0^{-s - \frac{1}{p} + \beta + \frac{1}{2} +}, 1 \right\} + N_0^{-s - \frac{1}{p} + \beta - \frac{3}{2} +} \right).$$

For simplicity, we denote  $A = -s - \frac{1}{p} + \beta$  and set  $N_0 = \tau^{-\frac{1}{2}}$ , then it follows that

$$\|v(t) - v(t_n)\|_{L_t^\infty H_x^\beta([t_n, t_{n+1}] \times \mathbb{T})} \leq C \tau \max \left\{ \tau^{-\frac{A}{2} - \frac{1}{4}}, 1 \right\},$$

which finishes the proof of the lemma.  $\square$

**Lemma 5.4** (Local error from  $R_1^n$ ). *Let  $\xi \in \hat{b}^{s,p}$  with the conditions of [Theorem 1.6](#) satisfied, so  $u, v \in L_t^\infty H_x^{s+\gamma p^-}([0, T])$ . Then, we have*

$$\|R_1^n\|_{L^2} \leq C\tau^{1+\min\{s+\frac{1}{p}+\frac{1}{4}, 1\}}, \quad (5.18)$$

where the constant  $C > 0$  only depends on  $\|\xi\|_{\hat{b}^{s,p}}$  and  $\|u\|_{L_t^\infty H_x^{s+\gamma p^-}}$ .

*Proof.* The analysis goes separately for the case  $s + \frac{1}{p} > 1$  and the case  $s + \frac{1}{p} \leq 1$ .

*Case I:*  $s + \frac{1}{p} > 1$ . Firstly, we consider the simpler case:  $s + \frac{1}{p} > 1$ . In this case, by Cauchy-Schwartz's inequality, we have

$$\|\xi\|_{L^\infty} \leq \|\hat{\xi}\|_{l^1} \leq \|\langle k \rangle^{-s}\|_{l^{p'}} \|\xi\|_{\hat{b}^{s,p}} \lesssim \|\xi\|_{\hat{b}^{s,p}}.$$

By this inequality, we find that

$$\begin{aligned} \|R_1^n\|_{L^2} &\lesssim \int_{t_n}^{t_{n+1}} \left\| e^{-i\rho\partial_x^2} \left[ \xi e^{i\rho\partial_x^2} (v(\rho) - v(t_n)) \right] \right\|_{L^2} d\rho \\ &\lesssim \int_{t_n}^{t_{n+1}} \left\| \xi e^{i\rho\partial_x^2} (v(\rho) - v(t_n)) \right\|_{L^2} d\rho \\ &\lesssim \tau \|\xi\|_{L^\infty} \|v(\rho) - v(t_n)\|_{L_t^\infty L_x^2([t_n, t_{n+1}])} \\ &\lesssim \tau \|\xi\|_{\hat{b}^{s,p}} \|v(\rho) - v(t_n)\|_{L_t^\infty L_x^2([t_n, t_{n+1}])}. \end{aligned}$$

By [Lemma 5.3](#), it infers that

$$\|v(\rho) - v(t_n)\|_{L_t^\infty L_x^2([t_n, t_{n+1}] \times \mathbb{T})} \leq C\tau,$$

and this further yields

$$\|R_1^n\|_{L^2} \leq C\tau^2.$$

This gives the desired estimate of the lemma.

*Case II:*  $s + \frac{1}{p} \leq 1$ . Now, we turn to consider the case:  $s + \frac{1}{p} \leq 1$ . By taking the Fourier transform, we write

$$\begin{aligned} \widehat{(R_1^n)}_k &= i \int_{t_n}^{t_{n+1}} \sum_{k_1+k_2=k} e^{i\rho(k^2-k_2^2)} \hat{\xi}_{k_1} (\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)) d\rho \\ &= i \int_{t_n}^{t_{n+1}} \sum_{\substack{k_1+k_2=k \\ |k_1| \lesssim |k_2|}} e^{i\rho(k^2-k_2^2)} \hat{\xi}_{k_1} (\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)) d\rho \\ &\quad + i \int_{t_n}^{t_{n+1}} \sum_{\substack{k_1+k_2=k \\ |k_1| \gg |k_2|}} e^{i\rho(k^2-k_2^2)} \hat{\xi}_{k_1} (\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)) d\rho \\ &\triangleq \widehat{(R_{11})}_k + \widehat{(R_{12})}_k. \end{aligned}$$

For  $R_{11}$ , by duality and Plancherel's identity we have

$$\begin{aligned} \|R_{11}\|_{L^2} &\lesssim \tau \left\| \sum_{\substack{k=k_1+k_2 \\ |k_1| \lesssim |k_2|}} |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)| \right\|_{L_t^\infty l_k^\infty([t_n, t_{n+1}])} \\ &\lesssim \tau \left\| \sup_{h: \|h\|_{l^2}=1} \sum_{\substack{k=k_1+k_2 \\ |k_1| \lesssim |k_2|}} |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)| |h_{k_1+k_2}| \right\|_{L_t^\infty([t_n, t_{n+1}])}. \end{aligned}$$

As before, we denote  $\frac{1}{r} = \frac{1}{2} - \frac{1}{p}$  and  $a = -s + \frac{1}{r} + \frac{1}{2} + = -s - \frac{1}{p} + 1 +$ . If  $s \leq \frac{1}{r}$ , the strategy here is to first sum up  $k_1$  and then sum up  $k_2$ . This gives

$$\begin{aligned} \|R_{11}\|_{L^2} &\lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \left\| \sum_{k_2} \|\langle k_1 \rangle^{-s}\|_{l^r(\{k_1: |k_1| \lesssim |k_2|\})} |\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)| \right\|_{L_t^\infty([t_n, t_{n+1}])} \\ &\lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \left\| \sum_{k_2} \langle k_2 \rangle^{-s+\frac{1}{r}+} |\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)| \right\|_{L_t^\infty([t_n, t_{n+1}])} \\ &\lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \|v(t) - v(t_n)\|_{L_t^\infty H_x^a([t_n, t_{n+1}])}. \end{aligned} \quad (5.19)$$

If  $s > \frac{1}{r}$ , now our strategy is to first sum up  $k_2$  and then sum up  $k_1$ . This gives

$$\begin{aligned} \|R_{11}\|_{L^2} &\lesssim \tau \left\| \sup_{h: \|h\|_{l^2}=1} \sum_{\substack{k=k_1+k_2 \\ |k_1| \lesssim |k_2|}} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle k_2 \rangle^a |\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)| |h_{k_1+k_2}| \langle k_1 \rangle^{-s} \langle k_2 \rangle^{-a} \right\|_{L_t^\infty} \\ &\lesssim \tau \|v(t) - v(t_n)\|_{L_t^\infty H_x^a([t_n, t_{n+1}] \times \mathbb{T})} \sum_{k_1} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle k_1 \rangle^{-s-a} \\ &\lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \|v(t) - v(t_n)\|_{L_t^\infty H_x^a([t_n, t_{n+1}] \times \mathbb{T})} \left\| \langle k_1 \rangle^{-s-a} \right\|_{l_{k_1}^{p'}}. \end{aligned}$$

Since  $s > \frac{1}{r}$ , we have that  $s + a > \frac{1}{p'}$ , and thus it gives

$$\|R_{11}\|_{L^2} \lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \|v(t) - v(t_n)\|_{L_t^\infty H_x^a([t_n, t_{n+1}] \times \mathbb{T})}.$$

Combining with (5.19), we have that for either  $s \leq \frac{1}{r}$  or  $s > \frac{1}{r}$ ,

$$\|R_{11}\|_{L^2} \lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \|v(t) - v(t_n)\|_{L_t^\infty H_x^a([t_n, t_{n+1}] \times \mathbb{T})}. \quad (5.20)$$

For  $R_{12}$ , we first consider the low-frequency part to have

$$\begin{aligned} \|P_{\leq N_0} R_{12}\|_{L^2} &\lesssim \tau \left\| \sum_{\substack{k=k_1+k_2 \\ |k_1| \gg |k_2|}} |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)| \right\|_{L_t^\infty l_k^2([t_n, t_{n+1}] \times \{k: |k| \leq N_0\})} \\ &\lesssim \tau \left\| \sup_{h: \|h\|_{l^2}=1} \sum_{\substack{|k_1| \gg |k_2| \\ |k_1+k_2| \leq N_0}} |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)| |h_{k_1+k_2}| \right\|_{L_t^\infty([t_n, t_{n+1}])} \\ &\lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \left\| \sum_{k_2} \|\langle k_1 \rangle^{-s}\|_{l_{k_1}^r(\{k_1: |k_2| \ll |k_1| \lesssim N_0\})} |\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)| \right\|_{L_t^\infty}. \end{aligned}$$

Notice that

$$\|\langle k_1 \rangle^{-s}\|_{l_{k_1}^r(\{k_1: |k_2| \ll |k_1| \lesssim N_0\})} \lesssim \begin{cases} N_0^{-s+1/r+}, & -s + 1/r \geq 0, \\ \langle k_2 \rangle^{-s+1/r}, & -s + 1/r < 0. \end{cases}$$

When  $s + \frac{1}{p} \leq \frac{1}{2} \Leftrightarrow s \leq \frac{1}{r}$ , we can then have

$$\begin{aligned} \|P_{\leq N_0} R_{12}\|_{L^2} &\lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \left\| \sum_{k_2} |\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)| \right\|_{L_t^\infty} \\ &\lesssim \tau N_0^{-s+\frac{1}{r}+} \|\xi\|_{\dot{b}^{s,p}} \|v(\rho) - v(t_n)\|_{L_t^\infty H_x^{\frac{1}{2}+}}. \end{aligned} \quad (5.21)$$

When  $s + \frac{1}{p} > \frac{1}{2} \Leftrightarrow s > \frac{1}{r}$ , noting that  $a = -s + \frac{1}{r} + \frac{1}{2} +$ , we then find

$$\begin{aligned} \|P_{\leq N_0} R_{12}\|_{L^2} &\lesssim \tau \|\xi\|_{\hat{b}^{s,p}} \left\| \sum_{k_2} \langle k_2 \rangle^{-s + \frac{1}{r}} |\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)| \right\|_{L_t^\infty} \\ &\lesssim \tau \|\xi\|_{\hat{b}^{s,p}} \|v(\rho) - v(t_n)\|_{L_t^\infty H_x^a}. \end{aligned} \quad (5.22)$$

Now we consider the high-frequency part of  $R_{12}$ . By integration-by-parts we have

$$\begin{aligned} (\widehat{R_{12}})_k &= i \sum_{\substack{k=k_1+k_2 \\ |k_1| \gg |k_2|}} \frac{e^{it_{n+1}(k^2-k_2^2)}}{i(k^2-k_2^2)} \hat{\xi}_{k_1} [\hat{v}_{k_2}(t_{n+1}) - \hat{v}_{k_2}(t_n)] \\ &\quad - i \int_{t_n}^{t_{n+1}} \sum_{\substack{k_1+k_2=k \\ |k_1| \gg |k_2|}} \frac{e^{i\rho(k^2-k_2^2)}}{i(k^2-k_2^2)} \hat{\xi}_{k_1} \partial_\rho [\hat{v}_{k_2}(\rho) - \hat{v}_{k_2}(t_n)] d\rho \\ &\triangleq (\widehat{R_{121}})_k + (\widehat{R_{122}})_k. \end{aligned}$$

For  $R_{121}$ , by duality and Plancherel's identity we have

$$\begin{aligned} \|P_{>N_0} R_{121}\|_{L^2} &\lesssim \sup_{h: \|h\|_{l^2}=1} \sum_{\substack{|k_1| \gg |k_2| \\ |k_1+k_2| > N_0}} \frac{1}{k_1^2} |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(t_{n+1}) - \hat{v}_{k_2}(t_n)| |h_{k_1+k_2}| \\ &= \sup_{h: \|h\|_{l^2}=1} \sum_{\substack{|k_1| \gg |k_2| \\ |k_1+k_2| > N_0}} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(t_{n+1}) - \hat{v}_{k_2}(t_n)| |h_{k_1+k_2}| \langle k_1 \rangle^{-s-2} \\ &\lesssim \|v(t_{n+1}) - v(t_n)\|_{L_x^2} \sum_{k_1: |k_1| \gtrsim N_0} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle k_1 \rangle^{-s-2} \\ &\lesssim \|\xi\|_{\hat{b}^{s,p}} \|v(t_{n+1}) - v(t_n)\|_{L_x^2} \|\langle k_1 \rangle^{-s-2}\|_{l^{p'}(\{|k_1|: |k_1| \gtrsim N_0\})}. \end{aligned}$$

Note that  $-s - 2 + \frac{1}{p'} = -s - \frac{1}{p} - 1 < 0$ . Then we further have

$$\|P_{>N_0} R_{121}\|_{L^2} \lesssim N_0^{-s - \frac{1}{p} - 1} \|\xi\|_{\hat{b}^{s,p}} \|v(t_{n+1}) - v(t_n)\|_{L_x^2}. \quad (5.23)$$

For  $R_{122}$ , by noting  $\partial_t v = ie^{-it\partial_x^2} [\xi u + |u|^2 u]$ , we have

$$\begin{aligned} (\widehat{R_{122}})_k &= \int_{t_n}^{t_{n+1}} \sum_{\substack{k_2+k_3=\tilde{k}_2 \\ |k_1| \gg |\tilde{k}_2|}} \frac{e^{i\rho(k^2-\tilde{k}_2^2)}}{i(k^2-\tilde{k}_2^2)} \hat{\xi}_{k_1} \hat{\xi}_{k_2} \hat{v}_{k_3}(\rho) d\rho \\ &\quad + \int_{t_n}^{t_{n+1}} \sum_{|k_1| \gg |k_2|} \frac{e^{i\rho k_1^2}}{i(k^2-k_2^2)} \hat{\xi}_{k_1} (\widehat{|u|^2 u})_{k_2}(\rho) d\rho \\ &\triangleq (\widehat{R_{1221}})_k + (\widehat{R_{1222}})_k, \end{aligned}$$

and we shall estimate the two separately.

For  $R_{1221}$ , similarly as before we have

$$\begin{aligned} &\|P_{>N_0} R_{1221}\|_{L^2} \\ &\lesssim \tau \left\| \sup_{h: \|h\|_{l^2}=1} \sum_{\substack{k_1, k_2, k_3 \\ |k_1| \gg |k_2+k_3|, |k_1+k_2+k_3| > N_0}} \frac{|\hat{\xi}_{k_1}| |\hat{\xi}_{k_2}|}{(k_1+k_2+k_3)^2} |\hat{v}_{k_3}(t)| |h_{k_1+k_2+k_3}| \right\|_{L_t^\infty([0, T])}. \end{aligned}$$

Now we change the variables and use the relationship  $|k_1 + k_2 + k_3| \sim |k_1|$  to obtain that

$$\begin{aligned}
& \|P_{>N_0} R_{1221}\|_{L^2} \\
& \lesssim \tau \left\| \sup_{h: \|h\|_{l^2}=1} \sum_{\substack{k_1, \tilde{k}_2, k_3 \\ |k_1| \gg |\tilde{k}_2|, |k_1| \gtrsim N_0}} \langle k_1 \rangle^{-2} |\hat{\xi}_{k_1}| |\hat{\xi}_{\tilde{k}_2 - k_3}| |\hat{v}_{k_3}| |h_{k_1 + \tilde{k}_2}| \right\|_{L_t^\infty([0, T])} \\
& = \tau \left\| \sup_{h: \|h\|_{l^2}=1} \sum_{\substack{k_1, \tilde{k}_2, k_3 \\ |k_1| \gg |\tilde{k}_2|, |k_1| \gtrsim N_0}} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle \tilde{k}_2 - k_3 \rangle^s |\hat{\xi}_{\tilde{k}_2 - k_3}| |\hat{v}_{k_3}| |h_{k_1 + \tilde{k}_2}| \right. \\
& \quad \left. \cdot \langle k_1 \rangle^{-2-s} \langle \tilde{k}_2 - k_3 \rangle^{-s} \right\|_{L_t^\infty([0, T])} \\
& \lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \sum_{\tilde{k}_2, k_3} \left\| \langle k_1 \rangle^{-2-s} \right\|_{l^r(\{k_1: |k_1| \gtrsim \max\{|\tilde{k}_2|, N_0\}\})} \langle \tilde{k}_2 - k_3 \rangle^{-s} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| |\hat{v}_{k_3}(t)|.
\end{aligned}$$

Note that

$$\left\| \langle k_1 \rangle^{-2-s} \right\|_{l^r(\{k_1: |k_1| \gtrsim \max\{|\tilde{k}_2|, N_0\}\})} \lesssim N_0^{-2s - \frac{2}{p} - \frac{1}{2} +} |\tilde{k}_2|^{s + \frac{1}{p} - 1 -}.$$

This gives

$$\begin{aligned}
\|P_{>N_0} R_{1221}\|_{L^2} & \lesssim \tau N_0^{-2s - \frac{2}{p} - \frac{1}{2} +} \|\xi\|_{\dot{b}^{s,p}} \sum_{\tilde{k}_2, k_3} \langle \tilde{k}_2 \rangle^{s + \frac{1}{p} - 1 -} \langle \tilde{k}_2 - k_3 \rangle^{-s} \langle \tilde{k}_2 - k_3 \rangle^s |\hat{\xi}_{\tilde{k}_2 - k_3}| |\hat{v}_{k_3}(t)| \\
& \lesssim \tau N_0^{-2s - \frac{2}{p} - \frac{1}{2} +} \|\xi\|_{\dot{b}^{s,p}}^2 \left\| \langle \tilde{k}_2 \rangle^{s + \frac{1}{p} - 1 -} \langle \tilde{k}_2 - k_3 \rangle^{-s} \right\|_{l_{\tilde{k}_2}^{p'}} \sum_{k_3} |\hat{v}_{k_3}(t)|.
\end{aligned}$$

Since now  $s + \frac{1}{p} \leq 1$ , we have

$$\left\| \langle \tilde{k}_2 \rangle^{s + \frac{1}{p} - 1 -} \langle \tilde{k}_2 - k_3 \rangle^{-s} \right\|_{l_{\tilde{k}_2}^{p'}} \lesssim 1.$$

This further implies

$$\begin{aligned}
\|P_{>N_0} R_{1221}\|_{L^2} & \lesssim \tau N_0^{-2s - \frac{2}{p} - \frac{1}{2} +} \|\xi\|_{\dot{b}^{s,p}}^2 \sum_{k_3} |\hat{v}_{k_3}(t)| \\
& \lesssim \tau N_0^{-2s - \frac{2}{p} - \frac{1}{2} +} \|\xi\|_{\dot{b}^{s,p}}^2 \|v\|_{L_t^\infty H_x^{\frac{1}{2}+}([0, T])}.
\end{aligned}$$

For  $R_{1222}$ , we have similarly

$$\begin{aligned}
\|P_{>N_0} R_{1222}\|_{L^2} & \lesssim \tau \left\| \sup_{h: \|h\|_{l^2}=1} \sum_{|k_1| \gg |k_2|, |k_1| \gtrsim N_0} \frac{1}{(k_1 + k_2)^2} |\hat{\xi}_{k_1}| |(\widehat{|u|^2 u})_{k_2}| |h_{k_1 + k_2}| \right\|_{L_t^\infty([0, T])} \\
& \lesssim \tau \left\| \sup_{h: \|h\|_{l^2}=1} \sum_{|k_1| \gg |k_2|, |k_1| \gtrsim N_0} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| |(\widehat{|u|^2 u})_{k_2}| |h_{k_1 + k_2}| \langle k_1 \rangle^{-2-s} \right\|_{L_t^\infty([0, T])} \\
& \lesssim \tau \|\xi\|_{\dot{b}^{s,p}} \left\| |k|^{-2-s} \right\|_{l^r(|k| \gtrsim N_0)} \sum_{k_2} |(\widehat{|u|^2 u})_{k_2}| \\
& \lesssim \tau N_0^{-2-s + \frac{1}{r}} \|\xi\|_{\dot{b}^{s,p}} \|u^3\|_{L_t^\infty H_x^{\frac{1}{2}+}} \lesssim \tau N_0^{-s - \frac{1}{p} - \frac{3}{2}} \|\xi\|_{\dot{b}^{s,p}} \|u\|_{L_t^\infty H_x^{\frac{1}{2}+}}^3.
\end{aligned}$$

Since  $s + \frac{1}{p} \leq 1$ , we have that  $-s - \frac{1}{p} - \frac{3}{2} \leq -2s - \frac{2}{p} - \frac{1}{2}$ , which implies that

$$\|P_{>N_0} R_{1222}\|_{L^2} \lesssim \tau N_0^{-2s - \frac{2}{p} - \frac{1}{2} +} \|\xi\|_{\hat{b}^{s,p}} \|u\|_{L_t^\infty H_x^{\frac{1}{2}+}}^3.$$

Collecting the estimates on  $R_{1221}$  and  $R_{1222}$ , we have

$$\|P_{>N_0} R_{122}\|_{L^2} \leq C\tau N_0^{-2s - \frac{2}{p} - \frac{1}{2} +}. \quad (5.24)$$

Together with (5.20), (5.21), (5.22), (5.23) and (5.24), we find in total for  $s + \frac{1}{p} \leq \frac{1}{2}$  that

$$\begin{aligned} \|R_1^n\| &\leq C\tau \|v(t) - v(t_n)\|_{L_t^\infty H_x^a([t_n, t_{n+1}] \times \mathbb{T})} + C\tau N_0^{-s - \frac{1}{p} + \frac{1}{2} +} \|v(t) - v(t_n)\|_{L_t^\infty H_x^{\frac{1}{2}+}([t_n, t_{n+1}] \times \mathbb{T})} \\ &\quad + C N_0^{-s - \frac{1}{p} - 1} \|v(t_{n+1}) - v(t_n)\|_{L^2} + C\tau N_0^{-2s - \frac{2}{p} - \frac{1}{2} +}, \end{aligned}$$

with  $a = -s - \frac{1}{p} + 1 +$ . For  $\frac{1}{2} < s + \frac{1}{p} \leq 1$ ,

$$\begin{aligned} \|R_1^n\| &\leq C\tau \|v(t) - v(t_n)\|_{L_t^\infty H_x^a([t_n, t_{n+1}] \times \mathbb{T})} + C N_0^{-s - \frac{1}{p} - 1} \|v(t_{n+1}) - v(t_n)\|_{L^2} \\ &\quad + C\tau N_0^{-2s - \frac{2}{p} - \frac{1}{2} +}. \end{aligned}$$

Now we can apply Lemma 5.3 to the above findings to get

$$\begin{aligned} \|v(t) - v(t_n)\|_{L_t^\infty H_x^a([t_n, t_{n+1}] \times \mathbb{T})} &\leq C \min \left\{ \tau^{\frac{1}{4} + s + \frac{1}{p} -}, \tau \right\}, \\ \|v(t) - v(t_n)\|_{L_t^\infty H_x^{\frac{1}{2}+}([t_n, t_{n+1}] \times \mathbb{T})} &\leq C \min \left\{ \tau^{\frac{1}{2}(s + \frac{1}{p}) -}, \tau \right\}, \\ \|v(t_{n+1}) - v(t_n)\|_{L^2} &\leq C \min \left\{ \tau^{\frac{3}{4} + \frac{1}{2}(s + \frac{1}{p}) -}, \tau \right\}. \end{aligned}$$

Therefore, by choosing  $N_0 = \tau^{-\frac{1}{2}}$ , we find that for  $s + \frac{1}{p} \leq \frac{1}{2}$ ,

$$\begin{aligned} \|R_1^n\|_{L^2} &\leq C\tau \tau^{\frac{1}{4} + (s + \frac{1}{p}) -} + C\tau \tau^{\frac{1}{2}(s + \frac{1}{p}) + \frac{1}{4} -} \tau^{\frac{1}{2}(s + \frac{1}{p}) -} \\ &\quad + C\tau \tau^{\frac{1}{2}(s + \frac{1}{p}) + \frac{1}{2}} \tau^{\frac{3}{4} + \frac{1}{2}(s + \frac{1}{p}) -} + C\tau \tau^{s + \frac{1}{p} + \frac{1}{4} -} \\ &\leq C\tau \tau^{s + \frac{1}{p} + \frac{1}{4} -}. \end{aligned}$$

For  $\frac{1}{2} < s + \frac{1}{p} \leq 1$ ,

$$\begin{aligned} \|R_1^n\|_{L^2} &\leq C\tau \min \left\{ \tau^{s + \frac{1}{p} + \frac{1}{4} -}, \tau \right\} + C\tau \tau^{\frac{1}{2}(s + \frac{1}{p}) + \frac{1}{2}} \min \left\{ \tau^{\frac{3}{4} + \frac{1}{2}(s + \frac{1}{p}) -}, \tau \right\} + C\tau \tau^{s + \frac{1}{p} + \frac{1}{4} -} \\ &\leq C\tau \min \left\{ \tau^{s + \frac{1}{p} + \frac{1}{4} -}, \tau \right\}. \end{aligned}$$

Together with the two estimates above, we obtain that for  $0 \leq s + \frac{1}{p} \leq 1$ ,

$$\|R_1^n\|_{L^2} \leq C\tau \min \left\{ \tau^{\frac{1}{4} + s + \frac{1}{p} -}, \tau \right\}.$$

This finishes the proof of the lemma.  $\square$

**Lemma 5.5** (Local error from  $R_2^n$ ). *Let  $\xi \in \hat{b}^{s,p}$  with the conditions of Theorem 1.6 satisfied, and  $u \in L_t^\infty H_x^{s + \gamma p^-}([0, T] \times \mathbb{T})$ . Then,*

$$\|R_2^n\|_{L^2} \leq C\tau^{1 + \min\{\frac{1}{4} + \frac{1}{p} + s -, 1\}}, \quad (5.25)$$

where the constant  $C > 0$  only depends on  $\|\xi\|_{\hat{b}^{s,p}}$  and  $\|u\|_{L_t^\infty H_x^{s + \gamma p^-}}$ .

*Proof.* Note that

$$\left| \widehat{(R_2^n)}_k \right| \lesssim \tau \sum_{k_1+k_2=k} \left| \mathcal{M}_\tau(e^{is(k^2-k_2^2)}) - \mathcal{M}_\tau(e^{isk^2})\mathcal{M}_\tau(e^{-isk_2^2}) \right| |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(t_n)|.$$

If  $k = 0$  or  $k_2 = 0$ , then

$$\mathcal{M}_\tau(e^{is(k^2-k_2^2)}) - \mathcal{M}_\tau(e^{isk^2})\mathcal{M}_\tau(e^{-isk_2^2}) = 0,$$

and thus  $R_2^n = 0$ . Therefore, we may assume that  $k \neq 0$  and  $k_2 \neq 0$  in the following. By [Lemma 2.1](#), we have that for any  $0 \leq a \leq 1$ ,

$$\left| \mathcal{M}_\tau(e^{is(k^2-k_2^2)}) - \mathcal{M}_\tau(e^{isk^2})\mathcal{M}_\tau(e^{-isk_2^2}) \right| \lesssim \tau^a \min\{|k|^2|k_2|^{2a-2}, |k_2|^2|k|^{2a-2}\}.$$

In the following, we set

$$a = \min \left\{ \frac{1}{4} + \frac{1}{p} + s, 1 \right\}.$$

Accordingly, we write

$$\left| \widehat{(R_2^n)}_k \right| \lesssim \tau \sum_{\substack{k_1+k_2=k \\ |k| \lesssim |k_2|}} \left| \mathcal{M}_\tau(e^{is(k^2-k_2^2)}) - \mathcal{M}_\tau(e^{isk^2})\mathcal{M}_\tau(e^{-isk_2^2}) \right| |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(t_n)| \quad (5.26a)$$

$$+ \tau \sum_{\substack{k_1+k_2=k \\ |k| \gg |k_2|}} \left| \mathcal{M}_\tau(e^{is(k^2-k_2^2)}) - \mathcal{M}_\tau(e^{isk^2})\mathcal{M}_\tau(e^{-isk_2^2}) \right| |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(t_n)|. \quad (5.26b)$$

For [\(5.26a\)](#), we apply the estimate:

$$\left| \mathcal{M}_\tau(e^{is(k^2-k_2^2)}) - \mathcal{M}_\tau(e^{isk^2})\mathcal{M}_\tau(e^{-isk_2^2}) \right| \lesssim \tau^a \langle k \rangle^2 |k_2|^{2a-2},$$

and then by duality, we have

$$\begin{aligned} \|(5.26a)\|_{l_k^2} &\lesssim \tau^{1+a} \left\| \sum_{\substack{k_1+k_2=k \\ |k| \lesssim |k_2|}} \langle k \rangle^2 \langle k_2 \rangle^{2a-2} |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(t_n)| \right\|_{l_k^2} \\ &\lesssim \tau^{1+a} \sup_{h: \|h\|_{l^2}=1} \sum_{\substack{k_1+k_2=k \\ |k_1| \lesssim |k_2|}} \langle k \rangle^2 \langle k_2 \rangle^{2a-2} |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(t_n)| |h_k| \\ &\lesssim \tau^{1+a} \sup_{h: \|h\|_{l^2}=1} \sum_{\substack{k_1, k_2 \\ |k| \lesssim |k_2|}} \langle k_1 \rangle^{-s} \langle k_1 + k_2 \rangle^2 \langle k_2 \rangle^{2a-2-s-\gamma_p-} \\ &\quad \cdot \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle k_2 \rangle^{s+\gamma_p-} |\hat{v}_{k_2}(t_n)| |h_{k_1+k_2}|. \end{aligned} \quad (5.27)$$

In the case when  $s + \frac{1}{p} \leq \frac{1}{2}$ , by Cauchy-Schwartz's inequality, we further get

$$\begin{aligned} \|(5.26a)\|_{l_k^2} &\lesssim \tau^{1+a} \sup_{h: \|h\|_{l^2}=1} \sum_{k_2} \sum_{k_1: |k_1| \lesssim |k_2|} \langle k_1 \rangle^{-s} \langle k_2 \rangle^{2a-s-\gamma_p-} \\ &\quad \cdot \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \langle k_2 \rangle^{s+\gamma_p-} |\hat{v}_{k_2}(t_n)| |h_{k_1+k_2}| \\ &\lesssim \tau^{1+a} \|\xi\|_{\dot{b}^{s,p}} \sum_{k_2} \left\| \langle k_1 \rangle^{-s} \right\|_{l^r\{|k_1: |k_1| \lesssim |k_2|\}} \langle k_2 \rangle^{2a-s-\gamma_p-} \langle k_2 \rangle^{s+\gamma_p-} |\hat{v}_{k_2}(t_n)| \\ &\lesssim \tau^{1+a} \|\xi\|_{\dot{b}^{s,p}} \sum_{k_2} \langle k_2 \rangle^{2a-2s-\gamma_p+\frac{1}{r}-} \langle k_2 \rangle^{s+\gamma_p-} |\hat{v}_{k_2}(t_n)|, \end{aligned}$$

where  $\frac{1}{r} = \frac{1}{2} - \frac{1}{p}$ . Note that

$$2a - 2s - \gamma_p + \frac{1}{r} < -\frac{1}{2},$$

we further find

$$\|(5.26a)\|_{l_k^2} \lesssim \tau^{1+a} \|\xi\|_{\hat{b}^{s,p}} \|v(t_n)\|_{H_x^{s+\gamma_p-}}.$$

In the case when  $s + \frac{1}{p} > \frac{1}{2}$ , noting that

$$2a - s - \gamma_p < 0 \quad \text{and} \quad 2a - 2s - \gamma_p + \frac{1}{p'} < 0,$$

by (5.27) and Cauchy-Schwartz's inequality, we get

$$\begin{aligned} \|(5.26a)\|_{l_k^2} &\lesssim \tau^{1+a} \sup_{h:\|h\|_{l^2}=1} \sum_{k_1} \sum_{k_2:|k_2|\gtrsim|k_1|} \langle k_1 \rangle^{2a-2s-\gamma_p-} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \\ &\quad \cdot \langle k_2 \rangle^{s+\gamma_p-} |\hat{v}_{k_2}(t_n)| |h_{k_1+k_2}| \\ &\lesssim \tau^{1+a} \|v(t_n)\|_{H_x^{s+\gamma_p-}} \sum_{k_1} \langle k_1 \rangle^{2a-2s-\gamma_p-} |\hat{\xi}_{k_1}| \\ &\lesssim \tau^{1+a} \|\xi\|_{\hat{b}^{s,p}} \|v(t_n)\|_{H_x^{s+\gamma_p-}} \|\langle k_1 \rangle^{2a-2s-\gamma_p-}\|_{l_{k_1}^{p'}} \\ &\lesssim \tau^{1+a} \|\xi\|_{\hat{b}^{s,p}} \|v(t_n)\|_{H_x^{s+\gamma_p-}}. \end{aligned}$$

Therefore, together with the estimates in the above two cases, we finally get that

$$\|(5.26a)\|_{l_k^2} \lesssim \tau^{1+a} \|\xi\|_{\hat{b}^{s,p}} \|v(t_n)\|_{H_x^{s+\gamma_p-}}.$$

For (5.26b), we apply the estimate:

$$\left| \mathcal{M}_\tau(e^{is(k^2-k_2^2)}) - \mathcal{M}_\tau(e^{isk^2}) \mathcal{M}_\tau(e^{-isk_2^2}) \right| \lesssim \tau^a |k|^{2a-2} |k_2|^2.$$

Again, by duality and noting that  $|\xi_1| \sim |\xi_2|$ , we have

$$\begin{aligned} \|(5.26b)\|_{l_k^2} &\lesssim \tau^{1+a} \left\| \sum_{\substack{k_1+k_2=k \\ |k|\gg|k_2|}} \langle k_1 \rangle^{2a-2} \langle k_2 \rangle^2 |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(t_n)| \right\|_{l_k^2} \\ &\lesssim \tau^{1+a} \sup_{h:\|h\|_{l^2}=1} \sum_{\substack{k_1+k_2=k \\ |k_1|\gg|k_2|}} \langle k_1 \rangle^{2a-2} \langle k_2 \rangle^2 |\hat{\xi}_{k_1}| |\hat{v}_{k_2}(t_n)| |h_k| \\ &\lesssim \tau^{1+a} \sup_{h:\|h\|_{l^2}=1} \sum_{\substack{k_1, k_2 \\ |k|\gg|k_2|}} \langle k_1 \rangle^{-s+2a-2} \langle k_2 \rangle^{-s-\gamma_p-} \langle k_1 \rangle^s |\hat{\xi}_{k_1}| \\ &\quad \cdot \langle k_2 \rangle^{s+\gamma_p-} |\hat{v}_{k_2}(t_n)| |h_{k_1+k_2}|. \end{aligned}$$

Note that

$$-s + 2a - 2 + \frac{1}{r} < 0 \quad \text{and} \quad -s - \gamma_p + \frac{1}{2} < 0,$$

then by Cauchy-Schwartz's inequality, we have

$$\begin{aligned} \|(5.26b)\|_{l_k^2} &\lesssim \tau^{1+a} \|\langle k_1 \rangle^{-s+2a-2}\|_{l_{k_1}^r} \|\langle k_2 \rangle^{-s-\gamma_p+}\|_{l_{k_2}^2} \|\xi\|_{\hat{b}^{s,p}} \|v(t_n)\|_{H_x^{s+\gamma_p-}} \\ &\lesssim \tau^{1+a} \|\xi\|_{\hat{b}^{s,p}} \|v(t_n)\|_{H_x^{s+\gamma_p-}}. \end{aligned}$$

Therefore, combining with two estimates on (5.26), we obtain

$$\left\| \widehat{(R_2^n)}_k \right\|_{L_k^2} \lesssim \tau^{1+a} \|\xi\|_{\hat{b}^{s,p}} \|v(t_n)\|_{H_x^{s+\gamma_p-}}.$$

This together with Plancherel's identity give the proof of the lemma.  $\square$

**Lemma 5.6** (Local error of splitting part). *Under assumptions of Theorem 1.6, for the truncation term  $R_3^n$  defined in (5.5), we have*

$$\|R_3^n\|_{L^2} \leq C\tau \left( \tau^{\min\{1, \frac{1}{2}(s+\gamma_p)-\}} + N^{-s-\gamma_p+} \right), \quad (5.28)$$

where the constant  $C > 0$  only depends on  $\|u\|_{L_t^\infty H_x^{s+\gamma_p-}}$ .

*Proof.* First of all, we rewrite (5.5) into five parts  $R_3^n = R_{31}^n + \dots + R_{35}^n$  with

$$\begin{aligned} R_{31}^n &\triangleq i \int_{t_n}^{t_{n+1}} e^{-i\rho\partial_x^2} \left[ |u(\rho)|^2 u(\rho) - |e^{i\tau\partial_x^2} u(\rho)|^2 e^{i\tau\partial_x^2} u(\rho) \right] d\rho, \\ R_{32}^n &\triangleq i \int_{t_n}^{t_{n+1}} (e^{-i\rho\partial_x^2} - e^{-it_{n+1}\partial_x^2}) \left( |e^{i\tau\partial_x^2} u(\rho)|^2 e^{i\tau\partial_x^2} u(\rho) \right) d\rho, \\ R_{33}^n &\triangleq i \int_{t_n}^{t_{n+1}} e^{-it_{n+1}\partial_x^2} \left[ |e^{i\tau\partial_x^2} u(\rho)|^2 e^{i\tau\partial_x^2} u(\rho) - |e^{i\tau\partial_x^2} u(t_n)|^2 e^{i\tau\partial_x^2} u(t_n) \right] d\rho, \end{aligned}$$

and

$$\begin{aligned} R_{34}^n &\triangleq i\tau e^{-it_{n+1}\partial_x^2} \left[ |e^{i\tau\partial_x^2} u(t_n)|^2 - |P_{\leq N} e^{i\tau\partial_x^2} u(t_n)|^2 \right] e^{i\tau\partial_x^2} u(t_n), \\ R_{35}^n &\triangleq e^{-it_{n+1}\partial_x^2} \left[ e^{i\tau\partial_x^2} u(t_n) + i\tau |P_{\leq N} e^{i\tau\partial_x^2} u(t_n)|^2 e^{i\tau\partial_x^2} u(t_n) - \mathcal{N}_\tau [e^{i\tau\partial_x^2} u(t_n)] \right]. \end{aligned}$$

For  $R_{31}^n$ , note that the inequality  $|e^{ix} - 1| \leq 2^{2-a}|x|^a$  holds for any  $x \in \mathbb{R}$  and  $0 \leq a \leq 1$ , and  $s + \gamma_p - > \frac{1}{2}$ . Then, by the triangle inequality and Sobolev inequality we have

$$\|R_{31}^n\|_{L^2} \lesssim \tau \|(e^{i\tau\partial_x^2} - 1)u\|_{L_t^\infty L_x^2} \|u\|_{L_t^\infty H_x^{\frac{1}{2}+}}^2 \leq C\tau^{1+\alpha},$$

for

$$\alpha = \begin{cases} 1, & s + \gamma_p > 2, \\ \frac{s + \gamma_p -}{2}, & s + \gamma_p \leq 2. \end{cases}$$

Here and after  $C > 0$  is some constant dependent on  $\|u\|_{L_t^\infty H_x^{s+\gamma_p-}}$ .

For  $R_{32}^n$ , we have

$$\begin{aligned} \|R_{32}^n\|_{L^2} &\leq \int_{t_n}^{t_{n+1}} \left\| (e^{-i(\rho-t_{n+1})\partial_x^2} - 1) \left( |e^{i\tau\partial_x^2} u(\rho)|^2 e^{i\tau\partial_x^2} u(\rho) \right) \right\|_{L^2} d\rho \\ &\lesssim \int_{t_n}^{t_{n+1}} (\rho - t_{n+1})^\alpha \|u(\rho)\|_{H^{s+\gamma_p-}}^3 d\rho \leq C\tau^{1+\alpha}. \end{aligned}$$

For  $R_{33}^n$ , by the triangle inequality and Sobolev inequality we have

$$\|R_{33}^n\|_{L^2} \lesssim \int_{t_n}^{t_{n+1}} \|u(\rho) - u(t_n)\|_{L^2} \|u\|_{H_x^{\frac{1}{2}+}}^2 d\rho. \quad (5.29)$$

Noting that

$$\begin{aligned}
& \|u(\rho) - u(t_n)\|_{L_t^\infty L_x^2([t_n, t_{n+1}] \times \mathbb{T})} \\
& \lesssim \|P_{\leq N}(u(\rho) - u(t_n))\|_{L_t^\infty L_x^2([t_n, t_{n+1}] \times \mathbb{T})} + \|P_{> N}(u(\rho) - u(t_n))\|_{L_t^\infty L_x^2([t_n, t_{n+1}] \times \mathbb{T})} \\
& \lesssim \tau \|P_{\leq N} \partial_t u\|_{L_t^\infty L_x^2([t_n, t_{n+1}] \times \mathbb{T})} + N^{-s-\gamma_{p^+}} \|u\|_{L_t^\infty H_x^{s+\gamma_{p^-}}} \\
& \lesssim \tau \|P_{\leq N}(\partial_{xx} u + \xi u - \lambda |u|^2 u)\|_{L_t^\infty L_x^2([t_n, t_{n+1}] \times \mathbb{T})} + N^{-s-\gamma_{p^+}} \|u\|_{L_t^\infty H_x^{s+\gamma_{p^-}}} \\
& \leq C(\tau N^{2-2\alpha} + N^{-s-\gamma_{p^+}}).
\end{aligned}$$

Hence, by Young's inequality,

$$\|u(\rho) - u(t_n)\|_{L_t^\infty L_x^2([t_n, t_{n+1}] \times \mathbb{T})} \leq C(\tau^\alpha + N^{-s-\gamma_{p^+}}).$$

Inserting this inequality into (5.29), we have

$$\|R_{33}^n\|_{L^2} \leq C\tau(\tau^\alpha + N^{-s-\gamma_{p^+}}).$$

For  $R_{34}^n$ , we have

$$\|R_{34}^n\|_{L^2} \leq C\tau \|u(t_n) - P_{\leq N} u(t_n)\|_{L^2} \leq C\tau N^{-s-\gamma_{p^+}}.$$

For  $R_{35}^n$ , we have

$$\begin{aligned}
\|R_{35}^n\|_{L^2} & \lesssim \left\| 1 + i\tau |P_{\leq N} e^{i\tau \partial_x^2} u(t_n)|^2 - \exp\left(i\tau |P_{\leq N} e^{i\tau \partial_x^2} u(t_n)|^2\right) \right\|_{L^2} \\
& \lesssim \tau^2 \left\| |P_{\leq N} e^{i\tau \partial_x^2} u(t_n)|^4 \right\|_{L^2} \leq C\tau^2.
\end{aligned}$$

Combining the estimates above, we obtain the assertion (5.28) and the proof is done.  $\square$

With the established stability results and the local error estimates, we are now ready to give the proof of [Theorem 1.6](#).

*Proof.* By [Theorem 1.1](#) and [Theorem 1.2](#), we have  $u \in L^\infty((0, T); H^{s+\gamma_{p^-}})$ . Moreover, from the proof of [Theorem 1.1](#), we have in fact

$$\|u\|_{L_t^\infty H_x^{s+\frac{3}{2}+\frac{1}{p^-}}([0, T] \times \mathbb{T})} \leq 2\|u_0\|_{H^{s+2}}.$$

Therefore, from [Lemmas 5.1, 5.2](#) and [5.4 to 5.6](#), we have that

$$\begin{aligned}
\|h^{n+1}\|_{L^2} & \leq \|\Phi(v^n) - \Phi(v(t_n))\|_{L^2} + \|\Psi(v^n) - \Psi(v(t_n))\|_{L^2} + \|R_1^n\|_{L^2} + \|R_2^n\|_{L^2} + \|R_3^n\|_{L^2} \\
& \leq (1 + C\tau)\|h^n\|_{L^2} + C\tau N\|h^n\|_{L^2}^3 + C\tau^{1+\min\{\frac{1}{4}+\frac{1}{p}+s-, 1\}} \\
& \quad + C\tau\left(\tau^{\min\{1, \frac{1}{2}(s+\gamma_{p^-})-\}} + N^{-s-\gamma_{p^+}}\right),
\end{aligned}$$

where the constant  $C > 0$  depends only on  $\|\xi\|_{\dot{H}^{s,p}}$ ,  $T$  and  $\|u_0\|_{H^{s+2}}$ . Noting that

$$\frac{1}{2}(s + \gamma_{p^-}) - \min\left\{\frac{1}{4} + \frac{1}{p} + s-, 1\right\} \geq \frac{1}{8},$$

and so by taking  $N = \tau^{-\frac{1}{2}+\varepsilon_0}$  for any fixed  $0 < \varepsilon_0 \leq \frac{1}{8(s+\gamma_{p^-})}$ , we can further get

$$\|h^{n+1}\|_{L^2} \leq C_1\tau^{1+\alpha} + (1 + C_2\tau)\|h^n\|_{L^2} + C_3\tau^{\frac{1}{2}+\varepsilon_0}\|h^n\|_{L^2}^3, \quad 0 \leq n < \frac{T}{\tau}, \quad (5.30)$$

where the constants  $C_j > 0, j = 1, 2, 3$  depend only on  $\|\xi\|_{\dot{H}^{s,p}}$ ,  $T$  and  $\|u_0\|_{H^{s+2}}$ . Here and after, we denote  $\alpha = \min\{\frac{1}{4} + \frac{1}{p} + s-, 1\}$  for short.

Now we claim that there exists some  $\tau_0 > 0$  (to be determined) such that for any  $\tau \in (0, \tau_0]$ ,

$$\|h^n\|_{H^\gamma} \leq C_1 \tau^{1+\alpha} \sum_{j=0}^n (1 + 2C_2 \tau)^j, \quad n = 0, 1, \dots, \frac{T}{\tau}. \quad (5.31)$$

We prove it by induction, see [42] for a similar process. Firstly, since  $h^0 \equiv 0$ , (5.31) trivially holds for  $n = 0$ . Now we assume that it holds till some  $0 \leq n_0 \leq \frac{T}{\tau} - 1$ , i.e.,

$$\|h^n\|_{H^\gamma} \leq C_1 \tau^{1+\alpha} \sum_{j=0}^n (1 + 2C_2 \tau)^j, \quad \forall 0 \leq n \leq n_0. \quad (5.32)$$

From (5.32), we have that for any  $0 \leq n \leq n_0$ ,

$$\|h^n\|_{H^\gamma} \leq C_4 \tau^\alpha, \quad (5.33)$$

where  $C_4 = C_1 C_2^{-1} e^{2C_2 T}$ . Then by (5.30), we find

$$\|h^{n_0+1}\|_{H^\gamma} \leq C_1 \tau^{1+\alpha} + \left(1 + C_2 \tau + C_3 C_4^2 \tau^{\frac{1}{2} + \varepsilon_0 + 2\alpha}\right) \cdot C_1 \tau^{1+\alpha} \sum_{j=0}^{n_0} (1 + 2C_2 \tau)^j.$$

Note that  $\frac{1}{2} + \varepsilon_0 + 2\alpha \geq 1 + \frac{1}{2}\varepsilon_0$ , then by choosing some  $\tau_0 \in (0, 1]$  such that  $C_3 C_4^2 \tau_0^{\frac{1}{2}\varepsilon_0} \leq C_2$ , we can further have that for any  $\tau \in (0, \tau_0]$ ,

$$\begin{aligned} \|h^{n_0+1}\|_{H^\gamma} &\leq C_1 \tau^{1+\alpha} + \left(1 + C_2 \tau + C_3 C_4^2 \tau^{1 + \frac{1}{2}\varepsilon_0}\right) \cdot C_1 \tau^{1+\alpha} \sum_{j=0}^{n_0} (1 + 2C_2 \tau)^j \\ &= C_1 \tau^{1+\alpha} + C_1 \tau^{1+\alpha} \sum_{j=0}^{n_0} (1 + 2C_2 \tau)^{j+1} \\ &= C_1 \tau^{1+\alpha} \sum_{j=0}^{n_0+1} (1 + 2C_2 \tau)^j. \end{aligned}$$

This finishes the induction and proves the claim (5.31).

Then by iteration, we have

$$\|h^n\|_{L^2} \leq C \tau^{\min\{\frac{1}{4} + \frac{1}{p} + s -, 1\}},$$

which finishes the proof of the theorem.  $\square$

## 6. NUMERICAL EXPERIMENTS

In this section, we will carry out some numerical tests to verify the given theoretical results on the regularity of the solution of (1.1) and on the accuracy of the proposed numerical scheme (1.6). The spatial discretization of (1.6) will be implemented here by the Fourier pseudo-spectral method [40]. In the end, some accuracy comparisons will be made between (1.6) and the existing schemes from the literature.

**6.1. On regularity of solution.** Firstly, we verify the regularity results on the solution of the PDE, i.e., [Theorem 1.1](#) and [Theorem 1.2](#). To do so, we solve the NLS model (1.1) by the numerical scheme (1.6) with very fine mesh so that the computations are accurate. We fix

$$\lambda = 1, \quad u_0(x) = \frac{\cos(x)}{2 + \sin(2x)}, \quad x \in (-\pi, \pi), \quad (6.1)$$

for (1.1) in this subsection, and we solve the equation till  $t = 2$  for the solution  $u(t, x)$ . The potential function in the following will be constructed as

$$\xi(x) = \operatorname{Re} \sum_{k=-N/2}^{N/2-1} \zeta_k e^{ik(x+\pi)}, \quad (6.2)$$

by choosing  $\zeta_k$  to determine its regularity and by taking an even integer  $N$  as the total number of Fourier frequencies which is also the number of spatial grid points.




FIGURE 1. Test of [Theorem 1.1](#) for  $s = 0, p = \infty$ : modulus of Fourier coefficient of the solution (left) and the generated potential  $\xi \in \hat{l}^\infty$  (right).

**Example 6.1.** (Test of [Theorem 1.1](#) for  $s = 0, p = \infty$ ) We begin with [Theorem 1.1](#). Construct a  $\xi(x) \in \hat{l}^\infty$  through (6.2) by randomly generating  $\zeta_k \in [-2, 2]$  based on the uniform distribution. To test the regularity result, we compute the Fourier coefficient of the solution:  $\hat{u}_k(t)$  at  $t = 2$ . If  $u \in H^{3/2-}$  as predicted by [Theorem 1.1](#) for  $s = 0, p = \infty$ , then  $\sum_k |k|^{3-} |\hat{u}_k|^2 < \infty$ , and so  $|k|^{3-} |\hat{u}_k|^2 \lesssim k^{-1-}$ . This implies that we expect to observe  $|\hat{u}_k| \lesssim k^{-2}$  in this case. With  $N = 1024$ , the modulus of Fourier coefficient of the solution together with that of the generated potential are plotted against the frequency in [Figure 1](#).

**Example 6.2.** (Test of [Theorem 1.1](#) for  $s = 0, p = 4$ ) To further test [Theorem 1.1](#), we then consider a  $\xi(x) \in \hat{b}^{0,4}$  through (6.2) by choosing

$$\zeta_k = \begin{cases} (\eta_{1,k} + i\eta_{2,k}) |k|^{-0.26}, & \text{if } k \neq 0, \\ 1, & \text{if } k = 0, \end{cases}$$

with  $\eta_{1,k}, \eta_{2,k} \in [-4, 4]$  randomly generated by the uniform distribution. [Theorem 1.1](#) in this case predicts  $u \in H^{7/4-}$ , which means  $\sum_k |k|^{7/2-} |\hat{u}_k|^2 < \infty$ . If so, then  $|k|^{7/2-} |\hat{u}_k|^2 \lesssim k^{-1-}$  and we are expecting  $|\hat{u}_k| \lesssim k^{-2.25}$  for this example. To verify it, again we compute  $\hat{u}_k(t)$  at  $t = 2$  and check its decaying rate with respect to  $k$ . With  $N = 1024$ , the corresponding numerical results are shown in [Figure 2](#).




FIGURE 2. Test of [Theorem 1.1](#) for  $s = 0, p = 4$ : modulus of Fourier coefficient of the solution (left) and the generated potential  $\xi \in \hat{l}^\infty$  (right).




FIGURE 3. Test of [Theorem 1.2](#) for  $s = 0, p = 2$ : modulus of Fourier coefficient of the solution (left) and the generated potential  $\xi \in L^2$  (right).

**Example 6.3.** (Test of [Theorem 1.2](#) for  $s = 0, p = 2$ ) To test [Theorem 1.2](#), we construct a  $\xi(x) \in L^2$  through (6.2) by choosing

$$\zeta_k = \begin{cases} (\eta_{1,k} + i\eta_{2,k})|k|^{-0.6}, & \text{if } k \neq 0, \\ 1, & \text{if } k = 0, \end{cases}$$

with  $\eta_{1,k}, \eta_{2,k} \in [-5, 5]$  randomly generated by the uniform distribution. Now [Theorem 1.2](#) tells that  $u \in H^2$ , i.e.,  $\sum_k |k|^4 |\hat{u}_k|^2 < \infty$ , and so  $|\hat{u}_k| \lesssim k^{-2.5-}$  is expected here. The Fourier coefficient at  $t = 2$  is computed to verify the decaying rate. With  $N = 1024$ , the numerical results are shown in [Figure 3](#).

All the numerical examples in this subsection illustrate that the decaying rate of the Fourier coefficient of the solution matches well with the expected value from the theorems. Thus, the theoretical results i.e., [Theorem 1.1](#) and [Theorem 1.2](#), on the regularity of the solution of (1.1), are valid and sharp. Note in addition that, the smoothness of the used initial data (6.1) in the tests did not provide more regularity for the solution than expected, and this illustrates [Remark 1.2](#).

**6.2. On accuracy of scheme.** Next, we test the theoretical result on the convergence order of the numerical scheme, i.e., [Theorem 1.6](#) for LRI (1.6). We construct an initial data

$u_0 \in H^2$  for (1.1) in this subsection as

$$u_0(x) = \sum_{k=-N/2}^{N/2-1} \widehat{(u_0)}_k e^{ik(x+\pi)}, \quad \widehat{(u_0)}_k = \begin{cases} \eta_k |k|^{-2.55}, & \text{if } k \neq 0, \\ 0, & \text{if } k = 0, \end{cases} \quad (6.3)$$

with  $\eta_k$  randomly sampled from the interval  $[0, 1]$  by the uniform distribution. We shall compute the numerical solution of (1.1) at  $t = t_n = 1$ , and we shall measure the relative error

$$\text{error} = \|u(t_n) - u^n\|_{L^2} / \|u(t_n)\|_{L^2} \quad (6.4)$$

of the scheme. The reference solution here is obtained by using very fine mesh size.




FIGURE 4. Results of Example 6.4: profiles of the potential  $\xi(x)$  and the solution  $|u_0(x)|$ ,  $|u(t=1, x)|$  (1st row); error (6.4) of LRI (1.6) (2nd row).

**Example 6.4.** (Lowest order in Theorem 1.6) We take  $\lambda = -2$  and the initial data (6.3) for (1.1). A potential function  $\xi \in \hat{l}^\infty$  is taken as

$$\xi(x) = -\frac{1}{5} \sum_{k=-N/2}^{N/2} e^{ikx}.$$

The profiles of the potential and the solution at  $t = 1$  are displayed in Figure 4. Theorem 1.6 in this case predicts the lowest convergence order for the LRI scheme (1.6) as  $\mathcal{O}(\tau^{1/4-})$ . With the number of spatial grids  $N = 2048$  fixed, the discretization error (6.4) of (1.6) at  $t = 1$  is shown in Figure 4 under different time steps. As shown by the error curve in Figure 4, the averaged decreasing rate is about  $1/4$ .




FIGURE 5. Results of **Example 6.5** under  $\xi \in \hat{b}^{0,2}$ : profiles of  $\xi(x)$ ,  $|u_0(x)|$  and  $|u(t=1, x)|$  (1st row); error (6.4) of LRI (1.6) (2nd row).

**Example 6.5.** (Other orders in **Theorem 1.6**) Generate a  $H^2$ -initial data as (6.3) for (1.1) and fix  $\lambda = 4$ . We construct the potential  $\xi \in \hat{b}^{s,p}$  via

$$\xi(x) = \operatorname{Re} \sum_{k=-N/2}^{N/2-1} \zeta_k e^{ik(x+\pi)}, \quad \zeta_k = \begin{cases} (\eta_{1,k} + i\eta_{2,k})|k|^{-\delta}, & \text{if } k \neq 0, \\ 1, & \text{if } k = 0, \end{cases}$$

with  $\eta_{1,k}, \eta_{2,k} \in [-5, 5]$  randomly generated by the uniform distribution. We consider

$$\delta = \begin{cases} 0.51, & \text{for } \xi(x) \in \hat{b}^{0,2}, \\ 0.76, & \text{for } \xi(x) \in \hat{b}^{1/4,2}, \end{cases}$$

and **Theorem 1.6** respectively predicts the accuracy of LRI (1.6) as  $\mathcal{O}(\tau^{3/4-})$  and  $\mathcal{O}(\tau^{1-})$ . The number of spatial grids  $N = 2048$  is again used and fixed for computations. For  $\delta = 0.51$ , the discretization error (6.4) of (1.6) at  $t = 1$  together with the profiles of the potential and solution in this case are shown in **Figure 5**. The corresponding results of  $\delta = 0.76$  are shown in **Figure 6**. The two error curves in **Figure 5** and **Figure 6** clearly decrease at the expected rates.

In total, the observed convergence results in this subsection all match well with **Theorem 1.6**. This verifies its validity and indicates the sharpness of the error estimate (1.8).

**6.3. Accuracy comparison.** At last, we conduct some numerical tests to compare the convergence/accuracy of the proposed LRI (1.6) with the existing schemes in the literature for (1.1). The concerned numerical schemes from the literature are listed below.

- The most traditional finite difference scheme:  $i\frac{1}{\tau}(u^{n+1} - u^n) + \partial_x^2 u^{n+1} + \xi u^{n+1} = \lambda |u^n|^2 u^n$ .




FIGURE 6. Results of [Example 6.5](#) under  $\xi \in \hat{b}^{1/4,2}$ : profiles of  $\xi(x)$ ,  $|u_0(x)|$  and  $|u(t=1, x)|$  (1st row); error (6.4) of LRI (1.6) (2nd row).

- The commonly used Lie-Trotter splitting scheme:  $u^{n+1} = e^{i\partial_x^2 \tau} e^{-i\tau(-\xi + \lambda|u^n|^2)} u^n$ .
- The recently analyzed exponential wave integrator (EWI) [7]:  $u^{n+1} = e^{i\partial_x^2 \tau} u^n - i\tau \mathcal{D}_\tau [(-\xi + \lambda|u^n|^2) u^n]$ .
- The recently proposed LRI from [4]:  $u^{n+1} = e^{i\partial_x^2 \tau} [u^n + i\tau u^n \mathcal{D}_{-\tau} \xi - i\tau \lambda (u^n)^2 \mathcal{D}_{-2\tau} \overline{u^n}]$ .

Note that the schemes presented above all have higher order versions, but their high order accuracy is achieved only when the setup of the NLS model is smooth enough. Here what we would like to address is the performance of schemes under the rough setup, so we choose to focus on the first order schemes for tests and comparisons. The reference solution and the measure of error are set the same as before.




FIGURE 7. Accuracy comparison of schemes for [Example 6.6](#): with the set of data (6.5) (left); with the set of data (6.6) (right).

**Example 6.6.** We first perform comparisons of schemes within the setup of **Theorem 1.6**:  $u_0 \in H^2, s = 0, p = \infty$ . Generate an initial data  $u_0 \in H^2$  for (1.1) as

$$u_0(x) = \sum_{k=-N/2}^{N/2-1} (\widehat{u_0})_k e^{ik(x+\pi)}, \quad (\widehat{u_0})_k = \begin{cases} \eta_k |k|^{-2.51}, & \text{if } k \neq 0, \\ 0, & \text{if } k = 0. \end{cases}$$

The coefficients  $\eta_k, \lambda$  and the potential  $\xi \in \hat{l}^\infty$  for (1.1) are taken as the following two sets of data. The first set considers the  $\delta$ -type potential function:

$$\eta_k \in [0, 1/3], \quad \lambda = -2, \quad \xi(x) = - \sum_{k=-N/2}^{N/2} \left( e^{ikx} + e^{ik(x+2)} + e^{ik(x-2)} \right). \quad (6.5)$$

The second set considers more general rough potential:

$$\text{Re}(\eta_k) \in [0, 1/3], \quad \text{Im}(\eta_k) \in [0, 1/5], \quad \lambda = -0.05, \quad \xi(x) = \text{Re} \sum_{k=-N/2}^{N/2-1} \zeta_k e^{ik(x+\pi)}, \quad (6.6)$$

with  $\zeta_k \in [0, 4]$  sampled by the uniform distribution. Under (6.5) or (6.6) with  $N = 2048$  fixed, the error of each concerned numerical scheme at  $t = 1$  is given in **Figure 7**.

It can be seen from the plots in **Figure 7** that the proposed LRI (1.6) is more accurate than all the other considered schemes. In contrast, the LRI [4] and the EWI [7] converge slowly and the accuracy order is unclear, while the Lie-Trotter splitting scheme is not working at all.




FIGURE 8. Accuracy comparison of schemes for **Example 6.7**.

**Example 6.7.** We then test the performance of the schemes under rougher initial data for (1.1) by generating a  $u_0 \in L^2$  as

$$u_0(x) = \sum_{k=-N/2}^{N/2-1} (\widehat{u_0})_k e^{ik(x+\pi)}, \quad (\widehat{u_0})_k = \begin{cases} \eta_k |k|^{-1.1}, & \text{if } k \neq 0, \\ 0, & \text{if } k = 0, \end{cases}$$

with the other parameters set as (6.5). With the number of spatial grid points fixed as  $N = 2048$ , the error of each scheme at  $t = 1$  is given in **Figure 8**. Although the tested setup is not covered by **Theorem 1.6**, we can see that the proposed LRI scheme (1.6) is still working well in this case, and its accuracy is much better than the others. This illustrates that (1.6) can have more advantages for solving the NLS equation (1.1) under rough setup.

## 7. CONCLUSION

We present new, sharp results of the cubic nonlinear Schrödinger equation (NLS) with a spatially rough potential, posed on a one-dimensional torus which is the mathematical model for nonlinear Anderson localization. We deal both with the well/ill-posedness analysis on the PDE level and its application for numerical discretization and convergence analysis. In the PDE analysis, we provide insights into how the regularity of the solution is impacted by the regularity of the potential, offering quantitative and explicit characterizations. Additionally, we establish ill-posedness results to demonstrate the sharpness of our regularity characterizations and to identify the minimum required regularity of the potential for the solvability of the NLS model. Based on our regularity results, we design a suitable numerical discretization for the model and demonstrate its convergence with an optimal error bound. The numerical experiments demonstrate the theoretical regularity results on the PDE level and also validate the established convergence rate of the proposed scheme. Furthermore, some comparisons with existing schemes are given, showcasing the superior accuracy of our scheme in the case of a rough potential.

## ACKNOWLEDGEMENTS

Financial support from the Austrian Science Fund (FWF) via the SFB project F65 is acknowledged. Y. Wu is partially supported by NSFC 12171356. X. Zhao is partially supported by NSFC 12271413 and the Natural Science Foundation of Hubei Province 2019CFA007. X. Zhao thanks the Wolfgang Pauli Institute Vienna for hospitality.

## REFERENCES

- [1] P.W. ANDERSON, Absence of diffusion in certain random lattices, *Physical Review* 109 (1958) 1492.
- [2] X. ANTOINE, W. BAO, C. BESSE, Computational methods for the dynamics of the nonlinear Schrödinger/Gross-Pitaevskii equations, *Computer Physics Communications* 184 (2013) 2621-2633.
- [3] J. BOURGAIN, *Global Solutions of Nonlinear Schrödinger Equations*, Colloquium Publications, American Mathematical Society, 1999.
- [4] Y.A. BRONSARD, Error analysis of a class of semi-discrete schemes for solving the Gross-Pitaevskii equation at low regularity, *Journal of Computational and Applied Mathematics* 418 (2023) 114632.
- [5] I. BEJENARU, T. TAO, Sharp well-posedness and ill-posedness results for a quadratic non-linear Schrödinger equation, *Journal of Functional Analysis* 233 (2006) 228-259.
- [6] W. BAO, Y. CAI, Mathematical theory and numerical methods for Bose-Einstein condensation, *Kinetic and Related Models* 6 (2013) 1-135.
- [7] W. BAO, C. WANG, Optimal error bounds on the exponential wave integrator for the nonlinear Schrödinger equation with low regularity potential and nonlinearity, *SIAM Journal on Numerical Analysis* 62 (2024) 93-118.
- [8] J. BOURGAIN, W. WANG, Quasi-periodic solutions of nonlinear random Schrödinger equations, *Journal of the European Mathematical Society* 10 (2008) 1-45.
- [9] A. DE BOUARD, A. DEBUSSCHE, A stochastic nonlinear Schrödinger equation with multiplicative noise, *Communications in Mathematical Physics* 205 (1999) 161-181.
- [10] T. CAZENAVE, *Semilinear Schrödinger Equations*, Courant Lecture Notes in Mathematics, American Mathematical Society, 2003.
- [11] C. CONTI, Solitonization of the Anderson localization, *Physical Review A* 86 (2012) 061801.
- [12] L. DUMAZ, C. LABBÉ, Localization of the continuous Anderson Hamiltonian in 1-D, *Probability Theory and Related Fields* 176 (2020) 353-419.
- [13] A. DEBUSSCHE, J. MARTIN, Solution to the stochastic Schrödinger equation on the full space, *Nonlinearity* 32 (2019) 1147-1174.
- [14] A. DEBUSSCHE, H. WEBER, The Schrödinger equation with spatial white noise potential, *Electronic Journal of Probability* 23 (2018) 1-16.
- [15] N. GHOFRANIHA, S. GENTILINI, V. FOLLI, E. DELRE, C. CONTI, Shock waves in disordered media, *Physical Review Letter* 109 (2012) 243902.

- [16] Y. GU, T. KOMOROWSKI, L. RYZHIK, The Schrödinger equation with spatial white noise: The average wave function, *Journal of Functional Analysis* 274 (2018) 2113-2138.
- [17] P. HENNING, D. PETERSEIM, Crank-Nicolson Galerkin approximations to nonlinear Schrödinger equations with rough potentials, *Mathematical Models and Methods in Applied Sciences* 27 (2017) 2147-2184.
- [18] J. HONG, X. WANG, Invariant Measures for Stochastic Nonlinear Schrödinger Equations: Numerical Approximations and Symplectic Structures, *Lecture Notes in Mathematics Book 2251*, Springer, 2019.
- [19] R. FUKUIZUMI, M. OHTA, T. OZAWA, Nonlinear Schrödinger equation with a point defect, *Annales de l'Institut Henri Poincaré, Analyse non Linéaire* 25 (2008) 837-845.
- [20] S. FISHMAN, Y. KRIVOLAPOV, A. SOFFER, The nonlinear Schrödinger equation with a random potential: results and puzzles, *Nonlinearity* 25 (2012) 53-72.
- [21] S. FLACH, D.O. KRIMER, CH. SKOKOS, Universal spreading of wave packets in disordered nonlinear systems, *Physical Review Letters* 102 (2009) 024101.
- [22] R.H. GOODMAN, P.J. HOLMES, M.I. WEINSTEIN, Strong NLS soliton-defect interactions, *Physica D* 192 (2004) 215-248.
- [23] F. GERMINET, A. KLEIN, A comprehensive proof of localization for continuous Anderson models with singular random potentials, *Journal of the European Mathematical Society* 15 (2013) 53-143.
- [24] I.Y. GOL'DSHTEIN, S.A. MOLCHANOV, L.A. PASTUR, A pure point spectrum of the stochastic one-dimensional Schrödinger operator, *Functional Analysis and Its Applications* 11 (1977) 1-8.
- [25] L.I. IGNAT, A splitting method for the nonlinear Schrödinger equation, *Journal of Differential Equations*, 250 (2011) 3022-3046.
- [26] L.I. IGNAT, E. ZUAZUA, Numerical dispersive schemes for the nonlinear Schrödinger equation, *SIAM Journal on Numerical Analysis* 47 (2009) 1366-1390.
- [27] S. JIN, P. MARKOWICH, C. SPARBER, Mathematical and computational methods for semiclassical Schrödinger equations, *Acta Numerica* 20 (2011) 121-209.
- [28] T. KACHMAN, S. FISHMAN, A. SOFFER, Numerical implementation of the multiscale and averaging methods for quasi periodic systems, *Computer Physics Communications* 221 (2017) 235-245.
- [29] B. LI, Y. WU, A fully discrete low-regularity integrator for the 1D periodic cubic nonlinear Schrödinger equation, *Numerische Mathematik* 149 (2021) 151-183.
- [30] B. LI, Y. WU, An unfiltered low-regularity integrator for the KdV equation with solutions below  $H^1$ , arXiv:2206.09320.
- [31] A. OSTERMANN, F. ROUSSET, K. SCHRATZ, Fourier integrator for periodic NLS: low regularity estimates via discrete Bourgain spaces, *Journal of the European Mathematical Society* 25 (2023) 3913-3952.
- [32] A. OSTERMANN, K. SCHRATZ, Low regularity exponential-type integrators for semilinear Schrödinger equations, *Foundations of Computational Mathematics* 18 (2018) 731-755.
- [33] M. PIRAUD, P. LUGAN, P. BOUYER, A. ASPECT, L. SANCHEZ-PALENCIA, Localization of a matter wave packet in a disordered potential, *Physical Review A* 83 (2011) 031603.
- [34] A.S. PIKOVSKY, D.L. SHEPELYANSKY, Destruction of Anderson localization by a weak nonlinearity, *Physical Review Letters* 100 (2008) 094101.
- [35] I. RODNIANSKI, W. SCHLAG, Time decay for solutions of Schrödinger equations with rough and time-dependent potentials, *Inventiones Mathematicae* 155 (2004) 451-513.
- [36] L. SANCHEZ-PALENCIA, D. CLÉMENT, P. LUGAN, P. BOUYER, G.V. SHLYAPNIKOV, A. ASPECT, Anderson localization of expanding Bose-Einstein condensates in random potentials, *Physical Review Letters* 98 (2007) 210401.
- [37] N. SERGEY, A. SOFFER, M.B. TRAN, On the wave turbulence theory for the nonlinear Schrödinger equation with random potentials, *Entropy* 21 (2019) 823.
- [38] CH. SKOKOS, D.O. KRIMER, S. KOMINEAS, S. FLACH, Delocalization of wave packets in disordered nonlinear chains, *Physical Review E* 79 (2009) 056211.
- [39] T. TAO, *Nonlinear Dispersive Equations. Local and Global Analysis*, American Mathematical Society, 2006.
- [40] L.N. TREFETHEN, *Spectral Methods in MATLAB*, SIAM, Philadelphia, 2000.
- [41] W. WANG, Z. ZHANG, Long time Anderson localization for nonlinear random Schrödinger equation, *Journal of Statistical Physics* 134 (2009) 953.
- [42] Y. WU, X. ZHAO, Embedded exponential-type low-regularity integrators for KdV equation under rough data, *BIT Numerical Mathematics* 62 (2022) 1049-1090.
- [43] X. ZHAO, Numerical integrators for continuous disordered nonlinear Schrödinger equation, *Journal of Scientific Computing* 89 (2021) 40.

N. MAUSER: RESEARCH PLATFORM MMM C/O FAK. MATHEMATIK, UNIV. WIEN, OSKAR-MORGENSTERN-PLATZ 1, A-1090, VIENNA.

*Email address:* [mauser@courant.nyu.edu](mailto:mauser@courant.nyu.edu)

*URL:* <https://www.wpi.ac.at/director.html>

Y. WU: CENTER FOR APPLIED MATHEMATICS, TIANJIN UNIVERSITY, 300072, TIANJIN, P. R. CHINA.

*Email address:* [yerfmth@gmail.com](mailto:yerfmth@gmail.com)

*URL:* <http://cam.tju.edu.cn/~yfw/>

X. ZHAO: SCHOOL OF MATHEMATICS AND STATISTICS & COMPUTATIONAL SCIENCES HUBEI KEY LABORATORY, WUHAN UNIVERSITY, 430072 WUHAN, CHINA

*Email address:* [matzhxf@whu.edu.cn](mailto:matzhxf@whu.edu.cn)

*URL:* <http://jszy.whu.edu.cn/zhaoxiaofei/en/index.htm>