ASYMPTOTICS OF PREDICTIVE DISTRIBUTIONS DRIVEN BY SAMPLE MEANS AND VARIANCES

SAMUELE GARELLI, FABRIZIO LEISEN, LUCA PRATELLI, AND PIETRO RIGO

ABSTRACT. Let $\alpha_n(\cdot) = P(X_{n+1} \in \cdot | X_1, \ldots, X_n)$ be the predictive distributions of a sequence (X_1, X_2, \ldots) of *p*-dimensional random vectors. Suppose

$$\alpha_n = \mathcal{N}(M_n, Q_n)$$

where $M_n = \frac{1}{n} \sum_{i=1}^{n} X_i$ and $Q_n = \frac{1}{n} \sum_{i=1}^{n} (X_i - M_n)(X_i - M_n)^t$. Then, there is a random probability measure α on the Borel subsets of \mathbb{R}^p such that $\|\alpha_n - \alpha\| \stackrel{a.s.}{\longrightarrow} 0$ where $\|\cdot\|$ is total variation distance. An explicit expression for α is provided and the convergence rate of $\|\alpha_n - \alpha\|$ is shown to be arbitrarily close to $n^{-1/2}$. Moreover, it is still true that $\|\alpha_n - \alpha\| \stackrel{a.s.}{\longrightarrow} 0$ even if $\alpha_n = \mathcal{L}(M_n, Q_n)$ where \mathcal{L} belongs to a class of distributions much larger than the normal. The predictives α_n are useful in various frameworks, including Bayesian predictive inference and predictive resampling. Finally, the asymptotic behavior of copula-based predictive distributions (introduced in [15]) is investigated and a numerical experiment is performed.

1. INTRODUCTION

Let $X = (X_1, X_2, ...)$ be a sequence of random variables, with values in a standard Borel space (S, \mathcal{B}) , and

$$X(n) = (X_1, \dots, X_n).$$

The predictive distributions of X are

$$\alpha_0(\cdot) = P(X_1 \in \cdot)$$
 and $\alpha_n(\cdot) = P[X_{n+1} \in \cdot | X(n)]$ for each $n \ge 1$.

Thus, α_0 is the marginal distribution of X_1 and α_n the conditional distribution of X_{n+1} given X(n). Note that α_0 is a fixed probability measure on \mathcal{B} while α_n is a random probability measure on \mathcal{B} for $n \geq 1$.

By the Ionescu-Tulcea theorem, in order to assess the probability distribution of X, it suffices to specify the sequence (α_n) ; see e.g. [6]. Precisely, suppose the X_n are defined on a set Ω and $X(\Omega) = S^{\infty}$. (The condition $X(\Omega) = S^{\infty}$ is just to avoid unnecessary complications; it is automatically true if $\Omega = S^{\infty}$ and the X_n are the canonical projections). Select any sequence $(\alpha_n : n \ge 0)$ such that:

- (i) $\alpha_0 \in \mathcal{P}$ and $\alpha_n : \Omega \to \mathcal{P}$ for $n \ge 1$, where \mathcal{P} denotes the collection of all probability measures on \mathcal{B} ;
- (ii) The map $\omega \in \Omega \mapsto \alpha_n(\omega, A)$ is measurable with respect to $\sigma[X(n)]$ for fixed $n \ge 1$ and $A \in \mathcal{B}$.

²⁰²⁰ Mathematics Subject Classification. 60B10, 60G57, 62F15, 62E20.

Key words and phrases. Bayesian predictive inference, Conditional identity in distribution, Convergence of probability measures, Predictive distribution, Predictive resampling, Total variation distance.

Then, there is a unique probability measure P on $\sigma(X)$ which admits the α_n as predictive distributions. Importantly, apart from conditions (i)-(ii), the α_n are *arbitrary*. Roughly speaking, to assign the distribution of X, it suffices to select (arbitrarily) the marginal distribution of X_1 , the conditional distribution of X_2 given X_1 , the conditional distribution of X_3 given (X_1, X_2) , and so on. This fact has various consequences.

For instance, consider a Bayesian forecaster who aims to predict X_{n+1} based on X(n) for every n. To this end, it is (necessary and) sufficient to select (α_n) . This suggests a few remarks.

- To make Bayesian predictions, the data sequence X is not forced to have any specific distributional form (such as exchangeable, stationary, Markov, and so on).
- However, in the special case where X is required to be exchangeable, it is possible to make predictions without explicitly selecting a prior. In fact, in the exchangeable case, selecting a prior or selecting the predictives (α_n) are two equivalent strategies to determine the distribution of X. Sometimes, the second strategy (i.e. assigning (α_n) directly) is more convenient.
- By choosing (α_n) , the forecaster is attaching probabilities to observable facts only. The value of X_{n+1} is actually observable, while the prior (being a probability on some random parameter) does not necessarily deal with observable facts.
- In some Bayesian problems, even if the primary goal is not prediction, it may be convenient to assign the distribution of X through the choice of (α_n) . This happens mainly in Bayesian nonparametrics.

In a nutshell, these are the basic ideas underlying the predictive approach to Bayesian inference; see [4], [5], [6] and references therein.

Finally, in addition to Bayesian predictive inference, there are other frameworks where the choice of (α_n) is the main task. Without any claim of being exhaustive, we mention machine learning [7], [12], [13], [16], causal inference [18], species sampling sequences [19], [20], Dawid's prequential approach [8], [9], and martingale posterior distributions [14], [15].

1.1. Convergence of predictive distributions. After selecting the sequence (α_n) of predictives, a natural question is whether it converges, in some sense, as $n \to \infty$. To formalize this question, suppose X is defined on the probability space $(\Omega, \sigma(X), P)$ and α is a random probability measure (r.p.m.) on \mathcal{B} defined on such a probability space. Recall also that the *total variation distance* between two probability measures on \mathcal{B} , say μ and ν , is

$$\|\mu - \nu\| = \sup_{A \in \mathcal{B}} |\mu(A) - \nu(A)|.$$

Then, α_n converges weakly to α a.s. if

 $\alpha_n(\omega,\cdot) \xrightarrow{weakly} \alpha(\omega,\cdot), \quad \text{as } n \to \infty, \text{ for } P\text{-almost all } \omega \in \Omega.$

Similarly, α_n converges in total variation to α a.s. if

 $\|\alpha_n(\omega,\cdot) - \alpha(\omega,\cdot)\| \longrightarrow 0, \quad \text{as } n \to \infty, \text{ for } P\text{-almost all } \omega \in \Omega.$

Clearly, $\alpha_n \to \alpha$ in total variation a.s. implies $\alpha_n \to \alpha$ weakly a.s. but not conversely.

Usually, convergence of α_n is a desirable property. This claim can be supported by various examples and remarks. Here, we report three of them.

Example 1. Suppose $\alpha_n \to \alpha$ weakly a.s. for some r.p.m. α on \mathcal{B} . Then, $\mu_n \to \alpha$ weakly a.s. where $\mu_n = \frac{1}{n} \sum_{i=1}^n \delta_{X_i}$ is the empirical measure. Moreover, X is asymptotically exchangeable, in the sense that $(X_n, X_{n+1}, \ldots) \xrightarrow{dist} Z$, as $n \to \infty$, for some exchangeable sequence $Z = (Z_1, Z_2, \ldots)$. As an aside, exploiting convergence of (α_n) , one obtains the following characterization of exchangeability

X exchangeable \iff X stationary and (α_n) converges weakly a.s.

In fact, " \Rightarrow " is very well known; see e.g. [3]. As to " \Leftarrow ", just note that stationarity of X implies $X \sim (X_n, X_{n+1}, \ldots)$ for each n. Since (α_n) converges weakly a.s., one also obtains $(X_n, X_{n+1}, \ldots) \xrightarrow{dist} Z$ for some exchangeable Z. Hence, $X \sim Z$.

Example 2. Fix a σ -finite measure λ on \mathcal{B} and suppose that X(n) has a density with respect to λ^n for each n. Suppose also that X is exchangeable or, more generally, conditionally identically distributed (in the sense of [2]). In this case, there is a r.p.m. α on \mathcal{B} such that $\alpha_n \to \alpha$ weakly a.s. and an obvious question is whether α still admits a density with respect to λ . Precisely, this means that

(1)
$$\alpha(\omega, dx) = f(\omega, x) \lambda(dx)$$

for some non-negative measurable function f and P-almost all $\omega \in \Omega$. By Theorem 1 of [3], the answer is

Condition (1) holds
$$\iff \alpha_n \to \alpha$$
 in total variation a.s.

Example 3. A new method for making Bayesian inference, referred to as predictive resampling, has been introduced by Fong, Holmes and Walker in [14]. A quick description of this method is provided in Section 3.1. Here, we just note that a fundamental step of predictive resampling is the choice of a converging sequence (α_n) of predictive distributions. Indeed, in a specific problem, predictive resampling is more or less effective depending on whether the choice of (α_n) is more or less suitable to the available data. In [14], convergence of (α_n) is proved via martingale arguments and the corresponding posteriors are called martingale posteriors. However, martingales are not mandatory. In order to apply predictive resampling, it is enough to select (α_n) and to show that it converges (in some sense).

1.2. Our contribution. This paper introduces and investigates a new sequence (α_n) of predictive distributions for the case

 $S = \mathbb{R}^p$.

The α_n are specified in Section 2.2. Here, we just note that each α_n depends on the data X(n) only through the sample mean and covariance matrix. A meaningful example is

$$\alpha_n = \mathcal{N}(M_n, Q_n)$$

where $M_n = \frac{1}{n} \sum_{i=1}^n X_i$, $Q_n = \frac{1}{n} \sum_{i=1}^n (X_i - M_n) (X_i - M_n)^t$ and $\mathcal{N}(a, B)$ denotes the *p*-dimensional Gaussian distribution with mean *a* and covariance matrix *B*.

Our main result is that α_n converges, in total variation a.s., to a r.p.m. α . An explicit formula for α is provided and the rate of convergence is shown to be arbitrarily close to $n^{-1/2}$. Moreover, it is still true that $\|\alpha_n - \alpha\| \xrightarrow{a.s.} 0$ even if $\alpha_n = \mathcal{L}(M_n, Q_n)$ where \mathcal{L} belongs to a class of distributions much larger than \mathcal{N} .

In case $S = \mathbb{R}$, a natural competitor to the previous α_n are the copula-based predictive distributions, introduced in [15] and denoted by β_n in this paper. Hence, the asymptotic behavior of β_n is investigated as well. In Section 2.3, it is shown that β_n converges weakly a.s. but not necessarily in total variation a.s. Moreover, conditions for β_n to converge in total variation a.s. are provided.

Finally, in Section 3, the empirical behavior of the α_n is tested via a numerical experiment. In particular, α_n is compared with β_n as regards the speed of convergence. Moreover, both α_n and β_n are used to implement the predictive resampling method for parameter estimation.

A last remark is that, to support the α_n , we focused mainly on their behaviour in parameter estimation via predictive resampling. However, the α_n 's scope is much larger. As an obvious example, in Bayesian predictive inference, the α_n may be used to predict future observations.

2. Results

This section includes our main results. Any consideration regarding their practical application is postponed to Section 3. Similarly, to make the paper more readable, all the proofs are deferred to a final Appendix.

2.1. **Preliminaries and notation.** From now on, $S = \mathbb{R}^p$ where $p \ge 1$. Each point $x \in \mathbb{R}^p$ is regarded as a column vector and $x^{(i)}$ denotes the *i*-th coordinate of x. Similarly, if B is any matrix, $B^{(i,j)}$ is the (i,j)-th entry of B. We denote by \mathcal{M} the collection of symmetric positive definite matrices of order $p \times p$. We write $\mathcal{L}(a, B)$ to indicate the probability distribution of $a + B^{1/2}Z$, where $a \in \mathbb{R}^p$, B is a symmetric non-negative definite matrix of order $p \times p$, and Z is a p-dimensional random vector such that

E(Z) = 0, $E(ZZ^t) = I$, $E\{(Z^tZ)^{u/2}\} < \infty$ for some u > 2, and Z has a density with respect to Lebesgue measure on \mathbb{R}^p .

In the previous notation, p is understood. We will write \mathcal{M}_p or $\mathcal{L}_p(a, B)$ instead of \mathcal{M} or $\mathcal{L}(a, B)$ every time that a mention of p is necessary.

Finally, given any sequence (α_n) of predictive distributions, convergence is always meant with respect to the probability measure P under which X has predictives α_n . As noted in Section 1, such a P exists and is unique by the Ionescu-Tulcea theorem.

2.2. Definition and asymptotics of the new predictive distributions. Let

$$M_n = \frac{1}{n} \sum_{i=1}^n X_i, \qquad Q_n = \frac{1}{n} \sum_{i=1}^n (X_i - M_n)(X_i - M_n)^t,$$

and

(2)
$$\alpha_0 = \mathcal{L}(0, I), \quad \alpha_1 = \mathcal{L}(X_1, I) \text{ and } \alpha_n = \mathcal{L}(M_n, Q_n) \text{ for } n \ge 2.$$

The predictives α_n have a simple form and can be easily handled in real problems. Among other things, they can serve in prediction problems and/or in applying the predictive resampling method; see Section 3. In addition, the α_n have a nice asymptotic behavior. This fact is formalized by the next two results, which are the main contributions of this paper.

Theorem 4. If the α_n are defined by (2), then $M_n \xrightarrow{a.s.} M$ and $Q_n \xrightarrow{a.s.} Q$ where M is a p-dimensional random vector and Q a random matrix of order $p \times p$. In addition, $Q \in \mathcal{M}$ a.s., so that

$$\|\alpha_n - \alpha\| \xrightarrow{a.s.} 0$$
 where $\alpha = \mathcal{L}(M, Q)$.

By Theorem 4, α_n converges in total variation a.s. to $\alpha = \mathcal{L}(M, Q)$, where M and Q are the a.s. limits of the sample means and the sample covariance matrices, respectively. As noted in Section 1.1, this asymptotic behavior of α_n is useful in various frameworks. To make Theorem 4 more effective, however, it is desirable to grasp some information on the convergence rate of α_n to α . This is actually possible if $\mathcal{L}(a, B) = \mathcal{N}(a, B)$.

Theorem 5. Let α_n , M and Q be as in Theorem 4. Suppose $\mathcal{L}(a, B) = \mathcal{N}(a, B)$, define $\alpha = \mathcal{N}(M, Q)$, and fix any sequence (d_n) of constants. Then,

$$d_n \|\alpha_n - \alpha\| \xrightarrow{P} 0$$
 whenever $\frac{d_n}{\sqrt{n}} \to 0$.

To better appreciate Theorem 5, it is worth noting that $\sqrt{n} \|\alpha_n - \alpha\|$ fails to converge to 0 in probability. We prove this fact for p = 1. In this case, for each $x \in \mathbb{R}$, one obtains

$$\|\alpha_n - \alpha\| \ge \left|\alpha_n \left((-\infty, x]\right) - \alpha \left((-\infty, x]\right)\right| = \left|\Phi\left(\frac{x - M_n}{\sqrt{Q_n}}\right) - \Phi\left(\frac{x - M}{\sqrt{Q}}\right)\right|$$

where Φ is the standard normal distribution function. Letting $x = M_n$, it follows that

$$\begin{split} \sqrt{n} \|\alpha_n - \alpha\| &\ge \sqrt{n} \left| \Phi(0) - \Phi\left(\frac{M_n - M}{\sqrt{Q}}\right) \right| = \frac{\sqrt{n}}{\sqrt{2\pi}} \int_0^{\frac{|M_n - M|}{\sqrt{Q}}} e^{-t^2/2} dt \\ &\ge \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{(M_n - M)^2}{2Q}\right) \frac{\sqrt{n}}{\sqrt{Q}} |M_n - M|. \end{split}$$

However, $\frac{\sqrt{n}}{\sqrt{Q}}(M_n - M)$ converges in distribution to the $\mathcal{N}(0, 1)$. Hence, the previous inequality implies that $\sqrt{n} \|\alpha_n - \alpha\|$ does not converge to 0 in probability.

2.3. Copula-based predictive distributions. A natural competitor of the predictives α_n of Section 2.2 are the copula-based predictive distributions of [15].

A bivariate *copula* is a distribution function on \mathbb{R}^2 whose marginals are uniform on (0, 1). The density of a bivariate copula, provided it exists, is said to be a *copula density*. (Henceforth, all densities are meant with respect to Lebesgue measure).

Let $S = \mathbb{R}$. Fix a density f_0 , a sequence (c_n) of bivariate copula densities, and a sequence (r_n) of constants in (0, 1]. For the sake of simplicity, assume $f_0 > 0$ and

 $\mathbf{6}$

 $c_n > 0$ for all $n \ge 0$. Then, a sequence (β_n) of predictive distributions may be built as follows. Call F_0 the distribution function corresponding to f_0 and define

$$f_1(x \mid y) = (1 - r_0) f_0(x) + r_0 f_0(x) c_0 \Big(F_0(x), F_0(y) \Big) \quad \text{for } x, y \in \mathbb{R}.$$

Note that $f_1 : \mathbb{R}^2 \to \mathbb{R}^+$ is a measurable function such that $f_1(\cdot | y)$ is a density for each fixed $y \in \mathbb{R}$. Hence, if $F_1(\cdot | y)$ denotes the distribution function corresponding to $f_1(\cdot | y)$, one can let

$$f_2(x \mid y, z) = (1 - r_1) f_1(x \mid y) + r_1 f_1(x \mid y) c_1 \Big(F_1(x \mid y), F_1(z \mid y) \Big)$$

for $x, y, z \in \mathbb{R}$. In general, given a measurable function $f_n : \mathbb{R}^{n+1} \to \mathbb{R}^+$ such that $f_n(\cdot | y)$ is a density for each fixed $y \in \mathbb{R}^n$, one can define

$$f_{n+1}(x \mid y, z) = (1 - r_n) f_n(x \mid y) + r_n f_n(x \mid y) c_n \Big(F_n(x \mid y), F_n(z \mid y) \Big)$$

for all $x, z \in \mathbb{R}$ and $y \in \mathbb{R}^n$. In the above formula, $F_n(\cdot | y)$ denotes the distribution function corresponding to $f_n(\cdot | y)$. Proceeding in this way, one obtains a collection of densities $\{f_n(\cdot | y) : n \ge 1, y \in \mathbb{R}^n\}$ and the β_n can be defined as

$$\beta_0(dx) = f_0(x) dx$$
 and $\beta_n(dx) = f_n(x \mid X(n)) dx$ for $n \ge 1$.

The predictives β_n have been introduced in [15] and then used in [14]. The construction given here is from [6].

If the predictive distributions of X are the β_n , then X is conditionally identically distributed. This is proved in [6] when $r_n = 1$ but the proof of [6] can be easily extended to any choice of r_n . Therefore, since X is conditionally identically distributed, there is a r.p.m. β such that

$$\beta_n \to \beta$$
 weakly a.s.

A natural question is whether $\beta_n \to \beta$ in total variation a.s., or equivalently

(3)
$$\beta(dx) = f(x) dx$$
 for some random density f ;

see Example 2. Among other things, condition (3) is tacitly assumed in [14] and [15]. As we now prove, however, it is not necessarily true.

Example 6. Condition (3) fails whenever

 $\limsup r_n > 0 \quad \text{and} \quad c_n = c \quad \text{for all } n \ge 0$

where c is any bivariate copula density such that $\int_0^1 |c(u, v) - 1| du > 0$ for each $v \in (0, 1)$. In fact, suppose the predictives of X are the β_n and define

$$D_n = \int_{-\infty}^{\infty} \left| f_{n+1} \left(x \mid X(n+1) \right) - f_n \left(x \mid X(n) \right) \right| dx.$$

Since X is conditionally identically distributed, condition (3) amounts to

$$\int_{-\infty}^{\infty} \left| f_n(x \mid X(n)) - f(x) \right| dx \xrightarrow{a.s.} 0 \quad \text{for some random density } f.$$

Hence, under (3), one obtains $D_n \xrightarrow{a.s.} 0$. However,

$$D_n = r_n \int_{-\infty}^{\infty} \left| c \Big(F_n(x \mid X(n)), F_n(X_{n+1} \mid X(n)) \Big) - 1 \Big| f_n(x \mid X(n)) \, dx \right|$$
$$= r_n \int_0^1 \left| c \Big(u, F_n(X_{n+1} \mid X(n)) \Big) - 1 \Big| \, du.$$

Conditionally on X(n), the random variable $F_n(X_{n+1} | X(n))$ is uniformly distributed on (0, 1). Hence, D_n is distributed as

$$r_n \int_0^1 |c(u, V) - 1| \, du$$

where V denotes any random variable with uniform distribution on (0, 1). Since $\int_0^1 |c(u, V) - 1| du > 0$ a.s. and $\limsup_n r_n > 0$, it follows that D_n does not converge to 0 a.s. Therefore, condition (3) fails.

Example 6 should be compared with Theorem 4. According to the latter, $\alpha_n \to \alpha$ in total variation a.s. where the α_n are given by (2) and $\alpha = \mathcal{L}(M,Q)$. Since $\mathcal{L}(0,I)$ admits a density, one obtains $\alpha(dx) = f(x) dx$ for some random density f.

Even if not generally true, however, condition (3) holds under some assumptions. We finally prove two results of this type. As probably expected, condition (3) holds provided $r_n \rightarrow 0$ fastly enough.

Theorem 7. Condition (3) holds whenever $\sum_{n} r_n < \infty$.

In the second result, the condition $\sum_n r_n < \infty$ is weakened at the price of a boundedness assumption on the copula densities c_n .

Theorem 8. Condition (3) holds whenever $\int_{-\infty}^{\infty} f_0(x)^2 dx < \infty$, $\sum_n r_n^2 < \infty$, and there is a constant b such that

 $c_n(u,v) \leq b$ for all $n \geq 0$ and $u, v \in [0,1]$.

Theorem 8 applies to any sequence c_n of bivariate copula densities, but it requires a strong boundedness condition on c_n . As expected, the latter condition can be dropped in some special cases. For instance, as proved in [14, Theorem 5], it can be dropped if $c_n = c_\rho$ for all n where c_ρ is a Gaussian copula density with correlation $\rho < 1/\sqrt{3}$.

3. Numerical illustrations

This section reports the results of an empirical comparison between the predictives α_n of Section 2.2 and the copula-based predictive distributions β_n . The comparison is based on the speed of convergence and the performance in Bayesian parametric inference when applying the *predictive resampling* (PR) method.

We let p = 1. Hence, $\alpha_0 = \mathcal{N}(0, 1)$, $\alpha_1 = \mathcal{N}(X_1, 1)$ and $\alpha_n = \mathcal{N}(M_n, Q_n)$ for n > 1. In turn, the β_n are built as in Section 2.3 with

 $f_0 =$ standard normal density and $c_n = c_\rho$ for all $n \ge 0$

where c_{ρ} is the copula density corresponding to a bivariate normal distribution with mean 0, variance 1 and correlation $\rho \in (0, 1)$; see [15, Example 2]. Note that $\alpha_0 = \beta_0 = \mathcal{N}(0, 1)$. To define the β_n , we also need a sequence (r_n) of weights. For reasons that are pointed out in Sections 3.2 and 3.3, (r_n) is specified in two different ways.

We begin with a brief recap on PR.

3.1. **Predictive resampling.** The PR method, introduced in [14], allows to make inference on a random parameter and/or to predict future observations. In a Bayesian framework, it is an alternative to the usual likelihood/prior scheme, and it has various similarities with the predictive approach to statistics [6]. In particular, the explicit assignment of a prior distribution is not required.

Suppose the object of inference is a function of the sequence X, say

$$\theta = f(X) = f(X_1, X_2, \ldots)$$

where f is a known measurable function. As an obvious example, think of θ as the mean of a population described by X, that is

(4)
$$\theta = \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} X_i$$
 (where the limit is assumed to exist a.s.)

More generally, the informal idea is that the uncertainty about θ would disappear if the whole data sequence X could be observed. The goal is to estimate the distribution of θ based on the available data.

To achieve this target, we first select a sequence (α_n) of predictive distributions. Moreover, we assume that a dataset

$$x = (x_1, \ldots, x_s)$$

has been observed, where s denotes the time when PR is applied. Then, through (α_n) and x, we draw N more observations $(X_{s+1}, \ldots, X_{s+N})$ and we use the data $(x, X_{s+1}, \ldots, X_{s+N})$ to estimate θ . This procedure is repeated a number B of times in order to end up with a sample of estimates of θ . More precisely, let $\hat{\theta} = \hat{\theta}(x, X_{s+1}, \ldots, X_{s+N})$ denote the adopted estimate of θ . For instance, if θ is given by (4), it is quite natural to let

$$\hat{\theta} = \frac{\sum_{i=1}^{s} x_i + \sum_{i=1}^{N} X_{s+i}}{N+s}.$$

Once $\hat{\theta}$ has been chosen, PR can be outlined as follows.

for i = 1, ..., B do for j = 1, ..., N do Sample $X_{s+j} \sim \alpha_{s+j-1}$ Update: $\{X_{s+j}, \alpha_{s+j-1}\} \mapsto \alpha_{s+j}$ end Evaluate $\theta_i = \hat{\theta}(x, X_{s+1}, ..., X_{s+N})$ end Return $\boldsymbol{\theta} = (\theta_1, ..., \theta_B)$.

Note that, since the data $x = (x_1, \ldots, x_s)$ are available, we start sampling from α_s and not from α_0 . Moreover, each x_i is implicitly assumed to have been drawn from α_{i-1} , for $i = 1, \ldots, s$.

The final output of PR is a random sample $(\theta_1, \ldots, \theta_B)$ from the probability distribution of $\hat{\theta}(x, X_{s+1}, \ldots, X_{s+N})$. In particular, $\theta_1, \ldots, \theta_B$ are i.i.d. and their common distribution $\Pi_N(\cdot \mid x)$ can be written as

$$\Pi_N(A \mid x) = \int_{\mathbb{R}^N} \mathbb{1}_A\{\hat{\theta}(x, x_{s+1}, \dots, x_{s+N})\} \prod_{j=1}^N \alpha_{s+j-1}(x, x_{s+1}, \dots, x_{s+j-1})(dx_{s+j})$$

 $\Pi_N(\cdot \mid x)$ can be regarded as the PR analogue of the usual posterior distribution and it is used to approximate the distribution of θ given x, i.e.

$$\Pi_{\infty}(A \mid x) = P(f \in A \mid x)$$

where $P(\cdot | x)$ denotes the probability law induced by the predictives $(\alpha_n : n \ge s)$ according to the Ionescu-Tulcea theorem; see Section 1.

So far, (α_n) is any sequence of predictives. However, for PR to make sense, α_n is required to converge (in some sense) as $n \to \infty$. Convergence is a sort of consistency condition. Roughly speaking, the population we are trying to reconstruct should mimic the observed sample. Essentially, it should be a version of the sample with similar shape and location but more uncertainty. This informal idea is realized by regarding such a population as distributed according to α , where α is the limit of the α_n . In addition, it is crucial that $\alpha_s(x)$ fits the observed data x or, at least, shares with x those features that we are estimating. Heuristically, these two requirements (convergence of α_n and good fit to x) guarantee that $\Pi_N(\cdot \mid x)$ approximates $\Pi_{\infty}(\cdot \mid x)$.

3.2. **Speed of convergence.** Let f_n^{α} and f_n^{β} denote the densities of α_n and β_n (with respect to Lebesgue measure). Following [14], to evaluate the speed of convergence, we computed the L^1 -distances $||f_0 - f_n^{\alpha}||_1$ and $||f_0 - f_n^{\beta}||_1$ where f_0 is the standard normal density. Roughly speaking, the informal idea is that, for any sequence (f_n) of random densities, the value of n at which $||f_0 - f_n||_1$ plateaus can be regarded as the index after which f_n is close to its limit (provided it exists).

Recall that, thanks to Theorem 4, $||f_n^{\alpha} - f||_1 \xrightarrow{a.s.} 0$ for some random density f. As for β_n , we focus on two different situations. In the first, β_n does not necessarily converge in total variation a.s. (at least to our knowledge) while in the second it does, i.e.

a) $r_n = r_n^{(a)} = \left(2 - \frac{1}{n+1}\right) \frac{1}{n+2}$ and $\rho = 0.9$. This choice of r_n , inspired by the Dirichlet process mixture model, is proposed in [14]. However, Theo-

rems 7-8 do not apply and the same happens for Theorem 5 of [14] since $\rho > 1/\sqrt{3}$. Hence, whether or not β_n converges in total variation a.s. is unknown.

unknown. b) $r_n = r_n^{(b)} = \left(2 - \frac{1}{n+2}\right) \frac{1}{(n+3)\left(\log(n+3)\right)^2}$ and $\rho = 0.9$. In this case, $\sum_n r_n < \infty$ so that β_n converges in total variation a.s. by Theorem 7.

To evaluate $||f_0 - f_n^{\alpha}||_1$, we ran PR starting from α_0 (and not from $\alpha_s(x)$ as in Section 3.1). In other terms, we just sampled $X_1 \sim \alpha_0$, $X_2 \sim \alpha_1(X_1)$, $X_3 \sim \alpha_2(X_1, X_2)$ and so on. Then, at each step of the algorithm, we evaluated f_n^{α} on a grid of points and computed $||f_0 - f_n^{\alpha}||_1$ numerically. Exactly the same procedure has been followed for $||f_0 - f_n^{\beta}||_1$.

Figure 1 shows the path of $||f_0 - f_n^{\alpha}||_1$. It takes about 150 iteration of PR for the L^1 -distance to stabilise.



FIGURE 1. α_n

Figure 2 shows the path of $||f_0 - f_n^\beta||_1$ for both $r_n^{(a)}$ and $r_n^{(b)}$ (with $\rho = 0.9$). In the first case, it takes about 4000 iteration of PR for the L^1 -distance to stabilise, while in the second the behaviour of $||f_0 - f_n^\beta||_1$ is similar to that of $||f_0 - f_n^\alpha||_1$. There might be two reasons for this discrepancy. Either β_n fails to converge in total variation a.s. when $r_n = r_n^{(a)}$ and $\rho = 0.9$. Or, $r_n^{(b)}$ approaches 0 so much faster than $r_n^{(a)}$ causing a faster convergence of $||f_0 - f_n^\beta||$.



FIGURE 2. β_n

3.3. **Parameter estimation.** We now compare α_n and β_n when used in PR to estimate the posterior distribution $\Pi_N(\cdot | x)$ of a parameter θ ; see Section 3.1. Such a θ is taken to be the mean or the variance of a reference population. In the sequel, $\hat{\theta}$ is short for $\hat{\theta}(x, X_{s+1}, \ldots, X_{s+N})$. Obviously, $\hat{\theta}$ is the sample mean of $(x, X_{s+1}, \ldots, X_{s+N})$ if θ is the population mean, while it is the sample variance of $(x, X_{s+1}, \ldots, X_{s+N})$ if θ is the population variance. Moreover, when necessary, we use the notations $\Pi_N^{\alpha}(\cdot | x)$ and $\Pi_N^{\beta}(\cdot | x)$ to stress that $\Pi_N(\cdot | x)$ has been obtained using the α_n or the β_n .

The simulation study has been organised as follows. We drew random samples of size s = 50, 100, 500, 1000, 1500, 2000 from a mixture of three shifted t-distributions with weights $w_1 = 0.3, w_2 = 0.1$ and $w_3 = 0.6$, shifting constants $\mu_1 = -5, \mu_2 = 0$ and $\mu_3 = 4$ and degrees of freedom $\nu = 3$. Then, we ran PR with N = 50, 500, 5000and B = 1000. Once a sample $\boldsymbol{\theta} = (\theta_1, \dots, \theta_B)$ from $\Pi_N(\cdot \mid x)$ was obtained, we applied kernel density estimation on $\boldsymbol{\theta}$ to estimate the posterior density $\pi_N(\cdot \mid x)$ of $\Pi_N(\cdot \mid x)$. We also kept track of the computational times. For the predictives β_n , we considered both sequences of weights $r_n^{(a)}$ and $r_n^{(b)}$ defined in Section 3.2. B was chosen following [14]. In general, we found B = 1000 allows to visualise well the posterior density and does not make PR too slow. As for s and N, we decided to consider different values in order to see how $\pi_N(\cdot \mid x)$ varies when more information is available (i.e. s gets larger) and when more uncertainty about the observed sample is reconstructed (i.e. N gets larger). When θ is the population mean, $\pi_N^{\beta}(\cdot \mid x)$ has been obtained with $r_n = r_n^{(a)}$. Instead, when θ is the population variance, $\pi_N^{\beta}(\cdot \mid x)$ has been obtained with $r_n = r_n^{(b)}$. There are no relevant differences between the effect of the two sequences of weights on $\pi_N^{\beta}(\cdot \mid x)$, but we wanted to provide evidence that both perform well.

Figure 3 shows the graph of $\pi_N(\cdot | x)$ when θ is the population mean. As a general pattern, note that the posterior variance of $\hat{\theta}$ increases as N gets larger. This might not be clear from the plots because of the different scaling of the axes in the sub-figures corresponding to different values of N. Therefore, as an example, in Table 2 we report the posterior variance of $\hat{\theta}$ for s = 2000. On the other hand, N has no effect on the posterior mean of $\hat{\theta}$. This can already be noted from the plots, but see also Table 1. Such a behaviour is precisely what we would expect from PR when the predictives are designed as required in Section 3.1, i.e. they converge and fit the observed data x. Recall that the goal of PR is to reconstruct uncertainty about the observed sample without changing its shape and location. As shown by Figure 3, this reflects into $\pi_N(\cdot | x)$ having larger variance as N increases, but keeping the same mean and number of modes. For α_n this behaviour can be motivated even more explicitly. Indeed, it is quite straightforward to see that

$$E_N^{\alpha}(\hat{\theta} \mid x) = \bar{x}_s \quad \text{and} \quad Var_N^{\alpha}(\hat{\theta} \mid x) = \hat{\sigma}_s^2 \frac{s}{s+1} \sum_{i=1}^N \frac{s+i}{(s+i-1)(s+i)^2},$$

where \bar{x}_s and $\hat{\sigma}_s^2$ are the mean and the variance of the observed dataset x. This shows that $\pi_N^{\alpha}(\cdot \mid x)$ is centred into the sample mean, while its variance increases in N.

The effect of the sample size s is quite different between $\pi_N^{\alpha}(\cdot \mid x)$ and $\pi_N^{\beta}(\cdot \mid x)$. In both cases, the posterior density that best grasps θ , being basically centred on its value, is the one corresponding to s = 1000. Nevertheless, for the other values of s, the former performs slightly better than the latter. Indeed, θ always falls in the main body of $\pi_N^{\alpha}(\cdot \mid x)$, while it is on the tails of $\pi_N^{\beta}(\cdot \mid x)$ or even quite far from it. Since $E_N^{\alpha}(\hat{\theta}|x) = \bar{x}_s \xrightarrow{a.s.} \theta$ as $s \to \infty$, using the α_n it is quite natural to expect that estimates are improved by a larger sample size. What is more subtle is the combined effect of s and N on $\pi_N^{\alpha}(\cdot \mid x)$. Indeed, since a bigger N implies a larger posterior variance, this causes the posterior to catch the true value of θ even for those values of s for which $\pi_N^{\alpha}(\cdot \mid x)$ is rather distant from θ when N is small. As an example, for s = 500, look at how $\pi_N^{\alpha}(\cdot \mid x)$ (green curve) changes as N increases.

A last remark on Figure 3 is in order. As mentioned in Section 3.1, for PR to perform well, the starting predictive distribution (i.e., $\alpha_s(x)$ or $\beta_s(x)$) should fit x well or at least preserve those features that we are interested in estimating. More explicitly, if we want to estimate the posterior distribution of the mean of a population, the predictive mean at time s should be close to \bar{x}_s . This is precisely the case of $\alpha_s(x)$. On the contrary, $\beta_s(x)$ is a much more complicated object. In particular, as shown extensively in [14], f_s^{β} is a very effective density estimator. Hence, Figure 3 is an evidence of the fact that to estimate the posterior of a parameter it is enough that the predictive distribution preserves the information about that parameter originating from the observed data. Indeed, both methods provide satisfying estimates (the α_n are even slightly better) and implementing the simpler one is even computationally more efficient. The latter statement will be clarified at the end of this section, but now we can already note that estimating two parameters (i.e. mean and variance) is obviously faster than estimating an entire density.

| N | 50 | 500 | 5000 | N | 50 | 500 | 5000 |
|------------|-------|-------|-------|------------|--------|--------|-------|
| α_n | 0.926 | 0.926 | 0.923 | α_n | 0.0002 | 0.0019 | 0.006 |
| β_n | 0.925 | 0.926 | 0.926 | β_n | 0.0002 | 0.0015 | 0.001 |

TABLE 1. $E_N(\hat{\theta} \mid x)$ for $\theta =$ population mean and s = 2000.

| TABLE 2. | $Var_N(\theta \mid$ | x) for $\theta =$ |
|------------|---------------------|-------------------|
| population | mean and | s = 2000. |

Figure 4 shows the posterior distribution when θ is the population variance. The effect of N and s on $\pi_N(\cdot | x)$ is basically the same as in Figure 3, as much as the comparison between $\pi_N^{\alpha}(\cdot | x)$ and $\pi_N^{\beta}(\cdot | x)$. Even in this case, the former is more precise in catching the value of θ . We just mention an additional detail about variance estimation when using the α_n . In this case, one obtains

$$E_N^{\alpha}(\hat{\theta} \mid x) = \hat{\sigma}_s^2 \frac{s}{s+1} \frac{s+N+1}{s+N}$$

and

$$Var_{N}^{\alpha}(\hat{\theta} \mid x) = \hat{\sigma}_{s}^{4} \left\{ \prod_{i=1}^{N} \left[1 + \frac{3}{(s+i)^{4}} - \frac{4}{(s+i)^{3}} \right] - \left[\frac{s}{(s+1)} \frac{s+N+1}{s+N} \right]^{2} \right\}.$$

Therefore, $\pi_N^{\alpha}(\cdot \mid x)$ is still centered in the sample statistic of the parameter of interest (with some bias that vanishes as $s \to \infty$) and its dispersion increases as $N \to \infty$ (at least numerically).







SAMUELE GARELLI, FABRIZIO LEISEN, LUCA PRATELLI, AND PIETRO RIGO

FIGURE 4. $\pi_N(\cdot | x)$ for θ = population variance = 19.2 and s = 50 (-), 100 (-), 500 (-), 1000 (-), 1500 (-), 2000 (-).

We next make some general comments on the posterior estimates obtained with α_n and β_n . As highlighted by Figures 3 and 4, $\pi_N^{\alpha}(\cdot \mid x)$ is usually more precise than $\pi_N^\beta(\cdot \mid x)$. The reason may be that in the first case we can easily control the centre of the posterior distribution. Nevertheless, for parameters different from mean and variance (e.g. skewness, kurtosis, quantiles, etc.) β_n is expected to provide better estimates. Indeed, α_n does not preserve information from the observed sample about any parameter other than mean and variance. On the contrary, since β_n is essentially a density estimator, it grasps the shape of the whole distribution. Furthermore, according to us, the performance of β_n could be enhanced in two (obvious) ways. Firstly, the correlation coefficient $\rho \in (0, 1)$, involved in the Gaussian copula density c_{a} , has to be estimated. To do so, following [14], we optimised the prequential log-likelihood for $\rho \in \{0.05, 0.1, \dots, 0.95\}$. By taking a finer grid, β_n might have a better fit on the observed data and this might make estimates more precise. The second possible improvement is related to the mechanism adopted to sample from β_n . Indeed, X_{s+1}, \ldots, X_{s+N} are generated via inverse sampling from a grid $\{y_1, \ldots, y_G\}$. Thus, once again, more accurate posterior estimates could be obtained by refining such grid. It is worth stressing that, if one wants only to update β_n into β_{n+1} , it is not necessary to sample $X_{n+1} \sim \beta_n$ and plug it in β_{n+1} . As explained in [14, Sec. 4.2], it is enough to draw an observation from a uniform distribution on (0, 1). But clearly, we cannot use this shortcut for we need to know the actual value of the sampled variables. Finally, we note that these improvements come at the cost of longer computational times. We explore this further in the next paragraph.

Tables 3 and 4 show the time needed to perform PR for each value of s and N. Algorithms for the implementation of PR have been coded in Rcpp. Computations have been carried out on a single-core version of R installed on a machine with an AMD Epyc 7763 2.45Ghz processor having 2TB RAM. As expected, in both cases PR is slower as N and s increase. The difference between the computational times of α_n and β_n , with the former being much faster than the latter, is quite evident. Reasons for this discrepancy have partially been pointed out already, so we summarise them and provide some more details. Sampling from α_n requires just to compute a mean and a variance and then draw an observation from a Gaussian distribution, for which efficient algorithms are available in any statistical software. On the other hand, sampling from β_n entails estimating the hyperparameter ρ and doing inverse sampling. The time needed for these tasks depends on the size of the grid of values taken to perform them. Such a grid cannot be too coarse, otherwise the accuracy of the estimates is worsened. Therefore, these two steps are likely to be the main reason why α_n is faster than β_n . Furthermore, the estimation of ρ is affected by the sample size s. Indeed, as explained in [14], optimizing the prequential log-likelihood requires to evaluate recursively $f_{i-1}^{\beta}(x_i \mid x_1, \ldots, x_{i-1})$, for each $i = 1, \ldots, s$ and each candidate value of ρ , and then pick the one that maximises $\sum_{i=1}^{s} \log f_{i-1}^{\beta}(x_i \mid x_1, \ldots, x_{i-1})$. This procedure is slower the larger the value of s is and has to be carried out before starting PR, because we cannot sample from β_n without specifying ρ . On the other hand, sampling from α_n does not require any such complicated initialisation step. Therefore, s has a much stronger effect on β_n than it has on α_n . To see this, look for instance at the increase in computational time from s = 50 to s = 2000, when N = 5000. In this case, PR with α_n is about 14 seconds slower while with β_n it is more than 2 minutes slower.

| | s = 50 | s = 100 | s = 500 | s = 1000 | s = 1500 | s = 2000 |
|----------|--------|---------|---------|----------|----------|----------|
| N = 50 | 0.0004 | 0.0005 | 0.001 | 0.0016 | 0.0032 | 0.0029 |
| N = 500 | 0.0074 | 0.0081 | 0.0133 | 0.0179 | 0.0241 | 0.0299 |
| N = 5000 | 0.3431 | 0.3498 | 0.4011 | 0.4591 | 0.521 | 0.5818 |

TABLE 3. Computational times for α_n (in minutes).

| | s = 50 | s = 100 | s = 500 | s = 1000 | s = 1500 | s = 2000 |
|----------|--------|---------|---------|----------|----------|----------|
| N = 50 | 0.0852 | 0.0988 | 0.3316 | 0.8326 | 1.5237 | 2.3548 |
| N = 500 | 0.6719 | 0.6313 | 0.8787 | 1.4696 | 2.1769 | 3.0014 |
| N = 5000 | 6.547 | 5.9677 | 6.0797 | 7.0695 | 7.9625 | 8.6769 |

TABLE 4. Computational times for β_n (in minutes).

4. Discussion

Let (X_n) be a sequence of p-dimensional random vectors and

$$\alpha_n(\cdot) = P(X_{n+1} \in \cdot \mid X_1, \dots, X_n)$$

the corresponding predictive distributions. In this paper, a new sequence (α_n) of predictives is introduced. The main feature of α_n is to depend on (X_1, \ldots, X_n) only through their mean and covariance matrix. Convergence in total variation of α_n is proved, along with an explicit formula for the limit r.p.m. In the (meaningful) special case where the α_n are Gaussian, the convergence rate is shown to be $n^{-1/2}$. A natural competitor of α_n is the copula-based predictive distribution β_n of [15]. Thus, β_n is investigated as well. It turns out that β_n converges weakly but not necessarily in total variation. Hence, sufficient conditions for β_n to converge in total variation are provided. Eventually, an empirical comparison between α_n and β_n is performed, assessing their speed of convergence and their ability to estimate, via PR, the posterior distribution of the mean and the variance of a reference population.

In all the numerical experiments, the kernel of α_n (denoted \mathcal{L}) is Gaussian. One reason is that, to estimate the posterior of a parameter θ via PR, it is enough for the predictive distribution to preserve the information about θ contained in the observed data. Hence, we chose $\mathcal{L} = \mathcal{N}$. In fact, when θ is the mean or the variance, it is straightforward to incorporate such information into a Gaussian distribution. Most probably, for estimating other parameters (e.g. skewness, kurtosis, quantiles, etc.) this simple model is no longer appropriate. Perhaps, the α_n should be enabled to capture the whole distribution of the observed data. Roughly speaking, a version of α_n more similar to β_n is likely to work better. A natural example could be to set \mathcal{L} as a mixture of Gaussian distributions, each of them centred in the main clusters of the observed data.

Moving forward from the idea of taking \mathcal{L} as a mixture of Gaussians, it would be reasonable to set α_0 as a mixture distribution (possibly data-dependent) but then update only the mean and the variance of the components from which the observations are drawn (along with the corresponding weights). It would be interesting to study the asymptotics of the resulting sequence of predictives. In addition, these predictives might result into more precise estimates, since the starting distribution and the updating mechanism preserve more of the information contained in the observed data than just their mean and variance. Hence, they would be relevant even from a statistical point of view.

We finally mention two further issues. The first is to give conditions on \mathcal{L} under which the convergence rate of α_n is still $n^{-1/2}$ (as it happens when $\mathcal{L} = \mathcal{N}$). The second is using the α_n in regression problems (as done in [14] with the β_n). In this case, the data are of the type $X_n = (Y_n, Z_n)$, where Y_n is a response variable and Z_n a vector of covariates. Then, the predictives α_n could be applied to estimate, via PR, the conditional distribution of the response given the observed values of the covariates.

APPENDIX

We close the paper by proving the results of Section 2. To this end, we first recall two known facts and we make a preliminary remark.

Lemma 9. If $x, y \in \mathbb{R}^p$ and B is a real non-singular matrix of order $p \times p$, then

$$det(B + xy^t) = det(B)(1 + x^t B^{-1}y).$$

Lemma 10. If $(V_n : n \ge 1)$ is an i.i.d. sequence of real random variables such that $E(V_1) = 0$ and $E(|V_1|^{1+\epsilon}) < \infty$, for some $\epsilon > 0$, then

the series
$$\sum_{n=1}^{\infty} \frac{V_n}{n}$$
 converges a.s.

Proof. Since

$$\sum_{n} P(|V_{n}| > n) = \sum_{n} P(|V_{1}| > n) \le E(|V_{1}|) < \infty,$$

the Borel-Cantelli lemma yields

 $P(|V_n| > n \text{ for infinitely many } n) = 0.$

Hence, it suffices to show that $\sum_n \frac{W_n}{n}$ converges a.s., where

$$W_n = V_n \, \mathbb{1}_{\{|V_n| \le n\}}.$$

To this end, first note that $\frac{W_n - E(W_n)}{n}$ are independent, centered, and

$$\sum_{n} E\left\{ \left(\frac{W_n - E(W_n)}{n}\right)^2 \right\} \le \sum_{n} E\left\{\frac{V_1^2 \, \mathbf{1}_{\{|V_1| \le n\}}}{n^2}\right\} \le \sum_{n} \frac{E\left(|V_1|^{1+\epsilon}\right)}{n^{1+\epsilon}} < \infty.$$

Hence, the series $\sum_{n} \frac{W_n - E(W_n)}{n}$ converges a.s. Finally, $E(V_1) = 0$ implies $E(W_n) = E\{V_1 \mid 1 \le r_1\} = -E\{V_1 \mid 1 \le r_2\}$.

$$E(W_n) = E\{V_1 \, \mathbb{1}_{\{|V_1| \le n\}}\} = -E\{V_1 \, \mathbb{1}_{\{|V_1| > n\}}\}.$$

Therefore,

$$\sum_{n} \frac{|E(W_n)|}{n} \le \sum_{n} \frac{E\{|V_1| \, \mathbb{1}_{\{|V_1| > n\}}\}}{n} \le \sum_{n} \frac{E(|V_1|^{1+\epsilon})}{n^{1+\epsilon}} < \infty.$$

This concludes the proof.

17

Both the previous lemmas are well known. A possible reference for Lemma 9 is [11], while Lemma 10 has been proved for we do not know of any explicit reference. Incidentally, in Lemma 10, the condition $E(|V_1|^{1+\epsilon}) < \infty$ can be possibly weakened but not completely removed. In fact, as shown in the next example, the series $\sum_{n=1}^{\infty} \frac{V_n}{n}$ may fail to converge a.s. even if (V_n) is i.i.d. and $E(V_1) = 0$.

Example 11. Let (V_n) be i.i.d. with

$$P(V_1 = -c/\log 2) = 1/2$$
 and $P(2 < V_1 \le x) = \frac{c}{2} \int_2^x \frac{dt}{(t \log t)^2}$ for $x > 2$,

where the constant c is given by $c = \left(\int_2^\infty \frac{dt}{(t \log t)^2}\right)^{-1}$. Then,

$$E(V_1) = -\left\{\frac{c}{2\log 2} + \frac{c}{2}\right\}, \int_2^\infty \frac{t}{(t\log t)^2} \, dt = 0.$$

Letting

$$W_n = \frac{V_n}{n} \, \mathbb{1}_{\{|V_n| \le n\}}$$

a direct calculation shows that $\sum_{n} E(W_n) = -\infty$ and $\sum_{n} \operatorname{Var}(W_n) < \infty$. Hence,

$$\sum_{n=1}^{\infty} W_n \stackrel{a.s.}{=} -\infty.$$

In turn, by the Borel-Cantelli Lemma, this implies $\sum_{n=1}^{\infty} \frac{V_n}{n} \stackrel{a.s.}{=} -\infty$. Finally, a remark is in order.

Remark 12. Fix an i.i.d. sequence Z_1, Z_2, \ldots of *p*-dimensional random vectors such that $Z_1 \sim \mathcal{L}(0, I)$. Define a sequence $Y = (Y_1, Y_2, \ldots)$ as

$$Y_1 = Z_1, \quad Y_2 = Y_1 + Z_2 \quad \text{and} \quad Y_{n+1} = N_n + R_n^{1/2} Z_{n+1} \quad \text{for } n \ge 2$$

where $N_n = \frac{1}{n} \sum_{i=1}^n Y_i$ and $R_n = \frac{1}{n} \sum_{i=1}^n (Y_i - N_n) (Y_i - N_n)^t.$

It is straightforward to see that

$$P(Y_1 \in \cdot) = \mathcal{L}(0, I), \quad P(Y_2 \in \cdot \mid Y_1) = \mathcal{L}(Y_1, I), \quad \text{and} \\ P(Y_{n+1} \in \cdot \mid Y_1, \dots, Y_n) = \mathcal{L}(N_n, R_n) \quad \text{for } n \ge 2.$$

Hence, $Y \sim X$ whenever the predictive distributions of X are the α_n of Theorems 4 and 5. In particular, since $Y \sim X$, we can use Y instead of X when proving Theorems 4 and 5. Equivalently, in the proofs of Theorems 4 and 5, we can suppose

 $X_1 = Z_1, \quad X_2 = X_1 + Z_2 \text{ and } X_{n+1} = M_n + Q_n^{1/2} Z_{n+1} \text{ for } n \ge 2$ where Z_1, Z_2, \ldots are i.i.d. and $Z_1 \sim \mathcal{L}(0, I)$.

We are now able to attack the results of Section 2.2. To this end, we let

$$\mathcal{F}_n = \sigma(Z_1, \dots, Z_n)$$
 with \mathcal{F}_0 = the trivial σ -field

Proof of Theorem 4. Suppose $M_n \xrightarrow{a.s.} M$, $Q_n \xrightarrow{a.s.} Q$ and $Q \in \mathcal{M}$ a.s. Let ϕ be a density of $\mathcal{L}(0, I)$ with respect to *p*-dimensional Lebesgue measure. Since $Q_n \in \mathcal{M}$ a.s. (as it is easily proved), one obtains

$$\|\alpha_n - \alpha\| = \|\mathcal{L}(M_n, Q_n) - \mathcal{L}(M, Q)\|$$

= $\frac{1}{2} \int_{\mathbb{R}^p} \left| \frac{\phi(Q_n^{-1/2}(x - M_n))}{\det(Q_n)^{1/2}} - \frac{\phi(Q^{-1/2}(x - M))}{\det(Q)^{1/2}} \right| dx.$

Given $\epsilon > 0$, take an integrable continuous function f on \mathbb{R}^p such that

$$\int_{\mathbb{R}^p} |\phi(x) - f(x)| \, dx < \epsilon$$

Define

$$I_n = \int_{\mathbb{R}^p} \left| \frac{f(Q_n^{-1/2}(x - M_n))}{\det(Q_n)^{1/2}} - \frac{f(Q^{-1/2}(x - M))}{\det(Q)^{1/2}} \right| dx$$

Since f is continuous, $I_n \xrightarrow{a.s.} 0$. Therefore,

$$\limsup_{n} \|\alpha_{n} - \alpha\| \le \epsilon + \frac{1}{2} \limsup_{n} I_{n} \stackrel{a.s.}{=} \epsilon.$$

This proves that $\|\alpha_n - \alpha\| \xrightarrow{a.s.} 0$ provided $M_n \xrightarrow{a.s.} M$, $Q_n \xrightarrow{a.s.} Q$ and $Q \in \mathcal{M}$ a.s. After noting this fact, the rest of the proof is split into three steps. Recall that, by Remark 12, we can assume

 $X_1 = Z_1, \quad X_2 = X_1 + Z_2 \quad \text{and} \quad X_{n+1} = M_n + Q_n^{1/2} Z_{n+1} \quad \text{for } n \ge 2$ where Z_1, Z_2, \ldots are i.i.d. and $Z_1 \sim \mathcal{L}(0, I)$.

Step 1: Q_n converges a.s.

Define

$$M_0 = 0$$
, $Q_0 = Q_1 = I$ and $L_n = Q_n^{1/2} Z_{n+1}$.

Then,

$$M_{n+1} - M_n = \frac{Q_n^{1/2} Z_{n+1}}{n+1} = \frac{L_n}{n+1}$$

After some algebra, this implies

(5)
$$Q_{n+1} = \frac{n}{n+1} Q_n + \frac{n}{(n+1)^2} L_n L_n^t$$

or equivalently

$$Q_{n+1}^{(i,j)} = \frac{n}{n+1} Q_n^{(i,j)} + \frac{n}{(n+1)^2} L_n^{(i)} L_n^{(j)} \quad \text{for all } i, j = 1, \dots, p.$$

The conditional distribution of L_n given \mathcal{F}_n is $\mathcal{L}(0, Q_n)$. Therefore,

$$E\left\{L_n^{(i)} L_n^{(j)} \mid \mathcal{F}_n\right\} = Q_n^{(i,j)}$$

and

$$E\{Q_{n+1}^{(i,j)} \mid \mathcal{F}_n\} = \frac{n}{n+1}Q_n^{(i,j)} + \frac{n}{(n+1)^2}Q_n^{(i,j)} = Q_n^{(i,j)}\left(1 - \frac{1}{(n+1)^2}\right).$$

For i = j, it follows that $(Q_n^{(i,i)} : n \ge 2)$ is a non-negative supermartingale. Hence,

$$E\{|Q_n^{(i,j)}|\} \le \frac{E\{Q_n^{(i,i)}\} + E\{Q_n^{(j,j)}\}}{2} \le \frac{E\{Q_2^{(i,i)}\} + E\{Q_2^{(j,j)}\}}{2}$$

and

$$\sum_{n} E \left| E \left\{ Q_{n+1}^{(i,j)} \mid \mathcal{F}_{n} \right\} - Q_{n}^{(i,j)} \right| = \sum_{n} \frac{E \left\{ |Q_{n}^{(i,j)}| \right\}}{(n+1)^{2}}$$
$$\leq \frac{E \left\{ Q_{2}^{(i,i)} \right\} + E \left\{ Q_{2}^{(j,j)} \right\}}{2} \sum_{n} \frac{1}{(n+1)^{2}} < \infty.$$

To sum up, for fixed *i* and *j*, the sequence $(Q_n^{(i,j)} : n \ge 2)$ is a quasi-martingale such that $\sup_n E\{|Q_n^{(i,j)}|\} < \infty$, and this implies $Q_n^{(i,j)} \xrightarrow{a.s.} Q^{(i,j)}$ for some random variable $Q^{(i,j)}$. Hence, $Q_n \xrightarrow{a.s.} Q$ where Q is the matrix $Q = (Q^{(i,j)} : 1 \le i, j \le p)$.

Step 2: M_n is a martingale and it is bounded in L_2 .

Just note that

$$E(M_{n+1} - M_n \mid \mathcal{F}_n) = E\left\{\frac{Q_n^{1/2}Z_{n+1}}{n+1} \mid \mathcal{F}_n\right\}$$
$$= \frac{Q_n^{1/2}E(Z_{n+1} \mid \mathcal{F}_n)}{n+1} = \frac{Q_n^{1/2}E(Z_{n+1})}{n+1} = 0.$$

Hence, M_n is a martingale. In addition,

$$E(M_n^t M_n) = E\left\{ \left(\sum_{i=1}^n \frac{Q_{i-1}^{1/2} Z_i}{i} \right)^t \left(\sum_{j=1}^n \frac{Q_{j-1}^{1/2} Z_j}{j} \right) \right\} = \sum_{i=1}^n \frac{E(Z_i^t Q_{i-1} Z_i)}{i^2}$$
$$= \sum_{i=1}^n \sum_{r,k=1}^p \frac{E(Z_i^{(r)} Z_i^{(k)}) E(Q_{i-1}^{(r,k)})}{i^2} = \sum_{i=1}^n \sum_{k=1}^p \frac{E(Q_{i-1}^{(k,k)})}{i^2}$$
$$\leq \sum_{i=1}^\infty \sum_{k=1}^p \frac{E(Q_{i-1}^{(k,k)})}{i^2} \le p + \frac{p}{4} + \sum_{k=1}^p E(Q_2^{(k,k)}) \sum_{i=3}^\infty \frac{1}{i^2}$$

where the last inequality is because $(Q_n^{(k,k)}: n \ge 2)$ is a supermartingale for fixed $k = 1, \ldots, p$. Therefore, M_n is a martingale such that $\sup_n E(M_n^t M_n) < \infty$, and this implies that $M_n \xrightarrow{a.s.} M$ for some *p*-dimensional random vector M.

Step 3: $Q \in \mathcal{M}$ a.s.

Since $Q \stackrel{a.s.}{=} \lim_{n \to \infty} Q_n$, the matrix Q is symmetric and non-negative definite a.s. Hence, to prove $Q \in \mathcal{M}$ a.s., it suffices to show that $\det(Q) > 0$ a.s.

Exploiting (5) and Lemma 9, one obtains

$$\det(Q_{n+1}) = \det\left(\frac{n}{n+1} Q_n^{1/2} \left(I + \frac{1}{(n+1)} Z_{n+1} Z_{n+1}^t\right) Q_n^{1/2}\right)$$
$$= \det(Q_n) \left(1 - \frac{1}{n+1}\right)^p \left(1 + \frac{1}{n+1} Z_{n+1}^t Z_{n+1}\right).$$

Letting $V_n = Z_n^t Z_n - p$, the above equation can be written as

$$\det(Q_{n+1}) = \det(Q_n) \sum_{j=0}^p {\binom{p}{j}} \left(\frac{-1}{n+1}\right)^j \left(1 + \frac{V_{n+1} + p}{n+1}\right)$$
$$= \det(Q_n) \left\{1 + \frac{V_{n+1}}{n+1} - p \frac{V_{n+1} + p}{(n+1)^2} + R_{n+1}\right\}$$

where

$$R_{n+1} = \sum_{j=2}^{p} \begin{pmatrix} p \\ j \end{pmatrix} \left(\frac{-1}{n+1}\right)^{j} \left(1 + \frac{V_{n+1} + p}{n+1}\right).$$

Therefore,

$$\det(Q_n) = \det(Q_2) \prod_{j=3}^n \frac{\det(Q_j)}{\det(Q_{j-1})} = \det(Q_2) \prod_{j=3}^n \left\{ 1 + \frac{V_j}{j} - p \frac{V_j + p}{j^2} + R_j \right\}.$$

Next, define

$$U_j = 1 + \frac{V_j}{j} - p \frac{V_j + p}{j^2} + R_j.$$

If $\sum_{j=3}^{n} \log U_j \xrightarrow{a.s.} T$ as $n \to \infty$, for some real random variable T, then

$$\det(Q) = \lim_{n} \det(Q_n) = \det(Q_2) \lim_{n} \exp\left(\sum_{j=3}^{n} \log U_j\right) = \det(Q_2) e^T > 0 \quad \text{a.s.}$$

Hence, it suffices to show that the series $\sum_j \log U_j$ converges a.s. In turn, this is true since each series involved in the definition of U_j converges a.s. Precisely, $\sum_j \frac{V_j}{j}$ converges a.s. because of Lemma 10. The series $\sum_j \frac{V_j + p}{j^2}$ converges a.s. since $V_j + p = Z_j^t Z_j \ge 0$ and

$$E\left\{\sum_{j}\frac{V_j+p}{j^2}\right\} = \sum_{j}\frac{E(Z_j^tZ_j)}{j^2} = \sum_{j}\frac{p}{j^2} < \infty.$$

Finally, $\sum_j R_j$ converges a.s. by exactly the same argument used for $\sum_j \frac{V_j + p}{j^2}$. Therefore, $\sum_j \log U_j$ converges a.s., and this concludes the proof.

Proof of Theorem 5. Let $\|\cdot\|_E$ be the Euclidean norm on \mathbb{R}^p and $\|\cdot\|_F$ the Frobenius norm on the matrices of order $p \times p$. The latter is defined as

$$||B||_F = \sqrt{\sum_{i,j} (B^{(i,j)})^2} = \sqrt{\operatorname{trace}(B^t B)} \quad \text{for any } p \times p \text{ matrix } B$$

and has the nice property that $||B_1B_2||_F \le ||B_1||_F ||B_2||_F$.

The total variation distance between two *p*-dimensional normal laws, say $\mathcal{N}(a, \Sigma_1)$ and $\mathcal{N}(b, \Sigma_2)$ with $a, b \in \mathbb{R}^p$ and $\Sigma_1, \Sigma_2 \in \mathcal{M}$, can be estimated as

$$\|\mathcal{N}(a,\Sigma_1) - \mathcal{N}(b,\Sigma_2)\| \le \|\mathcal{N}(a,\Sigma_1) - \mathcal{N}(b,\Sigma_1)\| + \frac{3}{2} \|\Sigma_1^{-1/2}\Sigma_2\Sigma_1^{-1/2} - I\|_F;$$

see e.g. [17, Theorem 2] and [10, Formula (2)]. In addition,

$$\|\Sigma_1^{-1/2}\Sigma_2\Sigma_1^{-1/2} - I\|_F = \|\Sigma_1^{-1/2}(\Sigma_2 - \Sigma_1)\Sigma_1^{-1/2}\|_F \le \|\Sigma_1^{-1/2}\|_F^2 \|\Sigma_2 - \Sigma_1\|_F$$

and

$$\|\mathcal{N}(a, \Sigma_1) - \mathcal{N}(b, \Sigma_1)\| \le \|b - a\|_E \|\Sigma_1^{-1/2}\|_F;$$

Collecting all these facts together

see [10, Page 5]. Collecting all these facts together,

$$\left\|\mathcal{N}(a,\Sigma_{1}) - \mathcal{N}(b,\Sigma_{2})\right\| \le C(\Sigma_{1}) \left\{ \|b - a\|_{E} + \|\Sigma_{2} - \Sigma_{1}\|_{F} \right\}$$

where $C(\Sigma_1)$ is a constant which depends on Σ_1 only.

Applying the previous inequality to our framework, one obtains

$$\|\alpha_n - \alpha\| = \|\mathcal{N}(M_n, Q_n) - \mathcal{N}(M, Q)\| \le C(Q) \left\{ \|M_n - M\|_E + \|Q_n - Q\|_F \right\}.$$

Hence, it suffices to show that

$$b := \sup_{n} \sqrt{n} E \Big\{ \|M_n - M\|_E + \|Q_n - Q\|_F \Big\} < \infty.$$

In fact, if $b < \infty$ and d_n is any sequence of constants such that $\frac{d_n}{\sqrt{n}} \to 0$, then

$$d_n E \left\{ \|M_n - M\|_E + \|Q_n - Q\|_F \right\} \le \frac{b \, d_n}{\sqrt{n}} \longrightarrow 0.$$

In particular, $d_n \left\{ \|M_n - M\|_E + \|Q_n - Q\|_F \right\} \xrightarrow{P} 0$, which in turn implies

$$d_n \|\alpha_n - \alpha\| \le d_n C(Q) \left\{ \|M_n - M\|_E + \|Q_n - Q\|_F \right\} \xrightarrow{P} 0.$$

To prove $b < \infty$, we first note that

$$M - M_n = \sum_{i=n}^{\infty} (M_{i+1} - M_i) = \sum_{i=n}^{\infty} \frac{Q_i^{1/2} Z_{i+1}}{i+1}$$

Therefore, for each $n \geq 2$,

$$E\left\{\|M_n - M\|_E\right\}^2 \le E\left\{\|M_n - M\|_E^2\right\} = E\left\{(M - M_n)^t (M - M_n)\right\}$$
$$= \sum_{i=n}^{\infty} \frac{E(Z_{i+1}^t Q_i Z_{i+1})}{(i+1)^2} = \sum_{i=n}^{\infty} \sum_{r,k=1}^p \frac{E(Z_{i+1}^{(r)} Z_{i+1}^{(k)}) E(Q_i^{(r,k)})}{(i+1)^2}$$
$$= \sum_{i=n}^{\infty} \sum_{k=1}^p \frac{E(Q_i^{(k,k)})}{(i+1)^2} \le \sum_{k=1}^p E(Q_2^{(k,k)}) \sum_{i=n}^{\infty} \frac{1}{(i+1)^2}$$

where the last inequality is because $\{Q_i^{(k,k)}: i \ge 2\}$ is a supermartingale for fixed k. On noting that $\sum_{i=n}^{\infty} \frac{1}{(i+1)^2} \le \frac{1}{n}$, one obtains

$$\sqrt{n} E \Big\{ \|M_n - M\|_E \Big\} \le \sqrt{\sum_{k=1}^p E(Q_2^{(k,k)})}.$$

Similarly, recalling that $Q_{n+1} = \frac{n}{n+1} Q_n^{1/2} \left(I + \frac{Z_{n+1} Z_{n+1}^t}{n+1} \right) Q_n^{1/2}$, one obtains

$$Q - Q_n = \sum_{i=n}^{\infty} (Q_{i+1} - Q_i)$$
$$= \sum_{i=n}^{\infty} \left\{ \frac{Q_i^{1/2} (Z_{i+1} Z_{i+1}^t - I) Q_i^{1/2}}{i+1} - \frac{Q_i^{1/2} Z_{i+1} Z_{i+1}^t Q_i^{1/2}}{(i+1)^2} \right\}$$

Define

$$q = \sup_{i} E\{\|Q_{i}^{1/2}\|_{F}^{4}\}$$
 and $K_{i} = \frac{Q_{i}^{1/2}(Z_{i+1}Z_{i+1}^{t} - I)Q_{i}^{1/2}}{i+1}.$

Since $E\{\|Q_i^{1/2}\|_F^2\} \leq \sqrt{q}$ for all i, $\sum_{i=n}^{\infty} \frac{E\{\|Q_i^{1/2}Z_{i+1}Z_{i+1}^tQ_i^{1/2}\|_F\}}{(i+1)^2} \leq \sum_{i=n}^{\infty} \frac{E\{\|Z_{i+1}Z_{i+1}^t\|_F\} E\{\|Q_i^{1/2}\|_F^2\}}{(i+1)^2}$ $\leq E\{\|Z_1Z_1^t\|_F\} \sum_{i=n}^{\infty} \frac{\sqrt{q}}{(i+1)^2} \leq \frac{\sqrt{q}}{n} E\{\|Z_1Z_1^t\|_F\}.$

Since K_i is symmetric and $E(K_iK_j) = 0$ for $i \neq j$,

$$E\left\{ \|\sum_{i=n}^{\infty} K_i\|_F^2 \right\} = E\left\{ \operatorname{trace}\left(\left(\sum_{i=n}^{\infty} K_i\right) \left(\sum_{j=n}^{\infty} K_j\right) \right) \right\} \\ = \operatorname{trace}\left(\sum_{i,j=n}^{\infty} E(K_iK_j)\right) = \operatorname{trace}\left(\sum_{i=n}^{\infty} E(K_iK_i)\right) \\ = \sum_{i=n}^{\infty} E\left\{ \frac{\|Q_i^{1/2}(Z_{i+1}Z_{i+1}^t - I)Q_i^{1/2}\|_F^2}{(i+1)^2} \right\} \\ \le E\left\{ \|Z_1Z_1^t - I\|_F^2 \right\} \sum_{i=n}^{\infty} \frac{E\left\{\|Q_i^{1/2}\|_F^4\right\}}{(i+1)^2} \le E\left\{\|Z_1Z_1^t - I\|_F^2\right\} \frac{q}{n} \right\}$$

Hence,

$$\sqrt{n} E\{\|Q_n - Q\|_F\} \le \sqrt{n} \sum_{i=n}^{\infty} \frac{E\{\|Q_i^{1/2} Z_{i+1} Z_{i+1}^t Q_i^{1/2}\|_F\}}{(i+1)^2} + \sqrt{n} E\{\|\sum_{i=n}^{\infty} K_i\|_F\} \\
\le \frac{\sqrt{q}}{\sqrt{n}} E\{\|Z_1 Z_1^t\|_F\} + \sqrt{q} E\{\|Z_1 Z_1^t - I\|_F^2\}.$$

Hence, $b < \infty$ provided $q < \infty$. We finally prove $q < \infty$.

Fix $i \geq 2, 1 \leq k \leq p$, and define $W_i = (Q_i^{(k,k)})^2$. As noted in the proof of Theorem 4, the conditional distribution of $L_j = Q_j^{1/2} Z_{j+1}$ given \mathcal{F}_j is $\mathcal{N}(0, Q_j)$. Hence,

$$E\{(L_j^{(k)})^2 \mid \mathcal{F}_j\} = Q_j^{(k,k)} \text{ and } E\{(L_j^{(k)})^4 \mid \mathcal{F}_j\} = 3(Q_j^{(k,k)})^2 = 3W_j.$$

Using (5), it follows that

$$E(W_{j+1} \mid \mathcal{F}_j) = W_j \left\{ \frac{j^2}{(j+1)^2} + \frac{3j^2}{(j+1)^4} + \frac{2j^2}{(j+1)^3} \right\} \le W_j.$$

Therefore,

$$E(W_i) = E\left\{W_2\prod_{j=2}^{i-1}\frac{W_{j+1}}{W_j}\right\} = E(W_2)\prod_{j=2}^{i-1}E\left\{\frac{W_{j+1}}{W_j}\right\}$$
$$= E(W_2)\prod_{j=2}^{i-1}E\left\{\frac{E(W_{j+1} \mid \mathcal{F}_j)}{W_j}\right\} \le E(W_2).$$

To sum up, $E\{(Q_i^{(k,k)})^2\} \le E\{(Q_2^{(k,k)})^2\}$ for all $i \ge 2$ and $1 \le k \le p$. Hence,

$$E\{\|Q_i^{1/2}\|_F^4\} = E\{\operatorname{trace}(Q_i)^2\} = \sum_{r,k=1}^p E\{Q_i^{(r,r)}Q_i^{(k,k)}\}$$
$$\leq \sum_{r,k=1}^p \sqrt{E\{(Q_i^{(r,r)})^2\}} E\{(Q_i^{(k,k)})^2\}$$
$$\leq p^2 \max_{1 \leq k \leq p} E\{(Q_i^{(k,k)})^2\} \leq p^2 \max_{1 \leq k \leq p} E\{(Q_2^{(k,k)})^2\}.$$

This proves that $q < \infty$ and concludes the proof of the theorem.

Let us turn to the results of Section 2.3. In such results, the predictives of X are the copula-based predictive distributions β_n . Hence, as noted in Section 2.3, X is conditionally identically distributed.

Proof of Theorem 7. As in Example 6, define

$$D_n = \int_{-\infty}^{\infty} \left| f_{n+1} \left(x \mid X(n+1) \right) - f_n \left(x \mid X(n) \right) \right| dx.$$

For n < m,

$$\int_{-\infty}^{\infty} \left| f_n(x \mid X(n)) - f_m(x \mid X(m)) \right| dx$$

= $\int_{-\infty}^{\infty} \left| \sum_{i=n}^{m-1} \left\{ f_{i+1}(x \mid X(i+1)) - f_i(x \mid X(i)) \right\} \right| dx \le \sum_{i=n}^{m-1} D_i.$

If $\sum_n D_n < \infty$ a.s., the sequence $f_n(\cdot | X(n))$ is Cauchy in the L^1 -norm a.s., so that it converges in the L^1 -norm a.s. Hence, it suffices to show that $\sum_n D_n < \infty$ a.s. To this end, recall that D_n is distributed as $r_n \int_0^1 |c_n(u, V) - 1| du$ where V is any random variable with uniform distribution on (0, 1); see Example 6. It follows that

$$E(D_n) = r_n E\left\{\int_0^1 |c_n(u, V) - 1| \, du\right\} = r_n \int_0^1 E\{|c_n(u, V) - 1|\} \, du$$
$$= r_n \int_0^1 \int_0^1 |c_n(u, v) - 1| \, dv \, du \le 2 \, r_n.$$

Therefore, $\sum_{n} E(D_n) \leq 2 \sum_{n} r_n < \infty$ which in turn implies $\sum_{n} D_n < \infty$ a.s. \Box

Proof of Theorem 8. Since X is conditionally identically distributed, by [3, Theorem 4], it suffices to show that

(6)
$$\sup_{n} E\left\{\int_{-\infty}^{\infty} f_n(x \mid X(n))^2 dx\right\} < \infty.$$

To prove (6), we note that, since X is conditionally identically distributed,

$$E\left\{f_{k+1}(x \mid X(k+1)) \mid X(k)\right\} = f_k(x \mid X(k))$$

for almost every $x \in \mathbb{R}$ (with respect to Lebesgue measure). This implies

$$E\left\{c_k\left(F_k\left(x \mid X(k)\right), F_k\left(X_{k+1} \mid X(k)\right)\right) \mid X(k)\right\} = 1$$

for almost every $x \in \mathbb{R}$. Since $c_k(u, v) \leq b$ for all $u, v \in [0, 1]$, it follows that

$$E\left\{\left(\frac{f_{k+1}(x \mid X(k+1))}{f_k(x \mid X(k))}\right)^2 \mid X(k)\right\} \le (1-r_k)^2 + r_k^2 b^2 + 2(1-r_k)r_k = 1 + r_k^2(b^2 - 1).$$

Finally, the previous inequality implies

$$E\left\{f_n\left(x \mid X(n)\right)^2\right\} = E\left\{\left(f_0(x)\prod_{k=0}^{n-1}\frac{f_{k+1}\left(x \mid X(k+1)\right)}{f_k\left(x \mid X(k)\right)}\right)^2\right\}$$
$$\leq f_0(x)^2\prod_{k=0}^{n-1}\left\{1 + r_k^2(b^2 - 1)\right\} \leq f_0(x)^2\exp\left\{(b^2 - 1)\sum_{k=0}^{n-1}r_k^2\right\}.$$

Thus, to check condition (6), it suffices noting that

$$E\left\{\int_{-\infty}^{\infty} f_n(x \mid X(n))^2 dx\right\} = \int_{-\infty}^{\infty} E\left\{f_n(x \mid X(n))^2\right\} dx$$
$$\leq \exp\{(b^2 - 1)\sum_{k=0}^{\infty} r_k^2\} \int_{-\infty}^{\infty} f_0(x)^2 dx.$$

References

- Azzalini A., Capitanio A. (2003) Distributions generated by perturbation of symmetry with emphasis on a multivariate skew t distribution, J. Royal Stat. Soc. B, 65, 367–389.
- [2] Berti P., Pratelli L., Rigo P. (2004) Limit theorems for a class of identically distributed random variables, Ann. Probab., 32, 2029-2052.
- [3] Berti P., Pratelli L., Rigo P. (2013) Exchangeable sequences driven by an absolutely continuous random measure, Ann. Probab., 41, 2090-2102.
- [4] Berti P., Dreassi E., Pratelli L., Rigo P. (2021) A class of models for Bayesian predictive inference, *Bernoulli*, 27, 702-726.
- [5] Berti P., Dreassi E., Leisen F., Pratelli L., Rigo P. (2023) Bayesian predictive inference without a prior, *Statistica Sinica*, 33, 2405-2429.
- [6] Berti P., Dreassi E., Leisen F., Pratelli L., Rigo P. (2024) A probabilistic view on predictive constructions for Bayesian learning, *Statist. Science*, to appear.
- [7] Clarke B., Fokoue E., Zhang H.H. (2009) Principles and theory for data mining and machine learning, Springer, New York.
- [8] Dawid A.P. (1984) Present position and potential developments: Some personal views: Statistical Theory: The prequential approach, J. Royal Stat. Soc. A, 147, 278-292.
- [9] Dawid A.P., Vovk V.G. (1999) Prequential probability: principles and properties, *Bernoulli*, 5, 125-162.
- [10] Devroye L., Mehrabian A., Reddad T. (2023) The total variation distance between highdimensional Gaussians with the same mean, arXiv:1810.08693v7
- [11] Ding J., Zhou A. (2007) Eigenvalues of rank-one updated matrices with some applications, *Appl. Math. Letters*, 20, 1223-1226.
- [12] Dutordoir V., Saul A., Ghahramani Z., Simpson F. (2023) Neural diffusion processes, Proc. 40th Intern. Conf. on Machine Learning, PMLR 202, 8990-9012.
- [13] Efron B. (2020) Prediction, estimation, and attribution, J.A.S.A., 115, 636-655.
- [14] Fong E., Holmes C., Walker S.G. (2023) Martingale posterior distributions (with discussion), J. Royal Stat. Soc. B, 85, 1357-1391.

26 SAMUELE GARELLI, FABRIZIO LEISEN, LUCA PRATELLI, AND PIETRO RIGO

- [15] Hahn P.R., Martin R., Walker S.G. (2018) On recursive Bayesian predictive distributions, J.A.S.A., 113, 1085-1093.
- [16] Hastie T., Tibshirani R., Friedman J. (2009) The elements of statistical learning: Data Mining, Inference, and Prediction, Springer, New York.
- [17] Kelbert M. (2023) Survey of distances between the most popular distributions, Analytics, 2, 225-245.
- [18] Li F., Ding P., Mealli F. (2023) Bayesian causal inference: A critical review, Phil. Trans. R. Soc. A, 381.
- [19] Pitman J. (1996) Some developments of the Blackwell-MacQueen urn scheme, Statistics, Probability and Game Theory, IMS Lect. Notes Mon. Series, 30, 245-267.
- [20] Pitman J., Yor M. (1997) The two-parameter Poisson-Dirichlet distribution derived from a stable subordinator, Ann. Probab., 25, 855-900.

SAMUELE GARELLI, DIPARTIMENTO DI SCIENZE STATISTICHE "P. FORTUNATI", UNIVERSITÀ DI BOLOGNA, VIA DELLE BELLE ARTI 41, 40126 BOLOGNA, ITALY *Email address*: samuele.garelli2@unibo.it

Fabrizio Leisen, Department of Mathematics, King's College, Strand WC2R 2LS, London, UK

 $Email \ address: \verb"fabrizio.leisen@gmail.com"$

LUCA PRATELLI, ACCADEMIA NAVALE, VIALE ITALIA 72, 57100 LIVORNO, ITALY *Email address:* luca_pratelli@marina.difesa.it

PIETRO RIGO (CORRESPONDING AUTHOR), DIPARTIMENTO DI SCIENZE STATISTICHE "P. FORTU-NATI", UNIVERSITÀ DI BOLOGNA, VIA DELLE BELLE ARTI 41, 40126 BOLOGNA, ITALY *Email address*: pietro.rigo@unibo.it