On the solutions of the local Zamolodchikov tetrahedron equation

M. Chirkov* and S. Konstantinou-Rizos[†]

Centre of Integrable Systems, P.G. Demidov Yaroslavl State University, Yaroslavl, Russia

Abstract

We study the solutions of the local Zamolodhcikov tetrahedron equation in the form of correspondences derived by 3×3 matrices. We present all the associated generators of 4-simplex maps satisfying the local tetrahedron equation. Moreover, we demonstrate that, from some of our solutions, we can recover the 4-simplex extensions of Kashaev–Korepanov–Sergeev and Hirota type tetrahedron maps. Finally, we construct several novel 4-simplex maps.

PACS numbers: 02.30.Ik, 02.90.+p, 03.65.Fd.

Mathematics Subject Classification 2020: 35Q55, 16T25.

Keywords: 4-simplex maps, tetrahedron maps, local Zamolodchikov tetrahedron equation, Yang–Baxter maps.

1 Introduction

The n-simplex equations are a family of important equations of Mathematical Physics. The first member of the family is trivial, while the second, third and fourth members are quite famous and have attracted the interest of many leading scientists from various fields of Mathematics and Physics.

In particular, the 2-simplex equation is the Yang–Baxter equation [8] which is one of the most fundamental equations of Mathematical Physics, the 3-simplex equation is the well-celebrated Zamolodchikov's tetrahedron equation [27, 28], and the 4-simplex equation is the Bazhanov–Stroganov equation [4].

Due to the importance of n-simplex maps, there is a need to develop methods for constructing and classifying them. Famous results towards the classification of n-simplex maps, include the classification of Yang-Baxter maps by Adler, Bobenko and Suris [2], as well as the classification of Zamolodchikov tetrahedron maps by Kashaev, Korepanov and Sergeev [11, 26]. There are several methods in the literature for constructing n-simplex maps; we indicatively refer to the derivation of n-simplex maps via the symmetries of lattice equations [25, 14], their construction via matrix refactorisation problems and their construction via Darboux transformations for integrable PDEs [20, 21, 23].

The most significant property that share all the n-simplex equations is that the local (n-1)-simplex equation is a generator of solutions to the n-simplex equation [5]. It is known that any unique solution of

^{*}mikhlchirkov@gmail.com

[†]skonstantin84@gmail.com

the local (n-1)-simplex equation satisfies the n-simplex equation. However, there is no proof of such a statement; thus, in order to prove that a map satisfying uniquely the local (n-1)-simplex is an n-simplex map, we either need to substitute it to the n-simplex equation, or use supplementary matrix conditions [16, 22].

Recently, it has become understood that non-unique solutions (in the form of correspondences) of the local (n-1)-simplex equation also give rise to n-simplex maps [6, 10, 15, 17, 19]. Moreover, this correspondence approach gives rise to novel n-simplex maps that do not belong to any of the known classification lists. Most known classification results focus on n-simplex maps themselves. It makes sense to classify the solutions of the local (n-1)-simplex equation which generate n-simplex maps which may lead to wider classification results.

In this paper, we study the solutions of the local Zamolodchikov tetrahedron equation. Specifically, we:

- Present all the possible solutions to the local tetrahedron equation in the form of correspondences generated by 3 × 3 matrices depending on free variables;
- We demonstrate that these solutions generate all known in the literature 4-simplex maps. As illustrative examples we use Kashaev–Sergeev–Korepanov and Hirota type 4-simplex maps [17, 18];
- We use the obtained local tetrahedron correspondences to construct new, noninvolutive, 4-simplex maps.

The rest of the text is organised as follows: In the next section, we provide all the necessary preliminary information which makes the text self-contained. In particular, we fix the notation that is used throughout the text, we explain what 4-simplex maps are, and what is their relation to the local tetrahedron equation. Section 3 deals with the study of all possible solutions of the local Zamolodchikov tetrahedron equation generated by 3×3 matrices. We show that the essentially different solutions to the local tetrahedron equation in the form of correspondences are generated by four simple 3×3 matrices. These correspondences are generators of 4-simplex maps. Finally, in section 4, we construct novel, noninvolutive, 4-simplex maps.

2 Preliminaries

2.1 Notation

Throughout the text:

- By \mathcal{X} we denote an arbitrary set, whereas by Latin italic letters (i.e. x, y, u, v etc.) the elements of \mathcal{X}
- By \mathcal{X}^n we denote the Cartesian product: $\mathcal{X}^n = \underbrace{\mathcal{X} \times \ldots \times \mathcal{X}}_n$.
- Matrices will be denoted by capital Latin straight letters (i.e. A, B, C) etc. Additionally, by $A^n_{i_1 i_2 \dots i_m}$ we denote the $n \times n$ extensions of the $m \times m$ (m < n) matrix A, where the elements of A are placed at the intersection of the $i_1, i_2 \dots i_m$ rows with the $i_1, i_2 \dots i_m$ columns of matrix $A^n_{i_1 i_2 \dots i_m}$. For example,

$$A_{12}^{3} = \begin{pmatrix} a_{11} & a_{12} & 0 \\ a_{21} & a_{22} & 0 \\ 0 & 0 & 1 \end{pmatrix}, A_{13}^{3} = \begin{pmatrix} a_{11} & 0 & a_{12} \\ 0 & 1 & 0 \\ a_{21} & 0 & a_{22} \end{pmatrix} \text{ and } A_{23}^{3} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & a_{11} & a_{12} \\ 0 & a_{21} & a_{22} \end{pmatrix} \text{ are } 3 \times 3 \text{ extensions of matrix } A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}.$$

2.2 Local Zamolodchikov tetrahedron equation and 4-simplex maps

As it was mentioned earlier, the local 3-simplex equation is a generator of solutions of the 4-simplex equation. In this section, we give the definitions of the local Zamolodchikov tetrahedron equation and a 4-simplex map.

Let \mathcal{X} be a set. A map $S: \mathcal{X}^4 \to \mathcal{X}^4$, namely

$$S: (x, y, z, t) \mapsto (u(x, y, z, t), v(x, y, z, t), w(x, y, z, t), r(x, y, z, t)), \tag{1}$$

is called a 4-simplex map if it satisfies the set-theoretical 4-simplex (or Bazhanov-Stroganov) equation

$$S^{1234} \circ S^{1567} \circ S^{2589} \circ S^{368,10} \circ S^{479,10} = S^{479,10} \circ S^{368,10} \circ S^{2589} \circ S^{1567} \circ S^{1234}. \tag{2}$$

Functions $S^{ijkl} \in \text{End}(\mathcal{X}^{10})$, $i, j, k, l = 1, \dots 10$, i < j < k < l, in (2) are maps that act as map S on the ijkl terms of the Cartesian product \mathcal{X}^{10} and trivially on the others. For instance,

$$S^{1567}(x_1, x_2, \dots x_{10}) = (u(x_1, x_5, x_6, x_7), x_2, x_3, x_4, v(x_1, x_5, x_6, x_7), w(x_1, x_5, x_6, x_7), r(x_1, x_5, x_6, x_7), x_8, x_9, x_{10}).$$

Now, let L = L(x) be a 3×3 square matrix depending on a variable $x \in \mathcal{X}$ and a parameter $\kappa \in \mathbb{C}$ of the form

$$L(x) = \begin{pmatrix} a(x) & b(x) & c(x) \\ d(x) & e(x) & f(x) \\ k(x) & l(x) & m(x) \end{pmatrix}, \tag{3}$$

where its entries a, b, c, d, e, f, k, l and m are scalar functions of x and k. Let $L_{ijk}^6(x; \kappa)$, $i, j, k = 1, \ldots 6$, i < j < k, be the 6×6 extensions of matrix (3), defined by

$$L_{123}^{6}(x) = \begin{pmatrix} a(x) & b(x) & c(x) & 0 & 0 & 0 \\ d(x) & e(x) & f(x) & 0 & 0 & 0 \\ k(x) & l(x) & m(x) & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad L_{145}^{6}(x) = \begin{pmatrix} a(x) & 0 & 0 & b(x) & c(x) & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ d(x) & 0 & 0 & e(x) & f(x) & 0 \\ k(x) & 0 & 0 & e(x) & f(x) & 0 \\ k(x) & 0 & 0 & l(x) & m(x) & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad L_{356}^{6}(x) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \quad L_{356}^{6}(x) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & a(x) & 0 & b(x) & c(x) \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & a(x) & 0 & b(x) & c(x) \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & d(x) & 0 & e(x) & f(x) \\ 0 & 0 & k(x) & 0 & l(x) & m(x) \end{pmatrix}. \tag{4b}$$

$$\mathbf{L}_{246}^{6}(x) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & a(x) & 0 & b(x) & 0 & c(x) \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & d(x) & 0 & e(x) & 0 & f(x) \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & k(x) & 0 & l(x) & 0 & m(x) \end{pmatrix}, \quad \mathbf{L}_{356}^{6}(x) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & a(x) & 0 & b(x) & c(x) \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & d(x) & 0 & e(x) & f(x) \\ 0 & 0 & k(x) & 0 & l(x) & m(x) \end{pmatrix}. \tag{4b}$$

We call the following matrix four-factorisation problem

$$L_{123}^{6}(u)L_{145}^{6}(v)L_{246}^{6}(w)L_{356}^{6}(r) = L_{356}^{6}(t)L_{246}^{6}(z)L_{145}^{6}(y)L_{123}^{6}(x)$$
(5)

local 3-simplex equation or local Zamolodchikov tetrahedron equation. The local tetrahedron equation is a generator of potential solutions to the 4-simplex equation. If map (1) satisfies equation (5), then the matrix refactorisation problem (5) is called a Lax representation for map (1).

Examples of 4-simplex maps can be found in [17, 9, 3].

3 Solutions of the local tetrahedron equation

In this section, we aim to study the solutions to the local tetrahedron equation in the form of correspondences generated by 3×3 matrices. Specifically, let $x_i \in \mathbb{C}$, $i = 1, \ldots, 9$, be free variables and

$$L(x_1, \dots, x_9) = \begin{pmatrix} x_1 & x_2 & x_3 \\ x_4 & x_5 & x_6 \\ x_7 & x_8 & x_9 \end{pmatrix}.$$
 (6)

Then, substitute (6) to the local tetrahedron equation:

$$L_{123}^{6}(u_{1},\ldots,u_{9})L_{145}^{6}(v_{1},\ldots,v_{9})L_{246}^{6}(w_{1},\ldots,w_{9})L_{356}^{6}(r_{1},\ldots,r_{9}) = L_{356}^{6}(t_{1},\ldots,t_{9})L_{246}^{6}(z_{1},\ldots,z_{9})L_{145}^{6}(y_{1},\ldots,y_{9})L_{123}^{6}(x_{1},\ldots,x_{9}).$$

$$(7)$$

We seek matrices (6) for which the matrix refactorisation problem (7) has solution for u_i , v_i , w_i and r_i , i = 1, ..., 9. In particular we have the following.

Theorem 3.1. Let $x_i \in \mathbb{C}$, i = 1, ..., 9, be free variables. Then, the essentially different solutions of the local Zamolodchikov tetrahedron equation (5) are given in the form of correspondences generated by matrices of the form:

$$A = \begin{pmatrix} x_1 & x_2 & 0 \\ x_3 & x_4 & 0 \\ 0 & 0 & x_5 \end{pmatrix}, \quad B = \begin{pmatrix} x_1 & x_2 & 0 \\ x_3 & x_4 & x_5 \\ 0 & 0 & 0 \end{pmatrix}, \quad C = \begin{pmatrix} 0 & x_1 & 0 \\ x_2 & 0 & x_3 \\ 0 & x_4 & 0 \end{pmatrix}, \quad D = \begin{pmatrix} 0 & 0 & 0 \\ x_1 & 0 & 0 \\ 0 & x_2 & x_3 \end{pmatrix}, \quad (8)$$

where $A = A(x_1, x_2, x_3, x_4, x_5)$, $B = B(x_1, x_2, x_3, x_4, x_5)$, $C = C(x_1, x_2, x_3, x_4)$ and $D = D(x_1, x_2, x_3)$. All the rest of solutions are generated by matrices (8), after:

- 1. Reflection of x_i 's with respect to the antidiagonal;
- 2. Rearranging x_i 's among themselves;
- 3. Setting any variable $x_i = 0$.

Proof. We substitute matrix A to the local tetrahedron equation:

$$A_{123}^{6}(u_{1}, \dots u_{5})A_{145}^{6}(v_{1}, \dots v_{5})A_{246}^{6}(w_{1}, \dots w_{5})A_{356}^{6}(r_{1}, \dots r_{5}) = A_{356}^{6}(t_{1}, \dots t_{5})A_{246}^{6}(z_{1}, \dots z_{5})A_{145}^{6}(y_{1}, \dots y_{5})A_{123}^{6}(x_{1}, \dots x_{5}).$$

The above matrix tetra-factorisation problem is equivalent to the system of polynomial equations

$$u_1v_1 = x_1y_1, \quad u_2w_1 + u_1v_2w_3 = x_2y_1 \quad u_2w_2 + u_1v_2w_4 = y_2, \quad u_3v_1 = x_3z_1 + x_1y_3z_2,$$

$$u_4w_1 + u_3v_2w_3 = x_4z_1 + x_2y_3z_2, \quad u_4w_2 + u_3v_2w_4 = y_4z_2, \quad r_1u_5 = t_1, x_5, \quad r_2u_5 = t_2y_5,$$

$$v_3 = x_3z_3 + x_1y_3z_4, \quad v_4w_3 = x_4z_3 + x_2y_3z_4, \quad v_4w_4 = y_4z_4,$$

$$r_3v_5 = t_3x_5, \quad r_4v_5 = t_4y_5, \quad r_5w_5 = t_5z_5.$$

$$(9)$$

The above system has infinitely many solutions for u_i , v_i , w_i and r_i , i = 1, ..., 5. For instance, it can be solved for u_1 , u_3 , u_4 , v_1 , v_4 , w_1 , w_2 , w_4 , r_1 , r_2 , r_3 , r_4 , r_5 in terms of u_5 , v_2 , v_3 , v_5 , w_3 and w_5 . Thus, system (9) defines a correspondence between \mathbb{C}^{20} and \mathbb{C}^{20} . However, we do not write this correspondence explicitly because of its length.

Now, we substitute matrix B to the local tetrahedron equation:

$$B_{123}^{6}(u_{1}, \dots u_{5})B_{145}^{6}(v_{1}, \dots v_{5})B_{246}^{6}(w_{1}, \dots w_{5})B_{356}^{6}(r_{1}, \dots r_{5}) = B_{356}^{6}(t_{1}, \dots t_{5})B_{246}^{6}(z_{1}, \dots z_{5})B_{145}^{6}(y_{1}, \dots y_{5})B_{123}^{6}(x_{1}, \dots x_{5}).$$

It follows that the above matrix tetra-factorisation problem is equivalent to the system of polynomial equations

$$u_1v_1 = x_1y_1, \quad u_2w_1 + u_1v_2w_3 = x_2y_1, \quad u_2w_2 + u_1v_2w_4 = y_2, \quad u_3v_1 = x_3z_1 + x_1y_3z_2,$$

$$u_4w_1 + u_3v_2w_3 = x_4z_1 + x_2y_3z_2, \quad u_4w_2 + u_3v_2w_4 = y_4z_2, \quad r_1u_5 = x_5z_1, \quad r_2u_5 = y_5z_2$$

$$v_3 = x_3z_3 + x_1y_3z_4, \quad v_4w_3 = x_4z_3 + x_2y_3z_4, \quad v_4w_4 = y_4z_4,$$

$$r_3v_5 = x_5z_3, \quad r_4v_5 = y_5z_4, \quad r_5v_5 = z_5.$$

$$(10)$$

The above system has infinitely many solutions for u_i , v_i , w_i and r_i , i = 1, ..., 5. For example, it can be solved for u_1 , u_3 , u_4 , v_1 , v_4 , w_1 , w_2 , w_4 , r_1 , r_2 , r_3 , r_4 , r_5 in terms of the rest variables u_5 , v_2 , v_3 , v_5 , w_3 and w_5 . Thus, system (10) defines a correspondence between \mathbb{C}^{20} and \mathbb{C}^{20} . We omit this correspondence because of its length.

Next, we substitute matrix C to the local tetrahedron equation:

$$C_{123}^{6}(u_{1}, \dots u_{4})C_{145}^{6}(v_{1}, \dots v_{4})C_{246}^{6}(w_{1}, \dots w_{4})C_{356}^{6}(r_{1}, \dots r_{4}) = C_{356}^{6}(t_{1}, \dots t_{4})C_{246}^{6}(z_{1}, \dots z_{4})C_{145}^{6}(y_{1}, \dots y_{4})C_{123}^{6}(x_{1}, \dots x_{4}).$$

It turns out that the above matrix tetra-factorisation problem is equivalent to the system of polynomial equations

$$u_1w_1 = y_1$$
, $u_2v_1w_2 = x_1y_2z_1$, $r_1u_3 + r_4u_2v_1w_3 = y_3z_1$, $u_4w_1 = t_1y_4$, $v_2 = x_2z_2$,
 $r_2v_3 = x_3z_2$, $r_3v_3 = z_3$, $v_4w_2 = t_2x_4 + t_3x_1y_2z_4$, $r_4v_4w_3 = t_3y_3z_4$, $w_4 = t_4y_4$. (11)

The above system has infinitely many solutions for u_i , v_i , w_i and r_i , i = 1, ..., 4. For instance, it can be solved for u_1 , u_4 , v_2 , v_4 , w_2 , w_4 , r_1 , r_2 , r_3 , r_4 , in terms of the rest variables u_2 , u_3 , v_1 , v_3 , w_1 and w_3 , namely

$$u_1 = \frac{y_1}{w_1}, \quad u_4 = \frac{t_1 y_4}{w_1}, \quad v_2 = x_2 z_2, \quad v_4 = \frac{u_2 v_1 (t_2 x_4 + t_3 x_1 y_2 z_4)}{x_1 y_2 z_1}, \quad w_2 = \frac{x_1 y_2 z_1}{u_2 v_1}, \quad w_4 = t_4 y_4,$$

$$r_1 = \frac{t_2 x_4 y_3 z_1}{u_3 (t_2 x_4 + t_3 x_1 y_2 z_4)}, \quad r_2 = \frac{x_3 z_2}{v_3}, \quad r_3 = \frac{z_3}{v_3}, \quad r_4 = \frac{t_3 x_1 y_2 y_3 z_1 z_4}{u_2 v_1 w_3 (t_2 x_4 + t_3 x_1 y_2 z_4)}.$$

Hence, system (11) defines the above correspondence between \mathbb{C}^{16} and \mathbb{C}^{16} .

Finally, we substitute matrix D to the local tetrahedron equation:

$$D_{123}^{6}(u_1, u_2, u_3)D_{145}^{6}(v_1, v_2, v_3)D_{246}^{6}(w_1, w_2, w_3)D_{356}^{6}(r_1, r_2, r_3) = D_{356}^{6}(t_1, t_2, t_3)D_{246}^{6}(z_1, z_2, z_3)D_{145}^{6}(y_1, y_2, y_3)D_{123}^{6}(x_1, x_2, x_3).$$

The above matrix tetra-factorisation problem is equivalent to the system of polynomial equations

$$v_1 = x_1 z_1, \quad v_2 w_1 = t_1 x_2, \quad r_1 v_3 = t_1 x_3, \quad w_2 = t_2 y_2, \quad r_2 w_3 = t_2 y_3, \quad r_3 w_3 = t_3 z_3.$$
 (12)

The above system has infinitely many solutions for u_i , v_i , w_i and r_i , i = 1, 2, 3. For instance, it can be solved for u_1 , u_4 , v_2 , v_4 , w_2 , w_4 , v_1 , v_2 , v_4 , v_1 , v_2 , v_3 , v_4 , v_4 , v_4 , v_5 , v_4 , v_5 , v_4 , v_5 , v_7 , v_8 , v_8 , v_8 , v_8 , v_9 , v

 $v_1 = x_1 z_1, \quad w_1 = \frac{t_1 x_2}{v_2}, \quad w_2 = t_2 y_2, \quad r_1 = \frac{t_1 x_3}{v_3}, \quad r_2 = \frac{t_2 y_3}{w_3}, \quad r_3 = \frac{t_3 z_3}{w_3}.$

That is, system (12) defines the above correspondence between \mathbb{C}^{12} and \mathbb{C}^{12} .

It can be readily verified that, setting any $x_i = 0$ in one of the matrices in (8), the resulting polynomial equations can be solved for the remaining nonzero u_i , v_i , w_i and r_i .

Furthermore, it can be verified that the reflection of the elements with respect to the antidiagonal in any of the matrices in (8) will leave systems (9), (10), (11) and (12) covariant. Specifically, changing A, B, C and D to

$$\tilde{\mathbf{A}} = \mathbf{P}\mathbf{A}^T\mathbf{P}, \quad \tilde{\mathbf{B}} = \mathbf{P}\mathbf{B}^T\mathbf{P}, \quad \tilde{\mathbf{C}} = \mathbf{P}\mathbf{C}^T\mathbf{P}, \quad \tilde{\mathbf{D}} = \mathbf{P}\mathbf{D}^T\mathbf{P}, \quad \text{where} \quad \mathbf{P} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix},$$

and substituting the latter to the local tetrahedron equation, we shall obtain exactly the same correspondences (9), (10), (11) and (12), respectively, after changing in them $(u, v, x, y) \longleftrightarrow (r, u, t, z)$. In Figure 1 we demonstrate this fact in a particular example.

Finally, after replacing any of the zero elements in (8) with a free variable, one can easily verify that the local tetrahedron equation has no solutions for u_i , v_i , w_i and r_i .




Figure 1: Reflection symmetry example.

Corollary 3.2. The maximum dimension of a map generated by substitution of matrix (6) to the local Zamolodchikov tetrahedron equation is 20.

4 Set-theoretical solutions to the 4-simplex equation

In this section, we use the solutions of the local Zamolodchikov tetrahedron equation in order to construct solutions to the Bazhanov–Stroganov 4-simplex equation.

The first, probably, 4-simplex maps in the literature appear in the work of Hietarinta [9], and also in [3]. The first 4-simplex maps generated by the local Zamolodchikov tetrahedron equation appeared in [17, 18]. In particular, in [17, 18] 4-simplex maps were constructed as extensions of well-known tetrahedron maps, using matrices of the form A in (8). Thus, all 4-simplex maps which appear in [17, 18] can be generated by system (9) after a change of variables.

4.1 Recovery of known 4-simplex maps

Here, we present some examples of known 4-simplex maps which can be recovered by the more general solutions to the local tetrahedron equation of the previous section.

Example 4.1. (Kashaev–Sergeev–Korepanov type 4-simplex maps) Changing in (9)

$$(x_1, y_1, z_1, t_1, u_1, v_1, w_1, r_1) \to (1, 1, 1, 1, 1, 1, 1, 1),$$

$$(x_2, y_2, z_2, t_2, u_2, v_2, w_2, r_2) \to (x_1, y_1, z_1, t_1, u_1, v_1, w_1, r_1),$$

$$(x_3, y_3, z_3, t_3, u_3, v_3, w_3, r_3) \to \left(\frac{k}{x_1}, \frac{k}{y_1}, \frac{k}{z_1}, \frac{k}{t_1}, \frac{k}{u_1}, \frac{k}{v_1}, \frac{k}{v_1}, \frac{k}{r_1}\right),$$

$$(x_4, y_4, z_4, t_4, u_4, v_4, w_4, r_4) \to (0, 0, 0, 0, 0, 0, 0, 0),$$

$$(x_5, y_5, z_5, t_5, u_5, v_5, w_5, r_5) \to (x_2, y_2, z_2, t_2, u_2, v_2, w_2, r_2),$$

and solving the associated equations for u_i , v_i , w_i and r_i , i = 1, 2, we obtain the following correspondence between \mathbb{C}^8 and \mathbb{C}^8 :

$$u_1 = \frac{x_1 y_1}{y_1 + x_1 z_1}, \quad u_2 = x_2, \quad v_1 = \frac{x_1 z_1}{k}, \quad v_2 = y_2, \quad w_1 = \frac{y_1 + x_1 z_1}{x_1}, \quad r_1 = \frac{t_1 y_2}{x_2}, \quad r_2 = \frac{t_2 z_2}{w_2}.$$

Fixing $w_2 = t_2$ or $w_2 = y_2$, we recover maps

$$S_{1}: (x_{1}, x_{2}, y_{1}, y_{2}, z_{1}, z_{2}, t_{1}, t_{2}) \rightarrow \left(\frac{x_{1}y_{1}}{y_{1} + x_{1}z_{1}}, x_{2}, \frac{x_{1}z_{1}}{k}, y_{2}, \frac{y_{1} + x_{1}z_{1}}{x_{1}}, t_{2}, \frac{t_{1}y_{2}}{x_{2}}, z_{2}\right),$$

$$S_{2}: (x_{1}, x_{2}, y_{1}, y_{2}, z_{1}, z_{2}, t_{1}, t_{2}) \rightarrow \left(\frac{x_{1}y_{1}}{y_{1} + x_{1}z_{1}}, x_{2}, \frac{x_{1}z_{1}}{k}, y_{2}, \frac{y_{1} + x_{1}z_{1}}{x_{1}}, y_{2}, \frac{t_{1}y_{2}}{x_{2}}, \frac{t_{2}z_{2}}{y_{2}}\right),$$

respectively, from [17].

Example 4.2. (Hirota type 4-simplex maps) Changing in (9) $(x_2, y_2, z_2, t_2, u_2, v_2, w_2, r_2) \rightarrow (1, 1, 1, 1, 1, 1, 1)$, and then

$$(x_1, y_1, z_1, t_1, u_1, v_1, w_1, r_1) \to \left(\frac{x_1}{x_2}, \frac{y_1}{y_2}, \frac{z_1}{z_2}, \frac{t_1}{t_2}, \frac{u_1}{u_2}, \frac{v_1}{v_2}, \frac{w_1}{w_2}, \frac{r_1}{r_2}\right),$$

$$(x_3, y_3, z_3, t_3, u_3, v_3, w_3, r_3) \to (1, 1, 1, 1, 1, 1, 1, 1, 1),$$

$$(x_4, y_4, z_4, t_4, u_4, v_4, w_4, r_4) \to (0, 0, 0, 0, 0, 0, 0, 0, 0),$$

$$(x_5, y_5, z_5, t_5, u_5, v_5, w_5, r_5) \to (x_2, y_2, z_2, t_2, u_2, v_2, w_2, r_2),$$

and solving the associated equations for u_i , v_i , w_i and r_i , i = 1, 2, we obtain the following correspondence between \mathbb{C}^8 and \mathbb{C}^8 :

$$u_1 = \frac{x_1y_1z_2}{x_2z_1 + x_1z_2}, \ u_2 = y_2, \ v_1 = \frac{x_2z_1 + x_1z_2}{z_2}, \ v_2 = x_2, \ w_1 = \frac{w_2x_2y_1z_1}{y_2(x_2z_1 + x_1z_2)}, \ r_1 = \frac{t_1x_2z_2}{w_2y_2}, \ r_2 = \frac{t_2z_2}{w_2}.$$

For the choices of the free variable $w_2 = t_2$ and $w_2 = z_2$ the above correspondence defines the following Hirota type 4-simplex maps:

$$S_{1}: (x_{1}, x_{2}, y_{1}, y_{2}, z_{1}, z_{2}, t_{1}, t_{2}) \rightarrow \left(\frac{x_{1}y_{1}z_{2}}{x_{2}z_{1} + x_{1}z_{2}}, y_{2}, \frac{x_{2}z_{1} + x_{1}z_{2}}{z_{2}}, x_{2}, \frac{t_{2}x_{2}y_{1}z_{1}}{y_{2}(x_{2}z_{1} + x_{1}z_{2})}, t_{2}, \frac{t_{1}x_{2}z_{2}}{t_{2}y_{2}}, z_{2}\right),$$

$$S_{2}: (x_{1}, x_{2}, y_{1}, y_{2}, z_{1}, z_{2}, t_{1}, t_{2}) \rightarrow \left(\frac{x_{1}y_{1}z_{2}}{x_{2}z_{1} + x_{1}z_{2}}, y_{2}, \frac{x_{2}z_{1} + x_{1}z_{2}}{z_{2}}, x_{2}, \frac{x_{2}y_{1}z_{1}z_{2}}{y_{2}(x_{2}z_{1} + x_{1}z_{2})}, z_{2}, \frac{t_{1}x_{2}}{y_{2}}, t_{2}\right),$$

which appeared in [18].

The above are illustrative examples, and after certain change of variables, one can recover all the 4-simplex maps which appeared in [17, 18]. Moreover, we can construct more general 4-simplex maps which can be restricted to the 4-simplex maps presented in [17, 18] at a particular limit. At the same time, as we shall see in the next section, we can construct new hierarchies of 4-simplex maps, other than the ones generated by matrix A in (8), by employing matrices B, C and D in (8).

4.2 Construction of novel 4-simplex maps

In section 3, we studied the solutions of the local Zamolodchikov tetrahedron equation that are obtained by 3×3 matrices. In this section, we employ the systems of polynomial equations (9), (10), (11) and (12), for the construction of novel 4-simplex maps. We will adopt the correspondence approach [10].

We start with correspondence (9). In (9) there are 14 independent equations. Indeed, for the system of equations

$$P_i(u_1, \dots u_5, v_1, \dots v_5, w_1, \dots w_5, r_1, \dots r_5, x_1, \dots x_5, y_1, \dots y_5, z_1, \dots z_5, t_1, \dots t_5) = 0, \quad i = 1, \dots, 14,$$

where

$$\begin{split} P_1 &= u_1v_1 - x_1y_1, \quad P_2 = u_2w_1 + u_1v_2w_3 - x_2y_1, \quad P_3 = u_2w_2 + u_1v_2w_4 - y_2, \quad P_4 = u_3v_1 - x_3z_1 - x_1y_3z_2, \\ P_5 &= u_4w_1 + u_3v_2w_3 - x_4z_1 - x_2y_3z_2, \quad P_6 = u_4w_2 + u_3v_2w_4 - y_4z_2, \quad P_7 = r_1u_5 - t_1, x_5, \\ P_8 &= r_2u_5 - t_2y_5, \quad P_9 = v_3 - x_3z_3 - x_1y_3z_4, \quad P_{10} = v_4w_3 - x_4z_3 - x_2y_3z_4, \quad P_{11} = v_4w_4 - y_4z_4, \\ P_{12} &= r_3v_5 - t_3x_5, \quad P_{13} = r_4v_5 - t_4y_5, P_{14} = r_5w_5 = t_5z_5, \end{split}$$

we have that the 14×20 matrix $[Q_1 \dots Q_{14}]$, consisting of vector rows $Q_i = \nabla P_i$, $i = 1, \dots, 14$ has rank 14. That is, in order to define a map, we need 6 (= 20 - 14) more equations.

In what follows, we shall see that for particular supplementary equations, the correspondence (9) implies a novel 4-simplex map.

Proposition 4.3. The map $S_A: \mathbb{C}^{20} \to \mathbb{C}^{20}$,

$$(x_1, \dots, x_5, y_1, \dots, y_5, z_1, \dots, z_5, t_1, \dots, t_5) \xrightarrow{S_A} (u_1, \dots, u_5, v_1, \dots, v_5, w_1, \dots, w_5, r_1, \dots, r_5),$$
 (13)

where

$$x_1 \mapsto u_1 = x_1 y_1$$

$$\begin{aligned} x_2 \mapsto u_2 &= \frac{(x_2x_3 - x_1x_4) \left[x_4y_2z_3 + x_2(y_2y_3 - y_1y_4)z_4\right]}{(x_3y_2z_3 + x_1y_1y_4z_4)x_4z_1 + (x_2x_3z_1z_4 + x_1x_2y_3z_2z_4 + x_1x_4z_2z_3)(y_2y_3 - y_1y_4)}, \\ x_3 \mapsto u_3 &= x_3z_1 + x_1y_3z_2, \\ x_4 \mapsto u_4 &= \frac{x_4y_4(x_1x_4 - x_2x_3)(z_1z_4 - z_2z_3)}{(x_3y_2z_3 + x_1y_1y_4z_4)x_4z_1 + (x_2x_3z_1z_4 + x_1x_2y_3z_2z_4 + x_1x_4z_2z_3)(y_2y_3 - y_1y_4)}, \\ x_5 \mapsto u_5 &= x_5, \\ y_1 \mapsto v_1 &= 1, \\ y_2 \mapsto v_2 &= \frac{x_4y_2z_1 + x_2(y_2y_3 - y_1y_4)z_2}{(x_1x_4 - x_2x_3)(z_1z_4 - z_2z_3)}, \\ y_3 \mapsto v_3 &= x_3z_3 + x_1y_3z_4, \\ y_4 \mapsto v_4 &= \frac{(x_3y_2z_3 + x_1y_1y_4z_4)x_4z_1 + (x_2x_3z_1z_4 + x_1x_2y_3z_2z_4 + x_1x_4z_2z_3)(y_2y_3 - y_1y_4)}{(x_1x_4 - x_2x_3)(z_1z_4 - z_2z_3)}, \\ y_5 \mapsto v_5 &= y_5, \\ z_1 \mapsto w_1 &= y_1z_1, \\ z_2 \mapsto w_2 &= \frac{x_1z_2(y_1y_4 - y_2y_3) - x_3y_2z_1}{x_1x_4 - x_2x_3}, \\ z_3 \mapsto w_3 &= \frac{(x_1x_4 - x_2x_3)(x_4z_3 + x_2y_3z_4)(z_1z_4 - z_2z_3)}{(x_3y_2z_3 + x_1y_1y_4z_4)x_4z_1 + (x_2x_3z_1z_4 + x_1x_2y_3z_2z_4 + x_1x_4z_2z_3)(y_2y_3 - y_1y_4)}, \\ z_4 \mapsto w_4 &= \frac{(x_1x_4 - x_2x_3)(x_4z_3 + x_2y_3z_4)(z_1z_4 - z_2z_3)}{(x_3y_2z_3 + x_1y_1y_4z_4)x_4z_1 + (x_2x_3z_1z_4 + x_1x_2y_3z_2z_4 + x_1x_4z_2z_3)(y_2y_3 - y_1y_4)}, \\ z_5 \mapsto w_5 &= z_5, \\ t_1 \mapsto t_1 &= t_1, \\ t_2 \mapsto t_2 &= \frac{t_2y_5}{x_5}, \\ t_4 \mapsto t_1 &= t_4, \\ t_5 \mapsto t_7 &= t_5, \end{aligned}$$

is a noninvolutive 4-simplex map and it admits the following functionally independent invariants:

$$I_1 = x_1 y_1, \quad I_2 = y_1 z_1, \quad I_3 = x_5, \quad I_4 = y_5, \quad I_5 = z_5,$$
 (14)

$$I_6 = t_1, \quad I_7 = t_5, \quad I_8 = t_2 t_3, \quad I_9 = y_4 z_4, \quad I_{10} = x_4 y_4.$$
 (15)

Proof. Map S_A can be obtained as a unique solution for u_i , v_i , w_i and r_i of the system of equations (9) supplemented by the following equations:

$$u_1 = x_1 y_1, \quad u_5 = x_5, \quad v_5 = y_5, \quad w_5 = z_5,$$

 $u_5(u_1 u_4 - u_2 u_3) = x_5(x_1 x_4 - x_2 x_3), \quad v_5(v_1 v_4 - v_2 v_3) = y_5(y_1 y_4 - y_2 y_3).$

The fact that S_A is a 4-simplex map can be verified by straightforward substitution of (13) to the 4-simplex equation (2). Moreover, we have, for instance, that $r_2 \circ S_A = \frac{t_2 y_5^2}{x_5^2} \neq t_2$, namely ${}_AS^2 \neq \mathrm{id}$, which means that S is noninvolutive.

Finally, it can be checked that $I_i \circ S_A = I_i$, i = 1, ..., 10, and that the vectors ∇I_i , i = 1, ..., 10, are linearly independent, which means that I_i , i = 1, ..., 10, are functionally independent invariants of map S_A .

Now, it is easy to show that (10) has 14 independent equations. In order to define a map, we need 6 supplementary equations. In particular, we have the following.

Proposition 4.4. The map $S_B: \mathbb{C}^{20} \to \mathbb{C}^{20}$,

$$(x_1, \dots, x_5, y_1, \dots, y_5, z_1, \dots, z_5, t_1, \dots, t_5) \xrightarrow{S_B} (u_1, \dots, u_5, v_1, \dots, v_5, w_1, \dots, w_5, r_1, \dots, r_5),$$
 (16)

where

$$\begin{aligned} x_1 &\mapsto u_1 = x_1 y_1, \\ x_2 &\mapsto u_2 = \frac{(x_2 x_3 - x_1 x_4) \left[x_4 y_2 z_3 + x_2 (y_2 y_3 - y_1 y_4) z_4 \right]}{(x_3 y_2 z_3 + x_1 y_1 y_4 z_4) x_4 z_1 + (x_2 x_3 z_1 z_4 + x_1 x_2 y_3 z_2 z_4 + x_1 x_4 z_2 z_3) (y_2 y_3 - y_1 y_4)}, \\ x_3 &\mapsto u_3 = x_3 z_1 + x_1 y_3 z_2, \\ x_4 &\mapsto u_4 = \frac{x_4 y_4 (x_1 x_4 - x_2 x_3) (z_1 z_4 - z_2 z_3)}{(x_3 y_2 z_3 + x_1 y_1 y_4 z_4) x_4 z_1 + (x_2 x_3 z_1 z_4 + x_1 x_2 y_3 z_2 z_4 + x_1 x_4 z_2 z_3) (y_2 y_3 - y_1 y_4)}, \\ x_5 &\mapsto u_5 = x_5, \\ y_1 &\mapsto v_1 = 1, \\ y_2 &\mapsto v_2 = \frac{x_4 y_2 z_1 + x_2 (y_2 y_3 - y_1 y_4) z_2}{(x_1 x_4 - x_2 x_3) (z_1 z_4 - z_2 z_3)}, \\ y_3 &\mapsto v_3 = x_3 z_3 + x_1 y_3 z_4, \\ y_4 &\mapsto v_4 = \frac{(x_3 y_2 z_3 + x_1 y_1 y_4 z_4) x_4 z_1 + (x_2 x_3 z_1 z_4 + x_1 x_2 y_3 z_2 z_4 + x_1 x_4 z_2 z_3) (y_2 y_3 - y_1 y_4)}{(x_1 x_4 - x_2 x_3) (z_1 z_4 - z_2 z_3)}, \\ y_5 &\mapsto v_5 &= y_5, \\ z_1 &\mapsto w_1 &= y_1 z_1, \\ z_2 &\mapsto w_2 &= \frac{x_1 z_2 (y_1 y_4 - y_2 y_3) - x_3 y_2 z_1}{x_1 x_4 - x_2 x_3}, \\ z_3 &\mapsto w_3 &= \frac{(x_1 x_4 - x_2 x_3) (x_4 z_3 + x_2 y_3 z_4) (z_1 z_4 - z_2 z_3)}{(x_3 y_2 z_3 + x_1 y_1 y_4 z_4) x_4 z_1 + (x_2 x_3 z_1 z_4 + x_1 x_2 y_3 z_2 z_4 + x_1 x_4 z_2 z_3) (y_2 y_3 - y_1 y_4)}, \\ z_4 &\mapsto w_4 &= \frac{(x_1 x_4 - x_2 x_3) (x_4 z_3 + x_2 y_3 z_4) (z_1 z_4 - z_2 z_3)}{(x_3 y_2 z_3 + x_1 y_1 y_4 z_4) x_4 z_1 + (x_2 x_3 z_1 z_4 + x_1 x_2 y_3 z_2 z_4 + x_1 x_4 z_2 z_3) (y_2 y_3 - y_1 y_4)}, \\ z_5 &\mapsto w_5 &= y_5 t_5, \\ t_1 &\mapsto t_1 &= z_1, \\ t_2 &\mapsto t_2 &= \frac{y_5 z_2}{x_5}, \\ t_3 &\mapsto t_3 &= \frac{x_5 z_3}{y_5}, \\ t_4 &\mapsto t_4 &= z_4, \\ t_5 &\mapsto t_7 &= \frac{z_5}{y_5}, \\ t_4 &\mapsto t_4 &= z_4, \\ t_5 &\mapsto t_7 &= \frac{z_5}{y_5}, \end{aligned}$$

is a noninvolutive 4-simplex map and it admits the following functionally independent invariants:

$$I_1 = x_1 y_1$$
, $I_2 = y_1 z_1$, $I_3 = x_5$, $I_4 = y_5$, $I_5 = y_4 z_4$, $I_6 = x_4 y_4$
 $I_7 = t_5 y_5 z_5$, $I_8 = x_1 x_4 - x_2 x_3$, $I_9 = y_1 y_4 - y_2 y_3$, $I_{10} = z_1 z_4 - z_2 z_3$.

Proof. Map S_B can be obtained as a unique solution for u_i , v_i , w_i and r_i of the system of equations (10) supplemented by the following equations:

$$u_1 = x_1 y_1, \quad u_5 = x_5, \quad v_5 = y_5, \quad w_5 r_5 = z_5 t_5,$$

 $u_5(u_1 u_4 - u_2 u_3) = x_5(x_1 x_4 - x_2 x_3), \quad v_5(v_1 v_4 - v_2 v_3) = y_5(y_1 y_4 - y_2 y_3).$

It can be verified by straightforward substitution that S_B satisfies the 4-simplex equation (2). Additionally, we have, for instance, that $w_1 \circ S_B = y_1 z_1 \neq y_1$, namely $S_B^2 \neq \text{id}$, i.e. S_B is noninvolutive.

Finally, $I_i \circ S_B = I_i$, i = 1, ..., 10, which means that I_i , i = 1, ..., 10, are invariants of S_B . Also, vectors ∇I_i , i = 1, ..., 10, are linearly independent, that is I_i , i = 1, ..., 10, are functionally independent. \square

Next, consider system (11). It is easy to verify that the latter has 10 independent equations. In order to define a map, we need 6 supplementary equations. Specifically, we have the following.

Proposition 4.5. The map $S_C: \mathbb{C}^{16} \to \mathbb{C}^{16}$,

$$(x_1, \dots, x_4, y_1, \dots, y_4, z_1, \dots, z_4, t_1, \dots, t_4) \xrightarrow{S_C} (u_1, \dots, u_4, v_1, \dots, v_4, w_1, \dots, w_4, r_1, \dots, r_4),$$
 (17)

$$\begin{aligned} x_1 &\mapsto u_1 = \frac{y_1}{t_1}, & z_1 &\mapsto w_1 = t_1, \\ x_2 &\mapsto u_2 = y_2, & z_2 &\mapsto w_2 = 1, \\ x_3 &\mapsto u_3 = y_3, & z_3 &\mapsto w_3 = t_3 y_3, \\ x_4 &\mapsto u_4 = y_4, & z_4 &\mapsto w_4 = t_4 y_4, \\ y_1 &\mapsto v_1 = x_1 z_1, & t_1 &\mapsto r_1 = \frac{t_2 x_4 z_1}{t_2 x_4 + t_3 x_1 y_2 z_4}, \\ y_2 &\mapsto v_2 = x_2 z_2, & t_2 &\mapsto r_2 = z_2, \end{aligned}$$

$$y_3 \mapsto v_3 = x_3,$$
 $t_3 \mapsto r_3 = \frac{z_3}{x_3},$ $y_4 \mapsto v_4 = t_2 x_4 + t_3 x_1 y_2 z_4,$ $t_4 \mapsto r_4 = \frac{z_4}{t_2 x_4 + t_3 x_1 y_2 z_4},$

is a noninvolutive 4-simplex map and it admits the following functionally independent invariants:

$$I_1 = x_3y_3$$
, $I_2 = x_2y_2z_2$, $I_3 = y_3z_3t_3$, $I_4 = x_1y_1z_1$, $I_5 = x_3 + y_3$, $I_6 = y_1 + x_1z_1$, $I_7 = z_3 + t_3y_3$, $I_8 = y_2 + x_2z_2$.

Proof. Supplementing the system of equations (11) by the following equations:

$$u_2v_1 = x_1y_2z_1$$
, $v_3 = x_3$, $w_3 = t_3y_3$, $u_3 = y_3$, $w_1 = t_1$, $u_2 = y_2$

we obtain a unique solution for u_i , v_i , w_i and r_i , i = 1, 2, 3, 4, which defines map (17).

Substitution that S_C to the 4-simplex equation (2) confirms that map (17) is a 4-simplex map. Moreover, we have, for instance, that $w_1 \circ S_C = \frac{t_2 x_4 z_1}{t_2 x_4 + t_3 x_1 y_2 z_4} \neq z_1$, namely $S_C^2 \neq \text{id}$, i.e. S_C is noninvolutive. Finally, $I_i \circ S_C = I_i$, $i = 1, \ldots, 8$, which means that I_i , $i = 1, \ldots, 8$, are invariants of S_C . Furthermore,

Finally, $I_i \circ S_C = I_i$, i = 1, ..., 8, which means that I_i , i = 1, ..., 8, are invariants of S_C . Furthermore, vectors ∇I_i , i = 1, ..., 8, are linearly independent, which implies that I_i , i = 1, ..., 8, are functionally independent.

Lastly, we consider system (12). It can be verified that the (12) has 6 independent equations. In order to define a map, we need 6 supplementary equations. In particular, we have the following.

Proposition 4.6. The map $S_D: \mathbb{C}^{12} \to \mathbb{C}^{12}$,

$$(x_1, x_2, x_3, y_1, y_2, y_3, z_1, z_2, z_3, t_1, t_2, t_3) \xrightarrow{S_C} \left(x_1, x_2, x_3, x_1 z_1, y_2, y_3, \frac{t_1 x_2}{y_2}, t_2 y_2, z_3, \frac{t_1 x_3}{y_3}, \frac{t_2 y_3}{z_3}, t_3\right), \quad (18)$$

is a noninvolutive 4-simplex map and it admits the following functionally independent invariants:

$$I_1 = x_1$$
, $I_2 = x_2$, $I_3 = x_3$, $I_4 = y_2$, $I_5 = y_3$, $I_6 = z_3$.

Proof. Supplementing system (12) by the following equations:

$$u_1 = x_1, \quad u_2 = x_2, \quad u_3 = x_3, \quad v_2 = y_2, \quad v_3 = y_3, \quad w_3 = z_3,$$

we obtain a unique solution for u_i , v_i , w_i and r_i , i = 1, 2, 3, which defines map (18).

After substitution of S_D to the 4-simplex equation (2) we obtain that map (18) is a 4-simplex map. Additionally, we have, for instance, that $r_1 \circ S_D = \frac{t_1 x_3^2}{y_3^2} \neq t_1$, namely $S_D^2 \neq \text{id}$, thus S_D is noninvolutive.

The invariants of the map are obvious and so is their functional independence.

5 Conclusions

In this paper we studied the set-theoretical solutions to the local Zamolodchikov tetrahedron equation that generate set-theoretical solutions to the Bazhanov–Stroganov 4-simplex equation. In particular, we studied the solutions of the local tetrahedron equation in the form of correspondences between \mathbb{C}^n and \mathbb{C}^n , generated by 3×3 matrices. Although, usually, in the literature n-simplex maps appear as unique solutions to the (n-1)-simplex equation, we adopt the correspondence approach [10] which gives rise to much wider hierarchies of solutions.

Specifically, we found that all the essentially different solutions to the Zamolodchikov tetrahedron equation generated by 3×3 matrices are given by matrices A, B, C and D in (8). In fact, these matrices generate all the possible solutions to the local 4-simplex equation in the form of correspondences (9), (10), (11) and (12). We demonstrated that (9) generates all known 4-simplex maps, using Kashaev–Korepanov–Sergeev and Hirota type 4-simplex maps as illustrative examples. Then, we employed the correspondences (9), (10), (11) and (12) to construct new 4-simplex maps, namely maps (13), (16), (17) and (18). All these maps are noninvolutive which, in terms of dynamics, are more interesting that involutive maps which have trivial dynamics.

Our results can be extended in the following ways:

 \bullet Study the Liouville integrability of the 4-simplex maps presented in this paper.

We demonstrated that all the 4-simplex maps constructed in this paper admit enough functionally independent first integrals; that is, half the dimension of the map in number. This already indicates their integrability, however their complete (Liouville) integrability is an open problem.

- Employ the correspondences (9), (10), (11) and (12) to construct parametric 4-simplex maps.

 We focused on non-parametric 4-simplex maps in this paper, however, supplementing systems (9), (10), (11) and (12) by equations containing parameters, one may construct new solutions to the parametric 4-simplex equation. Several parametric 4-simplex maps were presented in [17].
- Classify the solutions of the local tetrahedron equation.

Our motivation to study the solutions of the local Zamolodchikov tetrahedron equation is that such solutions generate 4-simplex maps. This, together with the fact that correspondences satisfying the local tetrahedron equation may give generate infinitely many solutions (and also give rise to solutions that do not belong to the known classification lists [10]), suggests that it makes sense to classify the solutions of the local tetrahedron equation. The classification of the latter solutions is an open problem.

In general, we propose to classify the generators of n-simplex maps, namely the solutions to the local n-simplex equation. This will give rise to wider from the already known classification lists even in the case of Yang-Baxter (2-simplex) maps.

• Study the solutions of the local pentagon equation.

The n-gon equations have equally rich algebro-geometric structure as the n-simplex equations [5, 7, 24]. The most famous member of the family of n-gon equations is the 5-gon or pentagon equation (see, e.g. [5, 7, 12, 24] and the references therein). Similar results to those obtained in this paper can be obtained for the local pentagon equation. Again, the motivation comes from the fact that the local (n-1)-gon equation is a generator of solutions to the n-gon equation.

• Extend the results to the noncommutative case.

Noncommutative n-simplex maps have been in the centre of interest for scientists from the mathematical physics community (see, e.g. [1, 6, 13, 16] and the references therein), due to the rapid development of noncommutative integrable systems. One could extend our results by considering that the elements of matrix (6) are matrices, or, more generally, free variables from a noncommutative division ring.

Acknowledgements

This work was funded by the Russian Science Foundation project No. 20-71-10110 (https://rscf.ru/en/project/23-71-50012/). Part of this work, namely the proof of Theorem 3.1, was carried out within the framework of a development programme for the Regional Scientific and Educational Mathematical Centre of the P.G. Demidov Yaroslavl State University with financial support from the Ministry of Science and Higher Education of the Russian Federation (Agreement on provision of subsidy from the federal budget No. 075-02-2023-948).

We would like to thank S. Igonin for several useful discussions.

References

[1] P. Adamopoulou and G. Papamikos Entwining Yang-Baxter maps over Grassmann algebras. arXiv:2311.18673 (2003).

- [2] V. Adler, A. Bobenko, and Y. Suris Classification of integrable equations on quad-graphs. The consistency approach. Comm. Math. Phys. **233** (2003) 513–543.
- [3] V. Bardakov, B. Chuzinov, I. Emel'yanenkov, M. Ivanov, T. Kozlovskaya, V. Leshkov, Settheoretical solutions of simplex equations arXiv:2206.08906 (2022).
- [4] V.V. Bazhanov and Yu.G. Stroganov. Conditions of commutativity of transfer matrices on a multidimensional lattice, Theor. Math. Phys. 52 (1982) 685–691.
- [5] A. Dimakis, F. Müller-Hoissen, Simplex and polygon equations, Symmetry, Integrability and Geometry: Methods and Applications (SIGMA) 11 (2015) 042.
- [6] A. Doliwa and R.M. Kashaev. Non-commutative bi-rational maps satisfying Zamolodchikov equation equation, and Desargues lattices, J. Math. Phys. 61 (2020) 092704.
- [7] A. Doliwa and S.M. Sergeev. The pentagon relation and incidence geometry, J. Math. Phys. 55 (2014) 063504.
- [8] V.G. Drinfeld, On some unsolved problems in quantum group theory, Lecture Notes in Math. 1510, Berlin: Springer (1992) 1–8.
- [9] J. Hietarinta, Permutation-type solutions to the Yang–Baxter and other n-simplex equations, J. Phys. A: Math. Gen. 30 (1997) 4757.
- [10] S. Igonin and S. Konstantinou-Rizos. Local Yang-Baxter correspondences and set-theoretical solutions to the Zamolodchikov tetrahedron equation. J. Phys. A: Math. Theor. 56 275202 (18pp) (2023).
- [11] R.M. Kashaev, I.G. Koperanov, and S.M. Sergeev, Functional Tetrahedron Equation, Theor. Math. Phys. 117 (1998) 370–384.
- [12] P. Kassotakis. Matrix factorizations and pentagon maps, Proc. R. Soc. A: Math. Phys. Eng. Sci. 479 (2023) 20230276.
- [13] P. Kassotakis and T. Kouloukas, On non-abelian quadrirational Yang–Baxter maps, J. Phys. A: Math. Theor. 55 (2022) 175203.
- [14] P. Kassotakis, M. Nieszporski, V. Papageorgiou, A. Tongas, Tetrahedron maps and symmetries of three dimensional integrable discrete equations, J. Math. Phys. 60 (2019) 123503.
- [15] S. Konstantinou-Rizos. Nonlinear Schrödinger type tetrahedron maps. Nuclear Phys. B 960 (2020) 115207
- [16] S. Konstantinou-Rizos. Noncommutative solutions to Zamolodchikov's tetrahedron equation and matrix six-factorisation problems. Physica D 440 (2022) 133466
- [17] S. Konstantinou-Rizos. Birational solutions to the set-theoretical 4-simplex equation. Physica D 448 (2023) 133696
- [18] S. Konstantinou-Rizos. Hirota type Bazhanov-Stroganov maps. arXiv:2303.13333 (2024)

- [19] S. Konstantinou-Rizos and T. Kouloukas, A noncommutative discrete potential KdV lift, J. Math. Phys. 59 (2018) 063506
- [20] S. Konstantinou-Rizos and A.V. Mikhailov. Darboux transformations, finite reduction groups and related Yang-Baxter maps. J. Phys. A: Math. Theor. 46 (2013) 425201
- [21] S. Konstantinou-Rizos, G. Papamikos, Entwining Yang-Baxter maps related to NLS type equations, Phys. A: Math. Theor. 52 (2019) 485201.
- [22] T.E. Kouloukas and V.G. Papageorgiou Yang–Baxter maps with first-degree-polynomial 2×2 Lax matrices J. Phys. A **42** 404012 (2009)
- [23] A.V. Mikhailov, G. Papamikos and J.P. Wang, Darboux Transformation for the Vector sine-Gordon Equation and Integrable Equations on a Sphere, Lett. Math. Phys. 106 (2016) 973–996.
- [24] F. Müller-Hoissen. On the structure of set-theoretic polygon equations, arXiv:2305.17974 (2023).
- [25] V.G. Papageorgiou, A.G. Tongas, A.P. Veselov, Yang–Baxter maps and symmetries of integrable equations on quad-graphs, J. Math. Phys. 47 (2006) 083502.
- [26] S.M. Sergeev, Solutions of the Functional Tetrahedron Equation Connected with the Local Yang–Baxter Equation for the Ferro-Electric Condition, Lett. Math. Phys. 45 (1998) 113–119.
- [27] A.B. Zamolodchikov, Tetrahedra equations and integrable systems in three-dimensional space, Sov. Phys. JETP 52 (1980) 325–336.
- [28] A.B. Zamolodchikov, Tetrahedron equations and the relativistic S matrix of straight strings in (2+1)-dimensions, Commun. Math. Phys. 79 (1981) 489–505.