

Stability distillation hypothesis for the origin of life

Cheng Bi

Email: cb236@foxmail.com

Abstract

The essence of life is a frequency distribution conversion process of molecules accompanied by changes in information. In wet-dry cycling hot springs, RNA of different sequences continuously undergoes polymerization and decomposition reactions, products with stable configurations will accumulate their frequency among all RNA molecules and finally become an important source of DNA coding and non-coding regions in primitive cells. This periodic extreme environmental change narrows the frequency distribution of macromolecules through the distillation process and new potential catalysts are searched. When enough macromolecules are accumulated and reliable reaction pathways are built, phospholipids randomly wrap the macromolecules in the hot springs which I would call it the pioneer pools, and these protocells become parallel calculators of molecule frequency distribution. Through the selective permeability of the cell membrane to different molecular weights and properties, cells with appropriate distribution have more opportunities to absorb small molecule substances and increase its intracellular frequencies of molecules. This will strongly induce the occurrence of macromolecules that can widely catalyze the synthesis of other macromolecules or themselves, such as ribosomes, etc. Rupture and fusion during cell division make all protocells share the same frequency distribution of molecules during the origin of life, thus ensuring that all present cells have very similar genetic materials and protein translation systems. This also suggests that viruses may have originated and evolved together with cells.

Different from the classic RNA world hypothesis, the original Earth environment assumed in the stability distillation hypothesis allow both DNA and RNA genetic materials as well as various macromolecules and small molecule substances, and it does not require any single macromolecule to be able to stably self-replicate and self-catalyze at the beginning of life, nor does it require a stable or mild environment. On the contrary, extreme environments and irregular cell division are key to the emergence of life in this hypothesis. According to this hypothesis, the most likely location for the origin of life is geysers that erupt with appropriate cycles on land. In this cyclically changing extreme environment, DNA and RNA can also be rapidly synthesized through methods similar to PCR and reverse transcription PCR, and DNA can track the distribution of RNA at all times through its own continuous decomposition and replication as well as reverse transcription of RNA. This ultimately allows all living macromolecules to become entangled and co-evolve in a form of life. This hypothesis can be verified in the laboratory through three possible experiments noted in this article,

and can also be used to partially restore the process of the origin of life through computer simulation.

Cell division and osmotic pressure

People usually believe that the self-replication of a few macromolecules such as nucleic acids is the key to the origin of life [1-2]. However, for organisms, it is not only DNA that ultimately replicates, but the distribution of all intracellular substances, including DNA, RNA, proteins, lipids and other substances, although these distributions are shifted sometimes in the progeny cells. The reproduction usually involves the generation and replication of a large number of macromolecular substances, and when the total amount of macromolecules increases, the osmotic pressure within the cells will objectively increase, causing the cell membrane to bear greater pressure. If this increase in intracellular osmotic pressure is also one of the driving forces for primitive cell division, this means that for cells during the origin of life, increasing the total amount of macromolecules may be an important evolutionary direction of primitive life. The replication of nucleic acids can be considered a special method of accumulating macromolecules. This means molecules or mechanisms that can broadly synthesize or catalyze small molecules into macromolecules can help cells to quickly accumulate macromolecular substances and become an important macromolecule of life. In other words, if the assumption is correct, it will be possible to naturally select the core molecules in today's cells during the origin of life, including ribosomes, tRNA, DNA polymerase, RNA polymerase, RNA ligase, ATP, etc.

Frequency distribution and information entropy

When early life macromolecules did not evolve the ability to replicate themselves, the production of macromolecules would rely on reliable reaction pathways or metabolic pathways rather than chaos reactions. This means that a desired product can be produced with certainty in large quantities under appropriate conditions and given sufficient substrate. As for randomly generated macromolecules, even if they have strong catalytic capabilities, they cannot become reliable life macromolecules, this is partially because when a macromolecule is randomly generated, its possible length and sequence combinations are close to infinite, the proportion of any randomly generated macromolecule in products will be extremely low, making it difficult to function in protocells. Even if there is a certain amount of this randomly generated macromolecule in the mother cell, when the cells divide, the frequency of this macromolecule in the daughter cells will rapidly decrease, making it impossible to copy the role it may have played in the mother cells to the offspring cells. This is also the reason why people are so obsessed with finding large molecules that could replicate themselves in the prebiotic Earth.

From a mathematical point of view, a reliable reaction pathway means that the generation of a macromolecular depends on the accumulation of its lower-level

substrate. For example, if RNA can be produced in prebiotic Earth, a reasonable conclusion is that sufficient amounts of ribonucleotides has accumulated in the environment. To generate ribonucleotides, people would expect that a large amount of ribose, phosphate and nucleobase or related intermediate products have accumulated in the environment rather than thinking that carbon, hydrogen, oxygen, nitrogen, phosphorus elements or other simple molecules can directly generate ribonucleotides or even RNA polymers through molecules' collisions. This is because the probability of two molecules combining to form a new product after collision in space is much greater than the probability of multiple molecules colliding at the same time and the same location. Theoretically, the combinations of compounds that can be formed by these chemical elements are infinite which can be easily ignored. When certain substrates are required to generate specific macromolecules, it also means that one or several combinations of these elements need to have a higher frequency than other random sequences. Increasing the frequency of one or more sequences in a variety of random combinations corresponds to a change in information. That is to say, compared to a random system, chemical reactions related to life are more likely to increase the information content of organism or reduce the information entropy. This is especially obvious in the polymerization reaction of monomers. When the polymer chain continues to grow in natural environments, its sequence and length are usually random, and you will end up with a mixture of macromolecules with different sequences and lengths, while the distribution of macromolecules produced by specific reactions within cells is usually much narrower.

Therefore, life can be seen as a conversion process of molecule frequency distribution accompanied by an increase in information content. Assuming that there are several small molecule substances {A, B, C, D...} in the environment, the possible sequence combinations that can be formed include {A-B-C, B-C-D, C-C-B-B-A, D-D, A-A-B-B-D,...}, the original frequency distribution is close to random {p1, p2, p3 , p4, p5,...}, under specific reaction conditions and catalyst, the frequency distribution of the products group becomes {q1, q2, q3, q4, q5}. Relative to randomly generated products, the information entropy change formula is:

$$\Delta Entropy = - \sum_i q(i) \log_2 q(i) - \left(\sum_i p(i) \log_2 p(i) \right)$$

Under the effect of suitable reaction conditions and catalysts, the distribution of the reaction products will become narrower, that is, the entropy value will be lower. This explains from another perspective why life feeds on negative entropy [3].

Therefore, the question becomes how to make a chemical reaction involving multiple reactants and potentially generating macromolecules of any sequence have a higher information content. This is not a problem for most simple reactions because the products are usually fixed, for example, the reaction between hydrogen and oxygen to form water. But for chemical reactions that generate large molecules or mixed reactions that may interfere with each other, most of the time the reactions can't be specific

enough, for example, in successive condensation reactions of multiple amino acids, peptides of any sequence may be generated. This makes the frequency distribution of any specific sequence of macromolecules extremely low and cannot have a reliable generation path or a repeatable generation path. Although suitable catalysts can increase the information content of the entire reaction by allowing substrates with a specific sequence to react or products to decompose from a specific sequence, these catalytic molecules themselves may also be large molecules, thus facing the same problem unless they happen to be able to catalyze themselves. However, the types of such catalytic macromolecules will be very limited, and the requirements for substrates and reaction conditions will be difficult to meet the characteristics of the early Earth. This will make related theoretical hypotheses inevitably fall into a chicken-and-egg paradox.

However, to increase the information content of a complex reaction, a catalyst or a pure single reaction substrate is not necessarily needed. A result-oriented approach is to continuously remove non-target macromolecules from the product, just like repeated purification during a distillation process. There are two possible methods: One is to construct a suitable membrane system that continuously removes by-products while mainly preserving reaction substrates and target products, thereby increasing the information content of the products within the membrane. This will strongly induce the cell membrane to evolve active transport capabilities to change the distribution within the membrane which is very common in today's cells. The other is to continuously decompose by-products into other substrates or small molecules, and retain only specific target macromolecules in each step. For the former, this can be easily associated with the complex membrane system in present cells which still plays an important role, while for the latter, since macromolecules of different sequences, such as RNA, may have different stabilities [4], extreme environments can be used to continuously decompose by-products and only retain macromolecules with stable configurations, thereby narrowing the frequency distribution of products. The aggregation and decomposition of macromolecules can be achieved through periodic external environments. Coincidentally, the existence of hot springs with wet-dry cycles in the early Earth could provide both extreme reaction conditions and periodic environmental changes which is an ideal place for the origin of life.

From another perspective, if we assume that life originated from a self-replication process of a very small number or even a single macromolecule, this requires the prebiotic Earth has sufficient substrates to replicate this specific macromolecule, for example, the establishment of the RNA world hypothesis requires the presence of a large number of ribonucleotides in the environment [32] any other molecules may interfere with the self-replication of this RNA. However, the self-replication of any single macromolecule is unlikely to produce a system complex enough to cope with the extreme environment of the prebiotic Earth. Even if such a fragile replication system can be established, after the introduction of new macromolecules, the original replication mechanism will be changed or even stopped. Therefore, every time it becomes complicated, it is a life and death test for this life form that starts from single-

molecule self-replication. The entire hypothesis will be highly limited by idealized environmental assumptions, and will need to explain every step of how life evolved from a few macromolecules to complex reaction networks.

Cross-catalytic networks and stability distillation hypothesis

It is known that in the hot spring environment with wet-dry cycles, there is an opportunity to form nucleotides [5] and polymerize into macromolecules of RNA [6]. These macromolecules have different folding structures, such as chainlike RNA, ring-shaped RNA [7], hairpin RNA, etc. In the prebiotic Earth environment, these macromolecules may have to face high temperatures, acidic environments and collisions with various other macromolecules, causing a large number of RNA molecules produced in dry environment become unstable and re-decompose into smaller RNA fragments or even ribonucleotides in the wet environment. Some RNA fragments have higher stability due to their folded structures or complexes formed with other macromolecules such as proteins, thereby existing longer and having a greater probability of participating in further reactions. When this wet-dry cycle is repeated, RNA molecules with stable configurations will accumulate and the whole RNA distribution will narrow. This distribution will eventually become the source of the RNA and DNA distribution of the primitive cells. This can also be viewed as a form of natural selection at the molecular level, when these molecules have not yet developed complex cross-catalytic networks and cellular structures.

And the number of paired bases and the sequence of bases will affect the stability of the RNA configuration [8-10]. In other words, RNA fragments with multiple complementary sequences are more likely to form a stable structure with lower energy, which makes randomly generated RNA molecules with stem-loop structures or other similar structures have higher frequencies in the prebiotic Earth due to their higher stability. The size of the loop in the hairpin-like RNA, the number of paired bases, pH value, high temperature and collisions of other macromolecules, etc. may all affect the stability of RNAs and change the frequency distribution of the corresponding RNA sequences in wet-dry cycles. When these RNA molecules with specific configurations and sequences can maintain a stable high frequency distribution in the environment, they may further form larger RNA molecules under the catalysis of other molecules because more numbers give them a greater chance of colliding with each other and reacting. As the new RNA molecules could contain more complementary sequences, this makes the reaction energetically favorable which means the system can search new macromolecules through this cyclic process. When two high-frequency RNA molecules with the same sequence and stem-loop structure are spliced together, it will create molecules that are highly similar to tRNA, and this tRNA-like molecule will exhibit central symmetry. After more tRNAs, stem-loop structured RNAs and other small RNAs are assembled together, it is possible to form the original rRNA and stabilize itself by filling in proteins, ultimately resulting in more stable ribosome-like RNA-protein complexes. There may be many types of ribosome-like complexes, but most

may only have the same high stability as ribosomes but no obvious catalytic activity.




Figure. (A) the green circles represent amino acids and proteins, and the blue circles represent small molecules of RNA and nucleotides. During the wet-dry cycle, RNA with an unstable structure will continue to decompose and the frequencies of RNA molecules with stable structures increase. The continuous accumulation of related substrates increases the frequency of macromolecules with stable structures, while proteins, amino acids and other RNA molecules with complementary sequences can help stabilize their structures. (B) When two RNA molecules with complementary sequences are connected, they will form a tRNA-like structure, which will exhibit centrosymmetric properties. As the stabilities of these tRNA-like molecules with complementary base pairs of different sequences are still different, this may result in only a few types of tRNA-like molecules being retained [8]. There is evidence that tRNA may come from the assembly of two hairpin-like structures of RNA [15].

When considering the presence of other organic matter in hot springs, such as proteins and DNA, and their interactions, the entire network becomes more complicated. The newly generated molecules in the network will become potential catalysts for all existing reactions, it could be small molecules, proteins, RNA, RNA-protein complexes and even DNA. All molecules in the network will also affect both the progress of new reactions and the stability of new products. These macromolecules with catalytic

capabilities, whether they catalyze decomposition reactions or synthesis reactions, may increase the information content of the corresponding products and expand new reactions, because of the specificity and efficiency of enzymes. When a large number of catalytic macromolecules have a reliable generation path and can be continuously generated by cross-catalytic networks, the entire stability distillation network gradually matures. It can be thought that the network is constantly looking for generation paths of new macromolecule in a way similar to distillation. Periodically changing extreme environments can be considered an important way to increase the content of reaction information in the origin of life, which is gradually being replaced by more complex membrane systems and an increasing number of enzymes.

Why RNA?

But not all macromolecules can access high-information content metabolic pathways through the stability distillation. The most important point is that if we treat molecules with different molecular weights and different reaction steps as different layers, it is necessary that among all possible molecules at each layer, there are considerable differences in the stability and there should be ones with high-stability configuration. And the stability needs to be able to increase with the condensation of smaller molecules which favors the generation of larger and stable catalytic macromolecules or the search capabilities of the network will be limited. The extreme environments are conducive to reflecting the difference in stability. On the other hand, the decomposition products in the system need to be able to frequently return to lower levels to retain partial information rather than complete decomposition. For example, proteins can be decomposed into peptides and amino acids, and amino acids can be regarded as stable molecules at the lowest level, however, if lots of proteins are completely decomposed into small molecules such as carbon dioxide, water, etc., the system will lose too much information and process will be slow down due to the lack of a reliable generation path for amino acids. Therefore, the catalyst can be considered as a perfect form that retains information as its frequency distribution does not change before and after the reaction and can catalyze other reactions. This allows it to efficiently accelerate the distillation process while retaining the necessary information. Catalysts also change reaction conditions, such as generally lower the activation energy and making the reaction easier to occur [33], which also objectively expands the areas and forms of life.

Compared with DNA and proteins, RNA-based macromolecules are more likely to change their original frequency distribution through such periodic extreme environmental changes. Whether different bases in an RNA sequence are paired can be considered an internal structure that can produce huge differences in stability. The condensation of two stable structures of RNA is easier to generate a more stable structure of RNA, showing some convergence in the stability of macromolecules. For DNA, its single strand lacks a stable folding configuration [11-12], while the stability difference between double helix DNA with high stability is small. For proteins, due to the wide variety of amino acids, it takes much more time to accumulate highly stable polypeptide fragments of a specific sequence through random reactions, and it cannot

be easily inferred that the larger proteins could have the same or higher stability than its reactants as there are many factors may influence the stability [13].

Since tRNA is a potential high-frequency RNA sequence, when the chemical energy change of the reaction allows, a large number of amino acids will have chance to connect to the corresponding tRNA and tRNA further connected to the template RNA through the pairing of codons and anticodons, then catalyzed by early ribosomes to continuously generate polypeptides and protein and finally evolved the protein translation system. Different from the previous chaotic generation reactions, this allows proteins to have a repeatable generation path due to the high information content matching rules established with RNA, and the distribution of proteins will also be highly dependent on the distribution of RNA in the system, which allows life to copy the distribution of RNA to achieve replication of protein distribution. This also implies that the coding of important proteins in early life will be highly dependent on the stable RNA sequences that exist frequently in the system. In other words, RNA containing a large number of complementary sequences, such as hairpin-like RNA, tRNA, and rRNA may encode a large number of important proteins including the proteins that constitute ribosomes in early stage of life [14].

On the other hand, the realization of this difference in stability requires that most of the sequence combinations can be decomposed. Otherwise, the information content of the generation path will be insufficient, and the system will accumulate too many useless macromolecules and affect the progress of other reactions. When separated from the extreme environment of the prebiotic Earth, various macromolecules will be more likely to exist stable, requires enzymes that can catalyze the decomposition of macromolecules containing specific sequences which objectively increase the information content of the reaction. On the other hand, this also gives some larger molecules the opportunity to form more stable structures because they have more possibilities hide vulnerable sequences by folding in three-dimensional space or combining with other molecules than their smaller substrates. This can also be considered as a compensation and continuation of the original stability distillation process after life escapes from extreme environments, and can explain why RNase and protease in present-day cells is so common. In other words, the origin of life required extreme environments to increase the information content of reactions, while present cells rely more on various enzymes.

The difference in the stability of RNA of different sequences make it become the core macromolecule in the origin of life. If RNA has only two complementary bases, a large number of random fragments will form a stable structure and the information content of the distillation will be insufficient. And if RNA has six or more paired bases, it will make it more difficult for RNA sequences that can form a stable structure to be generated through a random process, greatly prolonging the original frequency accumulation process, and may be overtaken by faster evolving life forms.

The pioneer pools and shared distribution

Based on the above discussion, life likely originated in one or several adjacent hot springs that could undergo wet and dry cycles, especially geyser that could erupt jets of water at appropriate cycles, which I would like to call them the pioneer pools or the pioneer pool.

It is known that a large amount of phospholipid substances may be formed in hot springs on the prebiotic Earth [16], so whether through wet-dry cycles [17] or the disturbance of water flow during geyser eruptions [18], these lipids may self-assemble into a lipid bilayer and then randomly wraps various substances in the pool, becoming protocells. Assuming original cell membrane has selective permeability, making it more difficult for macromolecular substances to pass through the cell membrane, while amino acids or nucleotides can more easily pass through the membrane. This makes these randomly formed cells become parallel molecule distribution calculators, increasing the osmotic pressure through the synthesis of macromolecule substances in the cells, thereby increasing the cells' absorption of various small molecule substances. In this way, substances with reliable production pathways in cells will increase due to the increase in small molecule nutrients, and their frequencies in the pioneer pools will also change accordingly. Self-replication of nucleic acid macromolecules can be considered a special form to change the frequency which can increase at an exponential rate.

Another benefit of geyser is that while maintaining a stable cycle, it will also continuously bring various organic and inorganic substances out of the underground [41], thereby replenishing the pool with various raw materials for life. This is important for the continued evolution of life, because when macromolecules continue to be generated and the number of cells continues to increase, the small molecule substances in the pool will inevitably decrease rapidly, and early life may not be able to directly convert large amounts of inorganic matter into organic matter like cyanobacteria or other bacteria. This periodic replenishment can also be used as a source of energy for protocells.




Figure. The most likely place for the origin of life is a geyser that erupts with a suitable period.

On the other hand, this process does not require perfect cell division or replication. When the cells rupture due to excessive growth of macromolecules and the resulting increase in osmotic pressure, the cell would return the contents of the cells to the pioneer pools, the frequencies of these molecules in the entire pool still change and they will be calculated again by the formation of new cells. This is equivalent to bootstrapping the molecule frequency distribution in the pool. Some cells with appropriate distribution will quickly absorb small molecule substances and increase the frequencies of large molecules in their cells which is equivalent to doing boosting. Changes in solution concentration, temperature changes and water flow disturbances during the wet-dry cycle will also increase the probability of early cell rupture. When there are multiple adjacent pools, the downstream pool can inherit the upstream molecule distribution while allowing cells to have a relatively stable replication environment. When cells can achieve continuous replication and keep the distribution of molecule frequency unchanged in daughter cells, the corresponding frequencies of these molecules in cells will increase exponentially in the pool which means perfect cell division and replication become an inevitable evolutionary direction because exponentially growing distributions can quickly replace other distributions. In this way, life can repeatedly extract molecule assemblies in the form of cells and increase the weight of the correct distributions through methods similar to bootstrapping and boosting.

This kind of parallel computing in the form of cells will be particularly conducive to increasing the frequency of two types of macromolecules. One is a macromolecule that can self-replicate, which is what people usually think of as the starting point of life. The other is a molecule that can catalyze or help the synthesis of a wide range of other macromolecules, such as RNA polymerase, ribosomes, tRNA, ATP, etc. For the cell as a whole, molecule frequency distribution in cells may migrate and integrate, or it may be compressed and then reconstructed before and after the division. Different methods of division and fusion will also produce different life forms, and because of DNA's powerful storage and restoration capabilities for distribution, it is easier for people to only consider the mutation and fusion of genes, and regard cells whose DNA remains unchanged but other substances change as the same type of cells.




Figure. Possible division and fusion situations of protocells, different colour circles represent different molecules:

A. After the cells grow rapidly, they directly rupture and return the molecules to the pool for wrapping new cells or the molecules are directly absorbed by other cells

B. A small number of vesicles produced by cell division do not have the ability to replicate independently. The material distribution after the fusion of two such cells or vesicles can continue to divide.

C. Cells rupture due to excessive growth of macromolecular substances, releasing a large number of vesicles containing partial substances of the whole distribution. These vesicles themselves cannot effectively replicate themselves. After integrating into new cells and releasing the substances they wrap, slow-growing cells may return to normal by supplementing these substances while normal cells may become too fast-growing and continue to rupture to produce these new vesicles again and make it a cycle.

D. Rapid growth and then division to generate two or more cells or vesicles, which can have the ability to continue to grow and divide.

E. One smaller cell mainly contains genetic material, and another cell contains other substances but only a small amount of genetic material. Neither can replicate itself independently. When the smaller cell enters the other cell, it ruptures and fuses into an ordinary complete cell and resume normal physiological activities. And if the cell does

not rupture, a new cooperative working mode may be created, in which a lipid bilayer appears in the inner cell.

An ultimately dominant distribution and corresponding proliferation mode should be the one that replicates itself most quickly and rapidly increases its molecule frequencies in the pool. From the above discussion, it can be seen that both the modern cell-like reproduction method (D) and the virus-like reproduction method (C) may become the dominant form of early life. For the former, cells can achieve continuous division and replication to increase the proportion of the corresponding distribution in pioneer pools, for the latter, when the virus is assembled in the cell, it can achieve the fastest macromolecule conversion speed and replicate a large number of offsprings with fewer individual substances. Specifically, the distribution that normal cells replicate is the distribution of macromolecule synthesis at an appropriate rate, while the distribution that virus-like cells replicate is the distribution after the fusion of the virus-like cell and the infected cell, so the completion of this distribution will rely on the virus's ability to find cells and complete fusion, and ultimately limit the growth of the virus's molecule frequencies. This may indicate that viruses and protocells originated at the same time and have evolved together. Three common hypotheses [19] about the evolutionary origin of viruses can be well explained at the same time through the analysis of this process.

On the other hand, methods similar to sexual reproduction may have been a common reproduction method in protocells (B), which contradicts the common belief that life initially only reproduced asexually [20] and then evolved sexual reproduction as the evolution from prokaryotes to eukaryotes [40]. A reasonable and possible explanation is that all prokaryotes were initially able to reproduce sexually, but the emergence of the bacterial cell wall made it difficult for the two daughter cells to fuse together, thus losing the ability to reproduce sexually. This suggests that if the cell walls of two bacteria are removed, they should be able to fuse and remain active, which is actually called protoplast fusion technology [21]. Similarly, from the discussion of various possible scenarios of cell fusion, eukaryotes may have appeared earlier than thought (E). Smaller cells containing large amounts of genetic material entered cells lacking genetic material and formed primitive nuclei. When this double-layered nuclear membrane can allow some genetic materials and proteins to pass through, cells may resume physiological activities. The small amount of genetic material originally in the cytoplasm may be lost during replication or continuously migrate to the nucleus during long-term evolution, just like the partial genes of chloroplasts or mitochondria in eukaryotic cells have been transferred to the nucleus [39]. The fact that ribosome subunits are assembled in the nucleolus can serve as indirect evidence [22] which implies a fact that the nucleus was originally an independent cell. When the division and replication cycle of the cell nucleus finally coincides with the division and replication cycle of the cell as a whole, we will get the familiar eukaryotic cell. If this hypothesis is true, then the evolution of nuclear envelope is more likely to directly reflect the time when eukaryotes appeared [23] rather than the evolution of cell

membranes. And this kind of similar fusion should appear repeatedly throughout the entire process of cell evolution, such as the generation of organelles mitochondria and chloroplasts [24-25] and constantly shaping the intracellular complex membrane system, and it may slow down when the prokaryotic cell walls and the defense system of multicellular organism [37-38] appeared which make this integration difficult to happen.

Therefore, cells in the pioneer pools continuously communicate and share their molecule frequency distribution through various methods, so that all protocells could have very similar distributions in the early stages of evolution. This extensive communication of molecule frequency distribution was later preserved by life, primarily in the form of sexual reproduction and the communication of genes. If the origin of life is produced by the duplication of a single macromolecule, then in the process of introducing new macromolecules, different evolutionary paths may produce completely different macromolecule frequency distributions, such as completely different ribosomes and tRNA, etc. This contradicts the fact that all modern organisms have almost the same protein translation system and genetic materials. When the cell membrane evolves new permeability by doping with different macromolecules, the optimal distribution within the cell membrane will also change. When a complex membrane system and organelles are formed through the fusion of various vesicles [25], which allows the same cell to have multiple different material distributions, and this will further enable the cells to perform more complex and varied physiological activities. The evolution of cell membranes may also help cells grow at a suitable rate and the proper ratios of growth rates of phospholipids and other macromolecules also contribute to avoid cell membrane rupture, making it easier for protocells to evolve continuously replicating life forms.

DNA is the compression and storage of frequency distributions

If the early division of cells is uneven and unstable, it means that a large number of daughter cells will reduce or even lose their ability to grow due to fluctuations in molecule frequency distribution. Since one DNA strand can transcribe a large amount of RNA and then translate a large amount of protein, which makes DNA equivalent to a kind of compression and storage of the entire distribution within the cell, so that only DNA and a small amount of RNAs and proteins need to be retained to have the opportunity to reconstruct the mother cells' distribution, thus the transmission of this distribution become more efficient, one of the extreme cases is viruses.

But whether DNA appeared in the earliest stages of life is uncertain. It is generally believed that early life used RNA as the only genetic material, and then was slowly replaced by the more stable DNA [26], but the specific time and process lack support from theory and evidence. There is evidence that deoxyribonucleic acid and ribonucleic acid are likely to coexist in the prebiotic Earth [27], if there are a large number of deoxyribonucleic acid molecules in the pioneer pools and the related DNA can be generated or saved through RNA reverse transcription or RNA-DNA hybridization

method, this means the DNAs may copy the RNA distribution. First of all, it is certain that double-stranded DNA cannot always exist in a stable form, because the stable double-helical structure DNA may prematurely lock the distribution of RNA in the early stage of evolution. The distribution of early RNAs would change rapidly in the cycle of stability distillation process, and its distribution at this moment was very wide and did not contain enough information to be used as a template for the genetic material of life. DNA can solve this problem by replicating dynamically. In an idealized geyser, DNA will continue to experience changes in temperature and solution concentration. When the geyser erupts into steam, the double helix DNA will be easier to unwind under the action of higher temperatures and various potential catalytic macromolecules. And some DNA single strands will break and lose the previously saved information in this condition. When the pool gradually dries out and the temperature is partially reduced but still high enough, this makes the polymerization reaction happen. Some RNAs generates new single-stranded DNAs through reverse transcription. Then, they replicate and forms stable double-stranded DNAs under the effect of early polymerase, so that the distribution of DNA will always follow the distribution of RNA. From another perspective, as the DNA and RNA in the pioneer pools are undergoing periodic environmental changes provided by the geyser, it can be considered as they are constantly performing defective PCR and reverse transcription PCR [28-29], which makes the nucleic acid macromolecules have the opportunity to be generated in large quantities. Therefore, DNA at this time can be considered as a real-time map of the distribution of RNA in the pioneer pools.

This means that high-frequency RNA sequences that have stable structures will appear repeatedly in DNA sequences, which include important gene sequences in current cells, such as genes that can be transcribed into tRNA or rRNA, but also contains a large number of stable but non-important RNA sequences accumulated in the pioneer pools. Even though cells may continuously change the overall RNA frequency distribution through parallel calculation and reduce the weight of unimportant RNA sequences, these products may still be continuously generated through the stability distillation process and maintain a certain frequency in the total distribution. This can well explain why there are a large number of non-coding regions in DNA, because RNAs with important roles account for only a small part of the original RNA distribution, and DNA that transcript unimportant RNAs will lose the ability to be transcribed due to the genetic mutations and lack of positive evolutionary pressure. This also shows that protocells may have complex genomes from the beginning of life, rather than gradually becoming more complex from a single gene sequence.




Figure. The distribution of DNA and RNA will be entangled with each other. High-frequency RNA sequences with stable structures will become the main source of the genome of primitive cells.

RNA and protein will affect the frequency distribution of DNA and its transcription, and DNA will also affect their respective distributions through the central dogma [30]. This makes the distributions of proteins, RNAs and DNAs entangled with each other. During the evolution process, the distribution of all substances will be changed simultaneously and a suitable solution will eventually be obtained. The huge impact of DNA transcription on RNA distribution also makes the RNA distribution in cells no longer rely solely on the stability of its configuration but more on the expression ability of the corresponding genes. And through the central dogma, the cell's search for protein configuration becomes much broader and easier.

On the other hand, if RNA is the only genetic material in protocells, you first need to solve the problem of unequal replication probabilities of different RNA sequences, which means that some RNA sequences with specific structures may be more difficult to replicate themselves [31], so that the RNA distribution after cell division may continue to shift and will never be stable. Secondly, when the original cells containing only RNA genetic material leave the original solution pool and enter the ocean, a large amount of RNA will quickly become stable due to the drop in temperature, and it will be more difficult to replicate itself. For example, tRNA and rRNA will find it difficult to copy itself under such conditions. Although dormant cells can theoretically be reactivated in seafloor hot springs and replace RNA with DNA as their new genetic material, the transcription of DNA will have a huge impact on the distribution of RNA already present in the cell, which will cause the cells no longer be able to function normally and will reenter dormant probably. Since it is difficult to gather a high concentration of macromolecules or a large number of cells in seawater, dormant cells cannot restore their functions by fusing new substances or other cells like they did in

terrestrial hot springs, and the process of life evolution will be interrupted. In other words, RNA played a central role in the beginning of evolution, but before life left the pioneer pools, DNA should have become the main genetic material in cells, rather than DNA replacing RNA gradually during evolution.

How to test this hypothesis

There are three possible experiments to test this hypothesis:

1. Simulating the dry-wet cycle and provide sufficient ribonucleotides. We should observe the frequency distribution changes of different configurations of RNA, and the more extreme the conditions are set, the more cycles there are, the narrower the distribution of RNA is likely to be. When more substances are added to the process, the changes in RNA frequency distribution will become more complex.
2. After establishing a sufficiently complex metabolic reaction network, let the phospholipids randomly encapsulate substances through stirring and wet-dry cycles and observe the growth and rupture of these vesicles, as well as the changes in the growth rate after the fusion of various vesicles.
3. Under complex and extreme conditions, perform PCR and reverse transcription PCR at the same time, and observe that the DNA distribution is dynamically tracking and coping the RNA distribution.

The stability distillation process can also be simulated by computer. First, a neural network needs to be pre-trained to estimate the stability and average existence time of molecules. At the same time, you need to train a new neural network to evaluate the collision between different molecules and the formation of new ones. In this way, these two neural networks can be used to simulate the intracellular metabolic reactions. All the molecules present will collectively determine the conduct of the new reaction and the stability of the products, while the newly formed macromolecules will act as potential catalysts for all known reactions and in turn affect the stability of all molecules. In this way, the frequency distribution of molecules in the system can be repeatedly updated in a manner similar to back propagation, and the new product is equivalent to an error, which will feedback to each layer in the system at the same time, update the weights, and then forward calculate all reactions to determine the change in formation of all products from each layer. As the stability of the product will determine the progress of the reaction, allowing simulations to calculate base pairing and nucleic acid self-replication. If the simulation of the entire origin of life process can be realized, then by adjusting the cycle period, temperature, pH value, types of organic matter, etc., it is likely that different protein translation systems or even gene systems can be obtained, such as a new configuration of ribosomes and its matching tRNA groups.




Figure. Simulate the stability distillation process through a method similar to back propagation in neural networks.

Discussion

In conclusion, the origin of life can be considered as a process of accumulating information, amplifying the stability differences of different macromolecules through extreme environments, and further narrowing the distribution of macromolecules in their generation paths with the help of catalysts and membrane systems. Life is a specific distribution of molecules that that can self-assembly, replicate itself, process molecules transformation, shift, fuse, and restore. The high stability difference and specificity generated by base pairing makes it possible to self-replicate nucleic acid macromolecules, and further improves the information content in the production process of protein and DNA by establishing the correspondence between amino acids and tRNA and the reverse transcription of RNA to DNA respectively. Life itself can be considered as a continuous conversion process of molecule frequency distribution, so the strength of vitality can be measured by this distribution conversion ability. For example, dormant seeds or prions have very weak vitality because they can only convert a small amount of substance or a single type of substance.

From the perspective of the direction of evolution, for macromolecules themselves, their evolutionary direction can be considered to have a higher frequency distribution, which usually requires high stability or high stability of their precursors, or the ability to self-catalytic and replicate, or the ability to join a cross-catalytic network. When the macromolecules in cells are constantly transforming different molecules, its evolutionary direction will deviate to catalyze the production of macromolecules, especially macromolecules and molecules that can widely catalyze or help the synthesis of a range of macromolecules, such as ribosomes, RNA ligases, and DNA polymerases, etc., and self-replication can be seen as a special case of it. The establishment of the central dogma expands the search for various catalytic macromolecules, allowing cells to find more catalytic macromolecules without requiring them to have high stability.

From the angle of cells, when the cell membrane evolves appropriately to expel proteins that are useless to the cell, it means that different cells have the possibility of rapid information exchange and can change the extracellular environment through massive aggregation to gain possible survival advantages, such as defense predation of other cells. On the other hand, the emergence of membrane systems and chromosomes also allows cells to store multiple distributions at the same time which means they may restore different distributions or utilize the properties of different distributions to perform new behaviors, such as cell differentiation. Together, they form the cornerstones of multicellular organisms. If the objective function of a single cell is to survive and replicate itself, then the emergence of multicellular organisms has further deviated from the objective function of one cell. Because in the multicellular organism, the replication of a single cell will depend on the overall survival and reproduction, which can be understood as the single cell establishing a causal chain with other cells, that is, the replication of A cell depends on B cell, and the replication of B cell depends on C cell, etc. The establishment of this causal relationship is ensured through the exchange of information between each other, such as communication through cell membrane proteins or extracellular membrane vesicles [34-36], so when cells mutate into cancer cells and lose the ability to communicate, it represents the collapse of this causal relationship. The cancer cell objective function backtracks to a function that targets individual survival and replication.

From the angle of organisms and populations, different external environmental changes will favor different intracellular molecule distributions. Due to the leading role of DNA in distribution inheritance and reconstruction, it will eventually be reflected in the filtering and selection of the population gene pool, which is manifested as natural selection. Therefore, more complex and diverse genomes may become a possible evolutionary direction, but the growth rate of its diversity will vary due to the intensity of environmental changes and different reproduction methods. When the genetic material within the cell remains unchanged, it is more beneficial to find a suitable environment to convert substances and reproduce itself. This essentially reflects the species' choice of the environment, so an important evolutionary direction of organisms is the ability to move and migrate, and may have different manifestations at different stages of the life cycle. Therefore, a more complete theory of natural selection should be understood as a process of mutual selection and change of nature and life.

REFERENCES

- [1] Joyce, Gerald F., and Jack W. Szostak. "Protocells and RNA self-replication." *Cold Spring Harbor Perspectives in Biology* 10.9 (2018): a034801.
- [2] Ma, Wentao. "What does "the RNA world" mean to "the origin of life"?" *Life* 7.4 (2017): 49.
- [3] Wu, Kun, Qiong Nan, and Tianqi Wu. "Philosophical analysis of the meaning and nature of entropy and negative entropy theories." *Complexity* 2020 (2020).
- [4] Nowakowski, Jacek, and Ignacio Tinoco Jr. "RNA structure and stability." *Seminars in virology*. Vol. 8. No. 3. Academic Press, 1997.

- [5] Becker, Sidney, et al. "Wet-dry cycles enable the parallel origin of canonical and non-canonical nucleosides by continuous synthesis." *Nature communications* 9.1 (2018): 163.
- [6] Hassenkam, Tue, and David Deamer. "Visualizing RNA polymers produced by hot wet-dry cycling." *Scientific reports* 12.1 (2022): 10098.
- [7] Hassenkam, Tue, et al. "AFM images of viroid-sized rings that self-assemble from mononucleotides through wet-dry cycling: Implications for the origin of life." *Life* 10.12 (2020): 321.
- [8] Serra, Martin J., et al. "RNA hairpin loop stability depends on closing base pair." *Nucleic acids research* 21.16 (1993): 3845-3849.
- [9] Vecenie, Christopher J., and Martin J. Serra. "Stability of RNA hairpin loops closed by AU base pairs." *Biochemistry* 43.37 (2004): 11813-11817.
- [10] Xu, Xiaojun, and Shi-Jie Chen. "A method to predict the structure and stability of RNA/RNA complexes." *RNA structure determination: methods and protocols* (2016): 63-72.
- [11] Bikard, David, et al. "Folded DNA in action: hairpin formation and biological functions in prokaryotes." *Microbiology and Molecular Biology Reviews* 74.4 (2010): 570-588.
- [12] Pal, Arumay, and Yaakov Levy. "Structure, stability and specificity of the binding of ssDNA and ssRNA with proteins." *PLoS computational biology* 15.4 (2019): e1006768.
- [13] Kumar, Sandeep, Chung-Jung Tsai, and Ruth Nussinov. "Factors enhancing protein thermostability." *Protein engineering* 13.3 (2000): 179-191.
- [14] Root-Bernstein, Robert, and Meredith Root-Bernstein. "The ribosome as a missing link in prebiotic evolution II: Ribosomes encode ribosomal proteins that bind to common regions of their own mRNAs and rRNAs." *Journal of Theoretical Biology* 397 (2016): 115-127.
- [15] Di Giulio, Massimo. "The origin of the tRNA molecule: Independent data favor a specific model of its evolution." *Biochimie* 94.7 (2012): 1464-1466.
- [16] Joshi, Manesh Prakash, et al. "Formation and stability of prebiotically relevant vesicular systems in terrestrial geothermal environments." *Life* 7.4 (2017): 51.
- [17] Damer, Bruce, and David Deamer. "The hot spring hypothesis for an origin of life." *Astrobiology* 20.4 (2020): 429-452.
- [18] Howes, Philip, et al. "Phospholipid encapsulated semiconducting polymer nanoparticles: their use in cell imaging and protein attachment." *Journal of the American Chemical Society* 132.11 (2010): 3989-3996.
- [19] Forterre, Patrick. "The origin of viruses and their possible roles in major evolutionary transitions." *Virus research* 117.1 (2006): 5-16.
- [20] Heitman, Joseph. "Evolution of sexual reproduction: a view from the fungal kingdom supports an evolutionary epoch with sex before sexes." *Fungal Biology Reviews* 29.3-4 (2015): 108-117.
- [21] Gokhale, D. V., U. S. Puntambekar, and D. N. Deobagkar. "Protoplast fusion: a tool for intergeneric gene transfer in bacteria." *Biotechnology advances* 11.2 (1993): 199-217.
- [22] Johnson, Arlen W., Elsebet Lund, and James Dahlberg. "Nuclear export of ribosomal subunits." *Trends in biochemical sciences* 27.11 (2002): 580-585.
- [23] Mans, Berend, et al. "Comparative genomics, evolution and origins of the nuclear envelope and nuclear pore complex." *Cell cycle* 3.12 (2004): 1625-1650.
- [24] Archibald, John M. "Endosymbiosis and eukaryotic cell evolution." *Current Biology* 25.19 (2015): R911-R921.

- [25] Gould, Sven B., Sriram G. Garg, and William F. Martin. "Bacterial vesicle secretion and the evolutionary origin of the eukaryotic endomembrane system." *Trends in microbiology* 24.7 (2016): 525-534.
- [26] Ribas de Pouplana, Lluís, et al. "Origin and evolution of DNA and DNA replication machineries." *The genetic code and the origin of life* (2004): 145-168.
- [27] Ribas de Pouplana, Lluís, et al. "Origin and evolution of DNA and DNA replication machineries." *The genetic code and the origin of life* (2004): 145-168.
- [28] Lauerman, Lloyd H. "Advances in PCR technology." *Animal health research reviews* 5.2 (2004): 247-248.
- [29] Bachman, Julia. "Reverse-transcription PCR (rt-PCR)." *Methods in enzymology*. Vol. 530. Academic Press, 2013. 67-74.
- [30] Shapiro, James A. "Revisiting the central dogma in the 21st century." *Annals of the New York Academy of Sciences* 1178.1 (2009): 6-28.
- [31] Repass, John F., and Shinji Makino. "Importance of the positive-strand RNA secondary structure of a murine coronavirus defective interfering RNA internal replication signal in positive-strand RNA synthesis." *Journal of virology* 72.10 (1998): 7926-7933.
- [32] Neveu, Marc, Hyo-Joong Kim, and Steven A. Benner. "The "strong" RNA world hypothesis: Fifty years old." *Astrobiology* 13.4 (2013): 391-403.
- [33] Alberts, Bruce, et al. "Catalysis and the Use of Energy by Cells." *Molecular Biology of the Cell*. 4th edition. Garland Science, 2002.
- [34] Zhou, Zhihang, et al. "The versatile roles of tetraspanins in cancer from intracellular signaling to cell–cell communication: cell membrane proteins without ligands." *Cell & Bioscience* 13.1 (2023): 59.
- [35] Turturici, Giuseppina, et al. "Extracellular membrane vesicles as a mechanism of cell-to-cell communication: advantages and disadvantages." *American Journal of Physiology-Cell Physiology* 306.7 (2014): C621-C633.
- [36] Ratajczak, Janina, et al. "Membrane-derived microvesicles: important and underappreciated mediators of cell-to-cell communication." *Leukemia* 20.9 (2006): 1487-1495.
- [37] Guryanova, Svetlana V., and Tatiana V. Ovchinnikova. "Innate immunity mechanisms in marine multicellular organisms." *Marine Drugs* 20.9 (2022): 549.
- [38] Medzhitov, Ruslan, and Charles A. Janeway. "Innate immunity: the virtues of a nonclonal system of recognition." *Cell* 91.3 (1997): 295-298.
- [39] Martin, William, and Reinhold G. Herrmann. "Gene transfer from organelles to the nucleus: how much, what happens, and why?." *Plant physiology* 118.1 (1998): 9-17.
- [40] Speijer, Dave. "What can we infer about the origin of sex in early eukaryotes?." *Philosophical Transactions of the Royal Society B: Biological Sciences* 371.1706 (2016): 20150530.
- [41] Gonsior, Michael, et al. "Yellowstone hot springs are organic chemodiversity hot spots." *Scientific reports* 8.1 (2018): 14155.