

Stability-distillation hypothesis for the origin of life

Cheng Bi

Email: cb236@foxmail.com

Abstract

Through in-depth thinking and reasoning about the conditions required for cells to maintain unchanged material distribution, it is concluded that life metabolic reactions require high information content. However, the self-replication of a few macromolecules such as RNA and DNA can only retain information, but cannot generate information. When thinking about the possible ways in which this information is generated, one possible way is to maintain the distribution of substances in the cell through selective permeability and active transport of membranes. This mechanism still plays an important role in today's cells but it is difficult to generate enough information. Another possible way is to generate information through stability-distillation, which continuously changes the molecular frequency distribution in the system through periodic changes in the environment and generates information from a consequentialist perspective. Unlike the RNA world hypothesis, under this hypothesis, life does not need to originate from a few self-replicating macromolecules, making the entire hypothesis more reasonable.

On the other hand, the structure of cells makes it easier for life to accumulate macromolecules and select molecules that can widely catalyze macromolecule production reactions and use energy forms that are conducive to production reactions. By further deducing the entire process of the origin of life, it is concluded by contradiction that when life originated, RNA, DNA, proteins, lipids and other substances all existed. The dry-wet cycle of hot springs produces a mechanism similar to PCR and reverse transcription PCR, which allows RNA and DNA to proliferate rapidly. In this process, DNA can continuously track the distribution of RNA in the pool and continuously retain the original information in RNA into the original genome. This gives an important inference that the genome of the original cell will have considerable complexity and insert a large number of repetitive RNA sequences with stable configurations. With the breakage, fusion and recombination of a large number of original cells in the hot spring, life will have two parallel evolutionary routes of viruses and cells, which can be regarded as two sides of life. At the end of the hypothesis, I discussed the possible reasons for the emergence of multicellular life and the conditions required for the birth of complex life and intelligent life.

Molecular frequency distribution and information entropy

When considering the macromolecules and cells in the origin of life, any macromolecule or cell has not yet evolved the ability to replicate itself. At this time, various chemical reactions may occur and generate any possible macromolecule. In this case, life still needs a reliable reaction pathway rather than a chaotic reaction. Macromolecules generated by random chemical reactions are difficult to become reliable life molecules. This is because after the self-organizing form of life is formed, the difference in molecular frequency distribution between its parents and offspring must be maintained within a certain range. The molecules generated by chance will quickly reduce their frequency distribution in the offspring as cells divide or mix with other molecules, and eventually lose their life functions. This is also the reason why people believe that life originated from large molecules that could replicate themselves in the prebiotic Earth [1-2]. However, for such macromolecules to appear on the primitive Earth, sufficient and pure substrates are also essential, which makes the entire hypothesis more and more bloated. In fact, life does not need to originate from a few self-replicating macromolecules.

From a mathematical point of view, a reliable reaction pathway means that the generation of a macromolecular depends on the accumulation of its lower-level substrate. For example, if RNA can be produced in prebiotic Earth, a reasonable conclusion is that sufficient amounts of ribonucleotides has accumulated in the environment. To generate ribonucleotides, people would expect that a large amount of ribose, phosphate and nucleobase or related intermediate products have accumulated in the environment rather than thinking that carbon, hydrogen, oxygen, nitrogen, phosphorus elements or other simple molecules can directly generate ribonucleotides or even RNA polymers through molecules' collisions. This is because the probability of two molecules combining to form a new product after collision in space is much greater than the probability of multiple molecules colliding at the same time and the same location. Theoretically, the combinations of compounds that can be formed by these chemical elements are infinite. When certain substrates are required to generate specific macromolecules, it also means that one or several combinations of these elements need to have a higher frequency than other random sequences. Increasing the frequency of one or more sequences in a variety of random combinations corresponds to a change in information. That is to say, compared to a random system, chemical reactions related to life are more likely to increase the information content of organism or reduce the information entropy. This is especially obvious in the polymerization reaction of monomers. When the polymer chain continues to grow in natural environments, its sequence and length are usually random, and you will end up with a mixture of macromolecules with different sequences and lengths, while the distribution of macromolecules produced by specific reactions within cells is usually much narrower.

Therefore, life can be seen as a conversion process of molecular frequency distribution

accompanied by an increase in information content. Assuming that there are several small molecule substances {A, B, C, D...} in the environment, the possible sequence combinations that can be formed include {A-B-C, B-C-D, C-C-B-B-A, D-D, A-A-B-B-D...}, the original frequency distribution is close to random {p1, p2, p3, p4, p5...}, under specific reaction conditions and catalyst, the frequency distribution of the products group becomes {q1, q2, q3, q4, q5}. Relative to randomly generated products, the information entropy change formula is:

$$\Delta Entropy = - \sum_i q(i) \log_2 q(i) - \left(\sum_i p(i) \log_2 p(i) \right)$$

Under the effect of suitable reaction conditions and catalysts, the distribution of the reaction products will become narrower, that is, the entropy value will be lower. This explains from another perspective why life feeds on negative entropy [3].

Therefore, the question becomes how to make a chemical reaction involving multiple reactants and potentially generating macromolecules of any sequence have a higher information content. This is not a problem for most simple reactions because the reactants and products are usually fixed, for example, the reaction between hydrogen and oxygen to form water. But for chemical reactions of polymers or mixed reactants that may interfere with each other, the reactions may be not converged enough, for example, in successive condensation reactions of multiple amino acids, peptides of any sequence may generate. This makes the frequency distribution of any specific sequence of macromolecules extremely low and cannot have a reliable generation path which means the whole process is not repeatable. Although suitable catalysts can increase the information content of the entire reactions by allowing substrates with a specific sequence to react or products to decompose from a specific sequence, these catalytic molecules themselves may also be large molecules, thus facing the same problem unless they happen to be able to catalyze themselves. This will make related theoretical hypotheses inevitably fall into a chicken-and-egg paradox.

However, to increase the information content of a complex reaction, a catalyst or single substrate is not necessarily needed. A result-oriented approach is to continuously remove non-target macromolecules from the product, just like repeated purification during a distillation process. There are two possible methods: One is to construct a suitable membrane system that continuously removes by-products while still preserving reaction substrates and target products, thereby increasing the information content of the products within the membrane. This will strongly prompt the cell membrane to evolve the ability to actively transport various molecules which is very common in today's cells. The other method is to continuously decompose by-products into original substrates and smaller molecules, and only retain few target macromolecules in each step. For the former, this can be easily associated with the complex membrane system in present cells which still plays an important role, while for the latter, macromolecules of different sequences, such as RNA, have different stabilities [4] and the extreme environment may help to amplify the stability differences, thereby narrowing the

frequency distribution of products. In this way, through repeated decomposition and synthesis, macromolecules with higher stability can be continuously accumulated, forming a reliable chemical reaction path from a result perspective. Coincidentally, the existence of hot springs with wet-dry cycles in the early Earth could provide both extreme reaction conditions and periodic environmental changes which is an ideal place for the origin of life.


On the other hand, if we assume that life originated from the self-replication of a few special macromolecules, this would require that the prebiotic Earth was filled with a large number of substrates that could support the self-replication of these macromolecules. For example, the RNA world hypothesis requires that the environment be filled with nucleotides [32]. However, this life form constructed by the self-replication of a few macromolecules is unlikely to have sufficient complexity to cope with the extreme environment of the prebiotic Earth. Even if there is a process that makes the simple self-replication of macromolecules gradually more complex, the introduction of any new molecule may interfere with or even completely interrupt the self-replication process of the original macromolecule. The entire hypothesis will be highly limited by idealized environmental assumptions, and will need to explain every step of how life evolved from a few macromolecules to complex reaction networks.

Cross-catalytic networks and the stability-distillation hypothesis

It is known that in certain hot spring environments with dry-wet cycles, nucleotides will naturally form [5] and polymerize into RNA macromolecules [6]. These macromolecules can different folding structures, such as chainlike RNA, ring-shaped RNA [7], hairpin RNA, etc., and have different stabilities. In the prebiotic Earth environment, these macromolecules may have to face high temperatures, acidic environments and collisions with various other macromolecules, causing a large number of long-chain RNA molecules produced in dry environment become unstable and re-decompose into smaller RNA sequences or even ribonucleotides in the wet environment. Some RNA sequences have higher stability due to their folded structures or complexes formed with other macromolecules such as RNA-protein complexes, making them more difficult to decompose and more likely to collide with other molecules. When this wet-dry cycle is repeated, RNA molecules with stable configurations will accumulate and the whole RNA distribution will narrow. This is essentially creating information, and is determined by factors such as the stability differences between macromolecules in the prebiotic earth environment and the number of times the cycle occurs. When enough macromolecules with similar sequences or even completely identical sequences are accumulated, this will make possible the continuous and deterministic generation of certain macromolecules of life.

When considering only the stability of RNA and RNA-protein complexes, the number of paired bases, related sequences, and bound proteins all affect the stability of RNA configuration [8-10]. RNA and RNA-protein complexes with multiple complementary

sequences are more likely to form stable structures with lower energy, which makes it possible that RNA molecules with stem-loop structures or other similar structures randomly generated in repeated polymerization and decomposition reactions have a higher proportion in the pre-biotic Earth due to their higher stability. The size of the loop in the hairpin RNA, the number of paired bases, pH, high temperature, and collisions with other macromolecules may all affect the stability of RNA and change the frequency distribution of RNA in the wet-dry cycle. These accumulated RNAs with stable configurations have more opportunities to generate larger RNA macromolecules in the cycle, and because the new RNA macromolecules may contain more complementary sequences, the reaction is energetically favorable. For example, when two stable RNA molecules with the same sequence and stem-loop structure are spliced together, a molecule highly similar to tRNA can be produced, and it can be predicted that the tRNA molecule will exhibit central symmetry. When more tRNA, stem-loop RNA and other small RNAs are assembled together, they can form a structure similar to ribosomal RNA, and stabilize the structure by filling proteins, eventually forming a more stable RNA-protein complex. The original ribosome should be included in these complexes. It can be inferred that these complexes have high stability in extreme environments, but only a few have strong catalytic ability like ribosomes. Even if this early ribosome has the ability to catalyze all its own substrates to increase its own generation, this effect may be significantly weakened when all small molecules, macromolecules, inorganic and organic substances are mixed together. This requires life to have more mechanisms to screen out these macromolecules with strong catalytic ability.






Figure. Figure. (A) Green circles represent amino acids and proteins, and blue circles represent small molecules such as RNA and nucleotides. During the dry-wet cycle, unstable RNA will continue to decompose, and the frequency of RNA molecules with stable structures will increase. The continuous accumulation of related substrates increases the frequency of stable macromolecules, while RNA molecules with complementary sequences such as proteins and amino acids help stabilize their structure. (B) Specifically, when two hairpin-shaped RNA molecules with complementary sequences are spliced together, different tRNA-like RNAs may be formed and exhibit central symmetry at the same time. Factors such as the loop sequence of the hairpin loop and the closing base pair will affect its stability [8] and indirectly affect the frequency distribution of different tRNAs. On the other hand, the interaction between the original tRNA and other molecules in the environment, such as binding to specific amino acids, will also affect its stability and change its final distribution.

When considering the entire cross-catalytic network, the newly generated macromolecules will interact with all other molecules in the system and become potential catalysts for all existing reactions, and the existing molecules in the network will also affect the stability of the newly generated macromolecules. In fact, this is very similar to a neural network with feedforward and feedback calculations. Whether the molecules in each layer of the network can maintain sufficient frequency determines whether the macromolecules in the next layer can be continuously generated. Therefore, when macromolecules in the system decompose, it is necessary to retain certain information. For example, when a ribosome dissociates into two stable subunits, the information is well preserved because the two subunits can quickly reassemble into a ribosome under appropriate conditions. When macromolecules such as DNA, RNA, and proteins decompose into nucleotides and amino acids, this information is still retained to some extent until these monomers further decompose into water and carbon dioxide, etc., and the information is drastically lost in the process. This mechanism also means that macromolecules with chain structures will appear more frequently in the network. Catalysts not only increase the information of the network, but also retain information because of their own invariance before and after the reaction. The

specificity of catalysts allows them to direct the reaction to yield a particular product in the mixture, thereby directly increasing the information and accelerating the calculations of the entire network. Decomposition reactions and synthesis reactions dominated by catalysts are equally important. The realization of this stability difference depends on the fact that most sequence combinations can be decomposed. Otherwise, the information content in the metabolic pathway will be insufficient, and too many useless macromolecules will accumulate in the network and occupy the substrates of other reactions or interfere with other reactions. When out of the extreme environment of the original earth, more intermediate macromolecules will become stable macromolecules and cause the cross-catalytic network to collapse, which requires more efficient enzymes to decompose these macromolecules in large quantities and help cells adapt to the new environment, which can explain why RNase and protease are so common now in cells. Therefore, the emergence of various enzymes can be considered to expand the boundaries of life [33], and the evolution of enzymes is crucial to the evolution of life.

Why RNA generated the original information?

Common macromolecules in life include RNA, DNA and proteins. Compared with other macromolecules, RNA and its complexes have great differences in stability in the extreme environment of the primitive earth. Base pairing in RNA can be regarded as a configuration change with great differences in stability. The stability of the structure formed by correct pairing is much greater than the stability of the structure combined with other incorrect bases or any macromolecules. This means that RNA with complementary sequences is more likely to form a stable configuration and further generate larger stable RNA macromolecules. This bias in the distribution of RNA sequences has become the original source of information for life. For DNA, its single strand lacks a stable folding configuration [11-12], while the stability difference between double-stranded DNA with higher stability is small, which makes it impossible for DNA itself to generate enough information, but it can amplify information through semi-conservative replication, but it does not completely exclude that some short sequences of DNA also contribute part of the original information. For another macromolecule protein that constitutes life, since the types of amino acids that constitute proteins are usually far more than four, the probability of generating a peptide chain of a specific sequence through random decomposition and polymerization becomes extremely low, so it takes much longer to accumulate proteins of a specific sequence, making the entire process almost impossible. This can also explain why RNA has only four main bases instead of six or more. On the other hand, the stability of peptide and protein configurations is determined by a variety of factors [13], so proteins with similar amino acid sequences may have completely different configurations and stabilities, making the entire stability-distillation process unable to converge and difficult to generate large amounts of macromolecular proteins.

Therefore, macromolecules with both stability convergence and stability differences can produce life capable of self-replication. The stability convergence of double-

stranded DNA itself and the stability difference caused by base pairing help it eventually evolve to self-replicate, but the stability differences of DNA of different sequences are small and cannot produce enough original information. These primitive genes may come from differences in the stability of RNA and RNA-protein complexes in extreme environments. Biomolecules commonly found in modern cells, such as double-stranded DNA, tRNA, ribosome subunits, etc., all have high stability, which can be considered to be the result of macromolecule selection and evolution.

This also means that proteins need to establish a connection with RNA to increase the information content of the reaction and become reliable macromolecules of life. The most direct way is that the combination of specific RNA and some proteins can produce highly stable complexes, thereby affecting the frequency distribution of all proteins, but a large number of complex proteins cannot be generated through random reactions, that is, RNA needs to be able to catalyze and regulate the protein generation process. This evolutionary process is difficult to obtain through simple reasoning, but a possible route is that some RNA or RNA-protein complexes first evolved the ability to catalyze the combination of two amino acids to form a peptide chain. Since the original information in the system is concentrated in the RNA sequence at this time, only reactions that can use this information can generate a large number of identical proteins. Through the difference in stability after the original tRNA combines with different amino acids, the early translation system eventually evolved. This also means that the encoding of the main proteins that existed in the early days of life will be highly dependent on a large number of stable and repetitive RNA sequences in the system. In other words, RNAs containing a large number of complementary sequences, such as hairpin RNA, tRNA, and rRNA, may encode a large number of important proteins, including proteins that constitute ribosomes in the early stages of life [14].


The pioneer pools, cells and shared distribution

Based on the above discussion, life most likely originated in one or several adjacent hot springs with wet-dry cycles, especially geyser that could erupt jets of water at a certain period. I would like to call these hot springs where life was born pioneer pools.

It is known that phospholipids [16] and other macromolecules essential for life [5, 6, 27] can form spontaneously in hot springs. Through other mechanisms such as wet-dry cycles [17] or the disturbance of water flow during geyser eruptions [18], these lipids may self-assemble into lipid bilayers and then randomly wrap molecules in the pool to become primitive cells. When the monomers of macromolecules can pass through the cell membrane more easily than their corresponding macromolecules, such as small amino acids that can enter cells more easily than proteins, the generation of new macromolecules inside the cell will increase the osmotic pressure, thereby driving small molecules into the cell and accumulating macromolecules inside the cell. Regardless of whether the cell divides normally or breaks and returns all the molecules to the hot spring, the corresponding macromolecules in the pool will still increase and re-enter the cycle with the formation of new cells or primitive predation behavior. In other words,

cells do not need to have evolved perfect division and replication mechanism at the origin of life, but more perfect division and replication can allow the corresponding molecules to increase their weight in the hot spring more quickly, thereby making this evolution path converge. In this way, the cells in the entire hot spring have similar molecular frequency distributions and form a parallel calculator for this distribution, which eventually converges to the calculation of genes. The key to the establishment of this calculation depends on two prerequisites. First, the cell membrane evolves appropriate selective permeability, so that the cell can accumulate macromolecules faster and replicate itself quickly by taking in appropriate smaller molecules and excreting waste. Second, there is sufficient communication between cells, because sufficient communication is the premise for the stability-distillation process to be established, so that the macromolecules in all cells as a whole can converge in evolution. In other words, primitive cells need to be easier to fuse or engulf each other than modern cells.

Specifically, the hot spring is likely to be a geyser that erupts periodically. In addition to being able to more easily maintain wet-dry cycles, the eruption of geysers also continuously brings in organic and inorganic substances [41]. Since primitive life is unlikely to be able to directly convert large amounts of inorganic substances into organic substances, replenishing the various life materials consumed by the generation of macromolecules and cell replication in the hot spring is crucial for the evolution of



life.

Figure. The most likely place for the origin of life is a geyser that erupts with a suitable period.




Figure. Possible division and fusion situations of protocells, different color circles represent different molecules:

A. After the cells grow rapidly, they directly rupture and return the molecules to the pool for wrapping new cells or the molecules are directly absorbed by other cells

B. A small number of vesicles produced by cell division do not have the ability to replicate independently. The material distribution after the fusion of two such cells or vesicles can continue to divide.

C. Cells rupture due to excessive growth of macromolecular substances, releasing a large number of vesicles containing partial substances of the whole distribution. These vesicles themselves cannot effectively replicate themselves. After integrating into new cells and releasing the substances they wrap, slow-growing cells may return to normal by supplementing these substances while normal cells may become too fast-growing and continue to rupture to produce these new vesicles again and make it a cycle.

D. Rapid growth and then division to generate two or more cells or vesicles, which can have the ability to continue to grow and divide.

E. One smaller cell mainly contains genetic material, and another cell contains other substances but only a small amount of genetic material. Neither can replicate itself independently. When the smaller cell enters the other cell, it ruptures and fuses into an ordinary complete cell and resume normal physiological activities. And if the cell does

not rupture, a new cooperative working mode may be created, in which a lipid bilayer appears in the inner cell.

Using cells as units, any group of molecules in the pioneer pools can be isolated and calculated independently. Through the continuous rupture and fusion of cells and the recombination of various molecules in the pioneer pools, life constantly searches for the optimal molecular frequency distribution in this process, and finally realizes a metabolic network that can replicate itself the fastest in this environment. This division and recombination may be the recombination of substances and cells, such as (A) in the picture, which may finally evolve into cell predation behavior, or it may be the recombination between cells, such as (B), the substances in the cell are completely released and recombined which is very similar to sexual reproduction. This seems to contradict the common view that life initially only reproduced asexually [20], and sexual reproduction did not appear until eukaryotic organisms evolved [40]. One possible explanation is that all prokaryotes were originally able to reproduce sexually, but the emergence of cell walls and the evolution of cell membranes made it difficult for the two daughter cells to fuse and recombine all the molecules in their cells. This also shows that if the cell walls of two bacteria are removed, they may be able to fuse and remain active, which is actually called protoplast fusion technology [21]. On the other hand, the exchange of substances such as DNA between bacteria still exists [37].

When cells grow and reproduce at a suitable rate, in order to achieve continuous and stable reproduction, the molecular frequency distribution of parents and offspring will tend to converge to the same, and eventually evolve into asexual reproduction in (D), which requires the cell membrane to evolve suitable selective permeability so that cells can grow at an appropriate rate in the environment. However, when the growth rate of macromolecules in cells is too fast and causes the cell membrane to rupture, the macromolecules that lose the suitable environment will likely lose the ability to replicate and grow. When some macromolecules or vesicles fuse into new cells again, the frequency distribution of newly generated molecules may restore the ability to grow rapidly and form a reproduction method similar to modern viruses. The replication rate of macromolecules that make up viruses in cells will be much greater than the total synthesis rate of macromolecules in normal cells, and the replication rate of these macromolecules after leaving cells will be much slower than that of cells, which puts viruses and cells in a strong competitive relationship. Using fewer macromolecules to form themselves, thereby producing more copies and stronger infection ability, naturally becomes the evolutionary direction of viruses. This also provides a reasonable explanation for the evolution of cell walls in prokaryotes: the high frequency of fusion and recombination between cells at the origin of life would strongly induce the generation and evolution of viruses, which means that the replication of some normal cells would become more dependent on the fused host and evolve to use fewer molecules to compose themselves, while the emergence of cell walls made sexual reproduction of prokaryotes difficult. For example, Chlamydia [38] can be considered to be in the process of evolving from bacteria to viruses. Eukaryotic cells may avoid

this evolutionary direction by producing reproductive cells with lower frequency but higher genetic accuracy or through other mechanisms such as evolving an effective immune system. This shows that viruses and cells may have originated and evolved together, and the reasoning and deduction of this process can satisfy the three main hypotheses related to the origin of viruses at the same time [19].

When a smaller cell is engulfed by a larger cell and still maintains its cell structure, the appropriate selective permeability of the cell membrane makes it possible for the small cell to maintain normal physiological activities. In other words, the evolution of the cell membrane may be the key to the establishment of the endosymbiotic theory [24]. This high frequency of fusion and engulfment between cells at the origin of life helped cells evolve organelles such as chloroplasts and mitochondria [24-25] and complex membrane systems. Just as some genes of chloroplasts or mitochondria in eukaryotic cells were transferred to the nucleus during evolution [39], the genetic material of larger cells may also be transferred to the nucleus or lost during replication. In other words, the nucleus of a eukaryotic cell may have originally been an independent cell with its own ribosomes and metabolic network. The evolution of the nuclear membrane allowed genetic material and some proteins to frequently enter and exit the nucleus, thus enabling small cells to acquire the ability to influence and regulate the activities of large cells. The fact that ribosomal subunits are assembled in the nucleolus can be seen as an indirect evidence [22]. From this perspective, the evolution of the nuclear membrane can be used to better estimate the time of the emergence of eukaryotes [23], and the emergence of eukaryotes should be the result of long-term evolution rather than accidental, which means that during the evolution process, there may be a large number of cells with multiple potential nuclei at the same time.

Therefore, the cells in the pioneer pool continuously communicate and share their molecular frequency distributions in various ways, so that in the early stages of evolution, all primitive cells can have very similar distributions. The sexual reproduction of modern organisms and the genetic exchange between bacteria can be regarded as the continuation of this behavior. As cells replicate, the distribution frequency of stable macromolecules with strong catalytic ability will continue to increase, thereby reducing the distribution frequency of stable but weak catalytic macromolecules produced by the stability-distillation process. If the origin of life is produced by the self-replication of a few macromolecules, then in the process of introducing new macromolecules and continuous complexity, different evolutionary paths may produce completely different combinations of life macromolecules, such as completely different translation systems composed of ribosomes and tRNA. This contradicts the fact that all modern organisms have almost the same protein translation system and genetic material.

When did DNA appear?

A very important question is whether DNA existed at the origin of life and functioned as genetic material, or whether RNA was the only genetic material at the origin of life

and was subsequently replaced by the more stable DNA [26]. If RNA was the only genetic material in primitive life, due to the unequal replication probabilities of different RNA sequences, for example, some RNAs with stable configurations may be more difficult to replicate [31]. This would cause the RNA distribution to shift after cell division and never stabilize, which would prevent cells from evolving stable decomposition and reproduction. When the environment changes, this shift in RNA distribution may become more obvious. On the other hand, since DNA can affect the distribution of RNA and protein through the central dogma [30], when the genetic material in the cell is replaced by DNA, the large amount of RNA and protein produced by DNA transcription will seriously interfere with the original distribution in the cell and cause the cell to hibernate or even die. This is like when a healthy cell is infected by a virus, the viral DNA will greatly affect the normal functioning of the cell. Even if this replacement can occur gradually at a very slow rate, it will cause the cell genome to have different evolutionary routes from the starting point and thus have no genetic similarities, which is inconsistent with the fact that different organisms are genetically consistent.

If the early division of cells is uneven and unstable, it means that a large number of daughter cells will reduce or even lose their ability to grow due to fluctuations in molecular frequency distribution. Since a single DNA chain can transcribe a large amount of RNA, which in turn translates into a large amount of protein, DNA can be viewed as a compression and storage of the entire distribution within the cell. Therefore, DNA is more efficient than RNA in retaining and transmitting information and this makes DNA look like a mechanism that evolved later in life to replace RNA. This may be one of the reasons why people are more willing to believe in the existence of the RNA world.

Therefore, DNA is more likely to have existed with RNA at the origin of life, which is supported by relevant evidence [27]. First, DNA can directly retain a large amount of RNA distribution in the pioneer pool through methods like RNA reverse transcription or RNA-DNA hybridization, and affect the distribution of molecules such as RNA and proteins through transcription and achieve a dynamic balance, thus ensuring that all primitive cells have similar genomes. Second, DNA can rapidly proliferate and continuously track RNA distribution through dynamic replication. For example, in an ideal geyser, DNA will constantly experience changes in temperature and solution concentration. When the geyser emits steam, under the condition of high temperature and catalytic macromolecules, the double helix DNA unwinds and some DNA single strands break, losing the previously stored information. When the pool gradually dries up and the temperature drops a little, RNA generates new single-stranded DNA through reverse transcription to store the current RNA distribution in the pioneer pool, and DNA replicates and forms stable double-stranded DNA again. This replication and proliferation in periodic environmental changes can be considered as RNA and DNA constantly undergoing imperfect PCR and reverse transcription PCR [28-29], which will rapidly increase the number of RNA and DNA macromolecules in the pool. From

this perspective, DNA can be regarded as a mapping of the distribution of RNA in the pool, and the genome of protocells should be quite large and complex.

This also means that a large number of RNA sequences with stable structures will appear repeatedly in the early genome, including RNAs that play important functions in the metabolic network, such as tRNA or ribosomal RNA sequences, as well as stable but non-functional RNA sequences and various short RNA sequences. These genes that store unimportant RNA sequences will lose their transcription ability due to gene mutations and lack of evolutionary pressure, which can well explain why there are a large number of non-coding regions in DNA. Moreover, due to the existence of the central dogma, DNA can be regarded as the compression and storage of the molecular frequency distribution in the entire cell. This means that the communication of material distribution between cells can be replaced by the communication of DNA, and the search for new catalysts in the cross-catalytic network has become the search for new gene sequences, which can be quickly obtained by simple gene mutations. All of these have greatly accelerated the evolution of organisms.




Figure. The distribution of DNA and RNA will be entangled with each other. High-frequency RNA sequences with stable configurations will become the main source of the genome of primitive cells.

How to test this hypothesis?

There are three possible experiments to test this hypothesis:

1. Simulating the dry-wet cycle. It is supposed to observe changes in the distribution of RNA with different sequences, and this distribution should become narrower after multiple wet-dry cycles in extreme environments.




Figure. Expected changes in the frequency distribution of RNA sequences.

2. After establishing a sufficiently complex metabolic reaction network, let the phospholipids randomly encapsulate substances through stirring and wet-dry cycles and observe the growth and rupture of these vesicles, as well as the changes in the growth rate after the fusion of various vesicles.
3. Under complex and extreme conditions, perform PCR and reverse transcription PCR at the same time, and observe that the DNA distribution is dynamically tracking and coping the RNA distribution.

The stability-distillation process can also be simulated by computer. First, a neural network needs to be pre-trained to estimate the stability and average existence time of molecules. At the same time, you need to train a new neural network to evaluate the collision between different molecules and the formation of new ones. In this way, these two neural networks can be used to simulate the intracellular metabolic reactions. All the molecules present will collectively determine the conduct of the new reaction and the stability of the products, while the newly formed macromolecules will act as potential catalysts for all known reactions and in turn affect the stability of all molecules. In this way, the frequency distribution of molecules in the system can be repeatedly updated in a manner similar to back propagation, and the new product is equivalent to an error, which will feedback to each layer in the system at the same time, update the weights, and then forward calculate all reactions to determine the change in formation of all products from each layer. As the stability of the product will determine the progress of the reaction, allowing simulations to calculate base pairing and nucleic acid self-replication. If the simulation of the entire origin of life process can be realized, then by adjusting the cycle period, temperature, pH value, types of organic matter, etc., it is likely that different protein translation systems or even gene systems can be obtained, such as a new configuration of ribosomes and its matching tRNA groups.




Figure. A possible training framework for simulating the stability-distillation process through a method similar to back propagation in neural networks.

Discussion

In conclusion, Life is a self-organizing and replicating molecule distribution that originated from terrestrial hot springs. From the perspective of macromolecules, a possible evolutionary goal is to generate or convert more macromolecules of the same type, which means that daughter cells will more easily have the same molecular frequency distribution as the mother, which is why people tend to believe that life originated from a few self-replicating macromolecules, but this life system constructed by very few self-replicating macromolecules, such as RNA, would be fragile and difficult to maintain in the early earth environment. According to this hypothesis, life should originate from a mixture of various organic and inorganic substances such as RNA, DNA, and proteins wrapped in lipids. In the hot springs of wet-dry cycles, macromolecules with more stable configurations are retained and aggregated into larger molecules. Through this stability-distillation, the molecular frequency distribution in the hot springs is narrowed until enough information is accumulated and the cross-catalytic network of primitive life is established. RNA and its complexes with stable configurations in extreme environments contributed to the initial gene source in early life, which also means that the earliest life already had a fairly complex and large genome, and it can be predicted that the macromolecules that existed at the origin of life should have high stability in extreme environments.

The structure of cells is also crucial to the origin and evolution of life. When a large number of macromolecules are gathered and wrapped by a selectively permeable lipid membrane, macromolecular monomers that can pass through the membrane more easily aggregate into macromolecules in the cell, and the increased intracellular osmotic pressure will favor cells to absorb more corresponding macromolecular monomers, thereby having the opportunity to synthesize more macromolecules and promote cell division. This means that this structure, usually called a cell, will have a strong

selectivity for substances that can widely catalyze the production of related macromolecules. The ribosome and its corresponding transcription system, RNA ligases, and DNA polymerases, etc., can be regarded as a catalyst for the production of a wide range of macromolecules, compounds that can provide energy for these polymerization reactions or membrane structures that can promote these reactions will also be selected at the same time, such as ATP and the inner membrane of chloroplasts. The unstable cell membrane and extreme environment during the origin of life caused the rupture, deformation, recombination, division and fusion of different cells to occur frequently, corresponding to the early sexual and asexual reproduction, which would also accelerate the search for the optimal molecular frequency distribution of life and produce two different evolutionary paths as cells and viruses at the same time. The recombination and separation of cells and cellular substances may evolve into the predation and reproduction behavior of cells respectively. The conclusions inferred from this hypothesis, such as the origin of life from terrestrial hot springs that can periodically erupt materials and are easily exposed to ultraviolet rays, the protocells genome has high complexity, and cells and viruses are two sides of life that originated at the same time, are consistent with the characteristics of the Luca genome of the earliest cells inferred from the microbial genome in the laboratory [42]. If we can simulate the hot springs with wet-dry cycle in the laboratory and test the changes in the distribution of different configurations of RNA and its complexes, and finally obtain a very narrow distribution of RNA sequences, then the hypothesis will be proven to be correct.

When a large number of cells gather, the contact and material exchange between cells [34-36] may change their interactions with each other and even the external environment around the cell, thereby gaining potential survival advantages. When the distribution of intracellular materials shifts after some cells divide and differentiate into cells with different adaptabilities, the aggregation of different cells also means more possible adaptability. These factors together may contribute to the emergence of multicellular organisms. From the perspective of the interaction between life and the environment, life can be regarded as a solution or confrontation to the complex environment, so life may adopt different survival and reproduction strategies to adapt to different environments. For example, the same individual can adapt to more complex environments as much as possible, which will allow organisms to evolve higher intelligence, longer lifespan, and even larger individuals. It is also possible to adopt strategies of different individuals or their different stages to adapt to different environments, such as insect eggs and their metamorphosis or plant seeds, etc. For any organism, appropriate movement ability is necessary, which means that it can use more limited adaptability to find a suitable environment and complete growth and development. So from this perspective, a more complete theory of natural selection should include a two-way selection of organisms and the environment.

According to this hypothesis, in the early Earth hot springs with wet-dry cycles under specific conditions, the time and process required for the origin of life should be a

definite process that can be simulated and calculated. However, the emergence of complex life, or even intelligent life, may be an uncertain process, because it is likely to require both appropriate marine and terrestrial environments. For example, the ocean can provide more nutrients and evolutionary space, while the land will have more complex environments. The emergence of complex life means that it needs to be able to develop from fewer cells or even single cells into complete individuals with different functional cells, which means that at least some cells need to be able to differentiate and migrate sequentially in a suitable environment and have multiple continuous attractors for the cells' molecule distribution. When the development process of these cells or embryos are not convergent enough, or the resulting organisms are not adaptable enough, it means that the entire evolutionary path would be started over or stagnant, this also implies the final complex life should have a highly conserved developmental path and each stage of development may exist as a separate species at some point in history. After this developmental path converges, local or terminal offsets and mutations can quickly produce a large number of new species without affecting the overall framework. And this continuous transfer of molecular frequency distribution in cells is very similar to the diffusion model in machine learning.

REFERENCES

- [1] Joyce, Gerald F., and Jack W. Szostak. "Protocells and RNA self-replication." *Cold Spring Harbor Perspectives in Biology* 10.9 (2018): a034801.
- [2] Ma, Wentao. "What does "the RNA world" mean to "the origin of life"?" *Life* 7.4 (2017): 49.
- [3] Wu, Kun, Qiong Nan, and Tianqi Wu. "Philosophical analysis of the meaning and nature of entropy and negative entropy theories." *Complexity* 2020 (2020).
- [4] Nowakowski, Jacek, and Ignacio Tinoco Jr. "RNA structure and stability." *Seminars in virology*. Vol. 8. No. 3. Academic Press, 1997.
- [5] Becker, Sidney, et al. "Wet-dry cycles enable the parallel origin of canonical and non-canonical nucleosides by continuous synthesis." *Nature communications* 9.1 (2018): 163.
- [6] Hassenkam, Tue, and David Deamer. "Visualizing RNA polymers produced by hot wet-dry cycling." *Scientific reports* 12.1 (2022): 10098.
- [7] Hassenkam, Tue, et al. "AFM images of viroid-sized rings that self-assemble from mononucleotides through wet-dry cycling: Implications for the origin of life." *Life* 10.12 (2020): 321.
- [8] Serra, Martin J., et al. "RNA hairpin loop stability depends on closing base pair." *Nucleic acids research* 21.16 (1993): 3845-3849.
- [9] Vecenie, Christopher J., and Martin J. Serra. "Stability of RNA hairpin loops closed by AU base pairs." *Biochemistry* 43.37 (2004): 11813-11817.
- [10] Xu, Xiaojun, and Shi-Jie Chen. "A method to predict the structure and stability of RNA/RNA complexes." *RNA structure determination: methods and protocols* (2016): 63-72.
- [11] Bikard, David, et al. "Folded DNA in action: hairpin formation and biological functions in prokaryotes." *Microbiology and Molecular Biology Reviews* 74.4 (2010): 570-588.
- [12] Pal, Arumay, and Yaakov Levy. "Structure, stability and specificity of the binding of ssDNA and ssRNA with proteins." *PLoS computational biology* 15.4 (2019): e1006768.

- [13] Kumar, Sandeep, Chung-Jung Tsai, and Ruth Nussinov. "Factors enhancing protein thermostability." *Protein engineering* 13.3 (2000): 179-191.
- [14] Root-Bernstein, Robert, and Meredith Root-Bernstein. "The ribosome as a missing link in prebiotic evolution II: Ribosomes encode ribosomal proteins that bind to common regions of their own mRNAs and rRNAs." *Journal of Theoretical Biology* 397 (2016): 115-127.
- [15] Di Giulio, Massimo. "The origin of the tRNA molecule: Independent data favor a specific model of its evolution." *Biochimie* 94.7 (2012): 1464-1466.
- [16] Joshi, Manesh Prakash, et al. "Formation and stability of prebiotically relevant vesicular systems in terrestrial geothermal environments." *Life* 7.4 (2017): 51.
- [17] Damer, Bruce, and David Deamer. "The hot spring hypothesis for an origin of life." *Astrobiology* 20.4 (2020): 429-452.
- [18] Howes, Philip, et al. "Phospholipid encapsulated semiconducting polymer nanoparticles: their use in cell imaging and protein attachment." *Journal of the American Chemical Society* 132.11 (2010): 3989-3996.
- [19] Forterre, Patrick. "The origin of viruses and their possible roles in major evolutionary transitions." *Virus research* 117.1 (2006): 5-16.
- [20] Heitman, Joseph. "Evolution of sexual reproduction: a view from the fungal kingdom supports an evolutionary epoch with sex before sexes." *Fungal Biology Reviews* 29.3-4 (2015): 108-117.
- [21] Gokhale, D. V., U. S. Puntambekar, and D. N. Deobagkar. "Protoplast fusion: a tool for intergeneric gene transfer in bacteria." *Biotechnology advances* 11.2 (1993): 199-217.
- [22] Johnson, Arlen W., Elsebet Lund, and James Dahlberg. "Nuclear export of ribosomal subunits." *Trends in biochemical sciences* 27.11 (2002): 580-585.
- [23] Mans, Berend, et al. "Comparative genomics, evolution and origins of the nuclear envelope and nuclear pore complex." *Cell cycle* 3.12 (2004): 1625-1650.
- [24] Archibald, John M. "Endosymbiosis and eukaryotic cell evolution." *Current Biology* 25.19 (2015): R911-R921.
- [25] Gould, Sven B., Sriram G. Garg, and William F. Martin. "Bacterial vesicle secretion and the evolutionary origin of the eukaryotic endomembrane system." *Trends in microbiology* 24.7 (2016): 525-534.
- [26] Ribas de Pouplana, Lluís, et al. "Origin and evolution of DNA and DNA replication machineries." *The genetic code and the origin of life* (2004): 145-168.
- [27] Ribas de Pouplana, Lluís, et al. "Origin and evolution of DNA and DNA replication machineries." *The genetic code and the origin of life* (2004): 145-168.
- [28] Lauerma, Lloyd H. "Advances in PCR technology." *Animal health research reviews* 5.2 (2004): 247-248.
- [29] Bachman, Julia. "Reverse-transcription PCR (rt-PCR)." *Methods in enzymology*. Vol. 530. Academic Press, 2013. 67-74.
- [30] Shapiro, James A. "Revisiting the central dogma in the 21st century." *Annals of the New York Academy of Sciences* 1178.1 (2009): 6-28.
- [31] Repass, John F., and Shinji Makino. "Importance of the positive-strand RNA secondary structure of a murine coronavirus defective interfering RNA internal replication signal in positive-strand RNA synthesis." *Journal of virology* 72.10 (1998): 7926-7933.
- [32] Neveu, Marc, Hyo-Joong Kim, and Steven A. Benner. "The "strong" RNA world hypothesis: Fifty years old." *Astrobiology* 13.4 (2013): 391-403.

- [33] Alberts, Bruce, et al. "Catalysis and the Use of Energy by Cells." *Molecular Biology of the Cell*. 4th edition. Garland Science, 2002.
- [34] Zhou, Zhihang, et al. "The versatile roles of testrapanins in cancer from intracellular signaling to cell–cell communication: cell membrane proteins without ligands." *Cell & Bioscience* 13.1 (2023): 59.
- [35] Turturici, Giuseppina, et al. "Extracellular membrane vesicles as a mechanism of cell-to-cell communication: advantages and disadvantages." *American Journal of Physiology-Cell Physiology* 306.7 (2014): C621-C633.
- [36] Ratajczak, Janina, et al. "Membrane-derived microvesicles: important and underappreciated mediators of cell-to-cell communication." *Leukemia* 20.9 (2006): 1487-1495.
- [37] Ibáñez de Aldecoa A L, Zafra O, González-Pastor J E. Mechanisms and regulation of extracellular DNA release and its biological roles in microbial communities[J]. *Frontiers in microbiology*, 2017, 8: 1390.
- [38] Elwell C, Mirrashidi K, Engel J. Chlamydia cell biology and pathogenesis[J]. *Nature Reviews Microbiology*, 2016, 14(6): 385-400. [39] Martin, William, and Reinhold G. Herrmann. "Gene transfer from organelles to the nucleus: how much, what happens, and why?." *Plant physiology* 118.1 (1998): 9-17.
- [40] Speijer, Dave. "What can we infer about the origin of sex in early eukaryotes?." *Philosophical Transactions of the Royal Society B: Biological Sciences* 371.1706 (2016): 20150530.
- [41] Gonsior, Michael, et al. "Yellowstone hot springs are organic chemodiversity hot spots." *Scientific reports* 8.1 (2018): 14155.
- [42] Moody E R R, Álvarez-Carretero S, Mahendrarajah T A, et al. The nature of the last universal common ancestor and its impact on the early Earth system[J]. *Nature Ecology & Evolution*, 2024: 1-13.