

# Sub-Dirac operators, spectral Einstein functionals and the noncommutative residue

Jin Hong<sup>a,b</sup>, Yuchen Yang<sup>a</sup>, Yong Wang<sup>a,\*</sup>

<sup>a</sup>*School of Mathematics and Statistics, Northeast Normal University, Changchun, 130024, China*

<sup>b</sup>*School of Mathematics and Statistics, Yili Normal University, Yining 835000, China*

---

## Abstract

In this paper, we define the spectral Einstein functional associated with the sub-Dirac operator for manifolds with boundary. A proof of the Dabrowski-Sitarz-Zalecki type theorem for spectral Einstein functions associated with the sub-Dirac operator on four-dimensional manifolds with boundary is also given.

*Keywords:* Sub-Dirac operator; noncommutative residue; spectral Einstein functional.

---

## 1. Introduction

The noncommutative residue which is found in [1, 2] by M. Wodzicki and V. W. Guillemin plays a prominent role in noncommutative geometry. Recently Dabrowski etc. [3] obtained the metric and Einstein functionals by two vector fields and Laplace-type operators over vector bundles, giving an interesting example of the spinor connection and square of the Dirac operator. An eminent spectral scheme is the small-time asymptotic expansion of the (localised) trace of heat kernel [4, 5] that generates geometric objects on manifolds such as residue, scalar curvature, and other scalar combinations of curvature tensors. The theory has very rich structures both in physics and mathematics.

In [6], Connes used the noncommutative residue to derive a conformal 4-dimensional Polyakov action analogy. Connes proved that the noncommutative residue on a compact manifold  $M$  coincided with Dixmier's trace on pseudodifferential operators of order  $-dim M$  [7]. Furthermore, Connes claimed the noncommutative residue of the square of the inverse of the Dirac operator was proportioned to the Einstein-Hilbert action in [7, 8]. Kastler provided a brute-force proof of this theorem in [9], while Kalau and Walze proved it in the normal coordinates system simultaneously in [10], which is called the Kastler-Kalau-Walze theorem now. Building upon the theory of the noncommutative residue introduced by Wodzicki, Fedosov etc. [11] constructed a noncommutative residue on the algebra of classical elements in Boutet de Monvel's calculus on a compact manifold with boundary of dimension  $n > 2$ . With elliptic pseudodifferential operators and noncommutative residue, it is natural way for investigating the Kastler-Kalau-Walze type theorem and operator-theoretic explanation of the gravitational action on manifolds with boundary. Concerning Dirac operators and signature operators, Wang performed computations of the noncommutative residue and successfully demonstrated the Kastler-Kalau-Walze type theorem for manifolds with boundaries [12–14].

Earlier Jean-Michel Bismut [15] proved a local index theorem for Dirac operators on a Riemannian manifold  $M$  associated with connections on  $TM$  which have non zero torsion. In [16], Ackermann and Tolksdorf proved a generalized version of the well-known Lichnerowicz formula for the square of the most general Dirac operator with torsion  $D_T$  on an even-dimensional spin manifold associated to a metric connection with torsion. Pfäffle and Stephan considered compact Riemannian spin manifolds without boundary equipped with orthogonal connections, and investigated the induced Dirac operators in [17]. Wang, Wang

---

\*Corresponding author.

*Email addresses:* jhong@nenu.edu.cn (Jin Hong), yangyc580@nenu.edu.cn (Yuchen Yang), wangy581@nenu.edu.cn (Yong Wang)

and Wu computed  $\widetilde{wres}[\pi^+ \tilde{\nabla}_X \tilde{\nabla}_Y (D_T^* D_T)^{-1} \circ \pi^+ (D_T^* D_T)^{-1}]$  and  $\widetilde{wres}[\pi^+ \tilde{\nabla}_X \tilde{\nabla}_Y D_T^{-1} \circ \pi^+ (D_T^* D_T D_T^*)]$  in [20], and provided an important reference for our paper calculation. In [25], in order to prove the Connes vanishing theorem, Liu and Zhang introduced the sub-Dirac operator. In [19] Wang and Wang defined lower dimensional volumes associated to sub-Dirac operators for foliations and compute these lower dimensional volumes. They also prove the Kastler-Kalau-Walze type theorems for foliations with or without boundary.

The purpose of this paper is to generalize the results in [3], [19], [20], and get spectral functionals associated with sub-Dirac operators on compact manifolds with boundary. For lower dimensional compact Riemannian manifolds with boundary, we compute the 4-dimensional residue of  $\tilde{\nabla}_X \tilde{\nabla}_Y D_F^{-4}$  and get Dabrowski-Sitarz-Zalecki theorems.

## 2. Spectral functionals for the sub-Dirac operator

In this section, we shall restrict our attention to the sub-Dirac operators for foliations. Let  $(M, F)$  be a closed foliation and  $M$  has spin leave,  $g^F$  be a metric on  $F$ . Let  $g^{TM}$  be a metric on  $TM$  which restricted to  $g^F$  on  $F$ . Let  $F^\perp$  be the orthogonal complement of  $F$  in  $TM$  with respect to  $g^{TM}$ . Then we have the following orthogonal splitting

$$TM = F \oplus F^\perp, \quad (2.1)$$

$$g^{TM} = g^F \oplus g^{F^\perp}, \quad (2.2)$$

where  $g^{F^\perp}$  is the restriction of  $g^{TM}$  to  $F^\perp$ .

Let  $P, P^\perp$  be the orthogonal projection from  $TM$  to  $F, F^\perp$  respectively. Let  $\nabla^{TM}$  be the Levi-Civita connection of  $g^{TM}$  and  $\nabla^F$  (*resp.*  $\nabla^{F^\perp}$ ) be the restriction of  $\nabla^{TM}$  to  $F$  (*resp.*  $F^\perp$ ). Without loss of generality, we assume  $F$  is oriented, spin and carries a fixed spin structure. Furthermore, we assume  $F^\perp$  is oriented and we do not assume that  $\dim F$  and  $\dim F^\perp$  are even. By assumption, we may write

$$\nabla^F = P \nabla^{TM} P, \quad (2.3)$$

$$\nabla^{F^\perp} = P^\perp \nabla^{TM} P^\perp. \quad (2.4)$$

Moreover, the connections  $\nabla^F$  ( $\nabla^{F^\perp}$ ) lift to  $S(F)$  ( $\wedge(F^{\perp,*})$ ) naturally denoted by  $\nabla^{S(F)}$  ( $\nabla^{\wedge(F^{\perp,*})}$ ). Then  $S(F) \otimes \wedge(F^{\perp,*})$  carries the induced tensor product connection

$$\nabla^{S(F) \otimes \wedge(F^{\perp,*})} = \nabla^{S(F)} \otimes \text{Id}_{\wedge(F^{\perp,*})} + \text{Id}_{S(F)} \otimes \nabla^{\wedge(F^{\perp,*})}. \quad (2.5)$$

Then we can define  $S \in \Omega(T^*M) \otimes \Gamma(\text{End}(TM))$ ,

$$\nabla^{TM} = \nabla^F + \nabla^{F^\perp} + S \quad (2.6)$$

For any  $X \in \Gamma(TM)$ ,  $S(X)$  exchanges  $\Gamma(F)$  and  $\Gamma(F^\perp)$  and is skew-adjoint with respect to  $g^{TM}$ . Let  $\{f_i\}_{i=1}^p$  be an oriented orthonormal basis of  $F$ ,  $\{h_s\}_{s=1}^q$  be an oriented orthonormal basis of  $F^\perp$ , we define

$$\tilde{\nabla}^F = \nabla^{S(F) \otimes \wedge(F^{\perp,*})} + \frac{1}{2} \sum_{j=1}^p \sum_{s=1}^q \langle S(\cdot) f_j, h_s \rangle c(f_j) c(h_s) \quad (2.7)$$

where the vector bundle  $F^\perp$  might well be non-spin.

**Definition 2.1.** [19] Let  $D_F$  be the operator mapping from  $\Gamma(S(F) \otimes \wedge(F^{\perp,*}))$  to itself defined by

$$D_F = \sum_{i=1}^p c(f_i) \tilde{\nabla}_{f_i}^F + \sum_{s=1}^q c(h_s) \tilde{\nabla}_{h_s}^F. \quad (2.8)$$

From (2.19) in [25], we shall make use of the Bochner Laplacian  $\Delta^F$  stating that

$$\Delta^F := - \sum_{i=1}^p \left( \tilde{\nabla}_{f_i} \right)^2 - \sum_{s=1}^q \left( \tilde{\nabla}_{h_s} \right)^2 + \tilde{\nabla}_{\sum_{i=1}^p \nabla_{f_i}^{TM}} f_i + \tilde{\nabla}_{\sum_{s=1}^q \nabla_{h_s}^{TM}} h_s. \quad (2.9)$$

Let  $r_M$  be the scalar curvature of the metric  $g^{TM}$ . Let  $R^{F^\perp}$  be the curvature tensor of  $F^\perp$ . From theorem 2.3 in [25], we have the following Lichnerowicz formula for  $D_F$ .

**Lemma 2.2.** [25] *The following identity holds*

$$\begin{aligned} D_F^2 = & \Delta^F + \frac{r_M}{4} + \frac{1}{4} \sum_{i=1}^p \sum_{r,s,t=1}^q \left\langle R^{F^\perp}(f_i, h_r) h_t, h_s \right\rangle c(f_i) c(h_r) \hat{c}(h_s) \hat{c}(h_t) \\ & + \frac{1}{8} \sum_{i,j=1}^p \sum_{s,t=1}^q \left\langle R^{F^\perp}(f_i, f_j) h_t, h_s \right\rangle c(f_i) c(f_j) \hat{c}(h_s) \hat{c}(h_t) \\ & + \frac{1}{8} \sum_{s,t,r,u=1}^q \left\langle R^{F^\perp}(h_r, h_l) h_t, h_s \right\rangle c(h_r) c(h_u) \hat{c}(h_s) \hat{c}(h_t). \end{aligned} \quad (2.10)$$

Let us now turn to compute the specification of  $D_F^3$ .

$$\begin{aligned} D_F^3 = & - \sum_l g^{tl}(x) c(\partial_t) g^{ij} \partial_i \partial_j \partial_l + \left[ \sum_l g^{tl}(x) c(\partial_t) \right] \left\{ \sum_{ij} (\partial_l g^{ij}) \partial_i \partial_j - 4 \sum_{ij} g^{ij} \sigma_i \partial_l \partial_j \otimes Id_{\wedge(F^\perp, *)} \right. \\ & - Id_{S(F)} \otimes 4 \sum_{ij} g^{ij} \tilde{\sigma}_i \partial_l \partial_j - 4 \sum_{ij} g^{ij} S(\partial_j) \partial_l \partial_i + 2 \sum_{ij} g^{ij} \sum_{kl} \Gamma_{ij}^k \partial_l \partial_k \\ & - \left. \left[ \frac{1}{4} \sum_{s,t=1}^p \left\langle \nabla_{\partial_j}^F f_s, f_t \right\rangle c(f_s) c(f_t) + \frac{1}{4} \sum_{l,\alpha=1}^q \left\langle \nabla_{\partial_j}^{F^\perp} h_l, h_\alpha \right\rangle [c(h_l) c(h_\alpha) - \hat{c}(h_l) \hat{c}(h_\alpha)] \right] \sum_{i,j} g^{ij} \partial_i \partial_j \right\} \\ & + \left[ \sum_l g^{tl}(x) c(\partial_t) \right] \left\{ -2 \sum_{ij} (\partial_l g^{ij}) \sigma_i \partial_j \otimes Id_{\wedge(F^\perp, *)} - 2 \sum_{ij} g^{ij} (\partial_l \sigma_i) \partial_j \otimes Id_{\wedge(F^\perp, *)} \right. \\ & - Id_{S(F) \otimes 2} (\partial_l g^{ij}) \tilde{\sigma}_i \partial_j - Id_{S(F)} \otimes 2 \sum_{ij} g^{ij} (\partial_l \tilde{\sigma}_i) \partial_j - 2 \sum_{ij} (\partial_l g^{ij}) S(\partial_j) \partial_i \\ & - 2 \sum_{ij} g^{ij} [\partial_l S(\partial_j)] \partial_i + \sum_{ij} (\partial_l g^{ij}) \sum_k \Gamma_{ij}^k \partial_k + \sum_{ij} g^{ij} \sum_k (\partial_l \Gamma_{ij}^k) \partial_k \left. \right\} \\ & + \left\{ \frac{1}{4} \sum_{s,t=1}^p \left\langle \nabla_{\partial_j}^F f_s, f_t \right\rangle c(f_s) c(f_t) + \frac{1}{4} \sum_{l,\alpha=1}^q \left\langle \nabla_{\partial_j}^{F^\perp} h_l, h_\alpha \right\rangle [c(h_l) c(h_\alpha) - \hat{c}(h_l) \hat{c}(h_\alpha)] \right\} \\ & \times \left\{ - \sum_{ij} g^{ij} \left[ 2 \sigma_i \partial_j \otimes Id_{\wedge(F^\perp, *)} + Id_{S(F)} \otimes 2 \tilde{\sigma}_i \partial_j + S(\partial_j) \partial_i + S(\partial_i) \partial_j - \sum_k \Gamma_{ij}^k \partial_k \right] + \frac{r_M}{4} \right. \\ & + \frac{1}{4} \sum_{i=1}^p \sum_{r,s,t=1}^q \left\langle R^{F^\perp}(f_i, h_r) h_t, h_s \right\rangle c(f_i) c(h_r) \hat{c}(h_s) \hat{c}(h_t) \\ & + \frac{1}{8} \sum_{i,j=1}^p \sum_{s,t=1}^q \left\langle R^{F^\perp}(f_i, f_j) h_t, h_s \right\rangle c(f_i) c(f_j) \hat{c}(h_s) \hat{c}(h_t) \\ & \left. + \frac{1}{8} \sum_{s,t,r,u=1}^q \left\langle R^{F^\perp}(h_r, h_l) h_t, h_s \right\rangle c(h_r) c(h_u) \hat{c}(h_s) \hat{c}(h_t) \right\}. \end{aligned} \quad (2.11)$$

In order to get a Kastler-Kalau-Walze type theorem for foliations, Liu and Wang [26] considered the noncommutative residue of the  $-n + 2$  power of the sub-Dirac operator, and got the following Kastler-Kalau-Walze type theorem for foliations.

The following lemma of Dabrowski etc.'s Einstein functional play a key role in our proof of the Einstein functional. Let  $V, W$  be a pair of vector fields on a compact Riemannian manifold  $M$ , of dimension  $n = 2m$ . Using the Laplace operator  $\Delta = -(\sum_{j=1}^n \tilde{\nabla}_{e_j} \tilde{\nabla}_{e_j} - \tilde{\nabla}_{\nabla_{e_j}^L e_j}) + E$  acting on sections of a vector bundle  $\overline{E}$  where  $\tilde{\nabla}$  is a connection on  $\overline{E}$ , which may contain both some nontrivial connections and torsion, the spectral functionals over vector fields defined by

**Lemma 2.3.** [3] *The Einstein functional equal to*

$$Wres(\tilde{\nabla}_V \tilde{\nabla}_W \Delta^{-m}) = \frac{v_{n-1}}{6} 2^m \int_M G(V, W) vol_g + \frac{v_{n-1}}{2} \int_M F(V, W) vol_g + \frac{1}{2} \int_M (\text{Tr} E) g(V, W) vol_g, \quad (2.12)$$

where  $G(V, W)$  denotes the Einstein tensor evaluated on the two vector fields,  $F(V, W) = \text{Tr}(V_a W_b F_{ab})$  and  $F_{ab}$  is the curvature tensor of the connection  $T$ ,  $\text{Tr} E$  denotes the trace of  $E$  and  $v_{n-1} = \frac{2\pi^m}{\Gamma(m)}$ .

The aim of this section is to prove the following.

**Theorem 2.4.** *For the Laplace (type) operator  $\Delta_F = D_F^2$ , the Einstein functional equal to*

$$Wres(\tilde{\nabla}_V^F \tilde{\nabla}_W^F (D_F)^{-2m}) = \frac{2^{\frac{p}{2}+q+1} \pi^{\frac{n}{2}}}{6\Gamma(\frac{p+q}{2})} \int_M G(V, W) dvol_g + 2^{\frac{p}{2}+q-3} \int_M s g(V, W) dvol_M, \quad (2.13)$$

where  $s$  is the scalar curvature.

*Proof.* By the definition of connection  $\tilde{\nabla}^F$ , we have

$$\begin{aligned} \tilde{\nabla}_X^F &= \nabla_X^{S(F) \otimes (F^\perp, *)} + \frac{1}{2} \sum_{j=1}^p \sum_{s=1}^q \langle S(X) f_j, h_s \rangle c(f_j) c(h_s) \\ &= X + \frac{1}{4} \sum_{j,l=1}^p \langle \nabla_X^F f_j, f_l \rangle c(f_j) c(f_l) + \frac{1}{4} \sum_{s,t=1}^q \langle \nabla_X^{F^\perp} h_s, h_t \rangle (c(h_s) c(h_t) - \hat{c}(h_s) \hat{c}(h_t)) \\ &\quad + \frac{1}{2} \sum_{j=1}^p \sum_{s=1}^q \langle S(X) f_j, h_s \rangle c(f_j) c(h_s) \\ &:= X + \overline{A}(X). \end{aligned} \quad (2.14)$$

Let  $V = \sum_{a=1}^n V^a e_a$ ,  $W = \sum_{b=1}^n W^b e_b$ , in view of that

$$F(V, W) = \text{Tr}(V_a W_b F_{ab}) = \sum_{a,b=1}^n V_a W_b \text{Tr}^{S(F) \otimes (F^\perp, *)}(F_{e_a, e_b}), \quad (2.15)$$

we obtain

$$F_{e_a, e_b} = e_a(\overline{A}(e_b)) - e_b(\overline{A}(e_a)) + \overline{A}(e_a)\overline{A}(e_b) - \overline{A}(e_b)\overline{A}(e_a) - \overline{A}([e_a, e_b]). \quad (2.16)$$

Also, straightforward computations yield

$$\begin{aligned} \text{Tr}(e_a(\overline{A}(e_b))) &= \text{Tr} \left[ e_a \left( \frac{1}{4} \sum_{j,l=1}^p \langle \nabla_X^F f_j, f_l \rangle c(f_j) c(f_l) + \frac{1}{4} \sum_{s,t=1}^q \langle \nabla_X^{F^\perp} h_s, h_t \rangle (c(h_s) c(h_t) - \hat{c}(h_s) \hat{c}(h_t)) \right. \right. \\ &\quad \left. \left. + \frac{1}{2} \sum_{j=1}^p \sum_{s=1}^q \langle S(X) f_j, h_s \rangle c(f_j) c(h_s) \right) \right] \\ &= 0, \end{aligned} \quad (2.17)$$

in the same vein

$$\mathrm{Tr}\overline{A}([e_a, e_b]) = 0, \quad (2.18)$$

so

$$F(V, W) = 0. \quad (2.19)$$

Let  $D_F^2 = \Delta^F + E$ , we have

$$\begin{aligned} E &= \frac{s}{4} + \frac{1}{4} \sum_{i=1}^p \sum_{r,s,t=1}^q \langle R^{F^\perp}(f_i, h_r)h_t, h_s \rangle c(f_i)c(h_r)\hat{c}(h_s)\hat{c}(h_t) \\ &\quad + \frac{1}{4} \sum_{i,j=1}^p \sum_{s,t=1}^q \langle R^{F^\perp}(f_i, f_j)h_t, h_s \rangle c(f_i)c(f_j)\hat{c}(h_s)\hat{c}(h_t) \\ &\quad + \frac{1}{4} \sum_{s,t,r,u=1}^q \langle R^{F^\perp}(h_r, h_u)h_t, h_s \rangle c(h_r)c(h_u)\hat{c}(h_s)\hat{c}(h_t), \end{aligned} \quad (2.20)$$

and

$$\mathrm{Tr}\langle R^{F^\perp}(f_i, h_r)h_t, h_s \rangle c(f_i)c(h_r)\hat{c}(h_s)\hat{c}(h_t) = 0, \quad (2.21)$$

$$\mathrm{Tr}\langle R^{F^\perp}(f_i, f_j)h_t, h_s \rangle c(f_i)c(f_j)\hat{c}(h_s)\hat{c}(h_t) = 0, \quad (2.22)$$

$$\mathrm{Tr}\langle R^{F^\perp}(h_r, h_u)h_t, h_s \rangle c(h_r)c(h_u)\hat{c}(h_s)\hat{c}(h_t) = 0. \quad (2.23)$$

Hence

$$\mathrm{Tr}E = \frac{s}{4}\mathrm{Tr}[Id] = \frac{s}{4}2^{\frac{p}{2}+q} = 2^{\frac{p}{2}+q-2}s, \quad (2.24)$$

which finishes the proof of Theorem 2.4.  $\square$

### 3. The residue for the sub-Dirac operator $\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}$ and $D_F^{-2}$

In this section, we compute the lower dimensional volume for 4-dimension compact manifolds with boundary and get a Dabrowski-Sitarz-Zalecki type formula in this case. Some basic knowledge such as boundary metric, frame, noncommutative residue of manifold with boundary and Boutet de Monvel's algebra can be referred to in [12], and we will not repeat it here. We will consider  $D_F^2$ . Since  $[\sigma_{-4}(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2} \circ D_F^{-2})]|_M$  has the same expression as  $[\sigma_{-4}(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2} \circ D_F^{-2})]|_M$  in the case of manifolds without boundary, so locally we can use Theorem 2.4 to compute the first term.

**Corollary 3.1.** *Let  $M$  be a 4-dimensional compact manifold without boundary and  $\tilde{\nabla}^F$  be an orthogonal connection. Then we get the volumes associated to  $\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}$  and  $D_F^2$  on compact manifolds without boundary*

$$Wres[\sigma_{-4}(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2} \circ D_F^{-2})] = \frac{8\pi^2}{3} \int_M G(X, Y) dvol_g + \int_M sg(X, Y) dvol_M, \quad (3.1)$$

where  $s$  is the scalar curvature.

Let  $p_1, p_2$  be nonnegative integers and  $p_1 + p_2 \leq n$ , denote by  $\sigma_l(\tilde{A})$  the  $l$ -order symbol of an operator  $\tilde{A}$ , an application of (3.5) and (3.6) in [12] shows that

**Definition 3.2.** *Spectral Einstein functional associated the sub-Dirac operator of manifolds with boundary is defined by*

$$\text{vol}_n^{\{p_1, p_2\}} M := \widetilde{\text{Wres}}[\pi^+(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F (D_F^2)^{-p_1}) \circ \pi^+(D_F^{-2})^{p_2}], \quad (3.2)$$

where  $\pi^+(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F (D_F^2)^{-p_1})$ ,  $\pi^+(D_F^{-2})^{p_2}$  are elements in Boutet de Monvel's algebra [13].

For the sub-Dirac operator  $\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}$  and  $D_F^{-2}$ , denote by  $\sigma_l(\tilde{A})$  the  $l$ -order symbol of an operator  $\tilde{A}$ . An application of (2.1.4) in [12] shows that

$$\begin{aligned} & \widetilde{\text{Wres}}[\pi^+(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F (D_F^2)^{p_1}) \circ \pi^+(D_F^2)^{-p_2}] \\ &= \int_M \int_{|\xi|=1} \text{tr}_{S(F) \otimes \wedge(F^\perp, *)} [\sigma_{-n}(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F (D_F^2)^{-p_1} \circ (D_F^2)^{-p_2}] \sigma(\xi) dx + \int_{\partial M} \Phi, \end{aligned} \quad (3.3)$$

where

$$\begin{aligned} \Phi &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+l}}{\alpha!(j+k+1)!} \text{tr}_{S(F) \otimes \wedge(F^\perp, *)} [\partial_{x_n}^j \partial_{\xi'}^\alpha \partial_{\xi_n}^k \sigma_r^+(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F (D_F^2)^{-p_1})(x', 0, \xi', \xi_n) \\ & \quad \times \partial_{x_n}^\alpha \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l((D_F^2)^{-p_2})(x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx', \end{aligned} \quad (3.4)$$

and the sum is taken over  $r - k - |\alpha| + \ell - j - 1 = -n, r \leq -p_1, \ell \leq -p_2$ .

So we only need to compute  $\int_{\partial M} \Phi$ . Recall the definition of the sub-Dirac operator  $D_F$  in [19].

$$D_F = \sum_{i=1}^p c(f_i) \tilde{\nabla}_{f_i} + \sum_{s=1}^q c(h_s) \tilde{\nabla}_{h_s}, \quad (3.5)$$

where  $c(f_i), c(h_s)$  denotes the Clifford action.

We define

$$\begin{aligned} \tilde{\nabla}_X^F &:= X + \frac{1}{2} \sum_{j=1}^p \sum_{s=1}^q \langle S(X) f_j, h_s \rangle c(f_j) c(h_s) + \frac{1}{4} \sum_{j,l=1}^p \langle \nabla_X^F f_j, f_l \rangle c(f_j) c(f_l) \\ & \quad + \frac{1}{4} \sum_{s,t=1}^q \langle \nabla_X^{F^\perp} h_s, h_t \rangle [c(h_s) c(h_t) - \hat{c}(h_s) \hat{c}(h_t)]. \end{aligned} \quad (3.6)$$

which is a connection on  $S(F) \otimes \wedge(F^\perp, *)$ .

Set

$$\begin{aligned} A(X) &= \frac{1}{2} \sum_{j=1}^p \sum_{s=1}^q \langle S(X) f_j, h_s \rangle c(f_j) c(h_s); \quad M(X) = \frac{1}{4} \sum_{j,l=1}^p \langle \nabla_X^F f_j, f_l \rangle c(f_j) c(f_l); \\ N(X) &= \frac{1}{4} \sum_{s,t=1}^q \langle \nabla_X^{F^\perp} h_s, h_t \rangle [c(h_s) c(h_t) - \hat{c}(h_s) \hat{c}(h_t)]. \end{aligned} \quad (3.7)$$

Let  $\tilde{\nabla}_X^F = X + A(X) + M(X) + N(X)$ , and  $\tilde{\nabla}_Y^F = Y + A(Y) + M(Y) + N(Y)$ , by (2.11), we obtain

$$\begin{aligned} \tilde{\nabla}_X^F \tilde{\nabla}_Y^F &= (X + A(X) + M(X) + N(X))(Y + A(Y) + M(Y) + N(Y)) \\ &= XY + M(Y)X + N(Y)X + A(Y)X + M(X)Y + N(X)Y + A(X)Y \\ & \quad + X[M(Y)] + X[N(Y)] + X[A(Y)] + M(X)M(Y) + M(X)N(Y) + M(X)A(Y) \\ & \quad + N(X)M(Y) + N(X)N(Y) + N(X)A(Y) + A(X)M(Y) + A(X)N(Y) + A(X)A(Y), \end{aligned} \quad (3.8)$$

where  $X = \sum_{j=1}^n X_j \partial_{x_j}$ ,  $Y = \sum_{l=1}^n Y_l \partial_{x_l}$ .

Let  $g^{ij} = g(dx_i, dx_j)$ ,  $\xi = \sum_k \xi_j dx_j$  and  $\nabla_{\partial_i}^L \partial_j = \sum_k \Gamma_{ij}^k \partial_k$ , we get

$$\begin{aligned} \sigma_k &= -\frac{1}{4} \sum_{k,l} \omega_{k,l}(\partial_k) c(f_k) c(f_l); \quad \tilde{\sigma}_k = \frac{1}{4} \sum_{r,t} \omega_{r,t}(\partial_k) [\bar{c}(h_r) \bar{c}(h_t) - c(h_r) c(h_t)]; \\ \xi^j &= g^{ij} \xi_i; \quad \Gamma^k = g^{ij} \Gamma_{ij}^k; \quad \sigma^j = g^{ij} \sigma_i; \quad \tilde{S}(\partial_i) = g^{ij} S(\partial_i). \end{aligned} \quad (3.9)$$

Then we have the following lemmas.

**Lemma 3.3.** *The following identities hold:*

$$\begin{aligned} \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) &= X[M(Y)] + X[N(Y)] + X[A(Y)] + M(X)M(Y) + M(X)N(Y) + M(X)A(Y) \\ &\quad + N(X)M(Y) + N(X)N(Y) + N(X)A(Y) + A(X)M(Y) + A(X)N(Y) + A(X)A(Y); \end{aligned} \quad (3.10)$$

$$\begin{aligned} \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) &= \sqrt{-1} \sum_{j,l=1}^n X_j \frac{\partial Y_l}{\partial x_j} \xi_l + \sqrt{-1} \sum_{j=1}^n [M(Y) + N(Y) + A(Y)] X_j \xi_j \\ &\quad + \sqrt{-1} \sum_{l=1}^n [M(X) + N(X) + A(X)] Y_l \xi_l; \end{aligned} \quad (3.11)$$

$$\sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) = - \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l. \quad (3.12)$$

**Lemma 3.4.** [19] *The following identities hold:*

$$\sigma_{-1}(D_F^{-1}) = \frac{ic(\xi)}{|\xi|^2}; \quad (3.13)$$

$$\sigma_{-2}(D_F^{-1}) = \frac{c(\xi)\sigma_0(D_F)c(\xi)}{|\xi|^4} + \frac{c(\xi)}{|\xi|^6} \sum_j c(dx_j) [\partial_{x_j}(c(\xi))|\xi|^2 - c(\xi)\partial_{x_j}(|\xi|^2)]; \quad (3.14)$$

$$\sigma_{-2}(D_F^{-2}) = |\xi|^{-2}; \quad (3.15)$$

$$\begin{aligned} \sigma_{-3}(D_F^{-2}) &= -\sqrt{-1}|\xi|^{-4} \xi_k (\Gamma^k - 2\sigma^k \otimes \text{Id}_{\wedge(F^\perp, \star)} - \text{Id}_{S(F)} \otimes 2\tilde{\sigma}^k \\ &\quad - \frac{1}{2} \sum_{i,j=1}^{2p} \sum_{s=1}^q \langle \nabla_{\partial_k}^{TM} f_j, h_s \rangle c(f_j) c(h_s) - \frac{1}{2} \sum_{s,t=1}^q \sum_{i=1}^{2p} \langle \nabla_{\partial_k}^{TM} f_j, h_s \rangle c(f_j) c(h_s) \\ &\quad - \sqrt{-1}|\xi|^{-6} 2\xi^j \xi_\alpha \xi_\beta \partial_j g^{\alpha\beta}. \end{aligned} \quad (3.16)$$

where,

$$\begin{aligned} \sigma_0(D_F) &= -\frac{1}{4} \sum_{i,k,l} \omega_{k,l}(f_i) c(f_i) c(f_k) c(f_l) \otimes \text{Id}_{\wedge(F^\perp, \star)} \\ &\quad - \frac{1}{4} \sum_{s,k,l} \omega_{k,l}(f_i) c(f_k) c(f_l) c(h_s) \otimes \text{Id}_{\wedge(F^\perp, \star)} \\ &\quad + \text{Id}_{S(F)} \otimes \frac{1}{4} \sum_{i,r,t} \omega_{r,t}(f_i) c(f_i) [\bar{c}(h_r) \bar{c}(h_t) - c(h_r) c(h_t)] \\ &\quad + \text{Id}_{S(F)} \otimes \frac{1}{4} \sum_{s,r,t} \omega_{r,t}(h_s) c(h_s) [\bar{c}(h_r) \bar{c}(h_t) - c(h_r) c(h_t)] \\ &\quad + \frac{1}{2} \sum_{i,j=1}^p \sum_{s=1}^q \langle \nabla_{f_i}^{TM} f_j, h_s \rangle c(f_i) c(f_j) c(h_s) + \frac{1}{2} \sum_{s,t=1}^q \sum_{i=1}^p \langle \nabla_{h_s}^{TM} h_t, f_i \rangle c(h_s) c(h_t) c(f_i). \end{aligned}$$

**Lemma 3.5.** *The following identities hold:*

$$\sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) = - \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l |\xi|^{-2}; \quad (3.17)$$

$$\begin{aligned} \sigma_{-1}(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) &= \sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-3}(D_F^{-2}) + \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-2}(D_F^{-2}) \\ &\quad + \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F)] D_{x_j} [\sigma_{-2}(D_F^{-2})]. \end{aligned} \quad (3.18)$$

Now we need to compute  $\int_{\partial M} \Phi$ . When  $n = p + q = 4$ , then  $\text{tr}_{S(F) \otimes (F^\perp, *)}[\text{id}] = 8$ , the sum is taken over  $r + l - k - j - |\alpha| = -3$ ,  $r \leq 0$ ,  $l \leq -2$ , then we have the following five cases:

**case a) I)**  $r = 0$ ,  $l = -2$ ,  $k = j = 0$ ,  $|\alpha| = 1$ .

By (3.4), we get

$$\Phi_1 = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{Tr}[\partial_{\xi'}^\alpha \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \times \partial_{x'}^\alpha \partial_{\xi_n} \sigma_{-2}(D_F^{-2})](x_0) d\xi_n \sigma(\xi') dx'. \quad (3.19)$$

By Lemma 2.2 in [13], for  $i < n$ , then

$$\partial_{x_i} \sigma_{-2}(D_F^{-2})(x_0) = \partial_{x_i} (|\xi|^{-2})(x_0) = - \frac{\partial_{x_i} (|\xi|^2)(x_0)}{|\xi|^4} = 0, \quad (3.20)$$

so  $\Phi_1 = 0$ .

**case a) II)**  $r = 0$ ,  $l = -2$ ,  $k = |\alpha| = 0$ ,  $j = 1$ .

By (3.4), we get

$$\Phi_2 = - \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \times \partial_{\xi_n}^2 \sigma_{-2}(D_F^{-2})](x_0) d\xi_n \sigma(\xi') dx'. \quad (3.21)$$

By Lemma 3.3, we have

$$\partial_{\xi_n}^2 \sigma_{-2}(D_F^{-2})(x_0) = \partial_{\xi_n}^2 (|\xi|^{-2})(x_0) = \frac{6\xi_n^2 - 2}{(1 + \xi_n^2)^3}. \quad (3.22)$$

It follows that

$$\partial_{x_n} \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2})(x_0) = \partial_{x_n} \left( - \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l |\xi|^{-2} \right) = \frac{\sum_{j,l=1}^n X_j Y_l \xi_j \xi_l h'(0) |\xi'|^2}{(1 + \xi_n^2)^2}. \quad (3.23)$$

By integrating formula, we obtain

$$\begin{aligned} \pi_{\xi_n}^+ \partial_{x_n} \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2})(x_0) &= \partial_{x_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \\ &= - \frac{i\xi_n + 2}{4(\xi_n - i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) - \frac{i\xi_n}{4(\xi_n - i)^2} X_n Y_n h'(0) \\ &\quad - \frac{i}{4(\xi_n - i)^2} \sum_{j=1}^{n-1} X_j Y_n \xi_j - \frac{i}{4(\xi_n - i)^2} \sum_{l=1}^{n-1} X_n Y_l \xi_l. \end{aligned} \quad (3.24)$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for



computing **case a) II**. From (3.32) and (3.34), we obtain

$$\begin{aligned}
& \text{Tr}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \times \partial_{\xi_n}^2 \sigma_{-2}(D_F^{-2})](x_0) \\
&= 4 \left[ \frac{i - 3i\xi_n^2}{(\xi_n - i)^4 (\xi_n + i)^3} + \frac{1 - 3\xi_n^2}{(\xi_n - i)^5 (\xi_n + i)^3} \right] \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \\
&+ 4 \left[ \frac{i - 3i\xi_n^2}{(\xi_n - i)^4 (\xi_n + i)^3} - \frac{1 - 3\xi_n^2}{(\xi_n - i)^5 (\xi_n + i)^3} \right] X_n Y_n h'(0) \\
&+ 4 \frac{(1 - 3\xi_n^2)i}{(\xi_n - i)^5 (\xi_n + i)^3} \sum_{j=1}^{n-1} X_j Y_n \xi_j + 4 \frac{(1 - 3\xi_n^2)i}{(\xi_n - i)^5 (\xi_n + i)^3} \sum_{l=1}^{n-1} X_n Y_l \xi_l.
\end{aligned} \tag{3.25}$$

Therefore, we get

$$\Phi_2 = \left( \frac{5}{24} \sum_{j=1}^{n-1} X_j Y_j - \frac{1}{8} X_n Y_n \right) h'(0) \pi \Omega_3 dx', \tag{3.26}$$

where  $\Omega_3$  is the canonical volume of  $S^2$ .

**case a) III**  $r = 0, l = -2, j = |\alpha| = 0, k = 1$ .

By (3.4), we get

$$\begin{aligned}
\Phi_3 &= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-2}(D_F^{-2})](x_0) d\xi_n \sigma(\xi') dx' \\
&= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \times \partial_{x_n} \sigma_{-2}(D_F^{-2})](x_0) d\xi_n \sigma(\xi') dx'.
\end{aligned} \tag{3.27}$$

By Lemma 4.4, we have

$$\partial_{x_n} \sigma_{-2}(D_F^{-2})(x_0)|_{|\xi'|=1} = -\frac{h'(0)}{(1 + \xi_n^2)^2}. \tag{3.28}$$

An easy calculation gives

$$\begin{aligned}
\pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2})(x_0)|_{|\xi'|=1} &= \frac{i}{2(\xi_n - i)} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - \frac{i}{2(\xi_n - i)} X_n Y_n \\
&- \frac{1}{2(\xi_n - i)} \sum_{j=1}^{n-1} X_j Y_n \xi_j - \frac{1}{2(\xi_n - i)} \sum_{l=1}^{n-1} X_n Y_l \xi_l.
\end{aligned} \tag{3.29}$$

Also, straightforward computations yield

$$\begin{aligned}
\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2})(x_0)|_{|\xi'|=1} &= \frac{i}{(\xi_n - i)^3} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - \frac{1}{(\xi_n - i)^3} X_n Y_n \\
&- \frac{1}{(\xi_n - i)^3} \sum_{j=1}^{n-1} X_j Y_n \xi_j - \frac{1}{(\xi_n - i)^3} \sum_{l=1}^{n-1} X_n Y_l \xi_l.
\end{aligned} \tag{3.30}$$

Therefore, we get

$$\begin{aligned}
\Phi_3 &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \times \partial_{x_n} \sigma_{-2}(D_F^{-2})](x_0) d\xi_n \sigma(\xi') dx' \\
&= \left( -\frac{5}{24} \sum_{j=1}^{n-1} X_j Y_j + \frac{5}{8} X_n Y_n \right) h'(0) \pi \Omega_3 dx'.
\end{aligned} \tag{3.31}$$

**case b)**  $r = 0$ ,  $l = -3$ ,  $k = j = |\alpha| = 0$ .

By (3.4), we get

$$\begin{aligned}\Phi_4 &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \times \partial_{\xi_n} \sigma_{-3}(D_F^{-2})](x_0) d\xi_n \sigma(\xi') dx' \\ &= i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \times \sigma_{-3}(D_F^{-2})](x_0) d\xi_n \sigma(\xi') dx'.\end{aligned}\quad (3.32)$$

By Lemma 3.4, we have

$$\begin{aligned}\sigma_{-3}(D_F^{-2})(x_0)|_{|\xi'|=1} &= \frac{-2ih'(0)\xi_n}{(1+\xi_n^2)^3} - \frac{i}{(1+\xi_n^2)^2} \left( \frac{3}{2}h'(0)\xi_n + \frac{1}{2} \sum_{k,l} \omega_{k,l}(\partial^k)(x_0)c(f_k)c(f_l) \otimes \text{Id}_{\wedge(F^{\perp,*})} \right. \\ &\quad \left. - \text{Id}_{S(F)} \otimes \frac{1}{2} \sum_{r,t} \omega_{r,t}(\partial^k)(x_0)[\hat{c}(h_r)\hat{c}(h_t) - c(h_r)c(h_t)] - \sum_{i,j=1}^p \sum_{s=1}^q \langle \nabla_{\partial^k}^{TM} f_j, h_s \rangle c(f_j)c(h_s) \right).\end{aligned}\quad (3.33)$$

$$\begin{aligned}\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2})(x_0)|_{|\xi'|=1} &= -\frac{i}{2(\xi_n - i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l + \frac{i}{2(\xi_n - i)^2} X_n Y_n \\ &\quad + \frac{1}{2(\xi_n - i)^2} \sum_{j=1}^{n-1} X_j Y_n \xi_j + \frac{1}{2(\xi_n - i)^2} \sum_{l=1}^{n-1} X_n Y_l \xi_l.\end{aligned}\quad (3.34)$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case b)**. By the trace identity  $\text{tr}(AB) = \text{tr}(BA)$ ,  $\text{tr}(A \otimes B) = \text{tr}(A) \cdot \text{tr}(B)$  and the relation of the Clifford action, we have

$$\text{Tr}[c(f_k)c(f_l)] = -\delta_k^l 2^{\frac{q}{2}}, \quad (3.35)$$

$$\text{Tr}[\hat{c}(h_r)\hat{c}(h_t) - c(h_r)c(h_t)] = 2\delta_r^t 2^q. \quad (3.36)$$

Then, we have

$$\begin{aligned}&\text{Tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \times \sigma_{-3} D_F^{-2}](x_0) \\ &\sim \left[ -\frac{8\xi_n}{(\xi_n - i)^5(\xi_n + i)^3} - \frac{6\xi_n}{(\xi_n - i)^4(\xi_n + i)^2} \right] h'(0) \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \\ &\quad + \left[ \frac{8\xi_n}{(\xi_n - i)^5(\xi_n + i)^3} + \frac{6\xi_n}{(\xi_n - i)^4(\xi_n + i)^2} \right] h'(0) X_n Y_n.\end{aligned}\quad (3.37)$$

Therefore, we get

$$\begin{aligned}\Phi_4 &= i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \times \sigma_{-3} D_F^{-2}](x_0) d\xi_n \sigma(\xi') dx' \\ &= \left( \frac{11}{24} \sum_{j=1}^{n-1} X_j Y_j - \frac{11}{8} X_n Y_n \right) h'(0) \pi \Omega_3 dx'.\end{aligned}\quad (3.38)$$

**case c)**  $r = -1$ ,  $\ell = -2$ ,  $k = j = |\alpha| = 0$ .

By (3.4), we get

$$\Phi_5 = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\pi_{\xi_n}^+ \sigma_{-1}(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \times \partial_{\xi_n} \sigma_{-2}(D_F^{-2})](x_0) d\xi_n \sigma(\xi') dx'. \quad (3.39)$$

By Lemma 4.4, we have

$$\partial_{\xi_n} \sigma_{-2}(D_F^{-2})(x_0)|_{|\xi'|=1} = -\frac{2\xi_n}{(\xi_n^2 + 1)^2}. \quad (3.40)$$

Since

$$\begin{aligned} \sigma_{-1}(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2})(x_0)|_{|\xi'|=1} &= \sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-3}(D_F^{-2}) + \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-2}(D_F^{-2}) \\ &+ \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F)] D_{x_j} [\sigma_{-2}(D_F^{-2})]. \end{aligned} \quad (3.41)$$

Explicit representation the first item of (3.41),

$$\begin{aligned} &\sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-3}(D_F^{-2})(x_0)|_{|\xi'|=1} \\ &= - \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l \times \left\{ \frac{-2ih'(0)\xi_n}{(1+\xi_n^2)^3} - \frac{i}{(1+\xi_n^2)^2} \left( \frac{3}{2} h'(0)\xi_n + \frac{1}{2} \sum_{k,l} \omega_{k,l} (\partial^k)(x_0) c(f_k) c(f_l) \otimes \text{Id}_{\wedge(F^{\perp,*})} \right. \right. \\ &\quad \left. \left. - \text{Id}_{S(F)} \otimes \frac{1}{2} \sum_{r,t} \omega_{r,t} (\partial^k)(x_0) [\hat{c}(h_r) \hat{c}(h_t) - c(h_r) c(h_t)] - \sum_{i,j=1}^p \sum_{s=1}^q \langle \nabla_{\partial^k}^T f_j, h_s \rangle c(f_j) c(h_s) \right) \right\}. \end{aligned} \quad (3.42)$$

Explicit representation the second item of (3.41),

$$\begin{aligned} &\sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-2}(D_F^{-2})(x_0)|_{|\xi'|=1} \\ &= \left( \sqrt{-1} \sum_{j,l=1}^n X_j \frac{\partial Y_l}{\partial x_j} \xi_l + \sqrt{-1} \sum_j [M(Y) + N(Y) + A(Y)] X_j \xi_j \right. \\ &\quad \left. + \sqrt{-1} \sum_l [M(X) + N(X) + A(X)] Y_l \xi_l \right) \times |\xi|^{-2}. \end{aligned} \quad (3.43)$$

Explicit representation the third item of (3.41),

$$\begin{aligned} &\sum_{j=1}^n \partial_{\xi_j} [\sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F)] D_{x_j} [\sigma_{-2}(D_F^{-2})](x_0)|_{|\xi'|=1} \\ &= \sum_{j=1}^n \partial_{\xi_j} \left[ - \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l \right] (-\sqrt{-1}) \partial_{x_j} (|\xi|^{-2}) \\ &= \sum_{j=1}^n \sum_{l=1}^n \sqrt{-1} (x_j Y_l + x_l Y_j) \xi_l \partial_{x_j} (|\xi|^{-2}). \end{aligned} \quad (3.44)$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case c**). Also, straightforward computations yield

$$\begin{aligned} &\text{Tr}[\pi_{\xi_n}^+ \sigma_{-1}(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \times \partial_{\xi_n} \sigma_{-2}(D_F^{-2})](x_0)|_{|\xi'|=1} \\ &= \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \times \frac{18i\xi_n - 10\xi_n^2}{(\xi_n - i)^5 (\xi_n + i)^2} h'(0) + X_n Y_n \times \left[ \frac{-20i\xi_n^3 - 36\xi_n^2 + 12i\xi_n}{(\xi_n - i)^5 (\xi_n + i)^2} + \frac{8\xi_n}{(\xi_n - i)^4 (\xi_n + i)^2} \right] h'(0), \end{aligned} \quad (3.45)$$

Hence in this case,

$$\begin{aligned}\Phi_5 &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\pi_{\xi_n}^+ \sigma_{-1}(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \times \partial_{\xi_n} \sigma_{-2} D_F^{-2}](x_0) d\xi_n \sigma(\xi') dx' \\ &= - \left( \frac{2}{3} \sum_{j=1}^{n-1} X_j Y_j + \frac{5}{8} X_n Y_n \right) h'(0) \pi \Omega_3 dx'.\end{aligned}\quad (3.46)$$

Now  $\Phi$  is the sum of the cases (a), (b) and (c). Let  $X = X^T + X_n \partial_n$ ,  $Y = Y^T + Y_n \partial_n$ , then we have  $\sum_{j=1}^{n-1} X_j Y_j(x_0) = g(X^T, Y^T)(x_0)$ . Therefore, we get

$$\Phi = \sum_{i=1}^5 \Phi_i = - \left[ \frac{5}{24} g(X^T, Y^T) + \frac{3}{2} X_n Y_n \right] h'(0) \pi \Omega_3 dx'. \quad (3.47)$$

Then we obtain following theorem

**Theorem 3.6.** *Let  $M$  be a 4-dimensional compact foliated manifold with boundary and spin leave,  $\tilde{\nabla}$  be an orthogonal connection with torsion. Then we get the volumes associated to  $\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}$  and  $D_F^{-2}$*

$$\begin{aligned}& \widetilde{\text{Wres}}[\pi^+(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \circ \pi^+(D_F^2)] \\ &= \frac{8\pi^2}{3} \int_M G(X, Y) d\text{vol}_g + \int_M s g(X, Y) d\text{vol}_M - \int_{\partial M} \left[ \frac{5}{24} g(X^T, Y^T) + \frac{3}{2} X_n Y_n \right] h'(0) \pi \Omega_3 dx'.\end{aligned}\quad (3.48)$$

where  $s$  is the scalar curvature.

By [13], we have the extrinsic curvature  $K = -\frac{3}{2} h'(0)$ , so when  $X_n = 0$  or  $Y_n = 0$  on boundary, we have

**Corollary 3.7.**

$$\begin{aligned}& \widetilde{\text{Wres}}[\pi^+(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-2}) \circ \pi^+(D_F^2)] \\ &= \frac{8\pi^2}{3} \int_M G(X, Y) d\text{vol}_g + \int_M s g(X, Y) d\text{vol}_M + \frac{5}{36} \int_{\partial M} g(X, Y)_{\partial M} K d\text{vol}_{\partial M}.\end{aligned}\quad (3.49)$$

So we have

**Definition 3.8.** *The spectral Einstein functional for manifolds with boundary*

$$E_H := \int_M G(X, Y) d\text{vol}_g + c_0 \int_{\partial M} g(X, Y)_{\partial M} K d\text{vol}_{\partial M}, \quad (3.50)$$

where  $c_0$  is a constant.

#### 4. The residue for sub-Dirac operators $\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}$ and $D_F^{-3}$

In this section, we compute the 4-dimension volume for sub-Dirac operators  $\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}$  and  $D_F^{-3}$ . Since  $[\sigma_{-4}(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1} \circ D_F^{-3})]_M$  has the same expression as  $[\sigma_{-4}(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1} \circ D_F^{-3})]_M$  in the case of manifolds without boundary, so locally we can use Theorem 2.4 to compute the first term.

**Corollary 4.1.** *Let  $M$  be a four dimensional compact manifold without boundary and  $\tilde{\nabla}$  be an orthogonal connection with torsion. Then we get the volumes associated to  $\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}$  and  $D_F^{-3}$  on compact manifolds without boundary*

$$\begin{aligned}& \text{Wres}[\sigma_{-4}(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1} \circ D_F^{-3})] \\ &= \frac{8\pi^2}{3} \int_M G(X, Y) d\text{vol}_g + \int_M s g(X, Y) d\text{vol}_M,\end{aligned}\quad (4.1)$$

where  $s$  is the scalar curvature.

Similar definition 3.2, we have

$$\begin{aligned} \tilde{\Phi} &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+l}}{\alpha!(j+k+1)!} \text{tr}_{S(F) \otimes \wedge(F^\perp, *)} [\partial_{x_n}^j \partial_{\xi'}^\alpha \partial_{\xi_n}^k \sigma_r^+ (\nabla_X^F \nabla_Y^F (D_F)^{-p_1}) (x', 0, \xi', \xi_n) \\ &\quad \times \partial_{x_n}^\alpha \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l ((D_F^3)^{-p_2}) (x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx', \end{aligned} \quad (4.2)$$

From lemma 3.3 and lemma 3.4, we have

**Lemma 4.2.** *The following identities hold:*

$$\sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}) = -\sqrt{-1} \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l c(\xi) |\xi|^{-2}; \quad (4.3)$$

$$\begin{aligned} \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}) &= \sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-2}(D_F^{-1}) + \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-1}(D_F^{-1}) \\ &\quad + \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F)] D_{x_j} [\sigma_{-1}(D_F^{-1})]. \end{aligned} \quad (4.4)$$

Write

$$D_x^\alpha = (-i)^{|\alpha|} \partial_x^\alpha; \quad \sigma(D_F^3) = p_3 + p_2 + p_1 + p_0; \quad \sigma(D_F^{-3}) = \sum_{j=3}^{\infty} q_{-j}. \quad (4.5)$$

By the composition formula of pseudodifferential operators, we have

$$\begin{aligned} 1 &= \sigma(D_F^3 \circ D_F^{-3}) = \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi'}^\alpha [\sigma(D_F^3) D_x^\alpha [\sigma(D_F^{-3})]] \\ &= (p_3 + p_2 + p_1 + p_0)(q_{-3} + q_{-4} + q_{-5} + \dots) \\ &\quad + \sum_j (\partial_{\xi_j} p_3 + \partial_{\xi_j} p_2 + \dots + \partial_{\xi_j} p_1 + \partial_{\xi_j} p_0) (D_{x_j} q_{-3} + D_{x_j} q_{-4} + D_{x_j} q_{-5} + \dots) \\ &= p_3 q_{-3} + (p_3 q_{-4} + p_2 q_{-3} + \sum_j \partial_{\xi_j} p_3 D_{x_j} q_{-3}) + \dots, \end{aligned} \quad (4.6)$$

so

$$q_{-3} = p_3^{-1}; \quad q_{-4} = -p_3^{-1} [p_2 p_3^{-1} + \sum_j \partial_{\xi_j} p_3 D_{x_j} (p_3^{-1})]. \quad (4.7)$$

Hence by Lemma 4.1 in [19] and (3.9), we have

**Lemma 4.3.** *The symbol of the sub-Dirac operator:*

$$\sigma_{-3}(D_F^{-3}) = \sqrt{-1} c(\xi) |\xi|^{-4}; \quad (4.8)$$

$$\begin{aligned} \sigma_{-4}(D_F^{-3}) &= \frac{c(\xi) \sigma_2(D_F^3) c(\xi)}{|\xi|^8} + \frac{\sqrt{-1} c(\xi)}{|\xi|^8} \left( |\xi|^4 c(dx_n) \partial_{x_n} c(\xi') \right. \\ &\quad \left. - 2h'(0) c(dx_n) c(\xi) + 2\xi_n c(\xi) \partial_{x_n} c(\xi') + 4\xi_n h'(0) \right), \end{aligned} \quad (4.9)$$

where,

$$\begin{aligned} \sigma_2(D_F^3) &= c(dx_l) \partial_l (g^{i,j}) \xi_i \xi_j + 2c(\xi) [2\sigma^i \otimes \text{Id}_{\wedge(F^\perp, *)} + \text{Id}_{S(F)} \otimes 2\tilde{\sigma}_i - \Gamma^k \\ &\quad + 2\tilde{S}(\partial_i) + (M(X) + N(X)) |\xi|^2] \xi_k. \end{aligned} \quad (4.10)$$

Now we need to compute  $\int_{\partial M} \tilde{\Phi}$ . When  $n = 4$ , then  $\text{tr}_{S(F) \otimes \wedge(F^{\perp, *})}[\text{id}] = 8$ , the sum is taken over  $r + l - k - j - |\alpha| = -3$ ,  $r \leq 0$ ,  $l \leq -2$ , then we have the following five cases:

**case a) I)**  $r = 1$ ,  $l = -2$ ,  $k = j = 0$ ,  $|\alpha| = 1$ .

By (4.2), we get

$$\tilde{\Phi}_1 = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{Tr}[\partial_{\xi'}^{\alpha} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}) \times \partial_{x'}^{\alpha} \partial_{\xi_n} \sigma_{-3}(D_F^{-3})](x_0) d\xi_n \sigma(\xi') dx'. \quad (4.11)$$

By Lemma 2.2 in [13], for  $i < n$ , then

$$\partial_{x_i} \sigma_{-3}(D_F^{-3})(x_0) = \partial_{x_i} (\sqrt{-1} c(\xi) |\xi|^{-4})(x_0) = \sqrt{-1} \frac{\partial_{x_i} c(\xi)}{|\xi|^4} (x_0) + \sqrt{-1} \frac{c(\xi) \partial_{x_i} (|\xi|^4)}{|\xi|^8} (x_0) = 0, \quad (4.12)$$

so  $\tilde{\Phi}_1 = 0$ .

**case a) II)**  $r = 1$ ,  $l = -3$ ,  $k = |\alpha| = 0$ ,  $j = 1$ .

By (4.2), we get

$$\tilde{\Phi}_2 = -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}) \times \partial_{\xi_n}^2 \sigma_{-3}(D_F^{-1})](x_0) d\xi_n \sigma(\xi') dx'. \quad (4.13)$$

By Lemma 4.3, we have

$$\partial_{\xi_n}^2 \sigma_{-3}(D_F^{-3})(x_0) = \partial_{\xi_n}^2 (c(\xi) |\xi|^{-4})(x_0) = \sqrt{-1} \frac{(20\xi_n^2 - 4)c(\xi') + 12(\xi_n^3 - \xi_n)c(dx_n)}{(1 + \xi_n^2)^4}, \quad (4.14)$$

and

$$\begin{aligned} \partial_{x_n} \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1})(x_0) &= \partial_{x_n} (-\sqrt{-1} \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l c(\xi) |\xi|^{-2}) \\ &= -\sqrt{-1} \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l \left[ \frac{\partial_{x_n} c(\xi')}{1 + \xi_n^2} - \frac{c(\xi) h'(0) |\xi'|^2}{(1 + \xi_n^2)^2} \right]. \end{aligned} \quad (4.15)$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case a) II)**. Then there is the following formula

$$\begin{aligned} &\text{Tr}[\partial_{x_n} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}) \times \partial_{\xi_n}^2 \sigma_{-3}(D_F^{-3})](x_0) \\ &= \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \frac{-8(3i\xi_n^2 + 2\xi_n - i)}{(\xi_n - i)^5 (\xi_n + i)^4} \\ &\quad + X_n Y_n h'(0) \left[ \frac{-16i(5\xi_n^2 - 1) + 48(\xi_n^3 - \xi_n)}{(\xi_n - i)^5 (\xi_n + i)^4} + \frac{8(5\xi_n^2 - 1) + 24i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right]. \end{aligned} \quad (4.16)$$

Therefore, we get

$$\begin{aligned} \tilde{\Phi}_2 &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left\{ \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \left[ \frac{-8(3i\xi_n^2 + 2\xi_n - i)}{(\xi_n - i)^5 (\xi_n + i)^4} \right] \right. \\ &\quad \left. + X_n Y_n h'(0) \left[ \frac{-16i(5\xi_n^2 - 1) + 48(\xi_n^3 - \xi_n)}{(\xi_n - i)^5 (\xi_n + i)^4} + \frac{8(5\xi_n^2 - 1) + 24i(\xi_n^3 - \xi_n)}{(\xi_n - i)^6 (\xi_n + i)^4} \right] \right\} d\xi_n \sigma(\xi') dx' \\ &= \left( \frac{5}{16} \sum_{j=1}^{n-1} X_j Y_j + \frac{1}{16} X_n Y_n \right) h'(0) \pi \Omega_3 dx', \end{aligned} \quad (4.17)$$

where  $\Omega_3$  is the canonical volume of  $S^2$ .

**case a) III)**  $r = 1, l = -3, j = |\alpha| = 0, k = 1$ .

By (4.2), we get

$$\begin{aligned}\tilde{\Phi}_3 &= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-3}(D_F^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\ &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}) \times \partial_{x_n} \sigma_{-3}(D_F^{-3})](x_0) d\xi_n \sigma(\xi') dx'.\end{aligned}\quad (4.18)$$

By Lemma 4.3, we have

$$\partial_{x_n} \sigma_{-3}(D_F^{-3})(x_0)|_{|\xi'|=1} = \frac{i \partial_{x_n} [c(\xi')]}{(1 + \xi_n^2)^4} - \frac{2ih'(0)c(\xi)|_{\xi'}|_{g^{\partial M}}^2}{(1 + \xi_n^2)^6}.\quad (4.19)$$

And we have

$$\begin{aligned}\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}) &= -\frac{c(\xi') + ic(dx_n)}{(\xi_n - i)^3} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l + \frac{c(\xi') + ic(dx_n)}{(\xi_n - i)^3} X_n Y_n \\ &\quad - \frac{ic(\xi') - c(dx_n)}{(\xi_n - i)^3} \sum_{j=1}^{n-1} X_j Y_n \xi_j - \frac{ic(\xi') - c(dx_n)}{(\xi_n - i)^3} \sum_{l=1}^{n-1} X_n Y_l \xi_l.\end{aligned}\quad (4.20)$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for computing **case a) III)**, then

$$\begin{aligned}&\text{Tr}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}) \times \partial_{\xi_n} \partial_{x_n} \sigma_{-3}(D_F^{-3})](x_0)|_{|\xi'|=1} \\ &= \left[ \frac{4i}{(\xi_n - i)^5 (\xi_n + i)^2} + \frac{16}{(\xi_n - i)^5 (\xi_n + i)^3} \right] h'(0) \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \\ &\quad - \left[ \frac{4i}{(\xi_n - i)^5 (\xi_n + i)^2} + \frac{16}{(\xi_n - i)^5 (\xi_n + i)^3} \right] h'(0) X_n Y_n.\end{aligned}\quad (4.21)$$

Therefore, we get

$$\begin{aligned}\tilde{\Phi}_3 &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \left( \left[ \frac{4i}{(\xi_n - i)^5 (\xi_n + i)^2} + \frac{16}{(\xi_n - i)^5 (\xi_n + i)^3} \right] h'(0) \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \right. \\ &\quad \left. - \left[ \frac{4i}{(\xi_n - i)^5 (\xi_n + i)^2} + \frac{16}{(\xi_n - i)^5 (\xi_n + i)^3} \right] h'(0) X_n Y_n \right) d\xi_n \sigma(\xi') dx' \\ &= \left( -\frac{25}{48} \sum_{j=1}^{n-1} X_j Y_j + \frac{25}{16} X_n Y_n \right) h'(0) \pi \Omega_3 dx'.\end{aligned}\quad (4.22)$$

**case b)**  $r = 0, l = -3, k = j = |\alpha| = 0$ .

By (4.2), we get

$$\tilde{\Phi}_4 = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\pi_{\xi_n}^+ \sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}) \times \partial_{\xi_n} \sigma_{-3}(D_F^{-3})](x_0) d\xi_n \sigma(\xi') dx'.\quad (4.23)$$

By Lemma 4.3, we obtain

$$\partial_{\xi_n} \sigma_{-3}(D_F^{-3})(x_0)|_{|\xi'|=1} = \frac{ic(dx_n)}{(1 + \xi_n^2)^2} - \frac{4\sqrt{-1}\xi_n c(\xi)}{(1 + \xi_n^2)^3}.\quad (4.24)$$

By Lemma 4.2, we have

$$\begin{aligned}
\sigma_0(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}) &= \sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-2}(D_F^{-1}) + \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-1}(D_F^{-1}) \\
&\quad + \sum_{j=1}^n \partial_{\xi_j} [\sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F)] D_{x_j} [\sigma_{-1}(D_F^{-1})] \\
&:= A + B + C.
\end{aligned} \tag{4.25}$$

(1) Explicit representation the first item of (4.25)

$$\begin{aligned}
\sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-2}(D_F^{-1})(x_0)|_{|\xi'|=1} &= - \sum_{j,l=1}^n X_j Y_l \xi_j \xi_l \sigma_{-2}(D_F^{-1})(x_0)|_{|\xi'|=1} \\
&= - \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \sigma_{-2}(D_F^{-1})(x_0)|_{|\xi'|=1} - X_n Y_n \xi_n^2 \sigma_{-2}(D_F^{-1})(x_0)|_{|\xi'|=1} \\
&\quad - \sum_{j=1}^{n-1} X_j Y_n \xi_j \xi_n \sigma_{-2}(D_F^{-1})(x_0)|_{|\xi'|=1} - \sum_{l=1}^{n-1} X_n Y_l \xi_n \xi_l \sigma_{-2}(D_F^{-1})(x_0)|_{|\xi'|=1},
\end{aligned} \tag{4.26}$$

we let

$$Q_1 = \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \sigma_{-2}(D_F^{-1})(x_0)|_{|\xi'|=1}, \quad Q_2 = X_n Y_n \xi_n^2 \sigma_{-2}(D_F^{-1})(x_0)|_{|\xi'|=1}. \tag{4.27}$$

By the trace identity  $\text{tr}(AB) = \text{tr}(BA)$ ,  $\text{tr}(A \otimes B) = \text{tr}(A) \cdot \text{tr}(B)$  and the relation of the Clifford action, we have

$$\text{Tr}[\sigma_0(D_F)c(h_q)] = 4 \sum_{i=1}^{p+q} \langle \nabla_{e_i}^{TM} e_i, h_q \rangle = -4 \sum_{i=1}^{n-1} \langle e_j, \nabla_{e_j}^{TM} \partial_{x_n} \rangle := -4 \text{div}_{\partial M}(\partial_{x_n}), \tag{4.28}$$

$$\text{Tr}[\sigma_0(D_F)c(\xi')] = 4 \sum_{i=1}^{p+q} \langle \nabla_{e_i}^{TM} e_i, \xi_i \rangle = 0. \tag{4.29}$$

Then

$$\begin{aligned}
&\text{Tr}[\pi_{\xi_n}^+ Q_1 \times \partial_{\xi_n} \sigma_{-3}(D_F^{-3})]|_{|\xi'|=1} \\
&= - \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \left[ \frac{3\xi_n^3 - 6i\xi_n^2 - 5\xi_n + 2i}{(\xi_n - i)^5(\xi_n + i)^3} - \frac{3\xi_n^4 - 9i\xi_n^3 - 17\xi_n^2 + 15i\xi_n + 4}{(\xi_n - i)^6(\xi_n + i)^3} \right] \\
&\quad - \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \frac{3\xi_n - i}{2(\xi_n - i)^4(\xi_n + i)^3} \text{Tr}[i\sigma_0(D_F)c(dx_n) + \sigma_0(D_F)c(\xi')] \\
&= -2 \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \frac{3\xi_n - i}{(\xi_n - i)^5(\xi_n + i)^3} + 2 \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \text{div}_{\partial M}(\partial_{x_n}) \frac{3i\xi_n + 1}{(\xi_n - i)^4(\xi_n + i)^3},
\end{aligned} \tag{4.30}$$



and

$$\begin{aligned}
& \text{Tr}[\pi_{\xi_n}^+ Q_2 \times \partial_{\xi_n}(D_F^{-3})]_{|\xi'|=1} \\
&= -X_n Y_n h'(0) \left[ \frac{3\xi_n^2 - 5i\xi_n}{(\xi_n - i)^4(\xi_n + i)^3} + \frac{3\xi_n^4 + 6i\xi_n^3 + 11\xi_n^2 - 6i\xi_n}{(\xi_n - i)^6(\xi_n + i)^3} \right] \\
&\quad - X_n Y_n h'(0) \frac{6i\xi_n^2 + 5\xi_n - i}{2(\xi_n - i)^4(\xi_n + i)^3} \text{Tr}[i\sigma_0(D_F)c(dx_n) + \sigma_0(D_F)c(\xi')] \\
&= -X_n Y_n h'(0) \frac{6\xi_n^4 - 5i\xi_n^3 - 2\xi_n^2 - i\xi_n}{(\xi_n - i)^6(\xi_n + i)^3} - 2X_n Y_n h'(0) \text{div}_{\partial M}(\partial x_n) \frac{6\xi_n^2 - 5i\xi_n - 1}{(\xi_n - i)^4(\xi_n + i)^3}. \tag{4.31}
\end{aligned}$$

We have,

$$\begin{aligned}
& -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\pi_{\xi_n}^+(Q_1 + Q_2) \times \partial_{\xi_n} \sigma_{-3}(D_F^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\
&= - \left( \frac{5}{16} \sum_{j=1}^{n-1} X_j Y_j + \frac{5}{16} X_n Y_n \right) h'(0) \pi \Omega_3 dx' + \left( \frac{1}{3} \sum_{j=1}^{n-1} X_j Y_j + \frac{1}{2} X_n Y_n \right) \text{div}_{\partial M}(\partial x_n) h'(0) \pi \Omega_3 dx'. \tag{4.32}
\end{aligned}$$

(2) Explicit representation the second item of (4.25)

$$\begin{aligned}
& \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-1}(D_F^{-1})(x_0)_{|\xi'|=1} \\
&= \left[ \sqrt{-1} \sum_{j,l=1}^n X_j \frac{\partial Y_l}{\partial x_j} \xi_l + \sqrt{-1} \sum_j (M(Y) + N(Y) + A(Y)) X_j \xi_j \right. \\
&\quad \left. + \sqrt{-1} \sum_l (M(X) + N(X) + A(X)) Y_l \xi_l \right] \frac{\sqrt{-1} c(\xi)}{|\xi|^2}; \tag{4.33}
\end{aligned}$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , then

$$\begin{aligned}
& \text{Tr} \left( \pi_{\xi_n}^+ \left( \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-1}(D_F^{-1}) \right) \times \partial_{\xi_n} \sigma_{-3}(D^{-3}) \right) (x_0) \\
&= \frac{4(3\xi_n - i)}{(1 + \xi_n^2)^3} \sum_{j,l=1}^n X_j \frac{\partial Y_l}{\partial x_j} \xi_l, \tag{4.34}
\end{aligned}$$

so we have

$$\begin{aligned}
& -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr}[\pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F) \sigma_{-1}(D^{-1}) \times \partial_{\xi_n} \sigma_{-3}(D^{-3})](x_0) d\xi_n \sigma(\xi') dx' \\
&= \frac{3i}{2} X(Y_n) \Omega_3 dx', \tag{4.35}
\end{aligned}$$

where  $X(Y_n)$  is  $\sum_{j=1}^n X_j \frac{\partial Y_n}{\partial x_j}$ .

(3) Explicit representation the third item of (4.25)

$$\sum_{j=1}^n \sum_{|\alpha|=1} \frac{1}{\alpha!} \partial_{\xi}^{|\alpha|} [\sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F)] D_{x_j} [\sigma_{-1}(D_F^{-1})](x_0)_{|\xi'|=1} = \sum_{j=1}^n \sum_{l=1}^n \sqrt{-1} (x_j Y_l + x_l Y_j) \xi_l \partial_{x_j} \left( \frac{\sqrt{-1} c(\xi)}{|\xi|^2} \right). \tag{4.36}$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , then

$$\begin{aligned}
& \text{Tr} \left( \pi_{\xi_n}^+ \left( \sum_{j=1}^n \sum_{|\alpha|=1} \frac{1}{\alpha!} \partial_{\xi}^{|\alpha|} [\sigma_2(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F)] D_{x_j} [\sigma_{-1}(D_F^{-1})] \right) \times \partial_{\xi_n} \sigma_{-3}(D^{-3}) \right) (x_0)_{|\xi'|=1} \\
&= 4X_n Y_n h'(0) \frac{3\xi_n^2 - i\xi_n}{(\xi_n - i)^4(\xi_n + i)^3}, \tag{4.37}
\end{aligned}$$

Substituting (4.37) into (4.23) yields

$$\begin{aligned}
& -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr} \left[ \pi_{\xi_n}^+ \left( \sum_{j=1}^n \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} [\sigma_2(\tilde{\nabla}_X \tilde{\nabla}_Y)] D_x^{\alpha} [\sigma_{-1}(D^{-1})] \right) \times \partial_{\xi_n} \sigma_{-3}(D^{-3}) \right] (x_0) d\xi_n \sigma(\xi') dx' \\
& = \frac{1}{2} X_n Y_n \pi h'(0) \Omega_3 dx'.
\end{aligned} \tag{4.38}$$

Summing up (1), (2) and (3) leads to the desired equality

$$\begin{aligned}
\tilde{\Phi}_4 & = \left( -\frac{5}{16} \sum_{j=1}^{n-1} X_j Y_j + \frac{3}{16} X_n Y_n \right) h'(0) \pi \Omega_3 dx' + \frac{3i}{2} X(Y_n) \Omega_3 dx' \\
& + \left( \frac{1}{3} \sum_{j=1}^{n-1} X_j Y_j + \frac{1}{2} X_n Y_n \right) \text{div}_{\partial M}(\partial x_n) h'(0) \pi \Omega_3 dx'.
\end{aligned} \tag{4.39}$$

**case c)**  $r = 1$ ,  $\ell = -4$ ,  $k = j = |\alpha| = 0$ .

By (4.2), we get

$$\begin{aligned}
\tilde{\Phi}_5 & = - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr} [\pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}) \times \partial_{\xi_n} \sigma_{-4} D_F^{-3}] (x_0) d\xi_n \sigma(\xi') dx' \\
& = \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{Tr} [\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1}) \times \sigma_{-4} D_F^{-3}] (x_0) d\xi_n \sigma(\xi') dx'.
\end{aligned} \tag{4.40}$$

By Lemma 4.3, we have

$$\begin{aligned}
\sigma_{-4}(D_F^{-3})(x_0)|_{|\xi'|=1} & = \frac{c(\xi) \sigma_2(D_F^3) c(\xi)}{(\xi_n^2 + 1)^4} + \frac{ic(\xi)}{(\xi_n^2 + 1)^4} [|\xi|^4 c(dx_n) \partial_{x_n} c(\xi') - 2h'(0) c(dx_n) c(\xi) \\
& + 2\xi_n c(\xi) \partial_{x_n} c(\xi') + 4\xi_n h'(0)],
\end{aligned} \tag{4.41}$$

where

$$\begin{aligned}
\sigma_2(D_F^3) & = \sum_{i,j,l} c(d_{x_i}) (\partial_l g^{ij}) + 2c(\xi) [2\sigma^i \otimes \text{Id}_{\wedge(F^{\perp,*})} + \text{Id}_{S(F)} \otimes 2\tilde{\sigma}^k - \Gamma^k + 2\tilde{S}(\partial_i)] \xi_k \\
& + [M(X) + N(X)] |\xi|^2,
\end{aligned} \tag{4.42}$$

and

$$\begin{aligned}
\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_1(\tilde{\nabla}_X^F \tilde{\nabla}_Y^F D_F^{-1})(x_0)|_{|\xi'|=1} & = \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2} \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l - \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2} X_n Y_n \\
& + \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)^2} \sum_{j=1}^{n-1} X_j Y_n \xi_j + \frac{ic(\xi') - c(dx_n)}{2(\xi_n - i)^2} \sum_{l=1}^{n-1} X_n Y_l \xi_l.
\end{aligned} \tag{4.43}$$

We note that  $i < n$ ,  $\int_{|\xi'|=1} \xi_{i_1} \xi_{i_2} \cdots \xi_{i_{2d+1}} \sigma(\xi') = 0$ , so we omit some items that have no contribution for

computing **case c**). Also, straightforward computations yield

$$\begin{aligned}
& \text{tr} \left[ \partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1} (\widetilde{\nabla}_X^F \widetilde{\nabla}_Y^F D_F^{-1}) \times \sigma_{-4} (D_F^{-3}) \right] (x_0) |_{|\xi'|=1} \\
&= \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l h'(0) \left[ \frac{2i\xi_n + 2}{(\xi_n - i)^4 (\xi_n + i)^2} + \frac{(24 + 4i)\xi_n^2 + (4 - 32i)\xi_n - 8}{(\xi_n - i)^6 (\xi_n + i)^4} - \frac{2\xi_n + 2i\xi_n^2}{(\xi_n - i)^5 (\xi_n + i)^3} \right] \\
&\quad - X_n Y_n h'(0) \left[ \frac{2i\xi_n + 2}{(\xi_n - i)^4 (\xi_n + i)^2} + \frac{(24 + 4i)\xi_n^2 + (4 - 32i)\xi_n - 8}{(\xi_n - i)^6 (\xi_n + i)^4} - \frac{2\xi_n + 2i\xi_n^2}{(\xi_n - i)^5 (\xi_n + i)^3} \right] \\
&\quad + \sum_{j,l=1}^{n-1} X_j Y_l \xi_j \xi_l \xi_k \xi_t g(dx^t, h_k) \left[ \frac{2(i-1)}{(\xi_n - i)^5 (\xi_n + i)^3} + \frac{(i-1)}{4(\xi_n - i)^4 (\xi_n + i)^2} \right] h'(0) \\
&\quad + X_n Y_n \xi_k \xi_t g(dx^t, h_k) \left[ \frac{2(1-i)}{(\xi_n - i)^5 (\xi_n + i)^3} + \frac{(1-i)}{4(\xi_n - i)^4 (\xi_n + i)^2} \right] h'(0). \tag{4.44}
\end{aligned}$$

We get

$$\widetilde{\Phi}_5 = \left[ \frac{129 - 44i}{320} \sum_{j=1}^{n-1} X_j Y_j - \frac{245 - 26i}{96} X_n Y_n \right] h'(0) \pi \Omega_3 dx'. \tag{4.45}$$

Let  $X = X^T + X_n \partial_n$ ,  $Y = Y^T + Y_n \partial_n$ , then we have  $\sum_{j=1}^{n-1} X_j Y_j(x_0) = g(X^T, Y^T)(x_0)$ . Now  $\Phi$  is the sum of the cases (a), (b) and (c). Combining with the five cases, this yields

$$\begin{aligned}
\widetilde{\Phi} = \sum_{i=1}^5 \widetilde{\Phi}_i &= - \left[ \frac{113 + 132i}{960} g(X^T, Y^T) + \frac{71 - 26i}{96} X_n Y_n \right] h'(0) \pi \Omega_3 dx' + \frac{3i}{2} X(Y_n) \Omega_3 dx' \\
&\quad + \left[ \frac{1}{3} g(X^T, Y^T) + \frac{1}{2} X_n Y_n \right] \text{div}_{\partial M}(\partial x_n) h'(0) \pi \Omega_3 dx'. \tag{4.46}
\end{aligned}$$

So, we are reduced to prove the following.

**Theorem 4.4.** *Let  $M$  be a 4-dimensional compact filiated manifold with boundary and spin leave,  $\widetilde{\nabla}$  be an orthogonal connection with torsion. Then we get the spectral Einstein functional associated to  $\widetilde{\nabla}_X^F \widetilde{\nabla}_Y^F D_F^{-1}$  and  $D_F^{-3}$  on  $M$*

$$\begin{aligned}
& \widetilde{\text{Wres}}[\pi^+(\widetilde{\nabla}_X^F \widetilde{\nabla}_Y^F D_F^{-1}) \circ \pi^+(D_F^{-3})] \\
&= \frac{8\pi^2}{3} \int_M G(X, Y) d\text{vol}_g + \int_M s g(X, Y) d\text{vol}_M \\
&\quad - \int_{\partial M} \left[ \frac{113 + 132i}{960} g(X^T, Y^T) + \frac{71 - 26i}{96} X_n Y_n \right] h'(0) \pi \Omega_3 dx' + \int_{\partial M} \frac{3i}{2} X(Y_n) \Omega_3 dx' \\
&\quad + \int_{\partial M} \left[ \frac{1}{3} g(X^T, Y^T) + \frac{1}{2} X_n Y_n \right] \text{div}_{\partial M}(\partial x_n) h'(0) \pi \Omega_3 dx'. \tag{4.47}
\end{aligned}$$

where  $s$  is the scalar curvature.

## Acknowledgements

This work is supported by the National Natural Science Foundation of China 11771070, 12061078 and the Youth Scientific Research Project of Yili Normal University 2023YSQN002. The authors thank the referee for his (or her) careful reading and helpful comments.

## References

- [1] M. Wodzicki.: Non-commutative residue I, Lecture Notes in Math., Springer, New York, Vol. 1289 320-399, (1987).
- [2] V. W. Guillemin.: A new proof of Weyl's formula on the asymptotic distribution of eigenvalues. *Adv. Math.* 55, no. 2, 131-160, (1985).
- [3] Dabrowski L, Sitarz A, Zalecki P. Spectral Metric and Einstein Functionals. *Advances in Mathematics*, 427, 109128, (2023).
- [4] P. B. Gilkey.: Invariance theory, the Heat equation, and the Atiyah-Singer Index theorem. Inc., USA, (1984).
- [5] H. Figueroa, J. Gracia-Bondía, and J. Várilly.: *Elements of Noncommutative Geometry*. Birkhäuser Boston, (2001).
- [6] A. Connes.: Quantized calculus and applications. XIth International Congress of Mathematical Physics(Paris,1994), Internat Press, Cambridge, MA, 15-36, (1995).
- [7] A. Connes.: The action functional in Noncommutative geometry. *Comm. Math. Phys.* 117, 673-683, (1988).
- [8] A. Connes, J. Lott.: Particle models and Non-commutative geometry. *Nucl. Phys. B Proc.Supp* 18B, 29-47, (1990).
- [9] D. Kastler.: The Dirac Operator and Gravitation. *Comm. Math. Phys.* 166, 633-643, (1995).
- [10] W. Kalau and M. Walze.: Gravity, Noncommutative geometry and the Wodzicki residue. *J. Geom. Physics.* 16, 327-344,(1995).
- [11] B. V. Fedosov, F. Gölse, E. Leichtnam, E. Schrohe.: The noncommutative residue for manifolds with boundary. *J. Funct. Anal.* 142, 1-31, (1996).
- [12] Y. Wang.: Differential forms and the Wodzicki residue for Manifolds with Boundary. *J. Geom. Physics.* 56, 731-753, (2006).
- [13] Y. Wang.: Gravity and the Noncommutative Residue for Manifolds with Boundary. *Letters in Mathematical Physics.* 80, 37-56, (2007).
- [14] Y. Wang.: Lower-Dimensional Volumes and Kastler-kalau-Walze Type Theorem for Manifolds with Boundary . *Commun. Theor. Phys.* Vol 54, 38-42, (2010).
- [15] J. M. Bismut.: A local index theorem for non Kähler manifolds. *Math. Ann.* 284, 681-699 (1989).
- [16] T. Ackermann and J. Tolksdorf.: A generalized Lichnerowicz formula, the Wodzicki residue and gravity. *J. Geom. Physics.* 19, 143-150,(1996).
- [17] F. Pfäffle and C. A. Stephan.: On gravity, torsion and the spectral action principle. *J. Funct. Anal.* 262, 1529-1565,(2012).
- [18] W. Greub, S. Halperin and R.Vanstone.: *Connections, Curvature and Cohomology, Vo.1* (Academic press, New York) (1976).
- [19] J. Wang and Y. Wang.: Noncommutative residue and sub-Dirac operators, *J. Math. Phys.* 54, 012501 (2013).
- [20] J. Wang, Y. Wang, T. Wu, Dirac operators with torsion, spectral Einstein functionals and the noncommutative residue. *J. Math. Phys.* 64, 102505(2023).
- [21] Y. Yu.: *The Index Theorem and The Heat Equation Method*, Nankai Tracts in Mathematics-Vol.2, World Scientific Publishing, (2001).
- [22] G. Grubb, E. Schrohe.: Trace expansions and the noncommutative residue for manifolds with boundary. *J. Reine Angew. Math.* 536, 167-207, (2001).
- [23] S. Alexandrov and D. Vassilevich.: Heat kernel for nonminimal operators on a Kähler manifold. *J. Math. Phys.* 37, 5715-5718, (1996).
- [24] B. Opozda.: Bochner- $\bar{\partial}$ 's technique for statistical structures. *Ann. Glob. Anal. Geom.* 48(4):357-395, (2015).
- [25] K. Liu, W. Zhang, Adiabatic limits and foliations. In: *Topology, Geometry, and Algebra: Interactions and New Directions*, Contemp Math, Vol. 279. Providence, RI: American Mathematical Society, 195C208, (2001).
- [26] K. Liu ,Y. Wang.: Adiabatic limits, vanishing theorems and the noncommutative residue. *Science in China Series A: Mathematics.* 52,12, 2699-2713,(2009).