

Identifying codes in graphs of given maximum degree. II. Triangle-free graphs*

Dipayan Chakraborty^{†‡} Florent Foucaud[†] Michael A. Henning[‡]
 Tuomo Lehtilä^{§¶†**}

March 27, 2024

Abstract

An *identifying code* of a closed-twin-free graph G is a set S of vertices of G such that any two vertices in G have a distinct intersection between their closed neighborhood and S . It was conjectured that there exists a constant c such that for every connected closed-twin-free graph G of order n and maximum degree Δ , the graph G admits an identifying code of size at most $(\frac{\Delta-1}{\Delta})n + c$. In a companion paper, we proved the conjecture for all trees. We show that the conjecture holds for all triangle-free graphs, with the same lists of exceptional graphs needing $c > 0$ as for trees: for $\Delta \geq 3$, $c = 1/3$ suffices and there is only a set of 12 trees requiring $c > 0$ for $\Delta = 3$, and when $\Delta \geq 4$ this set is reduced to the Δ -star only. Our proof is by induction, whose starting point is the above result for trees. Along the way, we prove a generalized version of Bondy's theorem on induced subsets [J. A. Bondy. Induced subsets. Journal of Combinatorial Theory, Series B, 1972] that we use as a tool in our proofs. We also use our main result for triangle-free graphs, to prove the upper bound $(\frac{\Delta-1}{\Delta})n + 1/\Delta + 4t$ for graphs that can be made triangle-free by the removal of t edges.

1 Introduction

In this article, we consider simple undirected loopless graphs. The *open neighborhood* of vertex u in graph G , $N_G(u)$, contains every vertex adjacent to vertex u while the *closed neighborhood* also contains vertex u itself. A set of vertices $D \subseteq V(G)$ is a *dominating set* if every vertex outside of D is adjacent to a vertex in D . A set $S \subseteq V(G)$ is a *separating set* if every vertex $u \in V(G)$ has a unique closed neighborhood in set S , that is, intersection $N_G[u] \cap S$ is unique for every vertex. Furthermore, a set $C \subseteq V(G)$ is an *identifying code* if it is a dominating set and a separating set [25]. See Figure 1 for examples of identifying codes. From the intuitive perspective, identifying codes allow us to locate or identify any vertex if we know which vertices in the set S are in its closed neighborhood. It is natural to ask what is the minimum number of vertices in an identifying code of graph G . This value is called the *identification number* of graph G and it is denoted by $\gamma^{\text{ID}}(G)$.

This paper concentrates on proving Conjecture 1 (stated below) on upper bounds for the minimum size of identifying codes of given maximum degree, for all triangle-free graphs. Previously, in the companion

*The two first authors were supported by the French government IDEX-ISITE initiative CAP 20-25 (ANR-16-IDEX-0001), the International Research Center "Innovation Transportation and Production Systems" of the I-SITE CAP 20-25, and the ANR project GRALMECO (ANR-21-CE48-0004). The research of Michael Henning was supported in part by the South African National Research Foundation (grant number 129265) and the University of Johannesburg.

[†]Université Clermont Auvergne, CNRS, Clermont Auvergne INP, Mines Saint-Etienne, LIMOS, 63000 Clermont-Ferrand, France.

[‡]Department of Mathematics and Applied Mathematics, University of Johannesburg, South Africa.

[§]University of Helsinki, Department of Computer Science, Helsinki, Finland.

[¶]Helsinki Institute for Information Technology (HIIT), Espoo, Finland.

[‡]University of Turku, Department of Mathematics and Statistics, Turku, Finland.

**Research partially supported by Business Finland project 10769/31/2022 6GTNF and Academy of Finland grant 338797.

paper [5], we have proved the conjecture for trees. Although the current paper is part II and builds on top of part I, both papers are designed so that they can be read independently.

A thorough treatise on domination in graphs can be found in [21, 22, 23]. Bounds on domination numbers for graphs with restrictions on their degree parameters are a natural and important line of research, see for example the influential result by Reed [31] that if G is a connected cubic graph of order n , then its domination number is at most $\frac{3}{8}n$. A detailed discussion on upper bounds on the domination number in terms of its order and degree parameters, as well as bounds with specific structural restrictions imposed, can be found in [23, Chapters 6, 7 and 10].

Originally, identifying codes were motivated by fault-detection in multiprocessor networks [25]. Numerous other applications have been discovered, such as threat location in facilities using sensor networks [34], logical definability of graphs [28] and canonical labeling of graphs for the graph isomorphism problem [1]. Moreover, multiple related concepts have been introduced since the 1960s, such as *separating systems*, *test covers* and *locating-dominating sets*, which have been independently discovered and studied, forming the general area of identification problems in graphs and other discrete structures. See for example [4, 27, 32, 29]. An extensive internet bibliography containing over 500 articles around these topics can be found at [8], while more information specifically about identifying codes can be found in the book chapter [26].

In this paper, we concentrate on connected identifiable triangle-free graphs. A graph is *identifiable* if it does not contain any *closed twins*, that is, vertices with the same closed neighborhoods. Moreover, a graph is *triangle-free* if no three vertices in it form a cycle. Similarly as closed twins, we define *open twins* as vertices with the same open neighborhoods. A graph without closed or open twins is called *twin-free*. We also define similarly *open-twin-free* and *closed-twin-free* graphs. Twins are significant for separating sets and identifying codes; if two vertices are closed twins, then the graph does not admit any separating set and hence no identifying code. Moreover, if t vertices have the same open neighborhood, then any separating set and hence also any identifying code, contains at least $t - 1$ of them. Note that every connected triangle-free graph on at least three vertices is identifiable.

As our main result, we prove the following conjecture for all identifiable triangle-free graphs.

Conjecture 1 ([13, Conjecture 1]). There exists a constant c such that for every connected identifiable graph on n vertices and of maximum degree $\Delta \geq 2$,

$$\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta - 1}{\Delta} \right) n + c.$$

For an easier presentation of the entire proof, we have divided the work into two parts. The current paper is the second part of that work. Although it can also be read independently. In the companion paper [5], we have proved Conjecture 1 for trees and have determined the exact set of trees requiring a positive constant c together with an exact value of c . In this part, we use our previous results for trees from the companion paper as a starting point for our proof of Conjecture 1 for all triangle-free graphs.

It is known that if true, the conjecture would be tight, that is, some graphs of maximum degree Δ have identification number at least $\frac{\Delta-1}{\Delta}n$. For $\Delta = 2$, the conjecture is tight for both paths and cycles with $c \leq 3/2$ (see corollary 7). For $\Delta = 3$, the conjecture is tight for example for trees presented in Figure 1 and for a path whose every vertex we join to a 2-path by a single edge (see [5]). For any $\Delta > 3$, the complete bipartite graph $K_{\Delta, \Delta}$ satisfies $\gamma^{\text{ID}}(K_{\Delta, \Delta}) = 2\Delta - 2 = \left(\frac{\Delta-1}{\Delta} \right) n$ and hence gives tight examples with $c = 0$. Furthermore, for any $\Delta > 3$ and an unbounded value of n , there are trees with identification number $\gamma^{\text{ID}}(T) > \frac{(\Delta-1)n}{\Delta} - \frac{n}{\Delta^2}$ [5]. Furthermore, there exist connected graphs of any maximum degree $\Delta \geq 3$ and arbitrarily large number n of vertices, with identification number $\left(\frac{\Delta-1}{\Delta} \right) n$, see [15, 17].

The condition on the maximum degree Δ is a necessary part of the conjecture. Without it, there are graphs on n vertices with identification number $n - 1$ [12, 18].

The best known general upper bounds for connected graphs with a maximum degree Δ and number of vertices n , when n is large enough, are of the form $n - \frac{n}{\Theta(\Delta^5)}$ [12] which has been improved to $n - \frac{n}{103(\Delta+1)^3}$ in [15] (for the sake of comparison, the conjectured bound can be rewritten as $n - \frac{n}{\Delta} + c$). When we

consider graph classes instead of general graphs, some improvements on these bounds are known. If every pair of closed neighborhoods in a graph differ by at least two vertices, then the general bound has been improved to $n - \frac{n}{103\Delta}$ and to $n - \frac{n}{f(k)\Delta}$ for graphs of clique number k [15]. For bipartite graphs, an upper bound of $n - \frac{n}{\Delta+9}$ has been proved [13]. In a short conference proceedings paper [6], we sketched a proof for Conjecture 1 for bipartite graphs without twins of degree 2 or greater (we did not include that proof in the current article, since the triangle-free result we present here is stronger). Moreover, in Part I [5], we gave the proof for all trees. Conjecture 1 also holds for line graphs of graphs of average degree at least 5 [11, Corollary 21] as well as graphs which have girth at least 5, minimum degree 2 and maximum degree at least 4 [2]. Furthermore, the conjecture holds in many cases for some graph products such as Cartesian and direct products [16, 24, 30]. See also the book chapter [26], where Conjecture 1 is presented.

Conjecture 1 has been considered also for triangle-free graphs previously. In [13], an upper bound of type $n - \frac{n}{\Delta+o(\Delta)}$ was presented for triangle-free graphs. When the triangle-free graph is also twin-free, this upper bound has been improved to $n - \frac{n}{3\Delta/(\ln \Delta - 1)}$. Note that the latter result implies that Conjecture 1 holds for triangle-free graphs without any open twins, whenever $\Delta \geq 55$ (because then, $3\Delta/(\ln \Delta - 1) \leq \Delta$). Note that the graphs containing open twins seem to be the toughest cases regarding Conjecture 1 (among triangle-free graphs). Indeed, we will see that every triangle-free graph requiring a positive constant c with $\Delta \geq 3$ contains open twins. Furthermore, in [14] it has been shown that every *twin-free* bipartite graph G on $n \geq 5$ vertices satisfies the upper bound $\gamma^{\text{ID}}(G) \leq \frac{2n}{3}$, while if we allow open twins, then there exist trees T with arbitrarily large numbers of vertices such that $\gamma^{\text{ID}}(T) > (\frac{\Delta-1}{\Delta})n - \frac{n}{\Delta^2}$ [5].

Our main result is to prove Conjecture 1 (in a strong form) for all triangle-free graphs. To state it, we define, for every integer $\Delta \geq 3$, a set \mathcal{F}_Δ of exceptional graphs of maximum degree at most Δ (see Section 2.3 for greater detail). For $\Delta = 3$, this set contains twelve trees (see Figure 1), the cycles on 4 and 7 vertices, and the path on 4 vertices. For every integer $\Delta > 3$, it contains exactly the Δ -star $K_{1,\Delta}$.

Theorem 2. Let $\Delta \geq 3$ be an integer, and let G be a connected triangle-free graph of order $n \geq 3$. If $G \in \mathcal{F}_\Delta$, then

$$\gamma^{\text{ID}}(G) = \left(\frac{\Delta-1}{\Delta}\right)n + \frac{1}{\Delta}.$$

On the other hand, if $G \notin \mathcal{F}_\Delta$ has maximum degree Δ , then

$$\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n.$$

Note that graphs in \mathcal{F}_Δ for $\Delta \geq 4$ have maximum degree Δ and the graphs in \mathcal{F}_3 have maximum degree either two or three.

In the companion paper [5], we have shown that Conjecture 1 holds for trees (see Theorem 12). It turns out that, for triangle-free graphs with maximum degree at least 3, the set of graphs requiring a positive constant c is exactly the same as the set of trees with maximum degree at least 3 needing a positive constant. Our proof uses the result for trees from [5] as a starting point for an induction. After that, we assume that a triangle-free graph contains at least one cycle containing some edge e . We remove that edge to construct a graph G' which, by induction, satisfies the conjecture. Hence, G' contains a *small* identifying code which we can use to construct another identifying code of the same size for G . One difficulty for proving the conjecture is the existence of the set of graphs requiring $c > 0$. Since \mathcal{F}_3 is the largest among the sets \mathcal{F}_Δ , the case $\Delta = 3$ requires a lot of special argumentation.

Structure of the paper. First in Section 2, we introduce some useful definitions and lemmas. In Subsection 2.1, we introduce terminology and results about a generalization of identifying codes, which are later used in the proof of our main result. We continue in Subsection 2.2 where we discuss Conjecture 1 when $\Delta = 2$. In Subsection 2.3, we introduce every connected triangle-free graph which requires a positive constant c for Conjecture 1. Understanding these graphs is crucial for the proof of the main theorem. In Section 3, we give the proof of Theorem 2. In Section 4, we use our bound to prove a weaker bound for graphs that have triangles, but can be made triangle-free by removing t edges. Finally, we conclude in Section 5.

2 Preliminaries and known results

In the following, we go through the notation we use throughout this article. We denote by $V(G)$ and $E(G)$ the vertex and edge sets of graph $G = (V(G), E(G))$. We usually denote $n = |V(G)|$. For a set of vertices S we denote $N(S) = \bigcup_{v \in S} N(v)$ and $N[S] = \bigcup_{v \in S} N[v]$. We denote by $\deg_G(v) = |N_G(v)|$ the *degree* of the vertex v in graph G . A *leaf* is a vertex with degree one and its only neighbor is called *support vertex*. In the literature, a leaf is also known as a *pendant vertex*. Naturally, any vertex of a graph G that is not a leaf of G is referred to as a *non-leaf* vertex of G . We denote the complement of a graph G by \overline{G} . We sometimes denote the maximum degree of graph G by $\Delta(G)$, its number of vertices by $n(G)$ and its number of edges by $m(G)$. The *girth* of the graph G refers to the number of vertices in the shortest cycle in graph G .

We say that vertex u (or vertex subset C) *separates* vertices u and v if vertex u is in exactly one of sets $N[u]$ and $N[v]$. Given a vertex subset C (often an identifying code), with *code neighborhood* of vertex $u \in V(G)$ we refer to the set $C \cap N[u]$.

On many occasions throughout this article, we look at a subgraph of a graph G obtained by deleting some vertices or edges. To that end, given a graph G and a set S containing some vertices and edges of G , we define $G - S$ as the subgraph of G obtained by deleting from G all vertices (and edges incident with them) and edges of G in S .

We use following lemma multiple times for arguing that some vertices are identified.

Lemma 3. Let G be a triangle-free graph and S , a subset of vertices of G . If three vertices u, v, w inducing a P_3 are in S , then each of them has a unique closed neighborhood in S .

Proof. Let $u, v, w \in S \subseteq V(G)$ induce a P_3 in G where v is the middle vertex of the path. Suppose first on the contrary that $N[u] \cap S = N[x] \cap S$ for some $x \in V(G) \setminus \{u\}$. However, now vertices x, u and v form a triangle or if $x = v$, then x, u and w form a triangle. The case with w is symmetric. Suppose then that $N[v] \cap S = N[x] \cap S$ for some $x \in V(G) \setminus \{v\}$. Now, v, w and x form a triangle or if $x = w$, then u, v and w form the triangle. Hence, the claim follows since G is triangle-free. \square

2.1 (X, Y) -separating codes and (X, Y) -identifying codes

We now introduce a generalization of identifying codes, and an upper bound for them that generalizes Bondy's theorem on induced subsets [4], that will be used several times in our proofs and, we believe, can be useful in many other settings as well.

Let $G = (V, E)$ be a graph and X and Y be two (not necessarily disjoint) vertex subsets of G . Then, Y induces a partition on X by the equivalence relation \sim defined by $u \sim v$ if and only if $N_G[u] \cap Y = N_G[v] \cap Y$ for any $(u, v) \in X \times X$. If the partition on X induced by Y is such that each part is a singleton set, that is, for each pair $u, v \in X$ there exists a vertex $w \in Y$ that separates u, v , then Y is called an (X, Y) -*separating set* in G and the set X is called Y -*separable*. Moreover, we call any vertex subset C of G an (X, Y) -*identifying code* of G if 1) C is an (X, Y) -separating set in G , and 2) C is a dominating set of X . If such an (X, Y) -identifying code of G exists, then the set X is called Y -*identifiable*. Notice that, if X is Y -identifiable and C is an (X, Y) -identifying code of G , then Y itself is an (X, Y) -identifying code of G . Furthermore, set X is also C -identifiable and C is an (X, C) -identifying code of G . In particular, if G is an identifiable graph and C is an identifying code of G , then the vertex set V is V -identifiable and C -identifiable and C is a (V, V) -identifying code and a (V, C) -identifying code of G . When X and Y are disjoint and induce a bipartite graph, an (X, Y) -identifying code has been called a *discriminating code* in the literature [7].

Lemma 4. Let G be a graph with vertex subsets X and Y such that X is Y -identifiable. Then, there is an (X, Y) -separating set in G of size at most $|X| - 1$, and an (X, Y) -identifying code of size at most $|X|$.

Proof. Assume that G is (X, Y) -separable. If $|X| = 1$, there is nothing to do. Otherwise, we inductively construct an (X, Y) -separating code C of G such that $|C| \leq |X| - 1$. To begin with, let u, v be an arbitrary pair of distinct vertices of X and let $c \in Y$ such that c separates the pair: c exists since G is

(X, Y) -separable. Then, we let $C = \{c\}$. Let \mathcal{P}_C be the partition induced by C on X , where two vertices of X are in the same part if and only if their closed neighbourhood in G intersects the same subset of C . Then, for as long as there exists a part P of \mathcal{P}_C such that $u', v' \in P$ for two distinct vertices u', v' of X , the construction of C follows inductively by selecting an element $c' \in Y$ such that c' separates u', v' , and letting $c' \in C$. At each step, since G is (X, Y) -separable, such c' exists. Moreover, we notice that at each inductive step, we have $|C| \leq |\mathcal{P}_C| - 1$, since at each step, we increase the number of parts by at least 1, and the size of C by exactly 1. This implies that we must have $|C| \leq |X| - 1$, since $|\mathcal{P}_C| \leq |X|$, affirming the first claim.

Now, if moreover G is (X, Y) -identifiable, we proceed as above to first build the (X, Y) -separating set C of G of size at most $|X| - 1$. Now, any two vertices of X are separated by C . Furthermore, there exists at most one vertex of X that is not dominated by C ; for otherwise, if there exist two distinct vertices $x, x' \in X$ not dominated by C , it would imply that $N_G[x] = N_G[x'] = \emptyset$ and so, C would not be an (X, Y) -separating set of X , a contradiction. Therefore, let $x \in X$ be not dominated by C (if such an x exists). Since Y dominates X , there exists $c'' \in N_G[x] \cap Y$. Then, we let $c'' \in C$, thus making C an (X, Y) -identifying code of G with $|C| \leq |X|$. This completes the proof. \square

We note that the core of Lemma 4 is similar to Bondy's celebrated theorem on "induced subsets" [4]. Indeed, in Bondy's theorem, one is given a set X of elements and a collection $\mathcal{A} = \{A_1, \dots, A_n\}$ of subsets of X ; it is proved by Bondy that there is a subset of at most $|X| - 1$ subsets of \mathcal{A} that form an (X, \mathcal{A}) -separating set, when viewing X and \mathcal{A} as the two partite sets of a bipartite graph (the incidence bipartite graph of the hypergraph (X, \mathcal{A})), provided this graph is \mathcal{A} -identifiable. Bondy's original proof uses an elegant graph-theoretic argument [4] (several proofs of algebraic nature have also been provided, see for example [35]). A similar statement, formulated in the language of graphs, is also proved by Gutin, Ramanujan, Reidl, and Wahlström in [20, Lemma 8], by an inductive argument similar to the one presented here. These prior results however are only concerned with separating sets (thus in their setting, one vertex may remain undominated), and with the special case where X and Y are disjoint. Hence, our result both generalizes and strenghtens these setups.

2.2 Paths and cycles

Our main result requires a precise understanding of graphs with $\Delta = 2$ and triangle-free graphs which need a positive constant c for Conjecture 1. Hence, in this subsection, we recall results on all connected graphs with $\Delta = 2$, that is, on paths and cycles. A path (cycle) on n vertices is denoted by P_n (C_n).

The identification number of all identifiable paths (that is, of all paths except P_2) was determined by Bertrand et al. [3]. Moreover, using an upper bound from [3] on even cycles of order at least 6, Gravier et al. [19] provided the exact values of the identification numbers of all identifiable cycles (that is, cycles of length at least 4). We summarize these results in the following theorems.

Theorem 5 ([3, Theorem 3]). If P_n is a path on n vertices, then we have

$$\gamma^{\text{ID}}(P_n) = \begin{cases} \frac{n}{2} + \frac{1}{2}, & \text{if } n \geq 1 \text{ is odd,} \\ \frac{n}{2} + 1, & \text{if } n \geq 4 \text{ is even.} \end{cases}$$

Theorem 6 ([19, Theorems 2 and 4]). If C_n is a cycle on n vertices, then we have

$$\gamma^{\text{ID}}(C_n) = \begin{cases} 3, & \text{if } n = 4, 5, \\ \frac{n}{2}, & \text{if } n \geq 6 \text{ is even,} \\ \frac{n}{2} + \frac{3}{2}, & \text{if } n \geq 7 \text{ is odd.} \end{cases}$$

Using Theorems 5 and 6, therefore, one has the following corollary.

Corollary 7. The following hold.

- (a) If G is a path, then $\gamma^{\text{ID}}(P_n) = \lfloor \frac{n}{2} \rfloor + 1$.
- (b) If $G = C_4$ or $G = C_5$, then $\gamma^{\text{ID}}(G) = \lfloor \frac{n}{2} \rfloor + 1$.
- (c) If $n \geq 6$ is even, then $\gamma^{\text{ID}}(C_n) = \frac{n}{2}$.
- (d) If $n \geq 7$ is odd, then $\gamma^{\text{ID}}(C_n) = \frac{n}{2} + \frac{3}{2}$.
- (e) If $n = 4$, then $\gamma^{\text{ID}}(P_n) = \frac{3}{4}n$, and if $n \geq 3$ and $n \neq 4$, then $\gamma^{\text{ID}}(P_n) \leq \frac{2}{3}n$.
- (f) If $n \in \{4, 7\}$, then $\frac{2}{3}n < \gamma^{\text{ID}}(C_n) \leq \frac{3}{4}n$, and if $n \geq 3$ and $n \notin \{4, 7\}$, then $\gamma^{\text{ID}}(C_n) \leq \frac{2}{3}n$.

We shall need the following elementary property of odd cycles with one edge added.

Observation 8 (Proposition 4.1 of [33]). If $n = 2k + 1 \geq 7$ is odd, $G = C_n$ and if $e \in E(\overline{G})$, then $\gamma^{\text{ID}}(G + e) \leq k + 1 \leq \frac{2}{3}n$.

2.3 Extremal triangle-free graphs

In this section, we define and discuss the exceptional triangle-free graphs of the statement of Theorem 2, that is, those in the set \mathcal{F}_Δ that require $c > 0$ in the bound of Conjecture 1. The notation for set \mathcal{T}_3 (already defined in the companion paper [5]) will be useful in our proof for the main theorem.

Definition 9. For $\Delta = 3$, we define $\mathcal{T}_3 = \{T_0, T_1, T_2, \dots, T_{11}\}$ to be the collection of 12 trees of maximum degree 3 as in Figure 1, and for $\Delta \geq 4$, we let $\mathcal{T}_\Delta = \{K_{1,\Delta}\}$.

For $\Delta = 3$, we let $\mathcal{F}_3 = \mathcal{T}_3 \cup \{P_4, C_4, C_7\}$ and for $\Delta \geq 4$, we let $\mathcal{F}_\Delta = \mathcal{T}_\Delta = \{K_{1,\Delta}\}$.

We note that $\gamma^{\text{ID}}(T_0) = 3$, $\gamma^{\text{ID}}(T_1) = \gamma^{\text{ID}}(T_2) = 5$, $\gamma^{\text{ID}}(T_3) = \gamma^{\text{ID}}(T_4) = \gamma^{\text{ID}}(T_5) = 7$, $\gamma^{\text{ID}}(T_6) = \gamma^{\text{ID}}(T_7) = 9$, $\gamma^{\text{ID}}(T_8) = \gamma^{\text{ID}}(T_9) = 11$, $\gamma^{\text{ID}}(T_{10}) = 13$, and $\gamma^{\text{ID}}(T_{11}) = 15$. Generally we have the following.

Proposition 10 ([5]). If $\Delta \geq 3$ is an integer and T is a tree of order n in \mathcal{T}_Δ , then $\gamma^{\text{ID}}(T) = \left(\frac{\Delta-1}{\Delta}\right)n + \frac{1}{\Delta}$.

By Corollary 7, if G has maximum degree $\Delta = 2$, then $\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n + \frac{3}{2}$. When $\Delta \geq 3$ we have the following.

Proposition 11. If $\Delta \geq 3$ is an integer and G is a graph of order n and maximum degree at most Δ in \mathcal{F}_Δ (possibly, if $\Delta = 3$, the maximum degree of G is 2), then $\gamma^{\text{ID}}(G) = \left(\frac{\Delta-1}{\Delta}\right)n + \frac{1}{\Delta}$.

Proof. If G is a tree in \mathcal{T}_Δ , then this follows from Proposition 10. Otherwise, $\Delta = 3$ and $G \in \{P_4, C_4, C_7\}$. If $G \in \{P_4, C_4\}$, by Corollary 7, $\gamma^{\text{ID}}(G) = 3 = \frac{2}{3}n + \frac{1}{3}$ and if $G = C_7$, $\gamma^{\text{ID}}(G) = 5 = \frac{2}{3}n + \frac{1}{3}$. \square

We have shown that trees satisfy Conjecture 1 in the companion paper [5], as follows.

Theorem 12 ([5]). If $T \notin \mathcal{T}_\Delta$ is a tree of order $n \geq 3$ with maximum degree $\Delta \geq 3$, then

$$\gamma^{\text{ID}}(T) \leq \left(\frac{\Delta-1}{\Delta}\right)n.$$

In the following proposition, we present some useful properties for the structure of identifying codes in trees in set \mathcal{T}_3 . The proof is provided in the companion paper [5].

Proposition 13 ([5]). If T is a tree in \mathcal{T}_3 , then the following properties hold.

- (i) If $T \neq T_2$, then T has an optimal identifying code $C(T)$ containing all vertices of degree at most 2.
- (ii) If $T \notin \{T_2, T_3\}$, then $C(T)$ can be chosen as an independent set. When we delete any code vertex v from T , set $C(T) \setminus \{v\}$ forms an optimal identifying code of the forest $T - v$.

We shall also need the following property of trees in the family \mathcal{T}_3 .

Lemma 14. Let $T \in \mathcal{T}_3$ be a tree of order n . If $e \in E(\overline{T})$ is such that $T + e$ is triangle-free and $\Delta(T + e) \leq 3$, then $\gamma^{\text{ID}}(T + e) < \frac{2}{3}n$.




Figure 1: The family \mathcal{T}_3 of trees of maximum degree 3 requiring $c > 0$ in Conjecture 1. The set of black vertices in each figure constitutes an identifying code of the tree.

Proof. Assume first that $T \notin \{T_2, T_3\}$. Let $C(T)$ be an optimal identifying code in T containing every vertex of degree at most 2 which is also an independent set. Such an identifying code exists by Proposition 13. Let us denote the end-points of edge e by u and v . Since $\Delta(T + e) = 3$, we have $u, v \in C(T)$. Observe that we may obtain another identifying code C' by shifting a single code vertex to any adjacent vertex. This is possible, since every vertex in $V(T) \setminus C(T)$ is dominated by exactly three vertices in $C(T)$. Thus, after shifting, the new code vertex belongs to a P_3 -component in $T[C']$ and is identified by Lemma 3. Furthermore, there now exists exactly one vertex w in $V(T) \setminus C(T)$ which is adjacent to one or two vertices in C' . Since the adjacent vertex in C' has at least three code vertices in its closed neighborhood, also vertex w is uniquely identified.

Let us now consider the shifted identifying code C' so that $u \notin C'$. By the previous considerations, C' is an identifying code in T . Furthermore, set C' is an identifying code also in $T + e$. Indeed, the only code neighborhood which is modified by the addition of edge e , is that of vertex u . we have $|N[u] \cap C'| \in \{2, 3\}$.

Note that u is the only vertex outside of C' which may have two code vertices in its closed neighborhood. Thus, if u is not identified, then we have $|N[u] \cap C'| = 3$. If $N[u] \cap C' = N[w] \cap C'$, then, by Lemma 3, we have $w \notin C'$ (otherwise u is identified by C'). Thus, there are at least two cycles in $T + e$, a contradiction.

Assume next that $T = T_2$. In this case, the identifying code depicted in Figure 1 is also an identifying code in $T + e$.

Finally, assume that $T = T_3$. In this case, the identifying code C depicted in Figure 1 is also an identifying code in $T + e$, unless the edge e is between two leaves at distance 4 in T . Let us call these two leaves u and v . Moreover, let u' be the leaf at distance 2 from u and let s be the support vertex adjacent to u and u' . Now, $\{s\} \cup C \setminus \{u'\}$ is an identifying code in $T + e$. \square

We have listed in Table 1 the graphs which are known (to us) to require a positive constant for Conjecture 1. Among these, are the extremal graphs discovered in [12] (those graphs of order n with identifying code number $n - 1$). Those are either stars, or can be built from any number of complements of half-graphs¹ by taking their complete join, and optionally, adding a single universal vertex. Note that the latter examples have large cliques and thus, are far from triangle-free. They have maximum degree $n - 1$ or $n - 2$, and so, they need $c = 1/\Delta$ or $c = 2/\Delta$ in the bound of Conjecture 1.

It is an interesting open question whether there exist any other such graphs and if the constant $c = \frac{3}{2}$ is enough for all graphs. Notice that by Theorem 2, if there exists a graph not listed in Table 1 that requires a positive constant c , then it must contain triangles.

Graph class	Δ	c	Reference
Odd paths	2	$1/\Delta = 1/2$	[3] (Theorem 5)
Even paths	2	$2/\Delta = 1$	[3] (Theorem 5)
Odd cycles C_n for $n \geq 7$	2	$3/\Delta = 3/2$	[19] (Theorem 6)
C_4	2	$2/\Delta = 1$	[19] (Theorem 6)
C_5	2	$1/\Delta = 1/2$	[19] (Theorem 6)
\mathcal{T}_3	3	$1/\Delta = 1/3$	[5]
$K_{1,\Delta}$	$\Delta \geq 3$	$1/\Delta$	[5]
Complements of half-graphs and their complete joins	even $\Delta \geq 2$	$2/\Delta$	[12]
Complements of half-graphs and their complete joins plus one universal vertex	odd $\Delta \geq 3$	$1/\Delta$	[12]

Table 1: Known graphs requiring a positive constant c for Conjecture 1.

3 Proof of the main result

In this section, we shall prove our main result, namely Theorem 2. Recall its statement.

Theorem 2. Let $\Delta \geq 3$ be an integer, and let G be a connected triangle-free graph of order $n \geq 3$. If $G \in \mathcal{F}_\Delta$, then

$$\gamma^{\text{ID}}(G) = \left(\frac{\Delta - 1}{\Delta} \right) n + \frac{1}{\Delta}.$$

On the other hand, if $G \notin \mathcal{F}_\Delta$ has maximum degree Δ , then

$$\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta - 1}{\Delta} \right) n.$$

¹A *half-graph* is a special bipartite graph with both parts of the same size, where each part can be ordered so that the open neighbourhoods of consecutive vertices differ by exactly one vertex [9]. Their complements can also be described as powers of paths of the form P_{2k}^{k-1} [12].

Proof of Theorem 2. The first part of the statement holds by Proposition 11.

For the second part, let G be a connected triangle-free graph of order n and size m with maximum degree $\Delta \geq 3$ such that $G \notin \mathcal{F}_\Delta$. Thus, $n \geq 5$. We proceed by induction on $n + m$ to show that $\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n$. Since G is connected, we note that $m \geq n - 1$, and so $n + m \geq 2n - 1 \geq 9$. If $n + m = 9$, then G is formed from a star by joining a pendant leaf to another leaf. By Theorem 12, we have $\gamma^{\text{ID}}(G) \leq \frac{2}{3}n$, in this case. Furthermore, if $n = m = 5$, then G is formed from a 4-cycle joined by a leaf. Observe that such a graph has $\gamma^{\text{ID}}(G) = 3 < \frac{2}{3}n$. This establishes the base cases. Let $n + m \geq 11$, where $n \geq 5$.

For the inductive hypothesis, assume that if G' is a connected triangle-free graph of order $n' \geq 3$ and size m' with $n' + m' < n + m$ and with maximum degree $\Delta' = \Delta(G') \geq 3$ such that $G' \notin \mathcal{F}_{\Delta'}$, then $\gamma^{\text{ID}}(G') \leq \left(\frac{\Delta'-1}{\Delta'}\right)n'$.

If $m = n - 1$, then G is a tree. Since $G \notin \mathcal{F}_\Delta$, by Theorem 12, we have $\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n$. Hence, we may assume that $m \geq n$, for otherwise the desired upper bound follows. Thus, the graph G contains a cycle edge, that is, an edge that belongs to a cycle in G . Moreover, since $\Delta \geq 3$, the graph G is not a cycle.

Among all cycle edges in G , let $e = uv$ be chosen so that the sum of the degrees of its ends is as large as possible, that is, $\deg_G(u) + \deg_G(v)$ is maximal. Since $\Delta(G) \geq 3$, we have for the edge e that

$$\deg_G(u) + \deg_G(v) \geq 5.$$

Let

$$G' = G - e.$$

Since e is a cycle edge of G , the graph G' is a connected triangle-free graph of order n . Let $\Delta(G') = \Delta'$, and so $\Delta' \geq \Delta - 1$. We note that $n(G') = n$ and $m(G') = m - 1$.

Suppose that $\Delta' = 2$. In this case, $\Delta = 3$ and G' is either a path or a cycle. Suppose firstly that G' is a cycle. By the triangle-free condition and our choice of the edge e , we infer that G is obtained from a cycle C_n where $n \geq 6$ by adding a chord between two non-consecutive vertices on the cycle in such a way as to create two cycles that both contain the edge e and both have length at least 4. Assume first that $n \neq 7$. By Corollary 7(f), $\gamma^{\text{ID}}(G') \leq \frac{2}{3}n$. By Observation 8, $\gamma^{\text{ID}}(G) \leq \gamma^{\text{ID}}(G') \leq \frac{2}{3}n$ if n is odd. Hence, we may consider the case with even $n \geq 6$. Let G' be the cycle $v_1v_2 \cdots v_nv_1$. Observe that both the set V_{even} of vertices with even subscript and the set V_{odd} of vertices with odd subscript are identifying codes of G' of size $n/2$. If the chord in G is between two vertices with even subscripts, then the set V_{odd} is an identifying code in G , and vice versa. Moreover, if the chord is between a vertex with even subscript and a vertex with odd subscript, then both sets V_{even} and V_{odd} are identifying codes of G . Hence, $\gamma^{\text{ID}}(G) \leq \frac{1}{2}n$. If $n = 7$, then $\gamma^{\text{ID}}(G) = \gamma^{\text{ID}}(G') - 1 = 4 < \frac{2}{3}n$ by Observation 8. Therefore, if G' is a cycle, then $\gamma^{\text{ID}}(G) \leq \frac{2}{3}n$, as desired. Suppose secondly that G' is a path. In this case, $n \geq 5$. For small values of n , namely $n \in \{5, 6, 7, 8, 10\}$, it can readily be checked that $\gamma^{\text{ID}}(G) \leq \gamma^{\text{ID}}(G') = \lfloor \frac{n}{2} \rfloor + 1 \leq \frac{2}{3}n$. Hence we may assume that $n \geq 9$ is odd or $n \geq 12$ is even. Any identifying code in the path G' can be extended to an identifying code in $G = G' + e$ by adding at most one vertex, and so, by Corollary 7(a) we have $\gamma^{\text{ID}}(G) \leq \gamma^{\text{ID}}(G') + 1 = \lfloor \frac{n}{2} \rfloor + 2 \leq \frac{2}{3}n$. Hence, we may assume that $\Delta' \geq 3$, for otherwise the desired bound holds.

Since G is triangle-free, we note that $G' \neq K_{1,\Delta'}$ (since otherwise adding back the deleted edge e would create a triangle in G). In particular if $\Delta' \geq 4$, then $G' \notin \mathcal{F}_{\Delta'}$. If $\Delta' = 3$ and $G' \in \mathcal{F}_{\Delta'}$, then by Observation 8 and Lemma 14 we infer that $\gamma^{\text{ID}}(G) < \frac{2}{3}n \leq \left(\frac{\Delta-1}{\Delta}\right)n$. Hence we may assume that $G' \notin \mathcal{F}_{\Delta'}$, for otherwise the desired bound holds. Applying the inductive hypothesis to the graph G' , we have $\gamma^{\text{ID}}(G') \leq \left(\frac{\Delta'-1}{\Delta'}\right)n \leq \left(\frac{\Delta-1}{\Delta}\right)n$.

For notational convenience, let $N_u = N_G(u) \setminus \{v\}$ and let $N_v = N_G(v) \setminus \{u\}$. Since G is triangle-free and $uv \in E(G)$, we note that $N_u \cap N_v = \emptyset$. Let A be the boundary of the set $\{u, v\}$, that is, $A = N_u \cup N_v$ is the set of vertices different from u and v that are adjacent to u or v . Further, let

$$A_{uv} = A \cup \{u, v\} = N_G[u] \cup N_G[v] \quad \text{and} \quad \overline{A}_{uv} = V(G) \setminus A_{uv}.$$

See Figure 2 for an illustration.




Figure 2: The general setting of the proof of Theorem 2. Since G is triangle-free, N_u and N_v are independent sets, but there can be edges across them.

Let C' be an optimal identifying code in G' , and so C' is an identifying code in G' and $|C'| = \gamma^{\text{ID}}(G')$. If C' is an identifying code in G , then $\gamma^{\text{ID}}(G) \leq |C'| = \gamma^{\text{ID}}(G') \leq \left(\frac{\Delta-1}{\Delta}\right)n$. Hence, we may assume that C' is not an identifying code in G , for otherwise the desired bound holds.

We will next proceed with proving a series of claims.

Claim A. If $V(G) = A_{uv}$, then

$$\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n.$$

Proof. Suppose that $V(G) = A_{uv}$. Recall that by our choice of the edge e , we have $\deg_G(u) \geq 2$, $\deg_G(v) \geq 2$, and $5 \leq \deg_G(u) + \deg_G(v) \leq 2\Delta$. In this case since $V(G) = A_{uv}$, we have $n = \deg_G(u) + \deg_G(v) \leq 2\Delta$. Let u' be an arbitrary neighbor of u in G different from v , and let v' be an arbitrary neighbor of v in G different from u . The code $C = V(G) \setminus \{u', v'\}$ is an identifying code in G , implying that

$$\begin{aligned} \gamma^{\text{ID}}(G) \leq |C| &= \deg_G(u) + \deg_G(v) - 2 \\ &= \left(\frac{\deg_G(u) + \deg_G(v) - 2}{\deg_G(u) + \deg_G(v)}\right)n \\ &\leq \left(\frac{2\Delta - 2}{2\Delta}\right)n \\ &= \left(\frac{\Delta - 1}{\Delta}\right)n, \end{aligned}$$

yielding the desired upper bound. \square

By Claim A, we may assume that $V(G) \neq A_{uv}$, for otherwise the desired result follows. Let

$$G_{uv} = G - N_G[u] - N_G[v],$$

and so $V(G_{uv}) = \bar{A}_{uv} = V(G) \setminus (A \cup \{u, v\})$. Since C' is an identifying code in G' but not in G , a pair of vertices in G is not identified by the code C' when adding back the deleted edge e to the graph G' to reconstruct G . Since G is triangle-free, the only possible pairs of vertices not identified by the code C' in G are the pairs $\{u, v\}$ or $\{u', v\}$ or $\{u, v'\}$ where u' is some neighbor of u different from v , and v' is some neighbor of v different from u .

Claim B. If for every optimal identifying code C' in G' the pair $\{u, v\}$ is the only pair not identified by the code C' in G , then

$$\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n.$$

Proof. Let C' be an optimal identifying code in G' , and suppose the pair $\{u, v\}$ is the only pair not identified by the code C' in G . Necessarily, $\{u, v\} \subseteq C'$, no neighbor of u different from v belongs to C' and no neighbor of v different from u belongs to C' ; that is, $C' \cap A = \emptyset$, and so no neighbor of u in G' and no neighbor of v in G' belongs to C' . Equivalently, $C' \setminus \{u, v\} = C' \cap \overline{A_{uv}}$. See Figure 2, where the black vertices belong to C' , and the other ones do not. We proceed further with a series of subclaims that we will need when proving Claim B. Since every neighbor of u (respectively, v) is identified by the code C' in the graph G' , we infer the following claim.

Claim B.1. Every neighbor of u (respectively, v) in G' has at least one neighbor that belongs to the set $C' \setminus \{u, v\}$.

We shall frequently use the following claim when obtaining structural properties of the graph G .

Claim B.2. If there exists a vertex $w \in C' \setminus \{u, v\}$ and a vertex $z \in A$ such that $(C' \setminus \{w\}) \cup \{z\}$ is an identifying code in the graph G , then $\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n$.

Proof. Consider the set $(C' \setminus \{w\}) \cup \{z\}$ be as defined in the statement of the claim. If this set is an identifying code in the graph G , then $\gamma^{\text{ID}}(G) \leq |C'| \leq \left(\frac{\Delta-1}{\Delta}\right)n$. \square

By Claim B.2, we may assume that there does not exist a vertex $w \in C' \setminus \{u, v\}$ and a vertex $z \in A$ such that $(C' \setminus \{w\}) \cup \{z\}$ is an identifying code in the graph G , for otherwise the desired bound holds. Recall that $C' \cap A = \emptyset$. In the following claim, we show that there are no P_2 -components in G_{uv} .

Claim B.3. No component in G_{uv} has order 2.

Proof. Suppose that the graph G_{uv} contains a component F of order 2, and so F is isomorphic to P_2 . As observed earlier, in the graph G' we have $C' \cap N(u) = C' \cap N(v) = \emptyset$. In this case, the two vertices in the component F are not separated by the code C' , a contradiction. \square

In the following subclaims we consider the case with $\Delta = 3$ separately. This is due to the more complex structure of set \mathcal{F}_3 compared to sets \mathcal{F}_i for $i \geq 4$.

Claim B.4. Let $\Delta = 3$. If C_7 is a component in G_{uv} , then $\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n$.

Proof. Let $F = C_7$ be a cycle component in G_{uv} . Since C' is an optimal identifying code in G' and since C' contains no vertices in the boundary A , the set $V(F) \cap C'$ is an identifying code in F , that is, $\gamma^{\text{ID}}(F) \leq |V(F) \cap C'|$. By Theorem 6, we have $\gamma^{\text{ID}}(C_7) = 5$. Let $V(F) = \{w_1, w_2, \dots, w_7\}$ and $E(F) = \{w_7w_1\} \cup \{w_iw_{i+1} \mid 1 \leq i \leq 6\}$. Observe that each vertex in F can be adjacent to at most one vertex in A since $\Delta = 3$. Note that, since $\Delta = 3$, we have $1 \leq |N_u| \leq 2$ and $1 \leq |N_v| \leq 2$, and so $|A| \leq 4$. We denote $N_u = \{u_1, u_2\}$ and $N_v = \{v_1, v_2\}$ (if these vertices exist). We further assume that $|N(F) \cap N_u| \geq |N(F) \cap N_v|$. Moreover, let w_2 be adjacent to $u_1 \in N_u$. Let us consider vertex sets

$$C_1 = (C' \setminus V(F)) \cup \{u_1, w_1, w_4, w_5, w_6\}$$

and

$$C_2 = (C' \setminus V(F)) \cup \{u_1, w_3, w_5, w_6, w_7\}.$$

In the following, we show that at least one of these two sets is an identifying code in G . Notice that $|C_1| = |C_2| \leq |C'|$.

If u_1 is the only vertex of A adjacent to a vertex in F , then C_1 is an identifying code in G . In particular, we may use Lemma 3 to see that u, v and u_1 have unique neighborhoods in C_1 , while $C_1 \cap V(F)$ forms an identifying code for $F \setminus \{w_2\}$, and w_2 is identified by vertices u_1 and w_1 . The remaining vertices are identified by the vertices in C' .

Assume then that there are two vertices of A (u_1 and, say, x) adjacent to vertices in F . If x is in N_u ($x = u_2$), then C_1 is an identifying code in G , even if x is not dominated by a vertex in $C_1 \cap V(F)$. Indeed, we have $u \in N[u_2] \cap C_1$ but $v, u_1 \notin N[u_2] \cap C_1$. Thus, u_2 is separated from all other vertices. If $x \in N_v$, then C_1 or C_2 is an identifying code in G . In this case, we choose such a set C_i ($i \in \{1, 2\}$) so that $N(x) \cap V(F) \cap C_i \neq \emptyset$.

Assume next that there are three vertices of A adjacent to vertices in F . In this case, due to our assumption that $|N_u \cap N(F)| \geq |N_v \cap N(F)|$, we have $|N_u \cap N(F)| = 2$ and hence, we may assume that $A \cap N(F) = \{u_1, u_2, v_1\}$. Again we choose a set C_i ($i \in \{1, 2\}$), such that $N(v_1) \cap V(F) \cap C_i \neq \emptyset$, as our identifying code. Note that as in the previous case, we do not need to dominate u_2 from F .

Finally, when (all) four vertices of A are adjacent to vertices in F , we again choose a set C_i ($i \in \{1, 2\}$) such that $N(v_1) \cap V(F) \cap C_i \neq \emptyset$ as our identifying code. As in the previous cases, we do not need to dominate vertices u_2 and v_2 from F . The argument for u_2 is similar as in the previous cases. Moreover, by Lemma 3, vertices v and u have unique neighborhoods in C_i , vertex v separates v_2 from vertices other than u, v and v_1 , while v_1 is separated from v_2 by a vertex in F .

The claim follows in these cases since $|C_1| = |C_2| \leq |C'|$ and since C_1 or C_2 is an identifying code in G . \square

Claim B.5. Let $\Delta = 3$. If C_4 is a component in G_{uv} , then $\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n$.

Proof. Let $F = C_4$ be a cycle component in G_{uv} . Since C' is an optimal identifying code in G' and since C' contains no vertices in the boundary A , the set $V(F) \cap C'$ is an identifying code in F , that is (by Theorem 6), $\gamma^{\text{ID}}(F) = 3 \leq |V(F) \cap C'|$. Let $V(F) = \{w_1, w_2, w_3, w_4\}$ and $E(F) = \{w_1w_2, w_2w_3, w_3w_4, w_4w_1\}$. Observe that each vertex in F can be adjacent to at most one vertex in A since $\Delta = 3$. Recall that by Claim B.1 every vertex in A is dominated by two vertices in C' .

Let us first assume that $V(F) \subseteq C'$. Then, we can apply Claim B.2 to any vertex in F and its neighbor in A and the claim follows. Thus, we can now assume that $|V(F) \cap C'| = 3$.

Let us first assume that there are one or three vertices in A adjacent to vertices in F . Note that each vertex in F can be adjacent to at most one vertex in A . If there is one vertex in A adjacent to a vertex in F , then we assume that the adjacent vertex in F is $w_2 \in F$. If there are three vertices in A adjacent to vertices in F , then we assume that they are adjacent to vertices w_1, w_2 and w_3 in F . Furthermore, we assume that $z \in A$ is adjacent to w_2 . We will consider vertex set $C = (C' \setminus V(F)) \cup \{w_1, w_3, z\}$. This is an identifying code in G since all vertices in A_{uv} are separated from other vertices by u and v and vertices u, v and z have unique code neighborhoods in C by Lemma 3. Moreover, vertices in F are separated from each other by z, w_1 and w_3 and finally vertices in $A \setminus \{z\}$ are separated from each other by the same vertices as in C' .

Assume then that there are four vertices in A adjacent to vertices in F . Let us assume without loss of generality that $\{w_1, w_2, w_3\} \subset C'$ and that $z \in A$ is adjacent to w_2 . As in the previous case, now $C = (C' \setminus \{w_2\}) \cup \{z\}$ is an identifying code in G .

Finally, we have the case where we have exactly two vertices in A adjacent to vertices in F . Assume first that these vertices in A are adjacent to two adjacent vertices of F , say to w_1 and w_2 . Let $z \in A$ be a vertex adjacent to w_2 . As in the previous cases, we let $C = (C' \setminus V(F)) \cup \{w_1, z, w_3\}$. With the same arguments as in the previous cases, C is an identifying code in G .

Let us next consider the case where there are two vertices in A adjacent to only non-adjacent vertices in F , say vertices w_1 and w_3 . Observe that now w_2 and w_4 are open twins in G' and we have exactly two edges between A and F . Let $z \in A$ be a neighbor of w_1 . Assume that $|N_u| \geq |N_v|$. Let us denote the neighbors of u in A by u_1 and u_2 and neighbors of v in A by v_1 and v_2 (if v_2 exists). Notice that if both edges from F to A are between F and N_u or F and N_v , then $C = \{z, w_2, w_4\} \cup (C' \setminus V(F))$ is an identifying code of G . Thus, we may assume without loss of generality, that there is an edge from w_1 to u_1 and from w_3 to v_1 . Assume next that for u_2 or v_2 there exists vertex $c \in C$ such that $c \in N(v_2) \setminus N(v_1)$ or $c \in N(u_2) \setminus N(u_1)$. If the former holds, then $C_u = \{u_1, w_2, w_4\} \cup (C' \setminus V(F))$ is an identifying code of G and if the latter holds, then $C_v = \{v_1, w_2, w_4\} \cup (C' \setminus V(F))$ is an identifying code of G . Hence, we next assume that c does not exist. Moreover, if v_2 does not exist, then we may again use identifying code C_u . Since each vertex in A is dominated by at least two vertices, there exist vertices $c_u \in C' \cap N(u_1) \cap N(u_2) \setminus \{u\}$ and $c_v \in C' \cap N(v_1) \cap N(v_2) \setminus \{v\}$. Observe that if there exists a vertex w_u such that $N[w_u] \cap C' = \{c_u\}$ or a vertex w_v such that $N[w_v] \cap C' = \{c_v\}$, then $w_u = c_u$ and $w_v = c_v$. Indeed, otherwise we would have $N[w_u] \cap C' = N[c_u] \cap C'$ or $N[w_v] \cap C' = N[c_v] \cap C'$. We may also observe that we have either $\{w_1, w_2, w_3\} \subseteq C'$ or $\{w_1, w_4, w_3\} \subseteq C'$. Let us assume, without loss of generality, the former. However, now $C = \{u_1\} \cup C' \setminus \{u\}$ is an identifying code in G . Indeed, only u, u_1

and u_2 have lost a codevertex from their neighborhoods compared to C' . Moreover, u is the only vertex adjacent to both v and u_1 , u_1 is the only vertex adjacent to w_1 and c_u and finally, $C \cap N[u_2] = \{c_u\}$ but now $C \cap N[c_u] \supseteq \{c_u, u_1\}$. Thus, also u_2 has a unique code neighborhood. Now, the claim follows. \square

Claim B.6. Let $\Delta = 3$. If P_4 is a component in G_{uv} , then $\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n$.

Proof. Let $F = P_4$ be a path component in G_{uv} . Since C' is an optimal identifying code in G' and since C' contains no vertices in the boundary A , the set $V(F) \cap C'$ is an identifying code in F , that is, by Theorem 5, $\gamma^{\text{ID}}(F) = 3 \leq |V(F) \cap C'|$. Let $V(F) = \{w_1, w_2, w_3, w_4\}$ and $E(F) = \{w_1w_2, w_2w_3, w_3w_4\}$.

Case 1: there is an edge between w_1 or w_4 and A . Without loss of generality, let it be w_1 . Consider now the graph $G_1 = G - \{w_2, w_3, w_4\}$. First observe that G_1 is connected. Second, if $G_1 \notin \mathcal{F}_3$, then by induction we have an identifying code $C_1 \subseteq V(G_1)$ of cardinality at most $\frac{2}{3}n(G_1)$. Moreover, if no neighbor (in G) of w_2, w_3 or w_4 is in C_1 , then $C \cup \{w_2, w_4\}$ is an identifying code of claimed cardinality in G . If on the other hand $C_1 \cap (N_G(w_2) \cup N_G(w_3)) \neq \emptyset$, then at least one of the sets $C_1 \cup \{w_2, w_4\}$ and $C_1 \cup \{w_3, w_4\}$ is an identifying code in G . The case with $C_1 \cap (N_G(w_4) \cup N(w_3)) \neq \emptyset$ is analogous. Moreover, these identifying codes have the claimed cardinality.

Let us next consider the case with $G_1 \in \mathcal{F}_3$. Furthermore, there is a vertex with degree 3 and there are at least six vertices in G_1 since $\deg(u) + \deg(v) \geq 5$ by our assumption. Thus, $G_1 \in \mathcal{F}_3 \setminus \{P_4, C_4, C_7, K_{1,3}\} = \mathcal{T}_3 \setminus \{K_{1,3}\}$. Let us assume first that $G_1 = T_2$. In this case, we have $n = 10$ and we need to find an identifying code containing at most six vertices. Notice that $\deg(u) + \deg(v) = 5$ and either u or v is the support vertex of degree 3 in T_2 and the other one is the adjacent vertex of degree 2. However, since w_1 is a leaf in T_2 , this means that it must have distance of either 1 or 3 to set $\{u, v\}$, a contradiction since that distance is actually 2.

Let us next assume that $G_1 = T_3$. Now, $n = 13$ and we need to find an identifying code containing nine vertices. Observe that in this case $\deg(u) + \deg(v) = 6$. Since w_1 has distance 2 to set $\{u, v\}$, vertex w_1 has to be one of the leaves adjacent to support vertices of degree 2 of T_3 . Moreover, the leaf of T_3 adjacent to the unique support vertex of degree 2 in T_3 as well as that support vertex are not in A and thus not adjacent to any vertex in F . However, in this case, to separate that support vertex and its adjacent leaf, the vertex of degree 2 adjacent to the only degree 3 non-support vertex (in T_3), must be in any identifying code. However, this vertex is in A , and this is against our assumption that $A \cap C' = \emptyset$, a contradiction.

Assume next that $G_1 \in \mathcal{T}_3 \setminus \{K_{1,3}, T_2, T_3\}$. Let C_1 be an optimal identifying code of G_1 such that every vertex with degree at most 2 in G_1 is in C_1 (such a code exists by Proposition 13, see Figure 1). Since G is not a tree, there are at least two edges between $A \cup \{w_1\}$ and $\{w_2, w_3, w_4\}$. Hence, there is an edge between w_1 and w_2 and between $G'_1 = G_1 - \{w_1\}$ and $\{w_2, w_3, w_4\}$.

Let us consider the case where there are no edges between A and w_2 or w_3 but there is an edge to w_4 . In this case, we may consider the identifying code $C = C_1 \cup \{w_3\}$ which has the claimed cardinality. Notice that this is indeed an identifying code since $w_1 \in C_1$ and w_4 is dominated by some vertex in C_1 .

Therefore, we may assume from now on that there is an edge between A and w_2 or w_3 . Let us assume that the edge is from A to w_2 (the case with an edge to w_3 is similar). Consider now graph $G''_1 = G - \{w_1, w_3, w_4\}$ together with an optimal identifying code C''_1 . Observe that G''_1 is a tree since G_1 is a tree and w_2 has only one edge to A . Moreover, we also notice that there are no edges between vertices in A since G_1 is a tree. Thus, if a 4-cycle contains exactly two vertices in F , then those vertices are w_1 and w_4 .

If $G''_1 \notin \mathcal{F}_3$, then we have two cases based on whether $w_2 \in C''_1$.

If $w_2 \in C''_1$, then $C''_1 \cup \{w_3, w_4\}$ is an identifying code in G . Indeed, by Lemma 3 vertices w_2, w_3 and w_4 have unique code neighborhoods. Moreover, also w_1 has a unique neighborhood in $C''_1 \cup \{w_2\}$ since any vertex of A adjacent to w_2 is also either in C''_1 or has another neighbor outside of A in C''_1 .

If $w_2 \notin C''_1$, then we use set $C_{23} = C''_1 \cup \{w_2, w_3\}$. Since C''_1 is an identifying code in G''_1 , the vertex adjacent to w_2 (call it z) in A is in C''_1 and z has another code neighbor. Hence, by Lemma 3, vertices z, w_2 and w_3 have unique code neighborhoods. Moreover, since w_1 is adjacent to w_2 and all other vertices in $N[w_2]$ have unique code neighborhoods, also w_1 has a unique neighborhood in C_{23} . Furthermore, also

w_4 has a unique neighborhood in C_{23} since the only other vertex x adjacent to w_3 which might not be separated from w_4 is in A and is in C_1'' or has another vertex in C_1'' adjacent to it. Since w_3 and w_4 cannot belong to the same 4-cycle in G and there are no triangles, vertices w_4 and x are separated. All the other vertices are pairwise separated by the set C_1'' . Hence, C_{23} is an identifying code in G .

Thus, we may assume that $G_1'' \in \mathcal{F}_3$. Observe that we may construct G_1'' from G_1 by removing a leaf and then adding a new leaf and vice versa. Hence, G_1, G_1'' are two trees of \mathcal{F}_3 (and thus, \mathcal{T}_3) with the same order of at least 6. Observe that for every i such that $i \in \{0, 1\}$ or $i \geq 4$, there are no support vertices of degree 2 in T_i ($T_i \in \mathcal{T}_3$). However, when we remove a leaf from any such tree of \mathcal{T}_3 and add a different leaf to a vertex of degree at most 2, there necessarily exists a support vertex of degree 2 in the resulting tree. Thus, at least one of G_1 and G_1'' must be in $\{T_2, T_3\}$. Note that we cannot obtain an isomorphic copy of T_3 by this operation when starting from T_3 , so at most one of G_1 and G_1'' is T_3 . Thus, $\{G_1, G_1''\} = \{T_1, T_2\}$, $\{G_1, G_1''\} = \{T_2, T_2\}$ or $\{G_1, G_1''\} = \{T_3, T_4\}$. Let us first assume that $G_1, G_1'' = \{T_3, T_4\}$. Consider the leaf adjacent to the support vertex of degree 2 in T_3 . Notice that the leaf has to belong to F . Moreover, its distance from set $\{u, v\}$ is exactly 2. However, now when we form T_4 by removing this leaf and attaching the new leaf, the new leaf is attached adjacent to u or v and hence, $A \cap V(F) \neq \emptyset$, a contradiction. Thus, $G_1, G_1'' \in \{T_1, T_2\}$.

We thus have $\{G_1, G_1''\} = \{T_1, T_2\}$ or $\{G_1, G_1''\} = \{T_2, T_2\}$. In both cases, we have $\deg(u) + \deg(v) = 5$. Hence, one of u or v is the degree 3 support vertex in T_2 and the other one is the adjacent degree 2 vertex. Moreover, the leaf adjacent to the support vertex of degree 2 in T_2 is in F . But this vertex should have distance 2 to set $\{u, v\}$, a contradiction. This finishes the proof of Case 1.

Case 2: there are no edges from w_1 or w_4 to A . Then, there is an edge from A to w_2 or w_3 ; without loss of generality, assume there is an edge from A to w_2 and denote $G_2 = G - \{w_1, w_3, w_4\}$. Observe that if $G_2 \notin \mathcal{F}_3$, then by induction, there exists an identifying code C_2 in G_2 with $|C_2| \leq \frac{2}{3}n(G_2)$ and $|C_2| + 2 \leq \frac{2}{3}n$. Moreover, either $C_2 \cup \{w_1, w_3\}$ or $C_2 \cup \{w_2, w_3\}$ is an identifying code in G depending on whether $w_2 \in C_2$ or not. Hence, we may assume that $G_2 \in \mathcal{F}_3$ and in fact $G_2 \in \mathcal{T}_3$ since G_2 has a vertex of degree 3. If we do not have an edge from w_3 to A , then G would be a tree, a contradiction. Hence, we may assume that there exists an edge from w_3 to A . Let us assume that $G_2 \neq T_2$ and C_2 contains all vertices of degree at most 2 in G_2 . This is possible by Proposition 13. Observe that in this case $C_2 \cup \{w_3\}$ is an identifying code of claimed size in G . Hence, we may assume that $G_2 = T_2$. Since $\deg(u) + \deg(v) \geq 5$, the only leaf of T_2 not in A is the leaf adjacent to the degree 2 support vertex. However, this vertex has distance 3 to set $\{u, v\}$. Thus $G_2 \neq T_2$, and the claim follows. \square

Claim B.7. Let $\Delta = 3$. If $T \in \mathcal{T}_3$ is a component in G_{uv} , then $\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n$.

Proof. Let $F = T \in \mathcal{T}_3$ be a tree component in G_{uv} . Since C' is an optimal identifying code in G' and since C' contains no vertices in the boundary A , the set $V(F) \cap C'$ is an identifying code in F , that is, $\gamma^{\text{ID}}(F) \leq |V(F) \cap C'|$. Let us call by T^2 the set of vertices of degree at most 2 in $G[F]$. Note that only vertices of T^2 can have a neighbor in A .

Let us first assume that $F \in \mathcal{T}_3 \setminus \{T_2, T_3\}$. By Proposition 13 we can choose an optimal identifying code C_T for T such that $T^2 \subseteq C_T$ (see Figure 1). Moreover, by Proposition 13(ii) $C_T \setminus \{t\}$ is an optimal identifying code in $T - \{t\}$ for any $t \in T^2$. Notice that $C_T' = (C' \setminus V(F)) \cup C_T$ is an identifying code in G' and $|C'| = |C_T'|$. Notice that Claim B.2 holds also for C_T' . Assume first that $z \in A$ is adjacent to $t \in T^2$. If $C = \{z\} \cup C_T' \setminus \{t\}$ is an identifying code in G , then the claim follows from Claim B.2. If C is not an identifying code, then we will show next that vertex t must be adjacent to a vertex $u_1 \in N_u$ and $v_1 \in N_v$ and t is the only vertex in C_T' separating u_1 from another vertex $u_2 \in N_u$, and similarly, vertex $v_2 \in N_v$ from $v_1 \in N_v$. Indeed, since $u, v \in C$, all vertices of A are separated from vertices in F . Moreover, since $C_T \setminus \{t\}$ is an identifying code in $F - t$, set C is an identifying code in $F - t$. Furthermore, vertex t is the only vertex which has exactly z in its code neighborhood. Consequently, z, u and v are identified by Lemma 3. Therefore, the only vertices which might not be separated belong to A . Since C_T' separated all vertices in A , we require t to separate some vertices which are not separated by z . Moreover, t can have at most two neighbors in A . If t is adjacent to only one vertex in A , then that vertex is z and $z \in C$ separates itself from other vertices in A . Thus, t is adjacent to two vertices in A . If both of these vertices are in N_u (or N_v), then u and v separate them from other vertices in A . Moreover, $z \in C$ separates it

itself from other neighbors of t . Hence, t is adjacent to a vertex $u_1 \in N_u$ and $v_1 \in N_v$. Since $u, v \in C$, the only vertex u_1 that can have the same code neighborhood with respect to C is vertex $u_2 \in N_u$. A similar statement holds for v_1 and $v_2 \in N_v$. Assume first that $C \setminus \{z\}$ separates v_1 and v_2 in G . In this case $(C \cup \{u_1\}) \setminus \{z\}$ is an identifying code in G . A similar argument holds for $C \setminus \{z\}$ separating u_1 and u_2 in G . Hence, we may assume that $C \setminus \{z\}$ does not separate pairs u_1, u_2 and v_1, v_2 . Thus, we require t to separate these two pairs in C'_T , as claimed.

Since v_2 and u_2 are dominated by a vertex in $C'_T \setminus \{u, v\}$ (Claim B.1), there is a code vertex $w_u \in (C'_T \setminus \{u, v\}) \cap N(u_1) \cap N(u_2)$ and a code vertex $w_v \in (C'_T \setminus \{u, v\}) \cap N(v_1) \cap N(v_2)$. In this case, we may consider graph G' and modify the identifying code C'_T into $C'' = \{u_1, v_1\} \cup C'_T \setminus \{u, v\}$ which is an identifying code in G' . This is a contradiction because C'' is an optimal identifying code of G' such that u, v are identified by C'' in G , contradicting the hypothesis of Claim B.

Assume next that $F = T_2$. Denote its unique support vertex of degree 3 by s . Let the leaves adjacent to s be denoted by l_1 and l_2 and the third one be l_3 . Finally, let f_1, f_2, f_3 be the three vertices on the path from s to l_3 (in that order). Assume first that there are no edges between A and f_2, f_3 or l_3 . Now consider the graph $G_f = G - \{f_2, f_3, l_3\}$. Assume first that $G_f \notin \mathcal{F}_3$ and let C_f be an optimal identifying code in G_f . In this case, if $f_1 \in C_f$, then $C_f \cup \{f_2, f_3\}$ is an identifying code of claimed cardinality in G . If $f_1 \notin C_f$, then $C_f \cup \{f_2, l_3\}$ is an identifying code in G of claimed cardinality. Assume then that $G_f \in \mathcal{F}_3$. Since there is a vertex of degree 3 in G_f , we have $G_f \in \mathcal{T}_3$. However, since there are no edges from A to $\{f_2, f_3, l_3\}$, graph G is a tree, a contradiction. Hence, we may assume that there is an edge from A to $\{f_2, f_3, l_3\}$.

Let us next consider graph $G_s = G - \{l_1, l_2, s\}$ and its optimal identifying code by C_s . Notice that G_s is connected. Assume first that $G_s \notin \mathcal{F}_3$. If one of the three vertices in $\{l_1, l_2, s\}$ is dominated by a vertex in C_s in G , then the set C_s together with two adjacent vertices from $\{l_1, l_2, s\}$ such that at least one of them is dominated by a vertex from C_s is an identifying code of claimed cardinality. If no vertex in $\{l_1, l_2, s\}$ is dominated by a vertex from C_s , then we consider $C_s \cup \{l_1, l_2\}$. This is an identifying code since s is separated from all other vertices. Indeed, if $u \in V(G) \setminus V(F)$ is adjacent to l_1 and l_2 , then it is adjacent also to some vertex in $C_s \cap (V(G) \setminus V(F))$ while vertex s cannot be (since it is not dominated by C_s). Hence, we may assume that $G_s \in \mathcal{F}_3$ and more specifically $G_s \in \mathcal{T}_3$ since there is a vertex of degree 3. Hence, there is a single edge between $\{f_1, f_2, f_3, l_3\}$ and A . Furthermore, if there is an edge from A to f_2 or f_3 , then $G_s \notin \mathcal{T}_3$, and if the edge is from A to f_1 or l_3 , then G_s has to be T_2 . However, there are at least five vertices in A_{uv} and hence $G_s \neq T_2$. Thus $G_s \notin \mathcal{F}_3$.

Let us finally consider the case where $F = T_3$. Denote by s_1 and s_2 its two support vertices of degree 3, and by f_2 the support vertex of degree 2. Let leaves l_1 and l_2 be adjacent to s_1 , and leaves l_3 and l_4 be adjacent to s_2 . Furthermore, let the leaf adjacent to f_2 be l_5 and the other vertex adjacent to f_2 be f_1 . Further denote the vertex of degree 3 adjacent to f_1 by f_s . Assume first that there are no edges from A to $\{f_1, f_2, l_5\}$. Thus, $G_f = G - \{f_1, f_2, l_5\}$ is a connected graph and let C_f be an optimal identifying code in G_f . Notice that if $G_f \in \mathcal{F}_3$, then $G_f \in \mathcal{T}_3$, and then G is a tree, a contradiction. Thus, $G_f \notin \mathcal{F}_3$ and C_f contains at most two-thirds of the vertices of G_f . In this case, if $f_s \in C_f$, then the set $C = C_f \cup \{f_1, f_2\}$ is an identifying code in G and if $f_s \notin C_f$, then the set $C = C_f \cup \{f_1, l_5\}$ is an identifying code in G . Moreover, both of these sets contain at most two-thirds of the vertices in G , and we are done. Hence, we may assume from now on that there is an edge from A to $\{f_1, f_2, l_5\}$. Assume then that there are no edges from A to $\{s_1, l_1, l_2\}$. Now, graph $G_s = G - \{s_1, l_1, l_2\}$ is connected, has a cycle and hence, by induction, also an optimal identifying code C_s satisfying the two-thirds upper bound. Moreover, set $C_s \cup \{l_1, l_2\}$ is an identifying code in G . Hence, there is an edge from A to $\{s_1, l_1, l_2\}$. By symmetry, a similar argument holds for $\{s_2, l_3, l_4\}$, so there is an edge from A to $\{s_2, l_3, l_4\}$. Hence we can assume next that there is an edge from A to sets $\{s_1, l_1, l_2\}$, $\{s_2, l_3, l_4\}$ and $\{l_5, f_1, f_2\}$. Let us consider graph $G'_f = G - \{s_2, l_3, l_4\}$ together with an optimal identifying code C'_f of G'_f . Notice that G'_f is connected, has a cycle, and maximum degree 3. Hence, $G'_f \notin \mathcal{F}_3$ and by induction, $|C'_f| \leq \frac{2}{3}n(G'_f)$. Consider set $C'_f \cup \{l_3, l_4\}$ if no vertex in C'_f dominates a vertex in $\{l_3, l_4\}$ and otherwise, set C'_f together with s_2 and a vertex in $\{l_3, l_4\}$ that is dominated by C'_f . Notice that in each case, the corresponding set is an identifying code of claimed cardinality for G . \square

By the above claims, we may assume that if $\Delta = 3$, then for any component F in G_{uv} we have

$$\gamma^{\text{ID}}(F) \leq \frac{2}{3}n(F).$$

Claim B.8. Let $\Delta \geq 4$. If T is a Δ -star component in G_{uv} , then $\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n$.

Proof. Let T be a Δ -star component in G_{uv} . Let $V(T) = \{w, w_1, \dots, w_\Delta\}$ and w be the center vertex of T . Observe that w is adjacent only to vertices in T . Let w_1 be adjacent to a vertex $z \in A$. Let $T' = T - \{w_1\}$ and let $G_T = G - T'$. Observe that G_T is a connected graph and at least one of u or v has degree at least 3 in G_T . Hence, there are at least six vertices in G_T . Let us assume first that $G_T \in \mathcal{F}_3$; then, $G_T \in \mathcal{T}_3$. In this case we have $\gamma^{\text{ID}}(G_T) \leq \frac{3}{4}n(G_T)$ and $\gamma^{\text{ID}}(T') \leq \left(\frac{\Delta-1}{\Delta}\right)n(T')$. Moreover, there are at least three leaves in T' . Let C_T be an optimal identifying code in G_T . Observe that $C = C_T \cup \{w, w_2, \dots, w_{\Delta-1}\}$ is an identifying code in G of cardinality at most $|C| \leq \frac{3}{4}n(G_T) + \left(\frac{\Delta-1}{\Delta}\right)n(T') \leq \left(\frac{\Delta-1}{\Delta}\right)n$. Furthermore, if $G_T \notin \mathcal{F}_3$, since also $G_T \notin \mathcal{F}_\Delta$, there exists an identifying code C_T of G_T such that $|C_T| = \gamma^{\text{ID}}(G_T) \leq \left(\frac{\Delta-1}{\Delta}\right)n(G_T)$ and hence, we may again consider $C = C_T \cup \{w, w_2, \dots, w_{\Delta-1}\}$ as our identifying code for G and $|C| \leq \left(\frac{\Delta-1}{\Delta}\right)n$ which completes the proof of the claim. \square

Let us denote the set of isolated vertices in G_{uv} by \mathcal{I} and let us further denote $G_{AI} = G[A_{uv} \cup \mathcal{I}]$.

Claim B.9. Set A_{uv} is an identifying code in G_{AI} .

Proof. Observe that $\mathcal{I} \subseteq C'$ (otherwise some vertex of \mathcal{I} would not be dominated by C' in G'). If set A_{uv} is not an identifying code in G_{AI} , then there are open twins w_1, w_2 in \mathcal{I} . Let $z \in N(w_1)$. Notice that $z \in A$ and hence, $z \notin C'$. Since vertices in $\{w_1, w_2\}$ are open twins, we may consider set $C = \{z\} \cup C' \setminus \{w_1\}$. Observe that C is an identifying code since w_2 separates all the same vertices as vertex w_1 . Hence, we obtain a contradiction from Claim B.2. Thus, there are no twins in \mathcal{I} and set A_{uv} is an identifying code in G_{AI} . \square

Let us next consider the minimum cardinality of an identifying code in G_{AI} . Assume that $|N_u| \geq |N_v|$. Since $|N_v| \leq |N_u| \leq \Delta - 1$, we have $|A_{uv}| \leq 2\Delta$. Thus, if $|A_{uv}| > \left(\frac{\Delta-1}{\Delta}\right)n(G_{AI}) = \left(\frac{\Delta-1}{\Delta}\right)(|A_{uv}| + |\mathcal{I}|)$, then $|A_{uv}| > (\Delta - 1)|\mathcal{I}|$. Hence, this implies that $|\mathcal{I}| \leq 2$. By Claim B.9, the set \mathcal{I} is A -identifiable. Hence, by Lemma 4, if $|\mathcal{I}| \leq 2$, then we require at most two vertices from A to separate and dominate the vertices in \mathcal{I} . Moreover, we have $|A| \geq 3$ due to the assumption $\deg(u) + \deg(v) \geq 5$. Notice that set $A_{uv} \setminus \{z_1, z_2\}$ remains an identifying code of G_{AI} when $z_1 \in N_u$ and $z_2 \in N_v$ and they are not required for dominating or separating vertices in \mathcal{I} . Indeed, these vertices will be the only ones which are adjacent to exactly u or v in the identifying code.

Hence, if $|\mathcal{I}| = 2$ and $|A_{uv}| = 2\Delta - 2 - a \leq 2\Delta - 2$ for some $a \geq 0$, then A_{uv} is an identifying code of G_{AI} that satisfies the conjectured bound for G_{AI} . Indeed,

$$|A_{uv}| = 2\Delta - 2 - a = \left(\frac{\Delta-1}{\Delta}\right)((2\Delta-2)+2) - a \leq \left(\frac{\Delta-1}{\Delta}\right)n(G_{AI}).$$

If $|\mathcal{I}| = 2$ and $|A_{uv}| \geq 2\Delta - 1$, then we may remove one well-chosen vertex from A since we require at most two vertices of A to identify vertices in \mathcal{I} while $|A| \geq 3$, and the resulting set remains an identifying code (as described above) while satisfying the conjectured bound in G_{AI} . Indeed, $2\Delta - 2 < \left(\frac{\Delta-1}{\Delta}\right)((2\Delta-2)+3)$ and $2\Delta - 1 < \left(\frac{\Delta-1}{\Delta}\right)((2\Delta-1)+3)$. If $|\mathcal{I}| = 1$ and $|A_{uv}| \leq 2\Delta - 1$, then we may remove one well-chosen vertex from N_u . If $|\mathcal{I}| = 1$ and $|A_{uv}| = 2\Delta$, then we may remove one well-chosen vertex from N_u and N_v . Finally, if $\mathcal{I} = \emptyset$, then we may just remove any single vertex from both N_u and N_v to obtain an identifying code satisfying conjectured bound.

We are now ready to complete the proof of Claim B. Observe that we have obtained above an identifying code, which contains u and v , satisfying the conjectured bound in G_{AI} . Denote by \mathcal{F}_{uv} the components in G_{uv} . Furthermore, observe that if $F \in \mathcal{F}_{uv}$ has size at least 2 in G_{uv} , then F is also a component in $G - G_{AI}$. Furthermore, by Claims B.3, B.4, B.5, B.6, B.7 and B.8, every component F of \mathcal{F}_{uv} admits an identifying code and we have $F \notin \mathcal{F}_\Delta$, and so, by the induction hypothesis, $\gamma^{\text{ID}}(F) \leq \left(\frac{\Delta-1}{\Delta}\right)n(F)$. Let us denote by C_F an optimal identifying code in component F and by C_{AI} an identifying code in G_{AI} which contains u and v , and has size at most $\left(\frac{\Delta-1}{\Delta}\right)n(G_{AI})$, as constructed

above. Since there are no edges between components of \mathcal{F}_{uv} and each edge from such a component F is to a vertex in A , which are separated from vertices in F by u and v , we have an identifying code

$$C = C_{AI} \cup \bigcup_{F \in \mathcal{F}_{uv}, n(F) \geq 3} C_F$$

and

$$\begin{aligned} |C| &\leq \left(\frac{\Delta-1}{\Delta}\right) n(G_{AI}) + \sum_{F \in \mathcal{F}_{uv}, n(F) \geq 3} \left(\frac{\Delta-1}{\Delta}\right) n(F) \\ &= \left(\frac{\Delta-1}{\Delta}\right) n. \end{aligned}$$

This completes the proof of Claim B. \square

By Claim B, we may assume that there exists an optimal identifying code C' in G' such that $\{u', v\}$ or $\{u, v'\}$ is not identified by C' in G , where u' is some neighbor of u different from v and v' is some neighbor of v different from u . Necessarily, in the case where $\{u', v\}$ is not identified, $u \in C'$ and $v \notin C'$, and if $X = N_{G'}(v) \cap C'$, then $X \neq \emptyset$ (in order for C' to dominate v in G') and $N_{G'}(u') \cap C' = X \cup \{u\}$. (We have the symmetric facts for the case where $\{u, v'\}$ is not identified.) Let v' be an arbitrary vertex in X , and let Q_{uv} be the cycle $uu'v'vu$ in G . Recall that $G_{uv} = G - N_G[u] - N_G[v]$. Further recall that $N_u = N_G(u) \setminus \{v\}$, $N_v = N_G(v) \setminus \{u\}$, and $A = N_u \cup N_v$. See Figure 3 for an illustration.




Figure 3: The setting of the proof of Theorem 2 after applying Claim B. Here, $\{u', v\}$ is not identified by C' (black vertices).

Claim C. If F is a component in G_{uv} and $F \in \mathcal{F}_\Delta$, then

$$\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right) n.$$

Proof. Suppose that F is a component in G_{uv} and $F \in \mathcal{F}_\Delta$. Suppose, firstly, that $F = K_{1,\Delta}$ where $\Delta \geq 3$. Let $V(F) = \{x, x_1, x_2, \dots, x_\Delta\}$ where x is the central vertex of F with leaf neighbors $x_1, x_2, \dots, x_\Delta$. Since G is connected, there is an edge f joining a vertex $z \in A$ and a vertex in $V(F)$. Renaming vertices if necessary, we may assume that $f = zx_\Delta$. Let $G^* = G - (V(F) \setminus \{x_\Delta\})$. Let G^* have order n^* , and so $n^* = n - \Delta$. Further, let $\Delta(G^*) = \Delta^*$. Since at least one of u and v has degree at least 3 in G , and since the degrees of u and v are the same in G and in G^* , we note that $\Delta^* \geq 3$. Thus, G^* is a connected triangle-free graph and $\Delta \geq \Delta^* \geq 3$. Moreover, G^* contains the cycle Q_{uv} and has order $n^* \geq 6$. These properties of G^* imply that $G^* \notin \mathcal{F}_{\Delta^*}$. Applying the inductive hypothesis to G^* , we have

$$\gamma^{\text{ID}}(G^*) \leq \left(\frac{\Delta^*-1}{\Delta^*}\right) n^* \leq \left(\frac{\Delta-1}{\Delta}\right) n^*.$$

Let C^* be an optimal identifying code in G^* , and so $|C^*| = \gamma^{\text{ID}}(G^*)$. The code $C^* \cup \{x, x_1, \dots, x_{\Delta-2}\}$ is an identifying code in G if $\Delta \geq 4$, or $\Delta = 3$ and if $N_G(x_1) \cap C^* \neq \emptyset$. Similarly, if $N_G(x_2) \cap C^* \neq \emptyset$ and $\Delta = 3$, then $C^* \cup \{x, x_2\}$ is an identifying code in G . Finally, if $\Delta = 3$ and $N_G(x_1) \cap C^* = N_G(x_2) \cap C^* = \emptyset$, then $C^* \cup \{x_1, x_2\}$ is an identifying code in G . Hence, we have

$$\begin{aligned} \gamma^{\text{ID}}(G) &\leq |C^*| + \Delta - 1 \\ &= \gamma^{\text{ID}}(G^*) + \Delta - 1 \\ &\leq \left(\frac{\Delta-1}{\Delta}\right)n^* + \left(\frac{\Delta-1}{\Delta}\right)(n-n^*) \\ &= \left(\frac{\Delta-1}{\Delta}\right)n, \end{aligned}$$

which yields the desired upper bound in the case when $F = K_{1,\Delta}$ where $\Delta \geq 3$. Hence, we may assume that $\Delta = 3$ and $F \in \mathcal{F}_\Delta \setminus \{K_{1,3}\}$. Thus, $F \in \{P_4, C_4, C_7\} \cup (\mathcal{T}_3 \setminus \{K_{1,3}\})$.

We next distinguish two cases.

Case 1: $F \neq P_4$. In this case, since the graph G is connected and $\Delta = 3$, one can check that there exists an induced path $P: x_1x_2x_3$ in F such that $G'' = G - V(P)$ is a connected graph. Let $n'' = n(G'')$, and so $n'' = n - 3$. Let $\Delta(G'') = \Delta''$. We note that G'' is a connected triangle-free graph and $\Delta \geq \Delta'' \geq 3$. Thus, $\Delta'' = 3$. Moreover, G'' contains the cycle Q_{uv} and has order $n'' \geq 6$. In particular, $G'' \notin \mathcal{F}_3$. Applying the inductive hypothesis to G'' , we have

$$\gamma^{\text{ID}}(G'') \leq \frac{2}{3}n''.$$

Let C'' be an optimal identifying code in G'' , and so, $|C''| = \gamma^{\text{ID}}(G'')$. If $N_G[x_1] \cap C'' = N_G[x_3] \cap C'' = \emptyset$, then we consider $C'' \cup \{x_1, x_3\}$ as a potential identifying code in G . Note that with this code, each of x_1 and x_3 is only dominated by itself, and no other vertex is in that case. If there exists some vertex y not separated from x_2 by $C'' \cup \{x_1, x_3\}$, then y must be adjacent to both x_1, x_3 . Since G is triangle-free, y is not adjacent to x_2 . Thus, $y \notin C''$ (otherwise y, x_2 would be separated). Thus, y is dominated by its third neighbor z , which is in C'' , and hence, z is adjacent to x_2 . Since $\Delta'' = \Delta = 3$, vertex y has no other neighbors in G . Thus, $y \notin A$, as in that case, y should be adjacent to u or v , but $z \notin \{u, v\}$ since z is adjacent to x_2 . Hence, $y \in F$ and F is C_4 . Thus, if $F \neq C_4$, we are done. If $F = C_4$, we show that $C'' \cup \{x_1, x_2\}$ is an identifying code in G . Indeed, z, x_1, x_2 are code vertices inducing a P_3 and we may apply Lemma 3 on them. Furthermore, x_3 is separated from all other neighbors of x_2 since they are in the identifying code. Furthermore, y is adjacent to z and x_1 . If this is true also for some vertex w , then $w \in A$ and z is adjacent to w, y, x_2 which is not possible since $\Delta = 3$ and z is adjacent to u or v . Hence, $C'' \cup \{x_1, x_2\}$ is an identifying code in G .

Assume next that for an index $j \in \{1, 3\}$, say $j = 1$, we have $N_G[x_1] \cap C'' \neq \emptyset$. Consider the set $C'' \cup \{x_1, x_2\}$. If it is an identifying code of G , then we are done. Thus, assume it is not the case. Vertices x_1, x_2 and the neighbor of x_1 in C'' induce a P_3 and thus by Lemma 3, they are uniquely identified. Hence, x_3 is not separated from some other vertex w . Thus, w is a neighbor of x_2 , and $w \notin C''$. As w is dominated by C'' , say by w' , vertices w and x_3 have $w' \in C''$ as a second common neighbor. Notice that in this case, vertices x_3, x_2, w, w' form a 4-cycle and $N(x_2) = \{x_1, x_3, w'\}$. However, now set $C'' \cup \{x_2, x_3\}$ is an identifying code in G . Indeed, x_2, x_3 and w' are separated from other vertices by Lemma 3. Moreover, vertex w' separates x_1 from the two other neighbors of x_2 . Therefore,

$$\begin{aligned} \gamma^{\text{ID}}(G) &\leq |C''| + 2 \\ &= \gamma^{\text{ID}}(G'') + 2 \\ &\leq \frac{2}{3}n'' + \frac{2}{3}(n-n'') \\ &= \frac{2}{3}n, \end{aligned}$$

which yields the desired upper bound.

Case 2: $F = P_4$. Let F be the path $x_1x_2x_3x_4$. If x_4 is adjacent to a vertex in the set A , then as before, there exists an induced path $P: x_1x_2x_3$ in F such that $G'' = G - V(P)$ is a connected graph of order at least 6 not in \mathcal{F}_3 . Then, the same arguments as in Case 1 apply and we obtain an identifying code of the desired size.

Hence, we may assume that x_4 has degree 1 in G (with x_3 as its unique neighbor in G). Analogously, we may assume that x_1 has degree 1 in G (with x_2 as its unique neighbor in G). Since G is connected, we may assume, renaming vertices if necessary, that x_2 is adjacent to a vertex in the set A .

We now consider the graph $G_F = G - \{x_3, x_4\}$. Let $n_F = n(G_F)$, and so $n_F = n - 2$. Let $\Delta(G_F) = \Delta_F$. We note that G_F is a connected triangle-free graph and $\Delta \geq \Delta_F \geq 3$. Thus, $\Delta_F = 3$. Moreover, G_F contains the cycle Q_{uv} and has order $n_F \geq 7$. In particular, $G_F \notin \mathcal{F}_3$. Applying the inductive hypothesis to G_F , we have

$$\gamma^{\text{ID}}(G_F) \leq \frac{2}{3}n_F.$$

Let C_F be an optimal identifying code in G_F , and so $|C_F| = \gamma^{\text{ID}}(G_F)$. If $x_2 \in C_F$, then in order to identify the vertices x_1 and x_2 , the code C_F contains at least one neighbor of x_2 in G_F . In this case, we let $C = C_F \cup \{x_3\}$. If $x_2 \notin C_F$, then $x_1 \in C_F$ and at least one neighbor of x_2 in the set A belongs to C_F . In this case, we let $C = (C_F \setminus \{x_1\}) \cup \{x_2, x_3\}$. In both cases (using Lemma 3 in the second case), C is an identifying code of G and $|C| = |C_F| + 1$, implying that

$$\begin{aligned} \gamma^{\text{ID}}(G) \leq |C| &= \gamma^{\text{ID}}(G_F) + 1 \\ &< \frac{2}{3}n_F + \frac{2}{3}(n - n_F) \\ &= \frac{2}{3}n. \end{aligned}$$

This completes the proof of Claim C. \square

By Claim C, we will from now on assume that if F is a component in G_{uv} of order at least 3, then $F \notin \mathcal{F}_\Delta$.

Let B be the set of all vertices that belong to a P_1 -component or to a P_2 -component in G_{uv} . Furthermore, let us denote $G^* = G[A \cup B \cup \{u, v\}]$. Note that it is possible that $G^* = G$ and hence we do not apply the induction hypothesis directly to G^* . Recall that $A = N_u \cup N_v$ and let $|A| = a$. Further, let $|B| = b$. Thus, $V(G^*) = A \cup B \cup \{u, v\}$ and $n^* = a + b + 2$. Next, in Claim D, we will show that G^* admits an identifying code of cardinality at most $\left(\frac{\Delta-1}{\Delta}\right)n^*$ that contains both vertices u and v . Notice that $G^* \notin \mathcal{F}_\Delta$ since G^* contains a 4-cycle and a vertex of degree at least 3.

Claim D. Graph G^* admits an identifying code containing vertices u and v of cardinality at most

$$\left(\frac{\Delta-1}{\Delta}\right)n^*.$$

Proof. Since G is a connected graph, every vertex in a P_1 -component of $G[B]$ is adjacent in G^* to at least one vertex in A . Moreover, by the connectivity of G^* , at least one vertex from every P_2 -component of $G[B]$ is adjacent in G^* to at least one vertex in A . However, it is possible that a P_2 -component of $G[B]$ contains exactly one vertex that has degree 1 in G^* (and is therefore not adjacent in G^* to a vertex from the set A). Let B_A be the set of all vertices in B that have at least one neighbor in A in the graph G^* , and let $B_L = B \setminus B_A$. Thus, each vertex in B_L is a vertex of degree 1 in G^* that belongs to a P_2 -component of $G[B]$ and is adjacent to no vertex of A in G^* . Possibly, $B_L = \emptyset$.

Let $B = (B_1, \dots, B_k)$ be a partition of the set B such that the following properties hold.

- All vertices in the set $B_i \cap B_A$ have the same neighborhood in the set A for all $i \in [k]$, that is, if $x, y \in B_i \cap B_A$, then $N_{G^*}(x) \cap A = N_{G^*}(y) \cap A \neq \emptyset$.
- Each vertex in the set $B_i \cap B_L$ has its unique neighbor (in G^*) in the set $B_i \cap B_A$ for all $i \in [k]$.
- Vertices in distinct sets $B_i \cap B_A$ and $B_j \cap B_A$ have different neighborhoods in A , that is, if $x \in B_i \cap B_A$ and $y \in B_j \cap B_A$ where $1 \leq i < j \leq k$, then $N_{G^*}(x) \cap A \neq N_{G^*}(y) \cap A$.




Figure 4: The setting of Claim D in the proof of Theorem 2. The aim is to construct a small identifying code containing both u and v .

An illustration is given in Figure 4. We note that $|B_i| = |B_i \cap B_A| + |B_i \cap B_L|$ for all $i \in [k]$. Moreover, $|B_i \cap B_L| \leq |B_i \cap B_A|$ and $|B_i \cap B_A| \geq 1$ for all $i \in [k]$. If $|B_i \cap B_L| = |B_i \cap B_A|$, then we observe that the subgraph $G[B_i]$ induced by the set B_i consists of vertex-disjoint copies of K_2 (and each such copy of K_2 contains exactly one vertex of degree 1 in G). Since each vertex in A is adjacent to either the vertex u or v , each vertex in A has at most $\Delta - 1$ neighbors in the set B , implying that $|B_i \cap B_A| \leq \Delta - 1$ for all $i \in [k]$. Let

$$|B_i| = b_i$$

for all $i \in [k]$. Thus, if $B_i \cap B_L = \emptyset$, then $b_i = |B_i \cap B_A| \leq \Delta - 1$, while if $B_i \cap B_L \neq \emptyset$, then $b_i \leq 2|B_i \cap B_A| \leq 2(\Delta - 1)$ for all $i \in [k]$. Recall that

$$b = |B| = \sum_{i=1}^k b_i.$$

Since G is triangle-free, we note that $B_i \cap B_A$ is an independent set in G^* for all $i \in [k]$. Thus, if two vertices w and z belong to the same P_2 -component in $G[B]$, then either $w \in B_i$ and $z \in B_j$ where $1 \leq i, j \leq k$ and $i \neq j$, or one of w and z belongs to $B_i \cap B_A$ and the other to $B_i \cap B_L$ for some $i \in [k]$. For $i \in [k]$, we now define the set B_i^* as follows. If $|B_i \cap B_L| = |B_i \cap B_A|$, then we define

$$B_i^* = B_i \cap B_L,$$

while if $|B_i \cap B_L| < |B_i \cap B_A|$, then we let w_i be an arbitrary vertex in the set $B_i \cap B_A$ that is isolated in the subgraph $G[B_i]$ induced by B_i (and so, w_i has no neighbor in B_i), and we define

$$B_i^* = (B_i \cap B_L) \cup \{w_i\}.$$

We now define

$$B^* = \bigcup_{i=1}^k B_i^*.$$

Moreover, let $|B_i^*| = b_i^*$ for all $i \in [k]$ and

$$b^* = \sum_{i=1}^k b_i^*.$$

An illustration of the set B^* is given in Figure 5.




Figure 5: The construction of the set B^* in the proof of Claim D. The vertices in B^* are shaded.

Claim D.1. The following hold.

- (a) $b_i^* \geq \frac{b_i}{\Delta-1}$ for all $i \in [k]$.
- (b) $b \leq (\Delta-1)b^*$.

Proof. By our earlier observations, if $B_i \cap B_L = \emptyset$, then $b_i \leq \Delta-1$ where $i \in [k]$. Moreover, $b_i \leq 2(\Delta-1)$ for all $i \in [k]$. If $b_i \leq \Delta-1$ for some $i \in [k]$, then

$$\frac{b_i - b_i^*}{b_i} \leq \frac{b_i - 1}{b_i} \leq \frac{\Delta - 2}{\Delta - 1}.$$

We note that if $b_i \geq \Delta$ for some $i \in [k]$, then $b_i^* \geq 2$, and so in this case

$$\frac{b_i - b_i^*}{b_i} \leq \frac{b_i - 2}{b_i} \leq \frac{2(\Delta - 1) - 2}{2(\Delta - 1)} = \frac{\Delta - 2}{\Delta - 1}.$$

Thus in both cases,

$$\frac{b_i - b_i^*}{b_i} \leq \frac{\Delta - 2}{\Delta - 1}.$$

Rearranging terms in the above inequality yields the inequality

$$b_i^* \geq \frac{b_i}{\Delta - 1}$$

for all $i \in [k]$. This proves property (a) in the statement of the claim. Hence,

$$b = \sum_{i=1}^k b_i \leq (\Delta - 1) \sum_{i=1}^k b_i^* = (\Delta - 1)b^*,$$

and so property (b) in the statement of the claim holds. \square

Let z_i be an arbitrary vertex in $B_i \cap B_A$, and so $z_i \in B_i$ and z_i has a neighbor in the set A for all $i \in [k]$. Let $Z = \{z_1, z_2, \dots, z_k\}$. By construction, the set A identifies the set Z since every pair of vertices in Z have distinct neighborhoods in A . Equivalently, the set Z is (Z, A) -identifiable by the set A . Let A^* be a subset of A of minimum cardinality that A -identifies Z . By Lemma 4, $1 \leq |A^*| \leq k$. Since $b_i^* \geq 1$ for all $i \in [k]$, we note that $b^* \geq k$, and so $|A^*| \leq b^*$. Recall that $Q_{uv} : uu'v'vu$ is a 4-cycle in G and hence in G^* , where u' is a neighbor of u in G different from v , and v' is a neighbor of v in G different from u .

Claim D.2. If $A = A^*$, then there exists an identifying code in G^* containing vertices u and v with cardinality at most

$$\left(\frac{\Delta - 1}{\Delta} \right) n^*.$$

Proof. Suppose that $A = A^*$. By the minimality of the set A^* , every vertex in A^* has a neighbor in B . Recall that $V(G^*) = A \cup B \cup \{u, v\}$ and $n^* = a + b + 2$. We now let

$$C^* = A \cup \{u, v\} \cup (B \setminus B^*).$$

The set C^* is an identifying code of G^* . Indeed, C^* is a dominating set which is connected in G^* . Thus, by Lemma 3, it separates vertices in C^* . Furthermore, each vertex in B^* has a unique neighborhood in $A \cup (B \setminus B^*)$.

Claim D.2(i). If $b^* \geq k + 2$, then $|C^*| \leq \left(\frac{\Delta-1}{\Delta}\right) n^*$.

Proof. Suppose that $b^* \geq k + 2$. Thus, $a = |A^*| \leq k \leq b^* - 2$. By Claim D.1(b) and our earlier observations, we have

$$\begin{aligned} |C^*| &= a + 2 + b - b^* \\ &= \left(\frac{\Delta-1}{\Delta}\right) (a + 2 + b) + \frac{1}{\Delta} (a + 2 + b - \Delta b^*) \\ &\leq \left(\frac{\Delta-1}{\Delta}\right) n^* + \frac{1}{\Delta} ((b^* - 2) + 2 + (\Delta - 1)b^* - \Delta b^*) \\ &= \left(\frac{\Delta-1}{\Delta}\right) n^*, \end{aligned}$$

yielding the desired upper bound. \square

By Claim D.2(i), we may assume that $b^* \leq k + 1$, implying that $b^* \in \{k, k + 1\}$. With this assumption, $b_i^* = 1$ for all $i \in [k]$, except for possibly exactly one value of i satisfying $b_i^* = 2$.

Recall that every vertex in A^* has a neighbor in B . Thus, since $A = A^*$, we note in particular that both vertices u' and v' have a neighbor in B . Since G^* is triangle-free, u' and v' do not have a common neighbor. Renaming the partitions in $B = (B_1, \dots, B_k)$ if necessary, we may assume that u' is adjacent to a vertex in B_1 and v' is adjacent to a vertex in B_2 . Since u' is adjacent to both u and v' , it has at most $\Delta - 2$ additional neighbors, implying that $|B_1 \cap B_A| \leq \Delta - 2$. Analogously, $|B_2 \cap B_A| \leq \Delta - 2$. Therefore,

$$1 \leq b_1^* \leq b_1 \leq 2(\Delta - 2) \quad \text{and} \quad 1 \leq b_2^* \leq b_2 \leq 2(\Delta - 2).$$

Let

$$B_{3,k} = B \setminus (B_1 \cup B_2) \quad \text{and} \quad B_{3,k}^* = B^* \setminus (B_1^* \cup B_2^*).$$

Further, let $b_{3,k} = |B_{3,k}|$ and $b_{3,k}^* = |B_{3,k}^*|$, and so $b = b_1 + b_2 + b_{3,k}$ and $b^* = b_1^* + b_2^* + b_{3,k}^*$. We note that

$$b_{3,k} = \sum_{i=3}^k b_i \quad \text{and} \quad b_{3,k}^* = \sum_{i=3}^k b_i^* \geq \sum_{i=3}^k 1 = k - 2.$$

We now define a partition (V_1, V_2, V_3) of $V(G^*)$ as follows. We let

$$\begin{aligned} V_1 &= B_1 \cup \{u, u'\}, \\ V_2 &= B_2 \cup \{v, v'\}, \\ V_3 &= B_{3,k} \cup (A \setminus \{u', v'\}). \end{aligned}$$

Further we define $n_i = |V_i|$ for $i \in [3]$, and so $n^* = n_1 + n_2 + n_3$. We note that $n_1 = b_1 + 2$, $n_2 = b_2 + 2$, and $n_3 = b_{3,k} + a - 2$. We again consider the identifying code C^* of G^* and we let $C_i^* = C^* \cap V_i$ for $i \in [3]$. Thus,

$$|C^*| = \sum_{i=1}^3 |C_i^*|.$$

Claim D.2(ii). $|C_3^*| \leq \left(\frac{\Delta-1}{\Delta}\right) n_3$.

Proof. By Claim D.1(a), we have

$$b_{3,k}^* = \sum_{i=3}^k b_i^* \geq \sum_{i=3}^k \frac{b_i}{\Delta-1} = \frac{b_{3,k}}{\Delta-1},$$

and so $(\Delta-1)b_{3,k}^* \geq b_{3,k}$, or, equivalently,

$$b_{3,k} - \Delta b_{3,k}^* \leq -b_{3,k}^* \leq -k + 2. \quad (1)$$

Recall that $n_3 = a - 2 + b_{3,k}$ and that $a \leq k$. Hence by Inequality (1), we have

$$\begin{aligned} |C_3^*| &= |A \setminus \{u', v'\}| + b_{3,k} - b_{3,k}^* \\ &= a - 2 + b_{3,k} - b_{3,k}^* \\ &= \left(\frac{\Delta-1}{\Delta}\right) (a - 2 + b_{3,k}) + \frac{1}{\Delta} (a - 2 + b_{3,k} - \Delta b_{3,k}^*) \\ &\leq \left(\frac{\Delta-1}{\Delta}\right) n_3 + \frac{1}{\Delta} (k - 2 - k + 2) \\ &= \left(\frac{\Delta-1}{\Delta}\right) n_3, \end{aligned}$$

yielding the desired upper bound. \square

Claim D.2(iii). If $b_i^* \geq \frac{b_i}{\Delta-2}$, then $|C_i^*| \leq \left(\frac{\Delta-1}{\Delta}\right) n_i$ for $i \in [2]$.

Proof. Let $i \in [2]$. Recall that $b_i^* \geq 1$ and $n_i = b_i + 2$. Suppose that $b_i^* \geq \frac{b_i}{\Delta-2}$. Thus, $(\Delta-2)b_i^* \geq b_i$, or, equivalently, $b_i - \Delta b_i^* \leq -2b_i^* \leq -2$ noting that $b_i^* \geq 1$. By definition, we have $C_1^* = \{u, u'\} \cup (B_1 \setminus B_1^*)$ and $C_2^* = \{v, v'\} \cup (B_2 \setminus B_2^*)$. Hence,

$$\begin{aligned} |C_i^*| &= 2 + b_i - b_i^* \\ &= \left(\frac{\Delta-1}{\Delta}\right) (2 + b_i) + \frac{1}{\Delta} (2 + b_i - \Delta b_i^*) \\ &\leq \left(\frac{\Delta-1}{\Delta}\right) n_i + \frac{1}{\Delta} (2 - 2) \\ &= \left(\frac{\Delta-1}{\Delta}\right) n_i, \end{aligned}$$

yielding the desired upper bound. \square

If $b_i^* \geq \frac{b_i}{\Delta-2}$ for $i \in [2]$, then by Claims D.2(ii) and D.2(iii), we have

$$|C^*| = \sum_{i=1}^3 |C_i^*| \leq \sum_{i=1}^3 \left(\frac{\Delta-1}{\Delta}\right) n_i = \left(\frac{\Delta-1}{\Delta}\right) n^*,$$

yielding the desired upper bound. Hence, we may assume that $b_i^* < \frac{b_i}{\Delta-2}$ for some $i \in [2]$. By symmetry, we may assume that $b_1^* < \frac{b_1}{\Delta-2}$. Let

$$|B_1 \cap B_A| = t_1 + t_2 \quad \text{and} \quad |B_1 \cap B_L| = t_2.$$

We note that $1 \leq t_1 + t_2 \leq \Delta - 2$.

If $t_1 \geq 1$ and $t_2 \geq 1$, then $b_1 = t_1 + 2t_2$ and $b_1^* = 1 + t_2 \geq 2$, implying that

$$\frac{b_1}{\Delta - 2} = \frac{t_2 + (t_1 + t_2)}{\Delta - 2} \leq \frac{t_2 + (\Delta - 2)}{\Delta - 2} = \frac{t_2}{\Delta - 2} + 1 \leq \frac{\Delta - 3}{\Delta - 2} + 1 < 2 \leq b_1^*.$$

If $t_1 \geq 1$ and $t_2 = 0$, then $b_1 = t_1 \leq \Delta - 2$ and $b_1^* = 1$, implying that

$$\frac{b_1}{\Delta - 2} \leq \frac{\Delta - 2}{\Delta - 2} = 1 = b_1^*.$$

If $t_1 = 0$ and $t_2 \geq 2$, then $b_1 = 2t_2$ and $b_1^* = t_2 \geq 2$. Moreover, $t_2 \leq \Delta - 2$. Thus,

$$\frac{b_1}{\Delta - 2} = \frac{2t_2}{\Delta - 2} \leq \frac{2(\Delta - 2)}{\Delta - 2} = 2 \leq b_1^*.$$

If $t_1 = 0$, $t_2 = 1$ and $\Delta \geq 4$, then $b_1 = 2$, $b_1^* = 1$ and

$$\frac{b_1}{\Delta - 2} = \frac{2}{\Delta - 2} \leq \frac{2}{2} = 1 = b_1^*.$$

In all the above four cases, we contradict our assumption that $b_1^* < \frac{b_1}{\Delta - 2}$. Hence, $t_1 = 0$, $t_2 = 1$, and $\Delta = 3$. In this case, B_1 induces a P_2 -component in G^* . Let $B_1 \cap B_A = \{u_1\}$ and let $B_1 \cap B_L = \{u_2\}$, and so $uu'u_1u_2$ is a path in G^* . We note that u_2 is a vertex of degree 1 in G^* . Since $\Delta = 3$ and v' is adjacent to v and u' , we note that either $B_2 = P_1$ or $B_2 = P_2$. We denote the vertex in $B_2 \cap N(v')$ by v_1 and if v_1 has another adjacent vertex outside of A , we denote it by v_2 . By our choice of u and v (maximizing $\deg_G(u) + \deg_G(v)$ for all adjacent u, v with uv a cycle edge), we note that both u and v have degree 3 in G^* (otherwise, $\deg_G(u') + \deg_G(v') > \deg_G(u) + \deg_G(v)$). We denote the third neighbor of u and v by u'' and v'' , respectively.

Let us assume next that $|N(u_1) \cap A| = 2$. Since G is triangle-free, the other vertex in $N(u_1) \cap A$ is not v' . Notice that in this case, we may remove u' from C_1^* , resulting in C_1^{**} , and the set $C^{**} = C_1^{**} \cup C_2^* \cup C_3^*$ is an identifying code in G^* . Indeed, C^* was an identifying code in G^* . Moreover, every neighbor of u' is in C^{**} and vertices in C^{**} induce a single component in $G[C^{**}]$. Hence, every vertex in C^{**} is separated by Lemma 3 while u' is the unique vertex not in C^{**} adjacent to vertices u and u_1 . Notice that $|C^{**}| \leq \frac{2}{3}n^*$. Indeed, we have $n_1 = 4$, $|C_1^{**}| = 2$, $(n_2, |C_2^*|) \in \{(3, 2), (4, 3)\}$ and $|C_3^*| \leq \frac{2}{3}n_3$ by Claim D.2(ii). Since $\frac{|C_1^{**}| + |C_2^*|}{n_1 + n_2} \leq \frac{5}{8} < \frac{2}{3}$, we have $|C^{**}| \leq \frac{2}{3}n^*$. Therefore, we may assume that one of B_1 or B_2 is a P_2 -component such that the vertex in B_A is adjacent to exactly one vertex in A . We may assume from now on without loss of generality, that $N(u_1) \cap A = \{u'\}$.

Recall that in graph G' , any minimum-size identifying code C' is such that it does not separate either vertices $\{u, v'\}$ or $\{v, u'\}$ in G . Thus, $C' \cap \{u, v, v', u'\}$ is either $\{u, v'\}$ or $\{v, u'\}$. Furthermore, since u_2 is a leaf and u_1 is adjacent only to u' , the only vertex which can separate u_1 and u_2 is u' . Therefore, $u' \in C'$. Thus, $C' \cap \{u, v, v', u'\} = \{u', v\}$ and $\{u, v'\}$ are not separated in G . Furthermore, we have $u'' \notin C'$ (otherwise $\{u, v'\}$ would be separated by C' in G). Notice that since C' separates u' and u , we have $u_1, u_2 \in C'$. However, now the set $C = \{u\} \cup (C' \setminus \{u_1\})$ is an identifying code of claimed cardinality in G . Indeed, vertices $u', u, v \in C$ are adjacent and are thus separated by Lemma 3. Moreover, also vertices u_1 and u_2 have unique code neighborhoods. This contradicts the properties of G and completes the proof of Claim D.2. \square

Recall that

$$1 \leq |A^*| \leq k \leq b^* \leq b.$$

By Claim D.2, we may assume that $A^* \subset A$, and so $1 \leq |A^*| < |A| = a \leq 2(\Delta - 1)$. Let $\overline{A^*} = A \setminus A^*$. Thus, $|\overline{A^*}| = |A| - |A^*| \geq 1$. We note that either $|\overline{A^*}| \leq \Delta - 2$ or $|\overline{A^*}| \geq \Delta - 1$. Further, either $\overline{A^*}$ contains a neighbor of u and a neighbor of v or $\overline{A^*} \subset N_{G^*}(u)$ or $\overline{A^*} \subset N_{G^*}(v)$. We proceed further with three claims.

Claim D.3. If $|\overline{A^*}| \leq \Delta - 2$, then there exists an identifying code in G^* containing vertices u and v with cardinality at most

$$\left(\frac{\Delta - 1}{\Delta}\right)n^*.$$

Proof. Suppose that $|\overline{A^*}| \leq \Delta - 2$. As observed earlier, $|A^*| \leq b^*$. Thus in this case, $a = |A| = |A^*| + |\overline{A^*}| \leq \Delta + b^* - 2$. By Claim D.1(b), we have $b - (\Delta - 1)b^* \leq 0$. Let x be an arbitrary vertex in $\overline{A^*}$ and let

$$C^* = V(G) \setminus (B^* \cup \{x\}).$$

The code C^* is an identifying code of G^* . Indeed, C^* is a dominating set in G^* . Furthermore, it is also connected and hence, by Lemma 3, separates every vertex in C^* . Furthermore, x is the only vertex in A which does not belong to set C^* and it is separated by $\{u, v\}$ from vertices in B . Finally, vertices in B^* are separated by $A^* \cup \{B \setminus B^*\}$. Next, we consider the cardinality of C^* :

$$\begin{aligned} |C^*| &= (a + b + 2) - (b^* + 1) \\ &= \left(\frac{\Delta - 1}{\Delta}\right)(a + b + 2) + \frac{1}{\Delta}(a + b - (\Delta - 2) - \Delta b^*) \\ &\leq \left(\frac{\Delta - 1}{\Delta}\right)n^* + \frac{1}{\Delta}((\Delta + b^* - 2) + b - \Delta + 2 - \Delta b^*) \\ &\leq \left(\frac{\Delta - 1}{\Delta}\right)n^* + \frac{1}{\Delta}(b - (\Delta - 1)b^*) \\ &\leq \left(\frac{\Delta - 1}{\Delta}\right)n^*, \end{aligned}$$

yielding the desired upper bound. \square

Claim D.4. If $\overline{A^*}$ contains a neighbor of u and a neighbor of v , then there exists an identifying code in G^* containing vertices u and v with cardinality at most

$$\left(\frac{\Delta - 1}{\Delta}\right)n^*.$$

Proof. Suppose that $\overline{A^*}$ contains a neighbor, u_1 , of u and a neighbor, v_1 , of v . By our earlier observations, $|\overline{A^*}| = a - |A^*| \leq 2(\Delta - 1) - 1 = 2\Delta - 3$. Hence, $a = |A^*| + |\overline{A^*}| \leq b^* + 2\Delta - 3$. By Claim D.1(b), we have $b - (\Delta - 1)b^* \leq 0$. We now let

$$C^* = V(G^*) \setminus (B^* \cup \{u_1, v_1\}).$$

The set C^* is an identifying code of G^* . Indeed, C^* is a dominating set in G^* . Furthermore, it is also connected and hence, by Lemma 3, separates every vertex in C^* . Furthermore, u_1 and v_1 are the only vertices in A which do not belong to set C^* and are separated by $\{u, v\}$ from vertices in B and from each

other. Finally, vertices in B^* are separated by $A^* \cup \{B \setminus B^*\}$. Next, we consider the cardinality of C^* :

$$\begin{aligned}
|C^*| &= (a + b + 2) - (b^* + 2) \\
&= \left(\frac{\Delta - 1}{\Delta}\right) (a + b + 2) + \frac{1}{\Delta} (a + b - 2(\Delta - 1) - \Delta b^*) \\
&\leq \left(\frac{\Delta - 1}{\Delta}\right) n^* + \frac{1}{\Delta} ((b^* + 2\Delta - 3) + b - 2(\Delta - 1) - \Delta b^*) \\
&\leq \left(\frac{\Delta - 1}{\Delta}\right) n^* + \frac{1}{\Delta} (b - (\Delta - 1)b^* - 1) \\
&\leq \left(\frac{\Delta - 1}{\Delta}\right) n^* - \frac{1}{\Delta} \\
&< \left(\frac{\Delta - 1}{\Delta}\right) n^*,
\end{aligned}$$

yielding the desired upper bound. \square

By Claim D.3, we may assume that $|\overline{A^*}| \geq \Delta - 1$, for otherwise there exists an identifying code of size at most $\left(\frac{\Delta - 1}{\Delta}\right) n^*$ in G^* containing vertices u and v .

Claim D.5. If $\overline{A^*} \subseteq N_u$ or $\overline{A^*} \subseteq N_v$, then there exists an identifying code in G^* containing vertices u and v with cardinality at most

$$\left(\frac{\Delta - 1}{\Delta}\right) n^*.$$

Proof. Suppose that either $\overline{A^*} \subseteq N_u$ or $\overline{A^*} \subseteq N_v$. Renaming u and v if necessary, we may assume that $\overline{A^*} \subseteq N_v$, and so $\overline{A^*}$ contains no neighbor of u . Since $\overline{A^*} \subseteq N_v$, we have $|\overline{A^*}| \leq |N_{G^*}(v)| - 1 \leq \Delta - 1$. However by our earlier assumption due to Claim D.3, $|\overline{A^*}| \geq \Delta - 1$. Therefore, $|\overline{A^*}| = \Delta - 1$, that is, $\deg_{G^*}(v) = \Delta$ and $\overline{A^*} = N_{G^*}(v) \setminus \{u\}$. Thus,

$$a = |A| = |A^*| + |\overline{A^*}| \leq b^* + \Delta - 1.$$

Recall that $u'v'$ is an edge in G^* , where u' is a neighbor of u in G^* different from v and v' is a neighbor of v in G^* different from u . Since $u' \in A^*$, the vertex u' is adjacent to at least one vertex in B , and therefore u' is adjacent to at most $\Delta - 2$ vertices in $\overline{A^*}$. Let v'' be a vertex in $\overline{A^*}$ that is not adjacent to the vertex u' . By Claim D.1(b), we have $b - (\Delta - 1)b^* \leq 0$. We now let

$$C^* = V(G^*) \setminus (B^* \cup \{v', v''\}).$$

The set C^* is an identifying code of G^* . Indeed, C^* is a dominating set in G^* . Furthermore, it is also connected and hence, by Lemma 3, separates every vertex in C^* . Furthermore, v' and v'' are the only vertices in A which do not belong to set C^* and are separated by v from vertices in B and by u' from each other. Finally, vertices in B^* are separated by $A^* \cup \{B \setminus B^*\}$. Next, we consider the cardinality of

C^* :

$$\begin{aligned}
|C^*| &= (a + b + 2) - (b^* + 2) \\
&= \left(\frac{\Delta - 1}{\Delta}\right)(a + b + 2) + \frac{1}{\Delta}(a + b - 2(\Delta - 1) - \Delta b^*) \\
&\leq \left(\frac{\Delta - 1}{\Delta}\right)n^* + \frac{1}{\Delta}((b^* + \Delta - 1) + b - 2(\Delta - 1) - \Delta b^*) \\
&\leq \left(\frac{\Delta - 1}{\Delta}\right)n^* + \frac{1}{\Delta}(b - (\Delta - 1)b^* - (\Delta - 1)) \\
&\leq \left(\frac{\Delta - 1}{\Delta}\right)n^* - \left(\frac{\Delta - 1}{\Delta}\right) \\
&< \left(\frac{\Delta - 1}{\Delta}\right)n^*,
\end{aligned}$$

yielding the desired upper bound. \square

By Claims D.2, D.3, D.4 and D.5, we have shown that graph G^* admits an identifying code of cardinality at most $\left(\frac{\Delta-1}{\Delta}\right)n^*$ containing vertices u and v . Hence, Claim D follows. \square

If $G^* = G$, then the theorem statement follows from Claim D. If $G^* \neq G$, then graph G_{uv} contains a component of cardinality at least 3. Furthermore, by Claim C, graph G_{uv} does not contain any components that belong to the set \mathcal{F}_Δ . Hence, we may assume that G_{uv} contains a component of order at least 3. Next, we finalize the proof by showing that also in this case, $\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n$.

Let us denote by \mathcal{K} the set of components of order at least 3 in G_{uv} . By Claim C, $K \notin \mathcal{F}_\Delta$ for any $K \in \mathcal{K}$. Applying the inductive hypothesis to a component $K \in \mathcal{K}$ of maximum degree Δ_K , component K satisfies

$$\gamma^{\text{ID}}(K) \leq \left(\frac{\Delta_K - 1}{\Delta_K}\right)n(K) \leq \left(\frac{\Delta - 1}{\Delta}\right)n(K). \quad (2)$$

Let C_K be an identifying code in K with minimum cardinality.

Observe that $G^* = G - \bigcup_{K \in \mathcal{K}} V(K)$. Let C^* be an identifying code of G^* with cardinality at most $\left(\frac{\Delta-1}{\Delta}\right)n^*$ containing vertices u and v , that exists by Claim D.

Let us next consider set $C = C^* \cup \left(\bigcup_{K \in \mathcal{K}} C_K\right)$. Notice that we have

$$\gamma^{\text{ID}}(G) \leq |C| = |C^*| + \sum_{K \in \mathcal{K}} |C_K| \leq \left(\frac{\Delta - 1}{\Delta}\right)n^* + \sum_{K \in \mathcal{K}} \left(\frac{\Delta - 1}{\Delta}\right)n(K) = \left(\frac{\Delta - 1}{\Delta}\right)n.$$

Set C is an identifying code in G since set C^* contains vertices u and v . Indeed, since C an union of multiple identifying codes, each identifying code dominates and pairwise separates the vertices within the corresponding component. Thus, the only problems might be between vertices of different components. In particular, in the case the component K contains vertex $z \in C_K$ such that $N[z] \cap C_K = \{z\}$, component G^* contains vertex $y \in C^*$ such that $N[y] \cap C^* = \{y\}$, and y and z are adjacent in G . However, this is not possible since $y \in A$ and thus, $N[y] \cap C^* \cap \{u, v\} \neq \emptyset$.

This completes the proof of Theorem 2. \square

As an immediate consequence of Theorem 2, we have the following corollary.

Corollary 15. If G is a connected, identifiable, triangle-free graph of order n , then the following holds.

- (a) If G is cubic, then $\gamma^{\text{ID}}(G) \leq \frac{2}{3}n$.
- (b) If G is subcubic and $n \geq 23$, then $\gamma^{\text{ID}}(G) \leq \frac{2}{3}n$.
- (c) If G has maximum degree Δ where $\Delta \geq 4$ is fixed and $n \geq \Delta + 2$, then $\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n$.

4 Beyond triangle-free graphs

We now apply Theorem 2 to graphs having triangles and obtain a bound weaker than the conjectured one, as follows.

Corollary 16. If G is a connected identifiable graph of order $n \geq 3$ with maximum degree $\Delta \geq 3$ such that G can be made triangle-free by deleting t edges, then

$$\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n + 4t + \frac{1}{\Delta}.$$

Proof. We prove the claim by induction on t . Let us assume that $E_t \subseteq E(G)$, with $|E_t| = t$, is a smallest set of edges which we can remove from G so that $G_t = G - E_t$ is triangle-free. Observe that G_t is connected and identifiable, since the only connected triangle-free graph that is not identifiable is K_2 , but we assume here that $n \geq 3$. By Table 1 and Theorem 2, the claim holds when G is triangle-free, that is, for $t = 0$. Assume next that the claim holds for every $t \leq t'$ and let $t = t' + 1$. Assume that G_t has maximum degree $\Delta_t \leq \Delta$. We have $\Delta_t \geq 2$ since G_t is connected and $n \geq 3$.

Let C_t be an optimal identifying code in G_t . By Theorem 2, if $\Delta_t \geq 3$, we have

$$|C_t| \leq \left(\frac{\Delta_t-1}{\Delta_t}\right)n + \frac{1}{\Delta_t} \leq \left(\frac{\Delta-1}{\Delta}\right)n + \frac{1}{\Delta}.$$

If $\Delta_t = 2$, by Corollary 7(e)-(f) and $n \notin \{4, 7\}$, then $|C_t| \leq \frac{2}{3}n \leq \left(\frac{\Delta-1}{\Delta}\right)n$ since $\Delta \geq 3$. If $n = 4$, $|C_t| = 3 \leq \left(\frac{\Delta-1}{\Delta}\right)n + \frac{1}{\Delta}$ and if $n = 7$, $|C_t| \leq 5 \leq \left(\frac{\Delta-1}{\Delta}\right)n + \frac{1}{\Delta}$ (again since $\Delta \geq 3$). Let edge $uv \in E_t$ and let us consider graph G_t together with edge uv , denoted by G'_t . Observe that if C_t is not an identifying code in G'_t , then the addition of edge uv modified some code-neighborhoods (this implies that u or v is in C_t). Therefore, there are at most four vertices which are no longer separated (possibly, vertex u together with some vertex u' , and possibly, vertex v with some vertex v'). As we add these edges back one at a time, each time we create at most four new vertices among which some vertex-pairs are not separated by C_t . Thus, in the graph G , there is a set S of at most $4t$ vertices in which some vertex-pairs are not separated by C_t . However, since G is identifiable, S is $V(G)$ -identifiable and thus, by Lemma 4, we can find an $(S, V(G))$ -identifying code C_S of size at most $|S| \leq 4t$. The set $C_t \cup C_S$ is an identifying code of G of the desired size, proving the claim. \square

5 Conclusion

We proved Conjecture 1 for all triangle-free graphs. In fact, we proved (see Theorem 2) the following stronger result. Let G be a connected, identifiable, triangle-free graph of order $n \geq 3$ with maximum degree Δ . If $\Delta \geq 4$, then we proved that $\gamma^{\text{ID}}(G) \leq \left(\frac{\Delta-1}{\Delta}\right)n$, except for one exceptional graph, namely the star $K_{1,\Delta}$. Moreover, if $\Delta \leq 3$, then $\gamma^{\text{ID}}(G) \leq \frac{2}{3}n$, unless G belongs to a forbidden family that contains fifteen graphs (all of order at most 22): P_4 , C_4 , C_7 and the twelve trees of maximum degree 3 from \mathcal{T}_3 .

In the special case when G is a triangle-free cubic graph, this implies that $\gamma^{\text{ID}}(G) \leq \frac{2}{3}n$ always holds. This establishes a best possible upper bound for triangle-free cubic graphs, since $\gamma^{\text{ID}}(K_{3,3}) = 4$ (we do not know if other cubic graphs for which this bound is tight exist). The previously best known upper bounds (prior to this paper) when G is triangle-free subcubic and cubic were, $\gamma^{\text{ID}}(G) \leq \frac{8}{9}n$ and $\gamma^{\text{ID}}(G) \leq \frac{5}{6}n$, respectively (see [10, Corollary 4.46]; the proof used the technique developed in [13]).

Towards a positive resolution of Conjecture 1, it would be interesting to prove it for all cubic graphs.

When it comes to general graphs, the list of exceptional graphs is larger than \mathcal{F}_Δ . Indeed, as mentioned in Table 1, the complements of half-graphs and related constructions defined in [12] do require $c > 0$. Nevertheless, those constructions have maximum degree Δ very close to the number n of vertices ($n - 1$ or $n - 2$), and, for any given Δ , there is only a finite number of such examples. Thus, it is possible that even for the general case, if the conjecture is true, the list of graphs requiring $c > 0$ is also finite for every fixed value of Δ .

As seen in Table 1, all graphs known to us that require $c > 0$ have $c \leq 3/2$ (which is reached only by odd cycles), and when $\Delta \geq 3$, in fact $c \leq 1/3$. Are there graphs that require higher values of c ? Another way to formulate these constants is in terms of the maximum degree. Do there exist any graphs that require the constant c to be larger than $3/\Delta$? By our results, such graphs would necessarily contain triangles. Note that it seems necessary to understand those graphs needing $c > 0$, in order to prove the conjecture.

References

- [1] L. Babai. On the complexity of canonical labeling of strongly regular graphs. *SIAM Journal on Computing*, 9(1):212–216, 1980.
- [2] C. Balbuena, F. Foucaud, and A. Hansberg. Locating-dominating sets and identifying codes in graphs of girth at least 5. *The Electronic Journal of Combinatorics*, 22(2):P2, 2015.
- [3] N. Bertrand, I. Charon, O. Hudry, and A. Lobstein. Identifying and locating-dominating codes on chains and cycles. *European Journal of Combinatorics*, 25(7):969–987, 2004.
- [4] J. A. Bondy. Induced subsets. *Journal of Combinatorial Theory, Series B*, 12(2):201–202, 1972.
- [5] D. Chakraborty, F. Foucaud, M. A. Henning, and T. Lehtilä. Identifying codes in graphs of given maximum degree. I. Trees. *arXiv preprint arXiv:2403.13172*, 2024.
- [6] D. Chakraborty, F. Foucaud, and T. Lehtilä. Identifying codes in bipartite graphs of given maximum degree. *Procedia Computer Science*, 223:157–165, 2023.
- [7] E. Charbit, I. Charon, G. D. Cohen, O. Hudry, and A. Lobstein. Discriminating codes in bipartite graphs: bounds, extremal cardinalities, complexity. *Advances in Mathematics of Communication*, 2(4):403–420, 2008.
- [8] J. Devin and A. Lobstein. Watching systems, identifying, locating-dominating and discriminating codes in graphs: a bibliography. *Published electronically at <https://dragazo.github.io/bibdom/main.pdf>*.
- [9] P. Erdős. Some combinatorial, geometric and set theoretic problems in measure theory. In *Measure Theory Oberwolfach 1983*, pages 321–327. Springer, 1984.
- [10] F. Foucaud. *Combinatorial and algorithmic aspects of identifying codes in graphs*. PhD thesis, Université Bordeaux 1, 2012.
- [11] F. Foucaud, S. Gravier, R. Naserasr, A. Parreau, and P. Valicov. Identifying codes in line graphs. *Journal of Graph Theory*, 73(4):425–448, 2013.
- [12] F. Foucaud, E. Guerrini, M. Kovše, R. Naserasr, A. Parreau, and P. Valicov. Extremal graphs for the identifying code problem. *European Journal of Combinatorics*, 32(4):628–638, 2011.
- [13] F. Foucaud, R. Klasing, A. Kosowski, and A. Raspaud. On the size of identifying codes in triangle-free graphs. *Discrete Applied Mathematics*, 160(10-11):1532–1546, 2012.
- [14] F. Foucaud and T. Lehtilä. Revisiting and improving upper bounds for identifying codes. *SIAM Journal on Discrete Mathematics*, 36:2619–2634, 2022.
- [15] F. Foucaud and G. Perarnau. Bounds for identifying codes in terms of degree parameters. *The Electronic Journal of Combinatorics*, 19(1):32, 2012.
- [16] W. Goddard and K. Wash. ID codes in Cartesian products of cliques. *Journal of Combinatorial Mathematics and Combinatorial Computing*, 85:97–106, 2013.

- [17] S. Gravier, M. Kovše, M. Mollard, J. Moncel, and A. Parreau. New results on variants of covering codes in Sierpiński graphs. *Designs, Codes and Cryptography*, 69(2):181–188, 2013.
- [18] S. Gravier and J. Moncel. On graphs having a $v \setminus \{x\}$ set as an identifying code. *Discrete Mathematics*, 307(3-5):432–434, 2007.
- [19] S. Gravier, J. Moncel, and A. Semri. Identifying codes of cycles. *European Journal of Combinatorics*, 27(5):767–776, 2006.
- [20] G. Z. Gutin, M. S. Ramanujan, F. Reidl, and M. Wahlström. Alternative parameterizations of metric dimension. *Theoretical Computer Science*, 806:133–143, 2020.
- [21] T. W. Haynes, S. T. Hedetniemi, and M. A. Henning, editors. *Topics in Domination in Graphs*, volume 64 of *Developments in Mathematics*. Springer, Cham, 2020.
- [22] T. W. Haynes, S. T. Hedetniemi, and M. A. Henning, editors. *Structures of Domination in Graphs*, volume 66 of *Developments in Mathematics*. Springer, Cham, 2021.
- [23] T. W. Haynes, S. T. Hedetniemi, and M. A. Henning. *Domination in Graphs: Core Concepts*. Springer Monographs in Mathematics. Springer, Cham, 2023.
- [24] V. Junnila, T. Laihonon, and T. Lehtilä. On a conjecture regarding identification in Hamming graphs. *The Electronic Journal of Combinatorics*, page P2, 2019.
- [25] M. G. Karpovsky, K. Chakrabarty, and L. B. Levitin. On a new class of codes for identifying vertices in graphs. *IEEE Transactions on Information Theory*, 44(2):599–611, 1998.
- [26] A. Lobstein, O. Hudry, and I. Charon. Locating-domination and identification. *Topics in Domination in Graphs*, pages 251–299, 2020.
- [27] B. M. E. Moret and H. D. Shapiro. On minimizing a set of tests. *SIAM Journal on Scientific and Statistical Computing*, 6(4):983–1003, 1985.
- [28] O. Pikhurko, H. Veith, and O. Verbitsky. The first order definability of graphs: Upper bounds for quantifier depth. *Discrete Applied Mathematics*, 154(17):2511–2529, 2006.
- [29] D. F. Rall and P. J. Slater. On location-domination numbers for certain classes of graphs. *Congressus Numerantium*, 45:97–106, 1984.
- [30] D. F. Rall and K. Wash. Identifying codes of the direct product of two cliques. *European Journal of Combinatorics*, 36:159–171, 2014.
- [31] B. Reed. Paths, stars and the number three. *Combinatorics, Probability and Computing*, 5(3):277–295, 1996.
- [32] A. Rényi. On random generating elements of a finite Boolean algebra. *Acta Scientiarum Mathematicarum Szeged*, 22:75–81, 1961.
- [33] R. D. Skaggs. *Identifying vertices in graphs and digraphs*. PhD thesis, 2007.
- [34] R. Ungrangsi, A. Trachtenberg, and D. Starobinski. An implementation of indoor location detection systems based on identifying codes. In F. A. Aagesen, C. Anutariya, and V. Wuwongse, editors, *Intelligence in Communication Systems*, pages 175–189, Berlin, Heidelberg, 2004. Springer Berlin Heidelberg.
- [35] A. J. Winter. Another algebraic proof of Bondy’s theorem on induced subsets. *Journal of Combinatorial Theory, Series A*, 89(1):145–147, 2000.