

Small planar hypohamiltonian graphs

Cheng-Chen Tsai*

Institute of Information Science, Academia Sinica, Taiwan

Abstract

A graph is hypohamiltonian if it is non-Hamiltonian, but the deletion of every single vertex gives a Hamiltonian graph. Until now, the smallest known planar hypohamiltonian graph had 40 vertices, a result due to Jooyandeh, McKay, Östergård, Pettersson, and Zamfirescu. That result is here improved upon by two planar hypohamiltonian graphs on 34 vertices. We exploited a special subgraph contained in two graphs of Jooyandeh et al., and modified it to construct the two 34-vertex graphs and six planar hypohamiltonian graphs on 37 vertices. Each of the 34-vertex graphs has 26 cubic vertices, improving upon the result of Jooyandeh et al. that planar hypohamiltonian graphs have 30 cubic vertices. We use the 34-vertex graphs to construct hypohamiltonian graphs of order 34 with crossing number 1, improving the best-known bound of 36 due to Wiener. Whether there exists a planar hypohamiltonian graph on 41 vertices was an open question. We settled this question by applying an operation introduced by Thomassen to the 37-vertex graphs to obtain several planar hypohamiltonian graphs on 41 vertices. The 25 planar hypohamiltonian graphs on 40 vertices of Jooyandeh et al. have no nontrivial automorphisms. The result is here improved upon by six planar hypohamiltonian graphs on 40 vertices with nontrivial automorphisms.

1 Introduction

A graph is *hypohamiltonian* if it is non-Hamiltonian, but the deletion of every single vertex gives a Hamiltonian graph. A graph $G = (V, E)$ is *almost hypohamiltonian* if there exists a $w \in V$ such that $G - w$ is non-Hamiltonian, but for every vertex $v \neq w$ in G , the graph $G - v$ is hamiltonian. We shall call w the *exceptional vertex* of G .

In 1973, Chvátal [2] raised the problem whether there exist planar hypohamiltonian graphs, and offered \$5 for its solution (Problem 19 of [3]). In 1974, Grünbaum [8] conjectured that there are no planar hypohamiltonian graphs. In 1976, Thomassen [14] gave infinitely many counterexamples, the smallest being of order 105. Various authors have improved this upper bound for the order of the smallest planar hypohamiltonian graph to order 57 (Hatzel [9] in 1979), 48

*Email address: u_e_smvdv@yahoo.com.tw

(C. T. Zamfirescu and T. Zamfirescu [25] in 2007), 42 (Araya and Wiener [20] in 2011), and 40 (Jooyandeh et al. [10] in 2017).

The lower bound for the order of the smallest planar hypohamiltonian graph was given by J. Goedgebeur and C. T. Zamfirescu [6].

Theorem 1.1 ([6, Theorem 3.2]). *The smallest planar hypohamiltonian graph has at least 23 vertices.*

Theorem 1.2 ([6, Theorem 3.3]). *The smallest planar hypohamiltonian graph with girth at least 4 has at least 27 vertices.*

J. Goedgebeur and C. T. Zamfirescu [6] defined h (h_g) to be the order of the smallest planar hypohamiltonian graph (of girth g). Combining [Theorem 1.1](#) and [Theorem 1.2](#) with the result of [10], we have $23 \leq h \leq 40$, $27 \leq h_4 \leq 40$, and $h_5 = 45$ [6, Corollary 3.4].

T. Zamfirescu [26] defined C_k^i and P_k^i to be the smallest order for which there is a planar k -connected graph such that every set of i vertices is disjoint from some longest cycle and path, respectively. The best bounds known so far were $C_3^1 \leq 40$, $C_3^2 \leq 2625$, $P_3^1 \leq 156$, $P_3^2 \leq 10350$, which were found based on a planar hypohamiltonian graph on 40 vertices [10], and a cubic planar hypohamiltonian graph on 70 vertices [1].

C. T. Zamfirescu [21] defined $\bar{\alpha}_0$ ($\bar{\alpha}_1$) to be the order of the smallest planar hypohamiltonian (smallest planar almost hypohamiltonian) graph. To emphasize the girth g , we use $\bar{\alpha}_{0,g}$ and $\bar{\alpha}_{1,g}$. By [Theorem 1.1](#), [Theorem 1.2](#) and the result of [10], we have $23 \leq \bar{\alpha}_0 \leq 40$ and $27 \leq \bar{\alpha}_{0,4} \leq 40$. By [7, Corollary 11], we have $\bar{\alpha}_1 \geq 22$ and $\bar{\alpha}_{1,4} \geq 26$. Wiener [19] showed that there exists a planar almost hypohamiltonian graph of order 31 and girth 4, which is the smallest known planar almost hypohamiltonian graph (of girth 4), so $\bar{\alpha}_1 \leq \bar{\alpha}_{1,4} \leq 31$. No planar almost hypohamiltonian graph of girth 3 is known [7].

Thomassen [15] showed that every planar hypohamiltonian graph contains a cubic vertex. C. T. Zamfirescu [24] proved that (i) every planar hypohamiltonian graph contains at least four cubic vertices, (ii) every planar almost hypohamiltonian graph contains a cubic vertex which is not the exceptional vertex (solving one of his problems in [21]), and (iii) every hypohamiltonian graph with crossing number 1 contains a cubic vertex. No planar hypohamiltonian graph with fewer than 30 cubic vertices was known [10, 24]. Thus, the minimum number of cubic vertices contained in planar hypohamiltonian graphs is between 4 and 30.

The smallest known cubic planar hypohamiltonian graphs have 70 vertices and girth 4. The first was found by Araya and Wiener [1], while six more were found by M. Jooyandeh and B. McKay [11, 17]. The cubic planar hypohamiltonian graphs must have girth at least 4 [4, 6]. McKay [12] showed that cubic planar hypohamiltonian graphs of girth 5 exist, and that the smallest ones are three graphs on 76 vertices. J. Goedgebeur and C. T. Zamfirescu [6, Theorem 3.5] proved that the smallest cubic planar hypohamiltonian graph has at least 54 and at most 70 vertices.

The paper is organized as follows. In [Section 2](#) we introduce the relations on some known planar hypohamiltonian graphs of order 40 and 42. These relations

allow us to construct the smaller ones. In [Section 3](#) we present planar hypohamiltonian graphs on 34, 37, 38, and 41 vertices. Hitherto it was unknown whether planar hypohamiltonian graphs of these orders exist. We exploit a special subgraph contained in two graphs of Jooyandeh et al. [\[10\]](#) to construct two planar hypohamiltonian graphs on 34 vertices. We further modify the 34-vertex graphs to obtain six planar hypohamiltonian graphs on 37 vertices, and rediscover Wiener’s planar almost hypohamiltonian graph on 31 vertices [\[19\]](#). No planar hypohamiltonian graph with fewer than 30 cubic vertices was known [\[24\]](#). The 34-vertex graphs have 26 cubic vertices, narrowing the gap between 4 and 30 [\[24\]](#). The smallest known hypohamiltonian graph with crossing number 1 is of order 36 due to Wiener [\[19\]](#). We reduce the best-known bound of 36 to 34 by the hypohamiltonian graphs with crossing number 1 obtained by adding an edge to the 34-vertex graphs. By applying an operation introduced by Thomassen [\[16\]](#) to the 37-vertex graphs, several planar hypohamiltonian graphs on 41 vertices are obtained, by which we settle the open question whether there exists a planar hypohamiltonian graph on 41 vertices [\[10\]](#). In [\[10\]](#), all the 25 planar hypohamiltonian graphs on 40 vertices have no nontrivial automorphisms. We present six planar hypohamiltonian graphs on 40 vertices with nontrivial automorphisms.

Our graphs make some progress on Holton’s question regarding the smallest natural number n_0 such that there exists a planar hypohamiltonian graph of order n for every $n \geq n_0$ [\[22, Problem 1.\(b\)\]](#). The best-known bound for n_0 was 42 [\[10, Theorem 4.3\]](#). By showing the existence of planar hypohamiltonian graphs on 41 vertices, we reduce n_0 to 40.

All of our graphs in this paper are constructed by hand. We use computer to check that they are hypohamiltonian and pairwise non-isomorphic. We did not exhaustively try all possible ways for constructing small planar hypohamiltonian graphs. It is possible that other new constructions yield even more graphs.

2 Equivalence relations and order relations

Planar hypohamiltonian graphs G_1, \dots, G_k are said to be *H-equivalent* if each of them can be obtained from a fixed graph H by adding vertices and edges such that for every $i, j = 1, \dots, k$, $|V(G_i) - V(H)| = |V(G_j) - V(H)|$ and $|E(G_i) - E(H)| = |E(G_j) - E(H)|$, denoted by $G_i =_H G_j$. Similarly, G is said to be *H-strictly bigger than* G' if $|V(G) - V(H)| > |V(G') - V(H)|$ and $|E(G) - E(H)| > |E(G') - E(H)|$, denoted by $G >_H G'$.

[Fig. 1](#), [Fig. 2](#), and [Fig. 3](#) are examples of such equivalence relations. These graphs can be found in the House of Graphs [\[5\]](#) by searching the corresponding House-of-Graphs ID (HoG ID), such as the Wiener-Araya Graph [\[20\]](#) with HoG ID 1431, which we will also call Graph 1431. In the figures, the number on the lower right of each graph is the corresponding HoG ID. In [Section 3](#) we will use these relations to construct two planar hypohamiltonian graphs on 34 vertices, as shown in [Fig. 6](#).




Figure 1: G_1 , G_2 , and G_3 are planar hypohamiltonian graphs of order 42. We have $G_1 =_{H_1} G_2 =_{H_1} G_3$.




Figure 2: G_1 , G_2 , G_3 and G_4 are planar hypohamiltonian graphs of order 42. We have $G_1 =_{H_2} G_2 =_{H_2} G_3 =_{H_2} G_4$.




Figure 3: G_5 and G_6 are planar hypohamiltonian graphs of order 40 [10]. We have $G_5 =_{H_3} G_6$.

3 Results

To construct planar hypohamiltonian graphs on fewer than 40 vertices, our approach is to modify H_3 to the graph in Fig. 4.




Figure 4: Graph 51085, denoted by H_4 .




Figure 5: An equivalent representation of H_4 . This will be used when constructing two planar hypohamiltonian graphs of order 40 with non-trivial automorphisms, Graph 51101 and 51102, as shown in Fig. 14.




Figure 6: Two planar hypohamiltonian graphs of order 34. We have $G_7 \cong_{H_4} G_8$.

C. T. Zamfirescu [22, Problem 1.(a)] asked, “Do planar hypohamiltonian graphs on less than 40 vertices exist?” By the existence of the graphs in Fig. 6, we settle the problem. No planar hypohamiltonian graph with fewer than 30 cubic vertices was known [10, 24]. We improve upon the result by the graphs in Fig. 6, which have 26 cubic vertices.

We tried to add vertices and edges to H_4 by hand in as many ways as possible. Not only can H_4 be used to construct G_7 and G_8 , but also two planar hypohamiltonian graphs of order 37 shown in Fig. 7.




Figure 7: Two planar hypohamiltonian graphs of order 37. We have $G_9 = G_{10} >_{H_4} G_7 =_{H_4} G_8$. The graphs have 28 cubic vertices.

We rediscover Wiener’s planar almost hypohamiltonian graph of order 31 shown in Fig. 8, which is the smallest known planar almost hypohamiltonian graph of girth 4. No planar almost hypohamiltonian graph of girth 3 is known [7].




Figure 8: Wiener’s planar almost hypohamiltonian graph of order 31. The exceptional vertex is the red vertex w . The HoG ID is 51072.

We tried by hand to modify H_4 , and found four more planar hypohamiltonian graphs on 37 vertices shown in Fig. 9 and Fig. 10.




Figure 9: (Left) Graph 51093. (Right) Graph 51094.




Figure 10: (Left) Graph 51095. (Right) Graph 51096.
These two graphs have no nontrivial automorphisms.

Theorem 3.1 (Grinberg's Theorem [18, Theorem 7.3.5]). *Let G be a simple plane graph. An i -face is a face of G which has size i . If G has a Hamiltonian cycle C and f_i (f'_i) is the number of i -faces of G inside (outside) of C , we have*

$$\sum_{i \geq 3} (i - 2)(f_i - f'_i) = 0.$$

Theorem 3.2. *The graphs shown in Fig. 6, Fig. 7, Fig. 9, and Fig. 10 are planar and hypohamiltonian.*

Proof. It is not hard to check that these graphs are planar. We follow the proof of [10, Theorem 4.1] to show that G_7 is non-Hamiltonian. For the other graphs, the proof is the same.

Assume to the contrary that G_7 contains a Hamiltonian cycle C , which must then satisfy Grinberg's Theorem. The graph G_7 contains five 4-faces and

18 5-faces. Then

$$\sum_{i \geq 3} (i-2)(f_i - f'_i) \equiv 2(f_4 - f'_4) \equiv 0 \pmod{3},$$

where $f_4 + f'_4 = 5$. So $f'_4 = 1$ and $f_4 = 4$, or $f'_4 = 4$ and $f_4 = 1$. Let Q be the 4-face on a different side from the other four.

Notice that an edge belongs to a Hamiltonian cycle if and only if the two faces it belongs to are on different sides of the cycle. Let the 4-face at the center of H_4 be q_c . Since q_c has edges in common with all other 4-faces and its edges cannot all be in a Hamiltonian cycle, q_c cannot be Q .

If Q is any of the other 4-faces, denote the edge belonging to Q and q_c by $e = \{a, b\}$ with $\deg(a) = 4$ and $\deg(b) = 3$. Denote the other two vertices of q_c by c and d with $\deg(c) = 4$ and $\deg(d) = 3$. Since the vertex a belongs to the edge e , the edge $\{d, a\}$ is not a part of C . The edge $\{d, c\}$ is not a part of C , either. It is because the two 4-faces that $\{d, c\}$ belongs to are on the same side of C . Thus, G_7 is non-Hamiltonian.

Finally, for each vertex it is routine to exhibit a cycle of length 33 (Fig. 6) or 36 (Fig. 7, Fig. 9, and Fig. 10) that avoids it. \square

The smallest known hypohamiltonian graph with crossing number 1 due to Wiener [19] has 36 vertices. We reduce the bound of 36 to 34 by the hypohamiltonian graphs with crossing number 1 obtained by adding an edge to the graphs in Fig. 6. We present six such graphs shown in Fig. 11 derived from G_7 in Fig. 6 by adding an edge (shown in red) to it.




Figure 11: Six hypohamiltonian graphs with crossing number 1 derived from G_7 in Fig. 6.

We use computer to check that the graphs in Fig. 11 are hypohamiltonian and pairwise non-isomorphic. It is not hard to check by hand that each red edge must cross over at least one black edge, so the crossing number is 1.

Whether there exists a planar hypohamiltonian graph on 41 vertices was an open question [10]. The Holton’s question is what is the smallest natural number n_0 such that there exists a planar hypohamiltonian graph of order n for every $n \geq n_0$ [22, Problem 1.(b)]. Motivated by Holton’s question, we now need an operation introduced by Thomassen [16] with which he showed that there exist infinitely many planar cubic hypohamiltonian graphs. Let G be a graph containing a 4-cycle $v_1v_2v_3v_4 = C$. We denote by $\text{Th}(G, C)$ the graph obtained from G by deleting the edges v_1v_2, v_3v_4 and adding a new 4-cycle $v'_1v'_2v'_3v'_4$ and the edges $v_iv'_i, 1 \leq i \leq 4$. Araya and Wiener [20] note that a result in [16] generalizes as follows, with the same proof.

Lemma 3.3 ([20, Lemma 4.4]). *Let G be a planar hypohamiltonian graph having a 4-cycle $v_1v_2v_3v_4 = C$ and suppose that the degrees of the vertices v_1, v_2, v_3, v_4 are 3. Then $\text{Th}(G, C)$ is also a planar hypohamiltonian graph.*

The assertion of Lemma 3.3 holds if one uses the condition that at least one of v_1, v_2, v_3, v_4 is cubic in G . It also holds if the edges v_1v_2 or v_3v_4 (possibly both) lie in G [24, Theorem 5].

Jooyandeh et al. [10, Theorem 4.3] used the operation Th to show that planar hypohamiltonian graphs of order n for every $n \geq 42$, which follows $n_0 \leq 42$. By applying the operation Th to the 37-vertex graphs in Fig. 7, Fig. 9, and Fig. 10, we obtain a number of planar hypohamiltonian graphs on 41 vertices, and thus settle the open question of Jooyandeh et al. [10]. Fig. 12 shows two such 41-vertex graphs obtained by applying the operation Th to the 4-cycle at the center of H_4 without deleting the edges v_1v_2 and v_3v_4 . The other graphs can be found in the House of Graphs [5] by searching for the text string “planar hypohamiltonian graph”.




Figure 12: Two planar hypohamiltonian graphs on 41 vertices, obtained by applying the operation Th to the graphs in Fig. 7.

By the graphs in Fig. 12, the result of Jooyandeh et al. [10, Theorem 4.3] is improved further in the next theorem.

Theorem 3.4. *There exist planar hypohamiltonian graphs of order n for every $n \geq 40$.*

By applying the operation Th to the graphs in Fig. 6, we have several planar hypohamiltonian graphs on 38 vertices, two of which are shown in Fig. 13. Whether there exists planar hypohamiltonian graphs on 35, 36, 39 vertices remains an open question.




Figure 13: Two planar hypohamiltonian graphs on 38 vertices, obtained by applying the operation Th to the graphs in Fig. 6.

The 25 planar hypohamiltonian graphs on 40 vertices in [10] have no nontrivial automorphisms, as mentioned in [10, Conclusions]. In this paper, we present six planar hypohamiltonian graphs on 40 vertices with nontrivial automorphisms shown in Fig. 14. Additionally, the planar hypohamiltonian graphs in Fig. 6, Fig. 7, Fig. 9, Fig. 12, and Fig. 13 have nontrivial automorphisms with the corresponding orders 34, 37, 41, and 38. In the Appendix, we also list some planar hypohamiltonian graphs on 43, 46, and 49 vertices with nontrivial automorphisms.

In 2011, Araya and Wiener [20, Corollary 3.1; Theorem 4.9] proved that there exist planar hypotractable graphs on $162 + 4k$ vertices for every $k \geq 0$, and on n vertices for every $n \geq 180$. In 2017, Jooyandeh et al. [10, Theorem 4.5] proved that there exist planar hypotractable graphs on 154 vertices, and on n vertices for every $n \geq 156$. In 2018, Wiener [19] constructed a planar hypotractable graph of order 138, improving the bound of 154 in [10]. To improve the best-known bound of 138, we make use of the following theorem, which is a slight modification of [13, Lemma 3.1].

Theorem 3.5 ([10, Theorem 4.4]). *Let $G_i = (V_i, E_i)$, $1 \leq i \leq 4$, be four planar hypohamiltonian graphs. Then there is a planar hypotractable graph of order $|V_1| + |V_2| + |V_3| + |V_4| - 6$.*

Combining Theorem 3.5 with Theorem 3.2 and Theorem 3.4, we obtain the following.

Theorem 3.6. *There exist planar hypotractable graphs on 130 vertices, and on n vertices for every $n \geq 154$.*




Figure 14: Six planar hypohamiltonian graphs on 40 vertices with non-trivial automorphisms. The two graphs on the rightmost contain H_4 .

Let H be a cubic graph and G be a graph containing a cubic vertex $w \in V(G)$. We say that we *insert* G into H , if we replace every vertex of H with $G - w$ and connect the endpoints of edges in H to the neighbours of w . We improve upon the previous results [6, 10, 21] as follows. Our arguments follow the same strategy as used in [10, Corollary 4.7] and [23, Theorem 4.1].

Corollary 3.7.

$$27 \leq h_4 \leq 34, \quad \bar{\alpha}_0 \leq 34, \quad \bar{C}_3^1 \leq 34, \quad \bar{C}_3^2 \leq 2205, \quad \bar{P}_3^1 \leq 132, \quad \bar{P}_3^2 \leq 8694.$$

Proof. The first inequality is obtained from [6, Theorem 3.3] and the graphs in Fig. 6, which improves upon the third inequality in [6, Corollary 3.4].

The second and third inequalities follow immediately from the graphs in Fig. 6. To obtain the last three inequalities, let G be one of the graphs in Fig. 6.

For the fourth inequality, insert G into the 70-vertex planar cubic hypohamiltonian graph—which we will call here G_0 —constructed by Araya and Wiener in [1]. (G_0 is the smallest known planar cubic hypohamiltonian graph.) This means that each vertex of G_0 is replaced by G minus some vertex of degree 3. We denote the resulting graph by G' . Araya and Wiener proved [1] (using a computer) that every pair of edges in G_0 is missed by a longest cycle. Combining this fact with the hypohamiltonicity of G and G_0 , we obtain that in G' any pair of vertices is avoided by a longest cycle. This property is not lost if all edges originally belonging to G_0 are contracted. By construction, the order of G' is $(34 - 1) \cdot 70 = 2310$. Since $|E(G_0)| = 105$, after contracting all edges originally belonging to G_0 , we obtain $\bar{C}_3^2 \leq 2310 - 105 = 2205$.

In order to prove the fifth inequality, insert G into K_4 . We obtain a graph in which every vertex is avoided by a path of maximal length. Thus, we have

$\overline{P}_3^1 \leq 4 \cdot (34 - 1) = 132$. The upper bound for \overline{P}_3^1 is not improved upon by [Theorem 3.6](#). The planar hypotractable graphs on 130 vertices are constructed by [\[13, Lemma 3.1; Fig. 2\]](#), which are 2-connected.

For the last inequality, consider the graph G_0 from the third paragraph of this proof and insert G_0 into K_4 to obtain H . Now insert G into H . Finally, contract all edges which originally belonged to H . \square

For the cubic planar case, the smallest known cubic planar hypohamiltonian graphs have 70 vertices and girth 4. The first one was found by Araya and Wiener [\[1\]](#), and the other six were found by Jooyandeh and McKay [\[11, 17\]](#). The cubic planar hypohamiltonian graphs must have girth at least 4 [\[4, 6\]](#). McKay [\[12\]](#) showed that cubic planar hypohamiltonian graphs of girth 5 exist, and that the smallest ones are three graphs on 76 vertices. Therefore the smallest cubic planar hypohamiltonian graph must have girth exactly 4. J. Goedgebeur and C. T. Zamfirescu [\[6, Theorem 3.5\]](#) proved that the smallest cubic planar hypohamiltonian graph has at least 54 and at most 70 vertices.

4 Conclusions

Based on the work of [\[10\]](#), the two planar hypohamiltonian graphs of order 34 with nontrivial automorphisms in this paper significantly decrease the gap between the lower and upper bounds for the order of the smallest planar hypohamiltonian graph. By the further study of the 34-vertex graphs, we obtain six planar hypohamiltonian graphs of order 37, four of which have nontrivial automorphisms. By applying the operation introduced by Thomassen [\[16\]](#) to the 34- and 37-vertex graphs, we obtain several planar hypohamiltonian graphs of order 38 and 41, some of which have nontrivial automorphisms.

By showing the existence of planar hypohamiltonian graphs on 41 vertices, we settle the open question in [\[10\]](#) whether there exist planar hypohamiltonian graphs on 41 vertices. Our 41-vertex graphs make some progress on Holton's question regarding the smallest natural number n_0 such that there exists a planar hypohamiltonian graph of order n for every $n \geq n_0$ [\[22, Problem 1.\(b\)\]](#). Our graphs gives $n_0 \leq 40$, improving upon the result of [\[10\]](#).

Quote the authors of [\[10\]](#), "An exhaustive study of graphs with prescribed automorphisms might lead to the discovery of new, smaller graphs." Their point of view was precise. Throughout the graphs in [Fig. 6](#), [Fig. 7](#), and [Fig. 9](#), each of them was constructed with some prescribed symmetries. Finally, new and smaller graphs were found. The further study of these graphs leads to six planar hypohamiltonian graphs of order 40 with nontrivial automorphisms.

Whether there exists planar hypohamiltonian graphs on 35, 36, and 39 vertices remains an open question.

Our graphs are based on two graphs of [\[10\]](#), Graph 17030 and 17052, shown in [Fig. 3](#). In future work one could try the other 23 graphs of [\[10\]](#) to see whether there are new planar hypohamiltonian graphs on fewer than 40 vertices. Additionally, all the graphs in this paper were constructed by hand by adding

vertices and edges to the graph in Fig. 4, or modifying it. We did not try all possible for constructing small planar hypohamiltonian graphs. An exhaustive study of our construction might lead to the discovery of new, smaller graphs.

We can hope that the current work inspires further progress in finding the smallest (cubic) planar hypohamiltonian or smallest planar almost hypohamiltonian graphs.

Acknowledgements

We thank Jan Goedgebeur and Carol T. Zamfirescu for their helpful comments. We thank Brendan McKay and Eric W. Weisstein for their comments on cubic planar hypohamiltonian graphs, and thank Eric W. Weisstein for the comment on Figure 11.

Appendix




Figure 15: Planar hypohamiltonian graphs on 43, 46, and 49 vertices. Note that Graph 51107 contains H_4 .

References

- [1] M. Araya and G. Wiener. On cubic planar hypohamiltonian and hypo-traceable graphs. *Electronic Journal of Combinatorics*, 18(1), 2011.
- [2] V. Chvátal. Flip-flops in hypohamiltonian graphs. *Canadian Mathematical Bulletin*, 16(1):33–41, 1973.

- [3] V. Chvátal, D. A. Klarner, and D. E. Knuth. Selected combinatorial research problems. *STAN-CS-72-292, Stanford University, CA, USA*, 1972.
- [4] J. B. Collier and E. F. Schmeichel. Systematic searches for hypohamiltonian graphs. *Networks*, 8(3):193–200, 1978.
- [5] K. Coolsaet, S. D’hondt, and J. Goedgebeur. House of graphs 2.0: A database of interesting graphs and more. *Discrete Applied Mathematics*, 325:97–107, 2023. URL <https://houseofgraphs.org>.
- [6] J. Goedgebeur and C. T. Zamfirescu. Improved bounds for hypohamiltonian graphs. *Ars Mathematica Contemporanea*, 13(2):235–257, 2017.
- [7] J. Goedgebeur and C. T. Zamfirescu. On almost hypohamiltonian graphs. *Discrete Mathematics & Theoretical Computer Science*, 21(4), 2019.
- [8] B. Grünbaum. Vertices missed by longest paths or circuits. *Journal of Combinatorial Theory, Series A*, 17(1):31–38, 1974.
- [9] W. Hatzel. Ein planarer hypohamiltonscher graph mit 57 knoten. *Mathematische Annalen*, 243:213–216, 1979.
- [10] M. Jooyandeh, B. D. McKay, P. R. J. Östergård, V. H. Pettersson, and C. T. Zamfirescu. Planar hypohamiltonian graphs on 40 vertices. *Journal of Graph Theory*, 84(2):121–133, 2017.
- [11] B. McKay. Plane graphs, n.d. URL <https://users.cecs.anu.edu.au/~bdm/data/planegraphs.html>.
- [12] B. D. McKay. Hypohamiltonian planar cubic graphs with girth 5. *Journal of Graph Theory*, 85(1):7–11, 2016.
- [13] C. Thomassen. Hypohamiltonian and hypotraceable graphs. *Discrete Mathematics*, 9(1):91–96, 1974.
- [14] C. Thomassen. Planar and infinite hypohamiltonian and hypotraceable graphs. *Discrete Mathematics*, 14(4):377–389, 1976.
- [15] C. Thomassen. Hypohamiltonian graphs and digraphs. In Y. Alavi and D. R. Lick, editors, *Theory and Applications of Graphs*, pages 557–571, Berlin, Heidelberg, 1978. Springer Berlin Heidelberg. ISBN 978-3-540-35912-8.
- [16] C. Thomassen. Planar cubic hypohamiltonian and hypotraceable graphs. *Journal of Combinatorial Theory, Series B*, 30(1):36–44, 1981.
- [17] E. W. Weisstein. ”Cubic Planar Hypohamiltonian Graph.” From MathWorld—A Wolfram Web Resource., n.d. URL <https://mathworld.wolfram.com/CubicPlanarHypohamiltonianGraph.html>.
- [18] D. B. West. *Introduction to Graph Theory, 2nd ed.* Prentice Hall, 2001.

- [19] G. Wiener. New constructions of hypohamiltonian and hypotraceable graphs. *Journal of Graph Theory*, 87(4):526–535, 2018.
- [20] G. Wiener and M. Araya. On planar hypohamiltonian graphs. *Journal of Graph Theory*, 67(1):55–68, 2011.
- [21] C. T. Zamfirescu. On hypohamiltonian and almost hypohamiltonian graphs. *Journal of Graph Theory*, 79(1):63–81, 2015.
- [22] C. T. Zamfirescu. Seven problems on hypohamiltonian and almost hypohamiltonian graphs. In K. Adiprasito, I. Bárány, and C. Vîlcu, editors, *Convexity and Discrete Geometry Including Graph Theory*, pages 253–255, Cham, 2016. Springer International Publishing. ISBN 978-3-319-28186-5.
- [23] C. T. Zamfirescu. *Hypohamiltonian and almost hypohamiltonian graphs*. PhD thesis, Ghent University, 2016.
- [24] C. T. Zamfirescu. Cubic vertices in planar hypohamiltonian graphs. *Journal of Graph Theory*, 90(2):189–207, 2019.
- [25] C. T. Zamfirescu and T. Zamfirescu. A planar hypohamiltonian graph with 48 vertices. *Journal of Graph Theory*, 55(4):338–342, 2007.
- [26] T. Zamfirescu. A two-connected planar graph without concurrent longest paths. *Journal of Combinatorial Theory, Series B*, 13(2):116–121, 1972.