

First Experiences with the Identification of People at Risk for Diabetes in Argentina using Machine Learning Techniques

Enzo Rucci^{1,2}[0000-0001-6736-7358] ✉, Gonzalo Tittarelli³, Franco Ronchetti^{1,2}[0000-0003-3173-1327] Jorge F. Elgart⁴[0000-0002-6101-1219], Laura Lanzarini¹[0000-0001-7027-7564], and Juan José Gagliardino⁴[0000-0001-7332-6536]

¹ III-LIDI, Facultad de Informática, UNLP – CIC. La Plata (1900), Argentina
 {erucci,fronchetti,laural}@lidi.info.unlp.edu.ar

² Comisión de Investigaciones Científicas (CIC). La Plata (1900), Argentina

³ Facultad de Informática, UNLP. La Plata (1900), Bs As, Argentina

⁴ CENEXA, Facultad de C. Médicas, UNLP-CONICET. La Plata (1900), Argentina
 {jelgart,jjgagliardino}@cenexa.org

⁵ CONICET. La Plata (1900), Argentina

Abstract. Detecting Type 2 Diabetes (T2D) and Prediabetes (PD) is a real challenge for medicine due to the absence of pathogenic symptoms and the lack of known associated risk factors. Even though some proposals for machine learning models enable the identification of people at risk, the nature of the condition makes it so that a model suitable for one population may not necessarily be suitable for another. In this article, the development and assessment of predictive models to identify people at risk for T2D and PD specifically in Argentina are discussed. First, the database was thoroughly preprocessed and three specific datasets were generated considering a compromise between the number of records and the amount of available variables. After applying 5 different classification models, the results obtained show that a very good performance was observed for two datasets with some of these models. In particular, RF, DT, and ANN demonstrated great classification power, with good values for the metrics under consideration. Given the lack of this type of tool in Argentina, this work represents the first step towards the development of more sophisticated models.

Keywords: public health · chronic disease · machine learning

This version of the contribution has been accepted for publication, after peer review (when applicable) but is not the

Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at:

https://doi.org/10.1007/978-3-031-62245-8_16. Use of this Accepted Version is subject to the publisher's Accepted Manuscript terms of use <https://www.springernature.com/gp/open-research/policies/accepted-manuscript-terms>

1 Introduction

Type 2 Diabetes (T2D) is a chronic disease characterized by high blood glucose levels; it manifests when the endocrine pancreas is unable to produce enough insulin for body tissues [11]. Due to its increasing prevalence in combination with its high cost of care [4,30], it represents a serious public health problem, and great efforts have been made to develop effective strategies for its prevention and timely treatment, as well as to avoid the development of its chronic complications.

It is worth noting that the negative consequences of this disease start in a previous stage known as prediabetes (PD), which is characterized by elevated blood glucose concentration higher than normal levels, but without reaching the diagnostic values for diabetes. PD involves Impaired Fasting Glucose (IFG), Impaired Glucose Tolerance (IGT), and a combination of both [1]. PD has a high risk of progressing into T2D — around 30% [7] and 70% [9] in the following 4 and 30 years, respectively.

The development of T2D is a slow and progressive process conditioned by genetic, environmental, and behavioral factors. Currently, there is no definitive cure for this disease. However, several studies have shown that its onset can be prevented or delayed in people with PD through the adoption of a healthy lifestyle (eating plan and regular physical activity) and/or associated with the intake of various drugs [7,27]. In Argentina, an initiative of this type is the Pilot Program for the Primary Prevention of Diabetes in the province of Buenos Aires (PPDBA), developed by CENEXA (UNLP-CONICET) and financed by the Ministry of Science and Technology of the Nation, the company SANOFI and CONICET [12].

Detecting T2D and PD is a real challenge for medicine due to the absence of pathogenic symptoms and the lack of known associated risk factors. This is why individuals may often go months (or even years) without knowing that they are at risk. In this sense, statistics published in 2018 by the International Diabetes Federation show that approximately 50% of the world's population is unaware of their disease [10]. This explains the need for a simple and accurate detection method. Thus, this article proposes the development and assessment of predictive models based on machine learning (ML) that will allow identifying people at risk of T2D and PD in Argentina. To this end, the PPDBA database will be used.

This paper is an extended and thoroughly revised version of [23]. The work has been extended by providing: (1) the proposal of a new segmentation that only considers clinical data from the PPDBA database and the training of several models that use it as input; and (2) a performance analysis of the new models and a cost-benefit analysis with the previous ones.

The remaining sections of this article are organized as follows: In Section 2, the background for this work is presented. In Section 3, database processing is described, while in Section 4 the results obtained are analyzed. Finally, in Section 5, conclusions and possible future work are presented.

2 Background

2.1 Diabetes and Prediabetes — Risk Factors and Diagnosis

The development of T2D is a slow and progressive process that is conditioned by genetic, environmental, and behavioral factors. Risk factors include gender, body mass index (BMI), waist circumference, eating habits, physical activity, age, family history of diabetes (including gestational), ethnicity, sleep disorders, and so forth. PD corresponds to a state before T2D, and its progression can be prevented and even reversed by adopting healthy lifestyles [28]. If we also consider that PD is not a pre-disease state, since it already presents metabolic dysfunctions, it is extremely important to identify people with PD as much as it is to do so with those who already have undiagnosed T2D.

Diagnosis is based on blood tests, and anyone presenting the associated symptoms or risk factors should be examined. In Argentina, the Oral Glucose Tolerance Test (OGTT) is usually used to determine if a person has T2D, PD, or neither ⁶.

2.2 Related Works

In the last decade, numerous models have been proposed to identify undiagnosed diabetes and/or PD using ML techniques. These are based on various clinical and laboratory variables linked to relevant risk factors. Most of them [15] [2] [13] [16] [19] [21] [16] [34] [6] [29] [22] [25] [18] [35] [31] [26] use a diabetes database known as PIMA Indian Diabetes (PID) from the repository of the University of California Irvine ⁷, USA. This dataset contains records of women from the Pima indigenous people (USA) and is made up of 8 attributes related to risk factors for developing the disease. It has a total of 786 records; however, this number is reduced by half if entries with null values are removed. These works are more of a proof of concept than a real implementation and, in general, have been aimed at improving the performance (accuracy) of diabetes classification algorithms.

Only a few of these works do not use the PID database. These include [14] [33] [8], whose authors proposed models to identify diabetes and prediabetes using data from the US NHANES Survey. Recently, the authors in [32] also studied the application of different ML techniques for detecting undiagnosed diabetes in the US population using the BRFSS database. Similarly, the authors in [20] presented various predictive models for these diseases for the Chinese population using an ad-hoc database. In [5], on the other hand, the development and validation of predictive models focused only on prediabetes.

As explained in Section 2.1, the onset of T2D is conditioned by factors that may vary from one population to another. This is why a predictive model that

⁶ For this test, a blood sample is drawn on a fasting state. Then, the subject is asked to drink a liquid containing a certain amount of glucose. After this, another blood sample is taken every 30 to 60 minutes after intake. The exam is financially expensive and can take up to 3 hours.

⁷ <https://archive.ics.uci.edu/ml/index.php>

is suitable for one population will not necessarily be suitable for another. This study is the first step toward building specific predictive models for the Argentine population.

3 Implementation

3.1 Dataset

Description Predictive models will be developed using the PPDBA program database [12], which has 1316 records. Each record corresponds to a person identified as diabetic, prediabetic or disease-free through OGTT. In addition to laboratory data (glycated hemoglobin, total cholesterol, HDL cholesterol, LDL cholesterol, triglycerides, and creatinine), there are clinical variables associated with the risk factors for these diseases such as sex, age, Body Mass Index (BMI), blood pressure, family history of diabetes, eating habits, physical activity, and so forth.

Characterization Out of the 1316 current records, 80 had to be discarded due to the lack of blood glucose data to calculate the OGTT (`class` could not be determined). Table 1 presents a brief statistical description of the 1236 remaining records. There are several null variables, which are analyzed in greater depth in the following section. As it can be noted, there are more women than men, most subjects are between 45-64 years old; most with a BMI greater than 30 kg/m²; most with a waist circumference greater than 102 cm (men) and 88 cm (women); most do physical activity; most eat vegetables and fruits; most do not take medications to control hypertension; most with a positive result for hyperglycemia at least once; most with at least one relative (first or second degree) with diabetes. Variables associated with baseline glycemia, HDL cholesterol, triglycerides, and baseline creatinine seem to have a wide dispersion; and as regards class, half of the subjects are not at risk for PD or T2D.

Cleanup

Noise To analyze the presence of noise in the variables, Tukey’s method was used to identify the ranges for outliers. They were detected in the following variables: `age` (4), `bmi` (10), `waist_circumference` (1), `baseline_glycemia` (27), `postprandial_glycemia` (46), `ldl_cholesterol` (7), `total_cholesterol` (9), `hdl_cholesterol` (11), `triglycerides` (21), and `baseline_creatinine` (8). Additionally, *far-out* outliers were found in `baseline_glycemia` (51), `postprandial_glycemia` (6), `triglycerides` (12), and `baseline_creatinine` (6).

Missing values Table 1 shows that there are no missing values in qualitative variables, except for `family_diagnosis`, with 3 records with null values. In the case of quantitative variables, the number of missing values is much greater.

Table 1. Statistical overview of the dataset

| Variable | # null | Metric | Result |
|---------------------------|--------|---------------------------------|----------------|
| sex | 0 | Male (%) | 395 (32%) |
| | | Female (%) | 841 (68%) |
| age | 564 | Mean+SD | 57.23±8.8428 |
| age_range | 0 | Less than 45 years (%) | 205 (17%) |
| | | 45-54 years (%) | 435 (35%) |
| | | 54-64 years (%) | 429 (34%) |
| | | More than 64 years (%) | 167 (14%) |
| bmi | 619 | Mean+SD | 31.65±6.314 |
| bmi_range | 0 | Less than 25kg/m2 | 93 (8%) |
| | | 25-30 kg/m2 | 319 (26%) |
| | | More than 30 kg/m2 | 824 (66%) |
| waist_circumference | 1181 | Mean+SD | 101.3091±13.55 |
| waist_circumference_range | 0 | Less than 94/80 cm (M/F) (%) | 56 (4%) |
| | | M: 94-102cm / F: 80-88cm (%) | 205 (17%) |
| | | M: More than 102/88cm (M/F) (%) | 975 (79%) |
| | | | |
| physical_activity | 0 | Yes (%) | 915 (74%) |
| | | No (%) | 321 (26%) |
| eat_fruit_vegetables | 0 | Yes (%) | 821 (66%) |
| | | No (%) | 415 (34%) |
| ht_drugs | 0 | Yes (%) | 497 (40%) |
| | | No (%) | 739 (60%) |
| hyperglycemia | 0 | Yes (%) | 999 (81%) |
| | | No (%) | 237 (19%) |
| family_diagnosis | 3 | No (%) | 395 (32%) |
| | | First degree (%) | 412 (33%) |
| | | Second degree (%) | 426 (34%) |
| baseline_glycemia | 0 | Mean+SD | 104.36±27.28 |
| postprandial_glycemia | 55 | Mean+SD | 119.59±42.51 |
| total_cholesterol | 705 | Mean+SD | 198.28±41.1 |
| ldl_cholesterol | 715 | Mean+SD | 119.79±36.82 |
| hdl_cholesterol | 706 | Mean+SD | 49.82±14.40 |
| triglycerides | 705 | Mean+SD | 151.4±95.59 |
| baseline_creatinine | 619 | Mean+SD | 1.117±5.8 |
| glycated_hemoglobine | 635 | Mean+SD | 5.61±0.43 |
| ogtt_result | 0 | No risk (%) | 620 (50%) |
| | | Prediabetes (%) | 480 (38%) |
| | | Diabetes (%) | 136 (12%) |

Fortunately, some of these quantitative variables have an associated qualitative variable that limits the range for that missing value. This occurs specifically in the cases of `age`, `bmi`, and `waist_circumference`. The other variables that have null values are: `postprandial_glycemia`, `total_cholesterol`,

`ldl_cholesterol`, `hdl_cholesterol`, `triglycerides`, `baseline_creatinine`, and `glycosylated_hemoglobin`.

Transformations

Treatment of outliers Medical experts were consulted about the occurrence of outliers in lab tests. It was concluded that, even if these values in the database are statistically atypical, they are within the possible range for their respective tests. There is only one exception — 2 values in `baseline_creatinine` (85 and 118) that are quite likely upload errors. Therefore, those two specific values were replaced with null values.

Treatment of missing values This task has two options: (1) removing variables with missing values; or (2) deleting records with null variables. The first option allows us to keep sample size at the cost of reducing the number of input variables for the models. The second option allows us to keep the number of features at the cost of reducing sample size. It was decided to select Option 2 only for the `family_diagnosis` variable, which has only 3 null values. Option 1 was applied to the variables `age`, `bmi` and `waist_circumference`, considering that there is an associated qualitative variable for each one of them that allows us to know its approximate value. All other variables will be discussed in Section 3.2.

Aggregation of class variable Table 1 shows that the distribution of classes is not balanced (`ogtt_result`). To minimize the impact of this issue, a new class variable is created that splits subjects at risk of having PD or T2D (records with a value of “PD” or “T2D”) and those who are risk-free (records with a value of “Normal”). As a result, the variable is balanced in terms of the number of occurrences of each value, simplifying subsequent analyses as it now becomes a binary classification problem. However, caution should be used when analyzing the results, especially concerning the prediction of T2D, as it is the least common.

Variable removal `ogtt_result` was discarded in the previous section, and `postprandial_glycemia` was excluded because any subject with that variable would require an OGTT and, in that case, using the proposed models would be meaningless.

Correlation Analyses Fig. 1 shows the correlation matrix obtained for the initial dataset. Clinically, it makes sense that there is a weak linear correlation between waist circumference range and BMI range. The weak correlation between age, BMI, and waist circumference ranges and their associated values for `age`, `bmi`, and `waist_circumference` is also expected. On the other hand, `total_cholesterol` is calculated from `ldl_cholesterol` [3], which explains the strong correlation between these two variables. Additionally, the relationship between `baseline_glycemia`, `postprandial_glycemia`, `ogtt_result` and, `class` is also to be expected, since the first two determine the value of the third, which in turn is grouped to generate the fourth.




Fig. 1. Correlation matrix for the initial dataset

3.2 Proposed Segmentations

Given the high percentage of missing values in lab variables, the possibility of generating several datasets from the original by applying different criteria for the treatment of null records was considered:

- **Clinical and Lab Dataset (CLD-bin)**. All complete records were eliminated, resulting in 16 variables with 503 examples (229 ⇒ No risk, 274 ⇒ At risk). This dataset keeps all available variables (clinical data and lab data) at the cost of losing many records.
- **Clinical and baseline Glycemia Dataset (CGD-bin)**. This dataset keeps all available clinical information and the only lab variable that does not have null values (`baseline_glycemia`); the rest of the variables were removed. Thus, this dataset has 10 variables and 1233 examples. In contrast to CLD, CGD preserves the number of records as opposed to lab variables value.
- **Clinical Dataset (CD-bin)**. This dataset keeps all available clinical information and removes all lab variables. Thus, this dataset includes 9 variables

and 1233 examples, preserving the number of records. By discarding all lab features, simpler and cost-free models can be trained.

4 Experimental Results

4.1 Experimental Design

An exploration, preprocessing, training, and evaluation process of various algorithms was carried out using *scikit-learn*. In particular, Logistic Regression (LR), Decision Tree (DT) with `max_depth = 5` and `class_weight = balanced`, k-Nearest Neighbor (kNN) with `n_neighbors = 7`, and Random Forest (RF) with `max_depth = 2` were considered. Likewise, TensorFlow was used to build Artificial Neural Networks (ANN) models with `epochs = 60` and `batch_size = 16`. Specifically, a single hidden layer ANN was built, with 100 neurons and the ReLU activation function, and L2 regularization = 0.1 to control overfitting. An Adamax optimizer was used with a learning rate of 0.001.

Following common guidelines in the area, the dataset was partitioned into two: training (70%) and evaluation (30%). To reduce possible biases, the cross-validation technique with stratified random sampling (`StratifiedShuffleSplit`, `n_splits=50` [24]) was applied. Since input variables are expressed in different magnitudes, a *min-max* normalization was applied for all models except for the ANN, where a standard normalization was used.

All experiments were carried out on a local computer using *Jupyter Notebook* [17]. The hardware used was an Intel Core i7 2.6 GHz Quad-Core CPU, 16 GB of RAM, and macOS operating system.

4.2 Models and Results for CLD-bin

Table 2 shows the performance metrics for the different models applied to the CLD-bin dataset, considering “At risk” as a positive class. As it can be seen, 3 of the 5 models yield accuracy values greater than 90%, which means that approximately 9 out of 10 of the total records evaluated were classified correctly (deviations are less than 2%). Of these 3 options, RF is the one that yields the best accuracy, followed by DT, and finally ANN. While the performance of LR is close to the previous ones with 86% accuracy, that of kNN is poor, reaching only 72%.

As regards precision, the values are consistent with those of accuracy. RF, DT and ANN yield values above 90% for the class of interest (“At risk”), which means that, with any of the options, more than 9 in 10 people are correctly told they are at risk. Similarly, these 3 models have recall values close to 91%, which means that only 1 in 10 people who are at risk is not identified.

Finally, when selecting a specific model, AUC values can be of interest, in addition to accuracy. In this case, it can be seen that RL displaces ANN as one of the 3 models with the highest AUC values, compared to those with the best accuracy.

Table 2. Results (evaluation) for the models applied to CLD-bin (positive class = “At risk”)

| Model | Accuracy | Precision | Recall | F-score | AUC |
|-------|--------------|--------------|--------------|--------------|------|
| RF | 94.58 ± 1.51 | 98.87 ± 1.3 | 91.1 ± 2.93 | 94.79 ± 1.53 | 0.95 |
| DT | 93.42 ± 1.59 | 96.02 ± 2.02 | 91.73 ± 2.65 | 93.79 ± 1.53 | 0.94 |
| ANN | 91.13 ± 1.98 | 92.29 ± 2.68 | 91.39 ± 2.77 | 91.79 ± 1.84 | 0.91 |
| LR | 85.56 ± 3.21 | 90.5 ± 3.11 | 82.1 ± 4.81 | 86.01 ± 3.3 | 0.93 |
| kNN | 71.62 ± 3.82 | 75.53 ± 3.87 | 70.83 ± 5.75 | 72.97 ± 4 | 0.72 |

Table 3. Results (evaluation) for the models applied to CGD-bin (positive class = “At risk”)

| Model | Accuracy | Precision | Recall | F-score | AUC |
|-------|--------------|--------------|--------------|--------------|------|
| RF | 93.23 ± 1.12 | 100 ± 0 | 86.38 ± 2.25 | 92.68 ± 1.3 | 0.93 |
| DT | 91.88 ± 1.67 | 96.68 ± 2.74 | 86.72 ± 2.2 | 91.39 ± 1.73 | 0.92 |
| ANN | 91.04 ± 1.63 | 94.45 ± 2.47 | 87.17 ± 2.31 | 90.64 ± 1.71 | 0.91 |
| LR | 75.66 ± 4.37 | 80.96 ± 4.44 | 66.7 ± 6.55 | 73.05 ± 5.3 | 0.83 |
| kNN | 69.97 ± 2.01 | 70.63 ± 2.32 | 67.92 ± 3.29 | 69.2 ± 2.23 | 0.7 |

4.3 Models and Results for CGD-bin

Table 3 shows the performance metrics for the different models applied to the CGD-bin dataset, considering “At risk” as a positive class. As with CLD-bin, RF, DT, and ANN yielded very good accuracy values, greater than 90% (deviations below 2%). kNN once again has a poor performance with 70% accuracy, while LR is between the previous 3 and kNN, although its performance is lower than with the previous dataset.

In terms of precision, RF overfits and obtains 100% for the “At Risk” class. DT and ANN yield values above 90% for the same class, with LR and kNN being well below this value. When analyzing recall values, RF, DT and, ANN are the models that achieve the best values, being close to 90% for the class of interest.

Finally, and unlike CLD-bin, the 3 models that obtain the highest AUC values are the same as the ones that achieve the best accuracy.

4.4 Models and Results for CD-bin

Table 4 shows the performance metrics for the different models applied to the CD-bin dataset, considering “At risk” as a positive class. Unlike the previous cases, the models present poor performances, reaching accuracy values from 54% to 57%. Moreover, the values presented for precision, recall, and F-score are along the same lines or worse. This means that the prediction is not much better than just flipping a coin.

Table 4. Results (evaluation) for the models applied to CD-bin (positive class = “At risk”)

| Model | Accuracy | Precision | Recall | F-score |
|-------|--------------|--------------|--------------|--------------|
| RF | 57.33 ± 1.93 | 63.77 ± 4.13 | 33.42 ± 4.84 | 43.61 ± 4.06 |
| DT | 54.98 ± 2.47 | 55.3 ± 3.07 | 51.16 ± 9.03 | 52.67 ± 5.01 |
| ANN | 57.55 ± 2.07 | 58.61 ± 3.06 | 51.05 ± 6.43 | 54.27 ± 3.71 |
| LR | 57.16 ± 2.32 | 60.88 ± 4.27 | 39.92 ± 5.25 | 47.94 ± 3.81 |
| kNN | 54.55 ± 2.45 | 54.45 ± 2.53 | 53.32 ± 4.79 | 53.77 ± 3.03 |

4.5 Discussion

Some of the models developed were able to achieve very good performances for both datasets. In particular, RF, DT, and ANN demonstrated great classification power, with high values in the considered metrics. In this sense, it should be noted that the proposed models are not intended to replace OGTT for the diagnosis of T2D and PD. As early detection is difficult for these diseases, the models are aimed at identifying individuals in the Argentine population who have a high probability of being affected and are unaware of their condition. To confirm the diagnosis, individuals identified as positive will eventually have to take an OGTT. The models would help identify those who should take the test and would make up for the lack of this type of tool.

It can be observed that there are no significant differences between the best accuracy and F-score values achieved for CLD-bin and CGD-bin datasets. Even though it is not (entirely) correct to compare results from models trained with different datasets, this issue could have an impact on the cost of putting the models into practice, considering that obtaining lab variables is neither free nor easy. To throw some light on this, a larger number of records without null values would be required.

Concerning the above, removing all lab features would lead to a simpler, cost-free model, that could be carried out at any time. Unfortunately, the performance results show that it is not feasible with the current CD-bin dataset. In that sense, it can be appreciated the great influence of the variable `baseline_glycemia` on the success of the prediction, probably due to the limited size of the dataset. Again, more records could compensate for the absence of lab variables.

Finally, it should also be taken into account that grouping PD and T2D into a single class favored balancing and simplified the problem by turning it into a binary classification. The cost of this decision is not being able to differentiate between PD and T2D. However, from a medical point of view, this would not be so relevant since, at the end of the day, the goal is to identify individuals at risk, regardless of which of the two conditions affects them.

5 Conclusions and Future Work

Considering that T2D and PD are difficult to detect, specific predictive models were developed and evaluated for the Argentine population based on the PPDBA database. Firstly, the database was carefully preprocessed, which led to the generation of three datasets (CLD-bin, CGD-bin, and CD-bin) with different approaches about the compromise between the number of variables and available records. Then, 5 different classification models were applied to each of them. The results obtained show that some of the models proposed were able to achieve very good performances for the first two datasets. In particular, RF, DT and ANN demonstrated great classification power, with high values for the metrics under consideration. In the opposite sense, removing all lab features led to poor performance results. Due to database limitations, the results are not conclusive, but they are promising. Given the lack of this type of tool in Argentina, this work represents the first step towards more sophisticated models.

Some of the future lines of work are:

- Obtaining more database records to improve its quality and representativeness, and then replicating the study carried out.
- Considering the usage of additional supervised learning methods (like Support Vector Machines) and the hyper-parameter tuning of all trained models.
- Analyzing the development of multiclass classification models to separate T2D and PD cases.

Funding. This study was partially supported by PICT-2020-SERIEA-00901.

References

1. Professional practice committee: Standards of medical care in diabetes—2021. *Diabetes Care* **44**(Supplement 1), S3–S3 (2021). <https://doi.org/10.2337/dc21-Sppc>, https://care.diabetesjournals.org/content/44/Supplement_1/S3
2. Al Jarullah, A.A.: Decision tree discovery for the diagnosis of type ii diabetes. In: 2011 International Conference on Innovations in Information Technology. pp. 303–307 (April 2011). <https://doi.org/10.1109/INNOVATIONS.2011.5893838>
3. Association, A.H.: What Your Cholesterol Levels Mean. <https://www.heart.org/en/health-topics/cholesterol/about-cholesterol/what-your-cholesterol-levels-mean> (2020), accedido: 2022-10-10
4. Bolin, K., Gip, C., Mörk, A.C., Lindgren, B.: Diabetes, healthcare cost and loss of productivity in sweden 1987 and 2005 - a register-based approach. *Diabetic medicine : a journal of the British Diabetic Association* **26**, 928–34 (10 2009). <https://doi.org/10.1111/j.1464-5491.2009.02786.x>
5. Choi, S.B., Kim, W.J., Yoo, T.K., Park, J.S., Chung, J.W., Lee, Y.h., Kang, E.S., Kim, D.W.: Screening for prediabetes using machine learning models. *Computational and mathematical methods in medicine* **2014** (2014)

6. Dey, S.K., Hossain, A., Rahman, M.M.: Implementation of a web application to predict diabetes disease: An approach using machine learning algorithm. In: 2018 21st International Conference of Computer and Information Technology (ICIT). pp. 1–5 (2018). <https://doi.org/10.1109/ICCITECHN.2018.8631968>
7. Diabetes Prevention Program Research Group: Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *New England Journal of Medicine* **346**(6), 393–403 (2002). <https://doi.org/10.1056/NEJMoa012512>
8. Dinh, A., Miertschin, S., Young, A., Mohanty, S.D.: A data-driven approach to predicting diabetes and cardiovascular disease with machine learning. *BMC Medical Informatics and Decision Making* **19**(1), 211 (Nov 2019). <https://doi.org/10.1186/s12911-019-0918-5>
9. Eddy, D.M., Schlessinger, L., Kahn, R.: Clinical Outcomes and Cost-Effectiveness of Strategies for Managing People at High Risk for Diabetes. *Annals of Internal Medicine* **143**(4), 251–264 (08 2005). <https://doi.org/10.7326/0003-4819-143-4-200508160-00006>
10. Federación Internacional de Diabetes: Atlas de la diabetes de la fid. Tech. rep., Federación Internacional de Diabetes (2017), <https://diabetesatlas.org/resources/2017-atlas.html>
11. Gagliardino, J.J., Assad, D., Gagliardino, G.G., Kronsbein, P., Lahera, E., Mercuri, N., Rizzuti, L., Zufriategui, Z.: *Cómo tratar mi diabetes*. Buenos Aires, Argentina, 3 edn. (11 2016)
12. Gagliardino, J.J., Etchegoyen, G., Bourgeois, M., Fantuzzi, G., García, S., González, L., Elgart, J.F., Ré, M., Ricart, A., Ricart, J.P., Spinedi, E.: Prevención primaria de diabetes tipo 2 en argentina: estudio piloto en la provincia de buenos aires. *Revista Argentina de Endocrinología y Metabolismo* **53**(4), 135 – 141 (2016). <https://doi.org/https://doi.org/10.1016/j.raem.2016.11.002>
13. Hashi, E.K., Zaman, M.S.U., Hasan, M.R.: An expert clinical decision support system to predict disease using classification techniques. In: 2017 International Conference on Electrical, Computer and Communication Engineering (ECCE). pp. 396–400 (Feb 2017). <https://doi.org/10.1109/ECACE.2017.7912937>
14. Heikes, K.E., Eddy, D.M., Arondekar, B., Schlessinger, L.: Diabetes risk calculator. *Diabetes Care* **31**(5), 1040–1045 (2008). <https://doi.org/10.2337/dc07-1150>
15. Ilango, B.S., Ramaraj, N.: A hybrid prediction model with f-score feature selection for type ii diabetes databases. In: Proceedings of the 1st Amrita ACM-W Celebration on Women in Computing in India. pp. 13:1–13:4. ACM, New York, NY, USA (2010). <https://doi.org/10.1145/1858378.1858391>
16. Jayanthi, N., Babu, B.V., Rao, N.S.: Survey on clinical prediction models for diabetes prediction. *Journal of Big Data* **4**(1), 26 (2017). <https://doi.org/10.1186/s40537-017-0082-7>
17. Jupyter Notebook Documentation: 7.0.0rc2 documentation. <https://jupyter-notebook.readthedocs.io/en/latest/index.html>, accedido: 2023-04-18
18. Kaur, H., Kumari, V.: Predictive modelling and analytics for diabetes using a machine learning approach. *Applied Computing and Informatics* (2018). <https://doi.org/https://doi.org/10.1016/j.aci.2018.12.004>
19. Maniruzzaman, M., Kumar, N., Menhazul Abedin, M., Shaykhul Islam, M., Suri, H.S., El-Baz, A.S., Suri, J.S.: Comparative approaches for classification of diabetes mellitus data: Machine learning paradigm. *CMPB* **152**, 23–34 (2017). <https://doi.org/https://doi.org/10.1016/j.cmpb.2017.09.004>

20. Meng, X.H., Huang, Y.X., Rao, D.P., Zhang, Q., Liu, Q.: Comparison of three data mining models for predicting diabetes or prediabetes by risk factors. *The Kaohsiung Journal of Medical Sciences* **29**(2), 93 – 99 (2013). <https://doi.org/https://doi.org/10.1016/j.kjms.2012.08.016>
21. Mercaldo, F., Nardone, V., Santone, A.: Diabetes mellitus affected patients classification and diagnosis through machine learning techniques. *Procedia Computer Science* **112**, 2519–2528 (2017). <https://doi.org/https://doi.org/10.1016/j.procs.2017.08.193>
22. Mir, A., Dhage, S.N.: Diabetes disease prediction using machine learning on big data of healthcare. In: 2018 Fourth International Conference on Computing Communication Control and Automation (ICCUBEA). pp. 1–6 (2018). <https://doi.org/10.1109/ICCUBEA.2018.8697439>
23. Rucci, E., Tittarelli, G., Ronchetti, F., , Elgart, J., Lanzarini, L., Gagliardino, J.J.: Primeras experiencias en la identificación de personas con riesgo de diabetes en la población argentina utilizando técnicas de aprendizaje automático. In: *Actas del XXIX Congreso Argentino de Ciencias de la Computación (CACIC 2023)*. pp. 146–158 (2023)
24. Scikit-Learn: 3.1. Cross-validation: evaluating estimator performance. https://scikit-learn.org/stable/modules/cross_validation.html, accedido: 2023-04-11
25. Sisodia, D., Sisodia, D.S.: Prediction of diabetes using classification algorithms. *Procedia Computer Science* **132**, 1578 – 1585 (2018). <https://doi.org/https://doi.org/10.1016/j.procs.2018.05.122>
26. Sneha, N., Gangil, T.: Analysis of diabetes mellitus for early prediction using optimal features selection. *Journal of Big Data* **6**(1), 13 (2019). <https://doi.org/10.1186/s40537-019-0175-6>
27. Tuomilehto, J., Lindström, J., Eriksson, J.G., Valle, T.T., Hämäläinen, H., Ilanne-Parikka, P., Keinänen-Kiukaanniemi, S., Laakso, M., Louheranta, A., Rastas, M., Salminen, V., Aunola, S., Cepaitis, Z., Moltchanov, V., Hakumäki, M., Mannelin, M., Martikkala, V., Sundvall, J., Uusitupa, M.: Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *New England Journal of Medicine* **344**(18), 1343–1350 (2001). <https://doi.org/10.1056/NEJM200105033441801>
28. Vistisen, D., Kivimäki, M., Perreault, L., Hulman, A., Witte, D.R., Brunner, E.J., Tabák, A., Jørgensen, M.E., Færch, K.: Reversion from prediabetes to normoglycaemia and risk of cardiovascular disease and mortality: the whitehall ii cohort study. *Diabetologia* **62**(8), 1385–1390 (Aug 2019). <https://doi.org/10.1007/s00125-019-4895-0>
29. Wei, S., Zhao, X., Miao, C.: A comprehensive exploration to the machine learning techniques for diabetes identification. In: 2018 IEEE 4th World Forum on Internet of Things (WF-IoT). pp. 291–295 (2018). <https://doi.org/10.1109/WF-IoT.2018.8355130>
30. Williams, R., Van Gaal, L., Lucioni, C.: Assessing the impact of complications on the costs of type ii diabetes. *Diabetologia* **45**(1), S13–S17 (Jul 2002). <https://doi.org/10.1007/s00125-002-0859-9>
31. Wu, H., Yang, S., Huang, Z., He, J., Wang, X.: Type 2 diabetes mellitus prediction model based on data mining. *Informatics in Medicine Unlocked* **10**, 100 – 107 (2018). <https://doi.org/https://doi.org/10.1016/j.imu.2017.12.006>
32. Xie, Z., Nikolayeva, O., Luo, J., Li, D.: Building risk prediction models for type 2 diabetes using machine learning techniques. *Preventing chronic disease* **16**, E130–E130 (Sep 2019). <https://doi.org/10.5888/pcd16.190109>

33. Yu, W., Liu, T., Valdez, R., Gwinn, M., Khoury, M.J.: Application of support vector machine modeling for prediction of common diseases: the case of diabetes and pre-diabetes. *BMC Medical Informatics and Decision Making* **10**(1), 16 (2010). <https://doi.org/10.1186/1472-6947-10-16>
34. Yuvaraj, N., SriPreethaa, K.R.: Diabetes prediction in healthcare systems using machine learning algorithms on hadoop cluster. *Cluster Computing* **22**(1), 1–9 (Jan 2019). <https://doi.org/10.1007/s10586-017-1532-x>
35. Zou, Q., Qu, K., Luo, Y., Yin, D., Ju, Y., Tang, H.: Predicting diabetes mellitus with machine learning techniques. *Frontiers in genetics* **9**, 515–515 (Nov 2018). <https://doi.org/10.3389/fgene.2018.00515>