

Two-sided Lieb-Thirring bounds

Sven Bachmann¹, Richard Froese¹, Severin Schraven¹

¹ Department of Mathematics, The University of British Columbia,
Vancouver, BC V6T 1Z2, Canada

March 29, 2024

Abstract

We prove upper and lower bounds for the number of eigenvalues of semi-bounded Schrödinger operators in all spatial dimensions. As a corollary, we obtain two-sided estimates for the sum of the negative eigenvalues of atomic Hamiltonians with Kato potentials. Instead of being in terms of the potential itself, as in the usual Lieb-Thirring result, the bounds are in terms of the landscape function, also known as the torsion function, which is a solution of $(-\Delta + V + M)u_M = 1$ in \mathbb{R}^d ; here $M \in \mathbb{R}$ is chosen so that the operator is positive. We further prove that the infimum of $(u_M^{-1} - M)$ is a lower bound for the ground state energy E_0 and derive a simple iteration scheme converging to E_0 .

1 Introduction

The Lieb-Thirring (LT) inequalities

$$\mathrm{tr}((-\Delta + V)_-^\gamma) \leq L_{\gamma,d} \int_{\mathbb{R}^d} V_-(x)^{\gamma + \frac{d}{2}} dx$$

are bounds on the moments of the negative eigenvalues of atomic Hamiltonians in terms of the negative part of the potential $V = V_+ - V_-$. In the context of the stability of matter for which it was originally developed [1], only an upper bound was needed and much of the vast amount of subsequent work extending and refining the estimates, see for example [2, 3], has similarly focussed on upper bounds. Among the few results about lower bounds, we mention [4, 5, 6] in special cases as well as [7, 8] for somewhat different estimates.

In this work, we give a simple proof of a two-sided estimate on the number of eigenvalues and their moments. The upper and lower bounds are matching in the sense that they differ only by a scaling and a multiplicative constant, but we are not able to conjecture the sharp values. The main difference with the LT inequality is that the bounds are not in terms of the potential V , but rather in terms of what we shall refer to as the *effective potential* $\frac{1}{u_M}$ where u_M is a solution of

$$(-\Delta + V + M \cdot \mathbf{1})u_M = 1. \tag{1.1}$$

Unlike in [9] where the potential V is assumed (among other hypotheses) to be a non-negative function, our assumption here is that the corresponding Schrödinger operator is semi-bounded below,

$$E_0 = \inf \text{spec}(-\Delta + V) > -\infty.$$

The constant in (1.1) is then $M > -E_0$. Among the auxiliary results of this paper, we will in particular show the existence of a weak solution u_M in the whole space, see also [10].

The equation (1.1) has been extensively studied without the potential term, i.e. for $-\Delta u_M = 1$, and its solution is known as the *torsion function*. It has been used to establish optimal Sobolev constant, estimate the ground state energy of the Laplacian and to obtain pointwise estimates for the eigenfunctions (and more general solutions) of elliptic operators [11, 12, 13, 14, 15, 16, 17, 18]. Following this line of work, the more general case with a potential term was considered only recently [19].

The connection of the torsion function with the phenomenon of localization for random Schrödinger operators was discovered in [20] in the case $M = 0$. In this context, it is mostly called the *landscape function* and it has been observed to yield excellent estimates on the density of states throughout the spectrum [21, 22, 23, 24] and to predict both the localization centres and exponential decay of localized eigenfunctions [25, 26, 27], see also [28]. As pointed out in [29] (or also implicitly in [30, page 2903]) the predicting power of the landscape function for the localization of eigenfunctions relies solely on the fact that the resolvent of the Schrödinger operator is positivity-preserving, a key property that we will use here as well.

The present work and our previous [9] focus on spectral estimates using the landscape function in the non-random case. In this context, the effective potential plays the role of a smoothing of the original potential at the right scale: It allows in particular for Cwikel-Lieb-Rozenblum (CLR)-type bounds [31, 32, 33] in cases where a strict CLR bound in terms of V fails. Specifically, the eigenvalue counting function can be estimated in terms of the volume of the sublevel sets of the effective potential. In this context, we mention [10, 34] for a magnetic version of the landscape function as well as the closely related approaches via Brownian motion [30], regularizing kernel [35], respectively maximal functions [36, 37, 38, 39, 40, 41, 42].

We shall see here that the key property for the effective potential to be useful in estimating the eigenvalues of the Schrödinger operator is that u_M satisfies a Harnack inequality, see (2.3) below. While we are not aware of the most general assumptions under which this holds, the results of [43, 44] yield that it does for atomic potentials, namely for negative V 's that are also in the Kato class (4.1), which are our main motivation here.

We state our main results in details in Section 2. In Section 3 we prove an abstract two-sided CLR-type bound conditional on the existence of a landscape function in the whole space and on a Harnack inequality. We then prove the validity of these assumptions in the case of Kato-class potentials in Section 4. In this case, the bounds can in fact be notably improved. In Section 5, we concentrate on the relationship between the infimum of the effective potential and the bottom of the spectrum: In particular, we propose an

iterative scheme which we converges to the ground state energy. Section 6 serves as both illustration of the results and analysis of their sharpness: On the one hand, in the case of the shallow square well in one dimension, we show how well the bottom of the effective potential approximates the ground state energy, and on the other hand we prove that our bounds match the asymptotic distribution of the eigenvalues at the bottom of the essential spectrum, in the examples of attractive radial potentials with a power law singularities. The proof of the two-sided Lieb-Thirring inequalities in terms of $\frac{1}{u_M} - M$ is provided in Section 7.

2 Results

One of the central objects of interest in this paper is the eigenvalue counting function for the Schrödinger operator $-\Delta + V$ on $L^2(\mathbb{R}^d)$, which is the rank of the spectral projection associated with $(-\infty, \mu]$, namely

$$\mathcal{N}^V(\mu) = \dim \text{Ran} \left(\mathbb{1}_{(-\infty, \mu]}(-\Delta + V) \right). \quad (2.1)$$

While most of the existing results in the literature are bounds on \mathcal{N}^V in terms of certain integrals of V , we aim here for estimates in terms of the effective potential, $\frac{1}{u_M}$ where u_M is, formally, the solution of (1.1). To start with, we shall assume its existence, namely that there is a $u_M \in H_{\text{loc}}^1(\mathbb{R}^d)$ such that

$$\int_{\mathbb{R}^d} (\nabla u_M(y) \cdot \nabla \varphi(y) + (V(y) + M)u_M(y)\varphi(y)) dy = \int_{\mathbb{R}^d} \varphi(y) dy \quad (2.2)$$

for all $\varphi \in C_c^\infty(\mathbb{R}^d)$. Our first bounds will be in terms of the volume $\mathcal{V}_M(\mu)$ of the sublevel sets of the effective potential

$$\mathcal{V}_M(\mu) = \int_{\{x \in \mathbb{R}^d : \frac{1}{u_M} \leq \mu\}} dx.$$

For the bound to hold, we assume what we shall refer to as a Harnack-Moser inequality, namely that there is $C_{HM} > 0$ such that

$$\sup_Q u_M \leq C_{HM} \left(\inf_Q u_M + \ell(Q)^2 \right) \quad (2.3)$$

for all cubes $Q \subseteq \mathbb{R}^d$ with sidelength $\ell(Q)$ (for exact definitions, see Section 3).

Theorem 2.1. *Let $V \in L_{\text{loc}}^1(\mathbb{R}^d)$ be such that*

$$E_0 = \inf \text{spec}(-\Delta + V) > -\infty.$$

Let $M > -E_0$ and let the solution u_M of (2.2) be positive, such that $u_M \in H_{\text{loc}}^1(\mathbb{R}^d) \cap L_{\text{loc}}^\infty(\mathbb{R}^d)$ and $\frac{1}{u_M} \in L_{\text{loc}}^\infty(\mathbb{R}^d)$. If (2.3) holds, then there exist constants $c_{0,d}, C_{0,d} > 0$ depending only on d, C_{HM} such that

$$(c_{0,d}\mu)^{d/2} \mathcal{V}_M(c_{0,d}\mu) \leq \mathcal{N}^{V+M}(\mu) \leq (C_{0,d}\mu)^{d/2} \mathcal{V}_M(C_{0,d}\mu), \quad (2.4)$$

for all $\mu \in \mathbb{R}$.

Note that the constants $c_{0,d}, C_{0,d}$ depend on M only via C_{HM} .

This bound in terms of the volume of the sublevel sets implies the following Lieb-Thirring bounds (see [45] for a study on how shifting affects the Lieb-Thirring inequalities in a similar setting).

Corollary 2.2. *Under the assumptions of Theorem 2.1 and if $E_0 < 0$, then*

$$c_{\gamma,d} \int_{\mathbb{R}^d} \left(|E_0| + \delta - \frac{1}{c_{0,d} u_M(x)} \right)_+^{\gamma + \frac{d}{2}} dx \leq \text{tr}((-\Delta + V)_-^\gamma) \quad (2.5)$$

for all $\gamma > 0$, and if $\gamma \geq 1$,

$$\text{tr}((-\Delta + V)_-^\gamma) \leq C_{\gamma,d} \int_{\mathbb{R}^d} \left(|E_0| + 2\delta - \frac{1}{C_{0,d} u_M(x)} \right)_+^{\gamma + \frac{d}{2}} dx. \quad (2.6)$$

Here, $\delta = M - |E_0|$ and

$$c_{\gamma,d} = \frac{c_{0,d}^{d/2} \gamma}{\frac{d}{2} + \gamma} \min \left\{ 1, \frac{\delta}{|E_0|} \right\}^{\frac{d}{2}}, \quad C_{\gamma,d} = \frac{C_{0,d}^{d/2} \gamma}{\frac{d}{2} + \gamma} \left(1 + \frac{|E_0|}{\delta} \right)^{\frac{d}{2}},$$

where $c_{0,d}, C_{0,d}$ are the constants of Theorem 2.1 for the potential $V + M$.

The Harnack-Moser inequality (2.3) controls the oscillations of u_M at all scales and is the natural choice in the present context. It originates in the PDE literature in slightly different settings. On finite domains with $V \equiv 0$ this inequality holds always true for nonnegative solutions of (2.2), see [46, Theorem 3.3]). On the other hand, for Kato-class potentials (see [44, Theorem 2.5]) it is known that nonnegative solutions of the homogeneous equation $(-\Delta + V)u = 0$ satisfy a Harnack inequality for balls with small radius, namely $\sup_{B(x,r)} u \leq C_H \inf_{B(x,r)} u$. The key difficulty to overcome here is the lack of compactness as we are working in \mathbb{R}^d and need the inequality to hold for arbitrary large cubes.

The assumptions of the above theorem, and in particular (2.3) can be verified for the physically relevant potentials in the Kato-class \mathcal{K}_d (see Section 4 for a precise definition). This includes all bounded potentials as well as power laws singularities $|x|^{-\rho}$ for $\rho \in [0, \min\{d, 2\})$. A ‘crystal’ of Coulomb singularities is allowed too.

Proposition 2.3. *Let V be such that $V_+ \in \mathcal{K}_d^{\text{loc}}, V_- \in \mathcal{K}_d$. Then $E_0 = \inf \text{spec}(-\Delta + V) > -\infty$. For all $M > -E_0$ there exists a positive solution u_M of (2.2) such that $u_M \in L^\infty(\mathbb{R}^d) \cap C^0(\mathbb{R}^d) \cap H_{\text{loc}}^1(\mathbb{R}^d)$ and $\frac{1}{u_M} \in L_{\text{loc}}^\infty(\mathbb{R}^d)$. If, additionally, $V_+ \in \mathcal{K}_d$, then $u_M^{-1} \in L^\infty(\mathbb{R}^d)$ and u_M satisfies (2.3). In fact,*

$$0 < A_M = \frac{\sup_{\mathbb{R}^d} u_M}{\inf_{\mathbb{R}^d} u_M} < \infty.$$

The fact that, with a Kato condition, the operator $-\Delta + V$ is bounded from below is well-known, see [47, Theorem A.2.7.]). Note that the very last claim of the theorem is

a global Harnack inequality which implies (2.3), although for a potentially suboptimal constant since $C_{HM} \leq A_M$.

The bound (2.4), while very general indeed, is not optimal as the spectral parameter is shifted by M . In other words, a sharp estimate would use $\frac{1}{u_M} - M$ instead of $\frac{1}{u_M}$ as the effective potential. The following theorem realizes this improvement for Kato-class potentials. Specifically, we consider the unshifted counting function $\mathcal{N}^V(\mu)$ and show (i) that a lower bound in terms of coarsened-grained volume of the sublevel sets of $\frac{1}{u_M} - M$ always holds, (ii) that the coarse-graining can be removed if $\frac{1}{u_M} - M$ satisfies a scale-invariant Harnack inequality and (iii) that a CLR-type upper bound, with V replaced by $\frac{1}{u_M} - M$, holds in dimensions 3 and higher.

Theorem 2.4. *Let $V \in \mathcal{K}_d$ and $M > -E_0$. Let u_M be the function given by Proposition 2.3.*

i) *Let $\mu < 0$. For any $c > 1$, let*

$$n_c(\mu) = \left| \left\{ Q \in \mathcal{Q}_{(C_c|\mu|)^{-1/2}} : \sup_Q \left(\frac{1}{u_M} - M \right) \leq c\mu \right\} \right|, \quad C_c = \frac{c-1}{2^d(5A_M)^2}, \quad (2.7)$$

where \mathcal{Q}_ℓ is a partition of \mathbb{R}^d into cubes of side length ℓ , see Section 3. Then

$$\mathcal{N}^V(\mu) \geq n_c(\mu). \quad (2.8)$$

ii) *Let $c > 1$. Assume there exists $\tilde{C}_H > 0$ such that for all $\ell > 0$ and all $Q \in \mathcal{Q}_\ell$ with $\sup_Q \left(\frac{1}{u_M} - M \right) \leq -\frac{c}{C_c \ell^2}$,*

$$\sup_Q \left(\frac{1}{u_M} - M \right) \leq \tilde{C}_H \inf_Q \left(\frac{1}{u_M} - M \right). \quad (2.9)$$

Then

$$\mathcal{N}^V(\mu) \geq (C_c|\mu|)^{d/2} \left| \left\{ x \in \mathbb{R}^d : \frac{1}{u_M(x)} - M \leq \frac{c\mu}{\tilde{C}_H} \right\} \right| \quad (2.10)$$

for all $\mu < 0$.

iii) *Let $d \geq 3$. There exist a constant $C_{\text{CLR}} > 0$ depending only on d such that*

$$\mathcal{N}^V(\mu) \leq C_{\text{CLR}} A_M^d \int_{\mathbb{R}^d} \left((\mu + \varepsilon) - \left(\frac{1}{u_M(x)} - M \right) \right)_+^{d/2} dx, \quad (2.11)$$

for all $\mu \in \mathbb{R}$ and all $\varepsilon > 0$.

Here again, these bounds yield upper and lower Lieb-Thirring inequalities. The difference with the bounds of Corollary 2.2 cannot be understated: Here, the effective potential appears as it really should, without an additional scaling and shift.

Corollary 2.5. *Under the assumptions of Theorem 2.4(ii), respectively (iii),*

$$k_{\gamma,d} \int_{\mathbb{R}^d} \left(\frac{1}{u_M(x)} - M \right)_-^{\gamma + \frac{d}{2}} dx \leq \text{tr}((-\Delta + V)_-^\gamma) \quad (2.12)$$

for all $\gamma > 0$, and if $\gamma \geq 1$,

$$\text{tr}((-\Delta + V)_-^\gamma) \leq L_{\gamma,d} A_M^{2\gamma+d} \int_{\mathbb{R}^d} \left(\frac{1}{u_M(x)} - M \right)_-^{\gamma + \frac{d}{2}} dx, \quad (2.13)$$

where $L_{\gamma,d}$ are the standard Lieb-Thirring constants and $k_{\gamma,d} = C_c^{\frac{d}{2}} \frac{\gamma}{\gamma + \frac{d}{2}} \left(\frac{\tilde{C}_H}{c} \right)^{\gamma + \frac{d}{2}}$.

While the above results are valid throughout the spectrum, their sharpness comes into focus at two critical points: the bottom of the essential spectrum and the bottom of the spectrum. The first proposition below shows that the infimum of $\frac{1}{u_M} - M$ is a lower bound for the ground state energy E_0 , see also [28] for nonnegative potentials as well as the previous work [13]. This in turn allows us to set up an iterative procedure converging to E_0 . Finally, we show that the CLR bound using $\frac{1}{u_M} - M$ yields the exact asymptotics of the negative eigenvalues accumulating at 0^- for ‘atomic’ potentials $V(x) = -|x|^{-\rho}$.

Proposition 2.6. *Let $V \in \mathcal{K}_d$ and $M > -E_0$. Let u_M be the function given by Proposition 2.3. We have*

$$E_0 \geq \inf_{\mathbb{R}^d} \left(\frac{1}{u_M} - M \right). \quad (2.14)$$

It may seem that the proposition is relatively useless without a priori information on the ground state energy. This is erroneous as the result can be bootstrapped to provide a convergent sequence converging to E_0 . As we shall see in Section 6, the approximation may already be very good even after one iteration, even if $M - E_0$ is very large.

Corollary 2.7. *Let $V \in \mathcal{K}_d$ and let*

$$F : (-E_0, \infty) \rightarrow (-E_0, \infty), \quad M \mapsto M - \inf_{\mathbb{R}^d} \frac{1}{u_M},$$

Let $M_0 \in (-E_0, \infty)$ and define recursively $M_{n+1} = F(M_n)$ for $n \in \mathbb{N}$. Then $M_{n+1} \leq M_n$ and $M_n \rightarrow -E_0$ as $n \rightarrow \infty$. Moreover,

$$\lim_{M \rightarrow (-E_0)^+} \|u_M\|_{L^\infty(\mathbb{R}^d)} = \infty. \quad (2.15)$$

We remark that (2.15) is in stark contrast to the random case. A shifted landscape function was considered in [48] in the case of a random potential. There, it is shown that the limit $M \rightarrow -E_0^+$ can be taken after averaging over the random potential. In the present case where no averaging is available, we will see explicitly in Section 6 in the case of a radial potential well that u_M blows up at every single point in the same limit.

Finally, we turn to the asymptotics of the eigenvalue counting function as $\mu \rightarrow 0^-$.

Proposition 2.8. *Let $\rho_d = \min\{d, 2\}$. For $\rho \in (0, \rho_d)$ the potential $V(x) = -|x|^{-\rho}$ is in the Kato class \mathcal{K}_d . Let $M > -E_0$ and let u_M be the function given by Proposition 2.3. Then as $\mu \rightarrow 0^-$*

$$\mathcal{N}^V(\mu) = (1 + o(1)) \left[(2\sqrt{\pi})^d \Gamma\left(\frac{d}{2} + 1\right) \right]^{-1} \int_{\mathbb{R}^d} \left(\mu - \left(\frac{1}{u_M(x)} - M \right) \right)_+^{d/2} dx \quad (2.16)$$

$$= (1 + o(1)) \mathcal{N}^{\frac{1}{u_M} - M}(\mu). \quad (2.17)$$

The first equality above shows the exactness of the CLR asymptotics when using the landscape function $\frac{1}{u_M} - M$, with the same constant as with the original potential V . In the second equality, we point out that $\mathcal{N}^{\frac{1}{u_M} - M}$ is the counting function for the operator $-\Delta + (\frac{1}{u_M} - M)$: This is non-trivial since $-\Delta + V$ is unitarily equivalent to $-\frac{1}{u_M^2} \nabla \cdot u_M^2 \nabla + (\frac{1}{u_M} - M)$, namely for a modified kinetic energy. This proves a conjecture of [22, Equation (1.5)] in this particular case.

To conclude this section, we point out that our proofs allow us to treat magnetic Schrödinger operators with suitably well-behaved magnetic fields.

3 Variational argument: Proof for Theorem 2.1

We follow the strategy of [9] and first estimate \mathcal{N}^V in terms of some coarse-grained volume of the sublevel sets of the effective potential. In a second step we will relate this to the measure of the sublevel set of $1/u_M$.

We first introduce some notation. A box of sidelength ℓ is a set of the form $\times_{i=1}^d [a_i, b_i]$ where $b_i - a_i = \ell$ and $a_i, b_i \in \ell\mathbb{Z}$. For any $\ell > 0$, we consider the collection \mathcal{Q}_ℓ of boxes of sidelength ℓ such that $\bigcup_{Q \in \mathcal{Q}_\ell} Q = \mathbb{R}^d$ and $\hat{Q} \cap \hat{Q}' = \emptyset$ whenever $Q \neq Q'$. We define for any $\mu > 0$

$$N(\mu) = \left| \left\{ Q \in \mathcal{Q}_{\mu^{-1/2}} : \inf_Q \frac{1}{u_M} \leq \mu \right\} \right|$$

and

$$n(\mu) = \left| \left\{ Q \in \mathcal{Q}_{\mu^{-1/2}} : \sup_Q \frac{1}{u_M} \leq \mu \right\} \right|.$$

The reason we are considering only cubes with corner on $\ell\mathbb{Z}^d$ is to make it possible to compare \mathcal{Q}_ℓ with $\mathcal{Q}_{n\ell}$ for $n \in \mathbb{N}$, the former being a refinement of the later.

For the class of potentials considered here, namely those satisfying the Harnack condition (2.3), both coarse-grained volumes are directly related to the measure $\mathcal{V}(\mu)$ of the sublevel set. The following lemma is essentially the same as [9, Lemma 4.1].

Lemma 3.1. *Let V satisfy the conditions of Theorem 2.1. Then*

$$n(\mu) \leq \mu^{d/2} \mathcal{V}_M(\mu) \leq N(\mu) \leq n\left(\left[\sqrt{2C_{HM}}\right]^2 \mu\right)$$

for all $\mu \in \mathbb{R}$. Here C_{HM} is the Harnack-Moser constant of (2.3).

Proof. The first two inequalities are immediate as, up to null sets, $n(\mu)/\mu^{d/2}$ is the measure of all boxes that are strictly contained in the sublevel set $\{1/u_M \leq \mu\}$ and $N(\mu)/\mu^{d/2}$ is the measure of all the boxes that intersect the sublevel set.

Let $K = \lceil \sqrt{2C_{HM}} \rceil^2$, then the cubes in $\mathcal{Q}_{(K\mu)^{-1/2}}$ are subcubes of exactly one cube in $\mathcal{Q}_{\mu^{-1/2}}$ since \sqrt{K} is an integer. To prove the last inequality, it is sufficient to show that every cube in $\mathcal{Q}_{\mu^{-1/2}}$ which contributes to $N(\mu)$ admits a subcube in $\mathcal{Q}_{(K\mu)^{-1/2}}$ which contributes to $n(K\mu)$. If $Q \in \mathcal{Q}_{\mu^{-1/2}}$ satisfies $\sup_Q \frac{1}{u_M} \leq \mu$ then all its subcubes contribute to $n(K\mu)$. The only other option for Q to contribute to $N(\mu)$ is if $\inf_Q \frac{1}{u_M} \leq \mu < \sup_Q \frac{1}{u_M}$. For those, we have in particular $\sup_Q u_M \geq \frac{1}{\mu}$. Now pick a subcube $\tilde{Q} \in \mathcal{Q}_{(K\mu)^{-1/2}}$ of Q such that $\sup_{\tilde{Q}} u_M \geq \frac{1}{\mu}$. Then by (2.3) we get, as $K = \lceil \sqrt{2C_{HM}} \rceil^2 \geq 2C_{HM}$,

$$\inf_{\tilde{Q}} u_M \geq \frac{1}{C_{HM}} \sup_{\tilde{Q}} u_M - \frac{1}{K\mu} \geq \frac{1}{C_{HM}\mu} - \frac{1}{2C_{HM}\mu} = \frac{1}{2C_{HM}\mu}.$$

Thus, $\sup_{\tilde{Q}} \frac{1}{u_M} \leq 2C_{HM}\mu \leq K\mu$, and so \tilde{Q} contributes to $n(K\mu)$. \square

We now structure the proof of Theorem 2.1 in two lemmas for the upper, respectively the lower bound.

Lemma 3.2. *Let V satisfy the conditions of Theorem 2.1. Then*

$$\mathcal{N}^{V+M}(\mu) \leq N(C\mu)$$

for all $C > \frac{4dC_H^2}{\pi^2}$ and all $\mu \in \mathbb{R}$.

Proof. For $\mu \leq 0$ we have $\mathcal{N}^{V+M}(\mu) = 0 = N(C\mu)$. Thus, we will assume for the rest of the proof that $\mu > 0$. In order to have that $\mathcal{N}^V(\mu) \leq N$ it suffices, by the Min-Max Principle (see [49, Theorem XIII.2]), to exhibit a subspace $\mathcal{H}_N \subseteq \text{dom}(H^{1/2})$ with codimension at most N such that

$$\int_{\mathbb{R}^d} (|\nabla v|^2 + (V + M)|v|^2) > \mu \int_{\mathbb{R}^d} |v|^2$$

for all $v \in \mathcal{H}_N$. Let \mathcal{F} be the collection of boxes such that

$$\mathcal{F} = \left\{ Q \in \mathcal{Q}_{(C\mu)^{-1/2}} : \inf_Q \frac{1}{u_M} \leq \mu \right\},$$

where $C > 0$ will be chosen later, and let

$$\mathcal{H}_N = \left\{ v \in \text{dom}(H^{1/2}) : \int_Q \frac{v}{u_M} = 0 \quad \forall Q \in \mathcal{F} \right\}.$$

The codimension of \mathcal{H}_N is equal to $|\mathcal{F}| = N(C\mu)$.

We first claim that

$$\int_{\mathbb{R}^d} (|\nabla\varphi|^2 + (V + M)|\varphi|^2) = \int_{\mathbb{R}^d} u_M^2 \left(\left| \nabla \left(\frac{\varphi}{u_M} \right) \right|^2 + \frac{1}{u_M} \left| \frac{\varphi}{u_M} \right|^2 \right) \quad (3.1)$$

for all $\varphi \in H^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$ with compact support. As $u_M \in H_{\text{loc}}^1(\mathbb{R}^d)$ and $1/u_M \in L_{\text{loc}}^\infty(\mathbb{R}^d)$ by assumption, we get that $\varphi^2/u_M \in H^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$ with compact support and thus, after a simple approximation argument, we have

$$\int_{\mathbb{R}^d} \left(\nabla u_M \cdot \nabla \left(\frac{|\varphi|^2}{u_M} \right) + (V + M)u_M \frac{|\varphi|^2}{u_M} \right) = \int_{\mathbb{R}^d} \frac{|\varphi|^2}{u_M},$$

since u_M solves (2.2). Furthermore, using the product rule for Sobolev functions [50, Lemma 7.4] yields

$$\nabla u_M \cdot \nabla \left(\frac{|\varphi|^2}{u_M} \right) = |\nabla\varphi|^2 - u_M^2 \left| \nabla \left(\frac{\varphi}{u_M} \right) \right|^2,$$

which readily implies (3.1). As $C_c^\infty(\mathbb{R}^d)$ is dense in the form domain of $-\Delta + V + M$ (see [47, Theorem A.2.8.]) the statement of the lemma follows if we can show that there exists $\varepsilon > 0$ such that for all non-zero $\varphi \in C_c^\infty(\mathbb{R}^d) \cap \mathcal{H}_N$,

$$\int_{\mathbb{R}^d} \left(u_M^2 \left| \nabla \left(\frac{\varphi}{u_M} \right) \right|^2 + \frac{|\varphi|^2}{u_M} \right) > (1 + \varepsilon)\mu \int_{\mathbb{R}^d} |\varphi|^2.$$

The additional ε is needed to ensure that the strict inequality (without ε) holds true on all of \mathcal{H}_N . We check this inequality using the partition into boxes. In any box $Q \notin \mathcal{F}$, the bound $\inf_Q 1/u_M > \mu$ and the nonnegativity of the first term yield the claim. If $Q \in \mathcal{F}$, we use that $\varphi \in \mathcal{H}_N$, namely that the integral of φ/u_M vanishes, together with Poincaré inequality on Q with optimal constant $\frac{\pi^2}{d}(C\mu)$, see [51], to conclude that

$$\int_Q u_M^2 \left| \nabla \left(\frac{\varphi}{u_M} \right) \right|^2 \geq \left(\frac{\inf_Q u_M}{\sup_Q u_M} \right)^2 \frac{\pi^2 C\mu}{d} \int_Q |\varphi|^2.$$

However, using the Harnack-Moser inequality (2.3) and the definition of \mathcal{F} we obtain

$$C_{HM} \inf_Q u_M \geq \sup_Q u_M - \frac{C_{HM}}{C\mu} \geq \left(1 - \frac{C_{HM}}{C} \right) \frac{1}{\mu}.$$

Thus, $\inf_Q u_M \geq \frac{1}{2C_{HM}\mu}$ for any $C \geq 2C_{HM}$ and hence, again by (2.3)

$$\sup_Q u_M \leq C_{HM} \left(\inf_Q u_M + (C\mu)^{-1} \right) \leq 2C_{HM} \inf_Q u_M.$$

This yields

$$\int_Q u_M^2 \left| \nabla \left(\frac{\varphi}{u_M} \right) \right|^2 \geq \frac{C\pi^2}{4dC_{HM}^2} \mu \int_Q |\varphi|^2$$

and so any choice of $C > \frac{4dC_{HM}^2}{\pi^2}$ will give the desired estimate. \square

Lemma 3.3. *Let V satisfy the conditions of Theorem 2.1. Then*

$$n(\mu) \leq \mathcal{N}^{V+M}(C\mu)$$

where $C = 1 + 2^{d+2}C_{HM}^2$ and for all $\mu \in \mathbb{R}$.

Proof. For a lower bound $N \leq \mathcal{N}^V(C\mu)$ it suffices, again by the Min-Max Principle, to find a subspace $\mathcal{H}_N \subseteq \text{dom}(H^{1/2})$ of dimension at least N such that

$$\int_{\mathbb{R}^d} (|\nabla v|^2 + (V + M)|v|^2) \leq C\mu \int_{\mathbb{R}^d} |v|^2.$$

We define

$$\mathcal{F} = \left\{ Q \in \mathcal{Q}_{\mu^{-1/2}} : \sup_Q \frac{1}{u_M} \leq \mu \right\}.$$

Furthermore, for a box Q we pick $\chi_Q \in C_c^\infty(Q)$ with $0 \leq \chi_Q \leq 1$, $\|\nabla \chi_Q\|_{L^\infty(\mathbb{R}^d)} \leq \tilde{C}\mu^{1/2}$ for some $\tilde{C} > 4$, $\chi_Q \equiv 1$ on $Q/2$. Since the functions $\chi_Q u_M$ are non-zero and orthogonal to each other, the space

$$\mathcal{H}_N = \text{span}\{\chi_Q u_M : Q \in \mathcal{F}\}$$

is of dimension $|\mathcal{F}| = n(\mu)$.

By assumption we have $u_M \in H_{\text{loc}}^1(\mathbb{R}^d) \cap L_{\text{loc}}^\infty(\mathbb{R}^d)$ and thus $\chi_Q u_M$ is in $H^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$ with compact support. By (3.1) and an approximation argument for $\chi_Q u_M$, we get

$$\int_{\mathbb{R}^d} (|\nabla(\chi_Q u_M)|^2 + (V + M)\chi_Q^2 u_M^2) = \int_{\mathbb{R}^d} (\chi_Q^2 u_M^2 + |\nabla \chi_Q|^2 u_M^2), \quad (3.2)$$

and hence, since $Q \in \mathcal{F}$,

$$\begin{aligned} \int_{\mathbb{R}^d} (|\nabla(\chi_Q u_M)|^2 + (V + M)\chi_Q^2 u_M^2) &\leq \left(\sup_Q \frac{1}{u_M} \right) \int_Q \chi_Q^2 u_M^2 + \tilde{C}^2 \mu \int_Q u_M^2 \\ &\leq \mu \left(\int_Q \chi_Q^2 u_M^2 + \tilde{C}^2 \int_Q u_M^2 \right). \end{aligned} \quad (3.3)$$

We are left to replace the integrand of the second term by $(\chi_Q u_M)^2$. As $Q \in \mathcal{F}$ we have $\inf_Q u_M \geq 1/\mu$ and hence, by the Harnack-Moser inequality (2.3) we get

$$\sup_Q u_M \leq C_{HM} \left(\inf_Q u_M + \frac{1}{\mu} \right) \leq 2C_{HM} \inf_Q u_M.$$

We conclude that

$$\int_Q u_M^2 \leq |Q| \sup_Q u_M^2 \leq |Q| (2C_{HM})^2 \inf_{Q/2} u_M^2 \leq 2^{d+2} C_{HM}^2 \int_{Q/2} u_M^2 \leq 2^{d+2} C_{HM}^2 \int_{\mathbb{R}^d} \chi_Q^2 u_M^2,$$

where the last inequality follows from the properties of χ_Q . This yields

$$n(\mu) \leq \mathcal{N}^{V+M}((1 + \tilde{C}2^{d+2}C_{HM}^2)\mu).$$

As $\mu \mapsto \mathcal{N}^{V+M}(\mu)$ is right-continuous and any $\tilde{C} > 4$ is admissible, we obtain the claim. \square

4 Kato-class potentials

We now turn to the case of potentials V that are in the Kato-class \mathcal{K}_d . We say that a measurable function $V : \mathbb{R}^d \rightarrow \mathbb{R}$ is in the Kato-class \mathcal{K}_d if

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0^+} \sup_{x \in \mathbb{R}^d} \int_{B(x, \varepsilon)} |x - y|^{-(d-2)} |V(y)| dy &= 0, & d \geq 3, \\ \lim_{\varepsilon \rightarrow 0^+} \sup_{x \in \mathbb{R}^d} \int_{B(x, \varepsilon)} \ln(|x - y|^{-1}) |V(y)| dy &= 0, & d = 2, \\ \sup_{x \in \mathbb{R}^d} \int_{B(x, 1)} |V(y)| dy &< \infty, & d = 1. \end{aligned} \quad (4.1)$$

We equip the sets \mathcal{K}_d with the following norms:

$$\|V\|_{\mathcal{K}_d} = \begin{cases} \sup_{x \in \mathbb{R}^d} \int_{B(x, 1)} |x - y|^{-(d-2)} |V(y)| dy, & d \geq 3, \\ \sup_{x \in \mathbb{R}^d} \int_{B(x, 1/2)} \ln(|x - y|^{-1}) |V(y)| dy, & d = 2, \\ \sup_{x \in \mathbb{R}^d} \int_{B(x, 1)} |V(y)| dy, & d = 1. \end{cases}$$

Furthermore, we say $V \in \mathcal{K}_d^{\text{loc}}$ if $\mathbb{1}_{B(0, R)} V \in \mathcal{K}_d$ for all $R > 0$.

We start by recalling some properties about the Green's function of $-\Delta + V$ in this particular case.

Theorem 4.1. [47, Theorem B.7.2.] *Suppose $V_+ \in \mathcal{K}_d^{\text{loc}}$, $V_- \in \mathcal{K}_d$. Let $0 < \alpha < d/2$ and $\text{Re}(z) < \inf \text{spec}(-\Delta + V)$ or $z \notin \text{spec}(-\Delta + V)$, α integral. Then $(-\Delta + V - z)^{-\alpha}$ is an integral operator with integral kernel $G^{(\alpha)}(x, y; z)$ obeying*

1. $G^{(\alpha)}(\cdot, \cdot, z)$ is continuous away from $x = y$ and bounded uniformly in each region $\{(x, y) \in \mathbb{R}^d \times \mathbb{R}^d : |x - y| \geq a\}$ for $a > 0$.

2. We have

$$|G^{(\alpha)}(x, y; z)| \leq C|x - y|^{-d+2\alpha}. \quad (4.2)$$

3. For $|x - y|$ sufficiently small and $|x| < R$

$$|G^{(\alpha)}(x, y; z)| \geq \tilde{C}_R |x - y|^{-d+2\alpha}. \quad (4.3)$$

If $V \in \mathcal{K}_d$, \tilde{C}_R can be chosen independent of R .

4. For $|x - y| \geq 1$

$$|G^{(\alpha)}(x, y; z)| \leq C_{\delta, \alpha, z} \exp(-\delta|x - y|) \quad (4.4)$$

for some $\delta > 0$ and if $\text{Re}(z) < \inf \text{spec}(-\Delta + V)$, for all δ with $\frac{1}{2}\delta^2 < \inf \text{spec}(-\Delta + V) - \text{Re}(z)$.

This allows us to prove next that the Green's function is nonnegative.

Lemma 4.2. *Under the assumptions of Proposition 2.3 we have for $d \geq 3$ that $G^{(1)}(x, y; -M) \geq 0$ and for all $f \in L^2(\mathbb{R}^d)$ we have that $f \geq 0$ implies $(-\Delta + V + M)^{-1}f \geq 0$.*

Proof. By Theorem 4.1 we know that $G^{(1)}(\cdot, \cdot; -M)$ is continuous away from the diagonal. Hence, it is enough to prove that $(-\Delta + V + M)^{-1}$ maps nonnegative L^2 functions to nonnegative L^2 functions (otherwise consider a positive bump function localize where the Green's function is negative to get a contradiction).

We note that for all $t > 0$ we have that $(-\Delta + t)^{-1}$ maps nonnegative L^2 functions to nonnegative L^2 functions, which can easily be seen from the fact that the corresponding explicit integral kernel is positive. As V_- is form bounded by the Laplacian with relative bound zero (see [47, Proposition A.2.3]) and $\mathcal{D}((-\Delta)^{1/2}) \cap \mathcal{D}(V_+^{1/2})$ is dense in $L^2(\mathbb{R}^d)$ (as $C_c^\infty(\mathbb{R}^d)$ is contained in this intersection), we get by [52, Lemma 2.1, Lemma 2.2] applied to V_+ and V_- that $(-\Delta + V + M)^{-1}$ maps nonnegative L^2 functions to nonnegative L^2 functions. \square

Proof of Proposition 2.3. We start with the case $d \geq 3$. In the end we will use Hadamard's method of descent to obtain all the properties for $d \in \{1, 2\}$ from the case $d = 3$.

By assumption $\inf \text{spec}(-\Delta + V + M) > 0$ and thus, the Green's function $G_M(x, y) := G^{(1)}(x, y; -M)$ satisfies the properties of Theorem 4.1. We define

$$u_M(x) = \int_{\mathbb{R}^d} G_M(x, y) dy$$

and will check that this function satisfies indeed all the properties of Proposition 2.3.

The boundedness of u_M follows immediately from (4.2) and (4.4). In order to check that u_M is a weak solution of (1.1) we follow the proof of [9, Proposition 3.3] and set $u_{M,L}$ to be the solutions of

$$(-\Delta + V + M)u_{M,L} = \mathbb{1}_{B(0,L)}. \quad (4.5)$$

Since $\mathbb{1}_{B(0,L)}$ is in $L^2(\mathbb{R}^d)$, Theorem 4.1 immediately implies that

$$u_{M,L}(x) = \int_{\mathbb{R}^d} G_M(x, y) \mathbb{1}_{B(0,L)}(y) dy.$$

Hence, by dominated convergence we have $\lim_{L \rightarrow \infty} u_{M,L}(x) = u_M(x)$. As $u_{M,L}$ is a weak solution of (4.5), one readily checks that u_M is a solution of (2.2). The fact that $1/u_M \in L_{\text{loc}}^\infty(\mathbb{R}^d)$. This follows from the lower bound (4.3) together with the fact that the Green's function G_M is nonnegative by Lemma 4.2. As $1/u_M \in L_{\text{loc}}^\infty(\mathbb{R}^d)$ we get $V + M + 1/u_M \in \mathcal{K}_d^{\text{loc}}$. The function u_M is nonnegative and is a distributional solution of $(-\Delta + V + M + 1/u_M)u_M = 0$. Thus, [43, Theorem 1.5] yields that u_M is continuous. Finally, $u_M \in H_{\text{loc}}^1(\mathbb{R}^d)$ follows from a Caccioppoli-type inequality (see [44, Lemma 1.1]).

The cases $d \in \{1, 2\}$ will now be covered by Hadamard's method of descent. Let's start with $d = 2$. We define $\tilde{V}(y_1, y_2, y_3) = V(y_1, y_2)$ and check that \tilde{V} satisfies the conditions of Proposition 2.3 for $d = 3$. Once we have done that, we can use the same proof as in [9, page 10] to conclude. Similarly, one deals with the case $d = 1$.

We first check that if $-\Delta_{\mathbb{R}^d} + V \geq M$, then we also have $-\Delta_{\mathbb{R}^3} + \tilde{V} \geq M$. To see this for $d = 2$, we take a Fourier transform in the variables and note that $-\Delta_{\mathbb{R}^3} + \tilde{V}$ is unitarily equivalent to the direct integral

$$\int_{\mathbb{R}}^{\oplus} (p^2 + (-\Delta_{\mathbb{R}^2} + V)) dp.$$

Thus, by [49, Theorem XIII.85] we have $\text{spec}(-\Delta_{\mathbb{R}^3} + \tilde{V}) = [\min \text{spec}(-\Delta_{\mathbb{R}^2} + V), \infty)$ and hence, establishes our first property.

We are left to show that for $d \in \{1, 2\}$ if $V \in \mathcal{K}_d$ (respectively $\mathcal{K}_d^{\text{loc}}$), then $\tilde{V} \in \mathcal{K}_3$ (respectively $\mathcal{K}_3^{\text{loc}}$). Let $d = 2$, $x \in \mathbb{R}^3$ and $0 < \varepsilon \leq 1$. We compute

$$\begin{aligned} \int_{B(x, \varepsilon)} \frac{|\tilde{V}(y)|}{|x - y|} dy &\leq 2 \int_{B((0,0), \varepsilon)} |V(z_1 + x_1, z_2 + x_2)| \left(\frac{\pi}{2} + \ln \left(\frac{\varepsilon}{\sqrt{z_1^2 + z_2^2}} \right)_+ \right) dz_2 dz_3 \\ &\leq (\pi + 2) \int_{B((x_1, x_2), \varepsilon)} \frac{|V(z_1, z_2)|}{|(x_1, x_2) - (z_1, z_2)|} dz_1 dz_2. \end{aligned}$$

For $d = 1$ we define $\tilde{V}(y_1, y_2) = V(y_1)$. We compute for $x \in \mathbb{R}^2$ and $0 < \varepsilon \leq 1$

$$\begin{aligned} \int_{B(x, \varepsilon)} |\tilde{V}(y)| \ln(|x - y|^{-1}) dy &\leq \int_{B(0, \varepsilon)} |V(x_1 + y_1)| \left(\int_{-1}^1 |\ln(|y_2|)| dy_2 \right) dy_1 \\ &\leq 2 \sup_{x_1 \in \mathbb{R}} \int_{B(x_1, \varepsilon)} |V(y_1)| dy_1. \end{aligned}$$

Let $d = 2$, let $\tilde{V}(y_1, y_2, y_3) = V(y_1, y_2)$ and let \tilde{u}_M be the solution of

$$(-\Delta_{\mathbb{R}^3} + \tilde{V} + M)\tilde{u}_M = 1$$

which we obtained in the case $d = 3$. One readily checks that $\widetilde{v_{M,L}}(x, t) = \widetilde{u_{M,L}}(x, t + \alpha)$ is a weak solution of $(-\Delta + \tilde{V} + M)\widetilde{v_{M,L}} = \mathbb{1}_{B((0,0,\alpha), L)}$. Thus, we have

$$\widetilde{u_{M,L}}(x, t + \alpha) = \int_{\mathbb{R}^3} G_M((x, t), y) \mathbb{1}_{B((0,0,\alpha), L)}(y) dy.$$

By dominated convergence we get

$$\widetilde{u}_M(x, t + \alpha) = \lim_{L \rightarrow \infty} \widetilde{u_{M,L}}(x, t + \alpha) = \int_{\mathbb{R}^3} G_M((x, t), y) dy = \widetilde{u}_M(x, t).$$

We define $u_M(x) = \widetilde{u}_M(x, 0)$. One readily checks that u_M inherits all the desired properties from \widetilde{u}_M . The same strategy works for $d = 1$.

Finally, we consider the case where $V_+ \in \mathcal{K}_d$. In this case the lower bound (4.3) holds true for all x and hence $\inf_{\mathbb{R}^d} u_M > 0$. This implies in turn that $0 < A_M < \infty$ since we have already established that $\sup_{\mathbb{R}^d} u_M < \infty$. \square

Next we turn to the proof of Theorem 2.4.

Proof of Theorem 2.4. (i) We adapt the proof in Lemma 3.3 to the potential $\frac{1}{u_M} - M$ and define

$$\mathcal{F} = \left\{ Q \in \mathcal{Q}_{|C_c\mu|^{-1/2}} : \sup_Q \left(\frac{1}{u_M} - M \right) \leq c\mu \right\}$$

Similarly, we pick a bump function χ_Q as before, but with $\|\nabla\chi_Q\|_{L^\infty(\mathbb{R}^d)} \leq 5|C_c\mu|^{1/2}$ and we let

$$\mathcal{H} = \text{span} \{ \chi_Q u_M : Q \in \mathcal{F} \},$$

for which $\dim(\mathcal{H}) = |\mathcal{F}| = n_c(\mu)$ holds. With this, (3.2,3.3) become

$$\begin{aligned} \int_{\mathbb{R}^d} (|\nabla(\chi_Q u_M)|^2 + V(\chi_Q u_M)^2) &= \int_Q |\nabla\chi_Q|^2 u_M^2 + \int_Q \left(\frac{1}{u_M} - M \right) (\chi_Q u_M)^2 \\ &\leq 5^2 C_c |\mu| \int_Q u_M^2 + c\mu \int_{\mathbb{R}^d} (\chi_Q u_M)^2. \end{aligned}$$

Here, we use

$$\int_Q u_M^2 \leq |Q| \sup_Q u_M^2 \leq |Q| A_M^2 \inf_{\mathbb{R}^d} u_M^2 \leq |Q| A_M^2 \inf_{Q/2} u_M^2 \leq 2^d (A_M)^2 \int_Q (\chi_Q u_M)^2$$

to conclude that

$$\int_{\mathbb{R}^d} (|\nabla(\chi_Q u_M)|^2 + V(\chi_Q u_M)^2) \leq \left(c - 2^d (5A_M)^2 C_c \right) \mu \int_{\mathbb{R}^d} (\chi_Q u_M)^2.$$

Recalling that $C_c = \frac{c-1}{2^d (5A_M)^2}$ we obtain that $\mathcal{N}^V(\mu) \geq |\mathcal{F}| = n_c(\mu)$.

(ii) With the additional assumption (2.9), we have that

$$n_c(\mu) \geq \left| \left\{ Q \in \mathcal{Q}_{(C_c|\mu|)^{-1/2}} : \inf_Q \left(\frac{1}{u_M} - M \right) \leq \frac{c\mu}{\tilde{C}_H} \right\} \right|.$$

Just as in the proof of Lemma 3.1, this is now lower bounded by the measure of the sublevel set.

(iii) Let $\mu \in \mathbb{R}$ and $\varepsilon > 0$. For every v in the form domain of $-\Delta + V$ we have by (3.1)

$$\begin{aligned} &\int_{\mathbb{R}^d} (|\nabla v(x)|^2 + (V(x) - \mu - \varepsilon)|v(x)|^2) dx \\ &= \int_{\mathbb{R}^d} \left(\left| \nabla \left(\frac{v(x)}{u_M(x)} \right) \right|^2 + \left(\frac{1}{u_M(x)} - M - \mu - \varepsilon \right) \left| \frac{v(x)}{u_M(x)} \right|^2 \right) u_M(x)^2 dx \\ &\geq \left(\inf_{\mathbb{R}^d} u_M \right)^2 \int_{\mathbb{R}^d} \left(\left| \nabla \left(\frac{v(x)}{u_M(x)} \right) \right|^2 - A_M^2 \left(\frac{1}{u_M(x)} - M - \mu - \varepsilon \right)_- \left| \frac{v(x)}{u_M(x)} \right|^2 \right) dx. \end{aligned}$$

If the integral on the RHS of (2.11) is infinite, then there is nothing to prove. Otherwise, we know from the standard CLR bound [33] that

$$N := \mathcal{N}^{-A_M^2(1/u_M - M - \mu - \varepsilon)_-} (0) \leq C_{\text{CLR}} A_M^d \int_{\mathbb{R}^d} \left(\frac{1}{u_M(x)} - M - \mu - \varepsilon \right)_-^{d/2} dx \quad (4.6)$$

and therefore, there exist linearly independent $\varphi_1, \dots, \varphi_N \in L^2(\mathbb{R}^d)$ such that for $0 \neq w \in \{f \in H^1(\mathbb{R}^d) \mid \forall j \in \{1, \dots, N\} : \langle f, \varphi_j \rangle_{L^2(\mathbb{R}^d)} = 0\}$ we have

$$\int_{\mathbb{R}^d} \left(|\nabla w(x)|^2 - A_M^2 \left(\frac{1}{u_M(x)} - M - \mu - \varepsilon \right)_- |w(x)|^2 \right) dx > 0.$$

Next, we define the codimension N space

$$\mathcal{K} = \left\{ v \in \text{dom}((-\Delta + V)^{1/2}) \mid \forall j \in \{1, \dots, N\} : \langle v, \varphi_j u_M \rangle_{L^2(\mathbb{R}^d)} = 0 \right\}.$$

For all $0 \neq v \in \mathcal{K} \cap C_c^\infty(\mathbb{R}^d)$ we have by the above that

$$\int_{\mathbb{R}^d} (|\nabla v(x)|^2 + (V(x) - \mu - \varepsilon)|v(x)|^2) dx > 0.$$

As $\mathcal{K} \cap C_c^\infty(\mathbb{R}^d)$ is dense \mathcal{K} , we conclude that

$$\int_{\mathbb{R}^d} (|\nabla v(x)|^2 + V(x)|v(x)|^2) dx > \mu \int_{\mathbb{R}^d} |v(x)|^2 dx$$

for all non-zero $v \in \mathcal{K}$. Thus, by the min-max principle we get that $\mathcal{N}^V(\mu) \leq N$. The claim follows from (4.6). \square

5 The ground state energy

In this Section we will establish the relationship between the bottom of the spectrum of $E_0 = \inf \text{spec}(-\Delta + V)$ and the infimum of the shifted landscape function. We follow [28, Lemma 5], where a similar result is proved for the torsion function in the case of nonnegative potentials.

Proof of Proposition 2.6. Let $-\Delta_L$ denote the Dirichlet Laplacian on $B(0, L)$ and let H_L be the operator $-\Delta_L + V + M$ on $L^2(B(0, L))$. Note that $\sigma_L := \inf \text{spec}(H_L) \geq \inf \text{spec}(-\Delta + V + M) > 0$. Thus

$$v_L = H_L^{-1} \mathbb{1}_{B(0, L)}$$

is well-defined. We remark that v_L is nonnegative (integration by parts yields that the Dirichlet Laplacian satisfies the Beurling-Deny condition [49, Theorem XIII.50] and hence $e^{t\Delta_L}$ is positivity-preserving for $t > 0$ which implies via functional calculus that H_L^{-1} is positivity-preserving [49, Theorem XIII.44]).

Kato-class potentials are infinitesimally form-bounded by the Dirichlet Laplacian (use [43, Theorem 4.7] to get that V is infinitesimally form-bounded by $-\Delta$ on \mathbb{R}^d , which implies that the same holds true on the ball) and as the Dirichlet Laplacian has compact resolvent ([49, Theorem XIII.73 and its corollary]), so does H_L (by [49, Theorem XIII.68]). In particular, σ_L is an eigenvalue. Let φ_L be an eigenfunction of H_L with eigenvalue σ_L . We can choose $\varphi_L > 0$ (see [50, Theorem 11.8]). Furthermore, as φ_L is square-integrable and $B(0, L)$ has finite Lebesgue measure, we get that φ_L is integrable too. Thus, we get

$$\int_{B(0,L)} \varphi_L = \int_{B(0,L)} \varphi_L (H_L v_L) = \sigma_L \int_{B(0,L)} \varphi_L v_L \leq \sigma_L \|v_L\|_{L^\infty(\mathbb{R}^d)} \int_{B(0,L)} \varphi_L.$$

This implies

$$1 \leq \sigma_L \|v_L\|_{L^\infty(\mathbb{R}^d)}.$$

Note that $(-\Delta + V + M)(u_M - v_L) = 0$ on $B(0, L)$ and $u_M - v_L = u > 0$ on $\partial B(0, L)$. Thus, by the weak maximum principle (see [53, Theorem 8.1]) we have $u_M > v_L$ on $B(0, L)$ and therefore,

$$1 \leq \sigma_L \|u_M\|_{L^\infty(\mathbb{R}^d)}. \quad (5.1)$$

Note that $\text{dom}(H_L^{1/2}) \subseteq \text{dom}(H_K^{1/2}) \subseteq \text{dom}((-\Delta + V + M)^{1/2})$ for all $K \geq L \geq 0$. Thus, $L \mapsto \sigma_L$ is a decreasing function and so

$$\lim_{L \rightarrow \infty} \sigma_L \geq \inf \text{spec}(-\Delta + V + M).$$

Since $C_c^\infty(\mathbb{R}^d)$ is dense in the form domain of $-\Delta + V + M$ (see [54, Theorem 8.2.1]), there is a sequence $(\xi_n)_{n \in \mathbb{N}} \subseteq C_c^\infty(\mathbb{R}^d)$, normalized in $L^2(\mathbb{R}^d)$, such that

$$\lim_{n \rightarrow \infty} \langle \xi_n, H \xi_n \rangle_{L^2(\mathbb{R}^d)} = \inf \text{spec}(-\Delta + V + M).$$

Now, for every n there exists L_n such that $\xi_n \in \text{dom}(H_{L_n}^{1/2})$ and so $\langle \xi_n, H \xi_n \rangle_{L^2(\mathbb{R}^d)} \geq \sigma_{L_n}$. It follows that

$$\lim_{L \rightarrow \infty} \sigma_L = \inf \text{spec}(-\Delta + V + M).$$

Taking the limit $L \rightarrow \infty$ in (5.1) yields (2.14). \square

We note that for nonnegative potentials, one can use the same proof as in [13, Theorem 1], respectively [28, Lemma 5] to get the upper bound

$$E_0 \leq (4 + d \log(2)) \inf_{\mathbb{R}^d} \frac{1}{u_M} - M.$$

Under additional assumption on the heat kernel of $-\Delta + V$ the dimensional constant can be further improved [19, Theorem 1.5].

The difference between the bottom of the spectrum of $-\Delta + V$ and the infimum of $1/u_M - M$ may a priori be large. However, the bound just proven allows us to set up the iteration procedure described in Corollary 2.7 which we now show to converges to the bottom of the spectrum.

Proof of Corollary 2.7. By (2.14) we have that the function

$$F : (-E_0, \infty) \rightarrow (-E_0, \infty)$$

$$M \mapsto M - \inf_{\mathbb{R}^d} \frac{1}{u_M}$$

is well defined. Furthermore, u_M is positive and bounded above and so

$$F(M) = M - \inf_{\mathbb{R}^d} \frac{1}{u_M} < M.$$

This readily implies that the sequence $\{M_n\}_n$ is convergent with $M_\infty = \lim_{n \rightarrow \infty} M_n \geq -E_0$. We claim that F is continuous, which implies that $F(M_\infty) = M_\infty$. If $M_\infty > -E_0$, then it is in the domain of F and so $F(M_\infty) < M_\infty$, which is a contradiction. The continuity of F follows from $u_{M+\delta} \leq u_M$ for all $\delta \geq 0$ (see [52, Lemma 2.1]) and the fact that $\lim_{\delta \rightarrow 0^+} u_{M+\delta}(x) = u_M(x)$ for all $x \in \mathbb{R}^d$ (using [52, Theorem 1.3] and the continuity of the shifted landscape functions).

Finally,

$$-E_0 < M_n - \inf_{\mathbb{R}^d} \frac{1}{u_{M_n}} < M_n$$

for all n . Since $M_n \rightarrow -E_0$, we immediately conclude that $\lim_{n \rightarrow \infty} \inf_{\mathbb{R}^d} \frac{1}{u_{M_n}} = 0$, which implies (2.15). \square

6 Examples

Since the constants that appear in the previous sections all depend on either Harnack constants C_{HM} or A_M , it is a priori unclear how sharp the estimates in terms of the landscape function really are. In this section, we provide a partial answer to that question by considering special cases. First of all, we show that for radially symmetric potentials of the form $-|v|^{-\rho}$ (in particular the Coulomb potential) whose eigenvalues accumulate at the bottom of the essential spectrum $\mu = 0$, the sharp constant in the bound (2.11) can be computed so that it becomes asymptotically exact. Secondly, we turn to the other end of the discrete spectrum and analyze the first few eigenvalues appearing in case of the spherical well potential $-\varepsilon \mathbb{1}_{\mathbb{R}^d \setminus B(0, \delta)}$ for $\varepsilon \ll 1$: The landscape function is explicit and its minimum approximates the ground state energy to order ε .

6.1 Asymptotics at the bottom of the essential spectrum

The asymptotic for power law potentials is known and we first recall a general result [55] in $d \geq 3$; the result extends to lower dimensions with appropriate assumptions. For other variants see also [49, Theorem XIII.82] or (with an oscillatory prefactor) [56, Theorem 3.1, Theorem 3.2].

Theorem 6.1. *Let $d \geq 3$. Let $1 < R < \infty$ and $V : \mathbb{R}^d \rightarrow \mathbb{R}$ such that V is C^1 for $|x| \geq R$ with $\lim_{|x| \rightarrow \infty} V(x) = 0$. There exists $q : \mathbb{R} \rightarrow (-\infty, 0]$ such that*

$$V(x) = q(|x|) + O(|x|^{-2}) \quad \text{as } |x| \rightarrow \infty. \quad (6.1)$$

Moreover, q is C^5 for $r \geq R$ and

$$M_1 r^{-1} \leq |q^{(k)}(r)/q^{(k-1)}(r)| \leq M_2 r^{-1} \quad (k \leq 5) \quad (6.2)$$

for all $r \geq R$ and for some $0 < M_1, M_2 < \infty$. Finally, we assume that there exists $\alpha > 0$ such that $d + \alpha \geq 2$ and such that

$$\int_{|x| \leq R} |V(x)|^{(d+\alpha)/2} dx < \infty. \quad (6.3)$$

Then, as $\mu \rightarrow 0^-$

$$\mathcal{N}^V(\mu) = (1 + o(1)) \left[(2\sqrt{\pi})^d \Gamma\left(\frac{d}{2} + 1\right) \right]^{-1} \int_{\mathbb{R}^d} (\mu - V(x))_+^{d/2} dx. \quad (6.4)$$

Proof of Proposition 2.8. Our goal is to establish pointwise estimates for u_M that match the assumptions (6.1,6.2). In particular, we will construct a radial function q such that $|u_M(x) - q(|x|)| \leq C_M/(1 + |x|^2)$.

By standard elliptic theory, we know that $u_M \in C^\infty(\mathbb{R}^d \setminus \{0\})$. The strategy is to compute successive approximations of u_M in terms of powers of the potential. We first note that

$$(-\Delta + V + M)(u_M - \frac{1}{M} + \frac{V}{M^2}) = -\frac{\Delta V}{M^2} + \frac{V^2}{M^2},$$

where ΔV decays fast, while V^2 does not. This formula suggests the correct expansion away from the origin where the potential is singular. Let $N \in \mathbb{N}$ be such that $N\rho < 2 \leq (N+1)\rho$. Then

$$(-\Delta + V + M) \left(u_M - \sum_{j=0}^N \frac{(-V)^j}{M^{j+1}} \right) = -\sum_{j=0}^N \Delta \left(\frac{(-V)^j}{M^{j+1}} \right) + (-1)^{N+1} \frac{V^{N+1}}{M^{N+1}}, \quad (6.5)$$

since $(V + M) \sum_{j=0}^N \frac{(-V)^j}{M^{j+1}}$ is telescopic. By assumption, the r.h.s. decays as $O(1/|x|^2)$. Although the resolvent is a bounded operator from $L^\infty(\mathbb{R}^d)$ to itself, this is not quite sufficient to conclude, both because of the singularity at the origin and the insufficient decay at infinity. We shall therefore introduce cutoffs in neighbourhoods of the origin and of infinity.

We choose a radial function $\chi \in C_c^\infty(\mathbb{R}^d)$ with $0 \leq \chi \leq 1$, $\chi(x) \equiv 1$ for $1/2 \leq |x| \leq 3/2$ and $\chi(x) \equiv 0$ for $|x| \in [0, 1/4] \cup [2, \infty)$. Let $x_0 \in \mathbb{R}^d \setminus \{0\}$ and let

$$\tilde{\chi}(x) = \chi(|x_0|^{-1}x).$$

With this, we define

$$\widetilde{u}_M = (-\Delta + V\tilde{\chi} + M)^{-1}\tilde{\chi}.$$

Let moreover

$$u_{M,L} = (-\Delta + V + M)^{-1}\mathbb{1}_{B(0,L)}.$$

Now,

$$(-\Delta + V\tilde{\chi} + M)\left(\widetilde{u}_M - \sum_{j=0}^N \frac{(-V)^j \tilde{\chi}^j}{M^{j+1}}\right) = -\sum_{j=1}^N \Delta\left(\frac{(-V)^j \tilde{\chi}^j}{M^{j+1}}\right) - \frac{(-V)^{N+1} \tilde{\chi}^{N+1}}{M^{N+1}}, \quad (6.6)$$

which readily implies that

$$\left\|\widetilde{u}_M - \sum_{j=0}^N \frac{(-V)^j \tilde{\chi}^j}{M^{j+1}}\right\|_{L^\infty(\mathbb{R}^d)} \leq \frac{C}{|x_0|^2}, \quad (6.7)$$

where C only depend on $\rho, d, M, \|\chi\|_{C^2(\mathbb{R}^d)}$ and $\|(-\Delta + V + M)^{-1}\|_{\infty, \infty}$, but not on x_0 . We now need to estimate the difference $u_{M,L}(x_0) - \widetilde{u}_M(x_0)$. We have

$$(-\Delta + V\tilde{\chi} + M)(u_{M,L} - \widetilde{u}_M) = V(\tilde{\chi} - 1)u_{M,L} + \mathbb{1}_{B(0,L)} - \tilde{\chi} \quad (6.8)$$

which is nonnegative whenever $L \geq 2|x_0|$. Since $(-\Delta + V\tilde{\chi} + M)^{-1}$ is positivity-preserving, this implies that $u_{M,L} - \widetilde{u}_M \geq 0$ for $L \geq 2|x_0|$. Furthermore, if additionally $|x_0| \geq 2$, then we have

$$(-\Delta + V\tilde{\chi} + M)(u_{M,L} - \widetilde{u}_M) = 0$$

on the annulus $B(0, |x_0| + 1) \setminus B(0, |x_0| - 1)$. Note that $u_{M,L} - \widetilde{u}_M$ is radial and, by the weak maximum principle, monotone in the radius and so

$$|u_{M,L}(x_0) - \widetilde{u}_M(x_0)| \leq |B(0, 1/2)|^{-1/2} \|u_{M,L} - \widetilde{u}_M\|_{L^2(B(x_0, 1))}. \quad (6.9)$$

Hence, we are left to obtain a local L^2 estimate in a neighbourhood of x_0 . For this we will use yet another cut-off function. Namely, we choose $\Xi \in C_c^\infty(\mathbb{R})$ such that $\frac{1}{e^2} \leq \Xi(r) \leq 1$ for $|r| \leq 2$, $\Xi(r) = 1$ for $|r| \leq 1$ and $\Xi(r) = e^{-r}$ for $r > 2$. Then we define

$$\Xi_\varepsilon(x) = \Xi(\varepsilon(|x - x_0|)),$$

where we will choose $0 < \varepsilon \leq 1$ later to be sufficiently small. We compute

$$\nabla \Xi_\varepsilon(x) = \Xi'(\varepsilon|x - x_0|)\varepsilon \frac{x - x_0}{|x - x_0|}$$

and

$$\Delta \Xi_\varepsilon(x) = \Xi''(\varepsilon|x - x_0|)\varepsilon^2 + \varepsilon \frac{d-1}{|x - x_0|} \Xi'(\varepsilon|x - x_0|).$$

Thus, for $|x - x_0| \geq 2/\varepsilon$ we have

$$|\nabla \Xi_\varepsilon(x)| + |\Delta \Xi_\varepsilon(x)| \leq \varepsilon \left[1 + \varepsilon \left(1 + \frac{d-1}{2} \right) \right] \Xi_\varepsilon(x).$$

As $1/e^2 \leq \Xi(r) \leq 1$ for $|r| \leq 2$, we get for all $x \in \mathbb{R}^d$

$$|\nabla \Xi_\varepsilon(x)| + |\Delta \Xi_\varepsilon(x)| \leq \varepsilon \left[e^2 (\|\Xi'\|_{L^\infty([0,2])} + \|\Xi''\|_{L^\infty([0,2])}) + \left(2 + \frac{d-1}{2} \right) \right] \Xi_\varepsilon(x) = C\varepsilon \Xi_\varepsilon(x), \quad (6.10)$$

with C independent of ε, x_0, x .

With these estimates in hand, we go back to (6.8). We will write $\|\cdot\|_{p,q}$ for the operator norm $\|\cdot\|_{\mathcal{L}(L^p(\mathbb{R}^d), L^q(\mathbb{R}^d))}$. Furthermore, we will repeatedly use the fact (see [52, Lemma 2.1]) that for all $V \leq W \leq 0$ and all $f \in L^2(\mathbb{R}^d)$,

$$(-\Delta + W + M)^{-1}|f| \leq (-\Delta + V + M)^{-1}|f|,$$

where the inequality has to be understood pointwise almost everywhere. This implies in particular the operator monotonicity of $V \mapsto (-\Delta + V + M)^{-1}$ in $L^2(\mathbb{R}^d)$. In turn, the same holds by density for $\|(-\Delta + V + M)^{-1}\|_{1,2}$.

Commuting the multiplication operator Ξ_ε through the resolvent yields

$$\begin{aligned} \Xi_\varepsilon(u_{M,L} - \widetilde{u}_M) &= (-\Delta + V\tilde{\chi} + M)^{-1} \Xi_\varepsilon(V(\tilde{\chi} - 1)u_{M,L} + \mathbb{1}_{B(0,L)} - \tilde{\chi}) \\ &\quad + (-\Delta + V\tilde{\chi} + M)^{-1} [-2\nabla \cdot (\nabla \Xi_\varepsilon) + (\Delta \Xi_\varepsilon)] (u_{M,L} - \widetilde{u}_M). \end{aligned} \quad (6.11)$$

Using (6.10) we can estimate the last term in L^2 by

$$\begin{aligned} &\|(-\Delta + V\tilde{\chi} + M)^{-1} [-2\nabla \cdot (\nabla \Xi_\varepsilon) + (\Delta \Xi_\varepsilon)] (u_{M,L} - \widetilde{u}_M)\|_{L^2(\mathbb{R}^d)} \\ &\leq [2\|(-\Delta + V\tilde{\chi} + M)^{-1}\nabla\|_{2,2} + \|(-\Delta + V\tilde{\chi} + M)^{-1}\|_{2,2}] C\varepsilon \|\Xi_\varepsilon(u_{M,L} - \widetilde{u}_M)\|_{L^2(\mathbb{R}^d)}. \end{aligned} \quad (6.12)$$

Since $V \leq V\tilde{\chi} \leq 0$, and with the fact that V is form bounded by $-\Delta$ with relative bound zero, we conclude that

$$\|(-\Delta + V\tilde{\chi} + M)^{-1}\nabla\|_{2,2} \leq \|(-\Delta + V + M)^{-1/2}\|_{2,2} \|(-\Delta + V + M)^{-1/2}\nabla\|_{2,2} < \infty.$$

Hence, the term of (6.12) in $[\dots]$ is bounded and there is $0 < \varepsilon \leq 1$ such that

$$\|(-\Delta + V\tilde{\chi} + M)^{-1} [-2\nabla \cdot (\nabla \Xi_\varepsilon) + (\Delta \Xi_\varepsilon)] (u_{M,L} - \widetilde{u}_M)\|_{L^2(\mathbb{R}^d)} \leq \frac{1}{2} \|\Xi_\varepsilon(u_{M,L} - \widetilde{u}_M)\|_{L^2(\mathbb{R}^d)}.$$

With this, (6.11) yields

$$\|\Xi_\varepsilon(u_{M,L} - \widetilde{u}_M)\|_{L^2(\mathbb{R}^d)} \leq 2\|(-\Delta + V\tilde{\chi} + M)^{-1} \Xi_\varepsilon(V(\tilde{\chi} - 1)u_{M,L} + \mathbb{1}_{B(0,L)} - \tilde{\chi})\|_{L^2(\mathbb{R}^d)}$$

We now split $V(\tilde{\chi} - 1)u_{M,L} = V(\tilde{\chi} - 1)(\mathbb{1}_{\mathbb{R}^d \setminus B(0,1)} + \mathbb{1}_{B(0,1)})u_{M,L}$. Using that

$$|V(\tilde{\chi} - 1)\mathbb{1}_{\mathbb{R}^d \setminus B(0,1)}u_{M,L} + \mathbb{1}_{B(0,L)} - \tilde{\chi}| \leq \mathbb{1}_{\mathbb{R}^d \setminus B(x_0, |x_0|/2)},$$

the definition of $\tilde{\chi}$ and the fact that $(-\Delta + V + M)^{-1}$ is bounded from $L^1(\mathbb{R}^d)$ to $L^2(\mathbb{R}^d)$ (see [47, Theorem B.2.2.]) we conclude that

$$\begin{aligned} \|\Xi_\varepsilon(u_{M,L} - \tilde{u}_M)\|_{L^2(\mathbb{R}^d)} &\leq 2\|(-\Delta + V + M)^{-1}\|_{2,2}\|\Xi_\varepsilon\mathbb{1}_{\mathbb{R}^d \setminus B(x_0, |x_0|/2)}\|_{L^2(\mathbb{R}^d)} \\ &\quad + 2\|(-\Delta + V + M)^{-1}\|_{1,2}\|\Xi_\varepsilon u_{M,L}\|_{L^\infty(B(0,1))}\|V\|_{L^1(B(0,1))} \\ &\leq Ce^{-\varepsilon|x_0|/4}, \end{aligned}$$

because of our choice of Ξ_ε , where C does not depend on x_0 or L . Using (6.9), the fact that $1/e^2 \leq \Xi_\varepsilon(x) \leq 1$ for $x \in B(x_0, 1)$, we finally obtain for $|x_0| \geq 4/\varepsilon$

$$|u_{M,L}(x_0) - \tilde{u}(x_0)| \leq Ce^{-\varepsilon|x_0|/4},$$

where ε, C do not depend on L or x_0 (but on M). Combining this with (6.7) and noting that $\tilde{\chi} = 1$ in a neighbourhood of x_0 we get for $|x_0|$ sufficiently large (independent of L)

$$|u_{M,L}(x_0) - \sum_{j=0}^N \frac{(-V(x_0))^j}{M^{j+1}}| \leq \frac{C}{|x_0|^2},$$

with C independent of L, x_0 .

As $u_{M,L}$ converges pointwise to u_M for $L \rightarrow \infty$, we conclude that

$$|u_M(x) - \sum_{j=0}^N \frac{(-V(x))^j}{M^{j+1}}| \leq \frac{C}{|x|^2},$$

for all $|x|$ sufficiently large. Recalling now that $-V(x) = |x|^{-\rho}$, this readily implies that there exists a polynomial P with real coefficients of degree at most N and $P(0) = 0 = P'(0)$ such that

$$\frac{1}{u_M(x)} - M + V(x) - P(|x|^{-\rho}) = O(|x|^{-2}) \quad (6.13)$$

for $|x| \rightarrow \infty$. Moreover, the boundedness of $1/u_M$ tells us that $\frac{1}{u_M(x)} - M$ satisfies (6.3).

Altogether, we obtain first that

$$\mathcal{N}^V(\mu) = (1 + o(1)) \left[(2\sqrt{\pi})^d \Gamma\left(\frac{d}{2} + 1\right) \right]^{-1} \int_{\mathbb{R}^d} (\mu - V(x))_+^{d/2} dx$$

as $\mu \rightarrow 0^-$, since V clearly satisfies the assumptions of Theorem 6.1. By (6.13), this further implies that

$$\mathcal{N}^V(\mu) = (1 + o(1)) \left[(2\sqrt{\pi})^d \Gamma\left(\frac{d}{2} + 1\right) \right]^{-1} \int_{\mathbb{R}^d} \left(\mu - \left(\frac{1}{u_M(x)} - M \right) \right)_+^{d/2} dx.$$

Finally, (6.13) also implies that Theorem 6.1 applies to $\frac{1}{u_M} - M$ and therefore that

$$\mathcal{N}^V(\mu) = (1 + o(1)) \mathcal{N}^{1/u_M - M}(\mu),$$

which concludes the proof. \square

In the case of the Coulomb potential, the proof above yields that the effective potential asymptotically equals the original potential itself, namely

$$\frac{1}{u_M(x)} - M = -\frac{1}{|x|} + O(|x|^{-2}).$$

6.2 Ground state energy for the square well

We now turn to $V(x) = -\varepsilon \mathbb{1}_{B(0,\delta)}$ for $\varepsilon, \delta > 0$ and focus on the case $d = 1$. The ground state energy of $-\Delta + \varepsilon \mathbb{1}_{\mathbb{R} \setminus (-\delta, \delta)}$ is, for ε small enough, given by the smallest positive solution of the transcendental equation

$$\sqrt{\varepsilon - x} = \sqrt{x} \tan(\sqrt{x}\delta).$$

This can be rewritten as

$$\varepsilon = F(x), \quad \text{where} \quad F(x) = x(1 + \tan(\sqrt{x}\delta))^2.$$

The function F has an analytic extension at the origin such that $F(x) = x + x^2\delta^2 + O(x^3)$, and hence

$$x = F^{-1}(\varepsilon) = \varepsilon - \delta^2\varepsilon^2 + O(\varepsilon^3).$$

This implies that as $\varepsilon \rightarrow 0^+$

$$\inf \text{spec}(-\Delta - \varepsilon \mathbb{1}_{(-\delta, \delta)}) = \inf \text{spec}(-\Delta + \varepsilon \mathbb{1}_{\mathbb{R} \setminus (-\delta, \delta)}) - \varepsilon = -\delta^2\varepsilon^2 + O(\varepsilon^3).$$

Next we compute the shifted landscape function, which is an even solution of

$$-f''(x) + cf(x) = 1$$

where the constant c depends on M and whether x is inside or outside of the well.

We start with the case $M > \varepsilon$, in which case

$$u_M(x) = \begin{cases} \frac{1}{M-\varepsilon} + a_2 \cosh(\sqrt{M-\varepsilon}x), & |x| < \delta, \\ \frac{1}{M} + b_1 e^{-\sqrt{M}|x|} + b_2 \cosh(\sqrt{M}x), & |x| > \delta. \end{cases}$$

As u_M is bounded, we get $b_2 = 0$. The other constants are set by the of u_M differentiability at $|x| = \delta$:

$$a_2 = \frac{\frac{1}{M} - \frac{1}{M-\varepsilon}}{\cosh(\sqrt{M-\varepsilon}\delta) + \frac{\sqrt{M-\varepsilon}}{\sqrt{M}} \sinh(\sqrt{M-\varepsilon}\delta)}$$

$$b_1 = -\frac{\sqrt{M-\varepsilon}}{\sqrt{M}} \sinh(\sqrt{M-\varepsilon}\delta) e^{\sqrt{M}\delta} a_2.$$

The maximum of u_M is attained at the origin and so

$$\inf_{\mathbb{R}} \frac{1}{u_M} - M = \varepsilon \frac{M-\varepsilon}{M} \frac{1}{\cosh(\sqrt{M-\varepsilon}\delta) + \frac{\sqrt{M-\varepsilon}}{\sqrt{M}} \sinh(\sqrt{M-\varepsilon}\delta)} - \varepsilon + o(\varepsilon).$$

Hence, for any $M > \epsilon$ — in particular arbitrarily large! — the first approximation of the ground state energy given by Corollary 2.7 is already of order ϵ .

For $M = \epsilon$, the computation in [9, Section 6] yields

$$\inf_{\mathbb{R}} \frac{1}{u_\epsilon} - \epsilon = \frac{-\delta\epsilon^{3/2} - \frac{\delta^2\epsilon^2}{2}}{1 + \delta\sqrt{\epsilon} + \frac{\delta^2\epsilon}{2}}.$$

Hence, for small ϵ , this yields an approximation of order $\epsilon^{3/2}$.

Finally if $M < \epsilon$, then

$$u_M(x) = \begin{cases} \frac{1}{M-\epsilon} + a \cos(\sqrt{\epsilon - M}x), & |x| < \delta, \\ \frac{1}{M} + be^{-\sqrt{M}|x|}, & |x| > \delta. \end{cases}$$

Differentiability yields now

$$a = \frac{\frac{1}{M} - \frac{1}{M-\epsilon}}{\cos(\sqrt{\epsilon - M}\delta) - \frac{\sqrt{\epsilon - M}}{\sqrt{M}} \sin(\sqrt{\epsilon - M}\delta)} \quad (6.14)$$

$$b = \frac{\sqrt{\epsilon - M}}{\sqrt{M}} \sin(\sqrt{\epsilon - M}\delta) e^{\sqrt{M}\delta} a \quad (6.15)$$

This allows us to exhibit explicitly the blowup of (2.15) in Corollary 2.7. For this we first note that $\epsilon + E_0$ is the ground state energy of the operator $-\Delta + \epsilon \mathbb{1}_{(-\infty, -\delta) \cup (\delta, \infty)}(x)$ and hence

$$\sqrt{-E_0} = \sqrt{\epsilon + E_0} \tan(\sqrt{\epsilon + E_0}\delta).$$

It follows from this and (6.14) that $\lim_{M \rightarrow (-E_0)^+} a = \infty$ and the same for b by (6.15). Hence $\lim_{M \rightarrow (-E_0)^+} u_M(x) = \infty$, or in other words for the effective potential

$$\lim_{M \rightarrow (-E_0)^+} \left(\frac{1}{u_M(x)} - M \right) = E_0$$

for all $x \in \mathbb{R}$.

7 Upper and lower Lieb-Thirring bounds

In this section we recall the standard argument for the derivation of the Lieb-Thirring inequality from estimates on the eigenvalue counting function. First we rewrite the sum of eigenvalues in terms of an integral.

Lemma 7.1. *For any $\gamma > 0$ we have*

$$\mathrm{tr}((-\Delta + V)_-^\gamma) = \gamma \int_0^\infty \lambda^{\gamma-1} \mathcal{N}^V(-\lambda) d\lambda. \quad (7.1)$$

Proof. If $\mathcal{N}^V(-\lambda) = \infty$ for any $\lambda > 0$, then both expressions in (7.1) are infinity. On the other hand, if $\mathcal{N}^V(-\lambda) < \infty$ for all $\lambda > 0$, then the negative spectrum consists of countably many eigenvalues of finite multiplicity. We label them as $\lambda_1 \leq \lambda_2 \leq \dots < 0$. Then we can write

$$\mathrm{tr}((-\Delta + V)_-^\gamma) = \sum_j |\lambda_j|^\gamma.$$

Since \mathcal{N}^V is a piecewise constant function, we have for any $N \in \mathbb{N}$

$$\gamma \int_{|\lambda_N|}^{\infty} \lambda^{\gamma-1} \mathcal{N}^V(-\lambda) d\lambda = \sum_{j=1}^{N-1} |\lambda_j|^\gamma + (N-1) |\lambda_N|^\gamma,$$

and the remainder term vanishes if $\sum_j |\lambda_j|^\gamma < \infty$, concluding the proof. \square

With this, it is now a simple calculation to derive two-sided Lieb-Thirring bounds from the estimates (2.4).

Proof of Corollary 2.2. Lemma 7.1 and the upper bound (2.4) yield immediately that

$$\begin{aligned} \mathrm{tr}((-\Delta + V)_-^\gamma) &= \gamma \int_0^{|E_0|} \lambda^{\gamma-1} \mathcal{N}^{V+\delta+|E_0|}(-\lambda + \delta + |E_0|) d\lambda \\ &\leq C_{0,d}^{\frac{d}{2}} \gamma \int_0^{|E_0|} \lambda^{\gamma-1} (-\lambda + \delta + |E_0|)^{\frac{d}{2}} \left| \left\{ x \in \mathbb{R}^d : \frac{1}{u_M(x)} \leq C_{0,d}(-\lambda + \delta + |E_0|) \right\} \right| d\lambda. \end{aligned} \quad (7.2)$$

Now since $\gamma \geq 1$, we use that $\lambda^{\gamma-1} \leq (\lambda + \delta)^{\gamma-1}$. We split the integral in two. On $[0, \frac{|E_0|}{2}]$, we use that $|E_0| - \lambda + \delta \leq |E_0| + \delta \leq \frac{|E_0| + \delta}{\delta}(\lambda + \delta)$, while $|E_0| - \lambda + \delta \leq \frac{|E_0|}{2} + \delta \leq \lambda + \delta$ holds on $[\frac{|E_0|}{2}, |E_0|]$. Writing the condition $\frac{1}{u_M(x)} \leq C_{0,d}(-\lambda + \delta + |E_0|)$ as $\lambda + \delta \leq |E_0| + 2\delta - \frac{1}{C_{0,d}u_M(x)}$ and changing variables to $s = \lambda + \delta$ yields that if $\gamma \geq 1$, then

$$\begin{aligned} \mathrm{tr}((-\Delta + V)_-^\gamma) &\leq C_{0,d}^{\frac{d}{2}} \gamma \left(1 + \frac{|E_0|}{\delta} \right)^{\frac{d}{2}} \int_\delta^{|E_0|+\delta} s^{\gamma-1+\frac{d}{2}} \left| \left\{ x \in \mathbb{R}^d : |E_0| + 2\delta - \frac{1}{C_{0,d}u_M(x)} \geq s \right\} \right| \\ &\leq C_{0,d}^{\frac{d}{2}} \frac{\gamma}{\gamma + \frac{d}{2}} \left(1 + \frac{|E_0|}{\delta} \right)^{\frac{d}{2}} \int_{\mathbb{R}^d} \left(|E_0| + 2\delta - \frac{1}{C_{0,d}u_M(x)} \right)_+^{\frac{d}{2}+\gamma} dx \end{aligned}$$

by the layer cake representation.

Similarly, for the lower bound, we start as in (7.2) but with $c_{0,d}$ instead of $C_{0,d}$. Here, $|E_0| - \lambda + \delta \geq \frac{|E_0|}{2} + \delta > \lambda$ on $[0, \frac{|E_0|}{2}]$ while $|E_0| - \lambda + \delta \geq \delta \geq \frac{\delta}{|E_0|} \lambda$ on $[\frac{|E_0|}{2}, |E_0|]$. We conclude that for $\gamma > 0$,

$$\begin{aligned} \mathrm{tr}((-\Delta + V)_-^\gamma) &\geq \min \left\{ 1, \frac{\delta}{|E_0|} \right\}^{\frac{d}{2}} c_{0,d}^{d/2} \gamma \int_0^{E_0} \lambda^{\frac{d}{2}+\gamma-1} \left| \left\{ x \in \mathbb{R}^d : \lambda \leq |E_0| + \delta - \frac{1}{c_{0,d}u_M(x)} \right\} \right| d\lambda \\ &= \min \left\{ 1, \frac{\delta}{|E_0|} \right\}^{\frac{d}{2}} c_{0,d}^{d/2} \frac{\gamma}{\gamma + \frac{d}{2}} \int_{\mathbb{R}^d} \left(|E_0| + \delta - \frac{1}{c_{0,d}u_M(x)} \right)_-^{\frac{d}{2}+\gamma} dx. \end{aligned}$$

For the last equality we used the bound (2.4) to conclude that the measure vanishes for all $\lambda > |E_0|$ since $\mathcal{N}^V(-\lambda) = 0$ for those λ . \square

Proof of Corollary 2.5. We follow the same strategy as above, using (2.10,2.11) instead of (2.4). For the lower bound, we immediately have

$$\begin{aligned} \operatorname{tr}((-\Delta + V)_-^\gamma) &= \gamma \int_0^{|E_0|} \lambda^{\gamma-1} \mathcal{N}^V(-\lambda) d\lambda \\ &\geq C_c^{\frac{d}{2}} \gamma \int_0^{|E_0|} \lambda^{\gamma-1+\frac{d}{2}} \left| \left\{ x \in \mathbb{R}^d : \lambda \leq \frac{\tilde{C}_H}{c} \left(M - \frac{1}{u_M(x)} \right) \right\} \right| \\ &= C_c^{\frac{d}{2}} \frac{\gamma}{\gamma + \frac{d}{2}} \left(\frac{\tilde{C}_H}{c} \right)^{\gamma + \frac{d}{2}} \int_{\mathbb{R}^d} \left(\frac{1}{u_M(x)} - M \right)_-^{\gamma + \frac{d}{2}} dx. \end{aligned}$$

For the upper bound, we follow the proof of Theorem 2.4(iii) to get

$$\mathcal{N}^V(\mu) \leq \mathcal{N}^{-A_M^2(1/u_M - M)_-}(\mu + \varepsilon).$$

With this,

$$\operatorname{tr}((-\Delta + V)_-^\gamma) \leq \gamma \int_0^{|E_0|} \lambda^{\gamma-1} \mathcal{N}^{-A_M^2(1/u_M - M)_-}(-\lambda + \varepsilon) d\lambda$$

Since the counting function is càdlàg by definition (2.1), we let $\varepsilon \rightarrow 0^+$ and conclude that

$$\operatorname{tr}((-\Delta + V)_-^\gamma) \leq \operatorname{tr}((-\Delta - A_M^2(1/u_M - M)_-)_-^\gamma)$$

and we use the standard Lieb-Thirring inequality to conclude. \square

Acknowledgements. The three authors were supported by NSERC of Canada. S.B. would like to thank S. Mayboroda and D. Arnold for inspiring discussions.

References

- [1] E.H. Lieb and W. Thirring. Inequalities for the moments of the eigenvalues of Schrödinger equations and their relations to Sobolev inequalities. *Studies in Mathematical Physics: Essays in Honor of Valentine Bergmann*, pages 269–303, 1976.
- [2] S.M. Birman and M.Z. Solomyak. Asymptotic behavior of the spectrum of differential equations. *Journal of Soviet Mathematics*, 12:247–283, 1979.
- [3] R.L. Frank, A. Laptev, and T. Weidl. *Schrödinger operators: Eigenvalues and Lieb–Thirring inequalities*, volume 200. Cambridge University Press, 2022.
- [4] U.W. Schmincke. On Schrödinger’s factorization method for Sturm-Liouville operators. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics*, 80(1-2):67–84, 1978.

- [5] A. Grigor'yan, Y. Netrusov, and S.-T. Yau. Eigenvalues of elliptic operators and geometric applications. *Surveys in differential geometry*, 9(1):147–217, 2004.
- [6] D. Damanik and C. Remling. Schrödinger operators with many bound states. *Duke Mathematical Journal*, 136(1):51 – 80, 2007.
- [7] O. Safronov and B. Vainberg. Estimates for negative eigenvalues of a random Schrödinger operator. *Proceedings of the American Mathematical Society*, 136(11):3921–3929, 2008.
- [8] S. Molchanov and B. Vainberg. Bargmann type estimates of the counting function for general Schrödinger operators. *Journal of Mathematical Sciences*, 184:457–508, 2012.
- [9] S. Bachmann, R. Froese, and S. Schraven. Counting eigenvalues of Schrödinger operators using the landscape function. *Journal of Spectral Theory*, 13(4):1445–1472, 2023.
- [10] B. Poggi. Applications of the landscape function for Schrödinger operators with singular potentials and irregular magnetic fields. *arXiv preprint arXiv:2107.14103*, 2021.
- [11] L.E. Payne. Bounds for solutions of a class of quasilinear elliptic boundary value problems in terms of the torsion function. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics*, 88(3-4):251–265, 1981.
- [12] R. Sperb. Bounds for the first eigenvalue of the elastically supported membrane on convex domains. *Zeitschrift für angewandte Mathematik und Physik ZAMP*, 54:879–902, 2003.
- [13] M. van den Berg and T. Carroll. Hardy inequality and L^p estimates for the torsion function. *Bulletin of the London Mathematical Society*, 41(6):980–986, 2009.
- [14] T. Giorgi and R.G. Smits. Principal eigenvalue estimates via the supremum of torsion. *Indiana University Mathematics Journal*, pages 987–1011, 2010.
- [15] M. van den Berg. Estimates for the torsion function and Sobolev constants. *Potential Analysis*, 36(4):607–616, 2012.
- [16] M. van den Berg and D. Bucur. On the torsion function with Robin or Dirichlet boundary conditions. *Journal of Functional Analysis*, 266(3):1647–1666, 2014.
- [17] A. Aghajani, A. Mosleh Tehrani, and N. Ghoussoub. Pointwise lower bounds for solutions of semilinear elliptic equations and applications. *Advanced Nonlinear Studies*, 14(4):839–856, 2014.
- [18] M. van den Berg. Spectral bounds for the torsion function. *Integral Equations and Operator Theory*, 88:387–400, 2017.

- [19] H. Vogt. L^∞ -estimates for the torsion function and L^∞ -growth of semigroups satisfying Gaussian bounds. *Potential Analysis*, 51:37–47, 2019.
- [20] M. Filoche and S. Mayboroda. Universal mechanism for Anderson and weak localization. *Proceedings of the National Academy of Sciences*, 109(37):14761–14766, 2012.
- [21] D.N. Arnold, G. David, M. Filoche, D. Jerison, and S. Mayboroda. Computing spectra without solving eigenvalue problems. *SIAM Journal on Scientific Computing*, 41(1):B69–B92, 2019.
- [22] G. David, M. Filoche, and S. Mayboroda. The landscape law for the integrated density of states. *Advances in Mathematics*, 390:107946, 2021.
- [23] P. Desforges, S. Mayboroda, S. Zhang, G. David, D.N. Arnold, W. Wang, and M. Filoche. Sharp estimates for the integrated density of states in Anderson tight-binding models. *Physical Review A*, 104(1):012207, 2021.
- [24] D. Arnold, M. Filoche, S. Mayboroda, W. Wang, and S. Zhang. The landscape law for tight binding Hamiltonians. *Communications in Mathematical Physics*, 396(3):1339–1391, 2022.
- [25] D.N. Arnold, G. David, M. Filoche, D. Jerison, and S. Mayboroda. Localization of eigenfunctions via an effective potential. *Communications in Partial Differential Equations*, 44(11):1186–1216, 2019.
- [26] M. Filoche, S. Mayboroda, and T. Tao. The effective potential of an M-matrix. *Journal of Mathematical Physics*, 62(4), 2021.
- [27] W. Wang and S. Zhang. The exponential decay of eigenfunctions for tight-binding Hamiltonians via landscape and dual landscape functions. In *Annales Henri Poincaré*, volume 22, pages 1429–1457. Springer, 2021.
- [28] M. van den Berg, D. Bucur, and T. Kappeler. On efficiency and localisation for the torsion function. *Potential Analysis*, 57(4):571–600, 2022.
- [29] D. Mugnolo. Pointwise eigenvector estimates by landscape functions: Some variations on the Filoche–Mayboroda–van den Berg bound. *Mathematische Nachrichten*, 2023.
- [30] S. Steinerberger. Localization of quantum states and landscape functions. *Proceedings of the American Mathematical Society*, 145(7):2895–2907, 2017.
- [31] E.H. Lieb. Bounds on the eigenvalues of the Laplace and Schrödinger operators. *Bulletin of the American Mathematical Society*, 82(5):751–753, 1976.
- [32] G.V. Rozenblum. Distribution of the discrete spectrum of singular differential operators. *Izvestiya Vysshikh Uchebnykh Zavedenii. Matematika*, 1:75–86, 1976.

- [33] M. Cwikel. Weak type estimates for singular values and the number of bound states of Schrödinger operators. *Annals of Mathematics*, 106:93–100, 1977.
- [34] J.G. Hoskins, H. Quan, and S. Steinerberger. Magnetic Schrödinger operators and landscape functions. *Communications in Partial Differential Equations*, 49(1-2):1–14, 2024.
- [35] J. Lu, C. Murphey, and S. Steinerberger. Fast localization of eigenfunctions via smoothed potentials. *Journal of Scientific Computing*, 90:1–18, 2022.
- [36] M. Otelbaev. Bounds for eigenvalues of singular differential operators. *Mathematical notes of the Academy of Sciences of the USSR*, 20(6):1038–1042, 1976.
- [37] M. Otelbaev. Imbedding theorems for spaces with a weight and their application to the study of the spectrum of a Schrödinger operator. *Trudy Matematicheskogo Instituta imeni V.A. Steklova, (Proceedings of the Steklov Mathematical Institute)*, 150:265–305, 1979.
- [38] Z. Shen. Eigenvalue asymptotics and exponential decay of eigenfunctions for Schrödinger operators with magnetic fields. *Transactions of the American Mathematical Society*, 348(11):4465–4488, 1996.
- [39] Z. Shen. On bounds of $N(\lambda)$ for a magnetic Schrödinger operator. *Duke Mathematical Journal*, 94(3):479–507, 1998.
- [40] L. Bugliaro, J. Fröhlich, G.M. Graf, J. Stubbe, and C. Fefferman. A Lieb-Thirring bound for a magnetic Pauli Hamiltonian. *Communications in Mathematical Physics*, 187:567–582, 1997.
- [41] Z. Shen. On moments of negative eigenvalues for the Pauli operator. *Journal of Differential Equations*, 149(2):292–327, 1998.
- [42] T.D. Do and L.X. Truong. Spectral asymptotics for generalized Schrödinger operators. *Annales Fennici Mathematici*, 48(2):703–727, 2023.
- [43] M. Aizenman and B. Simon. Brownian motion and Harnack inequality for Schrödinger operators. *Communications on Pure and Applied Mathematics*, 35(2):209–273, 1982.
- [44] F. Chiarenza, E. Fabes, and N. Garofalo. Harnack’s inequality for Schrödinger operators and the continuity of solutions. *Proceedings of the American Mathematical Society*, 98(3):415–425, 1986.
- [45] L. Read. Lieb—Thirring type bounds for perturbed Schrödinger operators with single-well potentials. *Journal of Mathematical Physics*, 64(6), 2023.
- [46] Q. Han and F. Lin. *Elliptic partial differential equations*, volume 1. American Mathematical Soc., 2011.

- [47] B. Simon. Schrödinger semigroups. *Bulletin of the American Mathematical Society*, 7(3):447–526, 1982.
- [48] G. David, A. Gloria, and S. Mayboroda. The landscape function on \mathbb{R}^d . *arXiv preprint arXiv:2307.11182*, 2023.
- [49] M. Reed and B. Simon. *IV: Analysis of Operators*, volume 4. Elsevier, 1978.
- [50] E.H. Lieb and M. Loss. *Analysis*, volume 14. American Mathematical Society, 2001.
- [51] L.E. Payne and H.F. Weinberger. An optimal Poincaré inequality for convex domains. *Archive for Rational Mechanics and Analysis*, 5(1):286–292, 1960.
- [52] M.A. Perelmuter. Positivity preserving operators and one criterion of essential self-adjointness. *Journal of Mathematical Analysis and Applications*, 82(2):406–419, 1981.
- [53] D. Gilbarg and N.S. Trudinger. *Elliptic partial differential equations of second order*, volume 224. Springer, 1977.
- [54] E.B. Davies. *Spectral theory and differential operators*, volume 42. Cambridge University Press, 1995.
- [55] F.H. Brownell and C.W. Clark. Asymptotic distribution of the eigenvalues of the lower part of the Schrödinger operator spectrum. *Journal of Mathematics and Mechanics*, pages 31–70, 1961.
- [56] G. Raikov. Discrete spectrum for Schrödinger operators with oscillating decaying potentials. *Journal of Mathematical Analysis and Applications*, 438(2):551–564, 2016.