TWISTS OF SUPERCONFORMAL ALGEBRAS

CHRIS ELLIOTT, OWEN GWILLIAM, AND MATTEO LOTITO

ABSTRACT. We develop the theory of "conformal twists" of superconformal field theories in dimension 3 to 6, extending the well-known analysis of twists for supersymmetric theories. The conformal twists describe all possible inequivalent choices of a nilpotent element in the superconformal algebra. Such twists can give rise to interesting subalgebras and protected sectors of operators, with the Donaldson–Witten topological field theory and the vertex operator algebras of 4-dimensional $\mathcal{N}=2$ SCFTs being prominent examples. We work mostly with the complexified superconformal algebras, unless explicitly stated otherwise; real forms of the superconformal algebra may have important physical implications, but we leave these subtleties to future work. To obtain mathematical precision, we explain how to extract vertex algebras and \mathbb{E}_n algebras from a twisted superconformal field theory using factorization algebras.

CONTENTS

1. Introduction	2
1.1. What Twisting Does to Superconformal Theories	3
Acknowledgements	6
2. Superconformal Algebras and Twists	6
2.1. The Conformal Case	7
2.2. The Superconformal Case	9
2.3. The Question of Classifying Twists	11
3. The Observables of a Twisted Superconformal Theory	15
3.1. Subalgebras of Q -Closed and Q -Exact Elements	16
3.2. Observables and BRST Operators	16
3.3. Structures on Observables on Affine Subspaces	18
3.4. Connections to the Physics Literature	21
4. 4d Superconformal Twists	22
4.1. Complex Nilpotence Variety for $\mathcal{N} \neq 4$	22
4.2. The $\mathcal{N} = 4$ Case	24
4.3. Complex Group Orbits	25
4.4. Closed and Exact Transformations	27
4.5. Real Group Orbits: The Case of a Chiral Twist	29
4.6. Case Study: 4d $\mathcal{N} = 2$ Chiral Supercharges	33
5. 3d Superconformal Twists	37
5.1. Complex Nilpotence Variety	37
5.2. Complex Group Orbits & Closed and Exact Transformations	38
5.3. Real Group Orbits	40
6. 5d and 6d Superconformal Twists	41

6.1.	5d Twists	41
6.2.	6d Twists	42
7.	Twisted Observables and Factorization Algebras	43
7.1.	Superconformal Prefactorization Algebras	43
7.2.	Algebras of Operator Products	44
7.3.	Obtaining Vertex Algebras or \mathbb{E}_k -Algebras	46
Refe	48	

1. INTRODUCTION

In this paper we explore the notion of supersymmetric twisting in the context of a superconformal field theory, with a focus on what can be seen directly from superconformal algebras. The questions we pursue in this paper are motivated by physics but connect with contemporary work in representation theory, and in this introduction we aim to situate our work in both subjects.

The idea of twisting a supersymmetric theory goes back to Witten [Wit88], who introduced it as a recipe for constructing topological quantum field theories, and it has provided a number of examples with rich consequences in physics and mathematics. We wish to emphasize a more abstract formulation of the twisting procedure: given a field theory with an action of a super Lie algebra g and an odd element $Q \in g$ satisfying [Q, Q] = 0, there is a deformation of the theory by the action of Q. For example, if H_Q is the Hamiltonian associated to Q, we may replace the algebra of local operators Op_{loc} by its Q-cohomology

$$Op_{loc}^{\mathcal{Q}} = H^{\bullet}(Op_{loc}, \{H_{\mathcal{Q}}, -\}),$$

where the condition [Q, Q] = 0 ensures that the differential does indeed square to zero. For us the pertinent super Lie algebra g will be a superconformal algebra (i.e., the relevant spacetime symmetries for a superconformal theory).

In the last two decades, representation theorists have analyzed the space of square-zero odd elements of super Lie algebras (see especially Jenkins [Jen22]). Of particular relevance to us is the work of Duflo and Serganova [DS05] (see also [Gor+22]), who studied *simple* super Lie algebras over the complex numbers and described the adjoint orbits of such square-zero odd elements of such algebras. Many superconformal algebras, after complexification, are simple, as we discuss in Section 2. For example, for spacetime dimensions three and six, the superconformal algebras correspond to the super Lie algebras osp(n|m), which were studied in detail by Gruson [Gru00] prior to Duflo and Serganova. In this paper we explain how these structural mathematical results interface with the role of these super Lie algebras as symmetries of superconformal field theories.

The more familiar case of supersymmetry (i.e. of a supersymmetric extension of the group of isometries rather than of conformal transformations) motivates our approach. As mentioned, Witten introduced twisting to obtain a topological quantum field theory out of a supersymmetric theory: if the stress-energy tensor $T_{\mu\nu}$ of a supersymmetric theories is exact for the action of Q,

then this stress-energy tensor vanishes in the Q-cohomology of observables ¹, and the associated twisted theory is topological. Thus, it is natural to study topological twisting in two stages:

- identify the square zero odd elements ("twists") of each super Poincaré algebra, possibly up to the adjoint action of the super Poincaré group, and
- describe the Q-cohomology of observables for supersymmetric theories, looking for those where the stress-energy tensor vanishes.

Recently, in [ES19; ESW22] the first author, together with Safronov and Williams, systematically described the topological twists of super Yang–Mills theories (formulated in the Batalin– Vilkovisky formalism) and identified the twists in super Poincaré algebras that would produce topological theories. Note that there are twists that lead to *non*-topological theories, where subtle structures emerge, such as theories that look holomorphic [Joh95; Bau11] or look topological in some directions and holomorphic in others [Kap06; KZ12; Aga+17]. In the present work our goal is to pursue analogues for *superconformal* field theories by working with the superconformal algebras that extend super Poincaré algebras.

Working in the superconformal case is not simply a matter of reproducing the calculations from earlier work in yet another example. There are substantially different features that allow us to construct interesting and novel algebras of operators upon twisting, and superconformal transformations produce fascinating relationships between different twists. An important motivation for us — demonstrating some novelties not apparent in the supersymmetric case — is the work of Beem, Lemos, Liendo, Peelaers, Rastelli and van Rees [Bee+15b], who showed how such superconformal twists lead to novel constructions of vertex algebras. Combinations of these authors wrote a series of subsequent papers that address more examples and explore their consequences [Bee+15a; BRR15; BPR17a; BR18]. In this paper we do not focus on specific theories, but rather analyze formal aspects of superconformal actions on algebras of observables and the structures that may arise upon twisting. Our paper might be seen as trying to provide some systematic explanation for the kinds of results found by Beem *et al.* In Section 3 we provide an overview with a physical flavor; in Section 7 we state precise mathematical results using the language of (pre)factorization algebras. (We discuss again the results of Beem et al in Section 4.6.)

We now turn to describing our questions and efforts in more detail.

1.1. What Twisting Does to Superconformal Theories. Let us provide a gloss for the geometric context before discussing field theories and their observables.

Let $\mathbb{R}^{p,q}$ denote the real vector space of dimension p + q = n with a pseudo-Riemannian structure of signature (p,q). Euclidean space is denoted $\mathbb{R}^{n,0}$ and Minkowski space is denoted $\mathbb{R}^{n-1,1}$ (or $\mathbb{R}^{1,n-1}$, depending on preference). There is a group $\operatorname{Conf}(p,q)$ of all conformal transformations, containing the isometries but also more. There is a natural conformal compactification $C(\mathbb{R}^{p,q})$ on which $\operatorname{Conf}(p,q)$ acts transitively, with $\mathbb{R}^{p,q}$ as a dense open submanifold, akin to the Riemann sphere compactifying the complex line. (For a fuller review, see Section 2.1.) Let $\operatorname{conf}(p,q)$ denote the real Lie algebra, and let $\operatorname{conf}(p+q,\mathbb{C})$ denote its complexification, which only depends on the total dimension, not the signature.

¹Note that such theories are sometimes called "Witten-type" topological theories, as opposed to "Schwarz-type" topological theories where $T_{\mu\nu}$ is exactly zero, as for instance in [Bir+91].

Almost tautologically, a conformal field theory lives on $C(\mathbb{R}^{p,q})$, but to compare with other theories (and for the sake of familiarity), one often restricts the theory to a dense open subset $\mathbb{R}^{p,q} \subset C(\mathbb{R}^{p,q})$, which one views as the relevant "spacetime." This choice of embedding is somewhat arbitrary, as any element g of the conformal group $\operatorname{Conf}(p,q)$ moves the "defining" $\mathbb{R}^{p,q}$ to another copy $g\mathbb{R}^{p,q}$. At first glance, a theory may look different on these two embeddings, especially when written in explicit coordinates.

A *super*conformal field theory has a super Lie group SConf(p,q|S) of symmetries, extending the action of Conf(p,q). (Here *S* just denotes data chosen to characterize the super extension.) For the most part we will focus on the super Lie algebra $\mathfrak{sconf}(p,q|S)$, which we call a *superconformal algebra*.

Let Obs denote the algebra of observables of a superconformal field theory (classical or quantum). The most natural place for a superconformal field theory to live is the conformal compactification $C(\mathbb{R}^{p,q})$, so our algebra Obs can be thought of as living over the conformal compactification. That is, for any open subset $U \subseteq C(\mathbb{R}^{p,q})$ there is a vector space Obs(U) of observables supported on the subset U. (The notion of prefactorization algebra encodes the dependence of observables on open sets.)

As a superconformal theory, the superconformal group acts equivariantly: for every $g \in$ SConf(p,q|S) and open set U, there is an isomorphism Obs $(U) \rightarrow$ Obs(gU). Similarly, the super Lie algebra $\mathfrak{sconf}(p,q|S)$ acts by derivations on every Obs(U). As the observables form a complex vector space, this action extends to an action by the complexified super Lie algebra, which we will denote $\mathfrak{sconf}(n|S_{\mathbb{C}},\mathbb{C})$ for now.

If Q is an odd element of the complexified superconformal Lie algebra such that [Q, Q] = 0, we may study the algebra of observables Obs^{Q} in the theory *twisted* by Q. This twisted algebra of observables still lives over $C(\mathbb{R}^{p,q})$, and a goal in this paper is to develop a systematic understanding of the properties of this twisted algebra determined purely by the superconformal algebra. To this end, let us describe a few natural structures determined by Q and independent of the choice of theory and ruminate on what they mean for the twisted theory.

A central fact is that not all symmetries of the superconformal theory are symmetries of a twisted theory. Instead, there is an interesting relationship between the symmetries of the superconformal theory and its twist.

Closed elements of the Lie algebra: Let

$$\mathfrak{z}_{\mathcal{O}}^{\mathbb{C}} = \{A \in \mathfrak{sconf}(n|S_{\mathbb{C}},\mathbb{C})_0 \colon [A,\mathcal{Q}] = 0\},\$$

where $\mathfrak{sconf}(n|S_{\mathbb{C}},\mathbb{C})_0$ denotes the even part of the superconformal algebra. The Lie subalgebra $\mathfrak{z}_{Q}^{\mathbb{C}}$ consists of even symmetries that commute with the supercharge Q. These symmetries remain as symmetries of the twisted theory Obs^{Q} . The action of non-closed symmetries is broken when we pass to the twist.

Closed elements of the Lie group: Let

$$Z_{\mathcal{Q}}^{\mathbb{C}} \subseteq \operatorname{SConf}(n|S_{\mathbb{C}},\mathbb{C})_{0}$$
⁴

denote the Lie group exponentiating $\mathfrak{z}_{\mathcal{Q}}^{\mathbb{C}}$, and let $Z_{\mathcal{Q}}$ denote its intersection with the real form SConf($p, q | S_{\mathbb{C}})_0$. This Lie subgroup consists of the even symmetries that continue to act on the twisted theory Obs^{\mathcal{Q}} at the group level, by genuine conformal transformations.

Example 1.1. If Z_Q contains the group of isometries of an affine subspace $\mathbb{R}^{a,b} \subseteq C(\mathbb{R}^{p,q})$ then we may study the observables in Obs^Q supported along this subspace: this algebra of restricted observables is now isometry-invariant.

• Exact elements of the Lie algebra: Let

$$\mathfrak{b}_{\mathcal{Q}}^{\mathbb{C}} = \{ A \in \mathfrak{sconf}(n|S_{\mathbb{C}},\mathbb{C})_0 \colon A = [\mathcal{Q},\mathcal{Q}'] \text{ for some } \mathcal{Q}' \}.$$

These provide even symmetries of the twisted theory that are explicitly trivialized. That is to say, the conserved current for a symmetry from $\mathfrak{b}_{\mathcal{Q}}^{\mathbb{C}}$ is exact in $Obs^{\mathcal{Q}}$, so it vanishes in cohomology.

Exact elements of the Lie group: Let

$$B_{\mathcal{Q}}^{\mathbb{C}} \subseteq \mathrm{SConf}(n|S_{\mathbb{C}},\mathbb{C})_0$$

denote the Lie group exponentiating $\mathfrak{b}_{Q}^{\mathbb{C}}$, and let B_{Q} denote its intersection with the real form $\mathrm{SConf}(p, q|S_{\mathbb{C}})_0$. This Lie subgroup provides the even symmetries of the twisted theory that act on spacetime by genuine conformal transformations but are trivialized on the twisted theory.

We can draw conclusions from this analysis most readily for specific types of symmetry occurring in the groups Z_Q and B_Q .

Example 1.2 (\mathbb{E}_k -algebras). Suppose there is an affine subspace $\mathbb{R}^k \subseteq C(\mathbb{R}^{p,q})$ with definite signature so that the group of isometries of the subspace lies in B_Q . Then if we restrict observables of Obs^Q to those supported along this affine subspace, we have an isometry action that is trivialized up to homotopy. If the dilation of \mathbb{R}^k also lies in B_Q , then these restricted observables have the structure of a *framed* \mathbb{E}_k -*algebra*: an algebra over the operad of framed little *k*-disks. This structure encodes many properties of a *k*-dimensional topological field theory, and such a twisted theory can reasonably be viewed as topological on this affine subspace.

Remark 1.3. When doing this calculation carefully, we will need the group of isometries of \mathbb{R}^k not only to be exact, but also we will need to choose the potentials coherently: it is necessary to have a subspace of odd symmetries (the potentials) that makes exact the full algebra of isometries, as in Definition 7.10.

Remark 1.4. One could just as well consider not only the subalgebra of *even* closed and exact elements, but the full sub super Lie algebra of *all* closed and exact elements. For instance, the full superalgebra of Q-closed elements will continue to act on the Q-twisted theory, and it could be useful to take advantage of the continued presence of certain odd symmetries. For simplicity we will not consider this further in the present paper.

Example 1.5 (Vertex Algebras). Suppose that there is a subspace $\mathbb{R}^2 \subseteq C(\mathbb{R}^{p,q})$ so that the full group of isometries of \mathbb{R}^2 lies in Z_Q , but only the complexified translation \overline{d}_z lies in $\mathfrak{b}_Q^{\mathbb{C}}$, where z denots a complex coordinate on $\mathbb{R}^2 \cong \mathbb{C}$. Then if we restrict observables of Obs^Q to those

supported along this subspace, we have an isometry action so that the complexified antiholomorphic translation is trivialized up to homotopy. In this case (and under some additional technical assumptions), the restricted observables can be equipped with the structure of a *vertex algebra*. This construction was originally considered by Beem et al [Bee+15b] in their derivation of vertex algebras from 4d $\mathcal{N} = 2$ superconformal field theory.

Our analysis also allows for a situation that lies in between these two extremes, where we extract a field theory on $\mathbb{R}^k \times \mathbb{C}^\ell$ whose observables are topological along the *k* real directions and holomorphic along the ℓ complex directions.

Remark 1.6. The Lie group $\text{SConf}(p,q|\mathcal{N})_0$ actssimultaneously on the space of nilpotent supercharges \mathcal{Q} by the adjoint action and on the spacetime $C(\mathbb{R}^{p,q})$ by conformal transformations. The resulting twisted theories are equivalent. We observe, however, that if we study the algebra of operators after restricting to an affine subspace, the conformal transformations will move this subspace around. Thus for instance, the intersection with a choice of subspace $\mathbb{R}^{p,q} \subseteq C(\mathbb{R}^{p,q})$ may look quite different for different points in the same orbit.

In the main body of this paper, we investigate the following questions.

- (1) For a given superconformal group SConf(*p*, *q*|*N*), what is the locus *N*ilp of square-zero odd elements (i.e., twists) of the complexified Lie algebra? What are the orbits for the adjoint action of the even part of the superconformal group?
- (2) For each orbit, what do the Lie groups and Lie algebras $\mathfrak{z}_{\mathcal{O}}^{\mathbb{C}}, Z_{\mathcal{Q}}, \mathfrak{b}_{\mathcal{O}}^{\mathbb{C}}, B_{\mathcal{Q}}$ look like?
- (3) Can we locate subspaces whose isometry groups are *Q*-closed? If so we can obtain nice isometry-invariant algebras of observables on R^k from the *Q*-twisted theory.
- (4) What does the subalgebra of exact isometries of such a subspace look like both real and complexified — and can we find potentials for these subalgebras? If so the algebra of twisted observables will have nice invariance properties that we can analyze.

Notice that all of these questions are totally independent from the specific choice of superconformal theory, and can be applied to draw conclusions in a completely uniform way. As a specific case study, we revisit the results of Beem et al in Section 4.6.

Acknowledgements. We thank Philip Argyres, Mitch Weaver, and Brian Williams for helpful discussions on this topic, and particularly Richard Eager, who offered detailed comments on the manuscript, especially in guiding us through the literature. The National Science Foundation supported O.G. through DMS Grant 2042052. ML is supported in part by the National Research Foundation of Korea (NRF) Grant 2021R1A2C2012350. C.E. and O.G. thank their REU students at Amherst College (Osha Jones and Ziji Zhou) and UMass Amherst (Nikos Orginos, Nathaniel Reid, and Sam Silver) for work on two- and three-dimensional versions of these problems, which gave helpful complementary insights.

2. SUPERCONFORMAL ALGEBRAS AND TWISTS

We rapidly review the conformal and superconformal groups and their associated (super) Lie algebras. For the conformal situation, we follow Schottenloher's conventions (see Chapter 2 of [Sch08]). 2.1. The Conformal Case. Throughout we will work with a pair (p, q) of natural numbers such that p + q > 2. The case p + q = 2 is strikingly different and so deserves a separate examination.

Let $\mathbb{R}^{p,q}$ denote the vector space \mathbb{R}^{p+q} equipped with the nondegenerate symmetric bilinear pairing of signature (p,q). We say it has dimension D = p + q. We denote the pairing by

$$\langle v,w\rangle = v_1w_1 + \cdots + v_pw_p - v_{p+1}w_{p+1} - \cdots - v_{p+q}w_{p+q},$$

and use $|v|^2 = \langle v, v \rangle$ for the associated "norm." A crucial role for us is played by a closely related space, as it leads to a clean definition of the conformal group and – later – will be the spacetime relevant to twisted superconformal field theories. Let \mathbb{RP}^{p+q+1} denote the p + q + 1-dimensional real projective space.

Definition 2.1. Let $C(\mathbb{R}^{p,q})$ denote the conformal compactification of $\mathbb{R}^{p,q}$. It is the projectivization of the real variety given by the vectors $v \in \mathbb{R}^{p+1,q+1}$ such that $|v|^2 = 0$. In other words, it is the hypersurface in the projective space $\mathbb{R}\mathbb{P}^{p+q+1}$ cut out by the quadratic equation

$$0 = x_0^2 + \dots + x_p^2 - x_{p+1}^2 - \dots - x_{p+q+1}^2$$

where $(x_0 : \cdots : x_{p+q+1})$ is a point given in homogeneous coordinates.

In concrete terms, there is an embedding ι : $\mathbb{R}^{p,q} \hookrightarrow \mathbb{R}^{p+1,q+1}$ by

$$\iota(v_1,\ldots,v_{p+q}) = \left(\frac{1-|v|}{2}, v_1,\ldots,v_{p+q},\frac{1+|v|}{2}\right).$$

Note that $\iota(v)$ is always a vector of norm zero, and that it is never the zero vector. We can thus map $\mathbb{R}^{p,q}$ into the projective space \mathbb{RP}^{p+q+1} by sending v to

$$\left(\frac{1-|v|}{2}:v_1:\ldots:v_{p+q}:\frac{1+|v|}{2}\right)\in\mathbb{RP}^{p+q+1}$$

using homogeneous coordinates. Then $C(\mathbb{R}^{p,q})$ is the natural completion of this copy of $\mathbb{R}^{p,q}$ in projective space.

It is straightforward to describe the symmetries of this compactification: we take the linear automorphisms of $\mathbb{R}^{p+1,q+1}$ that preserve the subspace of norm-zero vectors and then projectivize.

Definition 2.2. *The* conformal group Conf(p,q) *is the connected component of the identity in* $O(p + 1, q + 1)/{\pm 1}$.

There are familiar classes of transformations within this group:

- the orthogonal transformations SO(*p*, *q*),
- the translations \mathbb{R}^{p+q} ,
- the dilations $\mathbb{R}_{>0}$, and
- the "special conformal transformations." The subgroup consisting of such elements is isomorphic to a second copy of the additive group \mathbb{R}^{p+q} .

In fact, these subgroups collectively generate the full conformal group; in other words, we can always factor a conformal transformation as some composition of transformations of those four types. We refer to [Sch08, §2.2] and [DMS97, §4.1] for a thorough discussion ²

²Note that the action of the special conformal transformations on a point in $\mathbb{R}^{p,q}$ can be interpreted as an inversion (about the origin), followed by a translation, followed by another inversion.

Remark 2.3. The identification of these classes of generators of conformal transformations is associated to the chosen conformal embedding $\iota: \mathbb{R}^{p,q} \to C(\mathbb{R}^{p,q})$. For instance the orthogonal transformations, translations and dilations generate the subgroup of conformal transformations that preserve the image of ι . We obtain distinct but conjugate subgroups by choosing alternative conformal embeddings – from our construction of the conformal compactification we obtain such an embedding from every codimension two linear subspace of the form $\mathbb{R}^{p,q} \subseteq \mathbb{R}^{p+1,q+1}$, hence an embedding $SO(p,q) \hookrightarrow SO(p+1,q+1)$.

The Lie algebra conf(p,q) of conformal symmetries is thus isomorphic to $\mathfrak{so}(p+1,q+1)$.

Lemma 2.4. As a vector space, there is a direct sum decomposition

$$\mathfrak{conf}(p,q) \cong \mathfrak{so}(p,q) \oplus \mathbb{R}^{p+q} \oplus \mathbb{R} \oplus \mathbb{R}^{p+q},$$

where the summands correspond, in order, to the Lie algebras of the four subgroups listed above.

This Lie algebra has a representation in vector fields on $\mathbb{R}^{p,q}$, by differentiating the action of the group on the compactification, and the infinitesimal translations, orthogonal transformations, and dilations act in the familiar way, which we write out below in explicit detail. This infinitesimal action plays a central role in characterizing the superconformal Lie algebra below.

2.1.1. A more physical formulation. We now recall a standard notation in physics for generators of conf(p,q) and write the commutation relations. It explicitly realizes conf(p,q) as a Lie subalgebra of the vector fields on \mathbb{R}^{p+q} .

We use the customary notation from the physics literature where repeated indices are summed over, i.e. $\partial^{\mu} x_{\mu} = \sum_{\mu=1}^{d=p+q} \partial^{\mu} x_{\mu}$ Furthermore, indices are "raised" and "lowered" by contracting with the metric (tensor) $g_{\mu\nu}$ or its inverse, i.e.: $\partial_{\mu} = g_{\mu\nu}\partial^{\nu} = \sum_{\nu=1}^{d} g_{\mu\nu}\partial^{\nu}$. The metric is given by: $g_{\mu\nu} = \text{diag}(\mathbb{1}_{p}, -\mathbb{1}_{q})$.

Notation 2.5. Let μ , ν denote indices running from 1 to p + q, associated to the standard basis for \mathbb{R}^{p+q} . The generators of conf(p,q) are

$$M_{\mu\nu} = x_{\mu}\partial_{\nu} - x_{\nu}\partial_{\mu}$$

$$P_{\mu} = -\partial_{\mu}$$

$$D = -x_{\mu}\partial^{\mu}$$

$$K_{\mu} = x^{2}\partial_{\mu} - 2x_{\mu}x_{\nu}\partial^{\nu}$$

and they satisfy

$$\begin{split} [M_{\mu\nu}, M_{\rho\sigma}] &= g_{\rho\nu} M_{\mu\sigma} + g_{\mu\rho} g_{\sigma\nu} - g_{\mu\sigma} M_{\rho\nu} - g_{\sigma\nu} M_{\mu\rho} & [D, P_{\mu}] = P_{\mu} \\ [M_{\mu\nu}, P_{\rho}] &= g_{\nu\rho} P_{\mu} - g_{\mu\rho} P_{\nu} & [D, K_{\mu}] = -K_{\mu} \\ [M_{\mu\nu}, K_{\rho}] &= g_{\nu\rho} K_{\mu} - g_{\mu\rho} K_{\nu} & [K_{\mu}, P_{\nu}] = -2M_{\mu\nu} + 2g_{\mu\nu} D_{\mu\nu} \end{split}$$

with all other commutators vanishing.

Remark 2.6. As in Remark 2.3 the choice of decomposition of the conformal Lie algebra into these four subspaces is non-canonical: it depends on a choice of linear subspace $\mathbb{R}^{p,q} \subseteq \mathbb{R}^{p+1,q+1}$.

2.2. The Superconformal Case. We will focus on superconformal Lie *algebras*, as they play the key role in this paper and because they are simpler to describe in technical terms (In particular, we will not discuss super analogs of the conformal compactification.) Our approach is borrowed from Shnider's elegant and efficient discussion [Shn88]. We begin by recalling how one formulates the super versions of the spacetimes $\mathbb{R}^{p,q}$ and of their isometries. That is, we review what super-Minkowski space is and what the super Poincaré group is (in arbitrary signature) to elucidate the logic that motivates a formulation of superconformal algebras.

Definition 2.7. *The* Poincaré group *in signature* (*p*, *q*) *is the group*

$$\mathrm{ISO}(p,q) = \mathbb{R}^{p+q} \rtimes \mathrm{SO}(p,q)$$

or oriented isometries of $\mathbb{R}^{p,q}$. The Lorentz group is the subgroup $SO(p,q) \subseteq ISO(p,q)$ that fixes the origin in $\mathbb{R}^{p,q}$.

We can (and should) view $\mathbb{R}^{p,q}$ as the homogeneous space given by the quotient of the Poincaré group by the Lorentz group. There is a natural spin cover of the Poincaré group where we replace SO(p,q) by its spin cover. We will typically focus on the common Lie algebra $\mathfrak{so}(p,q)$ of these groups, which does not care about the spin cover. The Lie algebra $\mathfrak{so}(p,q)$ has an infinitesimal action on $\mathbb{R}^{p,q}$.

Super versions of these objects arise as extensions that depend on spinors as follows. Pick a real (Majorana) spinorial representation *S* of Spin(*p*, *q*) and a spin-equivariant symmetric bilinear map $\Gamma: S \times S \to \mathbb{R}^{p+q}$. (Here *S* need not be irreducible; it can be a direct sum of spinor representations. The pairing is unique up to rescaling if *S* is irreducible.)

Definition 2.8. The super Poincaré group is the super Lie group whose odd component is *S* and whose reduced (bosonic) Lie group is the Poincaré spin group; the group structure depends on the pairing Γ on spinors. We denote the super Poincaré group associated to this data by ISO(p,q|S).

There is a super manifold that is the quotient of this super group by the Lorentz spin group; we denote it by $\mathbb{R}^{p,q|s}$, where *s* will denote the dimension of *S*. The super Poincaré Lie algebra $\mathfrak{iso}(p,q|S)$ acts infinitesimally on this superspace. We will often wish to refer to the symmetries of *S* that commute with the spin action. We make the following definition.

Definition 2.9. The R-symmetry group G_R of the super Poincaré group ISO(p,q|S) is the group of outer automorphisms of ISO(p,q|S) that act trivially on the even part.

In looking for a super version of the conformal algebra, we want to construct a super Lie algebra $\mathfrak{sconf}(p, q|S)$ that depends on this choice of a Majorana spinor representation *S* and the bilinear form. It should include the super Poincaré Lie algebra as a subalgebra, just as the Poincaré Lie algebra is a subalgebra of $\mathfrak{conf}(p,q)$. Moreover, the reduced Lie algebra of $\mathfrak{sconf}(p,q|S)$ should be $\mathfrak{conf}(p,q)$. In other words, $\mathfrak{sconf}(p,q|S)$ should be a common generalization of usual supersymmetry algebra and usual conformal algebra.

Remark 2.10. When studying super Lie algebras \mathfrak{g} , we will use the notation [-, -] for the full graded skew-symmetric Lie bracket

$$[-,-]: \wedge^2(\mathfrak{g}) \to \mathfrak{g}$$

In particular we will not use a different notation for the (symmetric) bracket of two odd elements $\operatorname{Sym}^2(\mathfrak{g}_1) \to \mathfrak{g}_0$ compared to the notation the (skew-symmetric) bracket of two even elements. By contrast, in the physics literature these cases are sometimes distinguished in notation, for instance by writing the bracket of two odd elements \mathcal{Q} and \mathcal{Q}' as $\{\mathcal{Q}, \mathcal{Q}'\}$.

It is useful to ask for something stronger: a common generalization *in how they act as symmetries* of the supersymmetry algebra and the conformal algebra. We thus request super Lie algebras satisfying two properties:

Definition 2.11 (*cf.* [Shn88]). A super Lie algebra g is a superconformal algebra of signature p, q if g acts on super-Minkowski space $\mathbb{R}^{p,q|s}$ by infinitesimal derivations, and the following conditions hold:

- (1) There is an embedding $i: \mathfrak{iso}(p,q|S) \hookrightarrow \mathfrak{g}$ of super Lie algebras, and the restriction of the action along i coincides with the standard action of infinitesimal super-isometries.
- (2) There is an embedding j: conf(p,q) → g, and the restriction of the action along j coincides with the standard action of infinitesimal conformal transformations on ℝ^{p,q}, acting trivially on the odd coordinates.

The remarkable theorem is that such a super Lie algebra exists *only* when $p + q \le 6$, as discovered by Nahm [Nah78], although we emphasize here the clean mathematical approach of Shnider [Shn88], who articulates these properties as explicit hypotheses. Shnider begins by studying the complexification of the situation above, since any such real super Lie algebras has a complexification and there is no dependence on a choice of signature after complexifying. He then shows that if such a complex super Lie algebra g exists, there is a simple super Lie algebra, satisfying the same properties, which is a quotient or subalgebra of g. Moreover, its bosonic part splits as the sum of $\mathfrak{so}(p + q, \mathbb{C})$ and a complementary ideal. He then uses Kac's classification of simple super Lie algebras to show that such complex super Lie algebras do *not* exist above dimension 6. Identifying the relevant simple super Lie algebras in dimensions less than or equal to 6 is possible and already accomplished.

We will follow some of this strategy below, typically analyzing a problem in the complexification before turning to real forms. As in the bulk of the literature, we restrict our attention to the simple cases.

List 2.12. We record the simple super Lie algebras that appear as complexified superconformal algebras in dimensions 3 to 6.

- In dimension 3, the superconformal algebras are osp(k|4, C) for k ≥ 1. The even part is so(k, C) ⊕ sp(4, C) ≅ so(k, C) ⊕ so(5, C), containing the conformal algebra so(5, C) ≅ conf(3, C).
- In dimension 4, the superconformal algebras are $\mathfrak{sl}(4|k,\mathbb{C})$ for $k \ge 1, k \ne 4$, and $\mathfrak{psl}(4|4,\mathbb{C})$ in the special case k = 4. The even part is $\mathfrak{sl}(4,\mathbb{C}) \oplus \mathfrak{sl}(k,\mathbb{C}) \oplus \mathbb{C}$ if $k \ne 4$ and $\mathfrak{sl}(4,\mathbb{C}) \oplus \mathfrak{sl}(4,\mathbb{C})$ in the k = 4 case, containing the conformal algebra $\mathfrak{sl}(4,\mathbb{C}) \cong \mathfrak{so}(6,\mathbb{C}) \cong \mathfrak{conf}(4,\mathbb{C})$.
- In dimension 5, there is a unique superconformal algebra, the exceptional super Lie algebra f(4, C). The even part is so(7, C) ⊕ sl(2, C), containing the conformal algebra so(7, C) ≅ conf(5, C).
- In dimension 6, the superconformal algebras are osp(8|2k, C) for k ≥ 1. The even part is so(8, C) ⊕ sp(2k, C), containing the conformal algebra so(8, C) ≅ conf(6, C).

Remark 2.13. In each of these examples, the even part of the superconformal algebra includes $conf(n, \mathbb{C}) \cong \mathfrak{so}(n + 2, \mathbb{C})$ as a summand. We will denote the quotient algebra by \mathfrak{g}_R , and refer to it as the algebra of R-symmetries. In each case – as we can see directly from the classification – \mathfrak{g}_R coincides with the Lie algebra of the group G_R of outer automorphisms of the superconformal algebra that act trivially on the even part (the *group* of R-symmetries).

Remark 2.14. The classification of simple super Lie algebras has played an essential role in the physics literature on the structure and properties of superconformal theories [Mac77; Min98; BHP16]. In particular, Dolan and Osborn [DO03] developed the representation theory of 4d superconformal algebras, and Córdova, Dumitrescu and Intriligator [CDI19] provided a thorough analysis of the representation theory of superconformal algebras in dimensions 2 to 6 and its applications to superconformal field theory.

2.3. **The Question of Classifying Twists.** Topological field theories have had an enormous impact in both theoretical physics and mathematics, and the most important examples arise from supersymmetric theories by a twisting procedure, introduced by Witten [Wit88]. We recall here a less sophisticated version of that procedure before we turn to superconformal field theories., In between, we provide a quick survey of mathematical work on these notions, mostly growing out of the pioneering but unpublished paper of Duflo and Serganova [DS05], as it provides some important structural results.

2.3.1. *How Twists Appear in the Supersymmetric Case.* Loosely speaking, a supersymmetric field theory is a field theory on $\mathbb{R}^{p,q}$ with an action of a super Poincaré Lie algebra. (One typically asks for more, but we will focus on this kind of symmetry for now.) It is reasonable to explore how elements of the Lie algebra of supertranslations act on the theory. Let $\mathfrak{stran}(p,q|S)$ be this super Lie algebra, where *S* denotes the direct sum of spinor representations that provides the odd component and the non-trivial Lie bracket is given by a vector-valued pairing Γ on *S*.

Remark 2.15. For the moment, by $\mathfrak{stran}(p,q|S)$ we mean a *real* Lie algebra whose even part is $\mathbb{R}^{p,q}$. Later on, and in the parallel superconformal setting, we will consider the less refined setting of theories with an action of the complexification of such a real Lie algebra. The discussion of twists that follows makes sense either over the real or the complex numbers.

Definition 2.16. *A* translation twist *is an odd element* Q *such that* [Q, Q] = 0.

A choice of twist leads to some interesting structures:

(1) A subspace $\mathfrak{b}_{\mathcal{Q}} \subset \mathbb{R}^{p,q}$ defined by

$$\mathfrak{b}_{\mathcal{Q}} = \operatorname{Im}(\mathcal{Q})$$

where we view the twist as a map $Q: \Pi S \to \mathbb{R}^{p,q}$ sending an odd element *s* to the even element [Q, s]. Such a subspace determines a foliation \mathcal{F}^{Q} of $\mathbb{R}^{p,q}$ into affine leaves.

(2) A deformation of a supersymmetric theory by adding Q to its BRST differential (i.e., by adding a term to its Lagrangian that encodes how Q acts on the theory). This deformed theory is "trivial" along leaves of the foliation \mathcal{F}^{Q} in the sense that BRST cohomology classes of observables are isomorphic under translation along these leaves.

(3) A differential Z/2-graded Lie algebra (stran(p, q|S), [Q, −]), which can be seen as acting on the deformed field theory.

We call such a *Q*-deformation of a supersymmetric field theory a *twisted theory*, and we see that it admits a symmetry by this dg Lie algebra.

When $\mathfrak{b}_{\mathcal{Q}}$ contains all of $\mathbb{R}^{p,q}$, we call \mathcal{Q} a *topological* twist as the associated twisted theories must be "topological" in the sense that we can move insertion points around without changing expected values. But there are cases where $\mathfrak{b}_{\mathcal{Q}}$ is a proper subspace, leading to more subtle changes in the theory. For instance, in the Euclidean setting, twists lead to theories that can be holomorphic in some directions and topological in others.

Recently, there has been a systematic classification of the space of twists (at least for complexified supertranslation Lie algebras), motivated by the desire to organize and analyze the behavior of twisted supersymmetric field theories. For details of this classification, see work of the first author with Safronov and Williams [ES19; ESW22] and of Eager, Saberi and Walcher [ESW21].

The following definition when applied to the super Lie algebra $\mathfrak{g} = \mathfrak{stran}(p, q|S)$ characterizes the space of possible supertranslation twists.

Definition 2.17. Let $\mathfrak{g} = \mathfrak{g}_0 \oplus \Pi \mathfrak{g}_1$ be a super Lie algebra, where \mathfrak{g}_i for i = 0, 1 refer to its even and odd summands and Π is the parity shift operator. The (affine) nilpotence variety of \mathfrak{g} is the affine quadric subvariety of \mathfrak{g}_1 defined by

$$\operatorname{Nilp}_{\mathfrak{a}} = \{ \mathcal{Q} \in \mathfrak{g}_1 \colon [\mathcal{Q}, \mathcal{Q}] = 0 \}.$$

This affine variety is preserved by the rescaling action, and the projective nilpotence variety $\mathbb{P}Nilp_{\mathfrak{g}} \subseteq \mathbb{P}(\mathfrak{g}_1)$ of \mathfrak{g} is its projectivization.

The affine and projective nilpotence varieties both carry an action of the Lie algebra \mathfrak{g}_0 inherited from the adjoint action. If \mathfrak{g} is the super Lie algebra of a super Lie group *G*, then the even part G_0 of *G* also acts on the nilpotence varieties by the group-level adjoint action.

In the case where $\mathfrak{g} = \mathfrak{stran}(p,q|S)$ the nilpotence variety carries a natural action of the group $\operatorname{Spin}(p,q) \times G_R$ where G_R is the group of R-symmetries.

Definition 2.18. The moduli stack of twists for the supertranslation algebra is the quotient stack

$$\text{Twist}_{\mathfrak{g}} = \left[\text{Nilp}_{\mathfrak{g}} / (\text{Spin}(p,q) \times G_R) \right].$$

2.3.2. *Associated Varieties of Super Lie Algebras.* The notions we just mention were already explored by mathematicians who study super Lie algebras, so we quickly record the terminology used by this community and some of their pertinent results.

Definition 2.19. For a finite-dimensional super Lie algebra \mathfrak{g} over \mathbb{C} , its associated variety consists of the square-zero odd elements:

$$X_{\mathfrak{g}} = \{ x \in \mathfrak{g}_1 \mid [x, x] = 0 \}.$$

It is a Zariski-closed conical subset of g_1 , viewed as an affine space.

Hence the associated variety of a supertranslation algebra is precisely the affine nilpotence variety introduced in the preceding section. Note that a point $x \in X_g$ determines a dg Lie algebra

 $(\mathfrak{g}, [x, -])$, and Duflo and Serganova studied both its cohomology $\mathfrak{g}_x = \mathrm{H}^*(\mathfrak{g}, [x, -])$ and the *Duflo-Serganova* functor

$$DS_x: Mod(\mathfrak{g}) \to Mod(\mathfrak{g}_x),$$

which is nicely behaved. For instance, it is symmetric monoidal, preserves the superdimensions of finite-dimensional representations, and induces ring maps between the reduced Grothendieck groups of (well-behaved) abelian subcategories (such as finite-dimensional representations).

Let G_0 denote the simply-connected, connected algebraic group with Lie algebra \mathfrak{g}_0 . It acts on \mathfrak{g}_1 by the adjoint action and $X_\mathfrak{g}$ is preserved by this action. It is natural to study the stack quotient $[X_\mathfrak{g}/G_0]$, and in the case that \mathfrak{g} is a supertranslation algebra, it maps to the moduli stack of twists introduced in the preceding section. (That moduli space is a further quotient, taking into account the natural isometries, like rotation, of super-Minkowski space.)

When \mathfrak{g}_0 is reductive and \mathfrak{g}_1 is a semisimple \mathfrak{g}_0 -module, the super Lie algebra \mathfrak{g} is called *quasireductive*. In this case there is a Cartan subalgebra $\mathfrak{t} \subset \mathfrak{g}_0$. Let Δ_1 denote the roots of \mathfrak{t} acting on \mathfrak{g}_1 , and let Δ_0 denote the (usual, even) roots of \mathfrak{g}_0 .

There is a remarkable theorem showing that the set X_g/G_0 of G_0 -orbits in the associated variety is *finite* for an important class of Lie superalgebras (which does *not* include supertranslations). We will use this result in the superconformal setting.

To state the theorem, we must introduce more terminology.

Definition 2.20. An iso-set for \mathfrak{g} is a subset A of Δ_1 such that

- its elements are linearly independent and
- *if any pair of odd roots* $\alpha, \beta \in \Delta_1$ *satisfies* $\pm \alpha, \pm \beta \in A$ *, then* $\alpha + \beta \notin \Delta_0$ *.*

Let $S_{\mathfrak{g}}$ *denote the collection of iso-sets.*

The Weyl group *W* of G_0 preserves $S_{\mathfrak{g}}$.

Theorem 2.21 (Theorem 5.1, [Gor+22]). *There is a bijection between* X_g/G_0 *and* S_g/W *if* \mathfrak{g} *is a complex super Lie algebra from the list*

 $\{\mathfrak{gl}(m|n),\mathfrak{sl}(m|n) \text{ when } m \neq n, \mathfrak{osp}(m|2n), \mathfrak{d}(2|1;a), \mathfrak{f}(4), \mathfrak{g}(3)\}.$

In particular, there is a finite set of orbits.

In [Gor+22] they push further and for such classical super Lie algebras, they classify the stabilizers and normalizers for each point $x \in X_g$, as well as the dimensions of the orbits.

We want to raise a cluster of natural questions. We mention physical motivations below.

Question 2.22. For a real super Lie group (particularly the real simple ones), what are the real twists of its real super Lie algebra? That is, what is the associated variety? Moreover, what are the orbits?

2.3.3. *How Twists Appear in the Conformal Case.* We raise here the question of studying the analogue of twists by elements of a superconformal Lie algebra.

Let $\mathfrak{sconf}(p, q|S)$ be the super Lie algebra, where *S* denotes the spinor representation that provides the odd component and Γ is the pairing on *S*.

Definition 2.23. *A* conformal twist *is an odd element* $Q \in \Pi S$ *such that* $[Q, Q] = \Gamma(Q, Q) = 0$.

A twist Q determines a subspace of conf(p,q) and hence a foliation of the conformal compactification $C(\mathbb{R}^{p,q})$. If we view a superconformal field theory as living on $C(\mathbb{R}^{p,q})$, a twist then determines a deformation to a theory that is "trivial" along the leaves of this foliation. We call it the Q-twisted superconformal theory. Moreover, the dg Lie algebra $(\mathfrak{sconf}(p,q|S), [Q, -])$ acts on this twisted theory. It is to be hoped that such twisted theories admit phenomena at least as special and interesting as twists of supersymmetric theories. Indeed, since many supertranslation algebras may be embedded as subalgebras of superconformal algebras, many supersymmetric twists arise as special examples of superconformal twists. In this sense twisting in the superconformal context may be viewed as an extension of twisting in the usual context. Hence we encounter a basic question.

Question 2.24*. For each signature* (p,q) *with* p + q > 2*, what are the twists of the corresponding super-conformal Lie algebra?*

In the bulk of this paper we begin to explore this question by studying a few explicit examples and discuss how twisting affects theories in qualitative terms.

Following [ES19; ESW21], we phrase our problem in the following way. Note that for the rest of this section we will discuss nilpotent elements of the *complexified* superconformal algebra; we will discuss the role of real forms shortly.

Definition 2.25. The conformal nilpotence variety for $\mathfrak{sconf}(p,q|S)$ is the associated variety $X_{\mathfrak{sconf}}(p,q|S)$ of the complex super Lie algebra $\mathfrak{sconf}(p,q|S)_{\mathbb{C}}$. We denote it by $\operatorname{Nilp}_{(p,q|S)}(\mathbb{C})$.

Note that for each embedding of super Lie algebras from a supertranslation algebra to the corresponding superconformal algebra, there is a map of nilpotence varieties. Every translation twist determines a conformal twist, in other words.

There are natural symmetries to take into account.

• Let $SConf(p, q|S)_0$ is the simply-connected, connected algebraic group over \mathbb{C} whose Lie algebra is $\mathfrak{sconf}(p, q|S)$. It acts on the associated variety, and if two twists are in the same orbit, then the associated dg Lie algebras are isomorphic. Thus, it is natural to consider the quotient stack

$$[X_{\mathfrak{sconf}(p,q|S)}/\mathrm{SConf}(p,q|S)_0],$$

which we call the maximal quotient stack of conformal twists.

• On the other hand, this group $SConf(p, q|S)_0$ does *not* play nicely with the original superconformal geometry that motivated us. Instead it has various *real* forms that are symmetries of real super manifolds that "extend" the conformal compactification $C(\mathbb{R}^{p,q})$. If *G* denotes one of these real forms, it is natural to consider the quotient stack

$$[X_{\mathfrak{sconf}(p,q|S)}/G],$$

which we call the *G*-quotient stack of conformal twists.

One goal of this paper is to describe the geometric properties of these spaces, with the following possibility as motivation.

Question 2.26. Are there orbits among the conformal twists that do *not* come from the translation twists? These would leave to novel twisted field theories.

In light of List 2.12 and Theorem 2.21, Duflo and Serganova have given essentially a complete treatment of the maximal quotient stacks: they showed that for (nearly) every conformal nilpotence variety, its $SConf(p, q|S)_0$ -orbits consist of a finite set of points. That result implies each maximal quotient stack is a disjoint union of classifying stacks */Stab(x) where $x \in X_{sconf}(p,q|S)_0$ is a point on the associated orbit and where Stab(x) denotes its stabilizer inside $SConf(p,q|S)_0$.

The only open case is $\mathfrak{psl}(4|4)$, which we examine below. For this algebra, the orbits do *not* form a finite set; thus the stack is more interesting. This case is particularly interesting because it corresponds to $\mathcal{N}=4$ supersymmetric theories in four dimensions, which includes many special field theories.

But for physical questions, it is also important to examine the quotients by *real* groups. For example, the work of [Bee+15b] examines a complex twist but working in a fixed signature, and hence with an associated real form.

Finally, we wish to point out a third important situation. It is natural

- to consider twists in the *real* super Lie algebras arising from the real superconformal groups, (i.e., the real points Nilp_(p,q|S)(R) of the conformal nilpotence variety) and
- to identify the real orbits of these real twists.

In other words, we raise Question 2.22 here specifically for the real forms of the superconformal algebras. In this situation, we obtain Lie subalgebras of vector fields on the conformal manifold $C(\mathbb{R}^{p,q})$, namely $\mathfrak{z}_{\mathcal{Q}} = \text{Ker}(\mathcal{Q})$ and $\mathfrak{b}_{\mathcal{Q}} = \text{Im}(\mathcal{Q})$. These foliations have rich geometric content, and unpacking their geometry would undoubtedly be fruitful.

This third situation also has clear physical motivation: an action functional of a genuine physical theory should be real-valued, and a Hamiltonian should be self-adjoint. A superconformal field theory has an action of a real superconformal group, so it is natural to look for twists that determine deformations as a genuine theory. That pushes us to look at the real twists.

3. THE OBSERVABLES OF A TWISTED SUPERCONFORMAL THEORY

Given a superconformal theory with $\mathfrak{sconf}(p,q|S)$ as a symmetry algebra, a nilpotent supercharge Q can be added to the action as a BRST operator leading to a new theory that has the Q-cohomology of $\mathfrak{sconf}(p,q|S)$ as a symmetry algebra. For supersymmetric theories, this process (often dubbed "twisting") is well-known and produces new theories with remarkable properties, such as the topological field theories encoding Donaldson invariants of 4-manifolds [Wit88]. A crucial step is to understand the Q-cohomology of $\mathfrak{siso}(p,q|S)$ and its geometric meaning. For superconformal theories, the twists can have more exotic geometric consequences and the associated twisted theories have surprising behavior, including the localization discovered by Beem et al [Bee+15b].

Remark 3.1. We observe that while we introduced the notion of twisting as a modification of the action functional by the introduction of terms proportional to the nilpotent supercharge Q, we can, alternatively work directly with the algebra of operators that appear in the twisted theory by analyzing the Q-cohomology of the untwisted algebra of operators. As a consequence, our derivations will be valid also in the cases of inherently strongly coupled theories, defined in the absence of a classical action functional.

In this section we will give a physical description of how conformal twists affect a theory, and in section 7 we offer detailed mathematical formulations using the language of disc-algebras and factorization algebras.

3.1. **Subalgebras of** Q**-Closed and** Q**-Exact Elements.** Let Q be an odd square-zero element of $\mathfrak{sconf}(p,q|S)$. As we hinted above, the properties of the Q-twist of a superconformal field theory defined on $C(\mathbb{R}^{p,q}) - e.g.$, whether the twisted theory becomes topological or holomorphic upon restriction to certain affine subspaces – are determined by the cohomology of $\mathfrak{sconf}(p,q|S)$ under the differential [Q, -]. We will make this idea more precise as follows.

Definition 3.2. Let $Z_Q \subseteq \text{Conf}(p,q) \times G_R$ denote the stabilizer group of the element Q under the adjoint action. Let \mathfrak{z}_Q denote the (real) Lie algebra of Z_Q . In other words, it is the even part of the centralizer of Q in the superconformal algebra. We will also write $\mathfrak{z}_Q^{\mathbb{C}} = \mathfrak{z}_Q \otimes_{\mathbb{R}} \mathbb{C}$ for its complexification.

Definition 3.3. Let $\mathfrak{b}_{\mathcal{O}}^{\mathbb{C}}$ denote the image of the operator

$$[\mathcal{Q}, -]$$
: $\mathfrak{sconf}(n|S, \mathbb{C})_1 \to \mathfrak{sconf}(n|S, \mathbb{C})_0$,

arising from the adjoint action of Q.

These two Lie algebras are related, as the notation – drawn from homological algebra – suggests.

Proposition 3.4. The subspace $\mathfrak{b}_Q^{\mathbb{C}} \subseteq \mathfrak{z}_Q^{\mathbb{C}}$ is a (complex) Lie subalgebra. In fact, it is an ideal, and so the quotient $\mathfrak{z}_Q^{\mathbb{C}}/\mathfrak{b}_Q^{\mathbb{C}}$ is a Lie algebra.

As a result, we can identify $\mathfrak{b}_{\mathcal{Q}}^{\mathbb{C}}$ as the Lie algebra of a Lie subgroup $B_{\mathcal{Q}}^{\mathbb{C}} \subseteq Z_{\mathcal{Q}}^{\mathbb{C}}$ inside the complexification of $Z_{\mathcal{Q}}$. Write $B_{\mathcal{Q}}$ for the real Lie subgroup of $Z_{\mathcal{Q}}$ obtained with the Lie algebra $\mathfrak{b}_{\mathcal{Q}} = \mathfrak{b}_{\mathcal{Q}}^{\mathbb{C}} \cap \mathfrak{z}_{\mathcal{Q}}$.

Remark 3.5. As we discussed in Section 2.3.3 we will be particularly interested in the complexified Lie algebra (in the odd part of which the superconformal nilpotence variety is embedded), as well as real forms of the superconformal group (acting on the nilpotence variety. We define Z_Q intrinsically as a subgroup of the real superconformal group. On the other hand, for the image B_Q our must start with the (complex) Lie algebra, so we are forced to make our definition starting with the complex subgroup $B_Q^{\mathbb{C}}$ and then intersecting with a chosen real form.

Proof. Let $X \in \mathfrak{z}_Q$ and $Y = [Q, Q'] \in \mathfrak{b}_Q$. Then

$$[\mathcal{Q}, [X, \mathcal{Q}']] = [[\mathcal{Q}, X], \mathcal{Q}'] + [X, [\mathcal{Q}, \mathcal{Q}']] = [X, Y]$$

so [X, Y] is also in \mathfrak{b}_Q . This computation shows \mathfrak{b}_Q is an ideal.

3.2. **Observables and BRST Operators.** There are two major types of manifolds on which a superconformal field theory might live. It might be defined on $\mathbb{R}^{p,q}$ (as is usually implicit when a Lagrangian is given in coordinates), or it might be defined on the conformal compactification $C(\mathbb{R}^{p,q})$.

3.2.1. *The Maximal Case.* To start let us suppose the theory \mathcal{T} lives on the biggest possible space, namely $C(\mathbb{R}^{p,q})$. In that case, we may suppose that the superconformal group acts as symmetries of the theory; in particular, its even part provides a real Lie group acting as symmetries of the theory.

Let $\mathcal{T}^{\mathcal{Q}}$ denote the theory obtained from \mathcal{T} by twisting by the supercharge \mathcal{Q} . The full group of conformal transformations no longer acts on $\mathcal{T}^{\mathcal{Q}}$, but the group $Z_{\mathcal{Q}}$ and its subgroup $B_{\mathcal{Q}}$ continue to act. Similarly, there are the infinitesimal symmetries given by actions of the Lie algebras $\mathfrak{z}_{\mathcal{Q}}$ and $\mathfrak{b}_{\mathcal{Q}}$, which act on operators and hence also on operator products.

Note that group elements of SConf₀ that do *not* fix Q are *not* symmetries. Nor do non-Q-closed elements of \mathfrak{sconf}_0 act as infinitesimal symmetries. Twisting the theory changes the symmetries.

In the twisted theory, we focus on Q-closed operators (i.e., we only consider an operator O satisfying QO = 0). Moreover, Q-exact operators (i.e., O = QO' for some operator O') have trivial expected value. This means that in the twisted theory we reduce to studying cohomology classes of operators $[O]_Q$: that is, two operators differing by a Q-exact operator, i.e. O and O' = O + QO'' are identified in the twisted theory, $[O]_Q \simeq [O']_Q$. These properties have strong consequences for how the symmetries act on operators. Recall that in a topological twist of a supersymmetric theory, an *n*-point function

$$\langle \mathcal{O}_1(x_1)\mathcal{O}_2(x_2)\cdots\mathcal{O}_n(x_n)\rangle$$

is independent of the insertion points $\{x_i\}$ because translations are Q-exact. The same argument implies that for a conformal twist, the operator product is constant as we move an insertion point along a B_Q -orbit:

$$\langle \mathcal{O}_1(x_1)\mathcal{O}_2(x_2)\cdots\mathcal{O}_n(x_n)\rangle = \langle \mathcal{O}_1(b(x_1))\mathcal{O}_2(x_2)\cdots\mathcal{O}_n(x_n)\rangle$$

for any $b \in B_Q$. In formulas, we see this claim infinitesimally: let $\beta = [Q, Q']$ be a Q-exact element of the superconformal algebra and compute

$$\langle (\beta \mathcal{O}_1(x_1)) \mathcal{O}_2(x_2) \cdots \mathcal{O}_n(x_n) \rangle = \langle \mathcal{Q}(\mathcal{Q}' \mathcal{O}_1(x_1)) \cdots \mathcal{O}_n(x_n) \rangle + \langle \mathcal{Q}'(\mathcal{Q} \mathcal{O}_1(x_1)) \cdots \mathcal{O}_n(x_n) \rangle$$

= 0 + 0 = 0.

Thus, the twisted theory $\mathcal{T}^{\mathcal{Q}}$ will behave like it is "topological" along $B_{\mathcal{Q}}$ -orbits, in the sense that moving insertions along those orbits does not affect operator products.

In contrast, the action of Z_Q looks more like the action of translation in a supersymmetric theory before twisting. For instance, let \mathcal{O}_1 and \mathcal{O}_2 be Q-closed operators and $z \in Z_Q$. The operator product satisfies

$$\langle \mathcal{O}_1(x_1)\mathcal{O}_2(x_2)\rangle = \langle \mathcal{O}_1(z(x_1))\mathcal{O}_2(z(x_2))\rangle,$$

so that it is preserved by "translation" in the *z*-direction of *both* insertion points x_1 and x_2 . This is the same behaviour as we see in overall coordinate transformations in a Lorentz-invariant theory, the only difference here is that only transformations along a Z_Q -orbit are possible.

On the other hand, if we translate just one insertion, the expected value changes: $\langle O_1(z(x_1))O_2(x_2) \rangle$ varies with *z*, much as a 2-point function typically depends on the relative position of the insertion.

To summarize, we should analyze $C(\mathbb{R}^{p,q})$ (and configurations of multiple points in it) into Z_Q -orbits and, further, into B_Q -orbits. The twisted theory \mathcal{T}^Q will behave like it is "topological" along B_Q -orbits and equivariant along Z_Q -orbits.

Here we have emphasized the geometry of how these groups act on spacetime, but these groups also act, in some sense, "internally,", since the subgroups Z_Q and B_Q usually involve elements embedded diagonally in the product of the group of R-symmetries and the group of conformal symmetries.

3.2.2. Working on an Affine Patch. Now suppose we pick an "affine patch" $\mathbb{R}^{p,q} \subset C(\mathbb{R}^{p,q})$. We mean here that we pick the image $g\mathbb{R}^{p,q}$ of the defining $\mathbb{R}^{p,q}$ under the action of some element g of the conformal group. On such a patch, a theory has a nice coordinate description, and so it can be analyzed very explicitly. Note, however, that orbits of the groups Z_Q and B_Q are not contained in the affine patch.

Nonetheless, our above arguments about operators and their products carry over to this situation, with small modifications. For instance if \mathcal{O}_1 and \mathcal{O}_2 are \mathcal{Q} -closed operators and $z \in Z_{\mathcal{Q}}$, then the operator product satisfies

$$\langle \mathcal{O}_1(x_1)\mathcal{O}_2(x_2)\rangle = \langle \mathcal{O}_1(zx_1)\mathcal{O}_2(zx_2)\rangle,$$

so long as the points x_1, x_2 and their translates $z(x_1), z(x_2)$ all live in the affine patch. Similarly, the operator product is constant if we move an insertion point along a B_Q -orbit intersected with the affine patch.

3.3. **Structures on Observables on Affine Subspaces.** Let us summarize some of the structures that can be realized on the algebra of local operators of a twisted theory in an affine subspace (i.e., an affine subspace of an affine patch). For more detailed proofs we refer the reader to Section 7.

3.3.1. *Vertex Algebras.* We begin with the structure that originally motivated the authors of [Bee+15b] to study the observables in a conformal twist, that of a *vertex algebra*. This structure occurs when there is a two-dimensional affine subspace, which we identify with a copy of the complex plane $\mathbb{C} \hookrightarrow C(\mathbb{R}^{p,q})$ with holomorphic coordinate *z*. We require specific conditions on the twisting supercharge Q that we can describe as follows:

- (1) The group of isometries of \mathbb{C} should be closed (i.e. contained in $Z_{\mathcal{Q}}$).
- (2) The complexified translation ∂/∂z̄ in the antiholomorphic direction should be exact (i.e. contained in b^C_O).

Theorem 3.6 (See Theorem 7.15). If Q satisfies conditions (1) and (2) for an affine subspace \mathbb{C} , then the algebra of local operators in a Q-twisted theory \mathcal{T}^Q , restricted to this affine subspace, has the structure of a vertex algebra, under some additional functional analytic conditions.

We refer to Section 7.3.2 for a more precise statement.

Remark 3.7. In the introduction to [BRR15], it is observed that one way to obtain the desired structure of a vertex algebra in the twist is by choosing an embedding of $\mathfrak{su}(1,1|2)$ into the superconformal algebra, so that $\mathfrak{su}(1,1)$ acts on an affine subspace \mathbb{C} by antiholomorphic Möbius transformations. From this point of view, the supercharge \mathcal{Q} is identified with an odd element of

the complexification $\mathfrak{sl}(2|2,\mathbb{C})$, which is chiral of rank (2,0), and the subalgebra of \mathcal{Q} -exact elements is a copy of $\mathfrak{sl}(2,\mathbb{C})$ embedded diagonally in the even part of $\mathfrak{sl}(2|2,\mathbb{C})$. The authors of *loc. cit.* observe that such subalgebras only occur in a small number of examples of superconformal algebras (at least in Lorentzian signature), occurring in dimension 2, 4 and 6.

Remark 3.8. Given a vertex algebra of local operators along an affine subspace C in such a twisted theory, [ALW23] enhance the vertex algebra to a vertex algebra that includes *extended operators* in the directions transverse to the chosen complex plane (though not extended *within* the complex plane). We do not prove general results about such algebras of extended operators in this paper.

Remark 3.9. The examples of conformal twists studied by Beem *et al*, as discussed in Remark 3.7 above, are designed so that the vertex algebra carries an action of $\mathfrak{sl}(2, \mathbb{C})$. In fact, in their examples this action can be enhanced to an action of the Virasoro algebra (an infinite-dimensional Lie algebra). That is, the algebra of local operators is not just of a vertex algebra, but a vertex *operator* algebra. It would be interesting to analyze exactly what conditions on the superconformal theory and a chosen conformal twist guarantee that this enhancement is possible.

3.3.2. Topological Field Theories. In contrast, we can instead consider an affine subspaces $\mathbb{R}^k \hookrightarrow C(\mathbb{R}^{p,q})$ on which the Q-twisted theory is fully topological, i.e., the entire algebra of translations of \mathbb{R}^k is Q-exact. Examples of this nature generalize the topological twists of supersymmetric field theories first studied by Witten [Wit88], but conformal twists allow us to expand the range of possible supercharges to which we may apply the twisting construction.

The structure of the local observables in a topological twist can be described naturally using the structure of an \mathbb{E}_{k} -algebra. Let us briefly overview the idea behind these structure, though we will not give a complete definition here (see for instance [CG17, Chapter 3.6.4] and [MSS02, Chapter 4.1] for more details and context).

Let *A* be a complex vector space, or maybe more generally a cochain complex. To give *A* the structure of an \mathbb{E}_k -algebra is to define a multiplication operator $m_e \colon A^{\otimes j} \to A$ for each choice of embedding

$$e: B_k(x_1, r_1) \sqcup B_k(x_2, r_2) \sqcup \cdots \sqcup B_k(x_i, r_i) \hookrightarrow B_k(0, 1),$$

where $B_k(0, 1)$ is the *k*-dimensional open unit ball in \mathbb{R}^k , and $B_k(x_i, r_i)$ is a sub-ball of radius r_i and center x_i . These multiplication operators should vary smoothly as we vary the x_i and r_i , and they should be "associative," meaning compatible under concentric inclusions of *k*-balls.

Let us consider the simplest non-trivial example, that of an \mathbb{E}_1 -algebra. We will see that this is just a "homotopical" version of an associative algebra.

Example 3.10. Let k = 1, so $B_1(x, r)$ is the open interval $(x - r, x + r) \subseteq \mathbb{R}$. If we fix a positive integer j, the space of embeddings of j open intervals in the unit interval has connected components given by orderings on the set $\{1, ..., j\}$, with each component being contractible.

For example, if our multiplication operators m_e are constant on each connected component, then we may view m_2 as describing an ordinary *associative algebra* structure on A: the compatibility of the operations m_e under concentric inclusion is equivalent to the ordinary associativity condition.

Remark 3.11. More generally, one may think of an \mathbb{E}_k -algebra as an associative algebra that is "part-way" to commutative. In other words, \mathbb{E}_k interpolates between "associative" (corresponding to \mathbb{E}_1) and "commutative" (corresponding to \mathbb{E}_{∞}). For example, when we consider an \mathbb{E}_2 algebra, it has a binary product associated to a pair of two disjoint disks in the unit disk. There is a path in the space of embeddings that swaps these two disks, which can be seen as "commuting" the two inputs of the product. However, there is still a residue of non-commutativity: if one moves one disk all the way around the other (i.e. applies the swap twice), this path describes a loop in the space of embeddings that is non-contractible. (i.e. a non-trivial element of π_1 of the space of possible binary multiplications). More generally in an \mathbb{E}_k algebra the space of binary multiplications will have non-trivial topology represented by a non-trivial $(k - 1)^{st}$ homotopy group.

Remark 3.12. There is an enhanced version of an \mathbb{E}_k -algebra called a *framed* \mathbb{E}_k -algebra: instead of embeddings of *k*-balls, one studies framed embeddings. This amounts to keeping track of the action of the rotation group SO(*k*) on *A*. We will not go into detail here, except to point out that a framed \mathbb{E}_k -algebra may be used to construct a topological field theory on a general oriented *k*-manifold by the process of factorization homology as developed by Ayala and Francis [AF15]. See [Sch14] for a full development of this idea.

Theorem 3.13 (See Theorem 7.12 and Corollaries 7.13, 7.14). *If the algebra of translations of an affine* patch \mathbb{R}^k is \mathcal{Q} -exact and the action of dilations of \mathbb{R}^k is \mathcal{Q} -exact, then the local operators on \mathbb{R}^k have the structure of an \mathbb{E}_k -algebra.

If the metric restricted to the affine patch is positive definite and the action of rotations is also Q-exact, then the local operators on \mathbb{R}^k have the structure of a framed \mathbb{E}_k -algebra.

Example 3.14. For instance, if k = 1, so we are restricting to a line along which the translation is exact, then the local operators naturally have the structure of an ordinary associative algebra.

3.3.3. *Higher Dimensional Non-Topological Examples.* Finally, let us briefly consider a more expansive class of examples, where we study an affine patch of dimension > 2 that is nevertheless not fully topological. *Holomorphic* examples occur when there is an even real-dimensional affine subspace that we identify as a copy of $\mathbb{C}^d \subseteq C(\mathbb{R}^{p,q})$, all of whose isometries are Q-closed and whose anti-holomorphic complexified isometries are Q-exact. One may also study affine subspace of the form $\mathbb{C}^d \times \mathbb{R}^k \subseteq C(\mathbb{R}^{p,q})$ where the Q-twist is holomorphic in the first factor — in the sense we just outlined — and topological in the second factor — in the sense of the previous subsection.

Let us say more about one specific example, namely the case of an affine patch of the form $\mathbb{C} \times \mathbb{R} \hookrightarrow C(\mathbb{R}^{p,q})$ with one holomorphic dimension and one topological dimension. Let us use z for a holomorphic coordinate on \mathbb{C} and t for a coordinate on \mathbb{R} . Suppose we find a twisting supercharge \mathcal{Q} satisfying the following conditions:

- (1) The group of isometries of $\mathbb{C} \times \mathbb{R}$ should be closed (i.e. contained in $Z_{\mathcal{O}}$).
- (2) The complexified translation $\partial/\partial \overline{z}$ in the antiholomorphic direction of \mathbb{C} and the translation d/dt in the topological direction should both be exact (i.e. contained in $\mathfrak{b}_{O}^{\mathbb{C}}$).

We expect then — under the function-analytic conditions of Theorem 7.15 in the \mathbb{C} direction — that the local observables in the Q-twisted theory will have the structure of a *raviolo vertex algebra* in the sense recently described by Garner and Williams [GW23].

3.4. **Connections to the Physics Literature.** Twists of supersymmetric theories have been studied for a long time. One of the first examples of twists, giving rise to "topological" theories were introduced with the work of Witten [Wit88], which, as we mentioned, ignited the quantum field theoretic interpretation of Donaldson invariants of 4-manifolds. The simplest supersymmetric twists on \mathbb{R}^4 by a single nilpotent supercharge Q gives rise to a purely topological theory, since all the translation generators P^{μ} are in the exact subalgebra \mathfrak{b}_Q .

In superconformal theories, there are more complicated examples, identified by classes of square-zero elements Q in the superconformal algebra. These can be general combinations of ordinary and conformal supercharges. Thus, there is a more complicated structures of Q-closed and Q-exact generators in the even part $\mathfrak{sconf}(n|S_{\mathbb{C}},\mathbb{C})_0$, giving rise to more interesting subvarieties where the protected sector of operators lives.

Proposition 3.15. *The type of generators that appear in the closed and exact subalgebras determine how the twisted theory will look like and behave.*

A conformal twist of 4d $\mathcal{N}=2$ superconformal theories, described in [Bee+15b] has been the launching pad for several works works looking at algebraic structures corresponding to protected sectors of various superconformal theories. In this example, the nilpotent element is a combination of a super Poincaré charge and a superconformal one, so it can only exists in superconformal theories (but not in supersymmetric ones) The structure of the closed and exact subalgebras implies that only some of the translations are closed and/or exact, giving rise to an interesting "mixed" structure, where local operators from the 4d theory are restricted to a 2d complex plane inside space-time. The holomorphic coordinate on this plane corresponds to a closed (complex) translation, whereas the antiholomorphic coordinate, together with one of the real transverse translations, are in the exact subalgebra. The quotient $\mathfrak{z}_Q/\mathfrak{b}_Q$ is an $\mathfrak{s}\mathfrak{l}_2$ algebra, that is then extended to a Virasoro algebra by identifying a candidate stress-energy tensor, whose modes L_m give rise to an infinite chiral symmetry algebra. The restriction of local operators to the complex plane and the action of this algebra provides a correspondence between (protected sectors of) local operators of $4d\mathcal{N}=2$ SCFTs and 2d vertex operator algebras.

Since this seminal work, the standard lore in the community has been that the correspondence between $4d \ N=2$ SCFTs and vertex operator algebras is one-to-one, modulo restricting to only local operators [ALW23] or (not) allowing further restrictions of the spectrum [BJ23].

Remark 3.16. If one considers extended operators in this setting, it is also important to consider that these operators can stretch to infinity. In this case the signature of the underlying space-time may have a considerable impact on the behavior of the twisted theory, compared to the case when one only considers local operators [ALW23]. We will not focus on these subtle features in the present paper and work as much as possible with complex algebras and the largest embedding spaces possible. It is worth noting, however, that when writing explicitly a physical theory, these subtleties ought to be addressed.

Table 1 provides a sample of the conformal twists that have appeared in the physics literature. We exclude those twists of SCFTs that are generated by nilpotent elements constructed purely from Poincaré supercharges, since these are better known and have already been classified (for a similar table of examples in the super Poincaré case, see Tables 1–4 of [ESW22]).

Bulk theory	Name	Geometry of the twist	References
$3d \mathcal{N} = 4 \text{ on}$	1d protected associative algebra	$\mathbb{R} imes \mathbb{R}^2$	[BPR17b]
$4d \ \mathcal{N} = 2 \ \text{on} \ \mathbb{R}^4$	2d Vertex Operator Algebra	$\mathbb{C}\sim\mathbb{R}^2\subseteq\mathbb{R}^4$	[Bee+15b]
4d $\mathcal{N} = 2$ on $\mathbb{R}^{1,3}$	2d Extended VOA	$\mathbb{C}\sim\mathbb{R}^2\subseteq\mathbb{R}^{1,3}$	[ALW23]
6d $\mathcal{N} = (2,0)$ on \mathbb{R}^6	2d VOA	$\mathbb{C} \sim \mathbb{R}^2 \subseteq \mathbb{R}^6$	[BRR15]

TABLE 1. Occurrences of superconformal twists of superconformal theories in the prior literature, classified by the amount of superconformal symmetry and the class of superconformal supercharge by which the twisting is performed.

Remark 3.17. Notice that the conformal twists appearing in the table are all studied in the context of the algebra of operators after restriction to either a real line \mathbb{R} or a complex line \mathbb{C} . From the point of view taken in this paper there are also interesting structures to investigate on the operators of subspaces of larger dimension, for instance the structure of an \mathbb{E}_k algebra on the operators on a topological real *k*-plane.

4. 4D SUPERCONFORMAL TWISTS

In this section we explore the 4-dimensional case. We begin by describing each complex nilpotence variety and then analyze its orbits under the action of the associated complex superconformal group, including the closed and exact transformations for many twists. These results are available in [DS05] but we give the arguments here, both to demonstrate how concrete and accessible they are and also to describe them in a way convenient for later results.

Note, however, that the case of the N=4 superconformal algebra is not included in Theorem 2.21, it provides some novel twists of N=4 super Yang–Mills theory that might be fruitful to explore.

We then turn to analyzing the orbits under the action of real forms of the superconformal groups, which depend upon the signature of the spacetime. For these results we leverage work of Fels, Huckleberry, and Wolf.

4.1. **Complex Nilpotence Variety for** $N \neq 4$ **.** Consider the complexified 4d N = k superconformal algebra $\mathfrak{sconf}(4|k,\mathbb{C})$ for $k \neq 4$. (The k = 4 case is distinct, and we discuss in the next subsection.) The superconformal algebra can be identified with the simple super Lie algebra $\mathfrak{sl}(4|k)$. The even part of $\mathfrak{sconf}(4|k,\mathbb{C})$ is isomorphic to

$$(\mathfrak{sconf}(4|k,\mathbb{C}))_0 \cong \mathfrak{sl}(4,\mathbb{C}) \oplus \mathfrak{sl}(k,\mathbb{C}) \oplus \mathbb{C}$$
$$\cong \mathfrak{so}(6,\mathbb{C}) \oplus \mathfrak{sl}(k,\mathbb{C}) \oplus \mathbb{C}$$

where we use the exceptional isomorphism between $\mathfrak{sl}(4, \mathbb{C})$ and $\mathfrak{so}(6, \mathbb{C})$. The odd part of $\mathfrak{sconf}(4|k, \mathbb{C})$ is isomorphic to

$$(\mathfrak{sconf}(4|k,\mathbb{C}))_1 \cong V_4 \otimes W_k \oplus V_4^* \otimes W_k^*$$
$$\cong S_+ \otimes W_k \oplus S_- \otimes W_k^*$$

where W_k denotes the defining *k*-dimensional representation of the factor $\mathfrak{sl}(k, \mathbb{C})$ and V_4 denotes the four-dimensional defining representation of $\mathfrak{sl}(4, \mathbb{C})$, which is isomorphic to the positive

semispin (or Weyl spinor) representation S_+ of $\mathfrak{so}(6, \mathbb{C})$. The one-dimensional factor in the center of $(\mathfrak{sconf}(4|k, \mathbb{C}))_0$ acts trivially.

The Lie bracket of even elements is given by the direct sum of the natural brackets, and the even part acts on the odd part as determined by the representation structure just stated. It remains to describe the bracket between odd elements in explicit terms.

For any *k*, there is a $\mathfrak{sl}(k, \mathbb{C})$ -invariant decomposition

$$\hom(W_k, W_k) \cong W_k \otimes W_k^* \cong \mathfrak{sl}(k, \mathbb{C}) \oplus \mathbb{C}.$$

Denote the linear projections onto the two factors by

Tr:
$$W_k \otimes W_k^* \to \mathbb{C}$$
,

which correspond to matrix operation of trace $M \mapsto Tr(M)$, and

red:
$$W_k \otimes W_k^* \to \mathfrak{sl}(k, \mathbb{C})$$
,

which corresponds to "making a matrix trace-free" $M \mapsto M - (\text{Tr}(M)/k)$ Id. There is an analogous $\mathfrak{sl}(k, \mathbb{C})$ -invariant decomposition involving V_4 .

These decompositions lead to a convenient description of the bracket between odd elements.

Lemma 4.1. Under the isomorphism

$$V_4 \otimes W_k \oplus V_4^* \otimes W_k^* \cong \hom(V_4^*, W_k) \oplus \hom(W_k, V_4^*)$$

expressing an odd element Q as sum of matrices $Q_+ + Q_-$, the bracket is

$$\begin{split} [\mathcal{Q}, \mathcal{Q}'] &= [\mathcal{Q}_+ + \mathcal{Q}_-, \mathcal{Q}'_+ + \mathcal{Q}_-] \\ &= \operatorname{red}(\mathcal{Q}_- \circ \mathcal{Q}'_+ + \mathcal{Q}'_- \circ \mathcal{Q}_+) + \operatorname{red}(\mathcal{Q}_+ \circ \mathcal{Q}'_- + \mathcal{Q}'_+ \circ \mathcal{Q}_-) \\ &+ (\operatorname{Tr}((\mathcal{Q}_- \circ \mathcal{Q}'_+ + \mathcal{Q}'_- \circ \mathcal{Q}_+) + \operatorname{Tr}(\mathcal{Q}_+ \circ \mathcal{Q}'_- + \mathcal{Q}'_+ \circ \mathcal{Q}_-)) \end{split}$$

in $\mathfrak{sl}(4,\mathbb{C})\oplus\mathfrak{sl}(k,\mathbb{C})\oplus\mathbb{C}$.

Proof. Any odd element decomposes into a sum of terms from each summand. A pure tensor in the first summand has the form $Q_+ \otimes w_+$ with $Q_+ \in S_+$ and $w_+ \in W_k$. A pure tensor in the second summand has the form $Q_- \otimes w_-$ with $Q_+ \in S_+$ and $w_- \in W_k^*$. Pairs of elements in the same summand of $(\mathfrak{sconf}(4|k,\mathbb{C}))_1$ have trivial bracket. The Lie bracket between a pair of elements in opposite summands $[Q_+ \otimes w_+, Q'_- \otimes w'_-]$ is given by

$$\operatorname{red}(Q_+\otimes Q'_-)\operatorname{Tr}(w_+\otimes w'-) + \operatorname{Tr}(Q_+\otimes Q'_-)\operatorname{red}(w_+\otimes w'_-) + \operatorname{Tr}(Q_+\otimes Q'_-)\operatorname{Tr}(w_+\otimes w'_-).$$

It is then direct to verify the two descriptions of the bracket match under the isomorphism. \Box

It is now straightforward to compute the superconformal nilpotence variety.

Proposition 4.2. The complex nilpotence variety $\operatorname{Nilp}_{(4|k)}(\mathbb{C})$ for the 4d $\mathcal{N} = k$ superconformal algebra with $k \neq 4$ is

$$\{\mathcal{Q}_+ + \mathcal{Q}_- \in \hom(V_4^*, W_k) \oplus \hom(W_k, V_4^*) \colon \mathcal{Q}_+ \circ \mathcal{Q}_- = 0 \text{ and } \mathcal{Q}_- \circ \mathcal{Q}_+ = 0\}.$$
²³

Proof. In terms of the linear maps Q_+ , Q_- we have

$$[\mathcal{Q},\mathcal{Q}]=2[\mathcal{Q}_+,\mathcal{Q}_-],$$

which equals zero if

$$\operatorname{red}(\mathcal{Q}_{-}\circ\mathcal{Q}_{+}) = 0 \in \mathfrak{sl}(V_{4}^{*})$$

 $\operatorname{red}(\mathcal{Q}_{+}\circ\mathcal{Q}_{-}) = 0 \in \mathfrak{sl}(W_{k})$
 $\operatorname{Tr}(\mathcal{Q}_{-}\circ\mathcal{Q}_{+}) + \operatorname{Tr}(\mathcal{Q}_{+}\circ\mathcal{Q}_{-}) = 0 \in \mathbb{C}.$

The first two conditions require that $Q_- \circ Q_+$ and $Q_+ \circ Q_-$ are both multiples of an identity matrix, and the last condition requires the sum of their traces to be zero.

Now we use the condition that $k \neq 4$. This tells us that either $Q_- \circ Q_+$ has rank less than 4 (if k < 4) or $Q_+ \circ Q_-$ has rank less than k (if k > 4). In either case, the only scaled identity matrix with less than full rank is the zero matrix, so we must have $\text{Tr}(Q_- \circ Q_+) = -\text{Tr}(Q_+ \circ Q_-) = 0$, and hence $Q_- \circ Q_+$ and $Q_+ \circ Q_-$ both equal zero.

4.2. The $\mathcal{N} = 4$ Case. The 4d $\mathcal{N} = 4$ superconformal algebra is slightly different, because the super Lie algebra $\mathfrak{sl}(4|4)$ is not simple: it has a one-dimensional center generated by the element diag(1, 1, 1, 1|1, 1, 1, 1). In order to obtain a *simple* super Lie algebra, we should take the quotient by this element to obtain the super Lie algebra $\mathfrak{psl}(4|4)$. Thus, we identify the 4d $\mathcal{N} = 4$ complex superconformal algebra $\mathfrak{sconf}(4|4, \mathbb{C})$ as $\mathfrak{psl}(4|4)$.

Remark 4.3. If one is willing to work with non-simple superconformal algebras, then one can consider $\mathfrak{sl}(4|4)$, which we treated in the preceding section.

The even part of $\mathfrak{sconf}(4|4,\mathbb{C})$ can be identified with

$$(\mathfrak{sconf}(4|4,\mathbb{C}))_0 \cong \mathfrak{sl}(4,\mathbb{C}) \oplus \mathfrak{sl}(4,\mathbb{C}) \cong \mathfrak{so}(6,\mathbb{C}) \oplus \mathfrak{sl}(4,\mathbb{C}).$$

The odd part of $\mathfrak{sconf}(4|4, \mathbb{C})$ can be identified with

$$(\mathfrak{sconf}(4|k,\mathbb{C}))_1\cong V_4\otimes W_4\oplus V_4^*\otimes W_4^*\cong S_+\otimes W_4\oplus S_-\otimes W_4^*$$

identically to the description from the previous subsection. The Lie bracket between odd elements is similar to the $N \neq 4$ case, except that we do not need the sum of trace term.

Proposition 4.4. The complex nilpotence variety $\operatorname{Nilp}_{(4|4)}(\mathbb{C})$ for the 4d $\mathcal{N} = 4$ superconformal algebra *is*

$$C_{\mathrm{Tr}} \cup_{\{0\}} C_{\mathrm{red}},$$

the union of the two cones at the origin where

$$C_{\mathrm{Tr}} = \{Q_+ \otimes w_+ + Q_- \otimes w_- : \mathrm{Tr}(Q_+ \otimes Q_-) = \mathrm{Tr}(w_+ \otimes w_-) = 0\}$$

and

$$C_{\text{red}} = \{Q_+ \otimes w_+ + Q_- \otimes w_- : \operatorname{red}(Q_+ \otimes Q_-) = \operatorname{red}(w_+ \otimes w_-) = 0\}.$$

Proof. The right-hand side is contained in the left-hand side, this is immediate from the description of the Lie bracket. If $Q = Q_+ \otimes w_+ + Q_- \otimes w_-$ is a square zero element then $\text{Tr}(Q_+ \otimes Q_-)$ if and only if $\text{Tr}(w_+ \otimes w_-) = 0$ by the same argument as Proposition 4.2. If neither trace term equals zero and [Q, Q] = 0 then we must have $\text{red}(Q_+ \otimes Q_-) = \text{red}(w_+ \otimes w_-) = 0$.

Remark 4.5. Let us discuss the subspace of this nilpotence variety consisting of nilpotent elements in the super Poincaré algebra associated to a choice of embedding $\mathfrak{so}(4, \mathbb{C}) \subseteq \mathfrak{so}(6, \mathbb{C})$. Under this embedding the representation V_4 decomposes as $S_+ \oplus S_-$, where S_{\pm} are the two two-complexdimensional semispin representations of $\mathfrak{so}(4, \mathbb{C})$. The complex nilpotence variety for the $\mathcal{N} = 4$ super Poincaré algebra is isomorphic to

$$\operatorname{Nilp}_{(4|4)}^{\mathfrak{siso}}(\mathbb{C}) = \{Q'_{+} \otimes w_{+} + Q'_{-} \otimes w_{-} \colon \operatorname{Tr}(w_{+} \otimes w_{-}) = 0\} \subseteq C_{\operatorname{Tr}}$$

where $Q'_+ \in S_+ \subseteq V_4$ and $Q'_- \in S_- \cong S^*_- \subseteq V^*_4$. Note that the condition $\text{Tr}(Q'_+ \otimes Q'_-) = 0$ is automatically satisfied. The decomposition of this super Poincaré nilpotence variety into $\text{Spin}(4, \mathbb{C}) \times \text{SL}(4, \mathbb{C})$ -orbits is interesting, including for instance the one-parameter families of topological supercharges studied by Kapustin and Witten [KW07]. This decomposition is discussed in [ES19, Section 4.4] and [ESW21, Section 9].

4.3. **Complex Group Orbits.** The complex superconformal group is isomorphic to $SL(4|k, \mathbb{C})$ if $k \neq 4$, with even part isomorphic to

$$\operatorname{SConf}(4|k,\mathbb{C})_0 \cong \{(A,B) \in \operatorname{GL}(4,\mathbb{C}) \times \operatorname{GL}(k,\mathbb{C}) \colon \det(A) \det(B) = 1\}.$$

It might be helpful to see that there is a group map

$$SL(4,\mathbb{C}) \times SL(k,\mathbb{C}) \times \mathbb{C}^{\times} \to SConf(4|k,\mathbb{C})_0$$

sending (M, N, λ) to $(\lambda M, \lambda^{-1}N)$.

Remark 4.6. This map is locally a diffeomorphism. Indeed, working in a neighborhood of matrices (A, B) where det(A) lies in a small disk in \mathbb{C}^{\times} , and choose a local fourth root in this disk, we can define a local inverse sending (A, B) to $(A / \det(A), B / \det(B), \det(A)^{1/4})$.

In the k = 4 case the superconformal group is identified with $PSL(4|4, \mathbb{C})$, whose even part is isomorphic to $SL(4, \mathbb{C}) \times SL(4, \mathbb{C})$.

These super groups act on the super Lie algebra $\mathfrak{sconf}(4|k,\mathbb{C})$ by the adjoint action. In explicit terms, an element $(A, B, \lambda) \in SL(4, \mathbb{C}) \times SL(k, \mathbb{C}) \times \mathbb{C}^{\times}$ acts on

$$(\mathfrak{sconf}(4|k,\mathbb{C}))_1 \cong \hom(W_k^*,V_4) \oplus \hom(W_k,V_4^*)$$

by

$$(\mathcal{Q}_+, \mathcal{Q}_-) \mapsto (\lambda^{-2}B\mathcal{Q}_+A^{-1}, \lambda^2 A\mathcal{Q}_-B^{-1}).$$

This action manifestly preserves the *rank* r_{\pm} of each operator Q_{\pm} , and for $k \neq 4$, these invariants classify the orbits in Nilp_(4|k)(\mathbb{C}).

Proposition 4.7. *For* $k \neq 4$ *, there is a bijection*

$$\operatorname{Nilp}_{(4|k)}(\mathbb{C})/\operatorname{SConf}(4|k,\mathbb{C})_0 \cong \{(r_+,r_-) \in \mathbb{N}^2 : r_+ + r_- \le \min(4,k)\},\$$

where a twist (Q_+, Q_-) maps to $(\operatorname{rank}(Q_+), \operatorname{rank}(Q_-))$.

That is, the orbits of $\operatorname{Nilp}_{(4|k)}(\mathbb{C})$ are in bijection with pairs (r_+, r_-) of non-negative integers where $r_+ + r_- \leq \min(4, k)$. Note that we use $\operatorname{Nilp}_{(4|k)}(\mathbb{C})/\operatorname{SConf}(4|k, \mathbb{C})_0$ to denote the quotient set, in contrast to the quotient stack. (The quotient topology is not discrete (and so non-Hausdorff): it is given by the poset structure, as having rank $\geq k$ is an open condition.)

This result suggests the following notation.

Definition 4.8. Let $\mathcal{O}(4|k)_{r_+,r_-}$ denote the complex group orbit associated to a pair (r_+,r_-) of ranks.

Proof. Given a square-zero supercharge (Q_+, Q_-) , we assign to it a pair of non-negative integers by letting $r_{\pm} = \operatorname{rank}(Q_{\pm})$. Since $\operatorname{Im}(Q_+) \subseteq \ker(Q_-)$, the rank-nullity theorem tells us $r_+ \leq k - r_-$ so we have $r_+ + r_- \leq k$. Likewise, since $\operatorname{Im}(Q_-) \subseteq \ker(Q_+)$, we have $r_+ + r_- \leq 4$.

Now, we may choose bases (or apply row/column operations) for V_4^* and W_k so that Q_+ and Q_- are represented by matrices of the following form:

(1)
$$\mathcal{Q}_{+} = \begin{pmatrix} \mathrm{id}_{r_{+}} & 0 \\ 0 & 0 \end{pmatrix}, \ \mathcal{Q}_{-} = \begin{pmatrix} 0 & 0 \\ 0 & \mathrm{id}_{r_{-}} \end{pmatrix}$$

Note that these are not square matrices: we write 0 to refer to rectangular zero matrices of appropriate sizes.

In other words, the action of $GL(4, \mathbb{C}) \times GL(k, \mathbb{C})$ on the locus of square-zero supercharges of ranks (r_+, r_-) is transitive. To complete the proof, we just need to ensure that we can choose our two change of basis matrices to have the same determinant. Alternatively, it is enough to find an element $(X, Y) \in GL(4, \mathbb{C}) \times GL(k, \mathbb{C})$ stabilizing the diagonal matrices 1 where $\det(X) / \det(Y) = \lambda \in \mathbb{C}^{\times}$ for all values of λ .

To do this, again use the fact that $k \neq 4$. If k < 4 then $r_+ + r_- < 4$, and we let $X = \text{diag}(1, ..., 1, \lambda, 1, ..., 1)$ with λ in the $r_+ + 1^{\text{st}}$ position and $Y = \text{id}_k$. Likewise if k > 4 then $r_+ + r_- < k$, and we let $Y = \text{diag}(1, ..., 1, \lambda^{-1}, 1, ..., 1)$ with λ^{-1} in the $r_+ + 1^{\text{st}}$ position, and $X = \text{id}_4$.

Something interesting happens, however, when k = 4: the orbits are not a finite set, but instead come in continuous families. In other words, this case does *not* sit among the examples covered by Theorem 2.21 of Duflo and Serganova.

Proposition 4.9. There is an isomorphism

$$\operatorname{Nilp}_{(4|4)}(\mathbb{C})/\operatorname{SConf}(4|4,\mathbb{C})_0 \cong \{(r_+,r_-) \in \mathbb{Z}_+^2 : r_++r_- \leq 4\} \sqcup \mathbb{C}^{\times} \sqcup \mathbb{C}^{\times}.$$

When both terms Q_{\pm} have positive rank, the orbit of the twist (Q_+, Q_-) is an isolated point. When one term has rank 0 and the other has rank 4 (i.e., $(r_+, r_-) = (4, 0)$ or (0, 4)), then the orbit sits in a one-parameter family labeled by the determinant of the full rank term (in the (4, 0) case, det (Q_+)).

We will use $\mathcal{O}(4|4)_{r_+,r_-}$ to denote these components, but $\mathcal{O}(4|4)_{4,0} \cong \mathbb{C}^{\times}$, for instance. Note that the origin in a copy of \mathbb{C}^{\times} corresponds to the limit of a family where the determinant vanishes and hence the rank drops. Thus one can view the (4,0) case (or (0,4) case) as \mathbb{C} along with some points added that "thicken" the origin.

Proof. The argument in the k = 4 case carries over for matrices with positive ranks. In the case of ranks (4,0), the first matrix Q_+ is full rank, but the action of $SL(4, \mathbb{C})$ from the left (respectively, right) does not change the determinant. On the other hand, these actions by $SL(4, \mathbb{C}) \times SL(4, \mathbb{C})$ by row and column operations can be used to put Q_+ into a diagonal form with three entries of 1s and a final entry $det(Q_+)$.

4.4. **Closed and Exact Transformations.** Let us discuss some specific properties of the complex orbits in some of the low rank cases.

- (1) Rank (0,0): The zero supercharge always squares to zero. If *k* < 2 it is the only point in Nilp_(4|k)(ℂ). The kernel of [*Q*, −] is so(6, ℂ) ⊕ sl(*k*, ℂ) ⊕ ℂ (the entire even component), and the image of [*Q*, −] is just {0}.
- (2) Rank (r₊, 0) and (0, r_−): We refer to these as *chiral* supercharges, since either Q₊ or Q_− vanishes. Such supercharges automatically square to zero, so these strata in Nilp_(4|k)(C) are isomorphic to V₄ ⊗ W₄\{0} and V₄^{*} ⊗ W₄^{*}\{0}, respectively.

The stabilizer $Z_Q \subseteq SL(4, \mathbb{C}) \times SL(k, \mathbb{C}) \times \mathbb{C}^{\times}$ can be described in the following way. Let $P_{r,k} \subseteq SL(k, \mathbb{C})$ be the parabolic subgroup with block diagonal Levi subgroup $SL(r, \mathbb{C}) \times SL(k-r, \mathbb{C})$. Write

$$F_r: P_{r,4} \times P_{r,k} \to \mathrm{SL}(r,\mathbb{C})$$

for the homomorphism given by projection onto $SL(r, \mathbb{C}) \times SL(r, \mathbb{C})$ (i.e., projection onto the upper-left block in each factor) composed with the product map $SL(r, \mathbb{C}) \times SL(r, \mathbb{C}) \rightarrow SL(r, \mathbb{C})$.

Proposition 4.10. *If* Q *is a chiral supercharge of rank* (r, 0)*, then the stabilizer* Z_Q *is isomorphic to* $F_r^{-1}(I) \times \mathbb{C}^{\times}$.

Proof. We can check this by choosing a basis in which Q is represented by a block diagonal matrix of rank *r*. The stabilizer Z_Q is then readily computed by solving

$$\left(\begin{array}{c|c} A_1 & A_2 \\ \hline A_3 & A_4 \end{array}\right) \left(\begin{array}{c|c} I_r & 0 \\ \hline 0 & 0 \end{array}\right) \left(\begin{array}{c|c} B_1 & B_2 \\ \hline B_3 & B_4 \end{array}\right) = \left(\begin{array}{c|c} I_r & 0 \\ \hline 0 & 0 \end{array}\right).$$

Here $A = \begin{pmatrix} A_1 & A_2 \\ \hline A_3 & A_4 \end{pmatrix}$ is an element of SL(4, \mathbb{C}) and $B = \begin{pmatrix} B_1 & B_2 \\ \hline B_3 & B_4 \end{pmatrix}$ is an element of SL(*k*, \mathbb{C}). (For instance, B_1 is an $r \times r$ block, B_2 is an $r \times (k - r)$ block, and so on.) We find $A_1 = B_1^{-1}$ and $A_3 = B_2 = 0$. In other words (*A*, *B*) lies in F_r^{-1} .

In particular, the dimension of Z_Q is given as

$$\dim Z_{Q} = k^{2} - kr + r^{2} - 4r + 14$$

if $r \neq k$ and

$$\dim Z_{\mathcal{O}} = k^2 - kr + r^2 - 4r + 15$$

if r = k.

Similarly, we can describe the Lie algebra $\mathfrak{b}_{\mathcal{Q}} = \operatorname{Im}([\mathcal{Q}, -])$ as a subalgebra of $\mathfrak{z}_{\mathcal{Q}} = \operatorname{Lie}(Z_{\mathcal{Q}})$, by a similar direct computation. We find the following.

Proposition 4.11. Let Q be a chiral supercharge of rank r. The Lie algebra

$$\mathfrak{b}_{\mathcal{Q}} \subseteq \mathfrak{sl}(4,\mathbb{C}) \oplus \mathfrak{sl}(k,\mathbb{C}) \subseteq \mathfrak{sl}(4,\mathbb{C}) \oplus \mathfrak{sl}(k,\mathbb{C}) \oplus \mathbb{C}$$

is isomorphic to the Lie algebra of block matrices of the following form:

$$\mathfrak{b}_{r} = \left\{ \left(\left(\begin{array}{c|c} \alpha & \beta \\ \hline 0 & 0 \end{array} \right), \left(\begin{array}{c|c} \alpha & 0 \\ \hline \gamma & 0 \end{array} \right), \lambda \right) \in \mathfrak{sl}(4, \mathbb{C}) \oplus \mathfrak{sl}(k, \mathbb{C}) \oplus \mathbb{C} \colon \\ \alpha \in \mathfrak{sl}(r, \mathbb{C}), \beta \in \operatorname{Mat}_{r \times (4-r)}(\mathbb{C}), \gamma \in \operatorname{Mat}_{(k-r) \times r}(\mathbb{C}) \right\}.$$

In particular, we have

$$\dim \mathfrak{b}_{\mathcal{O}} = kr - r^2 + 4r.$$

For reference, we give a tabular description of the dependence of the dimension of \mathfrak{z}_{Q} and \mathfrak{b}_{Q} on r. Recall that $r \leq \min(4, k)$. Thus the dimensions for each k, in terms of r, are given in Table 2.

r	$\dim\mathfrak{z}_\mathcal{Q}$	$\dim \mathfrak{b}_\mathcal{Q}$	$\dim \mathfrak{z}_Q/\mathfrak{b}_Q$
1	$k^2 - k + 11 + \delta_{kr} - \delta_{k4}$	k + 3	$k^2 - 2k + 8 + \delta_{kr} - \delta_{k4}$
2	$k^2 - 2k + 10 + \delta_{kr} - \delta_{k4}$	2k + 4	$k^2 - 4k + 6 + \delta_{kr} - \delta_{k4}$
3	$k^2 - 3k + 11 + \delta_{kr} - \delta_{k4}$	3k + 3	$k^2 - 6k + 8 + \delta_{kr} - \delta_{k4}$
4	$k^2 - 4k + 14 + \delta_{kr} - \delta_{k4}$	4k	$k^2 - 8k + 14 + \delta_{kr} - \delta_{k4}$

TABLE 2. For chiral supercharges Q of rank r, we give the dimension of the kernel and image of [Q, -] as well as the dimension of the quotient.

We note from this computation that the inclusion $\mathfrak{z}_Q \subseteq \mathfrak{b}_Q$ is always proper, at least for $k \neq 4$. In the $k = \mathcal{N} = 4$ case the computation remains the same, but $\dim \mathfrak{z}_Q$ is decreased by one. The inclusion $\mathfrak{z}_Q \subseteq \mathfrak{b}_Q$ is still proper, with the exception of the maximal rank r = 4 case where $\mathfrak{b}_Q = \mathfrak{z}_Q$ and they span the diagonal copy of $\mathfrak{sl}(4, \mathbb{C}) \subseteq \mathfrak{sl}(4, \mathbb{C}) \oplus \mathfrak{sl}(4, \mathbb{C})$.

(3) Rank (1,1): For k > 1, let Q be a square zero supercharge of the form Q₊ ⊗ w₊ + Q₋ ⊗ w₋. The square-zero condition in this case is equivalent to the conditions Tr(w₊, w₋) = 0 = Tr(Q₊, Q₋). This determines an analysis of the stabilizer and the image of a super-charge similar to what we have done already.

Proposition 4.12. Let $Q = Q_+ \otimes w_+ + Q_- \otimes w_-$ be a rank (1, 1) square zero supercharge. The image $\mathfrak{b}_Q \subseteq \mathfrak{sl}(4, \mathbb{C}) \times \mathfrak{sl}(k, \mathbb{C}) \times \mathbb{C}$ of the linear map [Q, -] is isomorphic to

$$\mathfrak{b}_{\mathcal{O}} = \{(A, B) \in M_{21} \times M_{12} \colon \operatorname{Tr}(A) + \operatorname{Tr}(B) = 0\},\$$

where for $i \neq j$, $M_{ij} \subseteq \mathfrak{gl}(n, \mathbb{C})$ is the Lie subalgebra of matrices whose non-zero entries all lie in row *i* or column *j*.

Proposition 4.13. Let $Q = Q_+ \otimes w_+ + Q_- \otimes w_-$ be a rank (1, 1) square zero supercharge. The kernel \mathfrak{z}_Q is the Lie algebra of the stabilizer subgroup $Z_Q \subseteq SL(4,\mathbb{C}) \times SL(k,\mathbb{C}) \times \mathbb{C}^{\times}$, whose

elements are given in block form as

$$\left\{ \left(\begin{pmatrix} a & 0 & 0 & 0 \\ * & b & * & * \\ * & 0 & * & * \\ * & 0 & * & * \end{pmatrix}, \begin{pmatrix} a^{-1} & * & * & \cdots & * \\ 0 & b^{-1} & 0 & \cdots & 0 \\ 0 & * & * & \cdots & * \\ \vdots & \vdots & & & \vdots \\ 0 & * & * & \cdots & * \end{pmatrix}, \lambda \right\} : a, b \in \mathbb{C}^{\times} \right\}$$

One can check both propositions using the same setup. If $\{Q_1, Q_2, Q_3, Q_4\}$ and $\{w_1, \ldots, w_k\}$ are bases for S_+ and W, respectively, without loss of generality we may let $Q = Q_1 \otimes w_1 + Q_2^* \otimes w_2^*$. In this basis Q is represented by a $k \times 4$ matrix and a $4 \times k$ matrix with ones in the (1, 1) and (2, 2) positions respectively, and zeroes in all other positions. It is then an elementary computation to work out the image and kernel of the operator [Q, -].

Remark 4.14. Of course the classification does not need to end here: there are conformal twists of higher rank, including twists of rank (2, 2) generalizing the family of twists in the $\mathcal{N} = 4$ super Poincaré algebra studied by Kapustin and Witten [KW07]. We leave a detailed analysis of all cases for future work.

4.5. **Real Group Orbits: The Case of a Chiral Twist.** We now discuss the orbit structure of superconformal twisted theories under the action of real superconformal groups in various signatures. A key ingredient is to recognize that the complex orbits are fiber bundles of a nice kind. We begin by explaining this idea, and its power for studying real orbits, in the simplest example before gesturing at some general machinery. We stick to the chiral twists as the situation is already complicated there.

Recall that $\mathcal{O}(4|k)_{r_+,r_-}$ denotes the complex orbit of rank (r_+, r_-) . We will focus on *chiral* twists, so without loss of generality, set $r_- = 0$, use r to denote r_+ , and consider $\mathcal{O}(4|k)_{r,0}$.

Example 4.15. Let's start with the case that k = 2 and r = 1. (A moment's thought shows r = 0 is too simple.) There is a smooth map

$$\pi\colon \mathcal{O}(4|2)_{1,0}\to \operatorname{Gr}(1,2)\cong \mathbb{CP}^1$$

sending $Q: \mathbb{C}^2 \to \mathbb{C}^4$ to ker Q. An element (A, B) in

$$\operatorname{SConf}(4|2,\mathbb{C})_0 \cong \{(A,B) \in \operatorname{GL}(4,\mathbb{C}) \times \operatorname{GL}(2,\mathbb{C}) \colon \det(A) \det(B) = 1\}$$

acts on Q by AQB. There is a natural action of the factor $SL(2, \mathbb{C})$ on \mathbb{CP}^1 by Möbius transformations, so this map π is equivariant for the action of $SConf(4|2, \mathbb{C})_0$ since

$$\pi(A\mathcal{Q}B) = \ker(A\mathcal{Q}B) = \ker(\mathcal{Q}B) = \ker(\mathcal{Q}) \cdot B,$$

where on the far right, we use the natural action of $B \in GL(2, \mathbb{C})$ on the projective line. The action of the factor $A \in SL(4, \mathbb{C})$ preserves the fiber $\pi^{-1}(L)$ for each line $L \subset \mathbb{C}^2$, since AQB has the same kernel as Q. (The image of Q changes, however, under the action of $SConf(4|2, \mathbb{C})_0$.)

When we pick a real form $G_{\mathbb{R}}$ of $\mathrm{SConf}(4|2,\mathbb{C})_0$, we can analyze its action on $\mathcal{O}(4|2)_{1,0}$ by studying how $G_{\mathbb{R}} \cap \mathrm{SL}(2,\mathbb{C})$ acts on the base space \mathbb{CP}^1 and how $G_{\mathbb{R}} \cap \mathrm{SL}(4,\mathbb{C})$ acts on each fiber $\pi^{-1}(L)$ where *L* is in the image of π . For instance, recall that the real subgroup $\mathrm{SL}(2,\mathbb{R}) \subset$ $\mathrm{SL}(2,\mathbb{C})$ breaks \mathbb{CP}^1 into three orbits: the unit circle (or equator), the open hemisphere containing 0, and the open hemisphere containing ∞ . By contrast, the real subgroup $SU(2) \subset SL(2, \mathbb{C})$ has a single orbit on the base.

This mode of analysis generalizes nicely. When $k \ge 4$, it is perhaps more convenient to view Q as an element of hom(\mathbb{C}^4 , \mathbb{C}^k), so we take that perspective now. Thanks to our observations above, we have the following.

Proposition 4.16. For each rank $r \leq 4$, there is a smooth map

$$\pi\colon \mathcal{O}(4|k)_{r,0}\to \operatorname{Gr}(4-r,4)$$

sending Q to ker(Q). In terms of the map π , the group SConf($4|k, \mathbb{C}$) acts on the base through the SL($4, \mathbb{C}$) factor, and it acts on the fiber through the SL(k, \mathbb{C}) factor.

In words, the image of π is through the quotient $\mathcal{O}(4|k)_{r,0}/SL(k,\mathbb{C})$. We may also observe that π determines a fiber bundle in the base by the following computation.

Lemma 4.17. The fibers of π are all isomorphic as $SL(k, \mathbb{C})$ -spaces.

Proof. Given $X, Y \in Gr(4 - r, 4)$, choose $B \in GL(4, \mathbb{C})$ so that Y = BX. There is a natural isomorphism $\phi \colon \pi^{-1}(Y) \to \pi^{-1}(X)$ by $\phi(Q) = QB$. This map is equivariant for the left $SL(k, \mathbb{C})$ -action.

Example 4.18. In the case where k = 2, r = 1 discussed in Example 4.15 above we realized $\mathbb{O}(4|2)_{(1,0)}$ as a fiber bundle over \mathbb{CP}^1 with fiber isomorphic to the space of rank one 2×4 matrices with fixed kernel. We can identify this fiber with $\mathbb{C}^4 \setminus \{0\}$ with its natural action of SL(4, \mathbb{C}). We will discuss the real forms of SConf(4|2, \mathbb{C})₀ in general in a moment, but we may summarize here what happens in each of the three possible signatures for $\mathbb{R}^{p,4-p}$.

- (1) In signature (4,0) we have G_ℝ ∩ SL(4, C) ≅ SL(2, H), and G_ℝ ∩ SL(2, C) ≅ SU(2). The action of SU(2) on CP¹ is transitive, as is the action of SL(2, H) on C⁴ \{0} ≅ H² \{0}. In this case the real orbit space is a single point.
- (2) In signature (3,1) we have G_R ∩ SL(4, C) ≅ SU(2,2), and G_R ∩ SL(2, C) ≅ SU(2). The action of SU(2) on CP¹ is transitive, but the action of SL(2, H) on C⁴\{0} is more interesting: there is a continuous family of orbits parameterized by R[×] (corresponding to non-zero values of the split signature Hermitian form on C^{2,2}), but there is a family of orbits with a finite non-Hausdorff locus (corresponding to the null locus for the Hermitian form). (That is, the quotient space by the SL(2, H)-action is non-Hausdorff.)
- (3) In signature (2, 2) we have G_ℝ ∩ SL(4, C) ≅ SL(4, R), and G_ℝ ∩ SL(2, C) ≅ SL(2, R). The action of SL(2, R) on CP¹ has three orbits: two open orbits (the two open hemispheres) and one closed orbit (the equator), so the quotient space has three points with a connected non-Hausdorff topology in which a single point is dense. The action of SL(4, R) on C⁴\{0} is again interesting, because the quotient space parametrizing orbits is a connected non-Hausdorff space containing an open subspace isomorphic to R[×].

In general, consider a real form of the supergroup $SConf(4|k, \mathbb{C})$ containing the real form SO(p + 1, 5 - p) of the complex 4d conformal group $SO(6, \mathbb{C})$. The real form is labeled by its signature $\mathbb{R}^{p,4-p}$ so we will write $G_{p,4-p}$ for the even part of this real supergroup. Note that

 $G_{p,4-p}$ is a real form of a cover of $SL(4, \mathbb{C}) \times SL(k, \mathbb{C}) \times \mathbb{C}^{\times}$, if $k \neq 4$, or of $SL(4, \mathbb{C}) \times SL(4, \mathbb{C})$, if k = 4. Just as above, we will organize our analysis in terms of $SL(-, \mathbb{C})$ -factors, so let

$$G_{p,4-p}^{\text{base}} = G_{p,4-p} \cap \text{SL}(4,\mathbb{C})$$

and let

$$G_{p,4-p}^{\operatorname{fib}} = G_{p,4-p} \cap (\operatorname{SL}(k,\mathbb{C}) \times \mathbb{C}^{\times})$$

denote the corresponding real factors.

We now discuss the $G_{p,4-p}$ -orbits in the complex nilpotence variety for the possible choices of signature. In short, we want to understand the quotient space

$$\mathcal{O}(4|k)_{r,0}/G_{p,4-p}$$

of orbits for the action of the full group. It is convenient to approach this space in stages using the subgroups we just described. For instance we have maps



and we know that the fibers of π are all isomorphic with actions of $SL(k, \mathbb{C}) \times \mathbb{C}^{\times}$, and hence with actions of its real form $G_{v,4-v}^{\text{fib}}$.

In principle one can work out these orbit decompositions out by hand, but there are quite general results of Fels–Huckleberry–Wolf (going back to Wolf's work from the 1960s) that help here, notably in identifying the lower right corner of the diagram.

Theorem 4.19 ([Wol69, Theorem 2.6], [FHW06, Theorem 3.2.1]). Let *G* be a real form of $SL(n, \mathbb{C})$ of real rank ρ . Let *P* be a parabolic subgroup of $SL(n, \mathbb{C})$.

- There are finitely many G-orbits in the partial flag variety $SL(n, \mathbb{C})/P$.
- If the partial flag variety is a hermitian symmetric space, then the number of orbits is precisely $\binom{\rho+2}{2}$.
- For general partial flag varieties, an upper bound holds: there are at most k|W| orbits, where k is the number of conjugacy classes of Cartan subgroups in G.

In all cases there is a unique closed orbit. *The real dimension of this closed orbit equals the complex dimension (i.e. half the real dimension) of the partial flag variety.*

Note that a Grassmannian is a partial flag variety: $Gr(m, n) = SL(n, \mathbb{C}) / P_{m,n-m}$ where $P_{m,n-m}$ is the parabolic subgroup of block upper triangular matrices where the first diagonal block is $m \times m$ and whose second diagonal block is $(n - m) \times (n - m)$. (These matrices are the stabilizer of the *m*-dimensional subspace spanned by the first *m* elements of the standard basis.)

In our case of interest, n = 4 and m = 4 - r. The theorem implies that there are *finitely* many orbits in the bottom right corner:

$$D_{p,4-p} = \text{Gr}(4-r,4) / G_{p,4-p}^{\text{base}}$$

is a finite set with some interesting connected but non-Hausdorff topology. After identifying this finite set, we then need to understand the fibers of the map

$$\mathcal{O}(4|k)_{r,0}/G_{p,4-p} \to \mathrm{Gr}(4-r,4)/G_{p,4-p}^{\mathrm{base}} = D_{p,4-p}$$

and how $G_{p,4-p}^{\text{fib}}$ acts on them. The fibers are all isomorphic, so we see

$$\mathcal{O}(4|k)_{r,0}/G_{p,4-p} \cong D_{p,4-p} \times (\mathrm{GL}(r_+,\mathbb{C})/G_{p,4-p}^{\mathrm{hb}})$$

as a set.

Remark 4.20. The number *k* appearing in the upper bound for the number of orbits $|D_{*,*}|$ is computed for general real forms of $SL(n, \mathbb{C})$ by Sugiura [Sug59]. In particular we find k = 1 in Euclidean signature and k = 3 in split signature. The remaining Lorentzian signature case is a hermitian symmetric space so we obtain the exact number of orbits as being 6.

We now turn to discussing each signature in turn.

4.5.1. Signature (4, 0). We can identify

$$\mathrm{SConf}(4,0|2\ell,\mathbb{R})\cong\mathrm{SL}(2|\ell,\mathbb{H})$$

where \mathbb{H} denotes the quaternions. In general, when we write $SL(n|m, \mathbb{H})$, it refers to the set of quaternionic matrices such that

$$Nm(sdet(A)) = 1$$

where $\text{Nm}(q) \in \mathbb{R}_{\geq 0}$ denotes the norm of a quaternion q. In particular $\dim_{\mathbb{R}} \text{SL}(n, \mathbb{H}) = 4n^2 - 1$. In particular, the even part of this superconformal group can be identified with

$$\mathrm{SConf}(4,0|2\ell,\mathbb{R})_0 \cong \mathrm{SL}(2,\mathbb{H}) \times \mathrm{GL}(\ell,\mathbb{H}).$$

The real rank of $SL(2, \mathbb{H})$ is one, so in a partial flag variety of the form $SL(4, \mathbb{C})/P$, there are always precisely three orbits for the left-action of $SL(2, \mathbb{H})$.

Example 4.21. Consider the $\mathcal{N} = 2\ell$ superconformal algebra with $2\ell \neq 4$. In the space of chiral supercharges of rank $(r_+, 0)$, the space of orbits in signature (4, 0) can be identified with

$$(\operatorname{GL}(2\ell,\mathbb{C})/(\operatorname{GL}(\ell,\mathbb{H})) \times D)$$

for a finite topological space *D* with $|D| \leq 12$.

Example 4.22. If $2\ell = 2$, we can identify

$$\operatorname{GL}(2,\mathbb{C})/\operatorname{GL}(1,\mathbb{H}) \cong \operatorname{SL}(2,\mathbb{C})/\operatorname{SU}(2) \times \mathbb{C}^{\times}/\mathbb{R}^{\times} \cong \mathfrak{h}^{3} \times S^{1}$$

where \mathfrak{h}^3 refers to hyperbolic 3-space. Thus topologically, the space of orbits of rank (2,0) chiral supercharges in the $\mathcal{N} = 2$ superconformal algebra, in Euclidean signature, can be identified with

$$(\mathbb{R}^3 \times S^1)^{\sqcup m}.$$

4.5.2. *Signature* (3, 1). We can identify

$$SConf(3,1|k,\mathbb{R}) \cong SU(2,2|k).$$

In particular, the even part can be identified with

$$SConf(3,1|k,\mathbb{R})_0 \cong SU(2,2) \times SU(k) \times U(1).$$

The real rank of SU(2,2) is two, so in a partial flag variety of the form $SL(4, \mathbb{C})/P$ there are always at most six orbits for the left-action of SU(2,2).

Example 4.23. In the chiral case with $\mathcal{N} = 2$ the SU(2, 2)-action on the complex Grassmannian Gr(2, 4; \mathbb{C}) has orbits that we can describe quite concretely. They are given by the possible restrictions of the split signature Hermitian metric on $\mathbb{C}^{2,2}$ to a plane. There are six orbits: the restricted (possibly degenerate) Hermitian spaces are

$$\mathbb{C}^{2,0}, \mathbb{C}^{1,1}, \mathbb{C}^{0,2}, \mathbb{C}_{\text{null}} \times \mathbb{C}^{1,0}, \mathbb{C}_{\text{null}} \times \mathbb{C}^{0,1}, (\mathbb{C}_{\text{null}})^2,$$

where \mathbb{C}_{null} refers to \mathbb{C} with a trivial, completely degenerate metric.

4.5.3. *Signature* (2, 2). We can identify

$$\operatorname{SConf}(2,2|k,\mathbb{R}) \cong \operatorname{SL}(4|k,\mathbb{R}).$$

In particular, the even part can be identified with

$$\mathrm{SConf}(2,2|k,\mathbb{R})_0 \cong \mathrm{SL}(4,\mathbb{R}) \times \mathrm{SL}(k,\mathbb{R}) \times \mathbb{R}^{\times}$$

Example 4.24. In the chiral rank 2 case we can again identify the space of orbits. We find the product of a finite non-Hausdorff space and the homogeneous space

$$\operatorname{GL}(2,\mathbb{C})/(\operatorname{SL}(2,\mathbb{R})\times\mathbb{R}^{\times})\cong(\mathbb{R}^{2}\times(S^{1})^{2})$$

where now SL(2, \mathbb{C})/SL(2, \mathbb{R}) $\cong \mathbb{R}^2 \times S^1$ is a 3d hyperboloid of one sheet $\{x_1^2 + x_2^2 - x_3^2 - x_4^2 = 1\} \subseteq \mathbb{R}^4$.

Thus, overall, the set of orbits in this split signature case is identified with $\mathbb{R}^2 \times (S^1)^2 \times D$ for a finite topological space *D*.

4.6. **Case Study:** 4d $\mathcal{N} = 2$ **Chiral Supercharges.** In this section we will analyze the specific example considered by Beem et al [Bee+15b] in their derivation of vertex algebras from $\mathcal{N} = 2$ 4d superconformal field theory. Beem et al are concerned with a chiral supercharge of rank (2,0) that they dub the *twisted Schur operators*. The analysis above tells us that for this example

$$\dim \mathfrak{z}_{\mathcal{Q}}^{\mathbb{C}} = 11$$
$$\dim \mathfrak{b}_{\mathcal{Q}}^{\mathbb{C}} = 8.$$

Let us discuss such chiral supercharges in terms of the usual physics notation. We choose an affine patch $\mathbb{R}^{p,4-p} \hookrightarrow C(\mathbb{R}^{p,4-p})$, associated to which there is an embedding $\mathrm{ISO}(p,4-p) \hookrightarrow \mathrm{Conf}(p,4-p)$. Note that this embedding is non-canonical and so the decomposition that we will discuss below is also non-canonical.

The summands of the odd part of the superconformal algebra decompose as $\mathfrak{so}(4, \mathbb{C}) \subseteq \mathfrak{so}(6, \mathbb{C})$ representations as follows. If $V_4 \otimes W_2$ is the positive chirality piece of $\mathfrak{sconf}(4|2, \mathbb{C})_1$, we have the
decomposition

$$V_4 \otimes W_2 \cong S_+ \otimes W_2 \oplus S_- \otimes W_2$$

where S_{\pm} are the two semispin representations of $\mathfrak{so}(4, \mathbb{C}) \cong \mathfrak{sl}(2, \mathbb{C})_+ \oplus \mathfrak{sl}(2, \mathbb{C})_-$. The convention in physics is to choose bases for these summands: let

$$S_+ \otimes W_2 = \operatorname{Span}(\mathcal{Q}_{\alpha \mathcal{I}}), \quad S_- \otimes W_2 = \operatorname{Span}(\widetilde{\mathcal{S}}^{\dot{\alpha} \mathcal{I}})$$

where $\alpha \in \{+, -\}$, $\dot{\alpha} \in \{\dot{+}, \dot{-}\}$ and $\mathcal{I} \in \{1, 2\}$. Rank (2,0) elements contained in the first summand $S_+ \otimes W_2$ lie in an ordinary 4d $\mathcal{N} = 2$ super Poincaré algebra; this has the nature of the Donaldson–Witten twisting supercharge in [Wit88]. Beem et al [Bee+15b] refer to rank (2,0) elements not lying in either summand in terms of "Schur" supercharges, whose name comes from their connection to the Schur limit of the superconformal index of 4d $\mathcal{N} = 2$ superconformal theories, as Beem et al explain in the introduction and Section 4.4 of *loc. cit.* Following their terminology, a "Schur operator" in a superconformal theory is a local operator representing a non-trivial cohomology class for a fixed Schur supercharge.

4.6.1. *Relationship with Donaldson–Witten supercharges.* As we have seen, all rank (2,0) supercharges Q lie in the same orbit for the complex group SO(6, \mathbb{C}) × Sp(2, \mathbb{C}), and the complex Lie algebras $\mathfrak{z}_Q^{\mathbb{C}}, \mathfrak{b}_Q^{\mathbb{C}}$ are conjugate for different choices of Q. From the complexified point of view there is no distinction between twisting by a Donaldson–Witten supercharge and a Schur supercharge for any specific affine patch. This is not true at the level of a real form (i.e. for the actual group acting on the conformal compactification space in any specified signature), where there is an interesting orbit space as we saw in the preceding subsection.

Indeed, if we choose a specific affine patch $\iota: \mathbb{R}^{p,4-p} \hookrightarrow C(\mathbb{R}^{p,4-p})$, and therefore a subalgebra $\mathfrak{so}(4,\mathbb{C}) \subseteq \mathfrak{so}(6,\mathbb{C})$, as described above, the Schur supercharge does not lie in the same Spin(p, 4-p) orbit as the Donaldson–Witten supercharges. It is worth remarking, however, that this distinction relied on *fixing* an affine patch. On the other hand, given any rank (2,0) supercharge, we may always *choose* an affine patch for which it has Donaldson–Witten type.

Proposition 4.25. Let Q be a rank (2,0) nilpotent supercharge in a real form $\mathfrak{sconf}(p, 4-p|\mathcal{N}=2)$. Then there exists an affine patch $\iota: \mathbb{R}^{p,4-p} \hookrightarrow C(\mathbb{R}^{p,4-p})$ such that Q, viewed in the the complexification $\mathfrak{sconf}(4|\mathcal{N}=2,\mathbb{C})$, is a Donaldson–Witten supercharge, i.e., so that it lies in the span of the generators $Q_{\alpha I}$.

In other words, such a Schur supercharge on one "spacetime" (i.e., on one affine patch) looks like a Donaldson–Witten supercharge after moving to another "spacetime" by a conformal transformation.

Proof. View Q as a rank 2 linear map $V_4^* \to W_2$, and let $U = \ker(Q) \subseteq V_4^*$. Note that V_4 carries a canonical $\mathfrak{so}(6, \mathbb{C})$ -invariant inner product, allowing us to decompose $V_4 = U \oplus U^{\perp}$ into the sum of orthogonal two-dimensional subspaces. Consider now the Lie subalgebra of $\mathfrak{so}(6, \mathbb{C})$ that preserves this decomposition, which provides an embedding

$$\mathfrak{sl}(2,\mathbb{C})\oplus\mathfrak{sl}(2,\mathbb{C})\to\mathfrak{sl}(4,\mathbb{C})\cong\mathfrak{so}(6,\mathbb{C}),$$

or equivalently $\mathfrak{so}(4, \mathbb{C}) \subset \mathfrak{so}(6, \mathbb{C})$. Each summand U, U^{\perp} is identified with an $\mathfrak{so}(4, \mathbb{C})$ -irreducible summand S_{-}, S_{+} , respectively.

In terms of this decomposition, the odd part of the complexified superconformal algebra also decomposes as

$$\hom(V_4^*, W_2) = \hom(S_+, W_2) \oplus \hom(S_-, W_2)$$

and Q is a full rank element of the first summand. Thus it is a Donaldson-Witten twist.

Consider now the image of [Q, -] inside the even part of the complexified superconformal algebra. It is isomorphic, as a vector space, to \mathbb{C}^4 . In fact, it is an abelian Lie subalgebra, and we are free to view it as the complexified translations of an affine slice, so let us denote it by $T_{\mathbb{C}}$. (There is another copy of \mathbb{C}^4 , orthogonal to this image, which we can then view as the special conformal transformations for this affine slice.)

Even better, as Q comes from a real form, it is a fixed point of the corresponding involution (i.e., of the appropriate version of complex conjugation). The bracket is equivariant for this involution, so the image of [Q, -] restricted to the odd part of the real form gives a 4-dimensional real subspace $T_{\mathbb{R}}$ of $T_{\mathbb{C}}$. We define an affine patch $\iota: \mathbb{R}^{p,4-p} \hookrightarrow C(\mathbb{R}^{p,4-p})$ as the inclusion of the open orbit of the $T_{\mathbb{R}}$ -action on the conformal compactification $C(\mathbb{R}^{p,4-p})$.

4.6.2. *Real Forms of the Schur supercharges*. Let us now discuss the behavior of a Schur supercharge in various real forms, along with the real groups Z_Q and B_Q . For concreteness, let us work in the basis described above and consider the supercharge

$$\mathbb{Q} = \mathcal{Q}_{+1} + \mathcal{S}^{+2}.$$

Alternatively, viewing this supercharge as a 2 × 4 matrix via the given bases for S_+ , W, we can identify

$$\mathbb{Q} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

In this basis we can identify the complex Lie algebras of Q-closed and Q-exact elements explicitly. We now describe the projections of these algebras onto the Lie algebra of complexified infinitesimal conformal transformations:

$$\mathfrak{z}_{\mathbb{Q}}^{\mathbb{C}} \cap \mathfrak{sl}(4,\mathbb{C}) = \left\{ \left(\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right) : A, B, C, D \in \mathfrak{h} \text{ and } \operatorname{Tr}(A) + \operatorname{Tr}(D) = 0 \right\},$$
$$\mathfrak{b}_{\mathbb{Q}}^{\mathbb{C}} \cap \mathfrak{sl}(4,\mathbb{C}) = \left\{ \left(\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right) : A, B, C, D \in \mathfrak{k} \text{ and } \operatorname{Tr}(A) + \operatorname{Tr}(D) = 0 \right\},$$

with $\mathfrak{k} \subseteq \mathfrak{h} \subseteq \mathfrak{gl}(2,\mathbb{C})$ where \mathfrak{k} denotes the subalgebra consisting of matrices with non-zero elements in the left column only and where \mathfrak{h} denotes the subalgebra of lower triangular matrices.

Let us now discuss the real forms for these complex Lie algebras associated to each signature.

4.6.3. *Signature* (4,0). In Euclidean signature we are considering the real form $\mathfrak{sl}(2,\mathbb{H}) \subseteq \mathfrak{sl}(4,\mathbb{C})$. We may obtain it as the fixed points of the involution

$$\theta_{\text{Euc}}(X) = -JXJ$$
35

where

$$J = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}.$$

The fixed point locus of θ_{Euc} acting on $\mathfrak{z}_{\mathbb{Q}}^{\mathbb{C}} \cap \mathfrak{sl}(4,\mathbb{C})$ is three complex-dimensional, or six realdimensional. It consists of matrices of the form

$$\mathfrak{z}_{\mathbb{Q}} \cap \mathfrak{so}(1,5) = \left\{ \left(\begin{array}{c|c} a\mathbb{I} & b\mathbb{I} \\ \hline c\mathbb{I} & d\mathbb{I} \end{array} \right) : a, b, c, d \in \mathbb{C}, a+d=0 \right\}$$

where $\mathbb{I} = \text{diag}(1,1) \in \mathfrak{gl}(2,\mathbb{C})$. This subalgebra is isomorphic to $\mathfrak{sl}(2,\mathbb{C}) \cong \mathfrak{so}(1,3) \cong \text{conf}(2)$, and indeed it acts via conformal transformations on an embedded plane $\mathbb{R}^{2,0} \subseteq \mathbb{R}^{4,0}$. We may see this by observing that in $\mathfrak{sl}(2,\mathbb{H}) \cong \text{conf}(4)$, the subalgebras of infinitesimal translations and special conformal transformations, respectively, may be identified as the subalgebras of strict upper and lower triangular matrices.

The fixed point locus of θ_{Euc} acting on $\mathfrak{b}_{\mathbb{Q}}^{\mathbb{C}} \cap \mathfrak{sl}(4,\mathbb{C})$ is trivial, so there are no real \mathbb{Q} -exact conformal transformations in Euclidean signature.

4.6.4. *Signature* (3,1). In Lorentzian signature we are considering the real form $\mathfrak{su}(2,2) \subseteq \mathfrak{sl}(4,\mathbb{C})$. We may obtain it as the fixed points of the involution

$$\theta_{\rm Lor}(X) = -KX^{\dagger}K$$

where

$$K = diag(1, 1, -1, -1).$$

The fixed point locus of θ_{Lor} acting on $\mathfrak{z}_{\mathbb{Q}}^{\mathbb{C}} \cap \mathfrak{sl}(4,\mathbb{C})$ is seven real-dimensional. It consists of matrices of the form

$$\mathfrak{z}_{\mathbb{Q}} \cap \mathfrak{so}(2,4) = \left\{ \left(\begin{array}{c|c} A & B \\ \hline \overline{B} & D \end{array} \right) : A = \operatorname{diag}(\alpha,\beta), D = \operatorname{diag}(\gamma,\delta), B = \operatorname{diag}(a,b), \\ \text{where } a, b \in \mathbb{C} \text{ and } \alpha, \beta, \gamma, \delta \in i\mathbb{R}, \alpha + \beta + \gamma + \delta = 0 \right\}.$$

This subalgebra of $\mathfrak{su}(2,2)$ is a semisimple quotient isomorphic to $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$, and the solvable radical generates a dilation in the conformal group. The two factors of this semisimple quotient act on the two factors of $\mathbb{C} \oplus \mathbb{C} \cong \mathbb{R}^4$ by antiholomorphic vector fields.

The fixed point locus of θ_{Lor} acting on $\mathfrak{b}_Q^{\mathbb{C}} \cap \mathfrak{sl}(4,\mathbb{C})$ is three real-dimensional. It consists of matrices of the form

$$\mathfrak{b}_{\mathbb{Q}}\cap\mathfrak{so}(2,4)=\left\{\left(\begin{array}{c|c}A & B\\\hline \overline{B} & -A\end{array}\right): A=\operatorname{diag}(\alpha,0)B=\operatorname{diag}(a,0), a\in\mathbb{C}, \alpha\in i\mathbb{R}\right\}.$$

This Lie algebra is isomorphic to $\mathfrak{su}(2)$ (embedded in the $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$ above via the map $X \mapsto (X, 0)$), and therefore acts on an embedded complex line $\mathbb{C} \subseteq \mathbb{C}^2 \cong \mathbb{R}^4$ by antiholomorphic vector fields.

Remark 4.26. One can compare this discussion to the explicit description of the closed and exact transformations appearing in [ALW23, Section 2.3]. The generators of $\mathfrak{z}_{\mathcal{Q}}^{\mathbb{C}}$ are denoted there by $\mathfrak{sl}_{2} \oplus \widehat{\mathfrak{sl}_{2}} \oplus \mathcal{Z} \oplus \mathfrak{m}_{+}$, and the factor \mathfrak{m}_{+} is orthogonal to the fixed point locus of the involution $\theta_{\text{Lor.}}$.

4.6.5. *Signature* (2, 2). We will only discuss the split signature case very briefly. Recall that we must consider the real form $\mathfrak{sl}(4, \mathbb{R}) \subseteq \mathfrak{sl}(4, \mathbb{C})$, so the intersections $\mathfrak{z}_Q \cap \mathfrak{so}(3, 3)$ and $\mathfrak{b}_Q \cap \mathfrak{so}(3, 3)$ are very easy to understand: they are the subalgebras of $\mathfrak{z}_Q^C \cap \mathfrak{sl}(4, \mathbb{C})$ and $\mathfrak{b}_Q^C \cap \mathfrak{sl}(4, \mathbb{C})$ consisting of matrices with real coefficients. In particular, if we identify the upper-right block matrices with translations, then there are three closed real translations, of which two are exact.

5. 3D SUPERCONFORMAL TWISTS

5.1. **Complex Nilpotence Variety.** Over \mathbb{C} , the 3d $\mathcal{N} = k$ superconformal algebra $\mathfrak{sconf}(3|k,\mathbb{C})$ can be identified with the super Lie algebra $\mathfrak{osp}(k|4,\mathbb{C})$. The even part of $\mathfrak{sconf}(3|k,\mathbb{C})$ is isomorphic to

$$(\mathfrak{sconf}(3|k,\mathbb{C})_0 \cong \mathfrak{so}(k,\mathbb{C}) \oplus \mathfrak{sp}(4,\mathbb{C}) \cong \mathfrak{so}(k,\mathbb{C}) \oplus \mathfrak{so}(5,\mathbb{C}).$$

The odd part of $\mathfrak{sconf}(3|k, \mathbb{C})$ can be identified with

$$(\mathfrak{sconf}(3|k,\mathbb{C}))_1\cong W_k\otimes V_4\cong W_k\otimes \mathbb{S}_4,$$

where W_k is the *k*-dimensional fundamental representation of $\mathfrak{so}(k, \mathbb{C})$, and V_4 is the 4-dimensional fundamental representation of $\mathfrak{sp}(4, \mathbb{C})$, or equivalently the (Dirac) spinor representation of $\mathfrak{so}(5, \mathbb{C})$. The Lie bracket between two odd homogeneous elements $w \otimes Q$, $w' \otimes Q'$ is given as follows. Write $\omega(,)$ for the symplectic pairing on V_4 , and write g(,) for the symmetric pairing on W_{2k} . We have

$$[w \otimes Q, w' \otimes Q'] = (w \wedge w')\omega(Q, Q') + g(w, w')(Q \cdot Q')$$

using the isomorphism of $\mathfrak{so}(5,\mathbb{C})$ representations $\operatorname{Sym}^2(S_4) \to \mathfrak{so}(5,\mathbb{C})$ on the second factor.

Let us compute the associated superconformal nilpotence variety.

Proposition 5.1. *The complex nilpotence variety for the 3d* $\mathcal{N} = k$ *superconformal algebra is*

$$\mathcal{N}_{k}^{(3)} \cong (\{(w,w') \in W_{k}^{2} \colon g(w,w) = g(w',w') = g(w,w') = 0\} \times (\mathbb{CP}^{1})^{2}) / (((0,w'),(c,c')) \sim ((0,w'),(0,c')), ((w,0),(c,c')) \sim ((w,0),(c,0))).$$

Proof. Let Q_1, Q_2, Q_3, Q_4 be a Darboux basis for S_4 , such that $\omega(Q_1, Q_2) = \omega(Q_3, Q_4) = 1$. Let

$$Q = (w_1, w_2, w_3, w_4) = w_1 \otimes Q_1 + w_2 \otimes Q_2 + w_3 \otimes Q_3 + w_4 \otimes Q_4$$

be an arbitrary element of $(\mathfrak{sconf}(3|k,\mathbb{C}))_1$. Suppose this element squares to zero. If we first consider the $\mathfrak{so}(5,\mathbb{C})$ summand of $[\mathcal{Q},\mathcal{Q}]$, this means that

$$\sum_{i,j=1}^{4} g(w_i, w_j)(Q_i \cdot Q_j) = 0$$
$$\implies g(w_i, w_j) = 0 \text{ for all } i, j.$$

Now, considering the $\mathfrak{so}(2k, \mathbb{C})$ summand of $[\mathcal{Q}, \mathcal{Q}]$, we have

$$\sum_{i,j=1}^{4} (w_i \wedge w_j) \omega(Q_i, Q_j) = 0$$

$$\implies w_1 \wedge w_2 = 0$$

and $w_3 \wedge w_4 = 0.$

Thus let $w_1 = w$, $w_2 = cw$, $w_3 = w'$ and $w_4 = c'w'$ for complex constants c, c'.

Remark 5.2. How does the translation nilpotence variety embed inside the conformal nilpotence variety? Two copies of the 3d $\mathcal{N} = k$ supersymmetry algebra embed inside $\mathfrak{sconf}(3|k,\mathbb{C})$. We can understand the embedding by considering the restriction of the adjoint representation of $\mathfrak{sconf}(3|k,\mathbb{C})$ to a representation of the subalgebra $\mathfrak{so}(k,\mathbb{C}) \oplus \mathfrak{so}(3,\mathbb{C})$. At the level of super vector spaces, this restricted representation takes the form

$$(\mathfrak{so}(k,\mathbb{C})\oplus\mathfrak{so}(3,\mathbb{C})\oplus(\mathbb{C}^3)^2\oplus\mathbb{C})\oplus\Pi(W_k\otimes(\mathbb{S}_2)^2),$$

where S_2 is the (Dirac) spinor representation of $\mathfrak{so}(3, \mathbb{C})$. From this point of view, the usual nilpotence variety embeds either by considering the locus w' = 0 or the locus w = 0 inside the conformal nilpotence variety.

In the next sections we analyze the complex and real orbits in $\mathcal{N}_k^{(3)}$ under the action of the even part of the superconformal group.

5.2. **Complex Group Orbits & Closed and Exact Transformations.** The even part of the superconformal group

$$\operatorname{SConf}(3|k,\mathbb{C})_0 \cong \operatorname{Spin}(5,\mathbb{C}) \oplus \operatorname{SO}(k,\mathbb{C})$$

acts on $\mathcal{N}_k^{(3)}$, and each complex orbit has a well-defined *rank*: that is, the rank of a square-zero supercharge $\mathcal{Q} \in W_k \otimes V_4$ viewed as a linear map $W_k \to V_4^*$. The image of the linear map associated to \mathcal{Q} must be isotropic, so there will only be three complex orbits, associated to isotropic linear maps of rank 0, rank 1, and rank 2. (This classification is another example of the orbit classification of Duflo–Serganova [DS05, Theorem 4.2].)

For the rank *r* complex orbit we will describe the kernel and the image of the operator [Q, -] for a supercharge Q of rank *r*.

- **Rank 0**: The zero supercharge always squares to zero. If k < 2 this is the only point in $\mathcal{N}_{k}^{(3)}$. The kernel of $[\mathcal{Q}, -]$ consists of the entirety of $\mathfrak{so}(5, \mathbb{C}) \oplus \mathfrak{so}(k, \mathbb{C})$, and the image of $[\mathcal{Q}, -]$ is just $\{0\}$.
- Rank 1: Rank one nilpotent supercharges Q = Q ⊗ w exist for all k ≥ 2. The action of SConf(3|k, C)₀ on the space of rank one nilpotent supercharges is transitive.

Let us write $\mathfrak{p}_k \subseteq \mathfrak{so}(k, \mathbb{C})$ for the maximal parabolic subalgebra with block diagonal Levi quotient $\mathfrak{so}(2, \mathbb{C}) \oplus \mathfrak{so}(k-2, \mathbb{C})$. Write $f_k : \mathfrak{p}_k \to \mathbb{C} \cong \mathfrak{so}(2, \mathbb{C})$ for the projection onto the $\mathfrak{so}(2, \mathbb{C})$ summand in this quotient.

Proposition 5.3. The kernel $\mathfrak{z}_{\mathcal{Q}}$ of the operator $[\mathcal{Q}, -]$ is isomorphic to the kernel of the map

$$f_5 - f_k \colon \mathfrak{p}_5 \oplus \mathfrak{p}_k \to \mathbb{C}$$

as a subalgebra of $\mathfrak{so}(5,\mathbb{C})\oplus\mathfrak{so}(k,\mathbb{C})\cong\mathfrak{sp}(4,\mathbb{C})\oplus\mathfrak{so}(k,\mathbb{C}).$

Proof. Without loss of generality let $Q = Q_1 \otimes w$ where $w = (1, i, 0, ..., 0) \in W_k$. The subalgebra \mathfrak{z}_Q is given as

$$\mathfrak{z}_{\mathcal{Q}} = \{(A, B) \in \mathfrak{sp}(4, \mathbb{C}) \oplus \mathfrak{so}(k, \mathbb{C}) \colon AQ_1 = \lambda Q_1, Bw = -\lambda w \text{ for some } \lambda \in \mathbb{C}\}.$$

We first check that the reductive quotient l of ker $(f_5 - f_k)$ is contained in \mathfrak{z}_Q . Here l is the subalgebra

$$\mathfrak{l}=\mathfrak{so}(2,\mathbb{C})\oplus\mathfrak{so}(3,\mathbb{C})\oplus\mathfrak{so}(k-2,\mathbb{C})\subseteq\mathfrak{so}(5,\mathbb{C})\oplus\mathfrak{so}(k,\mathbb{C})$$

embedded via the map $(A, X, Y) \mapsto (\text{diag}(A, X), \text{diag}(-A, Y))$. If $\lambda \mapsto A_{\lambda}$ under the natural isomorphism $\mathbb{C}^{\times} \cong SO(2, \mathbb{C})$, certainly

$$\operatorname{diag}(-A_{\lambda}, Y)w = -\lambda w.$$

Let

$$X = \begin{pmatrix} a & b \\ c & -a \end{pmatrix} \in \mathfrak{sl}(2,\mathbb{C}) \cong \mathfrak{so}(3,\mathbb{C}).$$

Under the exceptional isomorphism $\mathfrak{so}(5,\mathbb{C}) \to \mathfrak{sp}(4,\mathbb{C})$, the matrix diag (A_{λ}, X) maps to

$$\begin{pmatrix} \lambda & 0 & 0 & 0 \\ 0 & a & 0 & b \\ 0 & 0 & -\lambda & 0 \\ 0 & c & 0 & -a \end{pmatrix} \in \mathfrak{sp}(4,\mathbb{C}).$$

Thus in particular diag(A_{λ} , X) $Q_1 = \lambda Q_1$ as required.

The parabolic subalgebra $\mathfrak{p}_k \subseteq \mathfrak{so}(k, \mathbb{C})$ has nilradical of dimension k - 2. To complete the argument we must show that the nilradical of $\mathfrak{p}_5 \oplus \mathfrak{p}_k$ is contained in \mathfrak{z}_Q . In other words, it is sufficient to show that the quotient $\mathfrak{z}_Q/\mathfrak{l}$ is an abelian subgroup of the same dimension as the nilradical of \mathfrak{p} , which is exactly (5-2) + (k-2) = k+1. We can compute this directly. The quotient $\mathfrak{z}_Q/\mathfrak{l}$ is spanned by matrices of the form

$$\begin{pmatrix} 0 & \alpha & \beta & \gamma \\ 0 & 0 & \gamma & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -\alpha & 0 \end{pmatrix} \in \mathfrak{sp}(4, \mathbb{C})$$

together with matrices of the form

$$\begin{pmatrix} 0 & 0 & x_1 & \cdots & x_{k-2} \\ 0 & 0 & ix_1 & \cdots & ix_{k-2} \\ -x_1 & -ix_1 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -x_{k-2} & -ix_{k-2} & 0 & \cdots & 0 \end{pmatrix} \in \mathfrak{so}(k, \mathbb{C}).$$

The ideal generated by these elements is indeed an abelian subalgebra of dimension k + 1 as required.

The image $\mathfrak{b}_{\mathcal{Q}} \subseteq \mathfrak{z}_{\mathcal{Q}}$ of $[\mathcal{Q}, -]$ is four-dimensional, spanned by a three-dimensional subspace of $\mathfrak{sp}(4, \mathbb{C})$ and a fourth generator contained in the diagonally embedded copy of $\mathfrak{so}(2, \mathbb{C})$ in the kernel described above.

• **Rank 2**: Rank two nilpotent supercharges exist if $k \ge 4$, and this is the maximal possible rank. There are two orbits in the space of rank two nilpotent supercharges. We can use the $\mathfrak{so}(5, \mathbb{C})$ action to scale the values of c, c' in Proposition 5.1 to zero, so we are studying the orbits in the space of pairs of orthogonal non-zero null vectors in W_k .

Proposition 5.4. *If* $k \ge 4$, *there is a one parameter family of* SO(k, \mathbb{C}) *orbits in the space of pairs of orthogonal non-zero null vectors in* W_k :

$$Orb_{x} = \{A \cdot ((1, i, 0, ..., 0), (x, ix, 1, i, 0, ..., 0)) \colon A \in SO(k, \mathbb{C})\}$$

where $x \in \mathbb{C}$.

Proof. Given a pair (w_1, w_2) of orthogonal null-vectors, we may use the transitive SO (k, \mathbb{C}) action to pass to a point in the orbit of the form ((1, i, 0, ..., 0), w), where $w \neq 0$ is orthogonal to (1, i, 0, ..., 0). Thus, this means that w = x(1, i, 0, ..., 0) + w', where $x \in \mathbb{C}$ and $w' = (0, 0, z_3, z_4, ..., z_k)$, but not all the z_i are equal to zero.

Let us now consider the SO(k - 2, \mathbb{C}) action on the last k - 2 coordinates. This action is transitive on the space of null vectors of the form $(0, 0, z_3, z_4, ..., z_k)$, so we may pass to a point in the same orbit where $z_3 = 1$, $z_4 = i$ and $z_j = 0$ for j > 4.

The subgroup SO($k - 2, \mathbb{C}$) is precisely the stabilizer of the element $(1, i, 0, ..., 0) \in W_k$, so these elements lie in different orbits for different values of $x \in \mathbb{C}$.

Example 5.5. In the $\mathcal{N} = 2$ case there is a single non-zero orbit of nilpotent supercharges given by elements $\mathcal{Q} = \mathcal{Q} \otimes w$, where \mathcal{Q} is a non-zero element of V_4 and w is a non-zero null vector in W_2 . A representative point is given by $\mathcal{Q} = (1, 0, 0, 0)$ and w = (1, i). The kernel $\mathfrak{z}_{\mathcal{Q}}$ is given by the subalgebra

$$\mathfrak{p} = \left\{ \begin{pmatrix} a & b & c & d \\ 0 & e & d & f \\ 0 & 0 & -a & 0 \\ 0 & g & -b & -e \end{pmatrix} \right\} \subseteq \mathfrak{sp}(4,\mathbb{C}) \oplus \mathfrak{so}(2,\mathbb{C}),$$

embedded via the map $X \mapsto (X, -a) \in \mathfrak{sp}(4, \mathbb{C}) \oplus \mathfrak{so}(2, \mathbb{C})$. The image $\mathfrak{b}_{\mathcal{Q}} \subseteq \mathfrak{z}_{\mathcal{Q}}$ is spanned by the four elements

ſ	(0	0	1	0)))	١	(0	0	0	1		١	((1)	0	0	0))		(0	1	0	0)	,)	
J		0	0	0	0				0	0	1	0		0,		0	0	0	0	1			0	0	0	0		
Ì		0	0	0	0	,0	′		0	0	0	0	,0			0	0	-1	0	,-1	′		0	0	0	0	,0	ĺ
l		0	0	0	0))		0	0	0	0)))	$\left(\right)$	0	0	0	0)	!,		(0	0	$^{-1}$	0)	- J	J
in	sp	(4,	C)∈	∋s¢	(2,	C).																						

5.3. **Real Group Orbits.** We may analyze the orbits in the complex nilpotence variety under the adjoint action of a real form of the even part of the superconformal group using the same technique as in four dimensions. In other words, for a complex orbit O_r , we first use the following fibration.

Proposition 5.6. There is a fibration

$$\pi\colon \mathcal{O}_r \to \operatorname{SpGr}(4-r,4)$$
⁴⁰

where $\operatorname{SpGr}(r, 4)$ is the Grassmannian of isotropic *r*-dimensional subspaces in $\mathbb{C}^4 \cong T^*\mathbb{C}^2$, equipped with *its standard holomorphic symplectic structure.*

Now, we pursue the same approach as in Section 4.5. That is, we observe that the two factors $SO(k, \mathbb{C})$ and $Sp(4, \mathbb{C})$ of the even part of the superconformal group act only on the fiber and only on the base of the fibration π , respectively. That is, $SO(k, \mathbb{C})$ acts on $GL(r, \mathbb{C})$ by left multiplication, viewing $GL(r, \mathbb{C})$ as the subgroup of $GL(k, \mathbb{C})$ consisting of matrices with zeroes outside the upper-left $r \times r$ block. Meanwhile $Sp(4, \mathbb{C})$ acts on the homogeneous space $SpGr(4 - r, 4) \cong Sp(4, \mathbb{C})/P$, again by left multiplication. Thus we may separate the question of orbits into the base and fiber directions. In the base we may use the analog of Theorem 4.19 for general semisimple groups.

Theorem 5.7 ([Wol69, Theorem 2.6]). Let $G_{\mathbb{C}}$ be a complex semisimple group, let G be a real form of $G_{\mathbb{C}}$ and let P be a parabolic subgroup of $G_{\mathbb{C}}$. Then there are finitely many G-orbits in the partial flag variety $G_{\mathbb{C}}/P$. There is a unique closed orbit of real dimension equal to dim_C $G_{\mathbb{C}}/P$.

We may therefore observe that there are finitely many orbits for any real form of $Sp(4, \mathbb{C})$.

In both signature (3,0) and (2,1), the real form of $SO(k, \mathbb{C})$ is just $SO(k, \mathbb{R})$. Thus we may conclude that the space of real sub-orbits of the complex orbit \mathcal{O}_r for the nilpotence variety in dimension 3 are isomorphic to finitely many copies of the homogeneous space

$$\operatorname{GL}(r,\mathbb{C})/\operatorname{SO}(r,\mathbb{R}).$$

For the case r = 0, this space is just a point. If r = 1, this homogeneous space is just \mathbb{C}^{\times} . If r = 2 it is the seven real-dimensional manifold $GL(2, \mathbb{C})/S^1 \cong \mathbb{C}^3 \times \mathbb{R}_{>0}$.

6. 5D AND 6D SUPERCONFORMAL TWISTS

6.1. **5d Twists.** In dimension 5 there is a unique complex superconformal algebra, namely the exceptional super Lie algebra $\mathfrak{sconf}(5|2,\mathbb{C}) = \mathfrak{f}(4)$. This algebra contains the 5d $\mathcal{N} = 1$ supersymmetry algebra as a subalgebra. Indeed, the even part of $\mathfrak{sconf}(5|2,\mathbb{C})$ is given by

$$(\mathfrak{sconf}(5|2,\mathbb{C}))_0 = \mathfrak{so}(7,\mathbb{C}) \oplus \mathfrak{sl}(2,\mathbb{C}),$$

and the odd part of $\mathfrak{sconf}(5|2,\mathbb{C})$ is given by

$$(\mathfrak{sconf}(5|2,\mathbb{C}))_1 = S \otimes W$$

where *S* is the 8-dimensional (Dirac) spinor representation of $\mathfrak{so}(7, \mathbb{C})$ and *W* is the 2-dimensional defining representation of $\mathfrak{sl}(2, \mathbb{C})$.

The bracket between two odd elements is given as follows. The 64-dimensional representation $S \otimes S$ of $\mathfrak{so}(7, \mathbb{C})$ decomposes as

$$S \otimes S \cong \wedge^2(S) \oplus \operatorname{Sym}^2(S)$$
$$\cong (V_7 \oplus \mathfrak{so}(7, \mathbb{C})) \oplus (\mathbb{C} \oplus \wedge^3 V_7),$$

where $\wedge^3 V_7$ is the 35-dimensional irreducible representation arising as the third exterior power of the defining representation. In particular, there is an anti-symmetric equivariant linear map

$$F\colon S\otimes S\to\mathfrak{so}(7,\mathbb{C}),$$

and a symmetric equivariant linear map

$$\langle,\rangle:S\otimes S\to\mathbb{C}.$$

Using the notation Tr, red from Section 4.1, the bracket is given by

$$[Q \otimes w, Q' \otimes w'] = (F(Q \otimes Q') \cdot \operatorname{Tr}(w \otimes w'), \langle Q, Q' \rangle \cdot \operatorname{red}(w \otimes w')).$$

When we restrict $(\mathfrak{sconf}(5|2,\mathbb{C}))_1$ to a representation of $\mathfrak{so}(5,\mathbb{C}) \subseteq \mathfrak{so}(7,\mathbb{C})$, we do indeed obtain two copies of $S_{(5)} \otimes W$, where $S_{(5)}$ is the four-dimensional (Dirac) spinor representation of $\mathfrak{so}(5,\mathbb{C})$. That is, two copies of the spinors in the 5d $\mathcal{N} = 1$ supersymmetry algebra.

Let us compute the associated superconformal nilpotence variety.

Proposition 6.1. The complex nilpotence variety for the 5d superconformal algebra is isomorphic to

$$\operatorname{Nilp}_{\mathbb{C}} \cong \{ (Q_1, Q_2) \in S^2 \colon F(Q_1 \otimes Q_2) = 0 \text{ and } \langle Q_i, Q_j \rangle = 0, \text{ for any } i, j = 1, 2 \}.$$

The complex adjoint orbits in the nilpotence variety are easy to analyze here: the action on $\text{Nilp}_{\mathbb{C}} - \{0\}$ is transitive, so there is only a single non-zero orbit in the nilpotence variety.

Proof. Let $\{w_1, w_2\}$ be a Darboux basis for W with respect to the trace pairing. A general element of $(\mathfrak{sconf}(5|2, \mathbb{C}))_1$ can be written as $Q = Q_1 \otimes w_1 + Q_2 \otimes w_2$ for some $(Q_1, Q_2) \in S^2$. Let us compute the bracket [Q, Q]. According to our description of the bracket, the $\mathfrak{so}(7, \mathbb{C})$ summand of [Q, Q] is

$$[Q,Q]_{\mathfrak{so}(7,\mathbb{C})} = \sum_{i,j=1}^{2} F(Q_i \otimes Q_j) \operatorname{Tr}(w_i \otimes w_j)$$
$$= 2F(Q_1 \otimes Q_2).$$

The $\mathfrak{sl}(2,\mathbb{C})$ summand of [Q,Q] is

$$[Q,Q]_{\mathfrak{sl}(2,\mathbb{C})} = \sum_{i,j=1}^{2} \langle Q_i, Q_j \rangle \operatorname{red}(w_i \otimes w_j)$$
$$= \langle Q_1, Q_1 \rangle e + \langle Q_2, Q_2 \rangle f + \langle Q_1, Q_2 \rangle h$$

where $\{e, f, h\}$ is the standard basis for $\mathfrak{sl}(2, \mathbb{C})$ Therefore [Q, Q] = 0 if and only if

$$F(Q_1 \otimes Q_2) = \langle Q_1, Q_1 \rangle = \langle Q_2, Q_2 \rangle = \langle Q_1, Q_2 \rangle = 0,$$

finishing the characterization of the nilpotent elements.

6.2. **6d Twists.** In dimension 6 there exist $\mathcal{N} = (k, 0)$ superconformal algebras for all positive integers *k*. We can identify superconformal algebras in dimension 6 with the simple super Lie algebras $\mathfrak{osp}(8|2k)$ using the triality isomorphism to identify the defining representation of $\mathfrak{so}(8, \mathbb{C})$ with a (Weyl) spinor representation. Thus, writing $\mathfrak{sconf}(6|k, \mathbb{C}) = \mathfrak{osp}(8|2k)$ for the $\mathcal{N} = k$ superconformal algebra, we can identify the even part with

$$(\mathfrak{sconf}(6|k,\mathbb{C}))_0 = \mathfrak{so}(8,\mathbb{C}) \oplus \mathfrak{sp}(2k,\mathbb{C}),$$

and the odd part with

$$(\mathfrak{sconf}(6|k,\mathbb{C}))_1 = V_8 \otimes W_{2k} \cong S_+ \otimes W_{2k}$$

where V_8 is the defining representation of $\mathfrak{so}(8, \mathbb{C})$, which is transformed into the Weyl spinor representation S_+ by the triality automorphism of $\mathfrak{so}(8, \mathbb{C})$. As before, W_{2k} denotes the 2*k*-dimensional defining representation of $\mathfrak{sp}(2k, \mathbb{C})$.

The Lie bracket between two odd elements of $\mathfrak{sconf}(6|k, \mathbb{C})$ is defined as follows, in a similar manner to the previous examples:

$$[v \otimes w, v' \otimes w'] = (v \wedge v' \omega(w, w'), g(v, v') F(w \otimes w')),$$

where $F: W_{2k} \otimes W_{2k} \to \mathfrak{sp}(2k, \mathbb{C})$ is the canonical projection.

Remark 6.2. It is interesting to note that the $\mathcal{N} = (2,0)$ complex superconformal algebra in six dimensions and the $\mathcal{N} = k$ complex superconformal algebra in three dimensions are actually isomorphic; the roles of the conformal and R-symmetry transformations are interchanged. Both are isomorphic to the simple super Lie algebra $\mathfrak{osp}(8|4,\mathbb{C})$. In particular, in the $\mathcal{N}=(2,0)$ case the classification of complex orbits for the even part of the complexified superconformal group is identical to the classification from Section 5.

7. TWISTED OBSERVABLES AND FACTORIZATION ALGEBRAS

There are many ways to approach QFT mathematically, and here we will use a framework that captures the spacetime-dependence of operators and operator products via operads. In fact, we will use two approaches drawn from [CG17; CG21], freely referencing those books below. The first approach uses prefactorization algebras and is highly flexible, allowing one to work on arbitrary manifolds and with operators whose support is an arbitrary open subspace. A key result of [CG21] is that a Lagrangian field theory on the manifold *M* naturally produces a prefactorization algebra on *M*, so we have a wealth of examples. (That book also examines the perturbative quantization of such theories in Euclidean signature and shows these also yield prefactorization algebras.) The second approach restricts to $M = \mathbb{R}^n$ and to open disks therein. It matches closely with standard manipulations in the physics literature and has the additional virtue that there are convenient comparison results letting one extract more familiar algebraic structures – like vertex algebras or associative algebras – efficiently. It may be more intuitive for most readers.

Both approaches let us make mathematical statements that are sharp versions of the discussion in Section 3. Our discussion here amounts, primarily, to mimicking ideas and results from [CG17; ES19], which examine chiral conformal and supersymmetric theories, respectively. The arguments carry over straightforwardly to superconformal theories, so we often state the result and merely indicate what needs to be changed.

7.1. **Superconformal Prefactorization Algebras.** We will follow the conventions of [CG17; CG21], so our prefactorization algebras take values in cochain complexes of differentiable vector spaces. This setting allows us to talk about smooth action of a Lie group on a prefactorization algebra, with a Lie algebra acting by differential operators. The experienced reader, however, will see how the idea of such symmetry could be ported to other settings.

Definition 7.1. A conformal (pre)factorization algebra *is a* (*pre*)*factorization algebra on a compactification* $C(\mathbb{R}^{p,q})$ *that is smoothly equivariant for the action of the conformal group* Conf(p,q).

Similarly, we can work in $\mathbb{Z}/2$ -graded cochain complexes of differentiable vector spaces for the superconformal case.

Definition 7.2. A superconformal (pre)factorization algebra is a conformal (pre)factorization algebra on a compactification $C(\mathbb{R}^{p,q})$ for which the Lie algebra action of conf(p,q) is extended to an action of a superconformal algebra sconf(p,q,|S).

A conformal twist Q for $\mathfrak{sconf}(p,q|S)$ then determines a deformation of such a superconformal factorization algebra.

Definition 7.3. Given a $\mathfrak{sconf}(p,q|S)$ -superconformal prefactorization algebra \mathcal{F} on $C(\mathbb{R}^{p,q})$ and a conformal twist \mathcal{Q} , let $\mathcal{F}^{\mathcal{Q}}$ denote the \mathcal{Q} -twisted prefactorization algebra with values in $\mathbb{Z}/2$ -graded cochain complexes of differentiable vector spaces whose underlying super vector space are those of \mathcal{F} but whose differential is $d_{\mathcal{F}} + \mathcal{Q}$.

This construction is a version of "adding Q as a BRST operator."

We note the following, which follows by inspection (at least for the reader of sections 3.7 and 4.8 of [CG17] or of section 2 of [ES19]).

Lemma 7.4. The groups Z_Q and B_Q act smoothly-equivariantly on \mathcal{F}^Q by restricting the action of $SConf(p,q)_0$ on \mathcal{F} . Moreover, the dg Lie algebra $(\mathfrak{sconf}(p,q|S), Q)$ acts by derivations on \mathcal{F}^Q .

This lemma already provides a precise encoding of much of the discussion in section 3. It says, for instance, that for any element $z \in Z_Q$ and for any disjoint open sets U_1, U_2 inside a larger open set $V \subset M = C(\mathbb{R}^{p,q})$, there is a commuting diagram

$$\begin{array}{ccc} \mathcal{F}^{\mathcal{Q}}(U_1) \otimes \mathcal{F}^{\mathcal{Q}}(U_2) & \longrightarrow & \mathcal{F}^{\mathcal{Q}}(V) \\ & & \downarrow^{\cong} & & \downarrow^{\cong} \\ \mathcal{F}^{\mathcal{Q}}(zU_1) \otimes \mathcal{F}^{\mathcal{Q}}(zU_2) & \longrightarrow & \mathcal{F}^{\mathcal{Q}}(zV) \end{array}$$

where zU_1 , for instance, denotes the translation of U_1 by z. (See definition 3.7.1 of [CG17].) If we take V to be the whole manifold and pick a conformal state (or expected value map) $\langle - \rangle$: $\mathcal{F}^{\mathcal{Q}}(M) \to \mathbb{C}$, this equivariance implies

$$\langle \mathcal{O}_1(x_1)\mathcal{O}_2(x_2)\rangle = \langle \mathcal{O}_1(zx_1)\mathcal{O}_2(zx_2)\rangle$$

where $x_i \in U_i$ and \mathcal{O}_i is a point-supported observable.

We will discuss the possibility of restricting to observables on a submanifold of $C(\mathbb{R}^{p,q})$ in section 7.3 below.

7.2. Algebras of Operator Products. We overview one way to encode the most important operators and their products for a field theory on the manifold \mathbb{R}^n . The essential idea is that one specifies the operators supported in any disk

$$D_r(x) = \{ y \in \mathbb{R}^n : |y - x| < r \}$$

and how operators from disjoint disks multiply to give operators on a bigger disk. We assume that the spaces of operators varies smoothly as one adjusts the location and radius of the disks, and

we assume that operator product varies smoothly in those parameters as well. More explicitly, let $\text{Disk}_{r_1,...,r_k;R}$ denote the open subset of $\mathbb{R}^{(k+1)n}$ given by points $(x_1,...,x_k;y)$ such that the disks $D_{r_i}(x_i)$ are pairwise disjoint but each sits inside $D_R(y)$. Let \mathcal{A} , which we will call an *n*-disks algebra, denote the following data:

- For each positive real number *r*, there is a cochain complex *A_r*, which we view as the operators supported in the disk *D_r*(0).
- For each point $p \in \text{Disk}_{r_1,...,r_k,R}$, there is a cochain map

$$m_p: A_{r_1} \otimes \cdots \otimes A_{r_k} \to A_R,$$

which we view as encoding the operator product. We require that m_p vary smoothly with the configuration p, i.e., there is a smooth function

$$m: \operatorname{Disk}_{r_1,\ldots,r_k;R} \to \operatorname{Hom}(A_{r_1} \otimes \cdots \otimes A_{r_k}, A_R)$$

from the space of configurations into the space of "multiplications."

These maps compose in a natural way based on embedding disks into bigger disks. To be precise, suppose we pick a point *p* ∈ Disk_{r1,...,rk}, and points *q_i* ∈ Disk_{s1,...,sij}, for every *i* such that "outgoing" disk of radius *r_i* from *q_i* equals the incoming disk of radius *i* from *p*. Then we have the equation

$$m_p \circ (m_{q_1} \otimes \cdots \otimes m_{q_k}) = m_F$$

where *P* denotes the point in $\text{Disk}_{s_1^1,...,s_{j_1}^1,s_1^2,...,s_{j_k}^k;R}$ arising by "composing" the disks. A detailed treatment of this notion appears in section 4.8 of [CG17] and in section 2 of [ES19].

Remark 7.5. A technical issue is to formulate in what sense *m* is a smooth function, which requires finding a setting where the codomain Hom has the necessary structure. In [CG17] it is shown that differentiable vector spaces provide a sufficient setting and it is shown that most topological vector spaces are differentiable vector spaces.

When a Lie group *G* acts smoothly by isometries on \mathbb{R}^n , it induces an action on the configuration spaces $\text{Disk}_{r_1,...,r_k;R}$, so one can talk about a *n*-disks algebra \mathcal{A} being *G*-equivariant. Similarly, one can formulate how a dg Lie algebra can act by derivations on \mathcal{A} .

Definition 7.6. An *n*-disks algebra A with values in $\mathbb{Z}/2$ -graded cochain complexes of differentiable vector spaces is superconformal if $\mathfrak{sconf}(p,q|S)$ acts by derivations on A, where n = p + q.

A conformal twist Q for $\mathfrak{sconf}(p,q|S)$ then determines a deformation of such a superconformal disks-algebra.

Definition 7.7. Given a sconf(p, q|S)-superconformal p + q-disks algebra A and a conformal twist Q, let A^Q denote the Q-twisted *n*-disks algebra whose underlying super vector space are those of A but whose differential is $d_A + Q$, where n = p + q.

A natural source of examples is to take a (well-behaved) superconformal prefactorization algebra (or twist thereof) and ask what it assigns to disks inside $\mathbb{R}^{p,q}$. Lagrangian theories with superconformal symmetry, whether classical or quantum, thus give many examples. 7.3. **Obtaining Vertex Algebras or** \mathbb{E}_k -**Algebras.** Under some reasonable hypotheses, one can obtain a vertex algebra or an \mathbb{E}_k algebra (i.e., algebra over the little *k*-disks operad) from the observables of a field theory, whether encoded as a prefactorization algebra or as a disk-algebra. Precise results appear in Chapter 5 of [CG17] and Section 2 of [ES19]. Here we will state their consequences.

We will explain the essential format by an example. Let Q be a square-zero element of the super Lie algebra $\mathfrak{sconf}(p,q|S)$. Suppose that the stabilizer Z_Q contains a subgroup of $\mathrm{ISO}(p,q)$ that is isomorphic to $\mathrm{ISO}(m,n)$. For simplicity, suppose this subgroup is the isometries of an isometric embedding $j: \mathbb{R}^{m,n} \hookrightarrow \mathbb{R}^{p,q}$.

We can consider observables supported along this submanifold: define $j^* \mathcal{F}^Q$ on $\mathbb{R}^{m,n}$ by

$$j^*\mathcal{F}^{\mathcal{Q}}(U) = \lim_{U \subset V \subset \mathbb{R}^{p,q}} \mathcal{F}^{\mathcal{Q}}(V),$$

where *V* runs over open subsets of $\mathbb{R}^{p,q}$ and lim denotes the limit in the appropriate category. (When working with cochain complexes, we mean an ∞ -categorical limit and hence, in practice, a homotopy limit.) This formula yields a precosheaf because if $U \subset U'$, then any open set $V \subset \mathbb{R}^{p,q}$ that contains U' also contains U, so we have a canonical map

$$j^* \mathcal{F}^{\mathcal{Q}}(U) \to \mathcal{F}^{\mathcal{Q}}(V)$$

and hence

$$j^*\mathcal{F}^{\mathcal{Q}}(U) \to \lim_{U' \subset V \subset \mathbb{R}^{p,q}} \mathcal{F}^{\mathcal{Q}}(V)$$

by taking limits. This formula also determines a prefactorization algebra on $\mathbb{R}^{m,n}$ where the structure map

$$j^*\mathcal{F}^{\mathcal{Q}}(U)\otimes j^*\mathcal{F}^{\mathcal{Q}}(U')\to j^*\mathcal{F}^{\mathcal{Q}}(U\sqcup U')$$

for any pair of disjoint open sets U, U' in $\mathbb{R}^{m,n}$ is defined as follows. Fix an open set $W \subset \mathbb{R}^{p,q}$ containing U but disjoint from U', and fix an open set $W' \subset \mathbb{R}^{p,q}$ containing U' but disjoint from W (and hence U). Given an open set $V \subset \mathbb{R}^{p,q}$ that contains $U \sqcup U'$, there is an inclusion of open sets

$$(W \cap V) \sqcup (W' \cap V) \subset V$$

and hence a map

$$j^*\mathcal{F}^{\mathcal{Q}}(U)\otimes j^*\mathcal{F}^{\mathcal{Q}}(U')\to \mathcal{F}^{\mathcal{Q}}(W\cap V)\otimes \mathcal{F}^{\mathcal{Q}}(W'\cap V)\to \mathcal{F}^{\mathcal{Q}}(V),$$

where the second map is the structure map of \mathcal{F}^{Q} as a prefactorization algebra on $\mathbb{R}^{p,q}$. Taking limits, we obtain a map

$$j^*\mathcal{F}^{\mathcal{Q}}(U) \otimes j^*\mathcal{F}^{\mathcal{Q}}(U') \to j^*\mathcal{F}^{\mathcal{Q}}(U \sqcup U') = \lim_{U \sqcup U' \subset V \subset \mathbb{R}^{p,q}} \mathcal{F}^{\mathcal{Q}}(V)$$

as claimed.

Remark 7.8. Even if $\mathcal{F}^{\mathcal{Q}}$ is a factorization algebra, this pullback $j^* \mathcal{F}^{\mathcal{Q}}$ is likely *not* a factorization algebra, as the limit need not play well with the colimits that appear in the local-to-global condition (i.e., codescent).

7.3.1. Obtaining \mathbb{E}_k -Algebras. Let us disscuss the circumstances under which the restricted twisted theory $j^* \mathcal{F}^Q$ on an affine subspace $\mathbb{R}^{m,n}$ given by the embedding j is topological. Corollary 2.30 of [ES19] gives two conditions that guarantee that $j^* \mathcal{F}^Q$ determines an algebra over the little m + n-disks operad – or more colloquially, that the twist looks like a topological field theory along $\mathbb{R}^{m,n}$. These conditions are

- the sub-Lie algebra of translations inside the Lie algebra of ISO(m, n) are Q-exact and
- any inclusion of disks in $\mathbb{R}^{m,n}$ determines a quasi-isomorphism of observables on the disks.

This second condition follows if dilation is also Q-exact.

Corollary 2.39 applies to the case that n = 0. It says that one gets a *framed* \mathbb{E}_m -algebra if, furthermore, the subalgebra $\mathfrak{so}(m)$ is \mathcal{Q} -exact. Such a situation appears not infrequently in Euclidean signature. (An analog of this result should hold for mixed signature as well, using the techniques of [ES19].)

Let us make this application of the results of [ES19] more precise.

Definition 7.9. A conformal embedding $j: \mathbb{R}^{a,b} \hookrightarrow C(\mathbb{R}^{p,q})$ defines a \mathcal{Q} -compatible subspace if

- Z_Q preserves the image of j and
- the group of translations of $j(\mathbb{R}^{a,b})$ is contained in Z_Q .

Let Obs be a prefactorization algebra on $C(\mathbb{R}^{p,q})$, and let Obs^Q denote its twist by the superconformal supercharge Q. Using a choice of Q-compatible subspace given by the embedding j, we obtain a prefactorization algebra j^*Obs^Q on $\mathbb{R}^{a,b}$ as discussed above. The conditions we have placed on the embedding j have been set up in order to ensure the existence of the following structure. We first give a general definition.

Definition 7.10. Let $\mathfrak{h} \subseteq \mathfrak{sconf}(p+q|S,\mathbb{C})_0$ be a Lie subalgebra. A \mathcal{Q} -potential for \mathfrak{h} is a linear subspace $\mathfrak{h}' \subseteq \mathfrak{sconf}(p+q|S,\mathbb{C})_1$ such that

- $[\mathcal{Q}, -]$ defines a linear isomorphism $\mathfrak{h}' \to \mathfrak{h}$.
- $[\mathfrak{h}',\mathfrak{h}']=0.$
- $[\mathfrak{h}, \mathfrak{h}'] \subseteq \mathfrak{h}'$, and as an \mathfrak{h} -representation \mathfrak{h}' is isomorphic to the adjoint representation.

Remark 7.11. If \mathfrak{h} is a Lie algebra, let \mathfrak{h}_{dR} denote the contractible dg Lie algebra with underlying graded Lie algebra $\mathfrak{h}[1] \oplus \mathfrak{h}$ and differential given by the identity. Definition 7.10 says that given $\mathfrak{h} \subseteq \mathfrak{sconf}(p+q|S,\mathbb{C})_0$ we can extend to a commutative triangle of dg Lie algebras



Now, Corollary 2.30 and Theorem 2.38 of [ES19] can be applied to the prefactorization algebra j^*Obs^{Q} in the presence of appropriate potentials. We obtain the following as a direct application of these results.

Theorem 7.12. Let $j: \mathbb{R}^{a,b} \hookrightarrow C(\mathbb{R}^{p,q})$ be a \mathcal{Q} -compatible subspace, and suppose that the group of translations of $j(\mathbb{R}^{a,b})$ is contained in $B_{\mathcal{Q}} \subseteq Z_{\mathcal{Q}}$. Let $G \subseteq B_{\mathcal{Q}}$ be a subgroup acting on the image of j. Let

 $\mathfrak{g} \subseteq \mathfrak{b}_{\mathcal{Q}}^{\mathbb{C}}$ denote the complexified Lie algebra of G, and let \mathfrak{g}' be a \mathcal{Q} -potential for \mathfrak{g} . Finally, suppose that the factorization map $j^*Obs^{\mathcal{Q}}(B_r(0)) \to j^*Obs^{\mathcal{Q}}(B_R(0))$ associated to the inclusion of concentric balls is a quasi-isomorphism for all 0 < r < R. Then $j^*Obs^{\mathcal{Q}}(B_1(0))$ canonically carries the structure of a $G \ltimes \mathbb{E}_{a+b}$ -algebra.

Corollary 7.13 (c.f. [ES19, Corollary 3.40]). If there is an embedding $\mathbb{R}^{\times} \hookrightarrow G$ whose image acts on the image of *j* by dilations, then the condition that $j^*Obs^{\mathcal{Q}}(B_r(0)) \to j^*Obs^{\mathcal{Q}}(B_R(0))$ is a quasi-isomorphism holds automatically.

In particular we obtain framed \mathbb{E}_a -algebras in the pure signature case.

Corollary 7.14 (c.f. [ES19, Corollary 2.39]). If b = 0, G = SO(a) acts on the image of j by rotations, and the hypotheses of Theorem 7.12 hold, then $j^*Obs^{\mathcal{Q}}(B_1(0))$ canonically carries the structure of a \mathbb{E}_a^{fr} -algebra.

7.3.2. *Obtaining Vertex Algebras.* Now let us address the circumstances under which we obtain a vertex algebra under twisting. Let Q be a nilpotent supercharge, and let $j: \mathbb{R}^2 \to C(\mathbb{R}^{p,q})$ be a Q-compatible subspace of definite signature. We will impose the following requirements.

- (1) The group ISO($j(\mathbb{R}^2)$) of isometries of the image of j is contained in Z_Q .
- (2) For a choice of complex coordinate z on \mathbb{R}^2 , the complexified translation $\partial_{\overline{z}}$ is in $\mathfrak{b}_{\mathcal{O}}^{\mathbb{C}}$.

Under these conditions we obtain the following result.

Theorem 7.15. Let $\mathcal{V} = j^* \text{Obs}^{\mathcal{Q}}(D_1(0))$ for j and \mathcal{Q} as above, where $D_1(0) \subseteq \mathbb{R}^2$ is the unit disk. Let

$$V = \bigoplus_{k \in \mathbb{Z}} \mathrm{H}^{\bullet}(\mathcal{V}^k),$$

where \mathcal{V}^k is the k-eigenspace of \mathcal{V} for the action of the group of rotations. Then V can be equipped with the structure of a vertex algebra under the following conditions:

- (1) For each 0 < r < R, the prefactorization map $j^*Obs^{\mathcal{Q}}(D_r(0)) \rightarrow j^*Obs^{\mathcal{Q}}(D_R(0))$ restricts to a quasi-isomorphism on every eigenspace for the S^1 -action.
- (2) $H^{\bullet}(\mathcal{V}^k) = 0$ for all k > K for some integer K.
- (3) The eigenspaces $H^{\bullet}(\mathcal{V}^k)$ satisfy a category-theoretic tameness condition.

This will be a direct consequence of [CG17, Theorem 2.2.1], and we refer the reader to that reference for a discussion of the meaning of the technical conditions that appear in the statement of this theorem. In particular, the "tameness" condition that we have not spelled out here is condition (iii) in *loc. cit*.

REFERENCES

- [AF15] David Ayala and John Francis. "Factorization homology of topological manifolds". Journal of Topology 8.4 (2015), pp. 1045–1084. arXiv: 1206.5522 [math.AT].
- [Aga+17] Mina Aganagic, Kevin Costello, Jacob McNamara, and Cumrun Vafa. "Topological Chern-Simons/Matter Theories" (2017). arXiv: 1706.09977 [hep-th].
- [ALW23] Philip C. Argyres, Matteo Lotito, and Mitch Weaver. "Vertex algebra of extended operators in 4d N=2 superconformal field theories. Part I". JHEP 10 (2023), p. 175. arXiv: 2211.04410 [hep-th].
- [Bau11] Laurent Baulieu. "SU(5)-invariant decomposition of ten-dimensional Yang–Mills supersymmetry". Physics Letters B 698.1 (2011), pp. 63–67. arXiv: 1009.3893 [hep-th].

- [Bee+15a] Christopher Beem, Wolfger Peelaers, Leonardo Rastelli, and Balt C. van Rees. "Chiral algebras of class S". Journal of High Energy Physics 2015.5 (May 2015). URL: https://doi.org/10.1007%2Fjhep05%282015%29020.
- [Bee+15b] Christopher Beem et al. "Infinite Chiral Symmetry in Four Dimensions". Communications in Mathematical Physics 336.3 (Jan. 2015), pp. 1359–1433. URL: https://doi.org/10.1007%2Fs00220-014-2272-x.
- [BHP16] Matthew Buican, Joseph Hayling, and Constantinos Papageorgakis. "Aspects of Superconformal Multiplets in D>4". JHEP 11 (2016), p. 091. arXiv: 1606.00810 [hep-th].
- [Bir+91] Danny Birmingham, Matthias Blau, Mark Rakowski, and George Thompson. "Topological field theory". Phys. Rept. 209 (1991), pp. 129–340.
- [B]23] Matthew Buican and Hongliang Jiang. "From Free Fields to Interacting SCFTs via Representation Theory" (Aug. 2023). arXiv: 2308.03194 [hep-th].
- [BPR17a] Christopher Beem, Wolfger Peelaers, and Leonardo Rastelli. "Deformation Quantization and Superconformal Symmetry in Three Dimensions". Communications in Mathematical Physics 354.1 (Feb. 2017), pp. 345–392. URL: https://doi.org/10.1007%2Fs00220-017-2845-6.
- [BPR17b] Christopher Beem, Wolfger Peelaers, and Leonardo Rastelli. "Deformation quantization and superconformal symmetry in three dimensions". Commun. Math. Phys. 354.1 (2017), pp. 345–392. arXiv: 1601.05378 [hep-th].
- [BR18] Christopher Beem and Leonardo Rastelli. "Vertex operator algebras, Higgs branches, and modular differential equations". Journal of High Energy Physics 2018.8 (Aug. 2018). URL: https://doi.org/10.1007%2Fjhep08%282018%29114.
- [BRR15] Christopher Beem, Leonardo Rastelli, and Balt C. van Rees. "W symmetry in six dimensions". JHEP 05 (2015), p. 017. arXiv: 1404.1079 [hep-th].
- [CDI19] Clay Cordova, Thomas T. Dumitrescu, and Kenneth Intriligator. "Multiplets of Superconformal Symmetry in Diverse Dimensions". JHEP 03 (2019), p. 163. arXiv: 1612.00809 [hep-th].
- [CG17] Kevin Costello and Owen Gwilliam. Factorization algebras in quantum field theory. Vol. 1. Vol. 31. New Mathematical Monographs. Cambridge University Press, Cambridge, 2017, pp. ix+387.
- [CG21] Kevin Costello and Owen Gwilliam. Factorization algebras in quantum field theory. Vol. 2. Vol. 41. New Mathematical Monographs. Cambridge University Press, Cambridge, 2021, pp. xiii+402. URL: https://doi.org/10.1017/9781316678664.
- [DMS97] P. Di Francesco, P. Mathieu, and D. Senechal. Conformal Field Theory. Graduate Texts in Contemporary Physics. New York: Springer-Verlag, 1997.
- [DO03] F. A. Dolan and H. Osborn. "On short and semi-short representations for four-dimensional superconformal symmetry". Annals Phys. 307 (2003), pp. 41–89. arXiv: hep-th/0209056.
- [DS05] Michel Duflo and Vera Serganova. "On associated variety for Lie superalgebras". *arXiv preprint math/0507198* (2005).
- [ES19] Chris Elliott and Pavel Safronov. "Topological twists of supersymmetric algebras of observables". Comm. Math. Phys. 371.2 (2019), pp. 727–786. arXiv: 1805.10806 [math-ph].
- [ESW21] Richard Eager, Ingmar Saberi, and Johannes Walcher. "Nilpotence varieties". Annales Henri Poincare 22.4 (2021), pp. 1319–1376. arXiv: 1807.03766 [hep-th].
- [ESW22] Chris Elliott, Pavel Safronov, and Brian R. Williams. "A taxonomy of twists of supersymmetric Yang-Mills theory". Selecta Math. 28.4 (2022), p. 73. arXiv: 2002.10517 [math-ph].
- [FHW06] Gregor Fels, Alan Huckleberry, and Joseph A. Wolf. Cycle spaces of flag domains. Vol. 245. Progress in Mathematics. A complex geometric viewpoint. Birkhäuser Boston, Inc., Boston, MA, 2006, pp. xx+339. URL: https://doi.org/10.1007/0-8170
- [Gor+22] Maria Gorelik, Crystal Hoyt, Vera Serganova, and Alexander Sherman. "The Duflo–Serganova functor, vingt ans aprés". *Journal of the Indian Institute of Science* 102.3 (2022), pp. 961–1000.
- [Gru00] Caroline Gruson. "Sur l'idéal du cône autocommutant des super algèbres de Lie basiques classiques et étranges". Ann. Inst. Fourier (Grenoble) 50.3 (2000), pp. 807–831. URL: http://www.numdam.org/item?id=AIF_2000__50_3_807_0.
- [GW23] Niklas Garner and Brian R Williams. "Raviolo vertex algebras" (2023). arXiv: 2308.04414.
- [Jen22]
 L. Andrew Jenkins. The Nilpotent Cone for Classical Simple Lie Superalgebras. Thesis (Ph.D.)–University of Georgia. ProQuest LLC, Ann Arbor, MI, 2022, p. 66. URL: http://gateway.proquest.com/openurl?url_ver=Z39.88-2004&rft_val_fmt=in
- [Joh95] A Johansen. "Twisting of $\mathcal{N} = 1$ SUSY gauge theories and heterotic topological theories". *International Journal*
- of Modern Physics A 10.30 (1995), pp. 4325–4357. arXiv: hep-th/9403017 [hep-th].
- [Kap06] Anton Kapustin. "Holomorphic reduction of N = 2 gauge theories, Wilson-'t Hooft operators, and S-duality" (2006). arXiv: hep-th/0612119 [hep-th].

- [KW07] Anton Kapustin and Edward Witten. "Electric-magnetic duality and the geometric Langlands program". *Communications in Number Theory and Physics* 1 (2007), pp. 1–236.
- [KZ12] Johan Källén and Maxim Zabzine. "Twisted supersymmetric 5D Yang-Mills theory and contact geometry". Journal of High Energy Physics 2012.5 (2012), p. 125. arXiv: 1202.1956 [hep-th].
- [Mac77] G. Mack. "All unitary ray representations of the conformal group SU(2,2) with positive energy". *Commun. Math. Phys.* 55 (1977), p. 1.
- [Min98] Shiraz Minwalla. "Restrictions imposed by superconformal invariance on quantum field theories". *Adv. Theor. Math. Phys.* 2 (1998), pp. 783–851. arXiv: hep-th/9712074.
- [MSS02] Martin Markl, Steve Shnider, and Jim Stasheff. *Operads in algebra, topology and physics*. Vol. 96. Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2002, pp. x+349.
- [Nah78] W. Nahm. "Supersymmetries and their Representations". Nucl. Phys. B 135 (1978), p. 149.
- [Sch08] M. Schottenloher. A mathematical introduction to conformal field theory. Second. Vol. 759. Lecture Notes in Physics. Springer-Verlag, Berlin, 2008, pp. xvi+249.
- [Sch14] Claudia Scheimbauer. "Factorization homology as a fully extended topological field theory". PhD thesis. ETH Zurich, 2014.
- [Shn88] Steven Shnider. "The superconformal algebra in higher dimensions". Lett. Math. Phys. 16.4 (1988), pp. 377– 383. URL: https://doi-org.amherst.idm.oclc.org/10.1007/BF00402046.
- [Sug59] Mitsuo Sugiura. "Conjugate classes of Cartan subalgebras in real semi-simple Lie algebras". J. Math. Soc. Japan 11 (1959), pp. 374–434. URL: https://doi.org/10.2969/jmsj/01140374.
- [Wit88] Edward Witten. "Topological Quantum Field Theory". Commun. Math. Phys. 117 (1988), p. 353.
- [Wol69] Joseph A Wolf. "The action of a real semisimple group on a complex flag manifold. I: Orbit structure and holomorphic arc components". *Bulletin of the American Mathematical Society* 75.6 (1969), pp. 1121–1237.