

# Half-century of Efros-Shklovskii Coulomb gap. Romance with Coulomb interaction and disorder.

B. I. Shklovskii

*School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA*

(Dated: April 1, 2024)

The Efros-Shklovskii (ES) Coulomb gap in the one-electron density of localized states and the ES law of the variable range hopping conductivity were coined 50 years ago. The theory and its first confirmations were reviewed in ES monograph published 40-years ago. This paper reviews the subsequent experimental evidence, theoretical advancements, and novel applications of the ES law. Out of hundreds of experimental validations of the ES law in a diverse range of materials, I focus on those where the dynamic range of conductivity exceeds four orders of magnitude. These include three- and two-dimensional semiconductors under both zero and high magnetic fields, localized phases in the quantum Hall effect, granular metals, nanocrystal arrays, and conducting polymers. Additionally, I discuss the non-ohmic ES law and the Coulomb gap near insulator-metal transition. Recent developments of other concepts of the ES book are also discussed.

## I. INTRODUCTION

50 years ago Efros and Shklovskii (ES) penned the paper [1] “Coulomb gap and low temperature conductivity of disordered systems”. It elucidated how Coulomb interaction between localized electrons in semiconductors creates the power law Coulomb gap of the one-electron density of states near the Fermi level and changes temperature dependence of Mott variable hopping conductivity to the ES law.

Localized electron states in lightly doped semiconductors are formed by randomly positioned donors. Typically semiconductors are compensated, i. e.  $n$ -type semiconductor with concentration of donors  $N_D$  may have a concentration  $N_A < N_D$  of acceptors. At low temperatures each acceptor captures an electron from a donor and becomes negative. As a result  $N_A$  donors become positive while  $N_D - N_A$  donors are still neutral. Charged donors and acceptors create random potential which shifts energy of the donor  $i$  up or down by a value  $\epsilon_i$ . This broadens originally  $\delta$ -functional donor peak of the density of states. At low temperature lower energy occupied donor levels are separated by the Fermi level (chemical potential)  $\mu$  from higher empty levels. Before 1975 it was believed that near the Fermi level  $\mu$  the density of states (DOS)  $g(\epsilon) = g(\mu) = Const$ . ES showed [1] that due to the Coulomb interaction between electrons

$$g(\epsilon) = \frac{\pi \kappa^3 (\epsilon - \mu)^2}{3 e^6}. \quad (1)$$

in a three-dimensional (3D) semiconductor and

$$g(\epsilon) = \frac{\pi \kappa^2 |\epsilon - \mu|}{2 e^4}. \quad (2)$$

in a two-dimensional (2D) one. Here  $\kappa$  is the dielectric constant of the semiconductor (and in 2D of its environment),  $e$  is the proton charge and coefficients  $\pi/3$  and  $\pi/2$  were found with help of self-consistent approximation [2]. Thus, the density of states has a power law gap at the Fermi level, which is termed the Coulomb gap by ES. One can say that the Coulomb gap originates due to the exciton effect, i. e. attraction between an electron excited from occupied to an empty in the ground state donor to the hole it left at the original donor. The major consequence of this gap is replacement of the Mott variable

range hopping conductivity law

$$\sigma = \sigma_0 \exp \left\{ - \left( \frac{T_M}{T} \right)^p \right\}, \quad (3)$$

with  $p = 1/4$  for three-dimensional (3D) and  $p = 1/3$  for 2D cases by the ES law

$$\sigma = \sigma_0 \exp \left\{ - \left( \frac{T_{ES}}{T} \right)^{1/2} \right\}, \quad (4)$$

valid for both 3D and 2D cases. Here  $T$  is the temperature in energy units,  $T_M = 22/g(\mu)\xi^3$  for 3D and  $T_M = 13/g(\mu)\xi^2$  for 2D,  $\xi$  is the localization length of localized states at the Fermi level (for a lightly doped semiconductors  $\xi$  is equal to the Bohr radius of a donor  $a$ ),

$$T_{ES} = \frac{C e^2}{\kappa \xi}, \quad (5)$$

where  $C = 2.8$  in 3D and  $C = 6$  in 2D [3, 4].

Derivations of Eqs. (1) and (4) and first experimental and numerical conformations of them are discussed at length in Shklovskii-Efros (SE) book [3]. The Russian original of SE book was published in 1979, while the updated and extended English translation was published in 1984, 40 years ago. SE book is still widely used.

I devote this article to the double 50/40 years anniversary. The main part of the article reviews new observations, generalizations and applications of Eqs. (1) and (4) while the second part briefly mentions developments of other ideas of SE book. I try to avoid repetitions of the SE book and refer to SE book chapters and figures starting from letters SE.

The plan of this short article is as follows. I dwell on major experimental confirmations of the ES Coulomb gap and ES law in Section II. Section III is devoted to non-Ohmic generalization of the ES law. In Section IV I discuss new advances of the Coulomb gap and ES law theory. I focus on the evolution of the Coulomb gap near metal-insulator transition in Section V. Section VI goes beyond the Coulomb gap and SE book and dwells on related problems of Coulomb interaction and disorder physics. I comment on developments of ideas of specific chapters of the SE book in Section VII.

## II. NEW EXPERIMENTAL CONFIRMATIONS OF COULOMB GAP AND ES LAW

An experimental verification of ES law Eq. (4) is done by plotting  $\ln \sigma$  as a function of  $T^{-1/2}$  and checking whether this plot can be fitted by a straight line. In order to reliably discriminate ES law from Eq. (3) with  $p = 1/4$  and particularly in 2D case with  $p = 1/3$  the dynamic range of measured conductivity should cover approximately 3 orders of magnitude. In this review I dwell only on the most spectacular data which cover at least 4 orders of magnitude.

I start from observations of the ES law in 3D elementary semiconductor crystals Ge and Si. First high quality data for compensated Ge were discussed Chapters 9 and 10 of the SE book. Later the ES law was observed in neutron-transmutation-doped  $p$ -Ge in 4 orders of magnitude range [5] and even in 6 orders of magnitude range [6]. Similar high quality data were obtained in Si samples in 5 orders of magnitude range [7]. It is remarkable that the data [5, 7] were obtained in the process of development of Ge and Si based bolometers for astrophysics and dark matter research.

Other spectacular 3D data for the ES law were obtained in CdSe nanocrystal arrays in 6 orders of magnitude range [8], in ZnO nanocrystal films in 4 orders of magnitude range [9], in magnetite nanocrystal arrays in 7 orders of magnitude range [10], in granular metals in insulator in 4-8 orders of magnitude ranges [11, 12], in different conducting polymers [13–19] in 4-10 orders of magnitude ranges, and in graphene oxide thin films [20, 21] in 4-5 orders of magnitude ranges.

Due to scattering of the tunneling electron on other non-resonant donors the ES law should work even in strong magnetic fields [22]. Indeed, the data obtained on InSb samples [23] and shown in SE Fig. 9.2b demonstrates excellent agreement with the ES law in 7 orders of magnitude range of resistance.

Let us now switch to 2D systems with localized electron states. The ES law was observed in 6 orders of magnitude range in the gated GaAs/GaAlAs low mobility heterostructures in strong a orthogonal magnetic fields [24] which shrinks localized state wave functions and makes electron hops shorter. At smaller magnetic fields, where the hop length becomes as large as the distance to the gate the authors observed the transition from ES law (4) to 2D Mott law (3) with  $p = 1/3$ . They interpreted this transition as a result of screening of electron-electron interactions in two-dimensional electron gas (2DEG) by the metallic gate. This observation confirms that ES law results from the Coulomb interaction between electrons.

Discovery of the Quantum Hall effect (QHE) in higher mobility samples lead to observations of ES law in 4 orders of magnitude range of the diagonal conductivity  $\sigma_{xx}$  at integer filling factors  $\nu$ , where the electron states at the Fermi level are localized [25, 26].

Ref. [27] argued that at low enough temperatures the ES law should be observed not only at integer  $\nu$ , but everywhere between two neighboring half-integer  $\nu_c$  where electron states are still localized. The localization length  $\xi$  diverges at a Landau level as  $\xi \propto |\nu - \nu_c|^{-\gamma}$  with  $\gamma = 7/3$  leading to

vanishing  $T_{ES}(\nu)$  at all half-integers. At a finite  $T$  equation  $T_{ES}(|\nu - \nu_c|) = T$  defines the temperature dependent half-width  $|\nu - \nu_c| \propto T^{1/\gamma}$  of a the peak of the resistivity  $\rho_{xx}$  near  $\nu_c$  in agreement with experimental data [28–30].

Predictions of Ref. 27 were also confirmed by comprehensive studies of ES variable range hopping in GaAs [31, 32] and graphene [33]. In these papers ES law for  $\sigma_{xx}$  was observed in 4-5 orders of magnitude dynamic range and the localization length  $\xi(\nu)$  was extracted from measured values of  $T_{ES}$ . It was found to diverge at  $\nu_c$  with the theoretical critical index  $7/3$ . Ref. 33 also studied transition from ES to Mott law due to screening of Coulomb interaction by the gate, again confirming the Coulomb origin of ES law. (In the absence of screening the ES law may yield to the Mott law only when bare (non-Coulomb) DOS  $g(\mu)$  is very small, so that the Coulomb gap width determined by equating Eqs. (1) and (2) to  $g(\mu)$  is narrower than the band energies around the Fermi level involved in the variable range hopping. Such a situation takes place in amorphous semiconductors, where the Mott law was originally observed. Even there the ES law should work at low enough temperatures, but this requires measuring unrealistically large resistances.)

So far we talked about the integer QHE. The ES law was observed also in the middles of the fractional QHE plateaus [34]. One can interpret such an observation assuming that fractional charges  $q$  (say  $q = -e/3$ ) bound to donors form the Coulomb gap of DOS. The variable range hopping of these fractional charges then obeys the ES law with  $T_{ES} = Cq^2/\kappa\xi$ , where  $\xi$  is the localization length of a charge  $q$ . Small values of  $q$  lead to a very small  $T_{ES}$  in agreement with experiment [34]. Then, a straightforward extension of the logic of Ref. 27 leads to the ES law for  $\sigma_{xx}$  everywhere in a fractional plateau. The divergence of the localization length near the filling factor border between two neighboring fractions leads to the prediction of the same behavior of widths  $\rho_{xx}$  peaks with growing temperature as in the integer QHE. This prediction agrees with experimental results [35–38].

## III. NON-OHMIC ES LAW

So far we discussed the Ohmic variable range hopping conductivity in small electric fields. It was shown theoretically [39, 40] that at very low temperatures the strong electric field  $E$  acts as the effective temperature  $e\xi E/2$ . As a result Eqs. (1) and (2) lead to the exponential growth of the current  $I$  with  $E$  (the non-Ohmic ES law)

$$I = I_0 \exp \left\{ - \left( \frac{E_0}{E} \right)^{1/2} \right\}, \quad (6)$$

where

$$E_0 = \frac{2T_{ES}}{e\xi} = \frac{2Ce}{\kappa\xi^2}. \quad (7)$$

Fitting the lowest temperature data with Eq. (7) and the temperature dependence of the Ohmic conductivity with Eq. (4) and using the resulting  $T_{ES}$  and  $E_0$  one can find unknown  $\xi$  and

$\kappa$ . This is particularly important in 3D cases where most measurements are done close to the insulator-metal transition, at which both  $\xi$  and  $\kappa$  diverge [27]. The non-Ohmic ES law was observed in large dynamic ranges of currents in Refs. 8, 20, and 21. These authors then found both  $\xi$  and  $\kappa$ .

Eq. (7) was also generalized for the low temperature variable range hopping in the quantum Hall effect, where the role of electric field  $E$  is played by the Hall electric field [27]. This allowed to predict that the half-width  $|\nu - \nu_c|$  of  $\rho_{xx}$  peaks at a half-integer  $\nu_c$  grows with the current  $I$  as  $I^{1/2\gamma}$ . This prediction was confirmed by Refs. 31–33, and 41. Temperature and current dependencies of the  $\rho_{xx}$  peaks widths based on the ES law [27] provide a microscopic interpretation to the scaling theory [42].

#### IV. COULOMB GAP AND ES LAW: NEW THEORETICAL DEVELOPMENTS

Multiple numerical studies of the Coulomb gap density of states confirmed power laws Eqs. (1), (2) for three and two-dimensional systems [43–48]. However, they did not find exponential depletion of the one-electron density of states at smallest energies due to the electron polaron effects predicted for three-dimensional case [2] and discussed in Chapter 10 of the SE book. This puzzle was resolved in Ref. 49. In two dimensions the ES law was confirmed by direct numerical modelling of the variable range hopping conductivity [50].

As we saw in Sec. II the ES law is observed in many granular metals and nanocrystal arrays [8, 9, 11, 12, 51]. Theory of the Coulomb gap and the ES law in such systems was developed later than for lightly doped semiconductors, because it required understanding of the origin of disorder. For the hopping conductivity in arrays of metal granules disorder is related to randomly positioned charged impurities in the insulator surrounding metallic granules [52–54]. The role of charged impurities in a system of isolated long metallic needles or flexible wires dispersed and frozen in an insulator was studied in Ref. 55 and 56. An interesting, but more complicated case of quasi-one-dimensional crystals with impurities was studied in Refs. 57 and 58.

On the other hand, in a periodic epitaxially connected nanocrystal array the disorder originates from random number of donors in individual nanocrystals [59–61].

As we saw above, the ES law data allows us to find the localization length of localized states with energy close to the Fermi level. This required the development of a new theory of localization length. It was shown that in granular metals long distance tunneling happens through a chain of neighboring granules or via relay co-tunneling [52–56, 62, 63]. In epitaxially connected nanocrystals similar relay co-tunneling happens through necks connecting nearest neighbor nanocrystals [59–61].

In doped semiconductors the related concept of tunneling with virtual scattering on impurities was developed in Ref. 22, 64–66. Later it was applied to theories of negative magnetoresistance, Aharonov-Bohm oscillations and mesoscopic transport in the range of ES law [67–71]. In QHE the




Figure 1. Schematic plot of DOS  $g(\epsilon)$  in the vicinity of critical concentration of the metal-insulator transition  $N_c$  [73]. DOS  $g(\epsilon)$  at  $N = N_c$  is shown by the thick line. Thin lines show  $g(\epsilon)$  at  $N = 2N_c$  and  $N = N_c/2$ . Dashed lines show  $g(\epsilon)$  for  $N = 1.25N_c$  and  $N = 0.8N_c$  at  $|\epsilon - \mu| < T_{ES}$  where they deviate from the thick line. On the metallic side the Altshuler-Aronov dip [74] at  $\epsilon = \mu$  is not shown.

localization length of localized states between two half-integers is formed by a chain of virtually scattered electron cyclotron orbits [72].

#### V. COULOMB GAP NEAR METAL-INSULATOR TRANSITION

The shape of the Coulomb gap of DOS in a doped 3D semiconductor with concentration of donors  $N$  close to the critical concentration  $N_c$  of the MIT was studied in Ref. 73 both theoretically and experimentally. It is known that on the insulator side of MIT the long wavelength dielectric constant  $\kappa$  diverges, so that according to Eq. (1) close to  $N_c$  DOS  $g(|\epsilon - \mu|)$  parabola becomes very steep. On the other hand, on the metallic side of MIT metal  $g(\epsilon)$  is large and weakly energy dependent at small  $|\epsilon - \mu|$ . In order to understand how crossover between these two limits happens it was assumed [73] that at MIT  $\kappa(q) = \kappa/qa$  and the Coulomb potential  $V(r) = e^2a/\kappa r^2$ , where  $a$  is the Bohr radius of the donor. (I use a simplified version of the scaling ansatz [73] with  $\eta = 2$  which is in good agreement with experiments).

Then, using ES arguments of Ref. 1 one arrives at the critical Coulomb gap DOS

$$g_c(\epsilon) = \frac{\kappa^{3/2}|\epsilon - \mu|^{1/2}}{a^{3/2}e^3}. \quad (8)$$

(see Fig. 1). In the vicinity of the MIT at  $|1 - N/N_c| \ll 1$  the large correlation (localization) length  $\xi = a|1 - N/N_c|^{-\nu}$  truncates the growth of the dielectric constant  $\kappa(q)$  at  $\kappa = \kappa(1/\xi)$  and defines the energy scale  $T_{ES} = e^2/\xi\kappa(1/\xi)$ . Then at  $|\epsilon - \mu| < T_{ES}$  on the insulating side of the MIT we recover Eq. (1) with  $\kappa = \kappa(1/\xi)$ , while on the metallic side  $g(\epsilon) = g_c(T_{ES}) = Const$  (see Fig. 1). ES law is valid at  $T < T_{ES}$  because it uses only the parabolic part of DOS.

Remarkably McMillan [75] arrived essentially at the same scaling ansatz but only for the metallic side of the MIT. The McMillan-Shklovskii scaling ansatz was supported by two recent theoretical papers [76, 77].

The Coulomb gap evolution near the MIT was studied by direct tunneling into Si crystals [73] with impurity concentrations  $0.9N_c < N < 1.1N_c$ . Results of this experiment are in agreement with McMillan-Shklovskii scaling ansatz.

The McMillan-Shklovskii scaling ansatz is based on the absence of the local screening of the Coulomb gap in a given realization of random coordinates of donors and acceptors discovered in numerical experiment Ref. 78, discussed in Ref. 73 and later supported by Refs. 48 and 79. Indeed, when we bring a new electron on an empty donor in the center of a lightly doped semiconductor sample it induces polarization either zero or small number of pairs, i. e. transitions of other electrons from occupied to empty donors. These pairs typically form a chain moving an electron charge away from the center in a particular direction, so that potential of the central new electron at distance  $r$  remains of the order  $e^2/\kappa r$ , but may change its sign. Thus, in a system of localized states the local screening is absent and does not affect the Coulomb gap. To arrive at the exponential Yukawa screening of the Coulomb potential we have to average over many realizations of random coordinates of surrounding central electron donors and acceptors. Such a procedure leads to averaging over many directions of polarization chains and makes the central electron polarization cloud spherically symmetric, but apparently it has nothing to do with ES law physics. On the other hand, when one calculates the thermodynamic DOS describing penetration of uniform electric field into orthogonal macroscopic slab of a lightly doped semiconductor, the averaging of local polarization over its surface makes perfect sense and leads to the finite DOS and the exponential Yukawa screening with the screening radius of the order of average distance between donors [78].

## VI. BEYOND COULOMB GAP

In this section I dwell on problems with the Coulomb interaction and disorder which are practically missing in the ES book. As we saw above, the Coulomb gap of the one-electron DOS determines the DC variable range hopping conductivity and the tunneling current into a semiconductor. On other hand, the high frequency the phononless hopping conductivity  $\sigma(\omega)$  is determined by electron hopping in compact resonance pairs of donors with small arms  $r_\omega$  populated by one electron only. The density of states of these pairs does not have a Coulomb gap and is (with logarithmic accuracy) constant at small excitation energies [1, 3]. So at the first sight the Coulomb interaction is not important here and the low temperature terahertz range phononless conductivity  $\sigma(\omega) \propto \omega^2$  as proposed by Mott [80] and Berezinskii [81]. However, we predicted [82, 83] that at gigahertz frequencies when the Coulomb energy of two electrons repulsion  $e^2/\kappa r_\omega$  exceeds  $\hbar\omega$  one power of  $\hbar\omega$  should be replaced by  $e^2/\kappa r_\omega$ , what leads to  $\sigma(\omega) \propto \omega$ . This happens because the concentration of pairs having one electron is proportional to  $e^2/\kappa r_\omega$ .

Multiple experiments [84–89] confirmed this prediction. Compact pairs are also responsible for the electron contribution to the specific heat of systems with localized states [90].

Far on the other side of the frequency spectrum, at very low frequencies there is the  $1/f$  noise of the hopping conductivity. It was studied experimentally [7] and theoretically [91–93].

The hopping conductivity in strong electric fields is another big field practically missing in the SE book. In Section III we dealt with very low temperature and very strong electric field where electric field leads to exponential growth of effective temperature. However, there are many intermediate cases. At a strong disorder, large concentrations of carriers and low temperatures the current always exponentially grows with the applied electric field [94–99]. This growth can be interpreted as the effect of reduction of large hopping energy barriers by the applied electric field.

However, at small concentrations of carriers and large enough temperatures the current decreases in a strong electric field. This happens because a very strong electric field does not allow electrons to use return parts in optimal tortuous percolation chains of hops of the ohmic conductivity [100, 101] and makes geometrical traps for electrons from them. Resulting negative differential conductivity was discovered in lightly doped and weakly compensated Si samples [102], where it led to new Gunn-like oscillations.

Hopping also plays important role in many related relaxation phenomena. The energy relaxation via hopping, the hopping photoconductivity, the impurity assisted recombination of electrons and holes, and the hopping relaxation of spin polarization were studied in Refs. 103–109.

Correlated arrangement of electrons in a lightly doped semiconductor leading to the Coulomb gap is similar to a pinned by disorder Wigner Crystal. Therefore, I returned to Wigner crystals several times [110–113]. While studying spin splitting of Landau levels in 2DEG [114] we were thinking whether electrons of a spin split high Landau level can form a Wigner crystal. We discovered that instead the peculiar box type of the electron-electron repulsion potential at Landau Levels with quantum numbers  $N \geq 2$  should lead to stripe and bubble phases [115, 116].

Predicted phases were discovered in Refs. 117 and 118 (see the review of the first 6 years of stripes and bubbles [119]). Multiple bubble phases with different number of electrons per bubble were observed in GaAs [120, 121] and graphene [122, 123]. They are direct relatives of the low density Wigner crystal.

Following ideas of the Ref. 124 we calculated anisotropy of the resistivity in a stripe phase and found a good agreement with the experimental data for cleanest GaAs samples where resistance anisotropy reaches 3000 [125]. We also studied what the increasing scattering by a short range disorder does to a stripe phase and its resistivity tensor. We discovered "hidden stripe phases" where the stripe order survives a growing short range disorder, but the resistivity is already isotropic [126, 127].

Theory [128] predicted that the compressibility of a low-density two-dimensional electron gas in the Wigner crystal state and Wigner-crystal-like liquid state is negative.

This counter-intuitive prediction was confirmed by experiments [129, 130]. When such a 2DEG is on both sides of a plane capacitor with a very small distance between plates its capacitance was predicted to become much larger than the geometrical one due to the Coulomb interplane electron correlations [131, 132]. It was also predicted that a 3D array of nanocrystals gated by a penetrating inside the array ionic liquid may abruptly get a finite charge at some critical voltage. At this critical voltage the differential capacitance of the array is infinite [133].

We would like to mention here only three experimental confirmations of predicted anomalously large capacitance [134–136]. Refs. 131–133 have more experimental references.

On the other hand, we used the concept of a classical Wigner crystal beyond solid state physics, namely in the chemistry and biology of water solutions with multivalent ions. We showed that the Wigner-crystal-like correlated liquid of multivalent counterions adsorbed at the surface of a negative macroion such as the double-helix DNA inverts the DNA net charge making DNA positive. A cell membrane is negative and repels negative naked DNA while a charge inverted positive DNA is allowed to enter the cell. Therefore, DNA charge inversion by positive polymers is used for the gene delivery [137, 138] and in modern mRNA vaccines [139].

Understanding of ion correlations in a narrow biological ion channel allows to solve the puzzle of deconfinement of pairs of positive and negative ions which leads to practically zero activation energy of the channel ion transport at ambient conditions [140, 141]. This deconfinement is similar to that of quarks in hot quark-gluon plasma [142].

## VII. SE BOOK AT 40. COMMENTS TO ITS CHAPTERS.

Chapter 2. Substantial progress in identifying the metal-insulator transition (MIT) point and its critical indexes is achieved by analyzing the distribution function  $P(s)$  of the level spacings  $s$ . It was discovered [143] that on the top of two known universal  $P(s)$ , Wigner-Dyson and Poisson ones, there is the third universal  $P(s)$  realized exactly at MIT. It behaves as Wigner-Dyson one at  $s < 1$  and is Poisson-like at  $s \gg 1$ .

We showed that for Anderson transition one can characterize

the function  $P(s)$  for each disorder strength  $W$  and the system size  $L$  by the area  $\gamma$  of its  $s > 2$  tail. Then doing the finite size analysis for the function  $\gamma(W, L)$  one easily finds the transition point and the index of correlation length. This method was widely used to identify different MIT transitions [144] until a new more convenient way of characterization of the function  $P(s)$  called  $r$ -method [145] replaced the use of  $\gamma(W, L)$ . One can also use the spectral rigidity to find an MIT threshold [146].

Later we extended  $r$ -method to the non-Hermitian analog of Anderson Hamiltonian where complex diagonal elements are distributed randomly in a square in the complex plane [147]. For this Hamiltonian we discovered the non-Hermitian Anderson transition at in three-dimensions and showed that there is no Anderson transition in two-dimensions [148, 149].

Chapter 3. Calculation of the activation energy of the nearest neighbor hopping conductivity of a 3D weakly compensated semiconductor is generalized for the 2D case [150] in order to describe surface conductivity of Si induced by ion liquid gating.

Chapter 5. A new application of percolation theory is proposed [151] to describe bulk magnetization induced by ion liquid gating of an oxide film.

Chapter 12. SE book section (12.3) deals with filtering of a random potential from its small scales via the wave function averaging. Recently a new way to do the filtering was suggested [152] and later compared with the conventional one [153].

Chapter 13. Theory of nonlinear screening of a long-range potential developed in Chapters 3 and 13 was applied to nonlinear screening of the random potential of remote impurities by 2DEG in QHE devices [114, 154–163]. The concepts of strongly or completely compensated semiconductor found an application to topological insulators [164–168], to other narrow gap semiconductors in subjected to a strong 3D Coulomb disorder [169, 170] and to the superconductor-insulator transition in high  $T_c$  superconductors [171].

## ACKNOWLEDGMENTS

I am grateful to all my coworkers and in particular to A. L. Efros for enjoyable collaborations.

- 
- [1] A. L. Efros and B. I. Shklovskii, J. Phys. C **8**, L49 (1975).
  - [2] A. L. Efros, Coulomb gap in disordered systems. J. Phys. C9, 2021 (1976).
  - [3] B. I. Shklovskii and A. L. Efros, "Electronic Properties of Doped Semiconductors" (Springer-Verlag, Berlin, 1984).
  - [4] Nguyen Van Lien, Fiz. Tech. Poluprovodn. **18**, 335 (1984) [Sov. Phys. Semicond. **18**, 207 (1984)].
  - [5] K. M. Itoh, W. L. Hansen, J. W. Beeman, E. E. Haller, J. W. Farmer, V. I. Ozogin, J. Low Temp. Phys. **93** 307 (1993).
  - [6] W. Schoepe, Z. Phys. B - Condensed matter **71** 455 (1988).
  - [7] Dan McCammon, in "Cryogenic Particle Detection", pages 35-62, Springer, Berlin, Heidelberg, (2005); arXiv:physics/0503086.
  - [8] Dong Yu, Congjun Wang, Brian L. Wehrenberg, Philippe Guyot-Sionnest, Phys. Rev. Lett. **92**, 216802-1 (2004).
  - [9] Benjamin L. Greenberg, Zachary L. Robinson, K. V. Reich, Claudia Gorynski, Bryan N. Voigt, Lorraine F. Francis, B. I. Shklovskii, Eray S. Aydil, Uwe R. Kortshagen, Nano Lett. **17**, 4634 (2017).
  - [10] Hao Zeng, C. T. Black, R. L. Sandstrom, P. M. Rice, C. B. Murray, Shouheng Sun, Magneto-transport of magnetite nanoparticle arrays, Phys. Rev. B **73**, 020402 (2006).
  - [11] B. Abeles, Ping Sheng, M. D. Coutts, Y. Arie, Structural and electrical properties of granular metal films. Adv. Phys. **24**, 407 (1975).
  - [12] T. Chui, G. Deutscher, P. Lindenfeld, W. L. McLean, Conduc-

- tion in granular aluminum near the metal-insulator transition. *Phys. Rev. B* 23, 6172 (1981).
- [13] L. Zuppiroli, M. N. Bussac, S. Paschen, O. Chauvet, L. Forro, *Phys. Rev. B* 50, 5196 (1994).
- [14] Fulin Zuo, Marie Angelopoulos, Alan G. MacDiarmid, Arthur J. Epstein, *Phys. Rev. B* 36, 3475 (1987).
- [15] B. Sixou, N. Mermilliod, J. P. Travers, *Phys. Rev. B* 53, 4509 (1996).
- [16] C.O. Yoon, M. Reghu, D. Moses, A.J. Heeger, Y. Cao, T.-A. Chen, X. Wu, R.D. Rieke, *Synthetic Metals* 75, 229 (1995).
- [17] Yongqiang Wang, S. S. Mohite, L. B. Bridwell, R. E. Giedd, C. J. Sofield, *J. Mater. Res.*, 8, 388 (1993).
- [18] A. B. Kaiser, *Rep. Prog. Phys.* 64, 1 (2001).
- [19] Shun Wang, Mingjing Ha, Michael Manno, C. Daniel Frisbie, C. Leighton, *Nature Communications* 3, 1210 (2013).
- [20] Daeha Joung, S. I. Khondaker, *Phys. Rev. B* 86, 235423 (2012).
- [21] R. M. Rudenko, O. O. Voitsihovska, A. A. Abakumov, I. B. Bychko, V. N. Poroshin, *Diamond and Related Materials*, 141, 110606 (2024).
- [22] B. I. Shklovskii, Variable-range hopping conduction in a strong magnetic field. *Zh. Eksp. Theor. Fiz. Pis. Red.* 36, 43 (1982). [English transl.: *Sov. Phys.-JETP Lett.* 36, 53 (1982)].
- [23] H. Tokumoto, R. Mansfield, M. J. Lea, Correlation effects in variable-range hopping in n-InSb. *Solid State Commun.* 35, 961 (1980).
- [24] F. W. Van Keuls, X. L. Hu, H. W. Jiang, A. J. Dahm, Screening of the Coulomb interaction in two-dimensional variable-range hopping. *Phys. Rev. B* 56, 1161 (1997).
- [25] G. Ebert, K. von Klitzing, C. Probst, E. Schuberth, K. Ploog, G. Weinmann, Two-dimensional magneto-quantum transport on GaAs-AlxGa(1-x)As heterostructures under non-ohmic conditions. *Solid State Commun.* 45, 625 (1983).
- [26] A. Briggs, Y. Gulder, J. P. Vierende, M. Voos, J. R. Hirtz, J. P. Razeghi, Low-temperature investigations of the quantum Hall effect in InxGa(1-x)As-InP heterojunctions. *Phys. Rev. B* 27, 6549 (1983).
- [27] D. G. Polyakov, B.I. Shklovskii, Conductivity peak broadening in the quantum Hall regime, *Phys. Rev. B* 48, 11167 (1993).
- [28] H. P. Wei, D. C. Paalanen, and A. M. M. Pruisken, *Phys. Rev. Lett.* 61, 1294 (1988).
- [29] H. P. Wei, S. Y. Lin, D. C. Tsui, and A. M. M. Pruisken, *Phys. Rev. B* 45, 3926 (1992).
- [30] S. Koch, R. J. Haug, K. v. Klitzing, and K. Ploog, *Phys. Rev. B* 46, 1596 (1992).
- [31] M. Furlan, Electronic transport and the localization length in the quantum Hall effect, *Phys. Rev. B* 57, 14818 (1998).
- [32] F. Hohls, U. Zeitler, and R. J. Haug, Hopping Conductivity in the Quantum Hall Effect: Revival of Universal Scaling, *Phys. Rev. Lett.* 88, 036802 (2002).
- [33] K. Bennaceur, P. Jacques, F. Portier, P. Roche, and D. C. Glattli, Unveiling quantum Hall transport by Efros-Shklovskii to Mott variable-range hopping transition in graphene, *Phys. Rev. B* 86, 085433 (2012).
- [34] G. S. Boebinger, H. L. Stormer, D. C. Tsui, A. M. Chang, J. C. M. Hwang, A. Y. Cho, C. W. Tu, G. Weimann, Activation energies and localization in the fractional quantum Hall effect, *Phys. Rev. B* 36, 7919 (1987).
- [35] L. Engel, H. Wei, D. Tsui, and M. Shayegan, *Surface science* 229, 13 (1990).
- [36] T. Machida, S. Ishizuka, S. Komiyama, K. Muraki, and Y. Hirayama, *Physica B: Condensed Matter* 298, 182 (2001), International Conference on High Magnetic Fields in Semiconductors.
- [37] P. T. Madathil, K. A. Villegas Rosales, C. T. Tai, Y. J. Chung, L. N. Pfeiffer, K. W. West, K. W. Baldwin, and M. Shayegan, *Phys. Rev. Lett.* 130, 226503 (2023).
- [38] Simrandeep Kaur, Tanima Chanda, Kazi Rafsanjani Amin, Kenji Watanabe, Takashi Taniguchi, Unmesh Ghorai, Yuval Gefen, G. J. Sreejith, Aveek Bid, Universality of Anderson Localization Transitions in the Integer and Fractional Quantum Hall Regime, arXiv:2312.06194
- [39] B.I. Shklovskii, Hopping conduction in semiconductors in a strong electric field, *Fiz. Tekh. Poluprov.* 6, 2335 (1973) - Engl. transl.: *Sov. Phys.-Semicond.* 6, 1964 (1973).
- [40] S. Marianer, B.I Shklovskii, Effective temperature of hopping electrons in a strong electric field, *Phys. Rev. B* 46, 13100 (1992).
- [41] H. P. Wei, L. W. Engel, D. C. Tsui, Current scaling in the integer quantum Hall effect, *Phys. Rev. B* 50, 14609 (1994).
- [42] S. L. Sondhi, S. M. Girvin, J. P. Carini, and D. Shahar, *Rev. Mod. Phys.* 69, 315 (1997).
- [43] H. Davies, P. A. Lee, and T. M. Rice, *Phys. Rev. B* 29, 4260 (1984).
- [44] A. Mobius, M. Richter, B. Drittler, *Phys. Rev. B* 45, 11568 (1992).
- [45] A. Mobius, P. Karmann, M. Schreiber, *J. Phys. Conf. Ser.* 150, 022057 (2009).
- [46] Mobius and M. Richter, *Phys. Rev. Lett.* 105, 039701 (2010).
- [47] M. Goethe and M. Palassini, *Phys. Rev. Lett.* 103, 045702 (2009).
- [48] M. Palassini, M. Goethe, *Journal of Physics: Conference Series* 376, 012009 (2012).
- [49] A. L. Efros, Brian Skinner, B. I Shklovskii, Coulomb gap in the one-particle density of states in three-dimensional systems with localized electrons. *Phys. Rev. B* 84, 064204 (2011).
- [50] D. N. Tsygankov, A. L. Efros, Variable Range Hopping in Two-Dimensional Systems of Interacting Electrons, *Phys. Rev. Lett.* 88, 176602 (2002).
- [51] I. S. Beloborodov, A. V. Lopatin, V. M. Vinokur, and K. B. Efetov, Granular electronic systems, *Rev. Mod. Phys.* 79, 469 (2007).
- [52] J. Zhang, B. I. Shklovskii, Density of States and Conductivity of Granular Metal or Array of Quantum Dots, *Phys. Rev. B*, 70, 115317 (2004).
- [53] Tianran Chen, Brian Skinner, B. I. Shklovskii, Coulomb gap triptych in a periodic array of metal nanocrystals, *Phys. Rev. Lett.* 109, 126805 (2012).
- [54] Tianran Chen, Brian Skinner, B. I. Shklovskii, Coulomb gap triptychs, square root of two effective charge, and hopping transport in periodic arrays of superconductor grains, *Phys. Rev. B* 86, 045135 (2012).
- [55] Tao Hu, B. I. Shklovskii, Hopping conductivity of a suspension of nanowires in an insulator, *Phys. Rev. B* 74, 054205 (2006).
- [56] Tao Hu, B. I. Shklovskii, Hopping conductivity of a suspension of flexible wires in an insulator, *Phys. Rev. B* 74, 174201 (2006).
- [57] M. M. Fogler, S. Teber, B. I. Shklovskii, Variable-range hopping in quasi-one-dimensional electron crystals, *Phys. Rev. B* 69, 035413 (2004).
- [58] M. Bacani, M. Novaka, I. Kokanovic, D. Babic, Hopping electron transport in doped polyaniline: An experimental verification of the Fogler-Teber-Shklovskii model, *Synthetic Metals* 172, 28 (2013).
- [59] Brian Skinner, Tianran Chen, B. I. Shklovskii, Theory of hopping conduction in arrays of doped semiconductor nanocrystals, *Phys. Rev. B* 85, 205316 (2012).
- [60] Han Fu, K. V. Reich, B. I. Shklovskii, Hopping conductivity and insulator-metal transition in films of touching semiconductor nanocrystals, *Phys. Rev. B* 93, 125430 (2016).

- [61] Ting Chen, K. V. Reich, N. J. Kramer, Han Fu, U. R. Kortshagen, B. I. Shklovskii, Metal-Insulator transition in doped semiconductor nanocrystal films, *Nature Materials*, 15, 299 (2016).
- [62] B. I. Shklovskii, New problem of percolation theory and conduction of granular metals, *Fiz.-Tv. Tela* 26, 585 (1984).
- [63] Feigel'man, M. V., Ioselevich, A.S. Variable-range cotunneling and conductivity of a granular metal, *Jetp. Lett.* 81, 277 (2005).
- [64] I. M. Lifshitz, V. Va. Kirpichenkov, Tunnel transparency of disordered systems. *Zh. Eksp. Theor. Fiz.* 77, 989 (1979) [English transl.: *Sov. Phys.-JETP* 50, 499 (1979)].
- [65] B. I. Shklovskii, A. L. Efros: Tunnel transparency of disordered systems in a magnetic field. *Zh. Eksp. Theor. Fiz.* 84, 81 (1983) [English transl.: *Sov. Phys.-JETP* 57, 470 (1983)].
- [66] A. B. Khaetskii, B. I. Shklovskii: Anomalous tunneling across magnetic field near semiconductor surface. *Zh. Eksp. Theor. Fiz.* 85, 721 (1983) [English transl.: *Sov. Phys.-JETP* 58, 421 (1983)].
- [67] B. Z. Spivak, V. A. Kharchenko, B. I. Shklovskii, Scattering effect in VRH conduction, *Fiz. Tekh. Poluprov.* 19, 493 (1985).
- [68] V. L. Nguyen, B. Z. Spivak, B. I. Shklovskii, Tunnel hops in disordered systems, *Zh. Eksp.Theor. Fiz.* 89, 1770 (1985) - Engl. transl.: *Sov. Phys.-JETP* 62, 1021 (1985).
- [69] V. L. Nguyen, B. I. Shklovskii, B. Z. Spivak, Aharonov-Bohm oscillations with normal and superconductive flux-quanta in hopping conduction, *Pis'ma Zh. Eksp. Teor. Fiz.* 41, 35 (1985) - Engl. transl.: *Sov. Phys. JETP Letters* 41, 42 (1985).
- [70] B. I. Shklovskii and B. Z. Spivak, Interference phenomena in variable range hopping conductivity in "Hopping and related phenomena", edited by H. Fritzsche and M. Pollak, *World Scientific* (1990).
- [71] B. I. Shklovskii, B. Z. Spivak, Scattering and interference phenomena in variable range hopping conduction, in: "Hopping conduction in semiconductors," ed. by M. Pollak and B. I. Shklovskii, North-Holland, Amsterdam (1990).
- [72] M. M. Fogler, A. Yu. Dobin, B. I. Shklovskii, Localization length at the resistivity minima of the quantum Hall effect, *Phys. Rev. B* 57, 4614 (1998).
- [73] Mark Lee, J. G. Massey, V. L. Nguyen, B. I. Shklovskii, Coulomb Gap in a Doped Semiconductor near the Metal-Insulator Transition: Tunneling Experiment and Scaling Ansatz, *Phys. Rev. B* 60, 1582 (1999). (see corrected misprints in arXiv:cond-mat/9902169).
- [74] B. L. Al'tshuler, A. G. Aronov, Contribution to the theory of disordered metals in heavily doped semiconductors. *Zh. Eksp. Theor. Fiz.* 77, 2028 (1979) [English transl.: *Sov. Phys.-JETP* 50, 968 (1979)].
- [75] W. L. McMillan, Scaling theory of the metal-insulator transition in amorphous materials, *Phys. Rev. B* 24, 2739 (1981).
- [76] M. Amini, V. E. Kravtsov, M. Mueller, Multifractality and quantum to classical crossover in the Coulomb anomaly at the Mott-Anderson metal-insulator transition, *New J. of Physics*, 16, 015022 (2014).
- [77] I. S. Burmistrov, I. V. Gornyi, and A. D. Mirlin, Tunneling into the localized phase near Anderson transitions with Coulomb interaction, *Phys. Rev. B* 89, 035430 (2014).
- [78] S. D. Baranovskii, B. I. Shklovskii, A. L. Efros, Screening in localized electron system, *Zh. Eksp. Teor. Fiz.* 1793 (1984) - Engl. transl.: *Sov. Phys. JETP* 60, 1031 (1984).
- [79] S.-R. Eric Yang, Ziqiang Wang, A. H. MacDonald, Thermodynamic and tunneling density of states of the integer quantum Hall critical state, *Phys. Rev. B* 65, 041302(R)(2001).
- [80] N. F. Mott, *Philos. Mag.* 22, 7 (1970); N. F. Mott and E. F. Davis, "Electronic Processes in Non-Crystalline Materials", Oxford (1971).
- [81] V. L. Berezinskiy, *Zh. Eksp. Teor. Fiz.* 65, 125 (1973) [*Sov. Phys. JETP* 38, 620 (1974)].
- [82] B. I. Shklovskii, A. L. Efros: Phononless hopping conduction in disordered systems. *Zh. Eksp. Theor. Fiz.* 81, 406 (1981) [English transl.: *Sov. Phys.-JETP* 54, 218 (1981)].
- [83] A. L. Efros, B. I. Shklovskii, Electron-Electron Interaction in Disordered systems with Localized States, in "Electron-Electron Interaction in Disordered Systems", ed. by A. L. Efros, M. Pollak, North-Holland, Amsterdam (1985).
- [84] R. M. Lewis, J. P. Carini, Frequency scaling of microwave conductivity in the Integer Quantum Hall Effect Minima, *Phys. Rev. B* 64, 073310 (2000).
- [85] F. Hohls, U. Zeitler, R. J. Haug, High frequency conductivity in the Quantum Hall regime, *Phys. Rev. Lett.* 86, 5124 (2001).
- [86] Mark Lee, M. L. Stutzmann, Microwave ac Conductivity Spectrum of a Coulomb Glass, *Phys. Rev. Lett.* 87, 056402-1 (2001).
- [87] E. Helgren, N. P. Armitage, and G. Gruner, Electrodynamic of a Coulomb Glass in n-Type Silicon, *Phys. Rev. Lett.* 89, 246601-1 (2002).
- [88] E. Helgren, N. P. Armitage, and G. Gruner, Frequency-dependent conductivity of electron glasses, *Phys. Rev. B* 69, 014201 (2004).
- [89] M. Hering, M. Scheffler, M. Dressel and H. v. Löhneysen, *Phys. Rev. B* 75, 205203 (2007).
- [90] S. D. Baranovskii, A. A. Uzakov, A. L. Efros: Thermodynamic properties of impurity band electrons. *Zh. Eksp. Theor. Fiz.* 83, 756 (1982) [English transl.: *Sov. Phys.-JETP* 56, 422 (1982)].
- [91] B. I. Shklovskii, Theory of  $1/f$  noise in hopping conduction, *Solid State Commun.* 33, 273 (1980).
- [92] B. I. Shklovskii,  $1/f$  noise in variable range hopping conduction, *Phys. Rev. B* 67, 045201 (2003).
- [93] A. L. Burin, Yu. M. Galperin, V. Kozub, V. Vinokur, B. I. Shklovskii, Many- electron theory of  $1/f$ -noise in hopping conductivity, *Phys. Rev. B* 74, 075205 (2006).
- [94] B. I. Shklovskii, Non-ohmic hopping conduction. *Fiz. Tekh. Poluprov.* 10, 1440 (1976) [English transl.: *Sov. Phys.-Semicond.* 10, 855 (1976)].
- [95] B. I. Shklovskii, Percolation conductivity in strong electric fields. *Fiz. Tekh. Poluprov.* 13, 93 (1979) [English transl.: *Sov. Phys.-Semicond.* 13, 53 (1979)].
- [96] Levin E. I., Percolative nonohmic conduction in polycrystalline semiconductors. *Fiz. Tekh. Poluprov.* 17, 255 (1983) [English transl.: *Sov. Phys.-Semicond.* 17, 158 (1983)].
- [97] E. I. Levin, Nguyen Van Lien, B. I. Shklovskii, Hopping electrical conduction in strong electric fields: A numerical computer experiment, *Fiz. Tekh. Poluprov.* 16, 815 (1982) [Engl. transl.: *Sov. Phys. Semicond.* 16, 523 (1982)].
- [98] E. I. Levin, B. I. Shklovskii, Low-temperature hopping conduction in strong electric field. Computer experiment, *Fiz. Tekh. Poluprov* 18, 856 (1984) [Engl. transl: *Sov. Phys.-Semicond.*, 11, 534 (1984)].
- [99] E. I. Levin, I. M. Ruzin, B. I. Shklovskii, Transverse hopping conductivity of amorphous films in strong electric fields. *Fiz. Tekh. Polup.* 22, 624 (1988) - Engl. Transl.: *Sov. Phys.-Semicond.* 22, 401 (1988).
- [100] Nguyen Van Lien, B. I. Shklovskii, Hopping conduction in strong electric fields and directed percolation, *Solid State Commun.* 38, 99 (1981).
- [101] E. I. Levin, B. I. Shklovskii, Negative differential conductivity of low density electron gas in random potential, *Solid State Comm.* 67, 233 (1988).
- [102] D. I. Aladashvili, Z. Adamia, K. Lavdovskii, E.I. Levin, B.I. Shklovskii, Negative differential resistance in hopping region

- in silicon, Pis'ma v Zh. Eksp. Teor. Fiz. 47. 390 (1988) - Engl. Transl.: Sov. Phys.-JETP Lett. 47. 466 (1988).
- [103] S. D. Baranovskii, E. I. Ivchenko, B. I. Shklovskii, New regime of recombination of photoinduced carriers in amorphous semiconductors, Zh. Eksp. Teor. Fiz. 22, 2234 (1987) - Engl. Transl.: Sov. Phys.-JETP 65, 1260 (1987).
- [104] S. D. Baranovskii, V. G. Karpov, B. I. Shklovskii, Theory of non-radiative recombination in amorphous semiconductors, J. Non-Cryst. Solids, 97, 487 (1987).
- [105] B. I. Shklovskii, H. Fritzsche and S. D. Baranovskii, Electronic Transport and Recombination in Amorphous semiconductors at Low Temperatures, Phys. Rev. Lett. 62, 2989 (1989).
- [106] S. D. Baranovskii, H. Fritzsche, E. I. Levin, I. M. Ruzin, B. I. Shklovskii, Low temperature electronic transport and recombination in amorphous semiconductors, Zh. Eksp. Teor. Fiz. 96, 1362 (1989) - Engl. Transl.: Sov. Phys. JETP 69, 773 (1990).
- [107] I. M. Ruzin, B. I. Shklovskii, Theory of hopping photoconductivity at low-frequency excitation, Fiz. Tekhn. Poluprov. 23. 1881 (1989) - Engl. Transl. Sov. Phys.- Semicond. 23, 1164 (1989).
- [108] B. I. Shklovskii, Model of surface recombination, Pis'ma Zh. Eksp. Teor. Fiz. 44, 95 (1986) - Engl. Transl: Sov. Phys.-JETP letters 14, 123 (1986).
- [109] B. I. Shklovskii, Dyakonov-Perel spin relaxation near metal-insulator transition and in hopping transport, Phys. Rev. B 73, 193201 (2006).
- [110] L. I. Glazman, I. M. Ruzin, B.I. Shklovskii, Quantum transport and pinning of one-dimensional Wigner crystal, Phys. Rev. B 45, 8454 (1992).
- [111] I. M. Ruzin, S. Marianer, B.I. Shklovskii, Pinning of two dimensional Wigner crystal by charged impurities, Phys. Rev. B 46, 3999 (1992).
- [112] A. A. Koulakov, B. I. Shklovskii, Charging Spectrum and Configurations of a Wigner Crystal Island, Phys. Rev. B 57, 2352 (1998).
- [113] B. I. Shklovskii, Coulomb gap and variable range hopping in a pinned Wigner crystal, Phys. Stat. Sol. (c) 1, 46 (2004).
- [114] M. M. Fogler, B. I. Shklovskii, Collapse of spin-splitting in the quantum Hall effect, Phys. Rev. B 52, 17366 (1995).
- [115] A. A. Koulakov, M. M. Fogler, and B. I. Shklovskii, Charge density wave in two-dimensional electron liquid in a weak magnetic field, Phys. Rev. Lett. 76, 499 (1996).
- [116] A. A. Koulakov, M. M. Fogler, and B. I. Shklovskii, The Ground State of a Two-Dimensional Electron Liquid in a Weak Magnetic Field, Phys. Rev. B 54, 1853 (1996).
- [117] Lilly, M. P., Cooper K. B., Eisenstein, J. P., Pfeiffer, L. N. and West, K. W., Evidence for an anisotropic state of two-dimensional electrons in high Landau levels. Phys. Rev. Lett. 82, 394 (1999).
- [118] Du, R. R., Tsui, D. C., Stormer, H. L., Pfeiffer, L. N., Baldwin, K.W. and West, K. W., Strongly anisotropic transport in higher two-dimensional Landau levels. Solid State Commun. 109, 389 (1999).
- [119] M. M. Fogler, Stripe and bubble phases in Quantum Hall systems, High Magnetic Fields: in "Applications in Condensed Matter Physics and spectroscopy" (Springer-Verlag, Berlin, 2002); ArXiv: cond-mat/0111001.
- [120] Fu, X., Shi, Q., Zudov, M. A., Gardner, G. C., Watson, J. D., Manfra, M. J., Two- and three-electron bubbles in  $\text{Al}_x\text{Ga}_{1-x}\text{As}/\text{Al}_{0.24}\text{Ga}_{0.76}\text{As}$  quantum wells. Phys. Rev. B 99, 161402 (2019).
- [121] Ro, D., Deng, N., Watson, J. D., Manfra, M. J., Pfeiffer, L. N., West, K. W. and Csáthy, G. A., Electron bubbles and the structure of the orbital wave function. Phys. Rev. B 99, 201111 (2019).
- [122] Chen, S., Ribeiro-Palau, R., Yang, K., Watanabe, K., Taniguchi, T., Hone, J., Goerbig, M. O. and Dean, C. R., Competing fractional quantum Hall and electron solid phases in graphene. Phys. Rev. Lett. 122, 026802 (2019).
- [123] Fangyuan Yang, Ruiheng Bai, Alexander A. Zibrov, Sandeep Joy, Takashi Taniguchi, Kenji Watanabe, Brian Skinner, Mark O. Goerbig, and Andrea F. Young, Cascade of multi-electron bubble phases in monolayer graphene at high Landau level filling, arXiv:2309.04319.
- [124] A. H. MacDonald and M. P. A. Fisher, Phys. Rev. B 61, 5724 (2000).
- [125] M. Sammon, X. Fu, Yi Huang, M. A. Zudov, B. I. Shklovskii, G. C. Gardner, J. D. Watson, M. J. Manfra, K. W. Baldwin, L. N. Pfeiffer, and K. W. West, Resistivity anisotropy of quantum Hall stripe phases, Phys. Rev. B 100, 241303(R) (2019).
- [126] Yi Huang, M. Sammon, M. A. Zudov, B. I. Shklovskii, Isotropically conducting (hidden) quantum Hall stripe phases in a two-dimensional electron gas, Phys. Rev. B 101, 161302 (2020).
- [127] X. Fu, Yi Huang, M. A. Zudov, B. I. Shklovskii, G. C. Gardner, M. J. Manfra, Hidden quantum Hall stripes in  $\text{Al}_x\text{Ga}_{1-x}\text{As}/\text{Al}_{0.24}\text{Ga}_{0.76}\text{As}$  quantum wells, Phys. Rev. Lett. 125, 236803 (2020).
- [128] M. S. Bello, E. I. Levin, B. I. Shklovskii, A. L. Efros, Density of localized states in the surface impurity band of a metal-insulator-semiconductor structure. Zh. Eksp. Teor. Fiz. 83, 1596 (1981) [English transl.: Sov. Phys.-JETP 53,822 (1981)].
- [129] S. V. Kravchenko, D. A. Rinberg, S. G. Semenchinsky, and V. M. Pudalov, Phys. Rev. B 42, 3741 (1990).
- [130] J. P. Eisenstein, L. N. Pfeiffer, and K. W. West, Phys. Rev. Lett. 68, 674 (1992).
- [131] B. Skinner, B. I. Shklovskii, Anomalously large capacitance of a plane capacitor with a two-dimensional electron gas, Phys. Rev. B 82, 155111 (2010).
- [132] Brian Skinner, B. I. Shklovskii, Giant capacitance of a plane capacitor with a two-dimensional electron gas in a magnetic field, Phys. Rev. B 87, 035409 (2013).
- [133] Brian Skinner, Tianran Chen, B. I. Shklovskii, Cooperative Charging in a Nanocrystal Assembly Gated By Ionic Liquid, Phys. Rev. B 84, 245304 (2011).
- [134] L. Li, C. Richter, S. Paetel, T. Kopp, J. Mannhart, and R. C. Ashoori, Science 332, 825 (2011).
- [135] Brian Skinner, G. L. Yu, A. V. Kretinin, A. K. Geim, K. S. Novoselov, B. I. Shklovskii, The effect of electron dielectric response on the quantum capacitance of graphene in a strong magnetic field. Phys. Rev. B 88, 155417 (2013).
- [136] Tingxin Li, Jiacheng Zhu, Yanhao Tang, Kenji Watanabe, Takashi Taniguchi, Veit Elser, Jie Shan, Kin Fai Mak, Charge-order-enhanced capacitance in semiconductor moiré superlattices, Nature Nanotechnology 16, 1068 (2021).
- [137] T. T. Nguyen, A. Yu. Grosberg, B. I. Shklovskii, Colloquium: Physics of charge inversion in chemical and biological systems, Rev. Mod. Phys. 74, 329 (2002).
- [138] T. T. Nguyen, B. I. Shklovskii, Inversion of DNA charge by a positive polymer via fractionalization of the polymer charge, Physica A 310, 197 (2002).
- [139] Xucheng Hou, Tal Zaks, Robert Langer, Yizhou Dong, Lipid nanoparticles for mRNA delivery, Nature Reviews Materials 6, 1078 (2021).
- [140] A. Kamenev, J. Zhang, B. I. Shklovskii, A. I. Larkin, Transport in one dimensional Coulomb gas: From ion channels to nanopores, Physica A, 359, 129 (2005).
- [141] A. Kamenev, J. Zhang, B. I. Shklovskii, Conductance of ion channels and nanopores with charged walls: a toy model, Phys.



- Rev. Lett. 95, 148101 (2005).
- [142] Leonard Susskind, Lattice models of quark confinement at high temperature, Phys. Rev. D 20, 2610 (1979).
- [143] B. I. Shklovskii, B. Shapiro, B. R. Sears, P. Lambrianidis, H. B. Shore, Statistics of spectra of disordered systems near the Metal-Insulator transition, Phys. Rev. B 47, 11487 (1993).
- [144] I. Kh. Zharekeshev and B. Kramer, Asymptotics of Universal Probability of Neighboring Level Spacings at the Anderson Transition, Phys. Rev. Lett. 79, 717 (1997).
- [145] Vadim Oganesyan and David A. Huse, Localization of interacting fermions at high temperature. Phys. Rev. B 75, 155111 (2007).
- [146] B. L. Altshuler, I. Zharekeshev, S. A. Kotochigova, B. I. Shklovskii, Repulsion between energy levels and the metal-insulator transition, Zh. Eksp. Teor. Fiz. 94, 343 (1988) - Engl. Transl.: Sov. Phys.-JETP 67, 625 (1988).
- [147] A. F. Tzortzakakis, K. G. Makris, and E. N. Economou, Phys. Rev. B 101, 014202 (2020).
- [148] Yi Huang, B. I. Shklovskii, Anderson transition in three-dimensional non-Hermitian disorder, Phys. Rev. B 101, 014204 (2020).
- [149] Yi Huang, B. I. Shklovskii, Spectral rigidity of non-Hermitian symmetric random matrices, Phys. Rev. B 102, 064212 (2020).
- [150] J. J. Nelson, K. V. Reich, M. Sammon, B. I. Shklovskii and A. M. Goldman, Hopping Conduction via Ionic Liquid Induced Silicon Surface States, Phys. Rev. B 92, 085524 (2015).
- [151] Peter P. Orth, Rafael M. Fernandes, Jeff Walter, C. Leighton, and B. I. Shklovskii, Percolation via Combined Electrostatic and Chemical Doping in Complex Oxide Films
- [152] D. N. Arnold, G. David, D. Jerison, S. Mayboroda, and M. Filoche, Effective confining potential of quantum states in disordered media, Phys. Rev. Lett. 116, 056602 (2016).
- [153] F. Gebhard, A. V. Nenashev, K. Meyerholz and S. D. Baranovskii, Quantum states in disordered media. I. Low-pass filter approach, Phys. Rev. B 107, 064206 (2023).
- [154] B. I. Shklovskii, A. L. Efros, Density of state oscillations of 2d electron gas in magnetic field, Pis'ma Zh. Eksp. Theor. Fiz. 44, 95 (1986) - Engl. Transl: Sov. Phys.-JETP Letters 44, 669 (1986).
- [155] A. L. Efros, Solid State Commun. 67, 1019 (1988).
- [156] A. L. Efros, Solid State Commun. 65, 1281 (1988).
- [157] A. L. Efros, Phys. Rev. B 45, 11354 (1992).
- [158] D. B. Chklovskii, B. I. Shklovskii and L. I. Glazman, Electrostatics of edge states, Phys. Rev. B 46, 4026 (1992).
- [159] D. B. Chklovskii, K. A. Matveev, B. I. Shklovskii, Ballistic conductance of interacting electrons, Phys. Rev. B 47, 12605 (1993).
- [160] M. M. Fogler, B. I. Shklovskii, Resistance of a long wire in the quantum Hall regime, Phys. Rev. B 50, 1656 (1994).
- [161] D. G. Polyakov, B. I. Shklovskii, Universal prefactor of activated conductivity in the Quantum Hall Effect, Phys. Rev. Lett. 74, 150 (1995).
- [162] M. Sammon, M. A. Zudov, and B. I. Shklovskii, Mobility and quantum mobility of modern GaAs/AlGaAs heterostructures. Phys. Rev. M 2, 064604 (2018).
- [163] M. Sammon, Tianran Chen, B. I. Shklovskii, Excess electron screening of remote donors and mobility in modern GaAs/AlGaAs heterostructures. Phys. Rev. M 2, 104001 (2018).
- [164] Brian Skinner, Tianran Chen, B. I. Shklovskii, Why is the bulk resistivity of topological insulators so small? Phys. Rev. Lett. 109, 176801 (2012).
- [165] T. Knispel, W. Jolie, N. Borgwardt, J. Lux, Zhiwei Wang, Yoichi Ando, A. Rosch, T. Michely, and M. Grüninger, Charge puddles in the bulk and on the surface of the topological insulator BiSbTeSe<sub>2</sub> studied by scanning tunneling microscopy and optical spectroscopy, Phys. Rev. B 96, 195135 (2017).
- [166] Brian Skinner, B. I. Shklovskii, Theory of the random potential and conductivity at the surface of a topological insulator, Phys. Rev. B 87, 0755454 (2013).
- [167] Yi Huang, B. I. Shklovskii, Disorder effects in topological insulator thin films, Phys. Rev. B 103, 165409 (2021).
- [168] Yi Huang, B. I. Shklovskii, Disorder effects in topological insulator nanowires, Phys. Rev. B 104, 054205 (2021).
- [169] Yi Huang, Yanjun He, Brian Skinner, B. I. Shklovskii, Conductivity of two-dimensional narrow gap semiconductors subjected to strong Coulomb disorder, Phys. Rev. B 105, 054206 (2022).
- [170] Yi Huang, Brian Skinner, B. I. Shklovskii, Conductivity of two-dimensional small gap semiconductors and topological insulators in strong Coulomb disorder, JETP 135, 409 (2022).
- [171] M. Mueller, B. I. Shklovskii, Compensation driven superconductor-insulator transition, Phys. Rev. B 79, 134504 (2009).