

Every locally compact group is the outer automorphism group of a II_1 factor

BY STEFAAN VAES¹

Abstract

We prove that every locally compact second countable group G arises as the outer automorphism group $\text{Out } M$ of a II_1 factor, which was so far only known for totally disconnected groups, compact groups and a few isolated examples. We obtain this result by proving that every locally compact second countable group is a *centralizer group*, a class of Polish groups that arise naturally in ergodic theory and that may all be realized as $\text{Out } M$.

1 Introduction and main results

The *outer automorphism group* $\text{Out } M$ of a II_1 factor M is the quotient of the group $\text{Aut } M$ of all $*$ -automorphisms of M by the normal subgroup $\text{Inn } M$ of inner automorphisms, of the form $\text{Ad } u$ with $u \in \mathcal{U}(M)$. In [IPP05] answering a question of Connes from the 1970s, it was proven that there exist II_1 factors M with trivial outer automorphism group. This provided at the same time the first complete computation of the outer automorphism group of any II_1 factor.

A II_1 factor M is called *full* if $\text{Inn } M$ is a closed subgroup of $\text{Aut } M$. This is equivalent with M not having property Gamma in the sense of Murray and von Neumann. If M is full and has separable predual, $\text{Out } M$ is a Polish group. It is a natural and by now well-studied question which Polish groups arise as the outer automorphism group of a full II_1 factor with separable predual. The following families of Polish groups could be realized in this way: compact abelian groups in [IPP05], countable discrete groups in [PV06, Vae07], arbitrary compact groups in [FV07] and several families of non locally compact groups in [PV21], including all closed subgroups of the group of all permutations of a countably infinite set.

All this however left open the question if all locally compact second countable groups, in particular Lie groups, can be realized as $\text{Out } M$. We solve this problem here and the following is our main result.

Theorem A. *Every locally compact second countable group G arises as the outer automorphism group $\text{Out } M$ of a full II_1 factor M with separable predual.*

Denote by (X, μ) the standard nonatomic probability space. We consider the Polish group $\text{Aut}(X, \mu)$ of all *nonsingular* automorphisms of (X, μ) , identifying two automorphisms that coincide almost everywhere. The *measure preserving* automorphisms form a closed subgroup $\text{Aut}_{\text{mp}}(X, \mu)$. In [Dep10, Theorem E], it was proven that for every countable subgroup $\Lambda < \text{Aut}_{\text{mp}}(X, \mu)$, the *centralizer* $C(\Lambda)$ of Λ inside $\text{Aut}_{\text{mp}}(X, \mu)$ is a Polish group that arises as the outer automorphism group $\text{Out } M$ of a full II_1 factor M with separable predual. A more direct proof of this result was given in [PV21, Proposition 8.5].

To prove Theorem A, it thus suffices to realize every locally compact second countable group as a centralizer inside $\text{Aut}_{\text{mp}}(X, \mu)$. Note here that a subgroup $G < \text{Aut}_{\text{mp}}(X, \mu)$ is of the

¹KU Leuven, Department of Mathematics, Leuven (Belgium), stefaan.vaes@kuleuven.be
Supported by FWO research project G090420N of the Research Foundation Flanders and by Methusalem grant METH/21/03 — long term structural funding of the Flemish Government.

form $C(\Lambda)$ for a subgroup $\Lambda < \text{Aut}_{\text{mp}}(X, \mu)$ if and only if G equals its double centralizer $C(C(G))$.

Definition B. We say that a Polish group G is a *centralizer group* if there exists an ergodic probability measure preserving action $\Lambda \curvearrowright (X, \mu)$ of a countable group Λ on a standard probability space (X, μ) such that G is isomorphic with the centralizer of Λ inside $\text{Aut}_{\text{mp}}(X, \mu)$.

So as explained above, by [Dep10, PV21] for every centralizer group G , there exists a full II_1 factor M with separable predual such that $G \cong \text{Out } M$. Since every locally compact group G is the centralizer of G acting on itself (by transformations that are not probability measure preserving), Theorem A is thus a consequence of the following result, which also shows that the class of centralizer groups is quite natural.

Theorem C. *The following Polish groups are centralizer groups.*

- (i) *The centralizer of any subgroup of the Polish group $\text{Aut}_{\text{mp}}(X, \mu)$ of measure preserving automorphisms of a possibly infinite σ -finite standard measure space (X, μ) .*
- (ii) *The centralizer of any subgroup of the Polish group $\text{Aut}(X, \mu)$ of nonsingular automorphisms of a standard probability space (X, μ) .*

To prove the first statement, we make use of *Poisson suspensions*, while we deduce the second statement from the first by using the *Maharam extension*. In Sections 4 and 5, we prove a few permanence properties for the class of centralizer groups, we reprove that every closed subgroup of the group S_∞ of all permutations of a countably infinite set is a centralizer group (cf. [PV21, Corollary 8.2]) and that $\text{Aut } K$ is a centralizer group for all compact second countable groups K .

Remark D. To summarize, we then get the following list of centralizer groups, which can all be realized as the outer automorphism group of a full II_1 factors with separable predual.

- (i) Locally compact second countable groups and their direct integrals (Theorem A and Proposition 4.2).
- (ii) Closed subgroups of S_∞ , the group of all permutations of a countably infinite set ([PV21, Corollary 8.2] and Theorem 5.1).
- (iii) The automorphism group $\text{Aut } K$ of a compact second countable group K (Proposition 5.3).
- (iv) The unitary group $\mathcal{U}(N)$ of a von Neumann algebra N with separable predual ([PV21, Corollary 8.2]).
- (v) A direct product of a countable family of centralizer groups (Proposition 4.1).

2 Preliminaries

We start by recalling the concept of *Poisson suspensions* and refer to [JRD20] for details and further references. Let (Y, η) be a standard σ -finite measure space. A measure on Y of the form $\sum_{i \in I} \delta_{y_i}$, where $(y_i)_{i \in I}$ is a countable set of points in Y , is called a *counting measure*. Write $\overline{\mathbb{N}} = \mathbb{N} \cup \{0, +\infty\}$. The set X of all counting measures ν on Y has a unique standard Borel structure such that the maps

$$N_A : X \rightarrow \overline{\mathbb{N}} : \nu \mapsto \nu(A)$$

are Borel for every Borel set $A \subset Y$. There is a unique probability measure μ on X such that for every Borel set $A \subset Y$ with $0 < \eta(A) < +\infty$, the random variable N_A has a Poisson distribution with intensity $\eta(A)$, i.e.

$$\mu(N_A^{-1}(\{k\})) = \exp(-\eta(A)) \frac{\eta(A)^k}{k!} \quad \text{for all } k \in \mathbb{N} \cup \{0\},$$

and such that the random variables N_{A_1}, \dots, N_{A_n} are mutually independent for all disjoint Borel sets $A_i \subset Y$ with $0 < \eta(A_i) < +\infty$. Also note that $N_A + N_B = N_{A \cup B}$ when $A, B \subset Y$ are disjoint Borel sets.

There then is a unique isometric real linear map

$$\theta : L_{\mathbb{R}}^2(Y, \eta) \rightarrow L_{\mathbb{R}}^2(X, \mu) : \theta(1_A) = N_A - \eta(A)1 \quad (2.1)$$

for all Borel sets $A \subset Y$ with $\eta(A) < +\infty$. One checks that

$$E_{\mu}(\exp(it\theta(\xi))) = \exp\left(\int_Y (\exp(it\xi) - it\xi - 1) d\eta\right) \quad \text{for all } \xi \in L_{\mathbb{R}}^2(Y, \eta), \quad (2.2)$$

where the right hand side is well-defined because $|\exp(ir) - ir - 1| \leq r^2$ for all $r \in \mathbb{R}$.

Denote by $L_{\mathbb{R}}^2(Y, \eta)^{\otimes_{\text{symm}} n}$ the closed subspace of symmetric vectors in $L_{\mathbb{R}}^2(Y, \eta)^{\otimes n}$. Consider the symmetric Fock space

$$\mathcal{F}_s(L_{\mathbb{R}}^2(Y, \eta)) = \mathbb{R}1 \oplus \bigoplus_{n=1}^{\infty} L_{\mathbb{R}}^2(Y, \eta)^{\otimes_{\text{symm}} n}$$

and define for every $\xi \in L_{\mathbb{R}}^2(Y, \eta)$ the vector $\text{Exp}(\xi) \in \mathcal{F}_s(L_{\mathbb{R}}^2(Y, \eta))$ by

$$\text{Exp}(\xi) = 1 \oplus \bigoplus_{n=1}^{\infty} \frac{1}{\sqrt{n!}} \xi^{\otimes n}.$$

Note that $\langle \text{Exp}(\xi), \text{Exp}(\zeta) \rangle = \exp(\langle \xi, \zeta \rangle)$.

Then $\theta : L_{\mathbb{R}}^2(Y, \eta) \rightarrow L_{\mathbb{R}}^2(X, \mu)$ uniquely extends to an isometric, bijective, real linear map

$$\Theta : \mathcal{F}_s(L_{\mathbb{R}}^2(Y, \eta)) \rightarrow L_{\mathbb{R}}^2(X, \mu) : \Theta(\text{Exp}(\exp(\xi) - 1)) = E_{\mu}(\exp(\theta(\xi)))^{-1} \exp(\theta(\xi)) \quad (2.3)$$

for all simple functions $\xi \in L_{\mathbb{R}}^2(Y, \eta)$, i.e. finite linear combinations of 1_A with $A \subset Y$ Borel and $\eta(A) < +\infty$.

To every $\alpha \in \text{Aut}_{\text{mp}}(Y, \eta)$ corresponds the Poisson suspension $\hat{\alpha} \in \text{Aut}_{\text{mp}}(X, \mu)$ defined by $\hat{\alpha}(\nu) = \alpha_*(\nu)$ for every counting measure ν on Y . Every measure preserving automorphism β of a σ -finite standard measure space (Z, ζ) , also gives rise to the canonical orthogonal transformation U_{β} of $L_{\mathbb{R}}^2(Z, \zeta)$. We then get that

$$U_{\hat{\alpha}} \circ \Theta = \Theta \circ \left(\text{id} \oplus \bigoplus_{n=1}^{\infty} U_{\alpha}^{\otimes_{\text{symm}} n} \right) \quad \text{for all } \alpha \in \text{Aut}_{\text{mp}}(Y, \eta). \quad (2.4)$$

3 Centralizer groups, proof of Theorems A and C

The following is the key technical lemma showing that in certain cases, the centralizer of a Poisson suspension equals the Poisson suspension of the centralizer.

Lemma 3.1. *Let (Y, η) be a standard σ -finite measure space and $\Lambda < \text{Aut}_{\text{mp}}(Y, \eta)$ a subgroup. Denote by (X, μ) the Poisson suspension of (Y, η) and consider $\widehat{\Lambda} < \text{Aut}_{\text{mp}}(X, \mu)$ given by $\widehat{\Lambda} = \{\widehat{\beta} \mid \beta \in \Lambda\}$. Assume that there exists a sequence $\beta_n \in \Lambda$ such that the orthogonal transformations U_{β_n} of $L_{\mathbb{R}}^2(Y, \eta)$ satisfy $U_{\beta_n} \rightarrow \rho \cdot \text{id}$ weakly for some $0 < \rho < 1$.*

Then $\widehat{\Lambda} < \text{Aut}_{\text{mp}}(X, \mu)$ is ergodic and the map $\alpha \mapsto \widehat{\alpha}$ is a bijection between the centralizer of Λ inside $\text{Aut}_{\text{mp}}(Y, \eta)$ and the centralizer of $\widehat{\Lambda}$ inside $\text{Aut}_{\text{mp}}(X, \mu)$.

Proof. Consider the symmetric Fock space $\mathcal{F}_s(L_{\mathbb{R}}^2(Y, \eta))$ and the isometry $\Theta : \mathcal{F}_s(L_{\mathbb{R}}^2(Y, \eta)) \rightarrow L_{\mathbb{R}}^2(X, \mu)$ given by (2.3). By (2.4) and the assumption that $U_{\beta_n} \rightarrow \rho \cdot \text{id}$ weakly,

$$\Theta^* \circ U_{\widehat{\beta}_n} \circ \Theta \rightarrow T := \text{id} \oplus \bigoplus_{n=1}^{\infty} \rho^n \cdot \text{id} \quad \text{weakly.}$$

So if $\xi \in L_{\mathbb{R}}^2(X, \mu)$ is $\widehat{\Lambda}$ -invariant, it follows that $\Theta^*(\xi) \in \mathbb{R}1$ and thus $\xi \in \mathbb{R}1$, proving the ergodicity of $\widehat{\Lambda} < \text{Aut}_{\text{mp}}(X, \mu)$.

Take $\varphi \in \text{Aut}_{\text{mp}}(X, \mu)$ centralizing $\widehat{\Lambda}$. It follows that $\Theta^* \circ U_{\varphi} \circ \Theta$ commutes with T , and thus globally preserves the eigenspace $L_{\mathbb{R}}^2(Y, \eta)$, so that U_{φ} globally preserves $\theta(L_{\mathbb{R}}^2(Y, \eta))$, with θ defined by (2.1). It follows from [Roy08, Proposition 4.4] that $\varphi = \widehat{\alpha}$ for a uniquely defined $\alpha \in \text{Aut}_{\text{mp}}(Y, \eta)$. For completeness, we provide the following easy argument to obtain this result.

Define the orthogonal transformation U of $L_{\mathbb{R}}^2(Y, \eta)$ such that $U_{\varphi} \circ \theta = \theta \circ U$. Take an arbitrary $\xi \in L_{\mathbb{R}}^2(Y, \eta)$. The characteristic function of the random variable $\theta(\xi)$ is given by (2.2). This is saying that $\theta(\xi)$ is infinitely divisible with Lévy measure ρ_{ξ} given by $\rho_{\xi}(\mathcal{U}) = \eta(\xi^{-1}(\mathcal{U}))$ for all Borel sets $\mathcal{U} \subset \mathbb{R} \setminus \{0\}$. Since φ is measure preserving, $\theta(\xi)$ and $U_{\varphi}(\theta(\xi))$ have the same distribution and hence, the same Lévy measure. Since $U_{\varphi}(\theta(\xi)) = \theta(U(\xi))$, it follows that $\rho_{\xi} = \rho_{U(\xi)}$ for all $\xi \in L_{\mathbb{R}}^2(Y, \eta)$.

Since $\xi \geq 0$ a.e. if and only if $\rho_{\xi}(-\infty, 0) = 0$ and since $\xi \leq 1$ a.e. if and only if $\rho_{\xi}(1, +\infty) = 0$, we conclude that the orthogonal transformation $U \in \mathcal{O}(L_{\mathbb{R}}^2(Y, \eta))$ maps positive functions to positive functions and maps functions $0 \leq \xi \leq 1$ to functions $0 \leq U(\xi) \leq 1$. This is only possible if $U = V_{\alpha}$ for some $\alpha \in \text{Aut}_{\text{mp}}(Y, \eta)$. So, $\varphi = \widehat{\alpha}$.

Since φ commutes with $\widehat{\Lambda}$, we get that α commutes with Λ and the lemma is proven. \square

Lemma 3.2. *Let Y be a standard Borel space and η an infinite, σ -finite, nonatomic Borel measure on Y . There exists a sequence $\beta_n \in \text{Aut}_{\text{mp}}(Y, \eta)$ such that the orthogonal transformations U_{β_n} of $L_{\mathbb{R}}^2(Y, \eta)$ satisfy $U_{\beta_n} \rightarrow (1/2) \cdot \text{id}$ weakly.*

Proof. Since all such (Y, η) are isomorphic, we may realize $(Y, \eta) = (X \times \mathbb{Z}, \mu \times \lambda)$ with $(X, \mu) = (\{0, 1\}^{\mathbb{N}}, \mu_0^{\mathbb{N}})$, $\mu_0(0) = \mu_0(1) = 1/2$, and λ the counting measure on \mathbb{Z} . We define $\beta_n \in \text{Aut}_{\text{mp}}(Y, \eta)$ by

$$\beta_n(x, k) = \begin{cases} (x, k + n) & \text{if } x_n = 0, \\ (x, k) & \text{if } x_n = 1. \end{cases}$$

Take $N \in \mathbb{N}$, $A_0, A_1 \in \{0, 1\}^N$ and $B_0, B_1 \subset [-N, N] \cap \mathbb{Z}$. Write $C_i = A_i \times \{0, 1\}^{>N} \subset X$. When $n > 2N$, we have that

$$\langle U_{\beta_n}(1_{C_0} \otimes 1_{B_0}), 1_{C_1} \otimes 1_{B_1} \rangle = \frac{1}{2} \langle 1_{C_0} \otimes 1_{B_0}, 1_{C_1} \otimes 1_{B_1} \rangle.$$

Since these vectors $1_C \otimes 1_B$ are total in $L_{\mathbb{R}}^2(Y, \eta)$, it follows that $U_{\beta_n} \rightarrow (1/2) \cdot \text{id}$ weakly. \square

Proof of Theorem C. To prove (i), take a standard σ -finite measure space (Y_0, η_0) and a subgroup $\Lambda_0 < \text{Aut}_{\text{mp}}(Y_0, \eta_0)$. Denote by G the centralizer of Λ_0 inside $\text{Aut}_{\text{mp}}(Y_0, \eta_0)$. We have to prove that G is a centralizer group in the sense of Definition B.

Take an arbitrary nonatomic, infinite, σ -finite standard measure space (Y_1, η_1) . By Lemma 3.2, take a sequence $\beta_n \in \text{Aut}_{\text{mp}}(Y_1, \eta_1)$ such that $U_{\beta_n} \rightarrow (1/2) \cdot \text{id}$ weakly. Denote $(Y, \eta) = (Y_0 \times Y_1, \eta_0 \times \eta_1)$. Define $\Lambda_1 < \text{Aut}_{\text{mp}}(Y, \eta)$ by $\Lambda_1 = \Lambda_0 \times \text{Aut}_{\text{mp}}(Y_1, \eta_1)$.

The centralizer of Λ_1 inside $\text{Aut}_{\text{mp}}(Y, \eta)$ equals $G \times \text{id}$. The sequence $\text{id} \times \beta_n \in \Lambda$ satisfies $U_{\text{id} \times \beta_n} \rightarrow (1/2) \cdot \text{id}$ weakly. Denote by (X, μ) the Poisson suspension of (Y, η) and denote by $\widehat{\Lambda}_1 < \text{Aut}(X, \mu)$ the Poisson suspension Λ_1 . By Lemma 3.1, the subgroup $\widehat{\Lambda}_1 < \text{Aut}_{\text{mp}}(X, \mu)$ is ergodic and its centralizer inside $\text{Aut}_{\text{mp}}(X, \mu)$ is isomorphic with G .

Since $\text{Aut}_{\text{mp}}(X, \mu)$ is a Polish group, we can choose a countable subgroup Λ of $\text{Aut}_{\text{mp}}(X, \mu)$ whose closure equals the closure of $\widehat{\Lambda}_1$. Since the centralizer of Λ equals the centralizer of $\widehat{\Lambda}_1$, it follows that G is a centralizer group in the sense of Definition B.

To prove (ii), take a standard probability space (Y_0, η_0) and a subgroup $\Lambda_0 < \text{Aut}(Y_0, \eta_0)$ of nonsingular transformations. Denote by G the centralizer of Λ_0 inside $\text{Aut}(Y_0, \eta_0)$. We have to prove that G is a centralizer group.

Denote by γ the measure on \mathbb{R} given by $d\gamma(t) = \exp(-t) dt$. Write $(Y_1, \eta_1) = (Y_0 \times \mathbb{R}, \eta_0 \times \gamma)$. For every $\beta \in \text{Aut}(Y_0, \eta_0)$, denote its Maharam extension by

$$\tilde{\beta} \in \text{Aut}_{\text{mp}}(Y_1, \eta_1) : \tilde{\beta}(y, t) = (\beta(y), t + \log \frac{d(\beta_*^{-1}(\eta))}{d\eta}(y)) .$$

For every $s \in \mathbb{R}$, denote by λ_s the measure scaling automorphism $\lambda_s(y, t) = (y, t + s)$. Then

$$\Psi : \text{Aut}(Y_0, \eta_0) \rightarrow \text{Aut}_{\text{mp}}(Y_1, \eta_1) : \Psi(\beta) = \tilde{\beta}$$

is a group homomorphism whose image is equal to

$$\{\zeta \in \text{Aut}_{\text{mp}}(Y_1, \eta_1) \mid \forall s \in \mathbb{R} : \zeta \circ \lambda_s = \lambda_s \circ \zeta \text{ a.e.}\} .$$

The image of Ψ is closed and Ψ is a homeomorphism between $\text{Aut}(Y_0, \eta_0)$ and the image of Ψ .

Then define $(Y, \eta) = (Y_1 \times \mathbb{R}, \eta_1 \times \gamma) = (Y_0 \times \mathbb{R} \times \mathbb{R}, \eta_0 \times \gamma \times \gamma)$. For every $s \in \mathbb{R}$, define $\rho_s \in \text{Aut}_{\text{mp}}(Y, \eta)$ by $\rho_s(y, t_1, t_2) = (y, t_1 + s, t_2 - s)$. Denote by $\Lambda < \text{Aut}_{\text{mp}}(Y, \eta)$ the subgroup generated by $\tilde{\beta} \times \text{id}$, $\beta \in \Lambda_0$, ρ_s , $s \in \mathbb{R}$, and $\text{id} \times \text{id} \times \text{Aut}_{\text{mp}}(\mathbb{R}, \gamma)$. Denote by G_1 the centralizer of Λ inside $\text{Aut}_{\text{mp}}(Y, \eta)$. Since we have already proven (i), it suffices to prove that $G \cong G_1$.

Take $\varphi \in G_1$. Since φ commutes with $\text{id} \times \text{id} \times \text{Aut}_{\text{mp}}(\mathbb{R}, \gamma)$, we get that $\varphi = \psi \times \text{id}$ for some $\psi \in \text{Aut}_{\text{mp}}(Y_1, \eta_1)$. Since φ commutes with all ρ_s , we get that ψ commutes with all λ_s . Since ψ is measure preserving, we get that $\psi = \tilde{\alpha}$ for a unique $\alpha \in G$. So the map $\alpha \mapsto \Psi(\alpha) \times \text{id}$ implements the isomorphism $G \cong G_1$. \square

Proof of Theorem A. Choose a probability measure μ that is equivalent with the Haar measure on G . View $G < \text{Aut}(G, \mu)$ by right translations. Then the centralizer of G equals the copy of G in $\text{Aut}(G, \mu)$ by left translations. By Theorem C.(ii), the group G is a centralizer group. By [Dep10, Theorem E] (see also [PV21, Proposition 8.5]), there exists a full II_1 factor M with separable predual such that $\text{Out } M \cong G$. \square

4 Permanence properties

When $(G_i)_{i \in I}$ is a countable family of Polish groups, the product $\prod_{i \in I} G_i$ with the topology of pointwise convergence is again a Polish group. We start with the following elementary

observation.

Proposition 4.1. *If $(G_i)_{i \in I}$ is a countable family of centralizer groups, the product $\prod_i G_i$ also is a centralizer group.*

Proof. Choose standard probability spaces (X_i, μ_i) and ergodic subgroups $\Lambda_i < \text{Aut}_{\text{mp}}(X_i, \mu_i)$ whose centralizers are isomorphic with G_i for all $i \in I$. Define the standard probability $(X, \mu) = \prod_{i \in I} (X_i, \mu_i)$. For every $i \in I$, denote by $\Gamma_i < \text{Aut}_{\text{mp}}(X, \mu)$ the group Λ_i acting in the i 'th coordinate. Define $\Gamma < \text{Aut}_{\text{mp}}(X, \mu)$ as the subgroup generated by all the Γ_i , $i \in I$. Since each $\Lambda_i < \text{Aut}_{\text{mp}}(X_i, \mu_i)$ is ergodic, any element $\alpha \in \text{Aut}_{\text{mp}}(X, \mu)$ that centralizes Γ must be of the form $\prod_i \alpha_i$ with $\alpha_i \in \text{Aut}_{\text{mp}}(X_i, \mu_i)$. From this the result follows immediately. \square

With some more care, we also have the following continuous variant of Proposition 4.1.

Recall from [VW24, Definition 7.3] (see also [Sut85]) the concept of a *measured field* $(G_x)_{x \in X}$ of Polish groups. In a nutshell, this means that (X, μ) is a standard probability space and that, after discarding from X a Borel set of measure zero, we are given a standard Borel structure on the disjoint union $G = \sqcup_{x \in X} G_x$ such that the following holds: the natural factor map $\pi : G \rightarrow X$ is Borel, there exists a dense sequence of Borel sections $\varphi_n : X \rightarrow G$ and, writing $G \times_{\pi} G = \{(g, h) \mid \pi(g) = \pi(h)\}$, the multiplication map $G \times_{\pi} G \rightarrow G$ and the inverse map $G \rightarrow G$ are Borel.

Given such a measured field $(G_x)_{x \in X}$ over (X, μ) , we consider the group \mathcal{G} of all Borel sections $\psi : X \rightarrow G$, where we identify sections that are equal a.e. With pointwise multiplication and with the topology of convergence in measure, where $\psi_k \rightarrow \psi$ if and only if $\mu(\{x \in X \mid d(\psi_k(x), \psi(x)) > \varepsilon\}) \rightarrow 0$ for all $\varepsilon > 0$, we get that \mathcal{G} is a Polish group.

Note that when X is a countable set and μ is an atomic probability measure that assigns a positive value to each atom, then $\mathcal{G} = \prod_{i \in X} G_i$ as above. In that sense, \mathcal{G} is a ‘‘direct integral’’ version of the usual product of Polish groups.

Proposition 4.2. *Let (X, μ) be a standard probability space and let $(G_x)_{x \in X}$ be a measured field of second countable locally compact groups. As above, denote by \mathcal{G} the Polish group of measurable sections of this field, identifying sections that are equal a.e. and using the topology of convergence in measure. Then \mathcal{G} is a centralizer group.*

Proof. We discard from X a Borel set of measure zero, so that $G = \sqcup_{x \in X} G_x$ is equipped with a standard Borel structure satisfying the conditions explained above (see [VW24, Definition 7.3]). Using [VW24, Proposition 10.2], we may choose probability measures γ_x on G_x such that for every $x \in X$, γ_x is equivalent with the Haar measure of G_x and such that for every positive Borel function $F : G \rightarrow [0, +\infty)$, the map

$$X \rightarrow [0, +\infty] : x \mapsto \int_{G_x} F d\gamma_x$$

is Borel. We then denote by γ the probability measure on the Borel σ -algebra of G given by

$$\gamma(\mathcal{U}) = \int_X \gamma_x(\mathcal{U} \cap G_x) d\mu(x) = \int_X \left(\int_{G_x} 1_{\mathcal{U}} d\gamma_x \right) d\mu(x) \quad \text{for every Borel set } \mathcal{U} \subset G.$$

By construction, $\pi_*(\gamma) = \mu$. For every Borel section $\varphi : X \rightarrow G$, define the Borel bijections

$$\alpha_{\varphi} : G \rightarrow G : \alpha_{\varphi}(g) = \varphi(\pi(g))g \quad \text{and} \quad \beta_{\varphi} : G \rightarrow G : \beta_{\varphi}(g) = g\varphi(\pi(g))^{-1}.$$

Since every γ_x is equivalent with the Haar measure of G_x , it follows that all α_φ and β_φ are nonsingular automorphisms of (G, γ) . By construction,

$$\alpha : \mathcal{G} \rightarrow \text{Aut}(G, \gamma) : \varphi \mapsto \alpha_\varphi \quad \text{and} \quad \beta : \mathcal{G} \rightarrow \text{Aut}(G, \gamma) : \varphi \mapsto \beta_\varphi$$

are faithful continuous group homomorphisms.

Note that whenever G_0 is a locally compact second countable group and γ_0 is a probability measure on G_0 that is equivalent with the Haar measure of G , then the embedding $G_0 \hookrightarrow \text{Aut}(G_0, \gamma_0)$ given by left translation has a closed image and is a homeomorphism of G_0 onto this image. It follows that the same holds for $\alpha : \mathcal{G} \rightarrow \text{Aut}(G, \gamma)$.

We claim that $\Delta \in \text{Aut}(G, \gamma)$ belongs to $\alpha(\mathcal{G})$ if and only if Δ commutes with $\beta(\mathcal{G})$ and satisfies $\pi(\Delta(g)) = \pi(g)$ for a.e. $g \in G$. One implication being obvious, assume that $\Delta \in \text{Aut}(G, \gamma)$ commutes with $\beta(\mathcal{G})$ and satisfies $\pi(\Delta(g)) = \pi(g)$ for a.e. $g \in G$. Choose a dense sequence of Borel sections $\varphi_n : X \rightarrow G$. Take a conull Borel set $X_0 \subset X$ such that for every $x \in X_0$, the following holds:

$$\begin{aligned} \pi(\Delta(g)) &= x \quad \text{for } \gamma_x\text{-a.e. } g \in G_x, \text{ and} \\ (\Delta \circ \beta_{\varphi_n})(g) &= (\beta_{\varphi_n} \circ \Delta)(g) \quad \text{for all } n \in \mathbb{N} \text{ and } \gamma_x\text{-a.e. } g \in G_x. \end{aligned}$$

So for every $x \in X_0$, there is a unique $\varphi(x) \in G_x$ such that $\Delta(g) = \varphi(x)g$ for γ_x -a.e. $g \in G_x$. It follows that $\varphi \in \mathcal{G}$ and $\Delta = \alpha(\varphi)$. So the claim is proven.

Take any nonatomic standard probability space (Y_1, η_1) . Choose an injective Borel map $\rho : X \rightarrow \text{Aut}_{\text{mp}}(Y_1, \eta_1)$. Write $(Y, \eta) = (G \times Y_1, \gamma \times \eta_1)$ and define $\delta \in \text{Aut}_{\text{mp}}(Y, \eta)$ by $\delta(g, y) = (g, \rho(\pi(g))(y))$.

Define $\Lambda < \text{Aut}(Y, \eta)$ as the subgroup generated by $\Lambda_0 \times \text{Aut}_{\text{mp}}(Y_1, \eta_1)$ and δ . We prove that the centralizer of Λ inside $\text{Aut}(Y, \eta)$ is given by $\alpha(\mathcal{G}) \times \text{id}$. The proposition then follows from Theorem C.(ii).

Choose $\Delta \in \text{Aut}(Y, \eta)$ that commutes with Λ . Since Δ commutes with $\text{id} \times \text{Aut}_{\text{mp}}(Y_1, \eta_1)$, we get that $\Delta = \Delta_0 \times \text{id}$ for some $\Delta_0 \in \text{Aut}(G, \gamma)$. Since Δ commutes with δ and ρ is injective, it follows that $\pi \circ \Delta_0 = \pi$. Since Δ_0 also commutes with Λ_0 , the result follows from the claim above. \square

Remark 4.3. It is plausible that for any measured field $(G_x)_{x \in X}$ of Polish groups with the property that a.e. G_x is a centralizer group, also the Polish group \mathcal{G} of measurable sections is a centralizer group. To prove such a result, one would need to make a measurable choice of $\Lambda_x < \text{Aut}(Y_x, \eta_x)$ such that G_x is the centralizer of Λ_x inside $\text{Aut}(Y_x, \eta_x)$. While it sounds plausible that such a measurable choice always exists, we were unable to prove this.

5 Closed groups of permutations of a countable set

We denote by S_∞ the Polish group of all permutations of \mathbb{N} , equipped with the topology of pointwise convergence. The following result was already proven in [PV21, Corollary 8.2], but with the spectral methods of Section 3, we can give a considerably easier proof.

Theorem 5.1. *Every closed subgroup of S_∞ is a centralizer group.*

The main point in proving Theorem 5.1 is the following lemma, which was obtained as [PV21, Lemma 8.6] and for which we give an easy spectral proof.

Lemma 5.2. *Let (X_0, μ_0) be a standard nonatomic probability space. Put $(X, \mu) = (X_0, \mu_0)^\mathbb{N}$. We view $S_\infty < \text{Aut}(X, \mu)$ by permuting the indices and we view $\text{Aut}_{\text{mp}}(X_0, \mu_0) < \text{Aut}(X, \mu)$ as diagonal transformations.*

In this way, S_∞ and $\text{Aut}_{\text{mp}}(X_0, \mu_0)$ are ergodic subgroups of $\text{Aut}(X, \mu)$ that are each other's centralizer inside $\text{Aut}(X, \mu)$.

Proof. For ever $\alpha \in \text{Aut}_{\text{mp}}(X_0, \mu_0)$, we denote by $\varphi_\alpha \in \text{Aut}_{\text{mp}}(X, \mu)$ the diagonal transformation $(\varphi_\alpha(x))_n = \alpha(x_n)$. When α is a mixing transformation, φ_α is mixing. So, $\text{Aut}_{\text{mp}}(X_0, \mu_0)$ is an ergodic subgroup of $\text{Aut}(X, \mu)$. Since $S_\infty < \text{Aut}_{\text{mp}}(X, \mu)$ contains a Bernoulli shift, also $S_\infty < \text{Aut}_{\text{mp}}(X, \mu)$ is an ergodic subgroup. To prove the lemma, it thus suffices to prove that S_∞ and $\text{Aut}_{\text{mp}}(X_0, \mu_0)$ are each other's centralizer inside $\text{Aut}_{\text{mp}}(X, \mu)$.

First assume that $\varphi \in \text{Aut}_{\text{mp}}(X, \mu)$ commutes with S_∞ . For every $k \in \mathbb{N}$, denote by $S_{\infty, k} < S_\infty$ the stabilizer of $k \in \mathbb{N}$. Since the action of $S_{\infty, k}$ on $(X_0, \mu_0)^{\mathbb{N} \setminus \{k\}}$ is ergodic and since φ commutes with all $S_{\infty, k}$, it follows that $\varphi = \prod_{k \in \mathbb{N}} \alpha_k$, where $\alpha_k \in \text{Aut}_{\text{mp}}(X_0, \mu_0)$. Since φ commutes with S_∞ , it then follows that all α_k are equal. So, $\varphi = \varphi_\alpha$ for some $\alpha \in \text{Aut}_{\text{mp}}(X_0, \mu_0)$.

Conversely, assume that $\psi \in \text{Aut}_{\text{mp}}(X, \mu)$ commutes with all φ_α , $\alpha \in \text{Aut}_{\text{mp}}(X_0, \mu_0)$. For every $\beta \in \text{Aut}_{\text{mp}}(X, \mu)$, we denote by $U_\beta \in \mathcal{U}(L^2(X, \mu))$ the corresponding unitary operator. Below we prove that we may choose a sequence $\alpha_n \in \text{Aut}_{\text{mp}}(X_0, \mu_0)$ such that

$$U_{\alpha_n} \rightarrow P_{\mathbb{C} \cdot 1} + \frac{1}{2} P_{(\mathbb{C} \cdot 1)^\perp} \quad \text{weakly in } B(L^2(X_0, \mu_0)). \quad (5.1)$$

Write $\varphi_n = \varphi_{\alpha_n}$. We identify $L^2(X, \mu)$ with the infinite tensor product of the Hilbert spaces $L^2(X_0, \mu_0)$ w.r.t. the vector $1 \in L^2(X_0, \mu_0)$. For every finite subset $\mathcal{F} \subset \mathbb{N}$, we define the closed subspace $H_{\mathcal{F}} \subset L^2(X, \mu)$ by

$$H_{\mathcal{F}} = \overline{\text{span}} \left(\bigotimes_{i \in \mathcal{F}} (L^2(X_0, \mu_0) \ominus \mathbb{C}1) \right) \subset L^2(X, \mu).$$

Note that $H_{\mathcal{F}} \perp H_{\mathcal{F}'}$ when $\mathcal{F} \neq \mathcal{F}'$. For every $k \in \mathbb{N}$, we write

$$L_k = \bigoplus_{\mathcal{F}: |\mathcal{F}|=k} H_{\mathcal{F}}$$

with the convention that $L_0 = \mathbb{C}1$. It then follows from (5.1) that

$$U_{\varphi_n} \rightarrow \sum_{k=0}^{\infty} 2^{-k} P_{L_k} \quad \text{weakly in } B(L^2(X, \mu)).$$

Since ψ commutes with all φ_α , we get that U_ψ commutes with all U_{φ_n} and hence also with its weak limit. It follows that $U_\psi(L_1) = L_1$.

Choose a Haar unitary $v : X_0 \rightarrow \mathbb{T}$ that generates $L^\infty(X_0, \mu_0)$, meaning that $v : X_0 \rightarrow \mathbb{T}$ is a measure preserving isomorphism from (X_0, μ_0) onto the circle with the Lebesgue measure. Denote by $\pi_i : L^2(X_0, \mu_0) \rightarrow L^2(X, \mu)$ the embedding as functions only depending on the i -th coordinate. Fix $i \in \mathbb{N}$. Define $w : X \rightarrow \mathbb{T}$ by $w(x) = v((\psi^{-1}(x))_i)$. So, $w = U_\psi(\pi_i(v))$ and because $U_\psi(L_1) = L_1$, we can uniquely write

$$w = \sum_{j \in \mathbb{N}} \pi_j(v_j) \quad \text{with } v_j \in L^2(X_0, \mu_0) \ominus \mathbb{C}1 \text{ and } L^2\text{-convergence.}$$

Since $w^2 = U_\psi(\pi_i(v^2)) \in L_1$, it follows that for all $j \neq k$, $\pi_j(v_j)\pi_k(v_k) = 0$. Taking the $\|\cdot\|_2$ -norm, we get that $\|v_j\|_2 \|v_k\|_2 = \|\pi_j(v_j)\pi_k(v_k)\|_2 = 0$ for all $j \neq k$. We conclude that at

most one of the v_j is nonzero. So there is a unique $j \in \mathbb{N}$ and $v_j \in L^2(X_0, \mu_0) \ominus \mathbb{C}1$ such that $U_\psi(\pi_i(v)) = w = \pi_j(v_j)$. Since v generates $L^\infty(X_0, \mu_0)$, we get that $U_\psi(\pi_i(L^\infty(X_0, \mu_0))) \subset \pi_j(L^\infty(X_0, \mu_0))$. Since a similar argument holds for ψ^{-1} , we find a permutation $\sigma \in S_\infty$ and a family $\beta_i \in \text{Aut}_{\text{mp}}(X_0, \mu_0)$ such that $U_\psi(\pi_i(v)) = \pi_{\sigma(i)}(U_{\beta_i}(v))$ for all $i \in \mathbb{N}$ and $v \in L^\infty(X_0, \mu_0)$.

Expressing that ψ commutes with all φ_α , it follows that every β_i commutes with every $\alpha \in \text{Aut}_{\text{mp}}(X_0, \mu_0)$. So, $\beta_i = \text{id}$ and we have proven that $\psi = \sigma$.

We still need to construct $\alpha_n \in \text{Aut}_{\text{mp}}(X_0, \mu_0)$ satisfying (5.1). We may realize (X_0, μ_0) as $(X_1 \times X_2, \mu_1 \times \mu_2)$, where $(X_1, \mu_1) = (\{0, 1\}^{\mathbb{Z}}, \nu^{\mathbb{Z}})$, $\nu(0) = \nu(1) = 1/2$ and (X_2, μ_2) is a nonatomic standard probability space. Choose a mixing $\gamma \in \text{Aut}_{\text{mp}}(X_2, \mu_2)$. Then define $\alpha_n \in \text{Aut}_{\text{mp}}(X_1 \times X_2, \mu_1 \times \mu_2)$ by

$$\alpha_n(x, y) = (x', y') \quad \text{with} \quad \begin{cases} (x', y') = (x, y) & \text{if } x_n = 0, \\ x'_k = x_{2n-k} & \text{for all } k \in \mathbb{Z} \text{ and } y' = \gamma^n(y) & \text{if } x_n = 1. \end{cases}$$

One easily checks (5.1). \square

We now repeat part of the proof of [PV21, Proposition 8.11] to prove Theorem 5.1 and use the opportunity to give a more self-contained argument.

Proof of Theorem 5.1. Fix a closed subgroup $G < S_\infty$. For every integer $n \geq 1$, denote by $I_{n,k} \subset \mathbb{N}^n$ an enumeration of the orbits of the diagonal action $G \curvearrowright \mathbb{N}^n$. Denote by λ the Lebesgue measure on the circle \mathbb{T} and define

$$(X, \mu) = (\mathbb{T}, \lambda)^{\mathbb{N}} \times \prod_{n,k} (\mathbb{T}, \lambda)^{I_{n,k}}.$$

Define the subgroup $\Lambda_0 < \text{Aut}_{\text{mp}}(X, \mu)$ consisting of the diagonal transformations $\alpha^{\mathbb{N}} \times \text{id}$, $\alpha \in \text{Aut}_{\text{mp}}(\mathbb{T}, \lambda)$, acting in the first factor of (X, μ) . Define $\Lambda_{n,k} < \text{Aut}_{\text{mp}}(X, \mu)$ as the subgroup of all diagonal transformations $\alpha^{I_{n,k}}$, with $\alpha \in \text{Aut}_{\text{mp}}(\mathbb{T}, \lambda)$, that only act in the factor $(\mathbb{T}, \lambda)^{I_{n,k}}$ and act as the identity elsewhere.

For every $x \in X$, we denote by $x_0 \in \mathbb{T}^{\mathbb{N}}$ and $x_{n,k} \in \mathbb{T}^{I_{n,k}}$ its natural coordinates. For every n, k and $1 \leq j \leq n$, define the transformation $\theta_{n,k,j} \in \text{Aut}_{\text{mp}}(X, \mu)$ by

$$\theta_{n,k,j}(x_0, (x_{n',k'})_{n',k'}) = (x_0, (y_{n',k'})_{n',k'})$$

$$\text{with} \quad (y_{n',k'})_i = \begin{cases} (x_0)_{i_j} \cdot (x_{n,k})_i & \text{if } (n', k') = (n, k), \\ (x_{n',k'})_i & \text{if } (n', k') \neq (n, k). \end{cases}$$

Define $\Lambda < \text{Aut}_{\text{mp}}(X, \mu)$ as the subgroup generated by Λ_0 , all $\Lambda_{n,k}$ and all $\theta_{n,k,j}$. We prove the centralizer of Λ inside $\text{Aut}_{\text{mp}}(X, \mu)$ is isomorphic with G .

Fix a $\psi \in \text{Aut}_{\text{mp}}(X, \mu)$ that commutes with Λ . Since ψ commutes with all Λ_0 and $\Lambda_{n,k}$, by Lemma 5.2, we find permutations σ_0 of \mathbb{N} and $\sigma_{n,k}$ of $I_{n,k}$ such that $\psi = \sigma_0 \times \prod_{n,k} \sigma_{n,k}$. Expressing that ψ commutes with all $\theta_{n,k,j}$, we get that $(\sigma_{n,k}(i))_j = \sigma_0(i_j)$ for all $i \in I_{n,k}$ and $1 \leq j \leq n$. This means that σ_0 is a permutation of \mathbb{N} with the property that the permutation σ_0^n of \mathbb{N}^n globally preserves all orbits of $G \curvearrowright \mathbb{N}^n$, for all $n \geq 1$, and that

$$\psi = \sigma_0 \times \prod_{n,k} \sigma_0^n|_{I_{n,k}}.$$

To conclude the proof of the theorem, it thus suffices to prove that such a permutation σ_0 must belong to G .

Take $i_1, \dots, i_n \in \mathbb{N}$. Since σ_0^n globally preserves the orbits of $G \curvearrowright \mathbb{N}^n$, there exists a $g \in G$ such that $\sigma_0^n(i) = g \cdot i$. This means that $\sigma_0(i_j) = g \cdot i_j$ for all $j \in \{1, \dots, n\}$. We have proven that σ_0 lies in the closure of G inside S_∞ . Since G is a closed subgroup, it follows that $\sigma_0 \in G$. \square

Lemma 5.2 also gives immediately the following result. Given a compact second countable group K , recall that the group $\text{Aut } K$ of automorphisms of K is a Polish group as a closed subgroup of the group of homeomorphisms of K .

Proposition 5.3. *For every compact second countable group K , the automorphism group $\text{Aut } K$ is a centralizer group.*

Proof. Denote by λ the Haar probability measure on K . Write $(X, \mu) = (K, \lambda)^\mathbb{N}$. Denote by $\Lambda < \text{Aut}_{\text{mp}}(X, \mu)$ the subgroup generated by S_∞ , acting by permuting the coordinates, and $\theta \in \text{Aut}_{\text{mp}}(X, \mu)$ given by

$$(\theta(x))_n = \begin{cases} x_1 x_2 & \text{if } n = 1, \\ x_n & \text{if } n \neq 1. \end{cases}$$

We prove that the centralizer of Λ inside $\text{Aut}_{\text{mp}}(X, \mu)$ is isomorphic with $\text{Aut } K$. Take $\psi \in \text{Aut}_{\text{mp}}(X, \mu)$ that commutes with Λ . For every $\alpha \in \text{Aut}_{\text{mp}}(K, \lambda)$ denote by $\varphi_\alpha \in \text{Aut}_{\text{mp}}(X, \mu)$ its diagonal product. Since ψ commutes with S_∞ by Lemma 5.2, we find $\alpha \in \text{Aut}_{\text{mp}}(K, \lambda)$ such that $\psi = \varphi_\alpha$. Since ψ commutes with θ , we get that $\alpha(xy) = \alpha(x)\alpha(y)$ for a.e. $(x, y) \in K \times K$. By [Zim84, Theorems B.2 and B.3], there is a unique $\beta \in \text{Aut } K$ such that $\alpha(x) = \beta(x)$ for a.e. $x \in K$. Since conversely every automorphism of K preserves λ , we have identified the centralizer of Λ with $\text{Aut } K$. \square

References

- [Dep10] S. Deprez, Explicit examples of equivalence relations and II_1 factors with prescribed fundamental group and outer automorphism group. *Trans. Amer. Math. Soc.* **367** (2015), 6837-6876.
- [FV07] S. Falguières and S. Vaes, Every compact group arises as the outer automorphism group of a II_1 factor. *J. Funct. Anal.* **254** (2008), 2317-2328.
- [IPP05] A. Ioana, J. Peterson and S. Popa, Amalgamated free products of weakly rigid factors and calculation of their symmetry groups. *Acta Math.* **200** (2008), 85-153.
- [JRD20] E. Janvresse, E. Roy and T. De La Rue, Dynamical systems of probabilistic origin: Gaussian and Poisson systems. In *Ergodic theory*, *Encycl. Complex. Syst. Sci.*, Springer, New York, 2023. pp. 217-232.
- [PV06] S. Popa and S. Vaes, Strong rigidity of generalized Bernoulli actions and computations of their symmetry groups. *Adv. Math.* **217** (2008), 833-872.
- [PV21] S. Popa and S. Vaes, W^* -rigidity paradigms for embeddings of II_1 factors. *Commun. Math. Phys.* **395** (2022), 907-961.
- [Roy08] E. Roy, Poisson suspensions and infinite ergodic theory. *Ergodic Theory Dynam. Systems* **29** (2009), 667-683.
- [Sut85] C.E. Sutherland, A Borel parametrization of Polish groups. *Publ. Res. Inst. Math. Sci.* **21** (1985), 1067-1086.
- [Vae07] S. Vaes, Explicit computations of all finite index bimodules for a family of II_1 factors. *Ann. Sci. Éc. Norm. Supér.* **41** (2008), 743-788.
- [VW24] S. Vaes and L. Wouters, Borel fields and measured fields of Polish spaces, Banach spaces, von Neumann algebras and C^* -algebras. *Preprint*. [arXiv:2405.16603](https://arxiv.org/abs/2405.16603)
- [Zim84] R.J. Zimmer, Ergodic theory and semisimple groups. *Monogr. Math.* **81**, Birkhäuser Verlag, Basel, 1984.