Generalized mixed product ideals whose powers have a linear resolution

MONICA LA BARBIERA AND ROYA MOGHIMIPOR

Abstract

In this paper we study classes of monomial ideals for which all of its powers have a linear resolution. Let $K[x_1, x_2]$ be the polynomial ring in two variables over the field K, and let L be the generalized mixed product ideal induced by a monomial ideal I. It is shown that, if $I \subset K[x_1, x_2]$ and the ideals substituting the monomials in I are Veronese type ideals, then L^k has a linear resolution for all $k \geq 1$. Furthermore, we compute some algebraic invariants of generalized mixed product ideals induced by a transversal polymatroidal ideal.

Key Words: Linear resolution, Monomial localization, Monomial ideals.2010 Mathematics Subject Classification: 13C13, 13D02

1 Introduction

The study of algebraic and homological properties of powers of a monomial ideal has been one of the main topics of combinatorial commutative algebra in recent years. Restuccia and Villarreal [13] introduced mixed product ideals, which form a particular class of squarefree monomial ideals, and they classified those among these ideals which are normal. In other words, they characterized the mixed product ideals whose Rees ring is normal.

In this paper we consider generalized mixed product ideals which were introduced by Herzog and Yassemi and which also include the so-called expansions of monomial ideals. The main objective of this paper is to study all powers of generalized mixed product ideals with linear resolution.

Let K be a field and $K[x_1, \ldots, x_n]$ the polynomial ring in n variables over K with each x_i of degree 1. Let $I \subset S$ be a monomial ideal and G(I) its unique minimal monomial generators.

Let K be a field and $S = K[x_1, \ldots, x_n, y_1, \ldots, y_m]$ be the polynomial ring over K in the variables x_i and y_j . A mixed product ideal is a monomial ideal of the form $(I_q J_r + I_{q'} J_{r'})S$, where for integers a and b, the ideal I_a (resp. J_b) is the ideal generated by all squarefree monomials of degree a in the polynomial ring $K[x_1, \ldots, x_n]$ (resp. of degree b in the polynomial ring $K[y_1, \ldots, y_m]$), and where $0 < q' < q \le n, 0 < r < r' \le m$. Thus, the ideal $L = (I_q J_r + I_{q'} J_{r'})S$ is obtained from the monomial ideal $I = (x^q y^r, x^{q'} y^{r'})$ by replacing x^q by I_q , $x^{q'}$ by $I_{q'}$, y^r by J_r and $y^{r'}$ by $J_{r'}$.

Together with Herzog and Yassemi [5] I introduced the generalized mixed product ideals, which are a far reaching generalization of the mixed product ideals introduced by Restuccia and Villarreal, and also generalizes the expansion construction by Bayati and Herzog [1]. For this construction we choose for each *i* a set of new variables $x_{i1}, x_{i2}, \ldots, x_{im_i}$ and replace each of the factor $x_i^{a_i}$ in each minimal generator $x_1^{a_1} x_2^{a_2} \cdots x_n^{a_n}$ of the monomial ideal *I* by a monomial ideal in $T_i = K[x_{i1}, x_{i2}, \ldots, x_{im_i}]$ generated in degree a_i .

The class of generalized mixed polymatroidal ideals is a special class of generalized mixed product ideals which for each i we replace each factor $x_i^{a_i}$ in each minimal generator $x_1^{a_1} x_2^{a_2} \cdots x_n^{a_n}$ of I by a polymatroidal ideal in T_i generated in degree a_i . In [12] we showed that powers of a generalized mixed polymatroidal ideal is generalized mixed polymatroidal ideal and monomial localizations of a generalized mixed polymatroidal ideal. A great deal of knowledge on the generalized mixed product ideal is accumulated in several papers [5, 7, 8, 9, 10, 11].

The present paper is organized as follows. In Section 2 we study all powers of generalized mixed product ideals with linear resolution. In Theorem 2.6 we compute the dimension of L in the case that $I \subset K[x_1, x_2]$ and the ideals who substitute the generators of I are the

ideals I_{ih_i} of Veronese type of degree h_i in the variables x_{i1}, \ldots, x_{im_i} . We also characterize the unmixed generalized mixed product ideals, see Theorem 2.8.

Our main result (Theorem 2.12) says that, if $I \subset K[x_1, x_2]$ is a monomial ideal and the ideals who substitute the generators of I are the ideals of Veronese type generated on degree h_i by the set $\{x_{i1}^{a_{i1}} \cdots x_{im_i}^{a_{im_i}} \mid \sum_{j=1}^{m_i} a_{ij} = h_i, 0 \leq a_{ij} \leq 2\}$, then L^k has a linear resolution for all $k \geq 1$.

Moreover, we compute the Castelnuovo-Mumford regularity of L^k for all $k \ge 1$, see Theorem 2.13.

In Section 3 we obtain an irredundant primary decomposition for any power of a generalized mixed product ideal L induced by the transversal polymatroidal ideal I, see Theorem 3.6. Let $\operatorname{astab}(I)$ and $\operatorname{dstab}(I)$ be the smallest integer k for which $\operatorname{Ass}_S(I^k)$ and $\operatorname{depth}(S/I^k)$ stabilize, respectively. In Theorem 3.5 we compute the stable set of associated prime ideals of L, and show that $\operatorname{astab}(L) = 1$ if I is a transversal polymatroidal ideal.

Furthermore, we study the depth of powers of L, and the index of depth stability of L is determined. In Theorem 3.9 it is shown that L is a Cohen–Macaulay transversal polymatroidal ideal if I is a Cohen–Macaulay transversal polymatroidal ideal.

2 Generalized mixed product ideals

Let R be a Noetherian ring and M a finitely generated R-module. A prime ideal $P \subset R$ is called an *associated prime ideal* of M, if there exists an element $x \in M$ such that P =Ann(x). Here Ann(x) is the *annihilator* of x, that is to say, $Ann(x) := \{f \in R : fx = 0\}$. The set of associated prime ideals of M is denoted $Ass_R(M)$.

Recall that an ideal I in a Noetherian ring R is P-primary, if $\operatorname{Ass}_R(R/I) = \{P\}$. In an abuse of notation, one often writes $\operatorname{Ass}_R(I)$ instead of $\operatorname{Ass}_R(R/I)$.

In [5] we introduced the generalized mixed product ideals. Let $S = K[x_1, \ldots, x_n]$ be the polynomial ring over a field K in the variables x_1, \ldots, x_n with the maximal ideal $\mathfrak{m} = (x_1, \ldots, x_n)$. Let $I \subset S$ be a monomial ideal with $I \neq S$ whose minimal set of generators is $G(I) = {\mathbf{x}^{\mathbf{a}_1}, \ldots, \mathbf{x}^{\mathbf{a}_m}}$. Here we set $\mathbf{x}^{\mathbf{a}} = x_1^{\mathbf{a}(1)} x_2^{\mathbf{a}(2)} \cdots x_n^{\mathbf{a}(n)}$ for $\mathbf{a} = (\mathbf{a}(1), \ldots, \mathbf{a}(n)) \in \mathbb{Z}_+^n$. In addition, we consider the polynomial ring T over K in the variables $x_{11}, \ldots, x_{1m_1}, x_{21}, \ldots, x_{2m_2}, \ldots, x_{n1}, \ldots, x_{nm_n}$.

For i = 1, ..., n and j = 1, ..., m let $L_{i,\mathbf{a}_j(i)}$ be a monomial ideal in the variables $x_{i1}, x_{i2}, ..., x_{im_i}$ such that

$$L_{i,\mathbf{a}_{j}(i)} \subset L_{i,\mathbf{a}_{k}(i)}$$
 whenever $\mathbf{a}_{j}(i) \ge \mathbf{a}_{k}(i)$. (1)

Given these ideals we define for j = 1, ..., m the monomial ideals

$$L_j = \prod_{i=1}^n L_{i,\mathbf{a}_j(i)} \subset T,$$
(2)

and set $L = \sum_{j=1}^{m} L_j$. The ideal L is called a *generalized mixed product ideal* induced by I.

Example 2.1. Let $T = K[x_{11}, \ldots, x_{1m_1}, x_{21}, \ldots, x_{2m_2}]$, and let $L = I_q J_r + I_p J_s \subset T$ be the generalized mixed product ideal induced by a monomial ideal $I = (x^q y^r, x^p y^s)$, where for integers α and β , the ideal I_α (resp. J_β) is the ideal generated by all squarefree monomials of degree α in the polynomial ring $K[x_{11}, \ldots, x_{1m_1}]$ (resp. of degree β in the polynomial ring $K[x_{21}, \ldots, x_{2m_2}]$), and where $0 , <math>0 < r < s \le m_2$. Ideals of this type are called squarefree Veronese ideals.

We want to compute the dimension of L in the case that $I \subset K[x_1, x_2]$ and the ideals who substitute the generators of I are the ideals I_{ih_i} of Veronese type of degree h_i in the variables x_{i1}, \ldots, x_{im_i} . Let \mathcal{F} be a non empty subset of $\{x_{11}, \ldots, x_{1m_1}, x_{21}, \ldots, x_{2m_2}, \ldots, x_{n1}, \ldots, x_{nm_n}\}$. We denote by $P_{\mathcal{F}}$ the prime ideal of T generated by the variables whose index is in \mathcal{F} .

Herzog and Hibi [4] proposed the concept of discrete polymatroid and they studied some algebraic and combinatorics properties related to it. Let I be a monomial ideal of S which is generated in a single degree. Then I is said to be *polymatroidal*, if for any two elements $u, v \in G(I)$ such that $\deg_{x_i} u > \deg_{x_i} v$ there exists an index j with $\deg_{x_j} u < \deg_{x_j} v$ such that $x_j(u/x_i) \in I$. The Veronese type ideals is a special class of polymatroidal ideals, introduced in [14].

Definition 2.2. Given a sequence of integers $(r_1, \ldots, r_n; d)$ such that $1 \le r_j \le d \le \sum_{i=1}^n r_i$ for all j, we define \mathcal{J} as the set of partitions

$$\mathcal{J} = \{ (\mathbf{a}(1), \dots, \mathbf{a}(n)) \in \mathbb{Z}^n \text{such that } \mathbf{a}(1) + \dots + \mathbf{a}(n) = d; \quad 0 \leq \mathbf{a}(i) \leq r_i \quad \forall i \}$$

and H as the set of monomials $H = {\mathbf{x}^{\mathbf{a}} \mid \mathbf{a} \in \mathcal{J}} = {f_1, \ldots, f_q}$. The ideal $I = (H) \subset S$ is said to be of Veronese type of degree d with defining sequence $(r_1, \ldots, r_n; d)$. If $r_i = 1$ for all i we call I the dth squarefree Veronese ideal, and if $r_i = d$ for all i we call I the dth veronese ideal.

In the following, we give a description of the associated prime ideals of L where the ideals L_{ij} are Veronese type ideals.

Theorem 2.3. Let $T = K[x_{11}, \ldots, x_{1m_1}, x_{21}, \ldots, x_{2m_2}]$ with $m_1, m_2 > 1$. Let

$$L = \sum_{1 \le h_i \le 2m_i, \sum_{i=1}^2 h_i = d} I_{1h_1} I_{2h}$$

where the ideals I_{ih_i} are the ideals of Veronese type generated on degree h_i by the set $\{x_{i1}^{a_{i1}} \cdots x_{im_i}^{a_{im_i}} \mid \sum_{j=1}^{m_i} a_{ij} = h_i, \quad 0 \le a_{ij} \le 2\}$. Then $P_{\mathcal{F}} \in \operatorname{Ass}_T(L)$ if and only if $|\mathcal{F}| \le 2$ for $d = 2m_1 + 2m_2 - 1$.

Proof. The ideal L is a generalized mixed product ideal induced by the ideal I generated by the monomials $x_1^{h_1}x_2^{h_2}$ with $\sum_{i=1}^2 h_i = d$ and $1 \leq h_i \leq 2m_i$. T. For a subset \mathcal{F} we denote by $P_{\mathcal{F}}$ the prime ideal of T generated by the variables whose index is in \mathcal{F} . We replace the set of variables $\{x_{11}, \ldots, x_{1m_1}\}$ with $\{y_1, \ldots, y_{m_1}\}$ and $\{x_{21}, \ldots, x_{2m_2}\}$ with $\{y_{m_1+1}, \ldots, y_{m_1+m_2}\}$. Suppose that $P_{\mathcal{F}} \in \operatorname{Ass}_T(L)$. Thus there exists a monomial $f \notin L$ such that $L : f = P_{\mathcal{F}}$. We prove that we can choose such a monomial f of degree d-1such that $L : f = P_{\mathcal{F}}$.

We assume that $f \notin L$, $L : f = P_{\mathcal{F}}$, $\deg(f) \geq d$ and $f = y_1^{a_1} \cdots y_{m_1+m_2}^{a_{m_1+m_2}}$. Hence, there exists an integer $r \in \{1, 2, \ldots, m_1 + m_2\}$ such that $a_r > 2$. Since $L : f = P_{\mathcal{F}}$, it follows that $y_t f \in L$ for all $t \in \mathcal{F}$ and $y_t f \notin L$ for all $t \notin \mathcal{F}$. Thus, there exists a monomial $v_t \in G(L)$ such that $v_t | (y_t f)$ for all $t \in \mathcal{F}$. Since $f \notin L$, we conclude that for all $t \in \mathcal{F}$, the variable y_t appears in v_t with exponent $a_t + 1$. This implies that $a_t < 2$ for all $t \in \mathcal{F}$. Therefore, $r \notin \mathcal{F}$.

Now let $g = f/y_r$. Since $f \notin L$, it follows that $g \notin L$. We claim that $L : g = P_{\mathcal{F}}$. In fact, since g divides f, it follows that $L : g \subseteq L : f$. Hence, $L : g \subseteq P_{\mathcal{F}}$. Now, we want to prove that $P_{\mathcal{F}} \subseteq L : g$. Let $y_t \in P_{\mathcal{F}}$. Since $P_{\mathcal{F}} = L : f$, it follows that $y_t f \in L$, so there exists $v_t \in G(L)$ such that $v_t \mid y_t f$. Now, since $a_r - 1 \ge 2$, we have that $v_t \mid y_t f/y_r$. Hence, $y_t \in L : f/y_r = L : g$. Therefore, $P_{\mathcal{F}} = L : g$

We proceed as before and obtain after a finite number of these reductions a monomial $f \notin L$ of degree d-1 such that $P_{\mathcal{F}} = L : f$. Then $fy_t \in L$ for all $t \in \mathcal{F}$ and $fy_t \notin L$ for all $t \notin \mathcal{F}$. Notice that $a_t + 1 \leq 2$ for all $t \in \mathcal{F}$, and $a_t \leq 2$ for all $t \notin \mathcal{F}$. Thus, $a_t = 2$ for all $t \notin \mathcal{F}$. Hence, $f = \prod_{t \in \mathcal{F}} y_t^{a_t} \prod_{t \notin \mathcal{F}} y_t^2$ with $0 \leq a_t < 2$ for all $t \in \mathcal{F}$. In particular we see that $\deg(\prod_{t \notin \mathcal{F}} y_t^2) = 2(m_1 + m_2 - |\mathcal{F}|) = q$. Then

$$2m_1 + 2m_2 \ge \sum_{t \in \mathcal{F}} (a_t + 1) + q = \sum_{t \in \mathcal{F}} a_t + |\mathcal{F}| + q = \deg(f) + |\mathcal{F}| = d - 1 + |\mathcal{F}|.$$

Hence $|\mathcal{F}| \leq 2$. Conversely, assume that $|\mathcal{F}| \leq 2$ for $d = 2m_1 + 2m_2 - 1$. Then for any monomial $v \in G(L)$ there exists an integer $l \in \mathcal{F}$ such that y_l divides v. Thus, $L \subset P_{\mathcal{F}}$. By using the condition $2m_1 + 2m_2 \geq d - 1 + |\mathcal{F}|$ we have $|\mathcal{F}| + 2m_1 + 2m_2 - 2|\mathcal{F}| \geq d - 1$, which together with $2m_1 + 2m_2 - 2|\mathcal{F}| \leq d - 1$ implies that there exists an integer $b_t < 2$ for all $t \in \mathcal{F}$ such that $b_t |\mathcal{F}| + 2m_1 + 2m_2 - 2|\mathcal{F}| = d - 1$. Therefore, the monomial $f = \prod_{t \in \mathcal{F}} y_t^{b_t} \prod_{t \notin \mathcal{F}} y_t^2$ has degree d - 1, and hence $f \notin L$. Thus $P_{\mathcal{F}} \subseteq L : f$.

Suppose, on the contrary, that $P_{\mathcal{F}}$ is a proper subset of L : f. Then there exists a monomial w, in the variables y_t with $t \notin \mathcal{F}$, of degree at least 1 such that $fw \in L$. Thus, we obtain that there exists a monomial $v = y_1^{a_1} \cdots y_{m_1+m_2}^{a_{m_1+m_2}} \in G(L)$ such that v divides fw. Observe that $a_t \leq b_t$ for all $t \in \mathcal{F}$, because $w \in K[y_t \mid t \notin \mathcal{F}]$. This implies that

$$d = \deg(v) = \sum_{t=1}^{m_1+m_2} a_t \le \sum_{t \in \mathcal{F}} b_t + 2m_1 + 2m_2 - 2|\mathcal{F}| = \deg(f) = d - 1;$$

a contradiction. Therefore, $P_{\mathcal{F}}$ is not a proper subset of L : f. This shows that $P_{\mathcal{F}} \in Ass_T(L)$.

Example 2.4. Let $T = K[x_{11}, x_{12}, x_{21}, x_{22}]$ be a polynomial ring over a field K, and let $L = \sum_{1 \le h_i \le 4, \sum_{i=1}^2 h_i = 7} I_{1h_1} I_{2h_2}$ where the ideals I_{ih_i} are the ideals of Veronese type generated on degree h_i by the set $\{x_{i1}^{a_{i1}} \cdots x_{im_i}^{a_{im_i}} \mid \sum_{j=1}^{m_i} a_{ij} = h_i, 0 \le a_{ij} \le 2\}$. This implies that

$$L = (x_{11}^2 x_{12} x_{21}^2 x_{22}^2, x_{11} x_{12}^2 x_{21}^2 x_{22}^2, x_{11}^2 x_{12}^2 x_{21}^2 x_{22}^2, x_{11}^2 x_{12}^2 x_{21}^2 x_{21}^2 x_{22}^2).$$

By using Theorem 2.3 one obtains

$$Ass_T(L) = \{(x_{11}), (x_{12}), (x_{21}), (x_{22}), (x_{11}, x_{12}), (x_{21}, x_{22}), (x_{11}, x_{21}), (x_{11}, x_{22}), (x_{12}, x_{21}), (x_{12}, x_{21}), (x_{12}, x_{22})\}.$$

Definition 2.5. A vertex cover of L is a subset C of

 $\{x_{11},\ldots,x_{1m_1},x_{21},\ldots,x_{2m_2},\ldots,x_{n1},\ldots,x_{nm_n}\}$

such that each $v \in G(L)$ is divided by some variables of C. Such a vertex cover is called *minimal* if no proper subset of C is vertex cover.

The minimal cardinality of the vertex covers of L is denoted by height(L). The reader can find more information in [14].

Theorem 2.6. Let $T = K[x_{11}, \ldots, x_{1m_1}, x_{21}, \ldots, x_{2m_2}]$ with $m_i = q > 1$ for i = 1, 2. Let $L = \sum_{1 \le h_i \le 2m_i, \sum_{i=1}^2 h_i = d} I_{1h_1} I_{2h_2}$ where the ideals I_{ih_i} are the ideals of Veronese type generated on degree h_i by the set $\{x_{i1}^{a_{i1}} \cdots x_{im_i}^{a_{im_i}} \mid \sum_{j=1}^{m_i} a_{ij} = h_i, 0 \le a_{ij} \le 2\}$.

- (i) If $2 \le d \le 2m_1 + 2$, then $\dim(T/L) = q$.
- (ii) If $d = 2(m_1 + m_2) 1$, then $\dim(T/L) = m_1 + m_2 1$.

Proof. (i) Let $L = \sum_{r=1}^{m} I_{1h_{1r}} I_{2h_{2r}}$ where the ideals $I_{1h_{1r}}$ in $K[x_{11}, x_{12}, \ldots, x_{1m_1}]$ and the ideals $I_{2h_{2r}}$ in $K[x_{21}, x_{22}, \ldots, x_{2m_2}]$ are Veronese type ideals of degree h_{1r} and h_{2r} , respectively. Assume further that

 $1 \le h_{1_1} < \dots < h_{1_{m_1}} \le 2m_1$ and $2m_2 \ge h_{2_1} > \dots > h_{2_{m_2}} \ge 1$,

where $h_{1_{r+1}} = h_{1_r} + 1$, $h_{2_{r-1}} = h_{2_r} + 1$ for all $r = 1, \dots, m-1$.

Let G(L) be the unique minimal set of monomial generators of L. A vertex cover of L is a subset C of $\{x_{11}, \ldots, x_{1m_1}, x_{21}, \ldots, x_{2m_2}\}$ such that each $u \in G(L)$ is divided by some variables of C. Thus, P is a minimal prime ideal of L if and only if P = (C) for some minimal vertex cover C of L.

We may suppose that $h_{1_r} + h_{2_r} = d$ for all r = 1, ..., m. Then the ideal L is a generalized mixed product ideal induced by the ideal

$$I = (x_1^{h_{1_1}} x_2^{h_{2_1}}, \dots, x_1^{h_{1_{m_1}}} x_2^{h_{2_{m_2}}}).$$

Then

$$I = (x_1^{h_{1_1}}) \cap \bigcap_{r=1}^{m-1} (x_1^{h_{1_{r+1}}}, x_2^{h_{2_r}}) \cap (x_2^{h_{2_{m_2}}}).$$

This is the irredundant decomposition of I as an intersection of irreducible ideals. It then follows from [8, Theorem 1.2] that

$$L = (I_{1h_{1_1}}) \cap \sum_{r=1}^{m-1} (I_{1h_{1_{r+1}}}, I_{2h_{2_r}}) \cap (I_{2h_{2_{m_2}}}).$$

Hence, [8, Corollary 1.7] implies that

$$\operatorname{height}(L) = \min_{r=1,\dots,m-1} \{\operatorname{height}(I_{1h_{1_1}}), \operatorname{height}(I_{2h_{2m_2}}), \operatorname{height}(I_{1h_{1_{r+1}}}) + \operatorname{height}(I_{2h_{2r}})\}.$$

Now we consider $2 \leq d \leq 2m_1 + 2$. Then the minimal vertex covers of L are $C_1 = \{x_{11}, \ldots, x_{1m_1}\}, C_2 = \{x_{21}, \ldots, x_{2m_2}\}$. Therefore,

$$\dim(T/L) = \dim T - \operatorname{height}(L) = 2q - \min\{m_1, m_2\}.$$

(ii) The minimal vertex covers of L are $C_l = \{x_{1l}\}$ for $l = 1, \ldots, m_1$ and $C'_p = \{x_{2p}\}$ for $p = 1, \ldots, m_2$. Hence, the minimal cardinality of the vertex covers of L is height(L) = 1, as desired.

Example 2.7. Let $T = K[x_{11}, x_{12}, x_{21}, x_{22}]$ be a polynomial ring over a field K, and let $L = \sum_{1 \le h_i \le 2, \sum_{i=1}^2 h_i = 3} I_{1h_1} I_{2h_2} \subset T$ where the ideals I_{ih_i} are the ideals of Veronese type generated on degree h_i by the set $\{x_{i1}^{a_{i1}} \cdots x_{im_i}^{a_{im_i}} \mid \sum_{j=1}^{m_i} a_{ij} = h_i, 0 \le a_{ij} \le 2\}$. The ideal L is a generalized mixed product ideal induced by the ideal I generated by the monomials $x_1^{h_1} x_2^{h_2}$ with $h_1 + h_2 = 3$ and $h_1, h_2 \ge 1$. Then $I = (x_1 x_2^2, x_1^2 x_2)$. One easily checks that

$$I = (x_1) \cap (x_1^2, x_2^2) \cap (x_2).$$

This is the irredundant decomposition of I as an intersection of irreducible ideals. It then follows that the minimal vertex covers of L are $C_1 = \{x_{11}, x_{12}\}, C_2 = \{x_{21}, x_{22}\}$. Therefore, height $(L) = |C_1| = |C_2| = 2$, and hence dim(T/L) = 2 by Theorem 2.6.

A monomial ideal is said to be *unmixed* if all its minimal vertex covers have the same cardinality. We recall the one-to-one correspondence between the minimal vertex covers of an ideal and its minimal prime ideals.

Now we assume that the generalized mixed product ideals induced by a monomial ideal $I \subset K[x_1, x_2]$, and give a criterion for its unmixedness in the case that the ideals I_{ih_i} are the ideals of Veronese type of degree h_i in the variables x_{i1}, \ldots, x_{im_i} .

Theorem 2.8. Let $T = K[x_{11}, \ldots, x_{1m_1}, x_{21}, \ldots, x_{2m_2}]$ with $m_i = q > 1$ for i = 1, 2. Let $L = \sum_{1 \le h_i \le 2m_i, \sum_{i=1}^2 h_i = d} I_{1h_1} I_{2h_2}$ where the ideals I_{ih_i} are the ideals of Veronese type generated on degree h_i by the set $\{x_{i1}^{a_{i1}} \cdots x_{im_i}^{a_{im_i}} \mid \sum_{j=1}^{m_i} a_{ij} = h_i, 0 \le a_{ij} \le 2\}$.

- (i) If $q \ge 3$ and $d = 2(m_1 + m_2) 1$, then L is unmixed.
- (ii) If q = 2 and $2 \le d \le 2(m_1 + m_2) 1$, then *L* is unmixed.

Proof. (i) The ideal L is a generalized mixed product ideal induced by the ideal I generated by the monomials $x_1^{h_1}x_2^{h_2}$ with $h_1 + h_2 = d$ and $h_1, h_2 \ge 1$. One has $C_p = \{x_{1p}\}, p = 1, \ldots, m_1, C'_{p'} = \{x_{2p'}\}, p' = 1, \ldots, m_2$ are the minimal vertex covers of L. Hence, the minimal cardinality of the vertex covers of L is one. Then the assertion follows.

(ii) The minimal vertex covers of L are $C_1 = \{x_{11}, x_{12}\}, C_2 = \{x_{21}, x_{22}\}$. Thus, all the minimal vertex covers have the same cardinality, as desired.

Example 2.9. Let $T = K[x_{11}, x_{12}, x_{13}, x_{14}, x_{21}, x_{22}, x_{23}, x_{24}]$ be a polynomial ring over a field K, and let $L = \sum_{1 \le h_i \le 14, \sum_{i=1}^2 h_i = 15} I_{1h_1} I_{2h_2}$ where the ideals I_{ih_i} are the ideals of Veronese type generated on degree h_i by the set $\{x_{i1}^{a_{i1}} \cdots x_{im_i}^{a_{im_i}} \mid \sum_{j=1}^{m_i} a_{ij} = h_i, 0 \le a_{ij} \le 2\}$. The minimal vertex covers of L are $C_l = \{x_{1l}\}$ for $l = 1, \ldots, 4$ and $C'_p = \{x_{2p}\}$ for $p = 1, \ldots, 4$. Therefore, height $(L) = |\mathcal{C}_l| = |\mathcal{C}'_p| = 1$ for $l = 1, \ldots, 4$ and $p = 1, \ldots, 4$. Hence, by applying Theorem 2.8 we obtain that L is unmixed.

In the following, we study all powers of generalized mixed product ideals induced by a monomial ideal $I \subset K[x_1, x_2]$, where the ideals substituting the monomials in I are Veronese type ideals. Note that a generalized mixed product ideal depends not only on Ibut also on the family L_{ij} . Then for a monomial ideal I with $G(I) = \{\mathbf{x}^{\mathbf{a}_1}, \ldots, \mathbf{x}^{\mathbf{a}_m}\}$ we set $L(I; \{L_{ij}\}) = \sum_{j=1}^m \prod_{i=1}^n L_{i,\mathbf{a}_j(i)}$.

Lemma 2.10. Let $T = K[x_{11}, ..., x_{1m_1}, x_{21}, ..., x_{2m_2}]$ be a polynomial ring over a field K with $m_1, m_2 > 1$. Let

$$L = \sum_{1 \le h_i \le 2m_i, \sum_{i=1}^2 h_i = d} I_{1h_1} I_{2h_2}$$

where the ideals I_{ih_i} are the ideals of Veronese type generated on degree h_i by the set $\{x_{i1}^{a_{i1}}\cdots x_{im_i}^{a_{im_i}} \mid \sum_{j=1}^{m_i} a_{ij} = h_i, \quad 0 \le a_{ij} \le 2\}$, and let

$$L' = \sum_{1 \le h'_i \le 2m_i, \sum_{i=1}^2 h'_i = d'} I_{1h'_1} I_{2h'_2}$$

where the ideals $I_{ih'_i}$ are the ideals of Veronese type generated on degree h'_i by the set $\{x_{i1}^{a'_{i1}} \cdots x_{im_i}^{a'_{im_i}} \mid \sum_{j=1}^{m_i} a'_{ij} = h'_i, \quad 0 \le a'_{ij} \le 2\}$. Assume further that $2 \le d \le 2m_1 + 2m_2 - 1$ and $2 \le d' \le 2m_1 + 2m_2 - 1$. Then LL' has a linear resolution.

Proof. Let $L(I; \{L_{ij}\}) = \sum_{1 \le h_i \le 2m_i, \sum_{i=1}^2 h_i = d} I_{1h_1} I_{2h_2}$ be the generalized mixed product ideal, where the ideals $I_{ih_i} \subset K[x_{i1}, \ldots, x_{im_i}]$ are the ideals of Veronese type generated on degree h_i by the set $\{x_{i1}^{a_{i1}} \cdots x_{im_i}^{a_{im_i}} | \sum_{j=1}^{m_i} a_{ij} = h_i, 0 \le a_{ij} \le 2\}$. Let \mathbb{Z}_+ be the set of non-negative integers and $\mathbb{Z}_+^{m_1+m_2}$ be the set of the vectors $(\mathbf{a}_1, \mathbf{a}_2)$ with $\mathbf{a}_1 \in \mathbb{Z}_+^{m_1}$ and $\mathbf{a}_2 \in \mathbb{Z}_+^{m_2}$, i.e., $(\mathbf{a}_1, \mathbf{a}_2) = (a_{11}, \ldots, a_{1m_1}, a_{21}, \ldots, a_{2m_2}) \in \mathbb{Z}_+^{m_1+m_2}$ with $a_{1j}, a_{2l} \ge 0$. Furthermore, let

$$\mathcal{A} = \{ (\mathbf{a}_1, \mathbf{a}_2) \in \mathbb{Z}_+^{m_1 + m_2} : \sum_{j=1}^{m_1} a_{1j} + \sum_{l=1}^{m_2} a_{2l} = d, |\mathbf{a}_1| = h_1, |\mathbf{a}_2| = h_2, \quad 0 \le a_{1j}, a_{2l} \le 2 \}$$

be the set of the vector exponents of the elements of $G(L(I; \{L_{ij}\}))$. The modulus of the vector $(\mathbf{a}_1, \mathbf{a}_2)$ is the number $|(\mathbf{a}_1, \mathbf{a}_2)| = |\mathbf{a}_1| + |\mathbf{a}_2| = \sum_{j=1}^{m_1} a_{1j} + \sum_{l=1}^{m_2} a_{2l}$. We assume that $(\mathbf{a}_1, \mathbf{a}_2), (\mathbf{b}_1, \mathbf{b}_2) \in \mathcal{A}$ with $a_{1r} > b_{1r}$ for some r. Hence, $a_{1r} - 1 \ge b_{1r}$. Then

$$|(a_{11}, \dots, a_{1r} - 1, \dots, a_{1m_1}, a_{21}, \dots, a_{2m_2})| = |(b_{11}, \dots, b_{1m_1}, b_{21}, \dots, b_{2m_2})| - 1 < 0$$

$$|(b_{11},\ldots,\ldots,b_{1m_1},b_{21},\ldots,b_{2m_2})|.$$

Thus, there exists r'' such that $(a_{11}, \ldots, a_{1r} - 1, \ldots, a_{1r''} + 1, \ldots, a_{1m_1}, a_{21}, \ldots, a_{2m_2}) \in \mathcal{A}$. Notice that $r \neq r''$. Then for $r \neq r''$, one has $a_{1r''} + 1 \leq b_{1r''}$. Hence, $a_{1r''} < b_{1r''}$. Then $L(I; \{L_{ij}\})$ is generated by all the monomials $\mathbf{X}_1^{\mathbf{a}_1} \mathbf{X}_2^{\mathbf{a}_2}$ with $(\mathbf{a}_1, \mathbf{a}_2) \in \mathcal{A}$, where $\mathbf{X}_1^{\mathbf{a}_1} \mathbf{X}_2^{\mathbf{a}_2}$ stands for $x_{11}^{a_{11}} \cdots x_{1m_1}^{a_{1m_1}} x_{21}^{a_{21}} \cdots x_{2m_2}^{a_{2m_2}}$.

Suppose that $u = \mathbf{X}_{1}^{\mathbf{a}_{1}} \mathbf{X}_{2}^{\mathbf{a}_{2}}$, $v = \mathbf{X}_{1}^{\mathbf{b}_{1}} \mathbf{X}_{2}^{\mathbf{b}_{2}} \in G(L(I; \{L_{ij}\}))$ with $a_{1r} > b_{1r}$ or $a_{2t} > b_{2t}$, for some $r'' \in \{1, \ldots, m_{1}\}$ with $a_{1r''} < b_{1r''}$ and $t'' \in \{1, \ldots, m_{2}\}$ with $a_{2t''} < b_{2t''}$, we have $(a_{11}, \ldots, a_{1r} - 1, \ldots, a_{1r''} + 1, \ldots, a_{1m_{1}}, a_{21}, \ldots, a_{2m_{2}}) \in \mathcal{A}$ and $(a_{11}, \ldots, a_{1m_{1}}, a_{21}, \ldots, a_{2t} - 1, \ldots, a_{2t''} + 1, \ldots, a_{2m_{2}}) \in \mathcal{A}$. Therefore, $x_{2r''}(u/x_{1r}) \in G(L(I; \{L_{ij}\}))$ or $x_{2t''}(u/x_{2t}) \in G(L(I; \{L_{ij}\}))$.

The ideal $L(I; \{L_{ij}\})$ is a generalized mixed product ideal induced by the ideal I generated by the monomials $x_1^{h_1} x_2^{h_2}$ with $\sum_{i=1}^2 h_i = d$, $1 \le h_i \le 2m_i$ and the ideal $L(I'; \{L_{ij}\})$ is a generalized mixed product ideal induced by the ideal I' generated by the monomials $x_1^{h'_1} x_2^{h'_2}$ with $\sum_{i=1}^2 h'_i = d$ and $1 \le h'_i \le 2m_i$. Then

$$I_{ih_i}I_{ih'_i} \subset I_{iq_i}I_{iq'_i}$$
 whenever $h_i + h'_i \ge q_i + q'_i$.

Thus, [12, Lemma 2.2] yields that $L(II'; \{L_{ij}\})$ is a generalized mixed product ideal, and $L(II'; \{L_{ij}\}) = L(I; \{L_{ij}\})L(I'; \{L_{ij}\})$. It then follows that the ideal $L(II'; \{L_{ij}\})$ is induced by the ideal II' generated by the monomials $x_1^{h_1+h'_1}x_2^{h_2+h'_2}$ with $1 \le h_i \le 2m_i$ and $1 \le h'_i \le 2m_i$. Since the ideals I and I' are polymatroidal ideals, it follows that II' is a polymatroidal ideal, by [4, Theorem 12.6.3]. Therefore, [4, Corollary 12.6.4] together with [4, Proposition 8.2.1] yield II' has a linear resolution. Hence by [12, Theorem 2.5] the desired follows.

Definition 2.11. The ideal $L \subset T$ has linear quotients if there is an ordering u_1, \ldots, u_r of the monomials belonging to G(L) with $\deg(u_1) \leq \cdots \leq \deg(u_r)$ such that for each $2 \leq j \leq r$, the colon ideal $(u_1, \ldots, u_{j-1}) : u_j$ is generated by a subset of

$$\{x_{11},\ldots,x_{1m_1},x_{21},\ldots,x_{2m_2},\ldots,x_{n1},\ldots,x_{nm_n}\}.$$

The reader can find more information in [14, Definition 6.3.45]. We are now come to the main result of present paper.

Theorem 2.12. Let $T = K[x_{11}, \ldots, x_{1m_1}, x_{21}, \ldots, x_{2m_2}]$ with $m_1, m_2 > 1$, and let

$$L = \sum_{1 \le h_i \le 2m_i, \sum_{i=1}^2 h_i = d} I_{1h_1} I_{2h_2}$$

where the ideals I_{ih_i} are the ideals of Veronese type generated on degree h_i by the set $\{x_{i1}^{a_{i1}} \cdots x_{im_i}^{a_{im_i}} \mid \sum_{j=1}^{m_i} a_{ij} = h_i, \quad 0 \le a_{ij} \le 2\}$. Furthermore, let $2 \le d \le 2m_1 + 2m_2 - 1$. Then L^k has a linear resolution for all $k \ge 1$.

Proof. We proceed with induction on k. For k = 1, we suppose that $L(I; \{L_{ij}\}) = \sum_{1 \le h_i \le 2m_i, \sum_{i=1}^2 h_i = d} I_{1h_1} I_{2h_2}$ be the generalized mixed product ideal, where the ideals $I_{ih_i} \subset K[x_{i1}, \ldots, x_{im_i}]$ are the ideals of Veronese type of degree h_i . Assume further that $v \in G(L(I; \{L_{ij}\}))$.

We define $\mathcal{I} = (w \in G(L(I; \{L_{ij}\})) : w \prec v)$ with \prec the lexicographical order on $x_{11}, \ldots, x_{1m_1}, x_{21}, \ldots, x_{2m_2}$ induced by $x_{11} \succ \cdots \succ x_{1m_1} \succ x_{21} \succ \cdots \succ x_{2m_2}$. We denote by (v, w) the greatest common divisor of v and w. It is well-known that $\mathcal{I} : v = (w/(v, w) \mid w \in \mathcal{I})$.

We claim that for any $w \prec v$, there exists a variable of T in $\mathcal{I}: v$ such that it divides w/(v, w). Suppose that $v = \mathbf{X}_1^{\mathbf{a}_1} \mathbf{X}_2^{\mathbf{a}_2}, w = \mathbf{X}_1^{\mathbf{b}_1} \mathbf{X}_2^{\mathbf{b}_2} \in G(L(I; \{L_{ij}\}))$. Indeed, since $w \prec v$, there exists an integer l such that $a_{1l} > b_{1l}$ and $a_{1q} = b_{1q}$ for $q = 1, \ldots, l-1$. Then there exists an integer l' with $a_{1l'} < b_{1l'}$ such that $x_{1l'}(v/x_{1l}) \in G(L(I; \{L_{ij}\}))$.

We know that l < l', it follows that $x_{1l'}(v/x_{1l}) \in \mathcal{I}$ and hence $x_{1l'} \in \mathcal{I} : v$. It is then clear that $x_{1l'}$ divides w/(v, w), because the l'-th component of the vector exponent of w/(v, w) is given by

$$b_{1l'} - \min\{b_{1l'}, a_{1l'}\} = b_{1l'} - a_{1l'} > 0.$$

If we assume that $a_{1q} = b_{1q}$ for all $q = 1, \ldots, m_1$, $a_{2l} > b_{2l}$ and $a_{2r} = b_{2r}$ for all $r = 1, \ldots, l-1$, then $x_{2l'} \in \mathcal{I} : v$ and $x_{2l'}$ divides w/(v, w). Therefore, $L(I; \{L_{ij}\})$ has linear quotients, and hence $L(I; \{L_{ij}\})$ has a *d*-linear resolution by [4, Proposition 8.2.1].

For k = 2 Lemma 2.10 implies that $L(I; \{L_{ij}\})^2$ has a 2d-linear resolution by [4, 1 toposition 0.2.1]. For k = 2 Lemma 2.10 implies that $L(I; \{L_{ij}\})^2$ has a 2d-linear resolution and $L(I^2; \{L_{ij}\}) = L(I; \{L_{ij}\})^2$. Now let k > 2. Thus, $L(I^k; \{L_{ij}\}) = L(I^{k-1}I; \{L_{ij}\})$ and we have as in the case k = 2 that $L(I^{k-1}I; \{L_{ij}\}) = L(I^{k-1}; \{L_{ij}\})L(I; \{L_{ij}\})$. By induction hypothesis and Lemma 2.10, $L(I; \{L_{ij}\})^{k-1}$ has a d(k-1)-linear resolution and $L(I; \{L_{ij}\})^{k-1} = L(I^{k-1}; \{L_{ij}\})$. Hence, $L(I; \{L_{ij}\})^k$ has a kd-linear resolution and

$$L(I; \{L_{ij}\})^k = L(I^k; \{L_{ij}\}).$$

Then the desired conclusion follows.

Together with Theorem 2.12 the following result is also useful to compute the Castelnuovo-Mumford regularity of powers of L in the case that all L_{ij} are Veronese type ideals.

Theorem 2.13. Let $T = K[x_{11}, \ldots, x_{1m_1}, x_{21}, \ldots, x_{2m_2}]$ with $m_1, m_2 > 1$, and let

$$L = \sum_{1 \le h_i \le 2m_i, \sum_{i=1}^2 h_i = d} I_{1h_1} I_{2h_2}$$

where the ideals I_{ih_i} are the ideals of Veronese type generated on degree h_i by the set $\{x_{i1}^{a_{i1}}\cdots x_{im_i}^{a_{im_i}} \mid \sum_{j=1}^{m_i} a_{ij} = h_i, \quad 0 \le a_{ij} \le 2\}$. Then $\operatorname{reg}(L^k) = kd$ for all $k \ge 1$.

Example 2.14. Let $T = K[x_{11}, x_{12}, x_{21}, x_{22}]$ be a polynomial ring over a field K in the variables $x_{11}, x_{12}, x_{21}, x_{22}$. Let $L = \sum_{1 \le h_i \le 4, \sum_{i=1}^2 h_i = 4} I_{1h_1} I_{2h_2}$, where the ideals I_{ih_i} are the ideals of Veronese type generated on degree h_i by the set $\{x_{i1}^{a_{i1}} \cdots x_{im_i}^{a_{im_i}} \mid \sum_{j=1}^{m_i} a_{ij} = h_i, 0 \le a_{ij} \le 2\}$. Then

$$L = (x_{11}^2 x_{12} x_{21}, x_{11}^2 x_{12} x_{22}, x_{11} x_{12}^2 x_{21}, x_{11} x_{12}^2 x_{22}, x_{11} x_{21}^2 x_{22}, x_{12} x_{21}^2 x_{22}, x_{11} x_{21} x_{22}^2, x_{12} x_{21}^2 x_{22}^2, x_{12} x_{21} x_{22}^2, x_{11} x_{12} x_{21} x_{22}^2,$$

Therefore, Theorem 2.13 yields $\operatorname{reg}(L^k) = 4k$ for all $k \ge 1$.

3 Stability of depth and Cohen-Macaulayness of generalized mixed product ideals induced by a transversal polymatroidal ideal

Let $I \subset S = K[x_1, \ldots, x_n]$ be a monomial ideal as in Section 1 with $G(I) = \{\mathbf{x}^{\mathbf{a}_1}, \ldots, \mathbf{x}^{\mathbf{a}_m}\}$, and L be defined as in (2).

The main goal of this section is to study algebraic properties of powers of generalized mixed product ideals induced by a transversal polymatroidal ideal. We denote the set of

monomial prime ideals of S by $\mathcal{P}(S)$. Let F be a non-empty subset of [n]. We denote by P_F the monomial prime ideal $(x_i)_{i \in F}$. We assume that $L(P_F; \{L_{ij}\})$ be the generalized mixed product ideal induced by P_F . Hence, by applying [8, Theorem 1.2] we obtain that

$$L(P_F; \{L_{ij}\}) = (L_{i,1})_{i \in F}.$$

We set $P'_F = (L_{i,1})_{i \in F}$, where the ideals $L_{i,1} = (x_{i1}, \ldots, x_{im_i})$ for all $i \in F$. We denote the set of monomial prime ideals of T by $\mathcal{P}(T)$.

A transversal polymatroidal ideal is an ideal I of the form

$$I = P_{F_1} P_{F_2} \cdots P_{F_r},\tag{3}$$

where F_1, \ldots, F_r is a collection of non-empty subsets of [n] with $r \ge 1$. By its definition, the product of transversal polymatroidal ideals is again a transversal polymatroidal ideal. Then by taking powers of the prime ideal factors of I which appear several times in (3), we have

$$I = \prod_{t=1}^{s} P_{G_t}^{a_t} \quad \text{with} \quad a_t \ge 1$$

$$\tag{4}$$

where $G_t \neq G_q$ for $t \neq q$.

In order to determine the associated prime ideals of L induced by the transversal polymatroidal ideal I we will need the following lemma.

Lemma 3.1. [12, Lemma 1.2] Let $L(I; \{L_{ij}\}) = \sum_{k=1}^{r} \prod_{i=1}^{n} L_{i,\mathbf{a}_{k}(i)}$ and $L(J; \{L_{ij}\}) = \sum_{l=1}^{s} \prod_{i=1}^{n} L_{i,\mathbf{b}_{l}(i)}$ be generalized mixed product ideals, respectively, induced by the monomial ideals I and J with $G(I) = \{\mathbf{x}^{\mathbf{a}_{1}}, \ldots, \mathbf{x}^{\mathbf{a}_{r}}\}$ and $G(J) = \{\mathbf{x}^{\mathbf{b}_{1}}, \ldots, \mathbf{x}^{\mathbf{b}_{s}}\}$. We assume that

$$L_{i,\mathbf{a}_{k}(i)}L_{i,\mathbf{b}_{l}(i)} \subset L_{i,\mathbf{a}_{p}(i)}L_{i,\mathbf{b}_{q}(i)} \text{ whenever } \mathbf{a}_{k} + \mathbf{b}_{l} \ge \mathbf{a}_{p} + \mathbf{b}_{q}.$$
(5)

Suppose that $G(IJ) = {\mathbf{x}^{\mathbf{c}_1}, \dots, \mathbf{x}^{\mathbf{c}_t}}$. Then given \mathbf{c}_j , there exist \mathbf{a}_k and \mathbf{b}_l such that $\mathbf{c}_j = \mathbf{a}_k + \mathbf{b}_l$. We set $L_{i,\mathbf{c}_j(i)} = L_{i,\mathbf{a}_k(i)}L_{i,\mathbf{b}_l(i)}$. Furthermore, let $L(IJ; {L_{ij}}) = \sum_{j=1}^t \prod_{i=1}^n L_{i,\mathbf{c}_j(i)}$. Then $L(IJ; {L_{ij}})$ is a generalized mixed product ideal, and

$$L(IJ; \{L_{ij}\}) = L(I; \{L_{ij}\})L(J; \{L_{ij}\})$$

Theorem 3.2. Let $I = P_{F_1}P_{F_2} \cdots P_{F_r} \subset S$ be a transversal polymatroidal ideal, where F_1, \ldots, F_r is a collection of non-empty subsets of [n] with $r \geq 1$. Let $L(I; \{L_{ij}\})$ be a generalized mixed product ideal induced by the monomial ideal I satisfying condition (5). Then $L(I; \{L_{ij}\})$ is a transversal polymatroidal ideal, and

$$L(I; \{L_{ij}\}) = L(P_{F_1}; \{L_{ij}\})L(P_{F_2}; \{L_{ij}\}) \cdots L(P_{F_r}; \{L_{ij}\}).$$

Proof. Let $I = P_{F_1} P_{F_2} \cdots P_{F_r}$. Thus, Lemma 3.1 implies that

$$L(I; \{L_{ij}\}) = L(P_{F_1}P_{F_2}\cdots P_{F_r}; \{L_{ij}\})$$

= $L(P_{F_1}; \{L_{ij}\})L(P_{F_2}; \{L_{ij}\})\cdots L(P_{F_r}; \{L_{ij}\}),$

is a generalized mixed product ideal induced by I, where $L(P_{F_d}; \{L_{ij}\}) = (L_{i,1})_{i \in F_d}$ for all d with $1 \leq d \leq r$. Since $L_{i,1} \subset K[x_{i1}, \ldots, x_{im_i}]$ is a monomial prime ideal, this implies that $L(I; \{L_{ij}\}) \subset T$ is a transversal polymatroidal ideal, as desired.

By Theorem 3.2, Lemma 3.1 together with [4, Theorem 12.6.2] yield

Corollary 3.3. Let $I = P_{F_1}P_{F_2} \cdots P_{F_r} \subset S$ be a transversal polymatroidal ideal, where F_1, \ldots, F_r is a collection of non-empty subsets of [n] with $r \geq 1$. Let L be a generalized mixed product ideal induced by the monomial I satisfying condition (5). Then L^k has a linear resolution for all $k \geq 1$.

Example 3.4. Let $F_1 = \{1, 2\}$, $F_2 = \{1, 2, 3, 4\}$, $F_3 = \{3, 5\}$, $F_4 = \{4, 5\}$ and $I = P_{F_1} \cdots P_{F_4}$ be the transversal polymetroidal ideal of $K[x_1, \ldots, x_5]$. Notice that

$$I = (x_1, x_2)(x_1, x_2, x_3, x_4)(x_3, x_5)(x_4, x_5)$$

Let

$$L = (L_{1,1}, L_{2,1})(L_{1,1}, L_{2,1}, L_{3,1}, L_{4,1})(L_{3,1}, L_{5,1})(L_{4,1}, L_{5,1})$$

be the generalized mixed product ideal induced by I in the polynomial ring

$$T = K[x_{11}, \dots, x_{1m_1}, x_{21}, \dots, x_{2m_2}, \dots, x_{51}, \dots, x_{5m_5}]$$

where the ideals $L_{i,1}$ are Veronese ideals of degree one in the variables $x_{i1}, x_{i2}, \ldots, x_{im_i}$ for $i = 1, \ldots, 5$. Therefore, Corollary 3.3 implies that L^k has a linear resolution for all $k \ge 1$.

Now we obtain the index of stability and the stable set of associated primes of L induced by a transversal polymatroidal ideal I. Brodmann showed [2] that there exists an integer k_1 such that $\operatorname{Ass}_S(I^k) = \operatorname{Ass}_S(I^{k_1})$ for all $k \ge k_1$. The smallest such number is called *index* of stability of I. We denote this number by $\operatorname{astab}(I)$. The stable set $\operatorname{Ass}_S(I^k)$ is denoted by $\operatorname{Ass}^{\infty}(I)$.

Theorem 3.5. Let L be the generalized mixed product ideal induced by a transversal polymatroidal ideal $I \subset S$ satisfying condition (5). Then $\operatorname{astab}(L) = 1$, that is

$$\operatorname{Ass}_T(L) = \operatorname{Ass}^\infty(L).$$

Proof. Let $I = P_{F_1}P_{F_2}\cdots P_{F_r} \subset S$ be a transversal polymatroidal ideal, where F_1, \ldots, F_r is a collection of non-empty subsets of [n] with $r \geq 1$. We assume that $k \geq 1$ be an integer. Furthermore, let $L(I; \{L_{ij}\})$ be the generalized mixed product ideal induced by I satisfying condition (5).

We denote I(P) the monomial localization of a monomial ideal I, and by S(P) the polynomial ring over K in the variables which belong to P. The monomial localization I(P)is the monomial ideal which is obtained from I as the image of the K-algebra homomorphism $\varphi: S \to S(P)$ which $\varphi(x_i) = x_i$ if $x_i \in P$, and $\varphi(x_i) = 1$, otherwise. According to Theorem 3.2 we have

$$L(I; \{L_{ij}\}) = L(P_{F_1}; \{L_{ij}\})L(P_{F_2}; \{L_{ij}\}) \cdots L(P_{F_r}; \{L_{ij}\}),$$

where $L(P_{F_d}; \{L_{ij}\}) = (L_{i,1})_{i \in F_d}$ for all $d = 1, \ldots, r$. Theorem 3.2 together with Lemma 3.1 now yields $L(I; \{L_{ij}\})^k$ is a transversal polymatroidal ideal induced by I^k , and $L(I; \{L_{ij}\})^k = L(I^k; \{L_{ij}\})$. By [6, Lemma 2.3], we obtain that $P \in \operatorname{Ass}_T(L(I; \{L_{ij}\})^k)$ if and only if depth $(T(P)/L(I; \{L_{ij}\})^k(P)) = 0$. We know that the localization of a transversal polymatroidal ideal is again a transversal polymatroidal ideal. Then $L(I; \{L_{ij}\})(P)$ is again a transversal polymatroidal ideal. Since powers of transversal polymatroidal ideals are again transversal polymatroidal ideal, and since by [4, Theorem 12.6.2] transversal polymatroidal ideals have linear resolutions. Thus, all powers of $L(I; \{L_{ij}\})(P)$ have a linear resolution. Therefore, [6, Proposition 2.2] yields depth $(T(P)/L(I; \{L_{ij}\})^q(P)) = 0$ for all $q \ge k$ and hence $P \in \operatorname{Ass}_T(L(I; \{L_{ij}\})^q)$ for all $q \ge k$. Thus, $\operatorname{Ass}_T(L(I; \{L_{ij}\})) \subset \operatorname{Ass}^{\infty}(L(I; \{L_{ij}\}))$.

For the converse inclusion, let $P \in \operatorname{Ass}^{\infty}(L(I; \{L_{ij}\}))$. Then there exists $k \geq 1$ such that $P \in \operatorname{Ass}_T(L(I; \{L_{ij}\})^k)$. Hence, by applying [6, Lemma 2.3] and [6, Corollary 4.5] we obtain that $P \in \operatorname{Ass}_T(L(I; \{L_{ij}\}))$. This yields the desired conclusion.

Next we study the irredundant primary decomposition of all powers of generalized mixed product ideals induced by a transversal polymatroidal ideal. Let $I = P_{F_1}P_{F_2}\cdots P_{F_r} \subset S$ be a transversal polymatroidal ideal. Without loss of generality we may assume that $\bigcup_{d=1}^r F_d = [n]$.

Let G_I be a graph attached to the ideal I on the vertex set $\{1, \ldots, r\}$ with $\{i, j\}$ an edge of G_I if $F_i \cap F_j \neq \emptyset$. In [6, Theorem 4.7] it is shown that the associated prime ideals of I corresponded to the trees of the graph G_I in the following way: $P_F \in \operatorname{Ass}_S(I)$ if and only if there exists a tree \mathcal{T} such that $P_F = P_{\mathcal{T}}$.

For each subgraph \mathcal{H} of G_I we associate the prime ideal $P_{\mathcal{H}} = \sum_{d \in \mathcal{V}(\mathcal{H})} P_{F_d}$. Furthermore, for a monomial prime ideal $P_{\mathcal{H}}$ we set

$$P'_{\mathcal{H}} = \sum_{d \in \mathcal{V}(\mathcal{H})} P'_{F_d}.$$

Notice that $P'_{\mathcal{H}}$ is a generalized mixed product ideal induced by the monomial prime $P_{\mathcal{H}}$, where the ideals $L_{i,1} = (x_{i1}, \ldots, x_{im_i})$ for all $i \in F_d$.

Theorem 3.6. Let I be a transversal polymatroidal ideal with $Ass_S(I) = \{P_1, \ldots, P_q\}$, and let L be the generalized mixed product ideal induced by I, where all $L_{i,1} = (x_{i1}, x_{i2}, \ldots, x_{im_i})$ for $d = 1, \ldots, r$. Consider $\mathcal{T}_1, \ldots, \mathcal{T}_q$ maximal trees of G_I such that $P_h = P_{\mathcal{T}_h}$ for all $h = 1, \ldots, q$. Then

$$L^k = \bigcap_{h=1}^q (P'_h)^{ka_h},$$

is an irredundant primary decomposition of L^k for any $k \ge 1$, where $a_h = |\mathcal{V}(\mathcal{T}_h)|$ for all h.

Proof. For a transversal polymatroidal ideal $I = P_{F_1}P_{F_2}\cdots P_{F_r} \subset S$, let $L(I; \{L_{ij}\})$ be the generalized mixed product ideal induced by I. By Theorem 3.2 together with Lemma 3.1 it follows that $L(I; \{L_{ij}\})^k$ is a transversal polymatroidal ideal induced by I^k , and $L(I; \{L_{ij}\})^k = L(I^k; \{L_{ij}\})$ for all $k \geq 1$.

Suppose that $\operatorname{Ass}_{S}(I) = \{P_1, \ldots, P_q\}$. Consider $\mathcal{T}_1, \ldots, \mathcal{T}_q$ maximal trees of G_I such that $P_h = P_{\mathcal{T}_h}$ for all $h = 1, \ldots, q$. Therefore, [6, Corollary 4.10] yields

$$I^k = \bigcap_{h=1}^q P_h^{ka_h},$$

is an irredundant primary decomposition of I^k , where $a_h = |\mathcal{V}(\mathcal{T}_h)|$ for all h. Hence, by [1, Proposition 1.2] we have

$$L(I; \{L_{ij}\})^{k} = L(I^{k}; \{L_{ij}\}) = L(\bigcap_{h=1}^{q} P_{h}^{ka_{h}}; \{L_{ij}\}) = \bigcap_{h=1}^{q} L(P_{h}; \{L_{ij}\})^{ka_{h}},$$

as desired.

Let I be a monomial ideal of S. We assume that t be a variable over S. The graded subalgebra $\mathcal{R}(I) := \bigoplus_{k=0}^{\infty} I^k t^k$ of S[t] is called the *Rees algebra* of I. The *K*-algebra $\mathcal{R}(I)/\mathfrak{m}\mathcal{R}(I)$ is called the *fiber ring* and its Krull dimension is called the *analytic spread* of I, denoted by $\ell(I)$.

By a theorem of Brodmann [3], depth (S/I^k) is constant for large k. We call this constant value the *limit depth* of I, and denote it by $\lim_{k\to\infty} \operatorname{depth}(S/I^k)$. The smallest integer k_0 such that depth $(S/I^k) = \operatorname{depth}(S/I^{k_0})$ for all $k \ge k_0$ is called the *index of depth stability* and is denoted by dstab(I).

In the following, we compute the analytic spread of L induced by a transversal polymatroidal ideal I.

Theorem 3.7. Let L be a generalized mixed product ideal induced by the transversal polymatroidal ideal I satisfying condition (5). Then $\operatorname{depth}(T/L) = \operatorname{depth}(T/L^k)$ for all $k \ge 1$. In particular, we have

$$\ell(L) = m_1 + \dots + m_m - \operatorname{depth}(T/L).$$

Proof. Let $I = P_{F_1}P_{F_2} \cdots P_{F_r} \subset S$ be a transversal polymatroidal ideal. Let $L(I; \{L_{ij}\})$ be a generalized mixed product ideal induced by I. Theorem 3.2 with Lemma 3.1 guarantees that $L(I; \{L_{ij}\})^k$ is a transversal polymatroidal ideal induced by I^k , and $L(I; \{L_{ij}\})^k =$ $L(I^k; \{L_{ij}\})$ for all $k \geq 1$. We denote the number of connected components of the graph G_I by $c(G_I)$. We know that $c(G_I) = c(G_{I^k})$ for any $k \geq 1$. Then [6, Theorem 4.12] implies that depth $(T/L(I; \{L_{ij}\})) = depth(T/L(I; \{L_{ij}\})^k)$. Therefore,

$$\lim_{k \to \infty} \operatorname{depth}(T/L(I; \{L_{ij}\})^k) = \operatorname{depth}(T/L(I; \{L_{ij}\}))$$

for all $k \geq 1$, and hence

$$\lim_{k \to \infty} \operatorname{depth}(T/L(I; \{L_{ij}\})^k) = m_1 + \dots + m_m - \ell(L(I; \{L_{ij}\}))$$

by [6, Corollary 3.5]. Thus, the desired conclusion follows.

Let k be a nonnegative integer. We define the k-th bracket power of I, to be the ideal $I^{[k]}$, generated by all monomials u^k , where $u \in I$ is a monomial. In particular, $I^{[0]} = S$, and $I^{[1]} = I$.

Note that if $G(I) = {\mathbf{x}^{\mathbf{a}_1}, \dots, \mathbf{x}^{\mathbf{a}_m}}$, then $G(I^{[k]}) = {\mathbf{x}^{k\mathbf{a}_1}, \dots, \mathbf{x}^{k\mathbf{a}_m}}$. Notice that $I^{[k]} \subset I^k$ for all k. When $k \ge 2$, the equality holds, if and only if I is principal. One can easily compute that $(I \cap J)^{[k]} = I^{[k]} \cap J^{[k]}$ for any monomial ideals $I, J \subset S$.

Let $I = P_{F_1}P_{F_2} \cdots P_{F_r} \subset S$ be a transversal polymatroidal ideal. Let G_1, \ldots, G_t be the connected components of G_I , with $t \geq 2$. Thus, $I = I_1 \cdots I_t$, where $I_a = \prod_{d \in \mathcal{V}(G_a)} P_{F_d}$. Furthermore, $G_a = G_{I_a}$ for all a. Note that $I = I_1 \cdots I_t = I_1 \cap \cdots \cap I_t$, we know that the ideals I_a are generated in pairwise disjoint sets of variables. Therefore, $I^{[k]} = I_1^{[k]} \cap \cdots \cap I_t^{[k]}$. Hence, [9, Lemma 2.3] implies that $\operatorname{Ass}_S(I^{[k]}) = \operatorname{Ass}_S(I_1) \cup \cdots \cup \operatorname{Ass}_S(I_t)$, where I_a is a transversal polymatroidal ideal with the associated connected graph G_{I_a} .

In the following, we want to determine the associated prime ideals of the k-th bracket power $L^{[k]}$ of generalized mixed product ideals L induced by a transversal polymatroidal ideal I.

Theorem 3.8. Let $I = P_{F_1}P_{F_2}\cdots P_{F_r}$ be a transversal polymatroidal ideal with the set of associated prime ideals $Ass_S(I) = \{P_1, \ldots, P_q\}$. Consider $\mathcal{T}_1, \ldots, \mathcal{T}_q$ maximal trees of G_I such that $P_h = P_{\mathcal{T}_h}$ for all $h = 1, \ldots, q$. Let L be a generalized mixed product ideal induced by I, where all $L_{i,1} = (x_{i1}, x_{i2}, \ldots, x_{im_i})$ for $d = 1, \ldots, r$. Then

$$\operatorname{Ass}_T(L^{[k]}) = \{ P'_h \mid h = 1, \dots, q \}.$$

for all $k \geq 1$.

Proof. The assertion follows by Theorem 3.2 and [9, Theorem 2.4].

Now we assume that the generalized mixed product ideals induced by a transversal polymatroidal ideal, and give a criterion for its Cohen-Macaulayness.

Theorem 3.9. Let L be the generalized mixed product ideal induced by a Cohen-Macaulay transversal polymatroidal ideal $I = P_{F_1}P_{F_2}\cdots P_{F_r}$ such that height(I) = t. Assume further that

$$L(P_{F_d}; \{L_{ij}\}) = (L_{i,1})_{i \in F_d}$$

where $L_{i,1} = (x_{i1}, \ldots, x_{im_i})$ for all d with $d = 1, \ldots, r$. If $m_1 = \cdots = m_n$ and $\operatorname{projdim}(T/L) = m_1 + \cdots + m_t$, then L is a Cohen-Macaulay transversal polymatroidal ideal.

Proof. Let $I = P_{F_1}P_{F_2}\cdots P_{F_r} \subset S$ be a Cohen-Macaulay transversal polymatroidal ideal. Then [6, Theorem 4.7] implies that

$$\dim S/I = n - \min\{|F_d| : d = 1, \dots, r\}.$$

We denote by t the minimal cardinality of a set F_d , where $d = 1, \ldots, r$. Thus, dim S/I = n-t. Let $L(I; \{L_{ij}\})$ be a generalized mixed product ideal induced by the monomial ideal I. We may assume that $\bigcup_{d=1}^{r} F_d = [n]$, and then it follows from [6, Theorem 4.12] that n-t=k-1, where k represents the number of connected components of G_I . We know that $n \ge kt$ it follows that $k-1=n-t \ge kt-t=(k-1)t$. Therefore we obtain that this inequality is valid either if k = 1 or t = 1. If k = 1, then t = n. This implies that $F_d = [n]$ for all d and thus $I = \mathfrak{m}^r$, the Veronese ideal. Otherwise t = 1 and hence k = n. In this case G_I has n connected components. Then r = n and $|F_d| = 1$ for all $d = 1, \ldots, r$. This yields that I is a principal ideal. Thus, Theorem 3.2, [4, Theorem 12.6.7] together with [8, Theorem 2.3] yield L is a Cohen-Macaulay transversal polymatroidal ideal, as desired.

Example 3.10. Let $I \subset S$ be Veronese ideal of degree r which is generated by all the monomials in the variables x_1, \ldots, x_n of degree r. Furthermore, let $L(\mathfrak{m}^r; \{L_{ij}\})$ be the generalized mixed product ideal induced by $I = \mathfrak{m}^r$, where the ideals substituting the monomials in I are Veronese ideals. It follows from [6, Lemma 2.3] that depth(S/I) = 0. We know that powers of Veronese ideals are again polymatroidal, see [4, Theorem 12.6.3], and since by [4, Theorem 12.6.2] Veronese ideals have linear resolutions, we obtain that all powers of I have a linear resolution. By [6, Proposition 1.3] we have depth $(S/I^k) = 0$ for all $k \geq 1$. Thus, Lemma 3.1 together with [1, Proposition 1.2] now yields $L(\mathfrak{m}; \{L_{ij}\}) \in Ass_T(L(\mathfrak{m}^r; \{L_{ij}\})^k)$ for all $k \geq 1$. Then $(x_{11}, x_{12}, \ldots, x_{nm_m}) \in Ass^{\infty}(L)$. Theorem 3.5 together with Theorem 3.7 implies that

$$\operatorname{astab}(L(\mathfrak{m}^r; \{L_{ij}\})) = \operatorname{dstab}(L(\mathfrak{m}^r; \{L_{ij}\})) = 1.$$

References

- S. Bayati and J. Herzog, Expansion of monomial ideals and multigraded modules, Rocky Mountain Journal of Mathematics, 44 (2014), 1781-1804.
- [2] M. Brodmann, Asymptotic stability of $Ass(M/I^nM)$, Proc. AMS 74 (1979), 16-18.
- [3] M. Brodmann, The asymptotic nature of the analytic spread, Math. Proc. Cambridge Philos. Soc. 86 (1979), no. 1, 35-39.
- [4] J. Herzog and T. Hibi, Monomial Ideals. GTM 260. Springer 2010.
- [5] J. Herzog, R. Moghimipor and S. Yassemi, *Generalized mixed product ideals*, Arch. Math. 103 (2014), 39-51.
- [6] J. Herzog, A. Rauf and M. Vladoiu, The stable set of associated prime ideals of a polymatroidal ideal, J. Algebraic Combinatorics, DOI 10.1007/s10801-012-0367-z.

- [7] M. La Barbiera, R. Moghimipor, Normality of Rees algebras of generalized mixed product ideals, International Electronic Journal of Algebra, DOI: 10.24330/ieja.1402961.
- [8] R. Moghimipor, Algebraic and homological properties of generalized mixed product ideals, Arch. Math, 114 (2020), 147-157.
- [9] R. Moghimipor, On the Cohen-Macaulayness of bracket powers of generalized mixed product ideals, Acta Mathematica Vietnamica, DOI: 10.1007/s40306-021-00460-z.
- [10] R. Moghimipor, On the normality of generalized mixed product ideals, Arch. Math. (Basel) 115 (2020), 147-157.
- [11] R. Moghimipor, Strong persistence property of generalized mixed product ideals, Bull. Math. Soc. Sci. Math. Roumanie Tome 66 (114), No 3, (2023), 281-294.
- [12] R. Moghimipor and A. Tehranian, Linear resolutions of powers of generalized mixed product ideals, Iran. J. Math. Sci. Inform., 14 (2019), 127-134.
- [13] G. Restuccia and R. Villarreal, On the normality of monomial ideals of mixed products, Commun. Algebra, 29 (2001), 3571-3580.
- [14] H. Villareal, Monomial algebras, Second Edition, Monographs and Research Notes in Mathematics, Chapman and Hall/CRC, 2015.

Department of Mathematics and Informatics, University of Messina, Viale Ferdinando Stagno d'Alcontres 31, 98166 Messina, Italy E-mail:monicalb@unime.it

> Department of Mathematics, Safadasht Branch, Islamic Azad University, Tehran, Iran E-mail: roya_moghimipour@yahoo.com