

# ON POLYNOMIAL SOLUTIONS TO THE MINIMAL SURFACE EQUATION

YIFAN GUO

ABSTRACT. We are interested in finding nonlinear polynomials  $P$  on  $\mathbb{R}^n$  that solves the minimal surface equation.

We first prove a structure theorem on such polynomials. We show that the highest degree term  $P_m$  must factor as  $P_0^k Q_m$  where  $k$  is odd,  $Q_m \geq 0$  on  $\mathbb{R}^n$  and  $P_0$  is irreducible, and that the level sets of  $P_m$  are all area minimizing. Moreover, the unique tangent cone of graph  $P$  at  $\infty$  is  $\{P_0 = 0\} \times \mathbb{R}$ . If  $k \geq 3$ , we know further that lower order terms down to some degree are divisible by  $P_0$ .

We also show that  $P$  must contain terms of both high and low degree. In particular, it cannot be homogeneous. As a corollary, we prove that there is no quadratic or cubic polynomial solution.

Using a general decay lower bound of Green's functions on minimal hypersurfaces we are able to show that  $\deg P_0 + k^{-1} \deg Q_m \leq n - 2$ .

Finally, we prove that a polynomial minimal graph cannot have  $C \times \mathbb{R}^l$  as its tangent cone at  $\infty$  where  $l \geq 1$  and  $C$  is any isoparametric cone. We also show that the existence of a nonlinear polynomial solutions on  $\mathbb{R}^8$  will imply the existence of an area minimizing but not strictly minimizing cubic cone in  $\mathbb{R}^8$ . These results indicate that finding an explicit polynomial solution can be hard.

## 1. INTRODUCTION

**1.1. Entire solutions to the minimal surface equation.** Let  $P$  be a function on  $\mathbb{R}^n$ . The graph of  $P$  has mean curvature 0 in  $\mathbb{R}^{n+1}$  is equivalent to  $P$  satisfying the minimal surface equation:

$$(1) \quad (1 + |\nabla P|)^2 \Delta P - \sum_{i,j=1}^n P_{x_i} P_{x_j} P_{x_i x_j} = 0.$$

Such function  $P$ , defined on all of  $\mathbb{R}^n$ , is called an entire solution to the minimal surface equation.

A well-known theorem of Bernstein [3] says that the only entire solutions to the minimal surface equation on  $\mathbb{R}^2$  are affine functions. The theorem was generalized to  $\mathbb{R}^7$  by the successive works of Fleming [13], De Giorgi [9], Almgren [2] and Simons [23].

On the other hand, Bombieri, De Giorgi, and Giusti [5] constructed nonlinear entire solutions to the minimal surface equation on  $\mathbb{R}^n$  for  $n \geq 8$  modeling on Simons' cone. Thus the restriction on  $n \leq 7$  in Bernstein theorem is sharp. Many other entire solutions have been constructed by Simon [21] modeling on isoparametric cones cf. subsection 1.5.

We are interested in finding entire solutions to the minimal surface equation which are *polynomials*. This is a problem raised in [22]. While graphs of complex polynomials are always minimal by a calibration argument, there is no known example of a real polynomial whose graph is minimal except affine functions. In fact, even though nonlinear entire minimal graphs are known to exist, none of them is explicit. In order to find an explicit example, we should try some special ansatz and polynomial seem to be a natural choice. The other motivation is the long standing conjecture that all entire solutions to the minimal surface equation have polynomial growth cf. [6, 22, 29]. This conjecture is verified by Simon [21] under some technical assumptions. Therefore, it is reasonable to believe that polynomials can be suitable candidates for solutions to the minimal surface equation.

**1.2. Structure of polynomial solutions.** We start by considering suitable blow-downs of graph  $P$  which will imply that a polynomial solution has a special structure. Neglecting the constant term, we write  $P$  as the sum of homogeneous polynomials

$$(2) \quad P = P_m + P_{m-1} + \cdots + P_2 + P_1$$

where  $P_i$  is homogeneous of degree  $i$ . We consider the graph obtained by scaling down graph  $P$  by  $\lambda$  and translating vertically by  $t\lambda^{-m+1}$  for some fixed  $t \in \mathbb{R}$ . We are able to show the following currents convergence

$$\lim_{\lambda \rightarrow 0} \eta_{\lambda\#} \text{graph } P - t\lambda^{-m+1} e_{n+1} = \partial[P_m < t] \times \mathbb{R}$$

where  $\eta_{\lambda} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is the map  $x \mapsto \lambda x$ . By Federer-Fleming compactness, this means that for all  $t \in \mathbb{R}$  the current  $\partial[P_m < t]$  is area minimizing i.e. the level sets of  $P_m$  are area minimizing. Using irreducibility of area minimizing currents as well as real Nullstellensatz, we obtain the following structure theorem for polynomial solutions.

**Theorem 1.1.** *Let  $P = P_m + P_{m-1} + \cdots + P_2 + P_1$  with  $m \geq 2$  solve the minimal surface equation where  $P_i$  is homogeneous of degree  $i$ . Then we have:*

- (i) *For every  $t \in \mathbb{R}$ , the current  $\partial[P_m < t]$  is area minimizing.*
- (ii) *There exist  $k \geq 1$  odd,  $P_0$  a nonlinear irreducible homogeneous polynomial that changes sign and  $Q_m$  an even degree homogeneous polynomial which is non-negative on  $\mathbb{R}^n$  and coprime with  $P_0$ , with  $\{Q_m = 0\}$  of dimension  $\leq n - 2$  so that*

$$(3) \quad P_m = P_0^k Q_m.$$

- (iii) *The unique tangent cone of graph  $P$  at  $\infty$  is  $\partial[P_0 < 0] \times \mathbb{R}$ .*
- (iv) *If  $k \geq 3$  in (3), then there exists  $2 \leq s \leq m - 1$  so that*

$$P = P_0^k Q_m + P_0^{k_{m-1}} Q_{m-1} + \cdots + P_0^{k_{s+1}} Q_{s+1} + P_0 Q_s + P_{s-1} + \cdots + P_1$$

where  $P_0$  and  $Q_i$  are coprime for  $l \leq i \leq m$  and  $k_i \geq 2$  for  $s + 1 \leq i \leq m - 1$ .

From (i) and (ii), we see that the zero locus of  $P_0$  defines an area-minimizing cone. Moreover,  $\{P_m = t\}$  for  $t \neq 0$  are smooth area-minimizing hypersurfaces lying on one side of  $\{P_0 = 0\}$  and thus they are the Hardt-Simon foliation of the cone  $\{P_0 = 0\}$ . The isoparametric cones introduced by Cartan [8] are perfect examples of such cones. However, we are going to show in Theorem 1.6 that all isoparametric cones *cannot* appear as  $\{P_0 = 0\}$  for any polynomial solution.

**1.3. No homogeneous solution to the minimal surface equation.** One convenience when we consider polynomial solutions to PDEs is that we can consider each degree terms separately. Notice that we write the minimal surface equation (1) in such a way that it maps a polynomial to a polynomial. To proceed, let  $L$  denote the 1-Laplacian:

$$LP := |\nabla P|^2 \Delta P - \sum_{i,j=1}^n P_{x_i} P_{x_j} P_{x_i x_j}.$$

Then the minimal surface equation becomes

$$\Delta P + LP = 0.$$

If  $P$  is homogeneous of degree  $m$ , then  $LP$  is homogeneous of degree  $3m - 4$  and  $\Delta P$  is homogeneous of degree  $m - 2$ . Thus equation (1) is equivalent to  $3m - 3$  equations for  $P_m, \dots, P_1$  which is obtained by expanding (1) according to (2). The first 2 equations are:

$$(4) \quad LP_m = 0 \quad (\text{degree } 3m - 4)$$

$$(5) \quad DL(P_m)P_{m-1} = 0 \quad (\text{degree } 3m - 5)$$

where (5) means that  $P_{m-1}$  satisfies the linearized equation of  $L$  at  $P_m$  or explicitly

$$|\nabla P_m|^2 \Delta P_{m-1} + 2\Delta P_m \nabla P_m \cdot \nabla P_{m-1} - \sum_{i,j=1}^n (P_{m,x_i} P_{m,x_i} P_{m-1,x_i x_j} + 2P_{m,x_j} P_{m,x_i x_j} P_{m-1,x_i}) = 0.$$

Essentially we will only use (4) explicitly in this article cf. Remark 3.2. Note that  $P_m$  solving  $LP_m = 0$  is equivalent to the fact that every level set of  $P_m$  has mean curvature 0 at each regular point. This is not surprising in view of Theorem 1.1 (i). Solving  $LP_m = 0$  is the first step in solving the full minimal surface equation and it is unknown whether there is any nonlinear homogeneous polynomial solution except powers of linear functions. One nontrivial solution to  $Lu = 0$  is the function  $F$  constructed in section III of [5] which gives the first proof of area minimality of Simons' cone. However, by Theorem 1.6, this function is not a polynomial.

It is natural to ask whether there is any homogeneous polynomial  $P$  solving the full minimal surface equation, not just the 1-Laplace equation. From equations (1) and (4), we see that this is equivalent to  $\Delta P = 0$  and  $LP = 0$ . With an elementary result Proposition 4.1 of [26] (see also Lemma 3.4), we show that such polynomials cannot be a solution. Actually we have a stronger theorem.

**Theorem 1.2.** *Polynomials of the form*

$$P = P_m + P_{m-1} + \dots + P_{\lceil \frac{m}{3} \rceil + 1}$$

*with  $m \geq 2$  and  $P_m \neq 0$  cannot solve the minimal surface equation. Here  $\lceil x \rceil$  is the least integer greater than or equal to  $x$ .*

*In particular, there is no homogeneous polynomial of degree  $m \geq 2$  solving the minimal surface equation.*

We also have a slight variant of the theorem above which will be an important ingredient in Theorem 1.8.

**Theorem 1.3.** *We have the following:*

- (i) *Polynomials of the form  $P = P_m + P_1$  for  $m \geq 2$  cannot solve the minimal surface equation. In particular, there is no quadratic polynomial solution to the minimal surface equation.*
- (ii) *There is no cubic polynomial solution to the minimal surface equation.*

**1.4. Jacobi fields on level sets and lower bound of Green's function.** Since the level sets of  $P_m$  are all minimal, the function  $u = \frac{1}{|\nabla(P_0 Q_m^{1/k})|}$  satisfies the Jacobi field equation on regular part of  $\{P_m = t\}$  for all  $t \in \mathbb{R}$  cf. Lemma 3.1. Since a positive Jacobi field is bounded below by some multiple of Green's function, we get estimates on  $\deg P_0 + k^{-1} \deg Q_m$  if we have a lower bound of the Green's function. This is obtained in the following result proved in [14].

**Theorem 1.4** ([14]). *Let  $n \geq 4$  and  $\Sigma \subset \mathbb{R}^n$  be an area-minimizing boundary i.e.  $\Sigma = \partial[U]$  for some  $U \subset \mathbb{R}^n$  open and is area minimizing as a current. Then there exists  $C_1(n), C_2(n) > 0$  so that for the Green's function  $G(x, y)$  on  $\Sigma$ ,  $x, y \in \text{spt } \Sigma$*

$$C_1(n)|x - y|^{3-n} \geq G(x, y) \geq C_2(n)|x - y|^{3-n}$$

where  $|x|$  denotes the length of the vector  $x \in \mathbb{R}^n$ .

Using the above theorem, we have

**Theorem 1.5.** *If  $P_m = P_0^k Q_m$  satisfies Theorem 1.1 (i) and (ii), then  $\deg P_0 + k^{-1} \deg Q_m \leq n - 2$ .*

**1.5. "Non-existence" theorems.** We have two "non-existence" theorems about polynomial solutions. We call them "non-existence" since they show that the existence of polynomial solutions will imply the existence of some algebraic area minimizing cones that are not known to exist.

To state the first theorem, we recall that a minimal hypercone  $C \subset \mathbb{R}^n$  is called isoparametric if the link  $C \cap S^{n-1}$  has constant principal curvature in  $S^{n-1}$  cf. [7]. Every isoparametric cone is the zero locus of a homogeneous polynomial called Cartan-Münzner polynomial [8], [17]. Our first "non-existence" theorem is as follows.

**Theorem 1.6.** *If  $P_m = P_0^k Q_m$  satisfies Theorem 1.1 (i) and (ii), then  $P_0$  is not a Cartan-Münzner polynomial.*

*In particular we must have  $\deg P_0 \geq 3$  since all quadratic minimal cones are isoparametric.*

In view of Theorem 1.1, this implies the following:

**Corollary 1.7.** *Polynomial minimal graph cannot have  $C \times \mathbb{R}^l$  as tangent cone at  $\infty$  where  $C$  is an isoparametric minimal cone and  $l \geq 1$ .*

To the author's best knowledge, isoparametric cones and their products with  $\mathbb{R}^l$  are the only known examples of area minimizing hypercones. Therefore, we need to construct new algebraic area-minimizing cones in order to find nonlinear polynomial solutions.

For the second "non-existence" theorem, we recall the Bernstein's theorem which says that the smallest dimension in which nonlinear entire solution could exist is  $n = 8$ . For polynomial solutions on  $\mathbb{R}^8$ , we have the following "non-existence" theorem.

**Theorem 1.8.** *If there is a nonlinear polynomial solution to the minimal surface equation on  $\mathbb{R}^8$ , then there exists a cubic, strictly stable, area minimizing cone in  $\mathbb{R}^8$  which is not strictly minimizing.*

We can rephrase it as a Bernstein type theorem.

**Corollary 1.9.** *If  $P$  is a polynomial on  $\mathbb{R}^8$  solving the minimal surface equation and the tangent cone of graph  $P$  at  $\infty$  is strictly minimizing, then  $P$  is an affine function.*

The idea is to combine the growth estimates of Simon [21] and no cubic solutions from Theorem 1.3. The strictly minimizing property comes from the assumption of [21]. Such assumption appears in many results of minimal hypersurfaces. Note that there is no known example of area minimizing but not strictly minimizing hypercones except for  $\mathbb{R}^2 \subset \mathbb{R}^3$  cf. [16]. Moreover, in  $\mathbb{R}^8$ , the only known area minimizing cones are cones over  $S^3 \times S^3$  and  $S^2 \times S^4$ , both of which are isoparametric and strictly minimizing.

The outline of the rest of the paper is as follows. In section 2, we prove the structure theorem Theorem 1.1. In section 3, we prove Theorem 1.2 and 1.3. In section 4, we use lower bound for Green's function to prove Theorem 1.5. In section 5, we prove the two "nonexistence" theorems: Theorem 1.6 and 1.8.

#### ACKNOWLEDGEMENT

The author would like to express his sincere gratitude to his advisor, Richard Schoen, for constant support and guidance. He would also like to thank Connor Mooney for helpful discussions on this problem.

## 2. STRUCTURE OF POLYNOMIAL SOLUTIONS

We introduce the following notations and definitions, mostly about geometric measure theory cf. [11], [19]:

$|x|$ : the length of the vector  $x \in \mathbb{R}^n$ ;

$B_r(x)$ : Euclidean ball radius  $r$  in  $\mathbb{R}^n$  centered at  $x$  (write  $B_r$  if  $x = 0$ );

$U$ : open set in  $\mathbb{R}^n$ ;

$R_k^{loc}(U)$ : integer multiplicity  $k$ -rectifiable currents in  $U$ ;

$I_k^{loc}(U)$ :  $k$ -integral currents in  $U$ ;

$[M]$ : the current associated to the smooth manifold  $M$ ;

$[f < 0]$ : the current defined by the open set  $\{f < 0\}$ ;

$\|S\|$ : Radon measure associated to  $S \in R_k^{loc}(U)$ ;

$S \llcorner V$ ,  $\mu \llcorner V$ : restriction of the current  $S \in R_k^{loc}(U)$  or a Radon measure  $\mu$  to a set  $V \subset U$ ;

$\nabla u$ : Euclidean gradient of a function  $u$  at  $x$

$\nabla_S u(x)$ : tangential component of  $\nabla u$  on  $T_x S$ ;

$\Delta_S u$ : Laplace-Beltrami operator of a smooth hypersurface  $S$ ;

For later reference, we quote here the Nullstellensatz over  $\mathbb{R}$  which can be found in [4, Theorem 4.5.1].

**Lemma 2.1** (Real Nullstellensatz). *Let  $P \in \mathbb{R}[x_1, \dots, x_n]$  be an irreducible polynomial. The following are equivalent.*

- (i) The set  $Z(P) := \{P = 0\}$  contains a regular point, i.e. there exists  $x \in \{P = 0\}$  so that  $\nabla P(x) \neq 0$ .
- (ii) The polynomial  $P$  changes sign.
- (iii)  $(P) = I(Z(P))$  where  $(P)$  is the ideal generated by  $P$  in  $\mathbb{R}[x_1, \dots, x_n]$  and  $I(Z(P))$  is the real radical i.e. the ideal generated by polynomials that vanishes on  $Z(P)$ .
- (iv)  $\dim Z(P) = n - 1$ .

In particular, if  $P$  satisfies any of the conditions above and  $Q$  is a polynomial that vanishes in a neighborhood of a regular point in  $\{P = 0\}$ , then  $P|Q$ .

**Remark 2.2.** Throughout the article, the dimension of a zero locus  $V$  of some polynomials means one of the following, the algebraic dimension, the largest dimension of the manifolds in the decomposition of  $V$  into finite union of smooth manifolds, or the Hausdorff dimension. They are all equivalent in this setting.

**Remark 2.3.** If  $P$  is a polynomial that does not satisfy the conditions in Theorem 2.1, then it cannot change sign,  $Z(P)$  is of high codimension and  $\nabla P \equiv 0$  on  $\{P = 0\}$ . One such example is the function  $x_1^2 + x_2^2$  on  $\mathbb{R}^3$ .

We first investigate the structure of the level sets of a homogeneous polynomial.

Let  $P_m$  be a homogeneous polynomial of degree  $m$  in  $\mathbb{R}^n$ . We write the irreducible factorization of  $P_m$  as

$$(6) \quad P_m = c_0 p_1^{k_1} \cdots p_a^{k_a} \cdots p_{a+b}^{k_{a+b}} \cdots p_{a+b+c}^{k_{a+b+c}}$$

where

$$(7) \quad \begin{aligned} & c_0 \neq 0 \text{ is some constant,} \\ & k_i, a, b, c \geq 0 \text{ are integers} \\ & k_i \text{ is odd for } 1 \leq i \leq a, \\ & k_i \text{ is even for } a+1 \leq i \leq a+b \\ & p_i \text{ are irreducible polynomials for } 1 \leq i \leq a+b+c, \\ & \gcd(p_i, p_j) = 1, 1 \leq i \neq j \leq a+b+c, \\ & p_i \text{ changes sign for } 1 \leq i \leq a+b, \\ & p_i \geq 0 \text{ for } a+b+1 \leq i \leq a+b+c. \end{aligned}$$

**Lemma 2.4.** *We have*

- (i) For any  $t \neq 0$ , the set  $\{P_m = t\}$  is a smooth hypersurface in  $\mathbb{R}^n$  with  $\nabla P_m$  nonvanishing on  $\{P_m = t\}$ .
- (ii) For  $1 \leq i \leq a+b$ , the set  $\{p_i = 0\}$  is  $n-1$ -dimensional and for  $a+b+1 \leq j \leq a+b+c$  the set  $\{p_j = 0\}$  is of dimension  $\leq n-2$ .
- (iii) For  $1 \leq i \neq j \leq a+b$ , the set  $\{p_i = 0\}$  intersects  $\{p_j = 0\}$  along a set of dimension  $\leq n-2$ .

*Proof.* (i) Since  $P_m$  is homogeneous of degree  $m$ , by Euler's identity, we have, for any  $x \in \{P_m = t\}$

$$x \cdot \nabla P_m(x) = mP_m(x) = mt \neq 0.$$

Thus  $\nabla P_m(x) \neq 0$  and  $\{P_m = t\}$  is a smooth hypersurface.

(ii) This is direct consequence of Lemma 2.1 and (7).

(iii) By Lemma 2.1,  $\dim\{p_i = 0\} = n - 1$ . If for  $1 \leq j \leq a + b$ ,  $j \neq i$ ,  $p_j$  vanishes on an  $n - 1$ -dimensional subset of  $\{p_i = 0\}$ , then  $p_i | p_j$ . This is not possible since they are coprime by assumption (7). Thus for  $i \neq j$ , the set  $\{p_i = p_j = 0\}$  have dimension  $\leq n - 2$ .  $\square$

Now let  $P = P_m + \dots + P_1$  be a polynomial on  $\mathbb{R}^n$  and graph  $P$  denote the graph of  $P$  with orientation given by the unit normal

$$\frac{(\nabla P(x), -1)}{\sqrt{1 + |\nabla P(x)|^2}}.$$

We consider the limit of graph  $P$  under appropriate blow down and translation. Let  $t \in \mathbb{R}$  be a number and  $\lambda$  be a parameter that tends to 0. We denote the map  $x \mapsto \lambda x$  by  $\eta_\lambda$ . We consider the current  $\eta_{\lambda\#} \text{graph } P - t\lambda^{-m+1}$  obtained by scaling down graph  $P$  by  $\lambda$  and translating vertically by  $t\lambda^{-m+1}$ . Then we have  $\eta_{\lambda\#} \text{graph } P - t\lambda^{-m+1} e_{n+1} = \text{graph } G_t^{(\lambda)}$  where

$$(8) \quad \begin{aligned} G_t^{(\lambda)}(x) &= \lambda P(x/\lambda) - t\lambda^{-m+1} \\ &= \lambda^{-m+1}(P_m(x) - t) + \lambda^{-m+2}P_{m-1}(x) + \dots + P_1(x) \end{aligned}$$

with the unit normal

$$(9) \quad \nu_{\text{graph } G_t^{(\lambda)}} = \frac{(\nabla G_t^{(\lambda)}(x), -1)}{\sqrt{1 + |\nabla G_t^{(\lambda)}(x)|^2}}.$$

We also denote  $G_0^{(\lambda)}$  by  $P^{(\lambda)}$ . We have the following lemma.

**Lemma 2.5.** *Let  $T \in R_n^{\text{loc}}(\mathbb{R}^{n+1})$  be the limit of graph  $G_t^{(\lambda_j)}$  for some  $\lambda_j \rightarrow 0$ . Then we have  $\text{spt } T \subset \{P_m = t\} \times \mathbb{R}$ .*

*Proof.* A simple calculation shows

$$\lim_{\lambda \rightarrow 0} |G_t^{(\lambda)}(x)| = \begin{cases} |P_1(x)| & \text{if } P_m(x) - t = P_{m-1}(x) = \dots = P_2(x) = 0 \\ \infty & \text{otherwise.} \end{cases}$$

Thus for any  $x$  with  $P_m(x) \neq t$ , there is a neighborhood  $U$  of  $x$  in  $\mathbb{R}^n$  so that  $|G_t^{(\lambda)}| \rightarrow \infty$  uniformly in  $U$  as  $\lambda \rightarrow 0$ . Thus  $(U \times \mathbb{R}) \cap \text{spt } T = \emptyset$  and we have  $\text{spt } T \subset \{P_m = t\} \times \mathbb{R}$ .  $\square$

Now let  $t \neq 0$  and  $z_0 \in \{P_m = t\}$ . We consider the convergence of graph  $G_t^{(\lambda)}$  in a neighborhood of  $\{z_0\} \times \mathbb{R}$ .

**Lemma 2.6.** *Let  $P = P_m + \dots + P_1$  be a polynomial. Let  $t \neq 0$  and  $G_t^{(\lambda)}$  as in (8). Then for any  $z_0 \in \{P_m = t\}$ , there exists a neighborhood  $U$  of  $z_0$  in  $\mathbb{R}^n$  so that*

$$\lim_{\lambda \rightarrow 0} \left[ \text{graph } G_t^{(\lambda)} \right] \llcorner (U \times \mathbb{R}) = (\partial[P_m < t] \llcorner U) \times \mathbb{R}$$

where the convergence is both in  $I_n^{\text{loc}}(U \times \mathbb{R})$  as currents and in  $C^1$  normal graphs over  $\{P_m = t\} \times \mathbb{R}$ .

*Proof.* Since  $t \neq 0$ , by Lemma 2.4 (i), we define  $\nu(z) = \frac{\nabla P_m(z)}{|\nabla P_m(z)|}$  for all  $z \in \{P_m = t\}$ . By implicit function theorem, we can choose a neighborhood  $U$  of  $z_0$  so that the following are satisfied:

- (i) We can write  $U$  as  $U = \psi(W \times (-\sigma, \sigma))$  where  $W = U \cap \{P_m = t\}$  and  $\psi : W \times (-\sigma, \sigma) \rightarrow U$  is a diffeomorphism given by  $\psi(z, r) = z + r\nu(z)$ .
- (ii) There exists  $\delta > 0$  so that for all  $(z, r) \in W \times (-\sigma, \sigma)$  we have

$$\nabla P_m(z + r\nu(z)) \cdot \nu(z) \geq 2\delta > 0.$$

Using (8), we compute for  $\lambda$  sufficiently small

$$\begin{aligned} \frac{d}{dr} G_t^{(\lambda)}(z + r\nu(z)) &= \nabla G_t^{(\lambda)}(z + r\nu(z)) \cdot \nu(z) \\ &\geq \lambda^{-m+1} \nabla P_m(z + r\nu(z)) \cdot \nu(z) - O(\lambda^{-m+2}) \\ &\geq \delta \lambda^{-m+1} > 0. \end{aligned}$$

Thus  $r \mapsto P^{(\lambda)}(z + r\nu(z))$  is increasing for all  $-\sigma < r < \sigma$ .

On  $W$ , we have

$$|G_t^{(\lambda)}(z)| = |\lambda^{-m+2} P_{m-1}(z) + \cdots + \lambda^{-1} P_2(z) + P_1(z)| \leq C \lambda^{-m+2}.$$

Therefore, the interval  $[-\sigma \delta \lambda^{-m+1} + C \lambda^{-m+2}, \sigma \delta \lambda^{-m+1} - C \lambda^{-m+2}]$  is in the range of  $G_t^{(\lambda)}$  on  $U$  by fundamental theorem of calculus. Note that this interval converge to  $\mathbb{R}$  as  $\lambda \rightarrow 0$ . Let  $R > 0$  be fixed and  $(z, y) \in W \times (-R, R)$ . By mean value theorem, there exists  $\lambda_0(R) > 0$  so that for all  $0 < \lambda < \lambda_0$ , there is a unique  $-\sigma < \varphi^{(\lambda)}(z, y) < \sigma$  so that

$$(10) \quad (x, G_t^{(\lambda)}(x)) = (z + \varphi^{(\lambda)}(z, y)\nu(z), y).$$

This means that we can write  $[\text{graph } G_t^{(\lambda)}]_{\perp}(U \times \mathbb{R})$  as a normal graph of  $\varphi^{(\lambda)}$  over  $W \times (-R, R)$ . Using (10), we compute the gradient of  $\varphi^{(\lambda)}$  as

$$(11) \quad \frac{\partial \varphi^{(\lambda)}}{\partial y}(z, y) = \frac{1}{\nabla G_t^{(\lambda)}(x) \cdot \nu(z)} = O(\lambda^{m-1})$$

$$(12) \quad \frac{\partial \varphi^{(\lambda)}}{\partial z_j}(z, y) = \frac{-1}{\nabla G_t^{(\lambda)}(x) \cdot \nu(z)} \frac{\partial G_t^{(\lambda)}}{\partial x_i}(x) (e_{ij} + \varphi^{(\lambda)}(z, y) \frac{\partial \nu^i}{\partial z_j}(z))$$

$$(13) \quad = \frac{-1}{\nabla P_m(x) \cdot \nu(z)} \frac{\partial P_m}{\partial x_i}(x) (e_{ij} + \varphi^{(\lambda)}(z, y) \frac{\partial \nu^i}{\partial z_j}(z)) + O(\lambda)$$

where  $x = z + \varphi^{(\lambda)}(z, y)\nu(z)$  and  $(e_{ij})_{i=1, \dots, n, j=1, \dots, n-1}$  is the projection matrix from  $\mathbb{R}^n$  to  $T_z\{P_m = t\} = \nu(z)^{\perp}$ .

From (8), we have

$$P_m(x) - t = \lambda^{m-1}y - \lambda P_{m-1}(x) - \cdots - \lambda^{m-1}P_1(x).$$

Hence with  $x = z + \varphi^{(\lambda)}(z, y)\nu(z)$ , we have  $P_m(x) \rightarrow t$  as  $\lambda \rightarrow 0$ . Thus  $d(x, \{P_m = t\}) \rightarrow 0$  and hence

$$\lim_{\lambda \rightarrow 0} \varphi^{(\lambda)}(z, y) = 0.$$



Since  $(e_{ij})$  is the projection onto  $T_z\{P_m = t\}$  we have  $\sum_{i=1}^n e_{ij} \partial_{x_i} P_m(z) = 0$ . Thus the first term in (13) converges to 0. Hence we see that graph  $G_t^{(\lambda)}$  converge to  $(U \cap \{P_m = t\}) \times \mathbb{R}$  in  $C^1$ . We observe that the unit normal of graph  $G_t^{(t)}$  as in (9) converges to  $\nu$  as  $\lambda \rightarrow 0$ . Thus  $[\text{graph } G_t^{(\lambda)}] \llcorner (U \times \mathbb{R}) \rightarrow (\partial[P_m < t] \llcorner U) \times \mathbb{R}$  as currents.  $\square$

Now we consider the convergence of  $\partial[P_m < t]$  as  $t \rightarrow 0$  in a neighborhood of  $z_0$  where  $z_0$  is a regular point of  $\{P_m = 0\}$  i.e. for some  $1 \leq i \leq a + b$

$$(14) \quad \begin{aligned} p_i(z_0) &= 0, \quad \nabla p_i(z_0) \neq 0, \quad \text{and} \\ p_j(z_0) &\neq 0 \quad \text{for all } j \neq i. \end{aligned}$$

In view of the factorization (6), we denote the polynomial  $c_0 p_1^{k_1} \cdots \widehat{p_i^{k_i}} \cdots p_{a+b+c}^{k_{a+b+c}}$  by  $P_m/p_i^{k_i}$  for  $1 \leq i \leq a + b$ .

**Lemma 2.7.** *Let  $z_0$  be as in (14). Then there exists a neighborhood  $U$  of  $z_0$  in  $\mathbb{R}^n$  so that the following hold.*

(i) *If  $1 \leq i \leq a$ , then*

$$\lim_{t \rightarrow 0} (\partial[P_m < t]) \llcorner U = \text{sgn} \left( \left( P_m/p_i^{k_i} \right) (z_0) \right) (\partial[p_i < 0]) \llcorner U$$

where the convergence is both in  $I_{n-1}^{loc}(U)$  and in  $C^\infty$  as smooth manifolds.

In particular

$$\lim_{t \rightarrow 0} \|\partial[P_m < t]\| \llcorner U = \mathcal{H}^{n-1} \llcorner (U \cap \{p_i = 0\})$$

(ii) *If  $a + 1 \leq i \leq a + b$  then as currents in  $I_{n-1}^{loc}(U)$*

$$\lim_{t \rightarrow 0} \partial[P_m < t] \llcorner U = 0.$$

(a) *If  $\text{sgn}((P_m/p_i^{k_i})(z_0)) > 0$ , then*

$$\begin{aligned} \lim_{t \rightarrow 0^+} \|\partial[P_m < t]\| \llcorner U &= 2\mathcal{H}^{n-1} \llcorner (U \cap \{p_i = 0\}), \\ \lim_{t \rightarrow 0^-} \|\partial[P_m < t]\| \llcorner U &= 0. \end{aligned}$$

(b) *If  $\text{sgn}((P_m/p_i^{k_i})(z_0)) < 0$ , then*

$$\begin{aligned} \lim_{t \rightarrow 0^+} \|\partial[P_m < t]\| \llcorner U &= 0, \\ \lim_{t \rightarrow 0^-} \|\partial[P_m < t]\| \llcorner U &= 2\mathcal{H}^{n-1} \llcorner (U \cap \{p_i = 0\}). \end{aligned}$$

*Proof.* (i) By (7)  $k_i$  is odd for  $1 \leq i \leq a + b$ . We have  $\{P_m = t\} = \{w = t^{1/k_i}\}$  where

$$(15) \quad w = \left( P_m/p_i^{k_i} \right)^{1/k_i} p_i.$$

Since  $\nabla p_i(z_0) \neq 0$  and  $p_j(z_0) \neq 0$  for  $j \neq i$ , we have  $\nabla w(z_0) \neq 0$ . By implicit function theorem, there exist a small neighborhood  $U$  of  $z_0$ ,  $\delta > 0$  and  $f : U \cap \{w = 0\} \times (-\delta, \delta) \rightarrow U$  a diffeomorphism so that  $w(f(z, t)) = t$  and  $f(z, 0) = z$  for all

$z \in U \cap \{w = 0\}$ ,  $t \in (-\delta, \delta)$ . Therefore,  $U \cap \{w = t^{1/k_i}\} = f(U \cap \{w = 0\}, t^{1/k_i})$ . Since  $f(\cdot, t) \rightarrow \text{id}_{U \cap \{w=0\}}$  in  $C^\infty$  as  $t \rightarrow 0$ , we have

$$U \cap \{P_m = t\} = U \cap \{w = t^{1/k_i}\} \rightarrow U \cap \{w = 0\} = U \cap \{p_i = 0\} \quad \text{in } C^\infty \text{ as } t \rightarrow 0.$$

Note that on  $\{P_m = t\}$  for  $t \neq 0$ , we have  $\frac{\nabla P_m}{|\nabla P_m|} = \frac{\nabla w}{|\nabla w|}$ . Thus we have

$$\frac{\nabla P_m}{|\nabla P_m|} = \frac{\nabla w}{|\nabla w|} \rightarrow \text{sgn}\left(\left(P_m/p_i^{k_i}\right)(z_0)\right) \frac{\nabla p_i}{|\nabla p_i|} \text{ as } t \rightarrow 0.$$

This is the desired convergence.

(ii) We only prove (a). Item (b) follows from applying (a) to  $-P_m$ .

Since  $\text{sgn}((P_m/p_i^{k_i})(z_0)) > 0$  and  $k_i$  is even, we have  $U \cap \{P_m = t\} = \emptyset$  for  $t < 0$ . Thus the convergence for  $t \rightarrow 0^-$  is obtained.

For  $t \rightarrow 0^+$ , we consider  $w = \left(P_m/p_i^{k_i}\right)^{1/k_i} p_i$  on  $U$ . This is well defined in  $U$  since  $\text{sgn}((P_m/p_i^{k_i})(z_0)) > 0$ . Now we have

$$U \cap \{P_m = t\} = (U \cap \{w = t^{1/k_i}\}) \cup (U \cap \{w = -t^{1/k_i}\}).$$

By the same argument as in (i), we have

$$U \cap \{w = \pm c^{1/k_i}\} \rightarrow \pm U \cap \{p_i = 0\}$$

both in  $I_{n-1}^{loc}(U)$  and in  $C^\infty$  as manifolds. Thus we have  $\partial[P_m < t] \llcorner U \rightarrow 0$  and  $\|\partial[P_m < t]\| \llcorner U \rightarrow 2\mathcal{H}^n|_{U \cap \{p_i=0\}}$  as  $t \rightarrow 0^+$ .  $\square$

**Corollary 2.8.** *We have the following convergence*

$$\begin{aligned} \lim_{t \rightarrow 0} \partial[P_m < t] &= \sum_{i=1}^a \text{sgn}((P_m/p_i^{k_i})(z_0)) \partial[p_i < 0] = \partial[P_m < 0] \\ \lim_{t \rightarrow 0^+} \|\partial[P_m < t]\| &= \sum_{i=1}^a \mathcal{H}^{n-1} \llcorner \{p_i = 0\} + \sum_{i=a+1}^{a+b} 2\mathcal{H}^{n-1} \llcorner \{p_i = 0, P_m/p_i^{k_i} > 0\} \\ \lim_{t \rightarrow 0^-} \|\partial[P_m < t]\| &= \sum_{i=1}^a \mathcal{H}^{n-1} \llcorner \{p_i = 0\} + \sum_{i=a+1}^{a+b} 2\mathcal{H}^{n-1} \llcorner \{p_i = 0, P_m/p_i^{k_i} < 0\}. \end{aligned}$$

*Proof.* For any  $z \in \{P_m \neq 0\}$ , the set  $\{P_m = t\}$  have positive distance to  $z$  for small  $t$ . Thus if  $\partial[P_m < t] \rightarrow T$  for some  $T \in I_{n-1}^{loc}(\mathbb{R}^n)$  then  $\text{spt} T \subset \{P_m = 0\}$ . Let  $z_0 \in \{P_m = 0\}$ . By Lemma 2.4, the set of singular points of  $\{P_m = 0\}$  has dimension  $\leq n - 2$ . Thus we can consider the limits near a regular point of  $\{P_m = 0\}$ . All 3 identities follows from summing over  $1 \leq i \leq a + b$  in Lemma 2.7.

It remains to show the second equality in the first equation. Let  $z_0 \in \{p_i = 0\}$  be a regular point and  $U$  be a small neighborhood of  $x$ . We define  $w$  in  $U$  as in (15).

For  $1 \leq i \leq a$ , we have  $[P_m < 0] \llcorner U = \text{sgn}((P_m/p_i^{k_i})(z_0)) [w < 0] \llcorner U$ . Thus  $\partial[P_m < 0] \llcorner U = \text{sgn}((P_m/p_i^{k_i})(z_0)) \partial[p_i < 0] \llcorner U$ .

For  $a + 1 \leq i \leq a + b$ , we have  $[P_m < 0] \llcorner U = [w \neq 0] \llcorner U = [U]$ . Thus  $\partial[P_m < 0] = 0$ . This finishes the proof.  $\square$

**Theorem 2.9.** *Let  $P = P_m + P_{m-1} + \dots + P_2 + P_1$  with  $m \geq 2$  solve the minimal surface equation where  $P_i$  is homogeneous of degree  $i$ . Then we have:*

- (i) For every  $t \in \mathbb{R}$ , the current  $\partial[P_m < t]$  is area minimizing.  
(ii) There exist  $k \geq 1$  odd,  $P_0$  a nonlinear irreducible homogeneous polynomial that changes sign and  $Q_m$  an even degree homogeneous polynomial which is non-negative on  $\mathbb{R}^n$  and coprime with  $P_0$ , with  $\{Q_m = 0\}$  of dimension  $\leq n - 2$  so that

$$(16) \quad P_m = P_0^k Q_m.$$

- (iii) The unique tangent cone of graph  $P$  at  $\infty$  is  $\partial[P_0 < 0] \times \mathbb{R}$ .  
(iv) If  $k \geq 3$  in (16), then there exists  $2 \leq s \leq m - 1$  so that

$$P = P_0^k Q_m + P_0^{k_{m-1}} Q_{m-1} + \cdots + P_0^{k_{s+1}} Q_{s+1} + P_0 Q_s + P_{s-1} + \cdots + P_1$$

where  $P_0$  and  $Q_i$  are coprime for  $l \leq i \leq m$  and  $k_i \geq 2$  for  $s + 1 \leq i \leq m - 1$ .

*Proof.* (i) By Lemma 2.6 and Federer-Fleming compactness [19], we have for any  $t \neq 0$ ,  $\partial[P_m < t]$  is area minimizing. By Corollary 2.8, we have  $\partial[P_m < t] \rightarrow \partial[P_m < 0]$  as  $t \rightarrow 0$ . Thus by Federer-Fleming compactness [19] again,  $\partial[P_m < 0]$  is area minimizing.

(ii) By convergence of area-minimizing currents [19], we have

$$\|\partial[P_m < t]\| \rightarrow \|\partial[P_m < 0]\| = \sum_{i=1}^a \mathcal{H}^{n-1} \llcorner \{p_i = 0\}.$$

Thus  $a \geq 1$  in (6). Moreover, in view of Corollary 2.8, we see that  $b = 0$  in (6). In other words, there is no factor  $p_i^{k_i}$  where  $p_i$  changes sign and  $k_i$  is even.

We claim that  $a = 1$ . Indeed, suppose  $a \geq 2$  for contradiction. We can assume  $c_0 > 0$  in (6) by adjusting  $p_1$  to be  $-p_1$  if necessary. We have

$$\{P_m < 0\} = \{p_1 < 0, p_2 \cdots p_a > 0\} \cup \{p_1 > 0, p_2 \cdots p_a < 0\}.$$

Observe that the two open sets on the right hand side are disjoint. This contradicts Theorem 1 of [6], which asserts that  $\{P_m < 0\}$  is connected.

Thus  $a = 1$ . We let  $P_0 = p_1$ ,  $k = k_1$  and  $Q_m = c_0 p_{a+b+1}^{k_{a+b+1}} \cdots p_{a+b+c}^{k_{a+b+c}}$ . Then the conclusion follows.

(iii) Let  $T$  be a tangent cone at  $\infty$  of graph  $P$ . Recall that we wrote  $\eta_{\lambda \#} \text{graph } P = \text{graph } P^{(\lambda)}$  in (8). By Lemma 2.5,  $\text{spt } T \subset \{P_m = 0\} \times \mathbb{R}$ . By [20, §2] we know that  $T = \partial[U] \times \mathbb{R}$  for some connected open set  $U \subset \mathbb{R}^n$ . Hence we have  $\text{spt } \partial[U] \subset \{P_m = 0\}$ . By (6), we have

$$\{P_m = 0\} = \{P_0 = 0\} \cup \left( \bigcup_{i=2}^{c+1} \{p_i = 0\} \right).$$

By Lemma 2.1, the set  $\{p_i = 0\}$  have dimension  $\leq n - 2$  for  $2 \leq i \leq c + 1$ . Thus  $\partial[P_m < 0] = \partial[P_0 < 0]$  and  $\text{spt } \partial[U] \subset \{P_0 = 0\}$ . If we take  $\phi \in C_c^\infty(\{P_0 < 0\})$  then  $\partial[U](\phi) = 0$ . Thus  $\partial[U] = 0$  in  $I_{n-1}^{loc}(\{P_0 < 0\})$ . Similarly  $\partial[U] = 0$  in  $I_{n-1}^{loc}(\{P_0 > 0\})$ . Thus by constancy theorem,  $[U]$  is constant on both  $\{P_0 < 0\}$  and  $\{P_0 > 0\}$ . Hence either  $[U] = [P_0 < 0]$  or  $[U] = [P_0 > 0]$ . Since we take the normal vector of graph  $P^{(\lambda)}$  as in (9), for  $x \notin \{P_0 = 0\}$ ,  $\nu_{\text{graph } P^{(\lambda)}}$  is close to  $\frac{\nabla P_0}{|\nabla P_0|}$  for  $\lambda$  small. Thus we have  $U \cap \{P_0 < 0\} \neq \emptyset$ . Therefore  $[U] = [P_0 < 0]$ .

(iv) Suppose  $k \geq 3$  in (16). Then  $\nabla P_m = 0$  on  $\{P_0 = 0\}$ . Since  $[\text{graph } P^{(\lambda)}] \rightarrow \partial[P_0 < 0] \times \mathbb{R}$  as  $\lambda \rightarrow 0$ , by Allard regularity, for any regular point  $(z_0, y_0) \in \{P_0 =$

$0\} \times \mathbb{R}$ , there is a neighborhood  $U$  of  $(z, y)$  so that  $\text{graph } P^{(\lambda)} \cap U$  can be written as normal graph of  $\varphi^{(\lambda)}$  over  $\{P_0 = 0\} \times \mathbb{R} \cap U$  and  $\varphi^{(\lambda)} \rightarrow 0$  smoothly. In particular, with  $x = z + \varphi^{(\lambda)}(z, y)\nu(z)$ , we have

$$\frac{(\nabla P^{(\lambda)}(x), -1)}{\sqrt{1 + |\nabla P^{(\lambda)}(x)|^2}} = \nu_{\text{graph } P^{(\lambda)}}(x) \rightarrow \nu(z) = \frac{(\nabla P_0(z), 0)}{|\nabla P_0(z)|} \text{ as } \lambda \rightarrow 0.$$

Since  $\frac{1}{\sqrt{1 + |\nabla P^{(\lambda)}|^2}} \rightarrow 0$  and  $\nabla P_m(z) = 0$ , there exists  $2 \leq s \leq m - 1$  so that

$$(17) \quad \nabla P_m(z) = \nabla P_{m-1}(z) = \cdots = \nabla P_{s+1}(z) = 0 \text{ and } \nabla P_s(z) \neq 0.$$

Since for every regular point  $z \in \{P_0 = 0\}$ , there exists  $s$  satisfying (17), we may choose the maximal  $s$  so that (17) holds on an open set  $V$  of  $\{P_0 = 0\}$ . By Lemma 2.1, we have  $P_0 | \nabla P_i$  for  $s+1 \leq i \leq m-1$ . By Euler's formula  $iP_i(x) = x \cdot \nabla P_i(x)$ , we have  $P_0 | P_i$ . Thus we can write  $P_i = P_0 \tilde{Q}_i$ . Since

$$0 = \nabla P_i = \tilde{Q}_i \nabla P_0 + P_0 \nabla \tilde{Q}_i = \tilde{Q}_i \nabla P_0 \quad \text{in } V,$$

we see that  $\tilde{Q}_i$  vanishes in  $V$ . Thus  $P_0 | \tilde{Q}_i$  and hence  $P_0^2 | P_i$ .

By (17), we have as  $\lambda \rightarrow 0$

$$\frac{(\nabla P^{(\lambda)}(x), -1)}{\sqrt{1 + |\nabla P^{(\lambda)}(x)|^2}} \rightarrow \frac{(\nabla P_s(z), 0)}{|\nabla P_s(z)|}.$$

Hence for all  $z \in V$ , we have

$$\frac{\nabla P_s(z)}{|\nabla P_s(z)|} = \frac{\nabla P_0(z)}{|\nabla P_0(z)|}.$$

By Euler's identity,

$$sP_s(z) = z \cdot \nabla P_s(z) = z \cdot \nabla P_0(z) \frac{|\nabla P_s(z)|}{|\nabla P_0(z)|} = \deg P_0 \cdot P_0(z) \frac{|\nabla P_s(z)|}{|\nabla P_0(z)|} = 0.$$

Thus we have  $P_s(z) = 0$  in an open set  $V$  of  $C$ . Thus by Lemma 2.1,  $P_0 | P_s$ . Since  $\nabla P_s \neq 0$  on  $V$ , we have  $P_0^2 \nmid P_s$ . This finishes the proof.  $\square$

### 3. NO HOMOGENEOUS POLYNOMIAL SOLUTION TO THE MINIMAL SURFACE EQUATION

In this section, we are going to show that polynomial solutions to the minimal surface equation must contain terms of both high and low degree. First we need a preparation lemma.

**Lemma 3.1.** *Let  $P = P_m + \cdots + P_1$  be a polynomial solution to the minimal surface equation. Then*

(i)  $P_m$  satisfies

$$LP_m = |\nabla P|^2 \Delta P - \sum_{i,j=1}^n P_{x_i} P_{x_j} P_{x_i x_j} = 0.$$

(ii) Let  $P_m = P_0^k Q_m$  be as in Theorem 2.9 (ii). Then  $u = \frac{1}{|\nabla(P_0 Q_m^{1/k})|}$  is the velocity of the family of minimal hypersurfaces  $(\{P_0 Q_m^{1/k} = t\})_{t \in \mathbb{R}}$ . Hence  $u$  satisfies the Jacobi field equation on regular part of  $\{P_m = t\}$  for all  $t \in \mathbb{R}$ .

(iii)  $P_{m-1}$  satisfies

$$(18) \quad \begin{aligned} DL(P_m)P_{m-1} &= |\nabla P_m|^2 \Delta P_{m-1} + 2\Delta P_m \nabla P_m \cdot \nabla P_{m-1} \\ &- \sum_{i,j=1}^n (P_{m,x_i} P_{m,x_i} P_{m-1,x_i x_j} + 2P_{m,x_j} P_{m,x_i x_j} P_{m-1,x_i}) = 0. \end{aligned}$$

(iv) The function  $v = -\frac{P_{m-1}}{|\nabla P_m|}$  is a Jacobi field on  $\{P_m = t\}$  with  $t \neq 0$ . If  $k = 1$  in (16), then  $v$  is also a Jacobi field on regular part of  $\{P_m = 0\}$ .

(v) Suppose  $t \neq 0$ . The function  $v$  viewed as a function on  $\{P_m = t\} \times \mathbb{R}$  is the velocity of the family of minimal hypersurfaces  $(\text{graph } G_t^{(\lambda)})_{\lambda \in [0, \epsilon]}$  at  $\lambda = 0$  where  $G_t^{(\lambda)}$  is defined as in (8).

**Remark 3.2.** If we have  $P_m$  solving  $LP_m = 0$ , then  $\partial_{x_i} P_m$  for  $1 \leq i \leq n$  are solutions to (18). This corresponds to translations of the level sets in  $\mathbb{R}^n$ . This indicates that (18) may not give too much information on the polynomial.

*Proof.* (i) This is the degree  $3m - 4$  term in the expansion of (1) according to (2).

(ii) By (i), Lemma 2.4 and Theorem 2.9 (ii) for any  $t \in \mathbb{R}$ , the level set  $\{P_0 Q_m^{1/k} = t\} = \{P_m = t^k\}$  is minimal. We take a  $\frac{1}{k}$  power of  $P_m$  since  $\nabla(P_0 Q_m^{1/k}) = Q_m^{1/k} \nabla P_0 \neq 0$  on regular part of  $\{P_0 = 0\}$ . Now the velocity of the family  $(\{P_0 Q_m^{1/k} = t\})_{t \in \mathbb{R}}$  is  $u = \frac{1}{|\nabla(P_0 Q_m^{1/k})|}$  and hence it satisfies the Jacobi equation on regular part of  $\{P_m = t\}$ .

(iii) This is the degree  $3m - 5$  term in the expansion of (1) according to (2).

(iv) For  $t \neq 0$ , Suppose  $\gamma(s)$  is a curve in  $\mathbb{R}^n$  so that  $\gamma(0) = x \in \{P_m = t\}$ ,  $\dot{\gamma}(0) = v(x) \frac{\nabla P_m(x)}{|\nabla P_m(x)|}$  and  $(P_m + sP_{m-1})(\gamma(s)) = 0$  for all  $s \in (-\epsilon, \epsilon)$ . Then

$$0 = \left. \frac{d}{ds} \right|_{s=0} (P_m + sP_{m-1})(\gamma(s)) = P_{m-1}(x) + \nabla P_m \cdot \left( v(x) \frac{\nabla P_m(x)}{|\nabla P_m(x)|} \right).$$

Thus  $v(x) = -\frac{P_{m-1}}{|\nabla P_m|}$  is the velocity field of the hypersurfaces  $(\{P_m + sP_{m-1} = t\})_{s \in (-\epsilon, \epsilon)}$  at  $s = 0$ . Equation (18) shows that the level set of  $P_m + sP_{m-1}$  has 0 mean curvature up to the first order in  $s$ , thus  $v$  is a Jacobi field on  $\{P_m = t\}$  for  $t \neq 0$ .

If  $k = 1$ , then  $\nabla P_m \neq 0$  on regular part of  $\{P_m = 0\}$ . The same argument as above shows that  $v$  is a Jacobi field on  $\{P_m = 0\}$ .

(v) Let  $t \neq 0$ . We write  $\text{graph } G_t^{(\lambda)}$  as a normal graph of  $\varphi^{(\lambda)}$  over  $\{P_m = t\} \times \mathbb{R}$  as in (10). Using the expression of  $G_t^{(\lambda)}$  in (8) and writing  $x = z + \varphi^{(\lambda)}(z, y)\nu(z)$ , we have

$$\begin{aligned} 0 &= \frac{d}{d\lambda} G_t^{(\lambda)} \left( z + \varphi^{(\lambda)}(z, y)\nu(z) \right) = \frac{\partial G_t^{(\lambda)}}{\partial \lambda}(x) + \nabla G_t^{(\lambda)}(x) \cdot \nu(z) \frac{\partial \varphi^{(\lambda)}}{\partial \lambda}(z) \\ &= \lambda^{-m}(1-m)(P_m(x) - t) + \lambda^{1-m} \left( (2-m)P_{m-1}(x) + \nabla P_m(x) \cdot \nu(z) \frac{\partial \varphi^{(\lambda)}}{\partial \lambda}(z) \right) \\ &\quad + O(\lambda^{2-m}) \\ &= \lambda^{1-m}(2-m) \left( P_{m-1}(x) + |\nabla P_m(z)| \frac{\partial \varphi^{(\lambda)}}{\partial \lambda}(z) \right) + O(\lambda^{2-m}) \end{aligned}$$

where in the last step, since  $P_m(x) - t \rightarrow 0$  as  $\lambda \rightarrow 0$ , we expand the first term as

$$\lambda^{-m}(1-m)(P_m(x) - t) = \lambda^{1-m}(1-m)|\nabla P_m(z)| \frac{\partial \varphi^{(\lambda)}}{\partial \lambda}(z) + O(\lambda^{2-m}).$$

Taking  $\lambda \rightarrow 0$ , we get (v).  $\square$

**Theorem 3.3.** *Polynomials of the form*

$$P = P_m + P_{m-1} + \cdots + P_{\lceil \frac{m}{3} \rceil + 1}$$

with  $m \geq 2$  and  $P_m \neq 0$  cannot solve the minimal surface equation. Here  $\lceil x \rceil$  is the least integer greater than or equal to  $x$ .

In particular, there is no homogeneous polynomial of degree  $m \geq 2$  solving the minimal surface equation.

*Proof.* Suppose there is such a polynomial solving the minimal surface equation. By Lemma 3.1 (i), we have  $LP_m = 0$ . The lowest degree term in  $LP$  is  $LP_{\lceil \frac{m}{3} \rceil + 1} = 0$  which has degree  $3\lceil \frac{m}{3} \rceil - 1 > m - 2 = \deg \Delta P_m$ . Thus the  $m - 2$  degree term in  $\Delta P + LP$  is  $\Delta P_m$  and we have  $\Delta P_m = 0$ . Combining it with  $LP_m = 0$ , we have a nonzero homogeneous polynomial  $P_m$  satisfying  $\Delta P_m = 0$  and the  $\infty$ -Laplacian equation

$$\sum_{i,j=1}^n P_{x_i} P_{x_j} P_{x_i x_j} = 0.$$

This is a contradiction in view of the next lemma.  $\square$

**Lemma 3.4** ([26, Proposition 4.1]). *Let  $P$  be a smooth function on  $\mathbb{R}^n \setminus \{0\}$  which is homogeneous of degree  $m \notin \{0, 1\}$ , and satisfies  $\Delta P = 0$  and  $\sum_{i,j=1}^n P_{x_i x_j} P_{x_i} P_{x_j} = 0$ . Then  $P = 0$ .*

**Remark 3.5.** Note that  $m$  is not required to be an integer. Hence the lemma actually shows that there is no homogeneous solution to the minimal surface equation on  $\mathbb{R}^n \setminus \{0\}$ .

We write down the proof of the lemma for the sake of convenience.

*Proof.* We can rewrite the  $\infty$ -Laplacian equation as

$$(19) \quad \frac{1}{2} \langle \nabla P, \nabla |\nabla P|^2 \rangle = 0.$$

Let  $x \in S^{n-1}$  so that  $P(x) = \max_{y \in S^{n-1}} P(y)$ . By the method of Lagrange multiplier, we have  $\nabla P(x) = \lambda x$ . Evaluating (19) at  $x$ , and using Euler's formula for the homogeneous function  $|\nabla P|^2$ , we have

$$0 = \frac{1}{2} \langle \lambda x, \nabla |\nabla P|^2(x) \rangle = \lambda(m-1) |\nabla P(x)|^2 = \lambda^3(m-1).$$

Since  $m \neq 1$ , we have  $\lambda = 0$  and thus  $\nabla P(x) = 0$ .

On the other hand, using Euler's formula for the homogeneous function  $P$ , we have

$$mP(x) = \langle x, \nabla P(x) \rangle = 0.$$

Hence  $\max_{y \in S^{n-1}} P(y) = 0$ . Similarly we have  $\min_{y \in S^{n-1}} P(y) = 0$  and thus  $P = 0$ .  $\square$

The following theorem is a slight variant of Theorem 3.3. This is important in later applications cf. Theorem 5.6.

**Theorem 3.6.** *We have*

- (i) *Polynomials of the form  $P = P_m + P_1$  for  $m \geq 2$  cannot solve the minimal surface equation. In particular, there is no quadratic polynomial solution to the minimal surface equation.*
- (ii) *There is no cubic polynomial solution to the minimal surface equation.*

*Proof.* (i) After rotating the coordinate in  $\mathbb{R}^n$ , we may assume that  $P_1(x) = ax_1$  with  $a \neq 0$ . Plugging  $P_1 = ax_1$  in (1), and considering degree  $m - 2$  and  $2m - 3$  terms, we obtain

$$(20) \quad (1 + a^2)\Delta P_m - a^2 P_{m,x_1 x_1} = 0$$

$$(21) \quad P_{m,x_1} \Delta P_m - \sum_{i=1}^n P_{m,x_i x_1} P_{m,x_i} = 0.$$

Eliminating  $\Delta P_m$  in (20) and (21), we get

$$\frac{a^2}{1 + a^2} P_{m,x_1 x_1} P_{m,x_1} - \sum_{i=1}^n P_{m,x_i x_1} P_{m,x_i} = 0$$

i.e.

$$\left( \frac{1}{1 + a^2} P_{m,x_1}^2 + P_{m,x_2}^2 + \cdots + P_{m,x_n}^2 \right)_{x_1} = 0.$$

Thus we have  $\frac{1}{1+a^2} P_{m,x_1}^2 + \sum_{i=2}^n P_{m,x_i}^2$  is independent of  $x_1$ .

We arrange the terms in  $P_m$  according to the degree in  $x_1$ :

$$P_m(x) = c_0 x_1^m + c_1(x') x_1^{m-1} + \cdots + c_m(x')$$

where  $x' = (x_2, \dots, x_n)$  and  $c_i$  is a polynomial in  $x'$  of degree  $i$ .

The term in  $\frac{1}{1+a^2} P_{m,x_1}^2 + \sum_{i=2}^n P_{m,x_i}^2$  with highest  $x_1$  degree is

$$\frac{1}{1 + a^2} c_0^2 m^2 x_1^{2m-2} + \sum_{i=2}^n c_{1,x_i}^2 x_1^{2m-2}.$$

Thus we must have  $c_0 = c_1 = 0$ .

Assume that  $j$  is the smallest index such that  $c_j \neq 0$ , then we have  $c_0 = \cdots = c_{j-1} = 0$ . The term in  $\frac{1}{1+a^2} P_{m,x_1}^2 + \sum_{i=2}^n P_{m,x_i}^2$  with highest  $x_1$  degree is

$$\sum_{i=2}^n c_{j,x_i} (x')^2 x_1^{2m-2j}.$$

Since  $c_j \neq 0$ , in order for  $\frac{1}{1+a^2} P_{m,x_1}^2 + \sum_{i=2}^n P_{m,x_i}^2$  to be independent of  $x_1$ , we must have  $m = j$ . Hence  $c_1 = \cdots = c_{m-1} = 0$  and  $P_m$  is independent of  $x_1$ .

By equation (20), we have  $\Delta P_m = 0$ . Since  $LP_m = 0$ , Lemma 3.4 implies that  $P_m = 0$  i.e.  $P = P_1$ .

(ii) Let  $P = P_3 + P_2 + P_1$  be a solution to the minimal surface equation with  $P_3 \neq 0$ .

By Theorem 2.9 (ii), we know that  $P_3 = P_0^k Q_3$  where  $Q_3 \geq 0$  has even degree. If  $\deg Q_3 = 2$ , then  $\deg P_0 = 1$  and  $k = 1$ . Thus by Theorem 2.9 (ii), the tangent cone at

infinity of graph  $P$  is  $\partial[P_0 < 0] \times \mathbb{R}$  which is a hyperplane. This implies that graph  $P$  is a hyperplane, a contradiction. If  $\deg Q_3 = 0$  i.e.  $Q_3 = 1$  then  $P_3 = P_0^k$ . If  $k = 3$ , then  $\deg P_0 = 1$  and  $\partial[P_0 < 0] \times \mathbb{R}$  is again a hyperplane which is not possible.

Thus we know that  $k = 1$  and  $P_3$  is irreducible. We have  $LP_3 = 0$ . By Theorem 1 of [25], we know that any irreducible cubic polynomial solving  $LP_3 = \lambda|x|^2P_3$  for some  $\lambda \in \mathbb{R}$  is harmonic. Thus  $\Delta P_3 = 0$ . Hence  $P_3 = 0$  by Lemma 3.4, a contradiction.  $\square$

#### 4. LOWER BOUNDS FOR GREEN'S FUNCTION AND ESTIMATING THE DEGREE OF $\deg P_0$

In this section we prove the following theorem using Theorem 1.4 proved in [14].

**Theorem 4.1.** *If  $P_m = P_0^k Q_m$  satisfies Theorem 2.9 (i) and (ii), then  $\deg P_0 + k^{-1} \deg Q_m \leq n - 2$ .*

*Proof.* Let  $u$  be the function  $\frac{1}{|\nabla(P_0 Q_m^{1/k})|}$ . By Lemma 3.1,  $u$  is a positive Jacobi field on  $\Sigma = \{P_m = 1\}$  i.e.

$$\Delta_\Sigma u + |A_\Sigma|^2 u = 0.$$

In particular,  $\Delta_\Sigma u \leq 0$  on  $\Sigma$ .

Let  $x \in \text{reg } \Sigma$  and  $r_0 > 0$  be fixed. By [14, Theorem 3.2], for any  $R > r_0$ , there exists  $G_R(x, y)$  the Dirichlet Green's function on  $\Sigma \cap B_R(x)$  satisfying  $G_R(x, y) \leq C_1(n)|x - y|^{3-n}$ . Thus have

$$\sup_{y \in \Sigma \cap \partial B_{r_0}(x)} G_R(x, y) \leq C_1 r_0^{3-n} \quad \forall R > r_0.$$

Since  $G_R(x, y) = 0$  for any  $y \in \partial B_R(x)$ , by weak maximum principle on  $\Sigma \cap B_R(x) \setminus B_{r_0}(x)$ , we have for any  $y \in \Sigma \cap B_R(x) \setminus B_{r_0}(x)$

$$(22) \quad u(y) \geq \left( \inf_{\Sigma \cap \partial B_{r_0}(x)} u \right) C_1^{-1} r_0^{n-3} G_R(x, y).$$

Letting  $R \rightarrow \infty$  in (22) and applying Theorem 1.4, we have for all  $y \in \Sigma$ ,  $|x - y| > r_0$

$$(23) \quad u(y) \geq \left( \inf_{\Sigma \cap \partial B_{r_0}(x)} u \right) C_1^{-1} r_0^{n-3} G(x, y) \geq c_1 |x - y|^{3-n}$$

where  $c_1 = \left( \inf_{\Sigma \cap \partial B_{r_0}(x)} u \right) C_2 C_1^{-1} r_0^{n-3} > 0$  is independent of  $y$ .

Now we want to transplant the lower bound of  $u$  on  $\Sigma$  to a lower bound of  $u$  on  $\{P_0 = 0\}$ . By Lemma 3.1,  $u$  is the velocity of the family  $(\{P_0 Q_m^{1/k} = t\})$ . Then there exists  $R_0 > 0$ , a conical neighborhood  $V$  in  $\mathbb{R}^n$  of a regular point of  $\{P_0 = 0\}$  and a function  $\phi$  over  $\{P_0 = 0\} \cap V \setminus B_{R_0}$  so that graph  $\phi \subset \{P_m = 1\}$  and  $\{P_m = 1\} \cap V \setminus \text{graph } \phi$  is compact. Moreover for some  $\alpha > 0$  and any  $|z| \geq R_0$ , we have

$$\phi(z) = c_2 |z|^{-\gamma} + O(|z|^{-\gamma-\alpha})$$

where  $\gamma = \deg P_0 + k^{-1} \deg Q_m - 1$  and  $c_2 > 0$ . Since  $\deg P_0 \geq 2$ , we have  $\phi \rightarrow 0$  as  $|z| \rightarrow \infty$ . Then by (23), we have

$$u(z + \phi(z)\nu(z)) \geq c_1 |z + \phi(z)\nu(z) - x|^{3-n} \geq c_3 |z|^{3-n}$$



for some  $c_3 > 0$  and all  $z \in \{P_0 = 0\} \cap V$  with  $|z|$  large. On the other hand, for  $z \in \{P_0 = 0\} \cap V$  with  $|z|$  large

$$|u(z + \phi(z)\nu(z)) - u(z)| \leq \sup_{t \in [0,1]} \|\nabla u(z + t\phi(z)\nu(z))\| |\phi(z)| \leq c_4 |z|^\gamma 2c_2 |z|^{-\gamma} = c_5.$$

Hence combining the above, we have for  $z \in \{P_0 = 0\} \cap V$  with  $|z|$  large

$$u(z) \geq c_2 |z|^{3-n} - c_5.$$

Letting  $|z| \rightarrow \infty$ , we obtain  $\deg P_0 + k^{-1} \deg Q_m \leq n - 2$ .  $\square$

## 5. "NON-EXISTENCE" THEOREMS

We first introduce some concepts related to minimal cones cf. [15].

**Definition 5.1.** A minimal hypercone  $C \subset \mathbb{R}^n$  is called *regular* if  $\text{sing } C \subset \{0\}$ . Let  $\Sigma = C \cap S^{n-1}$  and  $\lambda_1$  be the first eigenvalue of  $\Delta_\Sigma + |A_\Sigma|^2$ . The minimal cone  $C$  is called *stable* (*strictly stable*) if  $(n-3)^2 + 4\lambda_1 \geq 0$  ( $> 0$ ). There are 2 independent positive Jacobi fields on a stable cone  $C$ . If  $C$  is strictly stable, they are  $|x|^{-\gamma_\pm} \phi_1$ . If  $C$  is not strictly stable, they are  $|x|^{-(n-3)/2} \phi_1$  and  $|x|^{-(n-3)/2} (\log |x|) \phi_1$ . Here  $\phi_1$  is the first eigenfunction of  $\Delta_\Sigma + |A_\Sigma|^2$  on  $\Sigma$  and

$$(24) \quad \gamma_\pm = \frac{n-3 \pm \sqrt{(n-3)^2 + 4\lambda_1}}{2}.$$

By [15, Theorem 2.1] if  $C$  is regular area minimizing cone, then  $\mathbb{R}^n \setminus C$  consists of two components  $E_\pm$  and each  $E_\pm$  is foliated uniquely by homotheties of smooth minimal hypersurface  $S_\pm \subset E_\pm$ . There exists  $R(C) > 0, \alpha > 0$  so that  $S_\pm \setminus B_R(0) = \text{graph}_C(v_\pm)$  and as  $|x| \rightarrow \infty$

$$v_\pm(x) = w_\pm(x) + O(r^{-\gamma_+ - \alpha})$$

where  $\pm w_\pm > 0$  and  $\Delta_C w_\pm + |A_C|^2 w_\pm = 0$ .

By [15, Theorem 3.1], a regular area minimizing cone  $C$  is called *strictly minimizing* if  $w$  is the positive Jacobi field with slower decay rate near infinity.

**Definition 5.2.** A minimal hypercone  $C \subset \mathbb{R}^n$  is called *isoparametric* if the link  $C \cap S^{n-1}$  has constant principal curvature in  $S^{n-1}$ . Every isoparametric cone is the zero locus of a polynomial called Cartan-Münzner polynomial [8], [17].

**Lemma 5.3** ([1],[8],[17],[27]). *Let  $P_0$  be a Cartan-Münzner polynomial on  $\mathbb{R}^l$ . The following are all possibilities for the degree  $\deg P_0$ , number of variables  $l$  of  $P_0$  and the norm squared of the second fundamental form  $|A|^2$  of the link of  $\{P_0 = 0\}$ .*

- (i)  $\deg P_0 = 1, l \geq 2, |A|^2 = 0,$
- (ii)  $\deg P_0 = 2, l \geq 4, |A|^2 = l - 2,$
- (iii)  $\deg P_0 = 3, l = 5, 8, 14, 26, |A|^2 = 2(l - 2),$
- (iv)  $\deg P_0 = 4, l \geq 6$  is even,  $|A|^2 = 3(l - 2),$
- (v)  $\deg P_0 = 6, l = 8, 14, |A|^2 = 5(l - 2).$

Moreover,  $\{P_0 = 0\}$  is area minimizing if and only if  $l \geq 4 \deg P_0$  and  $\Sigma$  is not  $S^1 \times S^5$  or  $SO(8) \times SO(2)/SO(6) \times \mathbb{Z}_2$ .

*Proof.* By [17],  $\deg P_0$  is equal to the number of distinct principal curvatures of the link and it can only be 1, 2, 3, 4, 6. By [8], for  $\deg P_0 \leq 3$ , the isoparametric hypersurfaces are homogeneous and  $l$  can only take the value we mention. By [17], if the  $i$ -th principal curvature has multiplicity  $m_i$ , then  $m_i = m_{i+2}$  with index understood mod  $\deg P_0$ . Thus in case (iv), we have  $l = 2(m_1 + m_2) + 2 \geq 6$  is even. For  $\deg P_0 = 6$ , by [1],  $m_1 = m_2$  can only be 1 or 2 and thus  $l = 3(m_1 + m_2) + 2 = 8$  or 14. The value of  $|A|^2$  follows from exact values of principal curvatures of minimal isoparametric cones [17]. Calculation can be found in [24] in equations (48) (49) (50) and the formula below (53). The area minimizing property is a result of [27].  $\square$

**Theorem 5.4.** *If  $P_m = P_0^k Q_m$  satisfies Theorem 2.9 (i) and (ii), then  $P_0$  is not a Cartan-Münzner polynomial.*

*In particular  $\deg P_0 \geq 3$  since all quadratic minimal cones are isoparametric.*

In view of Theorem 2.9(iii) we have the following corollary.

**Corollary 5.5.** *If a polynomial  $P$  solves the minimal surface equation, then the tangent cone of graph  $P$  at  $\infty$  is not  $C \times \mathbb{R}^l$  where  $l \geq 1$  and  $C$  is any isoparametric cone.*

*Proof.* Suppose  $P_0$  is a Cartan-Münzner polynomial of degree  $\geq 2$ . Permuting the  $x_i$ 's if necessary, we assume that  $x_1, \dots, x_l$  are all variables involved in  $P_0$  for some  $1 \leq l \leq n$ .

We claim that  $P_m = P_0^k Q_m$  is independent of  $x_{l+1}, \dots, x_n$ . Indeed, by [27], the isoparametric cone  $\{P_0 = 0\}$  is strictly stable and strictly minimizing. By Lemma 2.4 and Theorem 2.9, for  $t \neq 0$ ,  $\{P_m = t\}$  is a smooth minimal hypersurface lying on one side of the cone  $\{P_m = 0\} = \{P_0 = 0\} \times \mathbb{R}^{n-l}$ . Moreover, the blow-down of  $\{P_m = t\}$  is  $\{P_0 = 0\} \times \mathbb{R}^{n-l}$  by Corollary 2.8. By the Liouville theorem of [10],  $\{P_m = t\}$  is of the form  $H \times \mathbb{R}^{n-l}$  where  $H$  is a leaf of the Hardt-Simon foliation of  $\{P_0 = 0\}$  in  $\mathbb{R}^l$ . Thus for all  $t \in \mathbb{R}$ ,  $\xi \in \mathbb{R}^{n-l}$ ,  $\{P_m = t\}$  is invariant under  $x \mapsto x + \xi$ . This show that  $P_m$  is independent of  $x_{l+1}, \dots, x_n$ .

Since  $\{P_0 = 0\}$  is strictly minimizing, the Hardt-Simon foliation  $\{P_m = t\}$  is a graph of  $v$  over  $\{P_0 = 0\}$  with  $v$  decays like  $|x|^{-\gamma} \phi_1$  near infinity. Since  $\frac{1}{|\nabla(P_0 Q_m^{1/k})|}$  is the velocity of the family  $(\{P_m = t^k\})_{t \in (-\epsilon, \epsilon)}$  at  $t = 0$  near regular points of  $\{P_m = 0\}$ , we have

$$(25) \quad \deg P_0 - 1 + \frac{\deg Q_m}{k} = \frac{l - 3 - \sqrt{(l-3)^2 + 4\lambda_1}}{2}.$$

Equation (25) implies

$$(26) \quad (l-3)^2 + 4\lambda_1 \text{ is a perfect square integer.}$$

To see this, since  $|A|^2$  is an *integer* for isoparametric cones by Lemma 5.3, we have  $\lambda_1 = -|A|^2$  is also an integer. Since (25) implies that the square root in (25) is rational, it follows that the integer  $(l-3)^2 + 4\lambda_1$  is a perfect square.

By Theorem 2.9 (ii), we also have

$$(27) \quad k \text{ is odd and } \deg Q_m \text{ is even.}$$

We claim that (25), (26), (27) cannot hold simultaneously if  $P_0$  is a Cartan-Münzner polynomial. Indeed, we show this for each case in Lemma 5.3.

**Case (i)**  $\deg P_0 = 1$ . This is not what we want by our assumption.

**Case (ii)**  $\deg P_0 = 2$ ,  $\lambda_1 = -(l - 2)$ .

In order for  $\{P_0 = 0\}$  to be area minimizing, we have  $l \geq 8$ . By (26), there is  $a \in \mathbb{N}$  so that

$$(l - 3)^2 - 4(l - 2) = a^2.$$

This is equivalent to

$$(l - 5 - a)(l - 5 + a) = 8.$$

The only nonnegative integer solutions to this equation is

$$(l, a) = (8, 1).$$

By (25), we have  $k^{-1} \deg Q_m = 1$ . This contradicts (27).

**Case (iii)**  $\deg P_0 = 3$ .

In order for  $\{P_0 = 0\}$  to be area minimizing, we have  $l = 14$  and  $l = 26$ .

If  $l = 14$ , then  $\lambda_1 = -24$  and thus  $(l - 3)^2 + 4\lambda_1 = 121 - 96 = 25$ . By (25)  $k^{-1} \deg Q_m = 1$ . This contradicts (27).

If  $l = 26$ , then  $\lambda_1 = -48$  and  $(n - 3)^2 + 4\lambda_1 = 337$  is not a perfect square. This contradicts (26).

**Case (iv)**  $\deg P_0 = 4$ .

In order for  $\{P_0 = 0\}$  to be area minimizing, we have  $l \geq 16$ . By (26), there is  $a \in \mathbb{N}$  so that

$$(l - 3)^2 - 12(l - 2) = a^2.$$

This is equivalent to

$$(l - 9 - a)(l - 9 + a) = 48.$$

The positive integer solutions to this equation are

$$(l, a) = (16, 1), (17, 4), (22, 11).$$

Since  $l$  must be even by Lemma 5.3, the solution  $(l, a) = (17, 4)$  is not possible.

If  $(l, a) = (16, 1)$ , from (25) we have  $k^{-1} \deg Q_m = 3$ .

If  $(l, a) = (22, 11)$ , from (25) we have  $k^{-1} \deg Q_m = 1$ . Both violate (27).

**Case (v)**  $\deg P_0 = 6$ .

By Lemma 5.3,  $\{P_0 = 0\}$  is not area minimizing in this case. This finishes the proof.  $\square$

Now we prove our second “non-existence” theorem.

**Theorem 5.6.** *If there is a nonlinear polynomial solution to the minimal surface equation on  $\mathbb{R}^8$ , then there exists a cubic, strictly stable, area minimizing cone in  $\mathbb{R}^8$  which is not strictly minimizing.*

*Proof.* We take  $n = 8$  throughout the proof. By Theorem 2.9, we have

$$P = P_0^k Q_m + P_{m-1} + \cdots + P_2 + P_1.$$

Since  $\{P_0 = 0\}$  is 7 dimensional, it is a regular cone.

By Lemma 3.1, the function  $u = \frac{1}{|\nabla_{(P_0 Q_m^{1/k})}|}$  is a positive Jacobi field on  $\{P_0 = 0\}$ .

By [23, Lemma 6.1.7], we have  $\lambda_1 \leq -(n - 2)$ . Thus

$$(28) \quad \deg P_0 \leq \gamma_+ + 1 - k^{-1} \deg Q_m \leq \frac{n - 1 + \sqrt{(n - 3)^2 - 4(n - 2)}}{2} = 4.$$

By Theorem 5.4, we know that  $\deg P_0 \geq 3$ .

If  $\deg P_0 = 4$ , then equality holds in (28) and in particular  $\lambda_1 = -(n-2) = -6$ . By the rigidity part of Theorem A in [28] (see also [18], [30]), we know that  $\Sigma$  must be the Clifford torus. Hence  $\{P_0 = 0\}$  is a quadratic cone. This contradicts  $\deg P_0 = 4$ .

Thus  $\deg P_0 = 3$ . We show that  $\{P_0 = 0\}$  is strictly stable and not strictly minimizing. Indeed suppose  $\{P_0 = 0\}$  were not strictly stable, then  $\lambda_1 = -\frac{25}{4}$ . Since  $u = \frac{1}{|\nabla(P_0 Q_m^{1/k})|}$  is a positive Jacobi field decaying without log factor, we have

$$\deg P_0 - 1 + k^{-1} \deg Q_m = \frac{5}{2}.$$

Hence  $k^{-1} \deg Q_m = \frac{1}{2}$ . This contradicts the fact that  $k$  is odd and  $\deg Q_m$  is even by Theorem 2.9. Now suppose  $\{P_0 = 0\}$  is strictly minimizing, then we can apply [21, Theorem 5] to see that

$$\deg P \leq \gamma_- + 1 \leq \frac{7}{2}.$$

This contradicts Theorem 3.6 where cubic polynomial solutions are ruled out.  $\square$

#### REFERENCES

- [1] U. Abresch. Isoparametric hypersurfaces with four or six distinct principal curvatures. *Math. Ann.* 264: 283–302, 1983.
- [2] F. Almgren. Some interior regularity theorems for minimal surfaces and an extension of Bernstein’s theorem. *Ann. of Math.*, 84(1): 277–292, 1966.
- [3] S. Bernstein. Über ein geometrisches Theorem und seine Anwendung auf die partiellen Differentialgleichungen vom elliptischen Typus. *Math. Z.*, 26: 551–558, 1927.
- [4] J. Bochnak, M. Coste and M. Roy. *Real algebraic geometry*. Springer Science & Business Media, 36, 2013.
- [5] E. Bombieri, E. De Giorgi and E. Giusti. Minimal cones and the Bernstein problem. *Invent. Math.*, 7: 243–268, 1969.
- [6] E. Bombieri and E. Giusti. Harnack’s inequality for elliptic differential equations on minimal surfaces. *Invent. Math.*, 15 (1): 24–46, 1972.
- [7] E. Cartan. Familles de surfaces isoparamétriques dans les espaces à courbure constante. *Annali di Mat.* 17: 177–191, 1938.
- [8] E. Cartan. Sur des familles remarquables d’hypersurfaces isoparamétriques dans les espaces sphériques. *Math. Z.*, 45: 335–367, 1939.
- [9] E. De Giorgi. Una estensione del teorema di Bernstein. *Ann. Sc. norm. super. Pisa - Cl. sci.*, 19 (1): 79–85, 1965.
- [10] N. Edelen and G. Székelyhidi. A Liouville-type theorem for cylindrical cones. *arXiv math.DG* 2301.05967.
- [11] H. Federer. *Geometric measure theory*. Springer, 2014.
- [12] D. Ferus, H. Karcher, and H. F. Münzner, Cliffordalgebren und neue isoparametrische Hyperflächen. *Math. Z.*, 177: 479–502, 1981.
- [13] W. Fleming. On the oriented Plateau problem. *Rend. Circ. Mat.*, 11 (1): 69–90, 1962.
- [14] Y. Guo. Green’s functions on minimal submanifolds *arxiv preprint*
- [15] R. Hardt and L. Simon. Area minimizing hypersurfaces with isolated singularities. *J. Reine Angew. Math.*, 362: 102–129, 1985.
- [16] F.H. Lin, Minimality and stability of minimal hypersurfaces in  $\mathbb{R}^n$ . *Bull. Aust. Math. Soc.*, 2: 209–214, 1987.
- [17] H. F. Münzner. Isoparametric hyperflächen in sphären, I and II. *Math. Ann.*, 251: 57–71, 1980, and 256: 215–232, 1981.

- [18] O. Perdomo. First stability eigenvalue characterization of Clifford hypersurfaces. Proc. Amer. Math. Soc., 130(11): 3379-3384, 2002.
- [19] L. Simon. Lectures on geometric measure theory. The Australian National University, Mathematical Sciences Institute, Centre for Mathematics & its Applications, 1983.
- [20] L. Simon. Asymptotic behaviour of minimal graphs over exterior domains. Ann. Henri Poincaré, 4 (3): 231-242, 1987.
- [21] L. Simon. Entire solutions of the minimal surface equation. J. Differ. Geom., 30(3): 643-688, 1989.
- [22] L. Simon. The minimal surface equation. Geometry V, 239-266, 1997.
- [23] J. Simons. Minimal varieties in Riemannian manifolds. Ann. of Math., 88: 62-105, 1968.
- [24] Z. Tang and Y. Zhang. Minimizing cones associated with isoparametric foliations. J. Differ. Geom., 115(2): 367-393, 2020.
- [25] V. Tkachev. On a classification of minimal cubic cones in  $\mathbb{R}^n$ . arXiv:1009. 5409 (2010)
- [26] V. Tkachev. On the non-vanishing property for real analytic solutions of the  $p$ -Laplace equation. Proc. Amer. Math. Soc., 144(6): 2375-2382, 2016.
- [27] Q.-M. Wang. On a class of minimal hypersurfaces in  $\mathbb{R}^n$ , Math. Ann., 298: 207-251, 1994.
- [28] C. Wu. New characterizations of the Clifford tori and the Veronese surface. Arch. Math. (Basel), 61(3): 277-284, 1993.
- [29] S.-T. Yau, Shing-Tung, ed. Seminar on differential geometry. No. 102. Princeton University Press, 1982.
- [30] J. Zhu. First stability eigenvalue of singular minimal hypersurfaces in spheres. Calc. Var. Partial. Differ. Equ., 57(5): 1-13, 2018.

DEPARTMENT OF MATHEMATICS, ROWLAND HALL, UNIVERSITY OF CALIFORNIA, IRVINE, IRVINE,  
CA 92697, USA

*Email address:* yifag15@uci.edu