LAPTH-015/24

# Spherical symmetric solutions of conformal Killing gravity: black holes, wormholes, and sourceless cosmologies

Gérard Clément<sup>a</sup>\*and Khireddine Nouicer<sup>b†</sup>

 <sup>a</sup> LAPTh, Université Savoie Mont Blanc, CNRS, F-74940 Annecy, France
 <sup>b</sup> LPTh, Department of Physics, University of Jijel, BP 98, Ouled Aissa, Jijel 18000, Algeria

March 30, 2024

#### Abstract

The most general set of static and spherically symmetric solutions for conformal Killing gravity coupled to Maxwell fields is presented in closed form. These solutions, depending on six parameters, include non-asymptotically flat black holes or naked singularities, nonasymptotically flat traversable wormholes, and (possibly singularityfree) closed universes. We also consider the inverse problem, showing that the most general energy-momentum tensor generating a given static spherically symmetric metric depends on three parameters. Sourceless time-dependent isotropic solutions are also given. These solutions depending on the curvature, the cosmological constant, and a new integration constant  $\alpha$ , present a rich variety, including singularity-free eternal cosmologies and universes evolving symmetrically from a big bang to a big crunch within a finite lapse of time.

<sup>\*</sup>Email: gclement@lapth.cnrs.fr

<sup>&</sup>lt;sup>†</sup>Email: khnouicer@univ-jijel.dz

# 1 Introduction

In the pursuit of understanding the universe's accelerated expansion [1, 2], numerous alternatives have been proposed, both within the framework of general relativity and beyond. One such alternative involves introducing a new matter component termed dark energy, characterized by negative pressure. While the cosmological constant  $\Lambda$ , originally introduced by Einstein, remains the primary candidate for dark energy due to its alignment with observational data, challenges such as the fine-tuning and coincidence problems persist [3, 4, 5, 6]. On the other hand, a different path follows the idea that general relativity fails to describe the universe at larger scales and hence it should be modified. These modifications often propose alterations to fundamental principles of physics, seeking to reconcile observational evidence with theoretical predictions [7, 8]. Consequently, alternative models like scalar tensor theories [9, 10], Einstein-æther theory [11, 12], and higher derivative theories of gravity like f(R) theories [13, 14] have been explored. Other kinds of modified theories are the modified teleparallel equivalent of general relativity, f(T) gravity where the Ricci scalar is replaced by the torsion scalar, and the theory grounded in Weyl geometry, termed f(Q) gravity [15]. The latter describes gravitational effects in terms of non-metricity, providing a broader framework than traditional Riemannian geometry. These theories offer different perspectives on gravity that may lead to new insights into cosmological and astrophysical phenomena.

Recently a new gravitational field theory was proposed by J. Harada [16], and subsequently termed "conformal Killing gravity" [17]. By construction, all solutions of the Einstein equations with or without cosmological constant (which arises as an integration constant of Harada's field equations) are also solutions of this new theory. But, the field equations of conformal Killing gravity being differential equations of third order, it also yields solutions beyond those of cosmological general relativity. The application of conformal Killing gravity to cosmology has led to solutions that shed light on the transition from decelerating to accelerating expansion, offering an explanation for the current cosmic acceleration [18]. Black hole solutions to conformal Killing gravity coupled to nonlinear electrodynamics have been explored in [19].

Very recently, A. Barnes [20, 21] considered the field equations of conformal Killing gravity coupled to Maxwell fields and obtained static solutions in the form of infinite power series using the Frobenius method. In this work, we revisit this problem and derive the most general static and spherically symmetric solutions to conformal Killing gravity field equations sourced by Maxwell field in closed and compact form. The structure of this paper is outlined as follows. In the next section, we introduce the field equations of conformal Killing gravity, and derive the solutions. In Sec. 3 we explore the geometric properties of the solutions, which contain black holes, wormholes, and singularity-free closed universes. In Sec. 4, we investigate the inverse problem, deriving the most general energy-momentum tensors which generate the metrics of Sec. 2. In Sec. 5 we consider the class of time-dependent Friedmann-Lemaître-Robertson-Walker (FLRW) cosmologies solving the sourceless equations of conformal Killing gravity and discuss them for flat, closed, and open space-times. Finally, we provide in Sec. 6 a summary and discussion of our results.

# 2 Static spherically symmetric solutions with Maxwell source

The gravitational field equations proposed by Harada [16] are expressed as follows:

$$H_{\mu\nu\rho} = 8\pi G T_{\mu\nu\rho},\tag{2.1}$$

where the totally symmetric tensor  $H_{\mu\nu\rho}$  is defined by

$$H_{\mu\nu\rho} \equiv \nabla_{\mu}R_{\nu\rho} + \nabla_{\nu}R_{\rho\mu} + \nabla_{\rho}R_{\mu\nu} - \frac{1}{3}\left(g_{\nu\rho}\partial_{\mu} + g_{\rho\mu}\partial_{\nu} + g_{\mu\nu}\partial_{\rho}\right)R, \quad (2.2)$$

with  $R_{\mu\nu}$  the Ricci tensor, R its trace, and  $\nabla$  the covariant derivative. The components of this three-tensor are constrained by the Bianchi identity

$$g^{\nu\rho}H_{\mu\nu\rho} \equiv 0. \tag{2.3}$$

The matter three-tensor  $T_{\mu\nu\rho}$  is similarly defined as:

$$T_{\mu\nu\rho} \equiv \nabla_{\mu}T_{\nu\rho} + \nabla_{\nu}T_{\rho\mu} + \nabla_{\rho}T_{\mu\nu} - \frac{1}{6}\left(g_{\nu\rho}\partial_{\mu} + g_{\rho\mu}\partial_{\nu} + g_{\mu\nu}\partial_{\rho}\right)T, \quad (2.4)$$

with  $T_{\mu\nu}$  the energy-momentum tensor, and T its trace. The conservation of energy-momentum is embodied in the equation

$$g^{\nu\rho}T_{\mu\nu\rho} = 0. (2.5)$$

It was shown in [17] that the Harada field equations (2.1) are equivalent to the Einstein equations

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = 8\pi G(T_{\mu\nu} + K_{\mu\nu})$$
(2.6)

modified by the addition to the matter energy-momentum tensor of a divergencefree ( $\nabla_{\nu} K^{\mu\nu} = 0$ ) conformal Killing tensor  $K_{\mu\nu}$  of trace K satisfying the equation

$$\nabla_{\mu}K_{\nu\rho} + \nabla_{\nu}K_{\rho\mu} + \nabla_{\rho}K_{\mu\nu} - \frac{1}{6}\left(g_{\nu\rho}\partial_{\mu} + g_{\rho\mu}\partial_{\nu} + g_{\mu\nu}\partial_{\rho}\right)K = 0.$$
(2.7)

This formulation will not be used in the following.

Let us now recall the derivation of the differential equations for static spherically symmetric metrics sourced by a Maxwell field [21]. The general static and spherically symmetric spacetime metric can be written as

$$ds^{2} = -e^{2\nu(r)}dt^{2} + e^{2\lambda(r)}dr^{2} + r^{2}d\Omega^{2}$$
(2.8)

 $(d\Omega^2 = d\theta^2 + \sin^2 \theta \, d\varphi^2)$ . The monopole solution of the Maxwell equations in this geometry leads to the energy-momentum tensor components

$$T^{t}_{t} = T^{r}_{r} = -T^{\theta}_{\theta} = -T^{\varphi}_{\varphi} = -\frac{q^{2}}{8\pi}r^{-4}, \qquad (2.9)$$

where  $q^2$  is the sum of the squared monopole electric and magnetic charges. The resulting components of the three-tensor (2.4) are

$$T^{t}_{tr} = -\frac{1}{3}T^{r}_{rr} = \frac{1}{2}T^{\theta}_{\theta r} = \frac{1}{2}T^{\varphi}_{\varphi r} = -\frac{q^{2}}{2\pi}r^{-5}.$$
 (2.10)

By virtue of the constraints (2.3) and (2.5) the only independent equations (2.1) are, for G = 1,

$$H^{t}_{tr} - 8\pi T^{t}_{tr} \equiv \frac{1}{3}e^{-2\lambda} \left[\nu''' + (2\nu' - 3\lambda')\nu'' - \nu'\lambda'' - 2(\nu' - \lambda')\nu'\lambda' + \frac{2}{r}(\nu'' + 2\lambda'' - 6\nu'^{2} - 8\nu'\lambda' - 4\lambda'^{2}) - \frac{2}{r^{2}}\nu' + \frac{4}{r^{3}}\right] - \frac{4}{3r^{3}} + \frac{4q^{2}}{r^{5}} = 0, \qquad (2.11)$$

where ' = d/dr, and the combination

$$H^{t}_{tr} - \frac{1}{3}H^{r}_{rr} = g_{tt}\frac{d}{dr}\left[g_{tt}^{-1}(R^{t}_{t} - R^{r}_{r})\right] = -2e^{2\nu}\left[e^{-2(\nu+\lambda)}\frac{(\nu'+\lambda')}{r}\right]' = 0.$$
(2.12)

This last equation is solved by

$$e^{-2(\lambda(r)+\nu(r))} = c + dr^2, \qquad (2.13)$$

where c and d are integration constants. Using this equation to eliminate  $\lambda(r)$  in terms of  $\nu(r)$  from equation (2.11), and putting  $e^{2\nu(r)} = y(r)$ , we arrive at the master linear differential equation

$$(c+dr^{2})r^{3}y^{\prime\prime\prime}(r) + (-2c+dr^{2})r^{2}y^{\prime\prime}(r) + (-2c-dr^{2})ry^{\prime}(r) + 8cy(r) - 8 + \frac{24q^{2}}{r^{2}} = 0$$
(2.14)

So the metric is

$$ds^{2} = -y(r) dt^{2} + \frac{dr^{2}}{(c+dr^{2})y(r)} + r^{2} d\Omega^{2}, \qquad (2.15)$$

where y(r) solves equation (2.14). For this metric to have the Lorentzian signature, the product  $g_{tt}g_{rr} = -(c + dr^2)$  must be negative definite. Three cases must be considered.

<u>Case 1: d = 0.</u> The metric has the Lorentzian signature provided c > 0. By rescaling the time coordinate, one can fix c = 1. The solution of the master equation (2.14) is [16]

$$y(r) = 1 - \frac{2m}{r} + \frac{q^2}{r^2} - \frac{\Lambda r^2}{3} - \frac{\lambda r^4}{5}.$$
 (2.16)

<u>Case 2:</u>  $d \neq 0, c \neq 0$ . The metric has the Lorentzian signature in three subcases: 2a) c > 0, d > 0, with r real; 2b) c < 0, d > 0, with  $r^2 > \mu^2$ ; 2c) c > 0, d < 0, with  $r^2 < \mu^2$ . In these three cases, the modulus of c can be rescaled to 1 by a rescaling of the time coordinate, so we can fix  $c = \epsilon, d = \eta/\mu^2$ , where  $\epsilon = \pm 1, \eta = \pm 1$ , and  $\mu$  is a constant with the dimension of a length. The generic solution for these three cases is

$$y(r) = \frac{q^2(2\eta r^2 + \epsilon\mu^2)}{\mu^2 r^2} + k_1 \frac{\sqrt{\eta r^2 + \epsilon\mu^2}}{r} + \epsilon + k_2 \left[\frac{\sqrt{\eta r^2 + \epsilon\mu^2}}{r}\psi\left(\frac{r}{\mu}\right) - 1\right] - \frac{\Lambda r^2}{3}, \quad (2.17)$$

where

$$a)\psi(x) = \operatorname{arsinh} x, \quad \epsilon = +1, \ \eta = +1,$$
 (2.18)

$$b)\psi(x) = \operatorname{arcosh} x, \quad \epsilon = -1, \ \eta = +1 \qquad (r^2 > \mu^2), \qquad (2.19)$$

$$c)\psi(x) = \arcsin x, \quad \epsilon = +1, \ \eta = -1 \qquad (r^2 < \mu^2).$$
 (2.20)

Expanding these solutions in Laurent series for the variable  $r/\mu$ , we recover the solutions given by Barnes [21].

<u>Case 3: c = 0.</u> In this case, the constant d must be positive. Putting again  $d = 1/\mu^2$ , the solution of equation (2.14) is [21]

$$y(r) = \frac{q^2 \mu^2}{4r^4} - \frac{\mu^2}{2r^2} + k_1 + k_2 \ln \left| \frac{r}{\mu} \right| - \frac{\Lambda r^2}{3}.$$
 (2.21)

# 3 Geometry

#### 3.1 Case 1

If  $\lambda \neq 0$ , the Ricci scalar [16]

$$R = 4\Lambda + 6\lambda r^2 \tag{3.22}$$

diverges on the sphere at spacelike infinity  $r \to \infty$ , which is a naked singularity of this spacetime. Furthermore, the coordinate transformation  $r = x^{-1}$  leads to the asymptotic form of the metric

$$ds^{2} \simeq \frac{\lambda}{5} x^{-4} dt^{2} - \frac{5}{\lambda} dx^{2} + x^{-2} d\Omega^{2} \quad (x \to 0), \qquad (3.23)$$

showing that "spacelike infinity" is actually at finite geodesic distance for all geodesics if  $\lambda > 0$ , and for spacelike radial geodesics if  $\lambda < 0$ . So the case  $\lambda \neq 0$  is unphysical. When  $\lambda = 0$  the solution reduces to the Reissner-Nordström-(anti-)de Sitter solution of general relativity.

#### 3.2 Case 2

In this case the Ricci scalar is

$$\mu^2 R = -6\eta \left[ \frac{\epsilon q^2}{r^2} + \left( k_1 + k_2 \psi \left( \frac{r}{\mu} \right) \right) \frac{\sqrt{\eta r^2 + \epsilon \mu^2}}{r} + \epsilon + \frac{2\eta q^2}{\mu^2} - \Lambda r^2 \right]$$
  
+2\eta k\_2 + 4\epsilon \Lambda \mu^2. (3.24)

The possible singularities are at  $r \to \infty$  and at the center r = 0. As in case 1,  $r \to \infty$  is a naked singularity at finite geodesic distance in subcases 2a or 2b unless  $\Lambda = 0$ . If  $\Lambda = 0$  but  $k_2 \neq 0$ , the logarithmic divergence of the function  $\psi(r/\mu)$  (an arsinh or arcosh) at  $r \to \infty$  in subcases 2a or 2b carries over to the curvature invariants. However, the coordinate transformation  $r = \mu e^x$  leads for  $d = 1/\mu^2$  to the asymptotic form of the metric

$$ds^2 \simeq -k_2 x dt^2 + \frac{dx^2}{k_2 x} + \mu^2 e^{2x} d\Omega^2 \quad (x \to \infty),$$
 (3.25)

showing that spacelike (for  $k_2 > 0$ ) or timelike (for  $k_2 < 0$ ) geodesics extend to  $r \to \infty$ . In other words, geodesics do not terminate at  $r \to \infty$  if  $\Lambda = 0$ but  $k_2 \neq 0$ . The invariants also diverge for  $r \to 0$ , unless q = 0 and  $k_1 = 0$ .

#### 3.2.1 Subcase 2a

We choose  $\Lambda = 0$  in (2.17) to avoid naked singularities for  $r \to \infty$ , and discuss first the simple case  $k_2 = 0$ . In the vacuum case q = 0,

$$y(r) = 1 + k_1 \frac{\rho}{r} \quad \left(\rho = \sqrt{r^2 + \mu^2}\right),$$
 (3.26)

with  $k_1 > -1$  for  $y(\infty) > 0$ . For  $-1 < k_1 < 0$ , the spacetime [2.15) is a black hole, with a single horizon shielding the singularity r = 0. For  $k_1 > 0$  the singularity is naked. For  $k_1 = 0$ , the singularity-free metric

$$ds^{2} = -dt^{2} + \frac{dr^{2}}{1 + r^{2}/\mu^{2}} + r^{2} d\Omega^{2}$$
(3.27)

is that of a negative density Einstein static universe [20].

In the electromagnetic case,  $q \neq 0$ , the lapse function is

$$y(r) = \bar{q}^2 \frac{\rho^2}{r^2} + k_1 \frac{\rho}{r} + \bar{q}^2 + 1 \quad (\bar{q} = q/\mu),$$
(3.28)

with  $1 + k_1 + 2\bar{q}^2 > 0$ . For  $-(1 + 2\bar{q}^2) < k_1 < -2|\bar{q}|\sqrt{1 + \bar{q}^2}$  the spacetime is a black hole with two horizons and a central singularity. It becomes an extreme black hole for  $k_1 = -2|\bar{q}|\sqrt{1 + \bar{q}^2}$ , with a double horizon at r = q. For  $k_1 > -2|\bar{q}|\sqrt{1 + \bar{q}^2}$ , the singularity is naked.

Consider now the general case  $k_2 > 0$  (the sign being chosen so that  $y(\infty) > 0$ ). The lapse function y is now

$$y(r) = y_0(r) + k_2 z(r), (3.29)$$

where  $y_0(r)$  is the function (3.26) or (3.28), and  $z(r) = (\rho/r) \operatorname{arsinh} (r/\mu) - 1$ is a positive, steadily increasing function, such that z(0) = z'(0) = 0. It follows that the results obtained above for  $k_2 = 0$  remain qualitatively valid (without a lower bound on  $k_1$ ). For q = 0 the spacetime is a black hole with a single horizon if  $k_1 < 0$ , it is geodesically complete if  $k_1 = 0$ , and the singularity is naked if  $k_1 > 0$ . For  $q \neq 0$  the spacetime corresponds either to a black hole with a double horizon, to an extreme black hole, or to a naked singularity, depending on the parameter values.

#### 3.2.2 Subcase 2b

We again choose  $\Lambda = 0$  to avoid naked singularities for  $r \to \infty$ . In this case, because  $r^2 > \mu^2$ , the metric is best parameterized in terms of the radial coordinate  $\rho = \sqrt{r^2 - \mu^2}$ :

$$ds^{2} = -y(\rho) dt^{2} + \frac{\mu^{2} d\rho^{2}}{(\mu^{2} + \rho^{2})y(\rho)} + (\mu^{2} + \rho^{2}) d\Omega^{2}, \qquad (3.30)$$

with

$$y(\rho) = \bar{q}^2 x^2 + k_1 x + \bar{q}^2 - 1 + k_2 [x \operatorname{artanh} x - 1], \quad x = \frac{\rho}{\sqrt{\mu^2 + \rho^2}} \quad (-1 < x < 1).$$
(3.31)

While the radial coordinate  $\rho$  has been defined to be positive, there is clearly no coordinate singularity for  $\rho = 0$ , so that the spacetime should be analytically extended to  $\rho < 0$ . The variable  $\rho$  then takes its values in the whole real axis, so that there are two points at infinity  $\rho \to \pm \infty$   $(x \to \pm 1)$ , and a sphere of minimal area  $4\pi\mu^2$  for  $\rho = 0$ .

We first discuss the case  $k_2 = 0$ . Because of the quadratic form of the metric function y, there are at most two horizons. The central singularity is absent, so that the spacetime will be geodesically complete after analytical extension. The solution is stationary for  $\rho \to +\infty$  provided

$$k_1 > 1 - 2\bar{q}^2, \tag{3.32}$$

which we now assume. It will also be stationary for  $\rho \to -\infty$  if  $k_1 < 2\bar{q}^2 - 1$ , in which case there will be two or no horizons. On the contrary, if  $k_1 > 2\bar{q}^2 - 1$ , the metric function  $y(\rho)$  changes sign between  $\rho \to +\infty$  and  $\rho \to -\infty$ , so that there is a single horizon.

We consider first the second possibility which, on account of (3.32), is realized if

$$\bar{q}^2 \le \frac{1}{2}$$
 or  $\bar{q}^2 > \frac{1}{2}$  and  $k_1 > 2\bar{q}^2 - 1.$  (3.33)

This case includes the vacuum solution q = 0. The spacetime is a singularity-free black hole with one horizon.

In the complementary case

$$\bar{q}^2 > \frac{1}{2}$$
 and  $k_1 < 2\bar{q}^2 - 1,$  (3.34)

there are three possibilities. For

$$k_1^2 > 4\bar{q}^2(\bar{q}^2 - 1), \tag{3.35}$$

the spacetime is a singularity-free black hole with two horizons. For

$$k_1^2 = 4\bar{q}^2(\bar{q}^2 - 1) \tag{3.36}$$

the spacetime is a singularity-free extreme black hole. And for

$$k_1^2 < 4\bar{q}^2(\bar{q}^2 - 1), \tag{3.37}$$

the spacetime is a traversable wormhole, which is symmetrical if  $k_1 = 0$ , the wormhole condition (3.37) implying  $\bar{q}^2 > 1$ . The existence of a traversable wormhole solution to conformal Killing gravity is a remarkable feature, since these are forbidden in general relativity by the null energy condition [22].

If  $k_2 > 0$ , the metric function y diverges for  $\rho \to \pm \infty$  as  $y \simeq k_2 \ln |2\rho/\mu|$ . Accordingly, there is necessarily an even number of horizons. It follows from  $y(0) = \bar{q}^2 - 1 - k_2$  that, if

$$\bar{q}^2 < 1 + k_2$$
 (3.38)

(including the vacuum solution q = 0), the spacetime is a singularity-free black hole with two (or possibly more) horizons. It can be a traversable wormhole if  $\bar{q}^2 > 1 + k_2$ . In the case  $k_1 = 0$ ,  $y(\rho)$  is even in  $\rho$  with a positive minimum at  $\rho = 0$  if

$$\bar{q}^2 > 1 + k_2,$$
 (3.39)

leading to a symmetrical traversable wormhole.

#### 3.2.3 Subcase 2c

This case is complementary to the preceding, with  $r^2 < \mu^2$ . The metric is, in terms of the radial coordinate  $\psi = \arcsin r/\mu \ (-\pi/2 < \psi < \pi/2)$ ,

$$ds^{2} = -y(\psi) dt^{2} + \mu^{2} \left[ \frac{d\psi^{2}}{y(\psi)} + \sin^{2} \psi \, d\Omega^{2} \right], \qquad (3.40)$$

with

$$y(\psi) = \bar{q}^2 \cot^2 \psi + k_1 \cot \psi + 1 - \bar{q}^2 + k_2(\psi \cot \psi - 1) - \frac{\Lambda \mu^2}{3} \sin^2 \psi. \quad (3.41)$$

As the spacetime does not extend to spacelike infinity, there is no constraint on the integration constant  $\Lambda$ . If q and/or  $k_1$  are different from 0, the metric is singular at the center  $\psi = 0$ . More interesting is the case of the vacuum solution, q = 0, with also  $k_1 = 0$ . The metric is then singularity-free. It is stationary at the center, where y(0) = 1, and is stationary everywhere  $(y(\psi) > 0)$  for all  $\psi \in [-\pi/2, \pi/2]$  provided

$$k_2 + \frac{\Lambda \mu^2}{3} < 1. \tag{3.42}$$

This case corresponds to a regular, closed static universe, with finite volume  $V = 2\pi^2 \mu^3$ . In the special subcase  $\Lambda = k_2 = 0$ , this is the Einstein static universe.

#### 3.3 Case 3

The Ricci scalar for this case is

$$\mu^2 R = -\frac{3q^2\mu^2}{2r^4} + \frac{3\mu^2}{r^2} - 4k_2 - 6\left(k_1 + k_2 \ln\left|\frac{r}{\mu}\right|\right) + 6\Lambda r^2.$$
(3.43)

As in subcases 2a and 2b, we choose  $\Lambda = 0$  to avoid naked singularities for  $r \to \infty$ . The metric function y is

$$y(r) = \frac{\bar{q}^2 \mu^4}{4r^4} - \frac{\mu^2}{2r^2} + k_1 + k_2 \ln \left| \frac{r}{\mu} \right|, \qquad (3.44)$$

with  $k_2 > 0$  or  $(k_2 = 0 \text{ and } k_1 > 0)$  for the solution to be stationary at spacelike infinity. The central singularity r = 0 is always present. In the vacuum case q = 0, the spacetime is a black hole with a single horizon. In the electromagnetic case  $q \neq 0$ , it can correspond either to a black hole with two horizons, an extreme black hole, or a naked singularity, depending on the parameter values.

# 4 The inverse problem

Let us recall that any solution of Einstein's field equations is, by construction, a solution of the equations of conformal Killing gravity. The static Einstein universe is a solution of general relativity sourced by a perfect fluid with constant density  $\rho$  and opposite pressure p. We have seen that the static Einstein universe is also a vacuum solution  $(T_{\mu\nu} = 0)$  of conformal Killing gravity, which implies that its perfect fluid source must satisfy the condition  $T_{\mu\nu\rho} = 0$ , and also generate the other vacuum solutions of Section 2. This prompts us to examine the inverse problem: What is the most general energy-momentum tensor generating a given metric (for instance the Schwarzschild solution)?

A static spherically symmetric matter source is characterized by a diagonal energy-momentum tensor  $T_{\mu\nu}$ , which may be thought of as a static anisotropic fluid, with components

$$T^{\nu}_{\mu} = \text{diag}\left(-\rho, p - 2\Pi, p + \Pi, p + \Pi\right).$$
 (4.1)

From (4.1) we obtain the components of the tensor  $T_{\mu\nu\rho}$ 

$$T_{100} \equiv e^{2\nu} \left[ \frac{5\rho'}{6} + \frac{p'}{2} - 2(\rho + p - 2\Pi)\nu' \right],$$
  

$$T_{111} \equiv 3e^{2\lambda} \left[ \frac{\rho'}{6} + \frac{p'}{2} - 2\Pi' \right],$$
  

$$T_{122} \equiv r^2 \left[ \frac{\rho'}{6} + \frac{p'}{2} + \Pi' - \frac{6\Pi}{r} \right], \quad T_{133} = \sin^2 \theta T_{122}, \quad (4.2)$$

satisfying the conservation law (2.5), which reads here

$$p' - 2\Pi' + (\rho + p - 2\Pi)\nu' - \frac{6\Pi}{r} = 0.$$
(4.3)

The most general energy-momentum tensor generating the vacuum (Q = 0) spacetime metrics of section 3 is such that the components (4.2) all vanish. We first consider the combination

$$-e^{-2\lambda}T_{111} + 3r^{-2}T_{122} \equiv 9\left(\Pi' - 2\frac{\Pi}{r}\right) = 0, \qquad (4.4)$$

which is solved by  $\Pi = (\alpha/3)r^2$ , with  $\alpha$  an integration constant. Inserting this in the equation  $T_{111} = 0$ , and integrating, we obtain

$$p = \frac{1}{3}(-\rho + 2\beta + 4\alpha r^2), \qquad (4.5)$$

with  $\beta$  a new integration constant. Finally, solving the equation  $T_{100} = 0$  leads to the characteristic functions of the most general energy-momentum tensor (4.1) generating a given Q = 0 spacetime metric (2.15):

$$\rho(r) = \gamma y(r) - \alpha r^2 - \beta, \ p(r) = \frac{1}{3} [-\gamma y(r) + 5\alpha r^2 + 3\beta], \ \Pi(r) = \frac{\alpha}{3} r^2, \ (4.6)$$

where  $\gamma$  is a third integration constant, and y(r) is the metric function for the spacetime under consideration. For instance, the most general energymomentum tensor generating the Schwarzschild metric of mass M in conformal Killing gravity corresponds to

$$\rho_S(r) = \gamma (1 - 2M/r) - \alpha r^2 - \beta, \ p_S(r) = \frac{1}{3} [-\gamma (1 - 2M/r) + 5\alpha r^2 + 3\beta], \ \Pi_S(r) = \frac{\alpha}{3} r^2.$$
(4.7)

For  $\alpha = \gamma = 0$ , this reduces to the energy-momentum generating the Einstein static universe in general relativity.

# 5 Time-dependent vacuum solutions

The occurrence of the Einstein static universe as a sourceless  $(T_{\mu\nu} = 0)$  solution of conformal Killing gravity suggests that this theory might admit other sourceless cosmological solutions. Let us first recall the derivation of the Friedmann-like equation given in [16]. Making the FLRW metric ansatz

$$ds^{2} = -dt^{2} + a^{2}(t) \left(\frac{dr^{2}}{1 - kr^{2}} + r^{2}d\Omega^{2}\right)$$
(5.1)

for an isotropic and spatially homogeneous universe, and assuming that the matter source is a perfect fluid with energy-momentum tensor

$$T^{\nu}_{\mu} = \operatorname{diag}\left(-\rho, p, p, p\right) \tag{5.2}$$

satisfying the conservation law

$$\dot{\rho} + 3(\rho + p)\frac{\dot{a}}{a} = 0 \tag{5.3}$$

(where  $\dot{=} d/dt$ ), Harada [16] showed that the equations of conformal Killing gravity may be first integrated to the second order equation

$$-\frac{\ddot{a}}{a} + \frac{2\dot{a}^2}{a^2} + \frac{2k}{a^2} - \frac{\Lambda}{3} = \frac{4\pi G}{3}(5\rho + 3p),$$
(5.4)

where the integration constant  $\Lambda$  may be identified with the cosmological constant.

Let us now integrate again this equation as follows. Putting  $a = b^{-1}$  and using the conservation law (5.3), equation (5.4) may be rewritten as

$$\frac{\ddot{b}}{b^3} + 2k - \frac{\Lambda}{3b^2} - \frac{4\pi G\dot{\rho}}{3b\dot{b}} - \frac{8\pi G\rho}{3b^2} = 0,$$
(5.5)

which (assuming  $\dot{b} \neq 0$ ) is first integrated by

$$\dot{b}^2 + kb^4 - \frac{\Lambda + 8\pi G\rho}{3}b^2 = \alpha,$$
 (5.6)

where  $\alpha$  is an integration constant, or

$$\frac{\dot{a}^2}{a^2} = \frac{8\pi G\rho}{3} + \frac{\Lambda}{3} - \frac{k}{a^2} + \alpha a^2.$$
(5.7)

This equation was previously obtained through an alternative method in [17] (our constant  $\alpha$  is equivalent to their constant -C/6). If  $\alpha = 0$  we recover the usual Friedmann equation. Thus, the integration constant  $\alpha$  measures the deviation of conformal Killing gravity from cosmological general relativity [17].

A more direct interpretation of this constant has been proposed in [18]. Consider the cosmological Einstein equations

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G (T_{\mu\nu} + T_{\text{eff}\,\mu\nu}), \qquad (5.8)$$

where  $T_{\mu\nu}$  is the sum of the energy-momentum tensors for matter and radiation, and  $T_{\text{eff}\mu\nu}$  is the energy-momentum tensor for a hypothetical matter component, which Harada identified with effective dark energy, with density  $\rho_{\text{eff}}$  and pressure  $p_{\text{eff}}$  satisfying the equation of state

$$5\rho_{\rm eff} + 3p_{\rm eff} = 0.$$
 (5.9)

For this matter component, the integration of the conservation law (5.3) leads to the solution  $\rho_{\text{eff}}(t) = Ca^2(t)$  (with C an integration constant). Thus, the contribution of this effective dark energy to the general-relativistic Friedmann equation

$$\frac{\dot{a}^2}{a^2} = \frac{8\pi G\rho}{3} + \frac{8\pi G\rho_{\text{eff}}}{3} + \frac{\Lambda}{3} - \frac{k}{a^2}$$
(5.10)

can be identified with the term  $\alpha a^2$  in (5.7), if  $C = \alpha/8\pi G$ . So, according to Harada, this term can account for an effective dark energy which appears in the theory without needing to be introduced in the physical energy-momentum tensor  $T_{\mu\nu}$ .

In vacuum ( $\rho = p = 0$ ), equation (5.7) reduces to

$$\dot{a}^2 + V(a) = 0, \quad V(a) = -\alpha a^4 - \frac{\Lambda}{3}a^2 + k.$$
 (5.11)

The general solution to this equation can be written in terms of Jacobi functions, but it is enough to discuss the behavior of the solutions depending on the relative values of the integration constants k,  $\Lambda$ ,  $\alpha$ , and on the sign of the discriminant of V(a), which is

$$\Delta = \frac{\Lambda^2}{9} + 4k\alpha. \tag{5.12}$$

Let us discuss the two cases  $\alpha > 0$ , and  $\alpha < 0$ .

#### **5.1** $\alpha > 0$

For  $\Delta > 0$ , the effective potential V(a) has two real roots  $a_{\pm}^2 = (-\Lambda/3 \pm \sqrt{\Delta})/2\alpha$ , with  $a_{\pm}^2 > 0$  if  $\Lambda < 0$  or k > 0, and  $a_{\pm}^2 > 0$  if  $\Lambda < 0$  and k < 0. The scale factor a(t) can vary from  $a_{\pm}$  to infinity, or from 0 to  $a_{\pm}$ , if  $a_{\pm}$  or  $a_{\pm}$  are real. If not, a(t) varies from zero to infinity. Near infinity, a(t) behaves as

(

$$a \simeq \pm \frac{1}{\sqrt{\alpha}} \frac{1}{t - t_0} \quad (t \to t_0), \tag{5.13}$$

for some finite  $t_0$ .

For  $\underline{k} > 0$  and all values of  $\Lambda$ , the evolution is time-symmetrical, with a minimum scale function a(t) at t = 0 and two asymptotes at  $t = \pm t_0$ .

For <u>k = 0</u>, equation (5.11) can be transformed by putting a(t) = 1/b(t) to  $\dot{b}^2 - (\Lambda/3)b^2 = \alpha$ , which can be solved in terms of elementary functions:

$$a = \pm \frac{\sqrt{\Lambda/3\alpha}}{\sinh\left(\sqrt{\Lambda/3}t\right)} \qquad (\Lambda > 0),$$
  

$$a = \pm \frac{1}{\sqrt{\alpha}t} \qquad (\Lambda = 0), \qquad (5.14)$$
  

$$a = \pm \frac{\sqrt{-\Lambda/3\alpha}}{\sin\left(\sqrt{-\Lambda/3}t\right)} \qquad (\Lambda < 0)$$

(up to a time translation). If  $\Lambda \ge 0$   $(a_+ = 0)$  the scale factor asymptotes to zero for  $t = \pm \infty$  and to infinity for t = 0. If  $\Lambda < 0$   $(a_+ = \sqrt{-\Lambda/3\alpha})$ , the evolution is the same as for k > 0.

For  $\underline{k} < 0$ ,  $\Lambda > -6\sqrt{-k\alpha}$ , V(a) is negative definite, so a(t) varies from zero to infinity, with the exact solution if  $\Lambda = 6\sqrt{-k\alpha}$ ,

$$a = \sqrt{\Lambda/6\alpha} |\tan(\sqrt{\Lambda/6t})|.$$
(5.15)

If  $\Lambda = -6\sqrt{-k\alpha}$ , we have again the two exact solutions

$$a = \sqrt{-\Lambda/6\alpha} |\tanh(\sqrt{\Lambda/6t})|, \text{ or } a = \sqrt{-\Lambda/6\alpha} |\coth(\sqrt{\Lambda/6t})|.$$
 (5.16)

If  $\Lambda < -6\sqrt{-k\alpha}$ , the two roots are real, so that the evolution is time symmetrical, with either the universe contracting to a minimum radius at  $a = a_+$  and then expanding again to infinity in a finite time interval  $2t_0$ ; or blowing up from a big bang at  $t = -t_0$  to a maximum radius at  $a = a_-$  and then recontracting to a big crunch at  $t = t_0$ .

## **5.2** $\alpha < 0$

This is possible only if  $\Delta > 0$ . The squared scale factor  $a^2(t)$  must remain between the two roots  $a_{\pm}^2 = (\Lambda/3 \pm \sqrt{\Delta})/(-2\alpha)$ , implying  $a_{\pm}^2 > 0$ .

For  $\underline{k} > 0$ ,  $a_{+}^{2}$  is positive only if  $\Lambda > 6\sqrt{-k\alpha}$ . In this case  $a_{-}^{2}$  is also positive, leading to an oscillating cosmology, with the scale factor varying periodically between  $a_{-}$  and  $a_{+}$ .

For  $\underline{k} = 0$ ,  $\Lambda > 0$ , we have the exact solution

$$a = \frac{\sqrt{\Lambda/(-3\alpha)}}{\cosh\left(\sqrt{\Lambda/3}t\right)}.$$
(5.17)

For  $\underline{k} < 0$ ,  $a_{+}^{2}$  is positive and  $a_{-}^{2}$  is negative for all values of  $\Lambda$ . The universe evolves symmetrically from a big bang to a big crunch in a finite lapse of time.

# 6 Discussion

We have obtained all the static spherically symmetric solutions, depending on six integration constants, of conformal Killing gravity with Maxwell source. These solutions fall in three classes, according to the values of the integration constants c and d. The Schwarzschild-like class d = 0 was previously given in [16]. The solutions of this class depend on the mass m, electric or magnetic charge q, cosmological constant  $\Lambda$ , and Harada's integration constant  $\lambda$ . However, when  $\lambda \neq 0$ , the metric is singular on the spheres  $r \to \infty$ , which are actually at finite geodesic distance. When  $\lambda = 0$ the solution reduces to the well-known Reissner-Nordström-(anti-)de Sitter solution. The solutions of the class c = 0, previously given in [21], also present for  $r \to \infty$  a naked singularity at finite geodesic distance, unless the cosmological constant  $\Lambda$  (an integration constant in this theory) vanishes. The spacetimes with  $c = 0, \Lambda = 0$  are not asymptotically flat, and can correspond to black holes or naked singularities, depending on the parameter values.

The class of solutions with non-zero integration constants c and  $d = \pm 1/\mu^2$  is the richest. These solutions are given here for the first time in closed form, enabling an analysis of the geometry of these spacetimes. Depending on the signs of c and d, these solutions can correspond to non-asymptotically flat black holes or naked singularities, to non-asymptotically flat traversable wormholes, or to (possibly singularity-free) closed universes. The traversable wormhole solutions occur only for overcharged solutions,  $q^2 > \mu^2$ . We can recall here that overcharged Reissner-Nordström-NUT solutions to the Einstein-Maxwell equations have been shown to correspond to traversable wormholes [23]. However these where actually generated by Misner-string line sources, violating the null energy condition [24]. Remarkably, the NUT-less wormhole solutions of conformal Killing gravity found here do not require the presence of exotic matter sources, the null energy condition being clearly satisfied by the monopole Maxwell configurations (2.9).

We have also examined the inverse problem. Because the field equations of conformal Killing gravity involve only the covariant derivatives of the matter energy-momentum tensor, this cannot be uniquely determined from the knowledge of the spacetime metric. We have found that the most general energy-momentum tensor generating a given static spherically symmetric metric, for instance the Schwarzschild metric, depends on three parameters  $\alpha$ ,  $\beta$  and  $\gamma$ , see equation (4.6). Moreover, all the energy-momentum tensors of (4.6) with  $\gamma = 0$  are physically equivalent, in the sense that they generate all the vacuum ( $\alpha = \beta = \gamma = 0$ ) solutions of section 2.

The existence of singularity-free closed universe vacuum solutions such as the Einstein static universe prompted us to examine the possibility of other sourceless cosmological solutions to this theory. We found again a rich variety of time-dependent FLRW solutions depending on the curvature k, the cosmological constant  $\Lambda$ , and a new integration constant  $\alpha$ , the value  $\alpha = 0$  corresponding to cosmological general relativity. These solutions range from universes with a finite timeline for  $\alpha > 0$  to eternal universes for  $\alpha < 0$ , and from singularity-free cosmologies for  $k \ge 0$  (including oscillating cosmologies for  $\alpha < 0, k \ge 0$ ) to universes with a big bang and/or a big crunch for k < 0.

The FLRW solutions we have found presumably do not exhaust the class of all time-dependent spherically symmetric sourceless solutions to conformal Killing gravity, which could arise from the metric ansatz

$$ds^{2} = -e^{2\nu(r,t)}dt^{2} + e^{2\lambda(r,t)}dr^{2} + r^{2}d\Omega^{2}$$
(6.1)

more general than (5.1). At any rate, it is clear that the Birkhoff theorem is not valid for conformal Killing gravity. We have shown that the Schwarzschild metric is not the unique static spherically symmetric solution to this theory. And there is at least a three-parameter manifold of time-dependent spherically symmetric solutions.

In conclusion, our analysis has shown that the third-order differential field equations of conformal Killing gravity admit a rich variety of solutions. Further investigations into the potentialities of this new theory are necessary in order to ascertain its power as a substitute for general relativity.

# Acknowledgments

One of us (K.N.) thanks the LAPTh Annecy, France, where this work was initiated, for warm hospitality and the University of Jijel, Algeria, for financial support. We are grateful to J. Harada and A. Barnes for valuable comments.

# References

- [1] A. G. Riess et al., Astron. Jour. **116**(1998) 1009.
- [2] S. Perlmutter et al. (Supernova Cosmology Project), Astrophys. Jour. 517 (1999) 565.
- [3] S. Weinberg, "The cosmological constant problem", Rev. Mod. Phys. 61 (1989) 1.
- [4] I. Zlatev, L.-M. Wang and P. J. Steinhardt, "Quintessence, cosmic coincidence, and the cosmological constant", Phys. Rev. Lett. 82 (1999) 896 [arXiv:astro-ph/9807002].
- [5] P. Bull, Y. Akrami, J. Adamek, T. Baker, E. Bellini, J. B. Jimenez, E. Bentivegna, S. Camera, S. Clesse, J. H. Davis, et al., "Beyond ΛCDM: Problems, solutions, and the road ahead", Physics of the Dark Universe 12 (2016) 56–99 [arXiv:1512.05356].
- [6] H. E. Velten, R. vom Marttens and W. Zimdahl, "Aspects of the cosmological coincidence problem", The European Physical Journal C 74 (2014) 1–8 [arXiv:1410.2509].
- [7] T. Clifton, P. G. Ferreira, A. Padilla and C. Skordis, "Modified gravity and cosmology", Physics Reports. 513 (1) (2012) 1–189 [ arXiv:1106.2476v].
- [8] E. N. Saridakis, R. Lazkoz, V. Salzano, P. V. Moniz, S. Capozziello, J. B. Jiménez, M. De Laurentis and G. J. Olmo, "Modified gravity and cosmology", Springer, (2021).
- [9] P. G. Bergmann, "Comments on the scalar tensor theory", Int. J. Theor. Phys. 1 (1968) 25–36.
- [10] R. Kase and S. Tsujikawa, "Dark energy in Horndeski theories after GW170817: A review", Int. J. Mod. Phys. D 28 (2019) no.05, 1942005 [arXiv:1809.08735].
- [11] T. Jacobson, "Einstein-aether gravity": a status report. PoS, QG-PH:020 (2007) 0801.1547.
- [12] T. Jacobson and D. Mattingly, "Gravity with a dynamical preferred frame", Phys. Rev.D 64 (2001) 024028 [arXiv:gr-qc/0007031].
- [13] T. P. Sotiriou and V. Faraoni. "f(R) theories of gravity", Rev. Mod. Phys. 82 (2010) 451 [ arXiv:0805.1726].

- [14] S. Nojiri and S. D. Odintsov, "Unified cosmic history in modified gravity: from F(R) theory to Lorentz non-invariant models," Phys. Rept. **505** (2011), 59-144 [arXiv:1011.0544].
- [15] R. Aldrovandi, and J. G. Pereira, Teleparallel Gravity: An Introduction, Springer Science & Business Media (2012).
- [16] J. Harada, "Gravity at cosmological distances: Explaining the accelerating expansion without dark energy", Phys. Rev. D 108 (2023) 044031[arXiv:2308.02115].
- [17] C.A. Mantica and L.G. Molinari, "Note on Harada's conformal Killing gravity", Phys. Rev. D 108 (2023) 124029 [arXiv:2308.06803].
- [18] J. Harada, "Dark energy in conformal Killing gravity", Phys. Rev. D 108 (2023) 104037 [arXiv:2308.07634].
- [19] J.S.S. Tarciso Junior, F.S.N. Lobo and M.N. Rodrigues, "(Regular) Black holes in conformal Killing gravity coupled to nonlinear electrodynamics and scalar fields", Class. Quantum Grav. 41 (2024) 055012 [arXiv:2310.19508].
- [20] A. Barnes, "Vacuum static spherically symmetric spacetimes in Harada's theory" [arXiv:2309.05336].
- [21] A. Barnes, "Harada-Maxwell static spherically symmetric spacetimes" [arXiv:2311.09171].
- [22] M. S. Morris, K. S. Thorne and U. Yurtsever, "Wormholes, Time Machines, and the Weak Energy Condition," Phys. Rev. Lett. 61 (1988), 1446-1449.
- [23] G. Clément, D. Gal'tsov and M. Guenouche, "NUT wormholes", Phys. Rev. D 93 (2016) 024048 [arXiv:1509.07854].
- [24] G. Clément and D. Gal'tsov, "Rotating traversable wormholes in Einstein-Maxwell theory", Phys. Lett. B 838 (2023) 137677 [arXiv:2210.08913[gr-qc]].