

Quadrupole Moment of a Magnetically Confined Mountain on an Accreting Neutron Star in General Relativity

PEDRO H. B. ROSSETTO ^{1,2} JÖRG FRAUENDIENER ³ AND ANDREW MELATOS ^{1,2}

¹*School of Physics, University of Melbourne, Parkville, VIC 3010, Australia*

²*ARC Centre of Excellence for Gravitational Wave Discovery (OzGrav), University of Melbourne, Parkville, VIC 3010, Australia*

³*Department of Mathematics & Statistics, University of Otago, 730 Cumberland Street, Dunedin 9016, New Zealand*

ABSTRACT

General relativistic corrections are calculated for the quadrupole moment of a magnetically confined mountain on an accreting neutron star. The hydromagnetic structure of the mountain satisfies the general relativistic Grad-Shafranov equation supplemented by the flux-freezing condition of ideal magnetohydrodynamics, as in previous calculations of the magnetic dipole moment. It is found that the ellipticity and hence the gravitational wave strain are 4% to 12% smaller than in the analogous Newtonian system. The direct contribution of the magnetic field to the nonaxisymmetric component of the stress-energy tensor is shown to be negligible in accreting systems such as low-mass X-ray binaries.

Keywords: Accretion — General relativity — Gravitational waves — Gravitational wave sources — LMXBs — Magnetic fields — Magnetohydrodynamics — Neutron stars — Relativistic fluid dynamics

1. INTRODUCTION

Fast-spinning deformed neutron stars are prime candidates for the emission of continuous gravitational waves (CWs) (Riles 2023). The Laser Interferometer Gravitational-Wave Observatory (LIGO), Virgo, and Kamioka Gravitational Wave Detector (KAGRA) routinely conduct targeted narrowband searches for CWs from known pulsars in the LIGO Scientific Collaboration data (Abbott 2022; Abbott et al. 2022a), directed searches in supernova remnants (Abbott et al. 2021) and all-sky searches for unknown CW sources (Abbott et al. 2022b). No CW signal has been found so far. It is important to model from first principles the formation mechanisms of non-axisymmetric stellar deformations, such as ‘mountains’, which are static in the star’s rotating frame, in order to predict the detectability of the associated CW signal. Proposed mechanisms include elastic stresses in the crust (Ushomirsky et al. 2000; Gittins et al. 2020; Gittins & Andersson 2021), single and repeated crustal fracture (Giliberti et al. 2019; Giliberti & Cambiotti 2022; Kerin & Melatos 2022), the Lorentz force in the magnetized interior (Bonazzola & Gourgoulhon 1996; Haskell et al. 2008) and the Lorentz force in the surface layers of polar mountains confined magnetically on accreting neutron stars (Melatos & Phinney 2001; Payne & Melatos 2004; Melatos & Payne 2005; Rossetto et al. 2023), the latter of which are the focus of this paper.

Magnetically confined mountains are formed by the accretion of conducting plasma onto the magnetic polar cap of a neutron star in systems like low-mass X-ray binaries (LMXBs). Several features have been incorporated previously into the theoretical model, including mountain stability (Vigelius & Melatos 2008; Mukherjee & Bhattacharya 2012; Mukherjee et al. 2013), Ohmic and thermal relaxation (Vigelius & Melatos 2009; Suvorov & Melatos 2019), core superconductivity (Passamonti & Lander 2014; Sur & Haskell 2021), mountain sinking (Choudhuri & Konar 2002; Wette et al. 2010), various equations of state (Priymak et al. 2011; Mukherjee 2017), triaxial configurations (Singh et al. 2020), higher magnetic multipole moments and toroidal fields (Suvorov & Melatos 2020; Fujisawa et al. 2022). Recently, Rossetto et al. (2023) formulated the problem of magnetically confined mountains on neutron stars in general relativity. They found that relativistic corrections change the hydromagnetic equilibrium of the accreted matter, diminishing the screening of the magnetic moment three-fold when compared to the Newtonian formulation.

In this paper, we discuss how general relativistic corrections modify the mass quadrupole moment of a magnetically confined mountain and hence affect the emission of gravitational radiation. In Section 2, we summarize the relativistic

model of the system and the numerical method. In Section 3, we calculate the ellipticity of the star, breaking down the contributions from various physical effects. We contrast our findings with the respective Newtonian results. Finally, in Section 4, we estimate the amplitude of the associated gravitational radiation and discuss its detectability.

2. HYDROMAGNETIC EQUILIBRIUM OF A POLAR MOUNTAIN

We employ the same theoretical framework presented by Rossetto et al. (2023). In this section, we review the equations that describe the hydromagnetic structure of a magnetically confined mountain produced by magnetic burial in general relativity. We assume that the neutron star generates a background Schwarzschild metric and that the accreted matter and magnetic field adjust to this curved spacetime. In the usual (t, r, θ, ϕ) coordinates, the metric is given by¹

$$ds^2 = -e^{2\Phi} dt^2 + e^{-2\Phi} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \quad (1)$$

with

$$\Phi(r) = \frac{1}{2} \ln \left(1 - \frac{2M_*}{r} \right), \quad (2)$$

where M_* is the gravitational mass of the neutron star. We assume the fiducial value $M_* = 1.4M_\odot$.

The hydromagnetic equilibrium structure of the mountain is determined by the balance between the Lorentz force, thermal pressure gradient, and gravity. That is, we look for motionless, steady-state configurations satisfying conservation of the energy-momentum tensor, $\nabla_a T^{ab} = 0$. The tensor T^{ab} has contributions from the accreted mass M_a , assumed to be a perfect fluid and a perfect conductor, and the magnetic field of the star $B^a = F^{ab}u_b$, where F_{ab} is the Faraday tensor and u^b is the fluid's four-velocity, with

$$T^{ab} = (e + B^2)u^a u^b + \frac{1}{2}g^{ab}B^2 + (g^{ab} + u^a u^b)p - B^a B^b, \quad (3)$$

where g^{ab} is the inverse metric, e is the energy density of the fluid, p is the pressure, and one has $B^2 = B_a B^a$.

Assuming that the magnetic field of the star has no toroidal component and that the whole system is axisymmetric, the magnetic field can be described by a scalar function $\psi = \psi(r, \theta)$ according to

$$B^a = \frac{1}{r^2 \sin^2 \theta} \epsilon^{abcd} u_d \nabla_b \phi \nabla_c \psi, \quad (4)$$

where ϵ^{abcd} is the completely anti-symmetric tensor. We also assume an isothermal equation of state, given by $p = c_s^2 e$ (Yabushita 1973; Chavanis 2008), with c_s being the sound speed. Additionally, we are interested in a static fluid configuration, in which the four-velocity of the fluid takes the form $u^a = e^{-\Phi} \partial_t^a$. Under these assumptions, $\nabla_a T^{ab} = 0$ reduces to the relativistic isothermal Grad-Shafranov equation

$$\Delta^* \psi = -F'(\psi) \exp[-(1 + c_s^{-2})(\Phi - \Phi_0)], \quad (5)$$

where Δ^* is the relativistic Grad-Shafranov operator given by

$$\Delta^* \psi = \frac{1}{r^2 \sin^2 \theta} \left\{ \frac{\partial}{\partial r} \left[\left(1 - \frac{2M_*}{r} \right) \frac{\partial \psi}{\partial r} \right] + \frac{\sin \theta}{r^2} \frac{\partial}{\partial \theta} \left(\frac{1}{\sin \theta} \frac{\partial \psi}{\partial \theta} \right) \right\}, \quad (6)$$

and $F(\psi)$ is a function related to the fluid pressure at the altitude of the reference potential Φ_0 .

Payne & Melatos (2004) developed a recipe to calculate the function $F(\psi)$ uniquely in the Newtonian theory, by connecting the pre- and post-accretion states through the flux freezing condition of ideal magnetohydrodynamics. Rossetto et al. (2023) adapted the recipe to general relativity, obtaining

$$F(\psi) = \left(\frac{dM}{d\psi} \right)^{1+c_s^2} \left(\frac{2\pi}{c_s^2} \int_C r \sin \theta |\nabla \psi|^{-1} e^{-(\Phi - \Phi_0)/c_s^2} ds \right)^{-(1+c_s^2)}. \quad (7)$$

In (7), $dM/d\psi$ specifies how accreted mass is distributed along flux tubes labelled uniquely by ψ . Its functional form is arbitrary and depends on the unknown details of the accretion process at the disk-magnetosphere boundary.

¹ In this paper we use geometrized units, i.e. $G = c = 1$. For electromagnetic quantities, we use Gaussian units, and we renormalise the magnetic field by a factor of $\sqrt{4\pi}$.




Figure 1. General relativistic versus Newtonian hydromagnetic equilibria for $M_a = 10^{-5} M_\odot$ and $b = \psi_*/\psi_a = 10$. (a) Magnetic field lines (contours of ψ). Solid lines are used for the relativistic solution and dashed for the Newtonian. (b) Residuals of the numerical solver. The dashed red line indicates a tolerance of 10^{-5} .

Qualitatively, however, it should express the fact, that accreting matter is funnelled preferentially onto field lines centered on the magnetic pole. One phenomenological functional form with this property is (Payne & Melatos 2004; Priymak et al. 2011)

$$M(\psi) = \frac{M_a}{2} \frac{1 - e^{-\psi/\psi_a}}{1 - e^{-\psi_*/\psi_a}}, \quad (8)$$

with $\psi_* = \psi(R_*, 0)$, where R_* is the radius of the neutron star and ψ_a is the flux surface that touches the inner edge of the accretion disk.

Numerical solutions to equations (5) and (7) are obtained iteratively. Initially, we choose the dipole ψ_d as a solution and numerically compute the integral in (7). With this value of $F(\psi)$, and thus $F'(\psi)$, we compute the source term of equation (5) and solve for ψ via a successive over-relaxation scheme. We recompute the integral in (7) using the updated solution for ψ , and the scheme proceeds iteratively until the difference between the solutions ψ^n and ψ^{n+1} obtained in two successive steps is less than a specified threshold, viz. $(\psi^n - \psi^{n+1})/\psi^n < 10^{-5}$ over the grid. The iterative nature of the numerical scheme resembles qualitatively, but is not the same as, the true temporal evolution of the system (Mouschovias 1974), in which an initial dipole evolves to a final distorted equilibrium configuration. Figure 1 shows an example of a hydromagnetic equilibrium generated by the solver, for $M_a = 10^{-5} M_\odot$ and $b = \psi_*/\psi_a = 10$. Panel 1(a) shows the general relativistic equilibrium state for the magnetic field lines (ψ level curves) as solid curves, and the Newtonian equivalent as dashed curves. The system reaches a deformed equilibrium state for both theories, but the deformation in the relativistic case is smaller. Panel 1(b) confirms that the numerical ψ residuals decrease with iteration number in a controlled fashion; that is, the numerical solution converges.

3. QUADRUPOLE MOMENT AND ELLIPTICITY

In the linear regime for the generation of gravitational waves (Maggiore 2007), the mass quadrupole source term is proportional to the second time derivative of the mass quadrupole moment, defined by

$$Q_{ij} = \int_{V'} T^{00} \left(x'_i x'_j - \frac{1}{3} x'^k x'_k \delta_{ij} \right) d^3 x', \quad (9)$$

where x'^i is a spatial position vector (assuming a small source), V' is the volume of the source and δ_{ij} is the identity tensor. Note that equation (9) is usually termed the mass quadrupole moment to distinguish it from the current quadrupole moment, which is proportional to the fluid velocity and generates current quadrupole gravitational radiation. However, it is important to recognise that T^{00} in equation (9) includes contributions from the energy density in




Figure 2. Hydromagnetic structure of the general relativistic equilibrium obtained for $M_a = 10^{-5}M_\odot$ and $b = \psi_*/\psi_a = 10$. (a) Rest-mass density ρ . (b) Magnitude of the magnetic field $B = \sqrt{B^2}$.

the magnetic field as well as the mass density of the fluid. We discuss this in Section 3.2. From equation (3), T^{00} is given by

$$T^{00} = e + \frac{B^2}{2}. \quad (10)$$

Furthermore, e can be decomposed as $e = \rho(1 + \varepsilon)$ where ρ is the rest-mass density of the fluid and ε is its specific internal energy. We keep both terms in (10) for the sake of generality and compare their magnitudes below.

In the triaxial model of a rigidly rotating neutron star, the gravitational wave amplitude is given by (Maggiore 2007)

$$h_0 = \frac{16\pi^2 I_0 f^2 \epsilon}{D}, \quad (11)$$

where $I_0 = (2/5)M_* R_*^2$ is the moment of inertia of the star before accretion, f is the rotation frequency, D is the distance to the source and ϵ is the (mass-energy) ellipticity of the star given by

$$\epsilon = \frac{\pi}{I_0} \int_{V'} T^{00} r^4 \sin \theta (3 \cos^2 \theta - 1) dr d\theta. \quad (12)$$

We also define the material and magnetic ellipticities ϵ_e and ϵ_B in terms of the same integral as in (12) but replacing T^{00} with e and $B^2/2$ respectively to give $\epsilon = \epsilon_e + \epsilon_B$. In equilibrium, e and B^2 can be related to the flux-function ψ output from the numerical solver, with

$$e = \frac{F(\psi)}{c_s^2} \exp[-(1 + c_s^{-2})(\Phi - \Phi_0)], \quad (13)$$

$$B^2 = \frac{1}{r^2 \sin^2 \theta} \left[\frac{1}{r^2} \left(\frac{\partial \psi}{\partial \theta} \right)^2 + e^{2\Phi} \left(\frac{\partial \psi}{\partial r} \right)^2 \right]. \quad (14)$$

3.1. Total ellipticity

Figure 2 shows the equilibrium density and magnitude of the magnetic field of the system, plotted in meridional cross-section. We see that the mass distribution is centered at the pole of the star, as expected. In contrast, the magnitude of the magnetic field is highest around the colatitude 25° . This happens because of the flux freezing condition. The frozen-in magnetic field is dragged equatorward with the fluid, as the accreted material spreads under its own weight (Payne & Melatos 2004; Rossetto et al. 2023). Both these configurations indicate a non-zero ellipticity of the star, as per equations (10)–(14).

Figure 3 displays calculations of the total ellipticity ϵ of the star. Panel 3(a) shows the convergence achieved by the numerical solver in both the relativistic and Newtonian scenarios. The ellipticity is plotted as a function of iteration




Figure 3. (a) Numerical convergence for $M_a = 10^{-5} M_\odot$ and $b = \psi_*/\psi_a = 3$: total ellipticity versus iteration number of the numerical scheme. (b) Total ellipticity versus accreted mass in the range $10^{-6} \leq M_a/M_\odot \leq 3 \times 10^{-5}$, for $b = \psi_*/\psi_a = 3$, in the Newtonian and relativistic scenarios. In both panels, the blue and orange curves correspond to the Newtonian and relativistic calculations respectively.

number, not as a function of time; the reader is reminded that the Grad-Shafranov equation applies in the steady state. We observe that ϵ relaxes over ~ 50 iterations for both scenarios, decreasing by $\approx 4\%$ and $\approx 2\%$ during the iterative process in the Newtonian and relativistic cases respectively.

Panel 3(b) shows the converged ellipticity for different values of accreted mass in the range $1 \times 10^{-6} \leq M_a/M_\odot \leq 3 \times 10^{-5}$. The upper limit of this mass range is set by the convergence of the numerical solver. For $M_a \gtrsim 10^{-5} M_\odot$, closed magnetic bubbles form, whose field lines do not connect to the stellar surface, and this phenomenon hinders the convergence of the solution. In the Newtonian regime, this happens for $M_a/M_\odot \gtrsim 10^{-4} b$ (Payne & Melatos 2004) and, in the relativistic regime, for $M_a/M_\odot \gtrsim 3 \times 10^{-4} b$. This difference is consistent with smaller magnetic deformation in the relativistic case. For $1 \times 10^{-6} \leq M_a/M_\odot \leq 3 \times 10^{-5}$, a simple linear regression on the curves shown in Panel 3(b) yields:

$$\epsilon_N = 0.57 M_a/M_\odot + 3 \times 10^{-7}, \quad (15)$$

$$\epsilon_{GR} = 0.54 M_a/M_\odot + 1 \times 10^{-7}, \quad (16)$$

where ϵ_N is the Newtonian ellipticity², displayed in blue in Figure 3, and ϵ_{GR} is the relativistic ellipticity, displayed in orange. Equations (15) and (16) reflect the reduction of the ellipticity in the general relativistic case. The simple regressions (15) and (16) must be modified for $M_a \geq 3 \times 10^{-5} M_\odot$, when the magnetic burial process saturates; cf. Equation (8) of Melatos & Payne (2005). In fractional terms, the relativistic result is $\sim 4\%$ to $\sim 12\%$ smaller than its Newtonian counterpart.

3.2. Purely magnetic ellipticity

The magnetic field contributes directly to the quadrupole moment (9) and the ellipticity (12) of the star through the B^2 term in equation (10). This contribution is distinct from the magnetically induced nonaxisymmetry in the accreted matter, which enters through the term ρ in (10). In the case of magnetically confined mountains, the accreted matter deforms the magnetic field and pushes it equatorwards (Payne & Melatos 2004; Rossetto et al. 2023), increasing the magnetic pressure $B^2/2$ locally. Figure 4 displays how the magnetic ellipticity increases with M_a . We find that the ellipticity generated directly by the magnetic field is of the order 10^{-11} , which is $\sim 10^6$ times smaller than the ellipticity generated by the accreted mass (cf. Figure 3).

Recent studies (Hacyan 2017; Nazari & Roshan 2020; Contopoulos et al. 2023) investigate the gravitational radiation emitted by a purely dipolar bar magnet rotating in vacuo about an axis inclined to its magnetic axis. They find that

² This agrees with Newtonian values in the literature, cf. Figure 2(a) in Melatos & Payne (2005).




Figure 4. Direct contribution of the magnetic field to the ellipticity of the star. Magnetic ellipticity ϵ_B versus accreted mass in the range $10^{-6} \leq M_a/M_\odot \leq 3 \times 10^{-5}$, for $b = \psi_*/\psi_a = 3$, in the Newtonian (blue curve) and relativistic (orange curve) scenarios.

the B^2 contribution is generally low but may be detectable by future-generation gravitational-wave detectors in the case of highly-magnetised fast-spinning neutron stars. In our simulations, we can recover the magnetic dipole solution in vacuo by taking the accreted mass to be low ($M_a \sim 10^{-9} M_\odot$). In the latter regime, we obtain $\epsilon_B \sim 10^{-15}$. This value is consistent with the literature, cf. equation (13) of [Hacyan \(2017\)](#), equation (46) of [Nazari & Roshan \(2020\)](#) for $B \sim 10^{12}$ G, and equation (57) of [Contopoulos et al. \(2023\)](#).

3.3. Compactness

The relativistic corrections to the mountain structure depend on the compactness M_*/R_* of the star. These corrections are greater for more compact objects. Figure 5 shows the effect of the compactness of the star for $M_a = 10^{-5} M_\odot$ and $b = 10$. Panel 5(a) shows the effects of varying M_* in the range $1 \leq M_*/M_\odot \leq 2.5$ with $R_* = 10$ km fixed, i.e. compactness in the range $0.15 \leq M_*/R_* \leq 0.37$. We can see from panel 5(a) that as the mass and compactness grow, the Newtonian and relativistic results deviate. The percentage difference reaches $\sim 30\%$ at $M_*/R_* = 2.50$.

Panel 5(b) shows the effects of varying R_* in the range $10 \text{ km} \leq R_* \leq 15 \text{ km}$ with $M_* = 1.4 M_\odot$ fixed, i.e. compactness in the range $0.20 \leq M_*/R_* \leq 0.31$. Panel 5(b) shows that the difference between the Newtonian and relativistic results ($\sim 4\%$ in fractional terms) stays practically unchanged for different values of compactness. This result indicates, that the compactness influences ϵ mainly through M_* .

4. DISCUSSION

The present work focuses on the modifications arising from including general relativity in models of the formation of magnetically confined mountains and hence the emission of continuous gravitational waves by accreting neutron stars. The baseline for comparison is the Newtonian model calculated by [Melatos & Payne \(2005\)](#) for an isothermal equation of state. The modifications brought by general relativity to the hydromagnetic structure of the mountain are discussed in detail by [Rossetto et al. \(2023\)](#). The main effect is to reduce the ellipticity in the relativistic case, as can be seen in equations (15) and (16).

The characteristic gravitational wave strain h_0 from a star with a static mountain with ellipticity ϵ can be written as

$$h_0 = 4.7 \times 10^{-26} \left(\frac{\epsilon}{10^{-6}} \right) \left(\frac{I_0}{1.1 \times 10^{38} \text{ kg m}^2} \right) \left(\frac{f}{100 \text{ Hz}} \right)^2 \left(\frac{1 \text{ kpc}}{r} \right). \quad (17)$$




Figure 5. Effect of the compactness on the star’s ellipticity for $M_a = 10^{-5}M_\odot$ and $b = 3$. (a) Ellipticity versus neutron star mass in the range $1 \leq M_*/M_\odot \leq 2.5$ for $R_* = 10$ km fixed. (b) Ellipticity versus neutron star radius in the range $10 \text{ km} \leq R_* \leq 15 \text{ km}$ for $M_* = 1.4M_\odot$ fixed. In both panels, the blue and orange curves correspond to the Newtonian and relativistic calculations respectively.

From the ellipticity results shown in Figure 3 we obtain strains between $\sim 10^{-26}$ and $\sim 10^{-25}$. The relativistic results are 4% to 12% smaller than the Newtonian ones, depending on M_a . The purely magnetic contribution from ϵ_B is negligible, generating a wave strain of the order 10^{-32} .

Several important effects already modeled in the Newtonian literature must also be incorporated into the relativistic model. A summary of these effects can be found in Section 1. The unmodeled effect that probably affects the system the most is the choice of the equation of state. In the Newtonian literature (Priymak et al. 2011), a polytropic or piecewise-polytropic equation of state reduced the quadrupole between three and four orders of magnitude compared to an isothermal equation of state. Furthermore, dynamical effects also need to be included. The Grad-Shafranov equation cannot model these effects, as it is an equilibrium equation. Dynamical effects are responsible for the stability of the accreted mountain (Payne & Melatos 2007; Vigelius & Melatos 2008; Mukherjee et al. 2013; Mukherjee 2017).

ACKNOWLEDGEMENTS

This work received financial support from the Australian Research Council Centre of Excellence for Gravitational Wave Discovery (OzGrav), through project number CE170100004. This work was partially supported financially by the Catalyst Fund provided by the New Zealand Ministry of Business, Innovation and Employment and administered by the Royal Society Te Apārangi; and the Division of Science of the University of Otago, New Zealand.

REFERENCES

- Abbott, R. 2022, *The Astrophysical Journal Letters*
- Abbott, R., Abbott, T. D., Abraham, S., et al. 2021, *The Astrophysical Journal*, 921, 80, doi: [10.3847/1538-4357/ac17ea](https://doi.org/10.3847/1538-4357/ac17ea)
- Abbott, R., Abbott, T. D., Acernese, F., et al. 2022a, *The Astrophysical Journal*, 932, 133, doi: [10.3847/1538-4357/ac6ad0](https://doi.org/10.3847/1538-4357/ac6ad0)
- Abbott, R., Abe, H., Acernese, F., et al. 2022b, *Physical Review D*, 106, 102008, doi: [10.1103/PhysRevD.106.102008](https://doi.org/10.1103/PhysRevD.106.102008)
- Bonazzola, S., & Gourgoulhon, E. 1996, *Astronomy and Astrophysics*, 312, 675, doi: [10.48550/arXiv.astro-ph/9602107](https://doi.org/10.48550/arXiv.astro-ph/9602107)
- Chavanis, P. H. 2008, *Astronomy & Astrophysics*, 483, 673, doi: [10.1051/0004-6361:20078287](https://doi.org/10.1051/0004-6361:20078287)
- Choudhuri, A., & Konar, S. 2002, *Monthly Notices of the Royal Astronomical Society*, 332, 933, doi: [10.1046/j.1365-8711.2002.05362.x](https://doi.org/10.1046/j.1365-8711.2002.05362.x)
- Contopoulos, I., Kazanas, D., & Papadopoulos, D. B. 2023, *Monthly Notices of the Royal Astronomical Society*, 527, 11198, doi: [10.1093/mnras/stad3913](https://doi.org/10.1093/mnras/stad3913)

- Fujisawa, K., Kisaka, S., & Kojima, Y. 2022, *Monthly Notices of the Royal Astronomical Society*, 516, 5196, doi: [10.1093/mnras/stac2585](https://doi.org/10.1093/mnras/stac2585)
- Giliberti, E., & Cambiotti, G. 2022, *Monthly Notices of the Royal Astronomical Society*, 511, 3365, doi: [10.1093/mnras/stac245](https://doi.org/10.1093/mnras/stac245)
- Giliberti, E., Cambiotti, G., Antonelli, M., & Pizzochero, P. M. 2019, *Monthly Notices of the Royal Astronomical Society*, stz3099, doi: [10.1093/mnras/stz3099](https://doi.org/10.1093/mnras/stz3099)
- Gittins, F., & Andersson, N. 2021, *Monthly Notices of the Royal Astronomical Society*, 507, 116, doi: [10.1093/mnras/stab2048](https://doi.org/10.1093/mnras/stab2048)
- Gittins, F., Andersson, N., & Jones, D. I. 2020, *Monthly Notices of the Royal Astronomical Society*, 500, 5570, doi: [10.1093/mnras/staa3635](https://doi.org/10.1093/mnras/staa3635)
- Hacyan, S. 2017, *Revista Mexicana de Física*, 3
- Haskell, B., Samuelsson, L., Glampedakis, K., & Andersson, N. 2008, *Monthly Notices of the Royal Astronomical Society*, 385, 531, doi: [10.1111/j.1365-2966.2008.12861.x](https://doi.org/10.1111/j.1365-2966.2008.12861.x)
- Kerin, A. D., & Melatos, A. 2022, *Monthly Notices of the Royal Astronomical Society*, 514, 1628, doi: [10.1093/mnras/stac1351](https://doi.org/10.1093/mnras/stac1351)
- Maggiore, M. 2007, *Gravitational Waves: Volume 1: Theory and Experiments*, 1st edn. (Oxford University Press/Oxford), doi: [10.1093/acprof:oso/9780198570745.001.0001](https://doi.org/10.1093/acprof:oso/9780198570745.001.0001)
- Melatos, A., & Payne, D. J. B. 2005, *The Astrophysical Journal*, 623, 1044, doi: [10.1086/428600](https://doi.org/10.1086/428600)
- Melatos, A., & Phinney, E. S. 2001, *Publications of the Astronomical Society of Australia*, 18, 421, doi: [10.1071/AS01056](https://doi.org/10.1071/AS01056)
- Mouschovias, T. C. 1974, *The Astrophysical Journal*, 192, 37, doi: [10.1086/153032](https://doi.org/10.1086/153032)
- Mukherjee, D. 2017, *Journal of Astrophysics and Astronomy*, 38, 48, doi: [10.1007/s12036-017-9465-6](https://doi.org/10.1007/s12036-017-9465-6)
- Mukherjee, D., & Bhattacharya, D. 2012, *Monthly Notices of the Royal Astronomical Society*, 420, 720, doi: [10.1111/j.1365-2966.2011.20085.x](https://doi.org/10.1111/j.1365-2966.2011.20085.x)
- Mukherjee, D., Bhattacharya, D., & Mignone, A. 2013, *MHD Instabilities in Accretion Mounds on Neutron Star Binaries*, arXiv, doi: [10.48550/ARXIV.1304.7262](https://doi.org/10.48550/ARXIV.1304.7262)
- Nazari, E., & Roshan, M. 2020, *Monthly Notices of the Royal Astronomical Society*, 498, 110, doi: [10.1093/mnras/staa2322](https://doi.org/10.1093/mnras/staa2322)
- Passamonti, A., & Lander, S. K. 2014, *Monthly Notices of the Royal Astronomical Society*, 438, 156, doi: [10.1093/mnras/stt2134](https://doi.org/10.1093/mnras/stt2134)
- Payne, D. J. B., & Melatos, A. 2004, *Monthly Notices of the Royal Astronomical Society*, 351, 569, doi: [10.1111/j.1365-2966.2004.07798.x](https://doi.org/10.1111/j.1365-2966.2004.07798.x)
- . 2007, *Monthly Notices of the Royal Astronomical Society*, 376, 609, doi: [10.1111/j.1365-2966.2007.11451.x](https://doi.org/10.1111/j.1365-2966.2007.11451.x)
- Priymak, M., Melatos, A., & Payne, D. J. B. 2011, *Monthly Notices of the Royal Astronomical Society*, 417, 2696, doi: [10.1111/j.1365-2966.2011.19431.x](https://doi.org/10.1111/j.1365-2966.2011.19431.x)
- Riles, K. 2023, *Living Reviews in Relativity*, 26, 3, doi: [10.1007/s41114-023-00044-3](https://doi.org/10.1007/s41114-023-00044-3)
- Rossetto, P. H. B., Frauendiener, J., Brunet, R., & Melatos, A. 2023, *Monthly Notices of the Royal Astronomical Society*, 526, 2058, doi: [10.1093/mnras/stad2850](https://doi.org/10.1093/mnras/stad2850)
- Singh, N., Haskell, B., Mukherjee, D., & Bulik, T. 2020, *Monthly Notices of the Royal Astronomical Society*, 493, 3866, doi: [10.1093/mnras/staa442](https://doi.org/10.1093/mnras/staa442)
- Sur, A., & Haskell, B. 2021, *Publications of the Astronomical Society of Australia*, 38, e043, doi: [10.1017/pasa.2021.39](https://doi.org/10.1017/pasa.2021.39)
- Suvorov, A. G., & Melatos, A. 2019, *Monthly Notices of the Royal Astronomical Society*, 484, 1079, doi: [10.1093/mnras/sty3518](https://doi.org/10.1093/mnras/sty3518)
- . 2020, *Monthly Notices of the Royal Astronomical Society*, 499, 3243, doi: [10.1093/mnras/staa3132](https://doi.org/10.1093/mnras/staa3132)
- Ushomirsky, G., Cutler, C., & Bildsten, L. 2000, *Monthly Notices of the Royal Astronomical Society*, 319, 902, doi: [10.1046/j.1365-8711.2000.03938.x](https://doi.org/10.1046/j.1365-8711.2000.03938.x)
- Vigelius, M., & Melatos, A. 2008, *Monthly Notices of the Royal Astronomical Society*, 386, 1294, doi: [10.1111/j.1365-2966.2008.13139.x](https://doi.org/10.1111/j.1365-2966.2008.13139.x)
- . 2009, *Monthly Notices of the Royal Astronomical Society*, 395, 1985, doi: [10.1111/j.1365-2966.2009.14698.x](https://doi.org/10.1111/j.1365-2966.2009.14698.x)
- Wette, K., Vigelius, M., & Melatos, A. 2010, *Monthly Notices of the Royal Astronomical Society*, 402, 1099, doi: [10.1111/j.1365-2966.2009.15937.x](https://doi.org/10.1111/j.1365-2966.2009.15937.x)
- Yabushita, S. 1973, *Monthly Notices of the Royal Astronomical Society*, 165, 17, doi: [10.1093/mnras/165.1.17](https://doi.org/10.1093/mnras/165.1.17)