

ORIENTED GRAPHS, FRATTINI-RESISTANCE, AND MAXIMAL PRO- p GALOIS GROUPS

CLAUDIO QUADRELLI

ABSTRACT. Let p be a prime. Following Snopce-Tanushevski, a pro- p group G is called Frattini-resistant if the function $H \mapsto \Phi(H)$, from the poset of all closed finitely-generated subgroups of G into itself, is a poset embedding. We prove that for an oriented right-angled Artin pro- p group (oriented pro- p RAAG) G associated to an oriented graph the following four conditions are equivalent: the associated oriented graph is of elementary type; G is Frattini-resistant; every topologically finitely generated closed subgroup of G is an oriented pro- p RAAG; G is the maximal pro- p Galois group of a field containing a root of 1 of order p . Also, we conjecture that in the \mathbb{Z}/p -cohomology of a Frattini-resistant pro- p group there are no essential triple Massey products.

1. INTRODUCTION

Let p be a prime number. In the paper [25], Ilir Snopce and Slobodan Tanushevski introduced the notion of *Frattini-resistant pro- p group*. Given a pro- p group G , let $\Phi(G)$ denote the Frattini subgroup of G . Then G is said to be Frattini-resistant if the following condition is satisfied: for every couple of topologically finitely generated closed subgroups H_1 and H_2 of G , one has

$$\Phi(H_1) \subseteq \Phi(H_2) \quad \text{if, and only if,} \quad H_1 \subseteq H_2$$

(cf. [25, § 4]). Besides being interesting on this own, this property is particularly relevant in Galois theory, as the *maximal pro- p Galois group* of a field containing a root of 1 of order p is Frattini-resistant — as shown by Snopce-Tanushevski (cf. [25, § 7]), and as we will recall below.

Our aim is to study Frattini-resistance for *oriented right-angled Artin pro- p groups* associated to *oriented graphs*. By an *oriented graph* Γ we mean a pair of sets $\Gamma = (\mathcal{V}, \mathcal{E})$ — we tacitly assume that $\mathcal{V} \cup \mathcal{E} = \emptyset$, and that \mathcal{V} is finite — where \mathcal{V} is the set of *vertices* of Γ , and \mathcal{E} is the set of *edges* of Γ , where

$$\mathcal{E} \subseteq \mathcal{V} \times \mathcal{V} \setminus \{(v, v) \mid v \in \mathcal{V}\}.$$


Given a pair of vertices $v, w \in \mathcal{V}$, if $(w, v) \in \mathcal{E}$ but $(v, w) \notin \mathcal{E}$ the edge (w, v) is said a *special edge*, while if both $(w, v), (v, w)$ lie in \mathcal{E} , we say that (w, v) and (v, w) are *ordinary edges*. A vertex which is the second coordinate of a special edge is said to be

Date: April 2, 2024.

2020 Mathematics Subject Classification. Primary 20E18; Secondary 20J06, 05C25, 12G05, 20E07.

Key words and phrases. Oriented right-angled Artin pro- p groups, Frattini-resistant pro- p groups, maximal pro- p Galois groups, Massey products.

special. For example, the diagrams



represent the same oriented graph with four vertices: in the first diagram every edge is represented as an arrow going from the first coordinate to the second, while in the second diagram we identify the ordinary edges joining the same pair of vertices, and we represent them as a single “headless” edge, so that only the special edges are represented as arrows. Moreover, the special vertex is the black one.

The *oriented right-angled Artin pro- p group* (oriented pro- p RAAG for short) associated to an oriented graph $\Gamma = (\mathcal{V}, \mathcal{E})$ and to a p -power $q = p^f$ with $f \geq 1$ — and $f \geq 2$ if $p = 2$ — is the pro- p group with pro- p presentation

$$G = \left\langle v \in \mathcal{V} \mid wuw^{-1} = \begin{cases} u^{1+q} & \text{if } (w, u) \text{ is special,} \\ u & \text{if } (w, u) \text{ is ordinary,} \end{cases} \quad \forall (u, w) \in \mathcal{E} \right\rangle.$$

The family of oriented pro- p RAAGs is very rich: inside it, one may find all finitely generated free — and free abelian — pro- p groups, pro- p completions of discrete RAAGs, certain families of p -adic analytic pro- p groups and even some finite p -groups (see, e.g., [3, § 1]). Recently, oriented pro- p RAAGs have been object of study in recent times, especially from a Galois-theoretic perspective (see, e.g., [2–4, 22, 27]). Our main goal is to characterize the oriented pro- p RAAGs which are Frattini-resistant, in terms of the associated oriented graph.

Theorem 1.1. *Let $\Gamma = (\mathcal{V}, \mathcal{E})$ be an oriented graph, and for $q = p^f$ (with $f \geq 2$ if $p = 2$) let G be the oriented pro- p RAAG associated to Γ and to q . The following are equivalent.*


- (i) Γ is an oriented graph of elementary type.
- (ii) G is Frattini-resistant.

An oriented graph $\Gamma = (\mathcal{V}, \mathcal{E})$ is said to be of *elementary type* if it satisfies the following conditions: every edge joining a special vertex and another vertex is an arrow pointing at the special vertex — roughly speaking, every special vertex is a “sinkhole” (for example, the only special vertex, v_1 , in the oriented graph (1.1), is a sinkhole); and for every induced subgraph Γ' of Γ , either

- (a) Γ' has at least two distinct connected components,
- (b) or Γ' has an ordinary vertex which is adjacent to all other vertices of Γ' .

For example, the oriented graph represented in (1.1) is not of elementary type, as the induced subgraph with vertices v_1, v_2, v_4 satisfies none of the conditions (a)–(b) above.

Also, consider the two oriented graphs with geometric representations



The first oriented graph is of elementary type, while the second is not as the special vertex v_1 is not a sinkhole. Oriented graphs of elementary type may be constructed starting from single vertices as “bricks”, and performing elementary operations, namely disjoint unions and “coning” (see [3, § 2.4] and § 3.2 below).

From Theorem 1.1 we may deduce some interesting corollaries. The first one may be seen as the translation, in terms of Frattini-resistance, of the “oriented pro- p version” of a famous result of C. Droms on discrete RAAGs (see [6]).

Corollary 1.2. *Let $\Gamma = (\mathcal{V}, \mathcal{E})$ be an oriented graph, and for $q = p^f$ (with $f \geq 2$ if $p = 2$) let G be the oriented pro- p RAAG associated to Γ and to q . The following are equivalent.*

- (i) *G is Frattini-resistant.*
- (ii) *Every topologically finitely generated closed subgroup of G occurs as an oriented pro- p RAAG associated to some oriented graph.*

The proof of Corollary 1.2 relies on the characterization of oriented pro- p RAAGs satisfying “Drom’s condition” (ii) proved by S. Blumer, Th.S. Weigel and the author in [3].

The second corollary we obtain from Theorem 1.1 involves the realizability of an oriented pro- p RAAG as the *maximal pro- p Galois group* of a field. Given a field \mathbb{K} , its maximal pro- p Galois group $G_{\mathbb{K}}(p)$ is the Galois group of the maximal pro- p Galois extension of \mathbb{K} — or, equivalently, the maximal pro- p quotient of the absolute Galois group of \mathbb{K} . Characterizing those pro- p groups which occur as the maximal pro- p Galois group of a field is one of the main open problems in Galois theory (see, e.g., [1, 3, 19, 20]).

Recall that a pro- p group G is said to complete into a *1-cyclotomic oriented pro- p group* if there exists a continuous G -module M , isomorphic to \mathbb{Z}_p as an abelian pro- p group, such that for every closed subgroup H of G , the natural cohomology maps

$$(1.3) \quad H^1(H, M/p^n M) \longrightarrow H^1(H, M/pM),$$

induced by the epimorphism of continuous H -modules $M/p^n M \rightarrow M/pM$, is surjective for every $n \geq 1$ — here we consider the continuous G -modules $M/p^n M$ as continuous H -modules in the obvious way —, see, e.g., [20, § 1]. It is well-known that the maximal pro- p Galois group of a field containing a root of 1 of order p , together with the *pro- p cyclotomic character*, completes into a 1-cyclotomic oriented pro- p group (see Example 2.9 below).

By [3, Thm. 1.1], an oriented pro- p RAAG completes into a 1-cyclotomic oriented pro- p group if, and only if, the associated oriented graph is of elementary type. Also, Snopce-Tanushevski proved that a pro- p group which completes into a 1-cyclotomic oriented pro- p group is Frattini-resistant (see [25, Thm. 1.11]). Hence, combining these two results with Theorem 1.1 yields the following.

Corollary 1.3. *Let $\Gamma = (\mathcal{V}, \mathcal{E})$ be an oriented graph, and for $q = p^f$ (with $f \geq 2$ if $p = 2$) let G be the oriented pro- p RAAG associated to Γ and to q . The following are equivalent.*

- (i) *G is Frattini-resistant.*
- (ii) *G completes into a 1-cyclotomic oriented pro- p group.*
- (iii) *G occurs as the maximal pro- p Galois group of a field containing a root of 1 of order p .*

Corollary 1.3 suggests that Frattini-resistance is a rather restrictive property for pro- p groups, and it may provide a powerful tool to study maximal pro- p Galois groups of fields. In particular, we believe that further investigations in this direction will lead to the discovery of new obstructions for the realization of pro- p groups as maximal pro- p Galois group.

Another cohomological property of maximal pro- p Galois groups which has been thoroughly studied in recent years is the presence of *essential Massey products* in the \mathbb{Z}/p -cohomology of these pro- p groups. Massey products in the \mathbb{Z}/p -cohomology of a pro- p group G are multi-valued maps which associate a sequence of elements of $H^1(G, \mathbb{Z}/p)$ to a subset of $H^2(G, \mathbb{Z}/p)$, and they generalize the cup-product: for an overview on Massey products in the \mathbb{Z}/p -cohomology of maximal pro- p Galois groups see, e.g., [16, 17] and references therein. E. Matzri proved that if \mathbb{K} is a field containing a root of 1 of order p , then there are no essential triple Massey products in the \mathbb{Z}/p -cohomology of $G_{\mathbb{K}}(p)$ — see [10, 13] —, i.e., whenever the subset of $H^2(G_{\mathbb{K}}(p), \mathbb{Z}/p)$ which is the value of a Massey product of a sequence of length 3 of elements of $H^1(G_{\mathbb{K}}(p), \mathbb{Z}/p)$ is not empty, it contains 0. Employing a result of W. Dwyer, it is possible to translate this property into purely group-theoretic terms, see § 7.1 below. This property was used to produce new examples of pro- p groups that do not occur as maximal pro- p Galois groups of fields containing a root of 1 of order p (see [17, § 7]).

We suspect that Frattini-resistance is strictly stronger than this cohomological property.

Conjecture 1.4. *Let G be a Frattini-resistant pro- p group. Then there are no essential triple Massey products in the \mathbb{Z}/p -cohomology of G .*

In § 7.2 we formulate Conjecture 1.4 in group-theoretic terms.

2. FRATTINI-RESISTANCE AND 1-CYCLOTOMICITY

2.1. Frattini-resistant pro- p groups. Let G be a pro- p group. From now on, every subgroup of G will be tacitly assumed to be closed with respect to the pro- p topology, and set of generators, and presentations, of pro- p groups will be intended in the topological sense.

Given two elements $x, y \in G$, we adopt the notation

$${}^x y = x \cdot y \cdot x^{-1} \quad \text{and} \quad [x, y] = {}^x y \cdot y^{-1} = xyx^{-1}y^{-1}.$$

Given a positive integer n , one has the normal subgroups

$$G^{p^n} = \langle x^{p^n} \mid x \in G \rangle \quad \text{and} \quad G' = \langle [x, y] \mid x, y \in G \rangle.$$

Finally, the *Frattini subgroup* of G is $\Phi(G) = G^p \cdot G'$. The quotient $G/\Phi(G)$ is a \mathbb{Z}/p -vector space, and a basis of this quotient yields a minimal generating set of G (cf., e.g., [5, Prop. 1.9]). We remark that if G is a finitely generated pro- p group, then also $\Phi(G)$ is finitely generated, and the dimension of $G/\Phi(G)$ is finite (cf., e.g., [5, Prop. 1.14 and Thm. 1.17]).

Thus, one may formulate the definition of Frattini-resistant pro- p group as follows: G is Frattini-resistant if the assignment $H \mapsto \Phi(H)$ yields a homomorphism of partially ordered sets from the partially ordered set of finitely generated subgroups of G into itself (cf. [26, Def. 1.1]). Another notion introduced by Snopce-Tanushevski, tightly related to Frattini-resistance, is the following (cf. [25, Def. 1.1]).

Definition 2.1. A pro- p group G is said to be *Frattini-injective* if, given two finitely generated subgroups H_1, H_2 of G , $\Phi(H_1) = \Phi(H_2)$ implies that $H_1 = H_2$.

It is easy to see that if G is Frattini-resistant, then it is also Frattini-injective (cf. [25, Cor. 4.3]). A first, easily checked, property enjoyed by Frattini-resistant (and Frattini-injective) pro- p groups is that they are torsion-free (cf. [25, § 1]).

Example 2.2. Let G be a pro- p group, and suppose that $g \neq 1$ is an element of G yielding non-trivial torsion, i.e., $g^{p^k} = 1$ and $g^{p^{k-1}} \neq 1$ for some $k \geq 1$. Then

$$1 = g^{p^k} = \left(g^{p^{k-1}}\right)^p \in \Phi(\{1\}) = \{1\} \quad \text{but} \quad 1 \neq g^{p^{k-1}} \notin \{1\},$$

and therefore G is not Frattini-resistant. Also, it is not Frattini-injective, as $\Phi(\langle g^{p^{k-1}} \rangle)$ and $\Phi(\{1\})$ are equal.

One has the following handy characterization of Frattini-resistant pro- p groups (cf. [26, Prop. 2.1]). For the reader's convenience, we recall briefly its proof, as presented by Snopce-Tanushevski.

Proposition 2.3. *Let G be a pro- p group. The following are equivalent:*

- (i) G is Frattini-resistant;
- (ii) if an element $x \in G$ satisfies $x^p \in \Phi(H)$ for some finitely generated subgroup $H \subseteq G$, then $x \in H$.

Proof. Suppose that G is Frattini-resistant, and let $x \in G$ and $H \subseteq G$ be such that $x^p \in \Phi(H)$. Then $\Phi(\langle x \rangle) = \langle x^p \rangle \subseteq \Phi(H)$, and hence $\langle x \rangle \subseteq H$.

Conversely, suppose that condition (ii) is satisfied, and let H_1, H_2 be two subgroups of G such that $\Phi(H_1) \subseteq \Phi(H_2)$. Then for every $x \in H_1$ one has $x^p \in \Phi(H_1) \subseteq \Phi(H_2)$, and therefore $x \in H_2$ by condition (ii). Thus $H_1 \subseteq H_2$. \square

Example 2.4. A free abelian pro- p group G is Frattini-resistant. Indeed, for any finitely generated subgroup $H \subseteq G$, one has

$$\Phi(H) = H^p = \{ h^p \mid h \in H \},$$

and hence if $g^p \in H$, for $g \in G$, then $g^p = h^p$ for some $h \in H$. Thus, $1 = g^p h^{-p} = (gh^{-1})^p$, which implies that $g = h$ as G is torsion-free.

Example 2.5. The *Heisenberg pro- p group*

$$\begin{aligned} G &= \langle x, y, z \mid [x, y] = z, [x, z] = [y, z] = 1 \rangle \\ &= \langle x, y \mid [x, [x, y]] = [[y, [x, y]] = 1 \rangle \end{aligned}$$

is not Frattini-resistant. Indeed, let H be the subgroup of G generated by x, y^p . Then $[x, y^p] = z^p$, and

$$H = \langle x, y^p, z^p \mid [x, y^p] = z^p, [x, z^p] = [y^p, z^p] = 1 \rangle,$$

so that $z^p \in \Phi(H)$, but $z \notin H$.

One may employ direct products with free abelian pro- p groups to produce new Frattini-resistant pro- p groups, but only under certain restrictions, as proved by Snopce-Tanushevski in [26, Thm. A] (we recall that a pro- p group G is said to be *absolutely torsion-free* if for every subgroup H , the abelianization H/H' is a free abelian pro- p group, cf. [28]).

Proposition 2.6. *Let G_1, G_2 be two non-trivial pro- p groups, and set $G = G_1 \times G_2$. Then G is Frattini-resistant if, and only if, and only if, both G_1, G_2 are absolutely torsion free, and at least one of the two factors is a free abelian pro- p group.*

2.2. 1-cyclotomic oriented pro- p groups. Let $1 + p\mathbb{Z}_p$ denote the multiplicative group of principal units of the ring of p -adic integers \mathbb{Z}_p — i.e.,

$$1 + p\mathbb{Z}_p = \{ 1 + p\lambda \mid \lambda \in \mathbb{Z}_p \}.$$

If $p \neq 2$ then $1 + p\mathbb{Z}_p \simeq \mathbb{Z}_p$ as an abelian pro- p group.

Given a pro- p group G , an *orientation* of G is a homomorphism of pro- p groups $\theta: G \rightarrow 1 + p\mathbb{Z}_p$, and the pair (G, θ) is called an *oriented pro- p group* (cf. [23]; oriented pro- p groups were introduced by I. Efrat in [8] with the name “cyclotomic pro- p pairs”). An oriented pro- p group (G, θ) comes endowed with a canonical continuous G -module $\mathbb{Z}_p(\theta)$, which is isomorphic to \mathbb{Z}_p as an abelian pro- p group, with the action given by

$$g \cdot \lambda = \theta(g) \cdot \lambda \quad \text{for every } g \in G, \lambda \in \mathbb{Z}_p(\theta).$$

Conversely, if G is a pro- p group and M is a continuous G -module, isomorphic to \mathbb{Z}_p as an abelian pro- p group, then the action of G on M induces an orientation $\theta: G \rightarrow 1 + p\mathbb{Z}_p$ by $\theta(g)m = g \cdot m$ for every $m \in M$, and $M \simeq \mathbb{Z}_p(\theta)$ as a continuous G -module.

Now consider the epimorphisms of continuous G -modules

$$\mathbb{Z}_p(\theta)/p^n\mathbb{Z}_p(\theta) \twoheadrightarrow \mathbb{Z}_p(\theta)/p\mathbb{Z}_p(\theta),$$

induced by the canonical projections $\mathbb{Z}/p^n \twoheadrightarrow \mathbb{Z}/p$, for every $n \geq 1$ — observe that the G -module $\mathbb{Z}_p(\theta)/p\mathbb{Z}_p(\theta)$ is isomorphic to the trivial G -module \mathbb{Z}/p . These epimorphisms induce in cohomology the natural maps

$$(2.1) \quad \mathrm{H}^1(G, \mathbb{Z}_p(\theta)/p^n\mathbb{Z}_p(\theta)) \longrightarrow \mathrm{H}^1(G, \mathbb{Z}/p).$$

An oriented pro- p group (G, θ) is said to be *1-cyclotomic* if the maps (2.1) are surjective for every $n \geq 1$, and also for every subgroup H of G , replacing G with H and $\mathbb{Z}_p(\theta)$ with $\mathbb{Z}_p(\theta|_H)$ in (2.1).

Remark 2.7. The cohomology group of degree 1 $\mathrm{H}^1(G, \mathbb{Z}/p)$ is the group of homomorphisms of pro- p groups $G \rightarrow \mathbb{Z}/p$. Hence, one has an isomorphism of discrete \mathbb{Z}/p -vector spaces $\mathrm{H}^1(G, \mathbb{Z}/p) \simeq (G/\Phi(G))^*$, where the latter is the \mathbb{Z}/p -dual of $G/\Phi(G)$ (cf., e.g., [24, Chap. I, § 4.2]).

We say that a pro- p group G may complete into a 1-cyclotomic oriented pro- p group if there exists an orientation $\theta: G \rightarrow 1 + p\mathbb{Z}_p$ such that the oriented pro- p group (G, θ) is 1-cyclotomic. In [25, Thm. 1.11], I. Snopce and S. Tanushevski prove that pro- p groups which may complete into 1-cyclotomic oriented pro- p groups are Frattini-resistant (with a condition if $p = 2$).

Proposition 2.8. *Let (G, θ) be a 1-cyclotomic torsion-free oriented pro- p group, and suppose that $\text{Im}(\theta) \subseteq 1 + 4\mathbb{Z}_2$ if $p = 2$. Then G is strongly Frattini-resistant.*

One of the most relevant examples of 1-cyclotomic oriented pro- p groups — and of Frattini-resistant pro- p groups — is provided by Galois theory (cf. [11, § 4] and [20, § 2.4]).

Example 2.9. Let \mathbb{K} be a field containing a root of 1 of order p (and also $\sqrt{-1}$ if $p = 2$), and consider its maximal pro- p Galois group $G_{\mathbb{K}}(p)$. The pro- p cyclotomic character $\theta_{\mathbb{K}}$ of $G_{\mathbb{K}}(p)$ is the orientation $\theta_{\mathbb{K}}: G_{\mathbb{K}}(p) \rightarrow 1 + p\mathbb{Z}_p$ satisfying

$$g(\zeta) = \zeta^{\theta_{\mathbb{K}}(g)} \quad \text{for every } g \in G \text{ and } \zeta \in \bar{\mathbb{K}}_s^\times \text{ of order } p^k$$

for some $k \geq 1$ (here $\bar{\mathbb{K}}_s$ denotes the separable closure of \mathbb{K}). Since we are assuming that $\sqrt{-1} \in \mathbb{K}$ if $p = 2$, then in this case $\text{Im}(\theta_{\mathbb{K}}) \subseteq 1 + 4\mathbb{Z}_2$. The oriented pro- p group $(G_{\mathbb{K}}(p), \theta_{\mathbb{K}})$ is 1-cyclotomic, and therefore the maximal pro- p Galois group $G_{\mathbb{K}}(p)$ is Frattini-resistant.

3. ORIENTED GRAPHS

3.1. Oriented graphs and specially oriented graphs. Let $\Gamma = (\mathcal{V}, \mathcal{E})$ be an oriented graph. Recall that the vertices that are the second coordinate of a special edge are special (and we represent them with black bullets); conversely, the vertices that are not special are called ordinary vertices (and we represent them with white bullets). One has the following notions for oriented graphs.

- (a) An *induced subgraph* of Γ is an oriented graph $\Gamma' = (\mathcal{V}', \mathcal{E}')$ such that $\mathcal{V}' \subseteq \mathcal{V}$, and $\mathcal{E}' = \mathcal{E} \cap (\mathcal{V}' \times \mathcal{V}')$; and moreover, a vertex $v \in \mathcal{V}'$ is special, respectively ordinary, whenever it is a special, respectively ordinary, vertex of Γ .
- (b) A special vertex $w \in \mathcal{V}$ is called a *sinkhole* if (u, w) is a special edge of Γ whenever $u \in \mathcal{V}$ is another vertex which is joined to w .
- (c) Γ is *specially oriented* if every special vertex is a sinkhole.

For example, the oriented graph represented in (1.1) and the left-side oriented graph in (1.2) are specially oriented, while the right-side oriented graph in (1.2) is not, as v_1 is not a sinkhole.

Remark 3.1. Henceforth, if $\Gamma = (\mathcal{V}, \mathcal{E})$ is an oriented graph and $(u, v), (v, u) \in \mathcal{E}$ are ordinary edges, then we will identify them, and we will say that u, v are joined by a single ordinary edge, with an abuse of notation. Moreover, we will call oriented graphs without special edges “unoriented graphs”.

It is straightforward to see that an oriented graph is specially oriented if, and only if, none of the following occurs:

$$(3.1) \quad \begin{array}{ccc} \textcircled{*} & \xrightarrow{x} & \bullet \\ & \text{---} & \bullet \\ z & \text{---} & y \end{array} \quad \text{or} \quad \begin{array}{ccc} \textcircled{*} & \xrightarrow{x} & \textcircled{*} \\ & \text{---} & \textcircled{*} \\ z & \text{---} & y \end{array}$$

— no matter whether the bottom vertices are joined or not (here we use $\textcircled{*}$ to represent vertices which are not necessarily ordinary nor special) — cf. [3, § 2.3] or [21, § 2.2]. Altogether, all possible cases of induced subgraphs with three vertices which prevent an oriented graph to be specially oriented are seven: namely, for each of the two representations in (3.1) one has four possible cases — z, y are disjoint; or joined by an ordinary edge; or joined by a special edge, either (z, y) or (y, z) —, and the two representations

$$\begin{array}{ccc} \circ & \xrightarrow{x} & \bullet \\ & \text{---} & \bullet \\ z & \text{---} & y \end{array} \quad \text{and} \quad \begin{array}{ccc} \bullet & \xrightarrow{x} & \circ \\ & \text{---} & \circ \\ z & \text{---} & y \end{array}$$

yield the same case, with the “roles” of the vertices x, y, z permuted ciclically (see § 5 below).

3.2. Oriented graphs of elementary type. One has the following two operations with oriented graphs.

- (a) Given two oriented graphs $\Gamma_1 = (\mathcal{V}_1, \mathcal{E}_1)$ and $\Gamma_2 = (\mathcal{V}_2, \mathcal{E}_2)$, their *disjoint union* is the oriented graph $\Gamma = (\mathcal{V}, \mathcal{E})$ with

$$\mathcal{V} = \mathcal{V}_1 \dot{\cup} \mathcal{V}_2 \quad \text{and} \quad \mathcal{E} = \mathcal{E}_1 \dot{\cup} \mathcal{E}_2.$$

- (b) Given an oriented graph $\Gamma = (\mathcal{V}, \mathcal{E})$, the *cone* $\nabla(\Gamma) = (\mathcal{V}_{\nabla(\Gamma)}, \mathcal{E}_{\nabla(\Gamma)})$ of Γ is the oriented obtained by adding a “new” ordinary vertex to Γ , and joining it with all ordinary vertices of Γ , via ordinary edges, and with all special vertices of Γ via special edges: namely,


$$\mathcal{V}_{\nabla(\Gamma)} = \{u\} \dot{\cup} \mathcal{V},$$

$$\mathcal{E}_{\nabla(\Gamma)} = \{(u, v), (v, u), (u, w) \mid v, w \in \mathcal{V}, v \text{ ordinary}, w \text{ special}\} \dot{\cup} \mathcal{E},$$

where u is the “new” vertex (which we call the tip of the cone).


Definition 3.2. An oriented graph $\Gamma = (\mathcal{V}, \mathcal{E})$ is of *elementary type* if it may be obtained by iterating disjoint union and cones starting from a subset \mathcal{V}_0 of \mathcal{V} .

Example 3.3. Let $\Gamma = (\mathcal{V}, \mathcal{E})$ be the left-side oriented graph in (1.2). Then Γ is of elementary type, and it may be constructed as follows: we start with the disjoint vertices v_1 and v_3 , which are special, then we make the cone with the ordinary vertex v_3 as the tip, and finally we make the cone of the resulting oriented graph with the ordinary vertex v_4 as tip.




In this case $\mathcal{V}_0 = \{v_1, v_3\}$.

Example 3.4. Consider the oriented graph $\Gamma = (\mathcal{V}, \mathcal{E})$ with geometric realization



Then Γ is of elementary type, and we construct it as follows: we start with $\mathcal{V}_0 = \{v_2, v_4, v_6\}$, first we make separately the cones $\nabla(\{v_4\}, \emptyset)$ and $\nabla(\{v_6\}, \emptyset)$ with tips respectively v_1 and v_3 , then we take the disjoint union of the two cones and of $(\{v_2\}, \emptyset)$, finally we make the cone of the disjoint union with tip v_5 .



The definition of oriented graph of elementary type given in the Introduction is equivalent to Definition 3.2.

Proposition 3.5. *An oriented graph $\Gamma = (\mathcal{V}, \mathcal{E})$ satisfies Definition 3.2 if, and only if, it is specially oriented and for every induced subgraph Γ' of Γ , either*

- (a) Γ' has at least two distinct connected components,
- (b) or Γ' has an ordinary vertex which is adjacent to all other vertices of Γ' .

Proof. First, suppose that Γ satisfies Definition 3.2. Since all new vertices added with the procedure described in the definitions are the tips of the cones, and thus ordinary vertices, all special vertices of Γ are in the starting set \mathcal{V}_0 . Moreover, these vertices are disjoint, and whenever we add the tip u of a cone, the edge joining u the special vertices in \mathcal{V}_0 are special edges with u as first coordinate. Hence, every special vertex of Γ is a sinkhole, and thus Γ is specially oriented.

Now let $\Gamma' = (\mathcal{V}', \mathcal{E}')$ be a proper induced subgraph of Γ . We proceed by induction on the number of vertices of Γ . Since Γ satisfies Definition 3.2, either $\Gamma = \Gamma_1 \dot{\cup} \Gamma_2$ for two proper subgraphs Γ_1, Γ_2 , or $\Gamma = \nabla(\bar{\Gamma})$ for some proper subgraph $\bar{\Gamma}$ of Γ . Clearly, in both cases the oriented graphs Γ_1, Γ_2 , and $\bar{\Gamma}$, satisfy Definition 3.2, and they have less vertices than Γ : therefore, by induction their induced subgraphs satisfy one of the two conditions (a)–(b).

- If $\Gamma = \Gamma_1 \dot{\cup} \Gamma_2$, either Γ' is the disjoint union of two non-trivial induced subgraphs respectively of Γ_1 and Γ_2 — and hence Γ' satisfies condition (a) —, or Γ' is an induced subgraph of Γ_i with $i \in \{1, 2\}$ — and hence it satisfies one of the two conditions (a)–(b) by induction;
- if $\Gamma = \nabla(\bar{\Gamma})$ with tip u , either $u \in \mathcal{V}'$ — and hence Γ' satisfies condition (b) with the ordinary vertex u —, or Γ' is an induced subgraph of $\bar{\Gamma}$ — and hence it satisfies one of the two conditions (a)–(b) by induction.

- (b) If Γ is unoriented, then every associated oriented pro- p RAAG does not depend on the choice of q , and it is isomorphic to the pro- p RAAG (i.e., the pro- p completion of the discrete RAAG) associated to Γ considered as a simplicial graph.

Lemma 4.2. *Let $\Gamma = (\mathcal{V}, \mathcal{E})$ be an oriented graph, and for some p -power q let G be the associated oriented pro- p RAAG.*


- (i) *If a vertex $v \in \mathcal{V}$ is an element of G yielding non-trivial torsion, then Γ is not specially oriented.*
 (ii) *If Γ is specially oriented, $v_1, v_2 \in \mathcal{V}$ are disjoint vertices, and there exists a third vertex $w \in \mathcal{V}$ which is joint to both v_1, v_2 , then the subgroup of G generated by v_1, v_2 is a 2-generated free pro- p group.*

Proof. (i) Suppose that Γ is specially oriented, and let $\Gamma_v = (\{v\}, \emptyset)$ be the induced subgraph of Γ whose only vertex is v . The oriented pro- p RAAG associated to Γ_v is the free pro- p -cyclic group $\langle v \rangle \simeq \mathbb{Z}_p$. By [3, Prop. 4.11], the inclusion $\{v\} \hookrightarrow \mathcal{V}$ induces a monomorphism of pro- p groups $\langle v \rangle \hookrightarrow G$, and therefore v is a torsion-free element of G .

- (ii) This statement is [3, Lemma 6.4]. □

4.2. Oriented pro- p RAAGs and torsion. The following examples, dealing with oriented pro- p RAAGs associated to oriented graphs that are not specially oriented, will be useful for the proof of Theorem 1.1.

Example 4.3. Let $\Gamma = (\mathcal{V}, \mathcal{E})$ be the non-specially oriented graph with geometric representation




The oriented pro- p RAAG associated to Γ and to a p -power q is

$$G = \langle x, y, z \mid [x, y] = y^q, [y, z] = z^q, [z, x] = x^q, \rangle,$$

which is a finite p -group, as shown by J. Mennike (cf. [24, Ch. I, § 4.4, Ex. 2(e)]). In particular, the oriented pro- p RAAG associated to a single vertex (which is isomorphic to \mathbb{Z}_p) is not subgroup of G (cf. [3, Ex. 4.7]).

Example 4.4. Let $\Gamma_1 = (\mathcal{V}, \mathcal{E}_1)$ and $\Gamma_2 = (\mathcal{V}, \mathcal{E}_2)$ be the non-specially oriented graph with geometric representation respectively



Given a p -power q , the oriented pro- p RAAG associated to q and to one of the two graphs is

$$G = \langle x, y, z \mid [x, y] = y^q, [y, z] = z^q, [x, z] = z^{\epsilon q} \rangle,$$

where $\epsilon = 1$ if G is associated to Γ_1 , and $\epsilon = 0$ if G is associated to Γ_2 . Then $[y^q, z] = z^{(1+q)^q - 1}$; on the other hand, one computes

$$\begin{aligned} [[y^q, z]] &= [xyx^{-1}y^{-1}, z] = x \left(y \left(x^{-1} \left(y^{-1}zy \right) x \right) y^{-1} \right) x^{-1} \cdot z^{-1} \\ &= \left(\left(\left(z^{(1+q)^{-1}} \right)^{\epsilon(1+q)^{-1}} \right)^{1+q} \right)^{\epsilon(1+q)} \cdot z^{-1} \\ &= z^1 \cdot z^{-1} = 1. \end{aligned}$$

Since $(1+q)^q - 1 = (1+q^2 + q^3(q-1)/2 + \dots) - 1 \neq 0$, in both cases z yields non-trivial torsion. In particular, the oriented pro- p RAAG associated to the oriented graph $\Gamma' = (\{z\}, \emptyset)$ is not a subgroup of G .

By Example 2.2, the oriented pro- p RAAGs of the above examples are not Frattini-injective nor Frattini-resistant.

Remark 4.5. One could prove that the oriented pro- p RAAGs in Examples 4.3–4.4 are not Frattini-injective using [25, Thm. 1.2], as these pro- p groups are *p -adic analytic* (cf., e.g., [25, § 3]).

4.3. Oriented pro- p RAAGs of elementary type. In analogy with Proposition 3.6, one may characterize oriented pro- p RAAGs associated to oriented graphs of elementary type in terms of subgroups (cf. [3, Prop. 6.3]).

Proposition 4.6. *Let $\Gamma = (\mathcal{V}, \mathcal{E})$ be a specially oriented graph, and for some p -power q let G be the associated oriented pro- p RAAG. Then Γ is of elementary type if, and only if, G has no subgroups isomorphic to the oriented pro- p RAAGs associated to an oriented graph with geometric realization as in (3.3), or*

$$(4.1) \quad \begin{array}{ccc} x & y & \text{or} & x & y & z & w \\ \circ & \text{---} & \circ & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \\ \mid & & \mid & & & & & & & \\ \circ & \text{---} & \circ & & & & & & & \\ w & & z & & & & & & & \end{array}$$

The main theorem of [3] states the following.

Theorem 4.7. *Let $\Gamma = (\mathcal{V}, \mathcal{E})$ be an oriented graph, and for $q = p^f$ (with $f \geq 2$ if $p = 2$) let G be the oriented pro- p RAAG associated to Γ and to q . The following are equivalent.*

- (i) Γ is an oriented graph of elementary type.
- (ii) Every finitely generated subgroup of G occurs as an oriented pro- p RAAG associated to some oriented graph.
- (iii) G may complete into a 1-cyclotomic oriented pro- p group.
- (iv) G occurs as the maximal pro- p Galois group of a field containing a root of 1 of order p , and θ_Γ coincides with the pro- p cyclotomic character.

Hence, from Theorem 4.7 and from Proposition 2.8, one deduces the implication (i) \Rightarrow (ii) of Theorem 1.1.

Corollary 4.8. *Let $\Gamma = (\mathcal{V}, \mathcal{E})$ be an oriented graph of elementary type, and for $q = p^f$ (with $f \geq 2$ if $p = 2$) let G be the oriented pro- p RAAG associated to Γ and to q . Then G is Frattini-resistant.*

Also, Corollary 1.2 follows by Theorem 1.1 and the equivalence (i) \Leftrightarrow (ii) of Theorem 4.7, while Corollary 1.3 follows by Theorem 1.1 and the equivalence (i) \Leftrightarrow (iii) \Leftrightarrow (iv) of Theorem 4.7.


To prove the implication (ii) \Rightarrow (i) of Theorem 1.1, we will proceed with a case-by-case analysis, showing — in the next two sections — that every induced subgraph which prevents an oriented graph from being of elementary type gives rise to pro- p groups that are not Frattini-resistant.

5. NON-SPECIALY ORIENTED GRAPHS AND FRATTINI-RESISTANCE


The goal of this section is to prove the following.

Proposition 5.1. *Let $\Gamma = (\mathcal{V}, \mathcal{E})$ be an oriented graph, and for a p -power q let G be the associated oriented pro- p RAAG. If G is Frattini-resistant, then Γ is specially oriented.*


So, let $\Gamma = (\mathcal{V}, \mathcal{E})$ be an oriented graph. Recall that the seven possible cases of induced subgraphs with three vertices that prevent Γ from being specially oriented are: the three triangle-graphs yielding torsion



(as shown in Examples 4.3–4.4); the two triangle-graphs



and the two line-graphs



We split the study of non-specially oriented graphs in these three cases.

5.1. Triangle subgraphs yielding non-trivial torsion. Suppose that Γ has an induced subgraph $\Gamma' = (\mathcal{V}', \mathcal{E}')$ as in (5.1). Let H be the oriented pro- p RAAG associated to q and to Γ' . Then by Examples 4.3–2.2, H yields non-trivial torsion. Now consider the homomorphism of pro- p groups $H \rightarrow G$ induced by the inclusion $\mathcal{V}' \hookrightarrow \mathcal{V}$. Then in all cases z is a non-trivial element of H — and hence also a non-trivial element of G — yielding non-trivial torsion. We have just proved the following.

Proposition 5.2. *If Γ has an induced subgraph as in (5.1), then G is not Frattini-resistant.*

5.2. Non-specially oriented triangles. Now suppose that Γ has an induced subgraph $\Gamma' = (\mathcal{V}', \mathcal{E}')$ as in (5.2). If G has non-trivial torsion, then clearly it is not Frattini-resistant by Example 2.2.

Therefore, we may suppose that G is torsion-free. Then the subgroup of G generated by x, y is a 2-generated free abelian pro- p group — i.e., it is isomorphic to \mathbb{Z}_p^2 —, and the subgroup generated by z is a 1-generated free abelian pro- p group — i.e., it is isomorphic to \mathbb{Z}_p — and it is normalized by x and y . Altogether, the subgroup of G generated by \mathcal{V}' is $\langle z \rangle \rtimes \langle x, y \rangle \simeq \mathbb{Z}_p \rtimes \mathbb{Z}_p^2$, and therefore every element g lying in it may be written in a unique way as

$$(5.4) \quad g = x^{\lambda_1} y^{\lambda_2} z^{\lambda_3} \quad \text{for some } \lambda_1, \lambda_2, \lambda_3 \in \mathbb{Z}_p.$$

Now set $t = yz$. Then

$$(5.5) \quad [x, t] = xyz y^{-1} x^{-1} z^{-1} = \begin{cases} z^q & \text{in the first case} \\ z^{q(1+q)} & \text{in the second case.} \end{cases}$$

Let H be the subgroup of G generated by x and t , and let N be the subgroup generated by t and z^q . Then N is isomorphic to \mathbb{Z}_p^2 in the first case, and to $\mathbb{Z}_p \rtimes \mathbb{Z}_p$ in the second case, and moreover by (5.5) it is a normal subgroup of H . In particular, one has $H = N \rtimes \langle x \rangle$. Thus, every element g of H may be written as

$$(5.6) \quad g = x^\lambda t^{\mu_1} z^{q\mu_2} = x^\lambda (yz)^{\mu_1} z^{q\mu_2} \quad \text{for some } \lambda, \mu_1, \mu_2 \in \mathbb{Z}_p.$$

By (5.5), one has that $z^{q(1+q)}$, and hence also z^q — recall that $1+q$ is invertible in \mathbb{Z}_p —, belongs to $\Phi(H)$. By (5.4), $z^{qp^{-1}}$ may not be written as in (5.6), and hence it does not belong to H . Therefore, G is not Frattini-resistant.

We have just proved the following.

Proposition 5.3. *If Γ has an induced subgraph as in (5.2), then G is not Frattini-resistant.*

Remark 5.4. The oriented pro- p RAAGs analyzed in this subsection are p -adic analytic (cf., e.g., [25, § 3] and Remark 4.5). Hence, one may use also [25, Thm. 1.2], to show that such an oriented pro- p RAAG is not Frattini-injective, and thus also neither Frattini-resistant.

5.3. Line-graphs. Finally, suppose that Γ has an induced subgraph $\Gamma' = (\mathcal{V}', \mathcal{E}')$ as in (5.3). If G has non-trivial torsion, then clearly it is not Frattini-resistant by Example 2.2.

Therefore, we may suppose that G is torsion-free. Let H be the subgroup of G generated by x, y, z , let H_1, H_2 , and V be the subgroups of H generated respectively by x, z , by x, y , and by y, z . Since we are assuming that G — and thus also H — is torsion-free, H_1 and H_2 are both isomorphic to $\mathbb{Z}_p \rtimes \mathbb{Z}_p$. Since

$$yx = \begin{cases} xy & \text{in the 1st case} \\ x^{1+q}y & \text{in the 2nd case} \end{cases} \quad \text{and} \quad zx = xz^{(1+q)^{-1}},$$

every element g of H may be written as $g = x^\lambda u$ for some $u \in V$ and $\lambda \in \mathbb{Z}_p$.

The following lemma is the “non-specially oriented analogous” of Lemma 5.5–(ii), and its proof follows the strategy of [3, Lemma 6.4].

Lemma 5.5. *The subgroup V is a 2-generated free pro- p group.*

Proof. Set $X = \langle x \rangle$. Since G is torsion-free, $X \simeq \mathbb{Z}_p$. Consider the second-countable pro- p tree $T = (\mathcal{V}_T, \mathcal{E}_T)$ where

$$\mathcal{V}_T = \{ gH_1, gH_2 \mid g \in H \}, \quad \mathcal{E}_T = \{ gX \mid g \in H \}.$$

Then H acts naturally on T by $h.(gH_i) = (hg)H_i$, with $i = 1, 2$, and $h.(gX) = (hg)X$ for every $h \in H$. The stabilizers in V of an edge gX and of a vertex gH_i are

$$\text{Stab}_V(gX) = V \cap {}^gX, \quad \text{and} \quad \text{Stab}_V(gH_i) = V \cap {}^gH_i, \quad i = 1, 2.$$

It is straightforward to see that $V \cap {}^gX = \{1\}$, and thus the stabilizer in V of any edge of T is trivial. Also, this implies that

$$V \cap {}^gH_1 = V \cap \langle {}^gz \rangle \quad \text{and} \quad V \cap {}^gH_2 = V \cap \langle {}^gy \rangle,$$

and thus these intersections are either trivial or isomorphic to \mathbb{Z}_p . Since $\text{Stab}_V(gX) = \{1\}$ for any $g \in H$, by [14, Thm. 5.6] H is isomorphic to the free pro- p product of some stabilizers $\text{Stab}_V(gH_i)$, $i \in \{1, 2\}$, and of a free pro- p group. Hence, altogether H is isomorphic to the free pro- p product of free pro- p groups — in fact, it cannot be but the free pro- p product of $\text{Stab}_V(H_1) = \langle z \rangle$ and $\text{Stab}_V(H_2) = \langle y \rangle$ —, and thus it is a free pro- p group. \square

Now set $t = zy$. Then

$$(5.7) \quad [x, t] = \begin{cases} [x, z] = z^q & \text{in the first case} \\ xzx^{-1-q}z^{-1} = x^{-q}z^{(1+q)^{1+q}-1} & \text{in the second case.} \end{cases}$$

Observe that $(1+q)^{1+q}-1$ is an element of \mathbb{Z}_p which is divisible by q but not by qp , as

$$\begin{aligned} (1+q)^{1+q} &= 1 + \binom{1+q}{1}q + \binom{1+q}{2}q^2 + \dots \\ &= 1 + (1+q)q + \frac{q(1+q)}{2}q^2 + \dots \end{aligned}$$

Hence, in the second case $[x, t] = x^{-q}z^{\lambda q}$, where λ is an invertible element of \mathbb{Z}_p . Let K be the subgroup of H generated by x, t . By (5.7), z^q belongs to $\Phi(K)$, as $[x, t], x^q \in \Phi(K)$.

We claim that $z^{qp^{-1}} \notin K$. Let W be the subgroup of K generated by t, z^q . Since V is a free pro- p group and $W \subseteq V$, also W is a free pro- p group, namely, the free pro- p group generated by t and z^q . It is easy to see that every element g of K may be written as

$$g = x^\lambda u \quad \text{for some } u \in W, \lambda \in \mathbb{Z}_p,$$

as done for the elements of H . Now, if $z^{qp^{-1}} \in K$, then $z^{qp^{-1}} = x^\lambda u$ with λ and u as above. Since $u, z^{qp^{-1}} \in V$, this implies that $x^\lambda \in V$, so that $\lambda = 0$, and $z^{qp^{-1}} = u \in W$. Consequently, $z^q = u^p$, which is impossible as W is the free pro- p group generated by t and z^q .

This proves the following.

Proposition 5.6. *If Γ has an induced subgraph as in (5.2), then G is not Frattini-resistant.*

Altogether, Propositions 5.2–5.3–5.6 prove Proposition 5.1.

6. ORIENTED GRAPHS OF ELEMENTARY TYPE AND FRATTINI-RESISTANCE

The goal of this section is to prove the following.

Proposition 6.1. *Let $\Gamma = (\mathcal{V}, \mathcal{E})$ be a specially oriented graph, and for a p -power q let G be the associated oriented pro- p RAAG. If G is Frattini-resistant, then Γ is of elementary type.*

Altogether, Proposition 6.1 and Proposition 5.1 yield implication (ii) \Rightarrow (i) of Theorem 1.1.

So, let $\Gamma = (\mathcal{V}, \mathcal{E})$ be a specially oriented graph. Recall that if Γ is specially oriented but not of elementary type, then by Proposition 3.6 G has a subgroup which is isomorphic to an oriented pro- p RAAG associated to the specially oriented graph (3.3), or to the oriented pro- p RAAG associated to one of the unoriented graphs as in (4.1).

As done in § 5, we proceed analyzing the two cases separately.

6.1. The specially oriented line-graph. Suppose that G has a subgroup which is isomorphic to the oriented pro- p RAAG associated to the specially oriented graph (3.3) and to q . Such a pro- p group is

$$\langle x, y, z \mid {}^x y = y^{1+q}, {}^x z = z^{1+q} \rangle = V \rtimes \langle x \rangle,$$

where $V = \langle y, z \rangle$, which is a 2-generated pro- p group by Lemma 4.2–(ii), and it is torsion-free.

If $q = p^f$ with $f \geq 2$ let H be the pro- p group generated by x, y, z . Otherwise, if $q = p$, let H be the subgroup of G generated by y, z and by x^λ where $\lambda \in \mathbb{Z}_p$ is such that $(1+p)^\lambda = 1+p^2$ (such λ exists because $1+p$ generates $1+p\mathbb{Z}_p$, and p divides λ). We write $\tilde{x} = x$ and $q' = q$ if $f \geq 2$, and $\tilde{x} = x^\lambda$ and $q' = p^2$ if $q = p$. Altogether, $H = V \rtimes \langle \tilde{x} \rangle$, and ${}^x y = y^{1+q'}$, ${}^x z = z^{1+q'}$.

Now set $t = yz^{-1}$ and $v = y^{q'p^{-1}}$, and let K be the subgroup of H generated by \tilde{x}, t, v . Since

$$\tilde{x}t = y^{1+q'}z^{-1-q'} = v^p t z^{-q'},$$

also $z^{q'} \in K$, and moreover

$$[\tilde{x}, t] = v^p t z^{-q'} t^{-1} = v^p [t, z^{-q'}] z^{-q'},$$

so that $z^{q'}$ belongs to $\Phi(K)$, too.

We claim that $z^{q'p^{-1}} \notin K$. To show this, first we prove the following.

Claim. *The subgroup W of H generated by $v, t, z^{q'}$ is the 3-generated free pro- p group on these three elements.*

Proof of the claim. Since V is a 2-generated free pro- p group, V is the free pro- p group generated by y and $t = yz^{-1}$.

Clearly, the subgroup $\langle v, t \rangle$ is not pro- p -cyclic, as t is not a power of y , and vice versa. Hence, it is a 2-generated free pro- p group. Now suppose that $\langle v, t \rangle = W$, so that $z^{q'} \in \langle v, t \rangle$. Then

$$z^{q'} = t^{\alpha_1} v^{\beta_1} t^{\alpha_2} v^{\beta_2} \dots = t^{\alpha_1} y^{q'p^{-1}\beta_1} t^{\alpha_2} y^{q'p^{-1}\beta_2} \dots$$

for some $\alpha_i, \beta_i \in \mathbb{Z}_p$. On the other hand,

$$z^{q'} = (t^{-1}y)^{q'} = \underbrace{t^{-1}yt^{-1}y \dots t^{-1}y}_{q' \text{ times}},$$

and thus one has two ways — which are distinct, as $q'p^{-1} \neq 1$ — to write $z^{q'}$ as an element generated by y and t . But this is a contradiction, as V is the free pro- p group generated by y and t . Thus, $z^{q'} \notin \langle v, t \rangle$, and W is the 3-generated free pro- p group generated by $v, t, z^{q'}$. This proves the claim. \square

To conclude, suppose for contradiction that $z^{q'p^{-1}} \in K$. Since $K = W \rtimes \langle \tilde{x} \rangle$, one may write $z^{q'p^{-1}} = \tilde{x}^\lambda w$ for some $\lambda \in \mathbb{Z}_p$ and $w \in W$, and therefore

$$z^{q'} = \left(z^{q'p^{-1}} \right)^p = \left(\tilde{x}^\lambda w \right)^p = \tilde{x}^{p\lambda} w' \quad \text{for some } w' \in W.$$

Since $z^{q'} \in W$, one deduces that $\lambda = 0$, and thus also $z^{q'p^{-1}} \in W$. But then $z^{q'}$ is a p -power of an element of W , which contradicts the fact that W is the free pro- p group on $v, t, z^{q'}$. Hence $z^{q'p^{-1}} \notin K$, and we have just proved the following.

Proposition 6.2. *If Γ is not of elementary type and G has a subgroup which is an oriented pro- p RAAG associated to the oriented graph (3.3), then G is not Frattini-resistant.*

6.2. The unoriented square-graph and line-graph. We are left with the two last cases, namely, G contains a subgroup which is the oriented pro- p RAAG associated to one of the two unoriented graphs (4.1).

Suppose first that G contains a subgroup H which is the oriented pro- p RAAG associated to the unoriented square graph with vertices x, y, z, w — namely,

$$\begin{aligned} H &= \langle x, y, z, w \mid [x, y] = [x, w] = [z, y] = [z, w] = 1 \rangle \\ &= \langle x, z \rangle \times \langle y, w \rangle. \end{aligned}$$

Since the two factors are 2-generated free pro- p groups by Lemma 4.2–(ii), none of them is free abelian, and hence H is not Frattini-resistant by Proposition 2.6.

Suppose now that G contains a subgroup which is the oriented pro- p RAAG associated to the unoriented line-graph with vertices x, y, z, w — namely, such a subgroup is

$$\langle x, y, z, w \mid [x, y] = [y, z] = [z, w] = 1 \rangle$$

— and let H be the subgroup of G generated by $y, z, {}^x z$ and t , where $t = xw$. Since $tzt^{-1} = {}^x z$, H is the HNN-extension of the subgroup $H_1 = \langle y, z, {}^x z \rangle$ with t , acting as an isomorphism $\langle z \rangle \simeq \langle {}^x z \rangle$ (cf. [27, Proof of Thm. 3.3]). In particular,

$$\begin{aligned} H &= \langle y, z, {}^z x, t \mid [y, z] = [y, {}^x z] = 1, [t, z] = {}^x z \cdot z^{-1} \rangle \\ &= \langle y, z, t \mid [y, z] = [[t, z], y] = 1 \rangle. \end{aligned}$$

Now consider the subgroup V of H (and of G) generated by t, yz, y^p, z^p . Then V is the HNN-extension of the subgroup $H_1 = \langle y^p, z^p, (xz)^p, yz \rangle$ with t , acting as an isomorphism $\langle z^p \rangle \simeq \langle (xz)^p \rangle$. One has

$$[t, z^p] = (xz)^p \cdot z^{-p} = (xz)^p (y^p \cdot y^{-p}) \cdot z^{-p} = (xz \cdot y)^p \cdot (yz)^{-p},$$

as y commutes with z and xz . Since $[t, z^p], (yz)^p \in \Phi(V)$, also $(xz \cdot y)^p$ lies in $\Phi(V)$. We claim that $xz \cdot y \notin V$.

Indeed, consider the normal subgroup N of H generated by y, z , and set $N_V = N \cap V$: namely, $H/N \simeq V/N_V \simeq \langle t \rangle$. Then

$$N = \left\langle t^k y, t^k z \mid k \in \mathbb{Z}, [t^k y, t^k z] = [t^k y, t^{k+1} z] = 1 \right\rangle,$$

and analogously

$$N_V = \left\langle t^k u \mid k \in \mathbb{Z}, [t^k u, t^k u'] = [t^k y^p, t^{k+1} z^p] = 1, u, u' = y^p, z^p, yz \right\rangle.$$

The abelianization N/N' is the free abelian pro- p group with basis

$$\mathcal{B}_N = \{ [t, (k), y]N', [t, (k), z]N' \mid k \in \mathbb{Z} \},$$

where

$$[a, (0), b] = b, \quad [a, (k), b] = \underbrace{[a, [a, \dots [a, [a, b]] \dots]]}_{k \text{ times}} \text{ for } k > 0,$$

and $[a, (k), b] = [a^{-1}, (-k), b]$ for $k < 0$; and analogously N_V/N'_V is the free abelian pro- p group with basis

$$\mathcal{B}_{N_V} = \{ [t, (k), u]N'_V \mid u = y^p, z^p, yz, k \in \mathbb{Z} \}.$$

We underline that \mathcal{B}_N and \mathcal{B}_{N_V} are subsets, of N/N' and N_V/N'_V respectively, converging to 1 — i.e., any open normal subgroup $U \subseteq H$ contains all but a finite number of elements of \mathcal{B}_N and \mathcal{B}_{N_V} —, as such a subgroup U contains all commutators of order bigger or equal to n , for some n (because H/U is a finite p -group, and hence it is nilpotent).

Within the free abelian pro- p group N/N' one has

$$(6.1) \quad {}^{xz}yN' = yN' \cdot zN' \cdot [t, z]N',$$

and therefore, (6.1) is the only way to express the coset of ${}^{xz}yN'$ using the basis \mathcal{B}_N . Now suppose that $xz \cdot y \in V$. Since $xz \cdot y \in N$, one has $xz \cdot y \in N_V$, and therefore one may write the coset of $xz \cdot y$ in N_V/N'_V as

$$(6.2) \quad {}^{xz}yN'_V = \prod_{k \in \mathbb{Z}} [t, (k), y^p]^{\alpha_k} N'_V \cdot [t, (k), z^p]^{\beta_k} N'_V \cdot [t, (k), yz]^{\gamma_k} N'_V$$

for some $\alpha_k, \beta_k, \gamma_k \in \mathbb{Z}_p$. Since $N' \supseteq N'_V$, and since

$$[t, (k), yz] \equiv [t, (k), y][t, (k), z] \quad \text{and} \quad [t, (k), u^p] \equiv [t, (k), u]^p$$

modulo N' for every $k \in \mathbb{Z}$ and $u \in N$, from (6.2) one obtains the following equality in the free abelian pro- p group N/N' :

$${}^{xz}yN' = y^{p\alpha_0 + \gamma_0} N' \cdot z^{p\beta_0 + \gamma_0} N' \cdot \prod_{k \neq 0} [t, (k), y]^{p\alpha_k + \gamma_k} N' \cdot [t, (k), z]^{p\beta_k + \gamma_k} N'.$$

By (6.1), one should have

$$p\alpha_1 + \gamma_1 = 0 \quad \text{and} \quad p\beta_1 + \gamma_1 = 1,$$

but then $p(\beta_1 - \alpha_1) = 1$, a contradiction, as $p \nmid 1$. Thus, ${}^x z \cdot y \notin V$. This completes the proof of the following.

Proposition 6.3. *If Γ is not of elementary type and G has a subgroup which is an oriented pro- p RAAG associated to one of the two unoriented graphs (4.1), then G is not Frattini-resistant.*

Altogether, Propositions 6.2–6.3 give Proposition 6.1.

7. MASSEY PRODUCTS AND FRATTINI-RESISTANT PRO- p GROUPS

7.1. Triple Massey products and upper-unitriangular representations. For $n \geq 2$ let \mathbb{U}_n denote the group of $n \times n$ upper unitriangular matrices with entries in \mathbb{Z}/p , namely,

$$\mathbb{U}_n = \left\{ \left(\begin{array}{cccc} 1 & a_{1,2} & a_{1,3} & \cdots \\ & 1 & a_{2,3} & \\ & & \ddots & \ddots \\ & & & 1 & a_{n-1,n} \\ & & & & 1 \end{array} \right) \mid a_{i,j} \in \mathbb{Z}/p \right\} \subseteq \mathrm{GL}_n(\mathbb{Z}/p).$$

The center $Z(\mathbb{U}_n)$ of \mathbb{U}_n consists of those matrices whose only non-0 entry — besides the main diagonal — is in the top-right corner, namely,

$$Z(\mathbb{U}_n) = \left\{ \left(\begin{array}{ccccc} 1 & 0 & \cdots & 0 & b \\ & 1 & & & 0 \\ & & \ddots & \ddots & \vdots \\ & & & 1 & 0 \\ & & & & 1 \end{array} \right) \mid b \in \mathbb{Z}/p \right\}.$$

The group \mathbb{U}_n is a finite p -group, and thus it is also a pro- p group.

Let G be a pro- p group, and consider \mathbb{Z}/p as a trivial G -module, as done in § 2.2. Recall that $H^1(G, \mathbb{Z}/p)$ is the \mathbb{Z}/p -vector space of all homomorphisms of pro- p groups $G \rightarrow \mathbb{Z}/p$ (cf. Remark 2.7). Let $\rho: G \rightarrow \mathbb{U}_n$ be a homomorphism of pro- p groups, and for $i = 1, \dots, n-1$, let $\rho_{i,i+1}$ denote the projection of ρ onto the $(i, i+1)$ -entry of ρ . Then $\rho_{i,i+1}: G \rightarrow \mathbb{Z}/p$ is a homomorphism of pro- p groups, and thus it is an element of $H^1(G, \mathbb{Z}/p)$. Analogously, if $\bar{\rho}: G \rightarrow \mathbb{U}_n$ is a homomorphism of pro- p groups, and for $i = 1, \dots, n-1$, $\bar{\rho}_{i,i+1}$ denotes the projection of $\bar{\rho}$ onto the $(i, i+1)$ -entry of $\bar{\rho}$, then $\bar{\rho}_{i,i+1}: G \rightarrow \mathbb{Z}/p$ is a homomorphism of pro- p groups, and thus an element of $H^1(G, \mathbb{Z}/p)$.

Given a sequence $\alpha_1, \dots, \alpha_n$ of length n of (non-necessarily distinct) elements of $H^1(G, \mathbb{Z}/p)$, the subset of $H^2(G, \mathbb{Z}/p)$ which is the value of the Massey product associated to the sequence $\alpha_1, \dots, \alpha_n$ is denoted by $\langle \alpha_1, \dots, \alpha_n \rangle$. One has the following “pro- p version” of the result of W. Dwyer in [7] (cf., e.g., [11, Lemma 9.3], see also [9, § 8]).

Proposition 7.1. *Let G be a pro- p group, and for $n \geq 2$ let $\alpha_1, \dots, \alpha_n$ a sequence of length n of (non-necessarily distinct) elements of $H^1(G, \mathbb{Z}/p)$.*

- (i) *The n -fold Massey product $\langle \alpha_1, \dots, \alpha_n \rangle$ is not empty if there exists a homomorphism*

$$\bar{\rho}: G \longrightarrow \mathbb{U}_{n+1}/\mathbb{Z}(\mathbb{U}_{n+1})$$

satisfying $\bar{\rho}_{i,i+1} = \alpha_i$ for all $i = 1, \dots, n$.

- (ii) *The n -fold Massey product $\langle \alpha_1, \dots, \alpha_n \rangle$ vanishes — i.e., contains 0 — if there exists a homomorphism*

$$\rho: G \longrightarrow \mathbb{U}_{n+1}$$

satisfying $\rho_{i,i+1} = \alpha_i$ for all $i = 1, \dots, n$.

Hence, there are no essential n -fold Massey products in the \mathbb{Z}/p -cohomology of G if for any sequence $\alpha_1, \dots, \alpha_n$ of length n of (non-necessarily distinct) elements of $\mathbb{H}^1(G, \mathbb{Z}/p)$, either there are no homomorphisms $\bar{\rho}: G \rightarrow \mathbb{U}_{n+1}/\mathbb{Z}(\mathbb{U}_{n+1})$ as in statement (i) of Proposition 7.1, or there is a homomorphism $\rho: G \rightarrow \mathbb{U}_{n+1}$ as in statement (ii) of Proposition 7.1.

Remark 7.2. The group \mathbb{U}_3 is isomorphic to the Heisenberg group modulo p , namely

$$(7.1) \quad \mathbb{U}_3 = \langle A, B, C \mid [A, B] = C, A^p = B^p = [A, C] = [B, C] = I_3 \rangle,$$

where

$$A = \begin{pmatrix} 1 & 1 & 0 \\ & 1 & 0 \\ & & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 & 0 \\ & 1 & 1 \\ & & 1 \end{pmatrix}, \quad C = \begin{pmatrix} 1 & 0 & 1 \\ & 1 & 0 \\ & & 1 \end{pmatrix}.$$

Now let G be a pro- p group. Given a sequence $\alpha_1, \alpha_2, \alpha_3$ of length 3 of elements of $\mathbb{H}^1(G, \mathbb{Z}/p)$, non necessarily distinct, then there exists a homomorphism $\bar{\rho}: G \rightarrow \mathbb{U}_4/\mathbb{Z}(\mathbb{U}_4)$ as in statement (i) of Proposition 7.1 if, and only if, there exist two homomorphisms $\tau, \tau': G \rightarrow \mathbb{U}_3$ satisfying

$$\tau(x) \equiv A^{\alpha_1(x)} B^{\alpha_2(x)} \pmod{\langle C \rangle} \quad \text{and} \quad \tau'(x) \equiv A^{\alpha_2(x)} B^{\alpha_3(x)} \pmod{\langle C \rangle}$$

for all $x \in G$.

7.2. Triple Massey products and Frattini-resistance. Let $\Gamma = (\mathcal{V}, \mathcal{E})$ be an oriented graph, and for a p -power q let G be the associated oriented pro- p RAAG. By [3, Cor. 1.2–(ii)], if Γ is of elementary type — and hence, if G is Frattini-resistant —, then there are no essential n -fold Massey products in the \mathbb{Z}/p -cohomology of G for every $n > 2$. In fact, by [21, Thm. 1.1], there are many more oriented graphs whose associated oriented pro- p RAAGs yield no essential n -fold Massey products, than oriented graphs yielding Frattini-resistant pro- p RAAGs: for example, for every unoriented graph — and thus also the unoriented graphs (4.1), which are not of elementary type — the associated oriented pro- p RAAGs yield no essential n -fold Massey products for any $n > 2$ (see also [2, Thm. 1.1]).

Therefore, within the family of oriented pro- p RAAGs associated to oriented graph, Frattini-resistance is far more restrictive than the absence of essential n -fold Massey products for every $n > 2$. This suggests to formulate Conjecture 1.4, which — by Proposition 7.1 and Remark 7.2 — may be formulated in group-theoretic terms as follows.

Conjecture 7.3 (Reformulation of Conjecture 1.4). *Let G be a Frattini-resistant pro- p group, and let $\alpha_1, \alpha_2, \alpha_3$ be (non-necessarily distinct) homomorphisms $G \rightarrow \mathbb{Z}/p$. If there exist two homomorphisms of pro- p groups $\tau, \tau' : G \rightarrow \mathbb{U}_3$ satisfying*

$$\tau(x) \equiv A^{\alpha_1(x)} B^{\alpha_2(x)} \pmod{\langle C \rangle} \quad \text{and} \quad \tau'(x) \equiv A^{\alpha_2(x)} B^{\alpha_3(x)} \pmod{\langle C \rangle}$$

for all $x \in G$, then there exists a homomorphism $\rho : G \rightarrow \mathbb{U}_4$ satisfying $\rho_{i,i+1} = \alpha_i$ for $i = 1, 2, 3$.

Remark 7.4. By Proposition 2.8, a positive answer to Conjecture 1.4 would provide a new proof of the recent result of A. Merkurjev and F. Scavia [15, Thm. 1.3], which states that pro- p groups which may complete into a 1-cyclotomic oriented pro- p group yield no essential triple Massey products in \mathbb{Z}/p -cohomology.

In fact, we suspect that Conjecture 1.4 may be true not only for triple Massey products, but also for n -fold Massey products for every $n \geq 3$... but maybe such a question is too daring — and a positive answer would provide a positive solution to the *Massey vanishing conjecture* for maximal pro- p Galois groups formulated by J. Minač and N.D. Tân, see [16, Conj. 1.1] —, so we do not formulate it as a conjecture on its own, but we just whisper it.

Acknowledgments. The author wishes to thank Ilir Snopce and Slobodan Tanushevski, for several inspiring discussions on Frattini-resistant pro- p groups (and in particular, for their talks on this topic the author attended). The author acknowledges his membership to the national group GNSAGA (Algebraic Structures and Algebraic Geometry) of the National Institute of Advanced Mathematics – a.k.a. INdAM – “F. Severi”.

REFERENCES

- [1] D. Benson, N. Lemire, J. Minač, and J. Swallow, *Detecting pro- p -groups that are not absolute Galois groups*, J. Reine Angew. Math. **613** (2007), 175–191.
- [2] S. Blumer, A. Cassella, and C. Quadrelli, *Groups of p -absolute Galois type that are not absolute Galois groups*, J. Pure Appl. Algebra **227** (2023), no. 4, Paper No. 107262.
- [3] S. Blumer, C. Quadrelli, and Th.S. Weigel, *Oriented right-angled Artin pro- ℓ groups and maximal pro- ℓ Galois groups*, Int. Math. Res. Not. (2024). In press, published on-line.
- [4] A. Cassella and C. Quadrelli, *Right-angled Artin groups and enhanced Koszul properties*, J. Group Theory **24** (2021), no. 1, 17–38.
- [5] J.D. Dixon, M.P.F. du Sautoy, A. Mann, and D. Segal, *Analytic pro- p groups*, 2nd ed., Cambridge Studies in Advanced Mathematics, vol. 61, Cambridge University Press, Cambridge, 1999.
- [6] C. Droms, *Subgroups of graph groups*, J. Algebra **110** (1987), no. 2, 519–522.
- [7] W.G. Dwyer, *Homology, Massey products and maps between groups*, J. Pure Appl. Algebra **6** (1975), no. 2, 177–190.
- [8] I. Efrat, *Small maximal pro- p Galois groups*, Manuscripta Math. **95** (1998), no. 2, 237–249.
- [9] ———, *The Zassenhaus filtration, Massey products, and representations of profinite groups*, Adv. Math. **263** (2014), 389–411.
- [10] I. Efrat and E. Matzri, *Triple Massey products and absolute Galois groups*, J. Eur. Math. Soc. (JEMS) **19** (2017), no. 12, 3629–3640.
- [11] I. Efrat and C. Quadrelli, *The Kummerian property and maximal pro- p Galois groups*, J. Algebra **525** (2019), 284–310.
- [12] M. D. Fried and M. Jarden, *Field arithmetic*, 4th ed., Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, vol. 11, Springer, Cham, 2023. Revised by M. Jarden.

- [13] E. Matzri, *Triple Massey products in Galois cohomology*, 2014. Preprint, available at [arXiv:1411.4146](https://arxiv.org/abs/1411.4146).
- [14] O. V. Mel'nikov, *Subgroups and the homology of free products of profinite groups*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 1, 97–120 (Russian); English transl., Math. USSR-Izv. **34** (1990), no. 1, 97–119.
- [15] A. Merkurjev and F. Scavia, *On the Massey Vanishing Conjecture and Formal Hilbert 90*, 2023. Preprint, available at [arXiv:2308.13682](https://arxiv.org/abs/2308.13682).
- [16] J. Minač and N.D. Tân, *The kernel unipotent conjecture and the vanishing of Massey products for odd rigid fields*, Adv. Math. **273** (2015), 242–270.
- [17] ———, *Triple Massey products and Galois theory*, J. Eur. Math. Soc. (JEMS) **19** (2017), no. 1, 255–284.
- [18] J. Neukirch, A. Schmidt, and K. Wingberg, *Cohomology of number fields*, 2nd ed., Grundlehren der Mathematischen Wissenschaften, vol. 323, Springer-Verlag, Berlin, 2008.
- [19] C. Quadrelli, *Two families of pro- p groups that are not absolute Galois groups*, J. Group Theory **25** (2022), no. 1, 25–62.
- [20] ———, *Chasing maximal pro- p Galois groups via 1-cyclotomicity*, Mediterranean J. Math. **21** (2024), Paper No. 56.
- [21] ———, *Oriented graphs, pro- p groups, and Massey products in Galois cohomology*, 2024. Preprint, available at [arXiv:2403.01464](https://arxiv.org/abs/2403.01464).
- [22] C. Quadrelli, I. Snopce, and M. Vannacci, *On pro- p groups with quadratic cohomology*, J. Algebra **612** (2022), 636–690.
- [23] C. Quadrelli and Th.S. Weigel, *Profinite groups with a cyclotomic p -orientation*, Doc. Math. **25** (2020), 1881–1916.
- [24] J-P. Serre, *Galois cohomology*, Corrected reprint of the 1997 English edition, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2002. Translated from the French by Patrick Ion and revised by the author.
- [25] I. Snopce and S. Tanushevski, *Frobenius-injectivity and maximal pro- p Galois groups*, 2020. Preprint, available at [arXiv:2009.09297](https://arxiv.org/abs/2009.09297).
- [26] ———, *Frobenius-resistant direct products of pro- p groups*, J. Algebra **622** (2023), 351–365.
- [27] I. Snopce and P. Zalesskiĭ, *Right-angled Artin pro- p groups*, Bull. Lond. Math. Soc. **54** (2022), no. 5, 1904–1922.
- [28] T. Würfel, *On a class of pro- p groups occurring in Galois theory*, J. Pure Appl. Algebra **36** (1985), no. 1, 95–103.

DEPARTMENT OF SCIENCE & HIGH-TECH, UNIVERSITY OF INSUBRIA, COMO, ITALY EU
 Email address: claudio.quadrelli@uninsubria.it