

Existence for noncoercive nonlinear elliptic equations with two lower-order terms

Thi Tam Dang*, Trung Hau Hoang †

April 2, 2024

Abstract

This paper considers a class of noncoercive nonlinear elliptic problems with coefficients defined in Marcinkiewicz and Lorentz spaces. We prove the existence of a solution for the corresponding Dirichlet problem and investigate the higher integrability properties of the solution.

1 Introduction

This paper considers a general noncoercive nonlinear elliptic problem of the form:

$$\begin{cases} -\operatorname{div}A(x, u, \nabla u) + B(x, u, \nabla u) + G(x, u) = F & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where Ω is a bounded open subset of \mathbb{R}^N with $N \geq 2$ and $F \in W^{-1,p'}(\Omega)$. The operator $A(x, z, \xi) : \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ is a Carathéodory vector field which meets the following conditions: for a.e. $x \in \Omega$, $z \in \mathbb{R}$ and $\xi, \xi_* \in \mathbb{R}^N$

$$\langle A(x, z, \xi), \xi \rangle \geq \alpha|\xi|^p - (b(x)|z|)^p - \varphi(x)^p, \quad \alpha > 0, \quad (1.2)$$

$$|A(x, z, \xi)| \leq \beta|\xi|^{p-1} + (b(x)|z|)^{p-1} + \varphi(x)^{p-1}, \quad \beta > 0, \quad (1.3)$$

$$\langle A(x, z, \xi) - A(x, z, \xi_*), \xi - \xi_* \rangle > 0 \quad \text{for } \xi \neq \xi_*, \quad (1.4)$$

where $b(x) \in L^{N,\infty}(\Omega)$, and $\varphi(x) \in L^p(\Omega)$ with $1 < p < N$.

The operator $B(x, z, \xi) : \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}$ is a Carathéodory function satisfies

$$|B(x, z, \xi)| \leq c(x)|\xi|^{p-1} + \phi(x), \quad (1.5)$$

for a.e. $x \in \Omega$, $z \in \mathbb{R}$, $\xi, \xi_* \in \mathbb{R}^N$, the coefficient $c(x) \in L^{N,1}(\Omega)$ and $\phi \in L^1(\Omega)$.

Finally, the vector field $G(x, z) : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function satisfying

$$G(x, z)z \geq 0, \quad (1.6)$$

$$|G(x, z)| \leq d(x)|z|^\lambda + \psi(x), \quad 0 \leq \lambda < \frac{N(p-1)}{N-p}, \quad (1.7)$$

where $d(x) \in L^{s',1}(\Omega)$ with $s = \frac{N(p-1)}{N-p} \frac{1}{\lambda}, \frac{1}{s} + \frac{1}{s'} = 1$ and $\psi(x) \in L^1(\Omega)$.

A simple example of the model problem (1.1) can be read as follows:

$$\begin{cases} -\Delta_p u + \operatorname{div}(K(x)|u|^{p-2}u) + c(x)|\nabla u|^{p-1} = -\operatorname{div}\left(\frac{x}{|x|^{N-p}}\right) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.8)$$

*Department of Mathematics, Leopold Franzens Universität Innsbruck, Austria, e-mail: Thi-Tam.Dang@uibk.ac.at

†Department of Mathematics, Leopold Franzens Universität Innsbruck, Austria, e-mail: Trung-Hau.Hoang@uibk.ac.at

for any $\frac{N}{2} - 1 < \nu \leq N - 1$. For a.e. $x \in \Omega$, we define the vector field $K : \Omega \rightarrow \mathbb{R}^N$ is a measurable function satisfying $|K(x)| \leq b(x)^{p-1}$ with $b(x) \in L^{N,\infty}(\Omega)$. Thus, the operator A is a combination of the following two terms:

$$A(x, z, \xi) := -\Delta_p u + \operatorname{div}(K(x)|u|^{p-2}u). \quad (1.9)$$

In the case the operator B and G vanishes, the problem (1.1) has been studied in [9] where the authors proved the existence of the solution for the quasilinear elliptic problems with singularity in the lower order term. The existence of the renormalized solutions has been studied in [13] in which the operator A is independent of u with no singularity behavior.

The problem (1.1) has two main features: first, the vector field A introduces a singular lower-order term influenced by u , described by the coefficient $b(x)$. The property that the coefficient $b(x)$ belongs to the Marcinkiewicz spaces $L^{N,\infty}(\Omega)$ leads to a lack of compactness of the operator $K(x, u)$. That is, the operator $K(x, u)$ does not belong to $L^{p'}(\Omega)$ in general, and the term $\operatorname{div}(K(x, u))$ is not an element of the dual space $W^{-1,p'}(\Omega)$. Second, the operator B is characterized by a coefficient, denoted $c(x)$, belonging to the Lorentz spaces $L^{N,1}(\Omega)$, which exhibit unboundedness. Typically, the presence of these singularity and unboundedness properties causes the problem to be noncoercive.

For any $\varepsilon \in [0, 1]$, we set

$$\zeta_\varepsilon(x) = \varepsilon c(x) + (1 - \varepsilon)b(x), \quad \forall x \in \Omega. \quad (1.10)$$

We further define $X_\varepsilon(\Omega)$ to be a convex subset of $L^{N,q}(\Omega)$, with $1 \leq q \leq \infty$ such that for $\zeta_\varepsilon(x) \in X_\varepsilon(\Omega)$ satisfies

$$\|\zeta_\varepsilon - T_k \zeta_\varepsilon\|_{N,q} \leq \theta, \quad \theta = \frac{1}{S_{N,p}} \left(\frac{\alpha}{2p} \right)^{\frac{1}{p}}, \quad (1.11)$$

for any $k > 0$, $S_{N,p}$ denotes the Sobolev constant given by Theorem (2.1) below. Our main existence result is stated in the next theorem.

Theorem 1.1. *Let the assumptions (1.2)-(1.7) be fulfilled. Furthermore, we assume that*

$$\zeta_\varepsilon \in X_\varepsilon(\Omega). \quad (1.12)$$

Then the problem (1.1) admits a solution.

The novelty of this study lies in establishing the existence theorem by an approximation method. To achieve this, a sequence of approximation problems is considered (see Section 3.1), and the existence of a solution to these problems is demonstrated using the Leray–Schauder fixed point theorem. The primary challenge is deriving a priori estimates for the sequence of solutions, which is facilitated by fulfilling our assumption (1.12). Subsequently, we establish the compactness of the sequence of solutions to the approximation problems through rigorous testing with an admissible test function. Then we show that the limit of the sequence is a solution to the original problem (see Section 3.2).

Consider for a moment the problem (1.1) in the linear case and the coefficient $c = 0$, Stampacchia’s results (see [29]) proved that $u \in L^{r^*}(\Omega)$ by assuming the coefficients of the lower order terms in appropriate Lebesgue spaces. For nonlinear elliptic problems, in [3] Stampacchia’s results were extended by assuming $b(x) \in L^{N/p-1}(\Omega)$. In [14] a similar result was achieved for nonlinear elliptic problems with lower order term coefficients in Marcinkiewicz spaces. Their analysis assumed that these coefficients are in $L^\infty(\Omega)$. In addition, in a recent publication [9], the authors extended the previous results to noncoercive quasilinear elliptic operators with a singular lower order term.

The intriguing question that arises is whether the aforementioned results can be extended to a broader class of noncoercive nonlinear elliptic problem, involving coefficients of lower order terms in both the Marcinkiewicz and Lorentz spaces. In our present work we aim to extend the above results to our problem. More precisely, we will prove the higher summability of a solution under the appropriate properties of the given data.

The structure of the paper is as follows: we start with the Preliminaries, where we recall the definition and some properties of Lorentz spaces, which play a fundamental role in our analytical framework. In Section 3 we will demonstrate the existence of a solution by an approximation procedure. We will show that the sequence of solutions to this approximation problem converges to a solution of the original problem. Finally, Section 4 investigates the regularity properties of a solution.

2 Preliminaries

This section recalls a definition and representative properties of Lorentz spaces, which are used extensively throughout this paper. For further information on Lorentz spaces, we refer the readers to [18].

Let $\Omega \subset \mathbb{R}^N$ be a bounded domain. The distribution function of f is given by

$$\omega_f(h) = |E_h| = |\{x \in \Omega : |f(x)| > h\}|, \quad (2.1)$$

where $|E_h|$ is the Lebesgue measure of E_h . For given $1 < r < \infty$ and $1 \leq s < \infty$, the Lorentz spaces denoted by $L^{r,s}(\Omega)$ consist of all measurable functions f defined on Ω which satisfy

$$\|f\|_{r,s}^r = \int_0^\infty (\omega_f(h))^{\frac{s}{r}} h^{s-1} dh < \infty. \quad (2.2)$$

We remark that $L^{r,s}(\Omega)$ becomes Banach space when it is endowed with the norm $\|\cdot\|_{r,s}$. In the case $r = s$, the Lorentz space simplifies to the Lebesgue space $L^r(\Omega)$. If $s = \infty$, the class $L^{r,\infty}(\Omega)$ consists of all measurable functions f defined on Ω which satisfy

$$\|f\|_{r,\infty}^r = \sup_{t>0} h^r \omega_f(h) < \infty, \quad (2.3)$$

and it coincides with the Marcinkiewicz class, the weak- $L^r(\Omega)$. For Lorentz spaces, the following continuous inclusions apply:

$$L^q(\Omega) \subset L^{r,s}(\Omega) \subset L^{r,q}(\Omega) \subset L^{r,\infty}(\Omega) \subset L^s(\Omega) \quad (2.4)$$

for $1 \leq s < r < q \leq \infty$. Furthermore, for any $f \in L^{r,s}(\Omega)$, $g \in L^{r',s'}(\Omega)$, where $1 < r < \infty$, $1 \leq s \leq \infty$, $\frac{1}{r} + \frac{1}{r'} = 1$, $\frac{1}{s} + \frac{1}{s'} = 1$, the following generalized Hölder inequality holds

$$\|fg\|_1 \leq \|f\|_{r,s} \|g\|_{r',s'}. \quad (2.5)$$

The distance of a function $f \in L^{r,\infty}(\Omega)$ to $L^\infty(\Omega)$ can be characterized as follows:

$$\lim_{k \rightarrow \infty} \|f - T_k f\|_{r,\infty} = \text{dist}_{L^{r,\infty}(\Omega)}(f, L^\infty(\Omega)), \quad (2.6)$$

where $T_k(s) : \mathbb{R} \rightarrow \mathbb{R}$ denotes the truncation operator at level $\pm k$, i.e. $T_k(s) = \frac{s}{|s|} \min\{|s|, k\}$. For more about the distance to $L^\infty(\Omega)$ and some applications, see [7].

The Sobolev embedding theorem in the framework of Lorentz spaces is stated in the next theorem.

Theorem 2.1. ([2, 9, 18]) *Let us assume that $1 < p < N$, $1 \leq q \leq p$. Then, any function $g \in W_0^{1,1}(\Omega)$ satisfying $|\nabla g| \in L^{p,q}(\Omega)$ belongs to $L^{p^*,p}(\Omega)$ where $p^* = \frac{Np}{N-p}$ and*

$$\|g\|_{p^*,q} \leq S_{N,p} \|\nabla g\|_{p,q}, \quad (2.7)$$

where $S_{N,p} = \omega_N^{-1/N} \frac{p}{N-p}$ and ω_N stands for the measure of the unit ball in \mathbb{R}^N .

We need the following the weak compactness lemma for the proof of the Theorem 1.1.

Lemma 2.1. (Lemma 3.1, [9]) Let \mathcal{B} be a nonempty subset of $W_0^{1,p}(\Omega)$. Assume that there exists a constant $C > 0$ such that

$$\|\nabla u\|_{L^p(\Omega \setminus E_k)}^p \leq C(1 + \|u\|_{L^p(\Omega \setminus E_k)}^p), \quad (2.8)$$

for any $k > 0$ and $u \in \mathcal{B}$, where $E_k := \{x \in \Omega : |u(x)| \geq k\}$. Then, there exists a constant $M > 0$ such that

$$\|u\|_{W^{1,p}(\Omega)} \leq M, \quad (2.9)$$

for any $u \in \mathcal{B}$.

We conclude this section with the well known Leray–Schauder fixed point theorem.

Theorem 2.2. (Leray–Schauder fixed point) ([9, 11]) Let \mathcal{F} be a compact mapping of a Banach space X into itself, and all suppose there exists a constant M such that $\|x\| < M$ for all $x \in X$ and $t \in [0, 1]$ satisfying $x = t\mathcal{F}(x)$. Then, \mathcal{F} has a fixed point.

3 Proof of theorem

We first consider the approximation problems, then showing that the approximation problems have a solution via the Leray-Schauder fixed-point theorem that converges to a solution of our original problem (1.1).

3.1 The approximation problems

For each $n \in \mathbb{N}$, for any $\varepsilon \in [0, 1]$, we set

$$\vartheta_n^\varepsilon(x) := \frac{T_n \zeta_\varepsilon(x)}{\zeta_\varepsilon(x)}, \quad \text{a.e. } x \in \Omega,$$

where ζ_ε is given by (1.10). We consider the following approximating problems:

$$-\operatorname{div} A_n(x, u_n, \nabla u_n) + B_n(x, u_n, \nabla u_n) + G_n(x, u_n) = F, \quad (3.1)$$

where the vector field $A_n : (x, z, \xi) \in \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ is defined by

$$A_n(x, z, \xi) = A(x, \vartheta_n^\varepsilon z, \xi).$$

The vector field A_n shares properties with A as follows:

$$\langle A_n(x, z, \xi), \xi \rangle \geq \alpha |\xi|^p - (T_n \zeta_0(x) |z|)^p - \varphi(x)^p, \quad (3.2)$$

$$|A_n(x, z, \xi)| \leq \beta |\xi|^{p-1} + (T_n \zeta_0(x) |z|)^{p-1} + \varphi(x)^{p-1}, \quad (3.3)$$

$$\langle A_n(x, z, \xi) - A_n(x, z, \xi_*), \xi - \xi_* \rangle > 0 \quad \text{for } \xi \neq \xi_*. \quad (3.4)$$

Moreover, we let $B_n(x, z, \xi) : \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ satisfy

$$|B_n(x, u, \xi)| \leq T_n \zeta_1(x) |\xi|^{p-1} + \phi(x). \quad (3.5)$$

Finally, the vector field $G_n(x, z) = T_n G(x, u)$ fulfils the following conditions:

$$G_n(x, z)z \geq 0, \quad (3.6)$$

$$|G_n(x, z)| \leq d(x) |z|^\lambda + \psi(x). \quad (3.7)$$

Our goal is to find $u_n \in W_0^{1,p}(\Omega)$, which solves the approximation problem (3.1). The existence of the solution u_n can be proven by applying the Leray-Schauder fixed point theorem. For this, we need the following technical lemmas, which provide an a priori estimate for u_n . The next lemma gives an a priori estimate for $\|\nabla S_k u_n\|_{N', \infty}^{p-1}$ where $S_k(u_n)$ is defined by (3.11) below.

Lemma 3.1. *Let Ω be an open subset of \mathbb{R}^N with finite measure and that $1 < p < N$. Let u be a measurable function satisfying $T_k(u_n) \in W_0^{1,p}(\Omega)$, for every positive $k > 0$, and such that*

$$\int_{\Omega} |\nabla T_{\sigma}(S_k u_n)|^p dx \leq Mk + L, \quad \forall k > 0, \quad (3.8)$$

where M and L are given constants. Then $|\nabla S_k u_n|^{p-1}$ belongs to $L^{N',\infty}(\Omega)$ and

$$\| |\nabla S_k u_n|^{p-1} \|_{N',\infty} \leq 2C(N,p) \left[\frac{\|\phi\|_1}{C_1} + |\Omega|^{\frac{1}{N'} - \frac{1}{p'}} L^{\frac{1}{p'}} \right] \quad (3.9)$$

where $C(N,p)$ is a constant depending only on N and p .

Proof. For $k > 0$, for all $s \in \mathbb{R}$, the remainder $S_k(s)$ of the truncation $T_k(s)$ is given by

$$S_k(s) = s - T_k(s), \quad (3.10)$$

that is

$$S_k(s) = \begin{cases} 0, & |s| \leq k, \\ (|s| - k)\text{sign}(s), & |s| > k. \end{cases} \quad (3.11)$$

For fixed $\sigma > 0$, using $T_{\sigma}(S_k(u_n))$ as a test function of the approximation equation (3.1), we obtain

$$\begin{aligned} \int_{\Omega} A_n(x, u_n, \nabla u_n) \cdot \nabla T_{\sigma}(S_k(u_n)) dx + \int_{\Omega} B_n(x, u_n, \nabla u_n) T_{\sigma}(S_k(u_n)) dx \\ + \int_{\Omega} G_n(x, u_n) T_{\sigma}(S_k(u_n)) dx = \int_{\Omega} F T_{\sigma}(S_k(u_n)) dx. \end{aligned} \quad (3.12)$$

By the definition of $S_k(u_n)$ given by (3.11) and the ellipticity condition (3.2) on A_n , we get

$$\begin{aligned} \int_{\Omega} A_n(x, u_n, \nabla u_n) \cdot \nabla T_{\sigma}(S_k(u_n)) dx \\ = \int_{\{k \leq |u_n| \leq k+\sigma\}} A_n(x, u_n, \nabla u_n) \cdot \nabla u_n dx \\ \geq \int_{\{k \leq |u_n| \leq k+\sigma\}} \left(\alpha |\nabla u_n|^p - (T_n(\zeta_0)|u_n|)^p - \varphi(x)^p \right) dx \\ \geq \alpha \int_{\Omega} |\nabla T_{\sigma}(S_k(u_n))|^p dx - \int_{\{k \leq |u_n| \leq k+\sigma\}} (T_n(\zeta_0)|u_n|)^p dx - \|\varphi\|_p^p. \end{aligned} \quad (3.13)$$

For any $n \geq m$, where m is a positive integer to be chosen later, we have

$$T_n \zeta_{\varepsilon} \leq T_m \zeta_{\varepsilon} + (\zeta_{\varepsilon} - T_m \zeta_{\varepsilon}), \quad \varepsilon = 0, 1,$$

which implies

$$\begin{aligned} \int_{\{k \leq |u_n| \leq k+\sigma\}} (T_n(\zeta_0)u_n)^p dx = \|T_n \zeta_0 |u_n| \chi_{\{k \leq |u_n| \leq k+\sigma\}}\|_p^p \\ \leq \|T_m(\zeta_0)u_n \chi_{\{k \leq |u_n| \leq k+\sigma\}}\|_p^p + \|(\zeta_0 - T_m(\zeta_0))u_n \chi_{\{k \leq |u_n| \leq k+\sigma\}}\|_p^p. \end{aligned} \quad (3.14)$$

Using the Hölder inequality (2.5), the generalized Sobolev embedding Theorem 2.1, and the assumption (1.12), we deduce that

$$\begin{aligned} \|T_m(\zeta_0)u_n \chi_{\{k \leq |u_n| \leq k+\sigma\}}\|_p^p &\leq m^p \|u_n \chi_{\{k \leq |u_n| \leq k+\sigma\}}\|_p^p \\ &\leq m^p \|1\|_{N,\infty}^p \|u_n \chi_{\{k \leq |u_n| \leq k+\sigma\}}\|_{p^*,p}^p \\ &\leq m^p |\Omega|^{\frac{p}{N}} S_{N,p}^p \|\nabla T_{\sigma}(S_k(u_n))\|_p^p, \end{aligned} \quad (3.15)$$

and

$$\begin{aligned}
\|(\zeta_0 - T_m(\zeta_0))u_n \chi_{\{k \leq |u_n| \leq k+\sigma\}}\|_p^p &\leq \|\zeta_0 - T_m(\zeta_0)\|_{N,\infty}^p \|u_n \chi_{\{k \leq |u_n| \leq k+\sigma\}}\|_{p^*,p}^p \\
&\leq S_{N,p}^p \|\zeta_0 - T_m \zeta_0\|_{N,\infty}^p \|\nabla T_\sigma(S_k(u_n))\|_p^p \\
&\leq \frac{\alpha}{2p} \|\nabla T_\sigma(S_k(u_n))\|_p^p.
\end{aligned} \tag{3.16}$$

By the definition of $S_k(u_n)$, $S_k(u_n) = 0$ for $|u_n| \leq k$, and the growth assumption (3.5) on B_n , we get

$$\begin{aligned}
&\left| \int_{\Omega} B_n(x, u_n, \nabla u_n) T_\sigma(S_k(u_n)) dx \right| \\
&\leq \sigma \int_{\Omega} |B_n(x, u_n, \nabla u_n)| dx \\
&\leq \sigma \left[\int_{\{|u_n| > k\}} T_n(\zeta_1) |\nabla u_n|^{p-1} dx + \int_{\Omega} \phi(x) dx \right] \\
&\leq \sigma \int_{\{|u_n| > k\}} T_n(\zeta_1) |\nabla u_n|^{p-1} dx + \sigma \|\phi\|_1.
\end{aligned} \tag{3.17}$$

For the first term on the right side of (3.17), we use the Hölder inequality and the assumption (1.12) to obtain

$$\begin{aligned}
&\sigma \int_{\{|u_n| > k\}} T_n(\zeta_1) |\nabla u_n|^{p-1} dx \\
&\leq \sigma \int_{\{|u_n| > k\}} T_m(\zeta_1) |\nabla u_n|^{p-1} dx + \sigma \int_{\{|u_n| > k\}} (\zeta_1 - T_m(\zeta_1)) |\nabla u_n|^{p-1} dx \\
&\leq \sigma m \int_{\Omega} |\nabla u_n \chi_{\{|u_n| > k\}}|^{p-1} dx + \sigma \|\zeta_1 - T_m(\zeta_1)\|_{N,1} \|\nabla S_k(u_n)\|_{N',\infty}^{p-1} \\
&\leq \sigma m \|1\|_{N,1} \|\nabla S_k(u_n)\|_{N',\infty}^{p-1} + \sigma \theta \|\nabla S_k(u_n)\|_{N',\infty}^{p-1} \\
&\leq \sigma (mN|\Omega|^{\frac{1}{N}} + \theta) \|\nabla S_k(u_n)\|_{N',\infty}^{p-1},
\end{aligned} \tag{3.18}$$

where θ is given by (1.12). The sign condition (3.6) on G_n leads to

$$\int_{\Omega} G_n(x, u_n) T_\sigma(S_k(u_n)) dx \geq 0. \tag{3.19}$$

Applying Young's inequality yields

$$\begin{aligned}
\int_{\Omega} F T_\sigma(S_k(u_n)) dx &\leq \|F\| \|\nabla T_\sigma(S_k(u_n))\|_p \\
&\leq \frac{\alpha}{2p} \|\nabla T_\sigma(S_k(u_n))\|_p^p + \frac{2^{p'/p}}{p' \alpha^{p'/p}} \|F\|_{p'}^{p'}.
\end{aligned} \tag{3.20}$$

We set

$$C_1 := \frac{\alpha}{p'} - m^p |\Omega|^{\frac{p}{N}} S_{N,p}^p. \tag{3.21}$$

In view of (3.13)-(3.20), we obtain

$$\|\nabla T_\sigma(S_k(u_n))\|_p^p \leq M\sigma + L, \quad \forall \sigma > 0, \tag{3.22}$$

where

$$\begin{aligned}
M &= \frac{1}{C_1} \left[(mN|\Omega|^{\frac{1}{N}} + \theta) \|\nabla S_k(u_n)\|_{N',\infty}^{p-1} + \|\phi\|_1 \right], \\
L &= \frac{1}{C_1} \left(\|\varphi\|_p^p + \frac{2^{p'/p}}{p' \alpha^{p'/p}} \|F\|_{p'}^{p'} \right).
\end{aligned} \tag{3.23}$$

By Lemma 4.1 of [13], one has

$$\begin{aligned}
& \| |\nabla S_k(u_n)|^{p-1} \|_{N', \infty} \\
& \leq C(N, p) \left[M + |\Omega|^{\frac{1}{N'} - \frac{1}{p'}} L^{\frac{1}{p'}} \right] \\
& \leq C(N, p) \left[\frac{mN|\Omega|^{\frac{1}{N}} + \theta}{C_1} \| |\nabla S_k(u_n)|^{p-1} \|_{N', \infty} + \frac{\|\phi\|_1}{C_1} + |\Omega|^{\frac{1}{N'} - \frac{1}{p'}} L^{\frac{1}{p'}} \right].
\end{aligned} \tag{3.24}$$

We can choose m to be large enough to guarantee that

$$C(N, p) \frac{mN|\Omega|^{\frac{1}{N}} + \theta}{C_1} \leq \frac{1}{2}. \tag{3.25}$$

Therefore we have

$$\| |\nabla S_k(u_n)|^{p-1} \|_{N', \infty} \leq 2C(N, p) \left[\frac{\|\phi\|_1}{C_1} + |\Omega|^{\frac{1}{N'} - \frac{1}{p'}} L^{\frac{1}{p'}} \right], \tag{3.26}$$

which completes the proof. \square

Lemma 3.2. *Let the assumptions of the Theorem 1.1 be fulfilled. Let u_n be a measurable function satisfying $T_k(u_n) \in W_0^{1,p}(\Omega)$, for every positive $k > 0$, there exist a constant C such that*

$$\| \nabla u \|_{L^p(\Omega \setminus E_k)}^p \leq C(1 + \| u \|_{L^p(\Omega \setminus E_k)}^p), \tag{3.27}$$

where E_k are defined by (2.1).

Proof. For $k > 0$, testing the equation (3.1) with $T_k(u_n)$, we get

$$\begin{aligned}
& \int_{\Omega} A_n(x, u_n, \nabla u_n) \cdot \nabla T_k(u_n) dx + \int_{\Omega} B_n(x, u_n, \nabla u_n) T_k(u_n) dx \\
& + \int_{\Omega} G_n(x, u_n) T_k(u_n) dx = \int_{\Omega} F T_k(u_n) dx.
\end{aligned} \tag{3.28}$$

The ellipticity condition (3.2) on A_n implies

$$\begin{aligned}
& \int_{\Omega} A_n(x, u_n, \nabla u_n) \cdot \nabla T_k(u_n) dx \\
& = \int_{\{|u_n| \leq k\}} A_n(x, u_n, \nabla u_n) \cdot \nabla u_n dx \\
& \geq \int_{\{|u_n| \leq k\}} \left(\alpha |\nabla u_n|^p - (T_n \zeta_0 |u_n|)^p - \varphi(x)^p \right) dx \\
& \geq \alpha \int_{\Omega} |\nabla T_k(u_n)|^p dx - \int_{\{|u_n| \leq k\}} \left(T_n(\zeta_0) |u_n| \right)^p dx - \|\varphi\|_p^p.
\end{aligned} \tag{3.29}$$

With the help of the Hölder inequality (2.5), the Sobolev embedding theorem (2.1), and the assumption (1.12), we gain

$$\begin{aligned}
& \int_{\{|u_n| \leq k\}} \left(T_n(\zeta_0) |u_n| \right)^p dx = \|(T_n(\zeta_0))u_n \chi_{\{|u_n| \leq k\}}\|_p^p \\
& \leq \|(T_m(\zeta_0))u_n \chi_{\{|u_n| \leq k\}}\|_p^p + \|(\zeta_0 - T_m(\zeta_0))u_n \chi_{\{|u_n| \leq k\}}\|_p^p \\
& \leq m^p \|u_n\|_{L^p(\Omega \setminus E_k)}^p + \|\zeta_0 - T_m(\zeta_0)\|_{N, \infty}^p \|T_k u_n\|_{p^*, p}^p \\
& \leq m^p \|u_n\|_{L^p(\Omega \setminus E_k)}^p + \|\zeta_0 - T_m(\zeta_0)\|_{N, \infty}^p S_{N, p}^p \|\nabla T_k u_n\|_p^p \\
& \leq m^p \|u_n\|_{L^p(\Omega \setminus E_k)}^p + \frac{\alpha}{2p} \|\nabla T_k u_n\|_p^p.
\end{aligned} \tag{3.30}$$

By the definition of $S_k(u_n)$ (3.11), the growth assumption (3.5) on B_n , and the Hölder inequality, one obtains

$$\begin{aligned}
& \left| \int_{\Omega} B_n(x, u_n, \nabla u_n) T_k(u_n) dx \right| \\
& \leq k \int_{\Omega} |B_n(x, u_n, \nabla u_n)| dx \\
& \leq k \left[\int_{\{|u_n| \leq k\}} T_n(\zeta_1) \cdot |\nabla u_n|^{p-1} dx + \int_{\{|u_n| > k\}} T_n(\zeta_1) \cdot |\nabla u_n|^{p-1} dx + \int_{\Omega} \phi(x) \right] \\
& \leq \frac{2^{p/p'} k^p}{p\alpha^{p/p'}} \|T_n(\zeta_1)\|_p^p + \frac{\alpha}{2p'} \|\nabla T_k(u_n)\|_p^p + k \|\phi\|_1 \\
& \quad + k \left[\int_{\{|u_n| > k\}} T_m(\zeta_1) \cdot |\nabla u_n|^{p-1} dx + \int_{\{|u_n| > k\}} (\zeta_1 - T_m(\zeta_1)) \cdot |\nabla u_n|^{p-1} dx \right] \quad (3.31) \\
& \leq \frac{2^{p/p'} k^p}{p\alpha^{p/p'}} \|T_n(\zeta_1)\|_p^p + \frac{\alpha}{2p'} \|\nabla T_k(u_n)\|_p^p + k \|\phi\|_1 \\
& \quad + kmN |\Omega|^{\frac{1}{N}} \|\nabla S_k u_n\|_{N', \infty}^{p-1} + k\theta \|\nabla S_k u_n\|_{N', \infty}^{p-1} \\
& \leq \frac{2^{p/p'} k^p}{p\alpha^{p/p'}} \|T_n(\zeta_1)\|_p^p + \frac{\alpha}{2p'} \|\nabla T_k(u_n)\|_p^p \\
& \quad + k \left[(mN |\Omega|^{\frac{1}{N}} + \theta) \|\nabla S_k u_n\|_{N', \infty}^{p-1} + \|\phi\|_1 \right],
\end{aligned}$$

where θ is given by (1.12). The sign condition (3.5) on G_n leads to

$$\int_{\Omega} G_n(x, u_n) T_k(u_n) dx \geq 0. \quad (3.32)$$

According to Young's inequality, we have the following

$$\begin{aligned}
\int_{\Omega} F T_k(u_n) dx & \leq \|F\| \|\nabla T_k(u_n)\|_p \\
& \leq \frac{2^{p'/p}}{p' \alpha^{p'/p}} \|F\|_{p'}^{p'} + \frac{\alpha}{2p} \|\nabla T_k(u_n)\|_p^p. \quad (3.33)
\end{aligned}$$

Therefore, using (3.29)-(3.33), we obtain

$$\begin{aligned}
\frac{\alpha}{2p'} \int_{\Omega} |\nabla T_k(u_n)|^p dx & \leq m^p \|u_n\|_{L^p(\Omega \setminus E_k)}^p + \frac{2^{p/p'} k^p}{p\alpha^{p/p'}} \|T_n(\zeta_1)\|_p^p + \frac{2^{p'/p}}{p' \alpha^{p'/p}} \|F\|_{p'}^{p'} \\
& \quad + k \left[(mN |\Omega|^{\frac{1}{N}} + \theta) \|\nabla S_k(u_n)\|_{N', \infty}^{p-1} + \|\phi\|_1 \right], \quad (3.34)
\end{aligned}$$

where the estimate of $\|\nabla S_k(u_n)\|_{N', \infty}^{p-1}$ is given by (3.9). This completes the proof. \square

The existence of a solution $u_n \in W_0^{1,p}(\Omega)$, which solves (3.1) can be directly achieved from the following Proposition.

Proposition 3.1. *Let (1.2)-(1.7) and (1.12) be in charge. The the approximation problem (3.1) admits a solution $u_n \in W_0^{1,p}$.*

Proof. For any fixed $F \in W^{-1,p'}(\Omega)$, let $\mathcal{F} : v_n \in W_0^{1,p}(\Omega) \mapsto u_n \in W_0^{1,p}(\Omega)$ be a mapping that takes v_n to the unique solution of the problem (3.1) is compact (see more Corollary 4.1, [9]). It is obvious that a fixed point of \mathcal{F} is a solution of the approximation (3.1). To apply the Leray-Schauder fixed point, we need to obtain an a priori estimate for the solution u_n by applying directly Lemma 2.1. A priori estimate for u_n can be obtained by first establishing a priori estimate for $|\nabla S_k(u_n)|^{p-1}$ (see Lemma 3.1), where the bound depends on both k and the

given data. We will then derive a priori estimate for $\nabla T_k(u_n)$, also depending on k and the data (see Lemma 3.2). By Lemma 2.1, we obtain

$$\|u_n\| \leq M, \quad (3.35)$$

where M is independent of n . The proof is concluded by applying the Leray–Schauder fixed point theorem (see Theorem 2.2). \square

We are now ready to proceed passing to the limit to conclude the proof of the Theorem 1.1.

3.2 Passing to the limit

The estimate of u_n given in (3.35) allows us to obtain

$$\begin{aligned} u_n &\rightharpoonup u && \text{in } W_0^{1,p}(\Omega) \text{ weakly,} \\ u_n &\rightarrow u && \text{in } L^q(\Omega) \text{ strongly for any } q < p^*, \end{aligned} \quad (3.36)$$

for some $u \in W_0^{1,p}(\Omega)$. We make use of $\gamma(u_n - u)$ (for simplicity, $\gamma(t) := \arctan t$) as a test function for approximation problem (3.1), we get

$$\begin{aligned} \int_{\Omega} A_n(x, u_n, \nabla u_n) \cdot \nabla \gamma(u_n - u) dx + \int_{\Omega} B_n(x, u_n, \nabla u_n) \gamma(u_n - u) dx \\ + \int_{\Omega} G_n(x, u_n) \gamma(u_n - u) dx = \int_{\Omega} F \gamma(u_n - u) dx, \end{aligned} \quad (3.37)$$

where $\nabla \gamma(u_n - u) = \gamma'(u_n - u)(\nabla u_n - \nabla u)$. Since $\gamma(0) = 0$, we have

$$\gamma(u_n - u) \rightharpoonup 0 \quad \text{in } W_0^{1,p}(\Omega) \text{ weakly,}$$

which implies that

$$\lim_{n \rightarrow \infty} \int_{\Omega} A_n(x, u_n, \nabla u_n) \cdot \nabla \gamma(u_n - u) dx = 0.$$

Arguing as in the proof of [9], we obtain

$$\nabla u_n \rightarrow \nabla u \quad \text{a.e. in } \Omega \quad (3.38)$$

and

$$A_n(x, u_n, \nabla u_n) \rightharpoonup A(x, u, \nabla u) \quad \text{in } L^{p'}(\Omega, \mathbb{R}^N) \text{ weakly.} \quad (3.39)$$

By the estimate (3.9), Lemma 3.1 of $|\nabla S_k(u_n)|^{p-1}$ together with the estimate (3.34) of $\nabla T_k(u_n)$, we arrive at

$$\begin{aligned} \|\nabla u_n\|^{p-1} \|_{N', \infty} &\leq \| |\nabla T_k(u_n)|^{p-1} \|_{N', \infty} + \| |\nabla S_k(u_n)|^{p-1} \|_{N', \infty} \\ &\leq C \| |\nabla T_k(u_n)|^{p-1} \|_p + \| |\nabla S_k(u_n)|^{p-1} \|_{N', \infty} \leq C. \end{aligned}$$

This results in

$$\begin{aligned} \|B_n(x, u_n, \nabla u_n)\|_1 &= \int_{\Omega} |B_n(x, u_n, \nabla u_n)| dx \\ &\leq \int_{\Omega} T_n(\zeta_1) |\nabla u_n|^{p-1} dx + \int_{\Omega} \phi(x) dx \\ &\leq \|T_n \zeta_1\|_{N, 1} \| |\nabla u_n|^{p-1} \|_{N', \infty} + \|\phi\|_1 \leq C. \end{aligned} \quad (3.40)$$

From (3.40) and the definition of B_n , we deduce that

$$B_n(x, u_n, \nabla u_n) \rightarrow B(x, u, \nabla u) \quad \text{a.e. in } \Omega. \quad (3.41)$$

For every measurable set $E \subset \Omega$, we have

$$\begin{aligned} \int_E |B_n(x, u_n, \nabla u_n)| dx &\leq \|T_n(\zeta_1)\|_{L^{N,1}(E)} \|\nabla u_n\|^{p-1}_{L^{N',\infty}(E)} + \|\phi\|_{L^1(E)} \\ &\leq \|T_n(\zeta_0)\|_{L^{N,1}(E)} + \|\phi\|_{L^1(E)} \leq C, \end{aligned} \quad (3.42)$$

which implies that

$$B_n(x, u_n, \nabla u_n) \text{ is equi-integrable.} \quad (3.43)$$

By employing the Vitali theorem, we can conclude that

$$B_n(x, u_n, \nabla u_n) \rightarrow B(x, u, \nabla u) \quad \text{in } L^1(\Omega) \text{ strongly.} \quad (3.44)$$

By the growth condition (3.5) on G_n and generalized Hölder inequality, we get

$$\begin{aligned} \|G_n(x, u_n)\|_1 &= \int_{\Omega} |G_n(x, u_n)| dx \leq \int_{\Omega} d(x) |u_n|^\lambda + \psi(x) dx \\ &\leq \|d\|_{\lambda',1} \| |u_n|^\lambda \|_{\lambda,\infty} + \|\psi\|_1 \leq C. \end{aligned} \quad (3.45)$$

Similarly, we get

$$G_n(x, u_n) \rightarrow G(x, u) \quad \text{in } L^1(\Omega) \text{ strongly.} \quad (3.46)$$

In view of (3.39), (3.44), and (3.46), we conclude that u is the solution of (1.1).

4 Regularity of the solution

In this section, we study the regularity properties of the solution of the problem (1.1). We generalize the result of [9, 14]. We show that $u \in L^{r^*}(\Omega)$ follows under appropriate assumptions on the given data.

Theorem 4.1. *Let us assume that the assumptions (1.2)-(1.7) hold with $\varphi \in L^r(\omega)$, $\phi \in L^r(\omega)$ and $F \in W^{-1, \frac{r}{p-1}}(\Omega)$ for $1 < p < r < N$. For any ζ_ε defined in (1.10), there exists $\eta = \eta(\alpha, N, p, r) > 0$ such that when*

$$\text{dist}_{L^{N,q}}(\zeta_\varepsilon, L^q(\Omega)) \leq \eta, \quad 1 \leq q \leq \infty, \quad (4.1)$$

then any solution $u \in W_0^{1,p}(\Omega)$ satisfies

$$|u|^{r^*/p^*} \in W_0^{1,p}(\Omega). \quad (4.2)$$

More precisely $u \in L^{r^*}(\Omega)$.

Proof. For $F \in W^{-1, \frac{r}{p-1}}(\Omega)$, we can take

$$F = \text{div}(|H|^{p-1}H).$$

For fixed $k > 0$, using in (1.1) the test function $S_k(u)$ defined in (3.11), we get

$$\alpha \int_{E_k} |\nabla u|^p dx \leq \int_{E_k} (\zeta_0^p |u|^p + \varphi^p) dx + \int_{E_k} (\zeta_1 |\nabla u|^{p-1} + \phi) dx + \int_{E_k} |H|^{p-1} |\nabla u| dx, \quad (4.3)$$

where E_k are defined by (2.1), $\zeta_\varepsilon, \varepsilon \in [0, 1]$ is given by (1.10).

The application of the Young's inequality gives

$$\begin{aligned} \alpha \int_{E_k} |\nabla u|^p dx &\leq \int_{E_k} (\zeta_0^p |u|^p + \varphi^p) dx + \int_{E_k} \phi dx \\ &\quad + \frac{2^{p'/p}}{\alpha^{p'/p} p'} \int_{E_k} |H|^p dx + \frac{\alpha}{2p} \int_{E_k} |\nabla u|^p dx \\ &\quad + \frac{2^{p/p'}}{\alpha^{p/p'} p} \int_{E_k} \zeta_1^p dx + \frac{\alpha}{2p'} \int_{E_k} |\nabla u|^p dx. \end{aligned} \quad (4.4)$$

Rearranging the terms on the right-hand side of (4.4), we get

$$\frac{\alpha}{2} \int_{E_k} |\nabla u|^p dx \leq \int_{E_k} \left[M_* \left(|H|^p + \zeta_1^p \right) + \zeta_0^p |u|^p + \varphi^p + \phi^p \right] dx \quad (4.5)$$

with $M_* = \max \left\{ \frac{2^{p'/p}}{\alpha^{p'/p}}, \frac{2^{p/p'}}{\alpha^{p/p'}} \right\}$.

We first multiply both sides of (4.5) by $k^{p\delta-1}$. Then, for any fixed $K > 0$, we integrate over the interval $[0, K]$ with respect to k to get

$$\frac{\alpha}{2} \int_{\Omega} |\nabla u|^p |T_K(u)|^{p\delta} dx \leq \int_{\Omega} \left[M_* \left(|H|^p + c^p \right) + b^p |u|^p + \varphi^p + \phi^p \right] |T_K(u)|^{p\delta} dx. \quad (4.6)$$

This can be rewritten as follows:

$$\begin{aligned} \left(\frac{\alpha}{2} \right)^{\frac{1}{p}} \|\nabla u |T_K(u)|^\delta\|_p &\leq M_* \|H |T_K(u)|^\delta\|_p + M_* \|\zeta_1 |T_K(u)|^\delta\|_p + \|\zeta_0 u |T_K(u)|^\delta\|_p \\ &\quad + \|\varphi |T_K(u)|^\delta\|_p + \|\phi |T_K(u)|^\delta\|_p. \end{aligned} \quad (4.7)$$

For $M > 0$, we have

$$\|\zeta_1 |T_K(u)|^\delta\|_p \leq \|(\zeta_1 - T_M \zeta_1) |T_K(u)|^\delta\|_p + M \| |T_K(u)|^\delta \|_p. \quad (4.8)$$

Moreover, we have $L^{N,1}(\Omega) \subset L^{N,\infty}(\Omega)$, thus $\zeta_1 \in L^{N,\infty}(\Omega)$. By using Hölder inequality and Sobolev embedding theorem, we get

$$\begin{aligned} \|(\zeta_1 - T_M(\zeta_1)) |T_K(u)|^\delta\|_p &\leq \|\zeta_1 - T_M(\zeta_1)\|_{N,\infty} \| |T_K(u)|^\delta \|_{p^*,p} \\ &\leq \|\zeta_1 - T_M(\zeta_1)\|_{N,\infty} S_{N,p} \delta \| |\nabla u |T_K(u)|^\delta \|_p \\ &\leq N \|\zeta_1 - T_M(\zeta_1)\|_{N,1} S_{N,p} \delta \| |\nabla u |T_K(u)|^\delta \|_p. \end{aligned} \quad (4.9)$$

Here we applied the following

$$\|\zeta_1 - T_M(\zeta_1)\|_{N,\infty} \leq N \|\zeta_1 - T_M(\zeta_1)\|_{N,1}.$$

Similarly, for $\|bu |T_K(u)|^\delta\|_p$, we have

$$\|\zeta_0 u |T_K(u)|^\delta\|_p \leq \|\zeta_1 - T_M(\zeta_1)\|_{N,\infty} S_{N,p} (1 + \delta) \| |\nabla u |T_K(u)|^\delta \|_p + M \|u |T_K(u)|^\delta\|_p. \quad (4.10)$$

We assume that

$$\|\zeta_1 - T_M(\zeta_1)\|_{N,\infty} S_{N,p} (1 + \delta) \leq \left(\frac{\alpha}{8} \right)^{\frac{1}{p}}. \quad (4.11)$$

We set

$$L^p = |H|^p + |u|^p + \varphi^p + \phi^p. \quad (4.12)$$

From (4.7) and using (4.9)-(4.10), we get

$$\|\nabla u |T_K u|^\delta\|_p \leq C \|L |T_K u|^\delta\|_p. \quad (4.13)$$

We apply the Hölder inequality on the right-hand side of (4.13). This leads to

$$\begin{aligned} \|\nabla u |T_K u|^\delta\|_p &\leq C \|L\|_r \| |T_K(u)|^\delta \|_{\delta \frac{rp}{r-p}} \\ &\leq C \|L\|_r \| |\nabla |T_K(u)|^{\frac{r^*}{p^*}} \|_p^{\delta \frac{p^*}{r^*}} \\ &\leq C \|L\|_r \| |\nabla u |T_K(u)|^\delta \|_p^{\frac{\delta}{\delta+1}}. \end{aligned} \quad (4.14)$$

Merging in (4.14), we get

$$\|\nabla u |T_K(u)|^\delta\|_p^{\frac{p^*}{\delta}} \leq C \|L\|_r. \quad (4.15)$$

Letting $K \rightarrow +\infty$ and taking into account the estimate (4.15), we have

$$\|\nabla u|u|^\delta\|_{\frac{p^*}{r^*}} \leq C\left(\|H\|_r + \|\varphi\|_r + \|\phi\|_r\right), \quad (4.16)$$

i.e.

$$\|\nabla|u|^{\frac{r^*}{p^*}}\|_p \leq C\left(\|H\|_r + \|\varphi\|_r + \|\phi\|_r\right)^{\frac{r^*}{p^*}}. \quad (4.17)$$

Therefore, as long as $u \in L^r(\Omega)$ implies $|u|^{\frac{r^*}{p^*}} \in W_0^{1,p}(\Omega)$. This concludes our proof. Clearly, the above argument works directly in the case $r < p^*$. For the case $r > p^*$, we use a bootstrap procedure. \square

References

- [1] P. Bénilan, L. Boccardo, T. Gallouët, R. Gariepy, M. Pierre, J. -L. Vazquez, *An L^1 - theory of existence and uniqueness of solutions of nonlinear elliptic equations*, Ann. Scuola Norm. Sup. Pisa 22 (1995) 241-273.
- [2] M. F. Betta, A. Mercado, F. Murat, M. M. Porzio, *Existence of renormalized solution to nonlinear elliptic equations with a lower-order term and right-hand side a measure*, J. Math. Pures Appl. 82 (2003) 90-124.
- [3] L. Boccardo, *Finite energy solutions of nonlinear Dirichlet problems with discontinuous coefficients*, Boll. Un. Mat. It. 5 (2012) 357-368.
- [4] L. Boccardo, D. Giachetti, J. -I. Diaz, F. Murat, *Existence and regularity of renormalized solutions to some elliptic problems involving derivations of nonlinear terms*, J. Differential Equations. 106 (1993) 215-237.
- [5] L. Boccardo, T. Gallouët, *Nonlinear elliptic equations with right-hand side measure*, Comm. Partial Differential Equations 17 (1992) 641-955.
- [6] F. E. Browder, *Existence theorems for nonlinear partial differential equations*, Proceedings of Symposia in Pure Mathematics, Vol. 16, S. S. CHERN and S. SMALE Eds., A.M.S., Providence, 1970, 1-60.
- [7] M. Carozza, C. Sbordone, *The distance to L^∞ in some function spaces and applications*, Differential Integral Equations 10 (1997) 599-607.
- [8] G. R. Cirmi, S. D'Asero, S. Leonardi, M. M. Porzio, *Local regularity results for solutions of linear elliptic equations with drift term*, Adv. Calc. Var. 15 1 (2022) 19-32.
- [9] F. Farroni, L. Greco, G. Moscariello, *Noncoercive quasilinear elliptic operators with singular lower-order terms*, Calc. Var. 60 83 (2021).
- [10] M. Giaquinta, E. Giusti, *Quasi-minima*, Ann. Inst. H. Poincaré Anal. Non Linéaire 1 2 (1984) 79-107.
- [11] D. Gilbarg, N.S. Trudinger, *Elliptic partial differential equations of second order*, 249. Springer, New York, (2014).
- [12] F. Giannetti, L. Greco, G. Moscariello, *Linear elliptic equations with lower-order terms*, Differential Integral Equations 26 5-6 (2013) 623-638.
- [13] O. Guibé, A. Mercado, *Existence of renormalized solutions to nonlinear elliptic equations with two lower-order terms and measure data*, Trans. Amer. Math. Soc. 360 2 (2008) 643-669.
- [14] L. Greco, G. Moscariello, G. Zecca, *Regularity for solutions to nonlinear elliptic equations*, Differential and Integral Equations 26 9-10 (2013) 1005-1113.

- [15] L. Greco, G. Moscarciello, *An embedding theorem in Lorentz–Zygmund spaces*, Potential Anal. 5 (1996) 581-590.
- [16] L. Greco, G. Moscarciello, G. Zecca, *An obstacle problem for noncoercive operators*, Abstr. Appl. Anal. (2015) 1-8.
- [17] L. Greco, G. Moscarciello, G. Zecca, *Very weak solutions to elliptic equations with singular convection term*, J. Math. Anal. 457 2 (2018) 1376-1387.
- [18] G. G. Lorentz, *Some new function spaces*, Ann. Math. 51 (1950) 37-55.
- [19] G. Moscarciello, *Existence and uniqueness for elliptic equations with lower-order terms*, Adv. Calc. Var. 4 4 (2011) 421-444.
- [20] H. Kim, T. P. Tsai, *Existence, uniqueness, regularity results for elliptic equation with drift terms in critical weak spaces*, SIAM J. Math. Anal. 52 2 (2020) 1146-1191.
- [21] J. Leray, J. L. Lions, *Quelques résultats de Visik sur le problèmes elliptiques non linéaires par les méthodes de Minty-Browder*, Bull. Soc. Math. France 93 (1965) 97-107.
- [22] G. D. Maso, F. Murat, L. Orsina, A. Prignet, *Renormalised solutions for elliptic equations with general measure data*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. 28 (1999) 741-808.
- [23] R. O’Neil, *Convolution operators and $L(p, q)$ spaces*, Duke math. J. 30 (1963) 129-142.
- [24] A. Porretta, *Existence results for nonlinear parabolic equations via strong convergence of truncations*, Ann. Mat. Pura Appl. 177 4 (1999) 143-172.
- [25] A. Porretta, *Weak solutions to Fokker–Planck equations and mean field games*, Arch. Ration. Mech. Anal. 216 1 (2015) 1–62.
- [26] M. M. Pozrio, *Existence, uniqueness and behavior of solutions for a class of nonlinear parabolic problems*, Nonlinear Anal. TMA 74 (2011) 5359-5382.
- [27] M. M. Pozrio, *On uniform and decay estimates for unbounded solutions of partial differential equations*, J. Differential Equations 259 (2015) 6960-7011.
- [28] T. Radice, G. Zecca, *Existence and uniqueness for elliptic equations with unbounded coefficients*, Ricerche Mat. 63 2 (2014) 355-367.
- [29] G. Stampacchia., *Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinuous*, Ann. Inst. Fourier (Grenoble) 15 (1965) 189-258.
- [30] G. Zecca, *Existence and uniqueness for nonlinear elliptic equations with lower-order terms*, Nonlinear Anal. Theory Methods Appl. 75 (2012) 899-912.