# Debiased calibration estimation using generalized entropy in survey sampling

Yonghyun Kwon

Department of Mathematics, Korea Military Academy, Seoul, Republic of Korea

and

Jae Kwang Kim

Department of Statistics, Iowa State University, Iowa, USA

and

Yumou Qiu

School of Mathematical Sciences, Peking University, Beijing, China

#### Abstract

Incorporating the auxiliary information into the survey estimation is a fundamental problem in survey sampling. Calibration weighting is a popular tool for incorporating the auxiliary information. The calibration weighting method of Deville and Särndal (1992) uses a distance measure between the design weights and the final weights to solve the optimization problem with calibration constraints. This paper introduces a novel framework that leverages generalized entropy as an objective function for optimization, where the design weights are employed in the debiasing constraint to ensure design consistency, rather than being part of the objective function. This innovative calibration framework is particularly attractive due to its generality and its ability to generate more efficient calibration weights compared to traditional methods based on Deville and Särndal (1992). Furthermore, under Poisson sampling, we identify the optimal choice of the generalized entropy function that achieves the minimum asymptotic variance across various choices of the generalized entropy function under the same constraints. Asymptotic properties, such as design consistency and asymptotic normality, are presented rigorously. The results from a limited simulation study are also presented. We demonstrate a real-life application using agricultural survey data collected from Kynetec, Inc.

*Keywords:* Empirical likelihood, generalized regression estimation, selection bias.

# 1 Introduction

Many scientific studies often rely on analyzing datasets consisting of samples from target populations. The fundamental assumption when using such data is that the sample accurately represents the target population, ensuring representativeness. Probability sampling is a classical tool for securing a representative sample from a target population. Once a probability sample is obtained, researchers can employ design consistent estimation methods, such as Horvitz-Thompson estimation, to estimate the parameters of the population. Statistical inferences, such as confidence intervals, can be made from the probability sample using the large sample theory (Fuller, 2009).

However, the Horvitz-Thompson (HT) estimator is not necessarily efficient as it does not incorporate all available information effectively. Improving the efficiency of the HT estimator is one of the fundamental problems in survey sampling. A classical approach to improving efficiency involves the use of additional information obtained from external sources, such as census data. To incorporate auxiliary information into the final estimate, the design weights are adjusted to meet the benchmarking constraints imposed by the auxiliary information. This weight modification method to satisfy the benchmarking constraint is known as calibration weighting. When the auxiliary variables used for calibration are correlated with the study variable of interest, the resulting calibration estimator is more efficient than the HT estimator. While calibration weighting also plays a role in mitigating selection bias in nonprobability samples, as highlighted by Dever and Valliant (2016) and Elliott and Valliant (2017), our discussion will focus on its application within probability samples.

The literature on calibration weighting is very extensive. Isaki and Fuller (1982) used a

linear regression superpopulation model to construct the regression calibration weights and showed that the resulting estimator is optimal in the sense that its anticipated variance achieves the lower bound of Godambe and Joshi (1965). Deville and Särndal (1992) developed a unified framework for calibration estimation and showed the asymptotic equivalence between the calibration estimator and the generalized regression (GREG) estimator. Wu and Sitter (2001) explicitly utilized the regression model to develop what is known as model calibration. Breidt and Opsomer developed a series of nonparametric regression estimators (Breidt et al., 2005; Breidt and Opsomer, 2017) that can be understood as nonparametric calibration weighting. Montanari and Ranalli (2005) also developed a framework for nonparametric model calibration using neural network models. Dagdoug et al. (2023) employ random forest to develop a nonparametric model calibration. See Wu and Lu (2016); Haziza and Beaumont (2017); Devaud and Tillé (2019) for a comprehensive review of calibration weighting methods in survey sampling.

Most existing calibration weighting methods are grounded in the framework established by Deville and Särndal (1992), which utilizes a distance measure between the design weights and the final weights to address the optimization problem within the calibration constraints. In this paper, we introduce a novel alternative framework that does not use the distance measure between two weights. Specifically, the proposed approach employs generalized entropy as an objective function for optimization, and the design weights are incorporated into the constraints rather than the objective function to solve the calibration optimization problem.

The constraint involving the design weights is designed to mitigate selection bias, whereas the benchmarking constraint aims to decrease variance. To differentiate between these two calibration constraints, the one that incorporates design weights is called the debiasing constraint, to emphasize its role in controlling selection bias. Although generalized entropy has been developed as a tool for efficient estimation (Newey and Smith, 2004), current literature lacks a discussion of the debiasing constraint under selection bias.

The idea of employing a debiasing constraint within the calibration weighting framework is not entirely new. Qin et al. (2002) is perhaps the first attempt to correct selection bias using empirical likelihood in the context of missing data. Berger and Torres (2016) employed debiasing constraints in empirical likelihood for survey sampling contexts. Chapter 2 of Fuller (2009) discussed the incorporation of the debiasing constraint into linear regression models.

This paper aims to construct the debiasing constraint in generalized entropy calibration and introduce a unified framework for debiased calibration that reduces selection bias and improves estimation efficiency in survey sampling scenarios. The proposed calibration estimator is asymptotically equivalent to a generalized regression estimator using an augmented regression model, where the additional covariate comes from the debiasing constraint. If the additional debiasing constraint in the augmented regression model is significant, then the resulting calibration estimator is more efficient than the calibration estimator of Deville and Särndal (1992). Our approach is based on generalized entropy, and the empirical likelihood approach of Qin et al. (2002) is a special case of our unified approach.

Furthermore, based on asymptotic theory, we formally show that the cross-entropy function is the optimal entropy function in the sense that it attains the minimum variance among the different choices of the generalized entropy function under the same constraints. Thus, in addition to generality, we also present an interesting result for asymptotic optimality under Poisson sampling in the design-based framework.

The debiasing constraint implicitly assumes that the design weights are available through-

out the finite population. When design weights are unavailable outside the sample, the debiasing constraint cannot be used directly. In this case, we can treat the population mean of the debiasing control as another unknown parameter and develop a modified generalized entropy approach that simultaneously estimates the population mean of the debiasing control and the calibration weights. Asymptotic linearization of the modified generalized entropy estimator is obtained under some special cases, and a comparison with Deville and Särndal (1992)'s estimator is discussed. The details are presented in Section 6.

The paper is organized as follows. In Section 2, the basic setup and the research problem are introduced. In Section 3, we present the proposed method using a generalized entropy calibration and give an illustration using empirical likelihood as a special case. In Section 4, the asymptotic properties of the proposed method are rigorously derived, and a consistent variance estimator is also presented. In Section 5, we show that the optimal entropy function in the context of calibration estimation is achieved with the cross-entropy function. In Section 6, we present a modification of the proposed method when the design weights are not available outside the sample. Results of a limited simulation study are presented in Section 7. In Section 8, we demonstrate a real-life application using agricultural survey data collected from Kynetec. Some concluding remarks are made in Section 9. All the technical proofs are relegated to the supplementary material (SM).

#### 2 Basic setup

Consider a finite population  $U = \{1, \dots, N\}$  of N units, with known N. A sample A of size n is selected from the finite population using a given probability sampling design. We

use  $\delta_i$  to denote the sampling indicator variable of unit i such that  $\delta_i = 1$  if  $i \in A$  and  $\delta_i = 0$  otherwise. We assume that first-order inclusion probabilities  $\{\pi_i : i \in U\}$  are available throughout the population. We will relax this assumption later in Section 6.

Let  $y_i$  be the study variable of interest, available only in sample A. Our goal is to estimate the finite population total  $\theta_N = \sum_{i \in U} y_i = \sum_{i=1}^N y_i$  from sample A. We consider a class of linear estimators

$$\hat{\theta} = \sum_{i \in A} \omega_i y_i \tag{2.1}$$

where  $\omega_i$  does not depend on  $\{y_i : i \in A\}$ . The Horvitz-Thompson (HT) estimator is a natural way to estimate  $\theta_N$  using  $\omega_i = \pi_i^{-1}$ , which is design-unbiased but can be inefficient.

In many practical situations, in addition to the study variable  $y_i$ , we observe *p*-dimensional auxiliary variables  $\boldsymbol{x}_i = (x_{i1}, \ldots, x_{ip})^{\top}$  with known population totals from external sources. In this case, to get external consistency, we often require that the final weights satisfy

$$\sum_{i \in A} \omega_i \boldsymbol{x}_i = \sum_{i \in U} \boldsymbol{x}_i.$$
(2.2)

Constraint (2.2) is often called a calibration constraint or a benchmarking constraint. Generally speaking, the calibration estimator is more efficient than the HT estimator when the study variable y of interest is related to the auxiliary variable  $\boldsymbol{x}$ . In particular, if  $y_i = \boldsymbol{x}_i^{\top} \boldsymbol{\beta}$ holds for some  $\boldsymbol{\beta}$ , then the resulting calibration is perfect in the sense that its mean squared error is zero. Thus,  $y_i = \boldsymbol{x}_i^{\top} \boldsymbol{\beta} + e_i$  can be viewed as a working regression model for calibration at the population level. However, the calibration condition itself does not guarantee design consistency.

To achieve the design consistency of the calibration estimator, Deville and Särndal (1992) proposed solving an optimization problem that minimizes the distance between the calibration weights  $\boldsymbol{\omega} = (\omega_i)_{i \in A}$  and the design weights  $\boldsymbol{d} = (\pi_i^{-1})_{i \in A}$ 

$$D(\boldsymbol{\omega}, \boldsymbol{d}) = \sum_{i \in A} d_i G(\omega_i / d_i)$$
(2.3)

subject to the calibration constraints in (2.2), where  $d_i = \pi_i^{-1}$  for  $i \in A$  and  $G(\cdot) : \mathcal{V} \to \mathbb{R}$  is a nonnegative function that is strictly convex, differentiable, and G'(1) = 0. The domain  $\mathcal{V}$  of  $G(\cdot)$  is an open interval in  $\mathbb{R}$ . The distance measure  $D(\boldsymbol{\omega}, \boldsymbol{d})$  in (2.3) serves as the divergence between two discrete measures  $\boldsymbol{\omega}$  and  $\boldsymbol{d}$ . For example,  $G(\omega) = \omega \log(\omega) - \omega + 1$ , with domain  $\mathcal{V} \subseteq (0, \infty)$ , corresponds to the Kullback-Leibler divergence, while  $G(\omega) = (\omega - 1)^2$ , with the domain  $\mathcal{V} \subseteq (-\infty, \infty)$ , corresponds to the Chi-squared distance from 1.

Let  $\hat{\omega}_{\mathrm{DS},i}$  be the solution to the above optimization problem, and let  $\hat{\theta}_{\mathrm{DS}} = \sum_{i \in A} \hat{\omega}_{\mathrm{DS},i} y_i$ be the resulting estimator. Under some conditions,  $\hat{\theta}_{\mathrm{DS}}$  is asymptotically equivalent to the generalized regression (GREG) estimator given by

$$\hat{\theta}_{\text{greg}} = \sum_{i \in U} \boldsymbol{x}_i^\top \hat{\boldsymbol{\beta}} + \sum_{i \in A} d_i (y_i - \boldsymbol{x}_i^\top \hat{\boldsymbol{\beta}})$$
(2.4)

where

$$\hat{\boldsymbol{\beta}} = \left(\sum_{i \in A} d_i \boldsymbol{x}_i \boldsymbol{x}_i^{\mathsf{T}}\right)^{-1} \sum_{i \in A} d_i \boldsymbol{x}_i y_i.$$
(2.5)

Note that the GREG estimator in (2.4) can be expressed as the sum of two terms; the prediction term and the bias correction term. The bias correction term is calculated from the sample using the HT estimation of the negative bias of the prediction estimator. The bias-corrected prediction estimator is also called a debiased prediction estimator in the causal inference literature (Athey et al., 2018). The debiasing property comes from the fact that the objective function in (2.3) is minimized at  $\omega_i = d_i$  if the calibration constraints are satisfied under these design weights. When the sample size is sufficiently large, by the law of large

numbers, the calibration constraints are nearly satisfied with the design weights. Thus, the final calibration weights should converge to the design weights as the sample size increases.

Although the bias correction term has the clear advantage of eliminating bias, it can have the disadvantage of increasing variance because the bias term is estimated by the HT estimation method. In the following sections, we propose another approach to the debiased calibration estimator that is often more efficient than the classical calibration estimator that uses (2.3). The basic idea is to incorporate bias correction into the calibration constraint, which will be called a debiasing constraint. That is, we can simply use debiasing and benchmarking constraints in the calibration weighting. The debiasing constraint is to control the selection bias, while the benchmarking constraint is to control the efficiency. The use of a debiasing constraint in calibration weighting is similar in spirit to interval bias calibration (IBC) in prediction estimation (Firth and Bennett, 1998). The debiasing constraint in the calibration estimator plays the role of IBC in the prediction estimator.

# 3 Methodology

Instead of minimizing the weight distance measure in (2.3), we now consider maximizing the generalized entropy (Gneiting and Raftery, 2007) that does not employ the design weights:

$$H(\boldsymbol{\omega}) = -\sum_{i \in A} G(\omega_i), \qquad (3.1)$$

where  $G(\cdot) : \mathcal{V} \to \mathbb{R}$  is a strictly convex and differentiable function with an open domain  $\mathcal{V}$  and  $\boldsymbol{\omega}$  denotes the vector of  $\{\omega_i : i \in A\}$ . Note that the function  $G(\cdot)$  in (3.1) is more general than that in (2.3). It does not need to be nonnegative or satisfy G'(1) = 0 as in

(2.3). The empirical likelihood is a special case of (3.1) with  $G(\omega_i) = -\log \omega_i$ , while the Shannon-entropy uses  $G(\omega_i) = \omega_i \log \omega_i$ .

To incorporate design information and guarantee design consistency, in addition to the benchmarking calibration constraints in (2.2), we propose including the following design calibration constraint

$$\sum_{i \in A} \omega_i g(d_i) = \sum_{i \in U} g(d_i), \tag{3.2}$$

where  $g(\omega) = dG(\omega)/d\omega$  denotes the first-order derivative of  $G(\cdot)$ . The constraint in (3.2) is the key constraint to make the proposed calibration estimator design consistent, which is called the *debiasing* calibration constraint. The mapping  $d_i \mapsto g(d_i)$  is called the debiasing transformation. Including the debiasing constraint in the generalized entropy function is our main proposal. While the primary reason for including the debiasing constraint is to achieve design consistency, we can also improve the efficiency of the resulting estimator as it incorporates an additional covariate in the working regression model for calibration.

Our goal is to find the calibration weights  $\boldsymbol{\omega}$  that maximize the generalized entropy in (3.1) under the calibration constraints in (2.2) and (3.2), so that the resulting weighted estimator is design consistent and efficient. The optimization problem of interest can be formulated as follows

$$\hat{\boldsymbol{\omega}} = \underset{\omega_i \in \mathcal{V}}{\operatorname{arg\,min}} \sum_{i \in A} G(\omega_i) \quad \text{subject to} \quad \sum_{i \in A} \omega_i \boldsymbol{z}_i = \sum_{i \in U} \boldsymbol{z}_i, \tag{3.3}$$

where  $\boldsymbol{z}_i^{\top} = (\boldsymbol{x}_i^{\top}, g(d_i)) \in \mathbb{R}^{p+1}$ . Note that  $\hat{\boldsymbol{\omega}}$  is the vector of  $\{\hat{\omega}_i : i \in A\}$ . Let  $\Omega_A = \{\boldsymbol{\omega} = (\omega_i)_{i \in A} : \sum_{i \in A} \omega_i \boldsymbol{z}_i = \sum_{i \in U} \boldsymbol{z}_i \text{ and } \omega_i \in \mathcal{V}\}$ . If  $\Omega_A$  is nonempty, a solution  $\hat{\boldsymbol{\omega}}$  to (3.3) exists, and the proposed calibration estimator of the population total  $\theta_N$  is constructed as

$$\hat{\theta}_{cal} = \sum_{i \in A} \hat{\omega}_i y_i. \tag{3.4}$$

The Karush-Kuhn-Tucker (KKT) conditions for the primal problem in (3.3) are

$$g(\omega_i) - \boldsymbol{\lambda}^{\top} \boldsymbol{z}_i = 0 \text{ and } \sum_{i \in A} \omega_i \boldsymbol{z}_i = \sum_{i \in U} \boldsymbol{z}_i,$$
 (3.5)

where  $\boldsymbol{\lambda} = (\boldsymbol{\lambda}_1^{\top}, \lambda_2)^{\top}$  are the Lagrange multipliers for the dual problem of (3.3). From Section 5 of Boyd and Vandenberghe (2004), if  $\Omega_A$  is nonempty, Slater's condition is satisfied and strong duality holds. Then,  $\hat{\boldsymbol{\omega}}$  is the solution to (3.3) if and only if

$$\hat{\omega}_i = \hat{\omega}_i(\hat{\boldsymbol{\lambda}}) = g^{-1} \{ \hat{\boldsymbol{\lambda}}_1^\top \boldsymbol{x}_i + \hat{\lambda}_2 g(d_i) \},$$
(3.6)

and  $\hat{\boldsymbol{\lambda}} = (\hat{\boldsymbol{\lambda}}_1^{\top}, \hat{\boldsymbol{\lambda}}_2)^{\top}$  satisfy the KKT conditions in (3.5). Let  $g(\mathcal{V}) = \{g(\omega) : \omega \in \mathcal{V}\}$  and  $\Lambda_A = \{\boldsymbol{\lambda} : \boldsymbol{\lambda}^{\top} \boldsymbol{z}_i \in g(\mathcal{V}) \text{ for all } i \in A\}$ . Let  $F(u) = -G(g^{-1}(u)) + g^{-1}(u)u$  for  $u \in g(\mathcal{V})$  be the convex conjugate function of  $G(\omega)$ , and f(u) = dF(u)/du be the first-order derivative of  $F(\omega)$ . By the chain rule, we can obtain  $f(u) = g^{-1}(u)$  for  $u \in g(\mathcal{V})$ . It is shown in the SM that  $\Lambda_A$  is non-empty and the solution to (3.5) can be obtained by the optimization problem

$$\hat{\boldsymbol{\lambda}} = \operatorname*{arg\,min}_{\boldsymbol{\lambda} \in \Lambda_A} \sum_{i \in A} F(\boldsymbol{\lambda}^{\top} \boldsymbol{z}_i) - \boldsymbol{\lambda}^{\top} \sum_{i \in U} \boldsymbol{z}_i$$
(3.7)

as  $n, N \to \infty$ . While  $\boldsymbol{\omega}$  is *n*-dimensional,  $\boldsymbol{\lambda}$  is (p+1)-dimensional, which implies that the dual problem in (3.7) is easier to compute compared to the primal problem of (3.3). Thanks to the existence of the solution to (3.7) with probability approaching to 1 (*w.p.a.1*), this implies that  $\Omega_A$  is non-empty and the solution to (3.3) exists *w.p.a.1*.

The expression of  $\hat{\omega}_i$  in (3.6) helps to understand the role of the debiasing constraint (3.2) in the proposed procedure. Since the calibration equations in (2.2) and (3.2) are satisfied with  $\omega_i = d_i = \pi_i^{-1}$  as  $n, N \to \infty$ , the difference between the solution  $\hat{\omega}_i$  and  $d_i$  will converge to zero for sufficiently large sample sizes. From the expression of  $\hat{\omega}_i(\hat{\lambda})$  in (3.6), note that  $|\hat{\omega}_i(\hat{\lambda}) - d_i| \to 0$  in probability if  $\hat{\lambda}_1 \to 0$  and  $\hat{\lambda}_2 \to 1$  in probability as  $n, N \to \infty$ . Therefore, as long as the above convergence of  $\hat{\lambda}$  is satisfied, the resulting calibration estimator  $\hat{\theta}_{cal}$  is design consistent. A rigorous theoretical justification of the design consistency of the proposed estimator is presented in the next section.

Deville and Särndal (1992) showed that the calibration weights  $\hat{\omega}_{\text{DS},i}$  using the divergence measure in (2.3) can be expressed as

$$\hat{\omega}_{\mathrm{DS},i} = \hat{\omega}_{\mathrm{DS},i}(\hat{\boldsymbol{\lambda}}_{\mathrm{DS},1}) = d_i g^{-1}(\hat{\boldsymbol{\lambda}}_{\mathrm{DS},1}^\top \boldsymbol{x}_i)$$
(3.8)

where  $\hat{\lambda}_{DS,1}$  is the Lagrange multiplier to satisfy the calibration constraints in (2.2). They also showed that  $\hat{\lambda}_{DS,1} \to \mathbf{0}$  in probability as  $n, N \to \infty$  under mild regularity conditions. By comparing (3.6) with (3.8), we can see that the main distinction between the divergence calibration method in Deville and Särndal (1992) and the proposed entropy calibration approach is in the way that the selection probabilities  $\{d_i\}$  are utilized. The divergence calibration method uses  $\{d_i\}$  of the sampled units only, which are incorporated directly into the objective function. However, the proposed method uses the selection probabilities in the entire population by imposing the debiasing calibration constraint in (3.2). An additional parameter  $\lambda_2$  is introduced to reflect the debiasing constraint in (3.2). In the case where the design weights  $\{\pi_i : i \notin A\}$  are not available, we can treat  $\sum_{i \in U} g(d_i)$  as an unknown parameter and estimate it from the sample. See Section 6 for more discussion.

Empirical Likelihood Example. The empirical likelihood (Owen, 1988) objective function of (3.1) uses  $G(\omega) = -\log(\omega)$ . Since  $g(\omega) = -\omega^{-1}$  in this case, the debiasing constraint in (3.2) takes the form  $\sum_{i \in A} \omega_i \pi_i = \sum_{i \in U} \pi_i = E(n)$ . When the sample size n is fixed,  $\sum_{i \in U} \pi_i$ is equal to n. The proposed calibration method using the empirical likelihood objective function solves the optimization problem

$$\hat{\boldsymbol{\omega}} = \operatorname*{arg\,max}_{\omega_i > 0} \ell^{\mathrm{EL}}(\boldsymbol{\omega}) = \operatorname*{arg\,max}_{\omega_i > 0} \sum_{i \in A} \log \omega_i \text{ subject to } \sum_{i \in A} \omega_i \pi_i = \sum_{i \in U} \pi_i \text{ and } (2.2).$$
(3.9)

The logarithm in  $\log(\omega_i)$  ensures that  $\omega_i$  are all positive. The use of (3.9) for complex survey design was considered by Berger and Torres (2016). For comparison, the empirical likelihood weight of Deville and Särndal (1992) is obtained by minimizing the Kullback-Leibler divergence  $D(\boldsymbol{w}, \boldsymbol{d}) = \sum_{i \in A} d_i \log (d_i/\omega_i)$ , which is equivalent to maximizing the pseudo empirical likelihood (PEL) proposed by Chen and Sitter (1999):

$$\ell^{\text{PEL}}(\boldsymbol{\omega}) = \sum_{i \in A} d_i \log(\omega_i)$$
(3.10)

subject to the calibration constraint in (2.2).

Another example is the shifted exponential tilting entropy  $G(\omega) = (\omega - 1) \log(\omega - 1) - \omega$ . In this case, the proposed calibration weights satisfy  $\hat{\omega}_i = 1 + \exp(\mathbf{z}_i^{\top} \hat{\boldsymbol{\lambda}})$  for the dual parameter  $\hat{\boldsymbol{\lambda}}$ , where the inverse weights have the logistic functional form. Other examples of generalized entropies and their debiasing transformation function  $g(d_i)$  can be found in Table 1.

Let  $n_0 = \mathbb{E}(n) = \sum_{i \in U} \pi_i$ . Under the case of  $\pi_i \to 0$  and  $n_0 = o(N)$  in the asymptotic setup, the sample size n would be much smaller than N and  $d_i \to \infty$  which makes  $g(d_i)$  a trivial value for all i asymptotically. To cover this case, we use a scaled weight  $d_i^* = n_0 d_i/N$ in the augmented calibration and modify the proposed approach in (3.3) as

$$\hat{\boldsymbol{\omega}} = \underset{n_0 \omega_i / N \in \mathcal{V}}{\arg\min} \sum_{i \in A} G\left(\frac{n_0}{N} \omega_i\right) \text{ subject to } \sum_{i \in A} \omega_i \boldsymbol{z}_i = \sum_{i \in U} \boldsymbol{z}_i, \quad (3.11)$$

where  $\boldsymbol{z}_i^{\top} = (\boldsymbol{x}_i^{\top}, g_i^*)$  and  $g_i^* = g(d_i^*)$  for the case of scaled weights. Note that we do not distinguish the notation  $\boldsymbol{z}_i$  for the unscaled and scaled approaches since we can define  $\tilde{G}(\omega) = G(n_0\omega/N)$  as a new entropy function for (3.11) if  $n_0$  and N are of the same order.

Entropy	$G(\omega)$	$g_i = g(d_i)$	$1/g'(d_i)$	Domain $\mathcal{V}$
Squared loss	$\omega^2/2$	$d_i$	1	$(-\infty,\infty)$
Empirical likelihood	$-\log \omega$	$-d_i^{-1}$	$d_i^2$	$(0,\infty)$
Exponential tilting	$\omega \log(\omega) - \omega$	$\log d_i$	$d_i$	$(0,\infty)$
Shifted Exp tilting	$(\omega - 1)\log(\omega - 1) - \omega$	$\log(d_i - 1)$	$d_i - 1$	$(1,\infty)$
Cross entropy	$(\omega - 1)\log(\omega - 1) - \omega\log(\omega)$	$\log(1-d_i^{-1})$	$d_i^2 - d_i$	$(1,\infty)$
Pseudo-Huber	$M^{2}\{1+\left(\omega/M\right)^{2}\}^{1/2}$	$d_i \{1 + (d_i/M)^2\}^{-1/2}$	$(d_i/g_i)^3$	$(-\infty,\infty)$
Hellinger distance	$-4\omega^{1/2}$	$-2d_i^{-1/2}$	$d_i^{3/2}$	$(0,\infty)$
Inverse	$1/(2\omega)$	$-d_{i}^{-2}/2$	$d_i^3$	$(0,\infty)$
Rényi entropy	$r^{-1}(r+1)^{-1}\omega^{r+1}$	$r^{-1}d_i^r$	$d_i^{-r+1}$	$(0,\infty)$

Table 1: Examples of generalized entropies with the corresponding  $G(\omega)$ , debiasing transformation function  $g_i = g(d_i) = g(\pi_i^{-1})$  and the regression weight  $1/g'(d_i)$  in (4.2), where  $g'(d_i)$  is the first-order derivative of  $g(d_i)$  with respect to  $d_i$  and the Rényi entropy requires  $r \neq 0, -1$ .

## 4 Statistical properties

To examine the asymptotic properties of the proposed entropy calibration estimator  $\hat{\theta}_{cal}$ , we consider an increasing sequence of finite populations and samples as in Isaki and Fuller (1982). In this section, we only consider the proposed estimator with the scaled weight in (3.11). The theoretical results also cover the case of unscaled weight in (3.3) if  $n_0$  and N are of the same order by setting  $\tilde{G}(\omega) = G(n_0\omega/N)$ . Let  $\lambda_0 = (\lambda_{10}^{\top}, \lambda_{20})^{\top}$ , where  $\lambda_{10} = \mathbf{0}$  and  $\lambda_{20} = 1$ . We make the following assumptions for our analysis. [A1]  $G(\omega)$  is strictly convex and twice continuously differentiable with an open domain  $\mathcal{V}$ .

[A2] There exist positive constants  $c_1, c_2 \in \mathcal{V}$  such that  $c_1 < N\pi_i/n_0 < c_2$  for  $i = 1, \dots, N$ .

[A3] Let  $\pi_{ij}$  be the joint inclusion probability of units *i* and *j* and  $\Delta_{ij} = \pi_{ij} - \pi_i \pi_j$ . Assume

$$\limsup_{N \to \infty} N^2 n_0^{-1} \max_{i,j \in U: i \neq j} \left| \Delta_{ij} \right| < \infty.$$

[A4] Assume  $\Sigma_{\boldsymbol{z}} = \lim_{N \to \infty} \sum_{i \in U} \boldsymbol{z}_i \boldsymbol{z}_i^\top / N$  exists and positive definite, the average 4th moment of  $(y_i, \boldsymbol{x}_i^\top)$  is finite such that  $\limsup_{N \to \infty} \sum_{i=1}^N ||(y_i, \boldsymbol{x}_i^\top)||^4 / N < \infty$ , and  $\Gamma(\boldsymbol{\lambda}) = \lim_{N \to \infty} \sum_{i \in U} f'(\boldsymbol{\lambda}^\top \boldsymbol{z}_i) \boldsymbol{z}_i \boldsymbol{z}_i^\top / N$  exists in a neighborhood around  $\boldsymbol{\lambda}_0$ .

Following Gneiting and Raftery (2007), we consider a generalized entropy that is strictly convex. Condition [A1] implies that the solution to the optimization problem (3.3) is unique when it exists. It also indicates that  $f(\cdot) = g^{-1}(\cdot)$  is differentiable (Deville and Särndal, 1992). Condition [A2] controls the behavior of the first-order inclusion probability (Fuller, 2009, Chapter 2.2). This condition avoids extremely large weights which may cause instability in estimation and prevents the random sample A from being concentrated on a few units in the population. Condition [A3] ensures that the mutual dependence of the two sampling units is not too strong, which is satisfied under many classical survey designs (Robinson and Särndal, 1983; Breidt and Opsomer, 2000). We also assume that the population has a finite average fourth moment, and the covariates  $\mathbf{z}$ 's are asymptotically of full rank in Condition [A4]. Note from (3.6) that  $f'(\mathbf{\lambda}_0^\top \mathbf{z}_i) = 1/g'(d_i^*)$  and  $\Gamma(\mathbf{\lambda}_0) = \lim_{N\to\infty} \sum_{i\in U} \{g'(d_i^*)\}^{-1} \mathbf{z}_i \mathbf{z}_i^\top/N$ , which is finite from Conditions [A1] and [A2] and the existence of  $\mathbf{\Sigma}_{\mathbf{z}}$ . Condition [A4] further assumes that  $\mathbf{\Gamma}(\mathbf{\lambda})$  is finite in a neighborhood of  $\mathbf{\lambda}_0$ . The following theorem presents the main asymptotic properties of the proposed entropy calibration estimator  $\hat{\theta}_{cal} = \sum_{i \in A} \hat{\omega}_i y_i$  where  $\hat{\omega}_i$  is the solution to (3.11).

**Theorem 1** (Design consistency). Suppose Conditions [A1]–[A4] hold. Then, the solution  $\hat{\omega}$  to (3.11) exists and is unique with probability approaching to 1. Furthermore, the proposed entropy calibration estimator  $\hat{\theta}_{cal} = \sum_{i \in A} \hat{\omega}_i y_i$  satisfies

$$\hat{\theta}_{\text{cal}} = \hat{\theta}_{\text{greg},\ell} + o_p(n_0^{-1/2}N), \qquad (4.1)$$

where

$$\hat{ heta}_{ ext{greg},\ell} = \sum_{i \in U} oldsymbol{z}_i^ op oldsymbol{\gamma}_N + \sum_{i \in A} rac{1}{\pi_i} (y_i - oldsymbol{z}_i^ op oldsymbol{\gamma}_N)$$

and

$$\boldsymbol{\gamma}_N = \left\{ \sum_{i \in U} \frac{\pi_i \boldsymbol{z}_i \boldsymbol{z}_i^\top}{g'(d_i^*)} \right\}^{-1} \sum_{i \in U} \frac{\pi_i \boldsymbol{z}_i y_i}{g'(d_i^*)}.$$
(4.2)

Since  $\hat{\theta}_{\text{greg},\ell}$  is design unbiased for  $\theta_N$ , the design consistency of the entropy calibration estimator is also established from Theorem 1. Note that  $g'(d_i^*) > 0$  under Condition [A1].

For the empirical likelihood example in (3.9), the asymptotic equivalence of  $\hat{\theta}_{cal}$  and  $\hat{\theta}_{greg,\ell}$ in (4.1) holds with  $\boldsymbol{\gamma}_N = \left(\sum_{i \in U} d_i \boldsymbol{z}_i \boldsymbol{z}_i^{\top}\right)^{-1} \left(\sum_{i \in U} d_i \boldsymbol{z}_i y_i\right)$ . On the contrary, if the pseudoempirical likelihood method (Chen and Sitter, 1999; Wu and Rao, 2006) is used, then we can obtain

$$\hat{\theta}_{\text{pel}} = \sum_{i \in U} \boldsymbol{x}_i^\top \boldsymbol{\beta}_N + \sum_{i \in A} \frac{1}{\pi_i} (y_i - \boldsymbol{x}_i^\top \boldsymbol{\beta}_N) + o_p(n_0^{-1/2} N), \qquad (4.3)$$

where  $\boldsymbol{\beta}_N = \left(\sum_{i \in U} \boldsymbol{x}_i \boldsymbol{x}_i^{\top}\right)^{-1} \left(\sum_{i \in U} \boldsymbol{x}_i y_i\right)$  is the probability limit of  $\hat{\boldsymbol{\beta}}$  used in the generalized regression estimator  $\hat{\theta}_{\text{greg}}$  in (2.4). Comparing (4.1) with (4.3), we can see that the proposed calibration estimator could be more efficient than the pseudo empirical likelihood estimator as the auxiliary variables  $\boldsymbol{z}_i = \left(\boldsymbol{x}_i^{\top}, g(d_i^*)\right)^{\top}$  use an augmented regression model with an additional covariate  $g(d_i^*)$  and the regression coefficients are estimated more efficiently. The efficiency gain with the additional covariate will be significant if the sampling design is informative in the sense of Pfeffermann and Sverchkov (2009), where the additional covariate  $g(d_i^*)$  may improve the prediction of  $y_i$  as the design weight is correlated with y after controlling on  $\boldsymbol{x}$ .

In order to construct a variance estimator and develop asymptotic normality, we need the following additional conditions.

[B1] The limit of the design covariance matrix of the HT estimator

$$oldsymbol{\Sigma} \coloneqq \lim_{N o \infty} rac{n_0}{N^2} \sum_{i,j \in U} rac{\pi_{ij} - \pi_i \pi_j}{\pi_i \pi_j} egin{pmatrix} y_i y_j & y_i oldsymbol{z}_j^{ op} \ oldsymbol{z}_i y_j & oldsymbol{z}_i oldsymbol{z}_j^{ op} \end{pmatrix}$$

exists and is positive-definite.

[B2] The HT estimator is asymptotically normal in the sense that

$$\left[\mathbb{V}\left\{\sum_{i\in U}\left(1-\frac{\delta_i}{\pi_i}\right)\boldsymbol{x}_i\right\}\right]^{-1/2}\sum_{i\in U}\left(1-\frac{\delta_i}{\pi_i}\right)\boldsymbol{x}_i\stackrel{d}{\to} N(0,\boldsymbol{I})$$

under the sampling design, if  $\limsup_{N \to \infty} N^{-1} \sum_{i \in U} \|\boldsymbol{x}_i\|^4 < \infty$  and

$$\frac{n_0}{N^2} \mathbb{V}\bigg\{\sum_{i \in U} \bigg(1 - \frac{\delta_i}{\pi_i}\bigg) \boldsymbol{x}_i\bigg\} = \frac{n_0}{N^2} \sum_{i,j \in U} \frac{\pi_{ij} - \pi_i \pi_j}{\pi_i \pi_j} \boldsymbol{x}_i \boldsymbol{x}_j^{\mathsf{T}}$$

is positive definite as  $N \to \infty$ , where  $\stackrel{d}{\to}$  stands for convergence in distribution.

Conditions [B1] and [B2] are standard conditions for survey sampling, which hold in many classical survey designs, including simple random sampling and stratified sampling (Fuller, 2009, Chapter 1). Under such conditions, the asymptotic normality of the entropy calibration estimator  $\hat{\theta}_{cal}$  can be established. Theorem 2 (Asymptotic normality). Under Conditions [A1]-[A4], [B1] and [B2], we have

$$\mathbb{V}^{-1/2}(\hat{\theta}_{\mathrm{cal}})(\hat{\theta}_{\mathrm{cal}}-\theta_{\mathrm{N}}) \xrightarrow{d} N(0,1)$$

where  $\mathbb{V}(\hat{\theta}_{cal}) = \mathbb{V}(\hat{\theta}_{greg,\ell})\{1 + o(1)\},\$ 

$$\mathbb{V}(\hat{\theta}_{\text{greg},\ell}) = \sum_{i,j \in U} (\pi_{ij} - \pi_i \pi_j) \frac{y_i - \boldsymbol{z}_i^\top \boldsymbol{\gamma}_N}{\pi_i} \frac{y_j - \boldsymbol{z}_j^\top \boldsymbol{\gamma}_N}{\pi_j}$$

and  $\gamma_N$  is defined in (4.2).

Theorem 2 implies that the design variance of  $\hat{\theta}_{cal}$  depends on the prediction error of the regression of y on z. It suggests that the proposed estimator will perform better if the debiased calibration covariate  $g(d_i^*)$  contains additional information on predicting y. By Theorem 2, the variance of  $\hat{\theta}_{cal}$  can be estimated by

$$\hat{\mathbb{V}}(\hat{\theta}_{cal}) = \sum_{i,j \in A} \frac{\pi_{ij} - \pi_i \pi_j}{\pi_{ij}} \frac{y_i - \boldsymbol{z}_i^\top \hat{\boldsymbol{\gamma}}}{\pi_i} \frac{y_j - \boldsymbol{z}_j^\top \hat{\boldsymbol{\gamma}}}{\pi_j}, \qquad (4.4)$$

where  $\hat{\boldsymbol{\gamma}} = \left\{ \sum_{i \in A} \boldsymbol{z}_i \boldsymbol{z}_i^\top / g'(d_i^*) \right\}^{-1} \left\{ \sum_{i \in A} \boldsymbol{z}_i y_i / g'(d_i^*) \right\}$ . It is shown in the SM that the ratio  $\hat{\mathbb{V}}(\hat{\theta}_{cal}) / \mathbb{V}(\hat{\theta}_{cal}) \xrightarrow{p} 1$  as  $n_0, N \to \infty$  under the conditions of Theorem 2 and a technical condition [A5] in the SM that regulates the dependence in high-order inclusion probabilities.

The asymptotic results in Theorems 1 and 2 are valid for any choice of the entropy function  $G(\cdot)$  in the debiased calibration estimation. In the next section, we discuss the optimal choice of  $G(\cdot)$  in the Poisson sampling design such that the asymptotic variance  $\mathbb{V}(\hat{\theta}_{cal})$  of  $\hat{\theta}_{cal}$  is minimized, where  $\{\delta_i\}_{i\in U}$  are independent.

# 5 Optimal entropy under Poisson sampling

According to Theorem 1, the entropy function  $G(\cdot)$  influences the proposed estimator  $\hat{\theta}_{cal}$ through  $g'(\cdot)$  in the weights of the regression coefficient  $\gamma_N$  in (4.2). In the following, we consider the optimal choice of the entropy function with the unscaled weights in (3.3). The same result applies to the procedure with the scaled weights. We consider a class of generalized regression estimators  $\mathcal{B} = \{\sum_{i \in U} \boldsymbol{z}_i^{\mathsf{T}} \boldsymbol{\xi} + \sum_{i \in A} \pi_i^{-1} (y_i - \boldsymbol{z}_i^{\mathsf{T}} \boldsymbol{\xi}) : \boldsymbol{\xi} \in \mathbb{R}^{p+1}\}$  for the population total  $\theta_N$  with the augmented covariates  $\boldsymbol{z}_i = (\boldsymbol{x}_i^{\mathsf{T}}, g(d_i))^{\mathsf{T}}$ , where p + 1 is the dimension of  $\boldsymbol{z}_i$ . Let

$$\boldsymbol{\gamma}_{\text{opt}} = \left\{ \mathbb{V}\left(\sum_{i \in A} d_i \boldsymbol{z}_i\right) \right\}^{-1} \mathbb{C}\left(\sum_{i \in A} d_i \boldsymbol{z}_i, \sum_{i \in A} d_i y_i\right),$$

where  $\mathbb{C}$  denotes the covariance operator. Then,

$$\hat{\theta}_{\text{greg,opt}} = \sum_{i \in U} \boldsymbol{z}_i^\top \boldsymbol{\gamma}_{\text{opt}} + \sum_{i \in A} \frac{1}{\pi_i} (y_i - \boldsymbol{z}_i^\top \boldsymbol{\gamma}_{\text{opt}})$$
(5.1)

is the design-optimal regression esitmator of  $\theta_{\rm N}$  with the smallest variance in the class  $\mathcal{B}$ . The optimal estimator in (5.1), also called as pseudo optimal estimator in Montanari (1987); Rao (1994); Berger et al. (2003), minimizes design variance among the class of design-unbiased estimators that are linear in  $\sum_{i \in A} d_i y_i$  and  $\sum_{i \in A} d_i z_i$ . It can also be interpreted as the projection of the HT estimator onto the orthogonal complement of the augmentation space generated by  $\sum_{i \in A} d_i z_i$  (Tsiatis, 2006).

Under Poisson sampling,  $\{\delta_i\}_{i\in U}$  are mutually independent, such that  $\pi_{ij} = \pi_i \pi_j$  for all  $i \neq j$ . In this case, we have  $\mathbb{V}(\hat{\theta}_{\text{greg},\ell}) = \sum_{i\in U} (1-\pi_i)\pi_i^{-1}(y_i - \boldsymbol{z}_i^\top \boldsymbol{\gamma}_N)(y_j - \boldsymbol{z}_j^\top \boldsymbol{\gamma}_N)$  and  $\boldsymbol{\gamma}_{\text{opt}} = (\sum_{i\in U} \pi_i q_i \boldsymbol{z}_i \boldsymbol{z}_i^\top)^{-1} \sum_{i\in U} \pi_i q_i \boldsymbol{z}_i y_i$ , where  $q_i = \pi_i^{-2} - \pi_i^{-1}$ . Now we wish to find the optimal choice of the generalized entropy function  $G(\cdot)$  such that the resulting debiased calibration estimator  $\hat{\theta}_{\text{cal}}$  is asymptotically equivalent to the design-optimal regression estimator  $\hat{\theta}_{\text{greg,opt}}$  under Poisson sampling.

To achieve this goal, using the asymptotic equivalence in (4.1), we only need to find a

special entropy function  $G(\cdot)$  such that

$$\frac{1}{g'(d_i)} = d_i^2 - d_i \tag{5.2}$$

where  $g(\omega) = \partial G(\omega)/\partial \omega$  is the first-order derivative of  $G(\omega)$ . The differential equation in (5.2) is satisfied with  $g(\omega) = \log(\omega - 1) - \log(\omega)$ . Therefore, the optimal entropy function is

$$G(\omega) = (\omega - 1)\log(\omega - 1) - \omega\log(\omega)$$
(5.3)

for  $\omega > 1$ , which is called the cross entropy between  $(\omega - 1) \log(\omega - 1)$  and  $\omega \log(\omega)$ . Note that the EL and exponential tilting (ET) approaches (Kim, 2010; Hainmueller, 2012) correspond to  $G(\omega)$  being  $-\log(\omega)$  and  $\omega \log(\omega)$ , respectively, and choosing  $G(\omega) = (\omega - 1) \log(\omega - 1)$ implies a logistic regression model for the inclusion probability  $\pi_i$ . In this view, the optimal entropy function in (5.3) can be regarded as a contrast between the logistic model and the exponential tilting model for the propensity scores.

Note that  $G(\omega)$  in (5.3) is strictly convex with a negative first derivative  $g(\omega) = \partial G(\omega)/\partial \omega = \log(\omega - 1) - \log(\omega) = \log(1 - \omega^{-1})$  and a positive second derivative  $g'(\omega) = \partial^2 G(\omega)/\partial^2 \omega = \{\omega(\omega - 1)\}^{-1}$  for  $\omega > 1$ . It takes negative values for  $\omega > 1$  with  $\lim_{\omega \downarrow 1} G(\omega) = 0$  and  $\lim_{\omega \to \infty} G(\omega) = -\infty$ . The proposed cross entropy calibration method can be described as the following constrained optimization problem

$$\hat{\boldsymbol{\omega}} = \operatorname{argmin}_{\omega_i > 1} \sum_{i \in A} \left\{ (\omega_i - 1) \log(\omega_i - 1) - \omega_i \log(\omega_i) \right\} \text{ subject to } (2.2) \text{ and } (5.4)$$

$$\sum_{i \in A} \omega_i \log(1 - \pi_i) = \sum_{i \in U} \log(1 - \pi_i),$$
(5.5)

where (5.5) is the debiasing calibration constraint, specifically designed for the cross entropy loss in (5.3), and (2.2) is the benchmarking calibration constraint for covariates. If the covariate  $\mathbf{z}_i$  in the regression estimator includes the debiasing covariate  $g(d_i) = \log(1 - \pi_i)$  for  $i \in U$ , Theorem 1 shows that the proposed entropy calibration estimator  $\hat{\theta}_{cal}$  using the cross entropy is asymptotically equivalent to the design-optimal regression estimator in (5.1). Note that the asymptotic equivalence of  $\hat{\theta}_{cal}$  and a generalized regression estimator does not need the condition of Poisson sampling, but the asymptotic optimality is justified under Poisson sampling. This result is summarized in the following corollary.

**Corollary 1.** Under the conditions of Theorem 1, the solution  $\hat{\omega}$  of the cross entropy weight in (5.4) exists and is unique with probability approaching to 1. Furthermore, if  $\{\delta_i\}_{i \in U}$  are independent, the proposed estimator  $\hat{\theta}_{cal}$  using the cross entropy weight in (5.4) satisfies  $\hat{\theta}_{cal} = \hat{\theta}_{greg,opt} + o_p(n_0^{-1/2}N)$ , where  $\hat{\theta}_{greg,opt}$  is defined in (5.1).

#### 6 Unknown population-level inclusion probabilities

To apply the proposed method, the population total  $\sum_{i \in U} g(d_i)$  must be known in the debiasing constraint in (3.2), which is possible if  $\{\pi_i\}$  are known throughout the finite population. If  $\{\pi_i\}$  is not available outside the sample, we cannot directly impose the constraint in (3.2). In this section, we consider the situation where  $\sum_{i \in U} g(d_i)$  is unknown and modify the proposed method to handle this situation.

In one approach, we can estimate  $\alpha_N = N^{-1} \sum_{i \in U} g(d_i)$  by  $\hat{\alpha} = N^{-1} \sum_{i=1}^N \hat{m}(\boldsymbol{x}_i)$ , where  $\hat{m}(\boldsymbol{x})$  is a nonparametric regression estimator of  $m(\boldsymbol{x}) = \mathbb{E}\{g(d) \mid \boldsymbol{x}\}$ . For example,

$$\hat{m}(\boldsymbol{x}) = \underset{m}{\operatorname{argmin}} \sum_{i \in A} d_i \left\{ g(d_i) - m \right\}^2 K_h(\boldsymbol{x}_i, \boldsymbol{x}),$$

by the kernel-smoothing regression, where  $K_h(\cdot, \cdot)$  is a kernel function with a bandwidth h. The entropy calibration estimator  $\hat{\theta}_{cal}$  can be constructed similarly to (3.4), but replacing the constraint in (3.2) by  $\sum_{i \in A} \omega_i g(d_i) = \sum_{i \in U} \hat{m}(\boldsymbol{x}_i)$ . However, such a nonparametric regression approach assumes that individual values of  $\boldsymbol{x}_i$  are available throughout the finite population, which may not hold in some practical situations. Furthermore, the performance of nonparametric regression might be poor when the sampling weights are highly variable.

An alternative approach is to modify the constrained optimization problem in (3.3) by treating  $\alpha_N$  as an additional unknown parameter, which can be expressed as follows

$$\left\{\hat{\boldsymbol{\omega}}_{\mathrm{m}},\hat{\boldsymbol{\alpha}}\right\} = \underset{\boldsymbol{\alpha}\in\mathcal{A}_{n},\boldsymbol{\omega}}{\arg\min}\sum_{i\in A}G(\omega_{i}) - N\hat{K}(\boldsymbol{\alpha}) \text{ subject to } (2.2) \text{ and } \sum_{i\in A}\omega_{i}g(d_{i}) = N\boldsymbol{\alpha}, \quad (6.1)$$

where  $\hat{\boldsymbol{\omega}}_{m} = {\{\hat{\omega}_{m,i} : \delta_i = 1\}}$ ,  $\mathcal{A}_n$  is a bounded open interval and  $\hat{K}(\alpha)$  may depend on the sample A. The proposed estimator for the population total is given by  $\hat{\theta}_{cal,m} = \sum_{i \in A} \hat{\omega}_{m,i} y_i$ . Once  $\alpha$  is given, the objective function in (6.1) is convex with respect to  $\boldsymbol{\omega}$ . Thus, the solution of  $\hat{\alpha}$  can be obtained by profiling  $\boldsymbol{\omega}$  out. See the detailed derivation in the SM. Under some regularity conditions, we can establish that

$$N^{-1}\hat{\theta}_{\text{cal,m}} = N^{-1}\hat{\theta}_{\text{greg},\ell} + (\hat{\alpha} - \alpha_N)\gamma_{N,p+1} + o_p(n_0^{-1/2}), \tag{6.2}$$

where  $\hat{\theta}_{\text{greg},\ell}$  is defined in (4.1) and  $\gamma_{N,p+1}$  is the last component in  $\gamma_N$  and  $\boldsymbol{z}_i^{\top} = (\boldsymbol{x}_i^{\top}, g(d_i))$ . The linearization of (6.2) has the same structure of (4.1) except for the additional term  $(\hat{\alpha} - \alpha_N)\gamma_{N,p+1}$  due to the estimation of  $\alpha_N$ . This term reflects the uncertainty in estimating the unknown  $\alpha_N$ . Note that  $\gamma_{N,p+1} = 0$  and  $\hat{\alpha}$  would not affect the linearization of  $\hat{\theta}_{\text{cal,m}}$  if the working regression model is correct and the sampling mechanism is non-informative. Furthermore, the linearization in (6.2) can be equivalently expressed as

$$N^{-1}\hat{\theta}_{\text{cal},\text{m}} = \frac{1}{N} \sum_{i \in U} (\boldsymbol{x}_i^{\top}, m_i) \boldsymbol{\gamma}_N + \frac{1}{N} \sum_{i \in A} d_i \{ y_i - (\boldsymbol{x}_i^{\top}, m_i) \boldsymbol{\gamma}_N \} + o_p(n_0^{-1/2}), \quad (6.3)$$

for some  $m_i$  that depends on the choice of  $\hat{K}(\cdot)$ . This linearization equation in (6.3) can be used for the estimation of the variance of  $\hat{\theta}_{cal,m}$ . In the SM, a sketched proof of (6.2), (6.3), the definition of  $m_i$ , and its variance estimation formula are presented.

**Remark 1.** One possible choice of  $\hat{K}(\alpha)$  is  $\hat{K}(\alpha) = \alpha$ . Under this choice of  $\hat{K}(\alpha)$ , we can profile out  $\alpha$  in (6.1), and the optimization problem reduces to

$$\hat{\boldsymbol{\omega}}_{\mathrm{m}} = \underset{\boldsymbol{\omega}}{\operatorname{arg\,min}} \sum_{i \in A} \{ G(\omega_i) - \omega_i g(d_i) \} \quad subject \ to \quad (2.2), \tag{6.4}$$

where  $\hat{\alpha} = N^{-1} \sum_{i \in A} \hat{\omega}_{m,i} g(d_i)$ , and the objective function in (6.4) is strictly convex with respect to  $\boldsymbol{\omega}$ . Using the Lagrangian multiplier method, the solution  $\hat{\boldsymbol{\omega}}_m$  to (6.4) satisfies  $\hat{\omega}_{m,i} = f(\hat{\boldsymbol{\lambda}}_1^T \boldsymbol{x}_i + g(d_i))$  for  $i \in A$ , which is similar to the solution  $\hat{\omega}_i$  of (3.3) in (3.6), but  $\hat{\lambda}_2$ is set to be 1. In this case, the entropy calibration estimator is asymptotically equivalent to

$$N^{-1}\hat{\theta}_{\mathrm{cal,m}} = \frac{1}{N}\sum_{i\in U} \boldsymbol{x}_i^{\top}\hat{\boldsymbol{\beta}}_{\mathrm{g}} + \frac{1}{N}\sum_{i\in A}\frac{1}{\pi_i}(y_i - \boldsymbol{x}_i^{\top}\hat{\boldsymbol{\beta}}_{\mathrm{g}}) + o_p(n_0^{-1/2})$$

where  $\hat{\boldsymbol{\beta}}_{g} = \left\{\sum_{i \in A} \boldsymbol{x}_{i} \boldsymbol{x}_{i}^{\top} / g'(d_{i})\right\}^{-1} \sum_{i \in A} \boldsymbol{x}_{i} y_{i} / g'(d_{i})$ . Thus, the effect of augmenting the covariates by adding  $g(d_{i})$  disappears for the choice of  $K(\alpha) = \alpha$ . However, this approach allows for different weights  $\{g'(d_{i})\}^{-1}$  in the regression coefficient  $\hat{\boldsymbol{\beta}}_{g}$  using different entropies compared to  $\hat{\boldsymbol{\beta}}$  in (2.5) for  $\hat{\theta}_{DS}$ .

**Remark 2.** Qin et al. (2002) proposed a method of estimating  $\alpha_N$  in the empirical likelihood framework. It turns out that their profile empirical likelihood method is a special case of our proposed method in (6.1) for  $G(\omega) = -\log(\omega)$  and  $\hat{K}(\alpha) = (\hat{\alpha}_{HT} + 1)\log(\alpha + 1)$  for  $\alpha \in (-1,0)$ , where  $\hat{\alpha}_{HT} = N^{-1} \sum_{i \in A} d_i g(d_i)$  which equals to -n/N under this case. The asymptotic properties of the profile EL estimator can also be found in Liu and Fan (2023). More generally, the entropy calibration estimator  $\hat{\theta}_{cal,m}$  with  $\hat{K}(\alpha) = (\hat{\alpha}_{HT} + 1) \log(\alpha + 1)$  in (6.1) can be linearized as (6.3) where

$$m_i = \frac{\alpha_N + 1}{\sum_{gg \cdot \boldsymbol{x}} + \alpha_N + 1} \boldsymbol{x}_i^{\top} \left( \sum_{i \in U} \frac{1}{d_i g'(d_i)} \boldsymbol{x}_i \boldsymbol{x}_i^{\top} \right)^{-1} \sum_{i \in U} \frac{g(d_i)}{d_i g'(d_i)} \boldsymbol{x}_i,$$

and  $\Sigma_{gg\cdot x}$  is defined in (A.14) in the SM. Observing that  $(\alpha_N + 1)/(\Sigma_{gg\cdot x} + \alpha_N + 1) \in (0, 1)$ ,  $m_i$  is the shrinkage linear projection of  $g(d_i)$  on the column space spanned by  $x_i$  for  $i \in U$ .

From the above analysis, we can make the following conclusions about the comparison between the proposed estimator  $\hat{\theta}_{cal,m}$  and  $\hat{\theta}_{DS}$  under unknown  $\alpha_N$ . First, if the working regression model implied by the calibration constraint is correct and the sampling mechanism is non-informative,  $\hat{\theta}_{cal,m}$  and  $\hat{\theta}_{DS}$  are asymptotically equivalent. Second, by choosing the cross entropy and  $\hat{K}(\alpha) = \alpha$ ,  $\hat{\theta}_{cal,m}$  is more efficient than  $\hat{\theta}_{DS}$  under misspecified regression model or informative sampling. This conclusion follows by combining the results in Remark 1 and Section 5. Third, from (6.3), for general choices of  $\hat{K}(\alpha)$  and  $G(\omega)$ , the linearization of  $\hat{\theta}_{cal,m}$ is not equivalent to the generalized regression estimator using the augmented covariates  $(\boldsymbol{x}_i, m_i)$ . There is no universal result on the efficiency gain of  $\hat{\theta}_{cal,m}$  over  $\hat{\theta}_{DS}$ .

# 7 Simulation study

To test our theory, we performed a limited simulation study. We consider a finite population of size N = 10,000. A vector of two auxiliary variables  $\boldsymbol{x}_i = (x_{1i}, x_{2i})^{\top}$  is available for  $i = 1, \dots, N$ , where  $x_{1i} \sim N(2, 1)$  follows the normal distribution with mean 2 and standard deviation 1, and  $x_{2i} \sim \text{Unif}(0, 4)$ , uniform distribution in [0, 4]. We consider two superpopulation models to generate the study variable y:  $y_i = x_{1i} + x_{2i} + e_i$  (Model 1) and  $y_i = x_{1i}/3 + x_{2i}/3 + x_{1i}x_{2i}^2/4 + e_i$  (Model 2), where  $e_i \sim N(0, 1)$ , independent of  $\boldsymbol{x}_i$ . From each of the finite populations, samples are selected using Poisson sampling with inclusion probability  $\pi_i = \min(\Phi_3(-x_{1i}/2 - x_{2i}/2 - 2), 0.7)$ , where  $\Phi_3(\cdot)$  is the cumulative distribution function of the *t* distribution with degree of freedom 3. The distribution of the design weights is right-skewed, resulting in some design weights being extremely large, as illustrated in Figure 1. The expected sample size is  $\mathbb{E}(n) \approx 1100$ . The model  $R^2$  and the partial correlation between  $\pi_i$  and  $y_i$  after regressing out  $\boldsymbol{x}_i$  were  $R^2 = 0.6973$  and pCor $(\pi, y \mid \boldsymbol{x}) = -0.0040$ , respectively, in Model 1. In Model 2,  $R^2 = 0.7893$  and pCor $(\pi, y \mid \boldsymbol{x}) = 0.6183$ . For a fixed realization of the population, samples are generated repeatedly 1,000 times. We are interested in estimating the population mean  $\mu_y = N^{-1} \sum_{i \in U} y_i$  from the sampled data. From each sample, we compare the following estimators.



Figure 1: Histogram and boxplot of the design weights. The red solid line is the mean of the design weights and the blue dashed line is the median of the design weights.

**Hájek** Hájek estimator:  $\hat{\mu}_{y,HT} = \left(\sum_{i \in A} d_i\right)^{-1} \left(\sum_{i \in A} d_i y_i\right).$ 

- **DS** Deville and Särndal (1992)'s divergence calibration estimator:  $\hat{\mu}_{y,DS} = N^{-1} \sum_{i \in A} \hat{w}_i y_i$ , where the calibration weight  $\hat{\boldsymbol{w}} = \{\hat{w}_i\}_{i \in A}$  minimizes the divergence measure  $D(\boldsymbol{w}, \boldsymbol{d}) = \sum_{i \in A} d_i G(\omega_i/d_i)$  defined in (2.3) subject to  $\sum_{i \in A} w_i = N$  and  $\sum_{i \in A} w_i \boldsymbol{x}_i = \sum_{i \in U} \boldsymbol{x}_i$ .
- **GEC** The proposed generalized entropy calibration estimator:  $\hat{\mu}_{y,cal} = N^{-1} \sum_{i \in A} \hat{w}_i y_i$ , where the calibration weight  $\hat{w} = \{\hat{w}_i\}_{i \in A}$  maximizes the entropy  $H(\omega) = -\sum_{i \in A} G(w_i)$ subject to  $\sum_{i \in A} w_i = N$ ,  $\sum_{i \in A} w_i x_i = \sum_{i \in U} x_i$  and  $\sum_{i \in A} w_i g(d_i) = \sum_{i \in U} g(d_i)$ (**GEC0**). If  $\alpha_N$  is unknown,  $\hat{w} = \{\hat{w}_i\}_{i \in A}$  and  $\hat{\alpha}$  maximize the adjusted entropy  $H(\omega, \alpha) = -\sum_{i \in A} G(w_i) + N\hat{K}(\alpha)$  subject to  $\sum_{i \in A} w_i = N$ ,  $\sum_{i \in A} w_i x_i = \sum_{i \in U} x_i$ , and  $\sum_{i \in A} w_i g(d_i) = N\alpha$ . We consider two candidates for  $\hat{K}(\alpha)$ :  $\hat{K}_1(\alpha) = \alpha$  (**GEC1**) and  $\hat{K}_2(\alpha) = (\hat{\alpha}_{HT} + 1) \log(\alpha + 1)$  (**GEC2**).

For each of **DS**, **GEC0**, **GEC1** and **GEC2** estimators, we consider the following entropy (divergence) functions  $G(\omega)$ : empirical likelihood (**EL**)  $G(\omega) = -\log \omega$ , exponential tilting (**ET**)  $G(\omega) = \omega \log \omega - \omega$ , cross entropy (**CE**)  $G(\omega) = (\omega - 1) \log (\omega - 1) - \omega \log \omega$ , and Hellinger distance (**HD**)  $G(\omega) = -4\sqrt{\omega}$ .

Table 2 presents the biases, standard errors (SE), and root mean squared errors (RMSE) of the estimators, along with the coverage rates (CR%) of their 95% confidence intervals, calculated from 1,000 Monte Carlo samples. When the sampling design is non-informative and the underlying working model of the calibration covariates is correct (Model 1), the **DS** estimator performs slightly better than the **GEC** estimators in terms of RMSE, as the additional debiasing covariate is unnecessary for predicting y when the true superpopulation model  $y_i$  is linear in the calibration covariates  $x_{1i}$  and  $x_{2i}$ . On the other hand, when the

		Model 1			Model 2				
		Bias	SE	RMSE	$\operatorname{CR}(\%)$	Bias	SE	RMSE	CR(%)
Háj	jek	0.51	7.96	7.97	96	-0.32	19.87	19.87	95
EL	DS	0.10	3.91	3.91	96	-0.23	8.28	8.29	93
	GEC0	0.13	3.92	3.92	96	-0.05	5.31	5.31	95
	GEC1	0.13	3.93	3.93	96	-0.02	7.09	7.09	94
	GEC2	0.13	3.93	3.93	96	-0.02	7.06	7.06	94
DS ET GEC0 GEC1 GEC2	0.10	3.91	3.91	96	-0.32	8.28	8.29	93	
	GEC0	0.10	3.91	3.91	96	0.20	5.15	5.16	95
	GEC1	0.10	3.91	3.91	96	-0.31	8.28	8.29	93
	GEC2	0.10	3.91	3.91	96	-0.31	8.28	8.28	93
	DS	0.10	3.91	3.91	96	-0.40	8.28	8.29	94
CE	GEC0	0.13	3.92	3.92	96	-0.08	5.42	5.43	95
CE	GEC1	0.13	3.93	3.93	96	0.01	7.06	7.06	95
	GEC2	0.13	3.93	3.93	96	0.01	7.03	7.03	95
	DS	0.10	3.91	3.91	96	-0.28	8.28	8.29	93
HD G G G	GEC0	0.11	3.91	3.91	96	-0.04	5.16	5.16	95
	GEC1	0.11	3.91	3.92	96	-0.21	7.45	7.45	93
	GEC2	0.11	3.91	3.92	96	-0.22	7.42	7.42	93

Table 2: Bias ( $\times 100$ ), standard error (SE,  $\times 100$ ), and root mean squared error (RMSE,  $\times 100$ ) of the estimators, and coverage rate (CR%) of their 95% confidence intervals under Model 1 (correct model) and Model 2 (incorrect model).

underlying working model for calibration is incorrect, we observe different results. Under Model 2, the **GEC** estimators demonstrate greater efficiency compared to the DS estimators. Unlike in the case of a linear superpopulation model, the debiasing constraint in **GEC** is helpful in reducing the residual variance, as the augmented regression model implicit in the proposed calibration method can have a better prediction power than the reduced model without  $g(d_i)$  in the covariates. Remarks 1 and 2 in Section 6 justify the **GEC1** and **GEC2** estimators when  $\alpha_N$  is not available. Meanwhile, since these two estimators are derived based solely on the population total of auxiliary variables,  $\sum_{i \in U} \boldsymbol{x}_i$ , their efficiency gain is less pronounced than that of the **GEC0** estimator. The simulation results in Table 2 indicate that the proposed **GEC2** estimator consistently outperforms the **DS** estimator in terms of RMSE under Model 2. The RMSE of the **GEC1** estimator falls between that of the **DS** estimator and the **GEC2** estimator. For the **ET** divergence, the performance of **DS** and **GEC1** is essentially identical, which can be attributed to their asymptotic equivalence.

Throughout Table 2, the bias of all estimators is negligible, with the contribution of the squared bias to the mean squared error being less than 1%. Generally, coverage rates are close to the nominal 95% level, within the bounds of experimental error. Under Model 2, coverage rates of the confidence intervals are slightly lower than the nominal coverage rates. This phenomenon aligns with the discussion by Rao et al. (2003) on the undercoverage property of the regression estimator under model misspecification.

#### 8 Real data analysis

We present an application of the proposed method using data from a proprietary pesticide usage survey collected from GfK Kynetec in 2020. Kynetec is a private company that specializes in agricultural information systems that involve data collection from farmers in the United States. One of the main objectives of this survey is to estimate the total amount (\$) spent by farm operations on pesticides in each state of the United States.

The survey was carried out by stratified sampling; the population was stratified by three

factors: 50 states, 60 crops, and the size of a farm (integers from 1 to 7). Since larger farms tend to use greater amounts of pesticides, the sampling design assigned a greater proportion of the sample to larger farms within each stratum to reduce variance. See Thelin and Stone (2013) for further details on the survey design.

For each farm *i*, the study variable  $y_i$  is the dollar amount spent on the pesticide, including the herbicide, insecticide, and fungicide produced by the five largest agrochemical companies: BASF, Bayer, Corteva Agriscience, FMC, and Syngenta. The auxiliary variables  $\boldsymbol{x}_i$  are the harvested areas(in acres) for each crop in each multi-county area, referred to as the Crop Reporting District(CRD). Total acres harvested for each crop-by-CRD combination are available from the USDA Census of Agriculture. Throughout the United States, there were more than 20,000 samples with more than 1,000 strata. We only report the results for four states for brevity.

For estimation, we compared the Horvitz-Thompson estimator (**HT**), generalized regression estimator (**Reg**) in (2.4), pseudo-empirical estimator (**PEL**) in (3.10), and the proposed generalized entropy calibration estimator using empirical likelihood (**EL**), crossentropy (**CE**), and Hellinger distance (**HD**) with  $\hat{K}(\alpha) = (\hat{\alpha}_{\text{HT}} + 1) \log(\alpha + 1)$ , denoted as **GEC2** in the previous section. Since the design weights and the auxiliary variables are not available in each population unit, the **GEC** method is not applicable to our data.

Table 3 summarizes the point estimates, standard errors, and 95 % confidence intervals of the estimators. All the calibration methods converge well and produce weights even when the number of auxiliary variables is greater than 30 as in Missouri. Incorporating auxiliary variables as in **Reg**, **PEL**, **EL**, **CE**, or **HD** dramatically improved performance compared to the Horvitz-Thompson estimator **HT**. The standard error of the proposed entropy calibration estimators using **EL** or **CE** was the smallest for all states reported. Although the entropy calibration estimators produce similar point estimates and standard errors as **Reg** or **PEL** estimators in Iowa and Florida, the difference between the estimators was significant in Missouri and Mississippi.

# 9 Concluding remarks

In this paper, we have proposed a novel approach that integrates generalized entropy with a debiasing constraint for calibration estimation. The proposed calibration method implicitly uses an augmented regression model with  $g(d_i)$  as an additional covariate, which not only achieves design consistency but also improves efficiency. The efficiency gain is significant when the working regression model is incorrect or when the sampling design is informative under the working regression model for calibration. Since we never know the true model in practice, the additional covariate proves useful, and the resulting calibration estimator is often more efficient than the traditional calibration method of Deville and Särndal (1992).

The proposed calibration weights can be applied for multiple outcome variables of interest, leading to a design-consistent estimator regardless of the outcome variable. The efficiency of this estimator for different outcome variables may vary, depending on the predictability of  $g(d_i)$  on  $y_i$ . Generally, if the regression coefficient for  $g(d_i)$  in the augmented regression is significant, then the proposed calibration estimator will be more efficient than the traditional method. However, it may increase the variance when the regression coefficient for  $g(d_i)$  is insignificant. A cautious user could perform a test before adopting the proposed method (Fuller, 1984). Nevertheless, the increase in variance using the proposed method when the

(N, n, p)	IA: (124947, 1197, 30)			MO: (85384, 677, 38)			
Method	Est	SE	CI	Est	SE	CI	
НТ	667.03	18.68	(630.42, 703.63)	353.55	18.55	(317.19, 389.90)	
Reg	660.27	10.59	(639.51, 681.03)	327.77	10.71	(306.79, 348.75)	
PEL	660.91	10.59	(640.15, 681.67)	335.61	10.71	(314.63, 356.60)	
EL	659.54	10.47	(639.02,  680.07)	327.52	10.64	(306.67, 348.36)	
CE	659.51	10.47	(638.98, 680.03)	327.47	10.64	(306.62, 348.31)	
HD	660.93	10.50	(640.35, 681.50)	327.78	10.66	(306.88, 348.67)	
	FL: (14573, 152, 13)			MS: (17072, 160, 19)			
(N, n, p)	F	L: (1457	73, 152, 13)	М	S: (1707	2, 160, 19)	
(N, n, p)Method	F Est	<sup>°</sup> L: (1457 SE	73, 152, 13) CI	M Est	S: (1707 SE	2, 160, 19) CI	
$\frac{(N, n, p)}{\text{Method}}$ HT	F Est 84.42	°L: (1457 SE 14.10	73, 152, 13) CI (56.79, 112.05)	M Est 114.01	S: (1707 SE 9.79	2, 160, 19) CI $(94.81, 133.20)$	
(N, n, p) Method HT Reg	F Est 84.42 55.92	L: (1457 SE 14.10 6.90	CI $(56.79, 112.05)$ $(42.40, 69.44)$	M Est 114.01 119.15	S: (1707 SE 9.79 5.99	$\begin{array}{c} 2, 160, 19) \\ \hline \\ CI \\ \hline \\ (94.81, 133.20) \\ (107.42, 130.89) \end{array}$	
(N, n, p) $Method$ $HT$ $Reg$ $PEL$	F Est 84.42 55.92 53.59	L: (1457 SE 14.10  6.90 6.90	CI $CI$ $(56.79, 112.05)$ $(42.40, 69.44)$ $(40.07, 67.11)$	M Est 114.01 119.15 115.96	S: (1707 SE 9.79 5.99 5.99	$\begin{array}{r} \hline 2, 160, 19) \\ \hline \\ $	
(N, n, p) $Method$ $HT$ $Reg$ $PEL$ $EL$	F Est 84.42 55.92 53.59 54.63	E: (1457 SE 14.10 6.90 6.90 6.80	$\begin{array}{c} \text{CI} \\ \hline \\ \hline \\ (56.79, 112.05) \\ \hline \\ (42.40, 69.44) \\ \hline \\ (40.07, 67.11) \\ \hline \\ (41.31, 67.95) \end{array}$	M Est 114.01 119.15 115.96 112.87	S: (1707 SE 9.79 5.99 5.99 5.99 5.93	$\begin{array}{c} 2, 160, 19) \\ \hline \\ CI \\ \hline \\ (94.81, 133.20) \\ (107.42, 130.89) \\ (104.22, 127.69) \\ (101.26, 124.49) \end{array}$	
(N, n, p) $Method$ $HT$ $Reg$ $PEL$ $EL$ $CE$	F Est 84.42 55.92 53.59 54.63 54.72	L: (1457 SE 14.10 6.90 6.90 6.80 6.80	$\begin{array}{c} \text{CI} \\ \hline \\ (56.79, 112.05) \\ (42.40, 69.44) \\ (40.07, 67.11) \\ (41.31, 67.95) \\ (41.40, 68.05) \end{array}$	M Est 114.01 119.15 115.96 112.87 112.82	S: (1707 SE 9.79 5.99 5.99 5.93 5.93	$\begin{array}{c} 2, 160, 19) \\ \hline \\ CI \\ \hline \\ (94.81, 133.20) \\ (107.42, 130.89) \\ (104.22, 127.69) \\ (101.26, 124.49) \\ (101.21, 124.44) \end{array}$	

Table 3: Point estimate (Est), standard error (SE), and confidence intervals (CI) of pesticide sales (×10<sup>6</sup>\$) in the four states, Iowa (IA), Missouri (MO), Florida (FL), and Mississippi (MS), where (N, n, p) stands for the number of farms, the number of samples, and the dimension of the auxiliary variable  $\boldsymbol{x}_i$  in a state, respectively.

working model for calibration is indeed correct is of a smaller order than the reduction in variance when the working model is incorrect. Thus, asymptotically, the potential efficiency gain outweighs the potential risk.

Although the optimal entropy function using cross-entropy, as discussed in Section 5, is asymptotically optimal under the same set of balancing functions, it may not produce the most efficient calibration estimate when different balancing functions are used, as demonstrated in our simulation study. In practice, one may fit augmented regression for y with different entropy functions and choose the one with the largest  $R^2$ . The selection of the optimal entropy function and  $\hat{K}(\alpha)$  in (6.1) presents an intriguing avenue for future research.

In practice, unit nonresponse often necessitates adjustments to design weights. For such scenarios, the debiasing constraint in the proposed method should incorporate the final weights after nonresponse adjustment. However, these adjusted weights introduce additional sampling error, complicating statistical inference. Also, when  $p = \dim(\mathbf{x})$  is large, we can apply generalized entropy with soft calibration using the  $L_2$  norm (Guggemos and Tillé, 2010) or the  $L_1$  norm (McConville et al., 2017; Wang and Zubizarreta, 2020). Once the debiasing constraint is satisfied, other benchmarking constraints can be relaxed to accommodate highdimensional auxiliary variables.

# Acknowledgement

The authors thank the AE and two anonymous referees for very constructive comments.

# References

- Athey, S., Imbens, G. W., and Wager, S. (2018). Approximate residual balancing: debiased inference of average treatment effects in high dimensions. *Journal of the Royal Statistical Society: Series B*, 80(4):597–623.
- Berger, Y. G., Tirari, M. E., and Tillé, Y. (2003). Towards optimal regression estimation in sample surveys. Australian & New Zealand Journal of Statistics, 45(3):319–329.
- Berger, Y. G. and Torres, O. D. L. R. (2016). Empirical likelihood confidence intervals for complex sampling designs. Journal of the Royal Statistical Society Series B: Statistical Methodology, 78(2):319–341.
- Boyd, S. P. and Vandenberghe, L. (2004). Convex optimization. Cambridge university press.
- Breidt, F. J., Claeskens, G., and Opsomer, J. D. (2005). Model-assisted estimation for complex surveys using penalised splines. *Biometrika*, 92(4):831–846.
- Breidt, F. J. and Opsomer, J. D. (2000). Local polynomial regression estimators in survey sampling. *Annals of Statistics*, 28(4):1026–1053.
- Breidt, F. J. and Opsomer, J. D. (2017). Model-assisted survey estimation with modern prediction techniques. *Statistical Science*, 32(2):190–205.
- Chen, J. and Sitter, R. R. (1999). A pseudo empirical likelihood approach to the effective use of auxiliary information in complex surveys. *Statistica Sinica*, 9(2):385–406.
- Dagdoug, M., Goga, C., and Haziza, D. (2023). Model-assisted estimation through random

forests in finite population sampling. *Journal of the American Statistical Association*, 118:1234–1251.

- Devaud, D. and Tillé, Y. (2019). Deville and Särndal's calibration: revisiting a 25-years-old successful optimization problem (with discussion). *Test*, 28:1033–1065.
- Dever, J. A. and Valliant, R. (2016). General regression estimation adjusted for undercoverage and estimated control totals. *Journal of Survey Statistics and Methodology*, 4:289–318.
- Deville, J.-C. and Särndal, C.-E. (1992). Calibration estimators in survey sampling. *Journal* of the American statistical Association, 87(418):376–382.
- Elliott, M. R. and Valliant, R. L. (2017). Inference for nonprobability samples. *Statistical Science*, 32(2):249–264.
- Firth, D. and Bennett, K. (1998). Robust models in probability sampling. Journal of the Royal Statistical Society: Series B (Statistical Methodology), 60(1):3–21.
- Fuller, W. A. (1984). Least squares and related analysis for complex survey designs. Survey Methodology, 10:97–118.
- Fuller, W. A. (2009). Sampling Statistics. Wiley series in survey methodology. Wiley, Hoboken, N.J.
- Gneiting, T. and Raftery, A. E. (2007). Strictly proper scoring rules, prediction, and estimation. Journal of the American statistical Association, 102(477):359–378.
- Godambe, V. P. and Joshi, V. M. (1965). Admissibility and Bayes estimation in sampling finite populations, 1. Annals of Mathematical Statistics, 36:1707–1722.

- Guggemos, F. and Tillé, Y. (2010). Penalized calibration in survey sampling: Design-based estimation assisted by mixed models. *Journal of statistical planning and inference*, 140:3199– 3212.
- Hainmueller, J. (2012). Entropy balancing for causal effects: A multivariate reweighting method to produce balanced samples in observational studies. *Political analysis*, 20(1):25– 46.
- Haziza, D. and Beaumont, J.-F. (2017). Construction of Weights in Surveys: A Review. Statistical Science, 32(2):206 – 226.
- Isaki, C. T. and Fuller, W. A. (1982). Survey design under the regression superpopulation model. Journal of the American Statistical Association, 77(377):89–96.
- Kim, J. K. (2010). Calibration estimation using exponential tilting in sample surveys. Survey Methodology, 36(2):145–155.
- Liu, Y. and Fan, Y. (2023). Biased-sample empirical likelihood weighting for missing data problems: an alternative to inverse probability weighting. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 85(1):67–83.
- McConville, K. S., Breidt, F. J., Lee, T. C. M., and Moisen, G. C. (2017). Model-assisted survey regression estimation with the LASSO. *Journal of Survey Statistics and Methodology*, 5:131–158.
- Montanari, G. E. (1987). Post-sampling efficient QR-prediction in large-sample surveys. International Statistical Review, 55(2):191–202.

- Montanari, G. E. and Ranalli, M. G. (2005). Nonparametric model calibration estimation in survey sampling. *Journal of the American Statistical Association*, 100(472):1429–1442.
- Newey, W. K. and Smith, R. J. (2004). Higher order properties of gmm and generalized empirical likelihood estimators. *Econometrica*, 72(1):219–255.
- Owen, A. B. (1988). Empirical likelihood ratio confidence intervals for a single functional. Biometrika, 75(2):237–249.
- Pfeffermann, D. and Sverchkov, M. (2009). Inference under informative sampling. Handbook of Statistics, vol. 29B. Amsterdam: Elsevier, pages 455–487.
- Qin, J., Leung, D., and Shao, J. (2002). Estimation with survey data under nonignorable nonresponse or informative sampling. *Journal of the American Statistical Association*, 97(457):193–200.
- Rao, J. (1994). Estimating totals and distribution functions using auxiliary information at the estimation stage. *Journal of official statistics*, 10(2):153.
- Rao, J. N. K., Jocelyn, W., and Hidiroglou, M. A. (2003). Confidence interval coverage properties for regression estimators in uni-phase and two-phase sampling. *Journal of Official Statistics*, 19:17–30.
- Robinson, P. and Särndal, C. E. (1983). Asymptotic properties of the generalized regression estimator in probability sampling. Sankhyā: The Indian Journal of Statistics, Series B, pages 240–248.

- Thelin, G. P. and Stone, W. W. (2013). Estimation of annual agricultural pesticide use for counties of the conterminous United States, 1992-2009. US Department of the Interior, US Geological Survey Sacramento, CA.
- Tsiatis, A. A. (2006). Semiparametric Theory and Missing Data. Springer.
- Wang, Y. and Zubizarreta, J. R. (2020). Minimal dispersion approximately balancing weights: asymptotic properties and practical considerations. *Biometrika*, 107:93–105.
- Wu, C. and Lu, W. W. (2016). Calibration weighting methods for complex surveys. International Statistical Review, 84(1):79–98.
- Wu, C. and Rao, J. N. K. (2006). Pseudo empirical likelihood ratio confidence intervals for complex surveys. *Canadian Journal of Statistics*, 34(3):359–375.
- Wu, C. and Sitter, R. R. (2001). A model-calibration approach to using complete auxiliary information from survey data. *Journal of the American Statistical Association*, 96(453):185–193.