

HERMITIAN NULL LOCI

QUANG-TUAN DANG

ABSTRACT. We establish a transcendental generalization of Nakamaye’s theorem to compact complex manifolds when the form is not assumed to be closed. We apply the recent analytic technique developed by Collins–Tosatti to show that the non-Hermitian locus of a nef and big $(1,1)$ -form, which is not necessarily closed, on a compact complex manifold equals the union of all positive-dimensional analytic subvarieties where the restriction of the form is not big (null locus). As an application, we can give an alternative proof of the Nakai–Moishezon criterion of Buchdahl and Lamari for complex surfaces and generalize this result in higher dimensions. This is also used for studying degenerate complex Monge–Ampère equations on compact Hermitian manifolds. Finally, we investigate finite time non-collapsing singularities of the Chern–Ricci flow, partially answering a question raised by Tosatti and Weinkove.

CONTENTS

1. Introduction	1
2. Preliminaries	6
3. Null loci and Nakai–Moishezon criteria	11
4. Singular Hermitian metrics and envelopes	20
5. Finite time singularities of the Chern–Ricci flow	26
References	32

1. INTRODUCTION

Characterizing the augmented base locus of a nef and big line bundle on a smooth algebraic variety over \mathbb{C} was initially motivated by applications to diophantine approximation problem [Nak15]. Nakamaye showed in [Nak00] that the augmented base locus of a nef and big line bundle on a projective manifold equals its null locus (which is the union of all irreducible subvarieties where the restriction of the bundle is not big). There are numerous generalizations of the latter result in \mathbb{R} -divisors [ELM⁺09], in possibly singular varieties [Bir17, CL14, BCL14], and in positive characteristics [CMM14]. We refer interested readers to [Tos18b] for a comprehensive overview.

Our primary goal in this paper is to extend Nakamaye’s theorem to the context of compact non-Kähler manifolds with non-closed differential forms. Let X be a compact complex manifold of dimension n , equipped with a Hermitian metric ω_X . We denote by $d^c = \frac{1}{2i}(\partial - \bar{\partial})$ so that $dd^c = i\partial\bar{\partial}$. Let θ be a smooth real $(1,1)$ form and let $[\theta]_{\partial\bar{\partial}}$ denote the $\partial\bar{\partial}$ -class of forms (resp. currents): $[\theta]_{\partial\bar{\partial}} = \{\theta + dd^c\varphi : \varphi \in C^\infty(X, \mathbb{R}) \text{ (resp. } u \in L^1(X))\}$. For simplicity, we omit the subscript $\partial\bar{\partial}$. When θ is closed, the $\partial\bar{\partial}$ -class $[\theta]$ coincides with its Bott–Chern cohomology class in

Date: April 2, 2024.

2020 Mathematics Subject Classification. 32W20, 53E30, 32U15.

Key words and phrases. Hermitian currents, null loci, Chern–Ricci flows, complex manifolds.

$H_{\text{BC}}^{1,1}(X, \mathbb{R})$. A $\partial\bar{\partial}$ -class $[\theta]$ is said to be *Hermitian* (or *positive*) if it contains at least a positive definite form, i.e., $\exists \varphi \in \mathcal{C}^\infty(X, \mathbb{R})$ such that $\theta + \text{dd}^c \varphi$ is a Hermitian metric. Although θ is not closed in general, one can still meaningfully define the concept (in the analytic sense) of nef $\partial\bar{\partial}$ -classes; cf. Def. 2.1. Similarly, a $\partial\bar{\partial}$ -class $[\theta]$ is said to be *big* if it contains a positive current dominating a Hermitian metric. It should be emphasized that the previous concepts can be defined when the underlying variety is possibly singular; cf. Sect. 2.3. By Demailly's regularization theorem, the Hermitian current in a big class can be taken to have (almost) analytic singularities (in the sense of [DP04, Theorem 3.2]). In particular, the Hermitian current is smooth in the complement of a Zariski closed subset of X . The smallest possible Zariski closed subset of X that one can take in this way is the non-Hermitian locus of a big $\partial\bar{\partial}$ -class $[\theta]$, denoted by $E_{\text{nH}}(\theta)$; cf. [Bou04, Definition 3.16].

A nef and big $\partial\bar{\partial}$ -class $[\theta]$, in general, may not be positive, and another subset to measure its non-positivity is its null locus, denoted by $\text{Null}(\theta)$, explicitly introduced by Nakamaye (possibly investigated earlier). The latter subset is defined as the union of all positive-dimensional irreducible analytic subvarieties $V \subset X$ where the restriction $\theta|_V$ is not big. Clearly, the null locus is a proper subset of X . Repeating line by line the proof of Collins–Tosatti, one can prove that given a nef and big $\partial\bar{\partial}$ -class $[\theta]$, its non-Hermitian locus and its null locus coincide. The proof does not require the closedness of the representative form.

Theorem 1.1 ([CT15, Theorem 1.1]). *Let X be a compact complex manifold. Let θ be a smooth real (1,1) form whose $\partial\bar{\partial}$ -class $[\theta]$ is nef and big. Then*

$$E_{\text{nH}}(\theta) = \text{Null}(\theta).$$

In the Kähler case (or the Fujiki class \mathcal{C} in general), the bigness of a closed nef form θ is equivalent to the positivity property of its self-intersection number, i.e., $\int_X \theta^n > 0$. Unlike the Kähler case, the difficulty we face is that the volume of two forms in the same $\partial\bar{\partial}$ -class might be different in general. Let X be a compact complex manifold of dimension n , equipped with a Hermitian metric ω_X . A question raised by Guedj–Lu is whether

$$v_+(X, \omega_X) = \sup \left\{ \int_X (\omega_X + \text{dd}^c \varphi)^n : \varphi \in \mathcal{C}^\infty(X), \omega_X + \text{dd}^c \varphi > 0 \right\} < +\infty;$$

$$v_-(X, \omega_X) = \inf \left\{ \int_X (\omega_X + \text{dd}^c \varphi)^n : \varphi \in \mathcal{C}^\infty(X), \omega_X + \text{dd}^c \varphi > 0 \right\} > 0.$$

We refer to [AGL23] for some classes of manifolds satisfying the condition above. We say that a Hermitian metric ω_X satisfies the volume condition (H_X) if:

$$(H_X) \quad v_+(X, \omega_X) < +\infty \quad \text{and} \quad v_-(V, \omega_X|_V) > 0$$

for all positive-dimensional closed submanifold $V \subset X$.

Remark 1.2. It has been shown in [AGL23, Theorem A] that for any closed submanifold $V \subset X$,

$$v_+(X, \omega_X) < +\infty \implies v_+(V, \omega_X|_V) < +\infty.$$

It is important to emphasize that the assumption (H_X) occurs in some contexts of complex differential geometry. For instance, if ω_X satisfies the following condition $\text{dd}^c \omega_X = 0$, $d\omega_X \wedge d^c \omega_X = 0$ then by [Chi16b], it is equivalent to the volume with respect to a form belonging to the $\partial\bar{\partial}$ -class $[\omega_X]$ is unchanged. That is, for any closed submanifold $V \subset X$,

$$\int_V (\omega_X|_V + \text{dd}^c \varphi)^{\dim V} = \int_V \omega_X|_V^{\dim V}, \forall \omega_X|_V + \text{dd}^c \varphi > 0.$$

These metrics were introduced by Guan–Li [GL10], which are called the Guan–Li metrics in this paper, although they were probably studied earlier; cf. [FT11].

Given θ a smooth real (1,1) form on X , if θ is nef, we can define

$$\widehat{v}_-(X, \theta) := \inf_{\varepsilon > 0} v_-(X, (1 - \varepsilon)\theta + \varepsilon\omega_X).$$

Although the form $(1 - \varepsilon)\theta + \varepsilon\omega_X$ is not positive, one could find by definition, for any $\varepsilon > 0$, a smooth function φ_ε such that $\theta + \text{dd}^c\varphi_\varepsilon \geq -\varepsilon\omega_X$ so that

$$(1 - \varepsilon)\theta + \varepsilon\omega_X + \text{dd}^c(1 - \varepsilon)\varphi_\varepsilon \geq \varepsilon^2\omega_X.$$

Hence the quantity $v_-(X, (1 - \varepsilon)\theta + \varepsilon\omega_X)$ can be understood as $v_-(X, (1 - \varepsilon)\theta + \varepsilon\omega_X + \text{dd}^c(1 - \varepsilon)\varphi_\varepsilon)$. Our main theorem is the following:

Theorem A. *Let X be a compact complex manifold equipped with a Hermitian metric ω_X . Assume that ω_X satisfies the condition (H_X). Let θ be a smooth real (1,1) form whose $\partial\bar{\partial}$ class is nef with $\widehat{v}_-(X, \theta) > 0$. Then $[\theta]$ is big and*

$$(1.1) \quad E_{nH}(\theta) = \text{Null}(\theta) = \bigcup_{\widehat{v}_-(V, \theta|_V) = 0} V$$

where the union is taken over all purely positive-dimensional irreducible analytic subvarieties V of X .

In particular, when θ is closed or merely satisfies the so-called Guan–Li condition: $\text{dd}^c\theta = 0$, $d\theta \wedge d^c\theta = 0$ (equivalent to $\text{dd}^c\theta^k = 0$ for any $k = 1, \dots, n - 1$), then we obtain $\widehat{v}_-(V, \theta|_V) = \int_V \theta^{\dim V}$ by [GL22, Lemma 4.4], provided that $v_+(X, \omega_X) < \infty$. This directly implies the result of Collins–Tosatti in the case of compact complex manifolds belonging to the Fujiki class \mathcal{C} .

A direct application of Theorem A is to establish the Nakai–Moishezon criterion for some compact non-Kähler manifolds, which generalizes the fundamental result of Demailly–Păun [DP04] for compact Kähler manifolds, and Buchdahl [Buc00] and Lamari [Lam99] for compact non-Kähler surfaces. Our proof relies essentially on the mass concentration result of Demailly–Păun, whose proof was simplified by Chiose [Chi16a] and Popovici [Pop16]. Using Theorem A allows one to avoid the inductive argument on dimension established in [DP04]. Inspired by the works of Collins–Tosatti [CT16], and Das–Hacon–Păun [DHP22], we establish the Nakai–Moishezon criterion on a possibly singular compact analytic normal variety, but the proof relies on the induction on dimension.

Theorem B. *Let X be a compact analytic normal variety of pure dimension n and ω_X be a Hermitian metric on X such that $\text{dd}^c\omega_X^k = 0$ for any $k = 1, \dots, n - 1$. Let θ be a smooth real (1,1) form on X such that*

- (1) $\text{dd}^c\theta^k = 0$, for any $k = 1, \dots, n - 1$;
- (2) for any $k = 1, \dots, n$,

$$\int_X \theta^k \wedge \omega_X^{n-k} > 0;$$

- (3) for every irreducible analytic subvariety $V \subset X$ with $\dim V > 0$,

$$\int_V \theta^k \wedge \omega_X^{\dim V - k} > 0, \quad \forall k = 1, \dots, \dim V.$$

Then the $\partial\bar{\partial}$ -class $[\theta]$ is Hermitian.

We remark that the condition (2) can be absorbed in the (3) when $V = X$. In the case of complex smooth surfaces, thanks to [Buc99, Proposition 5], we can relax the condition (3) to require only that $\int_V \theta > 0$ for every irreducible curve $V \subset X$ with strictly negative self-intersection $V \cdot V < 0$. We refer to Sect. 2.3 for

the definition of differential forms, Hermitian metrics, and integrals on singular analytic varieties.

The following application is to show the existence of solutions to degenerate complex Monge–Ampère equations in the context of nef and big form. We extend the result of Guedj–Lu [GL23, Theorem D] for semi-positive reference forms to nef ones. We precisely obtain the following.

Theorem C. *Let (X, ω_X) an n -dimensional compact complex manifold. Let θ be a smooth real $(1,1)$ form whose $\partial\bar{\partial}$ -class $[\theta]$ is nef and big. Let $f \in C^\infty(X, \mathbb{R})$. Then, there exists a constant $c > 0$ and an θ -psh function φ which is smooth on a Zariski open set U of X , $\sup_X \varphi = 0$ and*

$$(\theta + \text{dd}^c \varphi)^n = cf\omega_X^n$$

on U . More precisely, U is taken to be the complement $X \setminus \text{Null}(\theta)$ of the null locus of θ .

Extending the works of Berman [Ber19] to the non-Kähler case, we study the partial regularity of θ -plurisubharmonic envelope when the $\partial\bar{\partial}$ -class $[\theta]$ is assumed to be nef and big. Fixing a function $f \in C^\infty(X, \mathbb{R})$ and a Hermitian metric ω_X on X , for $\beta > 0$, we consider the following complex Monge–Ampère equation for a smooth solution φ_β :

$$(1.2) \quad (\theta + 2^{-\beta}\omega_X + \text{dd}^c \varphi_\beta)^n = e^{\beta(\varphi_\beta - f)}\omega_X^n.$$

By the fundamental works of Cherrier [Che87] and Tosatti–Weinkove [TW10a], the equation (1.2) admits a unique smooth solution. Indeed, since θ is nef so its $\partial\bar{\partial}$ -class $[\theta + 2^{-\beta}\omega_X]$ is Hermitian for any $\beta > 0$. This shows the existence of smooth solutions and the form $\omega_\beta := \theta + 2^{-\beta}\omega_X + \text{dd}^c \varphi_\beta$ defines a Hermitian metric in $[\theta + 2^{-\beta}\omega_X]$. Since $[\theta]$ is big, we can define the so-called θ -plurisubharmonic envelope $P_\theta(f)$ of f as the largest θ -plurisubharmonic function lying below f . We study the regularity of the envelope outside the null locus.

Theorem D. *Let θ be a smooth real $(1,1)$ form on a n -dimensional compact complex manifold X whose $\partial\bar{\partial}$ class $[\theta]$ is nef and big. Let $f \in C^\infty(X, \mathbb{R})$ and let $P_\theta(f)$ denote the corresponding θ -plurisubharmonic envelope of f . Let φ_β be the unique smooth solution of the complex Monge–Ampère equation (1.2). Then, as $\beta \rightarrow \infty$, the functions φ_β converge to $P_\theta(f)$ in $C_{\text{loc}}^{1,\alpha}(X \setminus \text{Null}(\theta))$ for any $\alpha \in]0, 1[$.*

The proof is an adaptation of [Ber19, Theorem 1.2] with slight changes. When θ is Hermitian, a strong result was proved in [CZ19] that $P_\theta(f)$ belongs to $C^{1,1}(X)$.

Finally, we study finite time singularities of the Chern–Ricci flow on compact complex manifolds, partially answering a question raised by Tosatti–Weinkove. Let $\omega = \omega(t)$ be a solution of the Chern–Ricci flow on X , for $t \in [0, T[$, starting at a Hermitian metric ω_0 ,

$$(1.3) \quad \frac{\partial \omega}{\partial t} = -\text{Ric}(\omega), \quad \omega(0) = \omega_0$$

where $\text{Ric}(\omega)$ is the Chern–Ricci form (also called first Chern form) of the Hermitian metric ω . It is classical that solving the Chern–Ricci flow boils down to solving a nonlinear parabolic scalar equation in $\varphi = \varphi(t)$ of the form

$$\frac{\partial \varphi}{\partial t} = \log \frac{(\theta_t + \text{dd}^c \varphi)^n}{\omega_0^n}, \quad \theta_t + \text{dd}^c \varphi > 0, \quad \varphi(0) = 0$$

where $\theta_t := \omega_0 - t \text{Ric}(\omega_0)$. Tosatti and Weinkove [TW15] characterized the maximal existence time T of the flow as

$$T = \sup\{t > 0 : \exists \psi \in C^\infty(X) \text{ with } \theta_t + \text{dd}^c \psi > 0\}$$

by this, we mean that T is such that for all $t < T$, the $\partial\bar{\partial}$ -class $[\theta_t]$ is Hermitian. Suppose the flow develops a finite time singularity $T < \infty$. We observe that the limiting $\partial\bar{\partial}$ class $[\theta_T]$ is nef but not Hermitian. We define the set of singularities of the flow, denoted by $\text{Sing}(\omega(t))$, to be the complement of the set of all points $x \in X$ such that there is a neighborhood $U \ni x$ on which $\omega(t)$ converge smoothly to a Hermitian metric ω_T as $t \rightarrow T^-$. Furthermore, we can show that the singularity set of the Chern–Ricci flow is equal to the set on which the Chern scalar curvature blows up. The latter was studied by Gill–Smith [GS15], generalizing Zhang’s analog result for Kähler–Ricci flows.

Tosatti and Weinkove [TW22, Question 6.1] posed the question of whether singularities of the Chern–Ricci flow develop precisely along analytic subvarieties of X ? In the Kähler case, this question was initially raised by Feldman–Ilmanen–Knopf and later affirmatively answered by Collins–Tosatti [CT15, Theorem 1.5]. Gill–Smith [GS15, Theorem 1.2] gave an affirmative answer for the case when X is a complex surface and ω_0 is Gauduchon. As observed in [TW22, Section 6], the answer is affirmative when the volume with respect to $\omega(t)$ shrinks to zero as $t \rightarrow T$. We investigate non-collapsing the case when $\hat{v}_-(X, \theta_T) > 0$ and provide an answer to this question on some underlying manifolds.

Theorem E. *Let (X, ω_0) be a compact Hermitian manifold of dimension n . Let $\omega(t)$ be a solution of the Chern–Ricci flow starting at ω_0 , which develops a singularity at finite time T . Assume that the limiting form $\theta_T = \omega_0 - T \text{Ric}(\omega_0)$ satisfies $\hat{v}_-(X, \theta_T) > 0$. Then if either*

- $n = 2$ and ω_0 is arbitrary,
- or $n \geq 3$, $\text{dd}^c \omega_0 = 0$, $d\omega_0 \wedge d^c \omega_0 = 0$,

then the metrics $\omega(t)$ develop singularities precisely along the analytic set $\text{Null}(\theta_T)$.

This theorem generalizes Gill–Smith’s result when the underlying manifold X is a surface, but the initial metric ω_0 is given to be arbitrary and also partially answers the question of Tosatti–Weinkove. With assumptions in this theorem, we have that the limiting form θ_T is big, so its null locus $\text{Null}(\theta_T)$ is a proper analytic subset of X (since $[\theta_T]$ is not Hermitian), and our theorem shows that no singularities develop on its complement. We have seen from Theorem A that $\text{Null}(\theta_T)$ is precisely the union of all purely positive-dimensional irreducible subvarieties V of X with $v_-(V, \omega(t))$ shrinking to zero as $t \rightarrow T^-$, so singularities of the flow develop at least there. We will see in Theorem 5.1 that the scalar curvature of $\omega(t)$ is locally uniformly bounded on compact sets outside singularities. Furthermore, it should be emphasized that in the setting of our theorem, as $t \rightarrow T^-$, the metrics $\omega(t)$ converge weakly to a positive current ω_T which is smooth on $X \setminus \text{Null}(\theta_T)$ and has minimal singularities.

A crucial case of such finite-time singularities occurs when the $\partial\bar{\partial}$ -class $[\theta_T]$ contains a pullback of a Hermitian metric via the blowdown of an exceptional divisor to a point. Interesting results were obtained by Tosatti–Weinkove [TW13], Tô [Tô18], and Nie [Nie17], in which they proved that the Chern–Ricci flow performs a *canonical surgical contraction*.

This paper is structured as follows. In Sect. 2, we introduce notation and definitions, and briefly review fundamental results extended to the non-Kähler context. In Sect. 3, we establish an extension-type theorem for Hermitian currents with analytic singularities, which enables us to prove Theorem A, and we also prove Theorem B there. Sect. 4 is dedicated to the study of degenerate complex Monge–Ampère equations, where we prove Theorems C and D. Finally, we investigate finite time singularities of the Chern–Ricci flow and prove Theorem E in Sect. 5.

Notation. Unless otherwise mentioned, X denotes a compact Hermitian manifold with a reference Hermitian metric ω_X . We interchangeably use the Hermitian metric ω_X and its associated Riemannian metric g . We denote by $d = \partial + \bar{\partial}$ and $d^c = \frac{i}{2}(\bar{\partial} - \partial)$ so that $dd^c = i\partial\bar{\partial}$. Throughout the paper, we use C to represent a positive constant, which may vary from line to line and can be uniformly controlled.

Acknowledgement. We would like to express our gratitude to V. Tosatti for suggesting this problem, his interest in this work, and his valuable comments. We are grateful to T.-D. Tô for insightful discussions and C. H. Lu for carefully reading the first draft. This work is partially supported by the ANR project PARAPLUI.

2. PRELIMINARIES

Let X be a n -dimensional compact complex manifold. Fix ω_X a reference Hermitian metric.

2.1. $\partial\bar{\partial}$ -classes. It is well-known that the $\partial\bar{\partial}$ -lemma does not hold on an arbitrarily complex manifold, so it is better to work with the $\partial\bar{\partial}$ -class of differential forms. We denote by $\mathcal{A}^{p,q}(X, \mathbb{C})$ (resp. $\mathcal{A}^{p,q}(X, \mathbb{R})$) the space of smooth (resp. real) forms of bidegree (p, q) . We say that two forms $\theta, \eta \in \mathcal{A}^{p,q}(X, \mathbb{C})$ are equivalent if they are modulo to the $\partial\bar{\partial}$ -exact ones, i.e., there exists a smooth form $\psi \in \mathcal{A}^{p-1, q-1}(X, \mathbb{C})$ such that

$$\theta - \eta = dd^c\psi.$$

Since then, we let $[\theta]_{\partial\bar{\partial}}$ denote the $\partial\bar{\partial}$ -class with smooth representative θ . For simplicity, we skip the subscript $\partial\bar{\partial}$, and unless we emphasize the $\partial\bar{\partial}$ class $[\theta]$ is interpreted as the set of forms $\theta + dd^c\psi$ for all $\psi \in \mathcal{C}^\infty(X)$.

We remark that a d -exact form is not $\partial\bar{\partial}$ -exact in the general context of non-Kähler manifolds. A compact complex manifold X is said to be a $\partial\bar{\partial}$ -manifold if it satisfies the exactness property: for any d -closed pure-type form u on X , u is d -exact $\iff u$ is $\partial\bar{\partial}$ -exact for any d -closed form u on X . In particular, a compact Kähler manifold is a $\partial\bar{\partial}$ -manifold.

Definition 2.1. A smooth real $(1,1)$ form θ is called *nef* if for any $\varepsilon > 0$ there exists a form $\theta_\varepsilon \in [\theta]$ such that $\theta_\varepsilon \geq -\varepsilon\omega_X$, i.e., $\exists \psi_\varepsilon \in \mathcal{C}^\infty(X, \mathbb{R})$ such that $\theta + dd^c\psi_\varepsilon \geq -\varepsilon\omega_X$. In this case, the $\partial\bar{\partial}$ -class $[\theta]$ is said to be nef.

If there exists a Hermitian metric $\omega \in [\theta]$, then we say that the $\partial\bar{\partial}$ -class $[\theta]$ is *Hermitian/positive*.

It should be emphasized that unlike in the Kähler case, even if two Hermitian metrics ω and ω' are in the same $\partial\bar{\partial}$ class, i.e., there is $\varphi \in \mathcal{C}^\infty(X)$ such that $\omega' = \omega + dd^c\varphi$, the volume with respect to ω does not equal the one with respect to ω' . It was shown by Chiose [Chi16b, Theorem 0.1] that the preservation of Monge–Ampère masses is equivalent to the condition introduced by Guan–Li [GL10]: $\int_X \omega^n = \int_X \omega'_X$ for all $\omega \in [\omega_X]$ Hermitian if and only if $dd^c\omega_X = 0$ and $d\omega_X \wedge d^c\omega_X = 0$. The latter condition is equivalent to $dd^c\omega_X^k = 0$ for all $k = 1, \dots, n-1$; see also [Chi16b, Theorem 0.2] for more equivalent conditions.

We recall some quantities on Monge–Ampère volume bounds introduced by Guedj–Lu [GL22].

Definition 2.2. We define

$$v_+(X, \omega_X) = \sup \left\{ \int_X \omega^n : \forall \omega \in [\omega_X], \omega \text{ Hermitian} \right\}$$

and

$$v_-(X, \omega_X) = \inf \left\{ \int_X \omega^n : \forall \omega \in [\omega_X], \omega \text{ Hermitian} \right\}.$$

A question raised by Guedj–Lu [GL22] is whether $v_+(X, \omega_X) < +\infty$ and/or $v_-(X, \omega_X) > 0$. We remark that these conditions somewhat depend on the complex structure, but they are independent of the choice of a Hermitian metric; see [GL22, Prop. 3.2]. Furthermore, they are bimeromorphic invariants, and so is the condition (H_X) .

Theorem 2.3 ([GL22, Thm. 3.7]). *Let $\pi : X \rightarrow Y$ be a bimeromorphic map between compact complex manifolds equipped with Hermitian metrics ω_X and ω_Y . Then ω_X satisfies the condition (H_X) if and only if ω_Y satisfies the condition (H_X) .*

Definition 2.4. If θ is nef, we can define

$$\widehat{v}_-(X, \theta) := \inf_{\varepsilon > 0} v_-(X, (1 - \varepsilon)\theta + \varepsilon\omega_X).$$

Although the form $(1 - \varepsilon)\theta + \varepsilon\omega_X$ is not positive, one could find by definition, for any $\varepsilon > 0$, a smooth function φ_ε such that $\theta + \text{dd}^c \varphi_\varepsilon \geq -\varepsilon\omega_X$ so that

$$(1 - \varepsilon)\theta + \varepsilon\omega_X + \text{dd}^c(1 - \varepsilon)\varphi_\varepsilon \geq \varepsilon^2\omega_X.$$

Hence the quantity $v_-(X, (1 - \varepsilon)\theta + \varepsilon\omega_X)$ is interpreted as $v_-(X, (1 - \varepsilon)\theta + \varepsilon\omega_X + \text{dd}^c(1 - \varepsilon)\varphi_\varepsilon)$. We expect that the definition above coincides with the one when θ is closed. The necessary condition is that θ satisfies the Guan–Li property.

Lemma 2.5 ([GL22, Lemma 4.4]). *If $v_+(X, \omega_X) < \infty$ and θ is a smooth nef $(1,1)$ form satisfying $\text{dd}^c \theta^k = 0$ for any $k = 1, \dots, n - 1$ then $\widehat{v}_-(X, \theta) = \int_X \theta^n$. Moreover, for any positive-dimensional closed submanifold $V \subset X$, $\widehat{v}_-(V, \theta|_V) = \int_V \theta^{\dim V}$.*

The proof is identical to that of [GL22, Lemma 4.4], which we omit here. Notice that the Guan–Li property is sufficient to guarantee the preservation of Monge–Ampère masses.

2.2. Quasi-plurisubharmonic functions and Hermitian currents. Recall that an upper semi-continuous function $\varphi : X \rightarrow \mathbb{R} \cup \{-\infty\}$ is *quasi-plurisubharmonic* (*quasi-psh* for short) if it is locally described as the sum of a plurisubharmonic (psh) function and a smooth function.

Definition 2.6. Let θ be a smooth $(1,1)$ form on X . One says that

- a quasi-psh function φ is *θ -plurisubharmonic* (*θ -psh* for short) if $\theta + \text{dd}^c \varphi \geq 0$ in the sense of currents. We let $\text{PSH}(X, \theta)$ denote the space of all θ -psh functions on X which is not identically $-\infty$;
- T is a positive $(1,1)$ current on X in the $\partial\bar{\partial}$ -class $[\theta]$ if $T = \theta + \text{dd}^c \varphi$ for some $\varphi \in \text{PSH}(X, \theta)$, and vice versa.

For any positive $(1,1)$ current T , we let $\text{Sing}(T)$ denote the *singular set* of T , namely, if $T = \theta + \text{dd}^c \varphi$ then $\text{Sing}(T) = \{\varphi = -\infty\}$.

We refer the reader to nice references [Dem12, GZ17] for more details about the basic properties of psh functions and positive currents.

The *complex Monge–Ampère measure* $(\theta + \text{dd}^c u)^n$ is well-defined for any θ -psh function u , which is *bounded* on X , as follows from Bedford–Taylor’s theory; cf. [DK12] or [GL22] for a brief recap. We recall the following maximum principle.

Lemma 2.7 (Maximum principle). *Let φ, ψ be bounded θ -psh functions in $U \subseteq X$ such that $\varphi \geq \psi$. Then*

$$\mathbf{1}_{\{\varphi=\psi\}}(\theta + \text{dd}^c \varphi)^n \geq \mathbf{1}_{\{\varphi=\psi\}}(\theta + \text{dd}^c \psi)^n.$$

Proof. The proof directly follows Bedford–Taylor’s maximum principle [GZ17, Theorem 3.27] in the local context. See also [GL22, Lemma 1.2] for a proof in the global one. \square

Definition 2.8. Let $\mathcal{I} \subset \mathcal{O}_X$ be an analytic coherent ideal sheaf and $c > 0$. A quasi-psh function φ on X is said to have *almost analytic singularities* of type (\mathcal{I}, c) if the following are satisfied

- (i) for any $x \in X$, there exists a neighborhood U of x such that on U we have

$$\varphi = c \log \left(\sum_j |f_j|^2 \right) + g$$

where the f_j ’s are local generators of a coherent sheaf \mathcal{I} , and g is a bounded function on U ,

- (ii) there is a (global) proper modification $\pi : X' \rightarrow X$ of X , obtained as a finite composition of blow-ups with smooth centers, and an effective divisor $\sum_\ell a_\ell D_\ell$ with normal crossings such that one can write $\varphi \circ \pi$ locally as

$$\varphi \circ \pi = c \sum_\ell a_\ell \log |g_\ell|^2 + h$$

where $(g_\ell = 0)$ are local equations of the divisors D_ℓ and h is a smooth function.

If a current $T = \theta + \text{dd}^c \varphi$ with φ a quasi-psh function having almost analytic singularities (along subscheme $V(\mathcal{I})$ induced by \mathcal{I}), then one says that T has *almost analytic singularities*.

Let us emphasize that the definition above is a bit lengthy but turns out to be the most suitable one for this paper. We will see below the $\partial\bar{\partial}$ -class with appropriate positivity contains plenty of currents with such singularities.

Definition 2.9. We say that a real (1,1)-current T is a *Hermitian current* if there exists a Hermitian metric ω such that $T \geq \omega$.

The $\partial\bar{\partial}$ -class $[\theta]$ is said to be *big* if it contains a Hermitian current, i.e., there exists $\varphi \in \text{PSH}(X, \theta)$ such that $T = \theta + \text{dd}^c \varphi \geq \omega$ for some Hermitian form ω . In this case, the smooth representative θ is also called a *big (1,1) form*.

We introduce the notion of psh envelopes, understood as the *least singularity* function lying below a prescribed one.

Definition 2.10. Given a Borel function $f : X \rightarrow \mathbb{R}$, we define the θ -psh envelope of f by

$$P_\theta(f) := (\sup\{u \in \text{PSH}(X, \theta) : u \leq f\})^*$$

where the $*$ means that we take the upper semi-continuous regularization.

The θ -psh envelope $P_\theta(f)$ might be $\equiv -\infty$ when the $\partial\bar{\partial}$ -class $[\theta]$ contains no positive currents. When θ is big, and f is bounded from below, $P_\theta(f)$ is well-defined as a θ -psh function.

We recall the basic type of regularization due to Demailly; cf. [DP04, Lemma 2.1, Theorem 3.2].

Theorem 2.11 (Demailly’s regularization theorem). *Let X be a compact complex manifold equipped with a Hermitian form ω . Let $T = \theta + \text{dd}^c \varphi$ be a real (1,1) current; i.e., θ is a smooth real (1,1) form and φ is quasi-psh. Suppose that $T \geq \gamma$ for some smooth (1,1) form γ . Then, there exists a decreasing sequence of quasi-psh functions φ_m such that*

- (1) $T_m = \theta + \text{dd}^c \varphi_m$ are currents with almost analytic singularities;
- (2) $T_m \rightarrow T$ weakly, and φ_m decreases to φ as $m \rightarrow \infty$;

(3) $T_m \geq \gamma - \varepsilon_m \omega$ for some sequence $\varepsilon_m \searrow 0$.

By replacing T (resp. γ) with $T - \theta$ (resp. $\gamma - \theta$) we can work with $T = \text{dd}^c \varphi$ a closed quasi-positive current. The proof relies on the appropriate technique using an asymptotic Bergman kernel procedure as we describe what happens locally. If φ is a psh function on the ball $B \subset \mathbb{C}^n$ one considers for any $m \geq 1$ the Hilbert space $\mathcal{H}(B, m\varphi)$ of L^2 holomorphic function on B such that $\int_B |f|^2 e^{-m\varphi} dV < \infty$. We choose a Hilbert basic $(f_k^m)_{k \in \mathbb{N}}$ of $\mathcal{H}(B, m\varphi)$ and put $\varphi_m := \frac{1}{m} \log(\sum_{k \in \mathbb{N}} |f_k^m|^2)$. One can check that φ_m is well-defined, psh, with analytic singularities and almost increasing. The difficult part is then to prove that φ_m converges to φ as m goes to infinity. To get the global approximations, one glues the local ones by using a partition of unity argument. In this gluing process, a small loss of positivity is established due to the change of holomorphic coordinates. This is where the ε_m 's come from. For each m , the corresponding modification μ_m is then obtained by principalization of singularities of the multiplier ideal sheaf $\mathcal{I}(m\varphi)$ so that $\mu_m^* \mathcal{I}(m\varphi)$ is the principal ideal sheaf associated with a normal crossing divisor, due to the fundamental result of Hironaka; cf. [Wlo09].

From the above regularization, we are able to approximate a positive current by Hermitian currents with almost analytic singularities, with reasonable control of singularities. In particular, a big $\partial\bar{\partial}$ -class $[\theta]$ contains various Hermitian currents with almost analytic singularities.

We introduce the definition of the non-Hermitian locus of a big $\partial\bar{\partial}$ -class, analog to the one in [Bou04].

Definition 2.12 (non-Hermitian locus). If θ is a big (1,1) form on X , we define its *non-Hermitian locus* to be the set

$$E_{\text{nH}}(\theta) = \bigcap_{T \in [\theta]} \text{Sing}(T)$$

where the intersection is taken over all Hermitian currents with almost analytic singularities in the $\partial\bar{\partial}$ -class $[\theta]$.

We investigate the bigness in some particular context of fixed Hermitian metric.

Theorem 2.13 (Guedj–Lu). *Let X be a compact complex manifold equipped with a Hermitian metric ω_X satisfying $v_+(\omega_X) < +\infty$. Let θ be a nef (1,1) form with $\widehat{v}_-(X, \theta) > 0$. Then the $\partial\bar{\partial}$ -class $[\theta]$ is big, i.e., there exists a Hermitian current T in $[\theta]$ with almost analytic singularities.*

Proof. This was proved in [GL22, Theorem 4.6], which was inspired by the one of Chiose [Chi16a] and Popovici [Pop16] relying on the characterization result of Lamari [Lam99]. For the reader's convenience, we give the proof here. According to the Hahn–Banach theorem as in [Lam99, Lemme 3.3], it thus suffices to show that there exists a constant $\delta_0 > 0$ such that

$$\int_X \theta \wedge \eta^{n-1} \geq \delta_0 \int_X \omega_X \wedge \eta^{n-1}$$

for all Gauduchon metrics η . Suppose by contradiction that for each $\varepsilon > 0$ there exists Gauduchon metrics η_ε such that

$$\int_X \theta \wedge \eta_\varepsilon^{n-1} \leq \varepsilon \int_X \omega_X \wedge \eta_\varepsilon^{n-1}.$$

We can normalize η_ε so that $\int_X \omega_X \wedge \eta_\varepsilon^{n-1} = 1$. We fix a function $\psi_\varepsilon \in C^\infty(X)$ such that $\theta_\varepsilon := \theta + \varepsilon \omega_X + \text{dd}^c \psi_\varepsilon$ is a Hermitian metric. By the main result of [TW10a] there exist $c_\varepsilon > 0$ and $u_\varepsilon \in \text{PSH}(X, \theta_\varepsilon) \cap C^\infty(X)$ such that $\sup_X u_\varepsilon = 0$ and

$$(\theta_\varepsilon + \text{dd}^c u_\varepsilon)^n = c_\varepsilon \omega_X \wedge \eta_\varepsilon^{n-1}.$$

By normalization we have

$$c_\varepsilon = \int_X (\theta_\varepsilon + \text{dd}^c u_\varepsilon)^n \geq \widehat{v}_-(X, \theta) > 0.$$

We apply [GL22, Lemma 4.13] which reformulates the one in [Pop16, Lemma 3.1] to obtain

$$(2.1) \quad \int_X (\theta_\varepsilon + \text{dd}^c u_\varepsilon) \wedge \eta_\varepsilon^{n-1} \times \int_X (\theta_\varepsilon + \text{dd}^c u_\varepsilon)^{n-1} \wedge \omega_X \geq \frac{c_\varepsilon}{n}.$$

The first term on the left-hand side can be written as $\int_X (\theta + \varepsilon \omega_X) \wedge \eta_\varepsilon^{n-1}$ since η_ε is Gauduchon and by assumption,

$$\int_X (\theta + \varepsilon \omega_X) \wedge \eta_\varepsilon^{n-1} \leq 2\varepsilon.$$

For the second term, it follows by the assumption that

$$(2.2) \quad \int_X (\theta_\varepsilon + \text{dd}^c u_\varepsilon)^{n-1} \wedge \omega_X \leq Mv_+(\omega_X)$$

which is bounded from above. Therefore we obtain

$$2\varepsilon Mv_+(\omega_X) \geq \frac{\widehat{v}_-(X, \theta)}{n} > 0$$

which is a contradiction as $\varepsilon \rightarrow 0$. The positive current can be taken to have almost analytic singularities thanks to Theorem 2.11. \square

Theorem 2.14. *Let θ be a big (1,1)-form. Then there exists a Hermitian current T_0 with almost analytic singularities such that $E_{\text{nH}}(\theta) = \text{Sing}(T_0)$. In particular, the non-Hermitian locus of a big class $[\theta]$ is an analytic subvariety of X .*

Proof. The proof is quite close to that of [Bou04, Theorem 3.17]. We denote by \mathcal{S} the set of all Hermitian currents with almost analytic singularities in the class $[\theta]$. Fix $T_j = \theta + \text{dd}^c \varphi_j \in \mathcal{S}$ for $j = 1, 2$. First, we claim that there exists a Hermitian current with almost analytic singularities T such that $\text{Sing}(T) \subset \text{Sing}(T_1) \cap \text{Sing}(T_2)$. Indeed, if we set $\varphi_3 = \max(\varphi_1, \varphi_2)$ then $T_3 = \theta + \text{dd}^c \varphi_3$ is a Hermitian current such that $(\varphi_3 = -\infty) \subset \text{Sing}(T_1) \cap \text{Sing}(T_2)$. By Theorem 2.11, we can find a Hermitian current $T = \theta + \text{dd}^c \varphi \in \mathcal{S}$ with almost analytic singularities such that $\varphi \geq \varphi_3$ so $\text{Sing}(T) \subset \text{Sing}(T_1) \cap \text{Sing}(T_2)$. This proves our claim.

Thanks to the claim above and Theorem 2.11 again, we can construct a sequence of currents $T_m = \theta + \text{dd}^c \varphi_m \in \mathcal{S}$ such that $\text{Sing}(T_m)$ is a decreasing sequence of analytic subsets. By the strong Noetherian property, $\text{Sing}(T_m)$ must be stationary for $m \geq m_0$. Therefore, we obtain $E_{\text{nH}}(\theta) = \text{Sing}(T_0)$ for $T_0 := T_{m_0}$, as desired.

Moreover, if $E_{\text{nH}}(\theta) = \emptyset$ then $\text{Sing}(T_0) = \emptyset$, it means that T_0 is smooth on X , so T_0 is in fact a Hermitian form. \square

To summarize, we have the following.

Theorem 2.15. *Let X be a compact complex manifold equipped with a positive form ω_X satisfying $v_+(\omega_X) < +\infty$. Let θ be a nef (1,1) form with $\widehat{v}_-(X, \theta) > 0$. Then there exists a Hermitian current T in the $\partial\bar{\partial}$ -class $[\theta]$ with almost analytic singularities such that*

$$\text{Sing}(T) = E_{\text{nH}}(\theta).$$

We recall the following lemma, which implies that the non-Hermitian locus of a big form has no isolated points.

Lemma 2.16. *If θ is a big (1,1)-form, then the analytic set $E_{\text{nH}}(\theta)$ has no isolated points.*

Proof. We refer to [CT15, Lemma 3.1]. The closeness is not involved. \square

2.3. Extension on complex analytic spaces. We give here a short recap of the extension of the pluripotential theory to compact complex spaces. We refer the interested readers to [Dem85] for more details.

Let X be a reduced complex analytic space of pure dimension n . We will let X_{reg} denote the locus of regular points of X , and $X_{\text{sing}} := X \setminus X_{\text{reg}}$ the locus of singular ones. We see that X_{reg} is a complex manifold of dimension n and X_{sing} is an analytic subset of X of complex codimension ≥ 1 . If $X = \bigcup_{\alpha} U_{\alpha}$ is an open covering, we consider local embeddings $j_{\alpha} : U_{\alpha} \hookrightarrow \mathbb{C}^N$ onto analytic subsets of \mathbb{C}^N for some $N \geq 1$. By these local embeddings, one can define the sheaves of smooth (p, q) -forms $\mathcal{A}^{p,q}$, and the space of currents is, by definition, the dual of the space of differential forms as in the smooth case. The differential operators $d, \partial, \bar{\partial}, d^c$, and dd^c satisfy the usual rules and are then well defined by duality. We can also define the concepts of Hermitian $\partial\bar{\partial}$ -classes, nef $\partial\bar{\partial}$ -classes, big $\partial\bar{\partial}$ -classes, etc.

We next move on to the notion of psh functions. Similarly, by local embeddings above, we say that a function $u : X \rightarrow \mathbb{R} \cup \{-\infty\}$ is psh on X if it is locally the restrictions of a psh function defined on an open set Ω_{α} of \mathbb{C}^N containing $j_{\alpha}(U_{\alpha})$.

Another way to define a psh function is introduced by Demailly: a function u is weakly psh on X if it is locally bounded from above on X and its restriction to the complex manifold X_{reg} is psh. One can extend it to X by

$$u^*(x) := \limsup_{X_{\text{reg}} \ni y \rightarrow x} u(y).$$

The function u^* is upper semi-continuous, locally integrable on X , and $dd^c u^*$ is a positive current. Since X is locally integrable, thanks to [Dem85, Théorème 1.10], u^* is psh on X . In the same way, one can define the notion of quasi-psh functions with almost analytic singularities, Hermitian currents, non-Hermitian loci, etc.

We end this section by discussing integration over analytic (irreducible) subvarieties of a smooth complex manifold X . If V is an analytic subvariety of X of pure dimension $k > 0$, for any smooth (k, k) form α on X , then the integral

$$\int_V \alpha := \int_{V_{\text{reg}}} \alpha$$

is finite thanks to the fundamental result of Lelong; see e.g. [GH78, page 32]. In the same way, we can define the quantities $v_-(V, \omega_V)$, $v_+(V, \omega_V)$ and $\hat{v}_-(V, \theta)$. Furthermore, for smooth real $(k-1, k-1)$ -form β , we have by [GH78, page 33],

$$\int_V dd^c \beta = 0.$$

In particular, we can extend Lemma 2.5 to the singular context, namely, for any nef $(1,1)$ form θ with $dd^c \theta^{\ell} = 0$ for all $1 \leq \ell \leq k$, then

$$\hat{v}_-(V, \theta|_V) = \int_V \theta^k.$$

3. NULL LOCI AND NAKAI-MOISHEZON CRITERIONS

3.1. Extension theorem. We define the null locus of a big form, following the algebraic version in [Nak00].

Definition 3.1 (Null locus). Let θ be a big $(1,1)$ form. We define the *null locus* of θ , denoted by $\text{Null}(\theta)$, to be the set

$$\text{Null}(\theta) = \bigcup_{\theta|_V \text{ not big}} V,$$

where the union is taken over all purely positive-dimensional irreducible analytic subvarieties $V \subset X$.

We describe the null locus with the following example.

Example 3.2. Let Y be a compact complex manifold of dimension n equipped with a positive (1,1) form ω_Y . Let $\pi : X \rightarrow Y$ be a blow-up of a point $y \in Y$ with the exceptional divisor $E = \pi^{-1}(y) \simeq \mathbb{C}P^{n-1}$. The form $\theta = \pi^*\omega_Y$ is semi-positive, so it is nef. It was shown (from e.g. [DP04, Lemma 3.5]) that there exists a smooth Hermitian metric h on the line bundle $\mathcal{O}_X(E)$ with curvature $\Theta_h(E)$ such that $\pi^*\omega_Y - \varepsilon\Theta_h(E)$ is positive definite on X for ε sufficiently small. In particular, let s be a section of $\mathcal{O}_X(E)$ vanishing along E of order 1. Then for any ε small, we have that $\pi^*\omega_Y + \varepsilon dd^c \log |s|_h^2$ is a Hermitian current on X that belongs to the $\partial\bar{\partial}$ -class $[\theta]$. One obviously sees that the latter current has almost analytic singularities precisely along E . Since θ is positive on $X \setminus E$, we also observe that

$$E = \bigcup_{\theta|_V \text{ not big}} V,$$

where the union is taken over all positive dimensional irreducible analytic subvarieties $V \subset X$.

The above example motivates us to prove our main theorem in this section.

Theorem 3.3. *Let X be a compact complex manifold of dimension n . Let θ be a nef and big (1,1) form. Then*

$$E_{nH}(\theta) = \text{Null}(\theta).$$

The key ingredient in the proof of the theorem above is an extension-type result for Hermitian current with almost analytic singularities in a nef and big $\partial\bar{\partial}$ -class (or merely big in the case of surfaces) on complex subvarieties. We mimic the main technical idea of [CT15, Theorem 3.2] in which they used Hironaka's resolutions of singularities and Richberg's gluing technique in order to glue psh functions with analytic singularities.

Theorem 3.4. *Let X be a compact complex manifold of dimension n . Let θ be a smooth real (1,1) form on X , whose $\partial\bar{\partial}$ -class is nef and big. Let $E = V \cup \cup_i Y_i$ be an analytic closed subset of X where V and Y_i are irreducible components of E , and V is a compact complex subvariety of purely positive dimension. Let $S = \theta + dd^c\psi$ be a Hermitian current with almost analytic singularities precisely along E . Let $T = \theta|_V + dd^c\varphi$ be a Hermitian current in the class $[\theta|_V]$ on V with almost analytic singularities. Then there exists a Hermitian current with almost analytic singularities \tilde{T} on X in the $\partial\bar{\partial}$ -class $[\theta]$ such that \tilde{T} is smooth in a neighborhood of the generic point of V .*

We have the following lemma, which is crucial in Richberg's gluing procedure.

Lemma 3.5. *Let $M \subset \mathbb{C}^N$ be a complex manifold equipped with a positive definite (1,1) form ω and let $E \subset M$ be an analytic subvariety. Assume θ is a continuous (1,1) form. Let $W'_j \Subset W_j \subset M$ be open subsets for $j = 1, 2$. Let $\varphi_j \in \text{PSH}(W_j, \theta)$ such that $\varphi_1|_{E \cap W_1 \cap W_2} = \varphi_2|_{E \cap W_1 \cap W_2}$ and $\varphi_1 - \varphi_2$ is continuous on $W_1 \cap W_2$. Then for any $\varepsilon > 0$, there exists an open neighborhood W of E and $\varphi \in \text{PSH}(M, \theta + \varepsilon\omega)$ such that*

- (1) $E \cap (\overline{W'_1} \cup \overline{W'_2}) \subset E \cap W$, $W \Subset W_1 \cup W_2$,
- (2) $\varphi|_{E \cap W \cap W_j} = \varphi_j|_{E \cap W \cap W_j}$, for $j = 1, 2$,
- (3) $\varphi|_{W \setminus \overline{W_2}} = \varphi_1|_{W \setminus \overline{W_2}}$, and $\varphi|_{W \setminus \overline{W_1}} = \varphi_2|_{W \setminus \overline{W_1}}$.

Moreover, if φ_j (for $j = 1, 2$) has almost analytic singularities of the same type, one also obtains φ to have almost analytic singularities.

Proof. The proof is quite similar to that of [Ric68, Satz 3.1], we provide the proof for reader's convenience. We consider two open sets U_j in M for $j = 1, 2$ such that

$W'_j \Subset U_j \Subset W_j$. We set $K := E \cap \overline{W'_1} \cup \overline{W'_2} \cap \overline{U_1} \cap \overline{U_2}$ and $K_j := K \cap \partial U_j$. We see that K , K_1 , and K_2 are compact subsets of $W_1 \cap W_2$. Since $K_1 \cap K_2 = \emptyset$, we can pick γ_1 a smooth nonnegative cutoff function being identically 1 in a neighborhood of K_2 in M and γ_2 a smooth nonnegative cutoff function being identically 1 in a neighborhood of K_1 in M such that the supports of γ_1 and γ_2 are disjoint. Next, we choose $\eta > 0$ so small that $\theta + \text{dd}^c(\varphi_j - \eta\gamma_j) \geq -\varepsilon\omega$. We define

$$\varphi_0 := \max(\varphi_1 - \eta\gamma_1, \varphi_2 - \eta\gamma_2).$$

Since $K \subset E \cap W_1 \cap W_2$, $\varphi_1|_{E \cap W_1 \cap W_2} = \varphi_2|_{E \cap W_1 \cap W_2}$, and $\varphi_1 - \varphi_2$ is continuous on $W_1 \cap W_2$ there exists an open neighborhood of K_2 such that $\varphi_1 - \varphi_2 < \eta$ there, hence

$$\varphi_1 - \eta\gamma_1 = \varphi_1 - \eta < \varphi_2 = \varphi_2 - \eta\gamma_2.$$

Similarly, there exists an open neighborhood of K_1 on which $\varphi_2 - \eta\gamma_2 < \varphi_1 - \eta\gamma_1$. Therefore, there exists an open neighborhood $W_0 \Subset W_1 \cap W_2$ of K such that

$$\varphi_0|_{W_0 \setminus \overline{W_2}} = \varphi_1|_{W_0 \setminus \overline{W_2}}, \quad \varphi_0|_{W_0 \setminus \overline{W_1}} = \varphi_2|_{W_0 \setminus \overline{W_1}}.$$

From the construction of W_0 and φ_0 , we can define W and φ_0 as follows. We set

$$W := W_0 \cup (U_1 \setminus \overline{U_2}) \cup (U_2 \setminus \overline{U_1})$$

which is a neighborhood of K in $W_1 \cap W_2$. We define the function φ on W such that

$$\varphi|_{W_0} = \varphi_0|_{W_0}, \quad \varphi|_{U_1 \setminus \overline{U_2}} = \varphi_1|_{U_1 \setminus \overline{U_2}}, \quad \varphi|_{U_2 \setminus \overline{U_1}} = \varphi_2|_{U_2 \setminus \overline{U_1}}.$$

The last statement follows from the construction above. \square

Proof of Theorem 3.4. The proof is quite close to that of [CT15, Theorem 3.2], which does not seem to require the closeness of the reference form θ . For the sake of completeness, we include the proof here.

We divide the proof into several steps, which we briefly organize as follows. First, we lift our problem to a modification \tilde{X} of X and \tilde{V} of V ; precisely, we use the resolution of singularities, allowing us to reduce the singularities of T to a simple normal crossing divisor on the strict transform \tilde{V} . Next, we construct local extensions on a finite covering of \tilde{V} in \tilde{X} . We then use the technique of Richberg to glue the local potentials to construct a local extension of Hermitian current in a neighborhood of \tilde{V} . The crucial point is that local potentials have the same singularity type on any non-empty intersection. Finally, after possibly reducing the Lelong numbers of the global Hermitian current S , we can glue the Hermitian current on the neighborhood to S to obtain the desired one.

Step 1: Lifting to the desingularization. Let $\mathcal{I}_T \subset \mathcal{O}_V$ be the coherent sheaf of ideals defining the singularities of T on V . It follows from Hironaka's theorem that we can find a principization of the sheaf of ideals \mathcal{I}_T (cf. [Hir77, Wło09]), that is, a finite sequence

$$p' : V' = V_r \xrightarrow{p_r} V_{r-1} \rightarrow \cdots \rightarrow V_1 \xrightarrow{p_1} V_0 = V$$

of blow-ups p_i with smooth centers $C_{i-1} \subset V_{i-1}$. Next, regarding the first center C_0 as a submanifold of X , we blow it up inside X and iterate r times to obtain a sequence

$$\pi : \tilde{X} = X_r \xrightarrow{\pi_r} X_{r-1} \xrightarrow{\pi_{r-1}} \cdots \xrightarrow{\pi_2} X_1 \xrightarrow{\pi_1} X_0 = X$$

of blow-ups $\pi_i : X_i \rightarrow X_{i-1}$ with smooth centers $C_{i-1} \subset X_{i-1}$. We observe that if \tilde{V} is the strict transform of V under π , then $\tilde{V} \rightarrow V'$ is a proper finite bimeromorphic morphism, so it is indeed an isomorphism since V' is smooth (due to Zariski's main theorem). We see, moreover, that the exceptional divisor E_i of π_i has simple

normal crossings with \tilde{V}_i and $E_i \cap \tilde{V}_i$ coincides with the exceptional divisor of p_i . Finally, on \tilde{X} , we get a divisor $D = \sum D_\ell$ which has simple normal crossings with the strict transform \tilde{V} of V via birational morphism π and

$$(\pi|_{\tilde{V}})^* \mathcal{I}_T = \mathcal{O}_{\tilde{V}} \left(- \sum_{\ell} a_{\ell} D_{\ell} |_{\tilde{V}} \right), \quad a_{\ell} > 0.$$

Note that since the analytic subvariety cutting out by \mathcal{I}_T has codimension at least 2 in X , it follows that all divisors D_{ℓ} are π -exceptional divisors.

By assumptions, we find a $\varepsilon > 0$ so small that

$$T = \theta|_V + \text{dd}^c \varphi \geq 3\varepsilon \omega|_V$$

in the sense of currents on V , where ω denotes a Hermitian metric on X . It follows from [DP04, Lemma 3.5] that there is a small $\delta > 0$ such that

$$\pi^* T + \delta \text{dd}^c \log |s_D|_{h_D}^2 \geq 3\tilde{\omega}|_{\tilde{V}}$$

for some Hermitian metric $\tilde{\omega}$ on \tilde{X} , where s_D is a holomorphic section defining $D = \sum_{\ell} D_{\ell}$ and h_D is a smooth Hermitian metric on the line bundle $\mathcal{O}_{\tilde{X}}(D)$. We set $\tilde{\varphi} := \varphi \circ \pi + \delta \log |s_D|_{h_D}^2$, defining a function on \tilde{V} such that

$$\pi^* \theta|_{\tilde{V}} + \text{dd}^c(\varphi \circ \pi) + \text{dd}^c(\delta \log |s_D|_{h_D}^2)|_{\tilde{V}} \geq 3\varepsilon \tilde{\omega}|_{\tilde{V}}.$$

In the next step, we will explain how to extend $\tilde{\varphi}$ to a small neighborhood of \tilde{V} .

Step 2: Gluing technique of Richberg. Since \tilde{V} is a closed submanifold of \tilde{X} and $\sum_{\ell} D_{\ell}|_{\tilde{V}}$ has simple normal crossing support, we can find a finite open covering $\{W_j\}_{j=1}^N$ of \tilde{V} such that on each local chart W_j there exist local coordinates $z_j = (z_j^1, \dots, z_j^n)$ with $\tilde{V} \cap W_j = \{z_j^1 = \dots = z_j^{n-k} = 0\}$, where $k = \dim \tilde{V}$, and with $\cup_{\ell} D_{\ell} \cap W_j = \{z_j^{i_1} \dots z_j^{i_p} = 0\}$, $n - k + 1 \leq i_1, \dots, i_p \leq n$. We denote by $z'_j = (z_j^1, \dots, z_j^{n-k})$ and $z''_j = (z_j^{n-k+1}, \dots, z_j^n)$. We define a function φ_j on each W_j by

$$\varphi_j(z'_j, z''_j) = (\varphi \circ \pi)(0, z''_j) + A|z'_j|^2 + \delta \log |s_D|_{h_D}^2$$

for some $A > 0$ being chosen so sufficiently large that

$$\pi^* \theta + \text{dd}^c \varphi_j \geq 2\varepsilon \tilde{\omega} \text{ on } W_j$$

holds for all j (we can shrink the W'_j 's if necessary). Since φ is smooth at the generic point of V , one sees immediately that all functions φ_j are smooth in the neighborhood of the generic point of $\tilde{V} \cap W_j$.

We are going to apply Lemma 3.5 to glue the functions φ_j together to obtain a function φ' defined in a neighborhood W of \tilde{V} in \tilde{X} such that $\varphi'|_{\tilde{V}} = \tilde{\varphi}$ and

$$\pi^* \theta + \text{dd}^c \varphi' \geq \varepsilon \tilde{\omega}.$$

This could be done if $\varphi_i - \varphi_j$ would be continuous on any overlap $W_i \cap W_j$, $\forall 1 \leq i, j \leq N$. Assuming this, we finish this step by induction of the number of open sets $\{W_j\}_{1 \leq j \leq N}$ covering \tilde{V} . We take open sets W'_j in \tilde{X} such that $W'_j \Subset W_j$ and $\tilde{V} \subset \cup_{j=1}^N W'_j$. If $N = 1$ we pick $W = W'_1$ and $\varphi := \varphi_1$. Otherwise, we start with two open sets W_1 and W_2 in the covering such that $W_1 \cap W_2 \cap \tilde{V} \neq \emptyset$. From Lemma 3.5, we can find a neighborhood $W_{1,2}$ of $(W'_1 \cup W'_2) \cap \tilde{V}$ in \tilde{X} and define a function $\varphi_{1,2}$ on $W_{1,2}$ such that

$$\varphi|_{\tilde{V} \cap W_{1,2} \cap W_j} = \varphi_j|_{\tilde{V} \cap W_{1,2} \cap W_j}, \quad j = 1, 2$$

and $\pi^*\theta + \text{dd}^c\varphi_{1,2} \geq 2\varepsilon(1 - 2^{-N})\tilde{\omega}$. We apply the inductive assumption with $\tilde{V} \subset W_{1,2} \cup \bigcup_{j=3}^n W_j$ and functions $\varphi_{1,2}, \varphi_j$ to procedure an open neighborhood W of \tilde{V} in \tilde{X} and a function φ' defined on W with having almost analytic singularities, which satisfies $\varphi'|_{\tilde{V}} = \tilde{\varphi}$ and $\pi^*\theta + \text{dd}^c\varphi' \geq \varepsilon\tilde{\omega}$ holds on W .

It remains to show that for any two open sets W_1, W_2 , with $W_1 \cap W_2 \cap \tilde{V} \neq \emptyset$, the difference $\varphi_1 - \varphi_2$ is continuous (smooth) on $W_1 \cap W_2$. We are done if none of D_ℓ intersects $W_1 \cap W_2$. Otherwise, up to relabelling local coordinates, we can write $D_\ell \cap W_1 = \{z_1^\ell = 0\}$ and $D_\ell \cap W_2 = \{z_2^\ell = 0\}$. We then see that on $W_1 \cap W_2$, $z_2^\ell = f^\ell \cdot z_1^\ell$ for some non-vanishing holomorphic function f^ℓ .

It follows from the construction that φ_j has almost analytic singularities along $\cup_\ell D_\ell \cap W_j$, for $j = 1, 2$. The sheaf of ideals defining the singularities of φ_j corresponds to an effective divisor $\sum_\ell a_j^\ell D_\ell$. We have on each W_j ,

$$\varphi_j = c_j \left(\sum_{r=1}^p a_j^{i_r} \log |z_j^{i_r}|^2 \right) + h_j$$

for some positive constants $c_j, a_j^{i_r}$, for some smooth functions h_j on W_j , and $\cup_\ell D_\ell \cap W_j = \{\prod_{r=1}^p |z_j^{i_r}|^{2a_j^{i_r}} = 0\}$. We remark that the term $\delta \log |s_D|_{h_D}^2$ is absorbed in the first one of the expression. Since $\varphi_1 = \varphi_2$ on $\tilde{V} \cap W_1 \cap W_2$ we must have $c_1 = c_2$ and $a_1^{i_r} = a_2^{i_r}$ there. The relation $z_2^\ell = f^\ell \cdot z_1^\ell$ implies that $\varphi_1 - \varphi_2$ is smooth on $W_1 \cap W_2$. Furthermore, the function $\max(\varphi_1, \varphi_2)$ defined on $W_1 \cap W_2$ has almost analytic singularities of the same type as φ_1 and φ_2 , and coincide with $\tilde{\varphi}$ on $\tilde{V} \cap W_1 \cap W_2$.

Step 3: Reducing Lelong numbers. From Step 2, we obtain a $\pi^*\theta$ -psh function φ' defined on W (a neighborhood of \tilde{V} in \tilde{X}) which has almost analytic singularities and $\pi^*\theta + \text{dd}^c\varphi' \geq \varepsilon\tilde{\omega}$ there. One moreover has that $\varphi' \equiv -\infty$ on all the components D_ℓ intersecting W , and φ' has the strictly positive generic Lelong numbers along each such component. Let \tilde{Y}_i be the strict transform of Y_i in \tilde{X} . Up to shrinking W slightly, we may assume that φ' is defined on an open neighborhood \tilde{W} of W . By [DP04, Lemma 2.1, Theorem 3.2], there exists a globally defined $C_0\tilde{\omega}$ -psh function ψ on \tilde{X} for some $C_0 > 0$, with almost analytic singularities along $\cup_i \tilde{Y}_i$. We then define $\varphi'' := \varphi' + \frac{\varepsilon}{2C_0}\psi$, which has almost analytic singularities along $(\cup_\ell D_\ell \cup \cup_i \tilde{Y}_i) \cap \tilde{W}$, and satisfies $\pi^*\theta + \text{dd}^c\varphi'' \geq \frac{\varepsilon}{2}\tilde{\omega}$ on \tilde{W} . Note that the generic Lelong number of φ'' along $(\cup_\ell D_\ell \cup \cup_i \tilde{Y}_i) \cap \tilde{W}$ is always strictly positive. We explain below how to reduce its generic Lelong number, for which we are able to extend φ'' in the whole \tilde{X} .

We work now with the function ψ in the assumption. One sees immediately that the function $\psi \circ \pi$ has singularities along $\cup_\ell D_\ell \cup \cup_i \tilde{Y}_i \cup \tilde{V}$. We now set $\tilde{\psi} := \psi \circ \pi + \delta \log |s_D|_{h_D}^2$ where s_D and h_D are defined in Step 1, $\delta > 0$ small enough. By [DP04, Lemma 3.5] again, we have

$$\pi^*\theta + \text{dd}^c\tilde{\psi} \geq \gamma\tilde{\omega}$$

for some small constant $\gamma > 0$. Since the form $\pi^*\theta$ is nef, for any small $\lambda > 0$ there exists a smooth function ρ_λ such that $\pi^*\theta + \text{dd}^c\rho_\lambda \geq -\lambda\pi^*\omega \geq -\lambda\tilde{\omega}$ on \tilde{X} . Up to adding a positive constant, we can assume that $\rho_\lambda > 0$. Therefore, we have

$$\pi^*\theta + \text{dd}^c(\lambda\tilde{\psi} + (1-\lambda)\rho_\lambda) \geq \lambda^2\gamma\tilde{\omega}.$$

Let us stress that this is the only place where the nefness of θ is used.

Fix any point $x \in (\cup_\ell D_\ell \cup \cup_i \tilde{Y}_i) \cap \partial W$, we choose a local chart U centered at x on which φ'' is also defined, $U \cap \tilde{V} = \emptyset$, with local coordinates (z^1, \dots, z^n) . Thanks to the fundamental theorem of Hironaka [Hir77], there is a principalization $q: \tilde{U} \rightarrow U$ of the sheaf of ideals of the singularities of φ'' and $\tilde{\psi}$ on U such that this sheaf of a simple normal crossing divisor. Namely, on \tilde{U} , we have

$$\begin{aligned} q^* \varphi'' &= c \left(\sum_p a^{i_p} \log |z^{i_p}|^2 \right) + g \\ q^* \tilde{\psi} &= d \left(\sum_p b^{i_p} \log |z^{i_p}|^2 \right) + h \end{aligned}$$

where c, d, a^{i_p}, b^{i_p} are positive constants and g, h are smooth functions on \tilde{U} . Therefore, we can choose $\lambda > 0$ so small that $\lambda \tilde{\psi} + (1 - \lambda) \rho_\lambda > \varphi''$ on a possibly smaller open neighborhood of x . Since x was taken arbitrarily, we can find an open neighborhood of $(\cup_\ell D_\ell \cup \cup_i \tilde{Y}_i) \cap \partial W$ so that $\lambda \tilde{\psi} + (1 - \lambda) \rho_\lambda > \varphi''$ there.

Step 4: Completing the proof. From the previous step, there exist an open neighborhood U of $\cup_\ell D_\ell \cup \cup_i \tilde{Y}_i$ and $\lambda \in]0, 1[$ so small such that $\lambda \tilde{\psi} + (1 - \lambda) \rho_\lambda > \varphi''$ on a neighborhood of $\partial W \cap U$. Since on $\partial W \setminus U$ we can find a constant $B > 0$ so large that

$$\lambda \tilde{\psi} + (1 - \lambda) \rho_\lambda > \varphi'' - B$$

hence $\lambda \tilde{\psi} + (1 - \lambda) \rho_\lambda > \varphi'' - B$ in a neighborhood of ∂W . Therefore, we can define a function Φ on \tilde{X} by

$$\Phi = \begin{cases} \max(\varphi'', \lambda \tilde{\psi} + (1 - \lambda) \rho_\lambda + B) & \text{on } \tilde{W} \\ \lambda \tilde{\psi} + (1 - \lambda) \rho_\lambda + B & \text{on } \tilde{X} \setminus \tilde{W}. \end{cases}$$

One easily sees that $T' = \pi^* \theta + \text{dd}^c \Phi \geq \varepsilon' \tilde{\omega}$ for some $0 < \varepsilon' < \min(\varepsilon, \lambda \gamma)$. We easily see that $\Phi = \varphi''$ in an open neighborhood of the generic point of \tilde{V} because $\tilde{\psi}|_{\tilde{V}} \equiv -\infty$. Therefore, $\Phi|_{\tilde{V}}$ is smooth in an open neighborhood of the generic point of \tilde{V} . Pushing forward T' on X , we obtain a Hermitian current $\tilde{T} = \pi_* T'$ in the $\partial\bar{\partial}$ -class $[\theta]$ which is an extension of T such that the restriction $\tilde{T}|_V$ is smooth in an open neighborhood of the generic point of V . The poof is complete. \square

Proof of Theorem 3.3. We first show that $\text{Null}(\theta) \subset E_{\text{nH}}(\theta)$. Let $x \in X$ and V an irreducible analytic subavriety of X of positive dimension such that $\theta|_V$ is not big. Suppose by contradiction that x does not belong to $E_{\text{nH}}(\theta)$, so there is a Hermitian current $T \in [\theta]$ with almost analytic singularities and smooth in an open neighborhood of x . We write $T = \theta + \text{dd}^c \varphi$ for some θ -psh function φ . However, $\theta|_V + \text{dd}^c \varphi|_V$ defines a Hermitian current with almost analytic singularities on V because $\varphi|_V \not\equiv -\infty$, a contradiction.

For the inverse implication $\text{Null}(\theta) \supset E_{\text{nH}}(\theta)$, suppose by contradiction that there exists a point $x \in E_{\text{nH}}(\theta) \setminus \text{Null}(\theta)$. Let V be an irreducible component of the closed analytic set $E_{\text{nH}}(\theta)$ containing x . Thanks to Lemma 2.16, the pure dimension of V is strictly positive.

Assume first that V is smooth. Since $\theta|_V$ is big, we can find a Hermitian current $T = \theta|_V + \text{dd}^c \varphi$ with almost analytic singularities (Theorm 2.11). We decompose

$$E_{\text{nH}}(\theta) = V \cup \bigcup_{i \in I} Y_i$$

for disjoint irreducible components V, Y_i of $E_{\text{nH}}(\theta)$ which are all purely positive dimensional as follows from Lemma 2.16. We can thus apply Theorem 3.4 with

$E = E_{\text{nH}}(\theta)$, with the current $S = \theta + \text{dd}^c \psi$ in Theorem 2.14 to obtain a Hermitian current $\tilde{T} = \theta + \text{dd}^c \Phi$ in the $\partial\bar{\partial}$ -class $[\theta]$ on X satisfying that $\tilde{T}|_V = T$ and T is smooth in a neighborhood of the generic point of V . In particular, we must have $V \not\subset \text{Sing}(\tilde{T})$, which is a contradiction since $V \subset E_{\text{nH}}(\theta) \subset \text{Sing}(\tilde{T})$.

If V is singular, we take an embedded desingularization of V , that is, $\pi : \tilde{X} \rightarrow X$, a finite composition of blow-ups with smooth centers. The strict transform \tilde{V} is smooth and is in simple normal crossings with the exceptional divisor. Since π is bimeromorphic, $\pi^*\theta$ is nef and big on \tilde{X} , has the same properties on \tilde{V} . Then there exists a Hermitian current $S \in [\pi^*\theta]$ with almost analytic singularities along an analytic subset $E_{\text{nH}}(\pi^*\theta)$. Thanks to [Tos18b, Proposition 2.5],

$$\tilde{V} \subset E_{\text{nH}}(\pi^*\theta) = \text{Exc}(\pi) \cup \pi^{-1}(E_{\text{nH}}(\theta)).$$

We can apply Theorem 3.4 with S and a Hermitian current $T = \pi^*\theta|_{\tilde{V}} + \text{dd}^c \varphi$ to obtain a Hermitian current $\tilde{T} = \pi^*\theta + \text{dd}^c \tilde{\Phi}$ in the $\partial\bar{\partial}$ -class $[\pi^*\theta]$ such that $\tilde{T}|_{\tilde{V}}$ is smooth in a neighborhood of the generic point of \tilde{V} . Hence, the direct image $\pi_*\tilde{T}$ is smooth near the generic point of V , which is again a contradiction. \square

The following example tells us that the nef condition in Theorem 3.3 is essential even in the algebraic version of Nakamaye's theorem; we would like to thank V. Tosatti for pointing out this example.

Example 3.6. Let $\pi_1 : X_1 \rightarrow \mathbb{C}\mathbb{P}^2$ be a blow-up at one point, with exceptional divisor F_1 . Let $\pi_2 : X \rightarrow X_1$ be the blow-up of a point $y \in F_1$ with exceptional divisor F_2 . The exceptional divisor E of $\pi = \pi_2 \circ \pi_1$ has two components $E = E_1 + E_2$ where E_1 is the strict transform of F_1 under π_2 while $E_2 = F_2$. We have that $E_1^2 = -2$, $E_2^2 = -1$ and $E_1 \cdot E_2 = 1$. We take $L = \pi^*H + E_1$ where H is a hyperplane bundle on $\mathbb{C}\mathbb{P}^2$. It is clear that L is big, and its positive part in the divisorial Zariski decomposition is π^*H , which is nef. Nakamaye's theorem yields

$$E_{\text{nH}}(L) = E_{\text{nH}}(\pi^*H) = E_1 \cup E_2.$$

However, since the restriction $L|_{E_2}$ is a line bundle on a (rational) curve with degree $E_1 \cdot E_2 = 1$, it is ample. This implies $E_{\text{nH}}(L) \neq \text{Null}(L)$.

3.2. Proof of Theorem A. Under some extra assumptions on the manifold, we can determine the null locus as follows.

Theorem 3.7. *Let X be a n -dimensional compact complex manifold equipped with a Hermitian metric ω_X . Assume that ω_X satisfies the volume condition (H_X) . Let θ be a nef $(1,1)$ form. Then θ is big if and only if $\hat{v}_-(X, \theta) > 0$ and moreover*

$$(3.1) \quad E_{\text{nH}}(\theta) = \text{Null}(\theta) = \bigcup_{\hat{v}_-(V, \theta|_V)=0} V$$

where the union is taken over all purely positive-dimensional irreducible analytic subvarieties of $V \subset X$.

Proof. We denote by N the set of the right-hand side of (3.1). The first statement is known in [GL22, Theorem 4.6]. It suffices to show that $E_{\text{nH}}(\theta) = N$.

We first show that $N \subset E_{\text{nH}}(\theta)$. Let $x \in X$ and V an irreducible analytic subvariety of X of dimension $k > 0$ such that $\hat{v}_-(V, \theta|_V) = 0$. Suppose by contradiction that x does not belong to $E_{\text{nH}}(\theta)$, so there is a Hermitian current $T \in [\theta]$ with almost analytic singularities and smooth in an open neighborhood of x . We write $T = \theta + \text{dd}^c \varphi$ for some θ -psh function φ . Set $\psi = \varphi|_V$.

Suppose first that V is smooth. Assume that $\theta|_V + \text{dd}^c \psi \geq \omega_V$ for some Hermitian metric ω_V , and $\sup_V \psi = 0$. Fix $\varepsilon > 0$ and $u \in \text{PSH}(V, \theta|_V + \varepsilon\omega_V) \cap \mathcal{C}^\infty(V)$. Set $v := P_{\omega_V}(u - \psi)$. We note that $u - \psi$ is merely well-defined outside a pluripolar

set and $v + \psi \leq u$ almost everywhere, hence everywhere. It follows from [GZ17, Theorem 9.17] that v is bounded from above. By the classical pluripotential theory (see e.g. [GL22, Theorem B]), v is a bounded ω -psh function and $(\omega_V + \text{dd}^c v)^k$ is supported on the contact set

$$D = \{v = u - \psi\} \subset \{\psi > -\infty\}$$

where the product $(\cdot)^k$ is understood as the Monge–Ampère product in the sense of Bedford–Taylor; cf. [GL22, Theorem 2.3]. By the maximum principle (Lemma 2.7), we have

$$\mathbf{1}_{D \cap \{\psi > -\infty\}} (\theta|_V + \varepsilon \omega_V + \text{dd}^c(v + \psi))^n \leq \mathbf{1}_{D \cap \{\psi > -\infty\}} (\theta|_V + \varepsilon \omega_V + \text{dd}^c u)^n.$$

Hence,

$$\int_V (\omega_V + \text{dd}^c v)^n = \int_D (\omega_V + \text{dd}^c v)^n \leq \int_V (\theta|_V + \varepsilon \omega_V + \text{dd}^c u)^n$$

since $D \cap \{\psi > -\infty\} = D$. Therefore,

$$0 < v_-(V, \omega_V) \leq v_-(V, \theta|_V + \varepsilon \omega_V)$$

by assumption. Taking the infimum for $\varepsilon > 0$ gives a contradiction.

If V is singular, we consider $\pi : X' \rightarrow X$ an embedded resolution of singularities obtained by a finite composition of blow-ups with smooth centers and assume V' is the strict transform of V . We set $\psi' = (\varphi \circ \pi)|_{V'} + \delta \log |s|_h^2$ where s is a section defining the exceptional divisor $\text{Exc}(\pi)$, and $\delta > 0$ small enough. Then $\pi^* \theta|_{V'} + \text{dd}^c \psi' \geq \omega_{V'}$ for some Hermitian metric $\omega_{V'}$. We apply the argument above with ψ' and $\omega_{V'}$ to obtain

$$0 < v_-(V', \omega_{V'}) \leq v_-(V', \pi^* \theta + \varepsilon \omega_{V'})$$

a contradiction.

Conversely, suppose by contradiction that there exists a point $x \in E_{\text{nH}}(\theta) \setminus N$. Let V be an irreducible component of the closed analytic set $E_{\text{nH}}(\theta)$ containing x . Thanks to Lemma 2.16, we have that the pure dimension of V is strictly positive, denoted by k .

Assume first that V is smooth. We see that $\widehat{v}_-(V, \theta|_V) > 0$ because $x \notin N$. Then, by the first statement, we can find a Hermitian current $T = \theta|_V + \text{dd}^c \varphi$ with almost analytic singularities (thanks to Demailly’s regularization theorem). We decompose

$$E_{\text{nH}}(\theta) = V \cup \bigcup_i Y_i$$

for disjoint irreducible components of $E_{\text{nH}}(\theta)$ which are all purely positive dimension by Lemma 2.16. We apply Theorem 3.4 with $E = E_{\text{nH}}(\theta)$, with the current $S = \theta + \text{dd}^c \psi$ in Theorem 2.14 to obtain a Hermitian current $\widetilde{T} = \theta + \text{dd}^c \Phi$ in the $\partial\bar{\partial}$ -class $[\theta]$ on X satisfying that $\widetilde{T}|_V = T$ and T is smooth in a neighborhood of the generic point of V . In particular, we must have $V \not\subset \text{Sing}(\widetilde{T})$, which is a contradiction because

$$V \subset E_{\text{nH}}(\theta) \subset \text{Sing}(\widetilde{T}).$$

If V is not smooth, we argue the same as in the proof of Theorem 3.3 to obtain again a contradiction; see also [CT15, page 1180]. \square

We end this section with some examples of the manifold X equipped with a Hermitian metric ω_X satisfying the volume assumption (H_X) .

Example 3.8. Obviously, if X is Kähler or belongs to the Fujiki class \mathcal{C} , then it always admits a metric ω_X satisfying the condition (H_X) .

When $\dim_{\mathbb{C}} X = 2$ or 3 and ω_X is pluriclosed, i.e., $\text{dd}^c \omega_X = 0$, it has been shown in [AGL23, Theorem 3.3] that the condition (H_X) is satisfied.

If ω_X satisfies the curvature condition given by Guan–Li, namely

$$\text{dd}^c \omega_X = 0, \quad d\omega_X \wedge d^c \omega_X = 0$$

which is equivalent to $\text{dd}^c \omega_X^k = 0$ for $k = 1, \dots, n-1$, then we have that

$$\int_V (\omega_X + \text{dd}^c \varphi)^{\dim V} = \int_V \omega_X^{\dim V}$$

for any compact submanifold (without boundary) of X . We can precisely choose X to be the product of a complex surface and a Kähler manifold. Namely, $X = M \times N$ where M is a complex surface and N is Kähler, and $\omega_X = p_1^* \omega_M + p_2^* \omega_N$ where ω_M is a dd^c -closed metric (it always exists thanks to Gauduchon’s theorem) and ω_N is Kähler metric.

Example 3.9. It was shown in [GL22, Theorem 3.7] that the condition (H_X) is bimeromorphic invariant. Therefore, we can find various compact complex manifolds with such conditions. For instance, we assume that a compact complex manifold X admits a Hermitian metric ω_X satisfying the condition (H_X) and $\pi : X' \rightarrow X$ a finite composition of blow-ups of X with smooth centers, then X' carries a Hermitian metric $\omega_{X'}$ with the same property.

3.3. Nakai–Moishezon’s criterion on some compact Hermitian manifolds. As a direct application of Theorem A, we discuss the following Nakai–Moishezon criterion for some compact non-Kähler manifolds.

Theorem 3.10. *Let X be a n -dimensional compact complex manifold equipped with a Hermitian metric ω_X . Assume that ω_X satisfies the volume condition (H_X) . Let θ be a nef $(1,1)$ form. Then the $\partial\bar{\partial}$ -class $[\theta]$ is Hermitian if and only if $\widehat{v}_-(V, \theta|_V) > 0$ for all irreducible analytic subvarieties $V \subset X$.*

Proof. If the $\partial\bar{\partial}$ class $[\theta]$ is Hermitian we can find a Hermitian metric ω such that $\omega = \theta + \text{dd}^c \varphi$ for $\varphi \in \mathcal{C}^\infty(X)$, and we have $\omega \geq \delta \omega_X$ for some $\delta > 0$. If $V \subset X$ is a closed smooth manifold, then by [GL22, Lemma 4.4],

$$\widehat{v}_-(V, \theta|_V) = v_-(V, \omega|_V) \geq \delta^n v_-(V, \omega_X|_V) > 0.$$

If V is not smooth, then thanks to Hironaka’s theorem [Hir77, Wlo09], we can take $\pi : X' \rightarrow X$ be an embedded resolution of singularities of V , obtained by composition of blow up with smooth centers. Let V' be the strict transform of V , which is now a closed smooth submanifold of X' . Thanks to [GL22, Theorem 3.7] the condition (H_X) is preserved. That theorem also yields

$$v_-(V', \omega') \geq \delta^n v_-(V', \pi^* \omega_X) > 0.$$

Conversely, with $V = X$ we observe that $\widehat{v}_-(X, \theta) > 0$, hence it follows from Theorem 2.15 that there exist a Hermitian current T in the $\partial\bar{\partial}$ class $[\theta]$ with almost analytic singularities along $E_{\text{nH}}(\theta)$. We apply Theorem 3.7 to get that the non-Hermitian (null) locus

$$E_{\text{nH}}(\theta) = \text{Null}(\theta) = \bigcup_{\widehat{v}_-(V, \theta|_V)=0} V$$

is empty. Therefore, the Hermitian current T is, indeed, a Hermitian metric. \square

Consequently, we can give an alternative proof of the Nakai–Moishezon criterion for compact non-Kähler surfaces [Buc00, Lam99]. We extend this result in higher dimensions, and also show that it holds on singular compact varieties. We are grateful to V. Tosatti for pointing out this to us.

Proof of Theorem B. We first treat the case when X is smooth. Obviously, the metric ω_X satisfies the condition (H_X) . Thanks to Theorem 3.10, it suffices to show that θ is nef. Indeed, if θ were nef, we would have that $\widehat{\nu}_-(V, \theta|_V) = \int_V \theta^{\dim V}$ by (1). We choose a number $t > 0$ so large that the $\partial\bar{\partial}$ -class $[\theta + t\omega]$ is Hermitian, and we denote by t_0 the minimum of such t . It thus remains to prove that $t_0 \leq 0$. Suppose by contradiction that $t_0 > 0$. We see that $[\theta + t_0\omega]$ is automatically nef, but not Hermitian. Since $t_0 > 0$, the conditions (1) and (2) yield

$$\widehat{\nu}_-(X, \theta + t_0\omega_X) = \int_X (\theta + t_0\omega_X)^n > 0$$

By Theorem 2.13, the $\partial\bar{\partial}$ -class $[\theta + t_0\omega_X]$ is big. On the other hand, the condition (3) and Theorem 3.10 imply that the $\partial\bar{\partial}$ -class $[\theta + t_0\omega_X]$ is Hermitian, a contradiction. Therefore, we must have $t_0 \leq 0$ so θ is nef.

If X is singular, thanks to Hironaka’s theorem, we can take $\pi : X' \rightarrow X$ to be an embedded resolution of singularities of X , obtained by finite compositions of blow-ups with smooth centers, with exceptional divisor D . On X' , $\omega_{X'} := \pi^*\omega_X + \text{dd}^c \delta \log |s_D|_h$ is a Hermitian metric for $\delta > 0$ sufficiently small, which satisfies $\text{dd}^c \omega_{X'}^k = 0$ for any $1 \leq k \leq n-1$. We apply the previous argument with $\pi^*\theta$ and $\omega_{X'}$ we can find a smooth function ψ on X' such that $\Theta = \pi^*\theta + \text{dd}^c \psi$ is a Hermitian metric on X' . The direct image $\pi_*\Theta$ is a Hermitian current with singularities along X_{sing} . In order to remove the singularities of $\pi_*\Theta$ we apply the induction technique on dimension used in [CT16, page 94] or [DHP22, pages 19–23] (probably used earlier in [DP04]), so we only give the main ideas. Suppose by induction that the theorem holds for any compact normal analytic variety of pure dimension smaller than n and that for any analytic subset $Y \subset X$, there exists an open neighborhood W_Y of Y and a smooth function ρ_Y on W_Y such that $\theta|_{W_Y} + \text{dd}^c \rho_Y \geq \varepsilon \omega_X|_{W_Y}$ for $\varepsilon > 0$. The latter can be obtained from the same arguments in the proof of Theorem 3.4 or from the aforementioned references. On the desingularization $p : Y' \rightarrow Y$, we can find a Hermitian metric $p^*\theta|_{Y'} + \text{dd}^c \psi_Y$, its push forward is a Hermitian current with singularities along $Z \subset Y$ ($p : Y \setminus p^{-1}(Z) \rightarrow Y \setminus Z$ is biholomorphic). By induction hypothesis, we can find a smooth function ρ_Z and a neighborhood W_Z of Z on which $\theta|_{W_Z} + \text{dd}^c \rho_Z \geq \delta \omega_X|_{W_Z}$ with $\delta > 0$. Hence, using the regularized maximum function; cf. [Dem12, page 43], we can glue $\pi_*\psi_Y$ with ρ_Z to get the desired one. Finally, we apply with $\pi_*\psi$ and $\rho_{X_{\text{sing}}}$ to conclude. \square

4. SINGULAR HERMITIAN METRICS AND ENVELOPES

4.1. Singular Hermitian metrics. This section is devoted to the study of degenerations of Hermitian metrics on compact complex manifolds. We first treat the case of smooth densities, proving the following.

Theorem 4.1. *Let X be a n -dimensional compact complex manifold equipped with a Hermitian metric ω_X . Let θ be a smooth real $(1,1)$ form whose $\partial\bar{\partial}$ -class is nef and big. Let $f \in C^\infty(X, \mathbb{R})$. Then, there exists a constant $c > 0$ and an θ -psh function φ which is smooth on a Zariski open U such that $\sup_X \varphi = 0$ and*

$$(\theta + \text{dd}^c \varphi)^n = c f \omega_X^n$$

on U . More precisely, U can be taken to be the complement of the null locus of θ .

Proof. Since the class $[\theta]$ is nef we have for any $t > 0$, there exists a smooth function u_t such that $\theta + t\omega_X + \text{dd}^c u_t$ is a positive Hermitian form. Set $\theta_t := \theta + t\omega_X$. Since θ is nef, the $\partial\bar{\partial}$ -class $[\theta_t]$ is Hermitian. Hence the main result of [TW10a] ensures that there exists a constant $c_t > 0$ and a smooth θ_t -psh function φ_t normalized by $\sup_X \varphi_t = 0$ such that

$$(\theta_t + \text{dd}^c \varphi_t)^n = c_t f \omega_X^n,$$

where c_t equals $\int_X \omega_t'^n / \int_X f \omega_X^n$ with $\omega_t' := \theta_t + \text{dd}^c \varphi_t$. We observe that ω_t' is the (unique) Hermitian metric in the $\partial\bar{\partial}$ -class $[\theta_t]$ with prescribed Chern–Ricci form. We will show that as $t \rightarrow 0^+$, ω_t' converges to some metric $\omega_0 = \theta + \text{dd}^c \varphi$ on $X \setminus E_{\text{nH}}(\theta)$ in $\mathcal{C}_{\text{loc}}^\infty(X \setminus E_{\text{nH}}(\theta))$. We divide the proof into several steps.

Step 1: \mathcal{C}^0 -estimates. Thanks to Theorem 2.14, we can find a θ -psh function ρ with almost analytic singularities precisely along $E_{\text{nH}}(\theta)$, normalized by $\sup_X \rho = 0$ such that

$$\theta + \text{dd}^c \rho \geq 2\delta_0 \omega_X.$$

In particular, ρ is smooth on the complement of the analytic subset $E_{\text{nH}}(\theta)$ of X . For t sufficiently small, we have

$$\theta_t + \text{dd}^c \rho \geq \delta_0 \omega_X.$$

We set $v_t = P_{\delta_0 \omega_X}(\varphi_t - \rho)$. Since the set $E = \{\rho > -1\}$ is not empty, it is non-pluripolar. We have that the global extremal function $V_{E, \delta_0 \omega_X}$ of E is not identically $+\infty$ (cf. [GZ17, Theorem 9.17]). On E , $v_t \leq \varphi_t - \rho \leq 1$, hence $v_t - 1 \leq V_{E, \delta_0 \omega_X}^*$ on X . Therefore $\sup_X v_t < +\infty$ independent of t . Since $\varphi_t - \rho$ is bounded from below and lower semi-continuous, it follows from [GL22, Theorem 2.3] that v_t is a bounded $(\delta_0 \omega_X)$ -psh function and $(\delta_0 \omega_X + \text{dd}^c v_t)^n$ is concentrated on the contact set

$$D_t := \{v_t = \varphi_t - \rho\} \subset \{\rho > -\infty\}.$$

Note that D_t is not empty and $\sup_X v_t$ is uniformly (in t) bounded from below since $\rho \leq 0$. Since $v_t + \rho \leq \varphi_t$ outside a pluripolar set, hence on X . It follows from the maximum principle (Lemma 2.7) that

$$\mathbf{1}_{D_t}(\theta_t + \text{dd}^c(v_t + \rho))^n \leq \mathbf{1}_{D_t}(\theta_t + \text{dd}^c \varphi_t)^n.$$

To summarize, we have

$$\begin{aligned} (\delta_0 \omega_X + \text{dd}^c v_t)^n &= \mathbf{1}_{D_t}(\delta_0 \omega_X + \text{dd}^c v_t)^n \\ &\leq \mathbf{1}_{D_t}(\theta_t + \text{dd}^c(v_t + \rho))^n \\ &\leq \mathbf{1}_{D_t}(\theta_t + \text{dd}^c \varphi_t)^n \\ &\leq c_t f \omega_X^n. \end{aligned}$$

We show that the constant c_t can be controlled, not being dependent on t . Let G be the Gaduchon factor of ω_X , i.e., $G \in \mathcal{C}^\infty(X)$ such that $\text{dd}^c(e^G \omega_X^{n-1}) = 0$. By the arithmetic-geometric inequality, we have

$$c_t^n f^n e^{\frac{(n-1)G}{n}} \omega_X^n \leq (\theta_t + \text{dd}^c \varphi_t) \wedge e^G \omega_X^{n-1}.$$

Integrating both sides on X we get an upper bound on c_t depending on a lower bound for $\int_X f^n e^{\frac{(n-1)G}{n}} \omega_X^n$ and $\int_X \omega_X^n$.

We have shown that $(\delta_0 \omega_X + \text{dd}^c v_t)^n \leq c_t f \omega_X^n$ with $c_t > 0$ uniformly bounded from above. By Dinew–Kolodziej–Nguyen’s uniform estimate (see also [GL23, Theorem 2.2]), we see that $v_t \geq -C_0$ for a uniform constant $C > 0$ only depending on n, p, δ_0, ω_X , and $\|f\|_p$ with $p > 1$. Here, we have used the fact that $\sup_X v_t$

is uniformly bounded, hence v_t belongs to a compact set of quasi-psh functions. Therefore, we have

$$\varphi_t \geq \rho - C_0.$$

For a lower bound of c_t , by [GL23, Lemma 2.1] we can find a bounded ω_X -psh function u and $c = c(n, \omega_X, \|f\|_p) > 0$, with $p > 1$ such that $(\omega_X + \text{dd}^c u)^n \geq cf\omega_X^n$. We have

$$c_t(\theta_t + \text{dd}^c(\rho + \delta_0 u))^n \geq \delta_0^n c_t (\omega_X + \text{dd}^c u)^n \geq c\delta_0^n (\theta_t + \text{dd}^c \varphi_t)^n$$

hence $c_t \geq c\delta_0^n > 0$ by the domination principle [GL23, Corollary 1.14].

Step 2: Laplacian estimates. In the sequel, we establish uniform bounds on $\Delta_{\omega_X} \varphi_t$ on compact subsets of $X \setminus E_{\text{NH}}(\theta)$. The difficulty in this part is dealing with the torsion term corresponding ω_X . We follow the trick due to Phong–Sturm [PS10]. We use C for positive constants whose values may change from line to line.

We set $u := \varphi - \rho + C_0 + 1$ so that $e^{-u} \in [0, 1]$, and $\omega' := \theta_t + \text{dd}^c \varphi_t$ (we omit the subscript t for simplicity). We denote by g (resp. g') the Riemann metric associated to ω_X (resp. ω'). We also denote by ∇ and T the Chern connection and torsion of g . Consider

$$Q := \log \text{tr}_g g' - Au + e^{-u}$$

with $A > 0$ to be determined hereafter. We are going to show that Q is uniformly bounded from above. If this would be done, then for each t and any compact subset $K \subset\subset X \setminus E_{\text{NH}}(\theta)$ we obtain a uniform bound for $\Delta_{\omega_X} \varphi_t$ on K .

Since $\rho \rightarrow -\infty$ near $E_{\text{NH}}(\theta)$ hence Q attains its maximum at some point $x_0 \in X \setminus E_{\text{NH}}(\theta)$. We assume that $\text{tr}_g g'(x_0) \geq 1$; otherwise we are done. We compute as in [TW15, Sect. 9] at the point x_0 ,

$$(4.1) \quad \begin{aligned} \Delta_{g'} \log \text{tr}_g g' &\geq \frac{2}{(\text{tr}_g g')^2} \text{Re} \left(g'^{\bar{q}k} T_{ik}^i \partial_{\bar{q}} \text{tr}_g g' \right) + \frac{\Delta_g \log c_t f}{\text{tr}_g g'} \\ &\quad - C \text{tr}_{g'} g - C \end{aligned}$$

with $C > 0$ only depending on ω_X and its torsion terms and Riemann curvature. Notice that we used $\text{tr}_g g'(x_0) \geq 1$. We observe that $|\Delta_g \log f| \leq B$ for a uniform constant $B > 0$. At the maximum point x_0 of Q we have $\partial_{\bar{q}} Q = 0$, hence

$$\frac{1}{\text{tr}_g g'} \partial_{\bar{q}} \text{tr}_g g' - (A + e^{-u}) \partial_{\bar{q}} u = 0.$$

The Cauchy–Schwarz inequality yields

$$(4.2) \quad \begin{aligned} \left| \frac{2}{(\text{tr}_g g')^2} \text{Re} \left(g'^{\bar{q}k} T_{ik}^i \partial_{\bar{q}} \text{tr}_g g' \right) \right| &\leq \left| \frac{2}{\text{tr}_g g'} \text{Re} \left((A + e^{-u}) \partial_{\bar{q}} u g'^{\bar{q}k} T_{ik}^i \right) \right| \\ &\leq e^{-u} |\partial u|_{g'}^2 + C(A+1)^2 e^u \frac{\text{tr}_{g'} g}{(\text{tr}_g g')^2} \end{aligned}$$

with $C > 0$ under control. If at x_0 , $(\text{tr}_g g')^2 \leq C(A+1)^2 e^u$ then

$$Q \leq C + \frac{1}{2}u - Au + e^{-u} \leq C + 1$$

we are done. Otherwise, from (4.1) together with (4.2), we have at x_0

$$\begin{aligned} 0 \geq \Delta_{g'} Q &\geq -e^{-u} |\partial u|_{g'}^2 - (B + C + 1) \text{tr}_{g'} g - C \\ &\quad - (A + e^{-u}) \text{tr}_{\omega'} (\omega' - (\theta_t + \text{dd}^c \rho)) + e^{-u} |\partial u|_{g'}^2 \\ &\geq (A\delta_0 - B - C - 1) \text{tr}_{\omega'} \omega_X - (A + 1)n - C \end{aligned}$$

where we used $\theta_t + \text{dd}^c \rho \geq \delta_0 \omega_X$. Choosing $A = (B + C + 2)\delta_0^{-1}$, at x_0 , we obtain

$$\text{tr}_{\omega'} \omega_X \leq C$$

Hence, at x_0

$$(4.3) \quad \mathrm{tr}_{\omega_X} \omega' \leq n \frac{\omega'^n}{\omega_X^n} (\mathrm{tr}_{\omega'} \omega_X)^{n-1} \leq n C^{n-1} c_t f$$

which is uniformly bounded from above since c_t is so. It therefore follows that

$$\mathrm{tr}_{\omega_X} \omega' \leq C e^{-A\rho} \quad \text{on } X.$$

Step 3: Higher order estimates and conclusion. For each compact subset $K \subset\subset X \setminus E_{\mathrm{nH}}(\theta)$, we obtain uniform bounds on $\|\Delta_{\omega_X} \varphi_t\|_{L^\infty(K)}$ as follows from the previous step. We apply a complex version of the Evans–Krylov–Trudinger estimate (see [TW10b, Sect. 4] in the non-Kähler context) and Schauder’s estimates to obtain for any $\alpha \in]0, 1[$, $j \in \mathbb{N}$,

$$\sup_t \|\varphi_t\|_{C^{j,\alpha}(K)} \leq C_{j,\alpha} < +\infty.$$

This guarantees that the family $(\varphi_t)_{t>0}$ is relatively compact in $C_{\mathrm{loc}}^\infty(X \setminus E_{\mathrm{nH}}(\theta))$. Up to exacting and relabelling, we can pass the limit to get

$$c = \lim_{t \rightarrow 0} c_t > 0, \quad \varphi = \lim_{t \rightarrow 0} \varphi_t$$

satisfying that on $X \setminus E_{\mathrm{nH}}(\theta)$, φ is smooth and $(\theta + \mathrm{dd}^c \varphi)^n = c f \omega_X^n$. By Hartogs’ lemma, we obtain $\sup_X \varphi = 0$. \square

The same result still holds when the density f is mildly singular.

Theorem 4.2. *Let θ be a smooth real (1,1) form whose $\partial\bar{\partial}$ -class is nef and big. Assume $f = e^{\psi^+ - \psi^-} \in L^p(X, \omega_X)$, $p > 1$, and f is smooth and positive in the complement of a divisor $X \setminus D$, with ψ^\pm quasi-psh functions on X . Then, there exists a constant $c > 0$ and $\varphi \in \mathrm{PSH}(X, \theta)$ such that $\sup_X \varphi = 0$ and on $X \setminus (D \cup E_{\mathrm{nH}}(\theta))$, φ is smooth and*

$$(\theta + \mathrm{dd}^c \varphi)^n = c f \omega_X^n.$$

Proof. The proof is almost identical to the previous one, so we only give the main idea. Without loss of generality, we assume that $\sup_X \psi^\pm \leq -1$. For a quasi-psh function ψ , the Demailly regularization $\psi_t := \rho_t(\psi)$ of ψ , with $t > 0$ so small, is defined by

$$\rho_t(\psi) := \frac{1}{t^{2n}} \int_{\zeta \in T_x X} \psi(\mathrm{exph}_x(\zeta)) \chi\left(\frac{|\zeta|^2}{t^2}\right) dV_{\omega_X}$$

where χ is a cut-off function normalized by $\int_{\mathbb{R}} \chi(t) = 1$ and exph is the formal holomorphic part of the Taylor expansion of the exponential map of the Chern connection on the tangent space TX associated with ω_X (see [Dem94, (3.1)]). We have that the corresponding regularization ψ_t^\pm is smooth on X and satisfies $\mathrm{dd}^c \psi_t^\pm \geq -A\omega_X$ where $A > 0$ is a uniform constant only depending on the Lelong numbers of ψ^\pm . Up to rescaling, we assume in the sequel that $A = 1$. We observe that $f_t := e^{\psi_t^+ - \psi_t^-}$ converges to f in L^p by the dominated convergence theorem. Up to extracting subsequence, we may assume that $f_t \leq F$ for some $F \in L^p(X, \omega_X)$.

As explained in the previous part, it suffices to obtain uniform estimates and Laplacian estimates of the potential φ_t solution of

$$(\theta + t\omega_X + \mathrm{dd}^c \varphi_t)^n = c_t f_t \omega_X^n, \quad \sup_X \varphi_t = 0.$$

We use the same trick as above with minor modifications. The same argument yields that c_t has a uniform upper bound only depending on ω_X , n and $\int_X f^{\frac{1}{n}} \omega_X^n$. From this we have

$$(\theta_t + \mathrm{dd}^c \varphi_t)^n \leq F \omega_X^n$$

with $F \in L^p(X)$, $p > 1$. It follows that $C_0 \geq \varphi \geq \rho - C_0$ for a constant $C_0 > 0$ depending on n, ω_X, p , and an upper bound for $\|F\|_p$.

For the Laplacian estimate, we consider

$$Q := \log \operatorname{tr}_{\omega_X} \omega' - Au + e^{-u}$$

where $A > 0$ sufficiently large and

$$\omega' := \theta + t\omega_X + \operatorname{dd}^c \varphi_t, \quad u := \varphi_t - \rho - 2\delta\psi^- + C_0 + 1, \quad \delta \in]0, \delta_0[.$$

By the same arguments as in Step 2 of the proof of Theorem 4.1 (see also [GL23, Theorem 4.2]), one obtains

$$\Delta_{\omega_X} \varphi_t \leq Ce^{-A(\rho+2\delta\psi^-)}.$$

The conclusion of the proof is then roughly the same as in the smooth case. \square

4.2. Regularity of envelopes. Our main theorem in this section is the following.

Theorem 4.3. *Let X be a n -dimensional compact complex manifold equipped with a Hermitian metric ω_X . Let θ be a smooth real $(1,1)$ form whose $\partial\bar{\partial}$ -class is nef and big. Let $f \in C^\infty(X, \mathbb{R})$. Then $P_\theta(f) \in \mathcal{C}^{1,\alpha}(X \setminus E_{\text{nH}}(\theta))$ with $\alpha \in]0, 1[$.*

Proof. The proof follows the PDE method of R. Berman [Ber19]. Up to replacing θ with $\theta_f = \theta + \operatorname{dd}^c f$, we may assume that $f = 0$. We consider, for $\beta > 1$ fixed positive numbers, the following complex Monge–Ampère equations for a smooth function

$$(\theta + 2^{-\beta}\omega_X + \operatorname{dd}^c \varphi_\beta)^n = e^{\beta\varphi_\beta} \omega_X^n.$$

Note that the $\partial\bar{\partial}$ -class $[\theta + 2^{-\beta}\omega_X]$ is Hermitian for $\beta > 0$. By the fundamental results of Cherrier [Che87] and Tosatti–Weinkove [TW10a] there exists indeed a unique smooth solution φ_β to the previous equation.

C^0 -estimates. Using the domination principle, we have

$$\left(1 - \frac{1}{\beta}\right) P_\theta(0) + \frac{1}{\beta}\rho - \frac{C \log \beta}{\beta} \leq \varphi_\beta \leq \frac{C}{\beta}$$

with $C > 0$ under control. In particular, φ_β converges to $P_\theta(0)$ in $\mathcal{C}_{\text{loc}}^0(X \setminus E_{\text{nH}}(\theta))$, as $\beta \rightarrow +\infty$.

Indeed, we assume that $\theta + 2^{-\beta}\omega_X \leq A\omega_X$ for a uniform constant $A > 0$. We have

$$(A\omega_X + \operatorname{dd}^c \varphi_\beta) \geq e^{\beta(\varphi - n \log A/\beta)} A^n \omega_X^n.$$

It follows from the domination principle that $\varphi_\beta \leq n \log A/\beta$. Conversely, fixing an θ -psh function $u \geq \rho$, we consider

$$v_\beta := \left(1 - \frac{1}{\beta}\right) u + \frac{1}{\beta}\rho - \frac{n \log(\delta_0^{-1}\beta)}{\beta}$$

where ρ is an θ -psh function such that $\sup_X \rho \leq 0$ and $\theta + \operatorname{dd}^c \rho \geq 2\delta_0\omega_X$ for small $\delta_0 > 0$. We have

$$(\theta + 2^{-\beta}\omega_X + \operatorname{dd}^c v_\beta)^n \geq (\delta_0/\beta)^n \omega_X^n \geq e^{\beta v} \omega_X^n.$$

It follows from the domination principle [GL23, Corollary 1.14] that $\varphi_\beta \geq v_\beta$. Taking the supremum for all candidates $u \geq \rho$, we obtain

$$\varphi_\beta \geq \left(1 - \frac{1}{\beta}\right) P_\theta(0) + \frac{1}{\beta}\rho - \frac{C \log \beta}{\beta}.$$

In particular, since $P_\theta(0)$ dominates all θ -psh function u with $\sup_X u \leq 0$ and $\beta > 1$, there exists a uniform constant $C_0 > 0$ independent of β such that

$$\varphi_\beta \geq \rho - C_0.$$

Laplacien estimates. Fixing $\beta > 1$ and setting $\omega' := \theta + 2^{-\beta}\omega_X + \text{dd}^c\varphi_\beta$, we let g denotes the Riemannian metric associated to ω_X and g' the one associated to ω' . Setting $u := \varphi_\beta - \rho - C_0 + 1$, we consider

$$Q := \log \text{tr}_g g' - Bu + e^{-u}$$

for $B > 0$ to be chosen hereafter. Since $u \rightarrow -\infty$ near $E_{\text{nH}}(\theta)$ hence Q attains its maximum at some point $x_0 \in X \setminus E_{\text{nH}}(\theta)$. At this point, we have $\partial_{\bar{q}}Q = 0$. Therefore, the same arguments yields (cf. (4.1), (4.2)), at x_0

$$0 \geq -C(A+1)^2 e^u \frac{\text{tr}_{g'} g}{(\text{tr}_g g')^2} + \frac{\Delta_g \beta \varphi_\beta}{\text{tr}_g g'} + (B\delta_0 - C) \text{tr}_g g' - C$$

If at x_0 , $(\text{tr}_g g')^2 \leq C(A+1)^2 e^u$ we are done. Otherwise, since $\Delta_g \varphi_\beta = \text{tr}_g g' - \text{tr}_g(\theta + 2^{-\beta}\omega_X)$, it thus follows that

$$0 \geq (B\delta_0 - C - 1) \text{tr}_{g'} g - C - (n+1)B + \beta - \beta \frac{\sup_X \text{tr}_g(\theta + 2^{-\beta}\omega_X)}{\text{tr}_g g'}.$$

Choosing $B = (C+1)\delta_0^{-1}$ gives

$$(C + (n+1)B) \text{tr}_g g' + \beta \sup_X \text{tr}_g(\theta + 2^{-\beta}\omega_X) \geq \beta \text{tr}_g g'$$

at x_0 . Hence for $\beta > C + (n+1)B =: \beta_0$,

$$\text{tr}_g g'(x_0) \leq \frac{1}{1 - \beta_0/\beta} M, \text{ with } M = \sup_X \text{tr}_g(\theta + \omega_X)$$

It follows that for $\beta > \beta_0$,

$$(4.4) \quad \text{tr}_{\omega_X}(\theta + 2^{-\beta}\omega_X + \text{dd}^c\varphi_\beta) \leq \exp\left(\frac{1}{1 - \beta_0/\beta} M\right) \delta_0^{-n} e^{B \sup_X \varphi_\beta} e^{-B\rho}.$$

We have known that $\sup_X \varphi_\beta$ is uniformly bounded in terms of $A > 0$. Therefore, for $\beta > \beta_0$, for any compact subset $K \subset X \setminus E_{\text{nH}}(\theta)$,

$$-C_K \leq \Delta_{\omega_X} \varphi_\beta \leq C_K.$$

After passing a subsequence, the basic compactness result yields that φ_β converges to a function φ in $\mathcal{C}^{1,\alpha}$ for any fixed $\alpha \in]0, 1[$. It follows from the \mathcal{C}^0 -estimates that $\varphi = P_\theta(0)$. This completes the proof. \square

Remark 4.4. We can relax the assumption of f by merely assuming that $f \in \mathcal{C}^0(X)$, $\sup_X |f| \leq C$, and $\text{dd}^c f \leq C\omega_X$ for a fixed constant $C > 0$. Indeed, we approximate f uniformly by smooth functions f_j with $\text{dd}^c f_j \leq (C+1)\omega_X$. Let $\varphi_{\beta,j}$ be the solutions of the complex Monge–Ampère equations obtained by replacing θ_f with $\theta_{f_j} = \theta + \text{dd}^c f_j$. We still have that $M_j := \sup_X \text{tr}_{\omega_X}(\theta_{f_j} + \omega_X) \leq M < \infty$ and $\sup_X \varphi_{\beta,j} \leq C$ independent of j , $\beta \geq \beta_0$. Hence, we obtain the precise Laplacian estimate as in (4.4)

$$\text{tr}_{\omega_X}(\theta_{f_j} + 2^{-\beta}\omega_X + \text{dd}^c \varphi_{\beta,j}) \leq \exp\left(\frac{1}{1 - \beta_0/\beta} M\right) \delta_0^{-n} e^{B \sup_X \varphi_{\beta,j}} e^{-B\rho}.$$

This shows that $\Delta_{\omega_X} P_\theta(f_j)$ is locally bounded on $X \setminus E_{\text{nH}}(\theta)$ as $\beta \rightarrow +\infty$. Letting $j \rightarrow +\infty$, since $P_\theta(f_j)$ uniformly converges to $P_\theta(f)$ we arrive at the conclusion.

5. FINITE TIME SINGULARITIES OF THE CHERN–RICCI FLOW

Let (X, ω_0) be a compact Hermitian manifold of dimension n . We consider a solution $\omega = \omega(t)$ of the Chern–Ricci flow starting from ω_0 is given by a smooth family of Hermitian metrics satisfying

$$(5.1) \quad \begin{cases} \frac{\partial \omega}{\partial t} = -\text{Ric}(\omega), & 0 \leq t < T \\ \omega(0) = \omega_0. \end{cases}$$

One observes that just like the Kähler–Ricci flow, the Chern–Ricci flow is equivalent to a parabolic equation for a scalar function. Namely, solving a solution to the Chern–Ricci flow starting at ω_0 boils down to solving the parabolic complex Monge–Ampère equation, for $\varphi = \varphi(t)$ with $t \in [0, T[$,

$$(5.2) \quad \frac{\partial \varphi}{\partial t} = \log \frac{(\theta_t + \text{dd}^c \varphi)^n}{\omega_0^n}, \quad \theta_t + \text{dd}^c \varphi > 0, \quad \varphi(0) = 0$$

where $\theta_t := \omega_0 - t \text{Ric}(\omega_0)$. Since X is compact, by the standard parabolic theory, the Chern–Ricci flow always admits a unique solution on some time interval $[0, T[$ for some $0 < T \leq \infty$; see e.g. [Tos18a, Theorem 3.5]. As follows from [TW15], the maximal existence time T can be characterized by

$$T = \sup\{t > 0 : \exists \psi \in \mathcal{C}^\infty(X) \text{ with } \theta_t + \text{dd}^c \psi > 0\}.$$

When ω_0 satisfies the Guan–Li condition $\text{dd}^c \omega_0^k = 0$ for all $1 \leq k \leq n-1$, we can check that this condition is preserved by the Chern–Ricci flow, also does θ_t . Thanks to the Nakai–Moishezon criterion (Theorem B), we obtain a geometric characterization of the maximal existence time

$$T = \sup \left\{ t > 0 : \int_V \theta_t^k \wedge \omega_0^{\dim V - k} > 0, \forall \text{ irreducible subvarieties } V \subset X \right\}.$$

This extends the result [TW15, Theorem 1.3] to higher dimensions.

In this section, we study finite time singularities of the Chern–Ricci flow, proving theorem E. Suppose that $T < \infty$, so the flow develops singularities at a finite time. Following [CT15] (see also [TW22]), the set on which singularities of the Chern–Ricci flow develop on M (depending on the initial metric ω_0) is defined by

$$(5.3) \quad \begin{aligned} \text{Sing}(\omega(t)) = X \setminus \{x \in X \mid U \ni x, \exists \omega_T \text{ Hermitian metric on } U \\ \text{s.t. } \omega(t) \xrightarrow{t \rightarrow T^-} \omega_T \text{ in } \mathcal{C}_{\text{loc}}^\infty(U)\}. \end{aligned}$$

Following Z. Zhang [Zha10], we define the singularity formation set of the flow Σ (which depends on the initial metric ω_0) to be the complement of the set of points $x \in X$ such that there exists an open neighborhood U of x a constant $C > 0$ with $|\text{Rm}(t)|_{\omega(t)} \leq C$ on $[0, T[\times U$, where $\text{Rm}(t)$ denotes the curvature tensor of $\omega(t)$. Also, we define the set Σ' to be the complement of the set of points $x \in X$ such that there exists an open neighborhood U of x a constant $C > 0$ with $|R(t)|_{\omega(t)} \leq C$ on $[0, T[\times U$, where $R(t)$ is the scalar curvature of $\omega(t)$.

It was shown by Gill–Smith [GS15] that the singularity set of the flow is equal to the region on which the Chern scalar curvature blows up. The latter set is the singularity formation locus of the flow due to [EMT11]. We give two equivalent definitions of the singularity set $\text{Sing}(\omega(t))$.

Theorem 5.1. *Assume that the Chern–Ricci flow (5.1) exists on the maximal time interval $[0, T[$ with $T < \infty$. Then*

$$\text{Sing}(\omega(t)) = \Sigma = \Sigma'.$$

Proof. This was proved by M. Gill and K. Smith. However, it is unclear how the derivative $\partial_i Q_1$ in (3.9) of [GS15] vanishes at the maximum point of Q_2 . We give the proof here, mimicking their estimates with minor modifications.

Since a uniform bound on the curvature tensor implies an upper bound on the scalar curvature, we have $\Sigma' \subset \Sigma$. If the metrics $\omega(t)$ converges smoothly to a Hermitian metric ω_U on $[0, T[\times U$ for some open subset U , then we have that $|\text{Rm}(t)|_{\omega(t)} \leq C$, hence the inclusion $\text{Sing}(\omega(t)) \subset \Sigma$. It remains to show that if for any $x \in X$ with an open neighborhood U of x and $C > 0$ satisfying $|R(t)| \leq C$ on $[0, T[\times U$, then on a possibly smaller open neighborhood U' of x we have that $\omega(t)$ converges smoothly to a Hermitian metric $\omega_{U'}$ as $t \rightarrow T^-$. We remark that most of the estimates from the proof of Theorem 5.3 are local and hence could be applied again here.

Fixing $x \in K \subset\subset U$ a compact subset, we will establish uniform estimates on $\omega(t)$ in K . It follows from the flow equation that

$$\frac{\partial \varphi}{\partial t} = \log \frac{\omega(t)^n}{\omega_0^n}, \quad \frac{\partial}{\partial t} \left(\frac{\partial \varphi}{\partial t} \right) = -R \leq C$$

on $[0, T[\times U$. Integrating in time implies that $|\partial_t \varphi| \leq C$ and $|\varphi| \leq C$ on $[0, T[\times U$. Therefore the quantity $H = t \partial_t \varphi - \varphi - nt$ is uniformly bounded on $[0, T[\times U$, also is $\partial_t H = -tR - n$, and

$$\left(\frac{\partial}{\partial t} - \Delta_\omega \right) H = -\text{tr}_\omega \omega_0.$$

We set

$$Q := \log \text{tr}_{\omega_0} \omega + AH + e^H$$

for $A > 0$ to be determined hereafter. The quantity Q is achieved its maximum at some point $(t_0, x_0) \in [0, T'] \times K$ with $T' < T$. It follows from [TW15, Proposition 3.1] (also [TW15, (4.2)]) that

$$(5.4) \quad \left(\frac{\partial}{\partial t} - \Delta_\omega \right) \log \text{tr}_{\omega_0} \omega \leq \frac{2}{(\text{tr}_{\omega_0} \omega)^2} \text{Re}(g^{\bar{q}k}(T_0)_{kp}^p \partial_{\bar{q}} \text{tr}_{\omega_0} \omega) + C \text{tr}_\omega \omega_0,$$

where $(T_0)_{kp}^p$ denote the torsion terms corresponding to ω_0 . At the maximum point (t_0, x_0) of Q we have $\partial_{\bar{q}} Q = 0$ hence

$$\frac{1}{\text{tr}_{\omega_0} \omega} \partial_{\bar{q}} \text{tr}_{\omega_0} \omega + A \partial_{\bar{q}} H + e^H \partial_{\bar{q}} H = 0.$$

Therefore, the Cauchy–Schwarz inequality yields

$$(5.5) \quad \left| \frac{2}{(\text{tr}_{\omega_0} \omega)^2} \text{Re}(g^{\bar{q}k}(T_0)_{kp}^p \partial_{\bar{q}} \text{tr}_{\omega_0} \omega) \right| \leq \left| \frac{2}{\text{tr}_{\omega_0} \omega} \text{Re}((A + e^H) g^{\bar{q}k}(T_0)_{kp}^p \partial_{\bar{q}} H) \right| \\ \leq e^H |\partial H|_\omega^2 + C(A + 1)^2 e^{-H} \frac{\text{tr}_\omega \omega_0}{(\text{tr}_{\omega_0} \omega)^2}$$

for uniform $C > 0$ only depending on the torsion term. From (5.4) and (5.5), we have at (t_0, x_0) ,

$$0 \leq \left(\frac{\partial}{\partial t} - \Delta_\omega \right) Q \leq C(A + 1)^2 e^{-H} \frac{\text{tr}_\omega \omega_0}{(\text{tr}_{\omega_0} \omega)^2} + C \text{tr}_\omega \omega_0 - (A + e^H) \text{tr}_\omega \omega_0.$$

If $(\text{tr}_\omega \omega_0)^2 \leq C(A + 1)^2 e^{-H}$ at (t_0, x_0) , then

$$Q \leq C - \frac{1}{2}H + AH + e^H \leq C_1$$

we are done. Otherwise, we choose $A = C + 1$, then at (t_0, x_0) ,

$$\text{tr}_\omega \omega_0 \leq C.$$

On the other hand, since $\omega^n / \omega_0^n = \partial_t \varphi \leq C$, hence

$$\mathrm{tr}_{\omega_0} \omega \leq C (\mathrm{tr}_{\omega} \omega_0)^{n-1} \frac{\omega^n}{\omega_0^n} \leq C.$$

This implies that Q , and thus $\mathrm{tr}_{\omega_0} \omega$ is bounded from above on $[0, T[\times K$ since T' was taken arbitrarily. Finally, the local higher-order estimates for ω on $[0, T[\times U'$ with $U' \subset U$ follow by Sherman–Weinkove [SW13]. The proof is complete. \square

Consequently, the scalar curvature blows up at a finite time singularity, generalizing Zhang’s analogous result [Zha10] in the Kähler case.

Corollary 5.2 ([GS15, Theorem 1.1]). *If the Chern–Ricci flow develops a finite time singularity, then the singularity formation set Σ is nonempty, and furthermore,*

$$\limsup_{t \rightarrow T} \left(\sup_X R(t) \right) = +\infty.$$

Proof. If $\Sigma = \emptyset$ then by Theorem 5.1, replacing K with X , we would have that $\omega(t)$ converges in C^∞ to a Hermitian metric ω_T on X , a contradiction. The blow-up of the supremum of the scalar curvature again follows from Theorem 5.1. \square

We next compare the singularity formation set of the flow $\mathrm{Sing}(\omega(t))$ with the null locus of the limiting form $\theta_T = \omega_0 - T \mathrm{Ric}(\omega_0)$ in some specific contexts. We see that the $\partial\bar{\partial}$ class $[\theta_T]$ is nef, but not Hermitian. Since the definition of $\widehat{v}_-(X, \theta_T)$ does not depend on the choice of reference Hermitian metric ω_X and for any $\varepsilon > 0$,

$$(1 - \varepsilon)\theta_T + \varepsilon\omega_0 = \theta_{(1-\varepsilon)T}$$

so we have

$$\widehat{v}_-(V, \theta_T|_V) = \inf_{t \rightarrow T^-} v_-(V, \omega(t)|_V)$$

for all irreducible analytic subvarieties $V \subset X$. We define the set-theoretic union of all such subvarieties for which the infimum volume shrinks to zero as $t \rightarrow T^-$,

$$(5.6) \quad N(\omega(t)) := \bigcup_{v_-(V, \omega(t)|_V) \xrightarrow{t \rightarrow T} 0} V.$$

In particular, the set $N(\omega(t))$ is a closed analytic subset of X .

If θ_T is big, we can find a θ_T -psh function ρ with almost analytic singularities along $E_{\mathrm{NH}}(\theta_T)$ such that

$$\theta_T + \mathrm{dd}^c \rho \geq 2\delta_0 \omega_0$$

for some $\delta_0 > 0$. Up to subtracting a positive constant, we can assume that $\sup_X \rho \leq 0$. We have

$$\theta_t + \mathrm{dd}^c \rho = \frac{1}{T} ((T-t)(\omega_0 + \mathrm{dd}^c \rho) + t(\theta_T + \mathrm{dd}^c \rho)) \geq \delta_0 \omega_0$$

if $t \in [T - \varepsilon, T[$ for some $\varepsilon > 0$.

We are interested in the behavior of the metrics $\omega(t)$ as $t \rightarrow T^-$, proving the following

Theorem 5.3. *Assume that the limiting form θ_T is big. Then there exists a positive (1,1) current ω_T on X in the $\partial\bar{\partial}$ class $[\theta_T]$, which is smooth precisely away from $E_{\mathrm{NH}}(\theta_T)$ and has minimal singularities, such that as $t \rightarrow T^-$ the metrics $\omega(t)$ converge to ω_T in both current and $C_{\mathrm{loc}}^\infty(X \setminus E_{\mathrm{NH}}(\theta_T))$ senses. In particular, $\mathrm{Sing}(\omega(t)) \subset E_{\mathrm{NH}}(\theta_T)$.*

Lemma 5.4. *There is a uniform constant $C_0 > 0$ such that on $[0, T[\times X$ we have*

- (i) $C_0 \geq \varphi \geq \rho - C_0$;
- (ii) $C_0 \geq \partial_t \varphi \geq C_0 \rho - C_0$.

Proof. The proof follows almost verbatim from the Kähler case [CT15, Sect. 4]. We also refer to [TW13, Sect. 2] or [Dan23, Sect. 5] for a proof. \square

Lemma 5.5. *There is a constant $C > 0$ such that on $[0, T] \times X$ we have*

$$\mathrm{tr}_{\omega_0} \omega \leq C e^{-C\rho}.$$

Proof. The proof relies on the trick of Phong–Sturm [PS10], we repeat the estimates derived in the proof of [TW15, Lemma 4.1] with minor modifications. Set $\psi = \varphi - \rho + C_0 \geq 0$, so $e^{-\psi} \in]0, 1]$. We compute the evolution of

$$Q = \log \mathrm{tr}_{\omega_0} \omega - A\psi + e^{-\psi}$$

for $A > 0$ to be determined hereafter. It suffices to show that Q is uniformly bounded from above. We observe that $Q \leq C$ on $[0, T - \varepsilon] \times X$ for a uniform $C > 0$. Fixing $T - \varepsilon < T' < T$, suppose that Q attains its maximum at some point $(t_0, x_0) \in [0, T'] \times X$ with $t \in]T - \varepsilon, T']$. In what follows, we compute at this point. At the maximum point (t_0, x_0) of Q we have $\partial_i Q = 0$ hence

$$\frac{1}{\mathrm{tr}_{\omega_0} \omega} \partial_i \mathrm{tr}_{\omega_0} \omega - A \partial_i \psi - e^{-\psi} \partial_i \psi = 0.$$

Therefore, the Cauchy–Schwarz inequality yields

$$(5.7) \quad \left| \frac{2}{(\mathrm{tr}_{\omega_0} \omega)^2} \mathrm{Re}(g^{\bar{q}k}(T_0)_{kp}^p \partial_{\bar{q}} \mathrm{tr}_{\omega_0} \omega) \right| \leq \left| \frac{2}{\mathrm{tr}_{\omega_0} \omega} \mathrm{Re}((A + e^{-\psi}) g^{\bar{q}k}(T_0)_{kp}^p \partial_{\bar{q}} \psi) \right| \\ \leq e^{-\psi} |\partial \psi|_{\omega}^2 + C(A + 1)^2 e^{\psi} \frac{\mathrm{tr}_{\omega} \omega_0}{(\mathrm{tr}_{\omega_0} \omega)^2}$$

for uniform $C > 0$ only depending on the torsion term. It thus follows from (5.4) and (5.7) that, at (t_0, x_0) ,

$$(5.8) \quad 0 \leq \left(\frac{\partial}{\partial t} - \Delta_{\omega} \right) Q \leq C(A + 1)^2 e^{\psi} \frac{\mathrm{tr}_{\omega} \omega_0}{(\mathrm{tr}_{\omega_0} \omega)^2} + C \mathrm{tr}_{\omega} \omega_0 \\ - (A + e^{-\psi}) \partial_t \varphi + (A + e^{-\psi}) \mathrm{tr}_{\omega} (\omega - (\alpha_t + \mathrm{dd}^c \rho)) \\ \leq C(A + 1)^2 e^{\psi} \frac{\mathrm{tr}_{\omega} \omega_0}{(\mathrm{tr}_{\omega_0} \omega)^2} + (C - A\delta_0) \mathrm{tr}_{\omega} \omega_0 + (A + 1) \log \frac{\omega_0^n}{\omega^n}$$

where we have used $\theta_t + \mathrm{dd}^c \rho \geq \delta_0 \omega_0$. If at (t_0, x_0) , $(\mathrm{tr}_{\omega_0} \omega)^2 \leq e^{\psi} C(A + 1)^2$ then at the same point we obtain

$$Q \leq C + \frac{1}{2} \psi - A\psi + e^{-\psi} \leq C + 1$$

since $\psi \geq 0$, we are done. Otherwise, we choose $A = \delta_0^{-1}(C + 2)$. Hence, from (5.8) one gets at the maximum of Q ,

$$\mathrm{tr}_{\omega} \omega_0 \leq C \log \frac{\omega_0^n}{\omega^n} + C,$$

so

$$\mathrm{tr}_{\omega_0} \omega \leq n (\mathrm{tr}_{\omega} \omega_0)^{n-1} \frac{\omega^n}{\omega_0^n} \leq C \frac{\omega^n}{\omega_0^n} \left(\log \frac{\omega_0^n}{\omega^n} \right)^{n-1} + C \leq C'$$

since $\omega^n / \omega_0^n \leq C_0$ by Lemma 5.4 and $y \mapsto y |\log y|^{n-1}$ is bounded from above as $y \rightarrow 0$. Thanks to Lemma 5.4 (iii), Q is bounded from above at its maximum, finishing the proof. \square

Proof of Theorem 5.3. We apply Lemmas 5.4 and 5.5 to get that for any compact set $K \subset X \setminus E_{\text{nH}}(\theta_T)$, $k \in \mathbb{N}$, there is a constant $C_{K,k} > 0$ such that

$$\|\varphi(t)\|_{C^k(K)} \leq C_{K,k}$$

using the complex parabolic Evans–Krylov theory together with Schauder’s estimate; see also Sherman–Weinkove’s local curvature estimates [SW13]. This implies that for some constant $C_K > 0$,

$$\frac{\partial}{\partial t}(\varphi(t) + C_K t) \geq 0$$

hence $\varphi(t) + C_K t$ is increasing in t as well as being bounded from above. This yields that $\varphi(t)$ has a limit as $t \rightarrow T^-$, denoted by φ_T . An elementary argument yields that $\varphi(t)$ converges in C^∞ on compact subsets of $X \setminus E_{\text{nH}}(\alpha)$. By weak compactness φ_T extends to a θ_T -psh function on X and $\varphi(t)$ converges to φ_T in $L^1(X)$. The current ω_T can be expressed as $\omega_T = \theta_T + \text{dd}^c \varphi_T$.

Next, we show that φ_T has minimal singularities, that is, for any θ_t -psh function u there is a constant $C > 0$ such that

$$\varphi \geq u - C \quad \text{on } [0, T] \times X.$$

Thanks to Demailly’s regularization theorem (Theorem 2.11), we can find functions with logarithmic singularities u_m decreasing to u such that

$$\theta_T + \text{dd}^c u_m \geq -\varepsilon_m \omega_0, \quad \varepsilon_m \searrow 0.$$

We set

$$G = (T - t)\partial_t \varphi + \varphi + nt - \varepsilon_m(t\partial_t \varphi - \varphi - nt) - u_m.$$

On $X \setminus \text{Sing}(u_m)$, G is smooth and

$$\left(\frac{\partial}{\partial t} - \Delta_\omega\right) G = \text{tr}_\omega(\theta_T + \varepsilon_m \omega_0 + \text{dd}^c u_m) \geq 0.$$

Since $\sup_X u_m \leq C$ uniformly, so G is achieved its minimum in $X \setminus \text{Sing}(u_m)$. Hence the minimum principle implies that $G \geq -C$ independent of m , so

$$(T - t - \varepsilon_m)\partial_t \varphi + (1 + \varepsilon_m)\varphi(t) \geq u_m - C \geq u - C.$$

We take $m \rightarrow +\infty$ and use the fact that $\partial_t \varphi \leq C_0$ to conclude. \square

Remark 5.6. The current ω_T has minimal singularities but is not a Hermitian current. Indeed, we write $\omega_T = \theta_T + \text{dd}^c \varphi_T$. If ω_T were Hermitian, we would have that the current $\theta_T + \text{dd}^c(1 - \varepsilon)\varphi_T$ is Hermitian for $\varepsilon > 0$ sufficiently small. By minimality, we have $\varphi_T + C \geq (1 - \varepsilon)\varphi_T$ for a constant $C > 0$ depending on φ_T , hence $\varphi_T \geq -C\varepsilon^{-1}$ is bounded. This implies that $\text{Sing}(\varphi_T) = \emptyset$, hence the $\partial\bar{\partial}$ -class $[\theta_T]$ is Hermitian, a contradiction.

Theorem 5.7. *Let (X^n, ω_0) be a compact Hermitian manifold. Let $\omega(t)$ be a solution of the Chern–Ricci flow starting at ω_0 , which develops a singularity at finite time T . Assume that the limiting form $\theta_T = \omega_0 - T \text{Ric}(\omega_0)$ satisfies $\hat{v}_-(X, \theta_T) > 0$. Then if either*

- $n = 2$,
- or $n \geq 3$, $\text{dd}^c \omega_0 = 0$, $d\omega_0 \wedge d^c \omega_0 = 0$,

then the metrics $\omega(t)$ develop singularities precisely along $N(\omega(t)) = E_{\text{nH}}(\theta_T)$.

Recall here that the set $N(\omega(t))$ is defined by (5.6). We will see in the proof that $N(\omega(t))$ is the set-theoretic union of all such subvarieties for which the volume shrinks to zero as $t \rightarrow T^-$. This, therefore, partially answers the question of Tosatti–Weinkove [TW22, Question 6.1].

Proof. By assumption, the metric ω_0 always satisfies the condition (H_X) . Since θ_T is nef and $\widehat{v}_-(X, \theta_T) > 0$, so θ_T is big. Thanks to Theorems 5.3 and 3.7, we have that

$$\text{Sing}(\omega(t)) \subset E_{\text{RH}}(\theta_T) = N(\omega(t)).$$

It suffices to show that the metrics $\omega(t)$ develop singularities along $N(\omega(t))$. Suppose by contradiction that there is $x \in N(\omega(t))$ with a neighborhood $U \ni x$ on which as $t \rightarrow T^-$, the metrics $\omega(t)$ converges smoothly to a Hermitian metric ω_U and there is a positive-dimension irreducible subvariety $V \subset X$ containing x and $v_-(V, \omega(t)|_V) \rightarrow 0$ as $t \rightarrow T^-$. We divide it into two cases.

If $n = 2$, we must have $\dim V = 1$ because $\widehat{v}_-(X, \theta_T) > 0$. Then

$$v_-(V, \omega(t)|_V) = \int_V \omega(t) \geq \int_{V \cap U} \omega(t) \xrightarrow{t \rightarrow T^-} \int_{V \cap U} \omega_U > 0,$$

a contradiction.

If $n \geq 3$ and $\text{dd}^c \omega_0 = 0$ and $d\omega_0 \wedge d^c \omega_0 = 0$ then $\text{dd}^c \omega_0^k = 0, \forall 1 \leq k \leq n-1$. The latter condition is preserved along the Chern–Ricci flow, i.e., $\text{dd}^c \omega(t)^k = 0$ for all $1 \leq k \leq n-1$. Then we have

$$v_-(V, \omega(t)|_V) = \int_V \omega(t)^{\dim V}$$

for every irreducible subvariety $V \subset X$, thanks to Lemma 2.5. On the other hand,

$$\int_V \omega(t)^{\dim V} \geq \int_{V \cap U} \omega(t)^{\dim V} \xrightarrow{t \rightarrow T^-} \int_{V \cap U} \omega_U^{\dim V} > 0,$$

a contradiction. \square

In the setup above, we show that the Chern–Ricci flow performs a canonical surgical contraction:

Theorem 5.8. *Let X^n be compact Hermitian manifolds equipped with a Hermitian metric ω_0 with $\text{dd}^c \omega_0 = 0$ and $d\omega_0 \wedge d^c \omega_0 = 0$. Assume that the solution $\omega(t)$ of the Chern–Ricci flow (5.1) starting at ω_0 exists on $[0, T[$ with $T < \infty$ and the limiting form $\theta_T = \omega_0 - T \text{Ric}(\omega_0)$ satisfies $\int_X \theta_T^n > 0$. Assume there exists a map $\pi : X \rightarrow Y$ blowing down disjoint irreducible exceptional divisors E_i on X to points $y_i \in Y$ for $i = 1, \dots, p$, with Y a compact Hermitian manifold, such that there exist a function $f \in C^\infty(X, \mathbb{R})$ and a smooth real (1,1)-form η on Y with*

$$(*) \quad \theta_T + \text{dd}^c f = \pi^* \eta.$$

Then we have the following

- (i) As $t \rightarrow T^-$, the metrics $\omega(t)$ converges to a smooth Hermitian metric ω_T on $X^\circ := X \setminus \bigcup_{i=1}^p E_i$ in $C_{\text{loc}}^\infty(X^\circ)$. Using π , ω_T can be regarded as a Hermitian metric on $Y^\circ := Y \setminus \{y_1, \dots, y_p\}$.
- (ii) Let d_{ω_T} be the distance function induced by ω_T on Y° . Then there exists a unique metric d_T on Y extending d_{ω_T} to be zero on $\{y_1, \dots, y_p\}$ such that (Y, d_T) is a compact metric space homeomorphic to Y .
- (iii) The metric space $(X, \omega(t)) \rightarrow (Y, d_T)$ as $t \rightarrow T^-$ in the Gromov–Hausdorff sense.
- (iv) A smooth maximal solution $g(t)$ of the Chern–Ricci flow on Y starting at ω_T exists for $t \in]T, T_Y[$ with $T < T_Y \leq \infty$, and $g(t)$ converges to ω_T in $C_{\text{loc}}^\infty(Y^\circ)$ as $t \rightarrow T^+$.
- (v) The metric space $(Y, g(t)) \rightarrow (Y, d_T)$ as $t \rightarrow T^+$ in the Gromov–Hausdorff sense.

Proof. We observe that $\text{dd}^c \omega_0^k = 0$ for all $1 \leq k \leq n-1$, this property is preserved along the Chern–Ricci flow. Since θ_T is nef and $\int_X \theta_T^n > 0$ we have that θ_T is big. By the previous analysis, together with assumptions, we know that

$$E_{\text{nH}}(\theta_T) = \text{Null}(\theta_T) = \bigcup_{1 \leq i \leq p} E_i.$$

The (i) follows immediately from Theorem 5.3.

We show that in the condition (*) the form η can be replaced with a Hermitian metric ω_Y on Y . Indeed, it is obvious that η is nef and for any irreducible subvariety $V \subset Y$ with $k = \dim V > 0$, we have that

$$\int_V \eta^k = \int_{\pi^{-1}(V)} (\pi^* \eta)^k = \int_{\pi^{-1}(V)} \theta_T^k > 0$$

Thanks to Nakai–Moishezon’s criterion (Theorem 3.10) the $\partial\bar{\partial}$ -class $[\eta]$ is Hermitian on Y . We thus obtain (ii) and (iii) as follows from [TW13, Theorem 1.3].

We have seen that ω_T is a positive current with minimal singularities, so it has bounded potentials. The push forward $\pi_* \omega_T$ is therefore a positive current on Y with bounded potentials. In this case, we have $\pi_* \omega_T = \omega_Y + \text{dd}^c \psi_T$ where ψ_T is a bounded ω_Y -psh function, which is smooth in Y° . With this condition, the (iv) and (v) was proved by Nie [Nie17] and Tô [Tô18] independently. \square

The major problems are whether the metric space (Y, d_T) is the metric completion of (Y°, d_{ω_T}) , and we can remove the condition (*). Those may require new techniques.

REFERENCES

- [AGL23] D. Angella, V. Guedj, and C. H Lu. Plurisigned hermitian metrics. *Trans. Amer. Math. Soc.*, 376(7):4631–4659, 2023. [2](#), [19](#)
- [BCL14] S. Boucksom, S. Cacciola, and A. F. Lopez. Augmented base loci and restricted volumes on normal varieties. *Math. Z.*, 278(3-4):979–985, 2014. [1](#)
- [Ber19] R. J. Berman. From Monge–Ampère equations to envelopes and geodesic rays in the zero temperature limit. *Math. Z.*, 291(1-2):365–394, 2019. [4](#), [24](#)
- [Bir17] C. Birkar. The augmented base locus of real divisors over arbitrary fields. *Math. Ann.*, 368(3-4):905–921, 2017. [1](#)
- [Bou04] S. Boucksom. Divisorial Zariski decompositions on compact complex manifolds. *Ann. Sci. École Norm. Sup. (4)*, 37(1):45–76, 2004. [2](#), [9](#), [10](#)
- [Buc99] N. Buchdahl. On compact Kähler surfaces. *Ann. Inst. Fourier (Grenoble)*, 49(1):vii, xi, 287–302, 1999. [3](#)
- [Buc00] N. Buchdahl. A Nakai–Moishezon criterion for non-Kähler surfaces. *Ann. Inst. Fourier (Grenoble)*, 50(5):1533–1538, 2000. [3](#), [20](#)
- [Che87] P. Cherrier. Équations de Monge–Ampère sur les variétés hermitiennes compactes. *Bull. Sci. Math. (2)*, 111(4):343–385, 1987. [4](#), [24](#)
- [Chi16a] I. Chiose. The Kähler rank of compact complex manifolds. *J. Geom. Anal.*, 26(1):603–615, 2016. [3](#), [9](#)
- [Chi16b] I. Chiose. On the invariance of the total Monge–Ampère volume of Hermitian metrics. [arXiv:1609.05945](#), 2016. [2](#), [6](#)
- [CL14] S. Cacciola and A. F. Lopez. Nakamaye’s theorem on log canonical pairs. *Ann. Inst. Fourier (Grenoble)*, 64(6):2283–2298, 2014. [1](#)
- [CMM14] P. Cascini, J. McKernan, and M. Mustață. The augmented base locus in positive characteristic. *Proc. Edinb. Math. Soc. (2)*, 57(1):79–87, 2014. [1](#)
- [CT15] T. C. Collins and V. Tosatti. Kähler currents and null loci. *Invent. Math.*, 202(3):1167–1198, 2015. [2](#), [5](#), [10](#), [12](#), [13](#), [18](#), [26](#), [29](#)
- [CT16] T. C. Collins and V. Tosatti. A singular Demailly–Păun theorem. *C. R. Math. Acad. Sci. Paris*, 354(1):91–95, 2016. [3](#), [20](#)
- [CZ19] J. Chu and B. Zhou. Optimal regularity of plurisubharmonic envelopes on compact Hermitian manifolds. *Sci. China Math.*, 62(2):371–380, 2019. [4](#)
- [Dan23] Q.-T. Dang. Singularities of the chern-ricci flow. [arXiv:2305.16461](#), 2023. [29](#)

- [Dem85] J.-P. Demailly. Mesures de Monge-Ampère et caractérisation géométrique des variétés algébriques affines. *Mém. Soc. Math. France (N.S.)*, (19):124, 1985. **11**
- [Dem94] J.-P. Demailly. Regularization of closed positive currents of type $(1, 1)$ by the flow of a Chern connection. In *Contributions to complex analysis and analytic geometry*, Aspects Math., E26, pages 105–126. Friedr. Vieweg, Braunschweig, 1994. **23**
- [Dem12] J.-P. Demailly. Complex analytic and differential geometry. *available at <https://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>*, 2012. **7, 20**
- [DHP22] O. Das, C. Hacon, and M. Păun. On the 4-dimensional minimal model program for Kähler varieties. *arXiv:2205.12205*, 2022. **3, 20**
- [DK12] S. Dinew and S. Kołodziej. Pluripotential estimates on compact Hermitian manifolds. In *Advances in geometric analysis*, volume 21 of *Adv. Lect. Math. (ALM)*, pages 69–86. Int. Press, Somerville, MA, 2012. **7**
- [DP04] J.-P. Demailly and M. Paun. Numerical characterization of the Kähler cone of a compact Kähler manifold. *Ann. of Math. (2)*, 159(3):1247–1274, 2004. **2, 3, 8, 12, 14, 15, 20**
- [ELM⁺09] L. Ein, R. Lazarsfeld, M. Mustață, M. Nakamaye, and M. Popa. Restricted volumes and base loci of linear series. *Amer. J. Math.*, 131(3):607–651, 2009. **1**
- [EMT11] J. Enders, R. Müller, and P. M. Topping. On type-I singularities in Ricci flow. *Comm. Anal. Geom.*, 19(5):905–922, 2011. **26**
- [FT11] A. Fino and A. Tomassini. On astheno-Kähler metrics. *J. Lond. Math. Soc. (2)*, 83(2):290–308, 2011. **3**
- [GH78] P. Griffiths and J. Harris. *Principles of algebraic geometry*. Wiley-Interscience [John Wiley & Sons], New York, 1978. Pure and Applied Mathematics. **11**
- [GL10] B. Guan and Q. Li. Complex Monge-Ampère equations and totally real submanifolds. *Adv. Math.*, 225(3):1185–1223, 2010. **3, 6**
- [GL22] V. Guedj and C. H. Lu. Quasi-plurisubharmonic envelopes 2: Bounds on Monge-Ampère volumes. *Algebr. Geom.*, 9(6):688–713, 2022. **3, 6, 7, 8, 9, 10, 17, 18, 19, 21**
- [GL23] V. Guedj and C. H. Lu. Quasi-plurisubharmonic envelopes 3: Solving Monge-Ampère equations on hermitian manifolds. *J. Reine Angew. Math.*, 800:259–298, 2023. **4, 21, 22, 24**
- [GS15] M. Gill and D. Smith. The behavior of the Chern scalar curvature under the Chern-Ricci flow. *Proc. Amer. Math. Soc.*, 143(11):4875–4883, 2015. **5, 26, 27, 28**
- [GZ17] V. Guedj and A. Zeriahi. *Degenerate complex Monge-Ampère equations*, volume 26 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2017. **7, 8, 18, 21**
- [Hir77] H. Hironaka. Bimeromorphic smoothing of a complex-analytic space. *Acta Math. Vietnam.*, 2(2):103–168, 1977. **13, 16, 19**
- [Lam99] A. Lamari. Courants kählériens et surfaces compactes. *Ann. Inst. Fourier (Grenoble)*, 49(1):vii, x, 263–285, 1999. **3, 9, 20**
- [Nak00] M. Nakamaye. Stable base loci of linear series. *Math. Ann.*, 318(4):837–847, 2000. **1, 11**
- [Nak15] M. Nakamaye. Roth’s theorem: an introduction to diophantine approximation. In *Rational points, rational curves, and entire holomorphic curves on projective varieties*, volume 654 of *Contemp. Math.*, pages 75–108. Amer. Math. Soc., Providence, RI, 2015. **1**
- [Nie17] X. Nie. Weak solution of the Chern-Ricci flow on compact complex surfaces. *Math. Res. Lett.*, 24(6):1819–1844, 2017. **5, 32**
- [Pop16] D. Popovici. Sufficient bigness criterion for differences of two nef classes. *Math. Ann.*, 364(1-2):649–655, 2016. **3, 9, 10**
- [PS10] D. H. Phong and Jacob Sturm. The Dirichlet problem for degenerate complex Monge-Ampère equations. *Comm. Anal. Geom.*, 18(1):145–170, 2010. **22, 29**
- [Ric68] R. Richberg. Stetige streng pseudokonvexe Funktionen. *Math. Ann.*, 175:257–286, 1968. **12**
- [SW13] M. Sherman and B. Weinkove. Local Calabi and curvature estimates for the Chern-Ricci flow. *New York J. Math.*, 19:565–582, 2013. **28, 30**
- [Tos18a] V. Tosatti. KAWA lecture notes on the Kähler-Ricci flow. *Ann. Fac. Sci. Toulouse Math. (6)*, 27(2):285–376, 2018. **26**
- [Tos18b] Valentino Tosatti. Nakamaye’s theorem on complex manifolds. In *Algebraic geometry: Salt Lake City 2015*, volume 97.1 of *Proc. Sympos. Pure Math.*, pages 633–655. Amer. Math. Soc., Providence, RI, 2018. **1, 17**
- [TW10a] V. Tosatti and B. Weinkove. The complex Monge-Ampère equation on compact Hermitian manifolds. *J. Amer. Math. Soc.*, 23(4):1187–1195, 2010. **4, 9, 21, 24**
- [TW10b] V. Tosatti and B. Weinkove. Estimates for the complex Monge-Ampère equation on Hermitian and balanced manifolds. *Asian J. Math.*, 14(1):19–40, 2010. **23**
- [TW13] V. Tosatti and B. Weinkove. The Chern-Ricci flow on complex surfaces. *Compos. Math.*, 149(12):2101–2138, 2013. **5, 29, 32**
- [TW15] V. Tosatti and B. Weinkove. On the evolution of a Hermitian metric by its Chern-Ricci form. *J. Differential Geom.*, 99(1):125–163, 2015. **4, 22, 26, 27, 29**

- [TW22] V. Tosatti and B. Weinkove. The Chern-Ricci flow. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.*, 33(1):73–107, 2022. [5](#), [26](#), [30](#)
- [Tô18] T. D. Tô. Regularizing properties of complex Monge-Ampère flows II: Hermitian manifolds. *Math. Ann.*, 372(1-2):699–741, 2018. [5](#), [32](#)
- [Wło09] J. Włodarczyk. Resolution of singularities of analytic spaces. In *Proceedings of Gökova Geometry-Topology Conference 2008*, pages 31–63. Gökova Geometry/Topology Conference (GGT), Gökova, 2009. [9](#), [13](#), [19](#)
- [Zha10] Z. Zhang. Scalar curvature behavior for finite-time singularity of Kähler-Ricci flow. *Michigan Math. J.*, 59(2):419–433, 2010. [26](#), [28](#)

THE ABDUS SALAM INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS (ICTP), STR. COSTIERA,
11, 34151 TRIESTE, TS, ITALY

Email address: `qdang@ictp.it`