

Towards a classification of 1-homogeneous distance-regular graphs with positive intersection number a_1

Jack H. Koolen^{a, b}, Mamoon Abdullah^a, Brhane Gebremichel^a, and Jae-Ho Lee^{*c}

^aSchool of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui, 230026, PR China.

^bCAS Wu Wen-Tsun Key Laboratory of Mathematics, University of Science and Technology of China, Hefei, Anhui, 230026, PR China.

^cDepartment of Mathematics and Statistics, University of North Florida, Jacksonville, FL 32224, U.S.A

Abstract

Let Γ be a graph with diameter at least two. Then Γ is said to be 1-homogeneous (in the sense of Nomura) whenever for every pair of adjacent vertices x and y in Γ , the distance partition of the vertex set of Γ with respect to both x and y is equitable, and the parameters corresponding to equitable partitions are independent of the choice of x and y . Assume Γ is 1-homogeneous distance-regular with intersection number $a_1 > 0$ and $D \geq 5$. Define $b = b_1/(\theta_1 + 1)$, where b_1 is the intersection number and θ_1 is the second largest eigenvalue of Γ . We show that if intersection number $c_2 \geq 2$, then $b \geq 1$ and one of the following (i)–(vi) holds: (i) Γ is a regular near $2D$ -gon, (ii) Γ is a Johnson graph $J(2D, D)$, (iii) Γ is a halved ℓ -cube where $\ell \in \{2D, 2D + 1\}$, (iv) Γ is a folded Johnson graph $\bar{J}(4D, 2D)$, (v) Γ is a folded halved $(4D)$ -cube, (vi) the valency of Γ is bounded by a function of b . Using this result, we characterize 1-homogeneous graphs with classical parameters and $a_1 > 0$, as well as tight distance-regular graphs.

Keywords: distance-regular graph, 1-homogeneous, local graph, classical parameters, tight graph

Mathematics Subject Classification: 05E30, 05C50

1 Introduction

In this paper, we study distance-regular graphs that have the 1-homogeneous property in the sense of Nomura [19]. To motivate our results, we recall some preliminaries and background on 1-homogeneous distance-regular graphs. For more details, refer to [1, 5, 19].

Throughout this paper, let Γ denote a finite, undirected, connected, and simple graph. Let $V(\Gamma)$ denote the vertex set of Γ . For two vertices $x, y \in V(\Gamma)$, the *distance* $d(x, y)$ is the length of a shortest path from x

*Corresponding author

E-mail addresses: koolen@ustc.edu.cn (J.H. Koolen), mamoonabdullah@hotmail.com (M. Abdullah), brhane@ustc.edu.cn (B. Gebremichel), jaeho.lee@unf.edu (J.-H. Lee)

to y in Γ . The *diameter* of Γ is the maximum value of $d(x, y)$ for all pairs of $x, y \in V(\Gamma)$. Let D denote the diameter of Γ . For an integer $0 \leq i \leq D$ and a vertex $x \in V(\Gamma)$, let $\Gamma_i(x)$ denote the set of vertices in Γ at distance i from x . Abbreviate $\Gamma(x) = \Gamma_1(x)$. The subgraph of Γ induced on the set $\Gamma(x)$ is called the *local graph* of Γ at x . For a pair of vertices $x, y \in V(\Gamma)$ with $d(x, y) = 2$, the subgraph of Γ induced on the set $\Gamma(x) \cap \Gamma(y)$ is called the μ -*graph* of x and y . The graph Γ is called *locally \mathcal{P}* whenever every local graph of Γ has the property \mathcal{P} (or belongs to the family \mathcal{P}). For example, we say Γ is locally connected or locally a strongly regular graph. For an integer $k \geq 0$, we say that Γ is *regular with valency k* (or *k -regular*) if $|\Gamma(x)| = k$ for every $x \in V(\Gamma)$. For an integer $0 \leq i \leq D$ and for a pair of vertices $x, y \in V(\Gamma)$ with $d(x, y) = i$ we define

$$C_i(x, y) := \Gamma_{i-1}(x) \cap \Gamma(y), \quad A_i(x, y) := \Gamma_i(x) \cap \Gamma(y), \quad B_i(x, y) := \Gamma_{i+1}(x) \cap \Gamma(y), \quad (1)$$

where $C_0(x, y) := \emptyset$ and $B_D(x, y) := \emptyset$. Observe that $\Gamma(y)$ is the disjoint union of the vertex sets $C_i(x, y)$, $A_i(x, y)$, $B_i(x, y)$. We say Γ is *distance-regular* whenever the cardinalities

$$c_i = |C_i(x, y)|, \quad a_i = |A_i(x, y)|, \quad b_i = |B_i(x, y)| \quad (0 \leq i \leq D) \quad (2)$$

are constants and do not depend on the choice of x and y . Note that $c_0 = a_0 = b_D = 0$, and $c_1 = 1$. Additionally, Γ is regular with valency $k = b_0$, and $a_i + b_i + c_i = k$ ($0 \leq i \leq D$). The numbers a_i, b_i, c_i in (2) are called the *intersection numbers* of Γ , and the array $\{b_0, b_1, \dots, b_{D-1}; c_1, c_2, \dots, c_D\}$ is called the *intersection array* of Γ . We note that a distance-regular graph with diameter D has exactly $D + 1$ distinct eigenvalues [5, Proposition 2.6].

Notation 1.1. Unless otherwise specified, whenever we denote Γ as a distance-regular graph, we use the following notation: Γ has diameter D , valency k , and distinct eigenvalues $\theta_0 > \theta_1 > \dots > \theta_D$. Moreover, the intersection numbers of Γ are denoted by $\{c_i\}_{i=1}^D$, $\{a_i\}_{i=0}^D$, $\{b_i\}_{i=0}^{D-1}$, as shown in (2).

Next, we recall the notion of the i -homogeneous property as introduced by Nomura [19]. Let Γ be a connected graph. We say Γ has the *i -homogeneous* property if, for every pair of vertices x and y at distance i , the partition of the vertex set of Γ according to the path-length distance to both x and y is equitable, and the parameters corresponding to equitable partitions are independent of the choice of x and y . Graphs with the i -homogeneous property are simply said to be i -homogeneous. Note that Γ is 0-homogeneous if and only if it is a distance-regular graph. Moreover, if Γ is 1-homogeneous, then it is a distance-regular graph and also locally strongly regular.

In this paper, we focus on 1-homogeneous distance-regular graphs. Examples of such graphs include the Johnson graphs $J(2D, D)$, the bipartite distance-regular graphs, and the regular near $2D$ -gons. Over the years, the 1-homogeneous property has received considerable attention and has been used in the study of distance-regular graphs across various contexts, including tight distance-regular graphs [12, 15], distance-regular graphs which support a spin model [3], the Terwilliger algebras [4], and Q -polynomial distance-regular graphs [17]. Jurišić and Koolen [7, 9, 11] explored 1-homogeneous graphs whose μ -graphs are a complete multipartite graph $K_{t \times n}$, $n \geq 1$ (i.e. the complement of t copies of the complete graph K_n). In [7], they introduced the CAB property to study the local structures of distance-regular graphs and used this

property to characterize 1-homogeneous graphs with $a_1 \neq 0$. Also, they classified 1-homogeneous graphs with $c_2 \geq 2$ whose μ -graphs are $K_{t \times 1}$, i.e., 1-homogeneous Terwilliger graphs. In [9], they classified 1-homogeneous graphs when $n = 2$, i.e., when the μ -graphs are $K_{t \times 2}$ (Cocktail Party graphs). In their subsequent study [10], they extended this work to distance-regular graphs whose μ -graphs are $K_{t \times n}$ ($n \geq 2$). Jurišić, Munemasa, and Tagami [13] investigated a more general case, namely, graphs (not necessarily distance-regular) whose μ -graphs are $K_{t \times n}$. Moreover, several studies examined distance-regular graphs whose μ -graphs are complete multipartite; see [8, 11, 15]. These studies have contributed to the research on classifying 1-homogeneous distance-regular graphs with complete multipartite μ -graphs, which is an important problem.

A significance of our result in this paper lies in providing a classification of 1-homogeneous distance-regular graphs with $a_1 \geq 1$. This classification extends to a broader context, covering 1-homogeneous graphs with complete multipartite μ -graphs discussed in the preceding paragraph. We now present the main result of this paper.

Theorem 1.2. *Let Γ be a 1-homogeneous distance-regular graph with diameter $D \geq 5$ and $a_1 \geq 1$. Define $b = b_1/(\theta_1 + 1)$. Then, either $c_2 = 1$, or $b \geq 1$ and one of the following holds:*

- (i) Γ is a regular near $2D$ -gon.
- (ii) Γ is a Johnson graph $J(2D, D)$.
- (iii) Γ is a halved ℓ -cube where $\ell \in \{2D, 2D + 1\}$.
- (iv) Γ is a folded Johnson graph $\bar{J}(4D, 2D)$.
- (v) Γ is a folded halved $(4D)$ -cube.
- (vi) The valency k of Γ is bounded by a function $F(b)$ of b , i.e., $k \leq F(b)$, where

$$F(b) = 16b^{10} + 80b^9 + 192b^8 + 256b^7 + 192b^6 + 72b^5 + 20b^4 + 24b^3 + 8b^2 + 1. \quad (3)$$

The proof of this theorem appears in Section 5.

Remark 1.3. (i) In Theorem 1.2, in the case where Γ is a regular near $2D$ -gon, we can further refine it under the following conditions: if $c_2 \geq 3$, then Γ is a dual polar graph; if $c_2 = 2$ and $c_3 = 3$, then Γ is a Hamming graph; cf. [5, Theorem 9.11].

(ii) In [15], Koolen et al. proposed a conjecture stating that for a tight distance-regular graph with $D \geq 3$ and $b = b_1/(1 + \theta_1) \geq 2$, the diameter D is bounded by a function of b [15, Conjecture 7.5]. We prove this conjecture in Section 7 using Theorem 1.2.

This paper is organized as follows. In Section 2, we review strongly regular graphs and their properties. We discuss a classification of strongly regular graphs with smallest eigenvalue ≤ -2 . In Section 3, we discuss distance-regular graphs that are locally strongly regular. We establish a bound on the intersection number c_2 for those graphs. We also show that when a distance-regular graph is locally a conference graph, it is a Taylor graph. In Section 4, we recall the CAB_i property of distance-regular graphs. We focus on the CAB_2 property and discuss cases where distance-regular graphs with the CAB_2 property are locally a Latin square

graph or a Steiner graph. In Section 5, we discuss 1-homogeneous distance-regular graphs with $a_1 \geq 1$. We prove our main result, Theorem 1.2. In Section 6, we give a classification of 1-homogeneous distance-regular graphs with $a_1 \geq 1$ that have classical parameters. Finally, we conclude the paper in Section 7 by providing comments on tight distance-regular graphs.

2 Strongly regular graphs with smallest eigenvalue $-m$

In this section, we review the properties of strongly regular graphs with smallest eigenvalue $-m$ ($m > 0$) and discuss their classification. We begin by recalling the definition of strongly regular graphs. Let Γ be a k -regular graph with v vertices. The graph Γ is called *strongly regular* with parameters (v, k, λ, μ) if each pair of distinct adjacent (resp. non-adjacent) vertices has exactly λ (resp. μ) common neighbors. Suppose Γ is a strongly regular graph with smallest eigenvalue s . It is well known that Γ satisfies $s \leq -2$, except in the cases where Γ is a disjoint union of cliques (with $s = -1$) or a pentagon (with $s = (-1 - \sqrt{5})/2$) [2, Section 1.1.10].

Let Γ be a strongly regular graph with parameters (v, k, λ, μ) and diameter two. We denote the eigenvalues of Γ as $k > r > s$. Note that k, r, s are integers except when Γ is a *conference* graph, i.e., a strongly regular graph with parameters $(4\mu + 1, 2\mu, \mu - 1, \mu)$ [6, Lemma 10.3.3]. The parameters v, k, λ of Γ can be expressed in terms of r, s , and μ as follows:

$$v = \frac{(k-r)(k-s)}{\mu}, \quad k = \mu - rs, \quad \lambda = \mu + r + s, \quad (4)$$

cf. [1, Theorem 1.3.1].

Example 2.1. A transversal design $TD(m; n)$ is a partial linear space with mn points and $m + n^2$ lines, with m lines (called groups) of size n forming a partition of the point set, and n^2 lines (called blocks) of size m , each meeting every group in a single point; cf. [2, Section 8.4.1]. The line graph of a transversal design $TD(m; n)$ with $2 \leq m \leq n$ is called a *Latin square* graph $LS_m(n)$; cf. [2, Section 8.4.2]. Note that $LS_m(n)$ is equivalent to the block graph of an orthogonal array $OA(m, n)$. A Latin square graph $LS_m(n)$ is strongly regular with parameters

$$(n^2, m(n-1), (m-1)(m-2) + n - 2, m(m-1)) \quad (5)$$

and eigenvalues $m(n-1) > n - m > -m$.

Example 2.2. A Steiner system $S(2, m, n)$ is a $2-(n, m, 1)$ design, that is, a collection of m -subsets of a n -set in which each pair of elements is contained in exactly one m -set. In this context, the elements of the n -set are referred to as points, and the m -sets are referred to as blocks of the system. The *block graph of a Steiner system* $S(2, m, n)$ is defined as the graph whose vertices are the blocks of the system, where two vertices are adjacent whenever they intersect at exactly one point. The block graph of a Steiner system $S(2, m, n)$ with $n > m \geq 2$ is strongly regular with parameters

$$\left(\frac{n(n-1)}{m(m-1)}, \frac{m(n-m)}{m-1}, (m-1)^2 + \frac{n-1}{m-1} - 2, m^2 \right). \quad (6)$$

The eigenvalues of this graph are $\frac{m(n-m)}{m-1} > \frac{n-m^2}{m-1} > -m$. In particular, the block graph of a Steiner system $S(2, m, mn + m - n)$ is called a *Steiner graph* $S_m(n)$. Note that a Steiner graph $S_m(n)$ has parameters

$$\left(\frac{(m + n(m-1))(n+1)}{m}, mn, m^2 - 2m + n, m^2 \right) \quad (7)$$

and eigenvalues $mn > n - m > -m$.

Next, we recall some known results concerning a classification of strongly regular graphs with smallest eigenvalue ≤ -2 . A strongly regular graph Γ is called *primitive* if both Γ and its complement are connected. For primitive strongly regular graphs with integral smallest eigenvalue $-m$, the parameter μ is bounded as a function of m .

Lemma 2.3 (cf. [18, Theorem 3.1]). *A primitive strongly regular graph with integral smallest eigenvalue $-m$, where $m \geq 2$ satisfies*

$$\mu \leq m^3(2m - 3). \quad (8)$$

We call (8) the μ -bound.

For a strongly regular graph with integral eigenvalues $k > r > s$, where $r = n - m$ and $s = -m$, the following lemma provides a bound on n in terms of the parameters μ and m , which is known as the *claw bound*.

Lemma 2.4 (cf. [18, Theorem 4.7], [2, Theorem 8.6.3]). *Let Γ be a primitive strongly regular graph with integral eigenvalues $r = n - m$ and $s = -m$ ($m \geq 2$). Let $f(m, \mu) = \frac{1}{2}m(m-1)(\mu+1) + m - 1$. Then*

- (i) *If $\mu = m(m-1)$ and $n > f(m, \mu)$, then Γ is a Latin square graph $LS_m(n)$.*
- (ii) *If $\mu = m^2$ and $n > f(m, \mu)$, then Γ is a Steiner graph $S_m(n)$.*
- (iii) (Claw bound) *If $\mu \neq m(m-1)$ and $\mu \neq m^2$, then $n \leq f(m, \mu)$.*

As a consequence of the μ -bound and the claw bound, the strongly regular graphs with integral smallest eigenvalue ≤ -2 are characterized as follows:

Lemma 2.5 (Sims, cf. [2, Theorem 8.6.4]). *The strongly regular graphs with integral smallest eigenvalue $-m$, where $m \geq 2$, are one of the following (i)–(iv) below.*

- (i) *complete multipartite graphs with classes of size m ;*
- (ii) *Latin square graphs $LS_m(n)$;*
- (iii) *Steiner graphs $S_m(n)$;*
- (iv) *finitely many further graphs.*

Remark 2.6. By applying the claw bound, we can replace (iv) in Lemma 2.5 with the following:

$$n \leq \frac{1}{2}m(m-1)(\mu+1) + m - 1. \quad (9)$$

We give a comment on Lemma 2.5. Let Γ be a strongly regular graph with integral smallest eigenvalue $-m$, where $m \geq 2$. Assume that Γ is none of a complete multipartite graph, a Latin square graph, or a Steiner graph. By Lemma 2.5 and Remark 2.6, Γ satisfies $n \leq \frac{1}{2}m(m-1)(\mu+1) + m - 1$. Since $n = r + m$, it follows

$$r \leq \frac{1}{2}m(m-1)(\mu+1) - 1. \quad (10)$$

Note that $n \neq m$ since Γ is not a complete multipartite graph. This implies $r \neq 0$, that is, $r \geq 1$. From the first equation in (4), we have $v = (k-r)(k+m)/\mu$. Substituting k with $\mu + rm$ and applying (10) to the result, we obtain

$$v = \mu + m - r + 2rm + \frac{rm(1-m)(1+r)}{\mu}. \quad (11)$$

Applying inequalities (8) and (10) to the right-hand side of (11) and expressing the result in terms of m , we obtain

$$v \leq m^3(2m-3) + m + (2m-1 + m^2(m-1)^2) \left(\frac{m(m-1)}{2} (m^3(2m-3) + 1) - 1 \right). \quad (12)$$

We abbreviate the right-hand side of (12) by $\varphi(m)$ and simplify it to have

$$\varphi(m) = m^{10} - \frac{9}{2}m^9 + \frac{15}{2}m^8 - \frac{7}{2}m^7 - 4m^6 + 4m^5 + m^4 - \frac{1}{2}m^3 - \frac{5}{2}m^2 - \frac{1}{2}m + 1. \quad (13)$$

Note that $\varphi(m) < m^{10}$ for all $m \geq 2$. By these comments, we restate Lemma 2.5 as follows:

Corollary 2.7. *Let Γ be a strongly regular graph with parameters (v, k, λ, μ) with integral smallest eigenvalue $-m$, where $m \geq 2$. Then one of the following holds:*

- (i) Γ is a complete multipartite graph with class of size m ,
- (ii) Γ is a Latin square graph $LS_m(n)$,
- (iii) Γ is a Steiner graph $S_m(n)$,
- (iv) The number of vertices of Γ is bounded by a function in m , i.e., $v \leq \varphi(m)$, where $\varphi(m)$ is from (13).

3 Distance-regular graphs that are locally strongly regular

In this section, we discuss distance-regular graphs whose local graphs are strongly regular and give a bound on their intersection number c_2 . Furthermore, we show that if a distance-regular graph is locally a conference graph, then it is a Taylor graph. We first recall some known results concerning distance-regular graphs that we shall need, along with references for further discussion.

Lemma 3.1 (cf. [1, Theorem 4.4.3]). *Let Γ be a distance-regular graph of diameter $D \geq 3$ with eigenvalues $k = \theta_0 > \theta_1 > \dots > \theta_D$ and intersection number b_1 . Let $b = b_1/(\theta_1 + 1)$. Then, $b > 0$. Moreover, for each vertex x in Γ , its local graph has the smallest eigenvalue $\geq -1 - b$.*

Let Γ be a graph with diameter at least two. The subgraph of Γ induced by the set of common neighbors of two vertices in Γ at distance two is called the c_2 -graph (or μ -graph) of Γ . If each c_2 -graph is a regular graph with valency κ , then we say that Γ is c_2 -graph-regular with parameter κ . The graph Γ is called a *Terwilliger graph* when it is c_2 -graph-regular and every c_2 -graph of Γ is complete.

Lemma 3.2 (cf. [8, Theorem 3.1]). *Let Γ be a distance-regular graph which is locally strongly regular with parameters (v', k', λ', μ') . Then Γ is c_2 -graph-regular with parameter μ' . Moreover, $c_2 \geq \mu' + 1$, with equality if and only if Γ is a Terwilliger graph.*

A *clique* in Γ is a subset of $V(\Gamma)$ such that every pair of distinct vertices is adjacent. A clique of size p is referred to as a complete graph K_p . A *coclique* of Γ is a subset of $V(\Gamma)$ such that no two vertices are adjacent. A *complete bipartite graph* $K_{p,q}$ is a graph whose vertex set can be partitioned into two cocliques, say a p -set V_1 and a q -set V_2 , where each vertex in V_1 is adjacent to all vertices in V_2 . A *complete multipartite graph* $K_{t \times p}$ is a graph whose vertex set can be partitioned into cocliques $\{V_i\}_{i=1}^t$ of size p , where each vertex in V_i is adjacent to all vertices in V_j ($1 \leq j \neq i \leq t$).

Lemma 3.3 (cf. [20, Lemma 3.7]). *Let Γ be a distance-regular graph with valency k , diameter $D \geq 5$ and second largest eigenvalue θ_1 . Assume that Γ contains an induced subgraph $K_{2,t}$ for some $t \geq 2$. Let $b = b_1/(\theta_1 + 1)$. Then $t \leq 4b^2 + 1$.*

We now establish a bound on the intersection number c_2 for distance-regular graphs that are locally strongly regular.

Proposition 3.4. *Let Γ be a distance-regular graph with diameter $D \geq 5$, valency k , intersection numbers b_1 and c_2 , and second largest eigenvalue θ_1 . Assume Γ is locally strongly regular with parameters (v', k', λ', μ') . Let $b = b_1/(\theta_1 + 1)$. Then*

$$c_2 \leq (4b^2 + 1)(\mu' + 1). \quad (14)$$

Proof. We assume $c_2 \geq 2$; otherwise, it is trivial. Since $c_2 \geq 2$, Γ contains an induced subgraph $K_{2,t}$ for some $t \geq 2$. Applying Lemma 3.3 to Γ , we obtain the bound $t \leq 4b^2 + 1$. Next, as Γ is locally strongly regular, by Lemma 3.2, Γ is c_2 -graph-regular with parameter μ' . Thus, we find that the number of vertices of the c_2 -graph is at most $t(\mu' + 1)$. By these comments, it follows that $c_2 \leq t(\mu' + 1) \leq (4b^2 + 1)(\mu' + 1)$. ■

Recall that a conference graph is a strongly regular graph with parameters $(4\mu + 1, 2\mu, \mu - 1, \mu)$, where $\mu > 0$. We discuss distance-regular graphs that are locally a conference graph. First, we recall a classification of distance-regular graphs that satisfy $a_1 \geq k/2 - 1$, as shown by Koolen and Park [16].

Lemma 3.5 ([16, Theorem 16]). *Let Γ be a distance-regular graph with diameter $D \geq 3$ and valency k . If $a_1 \geq \frac{1}{2}k - 1$, then one of the following holds:*

- (i) Γ is a polygon,
- (ii) Γ is the line graph of a Moore graph,
- (iii) Γ is the flag graph of a regular generalized D -gon of order (s, s) for some s ,
- (iv) Γ is a Taylor graph,
- (v) Γ is the Johnson graph $J(7, 3)$,
- (vi) Γ is the halved 7-cube.

Proposition 3.6. *Let Γ be a distance-regular graph with diameter $D \geq 3$. If Γ is locally a conference graph, then it is a Taylor graph.*

Proof. Let Δ denote a local graph of Γ . Then Δ is a conference graph, and we denote its parameters by $(4\mu + 1, 2\mu, \mu - 1, \mu)$, where $\mu > 0$. Thus, Γ has valency $k = 4\mu + 1$ and intersection number $a_1 = 2\mu$. We observe that k is odd, a_1 is nonzero and even, and Γ satisfies $a_1 > \frac{1}{2}k - 1$. By Lemma 3.5, Γ falls into one of the graphs (i)–(vi) listed therein. However, we observe that:

- In case (i), a polygon has $k = 2$ and $a_1 = 0$;
- In case (ii), the line graph of a Moore graph has the intersection array $\{2k' - 2, k' - 1, k' - 2; 1, 1, 4\}$, where $k' \in \{3, 7, 57\}$, implying that $a_1 = k'$ (an odd number);
- In case (iii), the flag graph of a regular generalized D -gon of order (s, s) , a generalized $2D$ -gon of order $(s, 1)$, has $k = 2s$ (an even number);
- In case (v), the Johnson graph $J(7, 3)$ has the intersection array $\{12, 6, 2; 1, 4, 9\}$, so $k = 12$ (even) and $a_1 = 5$ (odd).

From these observations, it follows that none of the graphs (i), (ii), (iii), (v) is locally a conference graph. Moreover, as the halved 7-cube has the intersection array $\{21, 10, 3; 1, 6, 15\}$ its local graph has 21 vertices and valency 10, but such a conference graph does not exist. Therefore, Γ is a Taylor graph. ■

We finish this section with a comment. The line graph of $K_{p,q}$ is called the $(p \times q)$ -grid. Note that the square $(p \times p)$ -grid, where $p \geq 2$, is a strongly regular graph and is equivalent to the Latin square graph $LS_2(p)$.

Lemma 3.7 (cf. [9, Proposition 4.1]). *Let Γ be a distance-regular graph with diameter $D \geq 2$. If Γ is locally the $(p \times q)$ -grid and $c_2 = 4$, then Γ is the Johnson graph $J(p + q, p)$, or $p = q$ and Γ is the folded Johnson graph $\bar{J}(2p, p)$.*

4 The CAB_i property

In this section, we review the CAB_i property of a distance-regular graph. We then discuss distance-regular graphs with the CAB_2 property, which are locally a Latin square graph or a Steiner graph. We begin by recalling the notion of an *equitable partition*. Let Γ be a connected graph. A partition $\pi = \{C_1, C_2, \dots, C_p\}$ of the vertex set $V(\Gamma)$ is called *equitable* if, for all $1 \leq i, j \leq p$ and for each $x \in C_i$, the number of neighbors of x in C_j is independent of the choice of the vertex x . In other words, for any pair of subsets C_i and C_j in π , the number $c_{ij} := |\Gamma(x) \cap C_j|$ is constant for all $x \in C_i$. These numbers $\{c_{ij}\}_{1 \leq i, j \leq p}$ are called the *parameters* of π . Suppose π is an equitable partition of Γ . The *quotient matrix* $Q = Q(\pi)$ associated with π is the $p \times p$ matrix defined by $Q_{ij} := c_{ij}$ for $1 \leq i, j \leq p$.

We now recall the notion of the CAB_i property. Let Γ be a distance-regular graph with diameter $D \geq 2$ and $a_1 \geq 1$. Note that every local graph of Γ has $k = b_0$ vertices and is regular with valency a_1 . For $0 \leq i \leq D$ and for $x, y \in V(\Gamma)$ with $d(x, y) = i$, we recall the subsets $C_i(x, y)$, $A_i(x, y)$, and $B_i(x, y)$ of $V(\Gamma)$ from (1). Additionally, we recall the intersection numbers c_i , a_i , and b_i , as defined in (2). Let $\Delta(y)$




Figure 1: The CAB_i partition and its parameters for $1 \leq i \leq D - 1$

denote the local graph of Γ at y . We observe that the set $\{C_i(x, y), A_i(x, y), B_i(x, y)\}$ partitions the vertex set of $\Delta(y)$. We denote this partition by $CAB_i(x, y)$ and call it the $CAB_i(x, y)$ partition of $\Delta(y)$. Observe that the $CAB_0(x, y)$ partition is $\Gamma(x)$, and the $CAB_D(x, y)$ partition is $\{C_D(x, y), A_D(x, y)\}$ if $a_D \neq 0$ and $\{C_D(x, y)\}$ if $a_D = 0$. For $1 \leq j \leq D$, we say that Γ has the CAB_j property if, for each $i \leq j$ and for every pair of vertices $x, y \in V(\Gamma)$ with $d(x, y) = i$, the partition $CAB_i = CAB_i(x, y)$ is equitable, and the parameters of the equitable partition CAB_i do not depend on the choice of vertices x and y . If Γ has the CAB_D property, then we simply say that it has the CAB property. We note that Γ has the CAB_1 property if and only if it is locally strongly regular. We say the (triple) intersection number γ of Γ exists if, for every triple of vertices (x, y, z) of Γ such that x and y are adjacent and z is at distance 2 from both x and y , the number of common neighbors of x, y , and z is constant and equal to γ . To avoid the degenerate case, we assume that there exists at least one such triple (x, y, z) in Γ (i.e., $a_2 \neq 0$) when we say γ exists. We note that if Γ has the CAB_2 property and $a_2 \neq 0$, then the intersection number γ exists in Γ , which is equal to the parameter α_2 (see Figure 1).

Lemma 4.1 (cf. [7, Proposition 2.1]). *Let Γ be a distance regular graph with $a_1 \geq 1$ and the CAB_1 property. Then all local graphs of Γ are either*

- (i) *connected strongly regular graphs with the same parameters, or*
- (ii) *disjoint unions of $(a_1 + 1)$ -cliques.*

We recall a regular near polygon. Let Γ be a distance-regular graph with diameter D . For integers s and t , we say that Γ is of order (s, t) if it is locally the disjoint union of $t + 1$ cliques of size s . The graph Γ of order (s, t) is called a *regular near polygon* if $a_i = c_i a_1$ for all $1 \leq i \leq D - 1$. If $a_D = c_D a_1$, we call Γ a *regular near $2D$ -gon*; otherwise it is called a *regular near $(2D + 1)$ -gon*.

Lemma 4.2 (cf. [5, Theorem 9.11]). *Let Γ be a regular near $2D$ -gon with $D \geq 4$. If $c_2 \geq 3$ or $c_i = i$ ($i = 2, 3$), then Γ is either a dual polar graph or a Hamming graph.*

Lemma 4.3 (cf. [7, Theorem 2.3]). *Let Γ be a locally disconnected distance-regular graph with diameter $D \geq 2$ and $a_1 \geq 1$. If Γ has the CAB property, then it is a regular near $2D$ -gon.*

Let Γ be a distance-regular graph with diameter $D \geq 2$ and $a_1 \geq 1$. Suppose Γ has the CAB_j property for some $1 \leq j \leq D$. For each $1 \leq i \leq j, i \neq D$, the quotient matrix Q_i associated with the CAB_i partition (see Figure 1) is

$$Q_i = \begin{pmatrix} \gamma_i & a_1 - \gamma_i & 0 \\ \alpha_i & a_1 - \beta_i - \alpha_i & \beta_i \\ 0 & \delta_i & a_1 - \delta_i \end{pmatrix} \quad (1 \leq i \leq j, i \neq D); \quad (15)$$

cf. [7, Theorem 2.4]. Note that Γ is locally strongly regular with the same parameters since Γ has the CAB_1 property. If Γ is locally connected, then the eigenvalues of Q_i are the same as the eigenvalues a_1, r, s of a local graph of Γ ; cf. [7, Lemma 2.6].

Lemma 4.4 (cf. [7, Theorem 2.7]). *Let Γ be a locally connected distance-regular graph with $D \geq 2$, $a_1 \geq 2$, and the CAB_j property for some $1 \leq j < D$. Let $\delta_0 = 0$. Then, for $1 \leq i \leq j$ the parameters $a_i, b_i, \alpha_i, \beta_i, \gamma_i, \delta_i$ of the CAB_i partition are expressed in terms of the eigenvalues $a_1 > r > s$ of a local graph of Γ and the parameters δ_{i-1} and c_i as follows:*

$$\gamma_i = \delta_{i-1}, \quad \alpha_i = \frac{c_i(a_1 - \delta_{i-1})}{k - c_i - b_i}, \quad \beta_i = \frac{\mu' b_i}{a_1 - \delta_{i-1}}, \quad \delta_i = \frac{\mu'(k - c_i)}{a_1 - \delta_{i-1}} - \beta_i, \quad (16)$$

$$b_i = k - c_i - \frac{c_i(a_1 - \delta_{i-1})^2}{(a_1 - \delta_{i-1})(a_1 + r + s - \delta_{i-1}) - \mu'(k - c_i)}, \quad (17)$$

and $a_i = k - b_i - c_i$, where $k = (a_1 - r)(a_1 - s)/\mu'$ and $\mu' = a_1 + rs$.

Proof. For (16), refer to the proof of [7, Theorem 2.7]. We will show (17) only. Observe that Γ has the CAB_1 property, so it is locally a strongly regular graph with parameters (k, a_1, λ', μ') and eigenvalues $a_1 > r > s$. Consider the quotient matrix Q_i associated with the CAB_i partition for $1 \leq i \leq j < D$. From (15), we have the trace $\text{tr}(Q_i) = 2a_1 - \alpha_i - \beta_i - \delta_i + \gamma_i$. Moreover, since the eigenvalues of Q_i equal the eigenvalues of a local graph of Γ , it follows that $\text{tr}(Q_i) = a_1 + r + s$. From these two equations for the trace of Q_i , we obtain

$$a_1 + r + s = \alpha_i + \beta_i + \delta_i - \gamma_i. \quad (18)$$

Eliminate $\alpha_i, \beta_i, \delta_i, \gamma_i$ in (18) using (16) and solve for b_i . Simplify the result to obtain (17). \blacksquare

Now, we discuss distance-regular graphs satisfying the CAB_2 property. Let Γ be a locally connected distance-regular graph with diameter $D \geq 2$ and $a_1 \geq 2$. We assume that Γ satisfies the CAB_2 property. By Lemma 4.1, every local graph of Γ is a connected strongly regular graph with the same parameters (v', k', λ', μ') and eigenvalues $a_1 > r > s$. Note that $v' = k$ and $k' = a_1$. Consider the CAB_2 partition. Evaluating (16), (17) at $i = 2$, the parameters of the CAB_2 partition are

$$\gamma_2 = \delta_1 = \mu', \quad \alpha_2 = \frac{c_2(a_1 - \mu')}{k - c_2 - b_2}, \quad \beta_2 = \frac{\mu' b_2}{a_1 - \mu'}, \quad \delta_2 = \frac{\mu'(k - c_2 - b_2)}{a_1 - \mu'}, \quad (19)$$

and

$$b_2 = k - c_2 - \frac{c_2(a_1 - \mu')^2}{(a_1 + 2\mu' - \lambda')(a_1 - \mu') - \mu'(k - c_2)}. \quad (20)$$

With reference to the above discussion, we will express the parameters of the CAB_2 partition in terms of $m = -s$ and $n = r - s$ when Γ is locally a Latin square graph or a Steiner graph. For our convenience, we will use the following notation.

Notation 4.5. Let $b \geq 1$. We define the polynomial G in the variable b as follows:

$$G(b) = 16b^{10} + 32b^9 + 56b^8 + 48b^7 + 17b^6 + 2b^5 - 11b^4 + 6b^3 + 6b^2 - 4b + 1. \quad (21)$$

Recall $F(b)$ from (3). Note that $G(b) < F(b)$ for all $b \geq 1$.

Recall the intersection number b_1 and the second largest eigenvalue θ_1 of Γ .

Lemma 4.6. *Let Γ be a distance-regular graph with diameter $D \geq 5$, valency k , $a_1 \geq 1$, and satisfying the CAB_2 property. Let $b = b_1/(\theta_1 + 1)$. Assume that $k > G(b)$, where $G(b)$ is from (21). If Γ is locally a Latin square graph $LS_m(n)$ with $m \geq 2$, then we have*

$$\alpha_2 = m, \quad \beta_2 = (m-1)(n-m^2+m), \quad \delta_2 = m^2(m-1), \quad (22)$$

$$a_2 = m^2(n-m), \quad b_2 = (n-m)(n-m^2+m), \quad c_2 = m^2. \quad (23)$$

Proof. From (5), we observe that the parameters (k, a_1, λ', μ') of $LS_m(n)$ are

$$k = n^2, \quad a_1 = m(n-1), \quad \lambda' = (m-1)(m-2) + n - 2, \quad \mu' = m(m-1). \quad (24)$$

First, we find the parameters α_2 and c_2 . In equation (18) at $i = 2$, eliminate β_2 and δ_2 using (19), then solve the resulting expression for α_2 to obtain $\alpha_2 = a_1 - \lambda' + 2\mu' - \mu'(k - c_2)/(a_1 - \mu')$. In this equation, eliminate the parameters k, a_1, λ', μ' using (24) and express the result in terms of n, m, c_2 to have

$$\alpha_2 = m + \frac{(m-1)(c_2 - m^2)}{n - m}. \quad (25)$$

Since α_2 is an integer, we have either $(m-1)(c_2 - m^2) = 0$ or $n - m$ divides $(m-1)(c_2 - m^2)$. Suppose $(m-1)(c_2 - m^2) \neq 0$, which implies $c_2 \neq m^2$ for $m \geq 2$. Then we have

$$n - m \leq (m-1)(c_2 - m^2), \quad (26)$$

Apply inequality (14) to c_2 in (26) and substitute $\mu' = m(m-1)$ into the result. Then, apply the condition $2 \leq m \leq 1 + b$ to the result and simplify it to obtain

$$n \leq 4b^5 + 4b^4 + 5b^3 + b^2 - 2b + 1. \quad (27)$$

Since $k = n^2$, squaring both sides of (27) implies $k \leq (4b^5 + 4b^4 + 5b^3 + b^2 - 2b + 1)^2 = G(b)$. This contradicts the assumption $k > G(b)$. Therefore, we have $c_2 = m^2$, and consequently, $\alpha_2 = m$.

Next, we find β_2 . In the third equation in (19), eliminate b_2 using (20) to obtain

$$\beta_2 = \frac{\mu'}{a_1 - \mu'} \left(k - c_2 - \frac{c_2(a_1 - \mu')^2}{(a_1 + 2\mu' - \lambda')(a_1 - \mu') - \mu'(k - c_2)} \right) \quad (28)$$

Express the right-hand side of (28) in terms of m and n using (24) and $c_2 = m^2$. Simplify the result to obtain $\beta_2 = (m-1)(n-m^2+m)$. To find b_2 , evaluate the right-hand side of (20) using (24) and $c_2 = m^2$. Simplify the result to obtain $b_2 = (n-m)(n-m^2+m)$. For δ_2 , solve the fourth equation in (19) using (24) and $b_2 = (n-m)(n-m^2+m)$ to obtain $\delta_2 = m^2(m-1)$. Finally, for a_2 , evaluate $k - b_2 - c_2$ using $k = n^2$, $b_2 = (n-m)(n-m^2+m)$, $c_2 = m^2$ to get $a_2 = m^2(n-m)$. The proof is now complete. \blacksquare

Lemma 4.7. *Let Γ be a distance-regular graph with diameter $D \geq 5$, valency k , $a_1 \geq 1$, and satisfying the CAB_2 property. Let $b = b_1/(\theta_1 + 1)$. Assume that $k > F(b)$, where $F(b)$ is from (3). If Γ is locally a Steiner graph $S_m(n)$ with $m \geq 2$, then we have*

$$\alpha_2 = m + 1, \quad \beta_2 = (m-1)(n-m^2+1), \quad \delta_2 = m^3, \quad (29)$$

$$a_2 = m^2(n-m), \quad b_2 = (m-1)(n-m)(n-m^2+1)/m, \quad c_2 = m(m+1). \quad (30)$$

Proof. From (7), we observe that the parameters (k, a_1, λ', μ') of $S_m(n)$ are

$$k = \frac{(m+n(m-1))(n+1)}{m}, \quad a_1 = mn, \quad \lambda' = m^2 - 2m + n, \quad \mu' = m^2. \quad (31)$$

In a similar manner to the proof of Lemma 4.6, we have

$$\alpha_2 = m + 1 + \frac{m(c_2 - m(m+1))}{n - m}. \quad (32)$$

Since α_2 is an integer, we have either $m(c_2 - m(m+1)) = 0$ or $n - m$ divides $m(c_2 - m(m+1))$. If $c_2 \neq m(m+1)$, then

$$n - m \leq m(c_2 - m(m+1)). \quad (33)$$

Apply the inequalities (33) and (14), along with the substitution $\mu' = m(m-1)$ and the condition $2 \leq m \leq 1 + b$, to the first equation in (31). Simplify the result to obtain

$$k = \frac{(m+n(m-1))(n+1)}{m} \leq F(b). \quad (34)$$

This contradicts the assumption $k > F(b)$. Therefore, $c_2 = m(m+1)$, and consequently, $\alpha_2 = m + 1$. The remaining parameters in (29) and (30) are obtained similarly to Lemma 4.6. The proof is complete. \blacksquare

Remark 4.8. (i) If Γ is locally a Latin square graph with smallest eigenvalue -2 (i.e., locally a square $(n \times n)$ -grid with $n \geq 3$), then by Lemma 4.6, we observe that the c_2 -graph of Γ is a quadrangle.

(ii) If Γ is locally a block graph of a Steiner $2-(v, 2, 1)$ design with smallest eigenvalue -2 (i.e., locally a triangular graph $T(n)$ with $n \geq 5$), then by Lemma 4.7, we observe that the c_2 -graph of Γ is an octahedron.

5 The 1-homogeneous distance-regular graphs

In this section, we discuss distance-regular graphs that have the 1-homogeneous property. We then prove our main result, Theorem 1.2, which classifies these graphs. We begin by recalling some known results regarding these graphs.

Let Γ be a connected graph with diameter D . For $0 \leq i \leq D$, fix a pair of vertices x and y in Γ with $d(x, y) = i$. For $0 \leq h, j \leq D$, we define the subset $D_j^h(x, y)$ of $V(\Gamma)$ by

$$D_j^h(x, y) := \Gamma_j(x) \cap \Gamma_h(y). \quad (35)$$

Let $\pi(x, y)$ denote the collection of nonempty sets $D_j^h(x, y)$ for $0 \leq h, j \leq D$. Observe that $\pi(x, y)$ forms a distance partition of $V(\Gamma)$. The graph Γ is said to be *i-homogeneous* whenever the partition $\pi(x, y)$ is equitable, and the parameters of $\pi(x, y)$ are independent of the choice of vertices x and y ; see [19]. Our discussion centers on the 1-homogeneous distance-regular graphs. We note that if Γ is 1-homogeneous distance-regular graph with $D \geq 2$ and $a_2 \neq 0$, then the intersection number γ exists. Jurišić and Koolen [7] studied these graphs in detail using the CAB property and characterized them as follows.

Lemma 5.1 (cf. [7, Theorem 3.1]). *Let Γ be a distance regular graph with diameter D and $a_1 \geq 1$. Then Γ is 1-homogeneous if and only if it has the CAB property.*

Consider the 1-homogeneous distance-regular graphs that are locally disconnected.

Corollary 5.2 (cf. [7, Corollary 3.3]). *A locally disconnected 1-homogeneous distance-regular graph with diameter $D \geq 2$ and $a_1 \geq 1$ is a regular near $2D$ -gon.*

Proof. By Lemmas 4.3 and 5.1. ■

Lemma 5.3 (cf. [5, Theorem 9.11]). *Let Γ be a locally disconnected 1-homogeneous distance-regular graph with $D \geq 4$ and $a_1 \geq 1$. If $c_2 \geq 3$ or $c_i = i$ ($i = 2, 3$), then Γ is either a dual polar graph or a Hamming graph.*

Proof. Immediate from Lemma 4.2 and Corollary 5.2. ■

We now discuss locally connected 1-homogeneous distance-regular graphs. There are some studies on 1-homogeneous distance-regular graphs whose c_2 -graphs are complete multipartite; cf. [9], [13]. In particular, Jurišić and Koolen [9] classified 1-homogeneous distance-regular graphs for which the c_2 -graphs are the Cocktail Party graph.

Theorem 5.4 ([9, Theorem 1.1]). *Let Γ be a 1-homogeneous distance-regular graph with diameter $D \geq 2$ which is locally strongly regular, and its c_2 -graph is a connected Cocktail Party graph. Then Γ is one of the following graphs:*

- (i) a Johnson graph $J(2D, D)$,
- (ii) a halved ℓ -cube where $\ell \in \{2D, 2D + 1\}$,
- (iii) a folded Johnson graph $\bar{J}(4D, 2D)$,
- (iv) a folded halved $(4D)$ -cube,
- (v) a Cocktail Party graph $K_{t \times 2}$ with $t \geq 3$.
- (vi) the Schläfli graph with intersection array $\{16, 5; 1, 8\}$,
- (vii) the Gosset graph with intersection array $\{27, 10, 1; 1, 10, 27\}$.

Recently, Koolen et al. [15] studied 1-homogeneous distance-regular graphs that are locally block graphs of Latin square graphs or Steiner systems. The main result they showed is as follows:

Lemma 5.5 (cf. [15, Theorem 1.2]). *Let Γ be a distance-regular graph with diameter $D \geq 3$ and valency k , satisfying the CAB_2 property.*

- (i) *If Γ is locally a Latin square graph with smallest eigenvalue $-m$ for $m \geq 3$ and $k > m^2$, then $c_2 \neq m^2$.*
- (ii) *If Γ is locally the block graph of a Steiner system $S(2, m, n)$ with smallest eigenvalue $-m$ for $m \geq 3$ and $k > m(m + 1)$, then $c_2 \neq m(m + 1)$.*

Remark 5.6. (i) Let Γ be a distance-regular graph that is locally a Latin square graph $LS_m(n)$ (resp. the block graph of a Steiner system $S(2, m, n)$). In [15], it is shown that if $c_2 = m^2$ (resp. $c_2 = m(m+1)$), then the c_2 -graph of Γ is a complete m -partite (resp. $(m+1)$ -partite) graph. This result, along with the findings of [13], is used to complete the proof of Lemma 5.5.

(ii) The icosahedron has $b_1 = 2$, $\theta_1 = \sqrt{5}$ and hence $b = \frac{b_1}{\theta_1+1} = \frac{1}{2}(-1 + \sqrt{5}) < 1$. The only locally connected distance-regular graph with the CAB_2 property that satisfies $b < 1$ is the icosahedron. This is because the only connected strongly regular graph with the smallest eigenvalue greater than -2 is the pentagon, and the only graph that is locally a pentagon is the icosahedron (cf. [1, p. 35]).

We are now ready to prove Theorem 1.2.

Proof of Theorem 1.2. We assume $c_2 \geq 2$. Our proof is divided into two cases: Γ is locally disconnected, and Γ is locally connected. First, we assume that Γ is locally disconnected. Since $c_2 \geq 2$, Γ contains an induced quadrangle. By the result of [21] (cf. [1, p. 170]), we obtain $\theta_1 \leq b_1 - 1$, which implies $b \geq 1$. Moreover, by Corollary 5.2 Γ is a regular near $2D$ -gon. Therefore, we have (i).

Next, we consider the case where Γ is locally connected. Since Γ satisfies the CAB_1 property, every local graph of Γ is connected strongly regular with the same parameters. Let $-m$ be the smallest eigenvalue of a local graph of Γ . Note that a connected strongly regular graph satisfies $-m < -1$ and the only strongly regular graph satisfying $-m > -2$ is a pentagon. Furthermore, the only graph that is locally pentagonal is the icosahedron. However, an icosahedron has diameter 3, which is less than 5. Therefore, we have $-m \leq -2$. By Lemma 3.1, it follows that $b \geq 1$.

Now, we show that one of the following (ii)–(vi) holds. We assume that $k > F(b)$; otherwise, we fall into the case (vi). Since Γ is locally a connected strongly regular graph with eigenvalues $a_1 > r > s$, it follows that either Γ is locally a conference graph or r and s are integers. If Γ is locally a conference graph, then it must be a Taylor graph by Proposition 3.6. However, Γ has $D \geq 5$ while a Taylor graph has diameter three. Therefore, a local graph of Γ is not a conference graph and hence has integral eigenvalues r and s . We put $s = -m$ and $r = n - m$, where $m \geq 2$. According to Corollary 2.7, a local graph of Γ satisfies one of (i)–(iv) listed therein. However, Case (iv) of Corollary 2.7 is ruled out as $b \geq m - 1 \geq 1$, and, by assumption, we have $k > F(b) > m^{10} > \varphi(m)$. Therefore, Γ is locally either a complete multipartite graph with classes of size m , a Latin square graph $LS_m(n)$, or a Steiner graph $S_m(n)$. Now, we consider each case.

First, consider the case where Γ is locally a complete multipartite graph with classes of size m , denoted as $K_{t \times m}$. By [1, Proposition 1.1.5], Γ is $K_{(t+1) \times m}$. This contradicts the assumption $D \geq 5$ since a complete multipartite graph has diameter two.

Next, consider the case where Γ is locally a Latin square graph $LS_m(n)$. Since $k > F(b) > G(b)$, by Lemma 4.6 we have $c_2 = m^2$, where $m \geq 2$. However, by Lemma 5.5(i), c_2 cannot be equal to m^2 for $m \geq 3$. By these comments, it follows $m = 2$. Thus, Γ is locally a Latin square graph $LS_2(n)$ with smallest eigenvalues -2 and $c_2 = 4$. Observe that $LS_2(n)$ is equivalent to the $(n \times n)$ -grid. By Lemma 3.7, Γ is the Johnson graph $J(2n, n)$ or a folded Johnson graph $\bar{J}(2n, n)$.

Lastly, consider the case where a local graph of Γ is a Steiner graph $S_m(n)$. Since $k > F(b)$, by Lemma 4.7 we have $c_2 = m(m+1)$, where $m \geq 2$. However, by Lemma 5.5(ii) c_2 cannot be equal to $m(m+1)$ for $m \geq 3$.

Consequently, it follows $m = 2$. Thus, Γ is locally a Steiner graph $S_2(n)$ with smallest eigenvalue -2 and $c_2 = 6$. Observe that the graph $S_n(2)$ is equivalent to a triangular graph $T(n+2)$ with $n \geq 3$. Thus, the c_2 -graph of Γ is a 4-regular graph on 6 vertices, that is a Cocktail Party graph $K_{3 \times 2}$. Hence, by Lemma 5.4 (also refer to the proof of [9, Theorem 5.3]) Γ is a halved ℓ -cube where $\ell \in \{2D, 2D+1\}$ or a folded halved $(4D)$ -cube. The proof is complete. \blacksquare

6 Distance-regular graphs with classical parameters

In this section, we discuss 1-homogeneous distance-regular graphs with classical parameters. For integers b and i , we recall the Gaussian binomial coefficient

$$\begin{bmatrix} i \\ 1 \end{bmatrix} = \begin{bmatrix} i \\ 1 \end{bmatrix}_b = \begin{cases} i & \text{if } b = 1; \\ (b^i - 1)/(b - 1) & \text{if } b \neq 1. \end{cases}$$

Let Γ be a distance-regular graph with diameter $D \geq 3$. We say that Γ has *classical parameters* (D, b, α, β) if its intersection numbers $\{b_i\}_{i=0}^{D-1}$ and $\{c_i\}_{i=1}^D$ satisfy

$$b_i = \left(\begin{bmatrix} D \\ 1 \end{bmatrix} - \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \left(\beta - \alpha \begin{bmatrix} i \\ 1 \end{bmatrix} \right), \quad c_i = \begin{bmatrix} i \\ 1 \end{bmatrix} \left(1 + \alpha \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \right). \quad (36)$$

Note that the parameter b is an integer, excluding 0 or -1 , since $b_2 \neq 0$ and $c_2 \neq 0$. Also, from $a_i + b_i + c_i = k = b_0$ ($0 \leq i \leq D$), it follows

$$a_i = \begin{bmatrix} i \\ 1 \end{bmatrix} \left(\beta - 1 + \alpha \left(\begin{bmatrix} D \\ 1 \end{bmatrix} - \begin{bmatrix} i \\ 1 \end{bmatrix} - \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \right) \right) \quad (0 \leq i \leq D). \quad (37)$$

Observe that $a_1 = 0$ if and only if $\beta = 1 - \alpha b \begin{bmatrix} D-1 \\ 1 \end{bmatrix}$. The eigenvalues of Γ are

$$\theta_i = \begin{bmatrix} D-i \\ 1 \end{bmatrix} \left(\beta - \alpha \begin{bmatrix} i \\ 1 \end{bmatrix} \right) - \begin{bmatrix} i \\ 1 \end{bmatrix} \quad (0 \leq i \leq D), \quad (38)$$

cf. [1, Corollary 8.4.2]. We note that the eigenvalues are in the natural ordering $\theta_0 > \theta_1 > \dots > \theta_D$ if $b > 0$. We recall a lower bound on the parameter β .

Lemma 6.1 ([14, Proposition 1]). *Let Γ be a distance-regular graph with classical parameters (D, b, α, β) and $D \geq 3$. If $b > 0$, then $\beta \geq 1 + \alpha \begin{bmatrix} D-1 \\ 1 \end{bmatrix}$, with equality if and only if $a_D = 0$.*

Now, we classify 1-homogeneous distance-regular graphs with classical parameters and $a_1 \geq 1$.

Theorem 6.2. *Let Γ be a 1-homogeneous distance-regular graph with classical parameters (D, b, α, β) and $a_1 \geq 1$, where $D \geq 5$, $b \geq 1$. Then one of the following holds:*

- (i) $\alpha = 0$,
- (ii) Γ is a Johnson graph $J(2D, D)$,
- (iii) Γ is a halved ℓ -cube where $\ell \in \{2D, 2D+1\}$,
- (iv) Γ is a folded Johnson graph $\bar{J}(4D, 2D)$,

(v) Γ is a folded halved $(4D)$ -cube,

(vi) $D \leq 9$, $\alpha > 0$, $b \geq 2$.

Proof. Let $\{\theta_i\}_{i=0}^D$ be the eigenvalues of Γ as in (38). Since $b \geq 1$, it follows $\theta_0 > \theta_1 > \dots > \theta_D$. We observe that $b = b_1/(\theta_1 + 1)$ by evaluating b_i in (36) and θ_i in (38) at $i = 1$. Moreover, $c_2 \geq 2$ since every distance regular graph with classical parameters and diameter at least three satisfies $c_i < c_{i+1}$ for $0 \leq i \leq D - 1$ (cf. [1, Theorem 6.1.2]). Therefore, by Theorem 1.2, Γ is either a regular near $2D$ -gon or falls into one of (ii)–(vi) therein.

We consider the parameter α . Evaluating c_i in (36) at $i = 2$ gives $c_2 = (1 + b)(1 + \alpha)$, which implies that $\alpha(1 + b)$ is an integer. Next, evaluating c_i in (36) at $i = 3$ gives $c_3 = (1 + b + b^2)(1 + \alpha(1 + b))$. Since $1 + b + b^2 > 0$ and $c_3 > 0$, it follows $1 + \alpha(1 + b) > 0$. As $\alpha(1 + b)$ is an integer, we have $\alpha(1 + b) \geq 0$. Since $1 + b > 0$, we obtain $\alpha \geq 0$.

We consider the case where $\alpha > 0$; otherwise, we obtain (i). Since $b \geq 1$, we divide the argument into two cases: either $b = 1$ or $b \geq 2$. If $b = 1$, all distance-regular graphs with classical parameters $(D, 1, \alpha, \beta)$ are known: the Gosset graph (with $\alpha = 4$), the Johnson graphs (with $\alpha = 1$) and the halved ℓ -cubes, where $\ell \in \{2D, 2D + 1\}$ (with $\alpha = 2$), folded Johnson graphs (with $\alpha = 1$), and folded halved $(4D)$ -cube (with $\alpha = 2$); cf [1, Sections 6.1, 6.3]. However, we rule out the Gosset graph since it has diameter $3 < 5$. Therefore, by Theorem 1.2, Γ is a Johnson graph $J(2D, D)$, a halved ℓ -cube with $\ell \in \{2D, 2D + 1\}$, a folded Johnson graph $\bar{J}(4D, 2D)$, or a folded halved $(4D)$ -cube.

Next, consider the case where $b \geq 2$ with $\alpha > 0$. Since $\alpha(1 + b)$ is a positive integer, it follows $\alpha \geq 1/(1 + b)$. Using this along with Lemma 6.1, we have $\beta > \alpha \begin{bmatrix} D-1 \\ 1 \end{bmatrix} \geq \frac{1}{1+b} \begin{bmatrix} D-1 \\ 1 \end{bmatrix}$. Note that $k = \beta \begin{bmatrix} D \\ 1 \end{bmatrix}$. By these comments, we have

$$k = \beta \begin{bmatrix} D \\ 1 \end{bmatrix} > \frac{1}{1+b} \begin{bmatrix} D \\ 1 \end{bmatrix} \begin{bmatrix} D-1 \\ 1 \end{bmatrix} = \frac{(b^D - 1)(b^{D-1} - 1)}{(b+1)(b-1)^2}.$$

Using computer, we find that $\frac{(b^D - 1)(b^{D-1} - 1)}{(b+1)(b-1)^2} > F(b)$ for $D \geq 10$, where $F(b)$ is from (3). In other words, for $D \geq 10$ we have $k > F(b)$. However, according to the classification in Theorem 1.2, it must be $k \leq F(b)$. Therefore, we have $D \leq 9$. The proof is complete. \blacksquare

Remark 6.3. (i) We give a comment on the case (i) of Theorem 6.2. In a subsequent paper, we will show that if Γ is a 1-homogeneous distance-regular graph with $a_1 \neq 0$ and classical parameters $(D, b, 0, \beta)$ where $D \geq 5$, then Γ is locally a disjoint union of cliques. Therefore, by Lemma 5.2, Γ is a regular near $2D$ -gon. Consequently, according to [1, Theorem 9.4.4], Γ is either a dual polar graph or a Hamming graph.

(ii) The unitary dual polar graphs $U(2D, r)$ ¹ have two different types of classical parameters:

$$(D, b, \alpha, \beta) = (D, q^2, 0, q), \tag{39}$$

$$(D, b, \alpha, \beta) = \left(D, -q, \frac{q(1+q)}{1-q}, \frac{q(1+(-q)^D)}{1-q} \right), \tag{40}$$

where q is a prime power and $r = q^2$. Note that in the case (40), we have $\alpha \neq 0$, but $b < 0$; see [1, Corollary 6.2.2].

¹This family of dual polar graphs is often denoted by ${}^2A_{2D-1}(r)$.

7 Tight distance-regular graphs

Let Γ be a distance-regular graph with $D \geq 3$ and eigenvalues $k = \theta_0 > \theta_1 > \cdots > \theta_D$. Jurišić, Koolen and Terwilliger established the following so-called fundamental bound for Γ :

$$\left(\theta_1 + \frac{k}{a_1 + 1}\right) \left(\theta_D + \frac{k}{a_1 + 1}\right) \geq -\frac{ka_1b_1}{(a_1 + 1)^2}, \quad (41)$$

cf. [12, Theorem 6.2]. The graph Γ is said to be *tight* whenever Γ is not bipartite and equality holds in (41). We note that if Γ is tight, then $a_1 \neq 0$; cf. [12, Corollary 6.3]. We recall some characterizations of tight distance-regular graphs.

Lemma 7.1 (cf. [12, Theorem 11.7, Theorem 12.6]). *Let Γ be a distance-regular graph with diameter $D \geq 3$. Then the following are equivalent.*

- (i) Γ is tight
- (ii) Γ is 1-homogeneous with $a_1 \neq 0$ and $a_D = 0$.
- (iii) Every local graph of Γ is connected strongly regular with eigenvalues $a_1 > r > s$, where

$$r = -1 - \frac{b_1}{\theta_D + 1}, \quad s = -1 - \frac{b_1}{\theta_1 + 1}.$$

Suppose Γ is tight. In [15, Theorem 1.3], Koolen et al. showed that if a local graph of Γ is neither the block graph of an orthogonal array nor the block graph of a Steiner system, then the valency k (and hence diameter D) of Γ is bounded by a function in b , where $b = b_1/(1 + \theta_1) \geq 2$. They then proposed a conjecture that generalizes this result: if Γ is a tight distance-regular graph with $D \geq 3$ and $b \geq 2$, then the diameter D of Γ is bounded by a function in b ; see [15, Conjecture 7.5]. In the following result, we prove this conjecture by using Theorem 1.2.

Corollary 7.2. *Let Γ be a tight distance-regular graph with diameter $D \geq 5$. Let $b = b_1/(\theta_1 + 1)$. Then $b \geq 1$ and one of the following holds:*

- (i) Γ is a Johnson graph $J(2D, D)$,
- (ii) Γ is a halved ℓ -cube where $\ell \in \{2D, 2D + 1\}$,
- (iii) Γ is locally connected with $k \leq F(b)$, where $F(b)$ is from (3).

Proof. By Lemma 7.1, the tight graph Γ is locally connected, which implies $c_2 \geq 2$. Moreover, Γ is 1-homogeneous with $a_1 \neq 0$ and $a_D = 0$. Therefore, by Theorem 1.2, it follows that $b \geq 1$, and Γ is either a regular near $2D$ -gon or falls into one of (ii)–(vi) in Theorem 1.2. However, Γ cannot be a regular near $2D$ -gon since $a_1 \neq 0$ and $a_D = 0$. Thus, Γ belongs to one of (ii)–(vi) therein.

Assume that $k > F(b)$; otherwise, it leads to (iii). By Lemma 7.1, the tight graph Γ is locally a connected strongly regular graph. We note that Γ has $a_D = 0$. Following the argument in the proof of Theorem 1.2, we find that Γ is locally either a Latin square graph or a Steiner graph.

If Γ is locally a Latin square graph, by Theorem 1.2, Γ is either a Johnson graph $J(2D, D)$ or a folded Johnson graph $\bar{J}(4D, 2D)$. However, we rule out the case of the graph $\bar{J}(4D, 2D)$ since it has $a_D \neq 0$. Therefore, Γ is a Johnson graph $J(2D, D)$.

If Γ is locally a Steiner graph, by Theorem 1.2 Γ is either a halved ℓ -cube ($\ell \in \{2D, 2D + 1\}$) or a folded halved $(4D)$ -cube. However, we rule out the case of a folded halved $(4D)$ -cube since it has $a_D \neq 0$. Therefore Γ is a halved ℓ -cube where $\ell \in \{2D, 2D + 1\}$. The desired result follows. ■

We remark that in the case of Corollary 7.2(i) and (ii), we have $b = 1$. Also, note that except for the halved ℓ -cubes ($\ell \in \{2D, 2D + 1\}$) and the Johnson graphs $J(2D, D)$, all known tight distance-regular graphs have diameter $D \leq 4$. We finish this section with a comment.

Corollary 7.3. *Let Γ be a tight distance-regular graph of diameter $D \geq 5$ with classical parameters (D, b, α, β) . Then $b \geq 1$ and one of the following holds:*

- (i) Γ is a Johnson graph $J(2D, D)$,
- (ii) Γ is a halved ℓ -cube where $\ell \in \{2D, 2D + 1\}$,
- (iii) $D \leq 9$, $\alpha > 0$, $b \geq 2$.

Proof. By Theorem 6.2 and Corollary 7.2. ■

Acknowledgements

J.H. Koolen is partially supported by the National Key R. and D. Program of China (No. 2020YFA0713100), the National Natural Science Foundation of China (No. 12071454), and the Anhui Initiative in Quantum Information Technologies (No. AHY150000). M. Abdullah is supported by the Chinese Scholarship Council at USTC, China. B. Gebremichel is supported by the National Key R. and D. Program of China (No. 2020YFA0713100) and the Foreign Young Talents Program (No. QN2022200003L).

References

- [1] A.E. Brouwer, A.M. Cohen, A. Neumaier, *Distance-Regular Graphs*, Springer-Verlag, Berlin, 1989.
- [2] A.E. Brouwer, H. Van Maldeghem, *Strongly regular graphs*, Cambridge University Press, Cambridge, 2022.
- [3] B. Curtin, K. Nomura, Homogeneity of a distance-regular graph which supports a spin model, *J. Alg. Combin.* 19 (2004), no. 3, 257–272.
- [4] B. Curtin, K. Nomura, 1-homogeneous, pseudo-1-homogeneous, and 1-thin distance-regular graphs, *J. Combin. Theory Ser. B* 93 (2005), no. 2, 279–302.
- [5] E.R. van Dam, J.H. Koolen, H. Tanaka, Distance-regular graphs, *Electron. J. Combin.*, (1983) DS22.

- [6] C. Godsil, G. Royle, *Algebraic Graph Theory*, Graduate Texts in Mathematics, Vol. 207, Springer-Verlag, New York, 2001.
- [7] A. Jurišić, J.H. Koolen, A local approach to 1-homogeneous graphs, *Des. Codes Cryptogr.*, 21 (2000) 127–147.
- [8] A. Jurišić, J.H. Koolen, Nonexistence of some antipodal distance-regular graphs of diameter four, *European J. Combin.*, 21 (2000) 1039–1046.
- [9] A. Jurišić, J.H. Koolen, 1-homogeneous graphs with Cocktail Party μ -graphs, *J. Alg. Combin.*, 18 (2003) 181–202.
- [10] A. Jurišić, J.H. Koolen, Distance-regular graphs with complete multipartite μ -graphs and the AT4 family, *J. Alg. Combin.*, 25 (2007) 459–471.
- [11] A. Jurišić, J.H. Koolen, Classification of the family $AT4(qs, q, q)$ of antipodal tight graphs, *J. Combin. Theory Ser. A.*, 118 (2011) 842–852.
- [12] A. Jurišić, J.H. Koolen, P. Terwilliger, Tight distance-regular graphs, *J. Alg. Combin.*, 12 (2000) 162–197.
- [13] A. Jurišić, A. Munemasa, Y. Tagami, On graphs with complete multipartite μ -graphs, *Discrete Math.*, 310 (2010) 1812–1819.
- [14] A. Jurišić, J. Vidali, Restrictions on classical distance-regular graphs, *J. Alg. Combin.*, 46 (2017) 571–588.
- [15] J.H. Koolen, J.-H. Lee, S.-D. Li, Y.-H. Li, X. Liang, Y.-Y. Tan, On the (non-)existence of tight distance-regular graphs: a local approach, *submitted*, preprint: [arXiv:2312.05595](https://arxiv.org/abs/2312.05595).
- [16] J.H. Koolen, J. Park, Distance-regular graphs with a_1 or c_2 at least half the valency, *J. Combin. Theory Ser. A*, 119 (2012) 546–555.
- [17] Š. Miklavič, Q -polynomial distance-regular graphs with $a_1 = 0$, *European J. Combin.* 25 (2004), no. 7, 911–920.
- [18] A. Neumaier, Strongly regular graphs with smallest eigenvalue $-m$, *Arch. Math.*, 33 (1979) 392–400.
- [19] K. Nomura, Homogeneous graphs and regular near polygons, *J. Combin. Theory Ser. B*, 60 (1994) 63–71.
- [20] Y.-Y. Tan, J.H. Koolen, M.-Y. Cao, J. Park, Thin Q -polynomial distance-regular graphs have bounded c_2 , *Graphs Combin.*, 38 (2022) #175.
- [21] P. Terwilliger, Distance-regular graphs with girth 3 or 4: I, *J. Combin. Theory Ser. B* 39 (1985), 265–281.