Convergence Guarantees for RMSProp and Adam in Generalized-smooth Non-convex Optimization with Affine Noise Variance

Qi Zhang¹ Yi Zhou² Shaofeng Zou¹

Abstract

This paper provides the first tight convergence analyses for RMSProp and Adam in non-convex optimization under the most relaxed assumptions of coordinate-wise generalized smoothness and affine noise variance. We first analyze RM-SProp, which is a special case of Adam with adaptive learning rates but without first-order momentum. Specifically, to solve the challenges due to dependence among adaptive update, unbounded gradient estimate and Lipschitz constant, we demonstrate that the first-order term in the descent lemma converges and its denominator is upper bounded by a function of gradient norm. Based on this result, we show that RM-SProp with proper hyperparameters converges to an ϵ -stationary point with an iteration complexity of $\mathcal{O}(\epsilon^{-4})$. We then generalize our analvsis to Adam, where the additional challenge is due to a mismatch between the gradient and first-order momentum. We develop a new upper bound on the first-order term in the descent lemma, which is also a function of the gradient norm. We show that Adam with proper hyperparameters converges to an ϵ -stationary point with an iteration complexity of $\mathcal{O}(\epsilon^{-4})$. Our complexity results for both RMSProp and Adam match with the complexity lower bound established in (Arjevani et al., 2023).

1. Introduction

Preprint.

RMSProp (Hinton et al., 2012) and Adam (Kingma & Ba, 2014) are among the most popular and powerful adaptive optimizers in training state-of-the-art machine learning models (Brock et al., 2018; Brown et al., 2020; Cutkosky & Mehta, 2020; Dosovitskiy et al., 2020). RM-

SProp and Adam only require first-order gradients with little memory requirement, and thus are efficient to use in practice. RMSProp is based on the idea of adaptive learning rates for each individual parameter, and Adam combines the benefits of RMSprop (Hinton et al., 2012) and AdaGrad (Duchi et al., 2011), which consists of two key components of adaptive learning rates and momentum. Despite their empirical success, theoretical understandings on the convergence and complexity, especially when optimizing non-convex loss functions, e.g., neural networks, still remain underdeveloped until very recently.

Recently, there have been a series of works in examining the convergence and complexity of RMSProp and Adam for non-convex loss functions (see Table 1 for a detailed review). However, these works do not completely explain the performance of RMSProp and Adam in training neural networks, as they rely on assumptions that may not necessarily hold. For example, (Zhang et al., 2019) pointed out that neural networks are not *L*-smooth, and instead satisfy the generalized (L_0, L_1) -smoothness, where the gradient Lipschitz constant increases linearly with the gradient norm. Furthermore, many of these works assumed that the stochastic gradient has a bounded norm/variance, which however does not even hold for linear regression (Wang et al., 2023b); and instead a relaxed affine noise condition shall be used.

In this paper, we derive the convergence guarantee and iteration complexity for RMSProp and Adam with coordinatewise generalized (L_0, L_1) -smooth loss function and affine noise variance. To the best of our knowledge, these are some of the most relaxed assumptions in the convergence analyses of RMSProp and Adam that best describes the training of neural networks. We prove that RMSProp and Adam with proper hyperparameters converge to an ϵ -stationary point with a complexity of $\mathcal{O}(\epsilon^{-4})$, which matches with the lower bound for first-order optimization in (Arjevani et al., 2023).

1.1. Challenges and Insights

Our theoretical analyses address several major challenges: (i) dependence between stepsize and gradient, (ii) potential unbounded gradients, (iii) mismatch between gradient and

¹Department of Electrical Engineering, University at Buffalo, NY, USA (email: qzhang48@buffalo.edu, szou3@buffalo.edu) ²Department of Electrical and Computer Engineering, University of Utah, Salt Lake City, USA (e-mail: yi.zhou@utah.edu).

first-order momentum, and (iv) additional bias terms due to affine variance and coordinate-wise (L_0, L_1) -smoothness. Prior research circumvented most of these challenges by introducing extra assumptions, whereas we provide several new insights and show that these assumptions may not be needed.

Algorithm 1 Adam

Initialize parameters: \boldsymbol{x}_1 , learning rate η , β_1 , β_2 , ζ , Iteration TInitialize first and second moment estimates: $\boldsymbol{v}_0 \in \mathbb{R}^{d+}$, $\boldsymbol{m}_0 = 0$ Initialize time step: t = 1while $t \leq T$ do Generate Stochastic gradient: \boldsymbol{g}_t Update first-order momentum estimate: $\boldsymbol{m}_t \leftarrow \beta_1 \cdot \boldsymbol{m}_{t-1} + (1 - \beta_1) \cdot \boldsymbol{g}_t$ Update second-order momentum estimate: $\boldsymbol{v}_t \leftarrow \beta_2 \cdot \boldsymbol{v}_{t-1} + (1 - \beta_2) \cdot \boldsymbol{g}_t \odot \boldsymbol{g}_t$ Update parameters: $\boldsymbol{x}_{t+1} \leftarrow \boldsymbol{x}_t - \eta \frac{1}{\sqrt{\boldsymbol{v}_t + \zeta}} \odot \boldsymbol{m}_t$ $t \leftarrow t + 1$ end while

In this paper, we have access to an unbiased estimate g of $\nabla f(x)$ such that $\mathbb{E}[g|x] = \nabla f(x)$ where $x \in \mathbb{R}^d$ is a *d*dimensional parameter and $f : \mathbb{R}^d \to \mathbb{R}$ is the objective function. Define $\mathcal{F}_t := \sigma(g_1, ..., g_{t-1})$ as the sigma field of the stochastic gradients up to t - 1. Consider the Adam algorithm in Algorithm 1, which reduces to RMSProp if $\beta_1 = 0$. For the coordinate-wise (L_0, L_1) -smooth objective functions, we have the following descent inequality (Lemma 1 in (Crawshaw et al., 2022)):

$$\underbrace{\mathbb{E}\left[\langle \nabla f(\boldsymbol{x}_{t}), \boldsymbol{x}_{t} - \boldsymbol{x}_{t+1} \rangle | \mathcal{F}_{t}\right]}_{\text{First Order}} \leq f(\boldsymbol{x}_{t}) - \mathbb{E}[f(\boldsymbol{x}_{t+1})| \mathcal{F}_{t}] \\
+ \underbrace{\sum_{i=1}^{d} \frac{L_{0}}{2\sqrt{d}} \mathbb{E}[\|\boldsymbol{x}_{t+1} - \boldsymbol{x}_{t}\| | \boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}| | \mathcal{F}_{t}]}_{\text{Second Order}} \\
+ \underbrace{\sum_{i=1}^{d} \frac{L_{1}|\partial_{i}f(\boldsymbol{x}_{t})|}{2} \mathbb{E}[\|\boldsymbol{x}_{t+1} - \boldsymbol{x}_{t}\| | \boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}| | \mathcal{F}_{t}]}_{\text{Additional Term}},$$
(1)

where the last term is the additional term due to the (L_0, L_1) -smooth assumption.

Challenge 1: dependence between stepsize and gradient. We use the RMSProp optimizer to explain our technical novelty. The challenge is the same for Adam. For RMSProp the optimized parameter x is updated: $x_{t+1} = x_t - \frac{\eta g_t}{\sqrt{v_t + \zeta}}$, where the adaptive stepsize $\eta_t = \frac{\eta}{\sqrt{v_t + \zeta}}$ depends on the current gradient estimate g_t , which makes it hard to bound the conditional expectation of the first-order term in (1). To address this challenge, studies on Adagrad (Ward et al., 2020; Défossez et al., 2020; Faw et al., 2022) and studies on RMSProp (Zaheer et al., 2018) propose a surrogate \tilde{v}_t of v_t which is independent of g_t , and then the first-order term is divided into two parts: the firstorder.a term $\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_t), \frac{-\eta g_t}{\sqrt{\tilde{v}_t + \zeta}} \right\rangle \middle| \mathcal{F}_t \right]$ and first-order.b term $\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_t), \frac{-\eta g_t}{\sqrt{\tilde{v}_t + \zeta}} \right\rangle \middle| \mathcal{F}_t \right]$. The main challenge lies in the first-order.b term (surrogate error). In (Zaheer et al., 2018), this term is bounded based on the assumption of bounded gradient norm, which does not hold in this paper.

Insight 1: reduce surrogate error. Recently (Wang et al., 2023b) shows that in the *t*-th iteration, the error term can be written as $\mathbb{E}[\phi(v_{t-1}) - \phi(v_t)|\mathcal{F}_t]$ where ϕ is some function. Thus, these terms cancel out with each other by taking sum from t = 1 to *T*. However, this method only works for Adagrad due to its non-increasing stepsize. We develop a novel approach, which involves a minor change to the adaptive stepsize from $\frac{\eta}{\sqrt{v_t+\zeta}}$ to $\frac{\eta}{\sqrt{v_t+\zeta}}$ but does not change the adaptive nature of the stepsize. In this way, we can cancel out terms by taking sum w.r.t. to *t* (details can be found in Lemma 3.3).

Challenge 2: unbounded gradient. Previous works, e.g., (De et al., 2018; Défossez et al., 2020; Zaheer et al., 2018) assumed that the gradient norm is bounded, based on which, they proved the convergence. However, with refined gradient noise, the gradient may be infinite, and thus those approaches do not apply.

Insight 2: recursive bounding technique. We first show that $\mathbb{E}\left[\frac{1}{T}\sum_{t=1}^{T}\frac{\|\nabla f(\boldsymbol{x}_t)\|^2}{\sqrt{\|\tilde{\boldsymbol{v}}_t\|+\zeta}}\right]$ is bounded. If the gradient norm is upper bounded, then $\tilde{\boldsymbol{v}}_t$ is bounded, and the convergence result directly follows. However, under refined gradient noise, the gradient norm may not be bounded. We develop a novel approach to bound $\mathbb{E}\left[\frac{1}{T}\sum_{t=1}^{T}\sqrt{\|\tilde{\boldsymbol{v}}_t\|+\zeta}\right]$ using $\mathbb{E}\left[\sum_{t=1}^{T}\|\nabla f(\boldsymbol{x}_t)\|\right]$ instead of a constant (see Lemma 3.5 for details). Applying Hölder's inequality (Hardy et al., 1952) we will obtain the convergence result. The complexity result matches with the lower bound in (Arjevani et al., 2023).

Challenge 3: mismatch between gradient and firstorder momentum. Compared with RMSProp, Adam employs the first momentum m_t . The momentum m_t is dependent on the surrogate stepsize $\frac{\eta}{\sqrt{v_t+\zeta}}$. Moreover, the momentum m_t is a biased estimate of the current true gradient. Both the above challenges make it hard to theoretically characterize the convergence rate of Adam.

Insight 3: bounding first-order term using
$$\mathbb{E}\left[\frac{1}{T}\sum_{t=1}^{T} \|\nabla f(\boldsymbol{x}_t)\|\right]$$
. Inspired by the analysis

for SGDM (Liu et al., 2020) and Adam (Wang et al., 2023a), we consider a potential function of $f(u_t)$ with $u_t = \frac{x_t - x_{t-1}\beta_1/\sqrt{\beta_2}}{1 - \beta_1/\sqrt{\beta_2}}$. We can show that $u_{t+1} - u_t$ is close to a function of $\frac{g_t}{\sqrt{v_t + \zeta}}$, which is much easier to analyze compared with $\frac{m_t}{\sqrt{v_t + \zeta}}$. Using the descent lemma of $f(u_t)$, we show the first-order term is also bounded by a constant plus a function of $\mathbb{E}\left[\frac{\sum_{t=1}^T ||\nabla f(x_t)||}{T}\right]$. Compared to RMSProp, this additional function is introduced due to the bias of m_t . Then via Hölder's inequality, we show Adam converges as fast as RMSProp.

Challenge 4: additional error terms due to affine variance and (L_0, L_1) -smoothness. Compared with *L*-smooth objectives, in the analysis for RMSProp with (L_0, L_1) -smooth objectives, there is an additional second-order error term: $\sum_{i=1}^{d} \frac{L_1 |\partial_i f(\boldsymbol{x}_t)|}{2} [||\boldsymbol{x}_{t+1} - \boldsymbol{x}_t|| |\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}|],$ which is hard to bound since $|\partial_i f(\boldsymbol{x}_t)|$ may be unbounded. Moreover, for RMSProp, since $\mathbb{E}[|\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}||\mathcal{F}_t] \leq \mathbb{E}\left[\frac{\eta|\boldsymbol{g}_{t,i}|}{\sqrt{\tilde{v}_{t,i}+\zeta}} |\mathcal{F}_t\right]$ and $\boldsymbol{g}_{t,i}$ is independent of $\tilde{\boldsymbol{v}}_{t,i}$ given \mathcal{F}_t , the affine noise assumption can be leveraged to bound the second-order error term directly. Nevertheless, for Adam due to the dependence between $\boldsymbol{m}_{t,i}$ and $\tilde{\boldsymbol{v}}_{t,i}$, the above approach cannot be applied directly.

Insight 4: bounding additional term by function of first-order term. For RMSProp, we can show that $\|x_{t+1} - x_t\| \leq \frac{\eta\sqrt{d}}{\sqrt{1-\beta_2}}$ and $|\partial_i f(x_t)| |x_{t+1,i} - x_{t,i}| \leq \frac{|\partial_i f(x_t)|^2 + \eta^2 g_{t,i}^2}{2\sqrt{\tilde{v}_{t,i} + \zeta}}$. According to the affine noise assumption, we have that $\mathbb{E}[g_{t,i}^2|\mathcal{F}_t]$ is upper bounded by a linear function of $(\partial_i f(x_t))^2$, thus we can bound the additional error term. However, we cannot directly apply this method to Adam since $\mathbb{E}[m_{t,i}^2|\mathcal{F}_t]$ is hard to bound. Instead, we provide a new upper bound on $\sum_{t=1}^T \frac{m_{t,i}^2}{\sqrt{v_{t,i}}}$ using the gradient norm.

1.2. Related work

1.2.1. RELAXED ASSUMPTIONS

Affine Noise Variance: In most of the studies on stochastic optimization, access to an unbiased estimate of the gradient with uniformly bounded variance is assumed (Nemirovskij & Yudin, 1983; Ghadimi & Lan, 2013; Bubeck et al., 2015; Foster et al., 2019). (Ghadimi & Lan, 2013) first showed that for *L*-smooth objectives, the SGD algorithm converges to a first-order ϵ -stationary point with an iteration complexity of $\mathcal{O}(\epsilon^{-4})$ if the stochastic gradient has uniformly bounded variance. Furthermore, (Arjevani et al., 2023) proved that for any first-order algorithm with uniformly bounded gradient variance, the iteration complexity of $\mathcal{O}(\epsilon^{-4})$ is optimal. For overparameterized neural networks, (Vaswani et al., 2019) considered another gradient noise assumption: the strong growth condition, where the upper bound on the second-order moment of the oracle norm scales with the gradient square norm. Both the uniformly bounded variance and strong growth condition are special cases of the affine noise variance. It was demonstrated in (Bottou et al., 2018) that for non-adaptive algorithms with affine noise variance, the optimal iteration complexity of $\mathcal{O}(\epsilon^{-4})$ can be achieved. The extension of affine noise assumption to adaptive algorithms is not straightforward and was studied in (Jin et al., 2021; Chen et al., 2022; 2023; Hong & Lin, 2023). In this paper, we study two adaptive optimizers: RMSProp and Adam with affine noise variance.

Generalized Smoothness: In stochastic optimization, the L-smooth objectives are widely assumed (Ghadimi & Lan, 2013; Ghadimi et al., 2016). However, it was demonstrated in (Zhang et al., 2019) that the L-smoothness does not hold for neural networks and polynomial functions with degree larger than 2. Then, extensive experiments were conducted to verify that these functions satisfy the generalized (L_0, L_1) -smoothness condition, where the gradient Lipschitz constant increases linearly with the gradient norm. Compared with L-smoothness, (L_0, L_1) -smoothness introduces extra second-order error terms, thus making the optimization problem hard to solve. The clipping algorithms for generalized smooth function were studied in (Zhang et al., 2019; 2020). However, they require the gradient norm to be bounded. A relaxed assumption on bounded gradient variance was studied in (Reisizadeh et al., 2023), where the SPIDER algorithm was applied. Furthermore, (Chen et al., 2023) showed that for generalized smooth objectives with affine noise variance, the SPIDER algorithm still finds a stationary point. Under the same assumption, (Jin et al., 2021) provided the convergence rate for a normalized momentum algorithm. With extensive experiments, (Crawshaw et al., 2022) showed that in the training of Transformer, the (L_0, L_1) -smoothness holds coordinate-wisely. In this paper, we consider functions that are coordinate-wise (L_0, L_1) -smooth.

1.2.2. Adaptive Optimizers

Adaptive optimizers are widely used in deep learning due to their ability to adjust to changing data and conditions. Adagrad (Duchi et al., 2011) is the first adaptive algorithm, which calculates the accumulated sum of the past gradient norms and uses the reciprocal of its square root to scale the current gradient. Recently, (Wang et al., 2023b; Faw et al., 2023) studied Adagrad under generalized smoothness and affine noise variance conditions. However, the training of the above Adagrad algorithm may stop in advance since the accumulated sum does not shrink, and thus the learning rate can be extremely close to 0. To overcome this problem, RMSProp (Hinton et al., 2012) was proposed, where a momentum method is employed to replace the accumulated sum. Thus, the adaptive learning rate can increase or decrease. There is a rich literature on the convergence analyses of RMSProp (Zaheer et al., 2018; De et al., 2018; Shi et al., 2020). However, all of them focus on Lsmooth objectives, and only (Shi et al., 2020) considered the affine noise variance. RMSProp is a special case of Adam, which only includes the second-order momentum, and is widely studied e.g., (Défossez et al., 2020; Zou et al., 2019; Chen et al., 2018; Zhang et al., 2022; Wang et al., 2022; Guo et al., 2021; Hong & Lin, 2023; Li et al., 2023; Wang et al., 2023a). In (Li et al., 2023), RMSProp for (L_0, L_1) -smooth objectives with sub-Gaussian norm was studied, where the gradient estimate bias follows a sub-Gaussian distribution. Under this assumption, based on the gradient estimate, the real gradient belongs to a bounded set with high probability, which converts the unbounded Lipschitz constant to a bounded one. The only results of Adam on (L_0, L_1) -smoothness with affine noise (for the special case of finite sum problems) were in (Wang et al., 2022). However, they only showed that Adam converges to the neighborhood of a stationary point with a constant learning rate. More details can be found in Table 1.

There are also two recent works (Hong & Lin, 2023; Wang et al., 2023a) that studied Adam for L-smooth objectives with affine noise variance. However, their methods can not be generalized to (L_0, L_1) smooth objectives due to the additional terms invalidating the key inequality or requirements of bounded Lipchitz constant in their key Lemma. In this paper, we study the more practical and challenging coordinate-wise (L_0, L_1) -smooth objectives. Moreover, we show that RMSProp and Adam converge to a stationary point with a complexity of $\mathcal{O}(\epsilon^{-4})$, which matches with the lower bound in (Arjevani et al., 2023). Our results improve those in (Hong & Lin, 2023) by logarithmic factors.

2. Preliminaries

Let $f : \mathbb{R}^d \to \mathbb{R}$ be a differentiable non-convex loss function. For a positive integer d, let [d] denote the set $\{1, 2, ..., d\}$. Let $x \in \mathbb{R}^d$ be an optimization variable. Our goal is to minimize the objective function f(x):

$$\min_{\boldsymbol{x}} f(\boldsymbol{x}).$$

For a differentiable function f, a point $x \in \mathbb{R}^d$ is called a first-order ϵ -stationary point if we have that $\|\nabla f(x)\| \leq \epsilon$. Denote by $x_t \in \mathbb{R}^d$ the optimization variable at the *t*-th iteration and we have access to an estimate g_t of $\nabla f(x_t)$. Define $\mathcal{F}_t := \sigma(g_1, ..., g_{t-1})$ is the sigma field of the stochastic gradients up to t - 1. We focus on the Adam algorithm shown in Algorithm 1, where \odot denotes the Hadamard product. For any $i \in [d]$, $\partial_i f(\boldsymbol{x}_t), \boldsymbol{g}_{t,i}, \boldsymbol{m}_{t,i}$ and $\boldsymbol{v}_{t,i}$ are the *i*-th element of $\nabla f(\boldsymbol{x}_t), \boldsymbol{g}_t, \boldsymbol{m}_t$ and \boldsymbol{v}_t , respectively.

Compared with the original Adam, we make a minor change in the adaptive stepsize from $\eta_t = \frac{\eta}{\sqrt{v_t+\zeta}}$ to $\frac{\eta}{\sqrt{v_t+\zeta}}$. This minor change does not influence the adaptivity of the algorithm but makes the analysis much easier.

2.1. Technical Assumptions

Throughout this paper, we make the following assumptions.

Assumption 2.1. f(x) is bounded from below such that $\inf_{x} f(x) > -\infty$.

Assumption 2.2 (Coordinate-wise affine noise variance (Wang et al., 2023a; Hong & Lin, 2023)). We have access to an unbiased gradient estimate g_t such that $\mathbb{E}[g_t|\mathcal{F}_t] = \nabla f(x_t)$ and for any $i \in [d] \mathbb{E}[g_{t,i}^2|\mathcal{F}_t] \leq D_0 + D_1(\partial_i f(x_t))^2$, where $D_0, D_1 \geq 0$ are some constants.

As discussed in (Hong & Lin, 2023), this assumption allows the magnitude of noise to scale with the corresponding gradient coordinate. Many widely used noise assumptions are special cases of this affine noise variance assumption. For example, when $D_0 = 0$, it is the bounded second-order moment assumption in (Zou et al., 2019) and for the finite sum formulation $f(x) = \frac{\sum_{j=1}^{N} f_j(x)}{N}$, when $D_1 = \frac{1}{N}$, it is equivalent to bounded gradient variance. However, as pointed out in (Wang et al., 2023b), these two assumptions of bounded second-order moment and bounded gradient variance do not even hold for linear regression problems. $D_0 = 0$ is called "strong growth condition" (Vaswani et al., 2019), which can be seen as a simple condition for overparameterized neural networks that interpolate all data points. Under Assumption 2.2, the norm of the gradient increases with the norm of the true gradient. This is important for model parameters that are multiplicatively perturbed by noise, e.g., multilayer network (Faw et al., 2022). In this paper, we study the coordinate-wise affine noise variance assumption, which was also used in (Hong & Lin, 2023; Wang et al., 2023a).

Though the *L*-smoothness assumption is widely used in optimization studies, recently it has been observed that in the training of neural networks, this assumption does not hold. Instead, it is numerically verified that the following generalized smoothness assumption better models the training of neural networks (Zhang et al., 2019): $\|\nabla f(x) - \nabla f(y)\| \leq (L_0 + L_1 \|\nabla f(x)\|) \|x - y\|$ for some positive L_0 and L_1 . This assumption is widely studied in the literature, e.g., (Jin et al., 2021; Chen et al., 2023; Li et al., 2023; Wang et al., 2022; 2023b; Faw et al., 2023). Compared with *L*-smooth functions, for the generalized smooth functions, the Lipschitz constant scales with the

Convergence Guarantees for RMSProp and Adam

Method	Smoothness	Algorithm	Stationary point ^a	Assumption	Batch size	Complexity
(De et al., 2018)	L-smooth	RMSProp	1	Bounded Norm ^b	$\mathcal{O}(1)$	$O(\epsilon^{-4})$
(Zaheer et al., 2018)	L-smooth	RMSProp	1	Bounded Norm	$\mathcal{O}(\epsilon^{-2})$	$\mathcal{O}(\epsilon^{-4})$
(Shi et al., 2020)	L-smooth	RMSProp	✓	Finite Sum Affine Noise	-	-
(Défossez et al., 2020)	L-smooth	Adam	1	Bounded Norm	$\mathcal{O}(1)$	$\mathcal{O}(\epsilon^{-4})$
(Zou et al., 2019)	L-smooth	Adam	1	Bounded Second-order Moment	$\mathcal{O}(1)$	$\mathcal{O}(\epsilon^{-4})$
(Chen et al., 2018)	L-smooth	Adam	×	Bounded Norm	-	-
(Zhang et al., 2022)	L-smooth	Adam	×	Finite Sum Affine Noise	-	-
(Wang et al., 2022)	Finite Sum (L_0, L_1) -smooth	Adam	×	Finite Sum Affine Noise	-	-
(Guo et al., 2021)	L-smooth	Adam	1	Affine Noise ^c	$\mathcal{O}(1)$	$\mathcal{O}(\epsilon^{-4})$
(Hong & Lin, 2023)	L-smooth	Adam	1	Coornidate-wise Affine Noise	$\mathcal{O}(1)$	$\mathcal{O}(ploy(\log(\epsilon^{-1}))\epsilon^{-4})$
(Wang et al., 2023a)	L-smooth	Adam	✓	Coornidate-wise Affine Noise	$\mathcal{O}(1)$	$\mathcal{O}(\epsilon^{-4})$
(Li et al., 2023)	(L_0, L_1) -smooth	Adam	1	Sub-Gaussian Norm	$\mathcal{O}(1)$	$\mathcal{O}(\epsilon^{-4})^{\mathrm{d}}$
Our method	Coornidate-wise (L_0, L_1) -smooth	Adam	1	Coornidate-wiseAffine Noise	$\mathcal{O}(1)$	$\mathcal{O}(\epsilon^{-4})$

Table 1. Comparison for existing RMSProp and Adam analyses. Only our paper focuses on the (L_0, L_1) -smooth objectives with affine noise variance, and we show convergence to a stationary point. For $\nabla f(\boldsymbol{x})$ with its estimate \boldsymbol{g} , the bounded norm assumption is $\|\boldsymbol{g}\| \leq G$ (almost surely), where G is some positive constant. The bounded second-order moment assumption is that $\mathbb{E}[\boldsymbol{g}] \leq G^2$. The bounded norm assumption is that $\|\boldsymbol{G} - \nabla f(\boldsymbol{x})\|$ follows a sub-Gaussian distribution, which is weaker than the bounded norm assumption but stronger than the bounded variance assumption.

Explanation on the upper footmarks: a: X indicates the algorithm only converges to the neighborhood of a stationary point, whose radius can not be made small. b: (De et al., 2018) requires the signs of the gradients to be the same across batches. c: (Guo et al., 2021) requires the upper bound on the gradient norm. d: A variance-reduced method is also investigated in (Li et al., 2023), and the complexity is $\mathcal{O}(\epsilon^{-3})$.

true gradient norm thus may not be bounded. For the training of Transformer models, (Crawshaw et al., 2022) finds the following coordinate-wise (L_0, L_1) -smoothness provides a more accurate characterization of the objective. It is generalized from the coordinate-wise *L*-smoothness (Richtárik & Takáč, 2014; Khaled & Richtárik, 2020; Bernstein et al., 2018). In this paper, we focus on coordinate-wise (L_0, L_1) -smooth functions as defined below:

Assumption 2.3 (Coordinate-wise (L_0, L_1) -smoothness). A function f is coordinate-wise (L_0, L_1) -smooth if for any $\boldsymbol{x}, \boldsymbol{y} \in \mathbb{R}^d$ and $i \in [d]$,

$$|\partial_i f(\boldsymbol{x}) - \partial_i f(\boldsymbol{y})| \le (L_0 + L_1 |\partial_i f(\boldsymbol{x})|) \|\boldsymbol{x} - \boldsymbol{y}\|.$$
(2)

The training of Adam enjoys adaptive learning rates for each parameter individually due to the coordinate-wise update process. Moreover, extensive experiments in (Crawshaw et al., 2022) show that the L_0 and L_1 for each coordinate varies a lot. Therefore, it is more accurate to leverage the coordinate-wise (L_0, L_1) -smoothness. In this paper, for the sake of simplicity and coherence with the coordinate-wise affine noise assumption, we assume the L_0 and L_1 to be identical for each coordinate. Our results can be easily adapted to the case with distinct L_0 and L_1 for different coordinates.

3. Convergence Analysis of RMSProp

To provide a fundamental understanding of Adam, in this section, we focus on RMSProp which consists of the design of adaptive learning rates for each individual parameter in Adam.

For RMSProp, the main challenges come from dependence between stepsize and gradient, potential unbounded gradients due to the affine noise, and additional error terms due to (L_0, L_1) -smoothness. The analysis can be extended to general Adam, which requires additional efforts to solve the first-order momentum. Define $c = \sqrt{\zeta} + d\sqrt{D_0 + \|\boldsymbol{v}_0\|}$. We then present our analysis of RMSProp in the following theorem.

Theorem 3.1 (Informal). Let Assumptions 2.1, 2.2 and 2.3 hold. Let $1 - \beta_2 \sim \mathcal{O}(\epsilon^2)$, $\eta \sim \mathcal{O}(\epsilon^2)$, and $T \sim \mathcal{O}(\epsilon^{-4})$. For small ϵ such that $\epsilon \leq \frac{\sqrt{5dD_0}}{\sqrt{D_1}\sqrt[4]{2}\zeta}$, we have that

$$\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}[\|\nabla f(\boldsymbol{x}_t)\|] \le \left(\frac{2d\sqrt{35D_0D_1}}{\sqrt[4]{\zeta}} + \sqrt{c}\right)\epsilon.$$
 (3)

To the best of our knowledge, this paper provides the first convergence analysis of RMSProp for (L_0, L_1) -smooth functions under the most relaxed assumption of affine noise variance. Existing studies mostly assume bounded gradient norm or variance (De et al., 2018; Zaheer et al., 2018) or only show the algorithm converges to the neighborhood of a stationary point (Shi et al., 2020). More importantly, our result matches the lower bound in (Arjevani et al., 2023), and thus is optimal.

The formal version of the theorem and detailed proof can be found in Appendix D. Below, we provide a proof sketch to highlight our major technical novelties.

Proof sketch. Our proof can be divided into three stages: Stage I: develop an upper bound of $\mathbb{E}\left[\frac{\|\nabla f(\boldsymbol{x}_t)\|^2}{\sqrt{\beta_2 \|\boldsymbol{v}_{t-1}\| + \zeta}}\right];$ Stage II: develop an upper bound on $\mathbb{E}[\sqrt{\beta_2} \| \boldsymbol{v}_{t-1} \| + \zeta]$; and Stage III: show $\mathbb{E}[\|\nabla f(\boldsymbol{x}_t)\|]$ converges using results from Stages I, II and Hölder's inequality.

Stage I: upper bound of $\mathbb{E}\left[\frac{\|\nabla f(\boldsymbol{x}_t)\|^2}{\sqrt{\beta_2 \|\boldsymbol{v}_{t-1}\| + \zeta}}\right]$. As discussed in Section 1, for coordinate-wise (L_0, L_1) -smooth functions, following the descent inequality (Lemma 1 in (Crawshaw et al., 2022)) we can get (1). We first obtain $\frac{\|\nabla f(\boldsymbol{x}_t)\|^2}{\sqrt{\beta_2 \|\boldsymbol{v}_{t-1}\| + \zeta}} \leq \sum_{i=1}^d \frac{\eta(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}}$. Therefore, in the following we will bound $\mathbb{E}\left[\sum_{i=1}^d \frac{\eta(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}}\right]$. Towards this goal, in Step 1.1, we will show the LHS of (1) is lower bounded by a function of $\sum_{i=1}^d \frac{\eta(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}}$. Combining the two steps, we will obtain an upper bound on $\sum_{i=1}^d \frac{\eta(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}}$.

Step 1.1: lower bound on the first-order term in (1). Since the adaptive stepsize and the gradient estimate are dependent, we design a surrogate $\tilde{v}_t = \beta_2 v_{t-1}$ to decompose the first-order term in (1) into two parts:

$$\underbrace{\mathbb{E}[\langle \nabla f(\boldsymbol{x}_{t}), \boldsymbol{x}_{t} - \boldsymbol{x}_{t+1} \rangle | \mathcal{F}_{t}]}_{\text{First Order}} = \underbrace{\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \frac{\eta \boldsymbol{g}_{t}}{\sqrt{\tilde{\boldsymbol{v}}_{t} + \zeta}}\right\rangle | \mathcal{F}_{t}\right]}_{\text{First Order.a}} + \underbrace{\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \frac{\eta \boldsymbol{g}_{t}}{\sqrt{\boldsymbol{v}_{t} + \zeta}} - \frac{\eta \boldsymbol{g}_{t}}{\sqrt{\tilde{\boldsymbol{v}}_{t} + \zeta}}\right\rangle | \mathcal{F}_{t}\right]}_{\text{First Order.b}}. \quad (4)$$

For the first-order a term in (4), we can show that

$$\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \frac{\eta \boldsymbol{g}_{t}}{\sqrt{\tilde{\boldsymbol{v}}_{t}+\zeta}} \right\rangle | \mathcal{F}_{t}\right] = \sum_{i=1}^{d} \frac{\eta(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}$$

We then bound the first-order.b term in (4).

Remark 3.2 (Importance of modified adaptive stepsize η). For the original RMSProp, (Zaheer et al., 2018) chose the same surrogate $\tilde{v}_t = \beta_2 v_{t-1}$ as in this paper. Then the first-order b term was lower bounded by a function of $\sum_{i=1}^{d} \mathbb{E} \left[\frac{-g_{t,i}^2}{\sqrt{\beta_2 v_{t-1,i} + \zeta}} \middle| \mathcal{F}_t \right] \times \mathbb{E} \left[\frac{(1-\beta_2)g_{t,i}^2}{(\sqrt{v_{t,i}} + \sqrt{\beta_2 v_{t-1,i}})^2} \middle| \mathcal{F}_t \right]$. Then they developed an upper bound on the second term $\mathbb{E} \left[\frac{(1-\beta_2)g_{t,i}^2}{(\sqrt{v_{t,i}} + \sqrt{\beta_2 v_{t-1,i}})^2} \middle| \mathcal{F}_t \right] \leq 1$ which is quite loose, thus they introduced an additional assumption on the upper bound of $|\boldsymbol{g}_{t,i}|$ (Zaheer et al., 2018).

In contrast, using our adaptive stepsize $\eta \frac{g_t}{\sqrt{v_t+\zeta}}$, we can show that the first-order.b term can be lower

bounded by a function of
$$\sum_{i=1}^{d} \mathbb{E} \left[\frac{-g_{t,i}^2}{\sqrt{\beta_2 v_{t-1,i}+\zeta}} \Big| \mathcal{F}_t \right] \times \mathbb{E} \left[\frac{(1-\beta_2)g_{t,i}^2}{(\sqrt{v_{t,i}+\zeta}+\sqrt{\beta_2 v_{t-1,i}+\zeta})^2} \Big| \mathcal{F}_t \right],$$

which can be further bounded by $\sum_{i=1}^{d} \mathbb{E} \left[-g_{t,i}^2 \Big| \mathcal{F}_t \right] \mathbb{E} \left[\frac{1}{\sqrt{\beta_2 v_{t-1,i}+\zeta}} - \frac{1}{\sqrt{v_{t,i}+\zeta}} \Big| \mathcal{F}_t \right].$

Applying the affine noise variance assumption in Åssumption 2.2, we obtain a lower bound on the first-order.b term in the following lemma.

Lemma 3.3 (Informal). Under Assumptions 2.2 and 2.3, we have that:

$$\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \frac{\eta \boldsymbol{g}_{t}}{\sqrt{\boldsymbol{v}_{t} + \zeta}} - \frac{\eta \boldsymbol{g}_{t}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1} + \zeta}}\right\rangle \middle| \mathcal{F}_{t}\right] \\
\geq -\sum_{i=1}^{d} \left[\mathcal{O}\left(\frac{\eta(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}}\right) \\
+ \mathcal{O}\left(\frac{\eta^{2}}{\sqrt{1 - \beta_{2}}} \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}}\right) \\
+ \mathcal{O}\left(\frac{\eta}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} - \mathbb{E}\left[\frac{\eta}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}}\middle| \mathcal{F}_{t}\right]\right) \\
+ \mathcal{O}\left(\frac{\eta(\partial_{i}f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} - \mathbb{E}\left[\frac{\eta(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}}\middle| \mathcal{F}_{t}\right]\right) \\
- small error.$$
(5)

The formal version and detailed proof of Lemma 3.3 can be found in the Appendix A, which is based on our modified update process on v_t , Hölder's inequality and Assumption 2.3.

In the RHS of (5), we have

$$\mathbb{E}\left[\frac{(\partial_i f(\boldsymbol{x}_{t-1}))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}} - \frac{(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}}\Big|\mathcal{F}_t\right].$$

Taking sum from t = 1 to T, the terms $\frac{(\partial_i f(\boldsymbol{x}_{t-1}))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta}}$ and $\frac{(\partial_i f(\boldsymbol{x}_{t-1}))^2}{\sqrt{\boldsymbol{v}_{t-1,i}+\zeta}}$ cancel with each other as $\beta_2 \to 1$. Similarly, $\frac{\eta}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta}} - \mathbb{E}\left[\frac{\eta}{\sqrt{\boldsymbol{v}_{-1,i}+\zeta}}\Big|\mathcal{F}_t\right]$ can also be bounded.

Step 1.2: upper bound on second-order and additional terms in (1). We first focus on the second-order term. Based on the update process of v_t and Assumption 2.2, we get that

$$\sum_{i=1}^{d} \frac{L_0}{2\sqrt{d}} \mathbb{E}[\|\boldsymbol{x}_{t+1} - \boldsymbol{x}_t\| | \boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}| | \mathcal{F}_t] \\ \leq \sum_{i=1}^{d} \frac{L_0 \eta^2}{2\sqrt{\zeta}} \frac{D_0 + D_1 (\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}}.$$
 (6)

We then focus on the additional term in (1) and we provide a new upper bound using $\sum_{i=1}^{d} \frac{(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta}}$ with some small errors: for any $\alpha_2 > 0$, we have that

$$\frac{L_{1}|\partial_{i}f(\boldsymbol{x}_{t})|}{2}\mathbb{E}[\|\boldsymbol{x}_{t+1} - \boldsymbol{x}_{t}\||\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}||\mathcal{F}_{t}] \\
\leq \frac{\eta^{2}\sqrt{d}}{2\sqrt{1 - \beta_{2}}} \left(L_{1}\sqrt{D_{1}} + \frac{1}{\alpha_{2}\sqrt{1 - \beta_{2}}}\right) \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} \\
+ \frac{\eta^{2}\sqrt{d}\alpha_{2}L_{1}^{2}D_{0}}{2\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}}.$$
(7)

Plugging Lemma 3.3, (6) and (7) in (1) we get the following lemma.

Lemma 3.4. Let Assumptions 2.1,2.2 and 2.3 hold. With the same parameters in Theorem 3.1, we have that

$$\frac{1}{T}\sum_{t=1}^{T} \mathbb{E}\left[\frac{\|\nabla f(\boldsymbol{x}_t)\|^2}{\sqrt{\beta_2 \|\boldsymbol{v}_{t-1}\| + \zeta}}\right] \le \epsilon^2.$$
(8)

The details of the proof can be found in the Appendix B. The major idea in the proof of Lemma 3.4 is to bound the first-order.b, the second-order term and the additional term

using
$$\sum_{i=1}^{d} \mathbb{E} \left[\frac{(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}} \right].$$

Stage II: upper bound of $\mathbb{E}[\sqrt{\beta_2 \|\boldsymbol{v}_{t-1}\| + \zeta}]$. With the bounded gradient norm assumption, $\frac{1}{T} \sum_{t=1}^{T} \mathbb{E}[\|\nabla f(\boldsymbol{x}_t)\|^2] \sim \mathcal{O}(\epsilon^2)$ follows directly from Lemma 3.4. However, under the refined gradient noise assumption in this paper, the gradient norm may not be bounded. Here, we establish a key observation that $\frac{1}{T} \sum_{t=1}^{T} \mathbb{E}[\sqrt{\beta_2 \|\boldsymbol{v}_{t-1}\| + \zeta}]$ can be bounded using $\frac{1}{T} \sum_{t=1}^{T} \mathbb{E}[\|\nabla f(\boldsymbol{x}_t)\|]$. By Hölder's inequality, we have

$$\left(\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}[\|\nabla f(\boldsymbol{x}_{t})\|]\right)^{2} \leq \left(\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}[\sqrt{\beta_{2}\|\boldsymbol{v}_{t-1}\|+\zeta}]\right)$$
$$\times \left(\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}\left[\frac{\|\nabla f(\boldsymbol{x}_{t})\|^{2}}{\sqrt{\beta_{2}\|\boldsymbol{v}_{t-1}\|+\zeta}}\right]\right).$$
(9)

It is worth noting that in the RHS of (9), the second term is bound in Lemma 3.4. If the first term is upper bounded using $\frac{1}{T} \sum_{t=1}^{T} \mathbb{E}[||\nabla f(\boldsymbol{x}_t)||]$, we then can prove an upper bound on $\frac{1}{T} \sum_{t=1}^{T} \mathbb{E}[||\nabla f(\boldsymbol{x}_t)||]$, and show the algorithm converges to a stationary point. In the following lemma, we show a novel bound on $\frac{1}{T} \sum_{t=1}^{T} \mathbb{E}[\sqrt{\beta_2 ||\boldsymbol{v}_{t-1}|| + \zeta}]$.

Lemma 3.5. Let Assumption 2.2 hold. Then, we have

$$\frac{1}{T} \sum_{t=1}^{T} \mathbb{E}[\sqrt{\beta_2 \|\boldsymbol{v}_{t-1}\| + \zeta}]$$

$$\leq c + \frac{2\sqrt{dD_1}}{\sqrt{(1-\beta_2)}} \frac{\sum_{t=1}^{T} \mathbb{E}[\|\nabla f(\boldsymbol{x}_t)\|]}{T}.$$
(10)

The detailed proof can be found in the Appendix C, which recursively applies Jensen's inequality (Jensen, 1906). The proof only depends on the affine noise variance and the update process on v_t . Thus, it works for both RMSProp and Adam with (L_0, L_1) -smooth objectives.

Stage III: upper bound of $\mathbb{E}[||\nabla f(\boldsymbol{x}_t)||]$. Now we show that the algorithm converges to a stationary point. Define $e = \frac{1}{T} \sum_{t=1}^{T} \mathbb{E}[||\nabla f(\boldsymbol{x}_t)||]$. By (9), Lemma 3.4 and Lemma 3.5 we have that

$$e^2 \le \epsilon^2 \left(c + \frac{2\sqrt{dD_1}}{\sqrt{1-\beta_2}} e \right). \tag{11}$$

Thus we have

$$e \leq \left(\frac{2d\sqrt{35D_0D_1}}{\sqrt[4]{\zeta}} + \sqrt{c}\right)\epsilon$$

if $\epsilon \leq \frac{\sqrt{5dD_0}}{\sqrt{D_1}\sqrt[4]{\xi^2}}$, which indicates the algorithm converges to a stationary point.

4. Convergence Analysis of Adam

In this section, we extend our convergence analysis of RM-SProp to Adam. Such a result is attractive since empirical results on complicated tasks show Adam may perform better, e.g., the mean reward is improved by 88% via RM-SProp and 110% via Adam for Atari games (Agarwal et al., 2020).

We present the convergence result for Adam as follows.

Theorem 4.1 (Informal). Let Assumptions 2.1, 2.2 and 2.3 hold. Let $1 - \beta_2 \sim \mathcal{O}(\epsilon^2)$, $0 < \beta_1 \leq \sqrt{\beta_2} < 1$, $\eta \sim \mathcal{O}(\epsilon^2)$, and $T \sim \mathcal{O}(\epsilon^{-4})$. For small ϵ such that $\epsilon \leq \frac{\sqrt{2C_2}}{\sqrt{7\alpha_0C_6D_1}}$, we have that

$$\frac{1}{T}\sum_{t=1}^{T} \mathbb{E}[\|\nabla f(\boldsymbol{x}_t)\|] \le \left(2c + \sqrt{2c} + \frac{4\sqrt{dD_1}}{\sqrt{C_6}}\right)\epsilon, \quad (12)$$

where C_6 is a positive constant defined in Appendix H.

To the best of our knowledge, Theorem 4.1 is the first convergence result of Adam to a stationary point under the most relaxed assumptions of (L_0, L_1) -smoothness and refined gradient noise. In (Li et al., 2023; Wang et al., 2023a), the authors show that the Adam converges to a stationary point with affine noise. However, their methods only work for *L*-smooth objectives, while in this paper we focus on the (L_0, L_1) -smooth functions. Moreover, we show that for Adam, our total computational complexity matches with the optimal results (Arjevani et al., 2023), while there is an additional logarithmic term in (Li et al., 2023). The normalized momentum algorithm (Jin et al., 2021) can also be viewed as a special case of the Adam family, which applies the momentum on the first-order gradient. However, in their algorithm, a mini-batch of samples is required in each training iteration, while we do not require such a mini-batch. Thus, in the distributed setting with heterogeneous data, where the data distributions under each computational node are different, Algorithm 1 can be used directly. However, the normalized momentum in (Jin et al., 2021) may require gradient information from many computational nodes, making the problem degrade to the centralized setting.

The formal version of the theorem and detailed proof can be found in Appendix H. Below, we provide the proof sketch to underscore our key technical novelties.

Proof Sketch. Similar to the proof of RMSProp, we divided our proof into three stages. The key difference lies in Stage I, which is because of the dependence between m_t and \tilde{v}_t .

Stage I: upper bound of $\mathbb{E}\left[\frac{||\nabla f(\boldsymbol{x}_t)||^2}{\sqrt{\beta_2}||\boldsymbol{v}_{t-1}||+\zeta}\right]$. For the Adam optimizer, the first-order a term $\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_t), \frac{-\eta \boldsymbol{m}_t}{\sqrt{\tilde{\boldsymbol{v}}_t+\zeta}}\right\rangle \middle| \mathcal{F}_t\right]$ is challenging to bound due to the dependence between \boldsymbol{m}_t and $\tilde{\boldsymbol{v}}_t$. Following the recent analyses of SGDM (Liu et al., 2020) and Adam (Wang et al., 2023a), we study a potential function $f(\boldsymbol{u}_t)$ with $\boldsymbol{u}_t = \frac{\boldsymbol{x}_t - \frac{\beta_1}{\sqrt{\beta_2}}\boldsymbol{x}_{t-1}}{1 - \frac{\beta_1}{\sqrt{\beta_2}}}$. The benefit of analyzing \boldsymbol{u}_t is that we have $\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_t), \boldsymbol{u}_t - \boldsymbol{u}_{t+1}\right\rangle | \mathcal{F}_t\right] \approx \frac{(1-\beta_1)}{1 - \frac{\beta_1}{\sqrt{\beta_2}}}\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_t), \frac{\eta \boldsymbol{g}_t}{\sqrt{\tilde{\boldsymbol{v}}_t+\zeta}}\right\rangle \middle| \mathcal{F}_t\right]$, where the numerator and denominator in the last term are independent.

In Step 1.1, we will show the LHS of the descent lemma for $f(u_t)$ (shown in (80)) is lower bounded by a function of $\sum_{i=1}^{d} \frac{\eta(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta}}$; and in Step 1.2 we will show the RHS of the descent lemma of $f(\boldsymbol{u}_t)$ is upper bounded by a function of $\sum_{i=1}^{d} \frac{\eta(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta}}$. Combining the two steps, we will obtain an upper bound on $\sum_{i=1}^{d} \frac{\eta(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta}}$.

Step 1.1: lower bound of the first-order term $\mathbb{E}[\langle \nabla f(\boldsymbol{u}_t), \boldsymbol{u}_t - \boldsymbol{u}_{t+1} \rangle | \mathcal{F}_t]$. We first divide the first-order terms into two parts:

$$egin{aligned} \mathbb{E}[\langle
abla f(oldsymbol{u}_t),oldsymbol{u}_t - oldsymbol{u}_{t+1}
angle |\mathcal{F}_t] \ &= \mathbb{E}[\langle
abla f(x_t),oldsymbol{u}_t - oldsymbol{u}_{t+1}
angle |\mathcal{F}_t] \ &+ \mathbb{E}\left[\langle
abla f(oldsymbol{u}_t) -
abla f(x_t),oldsymbol{u}_t - oldsymbol{u}_{t+1}
angle |\mathcal{F}_t]. \end{aligned}$$

The first part mimics the first-order term in the proof of RM-SProp and the second one is due to a mismatch between u_t and x_t . By bounding these two parts separately, we have a lemma similar to Lemma 3.3 but with two additional terms: $\mathbb{E}\left[\frac{m_{t,i}^2}{v_{t,i}+\zeta}\Big|\mathcal{F}_t\right]$ and $\mathbb{E}\left[\frac{m_{t,i}^2}{\sqrt{v_{t,i}+\zeta}}\Big|\mathcal{F}_t\right]$ due to the employment of the first-order momentum (see details in Lemma F.1). For the additional terms, (Wang et al., 2023a) showed that $\sum_{t=1}^{T} \frac{m_{t,i}^2}{v_{t,i}}$ can be bounded by a function of $v_{T,i}$ (shown in Lemma E.2). It is worth noting that due to the (L_0, L_1) -smoothness our objective is harder to bound and only Lemma E.2 is not enough. Here, we bound the $\sum_{t=1}^{T} \frac{m_{t,i}^2}{\sqrt{v_{t,i}}}$ term by a function of $\frac{\sqrt{v_{T,i}} - \sqrt{v_{0,i}}}{1-\beta_2} + \sum_{t=1}^{T} \sqrt{v_{t-1,i}}$ (the details can be found in Lemma E.3).

Step 1.2: upper bound on second-order and additional terms. Based on the update process of u_t and x_t , we bound the second-order and additional terms similarly to those in (6) and (7), but with g_t replaced by m_t (see details in (81) and (82)). Unlike in the proof of RMSProp, where $\mathbb{E}\left[\frac{g_{t,i}^2}{\sqrt{\beta_2 v_{t,i} + \zeta}} \middle| \mathcal{F}_t\right]$ can be bounded by $\frac{D_0}{\sqrt{\zeta}} + \frac{D_1(\partial_i f(x_t))^2}{\sqrt{v_{t,i} + \zeta}}$, Assumption 2.2 does not hold for m_t and this is the reason why two terms $\mathbb{E}\left[\frac{m_{t,i}^2}{v_{t,i} + \zeta} \middle| \mathcal{F}_t\right]$ and $\mathbb{E}\left[\frac{m_{t,i}^2}{\sqrt{v_{t,i} + \zeta}} \middle| \mathcal{F}_t\right]$ are kept in the upper bound of the second-order and additional terms. Then based on the descent lemma of $f(u_t)$, we can show $\mathbb{E}\left[\sum_{i=1}^d \frac{\eta(\partial_i f(x_t))^2}{\sqrt{\beta_2 v_{t-1,i} + \zeta}}\right]$ can be upper bounded by a function of $\mathbb{E}[\nabla f(x_t)]$. We then get the following lemma:

Lemma 4.2. Let Assumptions 2.1, 2.2 and 2.3 hold. With the same parameters as in Theorem 4.1, we have that

$$\frac{1}{T}\sum_{t=1}^{T} \mathbb{E}\left[\frac{\|\nabla f(\boldsymbol{x}_t)\|^2}{\sqrt{\beta_2 \|\boldsymbol{v}_{t-1}\| + \zeta}}\right] \le \epsilon^2 + \frac{\epsilon}{T}\sum_{t=1}^{T} \mathbb{E}\left[\|\nabla f(\boldsymbol{x}_t)\|\right].$$

The details of the proof can be found in the Appendix G.

Stage II is the same with the proof of RMSProp and thus is omitted here.

Stage III: upper bound of $\mathbb{E}[||\nabla f(\boldsymbol{x}_t)||]$. As we mentioned before, Lemma 3.5 and (9) hold for Adam. It is worth noting that in the RHS of (9), the first term is bounded in Lemma 4.2, which is more complicated than Lemma 3.4 and has an additional term. Let $e = \frac{1}{T} \sum_{t=1}^{T} \mathbb{E}[||\nabla f(\boldsymbol{x}_t)||]$. By (9), Lemma 4.2 and Lemma 3.5 we have that

$$e^{2} \leq c\epsilon^{2} + \frac{2\sqrt{dD_{1}}}{\sqrt{C_{6}\epsilon}}e\epsilon^{2} + ce\epsilon + \frac{e^{2}}{2}.$$
 (13)

Thus, $e \sim \mathcal{O}(\epsilon)$, which shows the algorithm converges to a stationary point.

5. Conclusion

In this paper, we provide the tightest convergence analyses for RMSProp and Adam for non-convex objectives under the most relaxed assumptions of generalized smoothness and refined gradient noise. The complexity to achieve an ϵ stationary point for both algorithms is shown to be $\mathcal{O}(\epsilon^{-4})$, which matches with the lower bound for first-order algorithms established in (Arjevani et al., 2023).

References

- Agarwal, R., Schuurmans, D., and Norouzi, M. An optimistic perspective on offline reinforcement learning. In *International Conference on Machine Learning*, pp. 104– 114. PMLR, 2020.
- Arjevani, Y., Carmon, Y., Duchi, J. C., Foster, D. J., Srebro, N., and Woodworth, B. Lower bounds for nonconvex stochastic optimization. *Mathematical Programming*, 199(1-2):165–214, 2023.
- Bernstein, J., Wang, Y.-X., Azizzadenesheli, K., and Anandkumar, A. signsgd: Compressed optimisation for non-convex problems. In *International Conference on Machine Learning*, pp. 560–569. PMLR, 2018.
- Bottou, L., Curtis, F. E., and Nocedal, J. Optimization methods for large-scale machine learning. *SIAM review*, 60(2):223–311, 2018.
- Brock, A., Donahue, J., and Simonyan, K. Large scale gan training for high fidelity natural image synthesis. *arXiv* preprint arXiv:1809.11096, 2018.
- Brown, T., Mann, B., Ryder, N., Subbiah, M., Kaplan, J. D., Dhariwal, P., Neelakantan, A., Shyam, P., Sastry, G., Askell, A., et al. Language models are few-shot learners. *Advances in Neural Information Processing Systems*, 33: 1877–1901, 2020.
- Bubeck, S. et al. Convex optimization: Algorithms and complexity. *Foundations and Trends*® *in Machine Learning*, 8(3-4):231–357, 2015.
- Chen, X., Liu, S., Sun, R., and Hong, M. On the convergence of a class of Adam-type algorithms for non-convex optimization. *arXiv preprint arXiv:1808.02941*, 2018.
- Chen, Z., Zhou, Y., Liang, Y., and Lu, Z. Generalizedsmooth nonconvex optimization is as efficient as smooth nonconvex optimization. *arXiv preprint arXiv:2303.02854*, 2023.
- Crawshaw, M., Liu, M., Orabona, F., Zhang, W., and Zhuang, Z. Robustness to unbounded smoothness of generalized signsgd. *Advances in Neural Information Processing Systems*, 35:9955–9968, 2022.
- Cutkosky, A. and Mehta, H. Momentum improves normalized SGD. In *International Conference on Machine Learning*, pp. 2260–2268. PMLR, 2020.
- De, S., Mukherjee, A., and Ullah, E. Convergence guarantees for RMSProp and Adam in non-convex optimization and an empirical comparison to Nesterov acceleration. *arXiv preprint arXiv:1807.06766*, 2018.

- Défossez, A., Bottou, L., Bach, F., and Usunier, N. A simple convergence proof of Adam and Adagrad. *arXiv* preprint arXiv:2003.02395, 2020.
- Dosovitskiy, A., Beyer, L., Kolesnikov, A., Weissenborn, D., Zhai, X., Unterthiner, T., Dehghani, M., Minderer, M., Heigold, G., Gelly, S., et al. An image is worth 16x16 words: Transformers for image recognition at scale. arXiv preprint arXiv:2010.11929, 2020.
- Duchi, J., Hazan, E., and Singer, Y. Adaptive subgradient methods for online learning and stochastic pptimization. *Journal of Machine Learning Research*, 12(7), 2011.
- Faw, M., Tziotis, I., Caramanis, C., Mokhtari, A., Shakkottai, S., and Ward, R. The power of adaptivity in SGD: Self-tuning step sizes with unbounded gradients and affine variance. In *Conference on Learning Theory*, pp. 313–355. PMLR, 2022.
- Faw, M., Rout, L., Caramanis, C., and Shakkottai, S. Beyond uniform smoothness: A stopped analysis of adaptive SGD. arXiv preprint arXiv:2302.06570, 2023.
- Foster, D. J., Sekhari, A., Shamir, O., Srebro, N., Sridharan, K., and Woodworth, B. The complexity of making the gradient small in stochastic convex optimization. In *Conference on Learning Theory*, pp. 1319–1345. PMLR, 2019.
- Ghadimi, S. and Lan, G. Stochastic first-and zeroth-order methods for nonconvex stochastic programming. SIAM Journal on Optimization, 23(4):2341–2368, 2013.
- Ghadimi, S., Lan, G., and Zhang, H. Mini-batch stochastic approximation methods for nonconvex stochastic composite optimization. *Mathematical Programming*, 155 (1-2):267–305, 2016.
- Guo, Z., Xu, Y., Yin, W., Jin, R., and Yang, T. A novel convergence analysis for algorithms of the Adam family and beyond. arXiv preprint arXiv:2104.14840, 2021.
- Hardy, G. H., Littlewood, J. E., and Pólya, G. *Inequalities*. Cambridge university press, 1952.
- Hinton, G., Srivastava, N., and Swersky, K. Lecture 6da separate, adaptive learning rate for each connection. *Slides of Lecture Neural Networks for Machine Learning*, 1:1–31, 2012.
- Hong, Y. and Lin, J. High probability convergence of Adam under unbounded gradients and affine variance noise. arXiv preprint arXiv:2311.02000, 2023.
- Jensen, J. L. W. V. Sur les fonctions convexes et les inégalités entre les valeurs moyennes. Acta Mathematica, 30:175 – 193,

1906. doi: 10.1007/BF02418571. URL https://doi.org/10.1007/BF02418571.

- Jin, J., Zhang, B., Wang, H., and Wang, L. Non-convex distributionally robust optimization: Non-asymptotic analysis. Advances in Neural Information Processing Systems, 34:2771–2782, 2021.
- Khaled, A. and Richtárik, P. Better theory for sgd in the nonconvex world. *arXiv preprint arXiv:2002.03329*, 2020.
- Kingma, D. P. and Ba, J. Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980*, 2014.
- Li, H., Jadbabaie, A., and Rakhlin, A. Convergence of Adam under relaxed assumptions. *arXiv preprint arXiv:2304.13972*, 2023.
- Liu, Y., Gao, Y., and Yin, W. An improved analysis of stochastic gradient descent with momentum. Advances in Neural Information Processing Systems, 33:18261– 18271, 2020.
- Nemirovskij, A. S. and Yudin, D. B. Problem complexity and method efficiency in optimization. 1983.
- Reisizadeh, A., Li, H., Das, S., and Jadbabaie, A. Variancereduced clipping for non-convex optimization. arXiv preprint arXiv:2303.00883, 2023.
- Richtárik, P. and Takáč, M. Iteration complexity of randomized block-coordinate descent methods for minimizing a composite function. *Mathematical Programming*, 144 (1-2):1–38, 2014.
- Shi, N., Li, D., Hong, M., and Sun, R. RMSProp converges with proper hyper-parameter. In *International Conference on Learning Representations*, 2020.
- Vaswani, S., Bach, F., and Schmidt, M. Fast and faster convergence of SGD for over-parameterized models and an accelerated perceptron. In *International Conference* on Artificial Intelligence and Atatistics, pp. 1195–1204. PMLR, 2019.
- Wang, B., Zhang, Y., Zhang, H., Meng, Q., Ma, Z.-M., Liu, T.-Y., and Chen, W. Provable adaptivity in Adam. arXiv preprint arXiv:2208.09900, 2022.
- Wang, B., Fu, J., Zhang, H., Zheng, N., and Chen, W. Closing the gap between the upper bound and the lower bound of Adam's iteration complexity. *arXiv preprint arXiv:2310.17998*, 2023a.
- Wang, B., Zhang, H., Ma, Z., and Chen, W. Convergence of Adagrad for non-convex objectives: Simple proofs and relaxed assumptions. In *Conference on Learning Theory*, pp. 161–190. PMLR, 2023b.

- Ward, R., Wu, X., and Bottou, L. Adagrad stepsizes: Sharp convergence over nonconvex landscapes. *The Journal of Machine Learning Research*, 21(1):9047–9076, 2020.
- Zaheer, M., Reddi, S., Sachan, D., Kale, S., and Kumar, S. Adaptive methods for nonconvex optimization. Advances in Neural Information Processing Systems, 31, 2018.
- Zhang, B., Jin, J., Fang, C., and Wang, L. Improved analysis of clipping algorithms for non-convex optimization. *Advances in Neural Information Processing Systems*, 33: 15511–15521, 2020.
- Zhang, J., He, T., Sra, S., and Jadbabaie, A. Why gradient clipping accelerates training: A theoretical justification for adaptivity. arXiv preprint arXiv:1905.11881, 2019.
- Zhang, Y., Chen, C., Shi, N., Sun, R., and Luo, Z.-Q. Adam can converge without any modification on update rules. *Advances in Neural Information Processing Systems*, 35: 28386–28399, 2022.
- Zou, F., Shen, L., Jie, Z., Zhang, W., and Liu, W. A sufficient condition for convergences of Adam and RMSProp. In *Proceedings of the IEEE/CVF Conference on computer vision and pattern recognition*, pp. 11127–11135, 2019.

A. Formal Version of Lemma 3.3 and Its Proof

Lemma A.1. Under Assumptions 2.2 and 2.3, for any $\alpha_0, \alpha_1 > 0, \mathbf{x}_0 = \mathbf{x}_1$ and t > 0, we have that:

$$\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \left[\frac{\eta}{\sqrt{\boldsymbol{v}_{t}+\zeta}} - \frac{\eta}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1}+\zeta}}\right] \odot \boldsymbol{g}_{t}\right\rangle \left|\mathcal{F}_{t}\right] \\
\geq -\left[\sum_{i=1}^{d} \frac{\eta(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{0}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} + \sum_{i=1}^{d} \frac{\eta\alpha_{0}D_{0}}{2}\mathbb{E}\left[\frac{1}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}\right]\mathcal{F}_{t}\right] \\
+ \sum_{i=1}^{d} \frac{\eta\alpha_{0}D_{1}}{2}\mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}\right]\mathcal{F}_{t}\right] \\
+ \sum_{i=1}^{d} \frac{\eta\alpha_{0}D_{1}}{2\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}\left(\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\alpha_{1}D_{1}} + \frac{\alpha_{1}dD_{1}L_{0}^{2}\eta^{2}}{1-\beta_{2}} + \frac{2\sqrt{d}\eta L_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{1-\beta_{2}}}\right)\right]$$
(14)

Proof. Since $v_t = \beta_2 v_{t-1} + (1 - \beta_2) g_t \odot g_t$, for any $i \in [d]$, we have that

$$\begin{pmatrix} -\frac{1}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}} + \frac{1}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta}} \end{pmatrix} \\ = \frac{(\boldsymbol{v}_{t,i}-\beta_2 \boldsymbol{v}_{t-1,i})}{(\sqrt{\boldsymbol{v}_{t,i}+\zeta})(\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta})(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta})} \\ = \frac{(1-\beta_2)\boldsymbol{g}_{t,i}^2}{(\sqrt{\boldsymbol{v}_{t,i}+\zeta})(\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta})(\sqrt{\boldsymbol{v}_{t,i}}+\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}})}.$$
(15)

Thus, the LHS of (14) can be bounded as

$$\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \left[\frac{-\eta}{\sqrt{\boldsymbol{v}_{t}+\zeta}} + \frac{\eta}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1}+\zeta}}\right] \odot \boldsymbol{g}_{t}\right\rangle |\mathcal{F}_{t}\right] \\
\leq \sum_{i=1}^{d} \mathbb{E}\left[\eta |\partial_{i}f(\boldsymbol{x}_{t})||\boldsymbol{g}_{t,i}| \left|\frac{-1}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}} + \frac{1}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}\right| |\mathcal{F}_{t}\right] \\
\leq \sum_{i=1}^{d} \mathbb{E}\left[\eta |\partial_{i}f(\boldsymbol{x}_{t})||\boldsymbol{g}_{t,i}| \left|\frac{(1-\beta_{2})\boldsymbol{g}_{t,i}^{2}}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}\right| |\mathcal{F}_{t}\right] \\
\leq \sum_{i=1}^{d} \eta \frac{|\partial_{i}f(\boldsymbol{x}_{t})|}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} \mathbb{E}\left[\left|\frac{\sqrt{1-\beta_{2}}\boldsymbol{g}_{t,i}^{2}}{(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta)}}\right| |\mathcal{F}_{t}\right],$$
(16)

where the second inequality is due to (15) and the last inequality is because $\frac{|g_{t,i}|}{\sqrt{v_{t,i}+\zeta}} \leq \frac{1}{\sqrt{1-\beta_2}}$. For any $a, b \in \mathbb{R}$, we have $ab \leq \frac{a^2+b^2}{2}$. For any $\alpha_0 > 0$, the RHS of (16) can be further bounded as

$$\eta \frac{|\partial_{i} f(\boldsymbol{x}_{t})|}{\sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta}} \mathbb{E}\left[\left\| \frac{\sqrt{1 - \beta_{2}} \boldsymbol{g}_{t,i}^{2}}{(\sqrt{\boldsymbol{v}_{t,i} + \zeta} + \sqrt{\beta_{2}} \boldsymbol{v}_{t-1,i} + \zeta)} \right\| |\mathcal{F}_{t} \right]$$

$$\leq \frac{\eta (\partial_{i} f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{0}\sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta}} + \frac{\eta \alpha_{0}}{2\sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta}} \left(\mathbb{E}\left[\frac{\sqrt{1 - \beta_{2}} \boldsymbol{g}_{t,i}^{2}}{(\sqrt{\boldsymbol{v}_{t,i} + \zeta} + \sqrt{\beta_{2}} \boldsymbol{v}_{t-1,i} + \zeta)} |\mathcal{F}_{t} \right] \right)^{2}.$$
(17)

For the last term of (17), due to Hölder's inequality, we have that

$$\frac{\eta}{2\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta}} \left(\mathbb{E}\left[\frac{\sqrt{1-\beta_2}\boldsymbol{g}_{t,i}^2}{(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta})}|\mathcal{F}_t\right] \right)^2$$

Convergence Guarantees for RMSProp and Adam

$$\leq \frac{(1-\beta_2)\eta}{2\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta}} \mathbb{E}[\boldsymbol{g}_{t,i}^2|\mathcal{F}_t] \mathbb{E}\left[\frac{\boldsymbol{g}_{t,i}^2}{(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta})^2}|\mathcal{F}_t\right].$$
(18)

Based on Assumption 2.2, the RKS of (18) can be further bounded as

$$\frac{(1-\beta_{2})\eta}{2\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}\mathbb{E}[\boldsymbol{g}_{t,i}^{2}|\mathcal{F}_{t}]\mathbb{E}\left[\frac{\boldsymbol{g}_{t,i}^{2}}{(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta})^{2}}|\mathcal{F}_{t}\right]$$

$$\leq \frac{\eta}{2}(D_{0}+D_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2})\times\mathbb{E}\left[\frac{(1-\beta_{2})\boldsymbol{g}_{t,i}^{2}}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta})}|\mathcal{F}_{t}\right]$$

$$\leq \frac{\eta}{2}(D_{0}+D_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2})\times\mathbb{E}\left[\frac{\boldsymbol{v}_{t,i}+\zeta\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta})}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta})}|\mathcal{F}_{t}\right]$$

$$\leq \frac{\eta}{2}(D_{0}+D_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2})\mathbb{E}\left[\frac{1}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}-\frac{1}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_{t}\right]$$
(19)

Thus for t = 1, combining (16), (17), (18) and (19), we have that

$$\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \left[\frac{-\eta}{\sqrt{\boldsymbol{v}_{t}+\zeta}} + \frac{\eta}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1}+\zeta}}\right] \odot \boldsymbol{g}_{t}\right\rangle |\mathcal{F}_{t}\right] \\
\leq \sum_{i=1}^{d} \frac{\eta(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{0}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} + \sum_{i=1}^{d} \frac{\eta\alpha_{0}}{2}(D_{0} + D_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2})\mathbb{E}\left[\frac{1}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_{t}\right]$$
(20)

For t > 1, (19) can be further bounded as

$$\frac{(1-\beta_2)\eta}{2\sqrt{\beta_2\boldsymbol{v}_{t-1,i}+\zeta}} \mathbb{E}[\boldsymbol{g}_{t,i}^2|\mathcal{F}_t] \mathbb{E}\left[\frac{\boldsymbol{g}_{t,i}^2}{(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_2\boldsymbol{v}_{t-1,i}+\zeta})^2}|\mathcal{F}_t\right] \\
\leq \frac{\eta D_0}{2} \mathbb{E}\left[\frac{1}{\sqrt{\beta_2\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_t\right] + \frac{\eta D_1}{2} \mathbb{E}\left[\frac{(\partial_i f(\boldsymbol{x}_{t-1}))^2}{\sqrt{\beta_2\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_t\right] \\
+ \frac{\eta D_1}{2} \mathbb{E}\left[\frac{(\partial_i f(\boldsymbol{x}_t))^2 - (\partial_i f(\boldsymbol{x}_{t-1}))^2}{\sqrt{\beta_2\boldsymbol{v}_{t-1,i}+\zeta}}|\mathcal{F}_t\right].$$
(21)

For the last term in (21) with (L_0, L_1) -smooth objectives and any $\alpha_1 > 0$, we have that

$$\frac{\eta D_{1}}{2} \mathbb{E} \left[\frac{(\partial_{i} f(\boldsymbol{x}_{t}))^{2} - (\partial_{i} f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta}} |\mathcal{F}_{t} \right] \\
\leq \frac{\eta D_{1}}{2} \mathbb{E} \left[\frac{2|\partial_{i} f(\boldsymbol{x}_{t})| ||\partial_{i} f(\boldsymbol{x}_{t})| - |\partial_{i} f(\boldsymbol{x}_{t-1})||}{\sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta}} |\mathcal{F}_{t} \right] \\
\leq \eta D_{1} \frac{|\partial_{i} f(\boldsymbol{x}_{t})| (L_{0} + L_{1} |\partial_{i} f(\boldsymbol{x}_{t})|) \eta \left\| \frac{1}{\sqrt{\boldsymbol{v}_{t-1} + \zeta}} \odot \boldsymbol{g}_{t-1} \right\|}{\sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta}} \\
\leq \frac{\eta D_{1}}{2\sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta}} \left(\frac{(\partial_{i} f(\boldsymbol{x}_{t}))^{2}}{\alpha_{1} D_{1}} + \alpha_{1} D_{1} L_{0}^{2} \eta^{2} \sum_{i=1}^{d} \frac{(\boldsymbol{g}_{t-1,i})^{2}}{\boldsymbol{v}_{t-1,i} + \zeta} \right) \\
+ \frac{\eta^{2} L_{1} D_{1} \sqrt{d} (\partial_{i} f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta}} \\
\leq \frac{\eta D_{1}}{2\sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta}} \left(\frac{(\partial_{i} f(\boldsymbol{x}_{t}))^{2}}{\alpha_{1} D_{1}} + \frac{\alpha_{1} d D_{1} L_{0}^{2} \eta^{2}}{1 - \beta_{2}} + \frac{2\sqrt{d} \eta L_{1} (\partial_{i} f(\boldsymbol{x}_{t}))^{2}}{\sqrt{1 - \beta_{2}}} \right).$$
(22)

where the first inequality is due to the fact that for any $a, b \ge 0$, we have $a^2 - b^2 \le 2a|a - b|$, the second inequality is by the (L_0, L_1) -smoothness and the third one is because $\frac{|g_{t,i}|}{\sqrt{v_{t,i}+\zeta}} \le \frac{1}{\sqrt{1-\beta_2}}$. Based on (16), (17), (18), (21) and (22), for t > 1 we can get that

$$\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \left[\frac{-\eta}{\sqrt{\boldsymbol{v}_{t}+\zeta}} + \frac{\eta}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1}+\zeta}}\right] \odot \boldsymbol{g}_{t}\right\rangle |\mathcal{F}_{t}\right] \\
\leq \sum_{i=1}^{d} \frac{\eta(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{0}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} + \sum_{i=1}^{d} \frac{\eta\alpha_{0}D_{0}}{2} \mathbb{E}\left[\frac{1}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_{t}\right] \\
+ \sum_{i=1}^{d} \frac{\eta\alpha_{0}D_{1}}{2} \mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_{t}\right] \\
+ \sum_{i=1}^{d} \frac{\eta\alpha_{0}D_{1}}{2\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} \left(\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\alpha_{1}D_{1}} + \frac{\alpha_{1}dD_{1}L_{0}^{2}\eta^{2}}{1-\beta_{2}} + \frac{2\sqrt{d}\eta L_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{1-\beta_{2}}}\right).$$
(23)

Since we set $x_0 = x_1$, (23) also holds for t = 0. We then completes the proof.

B. Proof of Lemma 3.4

For (L_0, L_1) -smooth objective functions, we have the following descent inequality (Lemma 1 in (Crawshaw et al., 2022)):

$$\underbrace{\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \frac{\eta}{\sqrt{\tilde{\boldsymbol{v}}_{t}+\zeta}} \odot \boldsymbol{g}_{t} \right\rangle |\mathcal{F}_{t}\right]}_{\text{First Order.a}} \leq f(\boldsymbol{x}_{t}) - \mathbb{E}[f(\boldsymbol{x}_{t+1})|\mathcal{F}_{t}] + \underbrace{\sum_{i=1}^{d} \frac{L_{0}}{2\sqrt{d}} \mathbb{E}[\|\boldsymbol{x}_{t+1} - \boldsymbol{x}_{t}\||\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}||\mathcal{F}_{t}]}_{\text{Second Order}} + \underbrace{\sum_{i=1}^{d} \frac{L_{1}|\partial_{i}f(\boldsymbol{x}_{t})|}{2} \mathbb{E}[\|\boldsymbol{x}_{t+1} - \boldsymbol{x}_{t}\||\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}||\mathcal{F}_{t}]}_{\text{Additional Term}} + \underbrace{\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \left[\frac{-\eta}{\sqrt{\boldsymbol{v}_{t}+\zeta}} + \frac{\eta}{\sqrt{\tilde{\boldsymbol{v}}_{t}+\zeta}}\right] \odot \boldsymbol{g}_{t}\right\rangle |\mathcal{F}_{t}\right]}_{\text{First Order.b}}.$$
(24)

For the first-order main item, given \mathcal{F}_t , we have \boldsymbol{g}_t independent of $\tilde{\boldsymbol{v}}_t$. It then follows that $\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_t), \frac{\eta}{\sqrt{\tilde{\boldsymbol{v}}_t + \zeta}} \odot \boldsymbol{g}_t \right\rangle | \mathcal{F}_t\right] = \sum_{i=1}^d \frac{\eta(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}}$. Based on Lemma 3.3, we can bound the first-order error term. For any $\alpha_0, \alpha_1 > 0$ and t > 1, we then have:

$$\begin{split} &\sum_{i=1}^{d} \frac{\eta(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}} \\ \leq & f(\boldsymbol{x}_t) - \mathbb{E}[f(\boldsymbol{x}_{t+1})|\mathcal{F}_t] + \underbrace{\sum_{i=1}^{d} \frac{L_0}{2\sqrt{d}} \mathbb{E}[\|\boldsymbol{x}_{t+1} - \boldsymbol{x}_t\| |\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}| |\mathcal{F}_t]}_{\text{Second Order}} \\ &+ \underbrace{\sum_{i=1}^{d} \frac{L_1 |\partial_i f(\boldsymbol{x}_t)|}{2} \mathbb{E}[\|\boldsymbol{x}_{t+1} - \boldsymbol{x}_t\| |\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}| |\mathcal{F}_t]}_{\text{Additional Term}} \\ &+ \underbrace{\sum_{i=1}^{d} \frac{\eta(\partial_i f(\boldsymbol{x}_t))^2}{2\alpha_0 \sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}} + \sum_{i=1}^{d} \frac{\eta\alpha_0 D_0}{2} \mathbb{E}\left[\frac{1}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} |\mathcal{F}_t\right]} \right] \end{split}$$

Convergence Guarantees for RMSProp and Adam

$$+\sum_{i=1}^{d} \frac{\eta \alpha_{0} D_{1}}{2} \mathbb{E} \left[\frac{(\partial_{i} f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta}} - \frac{(\partial_{i} f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} |\mathcal{F}_{t} \right] \\ +\sum_{i=1}^{d} \frac{\eta \alpha_{0} D_{1}}{2\sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta}} \left(\frac{(\partial_{i} f(\boldsymbol{x}_{t}))^{2}}{\alpha_{1} D_{1}} + \frac{\alpha_{1} d D_{1} L_{0}^{2} \eta^{2}}{1 - \beta_{2}} + \frac{2\sqrt{d} \eta L_{1} (\partial_{i} f(\boldsymbol{x}_{t}))^{2}}{\sqrt{1 - \beta_{2}}} \right).$$
(25)

Now we focus on the second-order term, which can be bounded as follows

$$\sum_{i=1}^{d} \frac{L_{0}}{2\sqrt{d}} \mathbb{E}[\|\boldsymbol{x}_{t+1} - \boldsymbol{x}_{t}\| \| \boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i} \| |\mathcal{F}_{t}]$$

$$\leq \sum_{i=1}^{d} \frac{L_{0}}{2\sqrt{d}} \mathbb{E}\left[\frac{\|\boldsymbol{x}_{t+1} - \boldsymbol{x}_{t}\|^{2}}{2\sqrt{d}} + \frac{\sqrt{d}}{2}|\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}|^{2}\mathcal{F}_{t}\right]$$

$$\leq \sum_{i=1}^{d} \frac{L_{0}}{2} \mathbb{E}\left[\eta^{2} \frac{\boldsymbol{g}_{t,i}^{2}}{\boldsymbol{v}_{t,i} + \zeta} |\mathcal{F}_{t}\right]$$

$$\leq \sum_{i=1}^{d} \frac{L_{0}\eta^{2}}{2} \mathbb{E}\left[\eta^{2} \frac{\boldsymbol{g}_{t,i}^{2}}{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta} |\mathcal{F}_{t}\right]$$

$$\leq \sum_{i=1}^{d} \frac{L_{0}\eta^{2}}{2} \frac{D_{0} + D_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}$$

$$\leq \sum_{i=1}^{d} \frac{L_{0}\eta^{2}}{2\sqrt{\zeta}} \frac{D_{0} + D_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}},$$
(26)

where the fourth inequality is due to Assumption 2.2. For the addition term due to (L_0, L_1) -smooth objectives, based on Assumption 2.2, for any $\alpha_2 > 0$ we have that

$$\frac{L_{1}|\partial_{i}f(\boldsymbol{x}_{t})|}{2}\mathbb{E}[\|\boldsymbol{x}_{t+1} - \boldsymbol{x}_{t}\||\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}||\mathcal{F}_{t}] \\
\leq \frac{\sqrt{d}L_{1}|\partial_{i}f(\boldsymbol{x}_{t})|}{2\sqrt{1 - \beta_{2}}}\mathbb{E}\left[\eta^{2}\frac{|\boldsymbol{g}_{t,i}|}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}}|\mathcal{F}_{t}\right] \\
\leq \frac{\eta^{2}\sqrt{d}L_{1}|\partial_{i}f(\boldsymbol{x}_{t})|}{2\sqrt{1 - \beta_{2}}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}}\sqrt{\mathbb{E}\left[\boldsymbol{g}_{t,i}^{2}|\mathcal{F}_{t}\right]} \\
\leq \frac{\eta^{2}\sqrt{d}L_{1}|\partial_{i}f(\boldsymbol{x}_{t})|(\sqrt{D_{0}} + \sqrt{D_{1}}|\partial_{i}f(\boldsymbol{x}_{t})|)}{2\sqrt{1 - \beta_{2}}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} + \frac{\eta^{2}\sqrt{d}(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\sqrt{1 - \beta_{2}}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} + \frac{\eta^{2}\sqrt{d}\alpha_{2}L_{1}^{2}D_{0}}{2\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}},$$
(27)

where the first inequality is due to $\|\boldsymbol{x}_{t+1} - \boldsymbol{x}_t\| \leq \frac{\sqrt{d\eta}}{\sqrt{1-\beta_2}}$, the second inequality is because that $\mathbb{E}\left[\boldsymbol{g}_{t,i}|\mathcal{F}_t\right] \leq \sqrt{\mathbb{E}\left[\boldsymbol{g}_{t,i}^2|\mathcal{F}_t\right]}$, the third inequality is due to Assumption 2.3 and the fact that for any $a, b \geq 0$, we have $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$. Plug (26) and (27) into (25), and it follows that

$$\begin{split} &\sum_{i=1}^{d} \frac{\eta(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} \\ &\leq f(\boldsymbol{x}_{t}) - \mathbb{E}[f(\boldsymbol{x}_{t+1})|\mathcal{F}_{t}] + \sum_{i=1}^{d} \frac{L_{0}\eta^{2}}{2\sqrt{\zeta}} \frac{D_{0} + D_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}. \\ &+ \sum_{i=1}^{d} \left(\frac{\eta^{2}\sqrt{d}L_{1}\sqrt{D_{1}}(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\sqrt{1-\beta_{2}}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} + \frac{\eta^{2}\sqrt{d}(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2(1-\beta_{2})\alpha_{2}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} + \frac{\eta^{2}\sqrt{d}\alpha_{2}L_{1}^{2}D_{0}}{2\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} \right) \end{split}$$

$$+\sum_{i=1}^{d} \frac{\eta(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{0}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} + \sum_{i=1}^{d} \frac{\eta\alpha_{0}D_{0}}{2}\mathbb{E}\left[\frac{1}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_{t}\right] \\ +\sum_{i=1}^{d} \frac{\eta\alpha_{0}D_{1}}{2}\mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_{t}\right] \\ +\sum_{i=1}^{d} \frac{\eta\alpha_{0}D_{1}}{2\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}\left(\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\alpha_{1}D_{1}} + \frac{\alpha_{1}dD_{1}L_{0}^{2}\eta^{2}}{1-\beta_{2}} + \frac{2\sqrt{d}\eta L_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{1-\beta_{2}}}\right).$$
(28)

which can be written as

$$\sum_{i=1}^{d} \left(\eta - \frac{\eta}{2\alpha_{0}} - \frac{\eta\alpha_{0}}{2\alpha_{1}} - \frac{L_{0}\eta^{2}D_{1}}{2\sqrt{\zeta}} - \frac{\eta^{2}L_{1}\sqrt{dD_{1}}}{2\sqrt{1-\beta_{2}}} - \frac{\eta^{2}\sqrt{d}}{2(1-\beta_{2})\alpha_{2}} - \frac{\eta^{2}\sqrt{d}\alpha_{0}L_{1}D_{1}}{\sqrt{1-\beta_{2}}} \right) \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}$$

$$\leq f(\boldsymbol{x}_{t}) - \mathbb{E}[f(\boldsymbol{x}_{t+1})|\mathcal{F}_{t}] + \frac{dL_{0}\eta^{2}D_{0}}{2\zeta} + \frac{\alpha_{0}\alpha_{1}d^{2}L_{0}^{2}\eta^{3}D_{1}^{2}}{2(1-\beta_{2})} \frac{1}{\sqrt{\zeta}} + \frac{\eta^{2}\alpha_{2}d^{1.5}L_{1}^{2}D_{0}}{2\sqrt{\zeta}}$$

$$+ \sum_{i=1}^{d} \frac{\eta\alpha_{0}D_{0}}{2} \mathbb{E}\left[\frac{1}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_{t}\right]$$

$$+ \sum_{i=1}^{d} \frac{\eta\alpha_{0}D_{1}}{2} \mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_{t}\right]$$
(29)

It is worth noting that the sum of last two terms from t = 1 to T can be further bounded. Specifically, for any $i \in [d]$ we have that

$$\sum_{t=1}^{T} \mathbb{E} \left[\frac{1}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} \right]$$
$$= \mathbb{E} \left[\frac{1}{\sqrt{\beta_2 \boldsymbol{v}_{0,i} + \zeta}} \right] + \sum_{t=1}^{T-1} \mathbb{E} \left[\frac{1}{\sqrt{\beta_2 \boldsymbol{v}_{t,i} + \zeta}} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} \right] - \mathbb{E} \left[\frac{1}{\sqrt{\boldsymbol{v}_{T,i} + \zeta}} \right]$$
$$\leq \mathbb{E} \left[\frac{1}{\sqrt{\zeta}} \right] + \sum_{t=1}^{T-1} \mathbb{E} \left[\frac{1}{\sqrt{\beta_2 \boldsymbol{v}_{t,i} + \zeta}} - \frac{\sqrt{\beta_2}}{\sqrt{\beta_2 \boldsymbol{v}_{t,i} + \zeta}} \right]$$
$$= \frac{1}{\sqrt{\zeta}} + \sum_{t=1}^{T-1} \mathbb{E} \left[\frac{1 - \sqrt{\beta_2}}{\sqrt{\beta_2 \boldsymbol{v}_{t,i} + \zeta}} \right]$$
$$\leq \frac{1}{\sqrt{\zeta}} + T \frac{1 - \sqrt{\beta_2}}{\sqrt{\zeta}}. \tag{30}$$

Similarly, for the last term, we have that

$$\mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{0}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{0,i}+\zeta}} - \frac{(\partial_{i}f(\boldsymbol{x}_{1}))^{2}}{\sqrt{\boldsymbol{v}_{1,i}+\zeta}}\right] + \sum_{t=2}^{T}\mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}\right] \\
= \mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{1}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{0,i}+\zeta}}\right] + \sum_{t=1}^{T-1}\mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t,i}+\zeta}} - \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}\right] - \mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{T,i}+\zeta}}\right] \\
\leq \mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{1}))^{2}}{\sqrt{\zeta}}\right] + \sum_{t=1}^{T-1}\mathbb{E}\left[\left(\frac{1}{\sqrt{\beta_{2}\boldsymbol{v}_{t,i}+\zeta}} - \frac{\sqrt{\beta_{2}}}{\sqrt{\beta_{2}\boldsymbol{v}_{t,i}+\zeta}}\right)(\partial_{i}f(\boldsymbol{x}_{t}))^{2}\right] \\
= \frac{(\partial_{i}f(\boldsymbol{x}_{1}))^{2}}{\sqrt{\zeta}} + \sum_{t=1}^{T-1}(1 - \sqrt{\beta_{2}})\mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t,i}+\zeta}}\right] \\\leq \frac{(\partial_{i}f(\boldsymbol{x}_{1}))^{2}}{\sqrt{\zeta}} + \sum_{t=1}^{T-1}(1 - \beta_{2})\mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}\right] \tag{31}$$

since $\boldsymbol{x}_0 = \boldsymbol{x}_1, \boldsymbol{g}_0 = 0$ and $\frac{1}{\beta_2} \leq 1 + \sqrt{\beta_2}$. By taking expectations and sums of (29) for t = 1 to T, based on (30) and (31), we have that

$$\sum_{t=1}^{T} \sum_{i=1}^{d} \left(\eta - \frac{\eta}{2\alpha_0} - \frac{\eta\alpha_0}{2\alpha_1} - \frac{L_0\eta^2 D_1}{2\sqrt{\zeta}} - \frac{\eta^2 L_1 \sqrt{dD_1}}{2\sqrt{1-\beta_2}} - \frac{\eta^2 \sqrt{d}}{2(1-\beta_2)\alpha_2} - \frac{\eta^2 \sqrt{d}\alpha_0 L_1 D_1}{\sqrt{1-\beta_2}} \right) \frac{(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}}$$

$$\leq f(\boldsymbol{x}_1) - \mathbb{E}[f(\boldsymbol{x}_{T+1})|\mathcal{F}_t] + \frac{\eta\alpha_0 D_1 ||\nabla f(\boldsymbol{x}_1)||^2}{2\sqrt{\zeta}} + \frac{\eta\alpha_0 dD_0}{2\sqrt{\zeta}} + T \frac{dL_0 \eta^2 D_0}{2\zeta} + T \frac{\alpha_0 \alpha_1 d^2 L_0^2 \eta^3 D_1^2}{2(1-\beta_2)} \frac{1}{\sqrt{\zeta}} + T \frac{\eta^2 \alpha_2 d^{1.5} L_1^2 D_0}{2\sqrt{\zeta}} + T \frac{\eta\alpha_0 dD_0 (1-\gamma\beta_2)}{2\sqrt{\zeta}} + T \frac{\eta\alpha_0 D_1 (1-\beta_2)}{2\sqrt{\zeta}} \sum_{i=1}^d \sum_{t=1}^T \mathbb{E} \left[\frac{||\partial_i f(\boldsymbol{x}_t)||^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}} \right]$$

$$(32)$$

Define $\Delta = f(\boldsymbol{x}_1) - f(\boldsymbol{x}^*) + \frac{\eta \alpha_0 dD_0}{2\sqrt{\zeta}} + \frac{\eta \alpha_0 D_1 \|\nabla f(\boldsymbol{x}_1)\|^2}{2\sqrt{\zeta}}$. If we set $\alpha_0 = 1, \alpha_1 = 7, \alpha_2 = 1$, obviously we can find some $1 - \beta_2 = \min(\frac{1}{7D_1}, \frac{\sqrt{\zeta}\epsilon^2}{35dD_0}) \sim \mathcal{O}(\epsilon^2)$, and $\eta = \min(\frac{\sqrt{\zeta}}{7L_0D_1}, \frac{\sqrt{1-\beta_2}}{\max(14L_1\sqrt{dD_1}, 7L_1\sqrt{dD_1})}, \frac{1-\beta_2}{7\sqrt{d}}, \frac{\zeta\epsilon^2}{35L_0dD_0}, \frac{\epsilon\sqrt{1-\beta_2}\sqrt{\zeta}}{7D_1L_0d\sqrt{5}}, \frac{\sqrt{\zeta}\epsilon^2}{35L_1^2d^{1.5}D_0}) \sim \mathcal{O}(\epsilon^2)$ such that

$$\frac{\eta}{14} \sum_{i=1}^{d} \sum_{t=1}^{T} \mathbb{E}\left[\frac{(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}}\right] \le \Delta + T \frac{\eta}{70} \epsilon^2 + T \frac{\eta}{70} \epsilon^2 + T \frac{\eta}{70} \epsilon^2 + T \frac{\eta}{70} \epsilon^2, \tag{33}$$

Set $T = \frac{70\Delta}{\eta\epsilon^2} \sim \mathcal{O}(\epsilon^{-4})$, and we have that

$$\frac{1}{T}\sum_{t=1}^{T} \mathbb{E}\left[\frac{\|\nabla f(\boldsymbol{x}_t)\|^2}{\sqrt{\beta_2}\|\boldsymbol{v}_{t-1}\|+\zeta}\right] \le \sum_{i=1}^{d} \frac{1}{T}\sum_{t=1}^{T} \mathbb{E}\left[\frac{(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2}\boldsymbol{v}_{t-1,i}+\zeta}\right] \le \epsilon^2.$$
(34)

This completes the proof.

C. Proof of Lemma 3.5

For any a, b > 0, we have that $\sqrt{a+b} \le \sqrt{a} + \sqrt{b}$. It then follows that

$$\mathbb{E}\left[\sqrt{\beta_2 \|\boldsymbol{v}_{t-1}\| + \zeta}\right] \le \sum_{i=1}^d \mathbb{E}\left[\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}}\right] + \sqrt{\zeta}.$$
(35)

For the first term, for t > 1 we have that

$$\mathbb{E}[\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}}] = \mathbb{E}_{\mathcal{F}_{t-1}}\left[\mathbb{E}\left[\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}}|\mathcal{F}_{t-1}\right]\right] = \mathbb{E}_{\mathcal{F}_{t-1}}\left[\mathbb{E}\left[\sqrt{\beta_2^2 \boldsymbol{v}_{t-2,i}} + \beta_2(1-\beta_2)(\boldsymbol{g}_{t-1,i})^2|\mathcal{F}_{t-1}\right]\right].$$
(36)

Give \mathcal{F}_{t-1} , the v_{t-2} is deterministic. By Jensen's inequality, we have that

$$\mathbb{E}_{\mathcal{F}_{t-1}} \left[\mathbb{E} \left[\sqrt{\beta_2^2 \boldsymbol{v}_{t-2,i} + \beta_2 (1 - \beta_2) (\boldsymbol{g}_{t-1,i})^2} | \mathcal{F}_{t-1} \right] \right] \\ \leq \mathbb{E}_{\mathcal{F}_{t-1}} \left[\sqrt{\mathbb{E} \left[\beta_2^2 \boldsymbol{v}_{t-2,i} + \beta_2 (1 - \beta_2) (\boldsymbol{g}_{t-1,i})^2 | \mathcal{F}_{t-1} \right]} \right] \\ \leq \mathbb{E}_{\mathcal{F}_{t-1}} \left[\sqrt{\beta_2^2 \boldsymbol{v}_{t-2,i} + \beta_2 (1 - \beta_2) (D_0 + D_1 (\partial_i f(\boldsymbol{x}_{t-1}))^2)} \right] \\ \leq \mathbb{E}_{\mathcal{F}_{t-1}} \left[\sqrt{\beta_2^2 \boldsymbol{v}_{t-2,i} + \beta_2 (1 - \beta_2) D_0} \right] + \mathbb{E} \left[\sqrt{\beta_2 (1 - \beta_2) D_1} | \partial_i f(\boldsymbol{x}_{t-1,i}) | \right], \tag{37}$$

where the second inequality is according to Assumption 2.2. By recursively applying (37) we have that

$$\mathbb{E}_{\mathcal{F}_{t-1}}\left[\sqrt{\beta_2^2 \boldsymbol{v}_{t-2} + \beta_2 (1-\beta_2) D_0}\right] \le \mathbb{E}_{\mathcal{F}_{t-2}}\left[\sqrt{\beta_2^3 \boldsymbol{v}_{t-3,i} + (\beta_2 + \beta_2^2)(1-\beta_2) D_0}\right] + \mathbb{E}\left[\sqrt{\beta_2^2 (1-\beta_2) D_1} |\partial_i f(\boldsymbol{x}_{t-2})|\right]$$
(38)

Specifically, we can get that

$$\mathbb{E}\left[\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}}\right] \leq \mathbb{E}_{\mathcal{F}_{t-1}}\left[\sqrt{\beta_{2}^{2}\boldsymbol{v}_{t-2,i} + \beta_{2}(1-\beta_{2})D_{0}}\right] + \mathbb{E}\left[\sqrt{\beta_{2}(1-\beta_{2})D_{1}}|\partial_{i}f(\boldsymbol{x}_{t-1})|\right] \\
\leq \mathbb{E}_{\mathcal{F}_{t-2}}\left[\sqrt{\beta_{2}^{3}\boldsymbol{v}_{t-3,i} + (\beta_{2}+\beta_{2}^{2})(1-\beta_{2})D_{0}}\right] + \mathbb{E}\left[\sqrt{\beta_{2}^{2}(1-\beta_{2})D_{1}}|\partial_{i}f(\boldsymbol{x}_{t-i})|\right] \\
+ \mathbb{E}\left[\sqrt{\beta_{2}(1-\beta_{2})D_{1}}|\partial_{i}f(\boldsymbol{x}_{t-1})|\right] \\\leq \dots \\
\leq \mathbb{E}\left[\sqrt{(\beta_{2}+\beta_{2}^{2}+\beta_{2}^{3} + \dots + \beta_{2}^{t})(1-\beta_{2})D_{0} + \boldsymbol{v}_{0,i}}\right] + \sum_{i=1}^{t-1}\mathbb{E}\left[\sqrt{\beta_{2}^{j}(1-\beta_{2})D_{1}}|\partial_{i}f(\boldsymbol{x}_{t-j})|\right] \\
\leq \sqrt{D_{0}+\|\boldsymbol{v}_{0}\|} + \sum_{j=1}^{t-1}\mathbb{E}\left[\sqrt{\beta_{2}^{j}(1-\beta_{2})D_{1}}|\partial_{i}f(\boldsymbol{x}_{t-j})|\right].$$
(39)

As a result, we have that

$$\frac{1}{T} \sum_{t=1}^{T} \mathbb{E} \left[\sqrt{\beta_2 \| \boldsymbol{v}_{t-1} \| + \zeta} \right]
\leq \sqrt{\zeta} + d\sqrt{D_0 + \| \boldsymbol{v}_0 \|} + \frac{1}{T} \sum_{i=1}^{d} \sum_{t=1}^{T} \sum_{j=1}^{t-1} \mathbb{E} \left[\sqrt{\beta_2^j (1 - \beta_2) D_1} | \partial_i f(\boldsymbol{x}_{t-j}) | \right]
\leq c + \sum_{i=1}^{d} \frac{\sum_{t=1}^{T} \mathbb{E} [|\partial_i f(\boldsymbol{x}_t)|]}{T} \sqrt{(1 - \beta_2) D_1} \left(\sum_{j=1}^{T} \sqrt{\beta_2^j} \right)
\leq c + \frac{2\sqrt{D_1}}{\sqrt{(1 - \beta_2)}} \sum_{i=1}^{d} \frac{\sum_{t=1}^{T} \mathbb{E} [|\partial_i f(\boldsymbol{x}_t)|]}{T}
\leq c + \frac{2\sqrt{dD_1}}{\sqrt{(1 - \beta_2)}} \frac{\sum_{i=1}^{T} \mathbb{E} [\| \nabla f(\boldsymbol{x}_t) \|]}{T},$$
(40)

where the last inequlity is due to $\sum_{i=1}^{d} \|\partial_i f(\boldsymbol{x})\| \leq \sqrt{d} \|f(\boldsymbol{x})\|$. We then complete the proof.

D. Formal Version of Theorem 3.1 and Its Proof

Define $c = \sqrt{\zeta} + d\sqrt{D_0 + \|\boldsymbol{v}_0\|}, \Delta = f(\boldsymbol{x}_1) - f(\boldsymbol{x}^*) + \frac{\eta \alpha_0 dD_0}{2\sqrt{\zeta}} + \frac{\eta \alpha_0 D_1 \|\nabla f(\boldsymbol{x}_1)\|^2}{2\sqrt{\zeta}}, \Lambda_1 = \frac{1}{\max(14L_1\sqrt{d}D_1, 7L_1\sqrt{d}D_1)}, \Lambda_2 = \min\left(\frac{\zeta}{35L_0dD_0}, \frac{\sqrt{\zeta}}{35L_1^2d^{1.5}D_0}\right) \text{ and } \Lambda_3 = \frac{4\sqrt{\zeta}}{7D_1L_0d\sqrt{5}}.$

Theorem D.1. Let Assumptions 2.1, 2.2 and 2.3 hold. Let $1 - \beta_2 = \min\left(\frac{1}{7D_1}, \frac{\sqrt{\zeta}\epsilon^2}{35dD_0}\right) \sim \mathcal{O}(\epsilon^2)$, $\eta = \min\left(\frac{\sqrt{\zeta}}{7L_0D_1}, \Lambda_1\sqrt{1-\beta_2}, \frac{1-\beta_2}{7\sqrt{d}}, \Lambda_2\epsilon^2, \Lambda_3\epsilon\sqrt{1-\beta_2}\right) \sim \mathcal{O}(\epsilon^2)$, and $T = \frac{70\Delta}{\eta\epsilon^2} \sim \mathcal{O}(\epsilon^{-4})$. For small ϵ such that $\epsilon \leq \frac{\sqrt{5dD_0}}{\sqrt{D_1}\sqrt[4]{\zeta}}$, we have that

$$\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}[\|\nabla f(\boldsymbol{x}_t)\|] \le \left(\frac{2d\sqrt{35D_0D_1}}{\sqrt[4]{\zeta}} + \sqrt{c}\right)\epsilon.$$
(41)

Proof. According to Lemma 3.4, we have that

$$\frac{1}{T}\sum_{t=1}^{T} \mathbb{E}\left[\frac{\|\nabla f(\boldsymbol{x}_t)\|^2}{\sqrt{\beta_2 \|\boldsymbol{v}_{t-1}\| + \zeta}}\right] \le \epsilon^2.$$
(42)

With bounded gradient norms assumption, it is easy to show that $\frac{1}{T}\sum_{t=1}^{T} \mathbb{E}[\|\nabla f(\boldsymbol{x}_t)\|^2] \sim \mathcal{O}(\epsilon^2)$ via Corollary 3.4. However, in this paper we focus on the bounded-free-assumptions thus the $\sqrt{\beta_2 \|\boldsymbol{v}_{t-1}\| + \zeta}$ is potentially infinite, making it hard to show the algorithm converges to a stationary point directly.

According to Lemma 3.5, we have that

$$\left(\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}[\sqrt{\beta_2 \|\boldsymbol{v}_{t-1}\| + \zeta}]\right) \leq c + \frac{2\sqrt{dD_1}}{\sqrt{(1-\beta_2)}} \frac{\sum_{t=1}^{T}\mathbb{E}[\|\nabla f(\boldsymbol{x}_t)\|]}{T}.$$
(43)

Define $e = \frac{1}{T} \sum_{t=1}^{T} \mathbb{E}[\|\nabla f(\boldsymbol{x}_t)\|]$. By Hölder's inequality, we have that

$$\left(\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}[\|\nabla f(\boldsymbol{x}_{t})\|]\right)^{2} \leq \left(\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}\left[\frac{\|\nabla f(\boldsymbol{x}_{t})\|^{2}}{\sqrt{\beta_{2}\|\boldsymbol{v}_{t-1}\|+\zeta}}\right]\right)\left(\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}[\sqrt{\beta_{2}\|\boldsymbol{v}_{t-1}\|+\zeta}]\right),\tag{44}$$

based on Lemma 3.5 and Lemma 3.4 which can be written as

$$e^2 \le \epsilon^2 (c + \frac{2\sqrt{dD_1}}{\sqrt{1-\beta_2}}e).$$
 (45)

Thus we have that

$$\frac{1}{T} \sum_{t=1}^{T} \mathbb{E}[\|\nabla f(x_t)\|] = e \le \frac{\sqrt{dD_1}\epsilon^2}{\sqrt{1-\beta_2}} + \frac{1}{2}\sqrt{\left(\frac{2\sqrt{dD_1}\epsilon^2}{\sqrt{1-\beta_2}}\right)^2 + 4c\epsilon^2}.$$

Since $1 - \beta_2 = \min(\frac{1}{7D_1}, \frac{\sqrt{\zeta}\epsilon^2}{35dD_0}) \sim \mathcal{O}(\epsilon^2)$, if $\epsilon \leq \frac{\sqrt{5dD_0}}{\sqrt{D_1}\sqrt[4]{\zeta}}$ we have $\frac{\epsilon^2}{\sqrt{1-\beta_2}} \leq \frac{\sqrt{35dD_0}}{\sqrt[4]{\zeta}}\epsilon$. It demonstrates that

$$\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}[\|\nabla f(x_t)\|] \le \left(\frac{2d\sqrt{35D_0D_1}}{\sqrt[4]{\zeta}} + \sqrt{c}\right)\epsilon,\tag{46}$$

which completes the proof.

E. Lemmas for Theorem 4.1

Here are some lemmas we need for the proof of Theorem 4.1.

Lemma E.1. (Lemma 6 in (Wang et al., 2023a)) For any $\{c_t\}_{t=0}^{\infty} \ge 0$ and $a_t = \beta_2 a_{t-1} + (1 - \beta_2)c_t^2$ and $b_t = \beta_1 b_{t-1} + (1 - \beta_1)c_t$, if $0 < \beta_1^2 < \beta_2 < 1$, we have that

$$\frac{b_t}{\sqrt{a_t + \zeta}} \le \frac{1 - \beta_1}{\sqrt{1 - \beta_2}\sqrt{1 - \frac{\beta_1^2}{\beta_2}}} \tag{47}$$

Proof. we have that

$$\frac{b_t}{\sqrt{a_t + \zeta}} \leq \frac{b_t}{\sqrt{a_t}} \leq \frac{\sum_{i=0}^{t-1} (1 - \beta_1) \beta_1^i c_{t-i}}{\sqrt{\sum_{i=0}^{t-1} (1 - \beta_2) \beta_2^i c_{t-i}^2 + a_0}} \\
\leq \frac{1 - \beta_1}{\sqrt{1 - \beta_2}} \frac{\sqrt{\sum_{i=0}^{t-1} \beta_2^i c_{t-i}^2} \sqrt{\sum_{i=0}^{t-1} \frac{\beta_1^{2i}}{\beta_2}}}{\sqrt{\sum_{i=0}^{t-1} \beta_2^i c_{t-i}^2}} \leq \frac{1 - \beta_1}{\sqrt{1 - \beta_2} \sqrt{1 - \frac{\beta_1^2}{\beta_2}}},$$
(48)

where the third inequality is due to Cauchy's inequality.

Lemma E.2. (Lemma 5 in (Wang et al., 2023a)) For any $\{c_t\}_{t=0}^{\infty} \ge 0$ and $a_t = \beta_2 a_{t-1} + (1 - \beta_2)c_t^2$ and $b_t = \beta_1 b_{t-1} + (1 - \beta_1)c_t$, if $0 < \beta_1^2 < \beta_2 < 1$ and $a_0 > 0$, we have that

$$\sum_{t=1}^{T} \frac{b_t^2}{a_t} \le \frac{(1-\beta_1)^2}{(1-\frac{\beta_1}{\sqrt{\beta_2}})^2(1-\beta_2)} \left(\ln\left(\frac{a_T}{a_0}\right) - T\ln(\beta_2) \right)$$
(49)

Proof. Due to the monotonicity of $\frac{1}{a}$ function and Lemma 5.2 in (Défossez et al., 2020), we have that

$$\frac{(1-\beta_2)c_t^2}{a_t} \le \int_{a=a_t-(1-\beta_2)c_t^2}^{a_t} \frac{1}{a} da = \ln\left(\frac{a_t}{a_t-(1-\beta_2)c_t^2}\right) = \ln\left(\frac{a_t}{a_{t-1}}\right) - \ln(\beta_2).$$
(50)

By telescoping, we have the $\sum_{t=1}^{T} \frac{c_t^2}{a_t} \leq \frac{1}{1-\beta_2} \ln\left(\frac{a_T}{a_0}\right) - T \frac{\ln(\beta_2)}{(1-\beta_2)}$. Moreover, for b_t we have that

$$\frac{b_t}{\sqrt{a_t}} \le (1 - \beta_1) \sum_{i=1}^t \frac{\beta_1^{t-i} c_i}{\sqrt{a_t}} \le (1 - \beta_1) \sum_{i=1}^t \left(\frac{\beta_1}{\sqrt{\beta_2}}\right)^{t-i} \frac{c_i}{\sqrt{a_i}}.$$
(51)

Moreover, due to the Cauchy's inequality, we get that

$$\frac{b_t^2}{a_t} \leq (1-\beta_1)^2 \left(\sum_{i=1}^t \left(\frac{\beta_1}{\sqrt{\beta_2}} \right)^{t-i} \frac{c_i}{\sqrt{a_i}} \right)^2 \\
\leq (1-\beta_1)^2 \left(\sum_{i=1}^t \left(\frac{\beta_1}{\sqrt{\beta_2}} \right)^{t-i} \right) \left(\sum_{i=1}^t \left(\frac{\beta_1}{\sqrt{\beta_2}} \right)^{t-i} \frac{c_i^2}{a_i} \right) \\
\leq \frac{(1-\beta_1)^2}{\left(1-\frac{\beta_1}{\sqrt{\beta_2}}\right)} \left(\sum_{i=1}^t \left(\frac{\beta_1}{\sqrt{\beta_2}} \right)^{t-i} \frac{c_i^2}{a_i} \right)$$
(52)

Thus we have that

$$\sum_{t=1}^{T} \frac{b_t^2}{a_t} \le \frac{(1-\beta_1)^2}{(1-\frac{\beta_1}{\sqrt{\beta_2}})^2(1-\beta_2)} \left(\ln\left(\frac{a_T}{a_0}\right) - T\ln(\beta_2) \right).$$
(53)

Lemma E.3. For any $\{c_t\}_{t=0}^{\infty} \ge 0$ and $a_t = \beta_2 a_{t-1} + (1 - \beta_2)c_t^2$ and $b_t = \beta_1 b_{t-1} + (1 - \beta_1)c_t$, if $0 < \beta_1^4 < \beta_2 < 1$ and $a_0 > 0$, we have that

$$\sum_{t=1}^{T} \frac{b_t^2}{\sqrt{a_t}} \le \frac{(1-\beta_1)^2}{(1-\frac{\beta_1}{\sqrt{\beta_2}})^2} \left(\frac{2}{1-\beta_2}(\sqrt{a_T}-\sqrt{a_0}) + \sum_{t=1}^{T} 2\sqrt{a_{t-1}}\right)$$
(54)

Proof. Due to the monotonicity of $\frac{1}{\sqrt{a}}$ function, we have that

$$\frac{(1-\beta_2)c_t^2}{\sqrt{a_t}} \le \int_{a=a_t-(1-\beta_2)c_t^2}^{a_t} \frac{1}{\sqrt{a}} da = 2\sqrt{a_t} - 2\sqrt{a_t - (1-\beta_2)c_t^2} = 2\sqrt{a_t} - 2\sqrt{\beta_2 a_{t-1}} \le 2\sqrt{a_t} - 2\sqrt{a_{t-1}} + 2(1-\beta_2)\sqrt{a_{t-1}}$$
(55)

By taking sum up from t = 1 to T, we have the

$$\sum_{t=1}^{T} \frac{(1-\beta_2)c_t^2}{\sqrt{a_t}} \le 2\sqrt{a_T} - 2\sqrt{a_0} + 2(1-\beta_2)\sum_{t=1}^{T} \sqrt{a_{t-1}}.$$
(56)

Moreover, for b_t we have that

$$\frac{b_t}{\sqrt[4]{a_t}} \le (1 - \beta_1) \sum_{i=1}^t \frac{\beta_1^{t-i} c_i}{\sqrt[4]{a_t}} \le (1 - \beta_1) \sum_{i=1}^t \left(\frac{\beta_1}{\sqrt[4]{\beta_2}}\right)^{t-i} \frac{c_i}{\sqrt[4]{a_i}}$$
(57)

Thus we have that

$$\frac{b_t^2}{\sqrt{a_t}} \leq (1-\beta_1)^2 \left(\sum_{i=1}^t \left(\frac{\beta_1}{\sqrt[4]{\beta_2}} \right)^{t-i} \frac{c_i}{\sqrt[4]{a_i}} \right)^2 \\
\leq (1-\beta_1)^2 \left(\sum_{i=1}^t \left(\frac{\beta_1}{\sqrt[4]{\beta_2}} \right)^{t-i} \right) \left(\sum_{i=1}^t \left(\frac{\beta_1}{\sqrt[4]{\beta_2}} \right)^{t-i} \frac{c_i^2}{\sqrt{a_i}} \right) \\
\leq \frac{(1-\beta_1)^2}{1-\frac{\beta_1}{\sqrt[4]{\beta_2}}} \left(\sum_{i=1}^t \left(\frac{\beta_1}{\sqrt[4]{\beta_2}} \right)^{t-i} \frac{c_i^2}{\sqrt{a_i}} \right)$$
(58)

Thus we have that

$$\sum_{t=1}^{T} \frac{b_t^2}{\sqrt{a_t}} \le \frac{(1-\beta_1)^2}{(1-\frac{\beta_1}{\sqrt[4]{\beta_2}})^2} \sum_{t=1}^{T} \frac{c_t^2}{\sqrt{a_t}}.$$
(59)

Plug (56) in (59), and we can complete the proof.

F. Lemma F.1 and Its Proof

Define $C_1 = 1 - \frac{\beta_1}{\sqrt{\beta_2}}, C_2 = \sqrt{1 - \frac{\beta_1^2}{\beta_2}}$, we have the following Lemma: Lemma F.1. For any $\alpha_0, \alpha_1, \alpha_3, \alpha_4 > 0$, $\beta_2 > 0.5$, and $0 < \beta_1^2 < \beta_2$, we have that

$$\mathbb{E}[\langle \nabla f(\boldsymbol{u}_{t}), \boldsymbol{u}_{t} - \boldsymbol{u}_{t+1} \rangle | \mathcal{F}_{t}] \\
\geq \left[\frac{\eta(1-\beta_{1})}{C_{1}} - \frac{\eta(1-\beta_{1})}{C_{1}C_{2}} \left(\frac{1}{2\alpha_{0}} + \frac{\alpha_{0}}{2\alpha_{1}} + \frac{\eta\alpha_{0}\sqrt{d}D_{1}L_{1}(1-\beta_{1})}{\sqrt{1-\beta_{2}}C_{2}} \right) - \frac{\eta\beta_{1}(1-\beta_{1})\sqrt{\zeta}}{2\alpha_{3}C_{1}C_{2}} - \frac{\eta L_{1}(1-C_{1})^{2}}{2\alpha_{4}C_{1}^{2}} - \frac{\eta L_{1}(1-C_{1})}{2\alpha_{4}C_{1}^{2}} \right] \\
\times \sum_{i=1}^{d} \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} \\
- \sum_{i=1}^{d} \frac{\eta(1-\beta_{1})}{C_{1}C_{2}} \left[\frac{\alpha_{0}D_{0}}{2} \left(\frac{1}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} - \mathbb{E}\left[\frac{1}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} | \mathcal{F}_{t} \right] \right] + \frac{\alpha_{0}D_{1}}{2} \mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} - \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} | \mathcal{F}_{t} \right] \right] \\
- \frac{\alpha_{0}\alpha_{1}D_{1}^{2}L_{0}^{2}\eta^{3}d^{2}(1-\beta_{1})^{3}}{2(1-\beta_{2})C_{1}C_{2}^{3}\sqrt{\zeta}} - \frac{\alpha_{3}\eta d\beta_{1}(1-\beta_{1})(1-\beta_{2})}{2C_{1}C_{2}} \\
- \sum_{i=1}^{d} \frac{\eta^{2}((1-C_{1})^{2}+0.5(1-C_{1}))\sqrt{d}L_{0}}{C_{1}^{2}} \frac{m_{t-1,i}^{2}}{\boldsymbol{v}_{t-1,i} + \zeta} - \sum_{i=1}^{d} \frac{\eta^{2}0.5(1-C_{1})\sqrt{d}L_{0}}{C_{1}^{2}} \mathbb{E}\left[\frac{m_{t,i}^{2}}{\boldsymbol{v}_{t,i} + \zeta} | \mathcal{F}_{t} \right] \\
- \sum_{i=1}^{d} \frac{\alpha_{4}\eta^{3}(1-\beta_{1})^{2}(1-C_{1})^{2}dL_{1}}{2(1-\beta_{2})C_{1}^{2}C_{2}^{2}} \frac{m_{t-1,i}^{2}}{\sqrt{\boldsymbol{v}_{t-1,i} + \zeta}} - \sum_{i=1}^{d} \frac{\alpha_{4}\eta^{3}(1-\beta_{1})^{2}(1-C_{1})dL_{1}}{2(1-\beta_{2})C_{1}^{2}C_{2}^{2}} \mathbb{E}\left[\frac{m_{t,i}^{2}}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} | \mathcal{F}_{t} \right].$$
(60)

Proof. Since $u_t = rac{x_t - rac{eta_1}{\sqrt{eta_2}} x_{t-1}}{1 - rac{eta_1}{\sqrt{eta_2}}}$, we then have that

$$u_{t+1} - u_t = \frac{x_{t+1} - x_t}{1 - \frac{\beta_1}{\sqrt{\beta_2}}} - \frac{\beta_1}{\sqrt{\beta_2}} \frac{x_t - x_{t-1}}{1 - \frac{\beta_1}{\sqrt{\beta_2}}} \\ = \frac{-\eta \odot \frac{m_t}{\sqrt{v_t + \zeta}}}{1 - \frac{\beta_1}{\sqrt{\beta_2}}} + \frac{\beta_1}{\sqrt{\beta_2}} \frac{\eta \odot \frac{m_{t-1}}{\sqrt{v_{t-1} + \zeta}}}{1 - \frac{\beta_1}{\sqrt{\beta_2}}}$$

$$= \frac{-\eta \odot \frac{m_{t}}{\sqrt{\beta_{2} v_{t-1} + \zeta}}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} + \frac{-\eta \odot \frac{m_{t}}{\sqrt{v_{t} + \zeta}} + \eta \odot \frac{m_{t}}{\sqrt{\beta_{2} v_{t-1} + \zeta}}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \\ + \frac{\eta \odot \frac{\beta_{1} m_{t-1}}{\sqrt{\beta_{2} v_{t-1} + \zeta}}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} + \frac{\eta \odot \frac{\beta_{1} m_{t-1}}{\sqrt{\beta_{2} v_{t-1} + \beta_{2}\zeta}} - \eta \odot \frac{\beta_{1} m_{t-1}}{\sqrt{\beta_{2} v_{t-1} + \zeta}}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \\ = \frac{-\eta \odot \frac{(1 - \beta_{1})g_{t}}{\sqrt{\beta_{2} v_{t-1} + \zeta}}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} + \frac{-\eta \odot \frac{m_{t}}{\sqrt{v_{t} + \zeta}} + \eta \odot \frac{m_{t}}{\sqrt{\beta_{2} v_{t-1} + \zeta}}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} + \frac{\eta \odot \frac{\beta_{1} m_{t-1}}{\sqrt{\beta_{2} v_{t-1} + \zeta}}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}}, \quad (61)$$

where the last equality is due to $m_t = \beta_1 m_{t-1} + (1 - \beta_1) g_t$. For (L_0, L_1) -smooth objectives, from Lemma 1 in (Crawshaw et al., 2022) we can get that

$$\underbrace{\mathbb{E}[\langle \nabla f(\boldsymbol{u}_{t}), \boldsymbol{u}_{t} - \boldsymbol{u}_{t+1} \rangle | \mathcal{F}_{t}]}_{\text{First Order}} \leq f(\boldsymbol{u}_{t}) - \mathbb{E}[f(\boldsymbol{u}_{t+1}) | \mathcal{F}_{t}] + \underbrace{\sum_{i=1}^{d} \frac{L_{0}}{2\sqrt{d}} \mathbb{E}[\|\boldsymbol{u}_{t+1} - \boldsymbol{u}_{t}\| | \boldsymbol{u}_{t+1,i} - \boldsymbol{u}_{t,i} | | \mathcal{F}_{t}]}_{\text{Second Order}} + \underbrace{\sum_{i=1}^{d} \frac{L_{1} |\partial_{i} f(\boldsymbol{u}_{t})|}{2} \mathbb{E}[\|\boldsymbol{u}_{t+1} - \boldsymbol{u}_{t}\| | \boldsymbol{u}_{t+1,i} - \boldsymbol{u}_{t,i} | | \mathcal{F}_{t}]}_{\text{Additional Term}}.$$
(62)

We then focus on the first-order term. Based on (61) we divide our first order terms into four parts

$$\mathbb{E}[\langle \nabla f(\boldsymbol{u}_{t}), \boldsymbol{u}_{t} - \boldsymbol{u}_{t+1} \rangle | \mathcal{F}_{t}] = \mathbb{E}[\langle \nabla f(\boldsymbol{x}_{t}), \boldsymbol{u}_{t} - \boldsymbol{u}_{t+1} \rangle | \mathcal{F}_{t}] + \mathbb{E}[\langle \nabla f(\boldsymbol{u}_{t}) - \nabla f(\boldsymbol{x}_{t}), \boldsymbol{u}_{t} - \boldsymbol{u}_{t+1} \rangle | \mathcal{F}_{t}]$$

$$= \mathbb{E}\begin{bmatrix} \left\langle \nabla f(\boldsymbol{x}_{t}), \frac{\eta \odot \frac{(1-\beta_{1})\boldsymbol{g}_{t}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1} + \zeta}} \right\rangle | \mathcal{F}_{t} \\ 1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}} \right\rangle | \mathcal{F}_{t} \end{bmatrix}$$
First Order Main
$$- \mathbb{E}\begin{bmatrix} \left\langle \nabla f(\boldsymbol{x}_{t}), \frac{\eta \odot \frac{\beta_{1}\boldsymbol{m}_{t-1}}{\sqrt{\beta_{2}}\boldsymbol{v}_{t-1} + \beta_{2}\zeta} - \eta \odot \frac{\beta_{1}\boldsymbol{m}_{t-1}}{\sqrt{\beta_{2}}\boldsymbol{v}_{t-1} + \zeta} \right\rangle | \mathcal{F}_{t} \end{bmatrix}$$
Error 1
$$- \mathbb{E}\begin{bmatrix} \left\langle \nabla f(\boldsymbol{u}_{t}), \frac{\eta \odot \frac{\beta_{1}\boldsymbol{m}_{t-1}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1} + \beta_{2}\zeta}} - \eta \odot \frac{\beta_{1}\boldsymbol{m}_{t-1}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1} + \zeta}} \right\rangle | \mathcal{F}_{t} \end{bmatrix}$$

$$- \mathbb{E}\begin{bmatrix} \left\langle \nabla f(\boldsymbol{u}_{t}), \frac{\eta \odot \frac{\beta_{1}\boldsymbol{m}_{t-1}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1} + \beta_{2}\zeta}} - \eta \odot \frac{\beta_{1}\boldsymbol{m}_{t-1}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1} + \zeta}} \right\rangle | \mathcal{F}_{t} \end{bmatrix}$$

$$- \mathbb{E}\begin{bmatrix} \left\langle \nabla f(\boldsymbol{u}_{t}), \frac{\eta \odot \frac{\beta_{1}\boldsymbol{m}_{t-1}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1} + \beta_{2}\zeta}} - \eta \odot \frac{\beta_{1}\boldsymbol{m}_{t-1}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1} + \zeta}} \right\rangle | \mathcal{F}_{t} \end{bmatrix}$$

$$- \mathbb{E}\begin{bmatrix} \left\langle \nabla f(\boldsymbol{u}_{t}), \frac{\eta \odot \frac{\beta_{1}\boldsymbol{m}_{t-1}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1} + \beta_{2}\zeta}} - \eta \odot \frac{\beta_{1}\boldsymbol{m}_{t-1}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1} + \zeta}} \right\rangle | \mathcal{F}_{t} \end{bmatrix}$$

$$- \mathbb{E}\begin{bmatrix} \nabla f(\boldsymbol{u}_{t}) - \nabla f(\boldsymbol{x}_{t}), \boldsymbol{u}_{t+1} - \boldsymbol{u}_{t} \rangle | \mathcal{F}_{t} \end{bmatrix}.$$

$$(63)$$

The first-order main is easy to bound. Since give $\mathcal{F}_t, \boldsymbol{g}_t$ is independent of \boldsymbol{v}_{t-1} and we have that

$$\underbrace{\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \frac{\eta \odot \frac{(1-\beta_{1})\boldsymbol{g}_{t}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1}+\zeta}}\right\rangle | \mathcal{F}_{t}\right]}_{\text{First Order Main}} = \sum_{i=1}^{d} \frac{\eta(1-\beta_{1})}{1-\frac{\beta_{1}}{\sqrt{\beta_{2}}}} \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}.$$
(64)

We then focus on the Error 1 term. Since $v_t = \beta_2 v_{t-1} + (1 - \beta_2) g_t \odot g_t$, we have (15) and it follows that

$$\underbrace{\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \frac{-\eta \odot \frac{\boldsymbol{m}_{t}}{\sqrt{\boldsymbol{v}_{t}+\zeta}} + \eta \odot \frac{\boldsymbol{m}_{t}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1}+\zeta}}\right\rangle |\mathcal{F}_{t}\right]}_{\text{Error 1}}_{\text{Error 1}} \\
\leq \underbrace{\sum_{i=1}^{d} \frac{\eta}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \mathbb{E}\left[\left|\partial_{i}f(\boldsymbol{x}_{t})\right| \frac{(1 - \beta_{2})\boldsymbol{g}_{t,i}^{2}\boldsymbol{m}_{t,i}}{(\sqrt{\boldsymbol{v}_{t,i}+\zeta})(\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta})(\sqrt{\boldsymbol{v}_{t,i}+\zeta} + \sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta})} |\mathcal{F}_{t}\right]. \tag{65}$$

According to Lemma E.1, we have $\frac{|\boldsymbol{m}_{t,i}|}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}} \leq \frac{1-\beta_1}{\sqrt{1-\beta_2}\sqrt{1-\frac{\beta_1^2}{\beta_2}}}$, which demonstrates that

$$\operatorname{Error} 1 \leq \sum_{i=1}^{d} \frac{\eta}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \frac{1 - \beta_{1}}{\sqrt{1 - \frac{\beta_{1}^{2}}{\beta_{2}}}} \frac{|\partial_{i} f(\boldsymbol{x}_{t})|}{\sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta}} \mathbb{E} \left[\left\| \frac{\sqrt{1 - \beta_{2}} \boldsymbol{g}_{t,i}^{2}}{(\sqrt{\boldsymbol{v}_{t,i} + \zeta} + \sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta})} \right\| |\mathcal{F}_{t} \right]$$
(66)

Similar to Appendix A, we have that for any $\alpha_0 > 0$ and $i \in [d]$

$$\frac{|\partial_{i}f(\boldsymbol{x}_{t})|}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}\mathbb{E}\left[\left\|\frac{\sqrt{1-\beta_{2}}\boldsymbol{g}_{t,i}^{2}}{(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i}+\zeta)}\right\||\mathcal{F}_{t}\right] \\
\leq \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{0}\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i}+\zeta} + \frac{\alpha_{0}}{2\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i}+\zeta}\left(\mathbb{E}\left[\frac{\sqrt{1-\beta_{2}}\boldsymbol{g}_{t,i}^{2}}{(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i}+\zeta)}|\mathcal{F}_{t}\right]\right)^{2}.$$
(67)

For the last term, due to Hölder's inequality, we have that

$$\frac{\alpha_{0}}{2\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} \left(\mathbb{E}\left[\frac{\sqrt{1-\beta_{2}}\boldsymbol{g}_{t,i}^{2}}{(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i}+\zeta)}|\mathcal{F}_{t}\right] \right)^{2} \\
\leq \frac{(1-\beta_{2})\alpha_{0}}{2\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i}+\zeta} \mathbb{E}[\boldsymbol{g}_{t,i}^{2}|\mathcal{F}_{t}] \mathbb{E}\left[\frac{\boldsymbol{g}_{t,i}^{2}}{(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i}+\zeta)^{2}}|\mathcal{F}_{t}\right] \\
\leq \frac{\alpha_{0}}{2}(D_{0}+D_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2}) \times \mathbb{E}\left[\frac{(1-\beta_{2})\boldsymbol{g}_{t,i}^{2}}{(\sqrt{\boldsymbol{v}_{t,i}+\zeta}\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i}+\zeta}(\sqrt{\boldsymbol{v}_{t,i}+\zeta}+\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i}+\zeta)}|\mathcal{F}_{t}\right] \\
\leq \frac{\alpha_{0}}{2}(D_{0}+D_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2}) \mathbb{E}\left[\frac{1}{\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i}+\zeta}-\frac{1}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_{t}\right] \tag{68}}$$

Similar to (21), we will show the sum of (68) from t = 1 to T can be bounded. For t > 1 we have that

$$\frac{\alpha_{0}}{2}(D_{0} + D_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2})\mathbb{E}\left[\frac{1}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_{t}\right] \\
= \frac{\alpha_{0}D_{0}}{2}\mathbb{E}\left[\frac{1}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_{t}\right] + \frac{\alpha_{0}D_{1}}{2}\mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}|\mathcal{F}_{t}\right] \\
+ \frac{\alpha_{0}D_{1}}{2}\mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2} - (\partial_{i}f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}|\mathcal{F}_{t}\right].$$
(69)

Note for the last term in RHS of (69), due to the different update of x_t , there is a little difference compared with (22). For the last term and any $\alpha_1 > 0$ and $i \in [d]$, we have that

$$\frac{\alpha_0 D_1}{2} \mathbb{E} \left[\frac{(\partial_i f(\boldsymbol{x}_t))^2 - (\partial_i f(\boldsymbol{x}_{t-1}))^2}{\sqrt{\beta \boldsymbol{v}_{t-1,i} + \zeta}} |\mathcal{F}_t \right] \\
\leq \frac{\alpha_0 D_1}{2} \mathbb{E} \left[\frac{2|\partial_i f(\boldsymbol{x}_t)| ||\partial_i f(\boldsymbol{x}_t)| - |\partial_i f(\boldsymbol{x}_{t-1})||}{\sqrt{\beta \boldsymbol{v}_{t-1,i} + \zeta}} |\mathcal{F}_t \right] \\
\leq \alpha_0 D_1 \frac{|\partial_i f(\boldsymbol{x}_t)| (L_0 + L_1 |\partial_i f(\boldsymbol{x}_t)|) \eta \left\| \frac{1}{\sqrt{\boldsymbol{v}_{t-1} + \zeta}} \odot \boldsymbol{m}_{t-1} \right\|}{\sqrt{\beta \boldsymbol{v}_{t-1,i} + \zeta}} \\
\leq \frac{\alpha_0 D_1}{2\sqrt{\beta \boldsymbol{v}_{t-1,i} + \zeta}} \left(\frac{(\partial_i f(\boldsymbol{x}_t))^2}{\alpha_1 D_1} + \alpha_1 D_1 L_0^2 \eta^2 \left\| \frac{1}{\boldsymbol{v}_{t-1} + \zeta} \odot \boldsymbol{m}_{t-1} \right\|^2 + 2\eta L_1 (\partial_i f(\boldsymbol{x}_t))^2 \left\| \frac{1}{\boldsymbol{v}_{t-1} + \zeta} \odot \boldsymbol{m}_{t-1} \right\| \right)$$

Convergence Guarantees for RMSProp and Adam

$$\leq \frac{\alpha_0 D_1}{2\sqrt{\beta \boldsymbol{v}_{t-1,i} + \zeta}} \left(\frac{(\partial_i f(\boldsymbol{x}_t))^2}{\alpha_1 D_1} + \alpha_1 D_1 L_0^2 \eta^2 \frac{d(1 - \beta_1)^2}{(1 - \beta_2)(1 - \frac{\beta_1^2}{\beta_2})} + 2\eta L_1 (\partial_i f(\boldsymbol{x}_t))^2 \frac{\sqrt{d}(1 - \beta_1)}{\sqrt{1 - \beta_2}\sqrt{1 - \frac{\beta_1^2}{\beta_2}}} \right), \tag{70}$$

where the first inequality is due to the fact that for any $a, b \ge 0$, we have $a^2 - b^2 \le 2a|a - b|$, the second inequality is by the (L_0, L_1) -smoothness and update process of x_t and the last inequality is due to Lemma E.1. As a result, for $\alpha_0, \alpha_1 > 0$, if t = 1 we have that

$$\operatorname{Error} 1 \leq \sum_{i=1}^{d} \frac{\eta}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \frac{1 - \beta_{1}}{\sqrt{1 - \frac{\beta_{1}^{2}}{\beta_{2}}}} \left(\frac{(\partial_{i} f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{0}\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i} + \zeta} + \frac{\alpha_{0}D_{0}}{2} \mathbb{E} \left[\frac{1}{\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i} + \zeta} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} |\mathcal{F}_{t}\right] \right]$$

$$+ \frac{\alpha_{0}D_{1}}{2} \mathbb{E} \left[\frac{(\partial_{i} f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i} + \zeta} - \frac{(\partial_{i} f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} |\mathcal{F}_{t}\right] \right),$$

$$(71)$$

and if t > 1 we have that

$$\operatorname{Error} 1 \leq \sum_{i=1}^{d} \frac{\eta}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \frac{1 - \beta_{1}}{\sqrt{1 - \frac{\beta_{1}^{2}}{\beta_{2}}}} \left[\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{0}\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i} + \zeta} + \frac{\alpha_{0}D_{0}}{2} \mathbb{E} \left[\frac{1}{\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i} + \zeta} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} |\mathcal{F}_{t} \right] \right. \\ \left. + \frac{\alpha_{0}D_{1}}{2} \mathbb{E} \left[\frac{(\partial_{i}f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i} + \zeta} - \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} |\mathcal{F}_{t} \right] \right. \\ \left. + \frac{\alpha_{0}D_{1}}{2\sqrt{\beta\boldsymbol{v}_{t-1,i} + \zeta}} \left(\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\alpha_{1}D_{1}} + \alpha_{1}D_{1}L_{0}^{2}\eta^{2} \frac{d(1 - \beta_{1})^{2}}{(1 - \beta_{2})(1 - \frac{\beta_{1}^{2}}{\beta_{2}})} + 2\eta L_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2} \frac{\sqrt{d}(1 - \beta_{1})}{\sqrt{1 - \beta_{2}}\sqrt{1 - \frac{\beta_{1}^{2}}{\beta_{2}}}} \right) \right].$$

$$(72)$$

Since we define $m_0 = 0$, then $x_0 = x_1$ so this inequality works for t = 1 too. Note many terms in the RHS of (72) can be reduced by telescoping, which will be demonstrated later. Now we move to the Error 2 term. For any $\alpha_3 > 0, 0 < \beta_1^2 < \beta_2$ and $\beta_2 > 0.5$, we have that

$$\underbrace{\mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \frac{\eta \odot \frac{\beta_{1}\boldsymbol{m}_{t-1}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1}+\beta_{2}\zeta}} - \eta \odot \frac{\beta_{1}\boldsymbol{m}_{t-1}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1}+\zeta}}\right\rangle |\mathcal{F}_{t}\right]}_{\text{Error 2}} \\
= \frac{\eta \beta_{1}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \mathbb{E}\left[\left\langle \nabla f(\boldsymbol{x}_{t}), \frac{(1 - \beta_{2})\zeta}{(\sqrt{\beta_{2}\boldsymbol{v}_{t-1}+\zeta}) \odot (\sqrt{\beta_{2}\boldsymbol{v}_{t-1}} + \beta_{2}\zeta)} \odot (\sqrt{\beta_{2}\boldsymbol{v}_{t-1}+\zeta} + \sqrt{\beta_{2}\boldsymbol{v}_{t-1}} + \beta_{2}\zeta)}\right] \odot \boldsymbol{m}_{t}\right\rangle |\mathcal{F}_{t}\right] \\
\leq \sum_{i=1}^{d} \frac{\eta \beta_{1}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} |\partial_{i}f(\boldsymbol{x}_{t})| \frac{(1 - \beta_{2})\zeta}{(\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta})(\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}} + \beta_{2}\zeta)}(\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta} + \sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}} + \beta_{2}\zeta)}} \\
\leq \sum_{i=1}^{d} \frac{\eta \beta_{1}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} |\partial_{i}f(\boldsymbol{x}_{t})| \frac{(1 - \beta_{2})\zeta}{(\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta})\sqrt{\zeta}} \frac{1 - \beta_{1}}{\sqrt{1 - \beta_{2}}\sqrt{1 - \frac{\beta_{1}^{2}}{\beta_{2}}}} \\
\leq \sum_{i=1}^{d} \frac{\eta \beta_{1}(1 - \beta_{1})\sqrt{\zeta}}{(1 - \frac{\beta_{1}}{\beta_{2}})\sqrt{1 - \frac{\beta_{1}^{2}}{\beta_{2}}}} \left(\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{3}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} + \frac{\alpha_{3}(1 - \beta_{2})}{2\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}\right), \tag{73}$$

where the second inequality is due to Lemmma E.1 and $2\beta_2 \ge 1$. For Error 3, we have that

$$\langle
abla f(oldsymbol{u}_t) -
abla f(oldsymbol{x}_t), oldsymbol{u}_t - oldsymbol{u}_{t+1}
angle$$

$$\leq \sum_{i=1}^{d} |\partial_{i}f(\boldsymbol{u}_{t}) - \partial_{i}f(\boldsymbol{x}_{t})||\boldsymbol{u}_{t,i} - \boldsymbol{u}_{t+1,i}|$$

$$\leq \sum_{i=1}^{d} (L_{0} + L_{1}|\partial_{i}f(\boldsymbol{x}_{t})|)||\boldsymbol{u}_{t} - \boldsymbol{x}_{t}|||\boldsymbol{u}_{t,i} - \boldsymbol{u}_{t+1,i}|$$

$$\leq \sum_{i=1}^{d} (L_{0} + L_{1}|\partial_{i}f(\boldsymbol{x}_{t})|)\frac{\frac{\beta_{1}}{\sqrt{\beta_{2}}}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}}||\boldsymbol{x}_{t} - \boldsymbol{x}_{t-1}|| \left|\frac{\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} - \frac{\beta_{1}}{\sqrt{\beta_{2}}}\frac{\boldsymbol{x}_{t,i} - \boldsymbol{x}_{t-1,i}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}}\right|, \quad (74)$$

where the second inequality is due to the Assumption 2.3. According to the update process of x_t and Lemmma E.2, it is easy to get that

$$\sum_{i=1}^{d} L_{0} \frac{\frac{\beta_{1}}{\sqrt{\beta_{2}}}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \|\boldsymbol{x}_{t} - \boldsymbol{x}_{t-1}\| \left\| \frac{\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} - \frac{\beta_{1}}{\sqrt{\beta_{2}}} \frac{\boldsymbol{x}_{t,i} - \boldsymbol{x}_{t-1,i}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \right\| \\
\leq \sum_{i=1}^{d} L_{0} \frac{\frac{\beta_{1}}{\sqrt{\beta_{2}}}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \left[\frac{1}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \left(\|\boldsymbol{x}_{t} - \boldsymbol{x}_{t-1}\| \| \boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i} \| \right) + \frac{\frac{\beta_{1}}{\sqrt{\beta_{2}}}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \left(\| \boldsymbol{x}_{t} - \boldsymbol{x}_{t-1,i} \| \right) \right] \\
\leq \sum_{i=1}^{d} L_{0} \frac{\frac{\beta_{1}}{\sqrt{\beta_{2}}}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \left[\frac{1}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \left(\frac{\| \boldsymbol{x}_{t} - \boldsymbol{x}_{t-1} \|^{2}}{2\sqrt{d}} + \frac{\sqrt{d} |\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i} |^{2}}{2\sqrt{d}} \right) + \frac{\frac{\beta_{1}}{\sqrt{\beta_{2}}}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \left(\frac{\| \boldsymbol{x}_{t} - \boldsymbol{x}_{t-1,i} \|^{2}}{2\sqrt{d}} + \frac{\sqrt{d} |\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i} |^{2}}{2\sqrt{d}} \right) \\
\leq L_{0} \eta^{2} \sum_{i=1}^{d} \left[\frac{\frac{\beta_{1}}{2\sqrt{\beta_{2}}} \sqrt{d} + \frac{\beta_{1}^{2}}{\beta_{2}} \sqrt{d}}{\left(1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}\right)^{2}} \frac{|\boldsymbol{m}_{t-1,i} |^{2}}{\boldsymbol{v}_{t-1,i} + \zeta} + \frac{\frac{\beta_{1}}{2\sqrt{\beta_{2}}} \sqrt{d}}{\left(1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}\right)^{2}} \frac{\boldsymbol{m}_{t,i}^{2}}{\boldsymbol{v}_{t,i} + \zeta}} \right]$$
(75)

However, it is hard to bound the remaining $|\partial_i f(\boldsymbol{x}_t)| \| \boldsymbol{x}_t - \boldsymbol{x}_{t-1} \| |\boldsymbol{u}_{t,i} - \boldsymbol{u}_{t+1,i}|$ term by directly applying Lemma E.1, which will induce a $\mathcal{O}(|\partial_i f(\boldsymbol{x}_t)| \frac{\eta^2}{1-\beta_2})$ term and in our affine variance setting we can only bound $\frac{(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i}+\zeta}}$ term. It is worth noting that this challenging additional term is due to the (L_0, L_1) -smoothness and the methods in (Wang et al., 2023a; Li et al., 2023) do not generalize to our case. To solve this challenge, we first bound the additional terms as follows:

$$\sum_{i=1}^{d} |\partial_{i}f(\boldsymbol{x}_{t})| \|\boldsymbol{x}_{t} - \boldsymbol{x}_{t-1}\| \|\boldsymbol{x}_{t,i} - \boldsymbol{x}_{t-1,i} \|$$

$$\leq \sum_{i=1}^{d} |\partial_{i}f(\boldsymbol{x}_{t})| \boldsymbol{x}_{t} - \boldsymbol{x}_{t-1}\| \| \frac{\eta |\boldsymbol{m}_{t-1,i}|}{\sqrt{\boldsymbol{v}_{t-1,i} + \zeta}}$$

$$\leq \sum_{i=1}^{d} \left[\frac{\eta (\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{4}\sqrt{\boldsymbol{v}_{t-1,i} + \zeta}} + \frac{\alpha_{4}\eta \boldsymbol{m}_{t-1,i}^{2}}{2\sqrt{\boldsymbol{v}_{t-1,i} + \zeta}} |\boldsymbol{x}_{t} - \boldsymbol{x}_{t-1}|^{2} \right]$$

$$\leq \sum_{i=1}^{d} \frac{\eta (\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{4}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} + \sum_{i=1}^{d} \frac{\alpha_{4}\eta^{3}\boldsymbol{m}_{t-1,i}^{2}}{2\sqrt{\boldsymbol{v}_{t-1,i} + \zeta}} \frac{d(1 - \beta_{1})^{2}}{(1 - \beta_{2})(1 - \frac{\beta_{1}^{2}}{\beta_{2}})}, \tag{76}$$

for all $\alpha_4 > 0$, where the last inequality is due to Lemma E.1. The motivation is that we can get a $\frac{(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}}$ term and the $\frac{\eta^3 \boldsymbol{m}_{t-1,i}^2}{2\sqrt{\boldsymbol{v}_{t-1,i} + \zeta}}$ can be bounded by Lemma E.3. Similarly, we have that

$$\sum_{i=1}^{d} |\partial_{i}f(\boldsymbol{x}_{t})| \|\boldsymbol{x}_{t} - \boldsymbol{x}_{t-1}\| \|\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}\|$$

$$\leq \sum_{i=1}^{d} \left[\frac{\eta(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{4}\sqrt{\boldsymbol{v}_{t,i} + \zeta}} + \frac{\alpha_{4}\eta\boldsymbol{m}_{t,i}^{2}}{2\sqrt{\boldsymbol{v}_{t,i} + \zeta}} |\boldsymbol{x}_{t} - \boldsymbol{x}_{t-1}|^{2} \right]$$

$$\leq \sum_{i=1}^{d} \frac{\eta(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{4}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} + \sum_{i=1}^{d} \frac{\alpha_{4}\eta^{3}\boldsymbol{m}_{t,i}^{2}}{2\sqrt{\boldsymbol{v}_{t,i} + \zeta}} \frac{d(1 - \beta_{1})^{2}}{(1 - \beta_{2})(1 - \frac{\beta_{1}^{2}}{\beta_{2}})}.$$
(77)

Combine (63), (64), (72), (73), (74), (75), (76) and (77), and we then have that

$$\begin{split} & \mathbb{E}[\langle \nabla f(\boldsymbol{u}_{t}), \boldsymbol{u}_{t+1} - \boldsymbol{u}_{t} \rangle | \mathcal{F}_{t}] \\ &\leq -\sum_{i=1}^{d} \frac{\eta(1-\beta_{1})}{1-\frac{\beta_{1}}{\sqrt{\beta_{2}}}} \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} \\ &+ \sum_{i=1}^{d} \frac{\eta}{1-\frac{\beta_{1}}{\sqrt{\beta_{2}}}} \frac{1-\beta_{1}}{\sqrt{1-\frac{\beta_{1}}{\beta_{2}}}} \left[\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{0}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} + \frac{\alpha_{0}D_{0}}{2} \mathbb{E} \left[\frac{1}{\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i} + \zeta} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} | \mathcal{F}_{t} \right] \\ &+ \frac{\alpha_{0}D_{1}}{2\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i} + \zeta} \left[\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{(\alpha_{1}D_{1})^{2}} + \alpha_{1}D_{1}L_{0}^{2}\eta^{2} \frac{d(1-\beta_{1})^{2}}{(1-\beta_{2})(1-\frac{\beta_{1}^{2}}{\beta_{2}})} + 2\eta L_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2} \frac{\sqrt{d}(1-\beta_{1})}{\sqrt{1-\beta_{2}}\sqrt{1-\frac{\beta_{1}^{2}}{\beta_{2}}}} \right] \\ &+ \frac{\alpha_{0}D_{1}}{2\sqrt{\beta\boldsymbol{v}_{t-1,i} + \zeta}} \left(\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{(\alpha_{1}D_{1})^{2}} + \alpha_{1}D_{1}L_{0}^{2}\eta^{2} \frac{d(1-\beta_{1})^{2}}{(1-\beta_{2})(1-\frac{\beta_{1}^{2}}{\beta_{2}})} + 2\eta L_{1}(\partial_{i}f(\boldsymbol{x}_{t}))^{2} \frac{\sqrt{d}(1-\beta_{1})}{\sqrt{1-\beta_{2}}\sqrt{1-\frac{\beta_{1}^{2}}{\beta_{2}}}} \right) \right] \\ &+ \sum_{i=1}^{d} \frac{\eta\beta_{1}(1-\beta_{1})\sqrt{\zeta}}{(1-\frac{\beta_{1}}{\sqrt{\beta_{2}}})\sqrt{1-\frac{\beta_{1}^{2}}{\beta_{2}}}} \left(\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{2\alpha_{3}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} + \frac{\alpha_{3}(1-\beta_{2})}{2\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} \right) \\ &+ L_{0}\eta^{2} \sum_{i=1}^{d} \left[\frac{\beta_{1}}{\frac{2\sqrt{\beta_{2}}}{\sqrt{d}}} \sqrt{\frac{\beta_{1}^{2}}{\sqrt{d}}} \frac{m_{t-1,i}^{2}}{\boldsymbol{v}_{t-1,i} + \zeta}} + \frac{\beta_{1}}{2\sqrt{\beta_{2}}\sqrt{d}} \frac{m_{t,i}^{2}}{2\sqrt{\boldsymbol{v}_{t,i} + \zeta}} \right] \\ &+ L_{1} \frac{\beta_{1}^{\frac{\beta_{2}}{\beta_{2}}}}{(1-\frac{\beta_{1}}{\sqrt{\beta_{2}}})^{2}} \left[\sum_{i=1}^{d} \frac{\eta(\partial_{i}f(\boldsymbol{x}_{i}))^{2}}{2\alpha_{4}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}}} + \sum_{i=1}^{d} \frac{\alpha_{4}\eta^{3}m_{t-1,i}^{2}}{2\sqrt{\boldsymbol{v}_{t-1} + \zeta}} \frac{d(1-\beta_{1})^{2}}{(1-\beta_{2})(1-\frac{\beta_{1}^{2}}{\beta_{2}})} \right] \\ &+ L_{1} \frac{\beta_{1}}{\sqrt{\beta_{2}}}} \frac{\beta_{1}}{(1-\frac{\beta_{1}}{\sqrt{\beta_{2}}})^{2}} \mathbb{E} \left[\sum_{i=1}^{d} \frac{\eta(\partial_{i}f(\boldsymbol{x}_{i}))^{2}}{2\alpha_{4}\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}}} + \sum_{i=1}^{d} \frac{\alpha_{4}\eta^{3}m_{t-1,i}^{2}}}{2\sqrt{\boldsymbol{v}_{t,i} + \zeta}} \frac{d(1-\beta_{1})^{2}}{(1-\beta_{2})(1-\frac{\beta_{1}^{2}}{\beta_{2}})} \right]. \end{split}$$

Since $C_1 = 1 - \frac{\beta_1}{\sqrt{\beta_2}}, C_2 = \sqrt{1 - \frac{\beta_1^2}{\beta_2}}$, (78) can be further bounded as follows

$$\mathbb{E}[\langle \nabla f(\boldsymbol{u}_{t}), \boldsymbol{u}_{t+1} - \boldsymbol{u}_{t} \rangle | \mathcal{F}_{t}] \\
\leq \left[-\frac{\eta(1-\beta_{1})}{C_{1}} + \frac{\eta(1-\beta_{1})}{C_{1}C_{2}} \left(\frac{1}{2\alpha_{0}} + \frac{\alpha_{0}}{2\alpha_{1}} + \frac{\eta\alpha_{0}\sqrt{d}D_{1}L_{1}(1-\beta_{1})}{\sqrt{1-\beta_{2}}C_{2}} \right) + \frac{\eta\beta_{1}(1-\beta_{1})\sqrt{\zeta}}{2\alpha_{3}C_{1}C_{2}} + \frac{\eta L_{1}(1-C_{1})^{2}}{2\alpha_{4}C_{1}^{2}} + \frac{\eta L_{1}(1-C_{1})}{2\alpha_{4}C_{1}^{2}} \right] \\
\times \sum_{i=1}^{d} \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} \\
+ \sum_{i=1}^{d} \frac{\eta(1-\beta_{1})}{C_{1}C_{2}} \left[\frac{\alpha_{0}D_{0}}{2} \left(\frac{1}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} - \mathbb{E}\left[\frac{1}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} | \mathcal{F}_{t} \right] \right) + \frac{\alpha_{0}D_{1}}{2} \mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i} + \zeta}} - \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} | \mathcal{F}_{t} \right] \right] \\
+ \frac{\alpha_{0}\alpha_{1}D_{1}^{2}L_{0}^{2}\eta^{3}d^{2}(1-\beta_{1})^{3}}{2(1-\beta_{2})C_{1}C_{2}^{3}\sqrt{\zeta}} + \frac{\alpha_{3}\eta d\beta_{1}(1-\beta_{1})(1-\beta_{2})}{2C_{1}C_{2}} \\
+ \sum_{i=1}^{d} \frac{\eta^{2}((1-C_{1})^{2}+0.5(1-C_{1}))\sqrt{d}L_{0}}{C_{1}^{2}} \frac{\boldsymbol{m}_{t-1,i}^{2}}{\boldsymbol{v}_{t-1,i} + \zeta} + \sum_{i=1}^{d} \frac{\eta^{2}0.5(1-C_{1})\sqrt{d}L_{0}}{C_{1}^{2}} \mathbb{E}\left[\frac{\boldsymbol{m}_{t,i}^{2}}{\boldsymbol{v}_{t,i} + \zeta} | \mathcal{F}_{t} \right] \\
+ \sum_{i=1}^{d} \frac{\alpha_{4}\eta^{3}(1-\beta_{1})^{2}(1-C_{1})^{2}dL_{1}}{2(1-\beta_{2})C_{1}^{2}C_{2}^{2}} \frac{\boldsymbol{m}_{t-1,i}^{2}}{\sqrt{\boldsymbol{v}_{t-1,i} + \zeta}} + \sum_{i=1}^{d} \frac{\alpha_{4}\eta^{3}(1-\beta_{1})^{2}(1-C_{1})dL_{1}}{2(1-\beta_{2})C_{1}^{2}C_{2}^{2}} \mathbb{E}\left[\frac{\boldsymbol{m}_{t,i}^{2}}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}} | \mathcal{F}_{t} \right].$$
(79)

This completes the proof.

G. Proof of Lemma 4.2

For (L_0, L_1) -smooth objective functions, we have the following descent inequality (Lemma 1 in (Crawshaw et al., 2022)):

$$\underbrace{\mathbb{E}[\langle \nabla f(\boldsymbol{u}_{t}), \boldsymbol{u}_{t} - \boldsymbol{u}_{t+1} \rangle | \mathcal{F}_{t}]}_{\text{First Order}} \leq f(\boldsymbol{u}_{t}) - \mathbb{E}[f(\boldsymbol{u}_{t+1}) | \mathcal{F}_{t}] + \underbrace{\sum_{i=1}^{d} \frac{L_{0}}{2\sqrt{d}} \mathbb{E}[\|\boldsymbol{u}_{t+1} - \boldsymbol{u}_{t}\| | \boldsymbol{u}_{t+1,i} - \boldsymbol{u}_{t,i} | | \mathcal{F}_{t}]}_{\text{Second Order}} + \underbrace{\sum_{i=1}^{d} \frac{L_{1} \| \partial_{i} f(\boldsymbol{u}_{t}) \|}{2} \mathbb{E}[\|\boldsymbol{u}_{t+1} - \boldsymbol{u}_{t}\| | \boldsymbol{u}_{t+1,i} - \boldsymbol{u}_{t,i} | | \mathcal{F}_{t}]}_{\text{Additional Term}}.$$
(80)

The first-order term is bounded by Lemma F.1, we then only need to bound the remaining two terms. For the second-order term, based on the definition of u_t and update process of x_t , we have that

$$\sum_{i=1}^{d} \frac{L_{0}}{2\sqrt{d}} \mathbb{E}[\|\boldsymbol{u}_{t+1} - \boldsymbol{u}_{t}\| | \boldsymbol{u}_{t+1,i} - \boldsymbol{u}_{t,i} | | \mathcal{F}_{t}] \\
\leq \sum_{i=1}^{d} \frac{L_{0}}{2\sqrt{d}} \mathbb{E}\left[\frac{1}{2\sqrt{d}} \left\|\frac{\boldsymbol{x}_{t+1} - \boldsymbol{x}_{t}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} - \frac{\beta_{1}}{\sqrt{\beta_{2}}} \frac{\boldsymbol{x}_{t} - \boldsymbol{x}_{t-1}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}}\right\|^{2} + \frac{\sqrt{d}}{2} \left|\frac{\boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}} - \frac{\beta_{1}}{\sqrt{\beta_{2}}} \frac{\boldsymbol{x}_{t,i} - \boldsymbol{x}_{t-1,i}}{1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}}}\right|^{2} | \mathcal{F}_{t} \right] \\
\leq \sum_{i=1}^{d} \frac{L_{0}}{2\sqrt{d}} \mathbb{E}\left[\frac{2\sqrt{d}}{(1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}})^{2}} | \boldsymbol{x}_{t+1,i} - \boldsymbol{x}_{t,i} |^{2} + \frac{2\sqrt{d}\frac{\beta_{1}^{2}}{\beta_{2}}}{(1 - \frac{\beta_{1}}{\sqrt{\beta_{2}}})^{2}} | \boldsymbol{x}_{t,i} - \boldsymbol{x}_{t-1,i} |^{2} | \mathcal{F}_{t} \right] \\
= \sum_{i=1}^{d} \frac{L_{0}}{2\sqrt{d}} \left(\frac{2\sqrt{d}}{C_{1}^{2}} \mathbb{E}\left[\frac{\eta^{2}\boldsymbol{m}_{t,i}^{2}}{\boldsymbol{v}_{t,i} + \zeta} | \mathcal{F}_{t}\right] + \frac{2\sqrt{d}(1 - C_{1})^{2}}{C_{1}^{2}} \frac{\eta^{2}\boldsymbol{m}_{t-1,i}^{2}}{\boldsymbol{v}_{t-1,i} + \zeta}\right).$$
(81)

Now we focus on the additional term. According to the definition of u_t and update process of x_t , for $\alpha_4 > 0$ we have that

$$\begin{split} &\sum_{i=1}^{d} \frac{L_1 \|\partial_i f(\mathbf{u}_t)\|}{2} \mathbb{E}[\|\mathbf{u}_{t+1} - \mathbf{u}_t\| \|\mathbf{u}_{t+1,i} - \mathbf{u}_{t,i}\| |\mathcal{F}_t] \\ &\leq \sum_{i=1}^{d} \frac{L_1}{2} (|\partial_i f(\mathbf{x}_t)| + (L_0 + L_1 |\partial_i f(\mathbf{x}_t)|) \| \mathbf{u}_t - \mathbf{x}_t\|) \mathbb{E}[\|\mathbf{u}_{t+1} - \mathbf{x}_t\| \|\mathbf{u}_{t+1,i} - \mathbf{u}_{t,i}\| |\mathcal{F}_t] \\ &\leq \sum_{i=1}^{d} \frac{L_1}{2} \left(|\partial_i f(\mathbf{x}_t)| + L_0 \frac{\frac{\beta_1}{\sqrt{\beta_2}}}{1 - \frac{\beta_1}{\sqrt{\beta_2}}} \| \mathbf{x}_t - \mathbf{x}_{t-1}\| + L_1 |\partial_i f(\mathbf{x}_t)| \frac{\frac{\beta_1}{\sqrt{\beta_2}}}{1 - \frac{\beta_1}{\sqrt{\beta_2}}} \| \mathbf{x}_t - \mathbf{x}_{t-1}\| \right) \\ &\times \mathbb{E}[\|\mathbf{u}_{t+1} - \mathbf{u}_t\| \| \mathbf{u}_{t+1,i} - \mathbf{u}_{t,i}\| |\mathcal{F}_t] \\ &\leq \sum_{i=1}^{d} \frac{L_1}{2} \left(1 + L_1 \frac{1 - C_1}{C_1} \eta \sqrt{d} \frac{1 - \beta_1}{\sqrt{1 - \beta_2 C_2}} \right) \\ &\times \left(\sqrt{d} \frac{2 - C_1}{C_1} \frac{\eta(1 - \beta_1)}{\sqrt{1 - \beta_2 C_2}} |\partial_i f(\mathbf{x}_t)| \left(\frac{1}{C_1} \mathbb{E} \left[\frac{\eta |\mathbf{m}_{t,i}|}{\sqrt{\mathbf{v}_{t,i} + \zeta}} |\mathcal{F}_t \right] + \frac{1 - C_1}{C_1} \frac{\eta |\mathbf{m}_{t-1,i}|}{\sqrt{\mathbf{v}_{t-1,i} + \zeta}} \right) \right) \\ &+ \sum_{i=1}^{d} \frac{L_1 L_0}{2} \frac{1 - C_1}{C_1} \eta \sqrt{d} \frac{1 - \beta_1}{\sqrt{1 - \beta_2 C_2}} \times \left(\frac{2\sqrt{d}}{C_1^2} \mathbb{E} \left[\frac{\eta^2 \mathbf{m}_{t,i}^2}{\mathbf{v}_{t,i} + \zeta} |\mathcal{F}_t \right] + \frac{2\sqrt{d}(1 - C_1)^2}{C_1^2} \frac{\eta^2 \mathbf{m}_{t-1,i}^2}{\mathbf{v}_{t-1,i} + \zeta} \right) \\ &\leq \sum_{i=1}^{d} \frac{L_1}{2} \left(1 + L_1 \frac{1 - C_1}{C_1} \eta \sqrt{d} \frac{1 - \beta_1}{\sqrt{1 - \beta_2 C_2}} \right) \left[\frac{2\sqrt{d}}{C_1^2} \left(\frac{\eta}{2\alpha_4} \frac{(\partial_i f(\mathbf{x}_t))^2}{\sqrt{\beta_2 \mathbf{v}_{t-1,i} + \zeta}} + \frac{\alpha_4 \eta^3(1 - \beta_1)^2}{2(1 - \beta_2) C_2^2} \mathbb{E} \left[\frac{\mathbf{m}_{t,i}^2}{\sqrt{\mathbf{v}_{t,i} + \zeta}} |\mathcal{F}_t \right] \right) \\ &= \frac{2\sqrt{d}(2 - C_1)^2}{C_1^2} \left(\frac{\eta}{2\alpha_4} \frac{(\partial_i f(\mathbf{x}_t))^2}{\sqrt{\beta_2 \mathbf{v}_{t-1,i} + \zeta}} + \frac{\alpha_4 \eta^3(1 - \beta_1)^2}{2(1 - \beta_2) C_2^2} \frac{\mathbf{m}_{t-1,i}^2}{\sqrt{\mathbf{v}_{t-1,i} + \zeta}} \right) \right] \end{aligned}$$

$$+\sum_{i=1}^{d} \frac{L_1 L_0}{2} \frac{1-C_1}{C_1} \eta \sqrt{d} \frac{1-\beta_1}{\sqrt{1-\beta_2} C_2} \left(\frac{2\sqrt{d}}{C_1^2} \mathbb{E}\left[\frac{\eta^2 \boldsymbol{m}_{t,i}^2}{\boldsymbol{v}_{t,i}+\zeta} | \mathcal{F}_t \right] + \frac{2\sqrt{d}(1-C_1)^2}{C_1^2} \frac{\eta^2 \boldsymbol{m}_{t-1,i}^2}{\boldsymbol{v}_{t-1,i}+\zeta} \right),$$
(82)

where the first inequality is due to the (L_0, L_1) -smoothness, the third inequality is due to Lemma E.1 and (81), and the last inequality is due to (76) and (77). Combine Lemma F.1, (80), (81) and (82) together, and we have that

$$\begin{split} & \left[\frac{\eta(1-\beta_{1})}{C_{1}}-\frac{\eta(1-\beta_{1})}{C_{1}C_{2}}\left(\frac{1}{2\alpha_{0}}+\frac{\alpha_{0}}{2\alpha_{1}}+\frac{\eta\alpha_{0}\sqrt{d}D_{1}L_{1}(1-\beta_{1})}{\sqrt{1-\beta_{2}C_{2}}}\right)-\frac{\eta\beta_{1}(1-\beta_{1})\sqrt{\zeta}}{2\alpha_{3}C_{1}C_{2}}-\frac{\eta L_{1}(1-C_{1})^{2}}{2\alpha_{4}C_{1}^{2}}-\frac{\eta L_{1}(1-C_{1})^{2}}{2\alpha_{4}C_{1}^{2}}\right)-\frac{\sqrt{d}L_{1}}{2\alpha_{4}C_{1}^{2}}+\frac{\eta(2-C_{1})^{2}}{\alpha_{4}C_{1}^{2}}\right)\right]\times \sum_{i=1}^{d}\frac{(\partial_{i}f(x_{i}))^{2}}{\sqrt{\beta_{2}v_{t-1,i}+\zeta}}\\ \leq f(u_{t})-\mathbb{E}[f(u_{t+1})]\mathcal{F}_{t}]\\ &+\sum_{i=1}^{d}\frac{\eta(1-\beta_{1})}{C_{1}C_{2}}\left[\frac{\alpha_{0}D_{0}}{2}\left(\frac{1}{\sqrt{\beta_{2}v_{t-1,i}+\zeta}}-\mathbb{E}\left[\frac{1}{\sqrt{v_{t,i}+\zeta}}|\mathcal{F}_{t}\right]\right)+\frac{\alpha_{0}D_{1}}{2}\mathbb{E}\left[\frac{(\partial_{i}f(x_{t-1}))^{2}}{\sqrt{\beta_{2}v_{t-1,i}+\zeta}}-\frac{(\partial_{i}f(x_{t}))^{2}}{\sqrt{v_{t,i}+\zeta}}|\mathcal{F}_{t}\right]\right]\\ &+\frac{\alpha_{0}\alpha_{1}D_{1}^{2}L_{0}^{2}\eta^{3}d^{2}(1-\beta_{1})^{3}}{2(1-\beta_{2})C_{1}C_{2}^{3}\sqrt{\zeta}}+\frac{\alpha_{3}\eta d\beta_{1}(1-\beta_{1})(1-\beta_{2})}{2C_{1}C_{2}}\\ &+\sum_{i=1}^{d}\left[\frac{\eta^{2}(2(1-C_{1})^{2}+0.5(1-C_{1}))\sqrt{d}L_{0}}{C_{1}^{2}}+\frac{dL_{1}L_{0}}{2}\frac{1-C_{1}}{C_{1}}\eta\frac{1-\beta_{1}}{\sqrt{1-\beta_{2}}C_{2}}\frac{2}{C_{1}^{2}}\eta^{2}\right]\mathbb{E}\left[\frac{m_{t,i}^{2}}{v_{t-1,i}+\zeta}\right]\\ &+\sum_{i=1}^{d}\left[\frac{\eta^{2}(0.5(1-C_{1})+1)\sqrt{d}L_{0}}{C_{1}^{2}}+\frac{dL_{1}L_{0}}{2}\frac{1-C_{1}}{C_{1}}\eta\sqrt{d}\frac{1-\beta_{1}}{\sqrt{1-\beta_{2}}C_{2}}\frac{2}{C_{1}^{2}}\eta^{2}\right]\frac{2\sqrt{d}(2-C_{1})^{2}}{C_{1}^{2}}\frac{\alpha_{4}\eta^{3}(1-\beta_{1})^{2}}{2(1-\beta_{2})C_{1}^{2}C_{2}^{2}}+\frac{L_{1}}{2}\left(1+L_{1}\frac{1-C_{1}}{C_{1}}\eta\sqrt{d}\frac{1-\beta_{1}}{\sqrt{1-\beta_{2}}C_{2}}\right)\frac{2\sqrt{d}(2-C_{1})^{2}}{C_{1}^{2}}\frac{\alpha_{4}\eta^{3}(1-\beta_{1})^{2}}{2(1-\beta_{2})C_{2}^{2}}\right]\mathbb{E}\left[\frac{m_{t,i}^{2}}{\sqrt{v_{t,i}+\zeta}}\right]\\ &\times\frac{m_{t-1,i}^{2}}{\sqrt{v_{t-1,i}+\zeta}}\\ &+\sum_{i=1}^{d}\left[\frac{\alpha_{4}\eta^{3}(1-\beta_{1})^{2}(1-C_{1})dL_{1}}{2(1-\beta_{2})C_{1}^{2}C_{2}^{2}}+\frac{L_{1}}{2}\left(1+L_{1}\frac{1-C_{1}}{C_{1}}\eta\sqrt{d}\frac{1-\beta_{1}}{\sqrt{1-\beta_{2}}C_{2}}\right)\frac{2\sqrt{d}}{C_{1}^{2}}\frac{\alpha_{4}\eta^{3}(1-\beta_{1})^{2}}{2(1-\beta_{2})C_{2}^{2}}}\right]\mathbb{E}\left[\frac{m_{t,i}^{2}}{\sqrt{v_{t,i}+\zeta}}\right]\\ &\times\frac{m_{t}^{2}}{\sqrt{v_{t-1,i}+\zeta}}\\ &+\sum_{i=1}^{d}\left[\frac{\alpha_{4}\eta^{3}(1-\beta_{1})^{2}(1-C_{1})dL_{1}}{2(1-\beta_{2})C_{1}^{2}}+\frac{L_{1}}{2}\left(1+L_{1}\frac{1-C_{1}}{C_{1}}\eta\sqrt{d}\frac{1-\beta_{1}}{\sqrt{1-\beta_{2}}C_{2}}\right)\frac{2\sqrt{d}}{C_{1}^{2}}\frac{\alpha_{4}\eta^{3}(1-\beta_{1})^{2}}{2(1-\beta_{2})C_{2}^{2}}\right]\mathbb{E}\left[\frac{m_{t}^{2}}{\sqrt{v_{t,i}+\zeta}}\right]\\ &+\sum_{i=1}^{d}\left[\frac{\alpha_{4}\eta^{3}(1-\beta_{1})$$

It is worth noting that (30) and (31) still hold for Adam since the update of v_t does not change. Specifically, for any $i \in [d]$ we have that

$$\sum_{t=1}^{T} \mathbb{E}\left[\frac{1}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}} - \frac{1}{\sqrt{\boldsymbol{v}_{t,i} + \zeta}}\right] \le \frac{1}{\sqrt{\zeta}} + T \frac{1 - \sqrt{\beta_2}}{\sqrt{\zeta}}$$
(84)

and since $x_0 = x_1$ we have that

$$\mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{0}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{0,i}+\zeta}} - \frac{(\partial_{i}f(\boldsymbol{x}_{1}))^{2}}{\sqrt{\boldsymbol{v}_{1,i}+\zeta}}\right] + \sum_{t=2}^{T}\mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t-1}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}} - \frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\boldsymbol{v}_{t,i}+\zeta}}\right] \\
\leq \frac{(\partial_{i}f(\boldsymbol{x}_{1}))^{2}}{\sqrt{\zeta}} + \sum_{t=1}^{T-1}(1-\beta_{2})\mathbb{E}\left[\frac{(\partial_{i}f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2}\boldsymbol{v}_{t-1,i}+\zeta}}\right]$$
(85)

By taking expectations and telescoping (83) for t = 1 to T, based on (84), (85), Lemma E.2 and Lemma E.3, we have that

$$\left[\frac{\eta(1-\beta_1)}{C_1} - \frac{\eta(1-\beta_1)}{C_1 C_2} \left(\frac{1}{2\alpha_0} + \frac{\alpha_0}{2\alpha_1} + \frac{\eta\alpha_0\sqrt{d}D_1L_1(1-\beta_1)}{\sqrt{1-\beta_2}C_2} \right) - \frac{\eta\beta_1(1-\beta_1)\sqrt{\zeta}}{2\alpha_3 C_1 C_2} - \frac{\eta L_1(1-C_1)^2}{2\alpha_4 C_1^2} - \frac{\eta L_1(1-C_1)}{2\alpha_4 C_1^2} - \frac{\sqrt{d}L_1(1-C_1)}{2\alpha_4 C_1^2} - \frac{\sqrt{d$$

$$\begin{split} &\leq f(\boldsymbol{u}_{1}) - \mathbb{E}[f(\boldsymbol{u}_{T+1})|\mathcal{F}_{t}] \\ &+ \sum_{i=1}^{d} \frac{\eta(1-\beta_{1})}{C_{1}C_{2}} \left[\frac{\alpha_{0}D_{0}}{2} \left(\frac{1}{\sqrt{\zeta}} + T\frac{1-\sqrt{\beta_{2}}}{\sqrt{\zeta}} \right) + \frac{\alpha_{0}D_{1}}{2} \left(\frac{(\partial_{i}f(\boldsymbol{x}_{1}))^{2}}{\sqrt{\zeta}} + \sum_{t=1}^{T-1} (1-\beta_{2})\mathbb{E} \left[\frac{(\partial_{i}f(\boldsymbol{x}_{1}))^{2}}{\sqrt{\beta_{2}}\boldsymbol{v}_{t-1,i} + \zeta} \right] \right) \right] \\ &+ T\frac{\alpha_{0}\alpha_{1}D_{1}^{2}L_{0}^{2}\eta^{3}d^{2}(1-\beta_{1})^{3}}{2(1-\beta_{2})C_{1}C_{3}^{2}\sqrt{\zeta}} + T\frac{\alpha_{3}\eta d\beta_{1}(1-\beta_{1})(1-\beta_{2})}{2C_{1}C_{2}} \\ &+ \left[\frac{\eta^{2}2(1-C_{1})^{2}\sqrt{d}L_{0}}{C_{1}^{2}} + \frac{\eta^{2}(2-C_{1})\sqrt{d}L_{0}}{C_{1}^{2}} + \frac{dL_{1}L_{0}}{2}\frac{1-C_{1}}{C_{1}}\eta \frac{1-\beta_{1}}{\sqrt{1-\beta_{2}}C_{2}} \frac{2(1-C_{1})^{2}+2}{C_{1}^{2}}\eta^{2} \right] \\ &\times \sum_{i=1}^{d} \mathbb{E} \left[\frac{(1-\beta_{1})^{2}}{(1-\frac{\beta_{1}}{\sqrt{\beta_{2}}})^{2}(1-\beta_{2})} \left(\ln\left(\frac{\boldsymbol{v}_{T,i}}{\boldsymbol{v}_{0,i}}\right) - T\ln(\beta_{2}) \right) \right] \\ &+ \left[\frac{\alpha_{4}\eta^{3}(1-\beta_{1})^{2}dL_{1}((1-C_{1})+(1-C_{1})^{2})}{2(1-\beta_{2})C_{1}^{2}C_{2}^{2}} + \frac{\sqrt{d}L_{1}}{2} \left(1+L_{1}\frac{1-C_{1}}{C_{1}}\eta\sqrt{d}\frac{1-\beta_{1}}{\sqrt{1-\beta_{2}}C_{2}} \right) \frac{2+2(2-C_{1})^{2}}{C_{1}^{2}}\frac{\alpha_{4}\eta^{3}(1-\beta_{1})^{2}}{2(1-\beta_{2})C_{2}^{2}} \\ &\times \sum_{i=1}^{d} \mathbb{E} \left[\frac{(1-\beta_{1})^{2}}{(1-\beta_{2}\sqrt{\beta_{2}})^{2}} \left(\frac{2}{1-\beta_{2}}(\sqrt{\boldsymbol{v}_{T,i}}-\sqrt{\boldsymbol{v}_{0,i}}) + \sum_{t=1}^{T} 2\sqrt{\boldsymbol{v}_{t-1,i}} \right) \right]. \end{aligned}$$

Moreover, for any a > 0 and $i \in [d]$ we have that $\ln(a) \le a - 1$. We then have that

$$\frac{1}{T}\mathbb{E}[\ln(\boldsymbol{v}_{T,i})] = \frac{2}{T}\mathbb{E}[\ln(\sqrt{\boldsymbol{v}_{T,i}})] \leq \frac{2}{T}\mathbb{E}[\sqrt{\boldsymbol{v}_{T,i}}] \leq \frac{2\sqrt{D_0 + \boldsymbol{v}_{0,i}}}{T} + \frac{2}{T}\sum_{i=0}^{T-1}\mathbb{E}\left[\sqrt{\beta_2^i(1 - \beta_2)D_1}|\partial_i f(\boldsymbol{x}_{T-i})|\right] \\
\leq \frac{2\sqrt{D_0 + \boldsymbol{v}_{0,i}}}{T} + \frac{2}{T}\sum_{i=0}^{T-1}\mathbb{E}\left[\sqrt{(1 - \beta_2)D_1}|\partial_i f(\boldsymbol{x}_{T-i})|\right], \quad (87)$$

where the second inequality is due to (39). In addition, for $\beta_2 \geq 0.5$ we have that

$$-\ln(\beta_2) = \ln\left(\frac{1}{\beta_2}\right) \le \frac{1}{\beta_2} - 1 \le 2(1 - \beta_2).$$
(88)

Combine (40), (86), (87) and (88), we then can show $\sum_{i=1}^{d} \sum_{t=1}^{T} \mathbb{E}\left[\frac{(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2 \boldsymbol{v}_{t-1,i} + \zeta}}\right]$ is upper bounded by a function of $\sum_{t=1}^{T} \mathbb{E}\left[\|\nabla f(\boldsymbol{x}_t)\|\right]$. Specifically, we can get

$$\begin{split} & \left[\frac{\eta(1-\beta_1)}{C_1} - \frac{\eta(1-\beta_1)}{C_1C_2} \left(\frac{1}{2\alpha_0} + \frac{\alpha_0}{2\alpha_1} + \frac{\eta\alpha_0\sqrt{d}D_1L_1(1-\beta_1)}{\sqrt{1-\beta_2C_2}} \right) - \frac{\eta\beta_1(1-\beta_1)\sqrt{\zeta}}{2\alpha_3C_1C_2} - \frac{\eta L_1(1-C_1)^2}{2\alpha_4C_1^2} - \frac{\eta L_1(1-C_1)}{2\alpha_4C_1^2} \right) \\ & - \frac{\sqrt{d}L_1}{2} \left(1 + L_1 \frac{1-C_1}{C_1} \eta\sqrt{d} \frac{1-\beta_1}{\sqrt{1-\beta_2C_2}} \right) \left(\frac{\eta}{\alpha_4C_1^2} + \frac{\eta(2-C_1)^2}{\alpha_4C_1^2} \right) - \frac{\eta(1-\beta_1)}{C_1C_2} \frac{\alpha_0D_1(1-\beta_2)}{2} \right] \times \sum_{t=1}^T \sum_{i=1}^d \frac{(\partial_i f(\boldsymbol{x}_t))^2}{\sqrt{\beta_2\boldsymbol{v}_{t-1,i} + \zeta}} \\ & \leq f(\boldsymbol{u}_1) - \mathbb{E}[f(\boldsymbol{u}_{T+1})|\mathcal{F}_t] + \frac{\eta\alpha_0dD_0(1-\beta_1)}{2C_1C_2\sqrt{\zeta}} + \frac{\eta\alpha_0D_1(1-\beta_1)||\nabla f(\boldsymbol{x}_1)||^2}{2C_1C_2\sqrt{\zeta}} \\ & + T \frac{\eta d\alpha_0D_0(1-\beta_1)(1-\beta_2)}{2C_1C_2\sqrt{\zeta}} + T \frac{\alpha_0\alpha_1D_1^2L_0^2\eta^3d^2(1-\beta_1)^3}{2(1-\beta_2)C_1C_2^3\sqrt{\zeta}} + T \frac{\alpha_3\eta d\beta_1(1-\beta_1)(1-\beta_2)}{2C_1C_2} \\ & + \left[\frac{\eta^22(1-C_1)^2\sqrt{d}L_0}{C_1^2} + \frac{\eta^2(2-C_1)\sqrt{d}L_0}{C_1^2} + \frac{dL_1L_0}{2} \frac{1-C_1}{C_1} \eta \frac{1-\beta_1}{\sqrt{1-\beta_2C_2}} \frac{2(1-C_1)^2+2}{C_1^2} \eta^2 \right] \\ & \times \sum_{i=1}^d \mathbb{E} \left[\frac{(1-\beta_1)^2}{(1-\frac{\beta_1}{\sqrt{\beta_2}})^2(1-\beta_2)} \left(2\sqrt{D_0+\boldsymbol{v}_{0,i}} + \sum_{t=1}^T \mathbb{E}[2\sqrt{(1-\beta_2)D_1}|\partial_i f(\boldsymbol{x}_t)|] + 2T(1-\beta_2) - \ln(\boldsymbol{v}_{0,i}) \right) \right] \\ & + \left[\frac{\alpha_4\eta^3(1-\beta_1)^2dL_1((1-C_1)+(1-C_1)^2)}{2(1-\beta_2)C_1^2C_2^2} + \frac{\sqrt{d}L_1}{2} \left(1 + L_1 \frac{1-C_1}{C_1} \eta \sqrt{d} \frac{1-\beta_1}{\sqrt{1-\beta_2C_2}} \right) \frac{2+2(2-C_1)^2}{C_1^2} \frac{\alpha_4\eta^3(1-\beta_1)^2}{2(1-\beta_2)C_2^2} \right] \right] \\ \end{split}$$

$$\times \sum_{i=1}^{d} \mathbb{E} \left[\frac{(1-\beta_{1})^{2}}{(1-\frac{\beta_{1}}{\sqrt{\beta_{2}}})^{2}} \left(\frac{2}{1-\beta_{2}} \left(\sqrt{D_{0} + \boldsymbol{v}_{0,i}} + \sum_{t=1}^{T} \mathbb{E} \left[\sqrt{(1-\beta_{2})D_{1}} |\partial_{i}f(\boldsymbol{x}_{t})| \right] \right) + 2T\sqrt{D_{0} + \boldsymbol{v}_{0,i}} + \frac{4\sqrt{D_{1}}}{\sqrt{1-\beta_{2}}} \sum_{t=1}^{T} \mathbb{E} \left[|\partial_{i}f(\boldsymbol{x}_{t})| \right] \right) \right].$$

$$(89)$$

If we have $\alpha_0 \geq \frac{21}{2C_2}, \alpha_1 \geq \frac{21\alpha_0}{2C_2}, \alpha_3 \geq \frac{7\beta_1\sqrt{\zeta}}{2C_2}, \alpha_4 \geq \frac{14L_1\sqrt{d}(2-C_1)^2}{C_1(1-\beta_1)}, \text{ for } \eta \leq \min\left(\frac{C_2^2\sqrt{1-\beta_2}}{21\alpha_0\sqrt{d}D_1L_1(1-\beta_1)}, \frac{C_1C_2\sqrt{1-\beta_2}}{\sqrt{d}L_1(1-C_1)(1-\beta_1)}\right), 1-\beta_2 \leq (\frac{2C_2}{7\alpha_0D_1}) \text{ we have that}$

$$\left[\frac{\eta(1-\beta_{1})}{C_{1}} - \frac{\eta(1-\beta_{1})}{C_{1}C_{2}} \left(\frac{1}{2\alpha_{0}} + \frac{\alpha_{0}}{2\alpha_{1}} + \frac{\eta\alpha_{0}\sqrt{d}D_{1}L_{1}(1-\beta_{1})}{\sqrt{1-\beta_{2}}C_{2}}\right) - \frac{\eta\beta_{1}(1-\beta_{1})\sqrt{\zeta}}{2\alpha_{3}C_{1}C_{2}} - \frac{\eta L_{1}(1-C_{1})^{2}}{2\alpha_{4}C_{1}^{2}} - \frac{\eta L_{1}(1-C_{1})}{2\alpha_{4}C_{1}^{2}} - \frac{\eta L_{1}(1-\beta_{1})}{2\alpha_{4}C_{1}^{2}} - \frac{\eta L_{1}(1-\beta_{1})}{2\alpha_{4}C_{1}^{2}}\right] - \frac{\sqrt{d}L_{1}}{2} \left(1 + L_{1}\frac{1-C_{1}}{C_{1}}\eta\sqrt{d}\frac{1-\beta_{1}}{\sqrt{1-\beta_{2}}C_{2}}\right) \left(\frac{\eta}{\alpha_{4}C_{1}^{2}} + \frac{\eta(2-C_{1})^{2}}{\alpha_{4}C_{1}^{2}}\right) - \frac{\eta(1-\beta_{1})}{C_{1}C_{2}}\frac{\alpha_{0}D_{1}(1-\beta_{2})}{2}\right] \ge \frac{\eta(1-\beta_{1})}{7C_{1}}.$$
(90)

Since
$$\Delta' = f(u_1) - f^* + \frac{\eta \alpha_0 dD_0(1-\beta_1)}{2C_1 C_2 \sqrt{\zeta}} + \frac{\eta \alpha_0 D_1(1-\beta_1) \|\nabla f(x_1)\|^2}{2C_1 C_2 \sqrt{\zeta}}, C_3 = \left[\frac{2(1-C_1)^2 \sqrt{d}L_0}{C_1^2} + \frac{(2-C_1)\sqrt{d}L_0}{C_1^2} + \sqrt{d}L_0 \frac{(1-C_1)^2+1}{C_1^2} \right],$$

$$C_4 = \left[\frac{\alpha_4(1-\beta_1)^2 dL_1((1-C_1)+(1-C_1)^2)}{2C_1^2 C_2^2} + \sqrt{d}L_1 \frac{2+2(2-C_1)^2}{C_1^2} \frac{\alpha_4(1-\beta_1)^2}{2C_2^2} \right],$$
 we then have that
$$\frac{\eta(1-\beta_1)}{7C_1} \sum_{i=1}^d \sum_{t=1}^T \mathbb{E} \left[\frac{(\partial_i f(x_t))^2}{\sqrt{\beta_2 v_{t-1,i} + \zeta}} \right]$$

$$\leq \Delta' + T \frac{\eta d\alpha_0 D_0(1-\beta_1)(1-\beta_2)}{2C_1 C_2 \sqrt{\zeta}} + T \frac{\alpha_0 \alpha_1 D_1^2 L_0^2 \eta^3 d^2(1-\beta_1)^3}{2(1-\beta_2) C_1 C_2^3 \sqrt{\zeta}} + T \frac{\alpha_3 \eta d\beta_1(1-\beta_1)(1-\beta_2)}{2C_1 C_2}$$

$$+ \frac{\eta^2 C_3(1-\beta_1)^2}{C_1^2(1-\beta_2)} \times \sum_{i=1}^d \mathbb{E} \left[\left(2\sqrt{D_0 + v_{0,i}} + \sum_{t=1}^T \mathbb{E} [2\sqrt{(1-\beta_2)D_1} |\partial_i f(x_t)|] + 2T(1-\beta_2) - \ln(v_{0,i}) \right) \right]$$

$$+ \frac{\eta^3 C_4}{(1-\beta_2)} \frac{(1-\beta_1)^2}{(1-\frac{\beta_1}{\sqrt{\beta_2}})^2}$$

$$\times \sum_{i=1}^d \mathbb{E} \left[\left(\frac{2}{1-\beta_2} \left(\sqrt{D_0 + v_{0,i}} + \sum_{t=1}^T \mathbb{E} [\sqrt{(1-\beta_2)D_1} |\partial_i f(x_t)|] \right) + 2T\sqrt{D_0 + v_{0,i}} + \frac{4\sqrt{D_1}}{\sqrt{1-\beta_2}} \sum_{t=1}^T \mathbb{E} [|\partial_i f(x_t)|] \right) \right]$$
(91)

It follows that

$$\frac{1}{T} \sum_{t=1}^{T} \mathbb{E} \left[\frac{\|\nabla f(\boldsymbol{x}_{t})\|^{2}}{\sqrt{\beta_{2} \|\boldsymbol{v}_{t-1}\| + \zeta}} \right] \\
\leq \frac{1}{T} \sum_{i}^{d} \sum_{t=1}^{T} \mathbb{E} \left[\frac{(\partial_{i} f(\boldsymbol{x}_{t}))^{2}}{\sqrt{\beta_{2} \boldsymbol{v}_{t-1,i} + \zeta}} \right] \\
\leq \frac{1}{T} \left(\frac{7C_{1}\Delta'}{\eta(1-\beta_{1})} + \frac{7\eta C_{3}(1-\beta_{1})}{C_{1}(1-\beta_{2})} (2d\sqrt{D_{0}} + \|\boldsymbol{v}_{0}\| - \sum_{i=1}^{d} \ln(\boldsymbol{v}_{0,i})) + \frac{14\eta^{2}C_{1}C_{4}(1-\beta_{1})d}{(1-\beta_{2})^{2}(1-\frac{\beta_{1}}{\sqrt{\beta_{2}}})^{2}} \sqrt{D_{0}} + \|\boldsymbol{v}_{0}\| \right) \\
+ \frac{7\alpha_{0}dD_{0}(1-\beta_{2})}{2C_{2}\sqrt{\zeta}} + \frac{7\alpha_{0}\alpha_{1}D_{1}^{2}L_{0}^{2}\eta^{2}d^{2}(1-\beta_{1})^{2}}{2(1-\beta_{2})C_{2}^{3}\sqrt{\zeta}} + \frac{7\alpha_{3}\beta_{1}(1-\beta_{2})d}{2C_{2}} + \frac{14\eta dC_{3}(1-\beta_{1})}{C_{1}} \\
+ \frac{14\eta^{2}C_{1}C_{4}(1-\beta_{1})d\sqrt{D_{0}} + \|\boldsymbol{v}_{0}\|}{(1-\beta_{2})(1-\frac{\beta_{1}}{\sqrt{\beta_{2}}})^{2}} \\
+ \left(\frac{14\eta C_{3}(1-\beta_{1})\sqrt{D_{1}}}{C_{1}\sqrt{1-\beta_{2}}} + \frac{42\eta^{2}C_{1}C_{4}(1-\beta_{1})\sqrt{D_{1}}}{(1-\beta_{2})^{1.5}(1-\frac{\beta_{1}}{\sqrt{\beta_{2}}})^{2}} \right) \left(\frac{\sum_{t=1}^{T}\sqrt{d}\mathbb{E}[\|\nabla f(\boldsymbol{x}_{t})\|]}{T} \right). \tag{92}$$

Convergence Guarantees for RMSProp and Adam

For
$$\eta \leq C_{5}(1-\beta_{2})$$
, (where $C_{5} > 0$ and will be introduced in Appendix H), $C_{6} = \min\left(\frac{C_{2}\sqrt{\zeta}}{21\alpha_{0}dD_{0}}, \frac{C_{2}^{3}\sqrt{\zeta}}{21\alpha_{0}d_{1}D_{1}^{2}L_{0}^{2}(1-\beta_{1})^{2}d^{2}C_{5}^{2}}, \frac{C_{2}}{21\alpha_{3}d\beta_{1}}, \frac{C_{1}}{84C_{3}C_{5}d(1-\beta_{1})}, \frac{(1-\frac{\beta_{1}}{\sqrt{\beta_{2}}})^{2}}{84C_{1}C_{4}C_{5}^{2}(1-\beta_{1})d\sqrt{D_{0}+\|\boldsymbol{v}_{0}\|}}, \frac{C_{1}^{2}}{784C_{3}^{2}C_{5}^{2}(1-\beta_{1})^{2}dD_{1}}, \frac{(1-\frac{\beta_{1}}{\sqrt{\beta_{2}}})^{4}}{7656C_{1}^{2}C_{4}^{2}C_{5}^{4}dD_{1}}\right), 1-\beta_{2} \leq C_{6}\epsilon^{2}, \text{ and } T \geq \max\left(\frac{126C_{1}\Delta'}{\eta(1-\beta_{1})\epsilon^{2}}, \frac{126C_{3}C_{5}(1-\beta_{1})}{C_{1}}(2d\sqrt{D_{0}+\|\boldsymbol{v}_{0}\|} - \sum_{i=1}^{d}\ln(\boldsymbol{v}_{0,i}))\epsilon^{-2}, \frac{252C_{5}^{2}C_{1}C_{4}(1-\beta_{1})d}{(1-\frac{\beta_{1}}{\sqrt{\beta_{2}}})^{2}}\sqrt{D_{0}+\|\boldsymbol{v}_{0}\|}\epsilon^{-2}\right), \text{ we then have that}$

$$\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}\left[\frac{\|\nabla f(\boldsymbol{x}_{t})\|^{2}}{\sqrt{\beta_{2}}\|\boldsymbol{v}_{t-1}\|+\zeta}\right] \leq \epsilon^{2} + \left(\frac{14\eta C_{3}(1-\beta_{1})\sqrt{dD_{1}}}{C_{1}\sqrt{1-\beta_{2}}} + \frac{42\eta^{2}C_{1}C_{4}(1-\beta_{1})\sqrt{dD_{1}}}{(1-\beta_{2})^{1.5}(1-\frac{\beta_{1}}{\sqrt{\beta_{2}}})^{2}}\right)\left(\frac{\sum_{t=1}^{T}\mathbb{E}[\|\nabla f(\boldsymbol{x}_{t})\|]}{T}\right).$$
(93)

Since we have that $\eta \leq C_5(1-\beta_2)$ and $1-\beta_2 \leq C_6\epsilon^2$, thus we have that

$$\frac{1}{T}\sum_{t=1}^{T} \mathbb{E}\left[\frac{\|\nabla f(\boldsymbol{x}_{t})\|^{2}}{\sqrt{\beta_{2}\|\boldsymbol{v}_{t-1}\|+\zeta}}\right] \leq \epsilon^{2} + \epsilon \left(\frac{\sum_{t=1}^{T} \mathbb{E}[\|\nabla f(\boldsymbol{x}_{t})\|]}{T}\right).$$
(94)

This completes the proof.

H. Formal Version of Theorem 4.1 and Its Proof

For
$$\alpha_0 \geq \frac{21}{2C_2}, \alpha_1 \geq \frac{21\alpha_0}{2C_2}, \alpha_3 \geq \frac{7\beta_1\sqrt{\zeta}}{2C_2}, \alpha_4 \geq \frac{14L_1\sqrt{d}(2-C_1)^2}{C_1(1-\beta_1)}, \text{ define } \Delta' = f(u_1) - f^* + \frac{\eta\alpha_0D_1(1-\beta_1)\|\nabla f(x_1)\|^2}{2C_1C_2\sqrt{\zeta}} + \frac{\eta\alpha_0D_1(1-\beta_1)\|\nabla f(x_1)\|^2}{2C_1C_2\sqrt{\zeta}}, C_3 = \left[\frac{2(1-C_1)^2\sqrt{d}L_0}{C_1^2} + \frac{(2-C_1)\sqrt{d}L_0}{C_1^2} + \sqrt{d}L_0\frac{(1-C_1)^2+1}{C_1^2}\right], C_4 = \left[\frac{\alpha_4(1-\beta_1)^2dL_1((1-C_1)+(1-C_1)^2)}{2C_1^2C_2^2} + \sqrt{d}L_1\frac{2+2(2-C_1)^2}{C_1^2}\frac{\alpha_4(1-\beta_1)^2}{2C_2^2}\right] C_5 = \min\left(\frac{C_1}{112C_3(1-\beta_1)dD_1}, \frac{1-\frac{\beta_1}{\sqrt{\beta_2}}}{168D_1C_1C_4(1-\beta_1)d}\right), C_6 = \min\left(\frac{C_2\sqrt{\zeta}}{21\alpha_0dD_0}, \frac{C_2^3\sqrt{\zeta}}{21\alpha_0\alpha_1D_1^2L_0^2(1-\beta_1)^2d^2C_5^2}, \frac{C_2}{21\alpha_3d\beta_1}, \frac{C_1}{84C_3C_5d(1-\beta_1)}, \frac{(1-\frac{\beta_1}{\sqrt{\beta_2}})^2}{84C_1C_4C_5^2(1-\beta_1)d\sqrt{D_0+||v_0||}}, \frac{C_1^2}{784C_3^2C_5^2(1-\beta_1)^2dD_1}, \frac{(1-\frac{\beta_1}{\sqrt{\beta_2}})^2}{C_1^2(1-\beta_1)^2dD_1}, \frac{(1-\frac{\beta_1}{\sqrt{\beta_2}})^2}{C_1^2(1-\beta_1)^2dD_1}, \frac{C_1^2}{21\alpha_0\sqrt{d}D_1L_1(1-\beta_1)}, \frac{C_1C_2}{\sqrt{d}L_1(1-C_1)(1-\beta_1)}\right), \Lambda_5 = \left(\frac{126C_3C_5(1-\beta_1)}{C_1}(2d\sqrt{D_0+||v_0||} - \sum_{i=1}^d \ln(v_{0,i}))\right) \text{ and } \Lambda_6 = \left(\frac{252C_5^2C_1C_4(1-\beta_1)d}{(1-\frac{\beta_1}{\sqrt{\beta_2}})^2} - C_2c^2\right) \approx \mathcal{O}(c^2), 0 < \beta_1 < \sqrt{\beta_2} < 1$$

Theorem H.1. Let Assumptions 2.1, 2.2 and 2.3 hold. Let $1 - \beta_2 = \min\left(\frac{2C_2}{7\alpha_0 D_1}, C_6 \epsilon^2\right) \sim \mathcal{O}(\epsilon^2), 0 < \beta_1 \le \sqrt{\beta_2} < 1, \eta = \min\left(\Lambda_4\sqrt{1-\beta_2}, C_5(1-\beta_2)\right) \sim \mathcal{O}(\epsilon^2), \text{ and } T \ge \max\left(\frac{126C_1\Delta'}{\eta(1-\beta_1)\epsilon^2}, \Lambda_5\epsilon^{-2}, \Lambda_6\epsilon^{-2}\right) \sim \mathcal{O}(\epsilon^{-4}).$ For small ϵ such that $\epsilon \le \frac{\sqrt{2C_2}}{\sqrt{7\alpha_0 C_6 D_1}}$, we have that

$$\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}[\|\nabla f(x_t)\|] \le \left(2c + \sqrt{2c} + \frac{4\sqrt{dD_1}}{\sqrt{C_6}}\right)\epsilon.$$
(95)

Proof. The proof follows Appendix D. According to (93) in the proof of Corollary 4.2, we have that

$$\frac{1}{T}\sum_{t=1}^{T} \mathbb{E}\left[\frac{\|\nabla f(\boldsymbol{x}_{t})\|^{2}}{\sqrt{\beta_{2}\|\boldsymbol{v}_{t-1}\|+\zeta}}\right] \leq \epsilon^{2} + \left(\frac{14\eta C_{3}(1-\beta_{1})\sqrt{dD_{1}}}{C_{1}\sqrt{1-\beta_{2}}} + \frac{42\eta^{2}C_{1}C_{4}(1-\beta_{1})\sqrt{dD_{1}}}{(1-\beta_{2})^{1.5}(1-\frac{\beta_{1}}{\sqrt[4]{\beta_{2}}})^{2}}\right) \left(\frac{\sum_{t=1}^{T} \mathbb{E}[\|\nabla f(\boldsymbol{x}_{t})\|]}{T}\right).$$
(96)

According to Lemma 3.5, we have that

$$\left(\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}[\sqrt{\beta_2 \|\boldsymbol{v}_{t-1}\| + \zeta}]\right) \leq c + \frac{2\sqrt{dD_1}}{\sqrt{(1-\beta_2)}} \frac{\sum_{t=1}^{T}\mathbb{E}[\|\nabla f(\boldsymbol{x}_t)\|]}{T}.$$
(97)

Define $e = \frac{1}{T} \sum_{t=1}^{T} \mathbb{E}[\|\nabla f(\boldsymbol{x}_t)\|]$. By Hölder's inequality, we have

$$\left(\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}[\|\nabla f(\boldsymbol{x}_{t})\|]\right)^{2} \leq \left(\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}\left[\frac{\|\nabla f(\boldsymbol{x}_{t})\|^{2}}{\sqrt{\beta_{2}\|\boldsymbol{v}_{t-1}\|+\zeta}}\right]\right)\left(\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}[\sqrt{\beta_{2}\|\boldsymbol{v}_{t-1}\|+\zeta}]\right),\tag{98}$$

based on Lemma 3.5 and Corollary 4.2 which can be written as

$$e^{2} \leq \left(\epsilon^{2} + \left(\frac{14\eta C_{3}(1-\beta_{1})\sqrt{dD_{1}}}{C_{1}\sqrt{1-\beta_{2}}} + \frac{42\eta^{2}C_{1}C_{4}(1-\beta_{1})\sqrt{dD_{1}}}{(1-\beta_{2})^{1.5}(1-\frac{\beta_{1}}{\sqrt{\beta_{2}}})^{2}}\right)e\right)\left(c + \frac{2\sqrt{dD_{1}}}{\sqrt{1-\beta_{2}}}e\right)$$
$$\leq c\epsilon^{2} + ce\epsilon + \frac{2\sqrt{dD_{1}}}{\sqrt{C_{6}\epsilon}}e\epsilon^{2} + \frac{e^{2}}{2},$$
(99)

where the second inequality is due to the that that $\left(\frac{14\eta C_3(1-\beta_1)\sqrt{dD_1}}{C_1\sqrt{1-\beta_2}} + \frac{42\eta^2 C_1 C_4(1-\beta_1)\sqrt{dD_1}}{(1-\beta_2)^{1.5}(1-\frac{\beta_1}{\sqrt[4]{\beta_2}})^2}\right) \leq \epsilon \text{ and }$

$$\left(\frac{14\eta C_3(1-\beta_1)\sqrt{dD_1}}{C_1\sqrt{1-\beta_2}} + \frac{42\eta^2 C_1 C_4(1-\beta_1)\sqrt{dD_1}}{(1-\beta_2)^{1.5}(1-\frac{\beta_1}{4\sqrt{\beta_2}})^2}\right)\frac{2\sqrt{dD_1}}{\sqrt{1-\beta_2}} \le \frac{1}{2} \text{ if } \eta \le C_5(1-\beta_2), \ 1-\beta_2 = \min\left(\frac{2C_2}{7\alpha_0 D_1}, C_6\epsilon^2\right) = C_6\epsilon^2$$

and $C_5 = \min\left(\frac{C_1}{112C_3(1-\beta_1)dD_1}, \frac{1-\frac{\beta_1}{4\sqrt{\beta_2}}}{168D_1C_1C_4(1-\beta_1)d}\right).$

Thus

$$\frac{1}{T}\sum_{t=1}^{T}\mathbb{E}[\|\nabla f(x_t)\|] = e \le \left(2c + \sqrt{2c} + \frac{4\sqrt{dD_1}}{\sqrt{C_6}}\right)\epsilon$$

which completes the proof.