

Conditionings to avoid points with various clocks for Lévy processes

Kohki IBA (Graduate School of Science, Osaka University)
Kouji YANO (Graduate School of Science, Osaka University)

Abstract

We discuss conditionings to avoid two points and one-point local time penalizations with conditioning to avoid another point, for which we adopt various clocks. We also give corrections to some of the previous results of Takeda–Yano for one-point local time penalizations.

1 Introduction

A penalization problem is to study the long-time limit of the form

$$\lim_{\tau \rightarrow \infty} \frac{\mathbb{P}_x[F_s \cdot \Gamma_\tau]}{\mathbb{P}_x[\Gamma_\tau]}, \quad (1.1)$$

where $(X = (X_t)_{t \geq 0}, (\mathcal{F}_t)_{t \geq 0}, (\mathbb{P}_x)_{x \in \mathbb{R}})$ is a Markov process, $(\Gamma_t)_{t \geq 0}$ is a non-negative process called a *weight*, $(F_s)_{s \geq 0}$ is a process of test functions adapted to $(\mathcal{F}_s)_{s \geq 0}$, and τ is a net of parametrized random times tending to infinity, called a *clock*.

We follow the notations and adopt the assumptions of Iba–Yano [2] (see Section 2 for the details). In [2], we considered the case where the weight process is given by

$$\Gamma_{a,b,t}^{\lambda_a, \lambda_b} := e^{-\lambda_a L_t^a - \lambda_b L_t^b}. \quad (1.2)$$

In this paper, we consider the case where the weight process is given by

$$\Gamma_{a,b,t}^{\lambda_a, \infty} := \lim_{\lambda_b \rightarrow \infty} \Gamma_{a,b,t}^{\lambda_a, \lambda_b} = e^{-\lambda_a L_t^a} \cdot 1_{\{L_t^b=0\}}, \quad (1.3)$$

$$\Gamma_{a,b,t}^{\infty, \infty} := \lim_{\lambda_a, \lambda_b \rightarrow \infty} \Gamma_{a,b,t}^{\lambda_a, \lambda_b} = \lim_{\lambda_a \rightarrow \infty} \Gamma_{a,b,t}^{\lambda_a, \infty} = 1_{\{L_t^a=L_t^b=0\}}, \quad (1.4)$$

where the limit processes may be formally obtained by the limit as $\lambda_a \rightarrow \infty$ or $\lambda_b \rightarrow \infty$ of those of (1.2), although the penalization limit should be proven independently.

Two-point local time penalization

For $-1 \leq \gamma \leq 1$, we say¹

$$(c, d) \xrightarrow{(\gamma)} \infty \text{ when } c \rightarrow \infty, d \rightarrow \infty, \text{ and } \frac{d-c}{c+d} \rightarrow \gamma. \quad (1.5)$$

¹To describe the penalization limits, our limit $(c, d) \xrightarrow{(\gamma)} \infty$ is more suitable than the limit $(c, d) \xrightarrow{\gamma} \infty$ of the equation (1.9) of Takeda–Yano [3]. See, Section 6.

Here for the random clock $\tau = (\tau_\lambda)$, we adopt one of the following:

1. exponential clock: $\tau = (e_q)$ as $q \rightarrow 0+$, where e_q has the exponential distribution with its parameter $q > 0$ and is independent of X ;
2. hitting time clock: $\tau = (T_c)$ as $c \rightarrow \pm\infty$, where T_c is the first hitting time at c ;
3. two-point hitting time clock: $\tau = (T_c \wedge T_{-d})$ as $(c, d) \xrightarrow{(\gamma)} \infty$;
4. inverse local time clock: $\tau = (\eta_u^c)$ as $c \rightarrow \pm\infty$, where $\eta^c = (\eta_u^c)_{u \geq 0}$ is an inverse local time.

Then, Iba–Yano [2] showed the following:

Theorem 1.1 (Iba–Yano [2]). *For distinct points $a, b \in \mathbb{R}$, for constants $\lambda_a, \lambda_b > 0$, and for a constant $-1 \leq \gamma \leq 1$, there exists a positive function $\varphi_{a,b}^{(\gamma), \lambda_a, \lambda_b}(x)$ such that the process*

$$\left(M_{a,b,s}^{(\gamma), \lambda_a, \lambda_b} := \varphi_{a,b}^{(\gamma), \lambda_a, \lambda_b}(X_s) e^{-\lambda_a L_t^a - \lambda_b L_t^b} \right)_{s \geq 0} \quad (1.6)$$

is a martingale, and the following assertions hold:

1. exponential clock: $\lim_{q \rightarrow 0+} r_q(0) \mathbb{P}_x \left[F_s \cdot \Gamma_{a,b,e_q}^{\lambda_a, \lambda_b} \right] = \mathbb{P}_x \left[F_s \cdot M_{a,b,s}^{(0), \lambda_a, \lambda_b} \right]$,
2. hitting time clock: $\lim_{c \rightarrow \pm\infty} h^B(c) \mathbb{P}_x \left[F_s \cdot \Gamma_{a,b,T_c}^{\lambda_a, \lambda_b} \right] = \mathbb{P}_x \left[F_s \cdot M_{a,b,s}^{(\pm 1), \lambda_a, \lambda_b} \right]$,
3. two-point hitting time clock: $\lim_{(c,d) \xrightarrow{(\gamma)} \infty} h^C(c, -d) \mathbb{P}_x \left[F_s \cdot \Gamma_{a,b,T_c \wedge T_{-d}}^{\lambda_a, \lambda_b} \right] = \mathbb{P}_x \left[F_s \cdot M_{a,b,s}^{(\gamma), \lambda_a, \lambda_b} \right]$,
4. inverse local time clock: $\lim_{c \rightarrow \pm\infty} h^B(c) \mathbb{P}_x \left[F_s \cdot \Gamma_{a,b,\eta_u^c}^{\lambda_a, \lambda_b} \right] = \mathbb{P}_x \left[F_s \cdot M_{a,b,s}^{(\pm 1), \lambda_a, \lambda_b} \right]$,

where r_q is the q -resolvent density (see (2.5)), h^B is defined by (2.16), and h^C is defined by (2.17).

We write for $-1 \leq \gamma \leq 1$,

$$h^{(\gamma)}(x) := h(x) + \frac{\gamma}{m^2} x,$$

where $m^2 = \mathbb{P}_0[X_1^2]$ and the function h is a renormalized zero resolvent (see Proposition 2.1). Then, the function $\varphi_{a,b}^{(\gamma), \lambda_a, \lambda_b}$ is given explicitly as follows:

$$\begin{aligned} \varphi_{a,b}^{(\gamma), \lambda_a, \lambda_b}(x) &= h^{(\gamma)}(x - a) - \mathbb{P}_x(T_b < T_a) h^{(\gamma)}(b - a) \\ &\quad + \mathbb{P}_x(T_a < T_b) \cdot \frac{h^{(\gamma)}(a - b)}{1 + \lambda_a h^B(a - b)} \end{aligned}$$

$$\begin{aligned}
& + \mathbb{P}_x(T_a < T_b) \cdot \frac{1}{1 + \lambda_a h^B(b-a)} \cdot \frac{1 + \lambda_a h^{(\gamma)}(b-a)}{\lambda_a + \lambda_b + \lambda_a \lambda_b h^B(a-b)} \\
& + \mathbb{P}_x(T_b < T_a) \cdot \frac{h^{(\gamma)}(b-a)}{1 + \lambda_b h^B(a-b)} \\
& + \mathbb{P}_x(T_b < T_a) \cdot \frac{1}{1 + \lambda_b h^B(b-a)} \cdot \frac{1 + \lambda_b h^{(\gamma)}(a-b)}{\lambda_a + \lambda_b + \lambda_a \lambda_b h^B(a-b)}. \tag{1.7}
\end{aligned}$$

Note that $\varphi_{a,b}^{(\gamma),\lambda_a,\lambda_b}(x)$ is symmetric with respect to a and b , i.e., for $x \in \mathbb{R}$,

$$\varphi_{a,b}^{(\gamma),\lambda_a,\lambda_b}(x) = \varphi_{b,a}^{(\gamma),\lambda_a,\lambda_b}(x). \tag{1.8}$$

When $m^2 = \infty$, we have

$$\varphi_{a,b}^{(\gamma),\lambda_a,\lambda_b}(x) \equiv \varphi_{a,b}^{(0),\lambda_a,\lambda_b}(x). \tag{1.9}$$

Therefore, when $m^2 = \infty$, we obtain

$$M_{a,b,t}^{(\gamma),\lambda_a,\lambda_b} = M_{a,b,t}^{(0),\lambda_a,\lambda_b} \tag{1.10}$$

for any $-1 \leq \gamma \leq 1$.

Main theorems about conditionings

We define

$$\varphi_{a,b}^{(\gamma),\lambda_a,\infty}(x) := h^{(\gamma)}(x-a) - \mathbb{P}_x(T_b < T_a)h^{(\gamma)}(b-a) + \mathbb{P}_x(T_a < T_b) \cdot \frac{h^{(\gamma)}(a-b)}{1 + \lambda_a h^B(a-b)}, \tag{1.11}$$

$$\varphi_{a,b}^{(\gamma),\infty,\infty}(x) := h^{(\gamma)}(x-a) - \mathbb{P}_x(T_b < T_a)h^{(\gamma)}(b-a). \tag{1.12}$$

When $m^2 = \infty$, we have

$$\varphi_{a,b}^{(\gamma),\lambda_a,\infty}(x) \equiv \varphi_{a,b}^{(0),\lambda_a,\infty}(x), \tag{1.13}$$

$$\varphi_{a,b}^{(\gamma),\infty,\infty}(x) \equiv \varphi_{a,b}^{(0),\infty,\infty}(x), \tag{1.14}$$

Note that $\varphi_{a,b}^{(\gamma),\infty,\infty}$ and $\varphi_{a,b}^{(\gamma),\lambda_a,\infty}$ are formally obtained by the limit as $\lambda_a \rightarrow \infty$ or $\lambda_b \rightarrow \infty$ of those of (1.7), i.e.,

$$\varphi_{a,b}^{(\gamma),\lambda_a,\infty}(x) = \lim_{\lambda_b \rightarrow \infty} \varphi_{a,b}^{(\gamma),\lambda_a,\lambda_b}(x), \tag{1.15}$$

$$\varphi_{a,b}^{(\gamma),\infty,\infty}(x) = \lim_{\lambda_a, \lambda_b \rightarrow \infty} \varphi_{a,b}^{(\gamma),\lambda_a,\lambda_b}(x) = \lim_{\lambda_a \rightarrow \infty} \varphi_{a,b}^{(\gamma),\lambda_a,\infty}(x). \tag{1.16}$$

Under the same assumption, our main theorems are as follows (see Sections 3 and 4 for the details):

Theorem 1.2. For distinct points $a, b \in \mathbb{R}$, for a constant $\lambda_a > 0$, and for a constant $-1 \leq \gamma \leq 1$, the process

$$\left(M_{a,b,s}^{(\gamma),\lambda_a,\infty} := \varphi_{a,b}^{(\gamma),\lambda_a,\infty}(X_s) e^{-\lambda_a L_s^a} 1_{\{L_s^b=0\}} \right)_{s \geq 0} \quad (1.17)$$

is a martingale, and the following assertions hold:

1. exponential clock: $\lim_{q \rightarrow 0^+} r_q(0) \mathbb{P}_x \left[F_s \cdot \Gamma_{a,b,e_q}^{\lambda_a,\infty} \right] = \mathbb{P}_x \left[F_s \cdot M_{a,b,s}^{(0),\lambda_a,\infty} \right],$
2. hitting time clock: $\lim_{c \rightarrow \pm\infty} h^B(c) \mathbb{P}_x \left[F_s \cdot \Gamma_{a,b,T_c}^{\lambda_a,\infty} \right] = \mathbb{P}_x \left[F_s \cdot M_{a,b,s}^{(\pm 1),\lambda_a,\infty} \right],$
3. two-point hitting time clock: $\lim_{(c,d) \xrightarrow{(\gamma)} \infty} h^C(c, -d) \mathbb{P}_x \left[F_s \cdot \Gamma_{a,b,T_c \wedge T_{-d}}^{\lambda_a,\infty} \right] = \mathbb{P}_x \left[F_s \cdot M_{a,b,s}^{(\gamma),\lambda_a,\infty} \right],$
4. inverse local time clock: $\lim_{c \rightarrow \pm\infty} h^B(c) \mathbb{P}_x \left[F_s \cdot \Gamma_{a,b,\eta_u^c}^{\lambda_a,\infty} \right] = \mathbb{P}_x \left[F_s \cdot M_{a,b,s}^{(\pm 1),\lambda_a,\infty} \right].$

Theorem 1.3. For distinct points $a, b \in \mathbb{R}$ and for a constant $-1 \leq \gamma \leq 1$, the process

$$\left(M_{a,b,s}^{(\gamma),\infty,\infty} := \varphi_{a,b}^{(\gamma),\infty,\infty}(X_s) 1_{\{L_s^a=L_s^b=0\}} \right)_{s \geq 0} \quad (1.18)$$

is a martingale, and the following assertions hold:

1. exponential clock: $\lim_{q \rightarrow 0^+} r_q(0) \mathbb{P}_x \left[F_s \cdot \Gamma_{a,b,e_q}^{\infty,\infty} \right] = \mathbb{P}_x \left[F_s \cdot M_{a,b,s}^{(0),\infty,\infty} \right],$
2. hitting time clock: $\lim_{c \rightarrow \pm\infty} h^B(c) \mathbb{P}_x \left[F_s \cdot \Gamma_{a,b,T_c}^{\infty,\infty} \right] = \mathbb{P}_x \left[F_s \cdot M_{a,b,s}^{(\pm 1),\infty,\infty} \right],$
3. two-point hitting time clock: $\lim_{(c,d) \xrightarrow{(\gamma)} \infty} h^C(c, -d) \mathbb{P}_x \left[F_s \cdot \Gamma_{a,b,T_c \wedge T_{-d}}^{\infty,\infty} \right] = \mathbb{P}_x \left[F_s \cdot M_{a,b,s}^{(\gamma),\infty,\infty} \right],$
4. inverse local time clock: $\lim_{c \rightarrow \pm\infty} h^B(c) \mathbb{P}_x \left[F_s \cdot \Gamma_{a,b,\eta_u^c}^{\infty,\infty} \right] = \mathbb{P}_x \left[F_s \cdot M_{a,b,s}^{(\pm 1),\infty,\infty} \right].$

Organization

This paper is organized as follows. In Section 2, we prepare some general results of Lévy processes. In Sections 3 and 4, we discuss the penalization results with $\Gamma_{a,b,t}^{\infty,\infty}$ and $\Gamma_{a,b,t}^{\lambda_a,\infty}$, respectively. In Section 5, we discuss the penalization results with inverse local time clock. In Section 6, we give some corrections to some results of Takeda–Yano [3].

2 Preliminaries

2.1 Lévy process and resolvent density

Let (X, \mathbb{P}_x) be the canonical representation of a real valued Lévy process starting from $x \in \mathbb{R}$ on the càdlàg path space. For $t > 0$, we denote by $\mathcal{F}_t^X = \sigma(X_s, 0 \leq s \leq t)$ the natural filtration of X and write $\mathcal{F}_t = \bigcap_{s>t} \mathcal{F}_s^X$. For $a \in \mathbb{R}$, let T_a be the hitting time of $\{a\}$ for X , i.e.,

$$T_a = \inf\{t > 0 : X_t = a\}. \quad (2.1)$$

For $\lambda \in \mathbb{R}$, we denote by $\Psi(\lambda)$ the characteristic exponent of X , which satisfies

$$\mathbb{P}_0 [e^{i\lambda X_t}] = e^{-t\Psi(\lambda)} \quad (2.2)$$

for $t \geq 0$. Moreover, by Lévy–Khinchin formula, it is denoted by

$$\Psi(\lambda) = i\nu\lambda + \frac{1}{2}\sigma^2\lambda^2 + \int_{\mathbb{R}} (1 - e^{i\lambda x} + i\lambda x 1_{\{|x|<1\}}) \nu(dx), \quad (2.3)$$

where $\nu \in \mathbb{R}$, $\sigma \geq 0$, and ν is a measure on \mathbb{R} , called a *Lévy measure*, with $\nu(\{0\}) = 0$ and $\int_{\mathbb{R}} (x^2 \wedge 1) \nu(dx) < \infty$.

Throughout this paper, we always assume (X, \mathbb{P}_0) is recurrent, i.e.,

$$\mathbb{P}_0 \left[\int_0^\infty 1_{\{|X_t - a| < \varepsilon\}} dt \right] = \infty \quad (2.4)$$

for all $a \in \mathbb{R}$ and $\varepsilon > 0$, and always assume the condition

$$\mathbf{(A)} \quad \int_0^\infty \left| \frac{1}{q + \Psi(\lambda)} \right| d\lambda < \infty \quad \text{for each } q > 0.$$

It is known that X has a bounded continuous resolvent density r_q :

$$\int_{\mathbb{R}} f(x) r_q(x) dx = \mathbb{P}_0 \left[\int_0^\infty e^{-qt} f(X_t) dt \right] \quad (2.5)$$

holds for $q > 0$ and non-negative measurable functions f . See, e.g., Theorems II.16 and II.19 of [1]. Moreover, there exists an equality that connects the hitting time of 0 and the resolvent density:

$$\mathbb{P}_x [e^{-qT_0}] = \frac{r_q(-x)}{r_q(0)} \quad (2.6)$$

for $q > 0$ and $x \in \mathbb{R}$. See, e.g., Corollary II.18 of [1].

2.2 Local time and excursion

We denote by \mathcal{D} the set of càdlàg paths $e : [0, \infty) \rightarrow \mathbb{R} \cup \{\Delta\}$ such that

$$\begin{cases} e(t) \in \mathbb{R} \setminus \{0\} & \text{for } 0 < t < \zeta(e), \\ e(t) = \Delta & \text{for } t \geq \zeta(e), \end{cases} \quad (2.7)$$

where the point Δ is an isolated point and ζ is the excursion length, i.e.,

$$\zeta = \zeta(e) := \inf\{t > 0 : e(t) = \Delta\}. \quad (2.8)$$

Let Σ denote the σ -algebra on \mathcal{D} generated by cylinder sets.

Assume the condition **(A)** holds. Then, we can define a local time at $a \in \mathbb{R}$, which we denote by $L^a = (L_t^a)_{t \geq 0}$. It is defined by

$$L_t^a := \lim_{\varepsilon \rightarrow 0^+} \frac{1}{2\varepsilon} \int_0^t 1_{\{|X_s - a| < \varepsilon\}} ds. \quad (2.9)$$

It is known that L^a is continuous in t and satisfies

$$\mathbb{P}_x \left[\int_0^\infty e^{-qt} dL_t^a \right] = r_q(a - x) \quad (2.10)$$

for $q > 0$ and $x \in \mathbb{R}$. See, e.g., Section V of [1]. In particular, from this expression, $r_q(x)$ is non-decreasing as $q \rightarrow 0^+$.

Let $\eta^a = (\eta_l^a)_{l \geq 0}$ be an inverse local time, i.e.,

$$\eta_l^a := \inf\{t > 0 : L_t^a > l\}. \quad (2.11)$$

It is known that the process (η^a, \mathbb{P}_a) is a possibly killed subordinator which has the Laplace exponent

$$\mathbb{P}_a \left[e^{-q\eta_l^a} \right] = e^{-\frac{l}{r_q(0)}} \quad (2.12)$$

for $l > 0$ and $q > 0$. See, e.g., Proposition V.4 of [1].

We denote ϵ_l^a for an excursion away from $a \in \mathbb{R}$ which starts at local time $l \geq 0$, i.e.,

$$\epsilon_l^a(t) := \begin{cases} X_{t+\eta_{l-}^a} & \text{for } 0 \leq t < \eta_l^a - \eta_{l-}^a, \\ \Delta & \text{for } t \geq \eta_l^a - \eta_{l-}^a. \end{cases} \quad (2.13)$$

Then, $(\epsilon_l^a)_{l \geq 0}$ is a Poisson point process, and we write n^a for the characteristic measure of ϵ^a . It is known that $(\mathcal{D}, \Sigma, n^a)$ is a σ -finite measure space. See, e.g., Section IV of [1]. For $B \in \mathcal{B}(0, \infty) \otimes \Sigma$, we define

$$N^a(B) := \#\{(l, e) \in B : \epsilon_l^a = e\}. \quad (2.14)$$

Then, N^a is a Poisson random measure with its intensity measure $ds \times n^a(de)$. It is known that the subordinator η^0 has no drift and its Lévy measure is $n^0(T_0 \in dx)$.

2.3 The renormalized zero resolvent

We define

$$h_q(x) := r_q(0) - r_q(-x) \quad (2.15)$$

for $q > 0$ and $x \in \mathbb{R}$. It is clear that $h_q(0) = 0$, and by (2.6), we have $h_q(x) \geq 0$. The following theorem plays a key role in our penalization results.

Proposition 2.1 (Theorem 1.1 of [3]). *If the condition (A) is satisfied, then for any $x \in \mathbb{R}$, $h(x) := \lim_{q \rightarrow 0^+} h_q(x)$ exists and is finite.*

We call h the *renormalized zero resolvent*. Moreover, we introduce the functions h_q^B and h^B .

Proposition 2.2 (Lemma 3.5 of [3]). *For $a \in \mathbb{R}$, it holds that*

$$h^B(a) := \lim_{q \rightarrow 0^+} h_q^B(a) = \mathbb{P}_0[L_{T_a}] = h(a) + h(-a). \quad (2.16)$$

Proposition 2.3 (Lemma 6.1 of [3]²). *For distinct points $a, b \in \mathbb{R}$, it holds that*

$$\begin{aligned} h^C(a, b) &:= \mathbb{P}_0[L_{T_a \wedge T_b}^0] \\ &= \frac{1}{h^B(a-b)} \left\{ \begin{aligned} &(h(b) + h(-a))h(a-b) + (h(a) + h(-b))h(b-a) \\ &+ (h(a) - h(b))(h(-b) - h(-a)) - h(a-b)h(b-a) \end{aligned} \right\}. \end{aligned} \quad (2.17)$$

3 Conditionings to avoid two points

We define

$$N_{a,b,t}^{q,\lambda_a,\lambda_b} := r_q(0) \mathbb{P}_x \left[\Gamma_{a,b,e_q}^{\lambda_a,\lambda_b}; t < e_q \mid \mathcal{F}_t \right], \quad (3.1)$$

$$N_{a,b,t}^{q,\infty,\infty} := r_q(0) \mathbb{P}_x (t < e_q < T_a \wedge T_b \mid \mathcal{F}_t), \quad (3.2)$$

$$M_{a,b,t}^{q,\infty,\infty} := r_q(0) \mathbb{P}_x (e_q < T_a \wedge T_b \mid \mathcal{F}_t) \quad (3.3)$$

for $q > 0$.

Theorem 3.1. *Let $x \in \mathbb{R}$. Then, $(M_{a,b,t}^{(0),\infty,\infty}, t \geq 0)$ is a non-negative $((\mathcal{F}_t), \mathbb{P}_x)$ -martingale, and it holds that*

$$\lim_{q \rightarrow 0^+} N_{a,b,t}^{q,\infty,\infty} = \lim_{q \rightarrow 0^+} M_{a,b,t}^{q,\infty,\infty} = M_{a,b,t}^{(0),\infty,\infty} \text{ a.s. and in } L^1(\mathbb{P}_x). \quad (3.4)$$

²In Takeda–Yano [3], there is an error in the assertion of Lemma 6.1. Proposition 2.3 corrects that error. See, Section 6.

Consequently, if $M_{a,b,0}^{(0),\infty,\infty} > 0$ under \mathbb{P}_x , it holds that

$$\lim_{q \rightarrow 0+} \mathbb{P}_x[F_t | \mathbf{e}_q < T_a \wedge T_b] = \mathbb{P}_x \left[F_t \cdot \frac{M_{a,b,t}^{(0),\infty,\infty}}{M_{a,b,0}^{(0),\infty,\infty}} \right] \quad (3.5)$$

for all bounded \mathcal{F}_t -measurable functionals F_t .

Proof. Note that

$$\lim_{\lambda_a, \lambda_b \rightarrow \infty} N_{a,b,t}^{q,\lambda_a,\lambda_b} = N_{a,b,t}^{q,\infty,\infty} \text{ a.s.} \quad (3.6)$$

We have by the lack of memory property of an exponential distribution and by the Markov property,

$$\begin{aligned} \lim_{\lambda_a, \lambda_b \rightarrow \infty} N_{a,b,t}^{q,\lambda_a,\lambda_b} &= \lim_{\lambda_a, \lambda_b \rightarrow \infty} r_q(0) e^{-qt} e^{-\lambda_a L_t^a - \lambda_b L_t^b} \mathbb{P}_{X_t} \left[e^{-\lambda_a L_{\mathbf{e}_q}^a - \lambda_b L_{\mathbf{e}_q}^b} \right] \\ &= e^{-qt} r_q(0) \mathbb{P}_{X_t} \left[\int_0^{T_a \wedge T_b} q e^{-qs} ds \right] 1_{\{t < T_a \wedge T_b\}} \text{ a.s.} \end{aligned} \quad (3.7)$$

Therefore, we obtain by the equation (3.24) of [2],

$$\lim_{q \rightarrow 0+} N_{a,b,t}^{q,\infty,\infty} = \lim_{q \rightarrow 0+} e^{-qt} r_q(0) \mathbb{P}_{X_t} \left[\int_0^{T_a \wedge T_b} q e^{-qs} ds \right] 1_{\{t < T_a \wedge T_b\}} = M_{a,b,t}^{(0),\infty,\infty} \text{ a.s.} \quad (3.8)$$

We have

$$\begin{aligned} M_{a,b,t}^{q,\infty,\infty} - N_{a,b,t}^{q,\infty,\infty} &= r_q(0) \mathbb{P}_x(\mathbf{e}_q < T_a \wedge T_b, \mathbf{e}_q \leq t | \mathcal{F}_t) \\ &= r_q(0) 1_{\{\mathbf{e}_q < T_a \wedge T_b\}} 1_{\{\mathbf{e}_q \leq t\}} \rightarrow 0 \text{ a.s.} \end{aligned} \quad (3.9)$$

as $q \rightarrow 0+$. Therefore, we obtain

$$\lim_{q \rightarrow 0+} M_{a,b,t}^{q,\infty,\infty} = M_{a,b,t}^{(0),\infty,\infty} \text{ a.s.} \quad (3.10)$$

Finally, using Theorem 15.2 of [4], the $L^1(\mathbb{P}_x)$ convergence holds. \square

We define

$$N_{a,b,t}^{c,\lambda_a,\lambda_b} := h^B(c) \mathbb{P}_x \left[\Gamma_{a,b,T_c}^{\lambda_a,\lambda_b}; t < T_c | \mathcal{F}_t \right], \quad (3.11)$$

$$N_{a,b,t}^{c,\infty,\infty} := h^B(c) \mathbb{P}_x(t < T_c < T_a \wedge T_b | \mathcal{F}_t), \quad (3.12)$$

$$M_{a,b,t}^{c,\infty,\infty} := h^B(c) \mathbb{P}_x(T_c < T_a \wedge T_b | \mathcal{F}_t) \quad (3.13)$$

for distinct points $a, b, c \in \mathbb{R}$.

Theorem 3.2. *Let $x \in \mathbb{R}$. Then, $(M_{a,b,t}^{(\pm 1),\infty,\infty}, t \geq 0)$ is a non-negative $((\mathcal{F}_t), \mathbb{P}_x)$ -martingale, and it holds that*

$$\lim_{c \rightarrow \pm\infty} N_{a,b,t}^{c,\infty,\infty} = \lim_{c \rightarrow \pm\infty} M_{a,b,t}^{c,\infty,\infty} = M_{a,b,t}^{(\pm 1),\infty,\infty} \text{ a.s. and in } L^1(\mathbb{P}_x). \quad (3.14)$$

Consequently, if $M_{a,b,0}^{(\pm 1),\infty,\infty} > 0$ under \mathbb{P}_x , it holds that

$$\lim_{c \rightarrow \pm\infty} \mathbb{P}_x[F_t | T_c < T_a \wedge T_b] = \mathbb{P}_x \left[F_t \cdot \frac{M_{a,b,t}^{(\pm 1),\infty,\infty}}{M_{a,b,0}^{(\pm 1),\infty,\infty}} \right] \quad (3.15)$$

for all bounded \mathcal{F}_t -measurable functionals F_t .

Proof. Note that

$$\lim_{\lambda_a, \lambda_b \rightarrow \infty} N_{a,b,t}^{c,\lambda_a,\lambda_b} = N_{a,b,t}^{c,\infty,\infty} \text{ a.s.} \quad (3.16)$$

By the strong Markov property, we have

$$\begin{aligned} \lim_{\lambda_a, \lambda_b \rightarrow \infty} N_{a,b,t}^{c,\lambda_a,\lambda_b} &= \lim_{\lambda_a, \lambda_b \rightarrow \infty} 1_{\{t < T_c\}} h^B(c) e^{-\lambda_a L_t^a - \lambda_b L_t^b} \mathbb{P}_{X_t} \left[e^{-\lambda_a L_{T_c}^a - \lambda_b L_{T_c}^b} \right] \\ &= 1_{\{t < T_c\}} h^B(c) 1_{\{t < T_a \wedge T_b\}} \mathbb{P}_{X_t}(T_c < T_a \wedge T_b). \end{aligned} \quad (3.17)$$

Therefore, we obtain by the equation (4.16) of [2],

$$\lim_{c \rightarrow \pm\infty} N_{a,b,t}^{c,\infty,\infty} = \lim_{c \rightarrow \pm\infty} 1_{\{t < T_c\}} h^B(c) 1_{\{t < T_a \wedge T_b\}} \mathbb{P}_{X_t}(T_c < T_a \wedge T_b) = M_{a,b,t}^{(\pm 1),\infty,\infty} \text{ a.s.} \quad (3.18)$$

The rest of the proof is the same as in Theorem 3.1, so we omit it. \square

We define

$$N_{a,b,t}^{c,d,\lambda_a,\lambda_b} := h^C(c, -d) \mathbb{P}_x \left[\Gamma_{a,b,T_c \wedge T_{-d}}^{\lambda_a,\lambda_b}; t < T_c \wedge T_{-d} \mid \mathcal{F}_t \right], \quad (3.19)$$

$$N_{a,b,t}^{c,d,\infty,\infty} := h^C(c, -d) \mathbb{P}_x(t < T_c \wedge T_{-d} < T_a \wedge T_b \mid \mathcal{F}_t), \quad (3.20)$$

$$M_{a,b,t}^{c,d,\infty,\infty} := h^C(c, -d) \mathbb{P}_x(T_c \wedge T_{-d} < T_a \wedge T_b \mid \mathcal{F}_t) \quad (3.21)$$

for $c, d > 0$.

Theorem 3.3. *Let $x \in \mathbb{R}$ and $-1 \leq \gamma \leq 1$. Then, $(M_{a,b,t}^{(\gamma),\infty,\infty}, t \geq 0)$ is a non-negative $((\mathcal{F}_t), \mathbb{P}_x)$ -martingale, and it holds that*

$$\lim_{(c,d) \xrightarrow{(\gamma)} \infty} N_{a,b,t}^{c,d,\infty,\infty} = \lim_{(c,d) \xrightarrow{(\gamma)} \infty} M_{a,b,t}^{c,d,\infty,\infty} = M_{a,b,t}^{(\gamma),\infty,\infty} \text{ a.s. and in } L^1(\mathbb{P}_x). \quad (3.22)$$

Consequently, if $M_{a,b,0}^{(\gamma),\infty,\infty} > 0$ under \mathbb{P}_x , it holds that

$$\lim_{(c,d) \xrightarrow{(\gamma)} \infty} \mathbb{P}_x[F_t | T_c \wedge T_{-d} < T_a \wedge T_b] = \mathbb{P}_x \left[F_t \cdot \frac{M_{a,b,t}^{(\gamma),\infty,\infty}}{M_{a,b,0}^{(\gamma),\infty,\infty}} \right] \quad (3.23)$$

for all bounded \mathcal{F}_t -measurable functionals F_t .

Proof. Note that

$$\lim_{\lambda_a, \lambda_b \rightarrow \infty} N_{a,b,t}^{c,d,\lambda_a,\lambda_b} = N_{a,b,t}^{c,d,\infty,\infty} \text{ a.s.} \quad (3.24)$$

By the strong Markov property, we have

$$\begin{aligned} \lim_{\lambda_a, \lambda_b \rightarrow \infty} N_{a,b,t}^{c,d,\lambda_a,\lambda_b} &= \lim_{\lambda_a, \lambda_b \rightarrow \infty} \mathbf{1}_{\{t < T_c \wedge T_{-d}\}} e^{-\lambda_a L_t^a - \lambda_b L_t^b} h^C(c, -d) \mathbb{P}_{X_t} \left[e^{-\lambda_a L_{T_c \wedge T_{-d}}^a - \lambda_b L_{T_c \wedge T_{-d}}^b} \right] \\ &= \mathbf{1}_{\{t < T_c \wedge T_{-d}\}} \mathbf{1}_{\{t < T_a \wedge T_b\}} h^C(c, -d) \mathbb{P}_{X_t}(T_c \wedge T_{-d} < T_a \wedge T_b). \end{aligned} \quad (3.25)$$

Therefore, we obtain by the equation (5.6) of [2],

$$\begin{aligned} \lim_{(c,d) \xrightarrow{(\gamma)} \infty} N_{a,b,t}^{c,d,\infty,\infty} &= \lim_{(c,d) \xrightarrow{(\gamma)} \infty} \mathbf{1}_{\{t < T_c \wedge T_{-d}\}} \mathbf{1}_{\{t < T_a \wedge T_b\}} h^C(c, -d) \mathbb{P}_{X_t}(T_c \wedge T_{-d} < T_a \wedge T_b) \\ &= M_{a,b,t}^{(\gamma),\infty,\infty} \text{ a.s.} \end{aligned} \quad (3.26)$$

The rest of the proof is the same as in Theorem 3.1, so we omit it. \square

We define

$$N_{a,b,t}^{c,u,\lambda_a,\lambda_b} := h^B(c) \mathbb{P}_x \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a,\lambda_b}; t < \eta_u^c \mid \mathcal{F}_t \right], \quad (3.27)$$

$$N_{a,b,t}^{c,u,\infty,\infty} := h^B(c) \mathbb{P}_x(t < \eta_u^c < T_a \wedge T_b \mid \mathcal{F}_t), \quad (3.28)$$

$$M_{a,b,t}^{c,u,\infty,\infty} := h^B(c) \mathbb{P}_x(\eta_u^c < T_a \wedge T_b \mid \mathcal{F}_t) \quad (3.29)$$

for $c \in \mathbb{R}$ and $u > 0$.

Theorem 3.4. *Let $x \in \mathbb{R}$. Then, it holds that*

$$\lim_{c \rightarrow \pm\infty} N_{a,b,t}^{c,u,\infty,\infty} = \lim_{c \rightarrow \pm\infty} M_{a,b,t}^{c,u,\infty,\infty} = M_{a,b,t}^{(\pm 1),\infty,\infty} \text{ a.s. and in } L^1(\mathbb{P}_x). \quad (3.30)$$

Consequently, if $M_{a,b,0}^{(\pm 1),\infty,\infty} > 0$ under \mathbb{P}_x , it holds that

$$\lim_{c \rightarrow \pm\infty} \mathbb{P}_x[F_t \mid \eta_u^c < T_a \wedge T_b] = \mathbb{P}_x \left[F_t \cdot \frac{M_{a,b,t}^{(\pm 1),\infty,\infty}}{M_{a,b,0}^{(\pm 1),\infty,\infty}} \right] \quad (3.31)$$

for all bounded \mathcal{F}_t -measurable functionals F_t .

Proof. Note that

$$\lim_{\lambda_a, \lambda_b \rightarrow \infty} N_{a,b,t}^{c,u,\lambda_a,\lambda_b} = N_{a,b,t}^{c,u,\infty,\infty} \text{ a.s.} \quad (3.32)$$

By the strong Markov property, we have

$$\begin{aligned} \lim_{\lambda_a, \lambda_b \rightarrow \infty} N_{a,b,t}^{c,u,\lambda_a,\lambda_b} &= \lim_{\lambda_a, \lambda_b \rightarrow \infty} \mathbf{1}_{\{t < T_c \wedge T_{-d}\}} e^{-\lambda_a L_t^a - \lambda_b L_t^b} h^B(c) \mathbb{P}_{X_t} \left[e^{-\lambda_a L_{\eta_u^c}^a - \lambda_b L_{\eta_u^c}^b} \right] \\ &= \mathbf{1}_{\{t < \eta_u^c\}} \mathbf{1}_{\{t < T_a \wedge T_b\}} h^B(c) \mathbb{P}_{X_t}(\eta_u^c < T_a \wedge T_b). \end{aligned} \quad (3.33)$$

Therefore, we obtain by the equation (6.8) and (6.17) of [2],

$$\begin{aligned} \lim_{c \rightarrow \pm\infty} N_{a,b,t}^{c,u,\infty,\infty} &= \lim_{c \rightarrow \pm\infty} \mathbf{1}_{\{t < \eta_u^c\}} \mathbf{1}_{\{t < T_a \wedge T_b\}} h^B(c) \mathbb{P}_{X_t}(\eta_u^c < T_a \wedge T_b) \\ &= M_{a,b,t}^{(\pm 1),\infty,\infty} \text{ a.s.} \end{aligned} \quad (3.34)$$

The rest of the proof is the same as in Theorem 3.1, so we omit it. \square

4 One-point local time penalization with conditioning to avoid another point

We define

$$N_{a,b,t}^{q,\lambda_a,\infty} := r_q(0) \mathbb{P}_x \left[e^{-\lambda_a L_{e_q}^a}; t < e_q < T_b \middle| \mathcal{F}_t \right], \quad (4.1)$$

$$M_{a,b,t}^{q,\lambda_a,\infty} := r_q(0) \mathbb{P}_x \left[e^{-\lambda_a L_{e_q}^a}; e_q < T_b \middle| \mathcal{F}_t \right] \quad (4.2)$$

for $q > 0$.

Theorem 4.1. *Let $x \in \mathbb{R}$. Then, $(M_{a,b,t}^{(0),\lambda_a,\infty}, t \geq 0)$ is a non-negative $((\mathcal{F}_t), \mathbb{P}_x)$ -martingale, and it holds that*

$$\lim_{q \rightarrow 0^+} N_{a,b,t}^{q,\lambda_a,\infty} = \lim_{q \rightarrow 0^+} M_{a,b,t}^{q,\lambda_a,\infty} = M_{a,b,t}^{(0),\lambda_a,\infty} \text{ a.s. and in } L^1(\mathbb{P}_x). \quad (4.3)$$

Consequently, if $M_{a,b,0}^{(0),\lambda_a,\infty} > 0$ under \mathbb{P}_x , it holds that

$$\lim_{q \rightarrow 0^+} \frac{\mathbb{P}_x[F_t \cdot e^{-\lambda_a L_{e_q}^a}; e_q < T_b]}{\mathbb{P}_x[e^{-\lambda_a L_{e_q}^a}; e_q < T_b]} = \mathbb{P}_x \left[F_t \cdot \frac{M_{a,b,t}^{(0),\lambda_a,\infty}}{M_{a,b,0}^{(0),\lambda_a,\infty}} \right] \quad (4.4)$$

for all bounded \mathcal{F}_t -measurable functionals F_t .

The proof is almost the same as that of Theorem 3.1, and so we omit it.

We define

$$N_{a,b,t}^{c,\lambda_a,\infty} := h^B(c) \mathbb{P}_x \left[e^{-\lambda_a L_{T_c}^a}; t < T_c < T_b \middle| \mathcal{F}_t \right], \quad (4.5)$$

$$M_{a,b,t}^{c,\lambda_a,\infty} := h^B(c) \mathbb{P}_x \left[e^{-\lambda_a L_{T_c}^a}; T_c < T_b \middle| \mathcal{F}_t \right] \quad (4.6)$$

for distinct points $a, b, c \in \mathbb{R}$.

Theorem 4.2. *Let $x \in \mathbb{R}$. Then, $(M_{a,b,t}^{(\pm 1),\lambda_a,\infty}, t \geq 0)$ is a non-negative $((\mathcal{F}_t), \mathbb{P}_x)$ -martingale, and it holds that*

$$\lim_{c \rightarrow \pm\infty} N_{a,b,t}^{c,\lambda_a,\infty} = \lim_{c \rightarrow \pm\infty} M_{a,b,t}^{c,\lambda_a,\infty} = M_{a,b,t}^{(\pm 1),\lambda_a,\infty} \text{ a.s. and in } L^1(\mathbb{P}_x). \quad (4.7)$$

Consequently, if $M_{a,b,0}^{(\pm 1),\lambda_a,\infty} > 0$ under \mathbb{P}_x , it holds that

$$\lim_{c \rightarrow \pm\infty} \frac{\mathbb{P}_x[F_t \cdot e^{-\lambda_a L_{T_c}^a}; T_c < T_b]}{\mathbb{P}_x[e^{-\lambda_a L_{T_c}^a}; T_c < T_b]} = \mathbb{P}_x \left[F_t \cdot \frac{M_{a,b,t}^{(\pm 1),\lambda_a,\infty}}{M_{a,b,0}^{(\pm 1),\lambda_a,\infty}} \right] \quad (4.8)$$

for all bounded \mathcal{F}_t -measurable functionals F_t .

The proof is almost the same as that of Theorem 3.1, and so we omit it.

We define

$$N_{a,b,t}^{c,d,\lambda_a,\infty} := h^C(c, -d) \mathbb{P}_x \left[e^{-\lambda_a L_{T_c \wedge T-d}^a}; t < T_c \wedge T-d < T_b \middle| \mathcal{F}_t \right], \quad (4.9)$$

$$M_{a,b,t}^{c,d,\lambda_a,\infty} := h^C(c, -d) \mathbb{P}_x \left[e^{-\lambda_a L_{T_c \wedge T-d}^a}; T_c \wedge T-d < T_b \middle| \mathcal{F}_t \right] \quad (4.10)$$

for $c, d > 0$.

Theorem 4.3. *Let $x \in \mathbb{R}$ and $-1 \leq \gamma \leq 1$. Then, $(M_{a,b,t}^{(\gamma),\lambda_a,\infty}, t \geq 0)$ is a non-negative $((\mathcal{F}_t), \mathbb{P}_x)$ -martingale, and it holds that*

$$\lim_{(c,d) \xrightarrow{(\gamma)} \infty} N_{a,b,t}^{c,d,\lambda_a,\infty} = \lim_{(c,d) \xrightarrow{(\gamma)} \infty} M_{a,b,t}^{c,d,\lambda_a,\infty} = M_{a,b,t}^{(\gamma),\lambda_a,\infty} \text{ a.s. and in } L^1(\mathbb{P}_x). \quad (4.11)$$

Consequently, if $M_{a,b,0}^{(\gamma),\lambda_a,\infty} > 0$ under \mathbb{P}_x , it holds that

$$\lim_{(c,d) \xrightarrow{(\gamma)} \infty} \frac{\mathbb{P}_x [F_t \cdot e^{-\lambda_a L_{T_c \wedge T-d}^a}; T_c \wedge T-d < T_b]}{\mathbb{P}_x [e^{-\lambda_a L_{T_c \wedge T-d}^a}; T_c \wedge T-d < T_b]} = \mathbb{P}_x \left[F_t \cdot \frac{M_{a,b,t}^{(\gamma),\lambda_a,\infty}}{M_{a,b,0}^{(\gamma),\lambda_a,\infty}} \right] \quad (4.12)$$

for all bounded \mathcal{F}_t -measurable functionals F_t .

The proof is almost the same as that of Theorem 3.1, and so we omit it.

We define

$$N_{a,b,t}^{c,u,\lambda_a,\infty} := h^B(c) \mathbb{P}_x \left[e^{-\lambda_a L_{\eta_u^c}^a}; t < \eta_u^c < T_b \middle| \mathcal{F}_t \right], \quad (4.13)$$

$$M_{a,b,t}^{c,u,\lambda_a,\infty} := h^B(c) \mathbb{P}_x \left[e^{-\lambda_a L_{\eta_u^c}^a}; \eta_u^c < T_b \middle| \mathcal{F}_t \right] \quad (4.14)$$

for $c \in \mathbb{R}$ and $u > 0$.

Theorem 4.4. *Let $x \in \mathbb{R}$. Then, it holds that*

$$\lim_{c \rightarrow \pm\infty} N_{a,b,t}^{c,u,\lambda_a,\infty} = \lim_{c \rightarrow \pm\infty} M_{a,b,t}^{c,u,\lambda_a,\infty} = M_{a,b,t}^{(\pm 1),\lambda_a,\infty} \text{ a.s. and in } L^1(\mathbb{P}_x). \quad (4.15)$$

Consequently, if $M_{a,b,0}^{(\pm 1),\lambda_a,\infty} > 0$ under \mathbb{P}_x , it holds that

$$\lim_{c \rightarrow \pm\infty} \frac{\mathbb{P}_x \left[F_t \cdot e^{-\lambda_a L_{\eta_u^c}^a}; \eta_u^c < T_b \right]}{\mathbb{P}_x \left[e^{-\lambda_a L_{\eta_u^c}^a}; \eta_u^c < T_b \right]} = \mathbb{P}_x \left[F_t \cdot \frac{M_{a,b,t}^{(\pm 1),\lambda_a,\infty}}{M_{a,b,0}^{(\pm 1),\lambda_a,\infty}} \right] \quad (4.16)$$

for all bounded \mathcal{F}_t -measurable functionals F_t .

The proof is almost the same as that of Theorem 3.1, and so we omit it.

5 The inverse local time clock

In this section, we consider the inverse local time clock with the limit as u tends to infinity and c being fixed. We define

$$N_{a,b,t}^{u,\lambda_a,\lambda_b,c} := \frac{1}{\mathbb{P}_c \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a,\lambda_b} \right]} \cdot \mathbb{P}_x \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a,\lambda_b}; t < \eta_u^c \middle| \mathcal{F}_t \right], \quad (5.1)$$

$$M_{a,b,t}^{u,\lambda_a,\lambda_b,c} := \frac{1}{\mathbb{P}_c \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a,\lambda_b} \right]} \cdot \mathbb{P}_x \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a,\lambda_b} \middle| \mathcal{F}_t \right], \quad (5.2)$$

for $c \in \mathbb{R}$ and $u > 0$. We further define

$$M_{a,b,t}^{\lambda_a,\lambda_b,c} := e^{L_t^c H_{a,b}^{c,\lambda_a,\lambda_b}} \Gamma_{a,b,t}^{\lambda_a,\lambda_b}, \quad (5.3)$$

where the constant $H_{a,b}^{c,\lambda_a,\lambda_b}$ is defined by (5.8). Note that the constant $H_{a,b}^{c,\lambda_a,\lambda_b}$ is independent of u .

Theorem 5.1. *Let $x \in \mathbb{R}$. Then, $(M_{a,b,t}^{\lambda_a,\lambda_b,c}, t \geq 0)$ is a non-negative $((\mathcal{F}_t), \mathbb{P}_x)$ -martingale, and it holds that*

$$\lim_{u \rightarrow \infty} N_{a,b,t}^{u,\lambda_a,\lambda_b,c} = \lim_{u \rightarrow \infty} M_{a,b,t}^{u,\lambda_a,\lambda_b,c} = M_{a,b,t}^{\lambda_a,\lambda_b,c} \text{ a.s. and in } L^1(\mathbb{P}_x). \quad (5.4)$$

Consequently, if $M_{a,b,0}^{\lambda_a,\lambda_b,c} > 0$ under \mathbb{P}_x , it holds that

$$\lim_{u \rightarrow \infty} \frac{\mathbb{P}_x \left[F_t \cdot \Gamma_{a,b,\eta_u^c}^{\lambda_a,\lambda_b} \right]}{\mathbb{P}_x \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a,\lambda_b} \right]} = \mathbb{P}_x \left[F_t \cdot \frac{M_{a,b,t}^{\lambda_a,\lambda_b,c}}{M_{a,b,0}^{\lambda_a,\lambda_b,c}} \right] \quad (5.5)$$

for all bounded \mathcal{F}_t -measurable functionals F_t .

Proof. By definition of η_u^c , we have

$$\begin{aligned} \eta_u^c &= \inf \{ s : L_s^c > u \} \\ &= \inf \{ s : L_s^c - L_t^c > u - v \} \Big|_{v=L_t^c} \\ &= \inf \{ s : L_{s+t}^c - L_t^c > u - v \} \Big|_{v=L_t^c} + t \\ &= \inf \{ s : L_s^c \circ \theta_t > u - v \} \Big|_{v=L_t^c} + t \\ &= \eta_{u-v}^c \circ \theta_t \Big|_{v=L_t^c} + t. \end{aligned} \quad (5.6)$$

By the Markov property at t , we have

$$\mathbb{P}_x \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a,\lambda_b}; t < \eta_u^c \middle| \mathcal{F}_t \right] = \mathbb{P}_x \left[e^{-\lambda_a(L_{\eta_{u-v}^c \circ \theta_t}^a - L_t^a) - \lambda_b(L_{\eta_{u-v}^c \circ \theta_t}^b - L_t^b)}; t < \eta_u^c \middle| \mathcal{F}_t \right] \Big|_{v=L_t^c}$$

$$\begin{aligned}
&= \mathbb{P}_x \left[e^{-\lambda_a(L_{\eta_{u-v}^c}^a \circ \theta_t + L_t^a) - \lambda_b(L_{\eta_{u-v}^c}^b \circ \theta_t + L_t^b)}; t < \eta_u^c \mid \mathcal{F}_t \right] \Big|_{v=L_t^c} \\
&= 1_{\{t < \eta_u^c\}} e^{-\lambda_a L_t^a - \lambda_b L_t^b} \mathbb{P}_{X_t} \left[e^{-\lambda_a L_{\eta_{u-v}^c}^a - \lambda_b L_{\eta_{u-v}^c}^b} \right] \Big|_{v=L_t^c}.
\end{aligned} \tag{5.7}$$

By (6.5) of [2], we can let

$$\mathbb{P}_c \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a, \lambda_b} \right] = e^{-uH_{a,b}^{c, \lambda_a, \lambda_b}} \tag{5.8}$$

for $u \geq 0$. Thus, we have

$$\frac{1}{\mathbb{P}_c \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a, \lambda_b} \right]} \cdot \mathbb{P}_c \left[e^{-\lambda_a L_{\eta_{u-v}^c}^a - \lambda_b L_{\eta_{u-v}^c}^b} \right] \Big|_{v=L_t^c} = e^{L_t^c H_{a,b}^{c, \lambda_a, \lambda_b}}. \tag{5.9}$$

Therefore, we obtain

$$\lim_{u \rightarrow \infty} N_{a,b,t}^{u, \lambda_a, \lambda_b, c} = M_{a,b,t}^{\lambda_a, \lambda_b, c} \text{ a.s.} \tag{5.10}$$

Since we have

$$\begin{aligned}
\lim_{u \rightarrow \infty} \left(M_{a,b,t}^{u, \lambda_a, \lambda_b, c} - N_{a,b,t}^{u, \lambda_a, \lambda_b, c} \right) &= \lim_{u \rightarrow \infty} \frac{1}{\mathbb{P}_c \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a, \lambda_b} \right]} \cdot \mathbb{P}_x \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a, \lambda_b}; \eta_u^c \leq t \mid \mathcal{F}_t \right] \\
&= \lim_{u \rightarrow \infty} \frac{\Gamma_{a,b,\eta_u^c}^{\lambda_a, \lambda_b}}{\mathbb{P}_c \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a, \lambda_b} \right]} \cdot 1_{\{\eta_u^c \leq t\}} = 0 \text{ a.s.},
\end{aligned} \tag{5.11}$$

we obtain

$$\lim_{u \rightarrow \infty} M_{a,b,t}^{u, \lambda_a, \lambda_b, c} = M_{a,b,t}^{\lambda_a, \lambda_b, c} \text{ a.s.} \tag{5.12}$$

Next, we show that the convergence of (5.10) and (5.12) hold also in the sense of $L^1(\mathbb{P}_x)$. Since $M_{a,b,t}^{\lambda_a, \lambda_b, c} \in L^1(\mathbb{P}_x)$ for all $t > 0$, by the dominated convergence theorem, we obtain

$$\lim_{u \rightarrow \infty} N_{a,b,t}^{u, \lambda_a, \lambda_b, c} = M_{a,b,t}^{\lambda_a, \lambda_b, c} \text{ in } L^1(\mathbb{P}_x) \tag{5.13}$$

for all $t > 0$. For $q > 0$, we have by (2.12),

$$\mathbb{P}_x(\eta_u^c \leq t) \leq \mathbb{P}_c(\eta_u^c \leq t) \leq \mathbb{P}_c \left[e^{q(t - \eta_u^c)}; \eta_u^c \leq t \right] \leq e^{qt} \mathbb{P}_c \left[e^{-q\eta_u^c} \right] = e^{qt} e^{-\frac{u}{r_q(0)}}. \tag{5.14}$$

Since $r_q(0) \rightarrow 0$ as $q \rightarrow \infty$, we may take $q > 0$ large enough so that $\frac{1}{r_q(0)} > H_{a,b}^{c, \lambda_a, \lambda_b}$. Thus, for such $q > 0$, we have

$$\mathbb{P}_x \left[M_{a,b,t}^{u, \lambda_a, \lambda_b, c} - N_{a,b,t}^{u, \lambda_a, \lambda_b, c} \right] = \frac{1}{\mathbb{P}_c \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a, \lambda_b} \right]} \cdot \mathbb{P}_x \left[\mathbb{P}_x \left[\Gamma_{\eta_u^c}; \eta_u^c \leq t \mid \mathcal{F}_t \right] \right]$$

$$\leq \frac{\mathbb{P}_x(\eta_u^c \leq t)}{\mathbb{P}_c \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a, \lambda_b} \right]} \leq e^{gt} e^{u(H_{a,b}^{c,\lambda_a, \lambda_b} - \frac{1}{r_q(0)})} \rightarrow 0 \quad (5.15)$$

as $u \rightarrow \infty$. This shows that for all $t > 0$, we obtain

$$M_{a,b,t}^{u,\lambda_a,\lambda_b,c} - N_{a,b,t}^{u,\lambda_a,\lambda_b,c} \rightarrow 0 \text{ in } L^1(\mathbb{P}_x), \quad (5.16)$$

which implies

$$\lim_{u \rightarrow \infty} M_{a,b,t}^{u,\lambda_a,\lambda_b,c} = M_{a,b,t}^{\lambda_a,\lambda_b,c} \text{ in } L^1(\mathbb{P}_x) \quad (5.17)$$

for all $t > 0$. □

We define

$$N_{a,b,t}^{u,\infty,\infty,c} := \frac{1}{\mathbb{P}_c \left[\Gamma_{a,b,\eta_u^c}^{\infty,\infty} \right]} \cdot \mathbb{P}_x \left(t < \eta_u^c < T_a \wedge T_b \mid \mathcal{F}_t \right), \quad (5.18)$$

$$M_{a,b,t}^{u,\infty,\infty,c} := \frac{1}{\mathbb{P}_c \left[\Gamma_{a,b,\eta_u^c}^{\infty,\infty} \right]} \cdot \mathbb{P}_x \left(\eta_u^c < T_a \wedge T_b \mid \mathcal{F}_t \right) \quad (5.19)$$

for $c \in \mathbb{R}$ and $u > 0$. We further define

$$M_{a,b,t}^{\infty,\infty,c} := e^{L_t H_{a,b}^{c,\infty,\infty}} \Gamma_{a,b,t}^{\infty,\infty}, \quad (5.20)$$

where $H_{a,b}^{c,\infty,\infty} := \lim_{\lambda_a, \lambda_b \rightarrow \infty} H_{a,b}^{c,\lambda_a, \lambda_b}$.

Theorem 5.2. *Let $x \in \mathbb{R}$. Then, $(M_{a,b,t}^{\infty,\infty,c}, t \geq 0)$ is a non-negative $((\mathcal{F}_t), \mathbb{P}_x)$ -martingale, and it holds that*

$$\lim_{u \rightarrow \infty} N_{a,b,t}^{u,\infty,\infty,c} = \lim_{u \rightarrow \infty} M_{a,b,t}^{u,\infty,\infty,c} = M_{a,b,t}^{\infty,\infty,c} \text{ a.s. and in } L^1(\mathbb{P}_x). \quad (5.21)$$

Consequently, if $M_{a,b,0}^{\infty,\infty,c} > 0$ under \mathbb{P}_x , it holds that

$$\lim_{u \rightarrow \infty} \mathbb{P}_x[F_t \mid \eta_u^c < T_a \wedge T_b] = \mathbb{P}_x \left[F_t \cdot \frac{M_{a,b,t}^{\infty,\infty,c}}{M_{a,b,0}^{\infty,\infty,c}} \right] \quad (5.22)$$

for all bounded \mathcal{F}_t -measurable functionals F_t .

The proof is almost the same as that of Theorem 3.1, and so we omit it.

We define

$$N_{a,b,t}^{u,\infty,\infty,c} := \frac{1}{\mathbb{P}_c \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a,\infty} \right]} \cdot \mathbb{P}_x \left[e^{-\lambda_a L_{\eta_u^c}^a}; t < \eta_u^c < T_b \mid \mathcal{F}_t \right], \quad (5.23)$$

$$M_{a,b,t}^{u,\infty,\infty,c} := \frac{1}{\mathbb{P}_c \left[\Gamma_{a,b,\eta_u^c}^{\lambda_a,\infty} \right]} \cdot \mathbb{P}_x \left[e^{-\lambda_a L_{\eta_u^c}^a}; \eta_u^c < T_b \mid \mathcal{F}_t \right] \quad (5.24)$$

for $c \in \mathbb{R}$ and $u > 0$. We further define

$$M_{a,b,t}^{\infty,\infty,c} := e^{L_t H_{a,b}^{c,\lambda_a,\infty}} \Gamma_{a,b,t}^{\lambda_a,\infty}, \quad (5.25)$$

where $H_{a,b}^{c,\lambda_a,\infty} := \lim_{\lambda_b \rightarrow \infty} H_{a,b}^{c,\lambda_a, \lambda_b}$.

Theorem 5.3. *Let $x \in \mathbb{R}$. Then, $(M_{a,b,t}^{\infty,\infty,c}, t \geq 0)$ is a non-negative $((\mathcal{F}_t), \mathbb{P}_x)$ -martingale, and it holds that*

$$\lim_{u \rightarrow \infty} N_{a,b,t}^{u,\infty,\infty,c} = \lim_{u \rightarrow \infty} M_{a,b,t}^{u,\infty,\infty,c} = M_{a,b,t}^{\infty,\infty,c} \text{ a.s. and in } L^1(\mathbb{P}_x). \quad (5.26)$$

Consequently, if $M_{a,b,0}^{\infty,\infty,c} > 0$ under \mathbb{P}_x , it holds that

$$\lim_{u \rightarrow \infty} \frac{\mathbb{P}_x[F_t \cdot e^{-\lambda_a L_{\eta_u^c}}; \eta_u^c < T_b]}{\mathbb{P}_x[e^{-\lambda_a L_{\eta_u^c}}; \eta_u^c < T_b]} = \mathbb{P}_x \left[F_t \cdot \frac{M_{a,b,t}^{\infty,\infty,c}}{M_{a,b,0}^{\infty,\infty,c}} \right] \quad (5.27)$$

for all bounded \mathcal{F}_t -measurable functionals F_t .

The proof is almost the same as that of Theorem 3.1, and so we omit it.

6 Corrections to some results of Takeda–Yano [3]

There is an error in the assertion of Lemma 6.1 of [3]. The correct assertion is as follows:

Proposition 6.1 (Lemma 6.1 of [3], corrected). *For distinct points $a \neq b$, it holds that*

$$\begin{aligned} h^C(a, b) &:= \mathbb{P}_0[L_{T_a \wedge T_b}^0] \\ &= \frac{1}{h^B(a-b)} \left\{ \begin{aligned} &(h(b) + h(-a))h(a-b) + (h(a) + h(-b))h(b-a) \\ &+ (h(a) - h(b))(h(-b) - h(-a)) - h(a-b)h(b-a) \end{aligned} \right\}. \end{aligned} \quad (6.1)$$

Proof. For $q > 0$, by the strong Markov property, we have

$$\begin{aligned} \mathbb{P} \left[\int_0^\infty e^{-qt} dL_t \right] &= \mathbb{P} \left[\int_0^{T_a \wedge T_b} e^{-qt} dL_t \right] \\ &\quad + \mathbb{P}[e^{-qT_a}; T_a < T_b] \mathbb{P}_a \left[\int_0^\infty e^{-qt} dL_t \right] \\ &\quad + \mathbb{P}[e^{-qT_b}; T_b < T_a] \mathbb{P}_b \left[\int_0^\infty e^{-qt} dL_t \right]. \end{aligned} \quad (6.2)$$

Using (2.10), we have

$$r_q(0) = \mathbb{P} \left[\int_0^{T_a \wedge T_b} e^{-qt} dL_t \right] + \mathbb{P}[e^{-qT_a}; T_a < T_b] r_q(-a) + \mathbb{P}[e^{-qT_b}; T_b < T_a] r_q(-b). \quad (6.3)$$

Thus, by Lemma 3.5 of [3], we have

$$\begin{aligned} &\mathbb{P} \left[\int_0^{T_a \wedge T_b} e^{-qt} dL_t \right] \\ &= r_q(0) - \mathbb{P}[e^{-qT_a}; T_a < T_b] r_q(-a) - \mathbb{P}[e^{-qT_b}; T_b < T_a] r_q(-b) \end{aligned}$$

$$\begin{aligned}
&= r_q(0) - \frac{h_q(b-a) + h_q(-b) - h_q(-a) - \frac{h_q(-b)h_q(b-a)}{r_q(0)}}{h_q^B(a-b)} \cdot r_q(-a) \\
&\quad - \frac{h_q(a-b) + h_q(-a) - h_q(-b) - \frac{h_q(-a)h_q(a-b)}{r_q(0)}}{h_q^B(a-b)} \cdot r_q(-b). \tag{6.4}
\end{aligned}$$

We consider only the numerator of the right-hand side. Then, we have

$$\begin{aligned}
&r_q(0)h_q^B(a-b) - \left(h_q(b-a) + h_q(-b) - h_q(-a) - \frac{h_q(-b)h_q(b-a)}{r_q(0)} \right) r_q(-a) \\
&\quad - \left(h_q(a-b) + h_q(-a) - h_q(-b) - \frac{h_q(-a)h_q(a-b)}{r_q(0)} \right) r_q(-b) \\
&= h_q(a-b)(r_q(0) - r_q(-b)) + h_q(b-a)(r_q(0) - r_q(-a)) - h_q(a-b)h_q(b-a) \\
&\quad + h_q(-a)(r_q(-a) - r_q(-b)) + h_q(-b)(r_q(-b) - r_q(-a)) \\
&\quad + \frac{h_q(-b)h_q(b-a)r_q(-a)}{r_q(0)} + \frac{h_q(-a)h_q(a-b)r_q(-b)}{r_q(0)} \\
&= h_q(a-b)h_q(b) + h_q(b-a)h_q(a) - h_q(a-b)h_q(b-a) \\
&\quad + h_q(-a)(h_q(b) - h_q(a)) + h_q(-b)(h_q(a) - h_q(b)) \\
&\quad + (h_q(-b)h_q(b-a))\mathbb{P}_a[e^{-qT_0}] + (h_q(-a)h_q(a-b))\mathbb{P}_b[e^{-qT_0}] \\
&\rightarrow h(a-b)h(b) + h(b-a)h(a) - h(a-b)h(b-a) \\
&\quad + h(-a)(h(b) - h(a)) + h(-b)(h(a) - h(b)) \\
&\quad + h(-b)h(b-a) + h(-a)h(a-b) \\
&= (h(b) + h(-a))h(a-b) + (h(a) + h(-b))h(b-a) \\
&\quad + (h(a) - h(b))(h(-b) - h(-a)) - h(a-b)h(b-a)
\end{aligned}$$

as $q \rightarrow 0+$. Therefore, letting $q \rightarrow 0+$ in the equation (6.4), we obtain the assertion. \square

Next, there are some errors in the assertion of Theorem 1.7 of [3]. The correct assertion is as follows ($(a, b) \xrightarrow{\gamma} \infty$ is replaced with $(a, b) \xrightarrow{(\gamma)} \infty$):

Theorem 6.2 (Theorem 1.7 of [3], corrected). *Suppose that the condition (A) is satisfied. Let f be a positive integrable function, $x \in \mathbb{R}$, and $a, b > 0$. Define*

$$N_t^{a,b} := h^C(a, -b)\mathbb{P}_x[f(L_{T_a \wedge T_b}^0)], \quad t < T_a \wedge T_b | \mathcal{F}_t, \tag{6.5}$$

$$M_t^{a,b} := h^C(a, -b)\mathbb{P}_x[f(L_{T_a \wedge T_b}^0) | \mathcal{F}_t]. \tag{6.6}$$

Then, it holds that

$$\lim_{(a,b) \xrightarrow{(\gamma)} \infty} N_t^{a,b} = \lim_{(a,b) \xrightarrow{(\gamma)} \infty} M_t^{a,b} = M_t^{(\gamma)}$$

in the sense of \mathbb{P}_x -a.s. and in $L^1(\mathbb{P}_x)$, where the process $(M_t^{(\gamma)})_{t \geq 0}$ is defined by (1.8) of [3]. Consequently, if $M_0^{(\gamma)} > 0$ under \mathbb{P}_x , it holds that

$$\lim_{(a,b) \xrightarrow{(\gamma)} \infty} \frac{\mathbb{P}_x[F_t \cdot f(L_{T_a \wedge T_b}^0)]}{\mathbb{P}_x[f(L_{T_a \wedge T_b}^0)]} = \mathbb{P}_x \left[F_t \cdot \frac{M_t^{(\gamma)}}{M_0^{(\gamma)}} \right] \tag{6.7}$$

for all bounded \mathcal{F}_t -measurable functionals F_t .

We omit the proof of this theorem because the proof of this theorem in [3] is correct.

References

- [1] Jean Bertoin. *Lévy processes*, volume 121 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1996.
- [2] Kohki Iba and Kouji Yano. Two-point local time penalizations with various clocks for Lévy processes. preprint, arXiv:2404.06759.
- [3] Shosei Takeda and Kouji Yano. Local time penalizations with various clocks for Lévy processes. *Electron. J. Probab.*, 28:Paper No. 12, 35, 2023.
- [4] Hiroshi Tsukada. A potential theoretic approach to Tanaka formula for asymmetric Lévy processes. In *Séminaire de Probabilités XLIX*, volume 2215 of *Lecture Notes in Math.*, pages 521–542. Springer, Cham, 2018.