Liouville Theory: An Introduction to Rigorous Approaches

Sourav Chatterjee¹ and Edward Witten²

¹Department of Statistics, Stanford University, Stanford CA 94305 USA ²School of Natural Sciences, Institute for Advanced Study, Princeton, NJ 08540 USA

Abstract: In recent years, a surprisingly direct and simple rigorous understanding of quantum Liouville theory has developed. We aim here to make this material more accessible to physicists working on quantum field theory.

Contents

1 Introduction

Since it was first proposed more than forty years ago, quantum Liouville theory, or just Liouville theory for short, has had many applications and has been studied from many points of view. Relevant papers and review articles include $[1-11]$ $[1-11]$, along with many others. Rigorous mathematical work relevant to Liouville theory also has a long history, and again it is not practical to give full references. Hoegh-Krohn rigorously constructed a two-dimensional theory of a massive scalar field with an exponential interaction, using the positivity of a normal-ordered real exponential [\[12\]](#page-40-2). Kahane [\[13\]](#page-40-3), developing ideas of Mandelbrot [\[14\]](#page-40-4), gave what can be interpreted as the first rigorous statistical analysis of the Liouville measure; the motivation for that work was the statistical theory of turbulence rather than relativistic field theory. Duplantier and Sheffield [\[15\]](#page-41-0) analyzed Liouville theory on a disc and demonstrated KPZ scaling from a rigorous point of view. Closer to our focus in the present article, David, Kupianen, Rhodes, and Vargas [\[16\]](#page-41-1) gave a rigorous definition of Liouville correlation functions on a sphere, and this was further developed to a proof of the DOZZ formula for the Liouville three-point function by Kupianen, Rhodes, and Vargas [\[17\]](#page-41-2), providing a mathematical framework for ideas of Teschner [\[9\]](#page-40-5). See [\[19–](#page-41-3)[24\]](#page-41-4) for further results and background and connections to some of the many other areas of rigorous mathematical physics that are related to Liouville theory but not discussed in the present article.

The present article has both a general and a specific purpose. The general purpose is to provide for physicists a gentle introduction to rigorous arguments concerning Liouville theory. The more specific purpose is to explain how this approach clarifies some otherwise obscure properties of Liouville correlation functions.

One preliminary point is that the Liouville action

$$
I = \int_{\Sigma} d^2 x \sqrt{g} \left(\frac{1}{4\pi} \partial_a \phi \partial^a \phi + \frac{Q}{4\pi} R \phi + \mu e^{2b\phi} \right), \qquad (1.1)
$$

is real-valued. Here Σ is a closed two-manifold with local coordinates x^a , $a = 1, 2$, metric g and scalar curvature $R; b > 0$ is the Liouville coupling,^{[1](#page-2-0)} and $Q = b + 1/b$; and $\mu > 0$ is the "cosmological constant," whose value is inessential as it can be adjusted by KPZ scaling [\[4\]](#page-40-6). Because I is real, the path integral measure $D\phi e^{-I}$ is positive-definite, and probabilistic methods can potentially be applied. However, this is hardly special to Liouville theory. Many other theories, such as ϕ^4 theory in any spacetime dimension, or four-dimensional gauge theory with vanishing theta-angle, likewise have a real action and a positive path integral measure.

Much more special to Liouville theory is that the interaction is positive, even quantum mechanically. Classically, the function $e^{2b\phi}$, with real b and ϕ , is positive, of course. Quantum mechanically, in the case of Liouville theory, the only renormalization that this interaction requires is normal-ordering with respect to an underlying free field theory measure. Normal-ordering of an exponential interaction such as $e^{2b\phi}$ gives a multiplicative renormalization, which preserves positivity. So the renormalized interaction is positive, and this plays a crucial role.

To appreciate how exceptional is the positivity of the Liouville interaction, let us consider some other possible perturbations of a free massless scalar field ϕ . A mass term $\frac{1}{2}m^2\phi^2$ can again be renormalized by normal-ordering. In this case, normal-ordering entails the subtraction of a divergent constant – an additive rather than multiplicative renormalization. After this subtraction, $\frac{1}{2}m^2\phi^2$ is a well-defined observable, but unbounded below: one cannot make it positive definite by adding a constant. Similarly any polynomial function $P(\phi)$ of degree greater than 1 is not bounded below after normal-ordering. The exponential interaction in Liouville theory is special.^{[2](#page-2-1)}

In Liouville theory, one usually wants to calculate the correlation function of a product of primary fields $e^{2\alpha_i \phi(x_i)}$, $i = 1, ..., n$. For our purposes in this article, the parameters α_i are real, to ensure that the primary fields $e^{2\alpha_i \phi(x_i)}$ are positive, as assumed in the probabilistic approach. But we do not assume that the α_i are positive. We will eventually impose the Seiberg bound $\alpha_i \leq Q/2$, after understanding why it is needed. We will concentrate

¹We follow conventions of $[8, 11]$ $[8, 11]$ $[8, 11]$, for example, but we should note that much of the rigorous literature is expressed in terms of $\gamma = 2b$. Similarly, as in [\[8,](#page-40-7) [11\]](#page-40-1) and much recent literature, we denote Liouville primary fields as $e^{2\alpha\phi}$; much of the rigorous literature is expressed in terms of $\alpha = 2\alpha$.

²Other theories of scalar fields in two dimensions with a potential that is a sum of real exponentials, such as sinh-Gordon theory with potential $V(\phi) = \mu \cosh 2b\phi$, have the same positivity. This class of theories was studied in [\[12\]](#page-40-2).

on correlation functions for the case that Σ has genus 0. As the genus 0 partition function does not converge, it is usual to define unnormalized correlation functions, without trying to divide by the partition function:

$$
\left\langle \prod_{i=1}^{n} e^{2\alpha_i \phi(x_i)} \right\rangle = \int D\phi \, e^{-I(\phi)} \prod_{i=1}^{n} e^{2\alpha_i \phi(x_i)}.
$$
\n(1.2)

Let us recall a few relevant facts. Goulian and Li $[5]$ showed that the path integral in eqn. (1.2) can be usefully studied by first integrating over the "zero-mode" of the Liouville field. One writes

$$
\phi = c + X,\tag{1.3}
$$

where c is a constant, and X is a real-valued field that is subject to one real constraint (which can be chosen in various ways). The integral over c can be done explicitly, reducing the evaluation of Liouville correlation functions to an integral over X . In genus 0, after changing variables from X and c to X and $t = e^{2bc}\mu \int d^2x \sqrt{g}e^{2bX}$ and integrating over t, and using the fact that $\int_{\Sigma} d^2 x \sqrt{g} R = 8\pi$, one gets

$$
\left\langle \prod_{i=1}^{n} e^{2\alpha_i \phi(x_i)} \right\rangle = \frac{\Gamma(-w)}{2b} \int DX \, e^{-I_{\text{red}}(X)} \prod_{i=1}^{n} e^{2\alpha_i X(x_i)} \left(\mu \int d^2 x \sqrt{g} e^{2bX(x)} \right)^w, \tag{1.4}
$$

where

$$
w = \frac{Q - \sum_{i} \alpha_i}{b} \tag{1.5}
$$

and the reduced action for X is a free field action

$$
I_{\rm red}(X) = \frac{1}{4\pi} \int d^2x \sqrt{g} \left(\partial_a X \partial^a X + QRX \right). \tag{1.6}
$$

Thus "all" we have to do to determine Liouville correlation functions is to compute the expectation value of $\prod_{i=1}^n e^{2\alpha_i X(x_i)} \left(\mu \int d^2 x \sqrt{g} e^{2bX(x)}\right)^w$ with respect to the free field measure $DX e^{-I_{\text{red}}(X)}$. The only problem is that standard methods suffice to understand and potentially to compute this expectation value only when w is a non-negative integer.

When w is a non-negative integer, the gamma function in eqn. (1.4) has a pole. We will call these poles "perturbative poles," as they can be understood by the simple semiclassical calculation that we have just explained, or alternatively since they can be understood in a perturbation expansion in powers of μ . Goulian and Li were able to compute the residue of the correlation function at a semiclassical pole, by using the fact that in this case the exponent w is a non-negative integer, and making use of integrals that had been computed earlier [\[26\]](#page-41-5).

The Liouville three-point function in genus 0 is particularly important as it is expected to determine the whole theory by bootstrap or factorization arguments. By looking for a relatively simple formula that incorporates the perturbative poles and residues, and also satisfies recursion relations that those poles and residues obey, Dorn and Otto [\[7\]](#page-40-9) and A. and Al. Zamolodchikov [\[8\]](#page-40-7) suggested an exact formula (the DOZZ formula) for this threepoint function. The DOZZ formula, which will be described in somewhat more detail in section [2.7,](#page-23-0) was put on a much more solid basis by Teschner [\[9\]](#page-40-5), helping provide the basis for a rigorous proof of this formula [\[17\]](#page-41-2).

There was guesswork in the original discovery of the DOZZ formula, and this discovery also created a puzzle: although the DOZZ formula for the Liouville three-point function was motivated by trying to reproduce the perturbative poles and their residues, it actually has many additional poles whose interpretation has been less obvious. We may call the poles that appear in the DOZZ formula and do not simply arise from the integral over the Liouville zero-mode "nonperturbative," as they cannot be understood in a simple perturbation expansion in powers of μ (or b). A qualitative or semiclassical understanding of the nonperturbative poles has been missing.

In this article, we will explain how the rigorous probabilistic approach to Liouville theory sheds some light on those nonperturbative poles. A detailed explanation, which will be given in the body of this article, requires understanding non-integer powers of the Liouville interaction $\mu \int_{\Sigma} d^2x \sqrt{g} e^{2b\phi}$, and related functions that appear in the analysis. In this introduction, however, we will give a heuristic explanation, in which we consider only integer powers.

Are there ultraviolet divergences in Liouville theory? Consider expanding Liouville correlation functions in powers of μ . In n^{th} order, we have to consider an n-fold integral of the normal-ordered interaction $:e^{2b\phi}$:, of the general form $\int d^2y_1 \cdots d^2y_n :e^{2b\phi(y_1)} \cdots :e^{2b\phi(y_n)}$. Let us first consider a possible ultraviolet divergence that occurs when the points y_1, \ldots, y_n over which we integrate coincide at a point $y_0 \in \Sigma$ that is disjoint from all the points x_i at which external vertex operators $:e^{2\alpha_i\phi(x_i)}$: are inserted. In looking for such a divergence, we can restrict the y_i to a compact region $A \subset \Sigma$ that does not contain any of the x_i and thus we consider the integral

$$
\int_{A^n} d^2 y_1 \cdots d^2 y_n : e^{2b\phi(y_1)} : e^{2b\phi(y_2)} : \cdots : e^{2b\phi(y_n)} : . \tag{1.7}
$$

At short distances, ϕ can be viewed as a free field with two-point function $\langle \phi(x)\phi(y)\rangle =$ $\log \frac{1}{|x-y|}$, and the expectation value of this integral becomes

$$
\int_{A^n} d^2 y_1 \cdots d^2 y_n \prod_{1 \le i < j \le n} \frac{1}{|y_i - y_j|^{4b^2}}.
$$
\n(1.8)

Along the diagonal $y_1 = y_2 = \cdots = y_n$, the integral scales as $\int_{|y| \leq 1} d^{2n-2}y |y|^{-2n(n-1)b^2}$, and diverges if and only if $n \geq 1/b^2$. (The behavior along subdiagonals where only some of the y's coincide is less singular.) From the point of view of the operator product expansion, this divergence can be understood as follows. On general grounds, one expects such an integral to diverge if and only if a relevant or marginal operator appears in the product of n copies of $:e^{2b\phi}$. The lowest dimension operator that appears in that n-fold product is $: e^{2bn\phi}$. Recalling that in Liouville theory, the (holomorphic or antiholomorphic) dimension of the operator $:e^{2\alpha\phi}$: is $\widehat{\Delta}_{\alpha} = \alpha(Q - \alpha)$, we see that the condition in Liouville theory to get an ultraviolet divergence in n^{th} order of perturbation theory in μ is $nb(Q - nb) \leq 1$, or $n \geq 1/b^2$. In the body of this article, we will explain how this analysis can be extended

rigorously to non-integer values of n , and that will account for one of the nonperturbative poles of the DOZZ formula, namely the pole at $Q - \sum_i \alpha_i = 1/b$ or $w = 1/b^2$.

We can also consider the case that multiple copies of the interaction collide with one of the external vertex operators. For this, we assume the region A to contain one of the points x at which an external vertex operator $:e^{2\alpha\phi(x)}$: is inserted. Now we consider the integral

$$
\int_{A^n} d^2y_1 d^2y_2 \cdots d^2y_n : e^{2\alpha\phi(x)} : : e^{2b\phi(y_1)} : : e^{2b\phi(y_2)} : \cdots : e^{2b\phi(y_n)} :
$$

and the expectation value of this expression in the free field approximation is

$$
\int_{A^n} d^2 y_1 d^2 y_2 \cdots d^2 y_n \prod_{k=1}^n \frac{1}{|x - y_k|^{4\alpha b}} \prod_{1 \le i < j \le n} \frac{1}{|y_i - y_j|^{4b^2}}.
$$
\n(1.9)

This integral diverges when all n points y_1, \ldots, y_n approach x if and only if $n > 1/b^2 + 1 2\alpha = (Q - 2\alpha)/b$. From the point of view of the operator product expansion, one expects such a divergence if and only if an operator whose dimension is no greater than that of $e^{2\alpha\phi}$ appears in the product of $e^{2\alpha\phi}$ with n copies of $e^{2b\phi}$. The lowest dimension operator that appears in that product is $e^{2(\alpha+nb)\phi}$, so the condition for a divergence is $\widehat{\Delta}_{\alpha+nb} \leq \widehat{\Delta}_{\alpha}$ or $n \ge (Q-2\alpha)/b$. When properly extended to non-integer values of n, this analysis accounts for another nonperturbative pole of the DOZZ formula.

This analysis will not explain all of the nonperturbative poles of the DOZZ formula, but only those that occur on the boundary of a certain accessible region. Actually for Σ of genus 0, the path integral [\(1.2\)](#page-3-0) that defines the Liouville correlation functions converges only if $w < 0$ or $\sum_i \alpha_i > Q$. Otherwise, the integral over c diverges, leading to a pole at $w = 0$ and (after analytic continuation past this pole) to the other perturbative poles at $w = 1, 2, 3, \ldots$. After integrating over c and extracting the explicit factor $\Gamma(-w)$ in eqn. (1.4) , the remaining integral over X has an extended region of convergence. But it is not obvious what is this extended region, since it is bounded by nonperturbative phenomena. In the body of this article, we will explain how the extended region of convergence has been determined rigorously by probabilistic arguments. Two boundaries of the extended region of convergence are associated with nonperturbative poles of the DOZZ formula. The locations of these boundaries can be guessed by assuming that the preceding analysis of ultraviolet divergences can be generalized to non-integer values of n . Perhaps surprisingly, the extended region of convergence has a third boundary that is associated with a zero rather than a pole of the DOZZ formula. This zero occurs when one of the α_i reaches the Seiberg bound at $Q/2$. One might be surprised that an ultraviolet effect can lead to a zero rather than a pole. What makes this possible is that when the exponent w is negative, an ultraviolet effect, a sort of divergence of the denominator in eqn. [\(1.4\)](#page-3-1), can lead to a vanishing of the correlation function. It is not clear to us whether based on heuristic arguments one should expect that the DOZZ formula would vanish when one of the α_i reaches the Seiberg bound. But at any rate this does happen, and the probabilistic analysis gives a nice explanation of why.

To more fully understand the nonperturbative behavior of the DOZZ formula, one would need to understand the analytic continuation of the path integral beyond its extended region of convergence. Even without that, the results that we have summarized may provide an interesting general hint about quantum gravity. In recent years, there has been much interest in possible unexpected breakdown of semiclassical behavior of gravity in complex situations with many particles. Bearing in mind that small b is the semiclassical regime of Liouville theory, ultraviolet divergences that begin at roughly order $1/b^2$ in perturbation theory may be a somewhat similar phenomenon.

In section [2](#page-6-0) of this article, we give an overview of aspects of Liouville theory correlators from a probabilistic point of view. In section [3,](#page-24-0) we explain more detailed arguments.

2 Liouville Theory Correlators: Overview

2.1 The Gaussian Free Field

Though Liouville theory is conformally invariant, to analyze Liouville theory correlators in genus 0, that is for the case that the two-manifold Σ is a sphere, it is convenient to pick a specific Kahler metric on Σ. An obvious choice would be a "round" metric, but it has been found that it is convenient to pick a metric in which Σ is built by gluing together two unit disks, each with a flat metric, along their common boundary. To describe such a metric explicitly, we can view Σ as the complex x-plane, plus a point at infinity, with the metric

$$
ds^{2} = dx d\overline{x} \cdot \begin{cases} 1 & |x| \le 1 \\ \frac{1}{|x|^{4}} & |x| > 1. \end{cases}
$$
 (2.1)

The region $|x| \leq 1$ is obviously a unit disc with a flat metric, and as the given metric is invariant under the inversion $x_i \to x_i/|x|^2$, the region $|x| \geq 1$ is another copy of the same thing. The scalar curvature R of this metric has delta function support on the unit circle $|x| = 1$, and is invariant under rotation of the circle. In polar coordinates $x = re^{i\theta}$,

$$
R = 4\delta(r - 1). \tag{2.2}
$$

It is also useful to make a specific choice of the constraint on X in the decomposition of the Liouville field as $\phi = c + X$, where c is a constant and X must satisfy one real constraint to make the decomposition unique. One obvious choice would be to say that the average value of X on the sphere should vanish. Another obvious choice would involve a constraint on the "area" $\int_{\Sigma} d^2x \sqrt{g} e^{2b\phi(x)}$. A rather less obvious choice is to ask that the average of X on the unit circle $|x| = 1$ should vanish:

$$
\int_0^{2\pi} d\theta \, X(e^{i\theta}) = 0. \tag{2.3}
$$

It turns out that this choice simplifies the analysis of the correlation functions. One pre-liminary reason for this is that, combining [\(2.2\)](#page-6-2) and [\(2.3\)](#page-6-3), we have $\int_{\Sigma} d^2x \sqrt{g}RX = 0$, and therefore the reduced action [\(1.6\)](#page-3-2) becomes purely quadratic

$$
I_{\rm red} = \frac{1}{4\pi} \int d^2x \sqrt{g} \partial_a X \partial^a X,\tag{2.4}
$$

so that X is a Gaussian free field, with mean zero.

This also happens if we choose a round metric on the sphere and constrain X so that its average on the sphere vanishes. However, the choice involving a circle average also leads to a useful simplification of the propagator or two-point function $G(x, y) = \langle X(x)X(y) \rangle$, which is

$$
G(x,y) = \log \frac{1}{|x-y|} + \log |x|_+ + \log |y|_+, \tag{2.5}
$$

where $|x|_+ = \max(1, |x|)$. In verifying this, and in later applications, it is useful to know the following formula for an integral over a circle of radius r centered at a point $x \in \mathbb{C}$:

$$
\frac{1}{2\pi} \int_0^{2\pi} d\theta \log|x + re^{i\theta} - y| = \begin{cases} \log|x - y| & |x - y| \ge r \\ \log r & |x - y| \le r. \end{cases} \tag{2.6}
$$

With $\log |x-y|$ as the two-dimensional analog of the Newtonian gravitational potential, this formula is the two-dimensional analog of Newton's discovery that (in modern language) the gravitational potential of a spherical shell is that of a point mass outside the shell, and is constant inside the shell. With the help of eqn. (2.6) , one can see that eqn. (2.5) is consistent with the constraint [\(2.3\)](#page-6-3), since it implies that $\int_0^{2\pi} d\theta \langle X(e^{i\theta})X(y)\rangle = \int_0^{2\pi} d\theta G(e^{i\theta}, y) = 0$ for all y. The formula [\(2.5\)](#page-7-2) follows from this along with the fact that $G(x, y)$ has the expected logarithmic singularity along the diagonal at $x = y$, and satisfies the appropriate differential equation $-\nabla_x^2 G(x, y) = 2\pi \delta^2(x - y) + \lambda \delta_{|x|=1}$ (where λ is a Lagrange multiplier related to the constraint at $|x|=1$.

Of course, placing the constraint on the circle $|x| = 1$ is arbitrary. For example, to place the constraint instead on a circle $|x| = T$, we would replace X with a new Gaussian free field

$$
\widetilde{X}(x) = X(x) - \frac{1}{2\pi} \int_0^{2\pi} d\theta X(Te^{i\theta}), \qquad (2.7)
$$

now satisfying

$$
\int_0^{2\pi} d\theta \widetilde{X}(Te^{i\theta}) = 0.
$$
 (2.8)

The propagator or two-point function $\langle \widetilde{X}(x) \widetilde{X}(y) \rangle = \widetilde{G}(x, y)$ is given by

$$
\widetilde{G}(x,y) = \log \frac{1}{|x-y|} + \log \max(T, |x|) + \log \max(T, |y|) - \log T.
$$
 (2.9)

2.2 Preliminary Steps

Our goal is to show that the formula (1.4) for the Liouville correlation functions makes sense (after imposing some restrictions on the α_i). A useful preliminary reduction, which mathematically can be viewed as an application of Girsanov's theorem, is the following. We can eliminate the factors $\prod_{i=1}^{n} e^{2\alpha_i X(x_i)}$ in eqn. [\(1.4\)](#page-3-1) by "completing the square" in the product

$$
e^{-I_{\text{red}}(X)} \prod_{i=1}^{n} e^{2\alpha_i X(x_i)} = \exp\left(-\frac{1}{4\pi} \int d^2 x \sqrt{g} \partial_a X \partial^a X\right) \prod_{i=1}^{n} e^{2\alpha_i X(x_i)}.
$$
 (2.10)

This is done by shifting $X(x) \to X(x) + 2\sum_i \alpha_i G(x, x_i)$, leading to^{[3](#page-8-0)}

$$
\left\langle \prod_{i=1}^{n} e^{2\alpha_i \phi(x_i)} \right\rangle
$$
\n
$$
= \frac{\Gamma(-w)\mu^w}{2b} \int DX \, e^{-I_{\text{red}}(X)} \prod_{i < j} \left(\frac{|x_i| + |x_j|}{|x_i - x_j|} \right)^{4\alpha_i \alpha_j} \left(\int_{\Sigma} d^2 x \sqrt{g} f(x) e^{2bX(x)} \right)^w,
$$
\n
$$
(2.11)
$$

where

$$
f(x) = \prod_{i=1}^{n} \left(\frac{|x| + |x_i|}{|x - x_i|} \right)^{4b\alpha_i}.
$$
 (2.12)

In other words, the Liouville correlator, up to some explicitly known factors, can be defined by the w^{th} moment of the observable

$$
M_f = \int_{\Sigma} d^2x \sqrt{g} f(x)e^{2bX(x)} \tag{2.13}
$$

with respect to the free field measure. More generally, we will consider the restriction of such an integral to a region $A \subset \Sigma$:

$$
M_f(A) = \int_A d^2x \sqrt{g} \, f(x)e^{2bX(x)}.
$$
\n(2.14)

If w is a non-negative integer, then the w^{th} moment of $M_f(A)$ is simply the expectation value in the free field measure of a w-fold integral of $f(x)e^{2bX(x)}$ over A. This can be evaluated, or at least reduced to an explicitly defined integral, by using the two-point function [\(2.5\)](#page-7-2) to evaluate the expectation value of a product of exponentials. That is what Goulian and Li did [\[5\]](#page-40-8), with a different choice of the Kahler metric and the condition on X. What can we do when w is not a non-negative integer? This is where probabilistic methods [\[16–](#page-41-1)[20\]](#page-41-6) have been useful, as we will aim to explain.

In studying $M_f(A)$ as a random variable, it turns out that if the Seiberg bound $\alpha_i \leq$ $Q/2$ is not satisfied, then $M_f(A) = +\infty$ with probability 1. In that case, probabilistic methods (or at least the arguments that we will explain) do not enable one to make sense of Liouville correlation functions. However, as long as all α_i satisfy the Seiberg bound, it is possible to define the M_f (or $M_f(A)$ for any A) as a positive random variable. Saying that M_f is a positive random variable means that there is a probability measure ρ on the positive real line that determines the expectation value of a function of M_f . By definition a probability measure on the half-line is non-negative and satisfies

$$
\int_0^\infty \mathrm{d}u \,\rho(u) = 1. \tag{2.15}
$$

The function ρ is not necessarily smooth; it might have delta function support at some values of u. In Liouville theory, there can be a delta function at $u = 0$ (but not elsewhere).

³The operators $e^{2\alpha_i X(x_i)}$ need to be normal-ordered, as we will do in a moment for e^{2bX} . As usual, taking this normal ordering into account eliminates divergent factors involving $G(x_i, x_i)$ that would otherwise appear in the following formula.

This happens precisely if $b > 1$, in which case it turns out that $M_f = 0$ with probability 1, corresponding to $\rho(u) = \delta(u)$. So for $b > 1$ the methods we will describe will again not be useful for defining Liouville correlation functions.

In general, the expectation value of a function $h(\Phi)$ of a random variable Φ with probability measure ρ is

$$
\mathbb{E}[h(\Phi)] = \int_0^\infty \mathrm{d}u \rho(u) h(u),\tag{2.16}
$$

assuming that this integral is well-defined. Here we switch to probabilistic notation and write $\mathbb{E}[h(\Phi)]$ rather than $\langle h(\Phi) \rangle$ for the expectation value of $h(\Phi)$. If the function h is not positive or negative-definite, the integral in eqn. [\(2.16\)](#page-9-0) may be ill-defined. However, in our application, h will be a positive function $h(u) = u^w$ for some real w. Given this, assuming that $w > 0$ or that $\rho(u)$ does not have a delta function at $u = 0$ (which would make the integral ill-defined if $w < 0$, the integral for $\mathbb{E}[M_f^w]$ converges to a positive real number or $+\infty$. When the integral converges to $+\infty$, this will mark a boundary of the extended region of convergence that was discussed in the introduction: on this boundary, the DOZZ formula will have a nonperturbative singularity. If $\mathbb{E}[M_f^w]$ vanishes for some $w > 0$, this will tell us that $\rho(u) = \delta(u)$ and that M_f vanishes with probability 1. This happens if $b > 1$. Finally, $\mathbb{E}[M_f^w]$ will converge for negative w if and only if $\rho(u)$ vanishes sufficiently rapidly near $u = 0$. In Liouville theory, the random variable M_f always has that property.

It will be convenient if we can take the two-point function of X to be simply $\log(1/|x$ y), not just because this is a simple function but more importantly because it has nice scale invariance properties and a useful relation to random walks. To reduce to this simple form of the propagator, we can note the following. A finite sum of positive random variables is again a positive random variable. To exploit this, we can decompose Σ as a union of two sets Σ_1 and Σ_2 , where Σ_1 is the unit disc $|x| \leq 1$, and Σ_2 is the set $|x| \geq 1$. Then $A = A \cap \Sigma_1 \cup A \cap \Sigma_2$, so

$$
M_f(A) = M_f(A \cap \Sigma_1) + M_f(A \cap \Sigma_2). \tag{2.17}
$$

Hence defining the $M_f(A \cap \Sigma_i)$ as positive random variables for $i = 1, 2$ will lead to such a definition for $M_f(A)$. Eqn. [\(2.5\)](#page-7-2) already has the desired form for $x, y \in A \cap \Sigma_1$, and the same is true in $A \cap \Sigma_2$ after a change of variables $x \to 1/x$. Another convenient fact is that in each of the two regions Σ_1 and Σ_2 , the metric is simply the Euclidean metric $dx_1^2 + dx_2^2$ (up to the inversion $x_i \to x_i/|x|^2$ in the case of Σ_2).

Of course, there is nothing special about the particular decomposition $\Sigma = \Sigma_1 \cup \Sigma_2$ that was used here. We could have used the variable \widetilde{X} instead of X, making a similar division at $|x| = T$, and we could similarly decompose Σ as the union of more than two pieces.

So finally, to define Liouville correlators, it suffices to consider a region $A \subset \Sigma$ in which the metric is flat and the two-point function of X is a simple logarithm, and to study the quantity

$$
M_f(A) = \int_A d^2x f(x)e^{2bX(x)},
$$
\n(2.18)

where the function $f(x)$ is either smooth and bounded in region A, or has just one singularity of the form $|x - x_0|^{-4b\alpha}$ for some α and some $x_0 \in A$.

2.3 Normal Ordered Interaction

The field $X(x)$ can be understood as a random Gaussian distribution. This statement means that if we pair it with a smooth function $f(x)$ (real-valued for simplicity, and necessarily of compact support as Σ is compact), we get a Gaussian random variable $X_f = \int d^2x f(x)X(x)$, with $\mathbb{E}[X_f] = 0$ and $\mathbb{E}[X_f^2] = \int d^2x d^2y f(x) f(y)G(x, y)$. This integral converges for any f. Without such smearing, although the "operator" $X(x)$ has well-defined correlation functions $\mathbb{E}[X(x)X(y)] = G(x, y)$ for $x \neq y$, it does not make sense as a random variable. Its variance would be $\mathbb{E}[X(x)^2] = G(x,x) = +\infty$, but a Gaussian measure $\rho(u) = \frac{1}{\sigma\sqrt{2\pi}}e^{-u^2/2\sigma^2}$ does not have a limit as the variance σ^2 goes to infinity. So one cannot assign probabilities to the values of $X(x)$; it does not really have values.

The random distribution that we want to define is not $X(x)$ but a renormalized version of $e^{2bX(x)}$. Here, renormalization is required just to define correlation functions of $e^{2bX(x)}$, even without trying to define it as a random variable. To accomplish this, we will first introduce a cutoff by smearing X slightly to get a cutoff version X_{ϵ} that is a Gaussian random variable, so that $e^{2bX_{\epsilon}(x)}$ will also make sense as a random variable. Then we will renormalize by normal-ordering. We will also describe another useful kind of cutoff in section [2.4.](#page-12-0)

In general, if Y is a centered Gaussian variable (the statement that it is "centered" means that its expected value is zero), the normal ordered version of Y is $e^{Y-\frac{1}{2}\mathbb{E}[Y^2]}$, which we will denote as : e^Y :. Note that $\mathbb{E}[e^Y] = 1$ for any Y. In particular, the normal-ordered version of $e^{2bX_{\epsilon}(x)}$ will be $:e^{2bX_{\epsilon}(x)} = e^{2bX_{\epsilon}(x)-2b^{2}\mathbb{E}[X_{\epsilon}^{2}]}$. Finally we will define

$$
M_{f,\epsilon}(A) = \int_A d^2x f(x) : e^{2bX_{\epsilon}(x)}:
$$
\n(2.19)

and investigate the behavior as $\epsilon \to 0$. The random variable that we want is

$$
M_f(A) = \lim_{\epsilon \to 0} M_{f,\epsilon}(A),\tag{2.20}
$$

if this limit exists. If the limit does exist, then it makes sense to define the moments $E[M_f(A)^w]$ of $M_f(A)$ with w not necessarily an integer, though in general these moments may equal $+\infty$. Similar we can take a linear combination of random variables $M_f(A)$ with positive coefficients and define their moments for general real w.

One obvious way to define a smeared version $X_{\epsilon}(x)$ of $X(x)$ would be to smear $X(x)$ over a disc of radius ϵ centered at x. However, it turns out that it is more convenient to instead average $X(x)$ over a circle of radius ϵ centered at x. Thus we define

$$
X_{\epsilon}(x) = \frac{1}{2\pi} \int_0^{2\pi} d\theta X(x + \epsilon e^{i\theta}).
$$
\n(2.21)

If $\langle X(x)X(y)\rangle = \log \frac{1}{|x-y|}$ in the relevant portion of Σ , then for $|x-y| > 2\epsilon$, we have simply $\langle X_{\epsilon}(x)X_{\epsilon}(y)\rangle = \log \frac{1}{|x-y|}$, just as at $\epsilon = 0$. For $|x-y| < 2\epsilon$, that is no longer true, since the circles of radius ϵ centered at x and y intersect. For example,

$$
\langle X_{\epsilon}(x)X_{\epsilon}(x)\rangle = \log\frac{1}{\epsilon},\tag{2.22}
$$

by use of eqn. (2.6) .

The general formula for $\langle X_{\epsilon}(x)X_{\epsilon}(y)\rangle$ is a little complicated, but what we will need to know can be summarized in a few simple facts. First, this function is monotonically increasing as ϵ becomes smaller. Equivalently,

$$
\frac{\partial}{\partial \epsilon} \langle X_{\epsilon}(x) X_{\epsilon}(y) \rangle \le 0 \tag{2.23}
$$

for all x, y. Indeed, more generally, for any $\epsilon, \epsilon' > 0$,

$$
\frac{\partial}{\partial \epsilon} \langle X_{\epsilon}(x) X_{\epsilon'}(y) \rangle \le 0, \tag{2.24}
$$

so $\langle X_{\epsilon}(x)X_{\epsilon'}(y)\rangle$ increases as either ϵ or ϵ' becomes smaller. To show that, first note from eqn. (2.6) that

$$
\langle X_{\epsilon}(x)X(y)\rangle = \min\left(\log\frac{1}{|x-y|}, \log\frac{1}{\epsilon}\right),\tag{2.25}
$$

and clearly this is monotonically increasing as ϵ decreases. Therefore, averaging over θ , $\langle X_{\epsilon}(x) X_{\epsilon'}(y) \rangle = \frac{1}{2\pi}$ $\frac{1}{2\pi} \int_0^{2\pi} d\theta \langle X_{\epsilon}(x) X(y + \epsilon' e^{i\theta}) \rangle$ is monotonically increasing as ϵ decreases. By symmetry, the same is true as ϵ' decreases. Setting $\epsilon = \epsilon'$, we deduce the inequality [\(2.23\)](#page-11-0), which will be used in section [3.](#page-24-0) A second useful fact is

$$
\langle X_{\epsilon}(x)X_{\epsilon}(y)\rangle \le \min\left(\log\frac{1}{|x-y|}, \log\frac{1}{\epsilon}\right) \tag{2.26}
$$

(with equality if and only if $|x - y| \ge 2\epsilon$ or $x = y$). This statement is equivalent to two inequalities: the left hand side is equal to or less than $\log \frac{1}{|x-y|}$ and equal to or less than $\log \frac{1}{\epsilon}$. The first inequality is true because it is true (as an equality) if $\epsilon = 0$, and therefore by eqn. [\(2.23\)](#page-11-0), it is also true for $\epsilon > 0$. Concerning the second inequality, by symmetry we can assume $x = 0$, $y = r$. By eqn. [\(2.22\)](#page-11-1), the second inequality holds as an equality if $r = 0$, so it suffices to show that

$$
\frac{\partial}{\partial r} \langle X_{\epsilon}(0) X_{\epsilon}(r) \rangle \le 0. \tag{2.27}
$$

We have

$$
\langle X_{\epsilon}(0)X_{\epsilon}(r)\rangle = \frac{1}{4\pi^2} \int_0^{2\pi} d\theta \int_0^{2\pi} d\theta' \langle X(\epsilon e^{i\theta})X(r + \epsilon e^{i\theta'})\rangle.
$$
 (2.28)

Integrating over θ via eqn. [\(2.6\)](#page-7-1), we get $\langle X_{\epsilon}(0)X_{\epsilon}(r)\rangle=\frac{1}{2i}$ $\frac{1}{2\pi} \int_0^{2\pi} d\theta' w(r, \theta')$, where

$$
w(r', \theta) = \min\left(\log \frac{1}{|r + \epsilon e^{i\theta'}|}, \log \frac{1}{\epsilon}\right). \tag{2.29}
$$

Then $\frac{\partial w(r,\theta')}{\partial r} \leq 0$ for all r, θ' , because $\frac{\partial}{\partial r}|r + \epsilon e^{i\theta'}|$ is positive except when $|r + \epsilon e^{i\theta'}| < \epsilon$, in which case $w = \log(1/\epsilon)$ and $\frac{\partial w}{\partial r} = 0$. Averaging the result $\frac{\partial w}{\partial r} \leq 0$ over θ' , we get the claimed inequality (2.27) . Eqn. (2.26) can be generalized to an upper bound on $\langle X_{\epsilon}(x)X_{\epsilon'}(y)\rangle$. The case of this that we will need is

$$
\langle X_{\epsilon}(x)X_{\epsilon'}(y)\rangle = \log \frac{1}{|x-y|} \quad \text{if} \quad |x-y| \ge 2\epsilon, 2\epsilon'. \tag{2.30}
$$

This follows from eqn. (2.25) together with (2.6) .

Since $\langle X_{\epsilon}(x)^{2} \rangle = \log(1/\epsilon)$, we have $:e^{2bX_{\epsilon}(x)} := e^{2bX_{\epsilon}(x)-2b^{2}\log(1/\epsilon)}$. Hence the cutoff version of $M_f(A)$ is

$$
M_{f,\epsilon}(A) = \int_A \mathrm{d}^2 x f(x) e^{2bX_{\epsilon}(x) - 2b^2 \log(1/\epsilon)}.
$$
\n(2.31)

This is a well-defined random variable, and we want to investigate its convergence as $\epsilon \to 0$.

2.4 Martingales

The circle cutoff that was just introduced will be useful in section [3.](#page-24-0) However, the simplest proof of existence of limits such as $\lim_{\epsilon \to 0} M_{f,\epsilon}(A)$ actually requires a different sort of cutoff.

In probability theory, a random variable $\Phi(t)$ that depends on a real (or integer-valued) parameter t is called a *martingale* if, conditional on a knowledge of the whole history of $\Phi(t)$ for $t \leq t_0$, the expected value of $\Phi(t)$ for any $t > t_0$ is precisely $\Phi(t_0)$. It does not matter when in the past the process started; $\Phi(t)$ is a martingale if the expected value of $\Phi(t)$ for any $t > t_0$ is precisely $\Phi(t_0)$, independent of what happened for $t < t_0$. For example, one's stake in a fair game of chance in which on the average one neither gains nor loses by playing is a martingale. Regardless of one's stake at time t_0 , the expected value of that stake at any future time is the same.

With a different regularization from the one that we described in section [2.3,](#page-10-0) the cutoff dependence of the random variable $M_f(A)$ is a martingale. This was exploited in the foundational work of Mandelbrot [\[14\]](#page-40-4) and Kahane [\[13\]](#page-40-3) on random variables such as $M_f(A)$.

Folllowing [\[15\]](#page-41-0), we will explain a variant of the construction that is slightly simpler to describe first for Liouville theory on a disc D , rather than on a two-sphere. In this case, one has to study the same Gaussian free field $X(x)$ but now with, for example, Dirichlet boundary conditions on the disc. Let $f_i(x)$ be the normalized eigenfunctions of the Dirichlet Laplacian on D, with increasing eigenvalues $0 < \lambda_1 \leq \lambda_2 \leq \cdots$. The field $X(x)$ can be expanded in modes:[4](#page-12-1)

$$
X(x) = \sum_{i=1}^{\infty} \frac{f_i(x)X_i}{\sqrt{\lambda_i}},
$$
\n(2.32)

where X_i are independent centered Gaussian variables, $\mathbb{E}[X_i X_j] = \delta_{ij}$. The normal ordered exponential of $X(x)$ is formally an infinite product:

$$
:e^{2bX(x)} := \prod_{i=1}^{\infty} e^{2bf_i(x)X_i - 2b^2f_i(x)^2}.
$$
 (2.33)

⁴Kahane's procedure was slightly more complicated, for a reason that will be explained in section [3.2.](#page-26-0)

For a regularization, we can simply truncate this as a finite product and define

$$
:e^{2bX(x)}:_{n} = \prod_{i=1}^{n} e^{2bf_i(x)X_i - 2b^2f_i(x)^2}.
$$
\n(2.34)

The sequence of random variables $:e^{2bX(x)}:n, n = 1, 2, 3, \ldots$, is a martingale, since conditioning on the values of X_1, X_2, \ldots, X_n does not give any information about X_{n+1} and $\mathbb{E}[e^{2bf_{n+1}(x)X_{n+1}-2b^2f_{n+1}(x)^2}] = 1$ for all n. So, conditional on the values of X_1, X_2, \ldots, X_n , the expected value of $:e^{2bX(x)}:_{n+1}$ is just the value of $:e^{2bX(x)}:_{n}$. Integration over a region $A \subset D$ does not disturb the martingale property, so likewise the sequence of random variables

$$
M_{f,n}(A) = \int_A d^2x f(x) : e^{2bX(x)} :_n
$$
\n(2.35)

is a martingale. Of course, this is a positive martingale as $:e^{2bX(x)}$; is a product of real exponentials. We have

$$
\mathbb{E}[M_{f,n}(A)] = \int_A d^2x f(x) \tag{2.36}
$$

for all n , as this is a universal property of the normal ordered exponential of a Gaussian random variable.

We can make the same construction for Liouville theory on a two-sphere, with only slightly more care. First we make a Fourier expansion $X(r, \theta) = \sum_{n \in \mathbb{Z}} X_n(r) e^{in\theta}$, as in eqn. [\(2.38\)](#page-14-0). The modes X_n for $n \neq 0$ can be expanded in eigenfunctions of the Laplacian on the sphere, just as before. For $n = 0$, we just have to note that because of the constraint $X_0(r) = 0$ at $r = 1$, the field $X_0(r)$ should be treated separately on the two discs $r \leq 1$ and $r \geq 1$. On each disc, $X_0(r)$ can be expanded in eigenmodes of the Dirichlet Laplacian on the disc.

A simple but rather striking result in probability theory is Doob's martingale con-vergence theorem (see [\[25\]](#page-41-7), p. 319), which says that if $\Phi(t)$ is a positive martingale then with probability 1, $\Phi(t)$ converges to a nonnegative real number. This limit is a random variable. An informal explanation of the proof is as follows. Think of $\Phi(t)$ as the price of a commodity at time t. The fact that $\Phi(t)$ is a martingale means that there is no way to make an expected profit by buying and selling the commodity, because at any time t_0 , the expected value of the price $\Phi(t_1)$ at any future time t_1 is just the current price $\Phi(t_0)$. The positivity of $\Phi(t)$ together with the martingale condition implies that $\Phi(t)$ does not have a positive probability to diverge to $+\infty$ for $t \to \infty$; if this occurs, the expectation value of $\Phi(t)$ would also diverge as $t \to \infty$, contradicting the martingale property. So with probability 1, $\Phi(t)$ does not diverge to $+\infty$. Suppose now that with probability $\delta > 0$, $\Phi(t)$ does not converge. This means that there are numbers $a < b$ such that with positive probability, $\Phi(t)$ oscillates infinitely many times between a and b. In this case, a strategy of selling when the price is b and buying when the price is a has an average gain of $+\infty$, contradicting the fact that there is no winning strategy. So it must be that $\Phi(t)$ converges with probability 1 and this limit is automatically a positive (or more precisely, non-negative) random variable.

Therefore, the random variables $M_{f,n}(A)$ have limits for $n \to \infty$. These limits are nonnegative random variables and it makes sense to define their non-integer moments (which in general are positive real numbers or $+\infty$). However, there is some fine print in the martingale convergence theorem. It is possible for the limit of a positive martingale to vanish. This is possible even if the martingale is normalized in the sense of eqn. [\(2.36\)](#page-13-0), which at first sight may appear to imply that the large n limit of $M_{f,n}(A)$ cannot possibly vanish.

It is not difficult to give an artificial example of this behavior.^{[5](#page-14-1)} However, an example based on Brownian motion is more relevant to Liouville theory and also gives some intuition about how the infinite product [\(2.33\)](#page-12-2) could converge to 0.

First let us explain how Brownian motion is related to Liouville theory. Set $\epsilon = e^{-t}$, and consider the dependence on t of $X_{\epsilon}(x) = X_{\epsilon^{-t}}(x)$, for some fixed value of x. We claim that the dependence on t of this variable is described by Brownian motion – the continuum version of a random walk. To see this, pick polar coordinates r, θ centered at x_0 , so $x = x_0 + re^{i\theta}$. Let $r = e^{-t}$. In terms of t and θ the Gaussian free field action for X (see eqn. (2.4)) becomes^{[6](#page-14-2)}

$$
I_{\rm red} = \frac{1}{4\pi} \int dt \int_0^{2\pi} d\theta \left(\left(\frac{\partial X}{\partial t} \right)^2 + \left(\frac{\partial X}{\partial \theta} \right)^2 \right).
$$
 (2.37)

Expand $X(t, \theta)$ in Fourier modes:

$$
X(t,\theta) = \sum_{n \in \mathbb{Z}} X_n(t)e^{in\theta}.
$$
\n(2.38)

The zero-mode $X_0(t)$ is precisely the variable $X_{e^{-t}}(x)$ whose dependence on t we claim is described by Brownian motion. The action I_{red} has the property that the $X_n(t)$ for different values of n are independent of each other. The zero-mode $X_0(t)$ is described by the action

$$
I_{\rm red,0} = \frac{1}{2} \int dt \left(\frac{dX_0(t)}{dt} \right)^2.
$$
 (2.39)

This is precisely the action that describes Brownian motion, confirming the claim that Brownian motion governs the dependence on t of $X_0(t) = X_{e^{-t}}(x_0)$.

⁵Let Φ_n , $n = 0, 1, 2, \ldots$ be a sequence of random variables, defined by saying that $\Phi_0 = 1$, and, inductively, if $\Phi_n = 0$ then $\Phi_{n+1} = 0$, but if $\Phi_n = 2^n$, then $\Phi_{n+1} = 2^{n+1}$ with probability $1/2$ and $\Phi_{n+1} = 0$ with probability 1/2. So Φ_n equals 2^n with probability 2^{-n} , and vanishes otherwise. We can think of Φ_n as the stake at time n of a gambler who starts with \$1 at time 0, and in each time step, risks the entire stake on the flip of a coin, doubling the stake if the coin comes up heads and losing everything (and leaving the game) if it comes up tails. Continuing to play the game does not change the expectation value of the gambler's stake, regardless of what that stake is at a given time, so this sequence of random variables is a martingale. The expectation value of the gambler's stake is \$1 at any time, but with probability 1, the gambler will ultimately lose the entire stake. So even though $\mathbb{E}[\Phi_n] = 1$ for all n, the large n limit of Φ_n is 0.

 6 This is a consequence of the fact that the Gaussian free field action is invariant under a Weyl rescaling of the metric $g \to e^h g$ for any real-valued function h, together with the fact that the change of variables from Euclidean coordinates on the plane to polar coordinates r, θ is a conformal mapping.

As a matter of terminology, standard Brownian motion is Brownian motion that starts at the origin at time 0, so $X_0(0) = 0$, and satisfies $\mathbb{E}[X_0(t)^2] = t$. Eqn. [\(2.39\)](#page-14-3) is normalized to lead to this time-dependence of the variance. If we choose $x = 0$, then the constraint $X_0(0) = 0$ is a restatement of eqn. [\(2.3\)](#page-6-3), so in that case $B_t = X_0(t)$ is described by a standard Brownian motion.

Now let us go back to martingales. The Brownian motion $X_0(t)$ is itself a fairly obvious example of a martingale. $X_0(t)$ could have any value at time $t = t_0$, but the change in X_0 for time $t > t_0$ is independent of what happened for $t < t_0$, and has average value zero, so conditional on knowing $X_0(t_0)$, the average value of $X_0(t_1)$, for any $t_1 > t_0$, is just $X_0(t_0)$.

Less obvious is that the normal-ordered exponential of $X_0(t)$ is again a martingale. This will be used in section [3.7.](#page-36-0) To show that $e^{\gamma X_0(t) - \frac{\gamma^2}{2}}$ $\frac{\gamma^2}{2} \mathbb{E}[X_0(t)^2]$ is a martingale for any γ , note that

$$
e^{\gamma X_0(t_1) - \frac{\gamma^2}{2} \mathbb{E}[X_0(t_1)^2]} = U \cdot e^{\gamma X_0(t_0) - \frac{\gamma^2}{2} \mathbb{E}[X_0(t_0)^2]},\tag{2.40}
$$

with

$$
U = e^{\gamma(X_0(t_1) - X_0(t_0)) - \frac{\gamma^2}{2} \left(\mathbb{E}[X_0(t_1)^2] - \mathbb{E}[X_0(t_0)^2] \right)}.
$$
\n(2.41)

So we have to show that conditional on a knowledge of $X_0(t_0)$ (which is equivalent to a knowledge of $e^{\gamma X_0(t_0) - \frac{\gamma^2}{2}}$ $\frac{\gamma^2}{2} \mathbb{E}[X_0(t_0)^2]$, the expected value of U is equal to 1. To show this, we observe that $\gamma(X_0(t_1) - X_0(t_0))$ is a centered Gaussian random variable, so

$$
\mathbb{E}[U] = e^{\frac{\gamma^2}{2}\mathbb{E}[(X_0(t_1) - X_0(t_0))^2] - \frac{\gamma^2}{2}(\mathbb{E}[X_0(t_1)^2] - \mathbb{E}[X_0(t_0)^2])} = e^{-\gamma^2 \mathbb{E}[(X_0(t_1) - X_0(t_0))X_0(t_0)]}. \tag{2.42}
$$

But $\mathbb{E}[(X_0(t_1) - X_0(t_0))X_0(t_0)] = 0$, since in Brownian motion, regardless of the value of $X_0(t_0)$, the expected value of the increment $X_0(t_1) - X_0(t_0)$ vanishes.

Since $e^{\gamma X_0(t) - \frac{1}{2}\gamma^2 \mathbb{E}[X_0(t)^2]} = e^{\gamma X_0(t) - \frac{1}{2}\gamma^2 t}$ is a positive martingale, it will have a large t limit. One might think that this limit would have to be nonzero, since $\mathbb{E}[e^{\gamma X_0(t)-\frac{1}{2}\gamma^2t}] = 1$ for all t . To see that this is wrong, one can ask what is the typical value of the exponent at large t. The first term $\gamma X_0(t)$ is typically of order $\gamma |t|^{1/2}$ while the second term $-\frac{\gamma^2}{2}$ $\frac{\gamma}{2}t$ is negative and proportional to t. So a typical value of $e^{\gamma X_0(t) - \frac{\gamma^2}{2}}$ $\frac{\gamma}{2}$ ^t is exponentially small at large t. Although it is true that $\mathbb{E}[e^{\gamma X_0(t) - \frac{\gamma^2}{2}}]$ $\sum_{i=1}^{T} t^{i} = 1$ for all t, this nonvanishing value at late times reflects the contribution of exponentially unlikely events in which $X_0(t)$ is unexpectedly large. For large t, with a probability that is exponentially close to 1, $e^{\gamma X_0(t) - \frac{\gamma^2}{2}}$ $\frac{\gamma}{2}^t$ is exponentially small, and the large t limit of this variable vanishes. The fact that the normal ordered exponential of Brownian motion decays exponentially in time will be important in section [3.6.](#page-35-0)

In more detail, one way to prove that a positive random variable Φ vanishes is to find $p > 0$ such that $\mathbb{E}[\Phi^p] = 0$. This implies that the probability density function ρ of Φ satisfies $\int_0^\infty dx \rho(x) x^p = 0$, forcing $\rho(x) = \delta(x)$ and $\Phi = 0$. Let us implement this with $\Phi(t) = e^{\gamma X_0(t) - \frac{\gamma^2}{2}}$ $\frac{\gamma^2}{2}$ ^t. We have $\Phi(t)^p = e^{p\gamma X_0(t) - p\frac{\gamma^2}{2}}$ $\frac{\gamma^2}{2}$ ^t, leading to $\mathbb{E}[\Phi(t)^p] = e^{-\frac{\gamma^2 t}{2}(p-p^2)}$. For $0 < p < 1$, the exponent is negative and $\lim_{t\to\infty} \mathbb{E}[\Phi(t)^p] = 0$, implying that $\lim_{t\to\infty} \Phi(t) = 0$ 0. Such behavior actually occurs in Liouville theory for $b > 1$, as we will show in sections [2.6](#page-18-0) and [3.3](#page-28-0) by studying the p^{th} moment of $M_f(A)$ for p slightly less than 1.

Although : $e^{2bX_{\epsilon}(x)}$: for fixed x can thus be viewed as a martingale in its dependence on ϵ , if we integrate this quantity over a region $A \subset \Sigma$, we do not get a martingale, because circles centered at different points in A intersect each other and the data on those circles are not independent. That is why we explained the alternative regularization based on a normal mode expansion, which does give a martingale. However, the regularization based on circle averaging has other advantages and is used in the rest of this article. Of course, physically one expects that the different regularizations will lead to equivalent results.

Thus a key issue in Liouville theory is to show that the random variables $M_f(A)$, defined by normal ordering, are actually nonzero for suitable b. As we explain next, there is a relatively simple proof of this for $b < 1/\sqrt{2}$ with a suitable condition on f. This leads to a simple rigorous definition of the Liouville correlation functions, for such b, at least if the Liouville momenta α_i of the operators considered are small enough. The proof that the $M_f(A)$ are nonzero for all $b < 1$ is considerably more difficult and is deferred to section [3.7.](#page-36-0)

 2.5 ² Convergence for $b < 1/\sqrt{ }$ 2

Going back to the circle regularization, here we will analyze the convergence of $M_{f,\epsilon}(A)$ as $\epsilon \to 0$ for the case $b < 1/\sqrt{2}$. We assume a region A in which the two-point function is just $\mathbb{E}[X(x)X(y)] = \log(1/|x-y|)$. To begin with, we assume that f is smooth and bounded; then we will consider singularities due to insertions of primary fields.

Real-valued random variables that have convergent second moments form a vector space, because as $(\Phi + \Phi')^2 \leq 2(\Phi^2 + \Phi'^2)$, it follows that if Φ and Φ' have convergent second moments, then so does $\Phi + \Phi'$: $\mathbb{E}[(\Phi + \Phi')^2] \leq \mathbb{E}[2(\Phi^2 + (\Phi')^2)] = 2\mathbb{E}[\Phi^2] + 2\mathbb{E}[(\Phi')^2] < \infty$. This vector space has a positive-definite inner product $(\Phi', \Phi) = \mathbb{E}[\Phi'\Phi]$. Completing it with respect to this inner product, we get a Hilbert space $\mathcal H$ of random variables that have finite second moments. We can think of this as the Hilbert space of square-integrable functions on some measure space. If Φ_{ϵ} , for $\epsilon > 0$, is a family of vectors in H, and the limit $\Phi = \lim_{\epsilon \to 0} \Phi_{\epsilon}$ exists in H, then we say that the random variables Φ_{ϵ} converge in L^2 to Φ as $\epsilon \to 0$. Since a Hilbert space is complete by definition, the family Φ_{ϵ} has such a limit as $\epsilon \to 0$ if and only if it is a Cauchy sequence in the L^2 sense, meaning that

$$
\lim_{\epsilon,\epsilon' \to 0} \mathbb{E}[(\Phi_{\epsilon} - \Phi_{\epsilon'})^2] = 0.
$$
\n(2.43)

We will show that for $b < 1/$ √ 2, the family $M_{f,\epsilon}(A)$ is such a Cauchy sequence, and therefore the limit

$$
M_f(A) = \lim_{\epsilon \to 0} M_{f,\epsilon}(A) \tag{2.44}
$$

exists in the L^2 sense. Consequently, in this range of b, the second moment of $M_f(A)$, which is its norm squared as a Hilbert space vector, exists and is the limit of the second moment of $M_{f,\epsilon}(A)$:

$$
\mathbb{E}[M_f(A)^2] = \lim_{\epsilon \to 0} \mathbb{E}[M_{f,\epsilon}(A)^2].\tag{2.45}
$$

That in particular will show that $M_f(A)$ is not zero in this range of b.

It should not come as a surprise that $M_f(A)$ has a convergent second moment if $b < 1/\sqrt{2}$, because in the introduction we explained that an observable such as $M_f(A)$

(with smooth bounded f) has a convergent nth moment, for integer n, if and only if $n < 1/b^2$, and here we are simply making this statement for $n = 2$. What we gain by defining $M_f(A)$ as a random variable is that we can deduce that its non-integer moments exist, at least in a suitable range of exponents. Indeed, if Φ is a positive random variable with $\mathbb{E}[\Phi^p] < \infty$ and associated to a probability density ρ on the positive half-line, then for $0 \leq q < p$, we have

$$
\mathbb{E}[\Phi^q] = \int_0^\infty \mathrm{d}u \rho(u) u^q = \int_0^1 \mathrm{d}u \rho(u) u^q + \int_1^\infty \mathrm{d}u \rho(u) u^q
$$

\n
$$
\leq \int_0^1 \mathrm{d}u \rho(u) + \int_1^\infty \mathrm{d}u \rho(u) u^p \leq 1 + \mathbb{E}[\Phi^p] < \infty. \tag{2.46}
$$

Setting $p = 2$, convergence of $\mathbb{E}[\Phi^2]$ implies that the q^{th} moment of Φ is always defined and convergent at least for $0 \le q \le 2$. In general, for $q < 0$, the q^{th} moment of a positive random variable Φ with convergent second moment will not necessarily converge; it will converge if and only if $\rho(u)$ behaves sufficiently well near $u = 0$. However, in the particular case of the random variables $M_f(A)$ in Liouville theory, in order for $M_f(A)$ to be extremely small requires $X(x)$ to be very negative throughout the region A. Because of the constraint that the average of X on the unit circle vanishes, the Gaussian free field action $I_{\text{red}}(X)$ becomes extremely large if $X(x)$ is very negative in a region A. As a result, the measure $\rho(u)$ vanishes faster than any power of u for $u \to 0$ and the moments $E[M_f(A)^q]$ are convergent also for negative q. Thus $M_f(A)$ has a q^{th} moment for all $q \leq 2$, including $q < 0$. To learn what happens for $q > 2$ requires more precise arguments that are presented in sections [2.6](#page-18-0) and [3.](#page-24-0)

In view of eqn. (2.26) , we have

$$
\mathbb{E}[:e^{2bX_{\epsilon}(x)} : :e^{2bX_{\epsilon}(y)} :] \leq \min(|x-y|^{-4b^2}, \epsilon^{4b^2}), \tag{2.47}
$$

with equality if $|x - y| \geq 2\epsilon$. This implies that

$$
\lim_{\epsilon \to 0} \mathbb{E}[M_{f,\epsilon}(A)^2] = \int d^2x d^2y f(x) f(y) \frac{1}{|x-y|^{4b^2}},
$$
\n(2.48)

since the region with $|x - y| < 2\epsilon$ makes a negligible contribution in the limit $\epsilon \to 0$. The right hand side is what one would naively write as the two-point function of $M_f(A)$ $\int_A d^2x f(x)$: $e^{2bX(x)}$. The more subtle fact that we are in the process of learning is that the quantity $M_f(A)$ that has this second moment can be defined as a random variable that has non-integer moments.

To show that the limit $\lim_{\epsilon \to 0} M_{f,\epsilon}(A)$ exists in the L^2 sense, we need to show that the family of random variables $M_{f,\epsilon}(A)$ is Cauchy in the L^2 sense, meaning that

$$
\lim_{\epsilon,\epsilon' \to 0} \mathbb{E}[(M_{f,\epsilon}(A) - M_{f,\epsilon'}(A))^2] = 0.
$$
\n(2.49)

Concretely the left hand side of eqn. [\(2.49\)](#page-17-0) is

$$
\int d^2x d^2y f(x) f(y) \mathbb{E}[(e^{2bX_{\epsilon}(x)}z - e^{2bX_{\epsilon'}(x)}z)(e^{2bX_{\epsilon}(y)}z - e^{2bX_{\epsilon'}(y)}z)],
$$
\n(2.50)

and we need to show that this vanishes for $\epsilon, \epsilon' \to 0$, say with $\epsilon > \epsilon'$. The integrand in [\(2.50\)](#page-17-1) vanishes if $|x - y| > 2\epsilon$ by virtue of eqn. [\(2.30\)](#page-12-3), while for $|x - y| < 2\epsilon$, this integrand is a sum of four terms, each of which is bounded above by $|x-y|^{-4b^2}$. Since $\int_{|x-y|<2\epsilon} d^2x |x-y|^{-4b^2} \sim \epsilon^{2-4b^2}$, the integral vanishes for $\epsilon, \epsilon' \to 0$ if $b < 1/\sqrt{2}$, as we aimed to show. The condition on b that makes the contribution from the region $|x-y| < 2\epsilon$ vanish as $\epsilon \to 0$ is the same as the condition that ensures convergence of the integral in [\(2.48\)](#page-17-2).

Up to this point, we have assumed that f is smooth and bounded. If instead we assume that f has a singularity $f(x) \sim \frac{1}{|x-x|}$ $\frac{1}{|x-x_0|^{4b\alpha}}$ for some $x_0 \in A$, only one thing changes in the preceding analysis. In order to have $\lim_{\epsilon \to 0} \mathbb{E}[M_{f,\epsilon}(A)^2] < \infty$, we need

$$
\int_{A \times A} d^2 x d^2 y \frac{1}{|x - x_0|^{4b\alpha}} \frac{1}{|y - x_0|^{4b\alpha}} \frac{1}{|x - y|^{4b^2}} < \infty.
$$
 (2.51)

Convergence of this integral for $x, y \to x_0$ is again equivalent to the condition that ensures that the family $M_{f,\epsilon}(A)$ is Cauchy for $\epsilon \to 0$. This convergence gives a new condition $8b\alpha + 4b^2 < 4$ or

$$
\alpha < \frac{1}{2b} - \frac{b}{2}.\tag{2.52}
$$

This is a stronger condition than the Seiberg bound $\alpha \leq \frac{Q}{2} = \frac{1}{2b} + \frac{b}{2}$ $\frac{b}{2}$. If the condition [\(2.52\)](#page-18-1) is satisfied, then the limit $M_f(A) = \lim_{\epsilon \to 0} M_{f,\epsilon}(A)$ exists in the L^2 sense and in particular

$$
\mathbb{E}[M_f(A)^2] = \lim_{\epsilon \to 0} \mathbb{E}[M_{f,\epsilon}(A)^2].
$$
\n(2.53)

We stress that the formulas for second moments obtained this way are not new. What we gain from this discussion is only a framework in which noninteger moments of $M_f(A)$ make sense.

Finally we can draw a conclusion about the existence of Liouville correlation functions. According to eqn. [\(2.11\)](#page-8-1), the Liouville correlator $\langle \prod_{i=1}^n e^{2\alpha_i \phi(x_i)} \rangle$ is the w^{th} moment of $M_f = \int_{\Sigma} d^2x f(x)e^{2bX(x)}$, with $w = Q - \sum_i \alpha_i$. The arguments that we have given up to this point show that the Liouville correlation functions are well-defined if $b < 1/\sqrt{2}$, $\alpha_i < \frac{1}{2b} - \frac{b}{2}$ $\frac{b}{2}$ for all *i*, and $w < 2$ or equivalently $Q - \sum_i \alpha_i < 2$. This is a nontrivial region of existence of the Liouville correlation functions, but it is not the best possible. To get the best possible results, one has to consider convergence in L^p for $p < 2$, rather than convergence in L^2 . We explain the relevant arguments in section [3.7.](#page-36-0)

First, though, we explain some relatively simple arguments showing that if certain inequalities are violated, these methods can not be used to define Liouville correlation functions. In section [3,](#page-24-0) it will become clear that these inequalities are optimal.

2.6 Where Things Break Down

In this section, we will explain that the probabilistic approach to Liouville theory that we have been describing fails if $b > 1$ or if certain inequalities on the moments considered are violated. The strategy we follow is similar to arguments in the mathematical literature – for example, see the proof of Proposition 3.1 in $[21]$ – except that, deferring the rigorous arguments to section [3,](#page-24-0) we explain how the key steps can be carried out using an operator product expansion.

We will need to make an operator product expansion in an unfamiliar situation with non-integer powers of the interaction, so perhaps we should recall the logic behind the operator product expansion. The idea of the operator product expansion is that any observable that can be defined in a small region A of spacetime can be expanded, in the limit that A becomes small, as an asymptotic series in local operators that can be defined in region A. The dominant contribution comes from the lowest dimension operator in region A whose symmetry properties enable it to appear in the expansion. In our application, the operators that can appear in the expansion are those with the right value of what is sometimes called the Liouville momentum, which is defined so that the operator $:e^{2\alpha X(x)}$. or the product of this with any polynomial in derivatives of X, has Liouville momentum α . The reasoning behind the operator product expansion is robust enough that one expects it to be valid, in a reasonable quantum field theory, for any reasonable observable that can be defined in region A, including the somewhat exotic observables that we will encounter.

As a first example of how the probabilistic approach to Liouville theory can break down, we will explain that the random distribution $:e^{2bX(x)} := \lim_{\epsilon \to 0} e^{2bX_{\epsilon}(x)}$: vanishes if $b > 1$. Of course, regardless of the value of b, the "operator" $:e^{2bX(x)} := \lim_{\epsilon \to 0} :e^{2bX_{\epsilon}(x)}$. has meaningful correlation functions, for instance $\mathbb{E}[e^{2bX(x)}\cdot] = 1$. If $b > 1$, however, the behavior of $:e^{2bX_{\epsilon}(x)}$: for $\epsilon \to 0$ is somewhat analogous to what happens in the toy example described in section [2.4](#page-12-0) of the normal ordered exponential of a standard Brownian motion. For small ϵ , a random variable $M_{f,\epsilon}(A) = \int_A d^2x f(x) :e^{2bX_{\epsilon}(x)}$: is close to zero with high probability and very large with a very small probability, in such a way that in the limit $\epsilon \to 0$, its correlation functions remain nonvanishing but the probability measure of $M_{f,\epsilon}(A)$ converges to a delta function at the origin. In such a situation, moments of $M_f(A)$ of noninteger order cannot be defined.

As discussed in section [2.4,](#page-12-0) to show that a positive random variable Φ vanishes, it suffices to find a positive number p such that $\mathbb{E}[\Phi^p] = 0$. In the case of $M_f(A) =$ $\int_A d^2x f(x)$: $e^{2bX(x)}$; we will show that the p^{th} moment vanishes for p slightly less than 1. The precise range of p for which the argument works will become clear. Of course, once one has $\mathbb{E}[M_f(A)^p] = 0$ (in the limit $\epsilon \to 0$) for some $p > 0$, it follows that $M_f(A) = 0$ as a random variable and all its moments vanish.

First of all, for $\Phi_1, \Phi_2, \ldots, \Phi_n > 0$ and $0 < p < 1$, one has

$$
\left(\sum_{i} \Phi_{i}\right)^{p} \le \sum_{i} \Phi_{i}^{p}.
$$
\n(2.54)

Applying this to $M_{f_i}(A)$ for a family of nonnegative functions f_i , and setting $f = \sum_i f_i$, we see that the p^{th} moment of $M_f(A)$ will vanish if that is true for each $M_{f_i}(A)$. So, writing any f as a sum of functions supported in small squares, we can reduce to the case that the support of f is a small square in Σ . We can also assume that f is a constant function, since replacing f by a constant $c > f$ will only make $M_f(A)$ bigger. Since $M_c(A)$ is just proportional to c, we also lose nothing essential if we set $c = 1$.

So finally the only case that we have to consider is the random variable $M(A)$ $\int_A d^2x e^{2bX(x)}$, where A is a square in the complex x-plane. Pick an integer q and decompose

A as the union of q^2 identical squares A_1, \cdots, A_{q^2} that are each smaller by a factor of q. Using eqn. (2.54) again, we get

$$
\mathbb{E}[M(A)^p] \le \sum_{i=1}^{q^2} \mathbb{E}[M(A_i)^p] = q^2 \mathbb{E}[M(A_{(q)})^p],\tag{2.55}
$$

where $A_{(q)}$ is any one of the little squares. In the last step we use the fact that $\mathbb{E}[M(A_i)^p]$ is independent of i.

We can choose $A_{(q)}$ so that as q increases, $A_{(q)}$ converges to a point $x_0 \in A$. This is the setting of the operator product expansion: asymptotically for $q \to \infty$, we can expect to expand any observable or product of observables supported in $A_{(q)}$, such as $M(A_{(q)})^p = \left(\int_{A_{(q)}} d^2x e^{2bX(x)}\right)^p$, in a series of local operators at the point x_0 . The operator of lowest dimension with the correct value of the Liouville momentum is simply $:e^{2bpX(x_0)}$; so the expansion takes the form

$$
M(A_{(q)})^p \sim C(q) : e^{2bpX(x_0)}: + \cdots,
$$
\n(2.56)

where the corrections are subleading for large q . A simple scaling argument determines the q dependence of the coefficient $C(q)$. The operator $:e^{2\alpha X(x)}$: in the Gaussian free field theory has holomorphic or antiholomorphic dimension^{[7](#page-20-0)} $\Delta_{\alpha} = -\alpha^2$ in energy units, so in two dimensions its overall scaling dimension is $-2\alpha^2$ in energy units or $+2\alpha^2$ in length units. So $M(A_{(q)}) = \int_{A_{(q)}} d^2x \; i e^{2bX(x)}$: has length dimension $2 + 2b^2$ (counting 2 for the integral) and $M(A_{(q)})^p$ has length dimension $(2+2b^2)p$. On the right hand side of eqn. [\(2.56\)](#page-20-1), the operator : $e^{2bpX(p)}$: has length dimension b^2p^2 . The difference in dimensions between the left and right determines the q-dependence of $C(q)$. Indeed, $1/q$ is a length parameter, since it determines the size of the region $A_{(q)}$. So the power of $1/q$ is the difference of length dimensions between the two sides, and a more precise form of the expansion [\(2.56\)](#page-20-1) is

$$
M(A_{(q)})^p = cq^{2b^2p^2 - (2+2b^2)p} : e^{2bpX(x_0)}: + \cdots
$$
\n(2.57)

where now the constant c is independent of q for $q \to \infty$ and again + \cdots denotes subleading terms. Now we take the expectation value of eqn. (2.57) , using $\mathbb{E}[:e^{2bpX(x_0)}:] = 1$. Replacing c with a slightly larger constant c' and taking q large enough that the subleading terms are negligible, we learn from eqn. [\(2.56\)](#page-20-1) that

$$
\mathbb{E}[M(A)^p] \le c' q^{2b^2 p^2 - (2+2b^2)p + 2} = c' q^{2(p-1)(b^2 p - 1)}.
$$
\n(2.58)

For $b > 1$ and p slightly less than 1, $2(p-1)(b^2p-1) < 0$, and therefore the validity of the inequality (2.58) for arbitrarily large q implies that

$$
\mathbb{E}[M(A)^p] = 0.\tag{2.59}
$$

⁷In the introduction, in a related calculation, we scaled using the Liouville theory formula for the dimensions of operators, namely $\hat{\Delta}_{\alpha} = -\alpha^2 + Q\alpha$. Here, as we have eliminated the Liouville zero-mode and X is just a Gaussian free field, it is more natural to scale using the Gaussian free field formula $\Delta_{\alpha} = -\alpha^2$. The two formulas differ by whether a scaling of spatial coordinates is accompanied by a shift of the field. Regardless, in either of the two calculations, the term linear in α cancels out and it does not matter which of the two formulas one uses.

As explained previously, this implies that $M(A) = 0$ for $b > 1$ and therefore that the methods that we have been reviewing in this article do not suffice to define Liouville theory for such values of b.

Liouville theory is believed to have a symmetry $b \leftrightarrow 1/b$, implying that the theory for $b > 1$ is actually equivalent to the theory for $b < 1$. But it requires a different approach to define the theory directly for $b \geq 1$. A short review of what has been said about this mathematically can be found in [\[21\]](#page-41-8), section 6. At $b = 1$, rather than a normal ordered version of $e^{2bX(x)}$, one considers [\[27\]](#page-41-9), as one might possibly expect from the reasoning in [\[6\]](#page-40-10), a normal ordered version of $-X(x)e^{2X(x)} = -\frac{1}{2}$ 2 ∂ $\frac{\partial}{\partial b}\Big|_{b=1}e^{2bX(x)}$. To be more precise, one considers $-\frac{1}{2}$ 2 ∂ $\frac{\partial}{\partial b}\Big|_{b=1}$: $e^{2bX(x)} := -\frac{1}{2}$ 2 ∂ $\frac{\partial}{\partial b}\big|_{b=1}$ $\lim_{\epsilon \to 0} e^{2bX_{\epsilon}(x)-2b^2\mathbb{E}[X_{\epsilon}^2]}$ = $\lim_{\epsilon \to 0} \left[\left(2\mathbb{E}[X_{\epsilon}(x)^{2}] - X_{\epsilon}(x) \right) e^{2X_{\epsilon}(x)-2\mathbb{E}[X_{\epsilon}(x)^{2}]} \right]$. Note that although $-\frac{1}{2}$ 2 $\frac{\partial}{\partial b}e^{2bX(x)}$ is negative classically, it turns out to be positive quantum mechanically at $b = 1$. The mathematical theory for $b > 1$ is more complicated.

Henceforth we assume $b < 1$. In section [2.5,](#page-16-0) we were able to define moments $\mathbb{E}[M_f(A)^p]$ for a certain range of p assuming $b < 1/\sqrt{2}$, and those results and limits will be improved in section [3.](#page-24-0) Here we will prove that the moments of $M_f(A)$ do not exist if p is too large. The results are best possible, as will become clear in section [3,](#page-24-0) and are related to a nonperturbative pole of the DOZZ formula.

First we will prove that $\mathbb{E}[M_f(A)^p]$ diverges if $p > 1/b^2$. This was already explained in the introduction for the case that p is an integer. The statement actually holds whether p is an integer or not.

Since $M_f(A)$ is nonnegative for any nonnegative f and can only increase as f increases, to show that it diverges, it suffices to show that it diverges for the case that A is a small square and f is a constant function, which we may as well take to be $f = 1$. Again we decompose A as the union of q^2 little squares A_i that are each translates of $[0,1/q]^2$. Since we assume $b < 1$ and we want to investigate what happens for $p > 1/b^2$, we can assume $p > 1$. Since

$$
\left(\sum_{i} \Phi_{i}\right)^{p} \ge \sum_{i} \Phi_{i}^{p} \tag{2.60}
$$

for $\Phi_i \geq 0$ and $p > 1$, we have an inequality similar to that of eqn. [\(2.55\)](#page-20-4) but going in the opposite direction:

$$
\mathbb{E}[M(A)^p] \ge \sum_{i=1}^{q^2} \mathbb{E}\left[\sum_i M(A_i)^p\right] = q^2 \mathbb{E}\left[M(A_{(q)})^p\right],\tag{2.61}
$$

where again $A_{(q)}$ is one of the little squares. For large q, we again make the operator product expansion [\(2.57\)](#page-20-2), leading to the same inequality as in [\(2.58\)](#page-20-3) but in the opposite direction:

$$
\mathbb{E}[M(A)^p] \ge c q^{2b^2 p^2 - (2+2b^2)p + 2}.
$$
\n(2.62)

From the fact that this inequality holds for arbitrarily large q, we learn that $\mathbb{E}[M(A)^p] =$ $+\infty$ if the exponent $2(p-1)(b^2p-1)$ is positive. Given that $p > 1$ (as assumed in the derivation), this inequality is satisfied precisely if $p > 1/b^2$, so the moments diverge in that range of p.

Now, let us assume that $p > 1$ is such that the moments converge when f is bounded, and ask what happens in the presence of a singularity^{[8](#page-22-0)} in f created by one of the Liouville vertex operators, say $f = |x|^{-4b\alpha}$ for some α near $x = 0$. For any $\eta > 0$, we can write

$$
M_f(A) = \int_{|x| \le \eta} d^2x f(x) : e^{2bX(x)}: + \int_{|x| \ge \eta} d^2x f(x) : e^{2bX(x)}: .
$$
 (2.63)

Because for $\Phi_1, \Phi_2 > 0$, $(\Phi_1 + \Phi_2)^p \leq 2^p(\Phi_1^p + \Phi_2^p)$, to get a divergence in $\mathbb{E}[M_f(A)^p]$, there must be a divergence if $M_f(A)$ is replaced by one of the two terms on the right hand side of eqn. [\(2.63\)](#page-22-1). Since we assume that p is such that the p^{th} moment of $M_f(A)$ converges if f is bounded, the second term on the right hand side of eqn. (2.63) has convergent p^{th} moment and any divergence will come from the first term. Similarly, adding to f a bounded function will not affect whether there is a new divergence in the p^{th} moment associated to the singularity in f. Hence, we can assume that f is precisely equal to $|x|^{-4b\alpha}$ in some disc around the origin, say of radius η_0 . Since restricting to a smaller region will make $M_f(A)$ smaller, we have for any $\eta \leq \eta_0$

$$
\mathbb{E}[M_f(A)^p] \ge \mathbb{E}[M_\eta^p],\tag{2.64}
$$

where

$$
M_{\eta} = \int_{|x| \le \eta} d^2 x |x|^{-4b\alpha} :e^{2bX(x)}: . \tag{2.65}
$$

So we will have $\mathbb{E}[M_f(A)^p] = +\infty$ if $\mathbb{E}[M^p_\eta]$ diverges for $\eta \to 0$ (in which case the same reasoning shows that $\mathbb{E}[M_{\eta}^p]$ is infinite even before taking $\eta \to 0$).

We can use the operator product expansion to determine the behavior of M_{η}^p for $\eta \to 0$. As $\eta \to 0$, M_{η}^{p} is an observable supported in a very small neighborhood of $x = 0$, so it has an asymptotic expansion in local operators at $x = 0$. The lowest dimension such operator that has the same Liouville momentum as M_{η}^{p} is $e^{2bpX(0)}$, with length dimension $2b^{2}p^{2}$. Since the length dimension of M_{η}^{p} is $p(2-4b\alpha+2b^{2})$, where we include 2 from the integral and $-4b\alpha$ from the explicit factor $|x|^{-4b\alpha}$, the form of the expansion is

$$
M_{\eta}^{p} \sim C\eta^{p(2-4b\alpha+2b^{2})-2p^{2}b^{2}}: e^{2bpX(0)}: + \cdots,
$$
\n(2.66)

where C is a constant, independent of η , and the omitted terms are subleading as $\eta \to 0$. Taking expectation values, using $\mathbb{E}[e^{2bpX(0)}\cdot] = 1$, and replacing C by a slightly larger constant so that subleading terms in eqn. (2.66) can be dropped for sufficiently small η , we get a more explicit version of the inequality [\(2.64\)](#page-22-3):

$$
\mathbb{E}[M_f(A)^p] \ge C\eta^{p(2-4b\alpha+2b^2)-2p^2b^2} = C\eta^{2p(1-2b\alpha+b^2-pb^2)}.
$$
\n(2.67)

So $\mathbb{E}[M_f(A)^p] = +\infty$ when the exponent on the right hand side of this inequality is negative, or in other words when

$$
p > \frac{1}{b^2} + 1 - \frac{2\alpha}{b}.\tag{2.68}
$$

⁸We will consider the case of just one such singularity; if there are multiple singularities, they can be treated independently by the same reasoning that we are about to explain.

For integer p, this was deduced in the introduction.

As asserted in eqn. [\(1.5\)](#page-3-3), Liouville correlation functions are moments of the Liouville interaction of order $w = (Q - \sum_i \alpha_i)/b$ (times some known factors), where $Q = b + 1/b$. So the preceding results imply that the Liouville correlators diverge, even after removing an explicit factor that contains the perturbative poles, unless

$$
\sum_{i} \alpha_i > b \tag{2.69}
$$

and

$$
\sum_{i} \alpha_i > 2\alpha_k \quad \text{for all } k. \tag{2.70}
$$

These constraints are the strongest possible, since in section [3,](#page-24-0) we will see that $\mathbb{E}[M_f(A)^p]$ < ∞ if they are satisfied along with the Seiberg bound $\alpha_i < Q/2$.

Note that in the case of the three-point function, eqn. [\(2.70\)](#page-23-1) is a triangle inequality, asserting that $\alpha_1 + \alpha_2 > \alpha_3$, and cyclic permutations. In the semiclassical regime of small b, the triangle inequality has a semiclassical interpretation, explained in [\[28\]](#page-41-10), in terms of complex saddle points of the classical Liouville equation.

2.7 Comparison to the DOZZ Formula

The results (2.69) and (2.70) can be compared to the DOZZ formula, whose general form is as follows. By virtue of the $SL(2,\mathbb{C})$ symmetry of a Riemann surface of genus 0, the Liouville three-point function in genus 0 has the general form

$$
\left\langle \prod_{i=1}^{3} :e^{2\alpha_i(x_i)}: \right\rangle = \frac{C(\alpha_1, \alpha_2, \alpha_3)}{|x_{12}|^{2(\widehat{\Delta}_1 + \widehat{\Delta}_2 - \widehat{\Delta}_3)}|x_{13}|^{2(\widehat{\Delta}_1 + \widehat{\Delta}_3 - \Delta_2)}|x_{23}|^{2(\widehat{\Delta}_2 + \widehat{\Delta}_3 - \widehat{\Delta}_1)}},\tag{2.71}
$$

where $\Delta_i = \alpha_i(Q - \alpha_i)$, $x_{ij} = x_i - x_j$, and $C(\alpha_1, \alpha_2, \alpha_3)$ does not depend on the x_i . The DOZZ formula is a formula for the function $C(\alpha_1, \alpha_2, \alpha_3)$:

$$
C(\alpha_1, \alpha_2, \alpha_3) = \left[\pi \mu \gamma (b^2) b^{2-2b^2}\right]^{(Q-\sum \alpha_i)/b}
$$

$$
\times \frac{\Upsilon_0 \Upsilon_b (2\alpha_1) \Upsilon_b (2\alpha_2) \Upsilon_b (2\alpha_3)}{\Upsilon_b (\alpha_1 + \alpha_2 + \alpha_3 - Q) \Upsilon_b (\alpha_1 + \alpha_2 - \alpha_3) \Upsilon_b (\alpha_2 + \alpha_3 - \alpha_1) \Upsilon_b (\alpha_1 + \alpha_3 - \alpha_2)}.
$$
(2.72)

Here $\Upsilon_b(x)$, for $\text{Re } b > 0$, is an entire function of x that has simple zeroes at (and only at) $x = -m/b - nb$ and $x = (m + 1)/b + (n + 1)/b$, for any non-negative integers m, n; $\Upsilon_0 = d\Upsilon_b(x)/dx|_{x=0}$; and $\gamma(x) = \Gamma(x)/\Gamma(1-x)$.

Zeroes of the denominator of the DOZZ formula lead to poles of the Liouville threepoint function. In particular, the perturbative poles found by Goulian and Li [\[5\]](#page-40-8) and described in the introduction result from the zeroes of $\Upsilon_b(\alpha_1 + \alpha_2 + \alpha_3 - Q)$ at $\sum_i \alpha_i - Q =$ $-nb$. The conditions that we have found in eqns. [\(2.69\)](#page-23-2) and [\(2.70\)](#page-23-1) correspond to some of the nonperturbative poles of the DOZZ formula. The condition [\(2.69\)](#page-23-2) is associated to the pole of the DOZZ formula at $\sum_i \alpha_i - Q = -\frac{1}{b}$ $\frac{1}{b}$, and the condition [\(2.70\)](#page-23-1) is associated to the poles of the DOZZ formula at $\alpha_1 + \alpha_2 - \alpha_3 = 0$ or permutations thereof.

After integrating over the Liouville zero-mode, the remaining integral over the Gaussian free field X has an extended region of convergence, as remarked in the introduction. This extended region is bounded by the conditions [\(2.69\)](#page-23-2) and [\(2.70\)](#page-23-1) along with the Seiberg bound $\alpha_i \leq Q/2$ for all i. What happens at the Seiberg bound is completely different from what happens at the other boundaries. At the Seiberg bound, the DOZZ formula has not a pole but a zero, resulting from the zero of $\Upsilon_b(2\alpha)$ at $\alpha = 0$. We have interpreted the conditions [\(2.69\)](#page-23-2) and [\(2.70\)](#page-23-1) in terms of ultraviolet divergences, which lead to poles of the DOZZ formula. What can account for a zero of the DOZZ formula?

In fact, suppose that we increase one of the Liouville momenta in the three-point function, say α_1 , until the Seiberg bound is violated. To avoid the pole at $\alpha_1 = \alpha_2 + \alpha_3$, we maintain the triangle inequality $\alpha_1 < \alpha_2 + \alpha_3$. So by the time one reaches the Seiberg bound at $\alpha_1 = Q/2$, we will have $\sum_i \alpha_i > Q$ and therefore $w = (Q - \sum_i \alpha_i)/b < 0$. This leads to vanishing of the DOZZ formula for the following reason. The Liouville three-point function is, as in eqn. [\(2.11\)](#page-8-1), a multiple of $\mathbb{E} \left[\left(\int d^2x f(x)e^{2bX(x)} \right)^w \right]$, with $f(x) \sim |x-x_1|^{-4b\alpha_1}$ for $x \to x_1$. In the probabilistic approach that is reviewed in the present article, the interpretation of the Seiberg bound is that if $\alpha \geq Q/2$, then, for any $\eta > 0$, the quantity $\int_{|x-x_1|<\eta} d^2x |x-x_1|^{-4b\alpha} e^{2bX(x)}$ is equal to $+\infty$ with probability 1. For the proof of this assertion, see section [3.6.](#page-35-0) Because this quantity equals $+\infty$ with probability 1, its negative powers vanish with probability 1, implying that $\mathbb{E}\left[\left(\int d^2x f(x)e^{2bX(x)}\right)^w\right]=0$ for $w<0$. This is a probabilistic interpretation of the vanishing of the DOZZ formula at the Seiberg bound.

The DOZZ formula shows that the Liouville correlation function can be analytically continued past the Seiberg bound (as was actually anticipated in [\[6\]](#page-40-10)). However, because the DOZZ formula has a simple zero at the Seiberg bound, the three-point function, continued so that one of the Liouville momenta slightly exceeds $Q/2$, becomes negative, contradicting the positivity that appears to follow from the definition of the correlation function as $\langle \prod_i :e^{2\alpha_i \phi(x_i)} \rangle$ and showing that the correlation functions analytically continued past the Seiberg bound do not have a simple probabilistic interpretation. Positivity is similarly lost upon analytic continuation past the simple poles that mark the other boundaries of the extended region of convergence of the path integral. Beyond the other boundaries of the extended region of convergence are the many other nonperturbative poles of the DOZZ formula, whose interpretation from a probabilistic point of view is not clear.

3 Liouville Theory Correlators: Detailed Arguments

In the rest of this article, we will sketch more detailed rigorous mathematical arguments about Liouville theory. The arguments are mostly adapted from the existing literature, though the treatment of certain singular integrals in section [3.5](#page-31-0) may be new.

First we explain two useful tools: the near scale invariance of the Gaussian free field (section [3.1\)](#page-25-0) and an inequality due to Kahane $[13]$ (section [3.2\)](#page-26-0). Then in section [3.3,](#page-28-0) we show that in the limit $\epsilon \to 0$ the random variables $M_{f,\epsilon}(A)$ of Liouville theory vanish if $b > 1$. Such a result was originally obtained in [\[13\]](#page-40-3) (with a different cutoff). Subsequently, we assume that $b < 1$ and we consider the behavior of the moments $\mathbb{E}[M_{f,\epsilon}(A)^p]$ with f

assumed initially to be bounded. In section [3.4,](#page-29-0) we show that these moments diverge in the limit $\epsilon \to 0$ if $p > 1/b^2$ but converge to finite, positive values if $1 \leq p < 1/b^2$ (also a result of [\[13\]](#page-40-3)). In section [3.5,](#page-31-0) we consider the case that $f(x)$ has a power law singularity^{[9](#page-25-1)} $1/|x-x_0|^{2\alpha b}$ for some α and again determine the range of p for which $\mathbb{E}[M_{f,\epsilon}(A)^p]$ converges for $\epsilon \to 0$, as first analyzed in [\[18\]](#page-41-11). In section [3.6,](#page-35-0) we show that $\lim_{\epsilon \to 0} M_f(A) = \infty$ with probability 1 if f has a singularity with α exceeding the Seiberg bound at $\alpha = Q$. As already explained in section [2.7,](#page-23-0) this accounts for the vanishing of the DOZZ formula at the Seiberg bound.

The results summarized up to this point do not address the question of whether, say for bounded f, the random variables $\mathbb{E}[M_{f,\epsilon}(A)]$ have nonzero limits for $\epsilon \to 0$. In general, a family of random variables $M_{f,\epsilon}(A)$ can converge to 0 for $\epsilon \to 0$, even while some moments remain nonzero. As explained in section [2.4,](#page-12-0) this can happen if, for very small ϵ , the nonzero moments of $M_{f,\epsilon}(A)$ reflect the contributions of very rare events in which $M_{f,\epsilon}(A)$ is unexpectedly large. In section [3.7,](#page-36-0) we show that this does not happen and that for all $b < 1$, a nontrivial Liouville probability measure does exist in the limit $\epsilon \to 0$. This is accomplished by showing that (for bounded f), the random variables $M_{f,\epsilon}(A)$ converge in L^1 to nonzero limits as $\epsilon \to 0$. In general, a positive martingale that converges in L^1 and has a convergent p^{th} moment also converges in L^p , so it follows^{[10](#page-25-2)} that (for bounded f), $M_{f,\epsilon}(A)$ converges in L^p .

3.1 Near Scale Invariance

In section [2,](#page-6-0) we defined the Gaussian free field so that for $|y|, |y'| < 1$, the two-point correlation function is just

$$
\mathbb{E}[X(y)X(y')] = \log \frac{1}{|y - y'|}.\tag{3.1}
$$

This immediately implies that for $q > 1$,

$$
\mathbb{E}[X(y/q)X(y'/q)] = \log \frac{q}{|y-y'|} = \mathbb{E}[X(y)X(y')] + \log q.
$$
 (3.2)

Thus, $X(y/q)$ has the same distribution as $X(y) + \Omega_q$, where Ω_q is a Gaussian random variable of mean zero and variance $\log q$, and independent of $X(y)$. Hence for any function $F,$

$$
\mathbb{E}[F(X(y/q))] = \mathbb{E}[F(X(y) + \Omega_q)].
$$
\n(3.3)

⁹Unfortunately, there is a clash of notation between what has become standard in the physics and mathematics literatures on Liouville theory. What has been called α in the math literature is 2α in the physics literature. To facilitate comparison of the following formulas with the mathematics literature, we define $\alpha = 2\alpha$, where the usual Liouville primary fields are denoted as $e^{2\alpha\phi(x)}$ in the physics literature and as $e^{\alpha\phi(x)}$ in the math literature. Up to this point, all formulas have been written in terms of α ; henceforth we use α . Another detail is that, as remarked in footnote [1,](#page-2-0) the mathematics literature is mostly written in terms of the variable $\gamma = 2b$.

¹⁰Here we are tacitly assuming that $\lim_{\epsilon \to 0} M_{f,\epsilon}(A)$ is equivalent to a similar limit defined using Kahane's cutoff, which gives the martingale property. This can plausibly be proved using Kahane's inequality, though we will not do so.

In general, we have to consider regularized versions of various expressions. If instead of $X(y)$ we consider its circle regularized version $X_{\epsilon}(y)$, then a formula similar to eqn. [\(3.1\)](#page-25-3) holds, except that we have to rescale ϵ along with the other lengths:

$$
\mathbb{E}[X_{\epsilon/q}(y/q)X_{\epsilon/q}(y'/q)] = \mathbb{E}[X_{\epsilon}(y)X_{\epsilon}(y')] + \log q.
$$
\n(3.4)

Therefore the cutoff version of eqn. [\(3.3\)](#page-25-4) is

$$
\mathbb{E}[F(X_{\epsilon/q}(y/q))] = \mathbb{E}[F(X_{\epsilon}(y) + \Omega_q)].
$$
\n(3.5)

In practice, since two different values of the cutoff parameter appear in eqn. [\(3.5\)](#page-26-1), to make use of this formula, we frequently need to know how to compare the theories with different cutoffs. It turns out that this can be done quite effectively using eqn. [\(2.23\)](#page-11-0), which asserts that $\mathbb{E}[X_{\epsilon}(x)X_{\epsilon}(y)]$ is monotonically increasing as ϵ becomes smaller.

Kahane [\[13\]](#page-40-3) considered a Gaussian free field that has a two-point function assumed to be of the form

$$
\mathbb{E}[X(y)X(y')] = \sum_{i=1}^{\infty} g_i(y, y'),\tag{3.6}
$$

where the functions $g_i(y, y')$ are positive pointwise and also are positive as kernels, in the sense that for real-valued functions f, the quadratic form $Q_i(f) = \int d^2y d^2y' f(y)g_i(y, y') f(y')$ is positive-definite. For suitable g_i , the logarithmic correlations of Liouville theory can be expressed as in eqn. [\(3.6\)](#page-26-2). We made a similar expansion in section [2.4,](#page-12-0) but in terms of the eigenfunctions $f_i(y)$ of the Laplace operator, which are not positive; in Kahane's approach, positivity is important. Positivity of the quadratic forms makes it possible to define Gaussian variables $X_{(i)}$ with $\mathbb{E}[X_{(i)}(y)X_{(i)}(y')] = g_i(y, y')$ (and with $X_{(i)}$ independent of $X_{(i')}$) for $i \neq i'$, and then if one sets $X_n = \sum_{i=1}^n X_{(i)}$, one has

$$
\mathbb{E}[X_n(y)X_n(y')] = \sum_{i=1}^n g_i(y, y').
$$
\n(3.7)

Pointwise positivity of the functions $g_i(y, y')$ ensures that $\mathbb{E}[X_n(y)X_n(y')]$ is an increasing function of n , analogous to the fact that with the cutoff based on circle averaging, $\mathbb{E}[X_{\epsilon}(x)X_{\epsilon}(y)]$ monotonically increases for $\epsilon \to 0$.

So in either approach, the two-point function increases as the cutoff is removed. This has important implications when combined with Kahane's inequality, which we describe next.

3.2 Kahane's Inequality

Let X_i and Y_i , for $i = 1, \ldots, n$ be two families of independent, centered Gaussian random Let X_i and Y_i , for $i = 1, ..., n$, be two families of independent, centered Gaussian variables and let c_i , $i = 1, ..., n$, be positive numbers.^{[11](#page-26-3)} Set $Z_i(t) = \sqrt{t}X_i +$ √ $1-tY_i,$ $0 \le t \le 1$. Of course, for each t the Z_i are again independent Gaussian random variables.

¹¹In explaining Kahane's inequality, we follow the appendix of $[21]$.

Let $\phi : \mathbb{R}_+ \to \mathbb{R}$ be a twice differentiable function with at most polynomial growth at infinity, and define

$$
\varphi(t) = \mathbb{E}\left[\phi\left(\sum_{i} c_i : e^{Z_i(t)}:\right)\right].\tag{3.8}
$$

Then a straightforward calculation reveals a surprisingly simple and useful result:

$$
\frac{d\varphi(t)}{dt} = \frac{1}{2} \sum_{i,j=1}^{n} c_i c_j (\mathbb{E}[X_i X_j] - \mathbb{E}[Y_i Y_j]) \mathbb{E}\left[:e^{Z_i(t) + Z_j(t)} \colon \phi''\left(\sum_{k=1}^{n} c_k :e^{Z_k(t)}: \right) \right].
$$
 (3.9)

Suppose then that

$$
\mathbb{E}[X_i X_j] \ge \mathbb{E}[Y_i Y_j], \quad 1 \le i, j \le n. \tag{3.10}
$$

Then if the function ϕ is convex, $\phi'' \geq 0$, we have $\varphi(1) \geq \varphi(0)$ or

$$
\mathbb{E}\left[\phi\left(\sum_{i}c_{i}:e^{X_{i}}\right)\right] \geq \mathbb{E}\left[\phi\left(\sum_{i}c_{i}:e^{Y_{i}}\right)\right],\tag{3.11}
$$

while if ϕ is concave, $\phi'' \leq 0$, we have the opposite inequality

$$
\mathbb{E}\left[\phi\left(\sum_{i}c_{i}:e^{X_{i}}\right)\right] \leq \mathbb{E}\left[\phi\left(\sum_{i}c_{i}:e^{Y_{i}}\right)\right].
$$
\n(3.12)

Specializing to moments,

$$
\mathbb{E}\left[\left(\sum_{i} c_{i} : e^{X_{i}}\right)^{p}\right] \geq \mathbb{E}\left[\left(\sum_{i} c_{i} : e^{Y_{i}}\right)^{p}\right], p \geq 1 \tag{3.13}
$$

and

$$
\mathbb{E}\left[\left(\sum_{i} c_{i} : e^{X_{i}}\right)^{p}\right] \leq \mathbb{E}\left[\left(\sum_{i} c_{i} : e^{Y_{i}}\right)^{p}\right], \quad 0 \leq p \leq 1. \tag{3.14}
$$

Equation [\(2.23\)](#page-11-0) implies that $\mathbb{E}[X_{\epsilon}(y)X_{\epsilon}(y')] \geq \mathbb{E}[X_{\epsilon}(x)X_{\epsilon}(y)]$ if $\epsilon' \leq \epsilon$. Hence a continuum limit of the inequalities [\(3.13\)](#page-27-0) and [\(3.14\)](#page-27-1) gives for $\epsilon' \leq \epsilon$, and any non-negative function f,

$$
\mathbb{E}\left[\left(\int_{A} d^{2}x f(x) \, :e^{2bX_{\epsilon'}(x)}:\right)^{p}\right] \geq \mathbb{E}\left[\left(\int_{A} d^{2}x f(x) \, :e^{2bX_{\epsilon}(x)}:\right)^{p}\right], \quad p \geq 1\tag{3.15}
$$

and

$$
\mathbb{E}\left[\left(\int_A \mathrm{d}^2 x f(x) \, : \! e^{2bX_{\epsilon'}(x)} \! : \right)^p\right] \le \mathbb{E}\left[\left(\int_A \mathrm{d}^2 x f(x) \, : \! e^{2bX_{\epsilon}(x)} \! : \right)^p\right], \quad 0 \le p \le 1. \tag{3.16}
$$

Similar inequalities hold with Kahane's cutoff, since the cutoff two-point function [\(3.7\)](#page-26-4) is an increasing function of n. Kahane's used these inequalities as follows. There are many possible choices of the positive functions $g_i(x, x')$ in the decomposition (3.6) and obviously the cutoff version based on truncating this decomposition to a finite sum does depend on the choice of these functions. Kahane used his inequality to prove that the large n limit is universal, independent of the choice of the $g_i(x, x')$.

3.3 Triviality For $b > 1$

Having developed the tools, our first goal is to show that Liouville theory, defined as we have assumed by simple normal ordering of the interaction, is trivial for $b > 1$. More precisely, with $M_{f,\epsilon}(A) = \int_A d^2x f(x) :e^{2bX_{\epsilon}(x)}$; we want to show that for any non-negative f and A, and for p slightly less than 1, $\lim_{\epsilon \to 0} \mathbb{E}[M_{f,\epsilon}(A)^p] = 0$. A priori, since $M_{f,\epsilon}(A) \ge 0$ and $\mathbb{E}[M_{f,\epsilon}(A)] = \int_A d^2x f(x)$, we have (with the help of eqn. [\(2.46\)](#page-17-3)) $0 \leq \mathbb{E}[M_{f,\epsilon}(A)^p] \leq$ $1 + \int_A d^2x f(x)$ for $0 \le p \le 1$. Moreover, $\mathbb{E}[M_{f,\epsilon}(A)^p]$ decreases as $\epsilon \to 0$ in view of eqn. [\(3.16\)](#page-27-2). Hence it has a limit for $\epsilon \to 0$ that is either positive or zero. We will show that the limit is 0.

The following proof that the limit vanishes is a rigorous counterpart of the explanation in section [2.6.](#page-18-0) As before, it is enough to consider the case that $f = 1$ and $A = [0, 1]^2$. We again pick an integer $q > 1$ and decompose A as the union of q^2 squares A_i that are all translates of $[0,1/q]^2$. Since $\mathbb{E}[M_{f,\epsilon}(A_i)^p]$ is certainly independent of i, we have for $p<1$

$$
\mathbb{E}\left[\left(\int_{[0,1]^2} d^2x :e^{2bX_{\epsilon}(x)}:\right)^p\right] \leq \sum_{i} \mathbb{E}\left[\left(\int_{A_i} d^2x :e^{2bX_{\epsilon}(x)}:\right)^p\right]
$$

$$
= q^2 \mathbb{E}\left[\left(\int_{[0,1/q]^2} d^2x :e^{2bX_{\epsilon}(x)}:\right)^p\right].
$$
(3.17)

Setting $x = y/q$ and using [\(3.5\)](#page-26-1), we get

$$
\mathbb{E}\left[\left(\int_{[0,1]^2} d^2x \, i e^{2bX_{\epsilon}(x)}\, \mathrm{d}x\right)^p\right] \leq q^{2-2p} \mathbb{E}\left[\left(\int_{[0,1]^2} d^2y \, i e^{2bX_{\epsilon}(y/q)}\, \mathrm{d}x\right)^p\right] \\
= q^{2-2p} \mathbb{E}\left[\left(\int_{[0,1]^2} d^2y \, i e^{2bX_{q\epsilon}(y)+2b\Omega_q}\, \mathrm{d}x\right)^p\right].\n\tag{3.18}
$$

Explicitly, $:e^{2bX_{q\epsilon}(y)+2b\Omega_q}$: $= e^{2b\Omega_q-2b^2\mathbb{E}[\Omega_q^2]}$ $:e^{2bX_{q\epsilon}(y)}$:, so

$$
\left(\int_{[0,1]^2} d^2 y : e^{2bX_{q\epsilon}(y) + 2b\Omega_q} : \right)^p = e^{2bp\Omega_q - 2b^2 p \mathbb{E}[\Omega_q^2]} \left(\int_{[0,1]^2} d^2 y : e^{2bX_{q\epsilon}(y)} : \right)^p, \tag{3.19}
$$

and therefore, as $\mathbb{E}[\Omega_q^2] = \log q$, eqn. [\(3.18\)](#page-28-1) becomes

$$
\mathbb{E}\left[\left(\int_{[0,1]^2} d^2x \, :e^{2bX_\epsilon(x)}: \right)^p\right] \le q^{2-2p+2b^2p^2-2b^2p} \mathbb{E}\left[\left(\int_{[0,1]^2} d^2x \, :e^{2bX_{q\epsilon}(x)}: \right)^p\right] \tag{3.20}
$$

The prefactor $q^{2-2p+2b^2p^2-2b^2p} = q^{2(p-1)(b^2p-1)}$ is familiar from eqn. [\(2.58\)](#page-20-3). As $q > 1$, for $b > 1$ one can pick $p < 1$ so that $q^{2(p-1)(b^2p-1)} < 1$. Having done so, take $\epsilon \to 0$ in eqn. (3.20) keeping q fixed. If the left hand side were to approach a positive limit as $\epsilon \to 0$, the expectation on the right hand side would approach the same limit, giving a contradiction. So the left hand side vanishes as $\epsilon \to 0$, as claimed.

3.4 Moments

Henceforth, we assume that $b < 1$, and we investigate the behavior of the moments $\mathbb{E}[M_{f,\epsilon}(A)^p]$ as $\epsilon \to 0$. In this section, we assume that the function f is smooth and bounded; the singular case is the subject of section [3.5.](#page-31-0)

Our first goal is to show, by a rigorous counterpart of the discussion in section [2.6,](#page-18-0) that the limit is $+\infty$ if $p > 1/b^2$. Since $b < 1$, $p > 1/b^2$ implies in particular that $p > 1$. So $\mathbb{E}[M_{f,\epsilon}(A)^p]$ increases as ϵ becomes smaller, according to eqn. [\(3.15\)](#page-27-3). Therefore for $\epsilon \to 0$, it converges either to a positive real number or to $+\infty$. We will show that the limit is $+\infty$ for $p > 1/b^2$ by a rigorous counterpart of the argument in section [2.6.](#page-18-0)

As before, since $M_{f,\epsilon}(A)$ becomes smaller if A is replaced by a smaller region or f by a smaller positive function, there is no loss in assuming that $f = 1$ and that A is a small square, which we may as well take to be the unit square $[0,1]^2$. Again we decompose A as the union of q^2 smaller squares all of size $1/q$. The first inequality in eqn. (3.17) now goes in the opposite direction, since now $p > 1$, so following the same steps now leads us to

$$
\mathbb{E}\left[\left(\int_{[0,1]^2} d^2x :e^{2bX_{\epsilon}(x)}:\right)^p\right] \geq q^{2-2p}\mathbb{E}\left[\left(\int_{[0,1]^2} d^2y :e^{2bX_{q\epsilon}(y)+2b\Omega_q}:\right)^p\right]
$$

$$
= q^{2(p-1)(b^2p-1)}\mathbb{E}\left[\left(\int_{[0,1]^2} d^2x :e^{2bX_{q\epsilon}(x)}:\right)^p\right].
$$
(3.21)

Since $q > 1$, if $p > 1/b^2$, we have $q^{2(p-1)(b^2p-1)} > 1$. Taking $\epsilon \to 0$ with fixed q, the inequality implies that the left hand side diverges for $\epsilon \to 0$, as claimed.

The next goal is to show that, with $b < 1$, the moments do converge for $p < 1/b^2$. If $p \leq 1$, eqn. [\(3.15\)](#page-27-3) shows that the p^{th} moment of $M_{f,\epsilon}(A)$ becomes smaller as ϵ decreases and therefore converges to a positive limit or zero. The fact that the limit is positive follows from the proof of L^1 convergence that we give in section [3.7.](#page-36-0) Here we consider the case $p > 1$, so that $\mathbb{E}[M_{f,\epsilon}(A)^p]$ increases as ϵ becomes smaller. We want to show that for $p < 1/b^2$, this moment approaches a finite limit rather than tending to infinity. The necessary argument, which is inspired by an argument by Bacry and Muzy [\[29\]](#page-41-12), is more elaborate than we have seen so far.

By replacing A with a possibly larger region and f with a possibly larger constant function, we can assume that f is a constant, which may as well be 1, and that A is a square, which may as well be the unit square $[0, 1]^2$. One preliminary point is that if p is an integer, the statement is true by direct calculation; the moment $\lim_{\epsilon \to 0} \mathbb{E}[M_{\epsilon}(A)^n]$ of integer order *n* is given by the integral in eqn. [\(1.7\)](#page-4-0), which converges if $n < 1/b^2$. Part of the proof will involve comparing what happens at p to what happens at an integer just less than p . So if p is not an integer, we let n be the integer that satisfies

$$
n - 1 < p < n,\tag{3.22}
$$

and the proof will make use of the fact that the $n-1$ th moment of $M_{\epsilon}(A)$ converges as $\epsilon \to 0$.

The first step is again to decompose A as the union of q^2 squares A_i each of size $1/q$, now taking q to be an even integer 2r. Let P be the set of $4r^2$ small squares. Partition P into four disjoint subsets \mathcal{P}_a , $1 \leq i \leq 4$, such that within each \mathcal{P}_a , the separation between any pair of small squares is positive. This can be done in such a way that each \mathcal{P}_a is a translate of \mathcal{P}_1 . Define

$$
S_{a,\epsilon} = \sum_{A_i \in \mathcal{P}_a} M_{\epsilon}(A_i),\tag{3.23}
$$

so that $M(A) = S_1 + S_2 + S_3 + S_4$. Then

$$
\mathbb{E}[M_{\epsilon}(A)^p] \le 4^{p-1} \sum_{a=1}^4 \mathbb{E}[S_a^p] = 4^p \mathbb{E}[S_{1,\epsilon}^p],\tag{3.24}
$$

where in the last step we use the fact that all S_a have the same distribution. Now with n as in eqn. (3.22) , we have

$$
S_{1,\epsilon}^p = \left(\sum_{A_i \in \mathcal{P}_1} M_{\epsilon}(A_i)\right)^p = \left[\left(\sum_{A_i \in \mathcal{P}_1} M_{\epsilon}(A_i)\right)^{p/n}\right]^n \le \left(\sum_{A_i \in \mathcal{P}_1} M_{\epsilon}(A_i)^{p/n}\right)^n, \quad (3.25)
$$

where in the last step we use that $p/n < 1$. Expanding this out, we get

$$
S_{1,\epsilon}^p \leq \sum_{A_1,\ldots,A_n \in \mathcal{P}_1} M_{\epsilon}(A_1)^{p/n} M_{\epsilon}(A_2)^{p/n} \cdots M_{\epsilon}(A_n)^{p/n}.
$$
 (3.26)

Decompose the right hand side as $U + V$, where U is the sum of all terms in which the A_i are all the same, and V is the sum of all other terms. As there are r^2 little squares in \mathcal{P}_1 and $M_{\epsilon}(A_i)$ is equally distributed for each one, we have

$$
\mathbb{E}[U] = r^2 \mathbb{E}\left[\left(\int_{[0,1/2r]^2} d^2x : e^{2bX_{\epsilon}(x)}:\right)^p\right]
$$

= $\frac{1}{4}(2r)^{2(p-1)(b^2p-1)} \mathbb{E}\left[\left(\int_{[0,1]^2} d^2y : e^{2bX_{2r\epsilon}(y)}:\right)^p\right].$ (3.27)

Here we have followed the same steps as before, setting $y = 2rx$, using eqn. [\(3.5\)](#page-26-1), and using $\mathbb{E}[e^{2b\Omega_{2r}}] = (2r)^{2bp^2 - 2bp}$. Because $p > 1$, Kahane's inequality (3.15) tells us that the right hand side of eqn. [\(3.27\)](#page-30-0) increases if we replace $X_{2r\epsilon}(y)$ by $X_{\epsilon}(y)$, and therefore

$$
\mathbb{E}[U] \le \frac{1}{4} (2r)^{2(p-1)(b^2p-1)} \mathbb{E}\left[\left(\int_{[0,1]^2} d^2 y : e^{2bX_{\epsilon}(y)}:\right)^p\right]
$$
\n
$$
= \frac{1}{4} (2r)^{2(p-1)(b^2p-1)} \mathbb{E}[M_{\epsilon}(A)^p].
$$
\n(3.28)

For $p \in (1, 1/b^2)$, the exponent in $(2r)^{2(p-1)(bp^2-1)}$ is negative, so we can choose a large r such that $C(b, p, r) = 4^{p-1}(2r)^{2(p-1)(b^2p-1)}$ is less than 1. As for the remainder V on the right hand side of eqn. [\(3.26\)](#page-30-1), since $p/n < 1$, we have $M_{\epsilon}(A_i)^{p/n} \leq 1 + M_{\epsilon}(A_i)$ for all i, and therefore

$$
V \le \sum_{A_1,\dots,A_n \in \mathcal{P}_1} \prod_{i=1}^n (1 + M_{\epsilon}(A_i)), \qquad (3.29)
$$

where the symbol \sum' represents a sum restricted to terms in which the A_i are not all the same. For each choice of A_1, A_2, \ldots, A_n , the corresponding contribution to the right hand side of eqn. (3.29) is the sum of 2^n terms, each of which is the product of at most n factors of $M_{\epsilon}(A_i)$ for various different $A_i \in \mathcal{P}_1$. Considering any one such term, let $A'_1, \ldots, A'_{n'}$, with $n' \leq n$, be the little squares that appear in that term. The contribution of that term to $\mathbb{E}[V]$ can be described as an explicit integral as in eqn. [\(1.7\)](#page-4-0):

$$
\mathbb{E}\left[\prod_{i=1}^{n'} M_{\epsilon}(A'_i)\right] = \int_{A'_1} \cdots \int_{A'_{n'}} d^2 x_1 \cdots d^2 x_{n'} \prod_{1 \le i < j \le n'} \prod_{i < j} \exp(4b^2 \mathbb{E}[X_{\epsilon}(x_i) X_{\epsilon}(x_j)]). \tag{3.30}
$$

We would like to have an upper bound on this integral that is independent of ϵ . Since $\mathbb{E}[X_{\epsilon}(x_i)X_{\epsilon}(x_j)]$ increases as ϵ becomes smaller, we can get an upper bound by simply setting $\epsilon = 0$, whereupon $\mathbb{E}[X_{\epsilon}(x_i)X_{\epsilon}(x_j)]$ reduces to log $\frac{1}{|x_i - x_j|}$ and eqn. [\(3.30\)](#page-31-1) becomes

$$
\mathbb{E}\left[\prod_{i=1}^{n'} M_{\epsilon}(A'_i)\right] \leq \int_{A'_1} \cdots \int_{A'_{n'}} d^2 x_1 \cdots d^2 x_{n'} \prod_{1 \leq i < j \leq n'} \frac{1}{|x_i - x_j|^{4b^2}}.\tag{3.31}
$$

As a consequence of the remark following eqn. [\(1.8\)](#page-4-1), the integral on the right hand side converges, because at most $n-1 < 1/b^2$ of the points x_i are contained in any one of the A'_{i} , and the distances between the different A'_{i} are positive. (It can happen that the A'_{i} are all the same, but this is only possible if $n' \leq n-1$.) Summing many such terms, V is then bounded above by a positive constant $D(b, p, r)$ that depends on b, p, and r but not on ϵ .

Finally, then, for $p \in (1, 1/b^2)$ we get an inequality of the form

$$
\mathbb{E}[M_{\epsilon}(A)^p] \le C(b, p, r)\mathbb{E}[M_{\epsilon}(A)^p] + D(b, p, r), \tag{3.32}
$$

where $C(b, p, r) < 1$ and both $C(b, p, r)$ and $D(b, p, r)$ depend only on b, p, and r and not ϵ . This implies an upper bound

$$
\lim_{\epsilon \to 0} \mathbb{E}[M_{\epsilon}(A)^p] \le \frac{D(b, p, r)}{1 - C(b, p, r)},
$$
\n(3.33)

showing that the left hand side remains bounded as $\epsilon \to 0$.

What happens for $p < 1$? For $0 < p \le 1$, the quantity $\mathbb{E}[M_{\epsilon}(A)^p]$ is bounded above by $1 + \mathbb{E}[M_{\alpha}(A)] = 2$ by virtue of eqn. [\(2.46\)](#page-17-3), and decreases as ϵ becomes smaller because of Kahane's inequality [\(3.16\)](#page-27-2). So its limit as $\epsilon \to 0$ certainly exists. In this range of p, the nontrivial question is whether the limit vanishes, as we know happens for $b > 1$. To show that this limit is nonvanishing for all $b < 1$ requires a fairly difficult argument that is explained in section [3.7.](#page-36-0)

3.5 Convergent Integrals of Singular Functions

In this section, we begin the analysis of the integral

$$
I(\alpha) = \int_{|x| \le 1} d^2 x |x|^{-2b\alpha} :e^{2bX(x)}; \qquad (3.34)
$$

with $\alpha > 0$. Note that this integral is always positive, but may potentially take the value $+\infty$. We will show that for $\alpha \in (b, Q)$ (where we recall that $Q = b + 1/b$), this does not happen, and in fact, for such α ,

$$
\mathbb{E}[I(\alpha)^p] < \infty \text{ for all } p \in (0, (Q - \alpha)/b). \tag{3.35}
$$

This was first shown in [\[18\]](#page-41-11). Before attempting a proof, we will first discuss some qualitative aspects of this statement. First of all, since the upper bound on α in this statement is $\alpha < b + 1/b$, in particular it is possible to have $\alpha \geq 1/b$. For such values of α , the $|x|^{-2b\alpha}$ singularity is not integrable in the usual sense: $\int_{|x|\leq 1} d^2x \frac{1}{|x|^2}$ $\frac{1}{|x|^{2b\alpha}} = \infty$. The claim in eqn. (3.35) only makes sense because including the factor $:e^{2bX(x)}$: improves the situation.

This is only true, however, if we interpret $I(\alpha)$ correctly. A naive imitation of what we have done up to this point would be to define

$$
I(\alpha) \stackrel{?}{=} \lim_{\epsilon \to 0} \int_{|x| \le 1} \mathrm{d}^2 x \, |x|^{-2b\alpha} \, :e^{2bX_{\epsilon}(x)}:, \tag{3.36}
$$

but this is in fact not correct. Indeed, the right hand side of eqn. [\(3.36\)](#page-32-1) is divergent if $\alpha \geq 1/b$. Once we replace X by X_{ϵ} , the random variable $:e^{2bX_{\epsilon}(x)}$: is typically a continuous and nonzero function near $x = 0$ and does not help with the convergence of the integral.

A simple and correct idea is to introduce a spatial cutoff η , restrict the integral defining $I(\alpha)$ to $\eta > 0$, and then take the limit $\eta \to 0$. Thus

$$
I(\alpha) = \lim_{\eta \to 0} \lim_{\epsilon \to 0} \int_{\eta \le |x| \le 1} d^2 x |x|^{-2b\alpha} :e^{2bX_{\epsilon}(x)}.
$$
 (3.37)

It is important that the limit $\epsilon \to 0$ is taken before the limit $\eta \to 0$. Then the moment that we want to study is

$$
\mathbb{E}[I(\alpha)^p] = \mathbb{E}\left[\lim_{\eta \to 0} \lim_{\epsilon \to 0} \left(\int_{\eta \le |x| \le 1} d^2 x |x|^{-2b\alpha} :e^{2bX_{\epsilon}(x)}: \right)^p \right]
$$

$$
\le \lim_{\eta \to 0} \lim_{\epsilon \to 0} \mathbb{E}\left[\left(\int_{\eta \le |x| \le 1} d^2 x |x|^{-2b\alpha} :e^{2bX_{\epsilon}(x)}: \right)^p \right].
$$
 (3.38)

where the inequality at the end is Fatou's inequality^{[12](#page-32-2)} from measure theory.

However, we should discuss the validity of eqn. (3.37) . In section [3.7,](#page-36-0) we will explain how to prove that the cutoff random measure $:e^{2bX_{\epsilon}(x)}$: d^2x converges as $\epsilon \to 0$ to a

¹²In general, the statement of Fatou's inequality involves liminf on both sides, but here that is not necessary as the limits exist on both sides. The limit as $\epsilon \to 0$ of the expectation in the second line exists because this expectation is monotonically increasing or decreasing (depending on p) for $\epsilon \to 0$, by virtue of Kahane's inequality, and then this expectation monotonically increases as $\eta \to 0$ since the integration region is becoming larger. The limit inside the expectation in the first line exists because of the convergence of the integral to a limit as $\epsilon \to 0$ and then by monotonicity as $\eta \to 0$. In section [2.4,](#page-12-0) we have seen examples in which Fatou's inequality is a strict inequality, namely martingales Φ_n such that $\mathbb{E}[\Phi_n] = 1$ for all n but $\lim_{n\to\infty} \Phi_n = 0$. Thus $0 = \mathbb{E}[\lim_{n\to\infty} \Phi_n] < \lim_{n\to\infty} \mathbb{E}[\Phi_n] = 1$. However, in eqn. [\(3.38\)](#page-32-4), it can be shown that the inequality is actually satisfied as an equality as long as p is in the range that makes the right hand side finite. But we will not need to show that; the inequality is sufficient for our purposes.

Liouville measure $M(x) d^2x = e^{2bX(x)}$: d^2x that is valued in positive random variables. Given this measure, the integral $\int d^2x M(x) f(x)$ makes sense for any positive function f that is measurable in the usual sense, and defines a positive random variable, though in general this random variable may have a positive probability (or even probability 1) to equal $+\infty$. In the present case, we want to carry out this construction with the positive measurable function $f(x)$ that equals $1/|x|^{2b\alpha}$ if $|x| \leq 1$ and vanishes otherwise.

The monotone convergence theorem of measure theory says that for any measure – in our case the Liouville measure $d^2x M(x) - if f_1, f_2, \ldots$ is an increasing sequence of nonnegative measurable functions that converges pointwise to f, then $\lim_{n\to\infty} \int d^2x M(x) f_n(x) =$ $\int d^2x M(x) f(x)$. The statement that eqn. [\(3.37\)](#page-32-3) is a valid way to define the random variable $I(\alpha)$ is an example of this idea with $f(x)$ as defined in the last paragraph and with $f_n(x)$ being the function that equals $f(x)$ for $|x| \geq \eta$ and vanishes for $|x| < \eta$. The functions $f_n(x)$ are increasing as $\eta \to 0$ and converge pointwise to $f(x)$. In eqn. [\(3.37\)](#page-32-3), the limit $\epsilon \to 0$ produces $\int d^2x M(x) f_\eta(x)$, and then the limit as $\eta \to 0$ produces $\lim_{\eta \to 0} \int d^2x M(x) f_\eta(x) =$ $\int d^2x M(x) f(x) = I(\alpha).$

We note that for $\alpha \geq 1/b$, the expectation value of $I(\alpha)$ is $+\infty$:

$$
\mathbb{E}[I(\alpha)] = \mathbb{E}\left[\int_{|x| \le 1} d^2 x : e^{2bX(x)} : \frac{1}{|x|^{2b\alpha}}\right] = \int_{|x| \le 1} d^2 x \frac{1}{|x|^{2b\alpha}} = +\infty.
$$
 (3.39)

For any $\alpha \in (b, 1/b+b)$, eqn. (3.35) asserts that there is a range of $p > 0$ with $\mathbb{E}[I(\alpha)^p] < \infty$, but if $\alpha \geq 1/b$, this range is limited to a subset of $p \in (0,1)$. The existence of any $p > 0$ with $\mathbb{E}[I(\alpha)^p] < \infty$ implies that the probability for $I(\alpha)$ to equal $+\infty$ is 0. Thus for $\alpha \in [1/b, 1/b + b), I(\alpha)$ is a random variable that is finite with probability 1, but whose expectation value is infinite.

We should also comment on the lower bound $\alpha > b$ in the statement [\(3.35\)](#page-32-0). For any $p \in (1, 1/b^2)$, there is $\alpha > b$ with $p < (Q - \alpha)/b$. So for any such p, the statement (3.35) implies that there is $\alpha > b$ with $\mathbb{E}[I(\alpha)^p] < \infty$. Making α smaller makes $1/|x|^{2b\alpha}$ smaller, and therefore makes $I(\alpha)$ smaller. So given that $\mathbb{E}[I(\alpha)^p] < \infty$ for some $\alpha > b$, it actually follows that the same is true as well for all $\alpha \leq b$ (including the possibility $\alpha < 0$). Making p smaller also improves the convergence, as long as p remains positive. So actually for $\alpha \leq b, E[I(\alpha)^p] < \infty$ for all $p \in (0, 1/b^2)$. The same remains true if $p < 0$. We will not prove that last assertion, but a heuristic explanation was given in section [2.5.](#page-16-0)

Finally we turn to a proof of the assertion (3.35) . First we consider the case $0 < p < 1$, which as we have already explained encompasses the most delicate situation with $\alpha \geq 1/b$. We use the formula [\(3.38\)](#page-32-4) for $\mathbb{E}[I(\alpha)^p]$, whose finiteness we wish to prove. In this formula, we take η to be of the form 2^{-n} with $n = 1, 2, 3, \ldots$, and we consider the limit $n \to \infty$. Thus for $k = 1, 2, 3, \ldots$ we define

$$
I_{k,\epsilon}(\alpha) = \int_{2^{-k} \le |x| \le 2^{-k+1}} d^2 x \, |x|^{-2b\alpha} \, :e^{2bX_{\epsilon}(x)}:, \tag{3.40}
$$

in terms of which

$$
I(\alpha) = \lim_{n \to \infty} \lim_{\epsilon \to 0} \sum_{k=1}^{n} I_{k,\epsilon}(\alpha),
$$
\n(3.41)

and so by Fatou's inequality,

$$
\mathbb{E}[I(\alpha)^p] \leq \lim_{n \to \infty} \lim_{\epsilon \to 0} \mathbb{E}\left[\left(\sum_{k=1}^n I_{k,\epsilon}(\alpha)\right)^p\right] \leq \lim_{n \to \infty} \lim_{\epsilon \to 0} \sum_{k=1}^n \mathbb{E}\left[I_{k,\epsilon}(\alpha)^p\right],\tag{3.42}
$$

where the last step is valid for $0 < p < 1$.

By familiar reasoning, we have

$$
\mathbb{E}\left[\left(\int_{\frac{c}{2}\leq|x|\leq c} d^2x|x|^{-2b\alpha} :e^{2bX_{\epsilon}(x)}:\right)^p\right]
$$
\n
$$
= c^{(2-2b\alpha)p}\mathbb{E}\left[\left(\int_{\frac{1}{2}\leq|y|\leq 1} d^2y|y|^{-2b\alpha} :e^{2bX_{\epsilon}(cy)}:\right)^p\right]
$$
\n
$$
= c^{(2-2b\alpha)p}\mathbb{E}\left[\left(\int_{\frac{1}{2}\leq|y|\leq 1} d^2y|y|^{-2b\alpha} :e^{2bX_{\epsilon/c}(y)+2b\Omega_{1/c}}:\right)^p\right]
$$
\n
$$
= c^{2p(1-b\alpha+b^2-b^2p)}\mathbb{E}\left[\left(\int_{\frac{1}{2}\leq|y|\leq 1} d^2y|y|^{-2b\alpha} :e^{2bX_{\epsilon/c}(y)}:\right)^p\right].
$$
\n(3.43)

In the following, we use the fact that for $c < 1$, $c^{2p(1-b\alpha+b^2-b^2p)} < 1$ precisely if $p <$ $(Q - \alpha)/b$.

Setting $c = 2^{-(k-1)}$, eqn. [\(3.43\)](#page-34-0) means that in eqn. [\(3.42\)](#page-34-1), we can replace $I_{k,\epsilon}(\alpha)^p$ with $2^{-2p(k-1)(1-b\alpha+b^2-b^2p)}I_{1,2^{k-1}\epsilon}(\alpha)$. Since we will be taking $\epsilon \to 0$ with fixed n and therefore with an upper bound on k, it makes no difference here to replace $I_{1,2^{k-1}\epsilon}(\alpha)$ with $I_{1,\epsilon}(\alpha)$. The sum over k on the right hand side of eqn. (3.42) is then just a geometric series and the $n \to \infty$ limit consists of continuing the geometric series up to infinity. So eqn. [\(3.42\)](#page-34-1) becomes an inequality

$$
\mathbb{E}[I(\alpha)^p] \le \sum_{k=1}^{\infty} 2^{-2p(k-1)(1-b\alpha+b^2-b^2p)} \lim_{\epsilon \to 0} \mathbb{E}[I_{1,\epsilon}(\alpha)^p]. \tag{3.44}
$$

The geometric series converges if $p < (Q-\alpha)/b$, and $\lim_{\epsilon \to 0} \mathbb{E}[I_{1,\epsilon}(\alpha)^p]$ is also finite, because $I_{1,\epsilon}(\alpha)$ is defined by an integral over the region $|x| > 1/2$, away from the singularity. Indeed, since we have assumed $p < 1$, we have for all ϵ , $\mathbb{E}[I_{1,\epsilon}(\alpha)^p] < 1 + \mathbb{E}[I_{1,\epsilon}(\alpha)] =$ $1 + \int_{\frac{1}{2} \leq |x| \leq 1} d^2x |x|^{-2b\alpha} < \infty$, though pending the analysis in section [3.7,](#page-36-0) we do not yet know that $\lim_{\epsilon \to 0} \mathbb{E}[I_{1,\epsilon}(\alpha)^p] \neq 0$. So eqn. [\(3.44\)](#page-34-2) implies that $\mathbb{E}[I(\alpha)^p] < \infty$ in the claimed range of p.

The other case $p > 1$ is less delicate as it only arises if $\alpha < 1/b$. Let $I_{n,\epsilon}(\alpha)$ be $I(\alpha)$ modified by the η and ϵ cutoffs, and for $\eta < c < 1$, define

$$
I_{\eta,\epsilon}(\alpha) = J_{\eta,\epsilon,c}(\alpha) + K_{\epsilon,c}(\alpha),\tag{3.45}
$$

with

$$
J_{\eta,\epsilon,c}(\alpha) = \int_{\eta \leq |x| \leq c} d^2 x |x|^{-2b\alpha} :e^{2bX_{\epsilon}(x)}:,\qquad K_{\epsilon,c}(\alpha) = \int_{c \leq |x| \leq 1} d^2 x |x|^{-2b\alpha} :e^{2bX_{\epsilon}(x)}:.
$$
\n(3.46)

Hence

$$
\mathbb{E}[I_{\eta,\epsilon}(\alpha)^p] = \mathbb{E}[(J_{\eta,\epsilon,c}(\alpha) + K_{\epsilon,c}(\alpha))^p] \le 2^{p-1} \left(\mathbb{E}[J_{\eta,\epsilon,c}(\alpha)^p] + \mathbb{E}[K_{\epsilon,c}(\alpha)^p] \right). \tag{3.47}
$$

The familiar scaling arguments show that

$$
\mathbb{E}[J_{\eta,\epsilon,c}(\alpha)^p] = c^{2p(1-b\alpha+b^2-b^2p)} \mathbb{E}[I_{\eta/c,\epsilon/c}(\alpha)^p].
$$
\n(3.48)

Replacing η/c by η and ϵ/c by ϵ makes $\mathbb{E}[I_{\eta/c,\epsilon/c}(\alpha)^p]$ larger; in the case of η , this is because the replacement makes the integration region larger, while in the case of ϵ , it is because of Kahane's inequality (3.15) . So the inequality (3.47) becomes

$$
\mathbb{E}[I_{\eta,\epsilon}(\alpha)^p] \le 2^{p-1} \left(c^{2p(1-b\alpha+b^2-b^2p)} \mathbb{E}[I_{\eta,\epsilon}(\alpha)^p] + \mathbb{E}[K_{\epsilon,c}(\alpha)^p] \right). \tag{3.49}
$$

On the right hand side, since the condition $p < (Q - \alpha)/b$ together with $\alpha > b$ implies that $p < 1/b^2$, the result of section [3.4](#page-29-0) implies that $\lim_{\epsilon \to 0} \mathbb{E}[K_{\epsilon,c}(\alpha)^p] < \infty$. Choosing c so that $2^{p-1}c^{2p(1-b\alpha+b^2-b^2p)} < 1$, the inequality (3.49) gives an upper bound on $\mathbb{E}[I_{\eta,\epsilon}(\alpha)^p]$ that is independent of the cutoff parameters η and ϵ . So $\mathbb{E}[I(\alpha)^p] < \infty$.

3.6 Divergent Integrals of Singular Functions

We will now investigate, somewhat informally, the counterpart of eqn. (3.35) , which says that $I(\alpha) = \infty$ with probability 1 if $\alpha \ge Q$. The analysis will also shed light on the convergence of $I(\alpha)$ for $\alpha < Q$.

We parametrize the x_1, x_2 plane with polar coordinates r, θ centered at $x = 0$ and set $r = e^{-s}$. For $s \ge 0$, let $B_s = X_{e^{-s}}(0)$ be the circle average of the Gaussian free field X on the circle $r = e^{-s}$. Also, define the "remainder"

$$
Y(s,\theta) = X(e^{-s}e^{i\theta}) - B_s.
$$
\n(3.50)

In terms of the Fourier expansion $X(s, \theta) = X_0(s) + \sum_{n \neq 0} e^{in\theta} X_n(s)$ of eqn. [\(2.38\)](#page-14-0), $B_s =$ $X_0(s)$ and $Y(s, \theta) = \sum_{n \neq 0} e^{in\theta} X_n(s)$. Since $X(s, \theta) = B_s + Y(s, \theta)$ and B_s and $Y(s, \theta)$ are independent, we have $:e^{2bX(s,\theta)}:=e^{2bB_s}$: $:e^{2bY(s,\theta)}$:. Here $:e^{2bB_s}=e^{2bB_s-2b^2\mathbb{E}[B_s^2]}$. We recall from section [2.4](#page-12-0) that B_s has the statistics of Brownian motion, normalized so that $B_0 = 0$ because of the condition^{[13](#page-35-3)} [\(2.3\)](#page-6-3) and with $\mathbb{E}[B_s^2] = s$ because of the normalization of the action [\(2.39\)](#page-14-3). So

$$
I(\alpha) = \int_{|x| < 1} d^2 x |x|^{-2b\alpha} :e^{2bX(x)}: = \int_0^\infty ds e^{-s(2-2b\alpha + 2b^2 s)} e^{2bB_s} \int_0^{2\pi} d\theta :e^{2bY(s,\theta)}: e^{2b^2 \mathbb{E}[B_0^2]} = \int_0^\infty ds e^{-2s(Q-\alpha)} e^{2bB_s} \int_0^{2\pi} d\theta :e^{2bY(s,\theta)}: \tag{3.51}
$$

The integral over s diverges exponentially if $\alpha > Q$. Indeed, the random variable B_s is typically of order $s^{1/2}$, too small to affect the convergence of the integral. Since the

¹³If we did not have the condition [\(2.3\)](#page-6-3), we would still have $\mathbb{E}[(B_s - B_0)^2] = s$ because of the normalization of the action. We would also have $\mathbb{E}[(B_s - B_0)B_0] = 0$, because in Brownian motion the increment $B_s - B_0$ is independent of B_0 . So we would get $\mathbb{E}[B_s^2] = s + \mathbb{E}[B_0^2]$. A constant factor $e^{2b^2\mathbb{E}[B_0^2]}$ would then appear in the following formulas; of course, this would not affect the discussion of convergence of integrals.

zero-mode of X has been removed in defining the random variable $Z_s = \int_0^{2\pi} d\theta \; i e^{2bY(s,\theta)}$; this random variable has a stationary distribution, invariant under constant shifts of s , reflecting the scale-invariance of Liouville theory. So on the average Z_s is not growing or decaying as s becomes large and does not help with the convergence of the integral.

If $\alpha = Q$, the integral does not diverge exponentially. But it does still diverge with probability 1, because Z_s has only short range correlations and is independent of B_s . So if we partition the half-line $s \geq 0$ as the union of infinitely many closed intervals $[n, n+1]$, then with probability 1, in infinitely many of those intervals one has $Z_s > \delta$ (for some chosen δ) and $B_s > 0$, ensuring a divergence of the integral.

For $\alpha < Q$, the integral in eqn. [\(3.51\)](#page-35-4) converges with probability 1, so $I(\alpha) < \infty$ with probability 1. This was already clear from section [3.5,](#page-31-0) where we showed that if $\alpha < Q$, then $\mathbb{E}[I(\alpha)^p] < \infty$ for some range of $p > 0$.

3.7 Construction of Liouville Measure for all $b < 1$

Finally, here we will sketch the proof of the convergence of the random variable $M_{f,\epsilon}(A)$ = $\int_A d^2x f(x)$: $e^{2bX_\epsilon(x)}$: to a nonzero limit for any $0 \le b < 1$ and assuming that f is smooth with nonzero integral over A. This is a much stronger claim than the statements about moments that have been discussed so far. The proof we give here is inspired by and very similar to an argument of Berestycki [\[30\]](#page-41-13). In contrast to section [2.5,](#page-16-0) which was based on convergence in the L^2 norm, here we will consider convergence in the L^1 norm, defined for any real-valued random variable Φ as the expectation of the absolute value of Φ . Random variables are complete in the L^1 norm just as in the L^2 norm. So if

$$
\lim_{\epsilon,\epsilon'\to\infty} \mathbb{E}[|M_{f,\epsilon}(A) - M_{f,\epsilon'}(A)|] = 0,
$$
\n(3.52)

which shows that $M_{f,\epsilon}(A)$ is a Cauchy sequence in the L^1 norm, then as $\epsilon \to 0$, the family of random variables $M_{f,\epsilon}(A)$ will converge in L^1 to a limit $M_f(A)$. Convergence in L^1 guarantees in particular that $\mathbb{E}[M_{f,\epsilon}(A)]$ converges to $\mathbb{E}[M_f(A)]$. But

$$
\mathbb{E}[M_{f,\epsilon}(A)] = \int_A \mathrm{d}^2 x f(x) \tag{3.53}
$$

for any ϵ , and therefore $\mathbb{E}[M_f(A)] \neq 0$. In particular, $M_f(A)$ has a nonzero chance of being nonzero. This compares to what happens for $b > 1$, where $\lim_{\epsilon \to 0} M_{f,\epsilon}(A) = 0$, as found in sections [2.6](#page-18-0) and [3.3.](#page-28-0)

To prove eqn. [\(3.52\)](#page-36-1), fix $0 < \epsilon'/2 < \epsilon < \epsilon'$. Let $I_{\epsilon} := M_{f,\epsilon}(A)$ and $I_{\epsilon'} := M_{f,\epsilon'}(A)$. We will define two related quantities J_{ϵ} and $J_{\epsilon'}$, and show that if ϵ and ϵ' are small, then

- I_{ϵ} is close to J_{ϵ} in L^{1} ,
- $I_{\epsilon'}$ is close to $J_{\epsilon'}$ in L^1 , and
- J_{ϵ} is close to $J_{\epsilon'}$ in L^2 .

For any random variable, the L^1 norm is smaller than or equal to the L^2 norm. Thus, if we are able to show the above, then by the triangle inequality for the L^1 norm, I_{ϵ} and $I_{\epsilon'}$ are close in the L^1 norm, proving eqn. (3.52) .

For each $x \in A$, we define a random variable $G(x)$, which is 1 if $X_a(x) \leq \zeta \log(1/a)$ for all $a \in [\epsilon', 3\epsilon']$, and 0 otherwise, where we will choose the number $\zeta \in (2b, 4b)$ later. We define

$$
J_{\epsilon} = \int_{A} d^{2}x \, i e^{2bX_{\epsilon}(x)} \colon G(x), \quad J_{\epsilon'} = \int_{A} d^{2}x \, i e^{2bX_{\epsilon'}(x)} \colon G(x). \tag{3.54}
$$

In particular, note that $J_{\epsilon} \leq I_{\epsilon}$ and $J_{\epsilon'} \leq I_{\epsilon'}$. Thus,

$$
\mathbb{E}|I_{\epsilon}-J_{\epsilon}|=\mathbb{E}[I_{\epsilon}-J_{\epsilon}]=\int_{A} \mathrm{d}^{2}x f(x)\mathbb{E}[\cdot e^{2bX_{\epsilon}(x)}\colon(1-G(x))].\tag{3.55}
$$

We will use Girsanov's theorem to evaluate the expectation on the right. The simplest version of Girsanov's theorem says the following. Suppose that B_t is a standard Brownian motion and $\widetilde{B}_t = B_t + \nu t$ is Brownian motion with drift ν . Let F be a function of the whole path $(B_t: 0 \le t \le T)$ for some T. Let F be the same function of $(B_t: 0 \le t \le T)$. Then^{[14](#page-37-0)}

$$
\mathbb{E}[\widetilde{F}] = \mathbb{E}[e^{\nu B_T - \frac{1}{2}\nu^2 \mathbb{E}[B_T^2]}F].\tag{3.56}
$$

Fixing any $x \in A$, let $B_t = X_{e^{-t}}(x)$, and let $\widetilde{B}_t = B_t + 2bt$. Recall that $G(x) = 1$ if $B_t \le \zeta t$ for all $t \in [\log(1/3\epsilon'), \log(1/\epsilon')],$ and 0 otherwise. Analogously, define $\widetilde{G}(x) = 1$ if $\widetilde{B}_t \leq \zeta t$ for all $t \in [\log(1/3\epsilon'), \log(1/\epsilon')]$ and 0 otherwise. Let $T = \log(1/\epsilon)$. Then by Girsanov's theorem,

$$
\mathbb{E}[\varepsilon e^{2bX_{\epsilon}(x)}: (1 - G(x))]
$$
\n
$$
= \mathbb{E}[e^{2bB_T - 2b^2 \mathbb{E}[B_T^2]}(1 - G(x))]
$$
\n
$$
= \mathbb{E}[1 - \widetilde{G}(x)] = \mathbb{P}(\widetilde{B}_t > \zeta t \text{ for some } t \in [\log(1/3\epsilon'), \log(1/\epsilon')]),
$$
\n
$$
(3.57)
$$

where we used the simple fact that if a random variable Y can only be 0 or 1, then $\mathbb{E}[Y] = \mathbb{P}(Y = 1)$ (with $\mathbb{P}(E)$ denoting the probability of an event E). But since $\zeta > 2b$ and B_t is a Brownian motion with drift 2b, it is very unlikely to have $B_t > \zeta t$ at late t, and more specifically it follows from standard facts about Brownian motion that

$$
\mathbb{P}(\widetilde{B}_t > \zeta t \text{ for some } t \in [\log(1/3\epsilon'), \log(1/\epsilon')]) \le e^{-C(\zeta, b)\log(1/\epsilon)},\tag{3.58}
$$

where $C(\zeta, b)$ is a positive constant that depends only on ζ and b. This shows that

$$
\mathbb{E}\left[|I_{\epsilon} - J_{\epsilon}|\right] \le e^{-C(\zeta, b)\log(1/\epsilon)},\tag{3.59}
$$

¹⁴The assertion that B_t is standard Brownian motion means that B_t is Brownian motion normalized so that $B_0 = 0$ and $\mathbb{E}[B_t^2] = t$. Thus B_t is governed by the distribution function $\frac{1}{Z} \exp\left(-\frac{1}{2} \int_0^T dt \left(\frac{dB_t}{dt}\right)^2\right)$, where Z is a normalization constant, and $B_t = B_t + \nu t$ is governed by the distribution function $\frac{1}{Z} \exp \left(-\frac{1}{2} \int_0^T \mathrm{d}t \left(\frac{\mathrm{d}\tilde{B}_t}{\mathrm{d}t} - \nu\right)^2\right)$. Hence $\mathbb{E}[F(\tilde{B}_t)] = \frac{1}{Z} \int D\tilde{B}_t F(\tilde{B}_t) \exp \left(-\frac{1}{2} \int_0^T \mathrm{d}t \left(\frac{\mathrm{d}\tilde{B}_t}{\mathrm{d}t} - \nu\right)^2\right) =$ $\frac{1}{Z} \int D\widetilde{B}_t F(\widetilde{B}_t) e^{\nu \widetilde{B}_t} e^{-\frac{1}{2}\nu^2 T} \exp\left(-\frac{1}{2}\int_0^T dt \left(\frac{d\widetilde{B}_t}{dt}\right)^2\right)$. Renaming the integration variable \widetilde{B}_t as B_t and using $T = \mathbb{E}[B_T^2]$, we conclude that $\mathbb{E}[F(\tilde{B}_t)] = \mathbb{E}[e^{\nu B_T - \frac{1}{2}\nu^2 \mathbb{E}[B_T^2]}F(B_T)]$, as claimed in the text, where the expectation on the right of the final statement is taken with respect to standard Brownian motion.

and a similar bound holds for $\mathbb{E}[|I_{\epsilon'} - J_{\epsilon'}|]$. Next, note that

$$
\mathbb{E}[(J_{\epsilon} - J_{\epsilon'})^2]
$$
\n
$$
= \int_{A \times A} d^2x d^2y f(x) f(y)
$$
\n
$$
\cdot \mathbb{E}[(e^{2bX_{\epsilon}(x)} \cdot - :e^{2bX_{\epsilon'}(x)} \cdot)(:e^{2bX_{\epsilon}(y)} \cdot - :e^{2bX'_{\epsilon}}(y) \cdot)G(x)G(y)].
$$
\n(3.60)

Take any $x, y \in A$ such that $|x - y| > 2\epsilon'$. Then, given the values of X outside the union of the two open balls of radius ϵ' centered at x and y, the conditional expected value of $:e^{2bX_{\epsilon}(x)}$: is $:e^{2bX_{\epsilon'}(x)}$; and the conditional expected value of $:e^{2bX_{\epsilon}(y)}$: is $:e^{2bX_{\epsilon'}(y)}$. (This follows from the fact that $G(x)$ and $G(y)$ only depend on data outside of the two balls of radius ϵ' , together with the martingale property of the normal ordered exponential of Brownian motion, which was explained in section (2.4) . Therefore, for such x and y, the integrand is zero.

Now, take any $x, y \in A$ such that $|x - y| \leq 2\epsilon'$. Let $B_t = X_{e^{-t}}(x)$ and $T_0 = \log(1/3\epsilon')$. Note that $G(x)G(y) \leq H$, where H is the random variable that is 1 if $B_{T_0} \leq \zeta T_0$ and 0 otherwise. Now, let us open up the parentheses inside the expectation on the right in eqn. [\(3.60\)](#page-38-0). Consider the first term:

$$
\mathbb{E}\left[\cdot e^{2bX_{\epsilon}(x)}\cdot e^{2bX_{\epsilon}(y)}\cdot G(x)G(y))\right] \leq \mathbb{E}[e^{2b(X_{\epsilon}(x)+X_{\epsilon}(y))-4b^2\log(1/\epsilon)}H].\tag{3.61}
$$

We will now use the following simple fact about Gaussian random variables. Suppose that X and Y are jointly Gaussian random variables with expected value zero. Take any ν , and let

$$
a = \nu \mathbb{E}[XY].\tag{3.62}
$$

Let c^2 be the variance of Y. Let Z be another Gaussian random variable, with expected value a and variance c^2 . Then for any function^{[15](#page-38-1)} F,

$$
\frac{\mathbb{E}[e^{\nu X}F(Y)]}{\mathbb{E}[e^{\nu X}]} = \mathbb{E}[F(Z)].
$$
\n(3.63)

Let us now apply this identity with $X = X_{\epsilon}(x) + X_{\epsilon}(y)$, $Y = B_{T_0}$, and $\nu = 2b$. Let F be the function such that $H = F(B_{T_0})$. Then

$$
\frac{\mathbb{E}[e^{2b(X_{\epsilon}(x)+X_{\epsilon}(y))}H]}{\mathbb{E}[e^{2b(X_{\epsilon}(x)+X_{\epsilon}(y))}]} = \mathbb{E}[F(Z)],
$$
\n(3.64)

where Z is a Gaussian random variable with expected value

$$
a = 2b\mathbb{E}[X_{3\epsilon'}(x)(X_{\epsilon}(x) + X_{\epsilon}(y))]
$$
\n(3.65)

and variance $c^2 = \text{Var}(X_{3\epsilon'}(x))$. Now, by eqn. [\(2.6\)](#page-7-1) and the fact that the circles of radius ϵ around x and y are both inside the disk of radius $3\epsilon'$ around x, we get that

$$
\mathbb{E}[X_{3\epsilon'}(x)X_{\epsilon}(x)] = \mathbb{E}[X_{3\epsilon'}(x)X_{\epsilon}(y)] = \log \frac{1}{3\epsilon'},
$$
\n(3.66)

¹⁵The following statement can be proved by integrating over X in the joint distribution of X and Y, which by hypothesis is Gaussian.

and thus,

$$
a = 4b \log \frac{1}{3\epsilon'}.\tag{3.67}
$$

Also by eqn. [\(2.6\)](#page-7-1), $c^2 = \log(1/3\epsilon')$. Thus, Z is a Gaussian random variable with mean $4b \log(1/3\epsilon')$ and variance $\log(1/3\epsilon')$. Consequently, if $\zeta < 4b$ and ϵ' is small, Z is very unlikely to be less than $\zeta \log(1/3\epsilon')$. By standard estimates for the tail of the Gaussian distribution,

$$
\mathbb{E}[F(Z)] = \mathbb{P}(Z \le \zeta \log(1/3\epsilon'))
$$

=
$$
\exp\left(-\frac{1}{2}(4b-\zeta)^2 \log \frac{1}{3\epsilon'} + o(\log(1/\epsilon'))\right).
$$
 (3.68)

On the other hand, by (2.26) ,

$$
\mathbb{E}[e^{2b(X_{\epsilon}(x)+X_{\epsilon}(y))}] = \exp(2b^2 \mathbb{E}[X_{\epsilon}(x)^2] + 2b^2 \mathbb{E}[X_{\epsilon}(y)^2] + 4b^2 \mathbb{E}[X_{\epsilon}(x)X_{\epsilon}(y))]
$$

\n
$$
\leq \exp(8b^2 \log(1/\epsilon)).
$$
\n(3.69)

Combining eqns. (3.61) , (3.64) , (3.68) and (3.69) , we get

$$
\mathbb{E}[:e^{2bX_{\epsilon}(x)} : :e^{2bX_{\epsilon}(y)} : G(x)G(y)]
$$
\n
$$
\leq \exp\left(4b^{2}\log\frac{1}{\epsilon} - \frac{1}{2}(4b-\zeta)^{2}\log\frac{1}{3\epsilon'} + o(\log(1/\epsilon'))\right)
$$
\n
$$
= \exp\left(\left(4b^{2} - \frac{1}{2}(4b-\zeta)^{2}\right)\log\frac{1}{\epsilon} + o(\log(1/\epsilon))\right).
$$
\n(3.70)

Dealing similarly with the other terms in the expansion of the integrand in eqn. [\(3.60\)](#page-38-0), we get

$$
\mathbb{E}[(J_{\epsilon}-J_{\epsilon'})^2)]
$$
\n
$$
\leq \exp\left(\left(4b^2 - \frac{1}{2}(4b-\zeta)^2\right)\log\frac{1}{\epsilon} + o(\log(1/\epsilon))\right)\text{Vol}(\{(x,y) \in A : |x-y| \leq 2\epsilon'\}) \quad (3.71)
$$
\n
$$
= \exp\left(\left(-2+4b^2 - \frac{1}{2}(4b-\zeta)^2\right)\log\frac{1}{\epsilon} + o(\log(1/\epsilon))\right).
$$

Now, if $b < 1$ and we choose $\zeta = 2b$, then

$$
-2 + 4b2 - \frac{1}{2}(4b - \zeta)2 = -2 + 4b2 - 2b2 = -2 + 2b2 < 0.
$$
 (3.72)

This shows that if $b < 1$, then we can choose ζ slightly bigger than 2b, such that we still have

$$
-2 + 4b^2 - \frac{1}{2}(4b - \zeta)^2 < 0. \tag{3.73}
$$

Thus, with such a choice of ζ , and combining the above with eqn. (3.59) , we get that for any $0 < \epsilon'/2 < \epsilon < \epsilon'$,

$$
\mathbb{E}|I_{\epsilon} - I_{\epsilon'}| \le e^{-C(\zeta, b)\log(1/\epsilon)},\tag{3.74}
$$

where $C(\zeta, b)$ is a positive constant that depends only on ζ and b. This almost proves eqn. [\(3.52\)](#page-36-1), except that we have the restriction that $\epsilon' > \epsilon > \epsilon'/2$. To remove this restriction, take any $0 < \epsilon < \epsilon'$, and find $\epsilon_1, \ldots, \epsilon_n$ such that $\epsilon = \epsilon_1 < \epsilon_2 < \cdots < \epsilon_n = \epsilon'$,

and $\epsilon_{k-1} > \epsilon_k/2$ for each k. Repeatedly applying eqn. [\(3.74\)](#page-39-2) to each $\mathbb{E}|I_{\epsilon_{k-1}} - I_{\epsilon_k}|$, and observing that $n = O(\log(1/\epsilon'))$, we get eqn. [\(3.52\)](#page-36-1).

This completes the proof of eqn. [\(3.52\)](#page-36-1) for smooth f. A consequence is that $M_{1,\epsilon}(A)$ converges to a nonzero limit as $\epsilon \to 0$, where 1 denotes the function that is 1 everywhere. We will denote this limit by $M(A)$. The function M associates a non-negative real number to every Borel set A, and it satisfies the axioms for being a measure, in the sense of measure theory. Thus, Liouville theory defines a random measure M on the Borel sets of \mathbb{R}^2 , which is nontrivial (i.e., nonzero) when $b < 1$.

Acknowledgements. Research of SC supported in part by NSF grants DMS-2113242 and DMS-2153654. Research of EW supported in part by NSF Grant PHY-2207584.

References

- [1] A. M. Polyakov, "Quantum Geometry of Bosonic Strings," Phys. Lett. 103B (1981) 207.
- [2] T. L. Curtright and C. B. Thorn, "Conformally Invariant Quantization of the Liouville Theory," Phys .Rev. Lett. 48 (1982) 1309.
- [3] J.-L. Gervais and A. Neveu, "New Quantum Treatment of Liouville Field Theory," Nucl. Phys. B224 (1983) 329-348.
- [4] V. Knizhnik, A. Polyakov, and A. Zamolodchikov, "Fractal Structure of 2d Quantum Gravity," Mod. Phys. Lett. A3 (1988) 819.
- [5] M. Goulian and M. Li, "Correlation Functions in Liouville Theory,"' Phys. Rev. Lett. 66 (1991) 2051.
- [6] N. Seiberg, "Notes on Liouville Theory and Quantum Gravity," Prog. Theor. Phys. Suppl. 102 (1990) 319-349.
- [7] H. Dorn and H.-J. Otto, "Two and Three Point Functions in Liouville Theory," Nucl. Phys. B429 (1994) 375–388, arXiv:hep-th/9403141.
- [8] A. Zamolodchikov and Al. Zamolodchikov, "Conformal Bootstrap in Liouville Field Theory," Nucl. Phys. B477 (1996) 577-605, arXiv:hep-th/9506136.
- [9] J. Teschner, "On the Liouville Three Point Function," Phys. Lett. B363 (1995) 65D70, hep-th/9507109.
- [10] J. Teschner, "Liouville Theory Revisited," Class. Quant. Grav. 18 (2001) R153-R222, arXiv:hep-th/0104158.
- [11] Yu. Nakayama, "Liouville Field Theory: A Decade After the Revolution," Int. J. Mod. Phys. A19 (2004) 2771-2930, arXiv:hep-th/0402009.
- [12] R. Hoegh-Krohn, " A General Class of Scalar Fields Without Cut-Offs in Two Spacetime Dimensions," Comm. Math. Phys. 21 (1971) 244-255.
- [13] J.-P. Kahane, "Sur le Chaos Multiplicatif," Ann. Sci. Math. Québec, 9 (1985), 105-150.
- [14] B. Mandelbrot, "Possible Refinement of the Lognormal Hypothesis Concerning the Distribution of Energy Dissipation in Intermittent Turbulence," in Statistical Models and

Turbulence, eds. M. Rosenblatt and C. Atta, Lecture Notes in Mathematics 12 (Springer, 1972), 331-51.

- [15] B. Duplantier and S. Sheffield, "Liouville Quantum Gravity and KPZ," Invent. Math. 185 (2011) 333-93.
- [16] F. David, A. Kupiainen, R. Rhodes, and V. Vargas, "Liouville Quantum Gravity on the Riemann Sphere," Commun. Math. Phys. 342 (2016) 869-907.
- [17] A. Kupiainen, R. Rhodes, and V. Vargas, "Integrability of Liouville Theory: Proof of the DOZZ Formula," Ann. Math. 191 (2020) 81, arXiv:1707.08785.
- [18] A. Kupiainen, R. Rhodes, and V. Vargas, "Local Conformal Structure of Liouville Quantum Gravity," Comm. Math. Phys. 371, 1005-69, arXiv:1512.01802.
- [19] V. Vargas, "Lecture Notes on Liouville Theory and the DOZZ Formula," arXiv:1712.00829.
- [20] N. Berestycki and E. Powell, Gaussian Free Field and Liouville Quantum Gravity (Cambridge University Press, to appear).
- [21] R. Robert and V. Vargas, "Gaussian Multiplicative Chaos Revisited," Ann. Probab. 38 (2010) 605-31, arXiv:0807.1030.
- [22] R. Rhodes and V. Vargas, "Gaussian Multiplicative Chaos and Applications: A Review," arXiv:1305.6221.
- [23] B. Duplantier, J. Miller, and S. Sheffield, "Liouville Quantum Gravity as a Mating of Trees," Asterisque (2021) 427, arXiv:1409.7055.
- [24] C. Guillarmou, A. Kupiainen, and R. Rhodes, "Review of the Probabilistic Construction and Conformal Bootstrap in Liouville Theory," arXiv:2403.12780.
- [25] J. L. Doob, *Stochastic Processes* (Wiley, New York, 1953).
- [26] V. S. Dotsenko and V. A. Fateev, "Four Point Correlation Functions and the Operator Algebra in the Two-Dimensional Conformal Invariant Theories with the Central Charge $c < 1$," Nucl. Phys. **B251** (1985) 691.
- [27] B. Duplantier, R. Rhodes, S. Sheffield, and V. Vargas, "Critical Gaussian Multiplicative Chaos: Convergence of the Derivative Martingale," Ann. Prob. 42 (2014) 1969-1808, arXiv:1206.1671.
- [28] D. Harlow, J. Maltz, and E. Witten, "Analytic Continuation of Liouville Theory," JHEP 12 (2011) 071, arXiv:1108.4417.
- [29] E. Bacry and J.-F. Muzy, "Log-infinitely Divisible Multifractal Process," Commun. Math. Phys. 236 (2003) 449-75.
- [30] N. Berestycki, "An Elementary Approach to Gaussian Multiplicative Chaos", Electr. Comm. Probab, 22 (2017) no. 27, 1-12.