

GENERALIZATION OF SPIVEY'S RECURRENCE RELATION

TAEKYUN KIM AND DAE SAN KIM

ABSTRACT. In 2008, Spivey found a recurrence relation for the Bell numbers ϕ_n . We consider the probabilistic r -Bell polynomials associated with Y , $\phi_{n,r}^Y(x)$, which are a probabilistic extension of the r -Bell polynomials. Here Y is a random variable whose moment generating function exists in some neighborhood of the origin and $\phi_n = \phi_{n,0}^1(1)$. The aim of this paper is to generalize the relation for the Bell numbers to that for the probabilistic r -Bell polynomials associated with Y .

1. INTRODUCTION

The Stirling number $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ counts the number of ways to partition a set of n objects into k nonempty subsets. The Bell polynomials are defined by $\phi_n(x) = \sum_{k=0}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} x^k$, with the Bell numbers given by $\phi_n = \phi_n(1)$. For $r \in \mathbb{N}$, the r -Stirling number $\left\{ \begin{smallmatrix} n+r \\ k+r \end{smallmatrix} \right\}_r$ enumerates the number of ways to partition a set of $n+r$ objects into $k+r$ nonempty subsets such that the first r elements are in distinct subsets. The r -Bell polynomials are defined by $\phi_{n,r}(x) = \sum_{k=0}^n \left\{ \begin{smallmatrix} n+r \\ k+r \end{smallmatrix} \right\}_r x^k$, with the r -Bell numbers given by $\phi_{n,r} = \phi_{n,r}(1)$. Assume that Y is a random variable whose moment generating function exists in some neighbourhood of the origin. Let $(Y_j)_{j \geq 1}$ be a sequence of mutually independent copies of the random variable Y , and let $S_k = Y_1 + Y_2 + \cdots + Y_k$, ($k \geq 1$), with $S_0 = 0$. Then we define the probabilistic r -Stirling numbers of the second kind associated with Y , $\left\{ \begin{smallmatrix} n+r \\ k+r \end{smallmatrix} \right\}_{r,Y}$, as a probabilistic extension of the r -Stirling numbers of the second kind $\left\{ \begin{smallmatrix} n+r \\ k+r \end{smallmatrix} \right\}_r$. The probabilistic r -Bell polynomials associated with Y are defined by $\phi_{n,r}^Y(x) = \sum_{k=0}^n \left\{ \begin{smallmatrix} n+r \\ k+r \end{smallmatrix} \right\}_{r,Y} x^k$, with the probabilistic r -Bell numbers associated with Y given by $\phi_{n,r}^Y = \phi_{n,r}^Y(1)$.

Spivey found the following recurrence relation on the Bell numbers:

$$\phi_{n+k} = \sum_{l=0}^k \sum_{j=0}^n j^{k-l} \binom{k}{l} \left\{ \begin{smallmatrix} n \\ j \end{smallmatrix} \right\} \phi_l.$$

The aim of this paper is to generalize the relation for the Bell numbers to that for the probabilistic r -Bell polynomials associated with Y . Indeed, we obtain

$$(1) \quad \phi_{j+n,r}^Y(y) = \sum_{l=0}^n \binom{n}{l} r^{n-l} \sum_{k=0}^l \sum_{m=0}^j \frac{y^k}{k!} \binom{j}{m} \sum_{l_1+\cdots+l_k=l} \binom{l}{l_1, \dots, l_k} E \left[S_k^{j-m} \prod_{i=1}^k Y_i^{l_i} \right] \phi_{m,r}^Y(y),$$

where l_1, l_2, \dots, l_k are positive integers.

By using the simple fact (16), we show (1), first for $\phi_n^Y = \phi_{n,0}^Y(1)$, second for

2010 *Mathematics Subject Classification.* 11B73; 11B83.

Key words and phrases. probabilistic r -Stirling numbers of the second kind; probabilistic r -Bell polynomials.

$\phi_n^Y(y) = \phi_{n,0}^Y(y)$, and finally for $\phi_{n,r}^Y(y)$. We note here that $\phi_n = \phi_{n,0}^1(1)$. For the rest of this section, we recall the facts that are needed throughout this paper.

For $n \geq 0$, the Stirling numbers of the second kind are defined by

$$(2) \quad x^n = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} (x)_k, \quad (\text{see [1–26]}),$$

where $(x)_0 = 1$, $(x)_k = x(x-1)\cdots(x-k+1)$, ($k \geq 1$).

The Bell numbers are given by

$$(3) \quad \phi_n = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}, \quad (n \geq 0), \quad (\text{see [15, 21]}).$$

The Bell polynomials are defined by

$$(4) \quad e^{x(e^t-1)} = \sum_{n=0}^{\infty} \phi_n(x) \frac{t^n}{n!}.$$

From (4), we note that

$$(5) \quad \phi_n(x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k, \quad (n \geq 0), \quad (\text{see [7, 8, 10, 15, 21]}).$$

Note that, for $x = 1$, $\phi_n = \phi_n(1)$ are the Bell numbers.

For $r \in \mathbb{N}$ and $n \geq 0$, the r -Stirling numbers of the second kind are given by

$$(6) \quad (x+r)^n = \sum_{k=0}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r (x)_k, \quad (\text{see [20]}).$$

From (6), we note that

$$(7) \quad \begin{aligned} \sum_{l=0}^n \left\{ \begin{matrix} n+r \\ l+r \end{matrix} \right\}_r (x)_l &= (x+r)^n = \sum_{k=0}^n \binom{n}{k} r^{n-k} x^k \\ &= \sum_{k=0}^n \binom{n}{k} r^{n-k} \sum_{l=0}^k \left\{ \begin{matrix} k \\ l \end{matrix} \right\} (x)_l \\ &= \sum_{l=0}^n \sum_{k=l}^n \binom{n}{k} \left\{ \begin{matrix} k \\ l \end{matrix} \right\} r^{n-k} (x)_l. \end{aligned}$$

Thus, by comparing the coefficients on both sides of (7), we get

$$(8) \quad \left\{ \begin{matrix} n+r \\ l+r \end{matrix} \right\}_r = \sum_{k=l}^n \binom{n}{k} \left\{ \begin{matrix} k \\ l \end{matrix} \right\} r^{n-k}, \quad (n \geq l \geq 0).$$

It is known that the Bell numbers satisfy the recurrence relation

$$(9) \quad \phi_{n+1} = \sum_{k=0}^n \binom{n}{k} \phi_k, \quad (n \geq 0), \quad (\text{see [14, 18]}).$$

In 2008, Spivey discovered the following remarkable relation

$$(10) \quad \phi_{n+k} = \sum_{l=0}^k \sum_{j=0}^n j^{k-l} \binom{k}{l} \left\{ \begin{matrix} n \\ j \end{matrix} \right\} \phi_l, \quad (\text{see [24]}).$$

The r -Bell numbers are defined by

$$(11) \quad \phi_{n,r} = \sum_{k=0}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r, \quad (n \geq 0), \quad (\text{see [13–20]}).$$

The r -Bell polynomials are given by

$$(12) \quad e^{rt} e^{x(e^t-1)} = \sum_{n=0}^{\infty} \phi_{n,r}(x) \frac{t^n}{n!}, \quad (\text{see [13-20]}).$$

Thus, by (12), we get

$$(13) \quad \phi_{n,r}(x) = \sum_{k=0}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r x^k, \quad (n \geq 0), \quad (\text{see [6, 8, 16, 17]}).$$

Note that, for $x = 1$, $\phi_{n,r} = \phi_{n,r}(1)$ are the r -Bell numbers.

For any integer $k \geq 0$, the partial Bell polynomials are given by

$$(14) \quad \frac{1}{k!} \left(\sum_{m=1}^{\infty} x_m \frac{t^m}{m!} \right)^k = \sum_{n=k}^{\infty} B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) \frac{t^n}{n!}, \quad (\text{see [10, 11, 14]}),$$

where

$$(15) \quad B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) \\ = \sum_{\substack{l_1 + \dots + l_{n-k+1} \\ l_1 + 2l_2 + \dots + (n-k+1)l_{n-k+1} = n}} \frac{n!}{l_1! l_2! \dots l_{n-k+1}!} \left(\frac{x_1}{1!} \right)^{l_1} \left(\frac{x_2}{2!} \right)^{l_2} \dots \left(\frac{x_{n-k+1}}{(n-k+1)!} \right)^{l_{n-k+1}}.$$

We will use the following simple fact

$$(16) \quad f(x+t) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x)}{n!} (x+t-x)^n = \sum_{n=0}^{\infty} \frac{f^{(n)}(x)}{n!} t^n \\ = \sum_{n=0}^{\infty} \frac{t^n D_x^n}{n!} f(x) = e^{tD_x} f(x),$$

where $D_x = \frac{d}{dx}$.

Assume that Y is a random variable such that the moment generating function of Y ,

$$E[e^{tY}] = \sum_{n=0}^{\infty} E[Y^n] \frac{t^n}{n!}, \quad (|t| < r),$$

exists for some $r > 0$ (see [3, 14, 22]). Here E denotes the mathematical expectation. Let $(Y_j)_{j \geq 1}$ be a sequence of mutually independent copies of the random variable Y , and let $S_k = Y_1 + Y_2 + \dots + Y_k$, ($k \geq 1$), with $S_0 = 0$.

Then we recall the probabilistic Stirling numbers of the second kind associated with Y , $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_Y$, which are defined in terms of the n -th moments of S_l , ($l = 0, 1, 2, \dots, k$):

$$(17) \quad \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_Y = \frac{1}{k!} \sum_{l=0}^k \binom{k}{l} (-1)^{k-l} E[S_l^n], \quad (0 \leq k \leq n), \quad (\text{see [3, 15]}).$$

The probabilistic Bell polynomials associated with Y are defined by

$$(18) \quad \phi_n^Y(x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_Y x^k, \quad (n \geq 0), \quad (\text{see [14, 23]}).$$

From [14], we note that the probabilistic Bell polynomials associated with Y satisfy the recurrence relation

$$(19) \quad \phi_{n+1}^Y(x) = x \sum_{k=0}^n \binom{n}{k} E[Y^{k+1}] \phi_{n-k}^Y(x).$$

2. GENERALIZATION OF SPIVEY'S RECURRENCE RELATION

Let $(Y_j)_{j \geq 1}$ be a sequence of mutually independent copies of the random variable Y , and let

$$S_0 = 0, \quad S_k = Y_1 + Y_2 + \cdots + Y_k, \quad (k \in \mathbb{N}).$$

For $r \in \mathbb{N}$, we define the *probabilistic r -Stirling numbers of the second kind associated with Y* by

$$(20) \quad \frac{1}{k!} \left(E[e^{tY}] - 1 \right)^k e^{rt} = \sum_{n=k}^{\infty} \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} \frac{t^n}{n!}, \quad (k \geq 0).$$

When $Y = 1$, $\left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} = \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r$, $(n \geq k \geq 0)$.

From (20), we note that

$$(21) \quad \begin{aligned} \sum_{n=k}^{\infty} \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} \frac{t^n}{n!} &= \frac{1}{k!} \left(E[e^{Yt}] - 1 \right)^k e^{rt} \\ &= \frac{1}{k!} \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} E[e^{(Y_1 + \cdots + Y_j)t}] e^{rt} \\ &= \frac{1}{k!} \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} E[e^{(S_j+r)t}] \\ &= \sum_{n=0}^{\infty} \left(\frac{1}{k!} \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} E[(S_j+r)^n] \right) \frac{t^n}{n!}. \end{aligned}$$

Therefore, by comparing the coefficients on both sides of (21), we obtain the following theorem.

Theorem 2.1. *For $n \geq k \geq 0$, we have*

$$\left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} = \frac{1}{k!} \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} E[(S_j+r)^n].$$

In view of (13), we define the *probabilistic r -Bell polynomials associated with Y* by

$$(22) \quad \phi_{n,r}^Y(x) = \sum_{k=0}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} x^k, \quad (n \geq 0).$$

When $Y = 1$, $\phi_{n,r}^Y(x) = \phi_{n,r}(x)$. For $x = 1$, $\phi_{n,r}^Y = \phi_{n,r}^Y(1)$ are called the *probabilistic r -Bell numbers associated with Y* .

From (22), we have

$$(23) \quad \begin{aligned} e^{x(E[e^{Yt}] - 1)} e^{rt} &= \sum_{k=0}^{\infty} \frac{x^k}{k!} \left(E[e^{Yt}] - 1 \right)^k e^{rt} \\ &= \sum_{k=0}^{\infty} x^k \sum_{n=k}^{\infty} \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} \frac{t^n}{n!} = \sum_{n=0}^{\infty} \sum_{k=0}^n x^k \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_{r,Y} \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \phi_{n,r}^Y(x) \frac{t^n}{n!}. \end{aligned}$$

Therefore, by (23), we obtain the following theorem.

Theorem 2.2. *The generating function of the probabilistic r -Bell polynomials associated with Y is given by*

$$(24) \quad e^{x(E[e^{Yt}]-1)} e^{rt} = \sum_{n=0}^{\infty} \phi_{n,r}^Y(x) \frac{t^n}{n!}.$$

From (24), we have

$$(25) \quad \begin{aligned} \sum_{n=0}^{\infty} \phi_{n,r}^Y(x) \frac{t^n}{n!} &= e^{x(E[e^{Yt}]-1)} e^{rt} = \sum_{k=0}^{\infty} \frac{1}{k!} \left(\sum_{j=1}^{\infty} xE[Y^j] \frac{t^j}{j!} \right)^k e^{rt} \\ &= \sum_{l=0}^{\infty} \sum_{k=0}^l B_{l,k} \left(xE[Y], xE[Y^2], \dots, xE[Y^{l-k+1}] \right) \frac{t^l}{l!} \sum_{m=0}^{\infty} r^m \frac{t^m}{m!} \\ &= \sum_{n=0}^{\infty} \sum_{l=0}^n \binom{n}{l} \sum_{k=0}^l B_{l,k} \left(xE[Y], xE[Y^2], \dots, xE[Y^{l-k+1}] \right) r^{n-l} \frac{t^n}{n!}. \end{aligned}$$

Therefore, by comparing the coefficients on both sides of (25), we obtain the following theorem.

Theorem 2.3. *For $n \geq 0$, we have*

$$\phi_{n,r}^Y(x) = \sum_{l=0}^n \binom{n}{l} \sum_{k=0}^l B_{l,k} \left(xE[Y], xE[Y^2], \dots, xE[Y^{l-k+1}] \right) r^{n-l}.$$

By (24), we get

$$(26) \quad \begin{aligned} \sum_{n=0}^{\infty} \phi_{n+1,r}^Y(x) \frac{t^n}{n!} &= \frac{d}{dt} \sum_{n=0}^{\infty} \phi_{n,r}^Y(x) \frac{t^n}{n!} = \frac{d}{dt} \left(e^{x(E[e^{Yt}]-1)} e^{rt} \right) \\ &= xE[Ye^{Yt}] e^{x(E[e^{Yt}]-1)} e^{rt} + r e^{x(E[e^{Yt}]-1)} e^{rt} \\ &= x \sum_{k=0}^{\infty} E[Y^{k+1}] \frac{t^k}{k!} \sum_{m=0}^{\infty} \phi_{m,r}^Y(x) \frac{t^m}{m!} + r \sum_{n=0}^{\infty} \phi_{n,r}^Y(x) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(x \sum_{k=0}^n \binom{n}{k} E[Y^{k+1}] \phi_{n-k,r}^Y(x) + r \phi_{n,r}^Y(x) \right) \frac{t^n}{n!}. \end{aligned}$$

Therefore, by comparing the coefficients on both sides of (26), we obtain the following theorem.

Theorem 2.4. *For $n \geq 0$, we have*

$$\phi_{n+1,r}^Y(x) = x \sum_{k=0}^n \binom{n}{k} E[Y^{k+1}] \phi_{n-k,r}^Y(x) + r \phi_{n,r}^Y(x).$$

From (18), we note that

$$(27) \quad e^{E[e^{Yx}]-1} = \sum_{n=0}^{\infty} \phi_n^Y \frac{x^n}{n!}.$$

By using (16), we have

$$(28) \quad \begin{aligned} e^{tD_x} e^{E[e^{Yx}]-1} &= e^{E[e^{Y(x+t)}]-1} \\ &= e^{E[e^{Yx}(e^{Yt}-1)+e^{Yx}]-1} \\ &= e^{E[e^{Yx}(e^{Yt}-1)]} e^{E[e^{Yx}]-1}. \end{aligned}$$

Now, we observe that

$$\begin{aligned}
(29) \quad e^{E[e^{Yx}(e^{Yt}-1)]} &= \sum_{k=0}^{\infty} \frac{1}{k!} \left(E[e^{Yx}(e^{Yt}-1)] \right)^k \\
&= \sum_{k=0}^{\infty} \frac{1}{k!} E \left[e^{(Y_1+\dots+Y_k)x} (e^{Y_1t}-1)(e^{Y_2t}-1)\dots(e^{Y_kt}-1) \right] \\
&= \sum_{k=0}^{\infty} \frac{1}{k!} \sum_{n=k}^{\infty} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} E \left[e^{S_k x} Y_1^{l_1} Y_2^{l_2} \dots Y_k^{l_k} \right] \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{1}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} E \left[e^{S_k x} \prod_{i=1}^k Y_i^{l_i} \right] \frac{t^n}{n!},
\end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers. From (27), (28) and (29), we note that

$$\begin{aligned}
(30) \quad e^{tD_x} e^{E[e^{Yx}]-1} &= e^{E[e^{Yx}(e^{Yt}-1)]} e^{E[e^{Yx}]-1} \\
&= \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{1}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} \frac{t^n}{n!} \sum_{j=0}^{\infty} E \left[S_k^j \prod_{i=1}^k Y_i^{l_i} \right] \frac{x^j}{j!} \sum_{m=0}^{\infty} \phi_m^Y \frac{x^m}{m!} \\
&= \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} \sum_{k=0}^n \sum_{m=0}^l \binom{l}{m} \frac{1}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} E \left[S_k^{l-m} \prod_{i=1}^k Y_i^{l_i} \right] \phi_m^Y \frac{t^n}{n!} \frac{x^l}{l!},
\end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers. On the other hand, by (27), we get

$$\begin{aligned}
(31) \quad e^{tD_x} e^{E[e^{Yx}]-1} &= e^{tD_x} \sum_{l=0}^{\infty} \phi_l^Y \frac{x^l}{l!} = \sum_{n=0}^{\infty} \frac{t^n}{n!} D_x^n \sum_{l=0}^{\infty} \phi_l^Y \frac{x^l}{l!} \\
&= \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} \phi_{l+n}^Y \frac{t^n}{n!} \frac{x^l}{l!}.
\end{aligned}$$

Therefore, by (30) and (31), we obtain the following theorem.

Theorem 2.5. *For $n, l \geq 0$, we have*

$$\phi_{l+n}^Y = \sum_{k=0}^n \sum_{m=0}^l \binom{l}{m} \frac{1}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} E \left[S_k^{l-m} \prod_{i=1}^k Y_i^{l_i} \right] \phi_m^Y,$$

where l_1, l_2, \dots, l_k are positive integers.

We note here that this reduces to the Spivey's relation (10) if $Y = 1$.

$$\begin{aligned}
\phi_{l+n} &= \sum_{k=0}^n \sum_{m=0}^l \binom{l}{m} \frac{1}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} k^{l-m} \phi_m \\
&= \sum_{k=0}^n \sum_{m=0}^l \binom{l}{m} \left\{ \begin{matrix} n \\ k \end{matrix} \right\} k^{l-m} \phi_m.
\end{aligned}$$

We extend Theorem 2.5 to the probabilistic Bell polynomials associated with Y , which are given by (see (18), Theorem 2.2)

$$(32) \quad e^{x(E[e^{Yt}]-1)} = \sum_{n=0}^{\infty} \phi_n^Y(x) \frac{t^n}{n!}.$$

Observe that

$$\begin{aligned}
 (33) \quad e^{tD_x} e^{Y(E[e^{Y^x}]-1)} &= e^{Y(E[e^{Y^{x+t}}]-1)} \\
 &= e^{Y(E[e^{Y^x}(e^{Yt}-1)+E[e^{Y^x}]-1]} \\
 &= e^{YE[e^{Y^x}(e^{Yt}-1)]} e^{Y(E[e^{Y^x}]-1)}.
 \end{aligned}$$

Now, we note that

$$\begin{aligned}
 (34) \quad e^{YE[e^{Y^x}(e^{Yt}-1)]} &= \sum_{k=0}^{\infty} \frac{y^k}{k!} \left(E[e^{Y^x}(e^{Yt}-1)] \right)^k \\
 &= \sum_{k=0}^{\infty} \frac{y^k}{k!} E \left[e^{(Y_1+Y_2+\dots+Y_k)x} (e^{Y_1t}-1)(e^{Y_2t}-1)\dots(e^{Y_kt}-1) \right] \\
 &= \sum_{k=0}^{\infty} \frac{y^k}{k!} E \left[e^{S_k x} (e^{Y_1t}-1)(e^{Y_2t}-1)\dots(e^{Y_kt}-1) \right] \\
 &= \sum_{k=0}^{\infty} \frac{y^k}{k!} \sum_{n=k}^{\infty} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} E \left[e^{S_k x} Y_1^{l_1} Y_2^{l_2} \dots Y_k^{l_k} \right] \frac{t^n}{n!} \\
 &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \frac{y^k}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} \right) E \left[e^{S_k x} \prod_{i=1}^k Y_i^{l_i} \right] \frac{t^n}{n!},
 \end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers. From (33) and (34), we have

$$\begin{aligned}
 (35) \quad e^{tD_x} e^{Y(E[e^{Y^x}]-1)} &= e^{YE[e^{Y^x}(e^{Yt}-1)]} e^{Y(E[e^{Y^x}]-1)} \\
 &= \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{y^k}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} \frac{t^n}{n!} \sum_{j=0}^{\infty} E \left[S_k^j \prod_{i=1}^k Y_i^{l_i} \right] \frac{x^j}{j!} \sum_{m=0}^{\infty} \phi_m^Y(y) \frac{x^m}{m!} \\
 &= \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} \sum_{k=0}^n \sum_{m=0}^l \frac{y^k}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} \binom{l}{m} E \left[S_k^{l-m} \prod_{i=1}^k Y_i^{l_i} \right] \phi_m^Y(y) \frac{t^n}{n!} \frac{x^l}{l!},
 \end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers. On the other hand, by (32), we get

$$\begin{aligned}
 (36) \quad e^{tD_x} e^{Y(E[e^{Y^x}]-1)} &= \sum_{n=0}^{\infty} \frac{D_x^n}{n!} t^n \sum_{l=0}^{\infty} \phi_l^Y(y) \frac{x^l}{l!} \\
 &= \sum_{n=0}^{\infty} \sum_{l=0}^{\infty} \phi_{l+n}^Y(y) \frac{t^n}{n!} \frac{x^l}{l!}.
 \end{aligned}$$

Therefore, by (35) and (36), we obtain the following theorem.

Theorem 2.6. For $n, l \geq 0$, we have

$$\phi_{l+y}^Y(y) = \sum_{k=0}^n \sum_{m=0}^l \binom{l}{m} \frac{y^k}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} E \left[S_k^{l-m} \prod_{i=1}^k Y_i^{l_i} \right] \phi_m^Y(y),$$

where l_1, l_2, \dots, l_k are positive integers.

When $Y = 1$, we note that

$$\begin{aligned}
 (37) \quad \phi_{l+n}(y) &= \sum_{k=0}^n \sum_{m=0}^l \binom{l}{m} y^k \frac{1}{k!} \sum_{l_1+\dots+l_k=n} \binom{n}{l_1, \dots, l_k} k^{l-m} \phi_m(y) \\
 &= \sum_{k=0}^n \sum_{m=0}^l \binom{l}{m} y^k \left\{ \begin{matrix} n \\ k \end{matrix} \right\} k^{l-m} \phi_m(y),
 \end{aligned}$$

where l, n are nonnegative integers. Note that (37) reduces to the Spivey's relation (10) by letting $y = 1$.

From (16) and (24), we note that

(38)

$$\begin{aligned}
e^{tD_x} \left(e^{y(E[e^{Yx}]-1)} e^{rx} \right) &= e^{y(E[e^{Y(x+t)}]-1)} e^{r(x+t)} \\
&= e^{y(E[e^{Yx}(e^{Yt}-1)])} e^{rt} e^{y(E[e^{Yx}]-1)} e^{rx} \\
&= \sum_{k=0}^{\infty} \frac{y^k}{k!} E \left[e^{(Y_1+Y_2+\dots+Y_k)x} (e^{Y_1t}-1) \dots (e^{Y_k t}-1) \right] e^{rt} e^{y(E[e^{Yx}]-1)} e^{rx} \\
&= \sum_{k=0}^{\infty} \frac{y^k}{k!} \sum_{l=k}^{\infty} \sum_{l_1+\dots+l_k=l} \binom{l}{l_1, \dots, l_k} E \left[e^{S_k x} \prod_{i=1}^k Y_i^{l_i} \right] \frac{t^l}{l!} e^{rt} e^{y(E[e^{Yx}]-1)} e^{rx} \\
&= \sum_{l=0}^{\infty} \sum_{k=0}^l \frac{y^k}{k!} \sum_{l_1+\dots+l_k=l} \binom{l}{l_1, \dots, l_k} \frac{t^l}{l!} e^{rt} E \left[e^{S_k x} \prod_{i=1}^k Y_i^{l_i} \right] e^{y(E[e^{Yx}]-1)} e^{rx} \\
&= \sum_{n=0}^{\infty} \sum_{l=0}^n \binom{n}{l} r^{n-l} \sum_{k=0}^l \frac{y^k}{k!} \sum_{l_1+\dots+l_k=l} \binom{l}{l_1, \dots, l_k} \frac{t^n}{n!} \sum_{j=0}^{\infty} E \left[S_k^{j-m} \prod_{i=1}^k Y_i^{l_i} \right] \frac{x^j}{j!} \sum_{m=0}^{\infty} \phi_{m,r}^Y(y) \frac{x^m}{m!} \\
&= \sum_{n=0}^{\infty} \sum_{j=0}^{\infty} \sum_{m=0}^j \sum_{l=0}^n \binom{n}{l} r^{n-l} \binom{j}{m} \sum_{k=0}^l \frac{y^k}{k!} \sum_{l_1+\dots+l_k=l} \binom{l}{l_1, \dots, l_k} E \left[S_k^{j-m} \prod_{i=1}^k Y_i^{l_i} \right] \phi_{m,r}^Y(y) \frac{t^n}{n!} \frac{x^j}{j!},
\end{aligned}$$

where l_1, l_2, \dots, l_k are positive integers. On the other hand, by (24), we get

$$\begin{aligned}
(39) \quad e^{tD_x} \left(e^{y(E[e^{Yx}]-1)} e^{rx} \right) &= \sum_{n=0}^{\infty} \frac{t^n}{n!} D_x^n \sum_{j=0}^{\infty} \phi_{j,r}^Y(y) \frac{x^j}{j!} \\
&= \sum_{n=0}^{\infty} \sum_{j=0}^{\infty} \phi_{j+n,r}^Y(y) \frac{t^n}{n!} \frac{x^j}{j!}.
\end{aligned}$$

Therefore, by (38) and (39), we obtain the following theorem.

Theorem 2.7. For $n, j \geq 0$, we have

$$\phi_{j+n,r}^Y(y) = \sum_{l=0}^n \binom{n}{l} r^{n-l} \sum_{k=0}^l \sum_{m=0}^j \frac{y^k}{k!} \binom{j}{m} \sum_{l_1+\dots+l_k=l} \binom{l}{l_1, \dots, l_k} E \left[S_k^{j-m} \prod_{i=1}^k Y_i^{l_i} \right] \phi_{m,r}^Y(y),$$

where l_1, l_2, \dots, l_k are positive integers.

When $Y = 1$, we have

$$\begin{aligned}
\phi_{j+n,r}(y) &= \sum_{l=0}^n \sum_{m=0}^j \binom{n}{l} \binom{j}{m} r^{n-l} \sum_{k=0}^l \frac{y^k}{k!} \sum_{l_1+\dots+l_k=l} \binom{n}{l_1, \dots, l_k} k^{j-m} \phi_{m,r}(y) \\
&= \sum_{l=0}^n \sum_{m=0}^j \sum_{k=0}^l \binom{n}{l} \binom{j}{m} r^{n-l} \phi_{m,r}(y) y^k k^{j-m} \left\{ \begin{matrix} l \\ k \end{matrix} \right\},
\end{aligned}$$

where j, n are nonnegative integers.

In particular, for $n = 1$, we have

$$\phi_{j+1,r}^Y(y) = y \sum_{k=0}^j \binom{j}{k} E[Y^{k+1}] \phi_{j-k,r}^Y(y) + r \phi_{j,r}^Y(y).$$

3. CONCLUSION

Let Y be a random variable such that the moment generating function of Y exists in a neighborhood of the origin. We introduced probabilistic extensions of the r -Stirling numbers of the second and the r -Bell polynomials, namely the probabilistic r -Stirling numbers of the second associated with Y , $\left\{ \begin{smallmatrix} n+r \\ k+r \end{smallmatrix} \right\}_{r,Y}$ and the probabilistic r -Bell polynomials associated with Y , $\phi_{n,r}^Y(y)$. Here the latter is a natural polynomial extension of the former. Spivey found a recurrence relation for the Bell numbers ϕ_n . In this paper, we generalized the recurrence relation for $\phi_n = \phi_{n,0}^1(1)$ to that for $\phi_{n,r}^Y(y)$ by utilizing the simple fact (16).

In more detail, we obtained for $\left\{ \begin{smallmatrix} n+r \\ k+r \end{smallmatrix} \right\}_{r,Y}$ an explicit expression in terms of the n th moments of $S_j + r$, ($0 \leq j \leq k$) in Theorem 2.1. We derived for $\phi_{n,r}^Y(x)$ the generating function, a finite sum expression involving the incomplete Bell polynomials and a recurrence relation, respectively in Theorem 2.2, Theorem 2.3 and Theorem 2.4. Then the relation for the Bell numbers in (10) was generalized to first that for $\phi_n^Y = \phi_{n,0}^Y(1)$, second that for $\phi_n^Y(y) = \phi_{n,0}^Y(y)$, and finally that for $\phi_{n,r}^Y(y)$, respectively in Theorem 2.5, Theorem 2.6 and Theorem 2.7.

As one of our future projects, we would like to continue to study probabilistic extensions of many special polynomials and numbers and to find their applications to physics, science and engineering as well as to mathematics.

REFERENCES

- [1] Abbas, M.; Bouroubi, S. *On new identities for Bell's polynomials*, Discrete Math. **293** (2005), no. 1-3, 5-10.
- [2] Abramowitz M.; Stegun, I.A. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Dover, New York, 1992.
- [3] Adell, J. A. *Probabilistic Stirling numbers of the second kind and applications*, J. Theoret. Probab. **35** (2022), no. 1, 636-652.
- [4] Belbachir, H.; Bousbaa, I. E. *Translated Whitney and r -Whitney numbers: a combinatorial approach*, J. Integer Seq. **16** (2013), no. 8, Article 13.8.6, 7 pp.
- [5] Boubellouta, K.; Boussayoud, A.; Araci, S.; Kerada, M. *Some theorems on generating functions and their applications*, Adv. Stud. Contemp. Math. (Kyungshang) **30** (2020), no. 3, 307-324.
- [6] Broder, A. Z. *The r -Stirling numbers*, Discrete Math. **49** (1984), no. 3, 241-259.
- [7] Carlitz, L. *Some remarks on the Bell numbers*, Fibonacci Quart. **18** (1980), no. 1, 66-73.
- [8] Carlitz, L. *Weighted Stirling numbers of the first and second kind II*, Fibonacci Quart. **18** (1980), no. 3, 242-257.
- [9] Carlitz, L. *Weighted Stirling numbers of the first and second kind I*, Fibonacci Quart. **18** (1980), no. 2, 147-162.
- [10] Comtet, L. *Advanced combinatorics. The art of finite and infinite expansions*, Revised and enlarged edition. D. Reidel Publishing Co., Dordrecht, 1974.
- [11] Graham, R. L.; Knuth, D. E.; Patashnik, O. *Concrete mathematics. A foundation for computer science*, Second edition. Addison-Wesley Publishing Company, Reading, MA, 1994.
- [12] Kilar, N.; Simsek, Y. *Combinatorial sums involving Fubini type numbers and other special numbers and polynomials: approach trigonometric functions and p -adic integrals*, Adv. Stud. Contemp. Math. (Kyungshang) **31** (2021), no. 1, 75-87.
- [13] Kim, D. S.; Kim, T. K. *Normal ordering associated with λ -Whitney numbers of the first kind in λ -shift algebra*, Russ. J. Math. Phys. **30** (2023), no. 3, 310-319.
- [14] Kim, T.; Kim, D. S. *Probabilistic degenerate Bell polynomials associated with random variables*, Russ. J. Math. Phys. **30** (2023), no. 4, 528-542.
- [15] Kim, T.; Kim, D. S.; Kwon, J. *Probabilistic degenerate Stirling polynomials of the second kind and their applications*, Math. Comput. Model. Dyn. Syst. **30** (2024), no. 1, 16-30.
- [16] Kim, T.; Kim, D. S. *Some identities on degenerate r -Stirling numbers via boson operators*, Russ. J. Math. Phys. **29** (2022), no. 4, 508-517.

- [17] Kim, T.; Kim, D. S. *Degenerate Whitney numbers of first and second kind of Dowling lattices*, Russ. J. Math. Phys. **29** (2022), no. 3, 358-377.
- [18] Kim, T.; Kim, D. S. *Degenerate r -Whitney numbers and degenerate r -Dowling polynomials via boson operators*, Adv. in Appl. Math. **140** (2022), Paper No. 102394, 21 pp.
- [19] Kim, T.; Kim, D. S. *Combinatorial identities involving degenerate harmonic and hyperharmonic numbers*, Adv. Appl. Math. **148** (2023), Paper No. 102535, 15 pp.
- [20] Kim, T.; Kim, D. S.; Lee, H.; Park, J.-W. *A note on degenerate r -Stirling numbers*, J. Inequal. Appl. **2020**, 2020:225.
- [21] Roman, S. *The umbral calculus*, Pure and Applied Mathematics, 111. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1984.
- [22] Ross, S. M. *Introduction to probability models*, Thirteenth edition. Academic Press, London, 2024.
- [23] Soni, R.; Vellaisamy, P.; Pathak, A. K. *A probabilistic generalization of the Bell polynomials*, J. Anal. (2023). <https://doi.org/10.1007/s41478-023-00642-y>
- [24] Spivey, M. Z. *A generalized recurrence for Bell numbers*, J. Integer Seq. **11** (2008), no. 2, Article 08.2.5, 3 pp.
- [25] Ta, B. Q. *Probabilistic approach to Appell polynomials*, Expo. Math. **33** (2015), no. 3, 269-294,
- [26] Teicher, H. *An inequality on Poisson probabilities*, Ann. Math. Statist. **26** (1955), 147-149.

DEPARTMENT OF MATHEMATICS, KWANGWOON UNIVERSITY, SEOUL 139-701, REPUBLIC OF KOREA

Email address: `tkkim@kw.ac.kr`

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 121-742, REPUBLIC OF KOREA

Email address: `dskim@sogang.ac.kr`