# Small diffusivity asymptotics for a linear parabolic SPDE in two space dimensions

Yozo Tonaki\*†‡

Yusuke Kaino<sup>§</sup>

Masayuki Uchida\*†¶

#### Abstract

We consider parameter estimation of the reaction term for a second order linear parabolic stochastic partial differential equation in two space dimensions driven by a *Q*-Wiener process under small diffusivity. We first construct an estimator of the reaction parameter based on continuous spatio-temporal data, and then derive an estimator of the reaction parameter based on high frequency spatio-temporal data by discretizing the estimator based on the continuous data. We show that the estimators have consistency and asymptotic normality. Furthermore, we give simulation results of the estimator based on high frequency data.

#### Keywords and phrases

High frequency data, linear parabolic stochastic partial differential equations, parametric estimation, reaction parameter, small diffusive parameter.

# 1 Introduction

Stochastic partial differential equations (SPDEs) combine partial differential equations with spatiotemporal noise to enable mathematical modeling of spatio-temporal phenomena and are used widely in natural sciences, medicine and economics. Second order parabolic SPDEs, which cover stochastic heat equations and stochastic reaction-diffusion equations, are important models and are applied in many fields such as geophysical fluid dynamics, mathematical finance, neurobiology and population genetics, see Piterbarg and Ostrovskii [29], Kusuoka [22], Tuckwell [36], Altmeyer et al. [1], and Dawson [10]. Since SPDEs are thus used in various fields and can describe complex phenomena, statistical inference for SPDEs has recently been studied as well as theories of SPDEs. Refer to Cialenco [7] for existing theories of statistical inference for SPDE models.

Let  $\mathbb{T} = [0,1]$  and  $D = (0,1)^2$ . We treat the following linear parabolic SPDE in two space dimensions

$$dX_t(y,z) = \left\{ \theta_2 \left( \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) + \theta_1 \frac{\partial}{\partial y} + \eta_1 \frac{\partial}{\partial z} + \theta_0 \right\} X_t(y,z) dt + \sigma dW_t^Q(y,z), \quad (t,y,z) \in \mathbb{T} \times D$$
(1.1)

with an initial value  $X_0$  and the Dirichlet boundary condition  $X_t(y, z) = 0$ ,  $(t, y, z) \in \mathbb{T} \times \partial D$ , where  $W_t^Q$  is a Q-Wiener process in a Sobolev space on D,  $X_0$  is an  $L^2(D)$ -valued random variable and independent of  $W_t^Q$  and  $(\theta_0, \theta_1, \eta_1, \theta_2, \sigma) \in \mathbb{R}^3 \times (0, \infty)^2$ . We consider parametric estimation

<sup>\*</sup>Graduate School of Engineering Science, Osaka University

 $<sup>^{\</sup>dagger}\mathrm{Center}$  for Mathematical Modeling and Data Science (MMDS), Osaka University

<sup>&</sup>lt;sup>‡</sup>e-mail: y.tonaki.es@osaka-u.ac.jp

<sup>&</sup>lt;sup>§</sup>Graduate School of Maritime Sciences, Kobe University

<sup>&</sup>lt;sup>¶</sup>CREST, Japan Science and Technology Agency

of  $\theta_0$  in SPDE (1.1) under small diffusivity, which means that  $\theta_2 \to 0$ ,  $\theta_1 \to 0$  and  $\eta_1 \to 0$ . Thus, we reparameterize the coefficients of SPDE (1.1) as follows.

$$\nu = \theta_2, \quad \kappa = \theta_1/\theta_2, \quad \eta = \eta_1/\theta_2.$$

Let  $\nu \in (0, 1)$  and  $(\kappa, \eta) \in \mathbb{R}^2$  be known parameters, and assume that  $\nu \to 0$  and  $(\kappa, \eta)$  is fixed. We then study estimation of the reaction parameter  $\theta_0$  in the parabolic SPDE

$$dX_t(y,z) = (-\nu \mathcal{A} + \theta_0)X_t(y,z)dt + \sigma dW_t^Q(y,z), \quad (t,y,z) \in \mathbb{T} \times D$$
(1.2)

under  $\nu \to 0$ , where the differential operator  $\mathcal{A}$  is given by

$$-\mathcal{A} = \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} + \kappa \frac{\partial}{\partial y} + \eta \frac{\partial}{\partial z}$$

 $(\theta_0, \sigma) \in \mathbb{R} \times (0, \infty)$  is an unknown parameter whose parameter space is a compact convex subset of  $\mathbb{R} \times (0, \infty)$ , and  $(\theta_0^*, \sigma^*)$  denotes the true value of  $(\theta_0, \sigma)$  and belongs to the interior of the parameter space.

Statistical inference for SPDEs has been studied by many researchers, see for example, Hübner et al. [17], Hübner and Rozovskii [18], Lototsky [23] and Cialenco and Glatt-Holtz [9]. As for discrete observations, see Markussen [24], Bibinger and Trabs [4], Cialenco et al. [8], Chong [6], Hildebrandt and Trabs [15, 16], Tonaki et al. [33, 34], Bibinger and Bossert [3], Bossert [5] and references therein. In the case of SPDE (1.1), which has a linear reaction term, the Fourier coefficients with respect to the eigenfunctions of its differential operator are Ornstein-Uhlenbeck dynamics. Using statistical inference for diffusion processes based on discrete observations, Tonaki et al. [33, 34] proposed an estimator of  $\theta_0$  of SPDE (1.1) when  $W_t^Q$  is a driving process with  $\theta_0$ . Tonaki et al. [35] proposed an estimator of  $\theta_0$  of SPDE (1.1) with small noise, which means that  $\sigma$  is known and  $\sigma \to 0$ . For parametric estimation for the linear reaction term of the second order parabolic SPDE with one space dimension driven by a cylindrical Brownian motion, see Kaino and Uchida [19, 20]. For statistical inference for diffusion processes based on discrete observations, see Kessler [21], Yoshida [41], Uchida and Yoshida [37], Sørensen and Uchida [32] and Gloter and Sørensen [13].

Recently, Gaudlitz and Reiss [12] treated estimation for the reaction term of SPDEs with *d*-space dimensions under small diffusivity in the case that the driving process is a cylindrical Brownian motion and the reaction term is nonlinear. However, their model does not encompass SPDE (1.2). Note that the  $L^2$ -estimate of the second order parabolic SPDE with two space dimensions whose driving process is a cylindrical Brownian motion is divergent, see Walsh [38]. For this reason, we study parametric estimation for the reaction term of SPDE (1.2) driven by a *Q*-Wiener process  $W_t^Q$  in line with the approach of [12]. The main purpose of this paper is to show that the estimator of  $\theta_0$  with the weight parameter  $\beta$  based on high frequency spatio-temporal data is bounded in probability or asymptotically normal with the rate  $\mathcal{R}_{\beta,\nu}^{-1}$  given in (2.5) below under  $\nu \to 0$ . We also show that the function  $\beta \mapsto \mathcal{R}_{\beta,\nu}$  is decreasing.

This paper is organized as follows. We state main results in Section 2. We first propose an estimator of  $\theta_0$  with the weight parameter  $\beta$  using continuous spatio-temporal data. We then derive an estimator of  $\theta_0$  based on high frequency spatio-temporal data by discretizing the estimator based on continuous spatio-temporal data. We discuss the optimal choice of the weight parameter  $\beta$ , and also address the estimation of the volatility parameter  $\sigma^2$ . Section 3 gives the simulation results of the estimator based on high frequency spatio-temporal data. Section 4 provides the proofs of our results.

# 2 Main results

#### 2.1 Setting and notation

Let  $(\Omega, \mathscr{F}, \{\mathscr{F}_t\}_{t \ge 0}, \mathbf{P})$  be a stochastic basis with usual conditions, and let  $\{w_{l_1, l_2}\}_{l_1, l_2 \in \mathbb{N}}$  be independent  $\mathbb{R}$ -valued standard Brownian motions on this basis.

Let  $\{\lambda_{l_1,l_2}, e_{l_1,l_2}\}_{l_1,l_2 \in \mathbb{N}}$  be the eigenpairs of the differential operator  $\mathcal{A}$ , where the eigenfunctions  $e_{l_1,l_2}$  and the corresponding eigenvalues  $\lambda_{l_1,l_2}$  are given by

$$\lambda_{l_1,l_2} = \pi^2 (l_1^2 + l_2^2) + \frac{\kappa^2 + \eta^2}{4}, \quad e_{l_1,l_2}(y,z) = 2\sin(\pi l_1 y)\sin(\pi l_2 z) e^{-(\kappa y + \eta z)/2}$$

for  $l_1, l_2 \in \mathbb{N}$  and  $(y, z) \in \overline{D}$ . Let  $\mathcal{H} = L^2(D)$  with the inner product

$$\langle u, v \rangle = \langle u, v \rangle_{\mathcal{H}} = \iint_{D} u(x, y) v(x, y) e^{\kappa x + \eta y} dx dy, \quad \|u\| = \|u\|_{\mathcal{H}} = \sqrt{\langle u, u \rangle_{\mathcal{H}}}.$$

Note that  $\{e_{l_1,l_2}\}_{l_1,l_2\in\mathbb{N}}$  is the complete orthonormal system of  $\mathcal{H}$  with the inner product  $\langle \cdot, \cdot \rangle$ .

Fix  $\alpha > 0$ , and define a trace class linear operator Q on a Hilbert space  $\mathcal{U} \supset \mathcal{H}$  with a norm  $\|\cdot\|_{\mathcal{U}}$  and the complete orthonormal system  $\{v_{l_1,l_2}\}_{l_1,l_2\in\mathbb{N}}, v_{l_1,l_2} = \|e_{l_1,l_2}\|_{\mathcal{U}}^{-1}e_{l_1,l_2}$  such that  $Qv_{l_1,l_2} = \mu_{l_1,l_2}^{-\alpha}\|e_{l_1,l_2}\|_{\mathcal{U}}^{2}v_{l_1,l_2}$ , where  $\mu_{l_1,l_2} = \pi^2(l_1^2 + l_2^2)$ . We consider the Q-Wiener process given by

$$W_t^Q = \sum_{l_1, l_2 \ge 1} \mu_{l_1, l_2}^{-\alpha/2} \|e_{l_1, l_2}\|_{\mathcal{U}} w_{l_1, l_2}(t) v_{l_1, l_2} = \sum_{l_1, l_2 \ge 1} \mu_{l_1, l_2}^{-\alpha/2} w_{l_1, l_2}(t) e_{l_1, l_2}.$$

In this paper, we assume that  $\alpha > 0$  is known. See [5] for the estimation of  $\alpha$ . Let  $\{S_t\}_{t \ge 0}$  be the semigroup generated by the differential operator  $-\mathcal{A}$ , that is,  $S_t = e^{-t\mathcal{A}}$ , where

$$\mathrm{e}^{-t\mathcal{A}}u = \sum_{l_1, l_2 \ge 1} \mathrm{e}^{-\lambda_{l_1, l_2} t} \langle u, e_{l_1, l_2} \rangle e_{l_1, l_2}, \quad u \in \mathcal{H}.$$

Notice that

$$G_t(\boldsymbol{x}, \boldsymbol{y}) = \sum_{l_1, l_2 \ge 1} e^{-\lambda_{l_1, l_2} t} e_{l_1, l_2}(\boldsymbol{x}) e_{l_1, l_2}(\boldsymbol{y}), \quad \boldsymbol{x}, \boldsymbol{y} \in D$$

is the kernel of the semigroup  $S_t$  and  $S_t u(\boldsymbol{x}) = \langle G_t(\boldsymbol{x}, \cdot), u \rangle, u \in \mathcal{H}$ . Let  $\widetilde{Q}$  be a linear operator such that  $\widetilde{Q}e_{l_1,l_2} = \mu_{l_1,l_2}^{-\alpha}e_{l_1,l_2}$ , and consider the fractional powers  $\widetilde{Q}^{\delta}$  for  $\delta \in \mathbb{R}$ :

$$\widetilde{Q}^{\delta}u = \sum_{l_1, l_2 \ge 1} \mu_{l_1, l_2}^{-\alpha\delta} u_{l_1, l_2} e_{l_1, l_2}$$
(2.1)

for  $u = \sum_{l_1, l_2 \ge 1} u_{l_1, l_2} e_{l_1, l_2}$ ,  $u_{l_1, l_2} \in \mathbb{R}$  such that  $\widetilde{Q}^{\delta} u \in \mathcal{H}$ . Note that  $\widetilde{Q}^{\delta} u \in \mathcal{H}$  for  $u \in \mathcal{H}$  if  $\delta \ge 0$ . Define  $\mathcal{U}_0 = \{ \widetilde{Q}^{1/2} v | v \in \mathcal{H} \}$  with the induced norm  $\| \widetilde{Q}^{-1/2} u \|, u \in \mathcal{U}_0$ . For a separable Hilbert space  $\mathcal{K}$ ,  $HS(\mathcal{K})$  denotes the space of all Hilbert-Schmidt operators from  $\mathcal{K}$  to  $\mathcal{H}$ . Since it holds from Proposition 4.1 that  $S_t \in \mathrm{HS}(\mathcal{U}_0)$  for t > 0 and  $\int_0^1 \|S_{\nu t}\|^2_{\mathrm{HS}(\mathcal{U}_0)} dt < \infty$  for  $\nu > 0$ , there exists a unique mild solution  $\{X_t\}_{t\in\mathbb{T}}$  of SPDE (1.2) such that the variation of constants formula

$$X_t = S_{\nu t} X_0 + \theta_0 \int_0^t S_{\nu(t-s)} X_s \mathrm{d}s + \sigma \int_0^t S_{\nu(t-s)} \mathrm{d}W_s^Q, \quad t \in \mathbb{T}$$
(2.2)

on  $L^2(\Omega; \mathcal{H})$ , and it follows from (4.19) that  $t \mapsto X_t$  is continuous **P**-a.s., see Theorem 7.5 in [30].  $\begin{aligned} \mathbf{P}_{\theta_0,\sigma} \text{ denotes the law of the solution } \{X_t\}_{t\in\mathbb{T}} \text{ of SPDE (1.2) on } C(\mathbb{T};\mathcal{H}). \\ \text{ For } \beta \in \mathbb{R}, \text{ let } \mathcal{L}_{\beta}^2 = \{u = \sum_{l_1, l_2 \geq 1} u_{l_1, l_2} e_{l_1, l_2} |||u||_{\mathcal{L}_{\beta}^2} < \infty\} \text{ with the weighted inner product} \end{aligned}$ 

$$\langle u, v \rangle_{\mathcal{L}^2_\beta} = \langle \widetilde{Q}^{\beta/2} u, \widetilde{Q}^{\beta/2} v \rangle, \quad \|u\|_{\mathcal{L}^2_\beta} = \sqrt{\langle u, u \rangle_{\mathcal{L}^2_\beta}}$$

Note that  $\mathcal{L}^2_{\beta_1} \subset \mathcal{L}^2_{\beta_2}$  for  $\beta_1 \leq \beta_2$  and  $\mathcal{L}^2_{\beta_-} \subset \mathcal{H} \subset \mathcal{L}^2_{\beta_+}$  for  $\beta_- \leq 0 \leq \beta_+$ . For families  $\{a_\lambda\}_\lambda, \{b_\lambda\}_\lambda \subset \mathbb{R}$ , we write  $a_\lambda \leq b_\lambda$  if  $|a_\lambda| \leq C|b_\lambda|$  for some universal constant C > 0 and any  $\lambda$ , and we write  $a_\lambda \sim b_\lambda$  if  $a_\lambda \leq b_\lambda$  and  $b_\lambda \leq a_\lambda$ . For a family  $\{a_\lambda\}_\lambda \subset \mathbb{R}$  and  $a \in \mathbb{R}$ , we write  $a_{\lambda} \equiv a$  if  $a_{\lambda} = a$  for all  $\lambda$ .

### **2.2** Estimation of $\theta_0$ based on continuous observations

Gaudlitz and Reiss [12] proposed the minimum likelihood estimator of the reaction term under small diffusive level. Since  $\sup_{t\in\mathbb{T}} \mathbb{E}[\|X_t\|_{\mathcal{L}^2_{-1}}^2] \gtrsim \sum_{l_1,l_2\geq 1} (l_1^2+l_2^2)^{-1} = \infty$  and the Novikov condition (2.3) in Moleriu [25] for  $f(t) = \widetilde{Q}^{-1}X_t$  is not satisfied in our model, we cannot construct a maximum likelihood estimator of  $\theta_0$  using the Girsanov theorem (Theorem 2.3 in [25]). Therefore, we propose the minimum contrast function based on the likelihood of [12] and provide the minimum contrast estimator of  $\theta_0$ . Let  $\mathbb{X}^{(\text{cont})} = \{X_t(y,z)\}_{t\in\mathbb{T},(y,z)\in\overline{D}}$  be continuous observations. Because it holds from Lemma 8 below that  $\sup_{t\in\mathbb{T}} \mathbb{E}[\|X_t\|_{\mathcal{L}^2_{\beta}}^2] < \infty$  for  $\beta > -1$ , we set the contrast function

$$U_{\beta}(\theta_{0}:\mathbb{X}^{(\text{cont})}) = \theta_{0} \int_{0}^{1} \langle X_{t}, \mathrm{d}X_{t} + \nu \mathcal{A}X_{t} \mathrm{d}t \rangle_{\mathcal{L}^{2}_{\beta}} + \frac{\theta_{0}^{2}}{2} \int_{0}^{1} \|X_{t}\|_{\mathcal{L}^{2}_{\beta}}^{2} \mathrm{d}t,$$
(2.3)

where  $\beta > -1$  can be chosen arbitrarily. Define the estimator  $\hat{\theta}_{0,\beta}^{(\text{cont})}$  as  $\theta_0$  which minimizes (2.3). That is,

$$\hat{\theta}_{0,\beta}^{(\text{cont})} = \hat{\theta}_{0,\beta,\nu}^{(\text{cont})} = \frac{\int_{0}^{1} \langle X_t, \mathrm{d}X_t + \nu \mathcal{A}X_t \mathrm{d}t \rangle_{\mathcal{L}^2_{\beta}}}{\int_{0}^{1} \|X_t\|_{\mathcal{L}^2_{\beta}}^2 \mathrm{d}t}.$$
(2.4)

The numerator of (2.4) is decomposed as follows:

$$\int_{0}^{1} \langle X_{t}, \mathrm{d}X_{t} + \nu \mathcal{A}X_{t} \mathrm{d}t \rangle_{\mathcal{L}^{2}_{\beta}}$$
  
= 
$$\sum_{l_{1}, l_{2} \geq 1} \left\{ \int_{0}^{1} \mu_{l_{1}, l_{2}}^{-\alpha\beta} \langle X_{t}, e_{l_{1}, l_{2}} \rangle \mathrm{d}\langle X_{t}, e_{l_{1}, l_{2}} \rangle + \nu \int_{0}^{1} \lambda_{l_{1}, l_{2}} \mu_{l_{1}, l_{2}}^{-\alpha\beta} \langle X_{t}, e_{l_{1}, l_{2}} \rangle^{2} \mathrm{d}t \right\}.$$

**Remark 1.** The estimator of  $\theta_0$  proposed by Gaudlitz and Reiss [12] in the SPDE driven by a cylindrical Brownian motion corresponds to our proposed estimator (2.4) for  $\beta = 0$  in the SPDE driven by the Q-Wiener process. If we choose  $\beta$  independent of the value of the damping parameter  $\alpha$  and spatio-temporal observation data, it may not be possible to construct a consistent estimator of  $\theta_0$  (see Theorem 2.1, 2.2 or Corollary 2.4 below). In fact, we observe that in our setting of Section 3, the consistent estimator of  $\theta_0$  based on discrete observations cannot be constructed unless  $\beta > 0.2$ . In order to get a consistent estimator of  $\theta_0$ , we choose a suitable  $\beta$  depending on the value of  $\alpha$  and spatio-temporal observation data and propose the estimator with the parameter  $\beta$ .

For  $\beta \geq -1$ , we make the following initial condition.

$$[\mathbf{A1}]_{\beta,2} \ \mathbb{E}[\|X_0\|_{\mathcal{L}^2_{\beta}}^2] < \infty.$$

For  $\beta > -1$ , define

$$\mathcal{R}_{\beta,\nu} = \frac{\int_0^1 \mathbb{E}\left[\|X_t\|_{\mathcal{L}^2_{\beta}}^2\right] \mathrm{d}t}{\sqrt{\int_0^1 \mathbb{E}\left[\|X_t\|_{\mathcal{L}^2_{2\beta+1}}^2\right] \mathrm{d}t}}.$$
(2.5)

We then obtain the following theorem.

**Theorem 2.1.** Let  $\alpha > 0$  and  $\beta > -1$ . Assume that  $[A1]_{\beta,2}$  holds.

(1) It follows that as  $\nu \to 0$ ,

$$\mathcal{R}_{\beta,\nu} \sim \begin{cases} \nu^{-1/2}, & -1 < \beta < \frac{1}{2\alpha} - 1, \\ (-\nu \log \nu)^{-1/2}, & \beta = \frac{1}{2\alpha} - 1, \\ \nu^{\alpha(\beta+1)-1}, & \frac{1}{2\alpha} - 1 < \beta < \frac{1}{\alpha} - 1, \\ -\log \nu, & \beta = \frac{1}{\alpha} - 1, \\ 1, & \frac{1}{\alpha} - 1 < \beta. \end{cases}$$
(2.6)

(2) If  $-1 < \beta \leq \frac{1}{2\alpha} - 1$ , then as  $\nu \to 0$ ,

$$\mathcal{R}_{\beta,\nu}(\hat{\theta}_{0,\beta}^{(\text{cont})} - \theta_0^*) \xrightarrow{d} N(0, (\sigma^*)^2).$$

(3) If  $\frac{1}{2\alpha} - 1 < \beta \leq \frac{1}{\alpha} - 1$ , then as  $\nu \to 0$ ,

$$\mathcal{R}_{\beta,\nu}(\hat{\theta}_{0,\beta}^{(\text{cont})} - \theta_0^*) = \mathcal{O}_{\mathbf{P}}(1).$$

**Remark 2.** When one estimates  $\theta_0$  based on continuous observations under a regular initial condition, it is sufficient to use the result of Theorem 2.1-(2) since one can choose  $\beta < \frac{1}{2\alpha} - 1$ . As mentioned in Remark 1, when one constructs the estimator of  $\theta_0$  based on discrete observations, one needs the result of Theorem 2.1-(3) since one chooses  $\beta$  depending on  $\alpha$  and the number of spatio-temporal observations, see Theorem 2.2-(3), Corollary 2.4 or Section 3.

The properties of  $\mathcal{R}_{\beta,\nu}$  and the optimal choice of the weight parameter  $\beta$  will be addressed in Subsection 2.4.

# **2.3** Estimation of $\theta_0$ based on discrete observations

Our main aim is to estimate  $\theta_0$  based on high frequency spatio-temporal data. To this end, we use discrete data and construct an estimator of  $\theta_0$  by discretizing the estimator given in (2.4).

Let  $\mathbb{X}_{N,M} = \{X_{t_i}(y_j, z_k)\}_{0 \le i \le N, 0 \le j \le M_1, 0 \le k \le M_2}$  be discrete observations, where  $t_i = i/N$ ,  $y_j = j/M_1$  and  $z_k = k/M_2$ . We define the estimator of  $\theta_0$  based on  $\mathbb{X}_{N,M}$  by

$$\hat{\theta}_{0,\beta} = \hat{\theta}_{0,\beta,\nu,N,M,L} = \sum_{i=1}^{N} \sum_{l_{1}=1}^{L} \sum_{l_{2}=1}^{L} \mu_{l_{1},l_{2}}^{-\alpha\beta} [X_{t_{i-1}}]_{M,l_{1},l_{2}} \Big( [X_{t_{i}}]_{M,l_{1},l_{2}} - e^{-\nu\lambda_{l_{1},l_{2}}/N} [X_{t_{i-1}}]_{M,l_{1},l_{2}} \Big) \\ \times \left\{ \frac{1}{N} \sum_{i=1}^{N} \sum_{l_{1}=1}^{L} \sum_{l_{2}=1}^{L} \mu_{l_{1},l_{2}}^{-\alpha\beta} [X_{t_{i-1}}]_{M,l_{1},l_{2}}^{2} \right\}^{-1},$$
(2.7)

where

$$h_{l}(x:a) = \frac{\sqrt{2}e^{ax/2}}{(a/2)^{2} + (\pi l)^{2}} \left(\frac{a}{2}\sin(\pi lx) - \pi l\cos(\pi lx)\right), \quad a, x \in \mathbb{R}, \ l \in \mathbb{N},$$
  
$$\delta_{j}^{[y]}h_{l}(a) = h_{l}(y_{j}:a) - h_{l}(y_{j-1}:a), \quad \delta_{k}^{[z]}h_{l}(a) = h_{l}(z_{k}:a) - h_{l}(z_{k-1}:a),$$
  
$$[X_{t}]_{M,l_{1},l_{2}} = \sum_{j=1}^{M_{1}} \sum_{k=1}^{M_{2}} X_{t}(y_{j-1}, z_{k-1})\delta_{j}^{[y]}h_{l_{1}}(\kappa)\delta_{k}^{[z]}h_{l_{2}}(\eta), \quad t \in \mathbb{T}.$$

For simplicity, we assume that there exist  $b_M > 1/2$  and  $b_L > (\alpha - 1)/4$  such that  $M_1 \wedge M_2 = N^{b_M}$ ,  $L = N^{b_L}$ . The derivation of this estimator can be found in Subsection 4.2.

For  $\beta \in \mathbb{R}$  and q > 2, define  $\mathcal{L}_{\beta}^{q} = \{ u \in \mathcal{L}_{\beta}^{2} | \| u \|_{\mathcal{L}_{\beta}^{q}} < \infty \}$  with

$$\|u\|_{\mathcal{L}^q_{\beta}} = \left(\iint_D |\widetilde{Q}^{\beta/2}u(x,y)|^q \mathrm{e}^{\kappa x + \eta y} \mathrm{d}x \mathrm{d}y\right)^{1/q}.$$

For  $\beta \in \mathbb{R}$  and  $q \geq 2$ , let  $\mathcal{L}^q = \mathcal{L}^q_0$  and  $\mathcal{L}^{\infty}_{\beta} = \bigcap_{q \geq 2} \mathcal{L}^q_{\beta}$ . For  $\alpha > 0, x \in (0, 1]$  and p > 4/x, we set

$$\begin{split} \rho_x^{(\text{time})} &= \begin{cases} 2x, & x \in (0, 1/2], \\ \frac{1}{2(1-x)}, & x \in (1/2, 1), \\ \infty, & x = 1, \end{cases} \\ \tau_{1,x,p}^{(\text{space})} &= 1 - \frac{2}{px-2}, \quad \tau_{2,x,p}^{(\text{space})} = \begin{cases} 1 - \frac{1}{px-1}, & x \in (0, 1/2], \\ 1 - \frac{2}{p}, & x \in (1/2, 1], \end{cases} \\ \rho_{1,x,p}^{(\text{space})} &= 1 - \frac{p+4}{2px+p-4}, \quad \rho_{2,x,p}^{(\text{space})} = \begin{cases} 1 - \frac{p+2}{2px+p-2}, & x \in (0, 1/2], \\ \frac{1}{2} - \frac{1}{p}, & x \in (1/2, 1], \end{cases} \\ \tau_{x,p}^{(\text{trunc})} &= 2\left(1 - \frac{2}{px-2}\right), \quad \rho_{x,p}^{(\text{trunc})} = 2\left(1 - \frac{p+2}{2px+p-4}\right), \end{cases} \\ \phi_{1,x,p}^{(\text{space})} &= \frac{p-1}{px-2}, \quad \phi_{2,x,p}^{(\text{space})} = \begin{cases} \frac{p}{px-1}, & x \in (0, 1/2], \\ 3 - 2x + \frac{2}{p}, & x \in (1/2, 1], \end{cases} \\ \psi_{1,x,p}^{(\text{space})} &= \frac{2(p-1)}{2px+p-4}, \quad \psi_{2,x,p}^{(\text{space})} = \begin{cases} \frac{2p}{2px+p-2}, & x \in (0, 1/2], \\ \frac{3}{2} - x + \frac{1}{p}, & x \in (1/2, 1], \end{cases} \\ \phi_{x,\alpha,p}^{(\text{trunc})} &= \frac{(\alpha-1)px+2(1-2\alpha+p)}{2(px-2)}, \quad \psi_{x,\alpha,p}^{(\text{trunc})} &= \frac{(\alpha-1)px+2(1-2\alpha+p)}{2px+p-4} \end{split}$$

and introduce the following conditions.

$$[\mathbf{A1}]_{\beta,p} \ \mathbb{E}[\|X_0\|_{\mathcal{L}^p_{\beta}}^2] < \infty$$

- $\begin{aligned} [\mathbf{B1}]_{\beta,p} \ \nu(M_1 \wedge M_2)^{2r_1} N^{-\phi_{1,\alpha(\beta+1),p}^{(\mathrm{space})}} \to \infty \text{ and } \nu(M_1 \wedge M_2)^{2r_2} N^{-\phi_{2,\alpha(\beta+1),p}^{(\mathrm{space})}} \to \infty \text{ as } \nu \to 0, \ N \to \infty \\ \text{and } M_1 \wedge M_2 \to \infty \text{ for some } r_1 < \tau_{1,\alpha(\beta+1),p}^{(\mathrm{space})} \text{ and } r_2 < \tau_{2,\alpha(\beta+1),p}^{(\mathrm{space})}. \end{aligned}$
- $[\mathbf{B2}]_{\beta,p} \ \nu L^s N^{-\phi_{\alpha(\beta+1),\alpha,p}^{(\operatorname{trunc})}} \to \infty \text{ as } \nu \to 0, \ N \to \infty \text{ and } L \to \infty \text{ for some } s < \tau_{\alpha(\beta+1),p}^{(\operatorname{trunc})}$
- $[\mathbf{C1}]_\beta \ \nu N^q \to \infty \text{ as } \nu \to 0 \text{ and } N \to \infty \text{ for some } q < \rho_{\alpha(\beta+1)}^{(\text{time})}.$
- $\begin{aligned} [\mathbf{C2}]_{\beta,p} \ \nu(M_1 \wedge M_2)^{2r_1} N^{-\psi_{1,\alpha(\beta+1),p}^{(\mathrm{space})}} \to \infty \text{ and } \nu(M_1 \wedge M_2)^{2r_2} N^{-\psi_{2,\alpha(\beta+1),p}^{(\mathrm{space})}} \to \infty \text{ as } \nu \to 0, \ N \to \infty \\ \text{and } M_1 \wedge M_2 \to \infty \text{ for some } r_1 < \rho_{1,\alpha(\beta+1),p}^{(\mathrm{space})} \text{ and } r_2 < \rho_{2,\alpha(\beta+1),p}^{(\mathrm{space})}. \end{aligned}$
- $[\mathbf{C3}]_{\beta,p} \ \nu L^s N^{-\psi_{\alpha(\beta+1),\alpha,p}^{(\operatorname{trunc})}} \to \infty \text{ as } \nu \to 0, \ N \to \infty \text{ and } L \to \infty \text{ for some } s < \rho_{\alpha(\beta+1),p}^{(\operatorname{trunc})}$

**Remark 3.** (1) For  $\beta \ge -1$  and p > 2,  $[A1]_{-1,p}$  is a stricter condition than  $[A1]_{\beta,2}$ .

- (2) Since  $\|u\|_{\mathcal{L}^p_{-1}} \sim \|\mathcal{A}^{\alpha/2}u\|_{\mathcal{L}^p}$  and  $\mathcal{A}$  is the second order elliptic operator, for example, if  $X_0$  is non-random and belongs to  $C^{\lceil \alpha \rceil}(D)$ , then  $[A1]_{-1,p}$  and especially  $[A1]_{-1,\infty}$  are satisfied.
- (3) For  $\beta > -1$  and p > 4,  $[C2]_{\beta,p}$  and  $[C3]_{\beta,p}$  are stricter conditions than  $[B1]_{\beta,p}$  and  $[B2]_{\beta,p}$ , respectively.

Propositions 4.1-4.3 below show that the estimators  $\hat{\theta}_{0,\beta}^{(\text{cont})}$  and  $\hat{\theta}_{0,\beta}$  are asymptotically equivalent as  $\nu \to 0$ ,  $N \to \infty$ ,  $M_1 \wedge M_2 \to \infty$  and  $L \to \infty$ . We obtain the following main theorem.

**Theorem 2.2.** Let  $\alpha > 0$ ,  $-1 < \beta \leq \frac{1}{\alpha} - 1$  and  $p > \frac{4}{\alpha(\beta+1)}$ . Assume that  $[A1]_{-1,p}$  holds.

(1) Under  $[B1]_{\beta,p}$  and  $[B2]_{\beta,p}$ , it holds that as  $\nu \to 0$ ,  $N \to \infty$ ,  $M_1 \wedge M_2 \to \infty$  and  $L \to \infty$ ,

 $\hat{\theta}_{0,\beta} \xrightarrow{\mathbf{P}} \theta_0^*.$ 

(2) If  $-1 < \beta \leq \frac{1}{2\alpha} - 1$ , then it holds that under  $[C1]_{\beta}$ ,  $[C2]_{\beta,p}$  and  $[C3]_{\beta,p}$ ,

$$\mathcal{R}_{\beta,\nu}(\hat{\theta}_{0,\beta} - \theta_0^*) \xrightarrow{d} N(0, (\sigma^*)^2)$$
(2.8)

as  $\nu \to 0$ ,  $N \to \infty$ ,  $M_1 \wedge M_2 \to \infty$  and  $L \to \infty$ .

(3) If  $\frac{1}{2\alpha} - 1 < \beta \leq \frac{1}{\alpha} - 1$ , then it holds that under  $[C1]_{\beta}$ ,  $[C2]_{\beta,p}$  and  $[C3]_{\beta,p}$ ,

$$\mathcal{R}_{\beta,\nu}(\hat{\theta}_{0,\beta} - \theta_0^*) = \mathcal{O}_{\mathbf{P}}(1) \tag{2.9}$$

as 
$$\nu \to 0$$
,  $N \to \infty$ ,  $M_1 \wedge M_2 \to \infty$  and  $L \to \infty$ .

See Subsection 2.4 for sufficient conditions for consistency and the asymptotic properties (2.8), (2.9) of the estimator  $\hat{\theta}_{0,\beta}$  under the given discrete data  $\mathbb{X}_{N,M}$ .

Finally, we give the sufficient conditions when  $p = \infty$ . Notice that

$$\rho_{2,x,p}^{(\text{space})} \le \rho_{1,x,p}^{(\text{space})}, \quad \phi_{1,x,p}^{(\text{space})} \le \phi_{2,x,p}^{(\text{space})}, \quad \psi_{2,x,p}^{(\text{space})} \le \psi_{1,x,p}^{(\text{space})}$$

for sufficiently large p. For  $\alpha > 0$  and  $x \in (0, 1]$ , define

$$\begin{split} \rho_x^{(\text{space})} &= 1 - \frac{1}{2x+1}, \quad \rho_x^{(\text{trunc})} = 2\bigg(1 - \frac{1}{2x+1}\bigg), \\ \phi_x^{(\text{space})} &= \frac{1}{x} \lor (3-2x), \quad \psi_{1,x}^{(\text{space})} = \frac{2}{2x+1}, \quad \psi_{2,x}^{(\text{space})} = \frac{3}{2} - x, \\ \phi_{x,\alpha}^{(\text{trunc})} &= \frac{(\alpha-1)x+2}{2x}, \quad \psi_{x,\alpha}^{(\text{trunc})} = \frac{(\alpha-1)x+2}{2x+1}. \end{split}$$

If  $p = \infty$ , the conditions  $[B1]_{\beta,p}$ ,  $[B2]_{\beta,p}$ ,  $[C2]_{\beta,p}$  and  $[C3]_{\beta,p}$  are respectively replaced by

$$[\mathbf{B1}]_{\beta,\infty} \ \nu(M_1 \wedge M_2)^{2r} N^{-\phi_{\alpha(\beta+1)}^{(\text{space})}} \to \infty \text{ as } \nu \to 0, \ N \to \infty \text{ and } M_1 \wedge M_2 \to \infty \text{ for some } r < 1,$$

$$[\mathbf{B2}]_{\beta,\infty} \ \nu L^s N^{-\phi_{\alpha(\beta+1),\alpha}^{(\mathrm{trunc})}} \to \infty \text{ as } \nu \to 0, N \to \infty \text{ and } L \to \infty \text{ for some } s < 2$$

$$\begin{split} [\mathbf{C2}]_{\beta,\infty} \ \nu(M_1 \wedge M_2)^{2r_1} N^{-\psi_{1,\alpha(\beta+1)}^{(\mathrm{space})}} \to \infty \text{ and } \nu(M_1 \wedge M_2)^{2r_2} N^{-\psi_{2,\alpha(\beta+1)}^{(\mathrm{space})}} \to \infty \text{ as } \nu \to 0, \ N \to \infty \\ \text{ and } M_1 \wedge M_2 \to \infty \text{ for some } r_1 < \rho_{\alpha(\beta+1)}^{(\mathrm{space})} \text{ and } r_2 < 1/2, \end{split}$$

$$[\mathbf{C3}]_{\beta,\infty} \ \nu L^s N^{-\psi_{\alpha(\beta+1),\alpha}^{(\mathrm{trunc})}} \to \infty \text{ as } \nu \to 0, \ N \to \infty \text{ and } L \to \infty \text{ for some } s < \rho_{\alpha(\beta+1)}^{(\mathrm{trunc})}$$

We now obtain the sufficient conditions required for consistency or the asymptotic properties (2.8), (2.9) of the estimator  $\hat{\theta}_{0,\beta}$  when  $[A1]_{-1,\infty}$  is satisfied.

#### Optimal choice of $\beta$ $\mathbf{2.4}$

In this subsection, we discuss the optimal choice of  $\beta$  for  $\alpha$ . More precisely, we fix  $\alpha > 0$  and investigate the superiority of the estimators  $\hat{\theta}_{0,\beta_1}$  and  $\hat{\theta}_{0,\beta_2}$  for  $-1 < \beta_1 < \beta_2 \le \frac{1}{\alpha} - 1$ . For fixed  $\alpha > 0$ , let  $\frac{1}{2\alpha} - 1 < \beta' < \frac{1}{\alpha} - 1$ . Because of (2.6) and the relation

$$1 \lesssim -\log\nu \lesssim \nu^{\alpha(\beta'+1)-1} \lesssim (-\nu\log\nu)^{-1/2} \lesssim \nu^{-1/2}, \quad \nu \to 0,$$

the convergence rate  $\mathcal{R}_{\beta,\nu}^{-1}$  of the estimator  $\hat{\theta}_{0,\beta}^{(\text{cont})}$  is faster as  $\beta$  gets smaller. However, the convergence rates of the estimators  $\hat{\theta}_{0,\beta_1}^{(\text{cont})}$  and  $\hat{\theta}_{0,\beta_2}^{(\text{cont})}$  are both  $\nu^{1/2}$  for  $-1 < \beta_1 < \beta_2 < \frac{1}{2\alpha} - 1$ , and we cannot determine the superiority of the estimators by only using (2.6). For this reason, we need to examine the properties of  $\mathcal{R}_{\beta,\nu}$  in more detail.

The following proposition shows the superiority of the estimators  $\hat{\theta}_{0,\beta_1}^{(\text{cont})}$  and  $\hat{\theta}_{0,\beta_2}^{(\text{cont})}$  for -1 < 1 $\beta_1 < \beta_2 < \frac{1}{2\alpha} - 1.$ 

**Proposition 2.3.** Assume that  $[A1]_{-1,2}$  holds. Then, for  $\alpha > 0$  and  $\nu \in (0,1)$ ,

$$\mathcal{R}_{\beta,\nu} = \frac{\sum_{l_1,l_2 \ge 1} f_{l_1,l_2}(2(\nu\lambda_{l_1,l_2} - \theta_0) : \alpha, \sigma^2, X_0)\mu_{l_1,l_2}^{-\alpha(\beta+1)}}{\sqrt{\sum_{l_1,l_2 \ge 1} f_{l_1,l_2}(2(\nu\lambda_{l_1,l_2} - \theta_0) : \alpha, \sigma^2, X_0)\mu_{l_1,l_2}^{-2\alpha(\beta+1)}}},$$

where  $f_{l_1,l_2}(x:\alpha,\sigma^2,X_0) = \mu_{l_1,l_2}^{\alpha} \mathbb{E}[\langle X_0,e_{l_1,l_2} \rangle^2] f_1(x) + \sigma^2 f_2(x),$ 

$$f_1(x) = \begin{cases} \frac{1 - e^{-x}}{x}, & x \neq 0, \\ 1, & x = 0, \end{cases} \quad f_2(x) = \begin{cases} \frac{x - 1 + e^{-x}}{x^2}, & x \neq 0, \\ \frac{1}{2}, & x = 0 \end{cases}$$

and the function  $\beta \mapsto \mathcal{R}_{\beta,\nu}$  is decreasing on  $(-1, \frac{1}{\alpha} - 1]$ .

Proposition 2.3 shows that for  $-1 < \beta_1 < \beta_2 < \frac{1}{2\alpha} - 1$ ,  $\hat{\theta}_{0,\beta_1}^{(\text{cont})}$  is superior to  $\hat{\theta}_{0,\beta_2}^{(\text{cont})}$  in the sense that the asymptotic variance is smaller. Indeed, by setting  $\mathcal{I}_{\beta} = \lim_{\nu \to 0} (\sigma^*)^{-2\nu} \mathcal{R}_{\beta,\nu}^2$  for  $-1 < \beta < \frac{1}{2\alpha} - 1$ , we see from Theorem 2.1 and Proposition 2.3 that  $\nu^{-1/2}(\hat{\theta}_{0,\beta}^{(\text{cont})} - \theta_0^*) \xrightarrow{d} N(0, \mathcal{I}_{\beta}^{-1})$ and  $\mathcal{I}_{\beta_1}^{-1} < \mathcal{I}_{\beta_2}^{-1}$ .

Theorem 2.2 indicates that the choice of  $\beta$  in the estimator  $\hat{\theta}_{0,\beta}$  depends on the high frequency

spatio-temporal data  $\mathbb{X}_{N,M}$ . We thus consider how to choose  $\beta$  from the given data  $\mathbb{X}_{N,M}$ . Because  $x \mapsto \rho_x^{(\text{time})}$ ,  $x \mapsto \tau_{j,x,p}^{(\text{space})}$  and  $x \mapsto \rho_{j,x,p}^{(\text{space})}$  are increasing and  $x \mapsto \phi_{j,x,p}^{(\text{space})}$  and  $x \mapsto \psi_{j,x,p}^{(\text{space})}$  are decreasing for  $p \ge 2$  and j = 1, 2, we see that the conditions required for consistency or the asymptotic properties (2.8), (2.9) of the estimator  $\hat{\theta}_{0,\beta}$  get more relaxed as  $\beta$ increases under a fixed  $\alpha$ . Therefore, we show the lower bound of  $\beta$  for the given data  $\mathbb{X}_{N,M}$ . For simplicity, let  $p = \infty$ . For  $\alpha$ , n, m > 0, define

$$\beta_{\alpha,n,m}^{(\text{cons})} = \begin{cases} \frac{n}{\alpha(m-1)} - 1, & m \ge 2n+1, \\ \frac{1}{2\alpha} (3 - \frac{m-1}{n}) - 1, & m < 2n+1, \end{cases}$$

and  $\beta_{\alpha,n,m}^{(asym)} = \beta_{\alpha,n}^{(as-t)} \vee \beta_{\alpha,n,m}^{(as-sp)}$ , where

$$\beta_{\alpha,n}^{(\text{as-t})} = \begin{cases} \frac{1}{2\alpha n} - 1, & n \ge 1, \\ \frac{1}{\alpha} (1 - \frac{n}{2}) - 1, & n < 1, \end{cases} \quad \beta_{\alpha,n,m}^{(\text{as-sp})} = \frac{1}{2\alpha} \left( \frac{1 + 2n}{m - 1} \vee \frac{3n - m + 2}{n} \right) - 1, \quad m > 1.$$

We then have the following corollary of Theorem 2.2 with  $p = \infty$ . Note that as mentioned in Remark 3-(2), the condition  $[A1]_{-1,\infty}$  is satisfied if the initial value  $X_0$  is deterministic and smooth.

**Corollary 2.4.** Let  $\alpha > 0$  and  $-1 < \beta \leq \frac{1}{\alpha} - 1$ . Assume that  $[A1]_{-1,\infty}$  holds. For m > n > 0 and  $\ell > \frac{n(\alpha-1)}{4} \vee 0$ , let  $N = \nu^{-n}$ ,  $(M_1 \wedge M_2)^2 = \nu^{-m}$  and  $L = \nu^{-\ell}$ .

(1) The estimator  $\hat{\theta}_{0,\beta}$  is consistent as  $\nu \to 0$  if  $\ell > \frac{1+n\phi_{\alpha(\beta+1),\alpha}^{(\text{trunc})}}{2}$  and  $\beta > \beta_{\alpha,n,m}^{(\text{cons})}$  for m > n+1.

(2) The estimator  $\hat{\theta}_{0,\beta}$  has the asymptotic property (2.8) or (2.9) as  $\nu \to 0$  if  $\ell > \frac{1+n\psi_{\alpha(\beta+1),\alpha}^{(\text{trunc})}}{\rho_{\alpha(\beta+1)}^{(\text{trunc})}}$ and  $\beta > \beta_{\alpha,n,m}^{(\text{asym})}$  for m > n + 2.

- **Remark 4.** (1) We find from Corollary 2.4 that the lower bounds of  $\beta$  required for consistency and the asymptotic properties (2.8), (2.9) are given by  $\beta_{\alpha,n,m}^{(cons)}$  and  $\beta_{\alpha,n,m}^{(asym)}$ , respectively, and from Proposition 2.3 that  $\beta$  close to the lower bound gives a better estimator if  $\nu$  is sufficiently small.
  - (2) Corollary 2.4 shows that we need to increase the number of spatial observations relatively more than the number of temporal observations in order to obtain the estimator  $\hat{\theta}_{0,\beta}$  with consistency or the asymptotic properties (2.8), (2.9) as well as Tonaki et al. [35] does.
  - (3) The lower bounds  $\beta_{\alpha,n,m}^{(cons)}$  and  $\beta_{\alpha,n,m}^{(asym)}$  are both decreasing for m. This fact means that the lower bounds get small as the number of spatial observations increases, and we can obtain a better estimator.

# **2.5** Estimation of $\sigma^2$

In this subsection, we give an estimator of the volatility parameter  $\sigma^2$  under  $\nu \to 0$  based on high frequency spatio-temporal data. We consider parametric estimation of  $\sigma^2$  in line with the approach of Tonaki et al. [34].

For a fixed  $b \in (0, 1/2)$ , let  $\mathbb{X}_{N,m}^{(\text{space})} = \{X_{t_i}(\widetilde{y}_j, \widetilde{z}_k)\}_{0 \le i \le N, 0 \le j \le m_1, 0 \le k \le m_2}$  be spatially thinned data of  $\mathbb{X}_{N,M}$  such that  $\widetilde{y}_0, \widetilde{z}_0 \ge b$ ,  $\widetilde{y}_{m_1}, \widetilde{z}_{m_2} \le 1 - b$ ,  $\widetilde{y}_j = \widetilde{y}_0 + j\delta$  and  $\widetilde{z}_k = \widetilde{z}_0 + k\delta$ , where  $\delta = \frac{\widetilde{y}_{m_1} - \widetilde{y}_0}{m_1} = \frac{\widetilde{z}_{m_2} - \widetilde{z}_0}{m_2}$ ,  $m_1 m_2 = \mathcal{O}(\nu^{-1}N)$  and  $N = \mathcal{O}(\nu m_1 m_2)$ . For the thinned data  $\mathbb{X}_{N,m}^{(\text{space})}$ , define the triple increment

$$\begin{split} T_{i,j,k}X &= X_{t_i}(\widetilde{y}_j, \widetilde{z}_k) - X_{t_i}(\widetilde{y}_{j-1}, \widetilde{z}_k) - X_{t_i}(\widetilde{y}_j, \widetilde{z}_{k-1}) + X_{t_i}(\widetilde{y}_{j-1}, \widetilde{z}_{k-1}) \\ &- X_{t_{i-1}}(\widetilde{y}_j, \widetilde{z}_k) + X_{t_{i-1}}(\widetilde{y}_{j-1}, \widetilde{z}_k) + X_{t_{i-1}}(\widetilde{y}_j, \widetilde{z}_{k-1}) - X_{t_{i-1}}(\widetilde{y}_{j-1}, \widetilde{z}_{k-1}). \end{split}$$

We denote the Bessel function of the first kind of order 0 by

$$J_0(x) = 1 + \sum_{k \ge 1} \frac{(-1)^k}{(k!)^2} \left(\frac{x}{2}\right)^{2k}$$

We assume  $\theta_0 \leq 0$  so that the operator  $\mathcal{A}_{\theta_0,\nu} = \nu \mathcal{A} - \theta_0$  is positive definite and self-adjoint. For simplicity, we assume the following initial condition as in [34].

**[A2]** The initial value  $X_0$  is non-random and  $\|\mathcal{A}^{(1+\alpha)/2}X_0\| < \infty$ .

Note that we can consider the general initial condition as in [33].

Let  $\Delta = 1/N$ . For  $\alpha \in (0,2)$  and  $\delta/\sqrt{\nu\Delta} \equiv r \in (0,\infty)$ , define

$$\psi_{r,\alpha} = \frac{2}{\pi} \int_0^\infty \frac{1 - e^{-x^2}}{x^{1+2\alpha}} \big( J_0(\sqrt{2}rx) - 2J_0(rx) + 1 \big) \mathrm{d}x.$$

In the same way as Proposition 2.1 in [34], we obtain the following result.

**Proposition 2.5.** Let  $\alpha \in (0,2)$  and  $\delta/\sqrt{\nu\Delta} \equiv r \in (0,\infty)$ . Under [A2], it follows that

$$\mathbb{E}[(T_{i,j,k}X)^2] = \nu^{\alpha-1} \Delta^{\alpha} \sigma^2 \mathrm{e}^{-\kappa(\widetilde{y}_{j-1}+\widetilde{y}_j)/2} \mathrm{e}^{-\eta(\widetilde{z}_{k-1}+\widetilde{z}_k)/2} \psi_{r,\alpha} + R_{i,j,k} + \mathcal{O}(\nu^{\alpha-1}\Delta^{1+\alpha}),$$

where  $\sum_{i=1}^{N} R_{i,j,k} = \mathcal{O}(\nu^{\alpha-1}\Delta^{\alpha})$  uniformly in j, k.

The estimator of  $\sigma^2$  is defined as

$$\hat{\sigma}^{2} = \hat{\sigma}_{\nu,N,m}^{2} = \frac{1}{N\nu^{\alpha-1}\Delta^{\alpha}} \sum_{k=1}^{m_{2}} \sum_{j=1}^{m_{1}} \sum_{i=1}^{N} (T_{i,j,k}X)^{2} e^{(-\kappa(\tilde{y}_{j-1}+\tilde{y}_{j})-\eta(\tilde{z}_{k-1}+\tilde{z}_{k}))/2} \\ \times \left(\psi_{r,\alpha} \sum_{k=1}^{m_{2}} \sum_{j=1}^{m_{1}} e^{-\kappa(\tilde{y}_{j-1}+\tilde{y}_{j})-\eta(\tilde{z}_{k-1}+\tilde{z}_{k})}\right)^{-1}.$$

Then, we have the following asymptotic property of the estimator  $\hat{\sigma}^2$ .

**Proposition 2.6.** Under [A2], as  $N \to \infty$ ,  $m_1 \wedge m_2 \to \infty$  and  $\nu \to 0$ ,

$$N(\hat{\sigma}^2 - (\sigma^*)^2) = \mathcal{O}_{\mathbf{P}}(1).$$

**Remark 5.** (1) The estimator  $\hat{\sigma}^2$  is defined as  $\sigma^2$  which minimizes the contrast function

$$K_{\nu,N,m}(\sigma^2) = \sum_{k=1}^{m_2} \sum_{j=1}^{m_1} \left\{ \frac{1}{N\nu^{\alpha-1}\Delta^{\alpha}} \sum_{i=1}^{N} (T_{i,j,k}X)^2 - \sigma^2 \psi_{r,\alpha} \mathrm{e}^{(-\kappa(\tilde{y}_{j-1}+\tilde{y}_j)-\eta(\tilde{z}_{k-1}+\tilde{z}_k))/2} \right\}^2.$$

This is because  $\sigma^2$  can be identified without using two rates  $r = \delta/\sqrt{\nu\Delta}$  and  $r' = \delta/\sqrt{2\nu\Delta}$  if the diffusive parameter  $\nu$  is known, see [34].

(2) The estimator  $\hat{\sigma}^2$  has the same rate as that in [34], even under  $\nu \to 0$ . This is due to the fact that the main term of  $\mathbb{E}[(T_{i,j,k}X)^2]$  is independent of  $\theta_0$ , see the proof of Lemma 4.1 in [34].

# 3 Simulations

The numerical solution of SPDE (1.2) is generated by

$$\tilde{X}_{t_i}(y_j, z_k) = \sum_{l_1=1}^{K_1} \sum_{l_2=1}^{K_2} x_{l_1, l_2}(t_i) e_{l_1, l_2}(y_j, z_k), \quad i = 1, \dots, N, j = 1, \dots, M_1, k = 1, \dots, M_2$$

with

$$\mathrm{d}x_{l_1,l_2}(t) = -\lambda_{l_1,l_2}^{(\theta_0)} x_{l_1,l_2}(t) \mathrm{d}t + \sigma \mu_{l_1,l_2}^{-\alpha/2} \mathrm{d}w_{l_1,l_2}(t), \quad x_{l_1,l_2}(0) = \langle X_0, e_{l_1,l_2} \rangle,$$

where  $\lambda_{l_1,l_2}^{(\theta_0)} = \nu \lambda_{l_1,l_2} - \theta_0$ . In this simulation, the true values of parameters  $(\theta_0^*, \kappa, \eta, \sigma^*) = (0, 1, 1, 1)$ . We set  $N = 10^2$ ,  $M_1 = M_2 = 200$ ,  $K_1 = K_2 = 10^4$ ,  $X_0 = 0$ ,  $\alpha = 0.5$ ,  $\nu = 0.1$ . We used R language to compute the estimators of Theorem 2.2 and Proposition 2.6. The number of iterations is 200.

First, we estimated  $\theta_0$ . Table 1 is the simulation results of the mean and the standard deviation (s.d.) of  $\hat{\theta}_{0,\beta}$ . We set that  $(N, M_1, M_2, \alpha, \nu, \beta, L) = (10^2, 200, 200, 0.5, 0.1, 0.6, 32)$ . In this case,  $n = 2, m \approx 4.6$  and  $\ell \approx 1.505$  in Corollary 2.4, thus  $\ell > \frac{1+n\phi_{\alpha(\beta+1),\alpha}^{(trunc)}}{2} = 1.5$  and  $\beta > \beta_{\alpha,n,m}^{(cons)} \approx 0.20$ . It seems from Table 1 that the bias of  $\hat{\theta}_{0,\beta}$  is very small and  $\hat{\theta}_{0,\beta}$  has a good performance.

Next, we estimated  $\sigma^2$ . Table 2 is the simulation results of the mean and the standard deviation (s.d.) of  $\hat{\sigma}^2$  with  $(N, m_1, m_2, \alpha, \nu) = (10^2, 30, 30, 0.5, 0.1)$ . We see from Table 2 that the bias of  $\hat{\sigma}^2$  is very small.

Table 1: Simulation results of  $\hat{\theta}_{0,\beta}$ 

	$\hat{ heta}_{0,eta}$
true value	0
mean	-0.029
s.d.	(0.274)

Table 2: Simulation results of  $\hat{\sigma}^2$ 

	$\hat{\sigma}^2$
true value	1
mean	0.987
s.d.	(0.008)

# 4 Proofs

We set the following notation.

- 1.  $\|\cdot\|_{\infty}$  denotes the uniform norm on D.
- 2. For a continuous square integrable martingale  $\{M_t\}_{t\geq 0}$  with  $M_0 = 0$ ,  $\langle\!\langle M \rangle\!\rangle_t$  stands for the quadratic variation of  $\{M_t\}_{t\geq 0}$ .
- 3. For  $\gamma \in [-1,\infty)$ , define  $\mathcal{U}_{\gamma} = \{\widetilde{Q}^{(\gamma+1)/2}v | v \in \mathcal{H}\}$  with the induced norm  $\|\widetilde{Q}^{-(\gamma+1)/2}u\|$ ,  $u \in \mathcal{U}_{\gamma}$ .
- 4. *Hilbert-Schmidt space*. For separable Hilbert spaces  $\mathcal{K}_1$  and  $\mathcal{K}_2$ , let  $\mathrm{HS}(\mathcal{K}_1; \mathcal{K}_2) = \{L : \mathcal{K}_1 \to \mathcal{K}_2 | L \text{ is bounded linear, } \|L\|_{\mathrm{HS}(\mathcal{K}_1; \mathcal{K}_2)} < \infty\}$ , where

$$||L||_{\mathrm{HS}(\mathcal{K}_1;\mathcal{K}_2)} = \sqrt{\sum_{j\geq 1} ||L\phi_j||_{\mathcal{K}_2}^2}$$

and  $\{\phi_j\}_{j\in\mathbb{N}}$  is a complete orthonormal system of  $\mathcal{K}_1$ . We write  $\operatorname{HS}(\mathcal{K}_1; \mathcal{K}_2)$  as  $\operatorname{HS}(\mathcal{K}_1)$  if  $\mathcal{K}_2 = \mathcal{H}$ .

5. Hölder space. For  $\gamma \in (0, 1]$ , define  $C^{\gamma} = \{u : D \to \mathbb{R} | ||u||_{C^{\gamma}} < \infty\}$ , where

$$\|u\|_{\mathcal{C}^{\gamma}} = \|u\|_{\infty} + \sup_{oldsymbol{x}
eq oldsymbol{y} \in D} rac{|u(oldsymbol{x}) - u(oldsymbol{y})|}{|oldsymbol{x} - oldsymbol{y}|^{\gamma}}.$$

6. Sobolev space. For  $\gamma \in (0,1)$  and  $q \geq 2$ , let  $\mathcal{W}^{\gamma,q} = \{u \in \mathcal{L}^q | \|u\|_{\mathcal{W}^{\gamma,q}} < \infty\}$ , where

$$\|u\|_{\mathcal{W}^{\gamma,q}} = \left(\|u\|_{\mathcal{L}^q}^q + \int_D \int_D \frac{|u(\boldsymbol{x}) - u(\boldsymbol{y})|^q}{|\boldsymbol{x} - \boldsymbol{y}|^{2+\gamma q}} \mathrm{d}\boldsymbol{x} \mathrm{d}\boldsymbol{y}\right)^{1/q}.$$

7. Let  $\bar{v}(y,z) = e^{\kappa y + \eta z}$  and  $v(y,z) = 1/\bar{v}(y,z)$ . Note that  $\bar{v}$  is the weight function in the inner product  $\langle \cdot, \cdot \rangle$ .

## 4.1 Proof of Theorem 2.1

For  $\beta > -1$  and t > 0, define

$$J_{\beta,\nu} = \int_0^1 \|X_t\|_{\mathcal{L}^2_{\beta}}^2 \mathrm{d}t, \quad K_{\beta,\nu} = \int_0^1 \langle X_t, \mathrm{d}X_t + \nu \mathcal{A}X_t \mathrm{d}t \rangle_{\mathcal{L}^2_{\beta}}$$
$$M_{\beta,\nu,t} = \int_0^t \langle X_s, \mathrm{d}W_s^Q \rangle_{\mathcal{L}^2_{\beta}}, \quad \mathcal{M}_{\beta,\nu,t} = \frac{M_{\beta,\nu,t}}{\sqrt{\mathbb{E}[\langle \langle M_{\beta,\nu} \rangle \rangle_t]}}.$$

From  $\mathcal{R}_{\beta,\nu} = \mathbb{E}[J_{\beta,\nu}]/\sqrt{\mathbb{E}[J_{2\beta+1,\nu}]}$  and Proposition 4.1 below, one can easily show the first claim of Theorem 2.1.

We next show (2) of Theorem 2.1. Note that  $\{M_{\beta,\nu,t}\}_{t\in\mathbb{T}}$  is continuous square integrable martingale,  $\hat{\theta}_{0,\beta}^{(\text{cont})} = K_{\beta,\nu}/J_{\beta,\nu}$  and

$$K_{\beta,\nu} = \int_0^1 \langle X_t, \theta_0^* X_t \mathrm{d}t + \sigma^* \mathrm{d}W_t^Q \rangle_{\mathcal{L}^2_\beta} = \theta_0^* J_{\beta,\nu} + \sigma^* M_{\beta,\nu,1}$$

under  $\mathbf{P}_{\theta_0^*,\sigma^*}$ . Since  $\langle\!\langle M_{\beta,\nu} \rangle\!\rangle_1 = J_{2\beta+1,\nu}$  and

$$\hat{\theta}_{0,\beta}^{(\text{cont})} - \theta_0^* = \frac{\sigma^* M_{\beta,\nu,1}}{J_{\beta,\nu}} = \sigma^* \times \frac{\mathbb{E}[J_{\beta,\nu}]}{J_{\beta,\nu}} \times \frac{\sqrt{\mathbb{E}[\langle\!\langle M_{\beta,\nu} \rangle\!\rangle_1]}}{\mathbb{E}[J_{\beta,\nu}]} \times \mathcal{M}_{\beta,\nu,1}$$

it holds that

$$\mathcal{R}_{\beta,\nu}(\hat{\theta}_{0,\beta}^{(\text{cont})} - \theta_0^*) = \sigma^* \times \frac{\mathbb{E}[J_{\beta,\nu}]}{J_{\beta,\nu}} \times \mathcal{M}_{\beta,\nu,1}.$$
(4.1)

Since it follows from Proposition 4.2 below that for  $-1 < 2\beta + 1 \leq \frac{1}{\alpha} - 1$ ,

$$\langle\!\langle \mathcal{M}_{\beta,\nu} \rangle\!\rangle_1 = \frac{\langle\!\langle M_{\beta,\nu} \rangle\!\rangle_1}{\mathbb{E}[\langle\!\langle M_{\beta,\nu} \rangle\!\rangle_1]} = \frac{J_{2\beta+1,\nu}}{\mathbb{E}[J_{2\beta+1,\nu}]} \xrightarrow{\mathbf{P}} 1$$

as  $\nu \to 0$ , we find from Theorem A.1 in [2] that  $\mathcal{M}_{\beta,\nu,1} \xrightarrow{d} N(0,1)$  as  $\nu \to 0$  for  $-1 < \beta \leq \frac{1}{2\alpha} - 1$ . With Slutsky's theorem and Proposition 4.2, we complete the proof of (2).

Finally, we verify (3) of Theorem 2.1. Using Proposition 4.1, we have  $\mathbb{E}[M_{\beta,\nu,1}^2] = \mathbb{E}[\langle M_{\beta,\nu} \rangle_1] = \mathbb{E}[J_{2\beta+1,\nu}] < \infty$  for any  $\nu > 0$  and  $-1 < \beta \leq \frac{1}{\alpha} - 1$ , and we obtain  $\mathbb{E}[\mathcal{M}_{\beta,\nu,1}^2] = 1$  for  $\nu > 0$  and  $-1 < \beta \leq \frac{1}{\alpha} - 1$ . For any  $\epsilon > 0$ , we can choose a constant  $C > \epsilon^{-1/2}$  such that

$$\sup_{\nu>0} \mathbf{P}(|\mathcal{M}_{\beta,\nu,1}| > C) \le C^{-2} \sup_{\nu>0} \mathbb{E}[\mathcal{M}_{\beta,\nu,1}^2] = C^{-2} < \epsilon,$$

which implies  $\mathcal{M}_{\beta,\nu,1} = \mathcal{O}_{\mathbf{P}}(1)$  for  $-1 < \beta \leq \frac{1}{\alpha} - 1$ . By (4.1) and Proposition 4.2, we get the desired result.

For  $\beta > -1$  and  $0 \le u \le t \le 1$ , let

$$\overline{X}_{t,u} = \sigma^* \int_u^t S_{\nu(t-s)} dW_s^Q, \quad \overline{X}_t = \overline{X}_{t,0}, \quad \widetilde{X}_t = X_t - \overline{X}_t$$
$$I_{\beta,\nu,t} = \int_0^t \|\overline{X}_s\|_{\mathcal{L}^2_{\beta}}^2 ds, \quad I_{\beta,\nu} = I_{\beta,\nu,1},$$
$$\varphi_{\beta}(\nu,t) = \int_0^t \|S_{\nu s}\|_{\mathrm{HS}(\mathcal{U}_{\beta})}^2 ds, \quad \varphi_{\beta}(\nu) = \varphi_{\beta}(\nu,1).$$

**Proposition 4.1.** Let  $\alpha > 0$  and  $\beta > -1$ . Then,  $S_t \in HS(\mathcal{U}_\beta)$  for t > 0 and

$$\varphi_{\beta}(\nu) \sim \begin{cases} \nu^{\alpha(\beta+1)-1}, & \alpha(\beta+1) < 1, \\ -\log\nu, & \alpha(\beta+1) = 1, \\ 1, & \alpha(\beta+1) > 1 \end{cases}$$
(4.2)

as  $\nu \to 0$ . Furthermore, if  $-1 < \beta \leq \frac{1}{\alpha} - 1$ , then under  $[A1]_{\beta,2}$ ,

$$\mathbb{E}[J_{\beta,\nu}] \sim \mathbb{E}[I_{\beta,\nu}] \sim \varphi_{\beta}(\nu)$$

 $as \ \nu \to 0.$ 

**Proposition 4.2.** Let  $\alpha > 0$ . If  $-1 < \beta \leq \frac{1}{\alpha} - 1$ , then under  $[A1]_{\beta,2}$ ,

$$\frac{\mathbb{E}[J_{\beta,\nu}]}{J_{\beta,\nu}} \xrightarrow{\mathbf{P}} 1$$

 $as \ \nu \to 0.$ 

The rest of this subsection is devoted to the proofs of these two propositions.

#### 4.1.1 Proof of Proposition 4.1

The proof consists of four steps. Specifically, we will show  $S_t \in \mathrm{HS}(\mathcal{U}_\beta)$  and (4.2) in Step 1, prove  $\mathbb{E}[I_{\beta,\nu}] \sim \varphi_\beta(\nu)$  in Step 2, verify  $\mathbb{E}[J_{\beta,\nu}] \lesssim \varphi_\beta(\nu)$  in Step 3 and show  $\mathbb{E}[J_{\beta,\nu}] \gtrsim \mathbb{E}[I_{\beta,\nu}]$  under  $\nu \to 0$  in Step 4.

Step 1. We can check immediately that  $S_t \in \mathrm{HS}(\mathcal{U}_\beta)$  for t > 0 since  $||S_t||_{\mathrm{HS}(\mathcal{U}_\beta)} = ||S_t \widetilde{Q}^{(\beta+1)/2}||_{\mathrm{HS}(\mathcal{H})}$ and

$$\|S_t \widetilde{Q}^{\gamma}\|_{\mathrm{HS}(\mathcal{H})}^2 = \sum_{l_1, l_2 \ge 1} \mathrm{e}^{-2t\lambda_{l_1, l_2}} \mu_{l_1, l_2}^{-2\alpha\gamma} \le \sum_{l_1, l_2 \ge 1} \frac{1}{2t\lambda_{l_1, l_2} \mu_{l_1, l_2}^{2\alpha\gamma}} < \infty$$

for  $\gamma > 0$  and t > 0. We next show (4.2). Note that  $\nu \in (0,1)$ . It follows from  $\frac{1-e^{-ax}}{x} \sim \frac{1}{x} \wedge a$ (a > 0) that for  $t \in (0,1]$  and  $\gamma > 0$ ,

$$\begin{split} \int_{0}^{t} \|S_{\nu s} \widetilde{Q}^{\gamma}\|_{\mathrm{HS}(\mathcal{H})}^{2} \mathrm{d}s &= \int_{0}^{t} \sum_{l_{1}, l_{2} \geq 1} \mathrm{e}^{-2\nu s \lambda_{l_{1}, l_{2}}} \mu_{l_{1}, l_{2}}^{-2\alpha \gamma} \mathrm{d}s \\ &= \sum_{l_{1}, l_{2} \geq 1} \frac{1 - \mathrm{e}^{-2\nu t \lambda_{l_{1}, l_{2}}}}{2\nu \lambda_{l_{1}, l_{2}}} \mu_{l_{1}, l_{2}}^{-2\alpha \gamma} \\ &\sim \sum_{l_{1}, l_{2} \geq 1} \left( \frac{1}{\nu (l_{1}^{2} + l_{2}^{2})} \wedge t \right) \frac{1}{(l_{1}^{2} + l_{2}^{2})^{2\alpha \gamma}} \\ &= \sum_{l_{1}^{2} + l_{2}^{2} > 1/\nu t} \frac{1}{\nu (l_{1}^{2} + l_{2}^{2})^{1+2\alpha \gamma}} + \sum_{1 < l_{1}^{2} + l_{2}^{2} \leq 1/\nu t} \frac{t}{(l_{1}^{2} + l_{2}^{2})^{2\alpha \gamma}} \\ &\sim \nu^{2\alpha \gamma - 1} t^{2\alpha \gamma} + \begin{cases} \nu^{2\alpha \gamma - 1} t^{2\alpha \gamma}, & 2\alpha \gamma < 1, \\ -t \log(\nu t), & 2\alpha \gamma = 1, \\ t, & 2\alpha \gamma > 1. \end{cases} \\ &\sim \begin{cases} \nu^{2\alpha \gamma - 1} t^{2\alpha \gamma}, & 2\alpha \gamma < 1, \\ -t \log(\nu t), & 2\alpha \gamma = 1, \\ t, & 2\alpha \gamma > 1. \end{cases} \end{split}$$

We therefore get the desired result.

Step 2. We verify that

$$\mathbb{E}[I_{\beta,\nu}] \sim \varphi_{\beta}(\nu), \tag{4.3}$$

and that for  $\tau \in (0, 1]$ ,

$$\mathbb{E}[I_{\beta,\nu}] \lesssim \int_0^\tau \varphi_\beta(\nu, s) \mathrm{d}s.$$
(4.4)

It is shown that

$$\varphi_{\beta}(\nu, t_1) \le \varphi_{\beta}(\nu, t_2) \le \frac{t_2}{t_1} \varphi_{\beta}(\nu, t_1), \quad 0 < t_1 \le t_2 \le 1$$
(4.5)

by using the facts that (i) for  $\gamma > 0$ , the function  $t \mapsto \|S_t \widetilde{Q}^{\gamma}\|_{\mathrm{HS}(\mathcal{H})}$  is decreasing in  $t \ge 0$ , and (ii) for a non-negative decreasing function  $f : [0, \infty) \to \mathbb{R}$ ,

$$\frac{1}{t_2} \int_0^{t_2} f(s) \mathrm{d}s \le \frac{1}{t_1} \int_0^{t_1} f(s) \mathrm{d}s, \quad 0 < t_1 \le t_2.$$

Since it follows from the Itô isometry (4.18) that

$$\mathbb{E}\left[\|\overline{X}_t\|_{\mathcal{L}^2_{\beta}}^2\right] = (\sigma^*)^2 \int_0^t \|S_{\nu(t-s)}\|_{\mathrm{HS}(\mathcal{U}_{\beta})}^2 \mathrm{d}s = (\sigma^*)^2 \varphi_{\beta}(\nu, t), \tag{4.6}$$

and from (4.5) that for  $\tau \in (0, 1]$ ,

$$\frac{\tau\varphi_{\beta}(\nu,\tau)}{2} \leq \int_{0}^{\tau} \varphi_{\beta}(\nu,s) \mathrm{d}s \leq \tau\varphi_{\beta}(\nu,\tau),$$

we find from (4.5) again that

$$\mathbb{E}[I_{\beta,\nu,\tau}] = \int_0^\tau \mathbb{E}\Big[\|\overline{X}_s\|_{\mathcal{L}^2_{\beta}}^2\Big] \mathrm{d}s \sim \tau \varphi_{\beta}(\nu,\tau), \quad \int_0^\tau \varphi_{\beta}(\nu,s) \mathrm{d}s \geq \frac{\tau^2 \varphi_{\beta}(\nu,1)}{2} \sim \tau^2 \mathbb{E}[I_{\beta,\nu}],$$

which imply that (4.3) and (4.4) hold.

**Step 3.** We show that  $\mathbb{E}[J_{\beta,\nu}] \lesssim \varphi_{\beta}(\nu)$ . Since

$$\widetilde{X}_t = S_{\nu t} X_0 + \theta_0^* \int_0^t S_{\nu(t-s)} (\widetilde{X}_s + \overline{X}_s) \mathrm{d}s$$

it holds that for  $t \in \mathbb{T}$ ,

$$\mathbb{E}\Big[\|\widetilde{X}_t\|_{\mathcal{L}^2_{\beta}}^2\Big] \leq 3\left\{\mathbb{E}\Big[\|S_{\nu t}X_0\|_{\mathcal{L}^2_{\beta}}^2\Big] + (\theta_0^*)^2 \left(\int_0^t \mathbb{E}\Big[\|S_{\nu(t-s)}\widetilde{X}_s\|_{\mathcal{L}^2_{\beta}}^2\Big] \mathrm{d}s + \int_0^t \mathbb{E}\Big[\|S_{\nu(t-s)}\overline{X}_s\|_{\mathcal{L}^2_{\beta}}^2\Big] \mathrm{d}s\right)\right\} \\
\leq 3\left\{\mathbb{E}\Big[\|X_0\|_{\mathcal{L}^2_{\beta}}^2\Big] + (\theta_0^*)^2 \left(\int_0^t \mathbb{E}\Big[\|\widetilde{X}_s\|_{\mathcal{L}^2_{\beta}}^2\Big] \mathrm{d}s + \int_0^t \mathbb{E}\Big[\|\overline{X}_s\|_{\mathcal{L}^2_{\beta}}^2\Big] \mathrm{d}s\right)\right\}.$$

Using Gronwall's inequality, we see from  $[A1]_{\beta,2}$ , (4.5) and (4.6) that

$$\mathbb{E}\Big[\|\widetilde{X}_t\|_{\mathcal{L}^2_{\beta}}^2\Big] \leq 3\mathrm{e}^{3(\theta_0^*)^2 t} \left(\mathbb{E}\Big[\|X_0\|_{\mathcal{L}^2_{\beta}}^2\Big] + (\theta_0^*)^2 \int_0^t \mathbb{E}\Big[\|\overline{X}_s\|_{\mathcal{L}^2_{\beta}}^2\Big] \mathrm{d}s\right) \\
\lesssim 1 + \varphi_{\beta}(\nu, t) \lesssim \varphi_{\beta}(\nu, t).$$
(4.7)

Therefore, we find from (4.5)-(4.7) that

$$\mathbb{E}[J_{\beta,\nu}] = \int_0^1 \mathbb{E}\Big[ \|X_t\|_{\mathcal{L}^2_{\beta}}^2 \Big] \mathrm{d}t \lesssim \int_0^1 \mathbb{E}\Big[ \|\overline{X}_t\|_{\mathcal{L}^2_{\beta}}^2 \Big] \mathrm{d}t + \int_0^1 \mathbb{E}\Big[ \|\widetilde{X}_t\|_{\mathcal{L}^2_{\beta}}^2 \Big] \mathrm{d}t$$

$$\lesssim \int_0^1 \varphi_\beta(\nu, t) \mathrm{d}t \le \varphi_\beta(\nu).$$

**Step 4.** We prove that  $\mathbb{E}[J_{\beta,\nu}] \gtrsim \mathbb{E}[I_{\beta,\nu}]$ . For any  $t \in \mathbb{T}$ , it follows from (4.5)-(4.7) that

$$\begin{split} \mathbb{E}\Big[\|X_t\|_{\mathcal{L}^2_{\beta}}^2\Big] &\geq \frac{\mathbb{E}\Big[\|\overline{X}_t\|_{\mathcal{L}^2_{\beta}}^2\Big]}{2} - \mathbb{E}\Big[\|\widetilde{X}_t\|_{\mathcal{L}^2_{\beta}}^2\Big] \\ &\geq \frac{\mathbb{E}\Big[\|\overline{X}_t\|_{\mathcal{L}^2_{\beta}}^2\Big]}{2} - 3\mathrm{e}^{3(\theta_0^*)^2 t} \Big(\mathbb{E}\Big[\|X_0\|_{\mathcal{L}^2_{\beta}}^2\Big] + (\theta_0^*)^2 \int_0^t \mathbb{E}\Big[\|\overline{X}_s\|_{\mathcal{L}^2_{\beta}}^2\Big] \mathrm{d}s\Big) \\ &= \frac{(\sigma^*)^2 \varphi_{\beta}(\nu, t)}{2} - 3\mathrm{e}^{3(\theta_0^*)^2 t} \Big(\mathbb{E}\Big[\|X_0\|_{\mathcal{L}^2_{\beta}}^2\Big] + (\theta_0^*\sigma^*)^2 \int_0^t \varphi_{\beta}(\nu, s) \mathrm{d}s\Big) \\ &\geq \varphi_{\beta}(\nu, t) \Big(\frac{(\sigma^*)^2}{2} - 3\mathrm{e}^{3(\theta_0^*)^2 t} (\theta_0^*\sigma^*)^2 t\Big) - 3\mathrm{e}^{3(\theta_0^*)^2 t} \mathbb{E}\Big[\|X_0\|_{\mathcal{L}^2_{\beta}}^2\Big] \\ &=: \varphi_{\beta}(\nu, t) C_1(t) - C_2(t). \end{split}$$

Fix  $\tau \in (0,1]$  such that  $C_1(\tau) > 0$ . Thanks to (4.2), we can choose a sufficiently small  $\nu$  to fulfill

$$\frac{C_1(\tau)}{2} \int_0^\tau \varphi_\beta(\nu, t) \mathrm{d}t \ge \tau C_2(\tau)$$

for  $-1 < \beta \leq \frac{1}{\alpha} - 1$ . Since  $C_1(t)$  (resp.  $C_2(t)$ ) is positive decreasing (resp. increasing) on  $[0, \tau]$ , we find from (4.4) that

$$\mathbb{E}[J_{\beta,\nu}] \ge \int_0^\tau \mathbb{E}\Big[ \|X_t\|_{\mathcal{L}^2_{\beta}}^2 \Big] \mathrm{d}t \ge \int_0^\tau \big(\varphi_{\beta}(\nu,t)C_1(t) - C_2(t)\big) \mathrm{d}t$$
$$\ge C_1(\tau) \int_0^\tau \varphi_{\beta}(\nu,t) \mathrm{d}t - \tau C_2(\tau)$$
$$\ge \frac{C_1(\tau)}{2} \int_0^\tau \varphi_{\beta}(\nu,t) \mathrm{d}t \gtrsim \mathbb{E}[I_{\beta,\nu}].$$

This completes the proof.

#### 4.1.2 Proof of Proposition 4.2

We use basic properties of the Malliavin calculus to obtain Proposition 4.2. To this end, we shall first give some properties of the Malliavin calculus.

For  $\beta > -1$ , we define  $X_{\beta,t} = \tilde{Q}^{\beta/2} X_t$ . By operating  $\tilde{Q}^{\beta/2}$  to the both sides of (2.2), it holds from Lemmas 4, 5 and 8 that under [A1]<sub> $\beta,2$ </sub>,

$$X_{\beta,t} = S_{\nu t} X_{\beta,0} + \theta_0 \int_0^t S_{\nu(t-s)} X_{\beta,s} \mathrm{d}s + \sigma \int_0^t \widetilde{Q}^{\beta/2} S_{\nu(t-s)} \mathrm{d}W_s^Q, \quad t \in \mathbb{T}$$
(4.8)

on  $L^2(\Omega; \mathcal{H})$ . Let  $\widehat{Q}_{\beta} = Q \widetilde{Q}^{\beta}$ . Since  $\widehat{Q}_{\beta}$  is a trace class operator on a Hilbert space and the  $\widehat{Q}_{\beta}$ -Wiener process  $W_t^{\widehat{Q}_{\beta}}$  has the representation

$$W_t^{\widehat{Q}_\beta} = \sum_{l_1, l_2 \ge 1} \mu_{l_1, l_2}^{-\alpha(\beta+1)/2} w_{l_1, l_2}(t) e_{l_1, l_2},$$

we can interpret (4.8) as

$$X_{\beta,t} = S_{\nu t} X_{\beta,0} + \theta_0 \int_0^t S_{\nu(t-s)} X_{\beta,s} \mathrm{d}s + \sigma \int_0^t S_{\nu(t-s)} \mathrm{d}W_s^{\widehat{Q}_\beta}, \quad t \in \mathbb{T}$$
(4.9)

by (4.21). For any  $(t, \mathbf{y}) \in \mathbb{T} \times D$ , the Malliavin derivative  $\mathcal{D}_{\beta}X_{\beta,t}(\mathbf{y})$  of  $X_{\beta,t}(\mathbf{y})$  given in (4.9) based on  $W_t^{\widehat{Q}_{\beta}}$  belongs to  $L^2(\Omega; L^2(\mathbb{T}; \mathcal{L}^2_{\beta+1}))$  and satisfies

$$\mathcal{D}_{\beta,\tau}X_{\beta,t}(\boldsymbol{y}) = \sigma G_{\nu(t-\tau)}(\boldsymbol{y},\cdot) + \theta_0 \int_0^t \int_D G_{\nu(t-s)}(\boldsymbol{y},\boldsymbol{z}) \mathcal{D}_{\beta,\tau}X_{\beta,s}(\boldsymbol{z}) \mathrm{d}\boldsymbol{z} \mathrm{d}\boldsymbol{s}$$
(4.10)

for  $\tau \in [0, t)$  and  $\mathcal{D}_{\beta, \tau} X_{\beta, t}(\boldsymbol{y}) = 0$  for  $\tau \in [t, 1]$  (see Theorem 7.1 in [31]). Then, we get

$$\sup_{(t,\boldsymbol{y})\in\mathbb{T}\times D}\mathbb{E}\left[\left(\int_{0}^{1}\|\mathcal{D}_{\beta,\tau}X_{\beta,t}(\boldsymbol{y})\|_{\mathcal{L}^{2}_{\beta+1}}^{2}\mathrm{d}\tau\right)^{2}\right]<\infty.$$
(4.11)

The proof of Proposition 4.2 is complete if we can show that for  $-1 < \beta \leq \frac{1}{\alpha} - 1$ ,

$$\mathbb{E}\left[\left(\int_{0}^{1} \|\mathcal{D}_{\beta,\tau}J_{\beta,\nu}\|_{\mathcal{L}^{2}_{\beta+1}}^{2} \mathrm{d}\tau\right)^{2}\right] \lesssim \sqrt{\mathbb{E}[J^{2}_{\beta,\nu}]}.$$
(4.12)

Indeed, it follows from the Poincaré inequality (Proposition 3.1 in [27]) that

$$\operatorname{Var}[J_{\beta,\nu}] \leq \mathbb{E}\left[\left(\int_0^1 \|\mathcal{D}_{\beta,\tau}J_{\beta,\nu}\|_{\mathcal{L}^2_{\beta+1}}^2 \mathrm{d}\tau\right)^2\right],$$

which together with (4.12) and  $\sqrt{\mathbb{E}[J^2_{\beta,\nu}]} \leq \sqrt{\operatorname{Var}[J_{\beta,\nu}]} + \mathbb{E}[J_{\beta,\nu}]$  yields

$$\operatorname{Var}[J_{\beta,\nu}] \lesssim \sqrt{\operatorname{Var}[J_{\beta,\nu}]} + \mathbb{E}[J_{\beta,\nu}].$$

Solving this quadratic inequality, one has

$$\operatorname{Var}[J_{\beta,\nu}] \lesssim 1 + \mathbb{E}[J_{\beta,\nu}] \lesssim \mathbb{E}[J_{\beta,\nu}].$$

Therefore, we see from the Chebyshev inequality and Proposition 4.1 that for any  $\epsilon > 0$ ,

$$\overline{\lim_{\nu \to 0}} \mathbf{P}\left( \left| \frac{J_{\beta,\nu}}{\mathbb{E}[J_{\beta,\nu}]} - 1 \right| > \epsilon \right) \le \epsilon^{-2} \overline{\lim_{\nu \to 0}} \operatorname{Var}\left[ \frac{J_{\beta,\nu}}{\mathbb{E}[J_{\beta,\nu}]} \right] \lesssim \epsilon^{-2} \overline{\lim_{\nu \to 0}} \frac{1}{\mathbb{E}[J_{\beta,\nu}]} = 0$$

if  $\alpha > 0$  and  $-1 < \beta \leq \frac{1}{\alpha} - 1$ . We prove (4.12). Note that the chain rule of Malliavin calculus (Proposition 1.2.3 in [28]) yields

$$\mathcal{D}_{\beta,\tau}J_{\beta,\nu} = 2\int_0^1\int_D X_{\beta,t}(\boldsymbol{y})\mathcal{D}_{\beta,\tau}X_{\beta,t}(\boldsymbol{y})\bar{v}(\boldsymbol{y})\mathrm{d}\boldsymbol{y}\mathrm{d}t.$$

By setting

$$\xi_1(X_{\beta,t}) = \|X_{\beta,t}\bar{v}\| \ (>0), \quad \xi_2(X_{\beta,t},\tau) = \left\| \int_D \frac{X_{\beta,t}(\boldsymbol{y})\bar{v}(\boldsymbol{y})}{\|X_{\beta,t}\bar{v}\|} \mathcal{D}_{\beta,\tau}X_{\beta,t}(\boldsymbol{y}) \mathrm{d}\boldsymbol{y} \right\|_{\mathcal{L}^2_{\beta+1}},$$

it holds from the Schwarz inequality and (4.16) that

$$\mathbb{E}\left[\left(\int_{0}^{1} \|\mathcal{D}_{\beta,\tau}J_{\beta,\nu}\|_{\mathcal{L}^{2}_{\beta+1}}^{2} \mathrm{d}\tau\right)^{2}\right] \\
\leq 4\mathbb{E}\left[\int_{0}^{1} \left(\int_{\tau}^{1} \left\|\int_{D} X_{\beta,t}(\boldsymbol{y})\mathcal{D}_{\beta,\tau}X_{\beta,t}(\boldsymbol{y})\bar{v}(\boldsymbol{y})\mathrm{d}\boldsymbol{y}\right\|_{\mathcal{L}^{2}_{\beta+1}}\mathrm{d}t\right)^{2}\mathrm{d}\tau\right]$$

$$\leq 4\mathbb{E}\left[\left(\int_{0}^{1}\xi_{1}(X_{\beta,t})^{2}\mathrm{d}t\right)^{2}\right]^{1/2}\mathbb{E}\left[\left(\int_{0}^{1}\int_{\tau}^{1}\xi_{2}(X_{\beta,t},\tau)^{2}\mathrm{d}t\mathrm{d}\tau\right)^{2}\right]^{1/2}$$
$$\leq 4\|\bar{v}\|_{\infty}^{2}\mathbb{E}[J_{\beta,\nu}^{2}]^{1/2}\sup_{t\in\mathbb{T}}\mathbb{E}\left[\left(\int_{0}^{1}\xi_{2}(X_{\beta,t},\tau)^{2}\mathrm{d}\tau\right)^{2}\right]^{1/2}.$$

If  $\varphi(\boldsymbol{y}) = \frac{X_{\beta,t}(\boldsymbol{y})\overline{v}(\boldsymbol{y})}{\|X_{\beta,t}\overline{v}\|}$ , then it follows from Lemma 1 below that

$$\begin{split} \sup_{t\in\mathbb{T}} \mathbb{E}\bigg[\left(\int_0^1 \xi_2(X_{\beta,t},\tau)^2 \mathrm{d}\tau\right)^2\bigg] &\leq \sup_{t\in\mathbb{T}, \|\varphi\|=1} \mathbb{E}\bigg[\left(\int_0^1 \left\|\int_D \varphi(\boldsymbol{y})\mathcal{D}_{\beta,\tau}X_{\beta,t}(\boldsymbol{y})\mathrm{d}\boldsymbol{y}\right\|_{\mathcal{L}^2_{\beta+1}}^2 \mathrm{d}\tau\right)^2\bigg] \\ &\lesssim (\sigma^* \mathrm{e}^{|\kappa|+|\eta|} \vee 1)^4 \exp\Big(4|\theta_0^*|\mathrm{e}^{|\kappa|+|\eta|}(\sigma^* \mathrm{e}^{|\kappa|+|\eta|} \vee 1)\Big), \end{split}$$

which yields the desired result.

#### 4.1.3Auxiliary results on Malliavin derivative

Here, we provide auxiliary results on Malliavin derivative  $\mathcal{D}_{\beta,\tau}X_{\beta,t}$  in order to show Proposition 4.2.

**Lemma 1.** For any  $\varphi \in \mathcal{H}$  with  $\|\varphi\| = 1$  and  $t, \tau \in \mathbb{T}$ , it holds that

$$\mathbb{E}\left[\left(\int_{0}^{1}\left\|\int_{D}\varphi(\boldsymbol{y})\mathcal{D}_{\boldsymbol{\beta},\tau}X_{\boldsymbol{\beta},t}(\boldsymbol{y})\mathrm{d}\boldsymbol{y}\right\|_{\mathcal{L}^{2}_{\boldsymbol{\beta}+1}}^{2}\mathrm{d}\tau\right)^{2}\right]\lesssim(\sigma^{*}\mathrm{e}^{|\boldsymbol{\kappa}|+|\boldsymbol{\eta}|}\vee1)^{4}\exp\left(4|\boldsymbol{\theta}_{0}^{*}|\mathrm{e}^{|\boldsymbol{\kappa}|+|\boldsymbol{\eta}|}(\sigma^{*}\mathrm{e}^{|\boldsymbol{\kappa}|+|\boldsymbol{\eta}|}\vee1)\right).$$

We approximate the Malliavin derivative  $\mathcal{D}_{\beta,\tau}X_{\beta,t}(\boldsymbol{y})$  by using the Picard iteration and show Lemma 1. Let  $\tau \in \mathbb{T}$ . For  $t \in (\tau, 1]$  and  $\boldsymbol{y} \in D$ , we define

$$\begin{cases} u_{t,\tau}^{(0)}(\boldsymbol{y}) = \sigma^* G_{\nu(t-\tau)}(\boldsymbol{y}, \cdot), \\ u_{t,\tau}^{(n)}(\boldsymbol{y}) = u_{t,\tau}^{(0)}(\boldsymbol{y}) + \theta_0^* \int_{\tau}^t \int_D G_{\nu(t-s)}(\boldsymbol{y}, \boldsymbol{z}) u_{s,\tau}^{(n-1)}(\boldsymbol{z}) \mathrm{d}\boldsymbol{z} \mathrm{d}s, \quad n \in \mathbb{N}. \end{cases}$$
(4.13)

Let  $u_{t,\tau}^{(n)}(\boldsymbol{y}) \equiv 0$  for  $t \in [0,\tau]$ ,  $\boldsymbol{y} \in D$  and  $n \in \mathbb{N} \cup \{0\}$ . It is easily shown that Lemma 1 holds from the following two lemmas.

**Lemma 2.** For  $\varphi \in \mathcal{H}$  with  $\|\varphi\| = 1$ ,  $\tau \in \mathbb{T}$ ,  $t \in (\tau, 1]$ ,  $\beta > -1$  and  $\{u_{t,\tau}^{(n)}\}_{n \in \mathbb{N}}$  given in (4.13), it follows that

$$\left\| \int_{D} \varphi(\boldsymbol{y}) u_{t,\tau}^{(n)}(\boldsymbol{y}) \mathrm{d}\boldsymbol{y} \right\|_{\mathcal{L}^{2}_{\beta+1}} \leq \sigma^{*} \|v\|_{\infty} + \sum_{k=1}^{n} \frac{(|\theta_{0}^{*}| \|v\|_{\infty}(\sigma^{*} \|v\|_{\infty} \vee 1))^{k}}{k!} (t-\tau)^{k}.$$
(4.14)

In particular,

$$\sup_{\substack{t,\tau\in\mathbb{T},\\\|\varphi\|=1,n\in\mathbb{N}}} \left\| \int_D \varphi(\boldsymbol{y}) u_{t,\tau}^{(n)}(\boldsymbol{y}) \mathrm{d}\boldsymbol{y} \right\|_{\mathcal{L}^2_{\beta+1}} \leq (\sigma^* \mathrm{e}^{|\kappa|+|\eta|} \vee 1) \exp\Big( |\theta_0^*|\mathrm{e}^{|\kappa|+|\eta|}(\sigma^* \mathrm{e}^{|\kappa|+|\eta|} \vee 1) \Big).$$

**Lemma 3.** For any  $t \in \mathbb{T}$ , it holds that

$$\lim_{n\to\infty}\sup_{\|\varphi\|=1}\mathbb{E}\left[\left(\int_0^1\left\|\int_D\varphi(\boldsymbol{y})\mathcal{D}_{\beta,\tau}X_{\beta,t}(\boldsymbol{y})\mathrm{d}\boldsymbol{y}-\int_D\varphi(\boldsymbol{y})u_{t,\tau}^{(n)}(\boldsymbol{y})\mathrm{d}\boldsymbol{y}\right\|_{\mathcal{L}^2_{\beta+1}}^2\mathrm{d}\tau\right)^2\right]=0.$$

### Proof of Lemma 2. Let

$$F_n(t,\tau,\varphi) = \left\| \int_D \varphi(\boldsymbol{y}) u_{t,\tau}^{(n)}(\boldsymbol{y}) \mathrm{d}\boldsymbol{y} \right\|_{\mathcal{L}^2_{\beta+1}}.$$

We prove (4.14) by induction.

(i) For n = 0, we have

$$F_0(t,\tau,\varphi) = \sigma^* \left\| \int_D \varphi(\boldsymbol{y}) G_{\nu(t-\tau)}(\boldsymbol{y},\cdot) \mathrm{d}\boldsymbol{y} \right\|_{\mathcal{L}^2_{\beta+1}} = \sigma^* \|S_{\nu(t-\tau)}\varphi v\|_{\mathcal{L}^2_{\beta+1}} \le \sigma^* \|v\|_{\infty}.$$

(ii) Let (4.14) be true for n. Then, we see from (4.16) that

$$\begin{aligned} F_{n+1}(t,\tau,\varphi) &\leq \left\| \int_{D} \varphi(\boldsymbol{y}) u_{t,\tau}^{(0)}(\boldsymbol{y}) \mathrm{d}\boldsymbol{y} \right\|_{\mathcal{L}^{2}_{\beta+1}} \\ &+ |\theta_{0}^{*}| \left\| \int_{D} \varphi(\boldsymbol{y}) \int_{\tau}^{t} \int_{D} G_{\nu(t-s)}(\boldsymbol{y},\boldsymbol{z}) u_{s,\tau}^{(n)}(\boldsymbol{z}) \mathrm{d}\boldsymbol{z} \mathrm{d}\boldsymbol{s} \mathrm{d}\boldsymbol{y} \right\|_{\mathcal{L}^{2}_{\beta+1}} \\ &\leq \sigma^{*} \|v\|_{\infty} + |\theta_{0}^{*}| \int_{\tau}^{t} \left\| \int_{D} u_{s,\tau}^{(n)}(\boldsymbol{z}) \int_{D} G_{\nu(t-s)}(\boldsymbol{y},\boldsymbol{z}) \varphi(\boldsymbol{y}) \mathrm{d}\boldsymbol{y} \mathrm{d}\boldsymbol{z} \right\|_{\mathcal{L}^{2}_{\beta+1}} \mathrm{d}\boldsymbol{s} \\ &= \sigma^{*} \|v\|_{\infty} + |\theta_{0}^{*}| \int_{\tau}^{t} \left\| \int_{D} u_{s,\tau}^{(n)}(\boldsymbol{z}) \widetilde{\varphi}_{s,t}(\boldsymbol{z}) \mathrm{d}\boldsymbol{z} \right\|_{\mathcal{L}^{2}_{\beta+1}} \mathrm{d}\boldsymbol{s} \\ &= \sigma^{*} \|v\|_{\infty} + |\theta_{0}^{*}| \int_{\tau}^{t} \| \widetilde{\varphi}_{s,t} \| \left\| \int_{D} u_{s,\tau}^{(n)}(\boldsymbol{z}) \widehat{\varphi}_{s,t}(\boldsymbol{z}) \mathrm{d}\boldsymbol{z} \right\|_{\mathcal{L}^{2}_{\beta+1}} \mathrm{d}\boldsymbol{s}, \end{aligned}$$

where

$$\widetilde{\varphi}_{s,t}(\cdot) = \int_{D} G_{\nu(t-s)}(\boldsymbol{y}, \cdot)\varphi(\boldsymbol{y}) \mathrm{d}\boldsymbol{y} = S_{\nu(t-s)}\varphi v(\cdot), \quad \widehat{\varphi}_{s,t} = \begin{cases} \widetilde{\varphi}_{s,t}/\|\widetilde{\varphi}_{s,t}\|, & \|\widetilde{\varphi}_{s,t}\| \neq 0, \\ 0, & \|\widetilde{\varphi}_{s,t}\| = 0. \end{cases}$$

Since  $\|\widetilde{\varphi}_{s,t}\| \leq \|v\|_{\infty}$  and  $\|\widehat{\varphi}_{s,t}\| = 1$  if  $\|\widetilde{\varphi}_{s,t}\| \neq 0$ , it follows from the induction hypothesis that

$$F_{n+1}(t,\tau,\varphi) \leq \sigma^* \|v\|_{\infty} + |\theta_0^*| \|v\|_{\infty} \int_{\tau}^t \left\{ \sigma^* \|v\|_{\infty} + \sum_{k=1}^n \frac{(|\theta_0| \|v\|_{\infty} (\sigma^* \|v\|_{\infty} \vee 1))^k}{k!} (s-\tau)^k \right\} ds$$
  
$$\leq \sigma^* \|v\|_{\infty} + |\theta_0^*| \|v\|_{\infty} (\sigma^* \|v\|_{\infty} \vee 1) (t-\tau) + \sum_{k=1}^n \frac{(|\theta_0^*| \|v\|_{\infty} (\sigma^* \|v\|_{\infty} \vee 1))^{k+1}}{(k+1)!} (t-\tau)^{k+1}$$
  
$$= \sigma^* \|v\|_{\infty} + \sum_{k=1}^{n+1} \frac{(|\theta_0^*| \|v\|_{\infty} (\sigma^* \|v\|_{\infty} \vee 1))^k}{k!} (t-\tau)^k.$$

Therefore, (4.14) is true for n + 1.

The last claim follows immediately from (4.14) and  $||v||_{\infty} \leq e^{|\kappa|+|\eta|}$ .

**Proof of Lemma 3.** For  $\varphi \in \mathcal{H}$  with  $\|\varphi\| = 1$ , let

$$\widetilde{F}_{n}(t,\tau,\varphi) = \left\| \int_{D} \varphi(\boldsymbol{y}) \mathcal{D}_{\beta,\tau} X_{\beta,t}(\boldsymbol{y}) \mathrm{d}\boldsymbol{y} - \int_{D} \varphi(\boldsymbol{y}) u_{t,\tau}^{(n)}(\boldsymbol{y}) \mathrm{d}\boldsymbol{y} \right\|_{\mathcal{L}^{2}_{\beta+1}}^{2}$$

•

It follows from (4.10), (4.13) and (4.16) that

$$\widetilde{F}_{n}(t,\tau,\varphi) = (\theta_{0}^{*})^{2} \left\| \int_{D} \varphi(\boldsymbol{y}) \int_{\tau}^{t} \int_{D} G_{\nu(t-s)}(\boldsymbol{y},\boldsymbol{z}) (\mathcal{D}_{\beta,\tau}X_{\beta,s}(\boldsymbol{z}) - u_{s,\tau}^{(n-1)}(\boldsymbol{z})) \mathrm{d}\boldsymbol{z} \mathrm{d}s \mathrm{d}\boldsymbol{y} \right\|_{\mathcal{L}^{2}_{\beta+1}}^{2}$$

$$\leq (\theta_0^*)^2 \left( \int_{\tau}^t \left\| \int_D (\mathcal{D}_{\beta,\tau} X_{\beta,s}(\boldsymbol{z}) - u_{s,\tau}^{(n-1)}(\boldsymbol{z})) \widetilde{\varphi}_{s,t}(\boldsymbol{z}) \mathrm{d}\boldsymbol{z} \right\|_{\mathcal{L}^2_{\beta+1}} \mathrm{d}s \right)^2 \\ \leq (\theta_0^*)^2 \|v\|_{\infty}^2 (t-\tau) \int_{\tau}^t \left\| \int_D (\mathcal{D}_{\beta,\tau} X_{\beta,s}(\boldsymbol{z}) - u_{s,\tau}^{(n-1)}(\boldsymbol{z})) \widehat{\varphi}_{s,t}(\boldsymbol{z}) \mathrm{d}\boldsymbol{z} \right\|_{\mathcal{L}^2_{\beta+1}}^2 \mathrm{d}s \\ \leq (\theta_0^*)^2 \|v\|_{\infty}^2 \int_{\tau}^t \widetilde{F}_{n-1}(s,\tau,\widehat{\varphi}_{s,t}) \mathrm{d}s.$$

When  $\|\widetilde{\varphi}_{s,t}\| = 0$ , it is obvious that Lemma 3 holds because  $\widetilde{F}_n(s,\tau,\widehat{\varphi}_{s,t}) = 0$ . Hereafter, let  $\|\widetilde{\varphi}_{s,t}\| \neq 0$ , that is,  $\|\widehat{\varphi}_{s,t}\| = 1$ . Let

$$\widehat{F}_n(t) = \sup_{\|\varphi\|=1} \mathbb{E}\left[\left(\int_0^1 \widetilde{F}_n(t,\tau,\varphi) \mathrm{d}\tau\right)^2\right].$$

In order to get the desired result, we prove that for any  $t \in \mathbb{T}$ ,  $\widehat{F}_n(t) \to 0$  as  $n \to \infty$ . Since

$$\begin{split} \widehat{F}_n(t) &= \sup_{\|\varphi\|=1} \mathbb{E}\left[ \left( \int_0^t \widetilde{F}_n(t,\tau,\varphi) \mathrm{d}\tau \right)^2 \right] \\ &\leq (\theta_0^*)^2 \|v\|_\infty^2 \sup_{\|\varphi\|=1} \mathbb{E}\left[ \left( \int_0^t \int_\tau^t \widetilde{F}_{n-1}(s,\tau,\varphi) \mathrm{d}s \mathrm{d}\tau \right)^2 \right] \\ &= (\theta_0^*)^2 \|v\|_\infty^2 \sup_{\|\varphi\|=1} \mathbb{E}\left[ \left( \int_0^t \int_0^s \widetilde{F}_{n-1}(s,\tau,\varphi) \mathrm{d}\tau \mathrm{d}s \right)^2 \right] \\ &\leq (\theta_0^*)^2 \|v\|_\infty^2 \int_0^t \sup_{\|\varphi\|=1} \mathbb{E}\left[ \left( \int_0^s \widetilde{F}_{n-1}(s,\tau,\varphi) \mathrm{d}\tau \right)^2 \right] \mathrm{d}s \\ &= (\theta_0^*)^2 \|v\|_\infty^2 \int_0^t \widehat{F}_{n-1}(s) \mathrm{d}s, \end{split}$$

it follows from Gronwall's lemma (Lemma 6.2 in [31]) that there exists a non-negative sequence  $\{a_n\}_{n\in\mathbb{N}}$  such that  $a_n \to 0$  and  $\widehat{F}_n(t) \leq a_n \sup_{t\in\mathbb{T}} \widehat{F}_0(t)$ . Noting that

$$\begin{split} \widehat{F}_{0}(t) &\lesssim \sup_{\|\varphi\|=1} \mathbb{E}\left[\left(\int_{0}^{t} \left\|\int_{D} \varphi(\boldsymbol{y}) \mathcal{D}_{\beta,\tau} X_{\beta,t}(\boldsymbol{y}) \mathrm{d}\boldsymbol{y}\right\|_{\mathcal{L}^{2}_{\beta+1}}^{2} \mathrm{d}\tau\right)^{2}\right] \\ &+ \sup_{\|\varphi\|=1} \left(\int_{0}^{t} \left\|\int_{D} \varphi(\boldsymbol{y}) G_{\nu(t-\tau)}(\boldsymbol{y},\cdot) \mathrm{d}\boldsymbol{y}\right\|_{\mathcal{L}^{2}_{\beta+1}}^{2} \mathrm{d}\tau\right)^{2} \\ &\leq \sup_{\|\varphi\|=1} \mathbb{E}\left[\left(\int_{0}^{t} \left(\int_{D} |\varphi(\boldsymbol{y})| \|\mathcal{D}_{\beta,\tau} X_{\beta,t}(\boldsymbol{y})\|_{\mathcal{L}^{2}_{\beta+1}}^{2} \mathrm{d}\boldsymbol{y}\right)^{2} \mathrm{d}\tau\right)^{2}\right] + (\sigma^{*} \|v\|_{\infty})^{4} \\ &\leq \sup_{\|\varphi\|=1} \left(\int_{D} \varphi(\boldsymbol{y})^{2} \mathrm{d}\boldsymbol{y}\right)^{2} \mathbb{E}\left[\left(\int_{0}^{t} \int_{D} \|\mathcal{D}_{\beta,\tau} X_{\beta,t}(\boldsymbol{y})\|_{\mathcal{L}^{2}_{\beta+1}}^{2} \mathrm{d}\boldsymbol{y} \mathrm{d}\tau\right)^{2}\right] + (\sigma^{*} \|v\|_{\infty})^{4} \\ &\leq \|v\|_{\infty}^{2} \int_{D} \mathbb{E}\left[\left(\int_{0}^{t} \|\mathcal{D}_{\beta,\tau} X_{\beta,t}(\boldsymbol{y})\|_{\mathcal{L}^{2}_{\beta+1}}^{2} \mathrm{d}\tau\right)^{2}\right] \mathrm{d}\boldsymbol{y} + (\sigma^{*} \|v\|_{\infty})^{4} \\ &\leq \|v\|_{\infty}^{2} \sup_{\boldsymbol{y}\in D} \mathbb{E}\left[\left(\int_{0}^{t} \|\mathcal{D}_{\beta,\tau} X_{\beta,t}(\boldsymbol{y})\|_{\mathcal{L}^{2}_{\beta+1}}^{2} \mathrm{d}\tau\right)^{2}\right] + (\sigma^{*} \|v\|_{\infty})^{4}, \end{split}$$

we find from (4.11) that

$$\sup_{t\in\mathbb{T}}\widehat{F}_0(t) \lesssim \|v\|_{\infty}^2 \sup_{(t,\boldsymbol{y})\in\mathbb{T}\times D} \mathbb{E}\left[\left(\int_0^1 \|\mathcal{D}_{\beta,\tau}X_{\beta,t}(\boldsymbol{y})\|_{\mathcal{L}^2_{\beta+1}}^2 \mathrm{d}\tau\right)^2\right] + (\sigma^*\|v\|_{\infty})^4 < \infty,$$

which yields  $\widehat{F}_n(t) \to 0$  as  $n \to \infty$ . This concludes the proof.

#### 4.1.4 Basic properties of closed linear operators

Here, we present some useful properties of closed operators. Note that the operator  $\widetilde{Q}^{\delta}$ ,  $\delta \in \mathbb{R}$  given in (2.1) is a closed linear operator.

Let  $(\mathcal{X}, \mathscr{A}, \mu)$  be a measure space, and  $\mathcal{B}_1, \mathcal{B}_2$  be Banach spaces. For a linear operator  $L, \mathscr{D}(L)$  denotes the domain of L.

**Lemma 4.** Let  $u : \mathcal{X} \to \mathcal{B}_1$  be a  $\mathcal{B}_1$ -valued Bochner  $\mu$ -integrable function.

(1) If  $L : \mathcal{B}_1 \to \mathcal{B}_2$  is a bounded linear operator, then Lu is  $\mathcal{B}_2$ -valued Bochner  $\mu$ -integrable function on  $\mathcal{X}$ , and

$$L \int_{\mathcal{X}} u(x) \mathrm{d}\mu(x) = \int_{\mathcal{X}} Lu(x) \mathrm{d}\mu(x).$$
(4.15)

(2) If  $L : \mathcal{B}_1 \supset \mathscr{D}(L) \to \mathcal{B}_2$  is a closed linear operator and Lu is  $\mathcal{B}_2$ -valued Bochner  $\mu$ -integrable function on  $\mathcal{X}$ , then  $\int_{\mathcal{X}} u(x) d\mu(x) \in \mathscr{D}(L)$  and (4.15) holds.

For the proof of this lemma, see Corollary V.5.2 in [40] and Theorem 3.10.16 in [11].

Let  $(\mathcal{X}, \mathscr{A}, \mu)$  be a finite measure space and  $p \geq 1$ . If  $u : \mathcal{X} \to \mathcal{B}_1$  is a  $\mathcal{B}_1$ -valued Bochner  $\mu$ -integrable function and L is the linear operator of Lemma 4 such that  $\|Lu(\cdot)\|_{\mathcal{B}_2}^p$  is Lebesgue  $\mu$ -integrable, then

$$\left\| L \int_{\mathcal{X}} u(x) \mathrm{d}\mu(x) \right\|_{\mathcal{B}_{2}}^{p} \lesssim \int_{\mathcal{X}} \|Lu(x)\|_{\mathcal{B}_{2}}^{p} \mathrm{d}\mu(x).$$
(4.16)

Indeed, one can find from (4.15), Corollary V.5.1 in [40] and the Hölder inequality that

$$\begin{split} \left\| L \int_{\mathcal{X}} u(x) \mathrm{d}\mu(x) \right\|_{\mathcal{B}_{2}}^{p} &= \left\| \int_{\mathcal{X}} Lu(x) \mathrm{d}\mu(x) \right\|_{\mathcal{B}_{2}}^{p} \\ &\leq \left( \int_{\mathcal{X}} \| Lu(x) \|_{\mathcal{B}_{2}} \mathrm{d}\mu(x) \right)^{p} \\ &\leq \mu(\mathcal{X})^{p-1} \int_{\mathcal{X}} \| Lu(x) \|_{\mathcal{B}_{2}}^{p} \mathrm{d}\mu(x) \end{split}$$

The interchange of an integral and a closed operator such as (4.15) also holds for a stochastic integral and a closed operator (see Proposition 4.30 in [30]), but we here consider the interchange of the stochastic integral  $\int_0^t (t-s)^{-\delta} S_{\nu(t-s)} dW_s^Q$  and the operator  $\tilde{Q}^{\beta/2}$ .

**Lemma 5.** Let  $t \in (0,1]$ ,  $\nu \in (0,1)$ ,  $\alpha > 0$  and  $\beta > -1$ . If  $\delta < \alpha(\beta + 1)/2$ , then  $\int_0^t (t - u)^{-\delta} S_{\nu(t-u)} dW_u^Q \in L^2(\Omega; \mathcal{L}^2_{\beta})$ ,  $\int_0^t (t - u)^{-\delta} \widetilde{Q}^{\beta/2} S_{\nu(t-u)} dW_u^Q \in L^2(\Omega; \mathcal{H})$  and

$$\widetilde{Q}^{\beta/2} \int_0^t (t-u)^{-\delta} S_{\nu(t-u)} \mathrm{d}W_u^Q = \int_0^t (t-u)^{-\delta} \widetilde{Q}^{\beta/2} S_{\nu(t-u)} \mathrm{d}W_u^Q \quad on \ L^2(\Omega; \mathcal{H}).$$
(4.17)

In particular, we obtain the Itô isometry

$$\mathbb{E}\left[\left\|\int_{0}^{t} S_{\nu(t-s)} \mathrm{d}W_{s}^{Q}\right\|_{\mathcal{L}^{2}_{\beta}}^{2}\right] = \int_{0}^{t} \|S_{\nu s}\|_{\mathrm{HS}(\mathcal{U}_{\beta})}^{2} \mathrm{d}s.$$
(4.18)

*Proof.* In the same way as the proof of Proposition 4.1, it follows that for  $\beta > -1$  and  $\delta < \alpha(\beta+1)/2$ ,

$$\int_{0}^{1} \|u^{-\delta} S_{u}\|_{\mathrm{HS}(\mathcal{U}_{\beta})}^{2} \mathrm{d}u = \sum_{l_{1}, l_{2} \ge 1} \mu_{l_{1}, l_{2}}^{-\alpha(\beta+1)} \int_{0}^{1} u^{-2\delta} \mathrm{e}^{-2u\lambda_{l_{1}, l_{2}}} \mathrm{d}u$$
$$\leq \sum_{l_{1}, l_{2} \ge 1} \mu_{l_{1}, l_{2}}^{-\alpha(\beta+1)} \int_{0}^{\infty} u^{-2\delta} \mathrm{e}^{-2u\lambda_{l_{1}, l_{2}}} \mathrm{d}u$$
$$= \frac{\Gamma(1 - 2\delta)}{2^{1 - 2\delta}} \sum_{l_{1}, l_{2} \ge 1} \frac{1}{\lambda_{l_{1}, l_{2}}^{1 - 2\delta} \mu_{l_{1}, l_{2}}^{\alpha(\beta+1)}} < \infty.$$
(4.19)

Since it follows that for  $\beta > -1$  and  $\delta < \alpha(\beta + 1)/2$ ,

$$\begin{split} \|u^{-\delta}S_{u}\|_{\mathrm{HS}(\mathcal{U}_{0};\mathcal{L}^{2}_{\beta})}^{2} &= \sum_{l_{1},l_{2}\geq1} \|u^{-\delta}S_{u}\widetilde{Q}^{1/2}e_{l_{1},l_{2}}\|_{\mathcal{L}^{2}_{\beta}}^{2} \\ &= \sum_{l_{1},l_{2}\geq1} \|\widetilde{Q}^{\beta/2}u^{-\delta}S_{u}\widetilde{Q}^{1/2}e_{l_{1},l_{2}}\|^{2} \ (=\|\widetilde{Q}^{\beta/2}u^{-\delta}S_{u}\|_{\mathrm{HS}(\mathcal{U}_{0})}^{2}) \\ &= \sum_{l_{1},l_{2}\geq1} \|u^{-\delta}S_{u}\widetilde{Q}^{(\beta+1)/2}e_{l_{1},l_{2}}\|^{2} \\ &= \|u^{-\delta}S_{u}\|_{\mathrm{HS}(\mathcal{U}_{\beta})}^{2}, \end{split}$$
(4.20)

the stochastic integrals in (4.17) can be defined and

$$\int_0^t (t-u)^{-\delta} S_{\nu(t-u)} \mathrm{d} W_u^Q \in L^2(\Omega; \mathcal{L}^2_\beta), \quad \int_0^t (t-u)^{-\delta} \widetilde{Q}^{\beta/2} S_{\nu(t-u)} \mathrm{d} W_u^Q \in L^2(\Omega; \mathcal{H}).$$

Therefore, we have the following representations

$$\int_{0}^{t} (t-u)^{-\delta} S_{\nu(t-u)} dW_{u}^{Q} = \sum_{l_{1}, l_{2} \ge 1} \mu_{l_{1}, l_{2}}^{-\alpha/2} e_{l_{1}, l_{2}} \int_{0}^{t} (t-u)^{-\delta} e^{-\nu(t-u)\lambda_{l_{1}, l_{2}}} dw_{l_{1}, l_{2}}(u),$$

$$\int_{0}^{t} (t-u)^{-\delta} \widetilde{Q}^{\beta/2} S_{\nu(t-u)} dW_{u}^{Q} = \sum_{l_{1}, l_{2} \ge 1} \mu_{l_{1}, l_{2}}^{-\alpha(\beta+1)/2} e_{l_{1}, l_{2}} \int_{0}^{t} (t-u)^{-\delta} e^{-\nu(t-u)\lambda_{l_{1}, l_{2}}} dw_{l_{1}, l_{2}}(u) \quad (4.21)$$

for  $t\in\mathbb{T}$  and obtain

$$\widetilde{Q}^{\beta/2} \int_0^t (t-u)^{-\delta} S_{\nu(t-u)} \mathrm{d}W_u^Q = \sum_{l_1, l_2 \ge 1} \mu_{l_1, l_2}^{-\alpha(\beta+1)/2} e_{l_1, l_2} \int_0^t (t-u)^{-\delta} \mathrm{e}^{-\nu(t-u)\lambda_{l_1, l_2}} \mathrm{d}w_{l_1, l_2}(u)$$
$$= \int_0^t (t-u)^{-\delta} \widetilde{Q}^{\beta/2} S_{\nu(t-u)} \mathrm{d}W_u^Q$$

on  $L^2(\Omega; \mathcal{H})$ . By using (4.17), the Itô isometry (4.30) in [30] and (4.20),

$$\mathbb{E}\left[\left\|\int_{0}^{t} S_{\nu(t-s)} \mathrm{d}W_{s}^{Q}\right\|_{\mathcal{L}_{\beta}^{2}}^{2}\right] = \mathbb{E}\left[\left\|\widetilde{Q}^{\beta/2} \int_{0}^{t} S_{\nu(t-s)} \mathrm{d}W_{s}^{Q}\right\|^{2}\right]$$
$$= \mathbb{E}\left[\left\|\int_{0}^{t} \widetilde{Q}^{\beta/2} S_{\nu(t-s)} \mathrm{d}W_{s}^{Q}\right\|^{2}\right]$$
$$= \int_{0}^{t} \|\widetilde{Q}^{\beta/2} S_{\nu s}\|_{\mathrm{HS}(\mathcal{U}_{0})}^{2} \mathrm{d}s.$$
$$= \int_{0}^{t} \|S_{\nu s}\|_{\mathrm{HS}(\mathcal{U}_{\beta})}^{2} \mathrm{d}s.$$

## 4.2 Proof of Theorem 2.2

In this subsection, we show in three steps that the estimators  $\hat{\theta}_{0,\beta}$  and  $\hat{\theta}_{0,\beta}^{(\text{cont})}$  are asymptotically equivalent as  $\nu \to 0$ ,  $N \to \infty$ ,  $M_1 \wedge M_2 \to \infty$  and  $L \to \infty$ . Auxiliary results on the discretization error can be found in the end of this subsection.

#### 4.2.1 Control of temporal discretization error

Consider a temporal discretization of the estimator  $\hat{\theta}_{0,\beta}^{(\text{cont})}$  given in (2.4). That is, we construct an estimator of  $\theta_0$  using discrete temporal data  $\mathbb{X}_N^{(\text{disc-cont})} = \{X_{t_i}(y,z)\}_{0 \le i \le N, (y,z) \in \overline{D}}$ . We use the property

$$-\mathcal{A}u = \lim_{h \downarrow 0} \frac{S_h u - u}{h}, \quad u \in \mathscr{D}(\mathcal{A})$$

and discretize the terms  $dX_t$  and  $\nu \mathcal{A}X_t dt$  in the numerator of the estimator  $\hat{\theta}_{0,\beta}^{(\text{cont})}$  with  $X_{ti} - X_{ti-1}$ and  $X_{ti-1} - S_{\nu(t_i-t_{i-1})}X_{ti-1}$  for  $t \in [t_{i-1}, t_i)$ , respectively. We then define the estimator of  $\theta_0$  based on  $\mathbb{X}_N^{(\text{disc-cont})}$  by

$$\hat{\theta}_{0,\beta}^{(\text{disc-cont})} = \hat{\theta}_{0,\beta,\nu,N}^{(\text{disc-cont})} = \frac{\sum_{i=1}^{N} \langle X_{t_{i-1}}, X_{t_i} - S_{\nu(t_i - t_{i-1})} X_{t_{i-1}} \rangle_{\mathcal{L}_{\beta}^2}}{\frac{1}{N} \sum_{i=1}^{N} \|X_{t_{i-1}}\|_{\mathcal{L}_{\beta}^2}^2}.$$
(4.22)

Let  $Y_{t,s} = X_t - S_{\nu(t-s)}X_s$ ,  $s \leq t$  and  $Y_i = Y_{t_i,t_{i-1}}$ ,  $i = 1, \ldots, N$ . For  $\beta > -1$ , define

$$\widehat{J}_{\beta,\nu} = \widehat{J}_{\beta,\nu,N} = \frac{1}{N} \sum_{i=1}^{N} \|X_{t_{i-1}}\|_{\mathcal{L}^{2}_{\beta}}^{2}, \quad \widehat{K}_{\beta,\nu} = \widehat{K}_{\beta,\nu,N} = \sum_{i=1}^{N} \langle X_{t_{i-1}}, Y_{i} \rangle_{\mathcal{L}^{2}_{\beta}},$$
$$\widehat{M}_{\beta,\nu} = \widehat{M}_{\beta,\nu,N} = \frac{\theta_{0}^{*}}{\sigma^{*}} \sum_{i=1}^{N} \int_{t_{i-1}}^{t_{i}} \langle X_{t_{i-1}}, S_{\nu(t_{i}-s)} X_{s} - X_{t_{i-1}} \rangle_{\mathcal{L}^{2}_{\beta}} \mathrm{d}s + \sum_{i=1}^{N} \int_{t_{i-1}}^{t_{i}} \langle X_{t_{i-1}}, S_{\nu(t_{i}-s)} \mathrm{d}W_{s}^{Q} \rangle_{\mathcal{L}^{2}_{\beta}}.$$

The difference between the estimators  $\hat{\theta}_{0,\beta}^{(\text{cont})}$  and  $\hat{\theta}_{0,\beta}^{(\text{disc-cont})}$  is controlled as follows.

**Proposition 4.3.** Let  $\alpha > 0$  and  $-1 < \beta \leq \frac{1}{\alpha} - 1$ . Assume that  $[A1]_{-1,2}$  holds.

- (1) It holds that  $\hat{\theta}_{0,\beta}^{(\text{cont})} \hat{\theta}_{0,\beta}^{(\text{disc-cont})} = \mathcal{O}_{\mathbf{P}}(1) \text{ as } \nu \to 0 \text{ and } N \to \infty.$
- (2) Under  $[C1]_{\beta}$ , it holds that  $\mathcal{R}_{\beta,\nu}(\hat{\theta}_{0,\beta}^{(\text{cont})} \hat{\theta}_{0,\beta}^{(\text{disc-cont})}) = \mathcal{O}_{\mathbf{P}}(1)$  as  $\nu \to 0$  and  $N \to \infty$ .

*Proof.* Since it follows that under  $\mathbf{P}_{\theta_0^*,\sigma^*}$ ,

$$X_{t} = S_{\nu(t-t_{i-1})} X_{t_{i-1}} + \theta_{0}^{*} \int_{t_{i-1}}^{t} S_{\nu(t-s)} X_{s} \mathrm{d}s + \sigma^{*} \int_{t_{i-1}}^{t} S_{\nu(t-s)} \mathrm{d}W_{s}^{Q}, \qquad (4.23)$$

we have

$$Y_{i} = \theta_{0}^{*} X_{t_{i-1}}(t_{i} - t_{i-1}) + \theta_{0}^{*} \int_{t_{i-1}}^{t_{i}} (S_{\nu(t_{i}-s)} X_{s} - X_{t_{i-1}}) \mathrm{d}s + \sigma^{*} \int_{t_{i-1}}^{t_{i}} S_{\nu(t_{i}-s)} \mathrm{d}W_{s}^{Q}.$$

Therefore, we find that  $\widehat{K}_{\beta,\nu} = \theta_0^* \widehat{J}_{\beta,\nu} + \sigma^* \widehat{M}_{\beta,\nu}$ , which together with  $\widehat{\theta}_{0,\beta}^{(\text{disc-cont})} = \widehat{K}_{\beta,\nu} / \widehat{J}_{\beta,\nu}$  yields

$$\hat{\theta}_{0,\beta}^{(\text{disc-cont})} - \theta_0^* = \sigma^* \frac{M_{\beta,\nu}}{\hat{J}_{\beta,\nu}}.$$

Let  $\widehat{\Delta}_{\beta,\nu}^{(1)} = |\widehat{J}_{\beta,\nu} - J_{\beta,\nu}|$  and  $\widehat{\Delta}_{\beta,\nu}^{(2)} = |\widehat{M}_{\beta,\nu} - M_{\beta,\nu}|$ . Notice that  $|\widehat{\theta}_{0,\beta}^{(\text{cont})} - \widehat{\theta}_{0,\beta}^{(\text{disc-cont})}| = \sigma^* \left| \frac{M_{\beta,\nu}}{J_{\beta,\nu}} - \frac{\widehat{M}_{\beta,\nu}}{\widehat{J}_{\beta,\nu}} \right| \le \sigma^* \left( \left| \frac{M_{\beta,\nu}}{J_{\beta,\nu}} \right| \left| 1 - \frac{J_{\beta,\nu}}{\widehat{J}_{\beta,\nu}} \right| + \frac{\widehat{\Delta}_{\beta,\nu}^{(2)}}{\widehat{J}_{\beta,\nu}} \right)$   $\le \frac{|\widehat{\theta}_{0,\beta}^{(\text{cont})} - \theta_0^*|\widehat{\Delta}_{\beta,\nu}^{(1)} + \sigma^*\widehat{\Delta}_{\beta,\nu}^{(2)}}{J_{\beta,\nu} - \widehat{\Delta}_{\beta,\nu}^{(1)}}.$ 

Since Theorem 2.1, Propositions 4.1 and 4.2 show

$$\hat{\theta}_{0,\beta}^{(\text{cont})} - \theta_0^* = \mathcal{O}_{\mathbf{P}}(\mathcal{R}_{\beta,\nu}^{-1}), \quad \frac{\varphi_{\beta}(\nu)}{J_{\beta,\nu}} = \mathcal{O}_{\mathbf{P}}(1),$$

it is sufficient to verify

$$\widehat{\Delta}_{\beta,\nu}^{(1)} = \mathcal{O}_{\mathbf{P}}(\varphi_{\beta}(\nu)), \qquad (4.24)$$

and show

$$\widehat{\Delta}_{\beta,\nu}^{(2)} = \mathcal{O}_{\mathbf{P}}(\varphi_{\beta}(\nu)) \tag{4.25}$$

for the proof of (1) and

$$\widehat{\Delta}_{\beta,\nu}^{(2)} = \mathcal{O}_{\mathbf{P}}(\varphi_{2\beta+1}(\nu)^{1/2}) \tag{4.26}$$

for (2), respectively. Indeed, we can deduce from (4.24) that

$$\frac{1}{J_{\beta,\nu} - \widehat{\Delta}_{\beta,\nu}^{(1)}} = \frac{\varphi_{\beta}(\nu)^{-1}}{J_{\beta,\nu}/\varphi_{\beta}(\nu) - \widehat{\Delta}_{\beta,\nu}^{(1)}/\varphi_{\beta}(\nu)} = \mathcal{O}_{\mathbf{P}}(\varphi_{\beta}(\nu)^{-1}),$$

from (4.24) and (4.25) that

$$|\hat{\theta}_{0,\beta}^{(\text{cont})} - \hat{\theta}_{0,\beta}^{(\text{disc-cont})}| = \mathcal{O}_{\mathbf{P}}(\varphi_{\beta}(\nu)^{-1}) \Big( \mathcal{O}_{\mathbf{P}}(\mathcal{R}_{\beta,\nu}^{-1}) \mathcal{O}_{\mathbf{P}}(\varphi_{\beta}(\nu)) + \mathcal{O}_{\mathbf{P}}(\varphi_{\beta}(\nu)) \Big) = \mathcal{O}_{\mathbf{P}}(1),$$

and from (4.24), (4.26) and  $\mathcal{R}_{\beta,\nu}\varphi_{\beta}(\nu)^{-1} \sim \varphi_{2\beta+1}(\nu)^{-1/2}$  that

$$\mathcal{R}_{\beta,\nu}|\hat{\theta}_{0,\beta}^{(\text{cont})} - \hat{\theta}_{0,\beta}^{(\text{disc-cont})}| = \mathcal{R}_{\beta,\nu}\mathcal{O}_{\mathbf{P}}(\varphi_{\beta}(\nu)^{-1}) \Big( \mathcal{O}_{\mathbf{P}}(\mathcal{R}_{\beta,\nu}^{-1})\mathcal{O}_{\mathbf{P}}(\varphi_{\beta}(\nu)) + \mathcal{O}_{\mathbf{P}}(\varphi_{2\beta+1}(\nu)^{1/2}) \Big) \\ = \mathcal{O}_{\mathbf{P}}(1).$$

**Step 1:** We first show (4.24). For  $0 < \gamma_{\beta} < \frac{\alpha(\beta+1)}{2}$ , it follows from Lemmas 8 and 9 that

$$\begin{split} \mathbb{E}\Big[\widehat{\Delta}_{\beta,\nu}^{(1)}\Big] &\leq \sum_{i=1}^{N} \int_{t_{i-1}}^{t_{i}} \mathbb{E}\Big[\big| \|X_{t_{i-1}}\|_{\mathcal{L}_{\beta}^{2}}^{2} - \|X_{s}\|_{\mathcal{L}_{\beta}^{2}}^{2}\big|\Big] \mathrm{d}s \\ &\leq \sum_{i=1}^{N} \int_{t_{i-1}}^{t_{i}} \mathbb{E}\Big[\big(\|X_{t_{i-1}}\|_{\mathcal{L}_{\beta}^{2}}^{2} + \|X_{s}\|_{\mathcal{L}_{\beta}^{2}}^{2}\big)\|X_{t_{i-1}} - X_{s}\|_{\mathcal{L}_{\beta}^{2}}^{2}\Big] \mathrm{d}s \\ &\lesssim \sum_{i=1}^{N} \int_{t_{i-1}}^{t_{i}} \mathbb{E}\Big[\|X_{t_{i-1}}\|_{\mathcal{L}_{\beta}^{2}}^{2} + \|X_{s}\|_{\mathcal{L}_{\beta}^{2}}^{2}\Big]^{1/2} \mathbb{E}\Big[\|X_{t_{i-1}} - X_{s}\|_{\mathcal{L}_{\beta}^{2}}^{2}\Big]^{1/2} \mathrm{d}s \\ &\lesssim \nu^{(\alpha(\beta+1)-1)/2} \left(\frac{\nu^{\alpha(\beta+1)-1}}{N^{2\gamma_{\beta}\wedge1}} + \left(\frac{\nu}{N}\right)^{2\gamma_{\beta}} + \frac{1}{N}\right)^{1/2} \\ &\lesssim \nu^{(\alpha(\beta+1)-1)/2} \left(\frac{\nu^{(\alpha(\beta+1)-1)/2}}{N^{\gamma_{\beta}\wedge1/2}} + \frac{1}{\sqrt{N}}\right). \end{split}$$

Since Proposition 4.1 implies that for  $0 < \alpha(\beta + 1) \le 1$ ,

$$\varphi_{\beta}(\nu)^{-1}\nu^{(\alpha(\beta+1)-1)/2} \left(\frac{\nu^{(\alpha(\beta+1)-1)/2}}{N^{\gamma_{\beta}\wedge 1/2}} + \frac{1}{\sqrt{N}}\right) \lesssim \frac{1}{N^{\gamma_{\beta}\wedge 1/2}} + \frac{\nu^{(1-\alpha(\beta+1))/2}}{\sqrt{N}} \to 0,$$

we get  $\mathbb{E}[\widehat{\Delta}_{\beta,\nu}^{(1)}] = \mathcal{O}(\varphi_{\beta}(\nu)).$ Step 2: We next estimate  $\widehat{\Delta}_{\beta,\nu}^{(2)}$ . We have

$$\begin{split} \widehat{\Delta}^{(2)}_{\beta,\nu} \lesssim \left| \sum_{i=1}^{N} \int_{t_{i-1}}^{t_i} \langle X_{t_{i-1}}, S_{\nu(t_i-s)} X_s - X_{t_{i-1}} \rangle_{\mathcal{L}^2_\beta} \mathrm{d}s \right| \\ + \left| \sum_{i=1}^{N} \int_{t_{i-1}}^{t_i} \langle S_{\nu(t_i-s)} X_{t_{i-1}} - X_s, \mathrm{d}W^Q_s \rangle_{\mathcal{L}^2_\beta} \right| \\ =: \widehat{\Delta}^{(2,1)}_{\beta,\nu} + \widehat{\Delta}^{(2,2)}_{\beta,\nu}. \end{split}$$

Note that the weighted Schwarz inequality:

$$|\langle u,v\rangle_{\mathcal{L}^2_{\beta}}| \leq \|u\|_{\mathcal{L}^2_{\beta+\delta'}} \|v\|_{\mathcal{L}^2_{\beta-\delta'}}, \quad u \in \mathcal{L}^2_{\beta+\delta'} \cap \mathcal{L}^2_{\beta}, \ v \in \mathcal{L}^2_{\beta-\delta'} \cap \mathcal{L}^2_{\beta} \ (\delta' \in \mathbb{R}).$$

For any  $0 \le \delta < \beta + 1$  and  $0 < \gamma_{\delta} < \frac{\alpha(\beta+\delta+1)}{2}$ , it holds from (4.36), Lemmas 8 and 9 that

$$\begin{split} \mathbb{E}\Big[\big(\widehat{\Delta}_{\beta,\nu}^{(2,1)}\big)^{2}\Big] &\leq \left(\sum_{i=1}^{N} \int_{t_{i-1}}^{t_{i}} \mathbb{E}\Big[\|X_{t_{i-1}}\|_{\mathcal{L}^{2}_{\beta-\delta}}\|S_{\nu(t_{i}-s)}X_{s} - X_{t_{i-1}}\|_{\mathcal{L}^{2}_{\beta+\delta}}\Big]\mathrm{d}s\right)^{2} \\ &\lesssim \sum_{i=1}^{N} \int_{t_{i-1}}^{t_{i}} \mathbb{E}\Big[\|X_{t_{i-1}}\|_{\mathcal{L}^{2}_{\beta-\delta}}^{2}\Big]\mathrm{d}s \\ &\qquad \times \sum_{i=1}^{N} \int_{t_{i-1}}^{t_{i}} \Big(\mathbb{E}\Big[\|S_{\nu(t_{i}-s)}X_{s} - S_{\nu(t_{i}-s)}X_{t_{i-1}}\|_{\mathcal{L}^{2}_{\beta+\delta}}^{2}\Big] \\ &\qquad + \mathbb{E}\Big[\|S_{\nu(t_{i}-s)}X_{t_{i-1}} - X_{t_{i-1}}\|_{\mathcal{L}^{2}_{\beta+\delta}}^{2}\Big]\mathrm{d}s \\ &\lesssim \sum_{i=1}^{N} \int_{t_{i-1}}^{t_{i}} \mathbb{E}\Big[\|X_{t_{i-1}}\|_{\mathcal{L}^{2}_{\beta-\delta}}^{2}\Big]\mathrm{d}s \\ &\qquad \times \sum_{i=1}^{N} \int_{t_{i-1}}^{t_{i}} \Big[\mathbb{E}\Big[\|X_{s} - X_{t_{i-1}}\|_{\mathcal{L}^{2}_{\beta+\delta}}^{2}\Big] + \Big(\frac{\nu}{N}\Big)^{2\gamma_{\delta}} \mathbb{E}\Big[\|X_{t_{i-1}}\|_{\mathcal{L}^{2}_{\beta+\delta-2\gamma_{\delta}/a}}^{2}\Big]\Big]\mathrm{d}s \\ &\lesssim \nu^{\alpha(\beta-\delta+1)-1}\Big(\frac{\nu^{\alpha(\beta+\delta+1)-1}}{N^{2\gamma_{\delta}\wedge1}} + \Big(\frac{\nu}{N}\Big)^{2\gamma_{\delta}} + \frac{1}{N} + \Big(\frac{\nu}{N}\Big)^{2\gamma_{\delta}}(1+\nu^{\alpha(\beta+\delta+1)-2\gamma_{\delta}-1})\Big) \\ &\lesssim \nu^{\alpha(\beta-\delta+1)-1}\Big(\frac{\nu^{\alpha(\beta+\delta+1)-1}}{N^{2\gamma_{\delta}\wedge1}} + \Big(\frac{\nu}{N}\Big)^{2\gamma_{\delta}} + \frac{1}{N}\Big) =: \mathcal{S}_{\beta,\delta,\gamma_{\delta}}(\nu, N). \end{split}$$

Similarly, it follows that for any  $0 < \gamma_1 < \frac{\alpha(\beta+2)}{2}$ ,

$$\mathbb{E}\Big[\big(\widehat{\Delta}_{\beta,\nu}^{(2,2)}\big)^2\Big] = \sum_{i=1}^N \int_{t_{i-1}}^{t_i} \mathbb{E}\Big[\|S_{\nu(t_i-s)}X_{t_{i-1}} - X_s\|_{\mathcal{L}^2_{\beta+1}}^2\Big] \mathrm{d}s$$
  
$$\lesssim \sum_{i=1}^N \int_{t_{i-1}}^{t_i} \Big(\mathbb{E}\Big[\|S_{\nu(t_i-s)}X_{t_{i-1}} - S_{\nu(t_i-s)}X_s\|_{\mathcal{L}^2_{\beta+1}}^2\Big] + \mathbb{E}\Big[\|S_{\nu(t_i-s)}X_s - X_s\|_{\mathcal{L}^2_{\beta+1}}^2\Big]\Big) \mathrm{d}s$$

$$\begin{split} &\lesssim \sum_{i=1}^{N} \int_{t_{i-1}}^{t_{i}} \left( \mathbb{E}\Big[ \|X_{t_{i-1}} - X_{s}\|_{\mathcal{L}_{\beta+1}}^{2} \Big] + \Big(\frac{\nu}{N}\Big)^{2\gamma_{1}} \mathbb{E}\Big[ \|X_{s}\|_{\mathcal{L}_{\beta+1-2\gamma_{1}/\alpha}}^{2} \Big] \Big) \mathrm{d}s \\ &\lesssim \Big(\frac{\nu^{\alpha(\beta+2)-1}}{N^{2\gamma_{1}\wedge 1}} + \Big(\frac{\nu}{N}\Big)^{2\gamma_{1}} + \frac{1}{N}\Big) + \Big(\frac{\nu}{N}\Big)^{2\gamma_{1}} (1 + \nu^{\alpha(\beta+2)-2\gamma_{1}-1}) \\ &\lesssim \frac{\nu^{\alpha(\beta+2)-1}}{N^{2\gamma_{1}\wedge 1}} + \Big(\frac{\nu}{N}\Big)^{2\gamma_{1}} + \frac{1}{N} =: \mathcal{T}_{\beta,\gamma_{1}}(\nu, N). \end{split}$$

Therefore, we see that for any  $0 \le \delta < \beta + 1$ ,  $0 < \gamma_{\delta} < \frac{\alpha(\beta+\delta+1)}{2}$  and  $0 < \gamma_1 < \frac{\alpha(\beta+2)}{2}$ ,

$$\mathbb{E}\Big[\big(\widehat{\Delta}_{\beta,\nu}^{(2)}\big)^2\Big] \lesssim \mathcal{S}_{\beta,\delta,\gamma_{\delta}}(\nu,N) + \mathcal{T}_{\beta,\gamma_1}(\nu,N).$$

**Step 3:** In order to show (4.25), we prove that there exist  $(\delta, \gamma_{\delta}, \gamma_1)$  such that

$$\varphi_{\beta}(\nu)^{-2}(\mathcal{S}_{\beta,\delta,\gamma_{\delta}}(\nu,N) + \mathcal{T}_{\beta,\gamma_{1}}(\nu,N)) \to 0.$$

By choosing  $\delta = 0$ ,  $\gamma_{\delta} > 0$  and  $\gamma_1 > 0$ , it follows from Proposition 4.1,  $1 - \alpha(\beta + 1) \ge 0$  and  $1 - \alpha\beta > 0$  that

$$\varphi_{\beta}(\nu)^{-2} \mathcal{S}_{\beta,\delta,\gamma_{\delta}}(\nu,N) \lesssim \frac{1}{N^{2\gamma_{\delta}\wedge 1}} + \frac{\nu^{1-\alpha(\beta+1)+2\gamma_{\delta}}}{N^{2\gamma_{\delta}}} + \frac{\nu^{1-\alpha(\beta+1)}}{N} \to 0,$$
  
$$\varphi_{\beta}(\nu)^{-2} \mathcal{T}_{\beta,\gamma_{1}}(\nu,N) \lesssim \frac{\nu^{1-\alpha\beta}}{N^{2\gamma_{1}\wedge 1}} + \frac{\nu^{2(1-\alpha(\beta+1)+\gamma_{1})}}{N^{2\gamma_{1}}} + \frac{\nu^{2(1-\alpha(\beta+1))}}{N} \to 0.$$

**Step 4:** To complete the proof of (4.26), we verify that under  $[C1]_{\beta}$ ,

$$\varphi_{2\beta+1}(\nu)^{-1}(\mathcal{S}_{\beta,\delta,\gamma_{\delta}}(\nu,N) + \mathcal{T}_{\beta,\gamma_{1}}(\nu,N)) \to 0$$

for some  $(\delta, \gamma_{\delta}, \gamma_1)$ .

(i) Consider the case that  $\alpha(\beta + 1) \leq 1/2$ . By noting that  $2\gamma_{\delta} < \alpha(\beta + \delta + 1) < 1$  and  $\gamma_1 > 0 \geq \alpha(\beta + 1) - 1/2$ , it follows from Proposition 4.1 that

$$\begin{split} \varphi_{2\beta+1}(\nu)^{-1} \mathcal{S}_{\beta,\delta,\gamma_{\delta}}(\nu,N) &\lesssim \nu^{-\alpha(\beta+\delta+1)} \left( \frac{\nu^{\alpha(\beta+\delta+1)-1}}{N^{2\gamma_{\delta}\wedge 1}} + \left(\frac{\nu}{N}\right)^{2\gamma_{\delta}} + \frac{1}{N} \right) \\ &\lesssim \frac{1}{\nu N^{2\gamma_{\delta}}} + \frac{1}{\nu^{\alpha(\beta+\delta+1)}N}, \\ \varphi_{2\beta+1}(\nu)^{-1} \mathcal{T}_{\beta,\gamma_{1}}(\nu,N) &\lesssim \frac{\nu^{-\alpha\beta}}{N^{2\gamma_{1}\wedge 1}} + \frac{\nu^{1-2\alpha(\beta+1)+2\gamma_{1}}}{N^{2\gamma_{1}}} + \frac{\nu^{1-2\alpha(\beta+1)}}{N} \\ &= \begin{cases} \mathcal{O}(1), & \beta \leq 0, \\ \frac{1}{\nu^{\alpha\beta}N^{2\gamma_{1}\wedge 1}} + \mathcal{O}(1), & \beta > 0. \end{cases} \end{split}$$

Hence, we prove that under  $\nu N^q \to \infty$  for some  $q < \rho_{\alpha(\beta+1)}^{(\text{time})} = 2\alpha(\beta+1)$ , there exist  $0 \le \delta < \beta+1$ and  $0 < \gamma_{\delta} < \frac{\alpha(\beta+\delta+1)}{2}$  such that

$$\nu N^{2\gamma_{\delta} \wedge \frac{1}{\alpha(\beta+\delta+1)}} \to \infty \tag{4.27}$$

and additionally show that there exists  $0 < \gamma_1 < \frac{\alpha(\beta+2)}{2}$  such that

$$\nu N^{\frac{2\gamma_1 \wedge 1}{\alpha\beta}} \to \infty \tag{4.28}$$

if  $\beta > 0$ .

Note that for a function  $f : \mathbb{R}^d \to \mathbb{R}$  and a domain  $E \subset \mathbb{R}^d$ , if  $\sup_{x \in E} f(x)$  exists and  $\nu N^q \to \infty$  for some  $q < \sup_{x \in E} f(x)$ , then  $\nu N^{f(x)} \to \infty$  for some  $x \in E$ .

Because  $2\gamma_{\delta} < \alpha(\beta + \delta + 1) < 1$ ,  $x \wedge \frac{1}{x} = x$  (0 < x < 1) and  $\delta < \beta + 1$ , we find that

$$\sup_{\delta} \sup_{\gamma_{\delta}} \left( 2\gamma_{\delta} \wedge \frac{1}{\alpha(\beta+\delta+1)} \right) = \sup_{\delta} \left( \alpha(\beta+\delta+1) \wedge \frac{1}{\alpha(\beta+\delta+1)} \right)$$
$$= \sup_{\delta} \alpha(\beta+\delta+1) = 2\alpha(\beta+1),$$
$$\sup_{\gamma_{1}} \frac{2\gamma_{1} \wedge 1}{\alpha\beta} = \left(1 + \frac{2}{\beta}\right) \wedge \frac{1}{\alpha\beta} > 1 \ge 2\alpha(\beta+1),$$

which imply that there exist  $(\delta, \gamma_{\delta}, \gamma_1)$  such that (4.27) and (4.28) under  $\nu N^q \to \infty$  for some  $q < 2\alpha(\beta + 1)$ .

(ii) Consider the case that  $1/2 < \alpha(\beta+1) \leq 1$ . By choosing  $\delta > \frac{1}{\alpha} - (\beta+1)$  and  $\gamma_{\delta} < \frac{\alpha(\beta+\delta+1)}{2}$ , it follows from Proposition 4.1 and  $2\gamma_{\delta} > 1$  that

$$\begin{split} \varphi_{2\beta+1}(\nu)^{-1} \mathcal{S}_{\beta,\delta,\gamma_{\delta}}(\nu,N) &\lesssim \nu^{\alpha(\beta-\delta+1)-1} \left( \frac{\nu^{\alpha(\beta+\delta+1)-1}}{N^{2\gamma_{\delta}\wedge 1}} + \left(\frac{\nu}{N}\right)^{2\gamma_{\delta}} + \frac{1}{N} \right) \\ &\lesssim \frac{1}{\nu^{1-\alpha(\beta-\delta+1)}N}. \end{split}$$

Furthermore, it holds that for  $\gamma_1 > 1/2$ ,

$$\varphi_{2\beta+1}(\nu)^{-1}\mathcal{T}_{\beta,\gamma_1}(\nu,N) \lesssim \frac{\nu^{\alpha(\beta+2)-1}}{N^{2\gamma_1\wedge 1}} + \left(\frac{\nu}{N}\right)^{2\gamma_1} + \frac{1}{N} = \frac{1}{\nu^{1-\alpha(\beta+2)}N} + \mathcal{O}(1).$$

Therefore, we see from

$$\sup_{\delta} \frac{1}{1 - \alpha(\beta - \delta + 1)} = \frac{1}{2(1 - \alpha(\beta + 1))},$$
$$\frac{1}{1 - \alpha(\beta + 2)} - \frac{1}{2(1 - \alpha(\beta + 1))} = \frac{1 - \alpha\beta}{2(1 - \alpha(\beta + 1))(1 - \alpha(\beta + 2))} > 0$$

that there exists  $\delta > \frac{1}{\alpha} - (\beta + 1)$  such that

$$\nu N^{\frac{1}{1-\alpha(\beta-\delta+1)}\wedge\frac{1}{1-\alpha(\beta+2)}} \to \infty$$

under  $\nu N^q \to \infty$  for some  $q < \rho_{\alpha(\beta+1)}^{(\text{time})} = \frac{1}{2(1-\alpha(\beta+1))}$ .

# 4.2.2 Control of spatial discretization error

Consider a spatial discretization of the estimator  $\hat{\theta}_{0,\beta}^{(\text{disc-cont})}$  given in (4.22). We set the estimator of  $\theta_0$  based on the discrete spatio-temporal data  $\mathbb{X}_{N,M}$  as follows.

$$\hat{\theta}_{0,\beta}^{(\text{disc})} = \hat{\theta}_{0,\beta,\nu,N,M}^{(\text{disc})} = \frac{\sum_{i=1}^{N} \langle \Psi_M X_{t_{i-1}}, \Psi_M X_{t_i} - S_{\nu(t_i - t_{i-1})} \Psi_M X_{t_{i-1}} \rangle_{\mathcal{L}^2_{\beta}}}{\frac{1}{N} \sum_{i=1}^{N} \|\Psi_M X_{t_{i-1}}\|_{\mathcal{L}^2_{\beta}}^2}, \quad (4.29)$$

where  $\Psi_M$  is the operator such that

$$\Psi_M f(y, z) = f(y_{j-1}, z_{k-1}), \quad (y, z) \in [y_{j-1}, y_j) \times [z_{k-1}, z_k)$$

for  $j = 1, ..., M_1$ ,  $k = 1, ..., M_2$  and a continuous function f on D.

For  $\beta > -1$ , define  $Y_i^{\Psi_M} = \Psi_M X_{t_i} - S_{\nu(t_i - t_{i-1})} \Psi_M X_{t_{i-1}}$ ,

$$\widetilde{J}_{\beta,\nu} = \widetilde{J}_{\beta,\nu,N,M} = \frac{1}{N} \sum_{i=1}^{N} \|\Psi_M X_{t_{i-1}}\|_{\mathcal{L}^2_{\beta}}^2, \quad \widetilde{K}_{\beta,\nu} = \widetilde{K}_{\beta,\nu,N,M} = \sum_{i=1}^{N} \langle \Psi_M X_{t_{i-1}}, Y_i^{\Psi_M} \rangle_{\mathcal{L}^2_{\beta}}.$$

Let  $M = (M_1 \wedge M_2)^2$ . The following proposition provides the asymptotic equivalences of the estimators  $\hat{\theta}_{0,\beta}^{(\text{disc-cont})}$  and  $\hat{\theta}_{0,\beta}^{(\text{disc})}$ .

**Proposition 4.4.** Let  $\alpha > 0$ ,  $-1 < \beta \leq \frac{1}{\alpha} - 1$  and  $p > \frac{4}{\alpha(\beta+1)}$ . Assume that  $[A1]_{-1,p}$  holds.

- (1) Under  $[B1]_{\beta,p}$ , it holds that  $\hat{\theta}_{0,\beta}^{(\text{disc-cont})} \hat{\theta}_{0,\beta}^{(\text{disc})} = \mathcal{O}_{\mathbf{P}}(1)$  as  $\nu \to 0$ ,  $N \to \infty$  and  $M \to \infty$ .
- (2) Under  $[C2]_{\beta,p}$ , it holds that  $\mathcal{R}_{\beta,\nu}(\hat{\theta}_{0,\beta}^{(\text{disc-cont})} \hat{\theta}_{0,\beta}^{(\text{disc})}) = \mathcal{O}_{\mathbf{P}}(1)$  as  $\nu \to 0$ ,  $N \to \infty$  and  $M \to \infty$ .

*Proof.* Recall  $\hat{\theta}_{0,\beta}^{(\text{disc-cont})} = \widehat{K}_{\beta,\nu}/\widehat{J}_{\beta,\nu}$ . Setting  $\widetilde{\Delta}_{\beta,\nu}^{(1)} = |\widetilde{J}_{\beta,\nu} - \widehat{J}_{\beta,\nu}|$  and  $\widetilde{\Delta}_{\beta,\nu}^{(2)} = |\widetilde{K}_{\beta,\nu} - \widehat{K}_{\beta,\nu}|$ , we have

$$\begin{split} |\hat{\theta}_{0,\beta}^{(\text{disc-cont})} - \hat{\theta}_{0,\beta}^{(\text{disc})}| &= \left|\hat{\theta}_{0,\beta}^{(\text{disc-cont})} - \frac{\widehat{K}_{\beta,\nu}}{\widetilde{J}_{\beta,\nu}} - \frac{\widetilde{K}_{\beta,\nu} - \widehat{K}_{\beta,\nu}}{\widetilde{J}_{\beta,\nu}}\right| \leq |\hat{\theta}_{0,\beta}^{(\text{disc-cont})}| \left|1 - \frac{\widehat{J}_{\beta,\nu}}{\widetilde{J}_{\beta,\nu}}\right| + \frac{\widetilde{\Delta}_{\beta,\nu}^{(2)}}{\widetilde{J}_{\beta,\nu}} \\ &\leq \frac{|\hat{\theta}_{0,\beta}^{(\text{disc-cont})}|\widetilde{\Delta}_{\beta,\nu}^{(1)} + \widetilde{\Delta}_{\beta,\nu}^{(2)}}{\widehat{J}_{\beta,\nu} - \widetilde{\Delta}_{\beta,\nu}^{(1)}}. \end{split}$$

Because it follows from Proposition 4.3 and (4.24) that

$$\hat{\theta}_{0,\beta}^{(\text{disc-cont})} = \mathcal{O}_{\mathbf{P}}(1), \quad \frac{\varphi_{\beta}(\nu)}{\widehat{J}_{\beta,\nu}} = \mathcal{O}_{\mathbf{P}}(1), \tag{4.30}$$

we show that

$$\widetilde{\Delta}^{(1)}_{\beta,\nu}, \widetilde{\Delta}^{(2)}_{\beta,\nu} = \mathcal{O}_{\mathbf{P}}(\varphi_{\beta}(\nu))$$
(4.31)

under  $[B1]_{\beta,p}$  for the proof of (1), and

$$\widetilde{\Delta}^{(1)}_{\beta,\nu}, \widetilde{\Delta}^{(2)}_{\beta,\nu} = \mathcal{O}_{\mathbf{P}}(\varphi_{2\beta+1}(\nu)^{1/2})$$

under  $[C2]_{\beta,p}$  for the proof of (2).

**Step 1:** We estimate  $\widetilde{\Delta}_{\beta,\nu}^{(1)}$  and  $\widetilde{\Delta}_{\beta,\nu}^{(2)}$ . Note that for any  $\delta_1, \delta_2, \delta_3 \in \mathbb{R}$ ,

$$\begin{aligned} |\langle \Psi_M f, \Psi_M g \rangle_{\mathcal{L}^2_{\beta}} - \langle f, g \rangle_{\mathcal{L}^2_{\beta}}| &\leq \|\Psi_M f - f\|_{\mathcal{L}^2_{\beta-\delta_1}} \|\Psi_M g - g\|_{\mathcal{L}^2_{\beta+\delta_1}} + \|f\|_{\mathcal{L}^2_{\beta-\delta_2}} \|\Psi_M g - g\|_{\mathcal{L}^2_{\beta+\delta_2}} \\ &+ \|\Psi_M f - f\|_{\mathcal{L}^2_{\beta-\delta_3}} \|g\|_{\mathcal{L}^2_{\beta+\delta_3}}. \end{aligned}$$

It holds from Lemmas 8, 10 and 11 that for any  $0 \leq \delta_1 < \beta + 1 - \frac{4}{\alpha p}, 0 \leq \delta_2, \delta_3 < \beta + 1,$  $\frac{1}{p} < \gamma_{\delta_j} < \left(\frac{\alpha(\beta+\delta_j+1)}{2} - \frac{1}{p}\right) \wedge \frac{1}{2} \text{ and } \frac{1}{p} < \gamma_{-\delta_1} < \frac{\alpha(\beta-\delta_1+1)}{2} - \frac{1}{p},$ 

$$\mathbb{E}\Big[\big|\|X_{t_{i-1}}\|^{2}_{\mathcal{L}^{2}_{\beta}} - \|\Psi_{M}X_{t_{i-1}}\|^{2}_{\mathcal{L}^{2}_{\beta}}\big|\Big] \\
\leq \mathbb{E}\Big[\|\Psi_{M}X_{t_{i-1}} - X_{t_{i-1}}\|^{2}_{\mathcal{L}^{2}_{\beta-\delta_{1}}}\Big]^{1/2} \mathbb{E}\Big[\|\Psi_{M}X_{t_{i-1}} - X_{t_{i-1}}\|^{2}_{\mathcal{L}^{2}_{\beta+\delta_{1}}}\Big]^{1/2} \\
+ 2\mathbb{E}\Big[\|X_{t_{i-1}}\|^{2}_{\mathcal{L}^{2}_{\beta-\delta_{2}}}\Big]^{1/2} \mathbb{E}\Big[\|\Psi_{M}X_{t_{i-1}} - X_{t_{i-1}}\|^{2}_{\mathcal{L}^{2}_{\beta+\delta_{2}}}\Big]^{1/2} \\
\lesssim \frac{\varphi_{\beta-\delta_{1}-2\gamma-\delta_{1}}/\alpha(\nu)^{1/2}\varphi_{\beta+\delta_{1}-2\gamma\delta_{1}}/\alpha(\nu)^{1/2}}{M^{\gamma\delta_{1}+\gamma-\delta_{1}-2/p}}$$

$$\begin{split} &+ \frac{\nu^{(\alpha(\beta-\delta_{2}+1)-1)/2}\varphi_{\beta+\delta_{2}-2\gamma_{\delta_{2}}/\alpha}(\nu)^{1/2}}{M^{\gamma_{\delta_{2}}-1/p}} \\ &= \frac{\nu^{(\alpha(\beta-\delta_{1}+1)-2\gamma-\delta_{1}-1)/2}\varphi_{\beta+\delta_{1}-2\gamma_{\delta_{1}}/\alpha}(\nu)^{1/2}}{M^{\gamma_{\delta_{1}}+\gamma-\delta_{1}-2/p}} \\ &+ \frac{\nu^{(\alpha(\beta-\delta_{2}+1)-1)/2}\varphi_{\beta+\delta_{2}-2\gamma_{\delta_{2}}/\alpha}(\nu)^{1/2}}{M^{\gamma_{\delta_{2}}-1/p}}, \\ &\mathbb{E}\Big[|\langle \Psi_{M}X_{t_{i-1}}, Y_{i}^{\Psi_{M}}\rangle_{\mathcal{L}_{\beta}^{2}} - \langle X_{t_{i-1}}, Y_{i}\rangle_{\mathcal{L}_{\beta}^{2}}|\Big] \\ &\leq \mathbb{E}\Big[||\Psi_{M}X_{t_{i-1}} - X_{t_{i-1}}||_{\mathcal{L}_{\beta-\delta_{1}}^{2}}\Big]^{1/2}\mathbb{E}\Big[||Y_{i}^{\Psi_{M}} - Y_{i}||_{\mathcal{L}_{\beta+\delta_{1}}^{2}}\Big]^{1/2} \\ &+ \mathbb{E}\Big[||\Psi_{M}X_{t_{i-1}} - X_{t_{i-1}}||_{\mathcal{L}_{\beta+\delta_{3}}}^{2}\Big]^{1/2}\mathbb{E}\Big[||Y_{i}||_{\mathcal{L}_{\beta+\delta_{2}}}^{2}\Big]^{1/2} \\ &\leq \frac{\nu^{(\alpha(\beta-\delta_{1}+1)-2\gamma-\delta_{1}-1)/2}\varphi_{\beta+\delta_{1}-2\gamma_{\delta_{1}}/\alpha}(\nu)^{1/2}}{M^{\gamma_{\delta_{1}}+\gamma-\delta_{1}-2/p}N^{\alpha(\beta+\delta_{1}+1)/2-\gamma_{\delta_{1}}}} \\ &\leq \frac{\nu^{(\alpha(\beta-\delta_{2}+1)-1)/2}\varphi_{\beta+\delta_{2}-2\gamma_{\delta_{2}}/\alpha}(\nu)^{1/2}}{M^{\gamma_{\delta_{2}}-1/p}N^{\alpha(\beta+\delta_{2}+1)/2-\gamma_{\delta_{2}}}} \\ &+ \frac{\varphi_{\beta+\delta_{3}-2\gamma_{\delta_{3}}/\alpha}(\nu)^{1/2}\nu^{(\alpha(\beta-\delta_{3}+1)-1)/2}}{M^{\gamma_{\delta_{3}}-1/p}N^{\alpha(\beta-\delta_{3}+1)/2}}. \end{split}$$

Therefore, for i = 1, 2, we have

$$\begin{split} \mathbb{E}\Big[\widetilde{\Delta}_{\beta,\nu}^{(i)}\Big] &\lesssim \frac{\nu^{(\alpha(\beta-\delta_{1}+1)-2\gamma_{-\delta_{1}}-1)/2}\varphi_{\beta+\delta_{1}-2\gamma_{\delta_{1}}/\alpha}(\nu)^{1/2}}{M^{\gamma_{\delta_{1}}+\gamma_{-\delta_{1}}-2/p}N^{\alpha(\beta+\delta_{1}+1)/2-\gamma_{\delta_{1}}-1}} \\ &+ \frac{\nu^{(\alpha(\beta-\delta_{2}+1)-1)/2}\varphi_{\beta+\delta_{2}-2\gamma_{\delta_{2}}/\alpha}(\nu)^{1/2}}{M^{\gamma_{\delta_{2}}-1/p}N^{\alpha(\beta+\delta_{2}+1)/2-\gamma_{\delta_{2}}-1}} \\ &+ \frac{\varphi_{\beta+\delta_{3}-2\gamma_{\delta_{3}}/\alpha}(\nu)^{1/2}\nu^{(\alpha(\beta-\delta_{3}+1)-1)/2}}{M^{\gamma_{\delta_{3}}-1/p}N^{\alpha(\beta-\delta_{3}+1)/2-1}} \\ &=: \mathcal{U}_{\beta,\delta_{1},\delta_{2},\delta_{3},\gamma_{\delta_{1}},\gamma_{-\delta_{1}},\gamma_{\delta_{2}},\gamma_{\delta_{3}}}(\nu,N,M). \end{split}$$

**Step 2:** In order to conclude this proof, we show that for some  $\delta_j$ ,  $\gamma_{\delta_j}$  (j = 1, 2, 3) and  $\gamma_{-\delta_1}$ ,

$$\varphi_{\beta}(\nu)^{-1}\mathcal{U}_{\beta,\delta_{1},\delta_{2},\delta_{3},\gamma_{\delta_{1}},\gamma_{-\delta_{1}},\gamma_{\delta_{2}},\gamma_{\delta_{3}}}(\nu,N,M)\to 0$$

under  $[\mathrm{B1}]_{\beta,p}$  and

$$\varphi_{2\beta+1}(\nu)^{-1/2}\mathcal{U}_{\beta,\delta_1,\delta_2,\delta_3,\gamma_{\delta_1},\gamma_{-\delta_1},\gamma_{\delta_2},\gamma_{\delta_3}}(\nu,N,M) \to 0$$

under [C2]<sub> $\beta,p$ </sub>, respectively. Recall

$$0 \le \delta_1 < \beta + 1 - \frac{4}{\alpha p}, \quad 0 \le \delta_2, \delta_3 < \beta + 1,$$
$$\frac{1}{p} < \gamma_{-\delta_1} < \frac{\alpha(\beta - \delta_1 + 1)}{2} - \frac{1}{p}, \quad \frac{1}{p} < \gamma_{\delta_j} < \left(\frac{\alpha(\beta + \delta_j + 1)}{2} - \frac{1}{p}\right) \land \frac{1}{2} \quad (j = 1, 2, 3).$$

(i) Consider the case that  $\alpha(\beta + 1) \leq 1/2$ . Since  $\varphi_{\beta+\delta_j-2\gamma_{\delta_j}/\alpha}(\nu) \sim \nu^{\alpha(\beta+\delta_j+1)-2\gamma_{\delta_j}-1}$  (j = 1, 2) and (1, 2, 3) and

$$\mathcal{U}_{\beta,\delta_1,\delta_2,\delta_3,\gamma_{\delta_1},\gamma_{-\delta_1},\gamma_{\delta_2},\gamma_{\delta_3}}(\nu,N,M)$$

$$\sim \nu^{\alpha(\beta+1)-1} \bigg( \frac{1}{\nu^{\gamma_{\delta_1}+\gamma_{-\delta_1}} M^{\gamma_{\delta_1}+\gamma_{-\delta_1}-2/p} N^{\alpha(\beta+\delta_1+1)/2-\gamma_{\delta_1}-1}} \\ + \frac{1}{\nu^{\gamma_{\delta_2}} M^{\gamma_{\delta_2}-1/p} N^{\alpha(\beta+\delta_2+1)/2-\gamma_{\delta_2}-1}} + \frac{1}{\nu^{\gamma_{\delta_3}} M^{\gamma_{\delta_3}-1/p} N^{\alpha(\beta-\delta_3+1)/2-1}} \bigg),$$

it suffices to verify that for c = 0, 1/2, there exist  $\delta_j$ ,  $\gamma_{\delta_j}$  (j = 1, 2, 3) and  $\gamma_{-\delta_1}$  such that

$$\begin{cases} \nu^{\gamma_{\delta_1}+\gamma_{-\delta_1}+c} M^{\gamma_{\delta_1}+\gamma_{-\delta_1}-2/p} N^{\alpha(\beta+\delta_1+1)/2-\gamma_{\delta_1}-1} \to \infty, \\ \nu^{\gamma_{\delta_2}+c} M^{\gamma_{\delta_2}-1/p} N^{\alpha(\beta+\delta_2+1)/2-\gamma_{\delta_2}-1} \to \infty, \\ \nu^{\gamma_{\delta_3}+c} M^{\gamma_{\delta_3}-1/p} N^{\alpha(\beta-\delta_3+1)/2-1} \to \infty \end{cases}$$

$$(4.32)$$

under  $[B1]_{\beta,p}$  and  $[C2]_{\beta,p}$ , respectively.

Let  $M = N^b$ ,  $b \ge 1$ . Notice that  $\frac{1}{p} < \gamma_{\delta_j} < \frac{\alpha(\beta+\delta_j+1)}{2} - \frac{1}{p}$  (j = 1, 2, 3). Since

$$\sup_{\delta_{1}} \sup_{\gamma_{\delta_{1}}} \sup_{\gamma_{-\delta_{1}}} \frac{b(\gamma_{\delta_{1}} + \gamma_{-\delta_{1}} - 2/p) + \alpha(\beta + \delta_{1} + 1)/2 - \gamma_{\delta_{1}} - 1}{\gamma_{\delta_{1}} + \gamma_{-\delta_{1}} + c}$$

$$= \sup_{\delta_{1}} \sup_{\gamma_{\delta_{1}}} \left( b + \frac{-b(c + 2/p) + \alpha(\beta + \delta_{1} + 1)/2 - \gamma_{\delta_{1}} - 1}{\gamma_{\delta_{1}} + \alpha(\beta - \delta_{1} + 1)/2 - 1/p + c} \right)$$

$$= b \left( 1 - \frac{c + 2/p}{\alpha(\beta + 1) - 2/p + c} \right) - \frac{1 - 1/p}{\alpha(\beta + 1) - 2/p + c} =: \Gamma_{\alpha(\beta + 1), p, c}^{(1)}$$

$$\sup_{\delta_{2}} \sup_{\gamma_{\delta_{2}}} \frac{b(\gamma_{\delta_{2}} - 1/p) + \alpha(\beta + \delta_{2} + 1)/2 - \gamma_{\delta_{2}} - 1}{\gamma_{\delta_{2}} + c}$$

$$= \sup_{\delta_{2}} \left( b - 1 + \frac{-b(c + 1/p) + c + \alpha(\beta + \delta_{2} + 1)/2 - 1}{\alpha(\beta + \delta_{2} + 1)/2 - 1/p + c} \right)$$

$$= b \left( 1 - \frac{c + 1/p}{\alpha(\beta + 1) - 1/p + c} \right) - \frac{1 - 1/p}{\alpha(\beta + 1) - 1/p + c} =: \widetilde{\Gamma}^{(2)}_{\alpha(\beta + 1), p, c},$$

and

$$\begin{split} \sup_{\delta_{3}} \sup_{\gamma_{\delta_{3}}} \frac{b(\gamma_{\delta_{3}} - 1/p) + \alpha(\beta - \delta_{3} + 1)/2 - 1}{\gamma_{\delta_{3}} + c} \\ &= \sup_{\delta_{3}} \left( b + \frac{-b(c + 1/p) + \alpha(\beta - \delta_{3} + 1)/2 - 1}{\alpha(\beta + \delta_{3} + 1)/2 - 1/p + c} \right) \\ &= b \left( 1 - \frac{c + 1/p}{\alpha(\beta + 1) - 1/p + c} \right) - \frac{1}{\alpha(\beta + 1) - 1/p + c} =: \Gamma^{(2)}_{\alpha(\beta + 1), p, c} \leq \widetilde{\Gamma}^{(2)}_{\alpha(\beta + 1), p, c}, \end{split}$$

we have (4.32) under  $\nu N^u \to \infty$  for some

$$u < \Gamma_{x,p,c}^{(1)} \wedge \Gamma_{x,p,c}^{(2)} = \begin{cases} (b\tau_{1,x,p}^{(\text{space})} - \phi_{1,x,p}^{(\text{space})}) \wedge (b\tau_{2,x,p}^{(\text{space})} - \phi_{2,x,p}^{(\text{space})}), & c = 0, \\ (b\rho_{1,x,p}^{(\text{space})} - \psi_{1,x,p}^{(\text{space})}) \wedge (b\rho_{2,x,p}^{(\text{space})} - \psi_{2,x,p}^{(\text{space})}), & c = 1/2, \end{cases}$$

where  $x = \alpha(\beta + 1) \in (0, 1/2]$ .

(ii) Consider the case that  $1/2 < \alpha(\beta + 1) \le 1$ . By choosing  $\gamma_{\delta_j} > \frac{\alpha(\beta + \delta_j + 1) - 1}{2}$  (j = 1, 2, 3), we have  $\varphi_{\beta+\delta_j-2\gamma_{\delta_j}/\alpha}(\nu) \sim \nu^{\alpha(\beta+\delta_j+1)-2\gamma_{\delta_j}-1}$ . Hence, it is enough to show that for c = 0, 1/2, there exist  $\delta_j$ ,  $\gamma_{\delta_j}$  (j = 1, 2, 3) and  $\gamma_{-\delta_1}$  such that (4.32) under [B1]\_{\beta,p} and [C2]\_{\beta,p}, respectively. Let  $M = N^b$ ,  $b \ge 1$  and  $E = \{\delta | \delta > \frac{1}{\alpha}(1 + \frac{2}{p}) - (\beta + 1)\}$ . Note that

$$\frac{1}{p} < \gamma_{\delta} < \begin{cases} \frac{\alpha(\beta+\delta+1)}{2} - \frac{1}{p}, & \delta \in E^{c}, \\ \frac{1}{2}, & \delta \in E. \end{cases}$$

It then follows from  $\delta < \frac{2\gamma_{\delta}+1}{\alpha} - (\beta+1)$   $(\delta \in E)$  that

$$\sup_{\delta_{1}} \sup_{\gamma_{\delta_{1}}} \sup_{\gamma_{-\delta_{1}}} \frac{b(\gamma_{\delta_{1}} + \gamma_{-\delta_{1}} - 2/p) + \alpha(\beta + \delta_{1} + 1)/2 - \gamma_{\delta_{1}} - 1}{\gamma_{\delta_{1}} + \gamma_{-\delta_{1}} + c} \\ = \Gamma_{\alpha(\beta+1),p,c}^{(1)} \lor \sup_{\delta_{1} \in E} \left( b - 1 + \frac{-b(c+2/p) + \alpha(\beta+1) - 1/p + c - 1}{1/2 + \alpha(\beta - \delta_{1} + 1)/2 - 1/p + c} \right) = \Gamma_{\alpha(\beta+1),p,c}^{(1)},$$

$$\begin{split} \sup_{\delta_{2}} \sup_{\gamma \delta_{2}} \frac{b(\gamma_{\delta_{2}} - 1/p) + \alpha(\beta + \delta_{2} + 1)/2 - \gamma_{\delta_{2}} - 1}{\gamma_{\delta_{2}} + c} \\ &= \sup_{\delta_{2} \in E^{c}} \left( b - 1 + \frac{-b(c + 1/p) + c + \alpha(\beta + \delta_{2} + 1)/2 - 1}{\alpha(\beta + \delta_{2} + 1)/2 - 1/p + c} \right) \\ &\lor \sup_{\gamma \delta_{2} \in (1/p, 1/2)} \left( b + \frac{-b(c + 1/p) - 1/2}{\gamma_{\delta_{2}} + c} \right) \\ &= b \left( 1 - \frac{c + 1/p}{1/2 + c} \right) - \frac{(1 - 1/p) \wedge 1/2}{1/2 + c} \\ &= b \left( 1 - \frac{c + 1/p}{1/2 + c} \right) - \frac{1/2}{1/2 + c} =: \widetilde{\Gamma}^{(3)}_{\alpha(\beta + 1), p, c}, \end{split}$$

and

$$\begin{split} \sup_{\delta_{3}} \sup_{\gamma_{\delta_{3}}} \frac{b(\gamma_{\delta_{3}} - 1/p) + \alpha(\beta - \delta_{3} + 1)/2 - 1}{\gamma_{\delta_{3}} + c} \\ &= \sup_{\delta_{3} \in E^{c}} \left( b + \frac{-b(c + 1/p) + \alpha(\beta - \delta_{3} + 1)/2 - 1}{\alpha(\beta + \delta_{3} + 1)/2 - 1/p + c} \right) \\ &\quad \lor \sup_{\delta_{3} \in E} \left( b + \frac{-b(c + 1/p) + \alpha(\beta - \delta_{3} + 1)/2 - 1}{1/2 + c} \right) \\ &= b \left( 1 - \frac{c + 1/p}{1/2 + c} \right) - \frac{3/2 - \alpha(\beta + 1) + 1/p}{1/2 + c} =: \Gamma^{(3)}_{\alpha(\beta + 1), p, c} \leq \widetilde{\Gamma}^{(3)}_{\alpha(\beta + 1), p, c}. \end{split}$$

Therefore, we get (4.32) under  $\nu N^u \to \infty$  for some

$$u < \Gamma_{x,p,c}^{(1)} \land \Gamma_{x,p,c}^{(3)} = \begin{cases} (b\tau_{1,x,p}^{(\text{space})} - \phi_{1,x,p}^{(\text{space})}) \land (b\tau_{2,x,p}^{(\text{space})} - \phi_{2,x,p}^{(\text{space})}), & c = 0, \\ (b\rho_{1,x,p}^{(\text{space})} - \psi_{1,x,p}^{(\text{space})}) \land (b\rho_{2,x,p}^{(\text{space})} - \psi_{2,x,p}^{(\text{space})}), & c = 1/2, \end{cases}$$

where  $x = \alpha(\beta + 1) \in (1/2, 1]$ .

# **4.2.3** Control of truncation error and derivation of $\hat{\theta}_{0,\beta}$

Because

$$\langle u, v \rangle_{\mathcal{L}^2_{\beta}} = \sum_{l_1, l_2 \ge 1} \mu_{l_1, l_2}^{-\alpha\beta} \langle u, e_{l_1, l_2} \rangle \langle v, e_{l_1, l_2} \rangle, \quad u, v \in \mathcal{L}^2_{\beta},$$

the estimator  $\hat{\theta}_{0,\beta}^{(\mathrm{disc})}$  given in (4.29) can be represented as

$$\hat{\theta}_{0,\beta}^{(\text{disc})} = \frac{\sum_{i=1}^{N} \sum_{l_{1},l_{2} \ge 1} \mu_{l_{1},l_{2}}^{-\alpha\beta} \langle \Psi_{M} X_{t_{i-1}}, e_{l_{1},l_{2}} \rangle \langle Y_{i}^{\Psi_{M}}, e_{l_{1},l_{2}} \rangle}{\frac{1}{N} \sum_{i=1}^{N} \sum_{l_{1},l_{2} \ge 1} \mu_{l_{1},l_{2}}^{-\alpha\beta} \langle \Psi_{M} X_{t_{i-1}}, e_{l_{1},l_{2}} \rangle^{2}}.$$

In practice, we need to truncate the infinite sums in the estimator  $\hat{\theta}_{0,\beta}^{(\text{disc})}$ . We thus consider the estimator

$$\hat{\theta}_{0,\beta} = \hat{\theta}_{0,\beta,\nu,N,M,L} = \frac{\sum_{i=1}^{N} \sum_{l_{1}=1}^{L} \sum_{l_{2}=1}^{L} \mu_{l_{1},l_{2}}^{-\alpha\beta} \langle \Psi_{M} X_{t_{i-1}}, e_{l_{1},l_{2}} \rangle \langle Y_{i}^{\Psi_{M}}, e_{l_{1},l_{2}} \rangle}{\frac{1}{N} \sum_{i=1}^{N} \sum_{l_{1}=1}^{L} \sum_{l_{2}=1}^{L} \mu_{l_{1},l_{2}}^{-\alpha\beta} \langle \Psi_{M} X_{t_{i-1}}, e_{l_{1},l_{2}} \rangle^{2}}.$$
(4.33)

For  $\beta > -1$ , define

$$\overline{J}_{\beta,\nu} = \overline{J}_{\beta,\nu,N,M,L} = \frac{1}{N} \sum_{i=1}^{N} \sum_{l_1=1}^{L} \sum_{l_2=1}^{L} \mu_{l_1,l_2}^{-\alpha\beta} \langle \Psi_M X_{t_{i-1}}, e_{l_1,l_2} \rangle^2,$$
  
$$\overline{K}_{\beta,\nu} = \overline{K}_{\beta,\nu,N,M,L} = \sum_{i=1}^{N} \sum_{l_1=1}^{L} \sum_{l_2=1}^{L} \mu_{l_1,l_2}^{-\alpha\beta} \langle \Psi_M X_{t_{i-1}}, e_{l_1,l_2} \rangle \langle Y_i^{\Psi_M}, e_{l_1,l_2} \rangle.$$

The following proposition shows the sufficient conditions for the asymptotic equivalence of the estimators  $\hat{\theta}_{0,\beta}^{(\text{disc})}$  and  $\hat{\theta}_{0,\beta}$ .

**Proposition 4.5.** Let  $\alpha > 0$ ,  $-1 < \beta \leq \frac{1}{\alpha} - 1$  and  $p > \frac{4}{\alpha(\beta+1)}$ . Assume that  $[A1]_{-1,p}$  and  $[B1]_{\beta,p}$  hold.

- (1) Under  $[B2]_{\beta,p}$ , it holds that  $\hat{\theta}_{0,\beta}^{(\text{disc})} \hat{\theta}_{0,\beta} = \mathcal{O}_{\mathbf{P}}(1)$  as  $\nu \to 0$ ,  $N \to \infty$ ,  $M \to \infty$  and  $L \to \infty$ .
- (2) Under  $[C3]_{\beta,p}$ , it holds that  $\mathcal{R}_{\beta,\nu}(\hat{\theta}_{0,\beta}^{(\text{disc})} \hat{\theta}_{0,\beta}) = \mathcal{O}_{\mathbf{P}}(1)$  as  $\nu \to 0$ ,  $N \to \infty$ ,  $M \to \infty$  and  $L \to \infty$ .

*Proof.* Recall  $\hat{\theta}_{0,\beta}^{(\text{disc})} = \widetilde{K}_{\beta,\nu} / \widetilde{J}_{\beta,\nu}$ . Let  $\overline{\Delta}_{\beta,\nu}^{(1)} = |\overline{J}_{\beta,\nu} - \widetilde{J}_{\beta,\nu}|$  and  $\overline{\Delta}_{\beta,\nu}^{(2)} = |\overline{K}_{\beta,\nu} - \widetilde{K}_{\beta,\nu}|$ . Note that

$$|\hat{\theta}_{0,\beta}^{(\text{disc})} - \hat{\theta}_{0,\beta}| \le \frac{|\hat{\theta}_{0,\beta}^{(\text{disc})}|\overline{\Delta}_{\beta,\nu}^{(1)} + \overline{\Delta}_{\beta,\nu}^{(2)}}{\widetilde{J}_{\beta,\nu} - \overline{\Delta}_{\beta,\nu}^{(1)}}.$$

Let i = 1, 2. Since it follows from Proposition 4.4, (4.30) and (4.31) that  $\hat{\theta}_{0,\beta}^{(\text{disc})} = \mathcal{O}_{\mathbf{P}}(1)$  and  $\varphi_{\beta}(\nu)/\widetilde{J}_{\beta,\nu} = \mathcal{O}_{\mathbf{P}}(1)$  under  $[\text{B1}]_{\beta,p}$ , it is enough to show that

$$\overline{\Delta}_{\beta,\nu}^{(i)} = \mathcal{O}_{\mathbf{P}}(\varphi_{\beta}(\nu)) \tag{4.34}$$

under  $[B2]_{\beta,p}$ , and

$$\overline{\Delta}_{\beta,\nu}^{(i)} = \mathcal{O}_{\mathbf{P}}(\varphi_{2\beta+1}(\nu)^{1/2})$$
(4.35)

under  $[C3]_{\beta,p}$ .

**Step 1:** We estimate  $\overline{\Delta}_{\beta,\nu}^{(1)}$  and  $\overline{\Delta}_{\beta,\nu}^{(2)}$ . It holds that for  $0 < \delta < \alpha(\beta+1) - \frac{4}{p}$ ,

$$\sum_{l_1^2+l_2^2>L^2} \mu_{l_1,l_2}^{-\alpha\beta} \langle \Psi_M X_t, e_{l_1,l_2} \rangle^2 \lesssim \frac{1}{L^{2\delta}} \sum_{l_1,l_2\geq 1} \mu_{l_1,l_2}^{\delta-\alpha\beta} \langle \Psi_M X_t, e_{l_1,l_2} \rangle^2 = \frac{\|\Psi_M X_t\|_{\mathcal{L}^2_{\beta-\delta/\alpha}}^2}{L^{2\delta}}$$
$$\sum_{l_1^2+l_2^2>L^2} \mu_{l_1,l_2}^{-\alpha\beta} |\langle \Psi_M X_t, e_{l_1,l_2} \rangle || \langle Y_i^{\Psi_M}, e_{l_1,l_2} \rangle || \lesssim \frac{\|\Psi_M X_t\|_{\mathcal{L}^2_{\beta-\delta/\alpha}}}{L^{2\delta}}.$$

Using Lemmas 8, 10 and 11, we see that for  $\frac{1}{p} < \gamma_{-\delta} < \frac{\alpha(\beta-\delta+1)}{2} - \frac{1}{p}$ ,

$$\begin{split} \mathbb{E}\Big[\|\Psi_M X_t\|^2_{\mathcal{L}^2_{\beta-\delta/\alpha}}\Big] \lesssim \mathbb{E}\Big[\|\Psi_M X_t - X_t\|^2_{\mathcal{L}^2_{\beta-\delta/\alpha}}\Big] + \mathbb{E}\Big[\|X_t\|^2_{\mathcal{L}^2_{\beta-\delta/\alpha}}\Big] \\ \lesssim \frac{\varphi_{\beta-\delta-2\gamma-\delta/\alpha}(\nu)}{M^{2(\gamma-\delta-1/p)}} + \nu^{\alpha(\beta+1)-\delta-1} \\ \lesssim \nu^{\alpha(\beta+1)-\delta-2\gamma_{\delta}-1}, \\ \mathbb{E}\Big[\|Y^{\Psi_M}_i\|^2_{\mathcal{L}^2_{\beta-\delta/\alpha}}\Big] \lesssim \frac{\nu^{\alpha(\beta+1)-\delta-2\gamma-\delta-1}}{N^{\alpha(\beta+1)-\delta-2\gamma-\delta}} \end{split}$$

and thus for i = 1, 2,

$$\mathbb{E}\Big[\overline{\Delta}_{\beta,\nu}^{(i)}\Big] \lesssim \frac{\nu^{\alpha(\beta+1)-\delta-2\gamma_{-\delta}-1}}{L^{2\delta}N^{(\alpha(\beta+1)-\delta-2\gamma_{-\delta})/2-1}} =: \mathcal{V}_{\beta,\delta,\gamma_{-\delta}}(\nu,N,L)$$

**Step 2:** In order to show (4.34) and (4.35), we verify that there exist  $\delta$  and  $\gamma_{-\delta}$  such that

$$(\nu^{1-c-\alpha(\beta+1)}\mathcal{V}_{\beta,\delta,\gamma_{-\delta}}(\nu,N,L))^{-1} = \nu^{\delta+2\gamma_{-\delta}+c}L^{2\delta}N^{\alpha(\beta-\delta+1)/2-\gamma_{-\delta}-1} \to \infty$$

for c = 0, 1/2, respectively. Let  $L = N^b$ ,  $b > \frac{\alpha - 1}{4}$ . Recall  $0 < \delta < \alpha(\beta + 1) - \frac{4}{p}$  and  $\frac{1}{p} < \gamma_{-\delta} < \frac{\alpha(\beta - \delta + 1)}{2} - \frac{1}{p}$ . Since

$$\begin{split} \sup_{\delta} \sup_{\gamma - \delta} \frac{2b\delta + \alpha(\beta - \delta + 1)/2 - \gamma_{-\delta} - 1}{\delta + 2\gamma_{-\delta} + c} \\ &= \sup_{\delta} \left( -\frac{1}{2} + \frac{(2b + (1 - \alpha)/2)\delta + \alpha(\beta + 1)/2 + c/2 - 1}{\delta + 2/p + c} \right) \\ &= 2b \left( 1 - \frac{2/p + c}{\alpha(\beta + 1) - 2/p + c} \right) - \frac{(\alpha - 1)\alpha(\beta + 1)/2 + (1 - 2\alpha)/p + 1}{\alpha(\beta + 1) - 2/p + c} \\ &= \begin{cases} b\tau_{x,p}^{(\text{trunc})} - \phi_{x,\alpha,p}^{(\text{trunc})}, & c = 0, \\ b\rho_{x,p}^{(\text{trunc})} - \psi_{x,\alpha,p}^{(\text{trunc})}, & c = 1/2, \end{cases}$$

we find that (4.34) and (4.35) hold under  $[B2]_{\beta,p}$  and  $[C3]_{\beta,p}$ , respectively.

From Propositions 4.3-4.5, we can prove that  $\hat{\theta}_{0,\beta}$  in (4.33) and  $\hat{\theta}_{0,\beta}^{(\text{cont})}$  are asymptotically equivalent. Finally, we conclude the proof of Theorem 2.2 by showing that  $\hat{\theta}_{0,\beta}$  in (4.33) equals that in (2.7). Since

$$\int_{b}^{c} \sqrt{2} \sin(\pi lx) \mathrm{e}^{ax/2} \mathrm{d}x = h_l(c:a) - h_l(b:a),$$

we have

$$\begin{split} \langle \Psi_M X_t, e_{l_1, l_2} \rangle &= \sum_{j=1}^{M_1} \sum_{k=1}^{M_2} \int_{z_{k-1}}^{z_k} \int_{y_{j-1}}^{y_j} 2X_t(y_{j-1}, z_{k-1}) \sin(\pi l_1 y) \sin(\pi l_2 z) \mathrm{e}^{(\kappa y + \eta z)/2} \mathrm{d}y \mathrm{d}z \\ &= \sum_{j=1}^{M_1} \sum_{k=1}^{M_2} X_t(y_{j-1}, z_{k-1}) \delta_j^{[y]} h_{l_1}(\kappa) \delta_k^{[z]} h_{l_2}(\eta) \\ &= [X_t]_{M, l_1, l_2} \end{split}$$

and

$$\langle Y_i^{\Psi_M}, e_{l_1, l_2} \rangle = \langle \Psi_M X_{t_i}, e_{l_1, l_2} \rangle - e^{\nu \lambda_{l_1, l_2}/N} \langle \Psi_M X_{t_{i-1}}, e_{l_1, l_2} \rangle$$
  
=  $[X_{t_i}]_{M, l_1, l_2} - e^{\nu \lambda_{l_1, l_2}/N} [X_{t_{i-1}}]_{M, l_1, l_2}.$ 

Therefore,  $\hat{\theta}_{0,\beta}$  in (4.33) can be expressed in the form of (2.7).

J

### 4.2.4 Auxiliary results for control of discretization errors

Here, we provide some lemmas which are useful in the proof of Theorem 2.2.

**Lemma 6.** Let  $\alpha > 0$ , q > 2 and  $\beta > \frac{2}{\alpha q} - 1$ . Then, for any  $0 \le u \le t \le 1$ ,

$$\mathbb{E}\Big[\|\overline{X}_{t,u}\|_{\mathcal{L}^q_{\beta}}^2\Big] \lesssim (t-u)^{\alpha(\beta+1)\wedge 1}\varphi_{\beta}(\nu).$$

Proof. Choose  $\frac{1}{q} < \gamma < \frac{\alpha(\beta+1) \wedge 1}{2}$  and define

$$Z_{\nu,\gamma}(t,u,\cdot) = \sigma \int_{u}^{t} (t-s)^{-\gamma} S_{\nu(t-s)} \mathrm{d}W_{s}^{Q}(\cdot).$$

Since

$$\begin{split} \|G_u(\boldsymbol{x},\cdot)\|_{\mathcal{L}^2_{\beta+1}}^2 &= \sum_{l_1,l_2 \ge 1} e^{-2\lambda_{l_1,l_2} u} \mu_{l_1,l_2}^{-\alpha(\beta+1)} e_{l_1,l_2}^2(\boldsymbol{x}) \\ &\lesssim \sum_{l_1,l_2 \ge 1} \frac{e^{-u(l_1^2+l_2^2)}}{(l_1^2+l_2^2)^{\alpha(\beta+1)}} \sim \iint_{\boldsymbol{x}^2+\boldsymbol{y}^2 \ge 1} \frac{e^{-u(\boldsymbol{x}^2+\boldsymbol{y}^2)}}{(\boldsymbol{x}^2+\boldsymbol{y}^2)^{\alpha(\beta+1)}} d\boldsymbol{x} d\boldsymbol{y} \\ &\lesssim \begin{cases} u^{\alpha(\beta+1)-1}, & \alpha(\beta+1) < 1, \\ -\log u, & \alpha(\beta+1) = 1, \\ 1, & \alpha(\beta+1) > 1 \end{cases} \end{split}$$

for  $u \in (0, 1]$ , we find from (4.17) that

$$\mathbb{E}\Big[\big(\widetilde{Q}^{\beta/2}Z_{\nu,\gamma}(t,u,\boldsymbol{x})\big)^2\Big] = \sigma^2 \int_u^t (t-s)^{-2\gamma} \|G_{\nu(t-s)}(\boldsymbol{x},\cdot)\|_{\mathcal{L}^2_{\beta+1}}^2 \mathrm{d}s$$

$$\lesssim \begin{cases} \nu^{\alpha(\beta+1)-1} \int_u^t (t-s)^{\alpha(\beta+1)-1-2\gamma} \mathrm{d}s, & \alpha(\beta+1) < 1, \\ -\int_u^t (t-s)^{-2\gamma} \log(\nu(t-s)) \mathrm{d}s, & \alpha(\beta+1) = 1, \\ \int_u^t (t-s)^{-2\gamma} \mathrm{d}s, & \alpha(\beta+1) > 1 \end{cases}$$

$$\sim (t-u)^{\alpha(\beta+1)\wedge 1-2\gamma} \times \begin{cases} \nu^{\alpha(\beta+1)-1}, & \alpha(\beta+1) < 1, \\ -\log(\nu(t-u)), & \alpha(\beta+1) = 1, \\ 1, & \alpha(\beta+1) > 1, \end{cases}$$

where the last estimation follows from

$$-\int_{u}^{t} (t-s)^{-2\gamma} \log(t-s) \mathrm{d}s = \frac{(t-u)^{1-2\gamma}}{(1-2\gamma)^{2}} (-(1-2\gamma)\log(t-u)+1).$$

Therefore, the Gaussianity of  $\widetilde{Q}^{\beta/2}Z_{\nu,\gamma}(t,u,\pmb{x})$  implies that

$$\begin{split} \mathbb{E}\Big[\|Z_{\nu,\gamma}(t,u)\|_{\mathcal{L}^{q}_{\beta}}^{q}\Big] &\lesssim \int_{D} \mathbb{E}\Big[\big|\widetilde{Q}^{\beta/2}Z_{\nu,\gamma}(t,u,\boldsymbol{x})\big|^{q}\Big] \mathrm{d}\boldsymbol{x} \\ &\sim \int_{D} \mathbb{E}\Big[\big(\widetilde{Q}^{\beta/2}Z_{\nu,\gamma}(t,u,\boldsymbol{x})\big)^{2}\Big]^{q/2} \mathrm{d}\boldsymbol{x} \\ &\lesssim (t-u)^{(\alpha(\beta+1)\wedge 1-2\gamma)q/2} \times \begin{cases} \nu^{(\alpha(\beta+1)-1)q/2}, & \alpha(\beta+1) < 1, \\ (-\log(\nu(t-u)))^{q/2}, & \alpha(\beta+1) = 1, \\ 1, & \alpha(\beta+1) > 1. \end{cases} \end{split}$$

Since it follows from Theorem 5.10 in [30] that

$$\overline{X}_{t,u} = \frac{\sigma \sin(\gamma \pi)}{\pi} \int_{u}^{t} (t-s)^{\gamma-1} S_{\nu(t-s)} Z_{\nu,\gamma}(s,u) \mathrm{d}s,$$

we see from (4.16), the Hölder inequality and

$$\int_0^t (-s^a \log(\nu s))^b \mathrm{d}s \lesssim t^{1+ab} (-\log(\nu t))^b$$

that for  $1/q + 1/q^* = 1$ ,

$$\mathbb{E}\left[\|\overline{X}_{t,u}\|_{\mathcal{L}_{\beta}^{q}}^{2}\right] \sim \mathbb{E}\left[\left\|\int_{u}^{t} (t-s)^{\gamma-1} S_{\nu(t-s)} Z_{\nu,\gamma}(s,u) \mathrm{d}s\right\|_{\mathcal{L}_{\beta}^{q}}^{2}\right]$$

$$\leq \mathbb{E}\left[\left(\int_{u}^{t} (t-s)^{\gamma-1} \|Z_{\nu,\gamma}(s,u)\|_{\mathcal{L}_{\beta}^{q}} \mathrm{d}s\right)^{2}\right]$$

$$\leq (t-u)^{2(q^{*}(\gamma-1)+1)/q^{*}} \mathbb{E}\left[\int_{u}^{t} \|Z_{\nu,\gamma}(s,u)\|_{\mathcal{L}_{\beta}^{q}}^{q} \mathrm{d}s\right]^{2/q}$$

$$\lesssim (t-u)^{2(\gamma-1/q)} (t-u)^{\alpha(\beta+1)\wedge 1-2\gamma+2/q}$$

$$\times \begin{cases} \nu^{\alpha(\beta+1)-1}, & \alpha(\beta+1) < 1, \\ -\log(\nu(t-u)), & \alpha(\beta+1) = 1, \\ 1, & \alpha(\beta+1) > 1 \end{cases}$$

$$\sim (t-u)^{\alpha(\beta+1)\wedge 1} \varphi_{\beta}(\nu).$$

**Lemma 7.** Let  $\alpha > 0$  and  $\beta > -1$ . Then, for any  $0 \le u \le t \le 1$ ,

$$\mathbb{E}\Big[\|\overline{X}_{t,u}\|_{\mathcal{L}^2_{\beta}}^2\Big] \sim \nu^{\alpha(\beta+1)-1}(t-u)^{\alpha(\beta+1)}.$$

*Proof.* The desired result follows from

$$\begin{split} \mathbb{E}\Big[\|\overline{X}_{t,u}\|_{\mathcal{L}^{2}_{\beta}}^{2}\Big] &= \sigma^{2} \int_{u}^{t} \|S_{\nu(t-s)}\|_{\mathrm{HS}(\mathcal{U}_{\beta})}^{2} \mathrm{d}s \\ &= \sigma^{2} \sum_{l_{1}, l_{2} \geq 1} \mu_{l_{1}, l_{2}}^{-\alpha(\beta+1)} \times \frac{1 - \mathrm{e}^{-2\lambda_{l_{1}, l_{2}}\nu(t-u)}}{2\lambda_{l_{1}, l_{2}}\nu} \\ &\sim \sum_{l_{1}, l_{2} \geq 1} (l_{1}^{2} + l_{2}^{2})^{-\alpha(\beta+1)} \Big\{ \frac{1}{(l_{1}^{2} + l_{2}^{2})\nu} \wedge (t-u) \Big\} \\ &= \nu^{-1} \sum_{l_{1}^{2} + l_{2}^{2} > (\nu(t-u))^{-1}} \frac{1}{(l_{1}^{2} + l_{2}^{2})^{1+\alpha(\beta+1)}} \\ &+ (t-u) \sum_{l_{1}^{2} + l_{2}^{2} \leq (\nu(t-u))^{-1}} \frac{\mathrm{d}x\mathrm{d}y}{(x^{2} + y^{2})^{1+\alpha(\beta+1)}} \\ &+ (t-u) \iint_{1 \leq x^{2} + y^{2} \leq (\nu(t-u))^{-1}} (x^{2} + y^{2})^{-\alpha(\beta+1)}\mathrm{d}x\mathrm{d}y \end{split}$$

$$\sim \nu^{\alpha(\beta+1)-1}(t-u)^{\alpha(\beta+1)}.$$

For  $\alpha > 0, \, \beta > -1$ , define

$$\widetilde{\varphi}_{\beta,q}(\nu) = \begin{cases} \nu^{\alpha(\beta+1)\wedge 1-1}, & q=2, \\ \varphi_{\beta}(\nu), & q>2. \end{cases}$$

**Lemma 8.** Let  $\alpha > 0$ . Assume that  $\beta$  and q satisfy any one of the following (i) and (ii).

- (i)  $q = 2, \beta > -1,$ (ii)  $q > 2, \beta > \frac{2}{\alpha q} - 1.$
- If  $[A1]_{\beta,q}$  holds, then

$$\mathbb{E}\Big[\|X_t\|_{\mathcal{L}^q_{\beta}}^2\Big] \lesssim 1 + \widetilde{\varphi}_{\beta,q}(\nu), \quad t \in \mathbb{T}.$$

Proof. It follows from (2.2), (4.16), [A1]\_{\beta,q} and Lemmas 6 and 7 that

$$\mathbb{E}\Big[\|X_t\|_{\mathcal{L}_{\beta}^q}^2\Big] \lesssim \mathbb{E}\Big[\|S_{\nu t}X_0\|_{\mathcal{L}_{\beta}^q}^2\Big] + \mathbb{E}\Big[\left\|\int_0^t S_{\nu(t-s)}X_s \mathrm{d}s\right\|_{\mathcal{L}_{\beta}^q}^2\Big] + \mathbb{E}\Big[\|\overline{X}_t\|_{\mathcal{L}_{\beta}^q}^2\Big] \\ \lesssim \mathbb{E}\Big[\|X_0\|_{\mathcal{L}_{\beta}^q}^2\Big] + \mathbb{E}\Big[\int_0^t \|S_{\nu(t-s)}X_s\|_{\mathcal{L}_{\beta}^q}^2 \mathrm{d}s\Big] + \widetilde{\varphi}_{\beta,q}(\nu) \\ \lesssim 1 + \widetilde{\varphi}_{\beta,q}(\nu) + \int_0^t \mathbb{E}\Big[\|X_s\|_{\mathcal{L}_{\beta}^q}^2\Big] \mathrm{d}s.$$

Using Gronwall's inequality, we have  $\mathbb{E}[\|X_t\|_{\mathcal{L}^q_{\beta}}^2] \lesssim 1 + \widetilde{\varphi}_{\beta,q}(\nu).$ 

**Lemma 9.** Let  $\alpha > 0$  and  $\beta > -1$ . Assume that  $[A1]_{\beta,2}$  holds. It holds that for  $0 \le \gamma < \frac{\alpha(\beta+1)}{2}$  and  $t \in [t_{i-1}, t_i], i = 1, \dots, N$ ,

$$\mathbb{E}\Big[\|X_t - X_{t_{i-1}}\|_{\mathcal{L}^2_{\beta}}^2\Big] \lesssim \frac{\nu^{\alpha(\beta+1)-1}}{N^{2\gamma\wedge 1}} + \left(\frac{\nu}{N}\right)^{2\gamma} + \frac{1}{N}.$$

*Proof.* Noting that

$$\|\mathcal{A}^{\gamma}u\|^{2} = \sum_{l_{1}, l_{2} \geq 1} \lambda_{l_{1}, l_{2}}^{2\gamma} \langle u, e_{l_{1}, l_{2}} \rangle^{2} \sim \sum_{l_{1}, l_{2} \geq 1} \mu_{l_{1}, l_{2}}^{-\alpha(-2\gamma/\alpha)} \langle u, e_{l_{1}, l_{2}} \rangle^{2} = \|\widetilde{Q}^{-\gamma/\alpha}u\|^{2} = \|u\|_{\mathcal{L}^{2}_{-2\gamma/\alpha}}^{2},$$

we see from Proposition 4.41 in [14], (4.23), (4.16), Lemmas 7 and 8 that for  $0 \le \gamma < \frac{\alpha(\beta+1)}{2}$ ,

$$\|S_t u - u\|_{\mathcal{L}^2_{\beta}} \lesssim t^{\gamma} \|\mathcal{A}^{\gamma} u\|_{\mathcal{L}^2_{\beta}} \sim t^{\gamma} \|u\|_{\mathcal{L}^2_{\beta-2\gamma/\alpha}}, \quad t \in \mathbb{T}$$

$$(4.36)$$

and

$$\begin{split} \mathbb{E}\Big[\|X_t - X_{t_{i-1}}\|_{\mathcal{L}^2_{\beta}}^2\Big] &\lesssim \mathbb{E}\Big[\|S_{\nu(t-t_{i-1})}X_{t_{i-1}} - X_{t_{i-1}}\|_{\mathcal{L}^2_{\beta}}^2\Big] \\ &+ \int_{t_{i-1}}^t \mathbb{E}\Big[\|S_{\nu(t-s)}X_s\|_{\mathcal{L}^2_{\beta}}^2\Big] \mathrm{d}s + \mathbb{E}\Big[\|\overline{X}_{t,t_{i-1}}\|_{\mathcal{L}^2_{\beta}}^2\Big] \\ &\lesssim \Big(\frac{\nu}{N}\Big)^{2\gamma} \mathbb{E}\Big[\|X_{t_{i-1}}\|_{\mathcal{L}^2_{\beta-2\gamma/\alpha}}^2\Big] + \int_{t_{i-1}}^t \mathbb{E}\Big[\|X_s\|_{\mathcal{L}^2_{\beta}}^2\Big] \mathrm{d}s + \mathbb{E}\Big[\|\overline{X}_{t,t_{i-1}}\|_{\mathcal{L}^2_{\beta}}^2\Big] \end{split}$$

$$\lesssim \frac{\nu^{\alpha(\beta+1)-1} + \nu^{2\gamma}}{N^{2\gamma}} + \frac{\nu^{\alpha(\beta+1)-1} + 1}{N} + \frac{\nu^{\alpha(\beta+1)-1}}{N^{\alpha(\beta+1)}} \\ \lesssim \frac{\nu^{\alpha(\beta+1)-1}}{N^{2\gamma\wedge 1}} + \left(\frac{\nu}{N}\right)^{2\gamma} + \frac{1}{N}.$$

Let  $M = (M_1 \wedge M_2)^2$ .

**Lemma 10.** Let  $\alpha > 0$ , q > 2 and  $\beta > \frac{4}{\alpha q} - 1$ . Assume that  $[A1]_{\beta,q}$  holds. It holds that for  $\frac{1}{q} < \gamma < (\frac{\alpha(\beta+1)}{2} - \frac{1}{q}) \wedge \frac{1}{2}$ ,

$$\mathbb{E}\Big[\|X_t - \Psi_M X_t\|_{\mathcal{L}^2_{\beta}}^2\Big] \lesssim \frac{\varphi_{\beta - 2\gamma/\alpha}(\nu)}{M^{2(\gamma - 1/q)}}, \quad t \in \mathbb{T}.$$

*Proof.* Since  $\|u\|_{\mathcal{L}^q_{-2\gamma/\alpha}} \sim \|\mathcal{A}^{\gamma}u\|_{\mathcal{L}^q}$  and Theorem 16.15 in [39] yield  $\mathcal{L}^q_{-2\gamma/\alpha} \hookrightarrow \mathcal{W}^{2\gamma,q}$ , we find from the Sobolev embedding theorem (Theorem 8.2 in [26]) that

$$\|u\|_{\mathcal{C}^{2(\gamma-1/q)}} \lesssim \|\mathcal{A}^{\gamma}u\|_{\mathcal{L}^{q}} \sim \|u\|_{\mathcal{L}^{q}_{-2\gamma/\alpha}}.$$

Therefore, it holds from  $\|\widetilde{Q}^{\beta/2}X_t\|_{\mathcal{C}^{2(\gamma-1/q)}} \lesssim \|X_t\|_{\mathcal{L}^q_{\beta-2\gamma/\alpha}}$  and Lemma 8 that for  $t \in \mathbb{T}$ ,

$$\begin{split} \|X_t - \Psi_M X_t\|_{\mathcal{L}^2_{\beta}}^2 &= \sum_{j=1}^{M_1} \sum_{k=1}^{M_2} \int_{z_{k-1}}^{z_k} \int_{y_{j-1}}^{y_j} (\widetilde{Q}^{\beta/2} X_t(y, z) - \widetilde{Q}^{\beta/2} X_t(y_{j-1}, z_{k-1}))^2 \bar{v}(y, z) \mathrm{d}y \mathrm{d}z \\ &\leq \sum_{j=1}^{M_1} \sum_{k=1}^{M_2} \int_{z_{k-1}}^{z_k} \int_{y_{j-1}}^{y_j} \|\widetilde{Q}^{\beta/2} X_t\|_{\mathcal{C}^2(\gamma-1/q)}^2 \left| \begin{pmatrix} y - y_{j-1} \\ z - z_{k-1} \end{pmatrix} \right|^{4(\gamma-1/q)} \bar{v}(y, z) \mathrm{d}y \mathrm{d}z \\ &\lesssim \left( \frac{\|X_t\|_{\mathcal{L}^q_{\beta-2\gamma/\alpha}}}{M^{\gamma-1/q}} \right)^2 \end{split}$$

and

$$\mathbb{E}\Big[\|X_t - \Psi_M X_t\|_{\mathcal{L}^2_{\beta}}^2\Big] \lesssim \frac{\mathbb{E}\Big[\|X_t\|_{\mathcal{L}^2_{\beta-2\gamma/\alpha}}^2\Big]}{M^{2(\gamma-1/q)}} \lesssim \frac{\varphi_{\beta-2\gamma/\alpha}(\nu)}{M^{2(\gamma-1/q)}}.$$

Lemma 11. Let  $\alpha > 0$ ,  $Y_i = X_{t_i} - S_{\nu(t_i - t_{i-1})} X_{t_{i-1}}$  and  $Y_i^{\Psi_M} = \Psi_M X_{t_i} - S_{\nu(t_i - t_{i-1})} \Psi_M X_{t_{i-1}}$ .

(1) For q and  $\beta$  satisfying either (i) or (ii) of Lemma 8, assume that  $[A1]_{\beta,q}$  holds. Then,

$$\mathbb{E}\Big[\|Y_i\|_{\mathcal{L}^q_{\beta}}^2\Big] \lesssim \frac{\widetilde{\varphi}_{\beta,q}(\nu)}{N^{\alpha(\beta+1)}}, \quad i=1,\ldots,N.$$

 $(2) \ \ For \ q>2 \ and \ \beta>\frac{4}{\alpha q}-1, \ assume \ that \ [A1]_{\beta,q} \ holds. \ It \ holds \ that \ for \ \frac{1}{q}<\gamma<(\frac{\alpha(\beta+1)}{2}-\frac{1}{q})\wedge\frac{1}{2},$ 

$$\mathbb{E}\Big[\|Y_i - Y_i^{\Psi_M}\|_{\mathcal{L}^2_{\beta}}^2\Big] \lesssim \frac{\varphi_{\beta-2\gamma/\alpha}(\nu)}{M^{2(\gamma-1/q)}N^{\alpha(\beta+1)-2\gamma}}, \quad i = 1, \dots, N.$$

Proof. (1) Using (4.23), (4.16) and Lemmas 6-8, we get

$$\mathbb{E}\Big[\|Y_i\|_{\mathcal{L}^q_{\beta}}^2\Big] \lesssim \mathbb{E}\Big[\left\|\int_{t_{i-1}}^{t_i} S_{\nu(t_i-s)} X_s \mathrm{d}s\right\|_{\mathcal{L}^q_{\beta}}^2\Big] + \mathbb{E}\Big[\|\overline{X}_{t_i,t_{i-1}}\|_{\mathcal{L}^q_{\beta}}^2\Big]$$

$$\begin{split} &\lesssim \mathbb{E}\bigg[\int_{t_{i-1}}^{t_i} \|S_{\nu(t_i-s)}X_s\|_{\mathcal{L}^q_{\beta}}^2 \mathrm{d}s\bigg] + \mathbb{E}\Big[\|\overline{X}_{t_i,t_{i-1}}\|_{\mathcal{L}^q_{\beta}}^2\Big] \\ &\lesssim \frac{1}{N} \int_{t_{i-1}}^{t_i} \mathbb{E}\Big[\|X_s\|_{\mathcal{L}^q_{\beta}}^2\Big] \mathrm{d}s + \mathbb{E}\Big[\|\overline{X}_{t_i,t_{i-1}}\|_{\mathcal{L}^q_{\beta}}^2\Big] \\ &\lesssim \frac{1+\widetilde{\varphi}_{\beta,q}(\nu)}{N^2} + \frac{\widetilde{\varphi}_{\beta,q}(\nu)}{N^{\alpha(\beta+1)}} \\ &\lesssim \frac{\widetilde{\varphi}_{\beta,q}(\nu)}{N^{\alpha(\beta+1)}}. \end{split}$$

(2) In the same way as Lemma 10, we see from (1) that

$$\mathbb{E}\Big[\|Y_i - Y_i^{\Psi_M}\|_{\mathcal{L}^2_{\beta}}^2\Big] \lesssim \frac{\mathbb{E}\Big[\|Y_i\|_{\mathcal{L}^q_{\beta-2\gamma/\alpha}}^2\Big]}{M^{2(\gamma-1/q)}} \lesssim \frac{\varphi_{\beta-2\gamma/\alpha}(\nu)}{M^{2(\gamma-1/q)}N^{\alpha(\beta+1)-2\gamma}}.$$

# 4.3 Proofs of Proposition 2.3 and Corollary 2.4

In this subsection, we provide proofs of the results in Subsection 2.4.

**Proof of Proposition 2.3.** Let  $\mathcal{A}_{\theta_0,\nu} = \nu \mathcal{A} - \theta_0$ ,  $T_{\nu,t} = e^{-t\mathcal{A}_{\theta_0,\nu}}$  and  $\lambda_{l_1,l_2}^{(\theta_0)} = \nu \lambda_{l_1,l_2} - \theta_0$ . Note that  $\mathcal{A}_{\theta_0,\nu}e_{l_1,l_2} = \lambda_{l_1,l_2}^{(\theta_0)}e_{l_1,l_2}$ . Moreover, let

$$f_1(x) = \begin{cases} \frac{1-e^{-x}}{x}, & x \neq 0, \\ 1, & x = 0, \end{cases} \quad f_2(x) = \begin{cases} \frac{x-1+e^{-x}}{x^2}, & x \neq 0, \\ \frac{1}{2}, & x = 0. \end{cases}$$

It follows from  $X_t = T_{\nu,t}X_0 + \sigma \int_0^t T_{\nu,t-s} dW_s^Q$  and the independence of  $X_0$  and  $W_t^Q$  that

$$\begin{split} \mathbb{E}[J_{\beta,\nu}] &= \int_{0}^{1} \mathbb{E}\Big[ \|T_{\nu,t}X_{0}\|_{\mathcal{L}^{2}_{\beta}}^{2} \Big] \mathrm{d}t + \sigma^{2} \int_{0}^{1} \int_{0}^{t} \|T_{\nu,t-s}\|_{\mathrm{HS}(\mathcal{U}_{\beta})}^{2} \mathrm{d}s \mathrm{d}t \\ &= \int_{0}^{1} \sum_{l_{1},l_{2} \geq 1} \mathrm{e}^{-2t\lambda_{l_{1},l_{2}}^{(\theta_{0})}} \mu_{l_{1},l_{2}}^{-\alpha\beta} \mathbb{E}[\langle X_{0}, e_{l_{1},l_{2}} \rangle^{2}] \mathrm{d}t \\ &\quad + \sigma^{2} \int_{0}^{1} \int_{0}^{t} \sum_{l_{1},l_{2} \geq 1} \mathrm{e}^{-2(t-s)\lambda_{l_{1},l_{2}}^{(\theta_{0})}} \mu_{l_{1},l_{2}}^{-\alpha(\beta+1)} \mathrm{d}s \mathrm{d}t \\ &= \sum_{l_{1},l_{2} \geq 1} f_{1}(2\lambda_{l_{1},l_{2}}^{(\theta_{0})}) \mu_{l_{1},l_{2}}^{-\alpha\beta} \mathbb{E}[\langle X_{0}, e_{l_{1},l_{2}} \rangle^{2}] + \sigma^{2} \sum_{l_{1},l_{2} \geq 1} f_{2}(2\lambda_{l_{1},l_{2}}^{(\theta_{0})}) \mu_{l_{1},l_{2}}^{-\alpha(\beta+1)} \\ &= \sum_{l_{1},l_{2} \geq 1} f_{l_{1},l_{2}}(2\lambda_{l_{1},l_{2}}^{(\theta_{0})}) \mu_{l_{1},l_{2}}^{-\alpha(\beta+1)}, \end{split}$$

where  $f_{l_1,l_2}(x) = f_1(x)\mu_{l_1,l_2}^{\alpha}\mathbb{E}[\langle X_0, e_{l_1,l_2}\rangle^2] + \sigma^2 f_2(x) > 0, x \in \mathbb{R}$ . Note that  $f_{l_1,l_2}(x) \lesssim 1/x$  under [A1]<sub>-1,2</sub>. Therefore, we obtain

$$\mathcal{R}_{\beta,\nu} = \frac{\mathbb{E}[J_{\beta,\nu}]}{\sqrt{\mathbb{E}[J_{2\beta+1,\nu}]}} = \frac{\sum_{l_1,l_2 \ge 1} f_{l_1,l_2}(2\lambda_{l_1,l_2}^{(\theta_0)})\mu_{l_1,l_2}^{-\alpha(\beta+1)}}{\sqrt{\sum_{l_1,l_2 \ge 1} f_{l_1,l_2}(2\lambda_{l_1,l_2}^{(\theta_0)})\mu_{l_1,l_2}^{-2\alpha(\beta+1)}}}$$

For x > 0, let

$$F(x) = \sum_{l_1, l_2 \ge 1} f_{l_1, l_2} (2\lambda_{l_1, l_2}^{(\theta_0)}) \mu_{l_1, l_2}^{-x}, \quad G(x) = \frac{F(x)^2}{F(2x)}.$$

Since F(x) is differentiable and

$$F'(x) = -\sum_{l_1, l_2 \ge 1} f_{l_1, l_2} (2\lambda_{l_1, l_2}^{(\theta_0)}) \mu_{l_1, l_2}^{-x} \log \mu_{l_1, l_2},$$

we have

$$\begin{split} F'(x)F(2x) &- F(x)F'(2x) \\ &= \sum_{l_1,\dots,l_4 \ge 1} f_{l_1,l_2}(2\lambda_{l_1,l_2}^{(\theta_0)}) f_{l_3,l_4}(2\lambda_{l_3,l_4}^{(\theta_0)}) \mu_{l_1,l_2}^{-2x} \mu_{l_3,l_4}^{-2x} (\mu_{l_3,l_4}^x - \mu_{l_1,l_2}^x) \log \mu_{l_1,l_2} \\ &= \sum_{\substack{l_1,\dots,l_4 \ge 1, \\ l_1^2 + l_2^2 < l_3^2 + l_4^2}} f_{l_1,l_2}(2\lambda_{l_1,l_2}^{(\theta_0)}) f_{l_3,l_4}(2\lambda_{l_3,l_4}^{(\theta_0)}) \mu_{l_1,l_2}^{-2x} \mu_{l_3,l_4}^{-2x} (\mu_{l_3,l_4}^x - \mu_{l_1,l_2}^x) \log \mu_{l_1,l_2} \\ &+ \sum_{\substack{l_1,\dots,l_4 \ge 1, \\ l_1^2 + l_2^2 > l_3^2 + l_4^2}} f_{l_1,l_2}(2\lambda_{l_1,l_2}^{(\theta_0)}) f_{l_3,l_4}(2\lambda_{l_3,l_4}^{(\theta_0)}) \mu_{l_1,l_2}^{-2x} \mu_{l_3,l_4}^{-2x} (\mu_{l_3,l_4}^x - \mu_{l_1,l_2}^x) \log \mu_{l_1,l_2} \\ &= \sum_{\substack{l_1,\dots,l_4 \ge 1, \\ l_1^2 + l_2^2 < l_3^2 + l_4^2}} f_{l_1,l_2}(2\lambda_{l_1,l_2}^{(\theta_0)}) f_{l_3,l_4}(2\lambda_{l_3,l_4}^{(\theta_0)}) \mu_{l_1,l_2}^{-2x} \mu_{l_3,l_4}^{-2x} (\mu_{l_3,l_4}^x - \mu_{l_1,l_2}^x) \log \mu_{l_1,l_2} \\ &= \sum_{\substack{l_1,\dots,l_4 \ge 1, \\ l_1^2 + l_2^2 < l_3^2 + l_4^2}} f_{l_1,l_2}(2\lambda_{l_1,l_2}^{(\theta_0)}) f_{l_3,l_4}(2\lambda_{l_3,l_4}^{(\theta_0)}) \mu_{l_1,l_2}^{-2x} \mu_{l_3,l_4}^{-2x} (\mu_{l_3,l_4}^x - \mu_{l_1,l_2}^x) \log \mu_{l_1,l_2} \\ &\leq 0 \end{split}$$

and hence

$$G'(x) = \frac{2F(x)}{F(2x)^2} (F'(x)F(2x) - F(x)F'(2x)) < 0,$$

which implies that the function  $\beta \mapsto \mathcal{R}_{\beta,\nu}$  is decreasing for  $\beta \in (-1, \frac{1}{\alpha} - 1]$ .

Proof of Corollary 2.4. Fix m > n > 0. (1) Since  $\nu (M_1 \wedge M_2)^{2r} N^{-\phi_x^{(\text{space})}} = \nu^{1-mr+n\phi_x^{(\text{space})}}$ , it follows from  $[B1]_{\beta,\infty}$  that

$$m > \frac{1 + n\phi_{\alpha(\beta+1)}^{(\text{space})}}{r} > 1 + n\phi_{\alpha(\beta+1)}^{(\text{space})}.$$
(4.37)

Because  $\phi_x^{(\text{space})} \ge \phi_1^{(\text{space})} = 1$ , there exists  $\beta$  such that (4.37) if m > n + 1. Solving the inequality (4.37) for  $\beta$ , we have  $\beta > \beta_{\alpha,n,m}^{(\text{cons})}$ . Noting that  $\nu L^s N^{-\phi_{x,\alpha}^{(\text{trunc})}} = \nu^{1-\ell s + n\phi_{x,\alpha}^{(\text{trunc})}}$ , we find from  $[B2]_{\beta,\infty}$ that for  $\beta > \beta_{\alpha,n,m}^{(\text{cons})}$ ,

$$\ell > \frac{1 + n\phi_{\alpha(\beta+1),\alpha}^{(trunc)}}{s} > \frac{1 + n\phi_{\alpha(\beta+1),\alpha}^{(trunc)}}{2}$$

(2) Notice that  $\nu N^q = \nu^{1-nq}$  and  $\nu (M_1 \wedge M_2)^{2r_j} N^{-\psi_{j,x}^{(\text{space})}} = \nu^{1-mr_j+n\psi_{j,x}^{(\text{space})}}$ . Hence, we see from  $[C1]_{\beta}$  and  $[C2]_{\beta,\infty}$  that

$$\begin{split} \mathbf{n} > \frac{1}{q} > \frac{1}{\rho_{\alpha(\beta+1)}^{(\text{time})}} = \begin{cases} \frac{1}{2\alpha(\beta+1)}, & \alpha(\beta+1) \le \frac{1}{2}, \\ 2(1-\alpha(\beta+1)), & \alpha(\beta+1) > \frac{1}{2}, \end{cases} \\ \mathbf{m} > \frac{1+\mathbf{n}\psi_{1,\alpha(\beta+1)}^{(\text{space})}}{r_1} \lor \frac{1+\mathbf{n}\psi_{2,\alpha(\beta+1)}^{(\text{space})}}{r_2} > \frac{1+\mathbf{n}\psi_{1,\alpha(\beta+1)}^{(\text{space})}}{\rho_{\alpha(\beta+1)}^{(\text{space})}} \lor 2(1+\mathbf{n}\psi_{2,\alpha(\beta+1)}^{(\text{space})}), \end{split}$$

which together with  $\psi_{1,x}^{(\text{space})} \ge \psi_{1,1}^{(\text{space})} = 2/3$ ,  $\psi_{2,x}^{(\text{space})} \ge \psi_{2,1}^{(\text{space})} = 1/2$  and  $\rho_x^{(\text{space})} \le \rho_1^{(\text{space})} = 2/3$ imply that  $\beta > \beta_{\alpha,n,m}^{(\text{asym})}$  for m > n + 2. Since  $\nu L^s N^{-\psi_{x,\alpha}^{(\text{trunc})}} = \nu^{1-\ell s + n\psi_{x,\alpha}^{(\text{trunc})}}$  and  $[\text{C3}]_{\beta,\infty}$ , we find

that for  $\beta > \beta_{\alpha,n,m}^{(asym)}$ ,

$$\ell > \frac{1 + \mathrm{n}\psi^{(\mathrm{trunc})}_{\alpha(\beta+1),\alpha}}{s} > \frac{1 + \mathrm{n}\psi^{(\mathrm{trunc})}_{\alpha(\beta+1),\alpha}}{\rho^{(\mathrm{trunc})}_{\alpha(\beta+1)}}.$$

#### Proofs of Propositions 2.5 and 2.6 **4.4**

This subsection gives proofs of Propositions 2.5 and 2.6. Since the proofs are analogous to those of Proposition 2.1 and Theorem 2.2 in [34], we shall focus on the different parts. For the detailed proofs, refer to [34].

Let  $e_l^{(1)}(x) = \sqrt{2} \sin(\pi l x) e^{-\kappa x/2}$  and  $e_l^{(2)}(x) = \sqrt{2} \sin(\pi l x) e^{-\eta x/2}$ . The second order moment of the triple increment  $T_{i,j,k}X$  is given by

$$\mathbb{E}[(T_{i,j,k}X)^2] = \sigma^2 F_{j,k}^{j,k} + R_{i,j,k},$$

where

$$F_{j',k'}^{j,k} = \sum_{l_1,l_2 \ge 1} \frac{1 - e^{\lambda_{l_1,l_2}^{(\theta_0)} \Delta}}{\lambda_{l_1,l_2}^{(\theta_0)} \mu_{l_1,l_2}^{\alpha}} (e_{l_1}^{(1)}(\widetilde{y}_j) - e_{l_1}^{(1)}(\widetilde{y}_{j-1})) (e_{l_2}^{(2)}(\widetilde{z}_k) - e_{l_2}^{(2)}(\widetilde{z}_{k-1})) \times (e_{l_1}^{(1)}(\widetilde{y}_{j'}) - e_{l_1}^{(1)}(\widetilde{y}_{j'-1})) (e_{l_2}^{(2)}(\widetilde{z}_{k'}) - e_{l_2}^{(2)}(\widetilde{z}_{k'-1}))$$

and  $R_{i,j,k}$  satisfies  $\sum_{i=1}^{N} R_{i,j,k} \lesssim F_{j,k}^{j,k}$ . For a function  $f: \mathbb{R}^2 \to \mathbb{R}$  and a positive number  $\epsilon$ , define  $D_{1,\epsilon}f(x,y) = f(x+\epsilon,y) - f(x,y)$ and  $D_{2,\epsilon}f(x,y) = f(x,y+\epsilon) - f(x,y)$ . For  $\alpha \in (0,2)$  and  $y, z \in [0,2)$ , let

$$F_{\alpha}(y,z) = \sum_{l_{1},l_{2}\geq 1} \frac{1 - e^{-\lambda_{l_{1},l_{2}}^{(\theta_{0})}\Delta}}{\lambda_{l_{1},l_{2}}^{(\theta_{0})}\mu_{l_{1},l_{2}}^{\alpha}} \cos(\pi l_{1}y)\cos(\pi l_{2}z)$$
$$= \nu^{\alpha}\Delta^{1+\alpha}\sum_{l_{1},l_{2}\geq 1} \frac{1 - e^{-\lambda_{l_{1},l_{2}}^{(\theta_{0})}\Delta}}{\lambda_{l_{1},l_{2}}^{(\theta_{0})}\Delta} \cdot \frac{1}{(\mu_{l_{1},l_{2}}\nu\Delta)^{\alpha}}\cos(\pi l_{1}y)\cos(\pi l_{2}z).$$

Under  $\nu \to 0$ , the following lemma holds instead of Lemmas 4.10 and 4.11 in [34].

**Lemma 12.** For  $\alpha \in (0,2)$  and  $\delta = r\sqrt{\nu\Delta}$ , it holds that

$$F_{j',k'}^{j,k} = e^{(-\kappa(\tilde{y}_{j-1}+\tilde{y}_{j'})-\eta(\tilde{z}_{k-1}+\tilde{z}_{k'}))/2} D_{1,\delta}^2 D_{2,\delta}^2 F_{\alpha}(\tilde{y}_{j'-1}-\tilde{y}_j, \tilde{z}_{k'-1}-\tilde{z}_k) + \mathcal{O}(\nu^{\alpha} \Delta^{1+\alpha}) + \mathcal{O}\left(\nu^{\alpha-1/2} \Delta^{1/2+\alpha} \left(\frac{\mathbf{1}_{\{j\neq j'\}}}{|j-j'|+1} + \frac{\mathbf{1}_{\{k\neq k'\}}}{|k-k'|+1}\right)\right), \\D_{1,\delta}^2 D_{2,\delta}^2 F_{\alpha}(\tilde{y}_{j'-1}-\tilde{y}_j, \tilde{z}_{k'-1}-\tilde{z}_k) = \begin{cases} \nu^{\alpha-1} \Delta^{\alpha} \psi_{r,\alpha} + \mathcal{O}(\nu^{\alpha-1} \Delta^{1+\alpha}), & (j',k') = (j,k) \\ \mathcal{O}\left(\nu^{\alpha-1} \Delta^{\alpha} \left(\nu \Delta + \frac{1}{(|j-j'|+1)(|k-k'|+1)}\right)\right), & (j',k') \neq (j,k) \end{cases}$$

Furthermore, it follows that under [A2],

$$\begin{aligned} \operatorname{Cov}[T_{i,j,k}X, T_{i',j',k'}X] &= \mathcal{O}\left(\frac{\nu^{\alpha-1}\Delta^{\alpha}}{|i-i'|+1} \left(\nu\Delta + \frac{1}{(|j-j'|+1)(|k-k'|+1)}\right)\right) \\ &+ \mathcal{O}\left(\frac{\nu^{\alpha-1/2}\Delta^{1/2+\alpha}}{|i-i'|+1} \left(\frac{\mathbf{1}_{\{j\neq j'\}}}{|j-j'|+1} + \frac{\mathbf{1}_{\{k\neq k'\}}}{|k-k'|+1}\right)\right),\end{aligned}$$

$$\begin{split} \sum_{i,i'=1}^{N} \operatorname{Cov}[T_{i,j,k}X, T_{i',j,k}X]^2 &= \mathcal{O}(\nu^{2\alpha-2}N\Delta^{2\alpha}) \quad uniformly \ in \ j, k, \\ \sum_{i,i'=1}^{m_2} \sum_{j,j'=1}^{m_1} \sum_{i,i'=1}^{N} \operatorname{Cov}[T_{i,j,k}X, T_{i',j',k'}X]^2 &= \mathcal{O}(\nu^{2\alpha-2}m_1m_2N\Delta^{2\alpha}), \\ \sum_{i,i'=1}^{N} \operatorname{Cov}[(T_{i,j,k}X)^2, (T_{i',j,k}X)^2] &= \mathcal{O}(\nu^{2\alpha-2}N\Delta^{2\alpha}) \quad uniformly \ in \ j, k, \\ \sum_{k,k'=1}^{m_2} \sum_{j,j'=1}^{m_1} \sum_{i,i'=1}^{N} \operatorname{Cov}[(T_{i,j,k}X)^2, (T_{i',j',k'}X)^2] &= \mathcal{O}(\nu^{2\alpha-2}m_1m_2\sqrt{m_1m_2N}\Delta^{2\alpha}) \end{split}$$

Proposition 2.5 is immediately obtained from Lemma 12. By the Taylor expansion, we have

$$-NK'_{\nu,N,m}((\sigma^*)^2) = \int_0^1 K''_{\nu,N,m}((\sigma^*)^2 + u(\hat{\sigma}^2 - (\sigma^*)^2)) \mathrm{d}u N(\hat{\sigma}^2 - (\sigma^*)^2).$$

Since Proposition 2.5 and Lemma 12 imply that  $NK'_{\nu,N,m}((\sigma^*)^2) = \mathcal{O}_{\mathbf{P}}(1)$  and there exists V > 0 such that  $\int_0^1 K''_{\nu,N,m}((\sigma^*)^2 + u(\hat{\sigma}^2 - (\sigma^*)^2)) du \xrightarrow{\mathbf{P}} V$ , we get Proposition 2.6.

# References

- R. Altmeyer, T. Bretschneider, J. Janák and M. Reiß. (2022). Parameter Estimation in an SPDE Model for Cell Repolarization. SIAM/ASA Journal on Uncertainty Quantification, 10(1):179– 199.
- [2] R. Altmeyer, A. Tiepner and M. Wahl. (2022). Optimal parameter estimation for linear SPDEs from multiple measurements. arXiv preprint arXiv:2211.02496.
- [3] M. Bibinger and P. Bossert. (2023). Efficient parameter estimation for parabolic SPDEs based on a log-linear model for realized volatilities. *Japanese Journal of Statistics and Data Science*, 6:407–429.
- [4] M. Bibinger and M. Trabs. (2020). Volatility estimation for stochastic PDEs using highfrequency observations. Stochastic Processes and their Applications, 130(5):3005–3052.
- [5] P. Bossert. (2023). Parameter estimation for second-order SPDEs in multiple space dimensions. arXiv preprint arXiv:2310.17828.
- [6] C. Chong. (2020). High-frequency analysis of parabolic stochastic PDEs. The Annals of Statistics, 48(2):1143–1167.
- [7] I. Cialenco. (2018). Statistical inference for SPDEs: an overview. Statistical Inference for Stochastic Processes, 21(2):309–329.
- [8] I. Cialenco, F. Delgado-Vences and H.J. Kim. (2020). Drift estimation for discretely sampled SPDEs. Stochastics and Partial Differential Equations: Analysis and Computations, 8:895–920.
- [9] I. Cialenco and N. Glatt-Holtz. (2011). Parameter estimation for the stochastically perturbed Navier-Stokes equations. Stochastic Processes and their Applications, 121(4):701-724.
- [10] D.A. Dawson. (1975). Stochastic evolution equations and related measure processes. Journal of Multivariate Analysis, 5(1):1-52.

- [11] Z. Denkowski, S. Migórski and N. S. Papageorgiou. (2003). An introduction to nonlinear analysis: theory. Kluwer Academic/Plenum, New York.
- [12] S. Gaudlitz and M. Reiss. (2023). Estimation for the reaction term in semi-linear SPDEs under small diffusivity. *Bernoulli* 29(4):3033-3058.
- [13] A. Gloter and M. Sørensen. (2009). Estimation for stochastic differential equations with a small diffusion coefficient. *Stochastic Processes and their Applications*, 119:679–699.
- [14] M. Hairer. (2009). An introduction to Stochastic PDEs. arXiv preprint arXiv:0907.4178.
- [15] F. Hildebrandt and M. Trabs. (2021). Parameter estimation for SPDEs based on discrete observations in time and space. *Electronic Journal of Statistics*, 15(1):2716–2776.
- [16] F. Hildebrandt and M. Trabs. (2023). Nonparametric calibration for stochastic reactiondiffusion equations based on discrete observations, *Stochastic Processes and their Applications*, 162:171–217.
- [17] M. Hübner, R. Khasminskii and B.L. Rozovskii. (1993). Two Examples of Parameter Estimation for Stochastic Partial Differential Equations, 149–160. Springer New York.
- [18] M. Hübner and B.L. Rozovskii. (1995). On asymptotic properties of maximum likelihood estimators for parabolic stochastic PDE's. *Probability Theory and Related Fields*, 103(2):143– 163.
- [19] Y. Kaino and M. Uchida. (2021). Parametric estimation for a parabolic linear SPDE model based on discrete observations. *Journal of Statistical Planning and Inference*, 211:190–220.
- [20] Y. Kaino and M. Uchida. (2021). Adaptive estimator for a parabolic linear SPDE with a small noise. Japanese Journal of Statistics and Data Science, 4:513–541.
- [21] M. Kessler. (1997). Estimation of an ergodic diffusion from discrete observations. Scandinavian Journal of Statistics, 24(2):211–229.
- [22] S. Kusuoka. (2000). Term structure and SPDE, In Advances in Mathematical Economics, 2:67–85, Springer.
- [23] S.V. Lototsky. (2003). Parameter estimation for stochastic parabolic equations: asymptotic properties of a two-dimensional projection-based estimator. *Statistical Inference for Stochastic Processes*, 6:65–87.
- [24] B. Markussen. (2003). Likelihood inference for a discretely observed stochastic partial differential equation. *Bernoulli*, 9:745–762.
- [25] R. Moleriu. (2009). A generalization of the Girsanov theorems on the Hilbert space. The Romanian Journal of Technical Sciences. Applied Mechanics, 54(2), 113–124.
- [26] E. Di Nezza, G. Palatucci and E. Valdinoci. (2012). Hitchhiker's guide to the fractional Sobolev spaces. Bulletin des Sciences Mathématiques 136:521-573.
- [27] I. Nourdin, G. Peccati and G. Reinert. (2009). Second order Poincaré inequalities and CLTs on Wiener space. *Journal of Functional Analysis*, 257:593-603.
- [28] D. Nualart. (2006). The Malliavin Calculus and Related Topics. Springer, 2nd edition.
- [29] L. Piterbarg and A. Ostrovskii. (1997). Advection and diffusion in random media: implications for sea surface temperature anomalies. Springer Science & Business Media.

- [30] G. Da Prato and J. Zabczyk. (2014). Stochastic Equations in Infinite Dimensions, 2nd edition. Encyclopedia of Mathematics and its Applications. Cambridge University Press.
- [31] M. Sanz-Soló. (2005). Malliavin calculus with applications to stochastic partial differential equations. Fundamental sciences. Mathematics. EPFL Press, Lausanne.
- [32] M. Sørensen and M. Uchida. (2003). Small-diffusion asymptotics for discretely sampled stochastic differential equations. *Bernoulli*, 9(6):1051–1069.
- [33] Y. Tonaki, Y. Kaino and M. Uchida. (2023). Parameter estimation for linear parabolic SPDEs in two space dimensions based on high frequency data. *Scandinavian Journal of Statistics*, 50(4):1568-1589.
- [34] Y. Tonaki, Y. Kaino and M. Uchida. (2023). Parametric estimation for linear parabolic SPDEs in two space dimensions based on temporal and spatial increments. arXiv preprint arXiv:2304.09441.
- [35] Y. Tonaki, Y. Kaino and M. Uchida. (2024). Parameter estimation for a linear parabolic SPDE model in two space dimensions with a small noise. *Statistical Inference for Stochastic Processes*, 27(1):123-179.
- [36] H.C. Tuckwell. (2013). Stochastic partial differential equations in neurobiology: linear and nonlinear models for spiking neurons, In *Stochastic biomathematical models*, 149–173. Springer.
- [37] M. Uchida and N. Yoshida. (2012). Adaptive estimation of an ergodic diffusion process based on sampled data. *Stochastic Processes and their Applications*, 122(8):2885–2924.
- [38] J.B. Walsh. (1986). An introduction to stochastic partial differential equations. In Ecole d'Eté de Probabilités de Saint Flour XIV-1984, 265–439, Springer.
- [39] A. Yagi. (2010). Abstract parabolic evolution equations and their applications. Springer monographs in mathematics. Springer.
- [40] K. Yoshida. (1980). Functional analysis, 6th edition. Springer, Berlin.
- [41] N. Yoshida. (2011). Polynomial type large deviation inequalities and quasi-likelihood analysis for stochastic differential equations. Annals of the Institute of Statistical Mathematics, 63:431– 479.