

Conditions on separability in multiqubit systems with an accelerating qubit using a conditional entropy

Harsha Miriam Reji, Hemant Shreepad Hegde, and R. Prabhu

Department of Physics, Indian Institute of Technology Dharwad, Dharwad, Karnataka, India - 580007

The separability in multiqubit pure and mixed Greenberger–Horne–Zeilinger (GHZ) and W states with an accelerating qubit has been characterized using Abe Rajagopal (AR) q -conditional entropy. We observe that the pure multiqubit GHZ and W states in the inertial : non-inertial bipartition with one of their qubits being accelerated will remain non-separable irrespective of the qubit's acceleration. In these systems, we effectively captured the variation of their non-separability with respect to the acceleration of the qubit and the AR q -conditional entropy parameter q . However, in the corresponding multiqubit mixed states obtained by introducing a noise to the above pure states, one could get stronger conditions on their separability in the inertial : non-inertial bipartition, in terms of the mixing parameter, acceleration of the qubit and the number of qubits in the system, in the asymptotic limit of parameter q . These conditions obtained from AR q -conditional entropy serves as a necessary conditions for separability in such multiqubit states with a relativistic qubit.

I. INTRODUCTION

A vast majority of quantum information tasks, like quantum teleportation [1–3], quantum dense coding [4], quantum key distribution, [5, 6] etc., consider certain essential properties present in quantum systems, such as quantum correlations, to be generally invariant during the realization of such quantum information protocols [7, 8]. However, it has been established that such properties exhibited by the quantum systems does not remain invariant when they or their subsystems are relativistically accelerated [9]. The studies involving the characterization of quantum properties in relativistically accelerated quantum system has lead to the field relativistic quantum information, which is a fusion of two previously disjoint fields, general relativity and quantum information. The primary objective of this scientific domain is to integrate relativistic effects into the quantum information tasks and study their performance. At present, under the umbrella of relativistic quantum information, studies involving characterization and quantification of entanglement in bosonic [9–14] and fermionic [15–24] modes when their subsystems are in non-inertial frames, entanglement near black holes[25–30], entanglement in expanding universe[31–35], relativistic quantum metrology[36–39], discord in relativistic states[40, 41], relativistic quantum teleportation [42–44], relativistic quantum speed limit [45–47], etc., have been explored.

In the inertial framework, the negativity of the von Neumann conditional entropy of a pure bipartite entangled state reveal a significant observation that the entangled systems tend to exhibit more local disorder than the global disorder, a distinction not present in separable states [48]. Consequently, there has been considerable interest in utilizing entropic measures to characterize separability in composite quantum systems [49–54]. To better understand and explore the valuable quantum information properties of mixed states, generalized entropic measures have been introduced [55–59]. The two prominent families of generalized entropies are Tsallis q -

entropy [60, 61] and α -Renyi entropy [62], which reduces to the von Neumann entropy in the limit as q approaches to 1. Their positivity provides a means to discern between global and local disorder in mixed states, thereby imposing stricter constraints the on separability compared to those derived from the von Neumann conditional entropy alone. However, the quantum systems with long range interactions, like entanglement, will not follow psuedo-additivity relation of Tsallis entropy. Therefore, a generalized form of conditional entropy, called the Abe-Rajagopal (AR) q -conditional entropy, derived from conditioning the Tsallis entropy was introduced [63, 64]. The AR q -conditional entropy assumes a negative value for non-separable states, suggesting its importance in the characterization of non-separability in mixed states. This conditional entropy has been applied to study the separability in single parameter family of mixed multiqubit states, Gaussian states, etc [65, 66]. Consequently, we employ this measure of conditional entropy to characterize the non-separability between two bipartitions of several multipartite quantum states with one of their bipartition being uniformly accelerated.

We initially consider a generalized pure 2-qubit Greenberger–Horne–Zeilinger (GHZ) state with one of its qubits under acceleration and characterize its non-separability in the inertial : non-inertial bipartition with respect system parameters, acceleration of the qubit, and parameter q . Since, the characterization of the non-separability using AR q -conditional entropy uses all the eigenvalues of the state and corresponding subsystem, we introduce a *eigenvalue truncation procedure* to numerically handle the infinite eigenvalues in such systems. Subsequently, we extend our non-separability characterization to the multiqubit pure GHZ and W states. Later, we explore the non-separability in mixed states generated by mixing a global noise to the above multiqubit pure GHZ and multiqubit pure W states. Since, these accelerated states do not posses a block structure to analytically obtain their eigenvalues, we introduce a *density matrix truncation procedure* to numerically evaluate the




FIG. 1: (Color online) The schematic representation of Rindler spacetime [42], where t is the proper time and $x = \text{constant}$ is the trajectory of a qubit with constant acceleration. The pink shaded region on the right of the vertical axis denote the region I and the green shaded region to the left of vertical axis denote the region II of Rindler spacetime.

AR q -conditional entropy. We derive conditions for separability in these mixed states which depends on their mixing parameter, acceleration of a qubit and number of qubits in the multiqubit state in the asymptotic limit of parameter q .

This paper is structured as follows: In Sec. II, we describe the bosonic quantum system in a non-inertial frame. A bosonic qubit, on acceleration, transforms as a two mode squeezed vacuum state, with the acceleration being related to its squeezing parameter. In Sec. III, we succinctly describe the quantification of non-separability using AR q -conditional entropy. In Sec. IV, the non-separability of multiqubit pure GHZ and W states with one of its qubits being accelerated using AR q -conditional entropy is described. In Sec. V, we give a detailed account of obtaining the non-separability of multiqubit mixed states obtained by adding a global noise to the pure states considered in Sec. IV. The strongest condition on separability for these states is identified in the asymptotic limit of q . In Sec. VI, the concluding remarks are presented.

II. RELATIVISTICALLY ACCELERATED QUANTUM STATE

Consider an arbitrary 2-qubit state ρ_{AB} in an inertial frame, shared between the two observers Alice and Bob. When Bob undergoes a relativistic acceleration, henceforth termed as Rob (relativistically accelerated Bob), his

qubit experiences a different spacetime around it. While Minkowski coordinates are commonly employed to depict the qubits in the inertial frames, they are unsuitable for describing the motion of an accelerated qubit. In non-inertial frames, Rindler coordinates prove to be more appropriate for describing the accelerating qubits. The schematic representation of the Rindler spacetime is provided in Fig. 1. The Rindler spacetime is divided into two distinct regions, region I (pink shade) and region II (green shade), which collectively contain all points mapped from the Minkowski coordinates. Using Bogoliubov transformations, the vacuum state and the single excited state in Minkowski coordinates will individually take the form of two mode squeezed state in Rindler spacetime [9, 42], i.e.,

$$|0_k\rangle_M = \frac{1}{\cosh r} \sum_{n=0}^{\infty} \tanh^n r |n_k\rangle_I \otimes |n_k\rangle_{II}, \quad (1)$$

and

$$|1_k\rangle_M = \frac{1}{\cosh^2 r} \sum_{n=0}^{\infty} \tanh^n r \sqrt{n+1} |(n+1)_k\rangle_I \otimes |n_k\rangle_{II} \quad (2)$$

respectively. Here $|n_k\rangle_I$ and $|n_k\rangle_{II}$ are the number states of k^{th} mode in the region I and region II of the Rindler spacetime respectively. The RHS of Eqs. (1) and (2) corresponds to a two mode squeezed states, with the squeezing parameter r is related to the acceleration a of the Rob's qubit as $\cosh r = (1 - e^{-2\pi|k|c/a})^{-1/2}$. Here, c is the speed of light in vacuum.

A composite state shared between Alice and Bob in the inertial frame ρ_{AB} transforms as $\rho_{AR_I R_{II}}$ when Bob's qubit is accelerated. However, the qubit moving in one of the regions of the Rindler spacetime cannot travel into the other region as they are causally disconnected. Without any loss of generality, one can trace out the modes of the region II of the accelerating qubit. Since a part of the accelerating state has been traced out, the final state between Alice and Rob's modes in region I is a *mixed* state and is in the basis $\{|0\rangle_M |n\rangle_I, |0\rangle_M |n+1\rangle_I, |1\rangle_M |n\rangle_I, |1\rangle_M |n+1\rangle_I\}$ with n going from zero to ∞ . Henceforth, we will drop all the subscripts M and I in the inertial basis, $\{|0\rangle, |1\rangle\}$, as well as accelerating qubit's basis, $\{|n\rangle, |n+1\rangle\}$, and denote the resultant mixed state of Alice and Rob's mode in region I by ρ_{AR} and Rob's subsystem as ρ_R .

III. ABE-RAJAGOPAL q -CONDITIONAL ENTROPY

A non-extensive generalization of Shannon entropy is given by the Tsallis q -entropy [60, 61], which is always positive and adheres to the non-additivity relationship for bipartite state ρ_{XY} , i.e., $S_q^T(\rho_{XY}) = S_q^T(\rho_X) + S_q^T(\rho_Y) + (1 - q) S_q^T(\rho_X) S_q^T(\rho_Y)$, where $S_q^T(\rho)$ is given

by

$$S_q^T(\rho) = \frac{\text{Tr}[\rho^q] - 1}{1 - q}. \quad (3)$$

Here Tr denotes the trace of the corresponding density matrix. However, this non-additivity condition is violated by the state ρ_{XY} when subsystems X and Y exhibit long-range interactions, such as strong entanglement, which persists even when they are spatially separated. To address this, a conditional form of Tsallis entropy was introduced by Abe et. al. [63]. This Abe-Rajagopal q -conditional entropy is defined as

$$S_q(X|Y) = \frac{S_q^T(\rho_{XY}) - S_q^T(\rho_Y)}{1 + (1 - q)S_q^T(\rho_Y)}, \quad (4)$$

where $S_q^T(\rho_{XY})$ and $S_q^T(\rho_Y)$ denote the Tsallis q -entropies associated with the composite system ρ_{XY} and its subsystem ρ_Y . The computable form of this conditional entropy for any bipartition $X : Y$ in a multiqubit state can be obtained by plugging in the Eq. (3) in Eq. (4) and is given by

$$\begin{aligned} S_q(X|Y) &= \frac{1}{q-1} \left[1 - \frac{\text{Tr}[\rho_{XY}^q]}{\text{Tr}[\rho_Y^q]} \right], \\ &= \frac{1}{q-1} \left[1 - \frac{\sum_n \lambda_n^q(\rho_{XY})}{\sum_m \lambda_m^q(\rho_Y)} \right], \end{aligned} \quad (5)$$

where λ_n and λ_m are the eigenvalues of the whole system ρ_{XY} and one of its subsystem ρ_Y respectively. This conditional entropy adheres to the non-additivity relation: $S_q^T(\rho_{XY}) = S_q^T(X|Y) + S_q^T(Y) + (1 - q)S_q^T(X|Y)S_q^T(\rho_Y)$. The AR q -conditional entropy yields negative values for the non-separable states and hence the positivity of AR q -conditional entropy forms a criterion for capturing the separability in bipartite states. From here onwards, we represent $S_q(X|Y)$ and $S_q(X|Y)$ as the AR q -conditional entropy for pure ρ_{XY} and mixed ρ_{XY} states with one of their qubit being accelerated respectively.

IV. MULTIQUBIT PURE STATES WITH AN ACCELERATING QUBIT

In this section, we consider pure multiqubit GHZ and W states with an accelerated qubit and characterize their non-separability in the inertial : non-inertial bipartition using AR q -conditional entropy given in Eq. (5) with respect to their system parameters, parameter q and acceleration of the qubit in the non-inertial frame.

A. Pure multiqubit GHZ states

Generalized pure 2-qubit GHZ state: A generalized pure 2-qubit state $|\text{GHZ}_2\rangle$ shared between Alice and Bob

in the inertial frame is given by

$$|\text{GHZ}_2\rangle = \cos\theta |00\rangle + \sin\theta |11\rangle; \quad \cos^2\theta + \sin^2\theta = 1, \quad (6)$$

which belongs to the Hilbert space $H_A \otimes H_B$. Let Bob's qubit be relativistically accelerated with respect to Alice's qubit and the resulting composite state, $\rho_{AR}^{\text{GHZ}_2}$, is obtained by using the transformations of $|0\rangle$ and $|1\rangle$ from Minkowski spacetime to Rindler spacetime as given in Eqs. (1) and (2). We now obtain the marginal mixed density matrix between Alice and Rob's modes in region I, by following the recipe given in Sec. II, as

$$\begin{aligned} \rho_{AR}^{\text{GHZ}_2} &= \frac{1}{\cosh^2 r} \sum_n \tanh^{2n} r \\ &\times \left[\cos^2\theta |0n\rangle \langle 0n| + \frac{n+1}{\cosh^2 r} \sin^2\theta |1n+1\rangle \langle 1n+1| \right. \\ &\left. + \frac{\sqrt{n+1}}{\cosh r} \cos\theta \sin\theta (|1n+1\rangle \langle 0n| + |0n\rangle \langle 1n+1|) \right]. \end{aligned} \quad (7)$$

The structure of this density matrix contains 2×2 blocks along the diagonal, with the remaining off diagonal elements being zero and is given by

$$\rho_{AR}^{\text{GHZ}_2} = \frac{1}{\cosh^2 r} \begin{pmatrix} 0 & & & & \\ & \Delta_0 & & & \\ & & \Delta_1 & & \\ & & & \ddots & \\ & & & & \Delta_n \\ & & & & & \ddots \end{pmatrix}, \quad (8)$$

where

$$\Delta_n(\rho_{AR}^{\text{GHZ}_2}) = \tanh^{2n} r \begin{pmatrix} \cos^2\theta & \alpha_n \cos\theta \sin\theta \\ \alpha_n \cos\theta \sin\theta & \alpha_n^2 \sin^2\theta \end{pmatrix} \quad (9)$$

with $\alpha_n = \frac{\sqrt{n+1}}{\cosh r}$. The eigenvalues of the n^{th} block of the state $\rho_{AR}^{\text{GHZ}_2}$ are 0 and

$$\Lambda_n = \frac{\tanh^{2n} r}{\cosh^2 r} \left(\cos^2\theta + \frac{n+1}{\cosh^2 r} \sin^2\theta \right), \quad (10)$$

such that $\text{Tr}(\rho_{AR}^{\text{GHZ}_2}) = \sum_{n=0}^{\infty} \Lambda_n = 1$.

The marginal density matrix $\rho_R^{\text{GHZ}_2}$ corresponding to the modes of Rob in region I of Rindler spacetime is computed by tracing out the Alice's subsystem A , i.e., $\rho_R^{\text{GHZ}_2} = \text{Tr}_A(\rho_{AR}^{\text{GHZ}_2})$ and is given by

$$\begin{aligned} \rho_R^{\text{GHZ}_2} &= \frac{1}{\cosh^2 r} \sum_{m=0}^{\infty} \tanh^{2m} r \left[\cos^2\theta |m\rangle \langle m| \right. \\ &\left. + \frac{m+1}{\cosh^2 r} \sin^2\theta |m+1\rangle \langle m+1| \right]. \end{aligned} \quad (11)$$

This is an infinite dimensional diagonal matrix whose m^{th} eigenvalue is given by

$$\Lambda_m = \frac{\tanh^{2m} r}{\cosh^2 r} \left(\cos^2\theta + \frac{m}{\sinh^2 r} \sin^2\theta \right), \quad (12)$$




FIG. 2: (Color online) (a) The variation of Abe-Rajagopal q -conditional entropy as a non-separability feature of a single qubit accelerated generalized pure 2-qubit GHZ state as a function of $\cos^2 \theta$ and λ for parameter $q = 2$ and 4. The upper blue solid mesh corresponds to $q = 2$ and the lower red solid mesh corresponds to $q = 4$. The $S_q^{\text{GHZ}_2}(A|R)$ is negative everywhere, implying that the state is non-separable. For this plot, we have chosen 100 values each of $\cos^2 \theta$ and λ to create the surface and the meshes are drawn for representation purpose. (b) The variation of AR q -conditional entropy of pure 2-qubit GHZ state with $\theta = \pi/4$ as a function of λ for different values of the parameter q is depicted in this figure.

where $\text{Tr}(\rho_R^{\text{GHZ}_2}) = \sum_{m=0}^{\infty} \Lambda_m = 1$.

To characterize the non-separability of the state $\rho_{AR}^{\text{GHZ}_2}$ given in Eq. (7), we substitute the eigenvalues Λ_n 's of $\rho_{AR}^{\text{GHZ}_2}$ (see Eq. (10)) and Λ_m 's of $\rho_R^{\text{GHZ}_2}$ (see Eq. (12)) in AR q -conditional entropy given in Eq. (5) to obtain

$$S_q^{\text{GHZ}_2}(A|R) = \frac{1}{q-1} \times \left[1 - \frac{\sum_{n=0}^{\infty} \left(\tanh^{2n} r \left(\cos^2 \theta + \sin^2 \theta \frac{n+1}{\cosh^2 r} \right) \right)^q}{\sum_{m=0}^{\infty} \left(\tanh^{2m} r \left(\cos^2 \theta + \sin^2 \theta \frac{m}{\sinh^2 r} \right) \right)^q} \right]. \quad (13)$$

The negative values of this AR q -conditional entropy $S_q^{\text{GHZ}_2}(A|R)$ characterize the non-separability between Alice's qubit and Rob's mode in region I of the Rindler spacetime. As the above expression contains summations running upto infinity, we have to evaluate $S_q^{\text{GHZ}_2}(A|R)$ numerically to study its behaviors with respect to state parameter, acceleration parameter r and parameter q .

Eigenvalue truncation procedure: To numerically compute this conditional entropy, we make use of the fact that the sum of the non-zero eigenvalues of $\rho_{AR}^{\text{GHZ}_2}$ and $\rho_R^{\text{GHZ}_2}$ tends to 1 as their respective number of eigenval-

ues n and m approaches infinity. We achieve this numerically by fixing a certain large value of $n \simeq k$ and $m \simeq k$ such that the sum of k eigenvalues approaches to 1. By inspection, we found that this is satisfied for both the states $\rho_{AR}^{\text{GHZ}_2}$ and $\rho_R^{\text{GHZ}_2}$ when $k \approx 10^4$. Now, using these k number of eigenvalues of $\rho_{AR}^{\text{GHZ}_2}$ and $\rho_R^{\text{GHZ}_2}$, we numerically compute the AR q -conditional entropy given in Eq. (13) and characterize the non-separability of the state in Eq. (7) with respect to the parameter q , the state parameter θ , and Rob's acceleration r .

We now introduce the modified acceleration parameter $\lambda = \tanh r$, which carries all the characteristics of Rob's acceleration r , i.e., we use $\lambda \in (0, 1)$ instead of $r \in (0, \infty)$. For simplicity, now onwards we call λ as the Rob's acceleration. The AR q -conditional entropy for the state given in Eq. (7), is then plotted as a function of its state parameter θ and acceleration of Rob λ in Fig. 2a for $q = 2$ (upper blue surface) and 4 (lower red surface). We observe that the $S_q^{\text{GHZ}_2}(A|R)$ always remains less than zero, implying that the state remains non-separable for any values of λ and θ . Note that, $S_q^{\text{GHZ}_2}(A|R)$ attains maximum non-separability at $\theta = \pi/4$, irrespective of the parameter q . When $\theta = \pi/4$, Eq. (6) corresponds to a maximally entangled 2-qubit Bell state in an inertial frame. Hence, in Fig. 2b, we plot the characterization of $S_q^{\text{GHZ}_2}(A|R)$ for the state $\rho_{AR}^{\text{GHZ}_2}$, given in Eq. (7), with the initial state chosen as the Bell state, with respect to

λ for various values of q . We observe that irrespective of the value of q , the non-separability of $\rho_{AR}^{\text{GHZ}_2}$ will be high at lower values of acceleration and decreases with the increase in acceleration, in particular, $S_q^{\text{GHZ}_2}(A|R)$ tends to 0 as λ approaches to 1. As the value of q increases, the AR q -conditional entropy suggests that the state $\rho_{AR}^{\text{GHZ}_2}$ may transition towards separability in the limit λ approaches to 1. However, $S_q^{\text{GHZ}_2}(A|R)$ approaching to 0 becomes more and more steeper at large values of both acceleration λ and parameter q .

Generalized pure 3-qubit GHZ state: Let us consider a generalized pure 3-qubit GHZ state given by

$$|\text{GHZ}_3\rangle = \cos\theta|000\rangle + \sin\theta|111\rangle; \quad \cos^2\theta + \sin^2\theta = 1, \quad (14)$$

shared between Alice and Bob, with first two of its qubits are in Alice's possession while the remaining qubit is in Bob's possession, i.e., they belong to the $H_{A_1} \otimes H_{A_2} \otimes H_B$ Hilbert space. By accelerating Bob's qubit, we study the non-separability between inertial Alice's qubits and modes of Rob in Rindler spacetime by exploring AR q -conditional entropy. Using the recipe given in Sec. II, we get the composite state of the inertial A_1A_2 and non-inertial R as $\rho_{A_1A_2R_I R_{II}}^{\text{GHZ}_3}$, and the state $\rho_{A_1A_2R}^{\text{GHZ}_3}$ as the mixed subsystem after tracing out the mode of Rob's qubit in region II. The density matrix of $\rho_{A_1A_2R}^{\text{GHZ}_3}$ is given by

$$\begin{aligned} \rho_{A_1A_2R}^{\text{GHZ}_3} &= \frac{1}{\cosh^2 r} \sum_{n=0}^{\infty} \tanh^{2n} r \left[\cos^2\theta |00n\rangle \langle 00n| \right. \\ &+ \frac{\sqrt{n+1}}{\cosh r} \cos\theta \sin\theta \left(|00n\rangle \langle 11n+1| + |11n+1\rangle \langle 00n| \right) \\ &\left. + \frac{n+1}{\cosh^2 r} \sin^2\theta |11n+1\rangle \langle 11n+1| \right]. \end{aligned} \quad (15)$$

Generally, there are two independent q -conditional entropies associated with 3-qubit states [56], which are given by

$$S_q(AB|C) = \frac{S_q^T(\rho_{ABC}) - S_q^T(\rho_C)}{1 + (1-q)S_q^T(\rho_C)} \quad (16)$$

and

$$S_q(A|BC) = \frac{S_q^T(\rho_{ABC}) - S_q^T(\rho_{BC})}{1 + (1-q)S_q^T(\rho_{BC})}, \quad (17)$$

where $S_q^T(\rho_i)$ is the Tsallis entropy of any state ρ_i . Since one of the qubits in the shared initial state is accelerating in a non-inertial frame, a simultaneous measurement cannot be performed on the inertial and the non-inertial qubits together. Hence, we choose the conditional entropy such that the inertial Alice's qubits A_1, A_2 are conditioned on the non-inertial Rob's qubit R . Therefore, we proceed to investigate the separability using AR q -conditional entropy $S_q^{\text{GHZ}_3}(A_1A_2|R)$ in the $A_1A_2 : R$ bipartition for the state $\rho_{A_1A_2R}^{\text{GHZ}_3}$ given in Eq. (15).

To get the eigenvalues of the states $\rho_{A_1A_2R}^{\text{GHZ}_3}$ and $\rho_R^{\text{GHZ}_3}$ to evaluate $S_q^{\text{GHZ}_3}(A_1A_2|R)$, we make use of the fact that




FIG. 3: (Color online) The variation of AR q -conditional entropy for pure 3-qubit W state as a function of acceleration parameter λ for various values of q . We have chosen $\theta = \cos^{-1}(1/\sqrt{3})$ and $\phi = \pi/4$ for this plot.

their density matrices have the same non-zero eigenvalues as that of $\rho_{AR}^{\text{GHZ}_2}$ and $\rho_R^{\text{GHZ}_2}$ respectively. Hence, the behavior of $S_q^{\text{GHZ}_3}(A_1A_2|R)$ with respect to state parameter θ , acceleration λ and q is exactly same as the behavior of AR q -conditional entropy for pure 2-qubit GHZ state with one of its qubit in acceleration $S_q^{\text{GHZ}_2}(A|R)$ as given in Fig. 2a and 2b. For logistic reasons which will be clear later, we now study the non-separability of pure N -qubit GHZ state instead of the generalized N -qubit GHZ state.

Pure N -qubit GHZ state: Here, we generalize the characterization of non-separability of the pure GHZ states to N -qubit GHZ state when one its qubits is accelerated using AR q -conditional entropy. Then, the pure N -qubit GHZ state is given by

$$|\text{GHZ}_N\rangle = \frac{|0\rangle^{\otimes N} + |1\rangle^{\otimes N}}{\sqrt{2}}, \quad (18)$$

belonging to Hilbert space $H_{A_1} \otimes H_{A_2} \otimes \dots \otimes H_{A_{N-1}} \otimes H_B$, i.e., the first $N-1$ qubits are with Alice and the last qubit is in Bob's possession. The choice of qubit, among pure N -qubit GHZ states, to be accelerated does not change the study of non-separability among the inertial and non-inertial qubits as the density matrix structure of the composite state and its subsystem remain the same. When Bob accelerates, the resulting composite state of inertial Alice and non-inertial Rob, denoted as $\rho_{A_1A_2\dots A_{N-1}R_I R_{II}}^{\text{GHZ}_N}$, is obtained by using the transformations from Minkowski to Rindler spacetime given in Eqs. (1) and (2). Then, $\rho_{A_1A_2\dots A_{N-1}R}^{\text{GHZ}_N}$ after tracing out one of the disjoint regions in Rindler spacetime (region




FIG. 4: (Color online) The variation of AR q -conditional entropy of pure N -qubit W state as a function of the acceleration λ for different values of N given in this figure. We have taken $q = 2$ for this plot.

II) is given by

$$\begin{aligned} \rho_{A_1 A_2 \dots A_{N-1} R}^{\text{GHZ}_N} &= \frac{1}{2 \cosh^2 r} \sum_{n=0}^{\infty} \tanh^{2n} r \\ &\times \left[|00 \dots 0n\rangle \langle 00 \dots 0n| \right. \\ &+ \frac{\sqrt{n+1}}{\cosh r} \left(|00 \dots 0n\rangle \langle 11 \dots 1n+1| \right. \\ &+ \left. |11 \dots 1n+1\rangle \langle 00 \dots 0n| \right) \\ &\left. + \frac{n+1}{\cosh^2 r} |11 \dots 1n+1\rangle \langle 11 \dots 1n+1| \right]. \end{aligned} \quad (19)$$

The Rob's subsystem $\rho_R^{\text{GHZ}_N}$ is equivalent to Eq. (11) with $\theta = \pi/4$. These two density matrices have the same non-zero eigenvalues as that of $\rho_{AR}^{\text{GHZ}_2}$ and $\rho_R^{\text{GHZ}_2}$ given in Eqs. (10) and (12) with $\theta = \pi/4$. Therefore, the behavior of the AR q -conditional entropy remains the same with respect to λ for various values of q as seen in Fig. 2b. Evidently, increasing the number of qubits in the inertial frame does not affect the behavior of non-separability of pure N -qubit GHZ state, when any one of its qubit is accelerated.

Hence, we can conclude that the pure multiqubit GHZ state with one of its qubits under acceleration always remain non-separable throughout the range of λ and choices of q . Importantly, the states are more non-separable when the acceleration is low and the non-separability reduces as the acceleration increases, irrespective of the value of the parameter q .

B. Pure multiqubit W states

Generalized pure 2-qubit W state: A generalized pure 2-qubit state $|W_2\rangle$ shared between Alice and Bob

in the inertial frame is given by

$$|W_2\rangle = \cos \theta |01\rangle + \sin \theta |10\rangle; \quad \cos^2 \theta + \sin^2 \theta = 1. \quad (20)$$

Let Bob's qubit be relativistically accelerated with respect to Alice's qubit and the resulting composite state is $\rho_{AR}^{W_2}$. Then, the mixed density matrix $\rho_{AR}^{W_2}$, by following the recipe given in Sec. II, is given by

$$\begin{aligned} \rho_{AR}^{W_2} &= \frac{1}{\cosh^2 r} \sum_n \tanh^{2n} r \\ &\times \left[\frac{n+1}{\cosh^2 r} \cos^2 \theta |0n+1\rangle \langle 0n+1| + \sin^2 \theta |1n\rangle \langle 1n| \right. \\ &\left. + \frac{\sqrt{n+1}}{\cosh r} \cos \theta \sin \theta \left(|0n+1\rangle \langle 1n| + |1n\rangle \langle 0n+1| \right) \right]. \end{aligned} \quad (21)$$

The structure of this state $\rho_{AR}^{W_2}$ is made up of 2×2 blocks along the diagonal, with the remaining off diagonal elements being zero. This structure is same as Eq. (8), however with the non-zero blocks given by

$$\Delta_n(\rho_{AR}^{W_2}) = \tanh^{2n} r \begin{pmatrix} \alpha_n^2 \cos^2 \theta & \alpha_n \cos \theta \sin \theta \\ \alpha_n \cos \theta \sin \theta & \sin^2 \theta \end{pmatrix}. \quad (22)$$

Here, $\alpha_n = \frac{\sqrt{n+1}}{\cosh r}$. The eigenvalues of this n^{th} block of the state $\rho_{AR}^{W_2}$ are 0 and

$$\Lambda'_n = \frac{\tanh^{2n} r}{\cosh^2 r} \left(\sin^2 \theta + \frac{n+1}{\cosh^2 r} \cos^2 \theta \right), \quad (23)$$

satisfying $\text{Tr}(\rho_{AR}^{W_2}) = \sum_{n=0}^{\infty} \Lambda'_n = 1$.

The marginal density matrix $\rho_R^{W_2}$ is computed by tracing out the other subsystem A and is given by

$$\begin{aligned} \rho_R^{W_2} &= \frac{1}{\cosh^2 r} \sum_{m=0}^{\infty} \tanh^{2m} r \left[\sin^2 \theta |m\rangle \langle m| \right. \\ &\left. + \frac{m+1}{\cosh^2 r} \cos^2 \theta |m+1\rangle \langle m+1| \right]. \end{aligned} \quad (24)$$

This is an infinite dimensional diagonal matrix whose m^{th} eigenvalue is given by

$$\Lambda'_m = \frac{\tanh^{2m} r}{\cosh^2 r} \left(\sin^2 \theta + \frac{m}{\sinh^2 r} \cos^2 \theta \right), \quad (25)$$

where $\text{Tr}(\rho_R^{W_2}) = \sum_{m=0}^{\infty} \Lambda'_m = 1$.

To characterize the non-separability of the state of inertial Alice and non-inertial Rob given in Eq. (21), we substitute the eigenvalues Λ'_n 's of $\rho_{AR}^{W_2}$ and Λ'_m 's of $\rho_R^{W_2}$

in AR q -conditional entropy given in Eq. (5) to obtain

$$S_q^{W_2}(A|R) = \frac{1}{q-1} \times \left[1 - \frac{\sum_{n=0}^{\infty} \left(\tanh^{2n} r \left(\sin^2 \theta + \cos^2 \theta \frac{n+1}{\cosh^2 r} \right) \right)^q}{\sum_{m=0}^{\infty} \left(\tanh^{2m} r \left(\sin^2 \theta + \cos^2 \theta \frac{m}{\sinh^2 r} \right) \right)^q} \right]. \quad (26)$$

Using the eigenvalue truncation procedure introduced earlier, we chose a sufficiently large $k \simeq 10^4$, and numerically compute the AR q -conditional entropy given in Eq. (26) and characterize the same with respect to the parameter q , the state parameter θ , and Rob's acceleration λ . The $S_q^{W_2}(A|R)$ always remains less than zero, implying the state remains non-separable for any values of λ , θ and q . We observe that irrespective of the value of q , the non-separability of $\rho_{AR}^{W_2}$ will be high at lower values of acceleration and decreases with the increase in acceleration. When $\theta = \pi/4$, Eq. (20) corresponds to a 2-qubit Bell state in an inertial frame and the corresponding $S_q^{W_2}(A|R)$ attains maximum non-separability irrespective of the parameter q . These observations are similar to that of the 2-qubit GHZ state given in Eq. (7), except the variation with respect to $\cos^2 \theta$ in Fig. 2a is shifted in case of $\rho_{AR}^{W_2}$. Evidently, the characterization of $S_q^{W_2}(A|R)$ for $\rho_{AR}^{W_2}$ in Eq. (21) with the initial state being the Bell state, with respect to λ for various values of q is same as given in Fig. 2b.

Generalized pure 3-qubit W state: A generalized pure 3-qubit W state, shared between Alice and Bob, is given by

$$|W_3\rangle = \sin \theta \cos \phi |001\rangle + \sin \theta \sin \phi |010\rangle + \cos \theta |100\rangle. \quad (27)$$

Let $x = \sin \theta \cos \phi$, $y = \sin \theta \sin \phi$ and $z = \cos \theta$ giving $x^2 + y^2 + z^2 = 1$, which normalizes the given state for any θ and ϕ . Consider a scenario where the first two qubits belong to Alice and Bob holds the third qubit. Subsequently, as Bob undergoes uniform acceleration (Rob) relative to Alice in an inertial frame, it gives rise to the composite state $\rho_{A_1 A_2 R_I}^{W_3}$. The final mixed state after tracing out the Rob's mode in region II is expressed as

$$\rho_{A_1 A_2 R}^{W_3} = \frac{1}{\cosh^2 r} \sum_{n=0}^{\infty} \tanh^{2n} r \left[\alpha_n^2 x^2 |00n+1\rangle \langle 00n+1| + xz\alpha_n (|00n+1\rangle \langle 10n| + |10n\rangle \langle 00n+1|) + xy\alpha_n (|00n+1\rangle \langle 01n| + |01n\rangle \langle 00n+1|) + z^2 |10n\rangle \langle 10n| + y^2 |01n\rangle \langle 01n| + yz (|10n\rangle \langle 01n| + |01n\rangle \langle 10n|) \right]. \quad (28)$$

The density matrix of this state $\rho_{A_1 A_2 R}^{W_3}$ assumes a block structure along the diagonal as given in Eq. (8) and the remaining off diagonal elements are zero. However, here,

the diagonal blocks are given by

$$\Delta_n(\rho_{A_1 A_2 R}^{W_3}) = \tanh^{2n} r \begin{pmatrix} y^2 & yz & 0 & xy\alpha_n \\ yz & z^2 & 0 & xz\alpha_n \\ 0 & 0 & 0 & 0 \\ xy\alpha_n & xz\alpha_n & 0 & x^2\alpha_n^2 \end{pmatrix}. \quad (29)$$

The non-zero eigenvalues of this n^{th} block of the state $\rho_{A_1 A_2 R}^{W_3}$ is given by

$$\delta_n = \frac{\tanh^{2n} r}{\cosh^2 r} \left(1 + \frac{n+1 - \cosh^2 r}{\cosh^2 r} x^2 \right). \quad (30)$$

The reduced density matrix $\rho_R^{W_3}$ corresponding to the modes of Rob in region I of the Rindler spacetime is an infinite dimensional diagonal matrix, with δ_m as its diagonal elements. Then, the m^{th} eigenvalue of $\rho_R^{W_3}$ is given by

$$\delta_m = \frac{\tanh^{2m} r}{\cosh^2 r} \left(1 + \frac{m - \sinh^2 r}{\sinh^2 r} x^2 \right). \quad (31)$$

To characterize the non-separability in the state $\rho_{A_1 A_2 R}^{W_3}$ given in Eq. (28), we substitute these eigenvalues δ_n 's and δ_m 's in the expression for AR q -conditional entropy for the bipartition $A_1 A_2 : R$ to obtain

$$S_q^{W_3}(A_1 A_2 | R) = \frac{1}{q-1} \times \left[1 - \frac{\sum_{n=0}^{\infty} \left(\tanh^{2n} r \left(1 + \frac{n+1 - \cosh^2 r}{\cosh^2 r} x^2 \right) \right)^q}{\sum_{m=0}^{\infty} \left(\tanh^{2m} r \left(1 + \frac{m - \sinh^2 r}{\sinh^2 r} x^2 \right) \right)^q} \right]. \quad (32)$$

We are considering the non-separability in the inertial : non-inertial bipartition as simultaneous measurements cannot be done in the other possible partitions. To numerically compute this conditional entropy, we use the eigenvalue truncation procedure and get $k \simeq 10^4$. Now, using these k eigenvalues of $\rho_{A_1 A_2 R}^{W_3}$ and $\rho_R^{W_3}$, we numerically compute the AR q -conditional entropy in Eq. (32) and characterize the same with respect to its state parameters θ , ϕ , acceleration λ and parameter q .

We observe that $S_q^{W_3}(A_1 A_2 | R)$ consistently remains below zero, suggesting the non-separability of the state for the entire range of system parameters, acceleration λ and parameter q . The maximum non-separability for generalized pure 3-qubit W states occurs for the choice of θ and ϕ such that $\sin \theta \cos \phi = 1/\sqrt{2}$. Moreover, to understand the behavior of $S_q^{W_3}(A_1 A_2 | R)$ with respect to the other numerous parameters involved, we choose a pure 3-qubit W state with equal coefficients, i.e., with state parameters $\theta = \cos^{-1}(1/\sqrt{3})$ and $\phi = \pi/4$. Subsequently, the AR q -conditional entropy for this state is then plotted as a function of the acceleration parameter λ in Fig. 3 for various q .

When Rob's acceleration is low, the non-separability of the state $\rho_{A_1 A_2 R}^{\text{W}_3}$ is most pronounced, as indicated by the minimum of $S_q^{\text{W}_3}(A_1 A_2 | R)$. However, with increasing acceleration, the conditional entropy rises and approaches 0 as λ tends to 1, across all values of q . As q increases, the $S_q^{\text{W}_3}(A_1 A_2 | R)$ becomes more negative for low λ , but converging to 0 as λ tends to 1. Although the choice of this θ and ϕ are not unique, the behavior of $S_q^{\text{W}_3}(A_1 A_2 | R)$ with respect to λ will follow the same trend. However, these values of conditional entropy are higher than the corresponding conditional entropy for the generalized pure 3-qubit GHZ state with one of its qubit in acceleration and hence indicating generalized pure 3-qubit W states having weaker non-separability than generalized pure 3-qubit GHZ states with one of its qubits being accelerated.

Pure N -qubit W state: A pure N -qubit W state, shared between Alice and Bob, is given by

$$|W_N\rangle = \frac{|00 \cdots 01\rangle + |00 \cdots 10\rangle + \cdots + |10 \cdots 00\rangle}{\sqrt{N}}, \quad (33)$$

where, say, the first $N-1$ qubits are with Alice, while the last qubit is in Bob's possession. Here, we again consider only the pure N -qubit W state instead of the generalized N -qubit W state for logistical reasons. The final Alice-Rob state $\rho_{A_1 A_2 \cdots A_{N-1} R}^{\text{W}_N}$, after accelerating Bob's qubit and tracing out the second Rindler region of Rob's spacetime using the recipe given in Sec. II, is given by

$$\begin{aligned} \rho_{A_1 A_2 \cdots A_{N-1} R}^{\text{W}_N} &= \frac{1}{N \cosh^2 r} \sum_{n=0}^{\infty} \tanh^{2n} r \\ &\times \left[\alpha_n^2 |00 \cdots 0n+1\rangle \langle 00 \cdots 0n+1| \right. \\ &+ \left(|00 \cdots 1n\rangle + \cdots + |01 \cdots 0n\rangle + |10 \cdots 0n\rangle \right) \\ &\times \left(\langle 00 \cdots 1n| + \cdots + \langle 01 \cdots 0n| + \langle 10 \cdots 0n| \right) \\ &+ \alpha_n \left(|00 \cdots 0n+1\rangle \left(\langle 00 \cdots 1n| + \cdots + \langle 01 \cdots 0n| \right. \right. \\ &\left. \left. + \langle 10 \cdots 0n| \right) + \text{c.c.} \right) \left. \right]. \quad (34) \end{aligned}$$

Since one of the qubits is accelerating in a non-inertial frame, a simultaneous measurement cannot be done on the inertial and the non-inertial qubits together. Hence we proceed our investigations by choosing the $A_1 A_2 \cdots A_{N-1} : R$ bipartition to calculate the AR q -conditional entropy $S_q^{\text{W}_N}(A_1 A_2 \cdots A_{N-1} | R)$. This conditional entropy is evaluated explicitly for the state given in Eq. (34), using the eigenvalues of the reduced system

$\rho_{A_1 A_2 \cdots A_{N-1} R}^{\text{W}_N}$ and its subsystem $\rho_R^{\text{W}_N}$ and is given by

$$\begin{aligned} S_q^{\text{W}_N}(A_1 A_2 \cdots A_{N-1} | R) &= \frac{1}{q-1} \\ &\times \left[1 - \frac{\sum_{n=0}^{\infty} \left(\tanh^{2n} r \left(N-1 + \frac{n+1}{\cosh^2 r} \right) \right)^q}{\sum_{m=0}^{\infty} \left(\tanh^{2m} r \left(N-1 + \frac{m}{\sinh^2 r} \right) \right)^q} \right]. \quad (35) \end{aligned}$$

This conditional entropy $S_q^{\text{W}_N}(A_1 A_2 \cdots A_{N-1} | R)$ can be numerically computed for various values of N , λ , and q using the eigenvalue truncation procedure given previously. With $k \simeq 10^4$, the $S_q^{\text{W}_N}(A_1 A_2 \cdots A_{N-1} | R)$ is found to be again non-separable throughout the ranges of system parameters, acceleration λ and q . The conditional entropy is plotted for various values of N and λ for $q = 2$ in Fig. 4. We observe that irrespective of N , the non-separability of $\rho_{A_1 A_2 \cdots A_{N-1} R}^{\text{W}_N}$ is high at lower values of acceleration and decreases with the increase in acceleration as $S_q^{\text{W}_N}(A_1 A_2 \cdots A_{N-1} | R)$ tends to 0 when λ reaches 1. However, as N increases, the variation of $S_q^{\text{W}_N}(A_1 A_2 \cdots A_{N-1} | R)$ with respect to λ becomes less steeper. Hence, we can conclude that the non-separability of the state $\rho_{A_1 A_2 \cdots A_{N-1} R}^{\text{W}_N}$ reduces as the number of qubits inertial subsystem increases. This observation remains true for all values of parameter q .

To consolidate, all the pure multiqubit GHZ and W states with one of its qubits under acceleration will remain non-separable in the inertial : non-inertial bipartition throughout the ranges of its state parameters, parameter q and acceleration λ . Moreover, the states are highly non-separable at low acceleration λ and they tend towards separability at very high acceleration of λ tending to 1.

V. MULTIQUBIT MIXED STATES WITH AN ACCELERATING QUBIT

In this section, we introduce a global noise to all the pure states considered in Sec. IV, when they have equal probability in their superposed states and study the non-separability of resultant mixed states, when one of its qubits is accelerated, using AR q -conditional entropy. The non-separability of the mixed states with an accelerating qubit thus obtained is characterized with respect to parameter q , acceleration parameter λ and mixing parameter p .

A. Mixed multiqubit GHZ states

Mixed 2-qubit GHZ state: A mixed 2-qubit GHZ state shared between Alice and Bob, obtained by adding a global noise to the pure 2-qubit GHZ state is given by

$$\varrho_{AB}^{\text{GHZ}_2} = \frac{p}{4} \mathbb{I}_d + (1-p) |\psi_{AB}\rangle \langle \psi_{AB}|, \quad (36)$$




FIG. 5: (Color online) (a) The variation of Abe-Rajagopal q -conditional entropy of a single qubit accelerated mixed 2-qubit GHZ state as a function of p and λ for $q = 2$. The black solid line corresponds to transition from negative to positive AR q -conditional entropy (b) The variation of the the mixing parameter p for which the $S_q^{\text{GHZ}_2}(A|R) = 0$, with respect to q for acceleration parameter λ . We can observe that the values of mixing parameter saturates to a values a $q \rightarrow \infty$ for a given acceleration λ . (c) The variation of the the mixing parameter p for which the $S_q^{\text{GHZ}_2}(A|R) = 0$, with respect to acceleration parameter λ for $q = 150$. The blue region denotes the region where the state remains non-separable, while the red region denotes the separability.

where $|\psi_{AB}\rangle = (|00\rangle + |11\rangle)/\sqrt{2}$, \mathbb{I}_d is the 4×4 identity matrix and p is the mixing parameter. Note that, p is a parameter in the range $[0, 1]$ and $p = 0$ denotes the pure 2-qubit GHZ state and $p = 1$ denotes the maximally mixed 2-qubit GHZ state. Let us consider that Bob is now relativistically accelerating with respect to Alice and the resulting composite state is $\varrho_{AR_I R_{II}}^{\text{GHZ}_2}$. We now obtain the reduced state of the Alice and Rob in region I, by tracing out Rob's mode in region II using the recipe given in Sec. II, as

$$\begin{aligned} \varrho_{AR}^{\text{GHZ}_2} &= \frac{1}{\cosh^2 r} \sum_{n=0}^{\infty} \tanh^{2n} r \\ &\times \left[\frac{2-p}{4} \left(|0n\rangle \langle 0n| + \frac{n+1}{\cosh^2 r} |1n+1\rangle \langle 1n+1| \right) \right. \\ &+ \frac{p}{4} \left(|1n\rangle \langle 1n| + \frac{n+1}{\cosh^2 r} |0n+1\rangle \langle 0n+1| \right) \\ &\left. + \frac{1-p}{2} \frac{\sqrt{n+1}}{\cosh r} \left(|0n\rangle \langle 1n+1| + |1n+1\rangle \langle 0n| \right) \right]. \end{aligned} \quad (37)$$

The Rob's subsystem in region I, by tracing out Alice's qubit, is given by

$$\begin{aligned} \varrho_R^{\text{GHZ}_2} &= \frac{1}{2 \cosh^2 r} \sum_{n=0}^{\infty} \tanh^{2n} r \\ &\times \left(|n\rangle \langle n| + \frac{n+1}{\cosh^2 r} |n+1\rangle \langle n+1| \right). \end{aligned} \quad (38)$$

These infinite dimensional density matrices does not possess a diagonal block structure; hence arriving at the an-

alytical form for their eigenvalues is not straightforward and one has to resort to numerical techniques. Below, we describe a procedure to obtain the eigenvalues of these two density matrices for evaluating the AR q -conditional entropy.

Density matrix truncation procedure: To numerically compute the AR q -conditional entropy, we make use of the fact that the sum of the eigenvalues of $\varrho_{AR}^{\text{GHZ}_2}$ and $\varrho_R^{\text{GHZ}_2}$ tends to 1 as n and m approaches infinity. However, unlike the case of pure states we do not have the analytical expression for $S_q^{\text{W}_2}(A|R)$ in terms of the eigenvalues of $\varrho_{AR}^{\text{GHZ}_2}$ and $\varrho_R^{\text{GHZ}_2}$. Therefore, we use the numerical technique in which the matrix dimension of infinity is approximated to an agreeable finite dimension $l \times l$ of the density matrix, such that $\text{Tr}(\varrho_{AR}^{\text{GHZ}_2})$ approaches to 1. For the density matrix given in Eq. (37), the dimension of $\varrho_{AR}^{\text{GHZ}_2}$ for which $\text{Tr}(\varrho_{AR}^{\text{GHZ}_2})$ approaches 1 is found to be around $l \approx 10^3$. Similarly, eigenvalues are also obtained for the Rob's subsystem in region I, $\varrho_R^{\text{GHZ}_2}$, by limiting its dimension to the finite $l \times l$, such that $\text{Tr}(\varrho_R^{\text{GHZ}_2})$ approaches to 1.

Now, the AR q -conditional entropy is numerically computed for the state given in Eq. (37) using density matrix truncation procedure given above, for various values of Rob's acceleration λ , mixing parameter p and parameter q . The AR q -conditional entropy for the initial mixed state whose subsystem is accelerated is given by $S_q^{\text{GHZ}_2}(A|R)$ and is plotted as a function of the acceleration λ and parameter p in Fig. 5a for $q = 2$. We can observe that in the plot of $S_q^{\text{GHZ}_2}(A|R)$ for the state given in Eq. (37), there exist regions of both separability

(red shades) and non-separability (blue shades) depending on the choices of λ and p . The $S_q^{\text{GHZ}_2}(A|R) = 0$ curve (black dotted) indicates the transition of $\varrho_{AR}^{\text{GHZ}_2}$ from non-separability to separability and it depends on values of λ and p . The whole transition curve shifts towards the right as q increases, indicating that with an increase in q value, the area of non-separability region with respect to λ and p keeps increasing. As q is increased to large values, say 50 and above, the transition curve becomes saturated with respect to λ and p as interpreted from Fig. 5b. Hence, the strongest condition on separability can be obtained in the asymptotic limit of q tending to ∞ .

For the asymptotic limit of $q \rightarrow \infty$, in Fig. 5c, we plot the transition point from non-separability to separability obtained numerically, i.e., $S_q^{\text{GHZ}_2}(A|R) = 0$ with respect to p and λ (dashed curve). Note that the saturation occurs beyond $q = 50$ and we chose a reasonably high value of $q = 150$ for the study of separability. The region with blue shade denotes the non-separable states, while the region with red shade denotes the separable states. These transition points for different acceleration can be fitted to a curve of the form

$$(p/b)^x + \lambda^y = 1. \quad (39)$$

The choice of the function for the fitted curve is made using the fact that the separability occurs at $p = b$ when the acceleration $\lambda = 0$ and $p = 0$ as the acceleration $\lambda \rightarrow 1$. Hence, the separability criteria for the state in Eq. (37), which relates the mixing parameter p and Rob's

acceleration λ , is given by

$$p > \frac{2}{3} (1 - \lambda^{2.2})^{(1/2.05)}. \quad (40)$$

When the Rob's acceleration $\lambda = 0$, the necessary and sufficient condition for separability of this state occurs at $p > 2/3$ [67], which is recovered from the Eq. (40). As the acceleration increases, the non-separable region keeps reducing and the non-separability vanishes only at acceleration λ approaching to 1. Hence, the Eq. (40) gives the strongest condition on separability in asymptotic limit of q for the mixed 2-qubit GHZ state with one of its qubits being accelerated.

Mixed N -qubit GHZ state: We add a global noise to pure N -qubit GHZ state to obtain mixed N -qubit GHZ state, which is shared between two parties Alice and Bob in the inertial frame, where Alice has the first $N - 1$ qubits, and Bob has N^{th} qubit in his possession. This state is given by

$$\varrho_{A_1 A_2 \dots A_{N-1} B}^{\text{GHZ}_N} = \frac{p}{2^N} \mathbb{I}_d + (1-p) |\psi_{A_1 A_2 \dots A_{N-1} B}\rangle \langle \psi_{A_1 A_2 \dots A_{N-1} B}|, \quad (41)$$

where $|\psi_{A_1 A_2 \dots A_{N-1} B}\rangle = (|0\rangle^{\otimes N} + |1\rangle^{\otimes N})/\sqrt{2}$, \mathbb{I}_d is the $2^N \times 2^N$ identity matrix and p is the mixing parameter. Let us now consider that Bob is uniformly accelerating with respect to Alice and the resulting state $\varrho_{A_1 A_2 \dots A_{N-1} R_I R_{II}}^{\text{GHZ}_N}$ is obtained by using the transformations from Minkowski to Rindler spacetime in Eqs. (1) and (2). The final state $\varrho_{A_1 A_2 \dots A_{N-1} R}^{\text{GHZ}_N}$ shared between Alice and Rob's mode in region I is obtained using the recipe given in Sec. II and is given by

$$\begin{aligned} \varrho_{A_1 A_2 \dots A_{N-1} R}^{\text{GHZ}_N} &= \frac{1}{\cosh^2 r} \sum_{n=0}^{\infty} \tanh^{2n} r \\ &\left[\frac{2-p}{2^N} \left((|00 \dots n\rangle + |01 \dots n\rangle + \dots + |11 \dots n\rangle) (\langle 00 \dots n| + \langle 01 \dots n| + \dots + \langle 11 \dots n|) \right) \right. \\ &+ \alpha_n^2 (|00 \dots n+1\rangle + |01 \dots n+1\rangle + \dots + |11 \dots n+1\rangle) (\langle 00 \dots n+1| + \langle 01 \dots n+1| + \dots + \langle 11 \dots n+1|) \left. \right) \\ &+ \frac{1-p}{2} \left(|00 \dots 0n\rangle \langle 00 \dots 0n| + \alpha_n (|00 \dots 0n\rangle \langle 11 \dots 1n+1| + |11 \dots 1n+1\rangle \langle 00 \dots 0n|) \right) \\ &\left. + \alpha_n^2 |11 \dots 1n+1\rangle \langle 11 \dots 1n+1| \right]. \quad (42) \end{aligned}$$

The Rob's subsystem is obtained by tracing out the inertial Alice's qubits, i.e.,

$$\varrho_R^{\text{GHZ}_N} = \text{Tr}_{A_1 A_2 \dots A_{N-1}} (\varrho_{A_1 A_2 \dots A_{N-1} R}^{\text{GHZ}_N}).$$

Using the density matrix truncation procedure given in last subsection, we numerically obtain the AR conditional entropy $S_q^{\text{GHZ}_N}(A_1 A_2 \dots A_{N-1} | R)$ in the inertial : non-inertial bipartition. For example, when a




FIG. 6: (Color online) The variation of the non-separable to separable transition points of AR q -conditional entropy for the (a) mixed N -qubit GHZ state and (b) mixed N -qubit W state, with one of its qubits in uniform acceleration, as a function of the acceleration parameter λ and mixing parameter p for different values of $N = 3, 4$ and 5 is given in this figure. The fitted curves are shown in the solid colored lines.

mixed 3-qubit GHZ state with one of its qubits under acceleration is considered (see Appendix 1), we observe a transition from non-separability to separability depending on the values of acceleration λ , parameter q and mixing parameter p . In the asymptotic limit of $q \rightarrow \infty$, the transition from non-separability to separability saturates to a curve, which is fitted to the function given in Eq. (39). Then, the separability criteria for the state $\varrho_{A_1 A_2 R}^{\text{GHZ}_3}$ (see Eq. (A.2)), in terms of the mixing parameter, is given as

$$p > 0.56(1 - \lambda^{2.47})^{(1/1.49)}. \quad (43)$$

Here, the separability occurs at $p = 0.56$ when the acceleration $\lambda = 0$ and it reduces to $p = 0$ when $\lambda = 1$ indicating the region of non-separability decreases with increase in acceleration for the 3-qubit mixed GHZ state with one of its qubits in acceleration.

Similar investigation is performed for multiqubit mixed GHZ states with $N = 4, 5$ when one of their qubits is accelerated. The non-separability to separability transition with respect to λ and p in the asymptotic limit of $q \rightarrow \infty$ is observed to be similar to the $N = 3$ case. To study the impact of N on the non-separability to separability transition, we plot $S_q^{\text{GHZ}_N}(A_1 A_2 \cdots A_{N-1} | R) = 0$ with respect to p and λ for $N = 3, 4$ and 5 in Fig. 6a. The strongest condition on separability obtained in the asymptotic limit of $q \rightarrow \infty$ for these states are fitted to the family of curves in Eq. (39). The non-separability region of mixed N -qubit GHZ states with one of its qubit under acceleration reduces as N increases. For all choices of N , the states tend to become separable as the acceleration λ approaches to 1. These separability conditions for the different mixed N -qubit GHZ states with one of

their qubits being accelerated ($N = 2, 3, 4$ and 5) are consolidated in the Table I.

B. Mixed multiqubit W states

Mixed 2-qubit W state: A mixed 2-qubit state shared between Alice and Bob, obtained by introducing a global noise to the pure 2-qubit W state is given by

$$\varrho_{AB}^{\text{W}_2} = \frac{p}{4} \mathbb{I}_d + (1-p) |\psi_{AB}\rangle \langle \psi_{AB}|, \quad (44)$$

where $|\psi_{AB}\rangle = (|01\rangle + |10\rangle)/\sqrt{2}$, \mathbb{I}_d is the 4×4 identity matrix and p is the mixing parameter. Let us consider that Bob is now relativistically accelerating with respect to Alice and the resulting composite state is $\varrho_{AR_I R_{II}}^{\text{W}_2}$. We now obtain the reduced state of the Alice and Rob in region I, by tracing out Rob's mode in region II using recipe given in Sec. II, as

$$\begin{aligned} \varrho_{AR}^{\text{W}_2} &= \frac{1}{\cosh^2 r} \sum_{n=0}^{\infty} \tanh^{2n} r \\ &\times \left[\frac{2-p}{4} \left(|1n\rangle \langle 1n| + \frac{n+1}{\cosh^2 r} |0n+1\rangle \langle 0n+1| \right) \right. \\ &+ \frac{p}{4} \left(|0n\rangle \langle 0n| + \frac{n+1}{\cosh^2 r} |1n+1\rangle \langle 1n+1| \right) \\ &\left. + \frac{1-p}{2} \frac{\sqrt{n+1}}{\cosh r} \left(|0n+1\rangle \langle 1n| + |1n\rangle \langle 0n+1| \right) \right]. \end{aligned} \quad (45)$$

The Rob's subsystem $\varrho_R^{\text{W}_2}$ is obtained by tracing out the inertial Alice's qubits from the state $\varrho_{AR}^{\text{W}_2}$ and is same as

$\varrho_R^{\text{GHZ}_2}$ given in Eq. (38). Now, using the density matrix truncation procedure given in Sec. V A, the conditional entropy $S_q^{\text{W}_2}(A|R)$ can be numerically evaluated for this state. The characterization of $S_q^{\text{W}_2}(A|R)$ for this state qualitatively remains the same as given in Figs. 5a and 5b for mixed 2-qubit GHZ state with one of its qubit in acceleration, but differ in values. The non-separability to separability transition for this state in the asymptotic limit of q can also be fitted to the function given in Eq. (39). Thus, the separability criteria in terms of mixing parameter and acceleration λ for $\rho_{AR}^{\text{W}_2}$ is given by

$$p > \frac{2}{3} (1 - \lambda^{1.57})^{(1/2.6)}, \quad (46)$$

indicating when $\lambda = 0$, we get $p > 2/3$ as the separable region and the separable region increases as λ increases. The mixing parameter p above which the state is separable tends to 0 as the acceleration λ tends to 1 in the asymptotic limit of q .

Mixed N -qubit W state: We get a mixed N -qubit W state by introducing a global noise to the pure N -qubit W state, and is given by

$$\varrho_{A_1 A_2 \dots A_{N-1} B}^{\text{W}_N} = \frac{p}{2^N} \mathbb{I}_d + (1-p) |\psi_{A_1 A_2 \dots A_{N-1} B}\rangle \langle \psi_{A_1 A_2 \dots A_{N-1} B}|, \quad (47)$$

where $|\psi_{A_1 A_2 \dots A_{N-1} B}\rangle = (|00 \dots 01\rangle + |00 \dots 10\rangle + \dots + |10 \dots 00\rangle) / \sqrt{N}$, \mathbb{I}_d is the $2^N \times 2^N$ identity matrix and p is the mixing parameter. This state is shared between the two parties Alice and Bob in the inertial frame, where Alice has the first $N-1$ qubits, and Bob has N^{th} qubit in his possession. Let us now consider that Bob is now uniformly accelerated with respect to Alice and the resulting state $\varrho_{A_1 A_2 \dots A_{N-1} R_I R_{II}}^{\text{W}_N}$ is obtained by using the transformations from Minkowski to Rindler spacetime given in Eqs. (1) and (2). The final state $\varrho_{A_1 A_2 \dots A_{N-1} R}^{\text{W}_N}$ between Alice and modes of Rob in region I using the recipe given in Sec. II is given by

$$\begin{aligned} \varrho_{A_1 A_2 \dots A_{N-1} R}^{\text{W}_N} &= \frac{1}{\cosh^2 r} \sum_{n=0}^{\infty} \tanh^{2n} r \\ &\left[\frac{2-p}{2^N} \left((|00 \dots n\rangle + |01 \dots n\rangle + \dots + |11 \dots n\rangle) (\langle 00 \dots n| + \langle 01 \dots n| + \dots + \langle 11 \dots n|) \right) \right. \\ &+ \alpha_n^2 (|00 \dots n+1\rangle + |01 \dots n+1\rangle + \dots + |11 \dots n+1\rangle) (\langle 00 \dots n+1| + \langle 01 \dots n+1| + \dots + \langle 11 \dots n+1|) \left. \right) \\ &+ \frac{1-p}{N} \left(\alpha_n^2 |00 \dots 0n+1\rangle \langle 00 \dots 0n+1| + \alpha_n (|00 \dots 0n+1\rangle (\langle 00 \dots 1n| + \dots + \langle 10 \dots 0n|) \right. \\ &+ (|00 \dots 1n\rangle + \dots + |10 \dots 0n\rangle) \langle 00 \dots 0n+1|) \\ &\left. + (|00 \dots 1n\rangle + \dots + |10 \dots 0n\rangle) (\langle 00 \dots 1n| + \dots + \langle 10 \dots 0n|) \right) \left. \right]. \quad (48) \end{aligned}$$

The Rob's subsystem $\varrho_R^{\text{W}_N}$ is obtained by tracing out the inertial Alice's qubits. We numerically obtain the AR q -conditional entropy $S_q^{\text{W}_N}(A_1 A_2 \dots A_{N-1} | R)$ in the inertial : non-inertial bipartition, using the density matrix truncation procedure given in Sec. V A. For example, when $N = 3$, a mixed 3-qubit W state with one of its qubits under acceleration is considered (see Appendix 2), depending on the values of acceleration λ , parameter q and mixing parameter p , we observe a transition from non-separability to separability. This transition saturates to a curve in the asymptotic limit $q \rightarrow \infty$, which is fitted to the function given in Eq. (39). Then, the separability criteria for the state $\varrho_{A_1 A_2 R}^{\text{W}_3}$ given in Eq. (A.4), in terms of the mixing parameter p and λ , is

$$p > 0.47(1 - \lambda^{1.86})^{(1/1.68)}, \quad (49)$$

implying that, $p > 0.47$ is separable region when $\lambda = 0$ and as λ tends to 1 the state remain separable irrespective of the value of mixing parameter p .

Further, we observe a similar non-separability to separability transition with respect to λ and p in the asymptotic limit of $q \rightarrow \infty$ for $N = 4$ and 5. In Fig. 6b, we plot $S_q^{\text{GHZ}_N}(A_1 A_2 \dots A_{N-1} | R) = 0$ with respect to p and λ for different values of $N = 3, 4$ and 5. These separability to non-separability transitions are fitted to the family of curves in Eq. (39). We can observe that the non-separability region reduces as N increases. Moreover, this reduction is much faster than its mixed N -qubit GHZ counterpart, indicating that mixed N -qubit GHZ states can sustain non-separability better than mixed N -qubit W states, when one of their qubit is uniformly accelerated

TABLE I: This table shows the conditions for separability obtained using the AR q -conditional entropy for multiqubit GHZ and W states. The multiqubit pure states with one of its qubit in acceleration remain non-separable (NS) everywhere. The conditions given for mixed GHZ and W states fit to the family of curves given in Eq. (39).

N	Pure $\rho_{AR}^{\text{GHZ}^N} / \rho_{AR}^{\text{W}^N}$	mixed $\varrho_{AR}^{\text{GHZ}^N}$	mixed $\varrho_{AR}^{\text{W}^N}$
2	NS	$\frac{2}{3}(1 - \lambda^{2.2})^{(1/2.05)}$	$\frac{2}{3}(1 - \lambda^{1.57})^{(1/2.6)}$
3	NS	$p > 0.56(1 - \lambda^{2.47})^{(1/1.49)}$	$p > 0.47(1 - \lambda^{1.86})^{(1/1.68)}$
4	NS	$p > 0.53(1 - \lambda^{1.92})^{(1/1.78)}$	$p > 0.36(1 - \lambda^{2.04})^{(1/1.32)}$
5	NS	$p > 0.51(1 - \lambda^{2.14})^{(1/1.55)}$	$p > 0.29(1 - \lambda^{2.28})^{(1/1.12)}$

with respect to others. The consolidated conditions of separability of single qubit accelerated multiqubit mixed GHZ and W states are presented in Table I.

VI. CONCLUSIONS

A shared state between two parties Alice and Bob ρ_{AB} , in an inertial frame can be represented using Minkowski coordinates. When Bob undergoes a uniform acceleration, the vacuum and excited states in Minkowski coordinates will transform to corresponding two-mode squeezed vacuum state representations in Rindler coordinates. The two modes of this accelerated Bob's (Rob) state belong to the two disjoint regions of the Rindler spacetime. Therefore, the final mixed state of inertial Alice and non-inertial Rob, ρ_{AR} , can be obtained by tracing out the region II of the Rindler spacetime. The non-separability of this relativistic reduced state is characterized using the AR q -conditional entropy. The AR q -conditional entropy takes negative values for a non-separable states and positive values for separable states. We capture the non-separability of pure multiqubit GHZ and W states, when one of their qubits is accelerated. Further, we study the non-separability of mixed states generated by adding a global noise to the same multiqubit GHZ and W states, when one of their qubits is in uniform acceleration.

The analytical form of AR q -conditional entropy for pure multiqubit GHZ and W states is evaluated using eigenvalues of the state of inertial Alice and Rob's mode in region I ($\rho_{A_1 A_2 \dots A_{N-1} R}$) and the Rob's subsystem (ρ_R). We then numerically study the non-separability using the *eigenvalue truncation procedure* performed on these infinite dimensional density matrices $\rho_{A_1 A_2 \dots A_{N-1} R}$ and ρ_R to get the required eigenvalues. The AR q -conditional entropy $S_q(A_1 A_2 \dots A_{N-1} | R)$ in the inertial : non-inertial bipartition always remains less than zero for both pure multiqubit GHZ and W states with one of their qubit under uniform acceleration, implying that they remain non-separable for any values of their state parameters and acceleration λ . Further, we observe that irrespective of the value of q , the non-separability of $\rho_{A_1 A_2 \dots A_{N-1} R}$ will be high at low values of acceleration and decreases with the increase in acceleration, in particular, the AR q -conditional entropy $S_q(A_1 A_2 \dots A_{N-1} | R)$ tends to 0 as

$\lambda \rightarrow 1$. Further, we observe that the non-separability of pure multiqubit GHZ state with one of its qubits in acceleration is not dependent on the N , i.e., it is independent of the number of qubits in the inertial frame. However, when the number of qubits in the inertial frame of the pure multiqubit W state with one of its qubits in acceleration is increased, the non-separability decreases as N increases.

We further extend our study to various mixed multiqubit states, $\varrho_{A_1 A_2 \dots A_{N-1} B}$, which are generated by mixing global noise to the pure multiqubit GHZ and W states. The non-separability using AR q -conditional entropy $S_q(A_1 A_2 \dots A_{N-1} | R)$ in the inertial : non-inertial bipartition is evaluated for these mixed multiqubit states, $\varrho_{A_1 A_2 \dots A_{N-1} R}$, with one of their qubits is accelerated. For these mixed multiqubit states, the AR q -conditional entropy is numerically calculated by employing the *density matrix truncation procedure*. We could observe that there exist regions of both separability and non-separability of the state $\varrho_{A_1 A_2 \dots A_{N-1} R}$ depending on the choices of acceleration λ , mixing parameter p and parameter q . As q is increased to large values, the transition curve $S_q(A_1 A_2 \dots A_{N-1} | R) = 0$ becomes saturated with respect to λ and p . Hence, the strongest condition on separability for these states, in terms of the acceleration and mixing parameter, is obtained in the asymptotic limit of q tending to ∞ . These transition points are fitted to the family of curves given by $(p/b)^x + \lambda^y = 1$. The separability conditions for all these multiqubit mixed GHZ and W states when one of their qubits is in uniform acceleration considered in our study are consolidated in Table I. For a mixed N -qubit GHZ and W states with one of their qubits is accelerated, we observe that the non-separable region reduces as the number of qubits in the inertial frame increases. The reduction of the non-separable region, as acceleration λ is increased, is much faster for the N -qubit mixed W state, $\varrho_{A_1 A_2 \dots A_{N-1} R}^{\text{W}^N}$, than its N -qubit mixed GHZ counterpart $\varrho_{A_1 A_2 \dots A_{N-1} R}^{\text{GHZ}^N}$, indicating that, in general, GHZ states can sustain the non-separability better than W states, when one of its qubit is uniformly accelerated with respect to others.

ACKNOWLEDGMENTS

We acknowledge the helpful discussions with P. Kiran. RP and HSH acknowledge financial support from the Science and Engineering Research Board (SERB), Government of India, under the project grant CRG/2021/008795. The authors acknowledge the computations performed using High Performance Computing facility *AnantGanak* at IIT Dharwad.

Appendix

1. Mixed 3-qubit GHZ state

A mixed 3-qubit GHZ state, with the first two qubits are with Alice and the third qubit in Bob's possession, is

$$\begin{aligned} \varrho_{A_1 A_2 R}^{\text{GHZ}_3} = & \frac{1}{\cosh^2 r} \sum_{n=0}^{\infty} \tanh^{2n} r \left[\frac{4-3p}{8} \left(|00n\rangle \langle 00n| + \frac{n+1}{\cosh^2 r} |11n+1\rangle \langle 11n+1| \right) \right. \\ & + \frac{1-p}{2} \frac{\sqrt{n+1}}{\cosh r} \left(|00n\rangle \langle 11n+1| + |11n+1\rangle \langle 00n| \right) + \frac{p}{8} \left(|11n\rangle \langle 11n| + |10n\rangle \langle 10n| + |01n\rangle \langle 01n| \right. \\ & \left. \left. + \frac{n+1}{\cosh^2 r} \left(|00n+1\rangle \langle 00n+1| + |01n+1\rangle \langle 01n+1| + |10n+1\rangle \langle 10n+1| \right) \right) \right]. \end{aligned} \quad (\text{A.2})$$

2. Mixed 3-qubit W state

A mixed 3-qubit inertial state, shared between Alice and Bob, is given by

$$\varrho_{A_1 A_2 B}^{\text{W}_3} = \frac{p}{8} \mathbb{I}_d + (1-p) |\psi_{A_1 A_2 B}\rangle \langle \psi_{A_1 A_2 B}| \quad (\text{A.3})$$

$$\begin{aligned} \varrho_{A_1 A_2 R}^{\text{W}_3} = & \frac{1}{\cosh^2 r} \sum_{n=0}^{\infty} \tanh^{2n} r \left[\frac{8-5p}{24} \left(|01n\rangle \langle 01n| + |10n\rangle \langle 10n| + \frac{n+1}{\cosh^2 r} |00n+1\rangle \langle 00n+1| \right) \right. \\ & + \frac{p}{8} \left(|11n\rangle \langle 11n| + |10n\rangle \langle 10n| + \frac{n+1}{\cosh^2 r} \left(|00n+1\rangle \langle 00n+1| + |01n+1\rangle \langle 01n+1| + |10n+1\rangle \langle 10n+1| \right) \right) \\ & \left. + \frac{1-p}{3} \left(|01n\rangle \langle 10n| + |10n\rangle \langle 01n| + \frac{\sqrt{n+1}}{\cosh r} \left(|01n\rangle \langle 00n+1| + |00n+1\rangle \langle 01n| + |00n+1\rangle \langle 10n| + |10n\rangle \langle 00n+1| \right) \right) \right]. \end{aligned} \quad (\text{A.4})$$

given by

$$\varrho_{A_1 A_2 B} = \frac{p}{8} \mathbb{I}_d + (1-p) |\psi_{A_1 A_2 B}\rangle \langle \psi_{A_1 A_2 B}|, \quad (\text{A.1})$$

where $|\psi_{A_1 A_2 B}\rangle = (1/\sqrt{2})(|000\rangle + |111\rangle)$, \mathbb{I}_d is the $2^3 \times 2^3$ identity matrix and p is the mixing parameter. Let us consider that Bob is now uniformly accelerating with respect to Alice and the resulting state, denoted as $\varrho_{A_1 A_2 R_I R_{II}}^{\text{GHZ}_3}$, is obtained by using the transformations from Minkowski to Rindler spacetime in Eqs. (1) and (2). Then, the mixed state $\varrho_{A_1 A_2 R}^{\text{GHZ}_3}$ after tracing out one of the disjoint regions in Rindler spacetime (region II) is given by

where $|\psi_{A_1 A_2 B}\rangle = (1/\sqrt{3})(|001\rangle + |010\rangle + |100\rangle)$, \mathbb{I}_d is the $2^3 \times 2^3$ identity matrix and p is the mixing parameter. Let us consider that Bob is now uniformly accelerating with respect to Alice and the resulting state $\varrho_{A_1 A_2 R}^{\text{W}_3}$, after tracing out the Rob's modes in region II of Rindler spacetime, is given by

[1] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Phys. Rev. Lett. **70**, 1895

(1993).

[2] S. Pirandola, J. Eisert, C. Weedbrook, A. Furusawa, and

- S. L. Braunstein, *Nature photonics* **9**, 641 (2015).
- [3] X.-M. Hu, Y. Guo, B.-H. Liu, C.-F. Li, and G.-C. Guo, *Nature Reviews Physics* **5**, 339 (2023).
- [4] C. H. Bennett and S. J. Wiesner, *Phys. Rev. Lett.* **69**, 2881 (1992).
- [5] C. Bennett and G. i. Brassard, in *Proc. of the IEEE Int. Conf. on Computers Systems and Signal Processing Bangalore India* (1984) p. 175.
- [6] P. W. Shor and J. Preskill, *Phys. Rev. Lett.* **85**, 441 (2000).
- [7] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Rev. Mod. Phys.* **81**, 865 (2009).
- [8] K. Modi, A. Brodutch, H. Cable, T. Paterek, and V. Vedral, *Rev. Mod. Phys.* **84**, 1655 (2012).
- [9] I. Fuentes-Schuller and R. B. Mann, *Phys. Rev. Lett.* **95**, 120404 (2005).
- [10] E. Martín-Martínez and J. León, *Phys. Rev. A* **81**, 052305 (2010).
- [11] B. Richter and Y. Omar, *Phys. Rev. A* **92**, 022334 (2015).
- [12] E. Martín-Martínez and J. León, *Phys. Rev. A* **81**, 032320 (2010).
- [13] D. E. Bruschi, A. Dragan, I. Fuentes, and J. Louko, *Phys. Rev. D* **86**, 025026 (2012).
- [14] F. Buscemi and P. Bordone, *Phys. Rev. A* **84**, 022303 (2011).
- [15] S. Moradi, *Phys. Rev. A* **79**, 064301 (2009).
- [16] E. Martín-Martínez and I. Fuentes, *Phys. Rev. A* **83**, 052306 (2011).
- [17] W.-C. Qiang, G.-H. Sun, Q. Dong, and S.-H. Dong, *Phys. Rev. A* **98**, 022320 (2018).
- [18] J. León and E. Martín-Martínez, *Phys. Rev. A* **80**, 012314 (2009).
- [19] S. Khan, N. A. Khan, and M. Khan, *Communications in Theoretical Physics* **61**, 281 (2014).
- [20] D. Ahn, Y. Moon, R. Mann, and I. Fuentes-Schuller, *Journal of High Energy Physics* **2008**, 062 (2008).
- [21] D. C. M. Ostapchuk and R. B. Mann, *Phys. Rev. A* **79**, 042333 (2009).
- [22] P. M. Alsing, I. Fuentes-Schuller, R. B. Mann, and T. E. Tessier, *Phys. Rev. A* **74**, 032326 (2006).
- [23] J. Wang and J. Jing, *Phys. Rev. A* **83**, 022314 (2011).
- [24] S. Khan, *Annals of Physics* **348**, 270 (2014).
- [25] E. Martín-Martínez and J. León, *Phys. Rev. A* **80**, 042318 (2009).
- [26] E. Martín-Martínez, L. J. Garay, and J. León, *Phys. Rev. D* **82**, 064028 (2010).
- [27] X.-H. Ge and S. P. Kim, *Classical and Quantum Gravity* **25**, 075011 (2008).
- [28] Q. Pan and J. Jing, *Phys. Rev. D* **78**, 065015 (2008).
- [29] S. Bhattacharya and N. Joshi, *Phys. Rev. D* **105**, 065007 (2022).
- [30] E. Martín-Martínez, L. J. Garay, and J. León, *Phys. Rev. D* **82**, 064006 (2010).
- [31] J. L. Ball, I. Fuentes-Schuller, and F. P. Schuller, *Physics Letters A* **359**, 550 (2006).
- [32] I. Fuentes, R. B. Mann, E. Martín-Martínez, and S. Moradi, *Phys. Rev. D* **82**, 045030 (2010).
- [33] G. V. Steeg and N. C. Menicucci, *Phys. Rev. D* **79**, 044027 (2009).
- [34] S.-M. Wu, H.-S. Zeng, and T. Liu, *Classical and Quantum Gravity* **39**, 135016 (2022).
- [35] N. Liu, J. Goold, I. Fuentes, V. Vedral, K. Modi, and D. E. Bruschi, *Classical and Quantum Gravity* **33**, 035003 (2016).
- [36] Z. Tian, J. Wang, H. Fan, and J. Jing, *Scientific reports* **5**, 7946 (2015).
- [37] M. Ahmadi, D. E. Bruschi, C. Sabín, G. Adesso, and I. Fuentes, *Scientific reports* **4**, 4996 (2014).
- [38] M. Ahmadi, D. E. Bruschi, and I. Fuentes, *Phys. Rev. D* **89**, 065028 (2014).
- [39] J. Wang, Z. Tian, J. Jing, and H. Fan, *Scientific reports* **4**, 7195 (2014).
- [40] J. Wang, J. Deng, and J. Jing, *Phys. Rev. A* **81**, 052120 (2010).
- [41] A. Datta, *Phys. Rev. A* **80**, 052304 (2009).
- [42] P. M. Alsing and G. J. Milburn, *Phys. Rev. Lett.* **91**, 180404 (2003).
- [43] E. Tjoa, *Phys. Rev. A* **106**, 032432 (2022).
- [44] P. T. Grochowski, G. Rajchel, F. Kiałka, and A. Dragan, *Phys. Rev. D* **95**, 105005 (2017).
- [45] S. Khan and N. A. Khan, *The European Physical Journal Plus* **130**, 1 (2015).
- [46] S. Haseli, *The European Physical Journal C* **79**, 616 (2019).
- [47] N. A. Khan, M. Jan, M. Shah, and D. Khan, *Annals of Physics* **440**, 168831 (2022).
- [48] M. A. Nielsen and J. Kempe, *Phys. Rev. Lett.* **86**, 5184 (2001).
- [49] O. Gühne and M. Lewenstein, *Phys. Rev. A* **70**, 022316 (2004).
- [50] V. Giovannetti, *Phys. Rev. A* **70**, 012102 (2004).
- [51] N. J. Cerf and C. Adami, *Phys. Rev. Lett.* **79**, 5194 (1997).
- [52] R. Horodecki and P. Horodecki, *Physics Letters A* **194**, 147 (1994).
- [53] R. Horodecki and M. Horodecki, *Phys. Rev. A* **54**, 1838 (1996).
- [54] R. Horodecki, P. Horodecki, and M. Horodecki, *Physics Letters A* **210**, 377 (1996).
- [55] C. Tsallis, S. Lloyd, and M. Baranger, *Phys. Rev. A* **63**, 042104 (2001).
- [56] S. Abe, *Phys. Rev. A* **65**, 052323 (2002).
- [57] R. Rossignoli, N. Canosa, and L. Ciliberti, *Journal of Russian Laser Research* **32**, 467 (2011).
- [58] J. Batle, A. R. Plastino, M. Casas, and A. Plastino, *Journal of Physics A: Mathematical and General* **35**, 10311 (2002).
- [59] J. Batle, A. Plastino, M. Casas, and A. Plastino, *The European Physical Journal B-Condensed Matter and Complex Systems* **35**, 391 (2003).
- [60] C. Tsallis, *Journal of statistical physics* **52**, 479 (1988).
- [61] C. Tsallis, R. Mendes, and A. R. Plastino, *Physica A: Statistical Mechanics and its Applications* **261**, 534 (1998).
- [62] A. Wehrl, *Rev. Mod. Phys.* **50**, 221 (1978).
- [63] S. Abe and A. K. Rajagopal, *Physica A: Statistical Mechanics and its Applications* **289**, 157 (2001).
- [64] S. Abe and A. Rajagopal, *Chaos, Solitons & Fractals* **13**, 431 (2002).
- [65] R. Prabhu, A. R. Usha Devi, and G. Padmanabha, *Phys. Rev. A* **76**, 042337 (2007).
- [66] Sudha, A. R. U. Devi, and A. K. Rajagopal, *Phys. Rev. A* **81**, 024303 (2010).
- [67] C. H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J. A. Smolin, and W. K. Wootters, *Phys. Rev. Lett.* **76**, 722 (1996).