

RECOVERING GENERALIZED HOMOLOGY FROM FLOER HOMOLOGY: THE COMPLEX ORIENTED CASE

LAURENT CÔTÉ AND YUSUF BARIŞ KARTAL

ABSTRACT. We associate an invariant called the completed Tate cohomology to a filtered circle-equivariant spectrum and a complex oriented cohomology theory. We show that when the filtered spectrum is the spectral symplectic cohomology of a Liouville manifold, this invariant depends only on the stable homotopy type of the underlying manifold. We make explicit computations for several complex oriented cohomology theories, including Eilenberg–MacLane spectra, Morava K-theories, their integral counterparts, and complex K-theory. We show that the result for Eilenberg–MacLane spectra depends only on the rational homology, and we use the computations for Morava K-theory to recover the integral homology (as an ungraded group). In a different direction, we use the completed Tate cohomology computations for the complex K-theory to recover the complex K-theory of the underlying manifold from its equivariant filtered Floer homotopy type. A key Floer theoretic input is the computation of local equivariant Floer theory near the orbit of an autonomous Hamiltonian, which may be of independent interest from the perspective of dynamics.

1. INTRODUCTION

1.1. **Context.** Given a Liouville manifold M , one can associate to it an invariant $SH(M)$ called *the symplectic cohomology*. This is a version of Hamiltonian the Floer cohomology for open manifolds. It depends implicitly on a choice of coefficient ring as well as (purely topological) grading and orientation data.

In contrast to Hamiltonian Floer cohomology on closed manifolds, symplectic cohomology does not in general remember any information about the homotopy type of M . For example, any subcritical Weinstein manifold has vanishing symplectic cohomology. In particular, if $M = N \times \mathbb{C}$ for a Liouville manifold N , then it follows from [Cie02] that $SH(M) = 0$. Therefore, the topology of M can be arbitrarily complicated, while the symplectic cohomology vanishes.

However, conjectures of Treumann [Tre19] inspired by string theory and mirror symmetry suggest that the topological K -theory of the underlying manifold should be determined by Floer theoretic invariants such as the wrapped Fukaya category. This is impossible as we have seen above: vanishing symplectic cohomology implies vanishing wrapped Fukaya category. However, Treumann suggests that once the extra data of an *action filtration* is remembered, one can reconstruct complex K -theory from these invariants.¹

Separately, the work of Albers–Cieliebak–Frauenfelder [ACF16] and independently Zhao [Zha19] shows that it is possible to recover rational information about the topology of M by remembering extra structures on symplectic cochains. The relevant structures are (i) the S^1 -action corresponding to “loop rotation” and (ii) the action filtration. [ACF16] and [Zha19] use these structures to build a version of “symplectic Tate (co)homology” with respect to the S^1 -action. They prove that the resulting theory recovers the *rational* cohomology of M ; however, it is insensitive to torsion information. In particular, most of the information about the homology and the complex K -theory of M is lost.

The goal of the present paper is to use methods of equivariant (Floer) homotopy theory ([CK23]) and chromatic homotopy theory to recover much more information about the stable homotopy type of a Liouville manifold M . Surprisingly, one can recover from the filtered S^1 -equivariant symplectic

¹Strictly speaking, Treumann’s conjecture is concerned only with cotangent bundles [Tre19, p. 15].

cohomology of M over the sphere spectrum, not only the integral homology and the complex K -theory, but also the Morava K -theories of M , which can be seen as the building blocks of the entire stable homotopy type of M (see [BB20] for a broad overview of the subject).

1.2. Extra structures on spectral symplectic cohomology. Classically, symplectic cohomology is constructed as a colimit of Floer cohomology groups associated to a sequence of Hamiltonians with slope increasing to infinity. In [Lar21], Large lifted this construction to the sphere spectrum \mathbb{S} under appropriate topological assumptions.

In [CK23], the authors further defined an S^1 -equivariant model $SH_S(M, \mathbb{S})$ for symplectic cohomology, where the S^1 -action corresponds to “loop rotation”. Analogously to [Lar21], the authors first construct an S^1 -equivariant spectrum $HF_S(H, \mathbb{S})$ associated to a (family of) Hamiltonian(s) linear at infinity, and take a colimit as the slope goes to infinity.

By choosing a cofinal sequence of Hamiltonians, one can naturally endow $SH_S(M, \mathbb{S})$ with the structure of a *filtered equivariant spectrum*, a notion which will be made precise in Section 2. While the filtration depends on the choice of the sequence, two such sequences produce equivalent filtered spectra. We emphasize that $SH_S(M, \mathbb{S})$, viewed as a filtered equivariant spectrum, is a true invariant of M as a Liouville manifold: it does not depend on any additional data such as e.g. a Hamiltonian or a contact-type hypersurface.

In summary, one can endow spectral symplectic cohomology with both an S^1 -action corresponding to loop rotation, and with a (suitably interpreted) action filtration.

1.3. Summary of results. To a complex oriented cohomology theory R and a filtered S^1 -equivariant spectrum X , we shall associate an invariant $\widehat{R}_{S^1}^*(X)$ which we call the *completed Tate cohomology*. The central result of this paper is the following computation:

Theorem 4.1. *If M is a stably framed Liouville manifold and R is a complex oriented ring spectrum,*

$$(1.1) \quad \widehat{R}_{S^1}^*(SH_S(M, \mathbb{S})) \simeq \widehat{R}_{S^1}^*(\Sigma^\infty(M/\partial_\infty M))$$

where $\Sigma^\infty(M/\partial_\infty M)$ is endowed with the trivial filtration and trivial S^1 action.

In other words, $\widehat{R}_{S^1}^*(SH_S(M, \mathbb{S}))$ only depends on the homotopy type of M . Moreover, as we will see, $\widehat{R}_{S^1}^*(\Sigma^\infty M_+)$ is a good approximation to $R_*(M)$ for many R , including Morava K -theories. We refer to Section 4.2 for a summary of the proof of Theorem 4.1.

Let us briefly explain the definition of $\widehat{R}_{S^1}^*(-)$, postponing the details to Section 2.3. First of all, recall that the complex orientation on R determines an isomorphism $R^*[[u]] = R^*(BS^1)$ (where $R^* = R^*(pt)$), and a formal group law $F_R \in R^*[[u_1, u_2]]$. The element $[n]_R(u) \in R^*[[u]]$ is, by definition, obtained by iterating u under F_R , n times. Given an S^1 -equivariant filtered spectrum X with filtration $X \simeq \text{colim}(X_1 \rightarrow X_2 \rightarrow \dots)$, the completed Tate cohomology $\widehat{R}_{S^1}^*(X)$ is defined as the limit of the $R^*((X_i)_{hS^1})$ localized at each $[n]_R(u)$ and u -adically completed.

We compute $\widehat{R}_{S^1}^*(SH_S(M, \mathbb{S}))$ explicitly in a variety of examples. For the Eilenberg–MacLane spectra $R = H\mathbb{Q}, H\mathbb{Z}, H\mathbb{F}_p$, we have the following analogues of [ACF16, Thm. 1.1(c) & Sec. 5.1] and [Zha19, Thm. 1.1 & Sec. 8]:

Corollary 4.2. *We have $\widehat{H}\widehat{\mathbb{Q}}_{S^1}^*(SH_S(M, \mathbb{S})) \simeq \widehat{H}\widehat{\mathbb{Z}}_{S^1}^*(SH_S(M, \mathbb{S})) \simeq H_{2n-*}(M, \mathbb{Q}((u)))$. On the other hand $\widehat{H}\widehat{\mathbb{F}}_{pS^1}^*(SH_S(M, \mathbb{S})) \simeq 0$.*

Note that for a commutative ring A , $\widehat{H}\widehat{A}_{S^1}^*(SH_S(M, \mathbb{S}))$ depends only on the S^1 -equivariant (co)filtered homotopy type of $SH^*(M, A)$, which does not require Floer homotopy theory to define. For $A = \mathbb{Q}$, this reaffirms the conclusion of [Zha19, ACF16] that the symplectic cohomology with its S^1 -action and filtration recovers the rational cohomology. For $A = \mathbb{Z}$ this can be interpreted as

a generalization of the computations for a disc and annulus in [Zha19, Sec. 8]. In this way, we lose the torsion information. See also [ACF16, Sec. 5.1] for related computations.

To access the torsion in the homology of M , we compute $\widehat{R}_{S^1}^*(SH_S(M, \mathbb{S}))$ for the (generalized) Morava K -theories $K_{p^k}(m)$:

Corollary 4.3. $\widehat{K}_{p^k}(m)_{S^1}^*(SH_S(M, \mathbb{S})) \simeq K_{p^k}(m)_{2n-*}(M)((u))$.

Corollary 4.2 and Corollary 4.3 also explain the term ‘‘completed Tate cohomology’’ (normally, to define Tate cohomology, one would localize $R^*((X_i)_{hS^1})$ at $[1]_R(u) = u$ only). For $R = H\mathbb{Q}$, $K_{p^k}(m)$, this produces the same answer.

This can be used to show

Theorem 4.4. *For $2(p^m - 1) > \dim_{\mathbb{R}}(M)$ (or for $4(p^m - 1) > \dim_{\mathbb{R}}(M)$ for Weinstein M), the groups $\widehat{K}_{p^k}(m)_{S^1}^*(SH_S(M, \mathbb{S}))$ and $H_{2n-*}(M, \mathbb{Z}[v_m, v_m^{-1}]/p^k)((u))$ are isomorphic.*

Combining Theorem 4.4 with the universal coefficients theorem, we show that the integral cohomology can be fully recovered:

Corollary 4.5. *The filtered S^1 -equivariant homotopy type of $SH_S(M, \mathbb{S})$ determines $H_*(M, \mathbb{Z})$.*

As noted, the invariant $\widehat{R}_{S^1}^*(SH(M, \mathbb{S}))$ depends on the filtration in a rather loose sense. Also, we choose to work with $SH(M, \mathbb{S})$ for simplicity, but the invariant $\widehat{R}_{S^1}^*(SH(M, \mathbb{S}))$ can be defined under milder assumptions on M , when R is complex oriented. Our results can likely be extended to this generality but we will not pursue this.

We also show we can recover complex K -theory:

Theorem 4.6. *After simultaneously completing at every prime p , the groups $\widehat{K}U_{S^1}^*(SH_S(M, \mathbb{S}))$ and $KU_{2n-*}(M)((u))$ become isomorphic. In other words, $\widehat{K}U_{S^1}^*(SH_S(M, \mathbb{S}))^\wedge \simeq KU_{2n-*}(M)((u))^\wedge$. As a result, the filtered S^1 -equivariant homotopy type of $SH_S(M, \mathbb{S})$ determines $KU_*(M)$ (and hence $KU^*(M)$ by the universal coefficients theorem).*

In other words Theorem 4.6 says that the (co)filtered equivariant homotopy type of ‘‘symplectic K -homology’’ determines the K -theory of the underlying manifold. (Recall that for an abelian group A , $A^\wedge := \lim_{n \in \mathbb{N}} A/nA$.) This result is consistent with Treumann’s proposal mentioned earlier; see Remark 4.7 for further discussions.

It is natural to ask whether it is possible to recover more information about M from symplectic cohomology with coefficients in various ring spectra.

Question 1.1. *Given a Liouville manifold M , does $SH_S(M; \mathbb{S})$ with its action filtration remember the stable homotopy type of M ?*

As we mentioned in the beginning, the Morava K -theories are the building blocks of spectra. As a result, it may already be possible to obtain an affirmative answer to Question 1.1 by combining the Floer theoretic input of this paper with other tools of chromatic homotopy theory, such as the chromatic tower and the chromatic fracture square.

Remark 1.2 (Potential dynamical applications). The constructions in this paper generalize to any autonomous Hamiltonian H that is linear at infinity (with respect to any choice of Liouville form), and associate to it an S^1 -equivariant spectrum. Moreover, the local Floer homology calculation Proposition 6.6 states that the S^1 -equivariant homotopy type in a small action window near a non-constant orbit $X \cong S^1/C_k$ of H is a Thom spectrum X^ν , where ν is an S^1 -equivariant virtual bundle over X . While the non-equivariant versions of this statement are straightforward (c.f. [CFHW96]), the S^1 -equivariant version requires the use of new machinery.

It seems reasonable to expect that the local equivariant Floer cohomology calculation could be useful in dynamical applications. Namely, an easy calculation shows that the S^1 -equivariant

R -cohomology $R_{S^1}^*(X^\nu) := R_{S^1}^*((X^\nu)_{hS^1})$ is equal (up to a shift) to (i) $R^*[[u]]/([k]_R(u))$ if ν is trivial (i.e. X is a *good orbit*), (ii) $R^*[[u]]/([k]_R(u)/[k/2]_R(u))$ if ν is non-trivial (i.e. X is a *bad orbit*, note that k is even in this case). Therefore, the local equivariant Floer cohomology retains information about the multiplicity k of the orbits, as well as the bad orbits. For instance, if one ignores the bad orbits, and considers the kernel of the localization at u , one discards all multiple orbits, and is left with the simple periodic orbits. One expects this to be helpful in answering questions related to simple periodic orbits (a major example of such a question is Hofer-Wysocki-Zehnder's two or infinity conjecture, which is recently proven in [CHHL23]). In a different direction, the same calculation shows that bad orbits do not contribute to the equivariant Floer cohomology when $2 \in R^*$ is invertible, and they behave differently than the good orbits when R is 2-local (e.g. $R = H\mathbb{F}_2$ or $K_2(n)$). Therefore, it is reasonable to anticipate that this calculation will be useful in addressing questions related to the existence of bad orbits.

Acknowledgements. The authors would like to thank Irakli Patchkoria, Tobias Barthel, Andrew Blumberg, Sanath Devalapurkar, Yash Deshmukh, Kaif Hilman, Jonas McCandless, Marco Volpe and Hiro Tanaka for useful discussions and comments. The authors would also like to thank Tim Large for a particularly inspiring seminar talk, from which they learned that equivariant localization works for Morava K-theory (see Example 2.13). This work was completed while the first author was visiting the Max Planck Institute for Mathematics, which he thanks for its hospitality. The first author was also partially supported by the National Science Foundation under Grant No. DMS-2305257 and by the Simons Collaboration on Homological Mirror symmetry; and the second author was supported by ERC Starting Grant 850713 (Homological mirror symmetry, Hodge theory, and symplectic topology), and the Simons Collaboration on Homological Mirror Symmetry.

2. FILTERED EQUIVARIANT SPECTRA AND COMPLETED TATE COHOMOLOGY

2.1. Reminders in homotopy theory.

2.1.1. *Conventions.* Let Sp be the infinity category of spectra. This is a symmetric monoidal category; we typically denote the monoidal product (i.e. the smash product) by $- \otimes -$ and denote the internal hom (i.e. the function spectrum) by $F(-, -)$. A *ring spectrum* (or *multiplicative cohomology theory*) is a spectrum $R \in \mathrm{Sp}$ equipped with a multiplication $R \otimes R \rightarrow R$ which is homotopically associative and unital. Following the usual convention, we will not distinguish between generalized cohomology theories and their representing spectra.

Let Top be the category of (compactly generated, weakly Hausdorff) topological spaces and let Top_* be the category of pointed spaces. There are natural functors $(-)_+ : \mathrm{Top} \rightarrow \mathrm{Top}_*$ and $\Sigma^\infty(-) : \mathrm{Top}_* \rightarrow \mathrm{Sp}$ (left adjoint to the forgetful functor, resp. to $\Omega^\infty(-)$). We typically denote their composition by $\Sigma^\infty(-)_+$.

Given a spectrum X , its R -cohomology is the graded abelian group $R^*(X) := \pi_{-*}F(X, R)$. If $X \in \mathrm{Top}_*$, we let $R^*(X) := R^*(\Sigma^\infty X)$ and let $\tilde{R}^*(X) := \ker(R^*(X) \rightarrow R^*(*))$. If $X \in \mathrm{Top}$, then $R^*(X) := R^*(X_+)$ and $\tilde{R}^*(X) := \tilde{R}^*(X_+)$. We let $R^* := R^*(\mathbb{S}) = \pi_{-*}R$ be the *coefficients* of the spectrum R . We adopt analogous conventions for R -homology.

2.1.2. *Borel G -spectra.* Let G be a compact Lie group. The functor category $\mathrm{Fun}(BG, \mathrm{Sp}) := \mathrm{Sp}^{BG}$ shall be called the category of *Borel G -spectra*. It inherits a symmetric monoidal structure from Sp , where the monoidal unit is the sphere with the trivial G -action. We will often refer to the objects of Sp^{BG} as *G -equivariant spectra*.²

Given a Borel G -spectrum $X : BG \rightarrow \mathrm{Sp}$, we let

$$(2.1) \quad X_{hG} := \mathrm{colim}_{BG} X \in \mathrm{Sp} \quad X^{hG} := \lim_{BG} X \in \mathrm{Sp}$$

²There are other non-equivalent objects in homotopy theory which also deserve this name, but they will never be considered in this paper.

be respectively the *homotopy quotient* (or *homotopy orbits*) and the *homotopy fixed points*. More concretely, X_{hG} can be defined as $X \wedge_G EG_+$, and X^{hG} can be defined as $Map^G(EG_+, X)$. It is often also useful to note that $(-)_hG$ and $(-)^{hG}$ are the left (resp. right) adjoints of the pullback $\pi^* : \mathrm{Sp} \rightarrow \mathrm{Sp}^{BG}$ of the map $\pi : BG \rightarrow *$ giving a spectrum the trivial G action. Note finally the identity $F(X_{hG}, R) \simeq F(X, R)^{hG}$, which follows from (2.1) by abstract nonsense.

Lemma 2.1. *If R is an E_k ring spectrum, $k \geq 1$, then $F(\mathbb{S}_{hG}, R) \simeq F(\Sigma^\infty BG_+, R) = R^*(BG)$ is also an E_k ring spectrum. For any $X \in \mathrm{Sp}^{BG}$, $F(X_{hG}, R)$ is a module over $F(\mathbb{S}_{hG}, R)$.*

Proof. By virtue of being the monoidal unit, the sphere \mathbb{S} with trivial G -action forms a co-commutative co-algebra. By virtue of being a left-adjoint of a lax monoidal functor, $(-)_hG$ is oplax monoidal, i.e. there are natural maps $(X \otimes Y)_{hG} \rightarrow X_{hG} \otimes Y_{hG}$ (satisfying some higher coherence conditions [HHLN23, Tor23], although for our purposes standard oplax monoidality in the homotopy category is sufficient). Hence \mathbb{S}_{hG} is a co-commutative co-algebra, and we have canonical maps $F(\mathbb{S}_{hG}, R) \otimes F(\mathbb{S}_{hG}, R) \rightarrow F(\mathbb{S}_{hG} \otimes \mathbb{S}_{hG}, R \otimes R) \rightarrow F(\mathbb{S}_{hG}, R \otimes R) \rightarrow F(\mathbb{S}_{hG}, R)$. Similarly, X_{hG} is a co-module over \mathbb{S}_{hG} , leading to a $F(\mathbb{S}_{hG}, R)$ -module structure on $F(X_{hG}, R)$. \square

By similar considerations, one can argue that if Y is a G -space, then the diagonal makes $\Sigma^\infty Y_+$ into a co-algebra, and hence $F(Y_{hG}, R)$ is an algebra.

The cohomology $R^*(X_{hG}) = \pi_* F(X_{hG}, R)$ is the R -valued *equivariant cohomology*. Note that if X is a *space*, then $\Sigma^\infty(X_{hG})_+ = (\Sigma^\infty X_+)_{hG}$ and we have $R^*((\Sigma^\infty X_+)_{hG}) = R^*(X_{hG}) = R^*(EG \times_G X)$.

2.1.3. Complex orientations.

Definition 2.2. A ring spectrum is said to be *complex orientable* if the pull-back $R^*(BS^1) = R^*(\mathbb{C}\mathbb{P}^\infty) \rightarrow R^*(\mathbb{C}\mathbb{P}^1)$ is surjective. A complex orientation is a lift of $1 \in \pi_0(\mathbb{C}\mathbb{P}^1) \cong \tilde{R}^2(\mathbb{C}\mathbb{P}^1)$ to $R^2(BS^1)$. A choice of such a lift specifies an isomorphism $R^*(BS^1) \cong R^*[[u]]$, where u is a variable of degree -2 .

A complex orientation defines a formal group law $F_R \in R^*[[u_1, u_2]]$. Concretely, F_R is the image of u under the pull-back $R^*[[u]] = R^*(BS^1) \rightarrow R^*[[u_1, u_2]] = R^*(BS^1 \times BS^1)$ of the classifying map $BS^1 \times BS^1 \rightarrow BS^1$ corresponding to the tensor product of universal line bundles. One can think of $u \in R^*[[u]] \cong R^*(BS^1)$ as the analogue of the first Chern class in the generalized cohomology theory R , and F_R is the formula for the Chern class of tensor products of line bundles.

Let $[n]_R(u) \in R^*[[u]]$ denote the n^{th} iterate of u under the formal group law F_R (i.e. $[2]_R(u) = F_R(u, u)$, $[3]_R(u) = F_R([2]_R(u), u)$, \dots).

Lemma 2.3. *Let $Y := S^1/C_k$ denote the homogeneous S^1 -space with stabilizer given by the cyclic group C_k , $k > 0$. Then $[k]_R(u) \in R^*(\mathbb{S}_{hS^1}) = R^*[[u]]$ acts by zero on $R^*(Y_{hS^1})$, under the $R^*(\mathbb{S}_{hS^1})$ -algebra structure induced by the map of G -spaces $Y \rightarrow *$.*

Proof. Note that $Y_{hS^1} \simeq BC_k$. It follows from the Gysin sequence

$$(2.2) \quad R^*[[u]] \xrightarrow{[k]_R(u)} R^*[[u]] \rightarrow R^*(BC_k) \rightarrow$$

associated to the fibration $S^1 \hookrightarrow BC_k \rightarrow BS^1$ that $[k]_R(u)$ pulls back to zero in $R^*(Y_{hS^1})$. \square

2.2. Filtered objects of stable ∞ -categories. In this section, we define the filtrations on objects of stable ∞ -categories. See in particular Definition 2.5 which will be crucial in the sequel.

Definition 2.4. . Let \mathcal{C} be a stable ∞ category. A *filtration* on an object $X \in \mathcal{C}$ is a sequence $X_1 \rightarrow X_2 \rightarrow \dots$ and an equivalence $X \simeq \mathrm{colim} X_i$. An object equipped with a filtration is said to be *filtered*. A morphism of filtered objects $X \rightarrow Y$ is a morphism of diagrams $X_i \rightarrow Y_i$ compatible with the identifications $X \simeq \mathrm{colim} X_i$ and $Y \simeq \mathrm{colim} Y_i$. We let $\tilde{\mathcal{C}}_{fil}$ be the category over \mathcal{C} consisting of filtered objects.

For a filtration $X \simeq \operatorname{colim}(X_1 \rightarrow X_2 \rightarrow \dots)$ and a sequence $1 \leq i_1 < i_2 < \dots$, there is a corresponding *subfiltration* $X \simeq \operatorname{colim}(X_{i_1} \rightarrow X_{i_2} \rightarrow \dots)$. We want to consider a subfiltration to be an equivalent filtration. To make this precise, consider *the tautological morphism* given at each level by $X_k \rightarrow X_{i_k}$.

Definition 2.5. Let \mathcal{C}_{fil} be obtained from $\tilde{\mathcal{C}}_{fil}$ by localizing at the tautological morphisms corresponding to every subfiltration. Two filtered objects are said to be *equivalent* if they are equivalent in \mathcal{C}_{fil} .

We call a filtration *trivial*, resp. *finite* if all of $X_k \rightarrow X_{k+1}$, resp. all but finitely many of $X_k \rightarrow X_{k+1}$ are equivalences. Observe that all finite filtrations are equivalent to the trivial one.

There is a forgetful functor $\mathcal{C}_{fil} \rightarrow \mathcal{C}$. However, objects in \mathcal{C} may have non-equivalent lifts to \mathcal{C}_{fil} . In other words, the objects may have non-equivalent filtrations.

Example 2.6. Consider the sequence of $\mathbb{Q}[u]$ -modules given by $\dots \rightarrow \mathbb{Q}[u]/(u^3) \rightarrow \mathbb{Q}[u]/(u^2) \rightarrow \mathbb{Q}[u]/(u)$. This is a filtration on $\mathbb{Q}[[u]]$ in the category $\mathcal{C} = \operatorname{Mod}(\mathbb{Q}[u])^{op}$. If this was equivalent to the trivial filtration, we would have

$$(2.3) \quad 0 = \lim_k (\mathbb{Q}[u]/(u^k) \otimes_{\mathbb{Q}[u]}^L \mathbb{Q}[u, u^{-1}]) = \lim_k (\mathbb{Q}[[u]] \otimes_{\mathbb{Q}[u]}^L \mathbb{Q}[u, u^{-1}]) \neq 0$$

Example 2.7. Let $\mathcal{C} = Sp$, and consider the filtration on $\Sigma^\infty \mathbb{C}P_+^\infty$ given by $\Sigma^\infty \mathbb{C}P_+^1 \rightarrow \Sigma^\infty \mathbb{C}P_+^2 \rightarrow \dots$. This is different from the trivial filtration $\Sigma^\infty \mathbb{C}P_+^\infty \rightarrow \Sigma^\infty \mathbb{C}P_+^\infty \rightarrow \dots$ as each level is a compact object of \mathcal{C} in the former, but not in the latter. A weaker property that an equivalence needs to respect is the compactness of $\operatorname{cofib}(X_i \rightarrow X_{i+1})$ for all $i \gg 0$.

A concrete way to show that two filtrations $X_1 \rightarrow X_2 \rightarrow \dots \rightarrow X$ and $X'_1 \rightarrow X'_2 \rightarrow \dots \rightarrow X$ are equivalent is to find sequences $1 \leq i_1 \leq i_2 \dots$, $1 \leq j_1 \leq j_2 \dots$, and maps $X'_k \rightarrow X_{i_k}$, $X_k \rightarrow X'_{j_k}$ of filtrations such that the two sided compositions are equivalent to the natural subfiltration maps $X_k \rightarrow X_{i_{j_k}}$ and $X'_k \rightarrow X'_{j_{i_k}}$. One can also show that, for any sequence $1 \leq i_1 \leq i_2 \leq \dots$ that is not strictly monotone, but $i_k \rightarrow \infty$, $X_{i_1} \rightarrow X_{i_2} \rightarrow \dots$ is still equivalent to $X_1 \rightarrow X_2 \rightarrow \dots$ (by a zigzag through a common subfiltration). Throughout the paper, unless stated otherwise, we only consider the filtrations $X_1 \rightarrow X_2 \rightarrow \dots \rightarrow X$ such that all of X_i are of finite type.

2.3. Completed Tate cohomology.

2.3.1. *The definition.* Given a complex-oriented cohomology theory R , we define

$$(2.4) \quad \mathfrak{F}_R := \{[1]_R(u), [2]_R(u), [3]_R(u), \dots\} \subset R^*[[u]].$$

By definition, $[k]_R(u)$ is homogeneous of degree -2 .

Definition 2.8. Let $X \in \operatorname{Sp}_{fil}^{BS^1}$ be a filtered S^1 -equivariant spectrum. Given a complex-oriented cohomology theory R , define the *completed Tate cohomology* of X to be the limit

$$(2.5) \quad \hat{R}_{S^1}^*(X) := \lim_k \mathfrak{F}_R^{-1} R^*((X_k)_{hS^1}) = \lim_k R^*((X_k)_{hS^1})[[1]_R(u)^{-1}, [2]_R(u)^{-1}, [3]_R(u)^{-1} \dots]$$

where the localization is completed with respect to u -adic topology.

Note 2.9. Recall that the completion of a filtered commutative group A is defined to be $\lim_p A/F^p A$. This is equivalent to the topological completion for the topology defined by the neighborhood basis $\{F^p A\}$. If A is graded, one takes the limit in the graded category too; thus, the completion is still graded. If A is a topological algebra and $S \subset A$, the localization $S^{-1}A$ inherits a topology. For example, $\mathbb{Q}[q]$ is filtered by $q^p \mathbb{Q}[q]$, $p \geq 0$, and this topology descends to $\mathbb{Q}[q, q^{-1}]$. Notice however, it is not given by $q^p \mathbb{Q}[q, q^{-1}]$, $p \geq 0$, but rather by $q^p \mathbb{Q}[q] \subset \mathbb{Q}[q, q^{-1}]$, which also is the topology corresponding to the u -adic norm. The completion with respect to this topology is given by $\mathbb{Q}((u))$.

Observe

- $\widehat{R}_{S^1}^*(X)$ carries the structure of a complete $\widehat{R}_{S^1}^*(\mathbb{S})$ -module.
- $\widehat{R}_{S^1}^*(X)$ is functorial in X and sends finite colimits to limits.
- $\widehat{R}_{S^1}^*(X)$ is functorial in R . In other words, if $R \rightarrow R'$ is an homomorphism of complex oriented ring spectra (respecting complex orientations), then there is a natural map $\widehat{R}_{S^1}^*(X) \rightarrow \widehat{R}'_{S^1}^*(X)$.

Be warned that localization does not commute with infinite limits, so one cannot in general pass the limit inside the localization in (2.5) (this is ultimately what makes the invariant useful). On the other hand, if the filtration is trivial/finite, then there is not need to pass to the limit. Hence we will also use the notation $\widehat{R}_{S^1}^*(X_k)$ to refer to the terms in the limit in (2.5).

2.3.2. Completed Tate cohomology of the building blocks. In this section, we compute the completed Tate cohomology of Thom spectra of bundles over homogeneous spaces over S^1 , as well as spaces with trivial S^1 -action. These are the basic building blocks of the spectra we are interested in. First

Lemma 2.10. *Consider S^1/C_k , the homogeneous S^1 -space with stabilizer given by the cyclic group $C_k, k > 0$. Consider a virtual equivariant vector bundle over S^1/C_k and let X denote the corresponding Thom spectrum. Then $\widehat{R}_{S^1}^*(X) = 0$.*

Proof. Write Y for the space S^1/C_k . A virtual equivariant vector bundle can be written as $E - \underline{V}$, where E is an equivariant vector bundle and \underline{V} is a topologically trivial equivariant bundle corresponding to a representation V of S^1 , see [Seg68]. The Thom spectrum is given by

$$(2.6) \quad X = \Sigma_V^{-1} Y^E = \mathbb{S}^{-V} \otimes Y^E$$

Similarly to Lemma 2.1, X is a comodule over $\Sigma^\infty Y_+$. More precisely, the diagonal $Y \rightarrow Y \times Y$ induces a map $Y^E \rightarrow Y^E \otimes \Sigma^\infty Y_+$ (to see this note that $E \boxplus 0$ pulls back to E under the diagonal and use the functoriality of Thom spectra). This immediately extends to a coassociative coproduct

$$(2.7) \quad X = \mathbb{S}^{-V} \otimes Y^E \rightarrow \mathbb{S}^{-V} \otimes Y^E \otimes \Sigma^\infty Y_+ = X \otimes \Sigma^\infty Y_+$$

and as in the proof of Lemma 2.1, this induces an $R^*(Y_{hS^1})$ -module structure on $R^*(X_{hS^1})$. On the other hand, $Y_{hS^1} = BC_k$ and Lemma 2.3 ensures that $[k]_R(u) \in R^*[[u]]$ acts on $R^*(Y_{hS^1})$ by 0. As a result, the $R^*[[u]]$ -module structure on $R^*(X_{hS^1})$ refines to a $R^*[[u]]/([k]_R(u))$ -module structure, and inverting $[k]_R(u)$ kills the group, and $\mathfrak{F}_R^{-1} R^*(X_{hS^1}) = 0$. \square

We also have

Lemma 2.11. *If X is a finite spectrum with trivial S^1 -action, then*

$$(2.8) \quad \widehat{R}_{S^1}^*(X) \simeq \mathfrak{F}_R^{-1} R^*(X)[[u]] = R^*(X)[[u]][[1]_R(u)^{-1}, [2]_R(u)^{-1}, [3]_R(u)^{-1} \dots]$$

where the localization is in the u -completed as before.

More concisely, we can write this as $\widehat{R}_{S^1}^*(X) \simeq \widehat{R}_{S^1}^*(\mathbb{S}) \otimes_{R^*} R^*(X)$. Since X is assumed to be finite, tensor product is automatically complete. Otherwise, one would need to use the completed tensor product.

Proof. It is easy to check that $R^*(X_{hS^1}) = R^*(X \wedge \Sigma^\infty BS^1_+) \simeq R^*(X)[[u]]$. The result follows from this. \square

2.3.3. Computations for the trivial action. In this section, we compute the completed Tate cohomology $\widehat{R}_{S^1}^*(X)$ of a finite spectrum X with trivial action and filtration for various R .

Example 2.12. Assume $R = HA$, the Eilenberg–Maclane spectrum of a commutative ring A . Then, $R^* = A$ and $F_R(u_1, u_2) = u_1 + u_2$, the additive formal group law. Therefore, $[n]_R(u) = nu$, and inverting it is the same as inverting n and u . As a result, if X is endowed with a trivial action

and filtration, then $\widehat{R}_{S^1}^*(X) = H^*(X, A \otimes_{\mathbb{Z}} \mathbb{Q})((u))$. In particular, if $n = 0$ in A , for some $n \geq 2$, then $\widehat{R}_{S^1}^*(X) = 0$. On the other hand, if $R = H\mathbb{Q}$ or $H\mathbb{Z}$, then $\widehat{R}_{S^1}^*(X) = H^*(X, \mathbb{Q})((u))$.

If in (2.5) we were not taking the u -adically completed localization, $\widehat{H\mathbb{Z}}_{S^1}^*(X)$ and $H^*(X, \mathbb{Q})((u))$ would match only after u -adically completing the former.

An important example is the following:

Example 2.13. Let $R = K_p(n)$ be a Morava K -theory. Then $R^* = \mathbb{F}_p[v_n, v_n^{-1}]$. As $[k]_R(u)$ is always of the form $ku + h.o.t.$, when k is coprime with p , $[k]_R(u)$ is u multiplied with a unit. Moreover, $[p]_R(u) = v_n u^{p^n}$ (see [Wur91, Theorem 1.3] for instance). Hence, by iteration, we see that $[k]_R(u)$ is the same as a power of u multiplied by a unit (if $p^l | k$, $[k]_R(u) = \frac{k}{p^l} v_n^{\frac{p^{nl}-1}{p-1}} u^{p^{nl}} + h.o.t.$). Therefore inverting all $[k]_R(u)$ is equivalent to inverting u . In other words, for X with a trivial action and trivial filtration $\widehat{K_p(n)}_{S^1}^*(X) = K_p(n)^*(X)((u))$.

Example 2.14. More generally, if p is nilpotent in R^* and F_R has height exactly k , i.e. $[p]_R(u)$ is u^{p^k} multiplied by a unit, then $\widehat{R}_{S^1}^*(X) = R^*(X)((u))$.

Example 2.15. Let $R = k_p(n)$ denote the connective Morava K -theory, which satisfies $R^* = \mathbb{F}_p[v_n]$ and $[p]_R(u) = v_n u^{p^n}$. The calculation of $[k]_R(u)$ is the same as in Example 2.13, but this time inverting these elements require inverting u and v_n . Therefore, $\widehat{k_p(n)}_{S^1}^*(X) = K_p(n)^*(X)((u))$, which is analogous to $H\mathbb{Z}$ leading to rational cohomology in Example 2.12.

The following example will also be important in the subsequent sections:

Example 2.16. Let $\widetilde{K}_p(n)$ be the integral Morava K -theory, i.e. the complex oriented spectrum satisfying $\widetilde{K}_p(n)^* = \mathbb{Z}_p[v_n, v_n^{-1}]$ and $[-p]_{\widetilde{K}_p(n)}(u) = -pu + v_n u^{p^n}$. Let $K_{p^k}(n)$ denote its quotient by p^k . See [AMS21, Proposition 5.14]. For $R = K_{p^k}(n)$, the same formula $[-p]_R(u) = -pu + v_n u^{p^n}$ holds. For $p \nmid m$, $[mp^l]_R(u)$ is of the form $u^{p^{nl}} + O(p) + o(u^{p^{nl}})$ times a unit, where $O(p)$ denotes a multiple of p and $o(u^{p^{nl}})$ denotes the sum of the terms such that the exponent of u is larger than p^{nl} . One can also write this as $u^{p^{nl}} + O(p)$ times a unit (as $u^{p^{nl}} + o(u^{p^{nl}})$ is $u^{p^{nl}}$ times a unit). As p is nilpotent, after inverting $u = [1]_R(u)$, elements of the form $u^{p^{nl}} + O(p)$ automatically become units. As a result, for X with a trivial action and filtration, $\widehat{R}_{S^1}^*(X) = R^*(X)((u))$. The same holds for $R = \widetilde{K}_p(n)$ after p -completing both sides, i.e. $\widehat{R}_{S^1}^*(X)_p^\wedge = R^*(X)((u))_p^\wedge$.

The following example will be important in recovering information about complex K -theory from Floer theory:

Example 2.17. Recall that KU , the complex K -theory spectrum, has $KU^* = \mathbb{Z}[\beta, \beta^{-1}]$, where $|\beta| = 2$ and $F_{KU}(u_1, u_2) = u_1 + u_2 + u_1 u_2 = (1 + u_1)(1 + u_2) - 1$, the *multiplicative formal group law*. Let p be a prime and $R = KU/p^k$ so that $R^* = \mathbb{Z}/p^k[\beta, \beta^{-1}]$ and $[p^l]_R(u) = \binom{p^l}{1}u + \binom{p^l}{2}u^2 + \cdots + u^{p^l}$, which implies $[mp^l]_R(u) = u^{p^l} + O(p)$ times a unit as in Example 2.16, for $p \nmid m$. As a result, inverting u suffices as before and for X with a trivial action and filtration, $\widehat{R}_{S^1}^*(X) = R^*(X)((u))$. The same holds for KU after simultaneous completion of both sides at each p , i.e. $\widehat{KU}_{S^1}^*(X)^\wedge = KU^*(X)((u))^\wedge$, where for an abelian group A , A^\wedge denotes the inverse limit of A/nA , $n \in \mathbb{N}$.

One can make such concrete calculations for other complex oriented spectra, such as Brown–Peterson spectrum (or its variants such as its localization at $v_1 \in BP^*$). This could possibly let one to recover information about MU from Floer theory.

3. S^1 -EQUIVARIANT SPECTRAL SYMPLECTIC COHOMOLOGY

3.1. Conventions. Let $(M, \omega = d\lambda)$ be an exact symplectic manifold. Set $S^1 = \mathbb{R}/\mathbb{Z}$. Given a Hamiltonian $H : S^1 \times M \rightarrow \mathbb{R}$, we write $H_t(-) = H(t, -)$. The Hamiltonian vector field is defined by $\omega(X_t, -) = -dH_t$. The H -perturbed action functional is the map

$$(3.1) \quad \mathcal{A}_H : \mathcal{L}M \rightarrow \mathbb{R}$$

$$(3.2) \quad x \mapsto \int_{S^1} (x^* \lambda - H_t(x(t))) dt$$

This is the opposite of conventions of [Abo15, Sei08]. Given an auxiliary (possibly time-dependent) almost-complex structure J , we shall consider solutions $u : \mathbb{R} \times S^1 \rightarrow M$ to Floer's equation

$$(3.3) \quad \partial_s u + J(\partial_t u - X_{H_t}) = 0$$

which are asymptotic to closed trajectories $x^\pm = \lim_{s \rightarrow \pm\infty} u(s, -)$ of the Hamiltonian H_t . We call x^+ the input and x^- the output. Solutions to Floer's equation (3.3) are formally negative gradient trajectories of (3.1), and we have the identity $\mathcal{A}_H(x_-) \leq \mathcal{A}_H(x_+)$. In other words, the Floer trajectories decrease the action. More generally, if $(H^s)_{s \in \mathbb{R}}$ is a family of Hamiltonians which is independent of s for $s \in \mathbb{R} - (0, 1)$ and satisfies $\partial_s H^s \leq 0$, we can generalize (3.3) to the *continuation equation* $\partial_s u + J(\partial_t u - X_{H_t^s}) = 0$ and we have $\mathcal{A}_{H_0}(x_-) \leq \mathcal{A}_{H_1}(x_+)$, i.e. the continuation trajectories also decrease the action.

3.2. Liouville manifolds. Recall that a *Liouville manifold* (M, λ) is an exact symplectic manifold with the property that there exists an embedding

$$(3.4) \quad \iota : (S^+(\partial_\infty M), \lambda_{can}) \hookrightarrow (M, \lambda), \iota^* \lambda = \lambda_{can}$$

of the positive symplectization of a co-oriented contact manifold $(\partial_\infty M, \xi_\infty)$ which covers a neighborhood of infinity.³ A choice of contact form $\ker \lambda_\infty = \xi_\infty$ on $\partial_\infty M$ induces a decomposition

$$(3.5) \quad M = \overline{M} \cup_{\partial \overline{M}} [1, \infty) \times \partial_\infty M,$$

where λ restricts on $[1, \infty) \times \partial_\infty M$ to $r\lambda_\infty$. The coordinate $r \in [1, \infty)$ shall be called the *Liouville parameter* and the vector field Z on M defined by the property $i_Z d\lambda = \lambda$ shall be called the *Liouville vector field*. Note that for a given Liouville manifold (M, λ) , the data of contact form $\ker \lambda_\infty = \xi_\infty$ is equivalent to the data of a contact hypersurface transverse to Z .

A Hamiltonian $H : M \rightarrow \mathbb{R}$ is said to be *linear at infinity* if $ZH = H$ outside a compact set. Any Hamiltonian which satisfies $H(r, m) = Cr$ outside a compact set, with respect to the (non-canonical) decomposition (3.5), is manifestly linear at infinity. The constant C is called the *slope* of H . We emphasize that the notion of slope only makes sense once we have fixed a decomposition (3.5), i.e. a contact form $\ker \lambda_\infty = \xi_\infty$. For instance, scaling λ_∞ scales the slope by a corresponding amount.

If R is the Reeb vector field on $(\partial_\infty M, \lambda_\infty)$ and $H(r, m) = h(r)$, then $X_H = h'(r)R$ on the collar $[1, \infty) \times \partial_\infty M$. So closed time-1 orbits of H on the level $\{r = r_0\} \subset [1, \infty) \times \partial_\infty M$ correspond to closed Reeb orbits of length $h'(r_0)$. The action of such an orbit is $r_0 h'(r_0) - h(r_0)$.

A Liouville manifold is said to be *stably framed* if it is equipped with a fixed isomorphism $T_M \oplus \mathbb{C}^k \cong \mathbb{C}^{N+k}$ of symplectic vector bundles, for some $k \geq 0$. *For the remainder of this paper, we restrict our attention to stably framed Liouville manifolds.* For this reason, we will typically drop the adjective ‘‘stably framed’’. The stable framing will be used to define $SH(M, \mathbb{S})$ and $SH_S(M, \mathbb{S})$. In fact, we will mostly need symplectic cohomology with coefficients in complex oriented spectra, an invariant which can be defined under weaker assumptions on M .

³Given a contact manifold (Y, ξ) , its symplectization $SY := \{\alpha \in T^*Y \mid \alpha(\xi) \neq 0\}$ is an exact symplectic manifold with respect to the canonical 1-form λ_{can} . We say that (Y, ξ) is co-orientable if SY is disconnected, and a co-orientation amounts to a choice of connected component S^+Y which we call the positive symplectization.

3.3. Review of (spectral) symplectic cohomology. The construction of symplectic cohomology over the sphere spectrum was first carried out by Large [Lar21]. However, our presentation will rather follow [CK23] which follows different conventions.

Let M be a Liouville manifold. Let $H : S^1 \times M \rightarrow \mathbb{R}$ be a Hamiltonian which is non-degenerate and linear at infinity. Let $J : S^1 \rightarrow \mathcal{J}(M)$ be a generic family of cylindrical almost-complex structures. To this data, one can associate a *flow category* $\mathcal{M}_{H,J}$ such that

- $ob(\mathcal{M}_{H,J})$ is the set of 1-periodic orbits of H
- the morphisms $\mathcal{M}_{H,J}(x, y)$ is given by the moduli of Floer trajectories from x to y and it forms a manifold with corners with a stable trivialization of its tangent bundle
- the composition maps are smooth embeddings into boundary strata, which is covered by the images of the composition maps

To the flow category $\mathcal{M}_{H,J}$, one associates a spectrum $HF(H; \mathbb{S}) := |\mathcal{M}_{H,J}|$ called the *geometric realization* (we omit J from the notation). To define this, one uses Pontryagin–Thom collapse and obtains a “chain complex in spectra”. Then there is a realization construction for such complexes.

Similarly to classical Floer theory, given a generic homotopy $(H_s, J_s)_{s \in \mathbb{R}}$ where $\partial_s H_s \leq 0$, $(H_s, J_s) = (H, J)$ for $s \gg 0$ and $(H_s, J_s) = (H', J')$ for $s \ll 0$, one has continuation maps $HF(H; \mathbb{S}) \rightarrow HF(H'; \mathbb{S})$. To define these maps, one assembles the continuation trajectories into a *framed flow bimodule* over $\mathcal{M}_{H,J} - \mathcal{M}_{H',J'}$, inducing a map of geometric realizations.

Consider a sequence of Hamiltonians H_1, H_2, \dots , such that $H_{i+1} > H_i$ outside a compact subset of M and such that the slope goes to infinity. We obtain a sequence $HF(H_1, \mathbb{S}) \rightarrow HF(H_2, \mathbb{S}) \rightarrow \dots$, and let the *spectral symplectic cohomology* $SH(M, \mathbb{S})$ be the homotopy colimit of this sequence. By construction, it is a filtered spectrum. Any other choice of data (such as Hamiltonians H_i , almost complex structures, continuation data) gives a homotopy equivalent spectrum, and an equivalent filtered spectrum in the sense of Definition 2.5.

3.4. Review of S^1 -equivariant spectral symplectic cohomology. In [CK23], the authors extended the realization construction for flow categories to the Morse–Bott and equivariant settings, and used this to define S^1 -equivariant versions of $HF(H, \mathbb{S})$ and $SH(M, \mathbb{S})$ by using what is often referred as the Borel construction.

3.4.1. Morse–Bott flow categories. A (framed) Morse–Bott flow category \mathcal{M} is similar to an ordinary flow category, except that one allows $ob(\mathcal{M})$ to be a finite disjoint union of smooth closed manifolds. If $X, Y \subset ob(\mathcal{M})$ denote two components and $\mathcal{M}(X, Y)$ denotes the morphisms from a point on X to one Y . We assume the domain and target maps $\mathcal{M}(X, Y) \rightarrow X, Y$ are smooth. The motivating example is the category associated to a Morse–Bott function $f : N \rightarrow \mathbb{R}$: the object space is the union of critical manifolds of f and the morphisms are given by the broken negative gradient trajectories (with respect to a generic metric). The stable framing condition is generalized as a framing of the relative tangent bundle $T_{\mathcal{M}(X,Y)} - T_X$ twisted by virtual bundles on X and Y . See [CK23] for more details.

Given the data of a framed flow category, [CK23] explains how to associate a spectrum $|\mathcal{M}|$, also called the *geometric realization*. If the category and the framings are equivariant with respect to a compact group action, $|\mathcal{M}|$ is also a genuine equivariant spectrum.

3.4.2. Data for the Borel construction. Let M be a Liouville manifold and fix a non-degenerate contact form λ_∞ on $\partial_\infty M$. Let $a > 0$ be a real number which is not equal to the length of a Reeb orbit for λ_∞ .

Let $S := S^\infty = \{(z_1, z_2, \dots) \mid \sum_i |z_i|^2 = 1, z_i \in \mathbb{C}\}$. Then S admits a free S^1 -action with quotient $\pi : S \rightarrow \mathbb{C}\mathbb{P}^\infty$. There is a standard Morse–Bott function $\tilde{f} : S \rightarrow \mathbb{R}$, $\tilde{f}(z_1, z_2, \dots) = \sum_i i|z_i|$. It descends to a Morse function on $\mathbb{C}\mathbb{P}^\infty$. We consider a generic metric on $\mathbb{C}\mathbb{P}^\infty$ and choose an equivariant lift to S^∞ . The lift determines a connection on the principal S^1 -bundle $S^\infty \rightarrow \mathbb{C}\mathbb{P}^\infty$

and we assume this is flat near the critical sets. Therefore, for every critical set $S^1 \cong X \subset S^\infty$, we have a canonical local trivialization $U \times X \subset S^\infty \rightarrow \mathbb{C}\mathbb{P}^\infty \supset U$.

Let $H : S \times S^1 \times M \rightarrow \mathbb{R}$ be a Hamiltonian having the following properties:

- H is invariant under the diagonal S^1 action.
- $H_z = H_z(-, -) : S^1 \times M \rightarrow \mathbb{R}$ has slope $a > 0$ for all $z \in S$, and H_z is non-degenerate for all $z \in \text{crit}(\tilde{f})$.
- in a canonically trivialized neighborhood $U \times X$ of a critical set H_z does not depend on $z \in U$. In other words, dH kills the horizontal directions.

Similarly, we let $J : S \times S^1 \rightarrow \text{End}(TM)$ be a generic family of almost-complex structures invariant under the diagonal S^1 action, cylindrical outside a (uniform) compact subset of M , and constant near the critical sets.

To this data, we associate an S^1 -equivariant (framed) Morse–Bott flow category $\mathcal{M}_{\tilde{f}, H}$ (J is omitted from the notation). The object space is given by pairs (a, x) , $a \in \text{crit}(\tilde{f})$, $x \in \text{orb}(H_a)$. They form a disjoint union of S^1 -torsors. The morphisms are given by the pairs $(\gamma, u) : \mathbb{R} \times S^1 \rightarrow S^\infty \times M$ satisfying the parameterized Floer equation

$$(3.6) \quad \begin{cases} \dot{\gamma} + \nabla_{\tilde{g}} \tilde{f}(\gamma) & = 0 \\ \partial_s u + J_{\gamma(s)}(\partial_t u - X_{H_{\gamma(s)}}) & = 0 \end{cases}$$

asymptotic to (a_\pm, x_\pm) . We define $HF_S(H, \mathbb{S}) := |\mathcal{M}_{\tilde{f}, H}|$.

Similar to before, there are equivariant continuation maps $HF_S(H, \mathbb{S}) \rightarrow HF_S(H', \mathbb{S})$, and we define $SH_S(M, \mathbb{S})$ to be colimit of $HF_S(H, \mathbb{S})$ as the slope of H goes to infinity. We obtain a filtered equivariant spectrum, whose filtration is well-defined up to equivalence.

Remark 3.1. Remembering both the filtration and the circle action on $SH_S(M; \mathbb{S})$ is crucial for our purposes. Indeed, $SH_S(M \times \mathbb{C}; \mathbb{S}) \simeq 0$ as a spectrum, which implies the same as a Borel equivariant spectrum. On the other hand, if we remember the filtration, but forget about the equivariant structure, we expect the filtration on $SH_S(M \times \mathbb{C}) \simeq 0$ to be equivalent to the trivial one. This can presumably be established by the second argument of [Sei08, (3f)]. In contrast, our main result will imply that $SH_S(M; \mathbb{S})$ is never zero as an object of $\text{Sp}_{fil}^{BS^1}$, for any Liouville manifold M .

4. APPROXIMATELY AUTONOMOUS HAMILTONIANS AND THE COMPLETED TATE COHOMOLOGY OF $SH_S(M, \mathbb{S})$

4.1. The main theorem and its applications. The goal of this section is to prove the following theorem:

Theorem 4.1. *Given a complex oriented ring spectrum R ,*

$$(4.1) \quad \widehat{R}_{S^1}^*(SH_S(M, \mathbb{S})) \simeq \widehat{R}_{S^1}^*(\Sigma^\infty(M/\partial_\infty M)) \simeq \mathfrak{F}_R^{-1}R^*(M, \partial_\infty M)[[u]] \simeq \mathfrak{F}_R^{-1}R_{2n-*}(M)[[u]]$$

where $\Sigma^\infty(M/\partial_\infty M)$ is endowed with the trivial filtration and trivial S^1 -action.

Recall that $\mathfrak{F}_R = \{[k]_R(u) : k \in \mathbb{N}\}$ and $\mathfrak{F}_R^{-1}R^*(M, \partial_\infty M)[[u]]$ is $R^*(M, \partial_\infty M)[[u]]$ localized at $[k]_R(u)$, for every $k \geq 1$. The second isomorphism follows from Lemma 2.11 and the third follows from Poincaré duality (which holds for R as TM is complex oriented). As we already noted, we could equivalently write the last term as $R_{2n-*}(M) \otimes_{R^*} \widehat{R}_{S^1}^*(\mathbb{S})$.

Before moving onto the proof, we discuss some corollaries in the light of examples given in Section 2.

Corollary 4.2. *We have $\widehat{H}\mathbb{Q}_{S^1}^*(SH_S(M, \mathbb{S})) \simeq \widehat{H}\mathbb{Z}_{S^1}^*(SH_S(M, \mathbb{S})) \simeq H_{2n-*}(M, \mathbb{Q}((u)))$. On the other hand $\widehat{H}\mathbb{F}_{pS^1}^*(SH_S(M, \mathbb{S})) \simeq 0$.*

This follows immediately by applying Theorem 4.1 to Example 2.12. Note that for a commutative ring A , $\widehat{HA}_{S^1}^*(SH_S(M, \mathbb{S}))$ depends only on the S^1 -equivariant (co)filtered homotopy type of $SH^*(M, A)$, which does not require Floer homotopy theory to define. For $A = \mathbb{Q}$, this reaffirms the conclusion of [Zha19, ACF16] that the symplectic cohomology with its S^1 -action and filtration recovers the rational cohomology. For $A = \mathbb{Z}$ this is a generalization of the computations for a disc and annulus in [Zha19, Sec. 8]. In this way, we lose the torsion information. See also [ACF16, Sec. 5.1] for related computations.

Similarly, by applying Theorem 4.1 to Example 2.13 and Example 2.16, we obtain

Corollary 4.3. $\widehat{K}_{p^k}(m)_{S^1}^*(SH_S(M, \mathbb{S})) \simeq K_{p^k}(m)_{2n-*}(M)((u))$.

Our second main statement is the following theorem, which follows from Corollary 4.3:

Theorem 4.4. *For $2(p^m - 1) > \dim_{\mathbb{R}}(M)$ (or for $4(p^m - 1) > \dim_{\mathbb{R}}(M)$ for Weinstein M), the groups $\widehat{K}_{p^k}(m)_{S^1}^*(SH_S(M, \mathbb{S}))$ and $H_{2n-*}(M, \mathbb{Z}[v_m, v_m^{-1}]/p^k)((u))$ are isomorphic.*

Proof. We borrow the idea from [AMS21, Lemma 5.15]. If $2(p^m - 1) > \dim_{\mathbb{R}}(M)$ or $4(p^m - 1) > \dim_{\mathbb{R}}(M)$ and M is Weinstein, $|v_m| = 2(p^m - 1)$ is larger than the range of the cohomology of M . Therefore the Atiyah–Hirzebruch spectral sequence degenerates and $K_{p^k}(m)^*(M, \partial_{\infty}M) \simeq H^*(M, \partial_{\infty}M, \mathbb{Z}[v_m, v_m^{-1}]/p^k)$. The theorem now follows from Corollary 4.3 and Poincaré duality. \square

In particular, $\widehat{K}_p(m)_{S^1}^*(SH_S(M, \mathbb{S})) \cong H_{2n-*}(M, \mathbb{F}_p[v_m, v_m^{-1}]((u)))$, whose rank over $\mathbb{F}_p[v_m, v_m^{-1}]((u))$ gives the total dimension of $H_{2n-*}(M, \mathbb{F}_p)$. However, one can recover more information: the isomorphism in Theorem 4.4 is natural in k ; hence, the inverse limits of $\widehat{K}_{p^k}(m)_{S^1}^*(SH_S(M, \mathbb{S}))$ and $H_{2n-*}(M, \mathbb{Z}[v_m, v_m^{-1}]/p^k)((u))$ in k are also isomorphic. Express $H_{2n-*}(M, \mathbb{Z})$ as a direct sum of cyclic groups. In

$$\lim_k H_{2n-*}(M, \mathbb{Z}[v_m, v_m^{-1}]/p^k)((u))$$

the \mathbb{Z} summands turn into $\mathbb{Z}_p[v_m, v_m^{-1}]((u))^\wedge$ and \mathbb{Z}/p^l summands turn into

$$\mathbb{Z}_p[v_m, v_m^{-1}]((u))^\wedge/p^l \oplus \mathbb{Z}_p[v_m, v_m^{-1}]((u))^\wedge/p^l$$

as modules over $\mathbb{Z}_p[v_m, v_m^{-1}]((u))^\wedge$ (the p -adic completion of $\mathbb{Z}_p[v_m, v_m^{-1}]((u))$). As a result

Corollary 4.5. *The filtered S^1 -equivariant homotopy type of $SH_S(M, \mathbb{S})$ determines $H_*(M, \mathbb{Z})$.*

Notice that we only need the (co)filtered, S^1 -equivariant) generalized cohomology of $SH_S(M, \mathbb{S})$ with respect to Morava K -theories. Therefore less information than $SH_S(M, \mathbb{S})$ is actually required.

Another implication of Theorem 4.1 is the following (recall that for an abelian group A , $A^\wedge := \lim_{n \in \mathbb{N}} A/nA$):

Theorem 4.6. *After simultaneously completing at every prime p , the groups $\widehat{KU}_{S^1}^*(SH_S(M, \mathbb{S}))$ and $KU_{2n-*}(M)((u))$ become isomorphic. In other words, $\widehat{KU}_{S^1}^*(SH_S(M, \mathbb{S}))^\wedge \simeq KU_{2n-*}(M)((u))^\wedge$. As a result, the filtered S^1 -equivariant homotopy type of $SH_S(M, \mathbb{S})$ determines $KU_*(M)$ (and hence $KU^*(M)$ by the universal coefficients theorem).*

This result can be interpreted as saying that the (co)filtered equivariant homotopy type of “symplectic K -homology” determines the K -theory of the underlying manifold.

Proof. The isomorphism statement follows from Theorem 4.1 and Example 2.17. Moreover, as M is of finite type, each $KU_i(M)$ is a finitely generated abelian group. $KU_*(M)((u))$ can be identified with $KU_0(M)((\beta^{-1}u))$ at even degrees and with $KU_1(M)((\beta^{-1}u))$ at odd degrees. Each of these groups split as $(T \oplus F)((\beta^{-1}u))$, where T is the torsion part of $KU_0(M)$, resp. $KU_1(M)$, and F is a finitely generated free abelian group. The completion procedure leaves $T((\beta^{-1}u))$ the same, and

$F((\beta^{-1}u))$ turns into a finitely generated free module over $\mathbb{Z}((\beta^{-1}u))^\wedge$ of the same rank (and the former is still the torsion part). It is clear that T and F are uniquely determined. \square

Remark 4.7. As we mentioned, Theorem 4.6 is closely related to a string theory and mirror symmetry inspired conjecture of Treumann, which states that the Fukaya category determines the complex K -theory, see [Tre19]. This is clearly false for wrapped Fukaya categories, as there can be subcritical Weinstein manifolds with different complex K -theories; however, Treumann conjectures filtered versions of this claim. In other words, the filtered wrapped Fukaya category recovers the complex K -theory. Theorem 4.6 provides evidence for this conjecture. When the Atiyah–Hirzebruch spectral sequence $H^*(SH_S(M, \mathbb{S}), KU^*) \Rightarrow KU^*(SH_S(M, \mathbb{S}))$ degenerates (e.g. when the ordinary symplectic cohomology is supported in even degrees), $KU^*(SH_S(M, \mathbb{S}))$ is uniquely determined by the symplectic cohomology. Assuming the filtration can be chosen so that the same degeneration claim holds uniformly at every level, $KU^*(SH_S(M, \mathbb{S}))$ is likely to be determined as a (co)filtered equivariant spectrum by the filtered equivariant symplectic cohomology. It is reasonable to expect that a filtered equivariant version of [Gan12, Gan19] holds, i.e. the filtered equivariant symplectic cohomology is determined by the filtered wrapped Fukaya category, and this would prove Treumann’s conjecture.

4.2. The strategy for the proof of Theorem 4.1. Our plan is to exhibit a cofinal sequence of suitably constructed linear Hamiltonians $H : S \times S^1 \times M \rightarrow \mathbb{R}$. For such Hamiltonians, we will show that $HF_S(H, \mathbb{S})$ –as an S^1 -equivariant spectrum– has the following building blocks:

- (i) a single copy of $\Sigma^\infty(M/\partial_\infty M)$
- (ii) Thom spectra of S^1 -equivariant virtual bundles $\nu \rightarrow S^1/C_k$ for various k

More precisely, $HF_S(H, \mathbb{S})$ admits a finite filtration such that the lowest level is equivalent to $\Sigma^\infty(M/\partial_\infty M)$ (with trivial action) and the subsequent subquotients are equivalent to spectra of the form $(S^1/C_k)^\nu$. By Lemma 2.10, the completed Tate cohomology of the latter vanishes, which implies that $\widehat{R}_{S^1}^*(HF_S(H, \mathbb{S})) \simeq \widehat{R}_{S^1}^*(\Sigma^\infty(M/\partial_\infty M))$. From this, we can conclude that $\widehat{R}_{S^1}^*(SH_S(M, \mathbb{S})) \simeq \widehat{R}_{S^1}^*(\Sigma^\infty(M/\partial_\infty M))$, as claimed.

Fix a generic large slope $a > 0$. One is tempted to use autonomous Hamiltonians $H \rightarrow \mathbb{R}$ of slope a that are small in the interior of M . Assuming that equivariant $HF_S(H, \mathbb{S})$ is defined, it can be filtered by action, where the constant orbits produce a building block equivalent to $\Sigma^\infty(M/\partial_\infty M)$ and others produce subquotients equivalent to some $(S^1/C_k)^\nu$. Even though our topological framework from [CK23] allows the use of such Hamiltonians, to avoid further gluing analysis, we will follow [Zha19] and use approximately autonomous Hamiltonians. In other words, we perturb autonomous Hamiltonians to non-autonomous ones near the non-constant orbits. The finite filtration we put on the corresponding spectra is a coarse version of the action filtration.

4.3. Constructing approximately autonomous Hamiltonians.

4.3.1. Perturbing an autonomous Hamiltonian. Fix a slope $a > 0$ that is different from the length of any Reeb orbit of λ_∞ . Following [Zha19, §5], it will be convenient to fix a particular class of Hamiltonians of slope a whose dynamics are simple to analyze. The following lemma is essentially proved in [Zha19, §5].

Lemma 4.8. *Given any $0 < \epsilon_a \ll 1$, there exists a Hamiltonian $\overline{H}^a : M \rightarrow \mathbb{R}$ with the following properties:*

$$(4.2) \quad \overline{H}^a(x) = \begin{cases} -F(x), & \text{for } x \in \text{int}(\overline{M}) \\ (r-1)^2/2, & x = (r, y) \in [1 + \epsilon_a, a + 1 - \epsilon_a] \times \partial\overline{M} \\ a(r-1) - a^2/2, & x = (r, y) \in [a + 1, \infty) \times \partial\overline{M}, \end{cases}$$

where $F : \overline{M} \rightarrow \mathbb{R}$ is Morse, bounded of size ϵ_a (up to a universal constant) in the C^2 norm, and its critical points are contained in the interior. We may assume that the time-1 orbits of \overline{H}^a all have different actions, and are of the following two types:

- (a) critical points of F , all of which are non-degenerate (see [AD14, Prop. 6.1.5])
- (b) of the form $\{\ell + 1\} \times \gamma_0$, where $\ell \in (0, a)$ and γ_0 is a length ℓ Reeb orbit. Such orbits are transversally non-degenerate and come in S^1 -families. Their action is $p_\ell := \ell^2/2 + \ell$. \square

We now introduce time-dependent perturbations of \overline{H}^a near each critical manifold of type (b). Let $h_0 : S^1 \rightarrow [0, 1]$ be a Morse function with a maximum at $1/2$ and minimum at 0 . For a non-constant k -fold orbit γ of \overline{H}^a , define a function h_t on $Im(\gamma)$ by $h_t(\gamma(t')) = h_0(k(t' - t))$ and extend h_t to a small tubular neighborhood V_γ of $Im(\gamma)$. Consider $\overline{H}_{\epsilon, \gamma}^a = \overline{H}^a + \epsilon h_t$, for a small $0 < \epsilon \ll \epsilon_a$. It is shown in [CFHW96, Prop. 2.2] that

- the Hamiltonian $\overline{H}_{\epsilon, \gamma}^a$ has two non-degenerate orbits within V_γ given by $\gamma^-(t) = \gamma(t)$ and $\gamma^+(t) = \gamma(t + 1/(2k))$
- $\mathcal{A}_{\overline{H}_\epsilon^a}(\gamma^-) = \mathcal{A}_{\overline{H}^a}(\gamma)$, $\mathcal{A}_{\overline{H}_\epsilon^a}(\gamma^+) = \mathcal{A}_{\overline{H}^a}(\gamma) - \epsilon$, and $ind(\gamma^-) = ind(\gamma^+) + 1$
- there are exactly two Floer trajectories from γ^- to γ^+ within V_γ

We make this perturbation for every non-constant orbit, and denote the resulting Hamiltonian by $\overline{H}_\epsilon^a : S^1 \times M \rightarrow \mathbb{R}$. More precisely, as non-constant orbits come in circle families, we make a choice of $\gamma : S^1 \rightarrow M$ for each of them. Then we apply the procedure above for each chosen orbit parametrization. For small enough ϵ , the perturbation will not create new 1-periodic orbits other than γ^\pm . In other words:

Lemma 4.9. *The time-1 orbits of \overline{H}_ϵ^a are of the following two types:*

- (a) the critical points of F
- (b) a pair of non-stationary orbits γ^+ and γ^- for every Reeb orbit γ . If γ has length ℓ , then γ^\pm have action arbitrarily close to $p_\ell := \ell^2/2 + \ell$. \square

In particular, the Hamiltonian \overline{H}_ϵ^a is non-degenerate and it has only finitely many orbits.

Choice of a generic almost complex structure $S^1 \rightarrow M$ allows one to define a flow category $\mathcal{M}_{\overline{H}_\epsilon^a, J}$ and its geometric realization $HF(\overline{H}_\epsilon^a, \mathbb{S})$. There is a finite, increasing action filtration on $\mathcal{M}_{\overline{H}_\epsilon^a, J}$ and $HF(\overline{H}_\epsilon^a, \mathbb{S})$. More precisely, let $F^p \mathcal{M}_{\overline{H}_\epsilon^a, J}$ denote the subcategory of $\mathcal{M}_{\overline{H}_\epsilon^a, J}$ spanned by orbits of action at most p , and let $F^p HF(\overline{H}_\epsilon^a, \mathbb{S})$ denote its geometric realization. For $p \leq p'$, $F^p \mathcal{M}_{\overline{H}_\epsilon^a, J} \hookrightarrow F^{p'} \mathcal{M}_{\overline{H}_\epsilon^a, J}$ is an inclusion of framed flow categories in the sense of [CK23], and it induces a map $F^p HF(\overline{H}_\epsilon^a, \mathbb{S}) \rightarrow F^{p'} HF(\overline{H}_\epsilon^a, \mathbb{S})$ of geometric realizations. We denote the homotopy cofiber of this map by $F^{(p, p']} HF(\overline{H}_\epsilon^a, \mathbb{S})$. It coincides with the geometric realization of the quotient flow category by [CK23, Cor. 2.36].

If p is the action of γ , there exists a $\delta > 0$ such that γ^\pm have action within the window $(p - \delta, p + \delta)$. We assume ϵ is small enough, so that there exists a $\delta > 0$ that works for every orbit and the action windows $(p - \delta, p + \delta)$ corresponding to different orbits do not intersect. We will show the following non-equivariant statements

- (1) $F^0 HF(\overline{H}_\epsilon^a, \mathbb{S}) \simeq \Sigma^\infty(M/\partial_\infty M)$ and the map $\Sigma^\infty(M/\partial_\infty M) \rightarrow HF(\overline{H}_\epsilon^a, \mathbb{S})$ is compatible with the continuation maps $HF(\overline{H}_\epsilon^a, \mathbb{S}) \rightarrow HF(\overline{H}_\epsilon^{a'}, \mathbb{S})$
- (2) if p is the action of a non-constant k -fold orbit γ of \overline{H}^a , then the spectrum $F^{(p - \delta, p + \delta]} HF(\overline{H}_\epsilon^a, \mathbb{S})$ is equivalent to $Im(\gamma)^\nu \simeq (S^1/C_k)^\nu$, for a virtual vector bundle ν on $Im(\gamma)$.

We will also establish equivariant versions of these two statements. For this, we first need to produce an equivariant version of $HF(\overline{H}_\epsilon^a, \mathbb{S})$ that admits a *coarse action filtration* that roughly matches the filtration on $HF(\overline{H}_\epsilon^a, \mathbb{S})$. More precisely, we shall construct approximately autonomous

Hamiltonians

$$(4.3) \quad H_\epsilon^a : S \times S^1 \times M \rightarrow \mathbb{R}$$

satisfying the required conditions and a filtration on $HF_S(H_\epsilon^a, \mathbb{S})$ such that

- (1') $F^0 HF_S(H_\epsilon^a, \mathbb{S}) \simeq \Sigma^\infty(S \times M/S \times \partial_\infty M) \simeq \Sigma^\infty(M/\partial_\infty M)$ as S^1 -equivariant spectra and the map $\Sigma^\infty(M/\partial_\infty M) \rightarrow HF_S(H_\epsilon^a, \mathbb{S})$ is compatible with the continuation maps $HF_S(H_\epsilon^a, \mathbb{S}) \rightarrow HF_S(H_\epsilon^{a'}, \mathbb{S})$
- (2') if p is the action of a non-constant k -fold orbit γ of \overline{H}^a , then the spectrum $F^{(p-\delta, p+\delta]} HF_S(H_\epsilon^a, \mathbb{S})$ is S^1 -equivariantly equivalent to $Im(\gamma)^\nu \simeq (S^1/C_k)^\nu$, for a virtual equivariant vector bundle ν on $Im(\gamma)$

We will prove (1') in Section 5 and (2') in Section 6. Assuming these, we can complete the proof of Theorem 4.1:

Proof of Theorem 4.1. Consider a sequence of real numbers $0 < a_1 < a_2 < \dots$ with $a_i \rightarrow \infty$ and each a_i is different from the lengths of non-constant Reeb orbits. Associated to this, we have a sequence $H_\epsilon^{a_k}$ of approximately autonomous Hamiltonians⁴ and we choose equivariant continuation data $H_\epsilon^{a_1} \rightsquigarrow H_\epsilon^{a_2} \rightsquigarrow \dots$

We can now use the filtration $HF_S(H_\epsilon^{a_1}, \mathbb{S}) \rightarrow HF_S(H_\epsilon^{a_2}, \mathbb{S}) \rightarrow \dots$ of $SH_S(M, \mathbb{S})$ to compute the completed Tate cohomology of $SH_S(M, \mathbb{S})$. This filtration is equivalent to

$$(4.4) \quad \Sigma^\infty(M/\partial_\infty M) \rightarrow HF_S(H_\epsilon^{a_1}, \mathbb{S}) \rightarrow HF_S(H_\epsilon^{a_2}, \mathbb{S}) \rightarrow \dots$$

where the composition $\Sigma^\infty(M/\partial_\infty M) \rightarrow HF_S(H_\epsilon^{a_k}, \mathbb{S})$ is equivalent to the PSS map itself by the compatibility statement in (1'). Therefore, to conclude the proof, we only need to show that this map induces an equivalence

$$(4.5) \quad \widehat{R}_{S^1}^*(HF_S(H_\epsilon^{a_k}, \mathbb{S})) \rightarrow \widehat{R}_{S^1}^*(\Sigma^\infty(M/\partial_\infty M))$$

for all k . We prove $\widehat{R}_{S^1}^*(F^p HF_S(H_\epsilon^{a_k}, \mathbb{S})) \rightarrow \widehat{R}_{S^1}^*(\Sigma^\infty(M/\partial_\infty M))$ is an equivalence for each p , which completes the proof as $F^p HF_S(H_\epsilon^{a_k}, \mathbb{S}) = HF_S(H_\epsilon^{a_k}, \mathbb{S})$ for $p \gg 0$. For $p = 0$, this follows from (1'). As one increases p , as long as one does not cross a δ neighborhood of the action on an orbit of \overline{H}^{a_k} , $F^p HF_S(H_\epsilon^{a_k})$ does not change. On the other hand, by (2'), the homotopy fiber of

$$(4.6) \quad \widehat{R}_{S^1}^*(F^{p+\delta} HF_S(H_\epsilon^{a_k})) \rightarrow \widehat{R}_{S^1}^*(F^{p-\delta} HF_S(H_\epsilon^{a_k}))$$

is given by $\widehat{R}_{S^1}^*((S^1/C_m)^\nu)$ for some $m \geq 1$ and for some S^1 -equivariant bundle ν on S^1/C_m . We have shown $\widehat{R}_{S^1}^*((S^1/C_m)^\nu) \simeq 0$ in Lemma 2.10. As a result $\widehat{R}_{S^1}^*(F^{p-\delta} HF_S(H_\epsilon^{a_k}))$ and $\widehat{R}_{S^1}^*(F^{p+\delta} HF_S(H_\epsilon^{a_k}))$ are the same for p in the action spectrum. Therefore,

$$(4.7) \quad \widehat{R}_{S^1}^*(F^p HF_S(H_\epsilon^{a_k})) \simeq \widehat{R}_{S^1}^*(F^0 HF_S(H_\epsilon^{a_k})) \simeq \widehat{R}_{S^1}^*(\Sigma^\infty(M/\partial_\infty M))$$

for $p \gg 0$. This finishes the proof. \square

4.3.2. Approximately autonomous Hamiltonians on the Borel construction and coarse action filtrations. Fix a slope $a > 0$ and let $\overline{H}^a : M \rightarrow \mathbb{R}$ be as before. Assume without loss of generality that the actions of non-constant orbits are positive and different from each other. As a result, there exists $\delta > 0$ such that δ is smaller than the action difference between any two non-constant orbits, as well the action difference between a constant and a non-constant orbit.

To begin with, we let

$$(4.8) \quad H^a : S \times S^1 \times M \rightarrow \mathbb{R}$$

be the pullback of \overline{H}^a under the map $S \times S^1 \times M \rightarrow S^1 \times M$ which forgets the first component.

⁴The subscript ϵ is an abuse of notation, as each of $H_\epsilon^{a_k}$ requires a sequence of small numbers, and these sequences can be different from each other; see Section 4.3.2.

We would like to input H^a into the Borel construction from Section 3.4.2. However, this is not allowed, because $H_z^a = H(z, -, -)$ fails to be non-degenerate over all $z \in \text{crit}(\tilde{f})$.

We will therefore deform H^a by a function $h : S \times S^1 \times M \rightarrow \mathbb{R}$ supported on a tubular neighborhood of the non-constant orbits. More precisely, for a non-constant k -fold orbit γ of \overline{H}^a , let V_γ denote a tubular neighborhood, and let $b_0, b_1, b_2, \dots \in S$ be a choice of a critical point from each component of $\text{crit}(\tilde{f})$. Consider a function $h : S \times S^1 \times M \rightarrow [0, 1/2]$ such that

- (1) h is equivariant with respect to the diagonal action of S^1
- (2) $h|_{\{b_i\} \times S^1 \times \text{Im}(\gamma)} : S^1 \times \text{Im}(\gamma) \rightarrow \mathbb{R}$ is equal to $\epsilon_i h_t$ where h_t is as before and $\epsilon_i > 0$
- (3) $h|_{\{b_i\} \times S^1 \times V_\gamma}$ is obtained by extending $h|_{\{b_i\} \times S^1 \times \text{Im}(\gamma)}$ as in [CFHW96] and $h|_{\{b_i\} \times S^1 \times M}$ is supported in the union of $\{b_i\} \times S^1 \times V_\gamma$
- (4) h is constant in the horizontal directions near the S^1 -orbit of b_i

The conditions so far ensure that the restriction of $H^a + h$ to $\{b_i\} \times S^1 \times M$ is a perturbed Hamiltonian of the form considered in Section 4.3.1; in particular, it has a constant orbit corresponding to each critical point of \overline{H}^a and two non-constant orbits corresponding to every non-constant orbit of \overline{H}^a . We also assume

- (5) the numbers ϵ_i are sufficiently small so that, for any i, j and pair of (possibly constant) orbits γ, γ' of \overline{H}^a of different action, the action gap between an orbit of $(H^a + h)|_{\{b_i\} \times S^1 \times M}$ obtained by perturbing γ and between an orbit of $(H^a + h)|_{\{b_j\} \times S^1 \times M}$ obtained by perturbing γ' is still strictly more than δ

In other words, the generators of $\mathcal{M}_{\tilde{f}, H^a + h, J}$ obtained by perturbing different non-constant orbits of \overline{H}^a still have an action gap of δ . Moreover, the action of every such orbit is more than δ . We can filter the objects of $\mathcal{M}_{\tilde{f}, H^a + h, J}$ and to ensure the morphisms decrease the action (up to a small error), we impose

- (6) along any gradient trajectory of \tilde{f} , the integral $\int_{\mathbb{R}} \sup_{S^1 \times M} |dh(\nabla_{\tilde{g}} \tilde{f})| ds$ is strictly less than $\delta/2$

Note that $\sup_{S^1 \times M} |dh(\nabla_{\tilde{g}} \tilde{f})|$ is compactly supported because of (4). More precisely, h is constant along horizontal directions and $\nabla_{\tilde{g}} \tilde{f}$ is horizontal.

Lemma 4.10. *Functions $h : S \times S^1 \times M \rightarrow [0, 1/2]$ satisfying (1)-(6) exist.*

Proof. One can construct such an h over each $S^{2i+1} \subset S$, and can guarantee (6) by rescaling it by a small constant. The main difficulty is to guarantee it over the entire S .

Let $U_i \subset S$ denote a small S^1 -equivariant neighborhood of the orbit of b_i , on which the connection is flat. Construct h over b_i , extend S^1 -equivariantly to its orbit and extend horizontally to U_i . We obtain a function over $\bigcup U_i$ satisfying (1)-(4). Using equivariant cutoff functions that have arbitrarily large support in U_i , extend h to all of S . We only have to check (6) holds. The function $\sup_{S^1 \times M} |dh(\nabla_{\tilde{g}} \tilde{f})|$ is supported near the boundary of U_i , and its integral along a partial negative gradient trajectory of \tilde{f} crossing the support near ∂U_i is bounded above by $2\epsilon_i$. (6) holds under the assumption that $2 \sum \epsilon_i < \delta/2$. See [Zha19, Lem. 5.4] for a similar computation. \square

We denote $H^a + h$ also by H_ϵ^a to stress its dependency on $\{\epsilon_i\}$. The purpose of conditions (1)-(6) is to guarantee:

Lemma 4.11. *The morphisms of $\mathcal{M}_{\tilde{f}, H_\epsilon^a, J}$ between objects obtained by perturbing different orbits of \overline{H}^a decrease the action. More generally, a morphism can increase the action by at most $\delta/2$.*

Proof. The first claim follows from the second: two such objects have an action gap of at least δ . Therefore, a morphism between them cannot increase the action.

To prove the second claim, consider a morphism (η, u) of $\mathcal{M}_{\tilde{f}, H_\epsilon^a, J} = \mathcal{M}_{\tilde{f}, H^a + h_{\eta(s)}, J}$. Here η is a negative gradient trajectory of \tilde{f} and u satisfies

$$(4.9) \quad \partial_s u + J_{\eta(s)}(\partial_t u - X_{H^a + h_{\eta(s)}}) = 0.$$

Let $H_s = H^a + h_{\eta(s)}$. The energy of such a cylinder is given by

$$(4.10) \quad E(u) = \mathcal{A}(x_{inp}) - \mathcal{A}(x_{out}) + \int_{\mathbb{R} \times S^1} \partial_s H_s(u(s, t)) ds dt.$$

Observe that $\partial_s H_s(u(s, t)) = -dh(\nabla_{\tilde{g}} \tilde{f})$. Therefore,

$$(4.11) \quad \left| \int_{\mathbb{R} \times S^1} \partial_s H_s(u(s, t)) ds dt \right| \leq \int_{\mathbb{R} \times S^1} |dh(\nabla_{\tilde{g}} \tilde{f})| ds dt \leq \int_{\mathbb{R}} \sup_{S^1 \times M} |dh(\nabla_{\tilde{g}} \tilde{f})| ds \leq \delta/2.$$

□

As a result, given p' that is at least δ away from any action of any orbit of \overline{H}^a , the span of objects of action at most p' defines a flow subcategory $F^{p'} \mathcal{M}_{\tilde{f}, H_\epsilon^a, J}$ (more precisely, there is an inclusion $F^{p'} \mathcal{M}_{\tilde{f}, H_\epsilon^a, J} \hookrightarrow \mathcal{M}_{\tilde{f}, H_\epsilon^a, J}$ in the sense of [CK23]). This may not be the case for p' closer to the action spectrum of \overline{H}^a . Clearly, varying p' continuously without getting closer to the action spectrum than δ does not change the subcategory or its realization. Therefore, we will be specifically interested in $F^{p \pm \delta} \mathcal{M}_{\tilde{f}, H_\epsilon^a, J}$, for p in the action spectrum of \overline{H}^a , their geometric realizations, and the homotopy cofiber

$$(4.12) \quad F^{(p-\delta, p+\delta]} HF_S(H_\epsilon^a) := \text{cofib}(|F^{p-\delta} \mathcal{M}_{\tilde{f}, H_\epsilon^a, J}| \rightarrow |F^{p+\delta} \mathcal{M}_{\tilde{f}, H_\epsilon^a, J}|).$$

5. AN EQUIVARIANT PSS MAP

The goal of this section is to prove

Proposition 5.1. *There is an S^1 -equivariant PSS map*

$$(5.1) \quad \Sigma^\infty(M/\partial_\infty M) \simeq \Sigma^\infty(S \times M/S \times \partial_\infty M) \rightarrow HF_S(H; \mathbb{S})$$

satisfying the following properties:

- Given $H \rightsquigarrow H'$, we have a homotopy-commutative diagram

$$(5.2) \quad \begin{array}{ccc} \Sigma^\infty(M/\partial_\infty M) & \longrightarrow & HF_S(H; \mathbb{S}) \\ & \searrow & \downarrow \\ & & HF_S(H'; \mathbb{S}) \end{array}$$

- when $H = H_\epsilon^a$, it factors through an equivalence $\Sigma^\infty(M/\partial_\infty M) \xrightarrow{\simeq} F^0 HF_S(H_\epsilon^a; \mathbb{S})$.

As a warm-up for the equivariant case, we prove the following non-equivariant version first

Lemma 5.2. *There is a PSS map*

$$(5.3) \quad \Sigma^\infty(M/\partial_\infty M) \rightarrow HF(H; \mathbb{S})$$

satisfying the following properties:

- Given $H \rightsquigarrow H'$, we have a homotopy-commutative diagram

$$(5.4) \quad \begin{array}{ccc} \Sigma^\infty(M/\partial_\infty M) & \longrightarrow & HF(H; \mathbb{S}) \\ & \searrow & \downarrow \\ & & HF(H'; \mathbb{S}) \end{array}$$

- when $H = \overline{H}_\epsilon^a$, it factors through an equivalence $\Sigma^\infty(M/\partial_\infty M) \xrightarrow{\simeq} F^0 HF(\overline{H}_\epsilon^a; \mathbb{S})$

To construct the non-equivariant map, fix a function f that is Morse, proper, with critical points only in the interior, and bounded above (also fix a Morse–Smale metric for f). The same argument as in [CK23, Thm. 3.11] shows that $|\mathcal{M}_{-f}| \simeq \Sigma^\infty M_+$, as $-f$ is bounded below; *mutatis mutandis* the argument also shows that $|\mathcal{M}_f| \simeq \Sigma^\infty(M/\partial_\infty M)$. We will eventually let $f = -\overline{H}^a$, which matches F in the interior.

Fix Floer data (H, J) . Given $x \in \text{crit}(f)$ and $y \in \text{orb}(H)$, (after a choice of data) define a bimodule over $\mathcal{M}_f\text{-}\mathcal{M}_{H,J}$ by associating to (x, y) the usual compactified moduli space of “spiked discs”. More precisely, this is the compactified moduli space of pairs (θ, u) where $\theta : (-\infty, 0] \rightarrow M$ is a half negative gradient trajectory of f from x to $\theta(0)$, $u : D = \mathbb{C}\mathbb{P}^1 \setminus \{0\} \rightarrow M$ satisfies the perturbed Cauchy–Riemann equation and it is asymptotic to y , and $\theta(0) = u(\infty)$. The construction of smooth charts on these moduli spaces can be done similarly to [Lar21, Sec. 6] (see also [PS24, Sec. 8]).

To define a map $|\mathcal{M}_f| \rightarrow |\mathcal{M}_{H,J}|$, we need to frame this bimodule. For this purpose, it is more natural to use the description of the framings on $\mathcal{M}_{H,J}$ in terms of negative caps (see [CK23, Lem. 5.5] and the preceding discussion). To explain this further in that context, if W_x denotes the framing bundle over $\{x\} \subset \text{crit}(f)$ and V_y^- is the bundle over $\{y\} \subset \text{orb}(H)$ defined as in loc. cit. then a point in the moduli of spiked discs has tangent space given by

$$(5.5) \quad W_x + V_y^- - \mathbb{R}^{2k} - T_M,$$

where k is the stabilization constant (i.e. we fix $T_M \oplus \mathbb{C}^k \simeq \mathbb{C}^{n+k}$). To see this, notice the moduli of spiked discs is the fibered product of moduli of half gradient trajectories from x and moduli of half discs to x over their evaluation maps to M . The former has tangent space that can be identified with W_x and the tangent space of the discs can be identified with $V_y^- - \mathbb{C}^k$. By subtracting T_M pulled back along the evaluation map at the spike, we obtain the asserted equivalence. Combining this with [CK23, Lem. 5.5] gives us the framings. Therefore, we have a map

$$(5.6) \quad \Sigma^\infty(M/\partial_\infty M) = |\mathcal{M}_f| \rightarrow |\mathcal{M}_{H,J}| = HF(H, \mathbb{S}).$$

The compatibility with the continuation map follows from standard Floer theory and [CK23, §3.4].

When $H = \overline{H}_\epsilon^a$, we let $f = -\overline{H}^a$, which matches F in the interior, and we impose an additional restriction that the Hamiltonian term of the Floer data parametrized by the disc matches \overline{H}^a and \overline{H}_ϵ^a outside small neighborhoods of non-constant orbits. It is easy to see that this map factors through $F^0 HF(\overline{H}_\epsilon^a, \mathbb{S})$. One can further filter \mathcal{M}_f and $F^0 \mathcal{M}_{\overline{H}_\epsilon^a, J}$ by the values of the Morse function $-F$ on the object (critical point resp. constant orbit). This filtration is compatible with the constructed bimodule above. Moreover, the bimodule has only one trajectory (the constant spiked disc) that does not strictly decrease this Morse filtration. As a result, $\Sigma^\infty(M/\partial_\infty M) \rightarrow F^0 HF(\overline{H}_\epsilon^a, \mathbb{S})$ induces isomorphisms on the subquotients. This proves Lemma 5.2.

We now turn to the construction of the S^1 -equivariant PSS map and the proof of Proposition 5.1. The proof essentially consists in rerunning the above arguments equivariantly. To do this, we will use the Morse–Bott techniques developed in [CK23].

Namely, let $f : M \rightarrow \mathbb{R}$ be Morse as above and let $\tilde{f} : S \rightarrow \mathbb{R}$ be the standard Morse–Bott function introduced in Section 3.4.2. Let us now consider the Morse–Bott flow category $\mathcal{M}_{\tilde{f}, f}$ where:

- $ob(\mathcal{M}_{\tilde{f}, f}) = \text{crit}(\tilde{f}) \times \text{crit}(f)$
- the morphisms are broken pairs of negative gradient trajectories for the Morse–Bott function \tilde{f} and the Morse function f (equivalently, negative gradient trajectories of $\tilde{f} + f$ with respect to the product metric)

This is an S^1 -equivariant Morse–Bott flow category. In fact, $\mathcal{M}_{\tilde{f}, f}$ is a trivial instance of the type of flow category considered in [CK23, §4]. Hence Theorem 4.6 of loc. cit. implies that the

geometric realization $|\mathcal{M}_{\tilde{f},f}|$ is S^1 -equivariantly equivalent to $\Sigma^\infty(M/\partial_\infty M)$ (strictly speaking we are appealing to an open manifold analog of the aforementioned theorem with the same proof).

To write the promised equivariant PSS map $\Sigma^\infty(M/\partial_\infty M) \rightarrow HF_S(H, \mathbb{S})$, we shall consider an equivariant $\mathcal{M}_{\tilde{f},f}$ - $\mathcal{M}_{\tilde{f},H,J}$ -bimodule \mathcal{N}^{PSS} , similarly to before. To define \mathcal{N}^{PSS} , the essential idea is to consider pairs given by a negative gradient η trajectory of \tilde{f} and a PSS-trajectory ‘‘above η ’’; see Figure 5.1. To implement this idea rigorously, we choose:

- an S^1 -equivariant function $S \times M \times (-\infty, 0] \rightarrow \mathbb{R}$ that agrees with $\tilde{f} + f$ near $-\infty$ and a family of S^1 -equivariant metrics on $S \times M$ parametrized by $(-\infty, 0]$ that agrees with the product metric near $-\infty$
- Floer data on M parametrized by $S \times D$ (where $D = \mathbb{C}\mathbb{P}^1 \setminus \{0\}$ with fixed negative cylindrical end) that agrees outside a compact subset of D with the $S \times S^1$ -parametric data chosen to define $HF_S(H, \mathbb{S})$ (in particular, outside a compact set, the data only depends on the argument of a point in D). We further assume that this data is circle equivariant with respect to the diagonal action on $S \times D$.

Let $\phi : D \rightarrow [0, \infty)$ denote an S^1 -equivariant smoothing of $q \mapsto |q|^{-1}$ near $q = \infty$, obtained by composing $q \mapsto |q|^{-1}$ with a monotone increasing function $[0, \infty) \rightarrow [0, \infty)$ that is equal to cx^2 near 0 and that is equal to the identity outside a small neighborhood of 0. In other words, ϕ is a smoothed projection function $D \rightarrow [0, \infty)$. See the right half of Figure 5.1.

We define a bimodule \mathcal{N}^{PSS} by assigning to $(p, x) \in ob(\mathcal{M}_{\tilde{f},f})$ and $(p', y) \in (\mathcal{M}_{\tilde{f},H,J})$ the set of (η, θ, u) such that

- η is a trajectory of $-\tilde{f}$ from p to p'
- $\theta : (-\infty, 0] \rightarrow M$ such that $(\eta|_{(-\infty, 0]}, \theta) : (-\infty, 0] \rightarrow S \times M$ is a negative (half) gradient trajectory with respect to the chosen Morse data above
- $u : D \rightarrow M$ satisfies the Cauchy–Riemann equation with respect to the chosen Floer data above. In other words, if we denote the Floer data (temporarily) by $(\mathbf{H}_{(s,q)}, J_{(s,q)})$ (where $\mathbf{H}_{(s,q)}$ denotes the Hamiltonian valued 1-form on D), then we have a Floer datum over D given by $(\mathbf{H}_{(\eta(\phi(q)), q)}, J_{(\eta(\phi(q)), q)})$ and u satisfies

$$(5.7) \quad (du - X_{\mathbf{H}_{(\eta(\phi(q)), q)}})^{0,1} = 0$$

- θ is asymptotic to x , u is asymptotic to y , and $\theta(0) = u(\infty)$

In other words, we are considering pairs of Morse trajectories η from p to p' and PSS-trajectories within $S \times M$ from (p, x) to (p', y) ‘‘above η ’’; see Figure 5.1.


By construction, this moduli space admits a *free* S^1 -action, so transversality can be achieved by combining the standard argument for the non-equivariant PSS map and the equivariant transversality arguments in [BO17]. The smooth structures are similarly induced from smooth structures on the S^1 -quotiented moduli spaces, as explained in [CK23, Sec. 7]. We have natural framings analogous to (5.5).

As a result, [CK23, Prop. 3.4] furnishes an equivariant PSS-map

$$(5.8) \quad \Sigma^\infty(M/\partial_\infty M) \simeq \Sigma^\infty(S \times M/S \times \partial_\infty M) \rightarrow HF_S(H, \mathbb{S}).$$

Similarly to the non-equivariant case, the compatibility with the continuation map follows from standard Floer theory and [CK23, §3.4].

Assume that $H = H_\epsilon^a$. Next we prove the equivalence $\Sigma^\infty(M/\partial_\infty M) \xrightarrow{\simeq} F^0 HF_S(H_\epsilon^a; \mathbb{S})$. The bimodule constructed above is non-empty only when $(p', y) \in ob(F^0 \mathcal{M}_{H_\epsilon^a, J})$. Therefore, the induced map factors through $F^0 HF_S(H_\epsilon^a, \mathbb{S})$. The objects of $F^0 \mathcal{M}_{H_\epsilon^a, J}$ consists of pairs (p', y) , where y is a constant trajectory and hence a critical point of F . Therefore, one can identify the objects of $\mathcal{M}_{\tilde{f},f}$ and $F^0 \mathcal{M}_{H_\epsilon^a, J}$. Moreover, both categories are filtered by the value of $\tilde{f} + f$ and the bimodule respects this filtration. A similar argument shows that the induced map is an equivalence.

FIGURE 5.1. A spiked disc “over η ”

6. LOCAL FLOER HOMOLOGY COMPUTATION

The goal of this section is to prove the claims (2) and (2') that were stated in Section 4.3.1. We carry over the notation from the previous sections. In particular, we fix a slope $a > 0$ and let γ be a k -fold orbit of the autonomous Hamiltonian \overline{H}^a of action p . Let $0 < \delta \ll 1$ be as in Section 4.3.2 (in particular, smaller than the action gap between orbits of \overline{H}^a). Let ϵ be as in Section 4.3.2.⁵

First observe

Lemma 6.1. *A framed flow category \mathcal{M} with two objects x, y , and two morphisms in $\mathcal{M}(x, y)$ has geometric realization equivalent to either $\Sigma^\infty(S^1)_+$ or $\Sigma^\infty\mathbb{R}P^2$ up to shift. In other words, it is the same as a Thom spectrum over S^1 .*

Proof. \mathcal{M} is the flow category of the standard Morse function on S^1 , and different framings correspond to the standard framing twisted by a virtual bundle ν over S^1 . By [CK23, Theorem 4.6'], $|\mathcal{M}| \simeq (S^1)^\nu$. The virtual bundles on S^1 are equivalent to the trivial bundle or the Möbius strip (up to shift). In the former case, $|\mathcal{M}|$ is equivalent to $\Sigma^\infty(S^1)_+$, and in the latter to $\Sigma^\infty\mathbb{R}P^2$ (up to shift). \square

As explained in Section 4.3.1, it is a consequence of [CFHW96, Lem. 2.2] that the spectrum $F^{(p-\delta, p+\delta]}HF(\overline{H}_\epsilon^a, \mathbb{S})$ is the geometric realization of a flow category satisfying the conditions of Lemma 6.1. This implies (2). However, this argument is harder to implement in the equivariant case, so instead we will define a map using the formalism of P -relative modules, which we introduced in [CK23]. This will give an alternative proof of (2), which we will then rerun equivariantly to produce the map in (2').

We begin by considering the (non-equivariant) flow category $\mathcal{M}_{\overline{H}_\epsilon^a, J}$. Let $X_\gamma \simeq S^1/C_k \simeq S^1$ denote the image of γ . We define a X_γ -relative module on $F^{p+\delta}\mathcal{M}_{\overline{H}_\epsilon^a, J}$ by using the moduli space of continuation trajectories with output given by (some rotation of) γ . Note that, heuristically, this is like using continuation maps from $HF(\overline{H}_\epsilon^a, \mathbb{S})$ to “ $HF(\overline{H}^a, \mathbb{S})$ ”. We prefer to avoid formally defining $HF(\overline{H}^a, \mathbb{S})$ as \overline{H}^a is autonomous and we do not wish to discuss gluing of trajectories along the orbit of an autonomous Hamiltonian.

Choose generic Floer data parametrized by the cylinder that is equal to $(\overline{H}_\epsilon^a, J)$ on the input (positive) end and whose Hamiltonian term is the same as $\overline{H}^a + \epsilon$ on the output end. We also assume the Hamiltonian has negative derivative in the s -direction. Given $x \in \text{ob}(F^{p+\delta}\mathcal{M}_{\overline{H}_\epsilon^a, J})$, define $\mathcal{N}(x)$ to be the moduli space of broken trajectories from x to an orbit of \overline{H}^a supported at X_γ (i.e. some $\gamma(t + \theta)$).

⁵Recall from Section 4.3.1 that $0 < \epsilon \ll \delta$ controls the size of the perturbation of the autonomous Hamiltonian \overline{H}^a . When we extend this construction to the Borel-equivariant setting in Section 4.3.2, then recall that ϵ is secretly shorthand for a decreasing sequence $\{\epsilon_i\}$, where ϵ_i controls the size of the perturbation over the unique index $2i$ critical points of $f : \mathbb{C}P^\infty \rightarrow \mathbb{R}$.

Lemma 6.2. \mathcal{N} is a X_γ -relative module over $F^{p+\delta}\mathcal{M}_{\overline{H}_\epsilon^a, J}$.

Proof. The key point is compactness. Given $x \in ob(F^{p+\delta}\mathcal{M}_{\overline{H}_\epsilon^a, J})$, we are asserting that a sequence of trajectories from x to an orbit of $t \mapsto \gamma(t + \theta)$ of \overline{H}^a breaks into a broken Floer trajectory \overline{H}_ϵ^a from x to some $x' \in ob(F^{p+\delta}\mathcal{M}_{\overline{H}_\epsilon^a, J})$, followed by an unbroken trajectory from x' to some $\gamma(t + \theta)$. In other words, breakings on the \overline{H}^a side cannot occur. This is true for elementary action reasons: if our sequence of curves broke on the \overline{H}^a side, then we would have a trajectory from an object of $F^{p+\delta}\mathcal{M}_{\overline{H}_\epsilon^a, J}$ to a higher action orbit of \overline{H}^a . This is not possible due to the negative derivative condition in s -direction.

Smoothness of the moduli spaces is achieved by choosing our Floer data generically. One can use gluing arguments as in [Lar21, Sec. 6] to show that \mathcal{N} defines a module over $F^{p+\delta}\mathcal{M}_{\overline{H}_\epsilon^a, J}$. Moreover, there is a compatible evaluation map $\mathcal{N}(x) \rightarrow X_\gamma$: send a trajectory that is asymptotic to $\gamma(t + \theta)$ at the negative end to $\gamma(\theta) \in X_\gamma$ (one can identify X_γ with these orbits, we are sending a trajectory to the orbit it is asymptotic to). \square

We now discuss framings, which are similar to the standard continuation maps. More precisely, recall that we fix an identification $TM \oplus \mathbb{C}^k \cong \mathbb{C}^{n+k}$, and consider the space \mathcal{U}_γ of abstract caps given by the tuples (y, L, J, Y, g) where

- $y \in X_\gamma$, also thought of as an orbit of \overline{H}^a of the form $t \mapsto \gamma(t + \theta_0)$
- $\epsilon : [0, \infty) \times S^1 \rightarrow S_- := \mathbb{C}$ is a tubular end
- $L > 0$ is a real number
- (J, Y) is a Floer datum on $S_- \times \mathbb{C}^{n+k}$ that matches over $\epsilon([L, \infty) \times S^1)$ with the datum for the continuation trajectories under stabilization and the identification $TM \oplus \mathbb{C}^k \cong \mathbb{C}^{n+k}$
- g is a metric on S_- that matches over $\epsilon([L, \infty) \times S^1)$ with the standard metric

This is an autonomous version of [CK23, Definition 5.2] (see also [CK23, §5.3] for more details). We have

- (1) an homotopy equivalence $\mathcal{U}_\gamma \rightarrow X_\gamma$ given by $(y, L, J, Y, g) \mapsto y$
- (2) a map $\mathcal{U}_\gamma \rightarrow Fred$, the space of Fredholm operators

Moreover, \mathcal{U}_γ carries a circle action, and both the equivalence and the map into $Fred$ are compatible with the circle action. As a result, we have an S^1 -equivariant index bundle over \mathcal{U}_γ and over X_γ ; see also [CK23, Remark 5.7]. We denote this bundle by ν .

Note 6.3. In [CK23], $Fred$ is identified with the space of Fredholm operators on a Sobolev space. Since \overline{H}^a is autonomous, the action functional has Morse–Bott degeneracies and we must therefore use weighted Sobolev spaces; see [BO09]. We do not spell out the details of this generalization and suppress the distinction from our notation.

Similarly to [Lar21, CK23], we have isomorphisms

$$(6.1) \quad V_x = T_{\mathcal{N}(x)} \oplus \nu$$

giving us a framing in the sense of [CK23, Definition 3.5].

Therefore, we get a map

$$(6.2) \quad F^{p+\delta}HF(\overline{H}_\epsilon^a, \mathbb{S}) := |F^{p+\delta}\mathcal{M}_{\overline{H}_\epsilon^a, J}| \rightarrow (X_\gamma)^\nu.$$

By the same action considerations as in the proof of Lemma 6.2, given any $x \in ob(F^{p-\delta}\mathcal{M}_{\overline{H}_\epsilon^a, J})$, the set $\mathcal{N}(x)$ is empty. Therefore (6.2) factors through

$$(6.3) \quad F^{(p-\delta, p+\delta)}HF(\overline{H}_\epsilon^a, \mathbb{S}) \rightarrow (X_\gamma)^\nu.$$

Before moving on to the equivariant case, we show

Proposition 6.4. (6.3) is an equivalence.

Before starting the proof, observe that one can see (X_γ, ν) as a framed Morse–Bott flow category with no morphisms (where ν is the framing bundle). We denote it by \mathcal{M}_γ . Similarly, the module \mathcal{N} can be seen as a framed $F^{(p-\delta, p+\delta]} \mathcal{M}_{\overline{H}_\epsilon^a, J} \mathcal{M}_\gamma$ -bimodule. If we had formally constructed the category “ $\mathcal{M}_{\overline{H}^a, J}$ ” associated to the autonomous Hamiltonian \overline{H}^a , then \mathcal{M}_γ would be $F^{(p-\delta, p+\delta]} \mathcal{M}_{\overline{H}^a, J}$. In particular, \mathcal{M}_γ can be interpreted as the local Floer homology of open set $V_\gamma \subset M$.

We now introduce a \mathcal{M}_γ - $F^{(p-\delta, p+\delta]} \mathcal{M}_{\overline{H}_\epsilon^a, J}$ -bimodule \mathcal{N}' that will be a partial inverse to \mathcal{N} . Fix Floer data over the cylinder whose Hamiltonian term is \overline{H}^a on the positive (input) end and \overline{H}_ϵ^a on the negative (output) end. Given $y \in \text{ob}(\mathcal{M}_\gamma)$, $x \in F^{(p-\delta, p+\delta]} \mathcal{M}_{\overline{H}_\epsilon^a, J}$ define $\mathcal{N}'(y, x)$ to be the compactified moduli space of cylinders that is asymptotic to y on the positive (input) end (i.e. to some $\gamma(t + \theta)$) and to x on the negative (output) end. This varies continuously in y and defines a \mathcal{M}_γ - $F^{(p-\delta, p+\delta]} \mathcal{M}_{\overline{H}_\epsilon^a, J}$ -bimodule. The framing of \mathcal{N}' is analogous to that of \mathcal{N} .

We would like to warn the reader that \mathcal{N}' does not extend to a \mathcal{M}_γ - $F^{p+\delta} \mathcal{M}_{\overline{H}_\epsilon^a, J}$ -bimodule. Indeed, to extend to such a bimodule, we would need to add trajectories from $y \in X_\gamma$ to lower action orbits of \overline{H}_ϵ^a , as they naturally appear in composition. But the moduli space of such trajectories is not compact, as it does not include the breakings on the autonomous side. Morally, \mathcal{N}' extends to a “ $F^{p+\delta} \mathcal{M}_{\overline{H}_\epsilon^a, J}$ ”- $F^{p+\delta} \mathcal{M}_{\overline{H}_\epsilon^a, J}$ -bimodule; however, we avoid this as we have not defined the former category.

Proof of Proposition 6.4. \mathcal{N}' gives us a map $X_\gamma^\nu \rightarrow F^{(p-\delta, p+\delta]} HF(\overline{H}_\epsilon^a, \mathbb{S})$. By composing this with \mathcal{N} , we obtain a map $X_\gamma^\nu \rightarrow X_\gamma^\nu$. This map is homotopic to a map induced by a \mathcal{M}_γ - \mathcal{M}_γ -bimodule \mathcal{N}'' that associates to (y, y') the moduli of trajectories from y to y' with respect to continuation data that is equal to \overline{H}^a on the input end, and to $\overline{H}^a + \epsilon$ on the output end.

This map is homotopic to identity, hence an isomorphism. More precisely, a \mathcal{M}_γ - \mathcal{M}_γ -bimodule is simply a smooth manifold with two evaluation maps to X_γ . By letting ϵ vary, we see that \mathcal{N}'' is cobordant to a similarly defined bimodule for $\epsilon = 0$. But the latter is clearly equal to X_γ with evaluation maps given by the identity. The cobordism is compatible with the framings, which tells us the induced maps are homotopic.

In summary, the composition

$$(6.4) \quad (X_\gamma)^\nu \rightarrow F^{(p-\delta, p+\delta]} HF(\overline{H}_\epsilon^a, \mathbb{S}) \xrightarrow{(6.3)} (X_\gamma)^\nu$$

is homotopic to the identity.

As both sides of (6.3) are connective, it suffices by the stable Hurewicz theorem to prove the map induced on integral homology is an isomorphism. In other words, if the induced map is an isomorphism, then the cone of (6.3) is a connective spectrum with vanishing homology; therefore it is zero.

It follows from (6.4) that the composition

$$(6.5) \quad \tilde{H}_*(X_\gamma^\nu, \mathbb{Z}) \rightarrow \tilde{H}_*(F^{(p-\delta, p+\delta]} HF(\overline{H}_\epsilon^a, \mathbb{S}), \mathbb{Z}) \xrightarrow{(6.3)_*} \tilde{H}_*(X_\gamma^\nu, \mathbb{Z})$$

is the identity. In other words, $\tilde{H}_*(X_\gamma^\nu, \mathbb{Z})$ is a direct summand of $\tilde{H}_*(F^{(p-\delta, p+\delta]} HF(\overline{H}_\epsilon^a, \mathbb{S}), \mathbb{Z})$.

Observe that both are isomorphic to a shift of $\mathbb{Z} \oplus \mathbb{Z}[1]$ or $\mathbb{Z}/2\mathbb{Z}$. Indeed, being a virtual bundle on the circle, ν is either a trivial bundle, or a Möbius bundle plus a trivial bundle. In the first case, $(X_\gamma)^\nu$ is a shift of $\Sigma^\infty S_+^1$; in the second case it is a shift of $\Sigma^\infty \mathbb{R}P^2$. The same is true for $F^{(p-\delta, p+\delta]} HF(\overline{H}_\epsilon^a, \mathbb{S})$: being the realization of a framed flow category $F^{(p-\delta, p+\delta]} \mathcal{M}_{\overline{H}_\epsilon^a, J}$ with two objects, and two morphisms from one to the other, it is either $\Sigma^\infty S_+^1$ or $\Sigma^\infty \mathbb{R}P^2$, up to a shift by Lemma 6.1. This proves the claim about their homology groups.

The only way for a group isomorphic to $\mathbb{Z} \oplus \mathbb{Z}[1]$ or $\mathbb{Z}/2\mathbb{Z}$ to be a direct summand of another such group is that they are the same. This implies $(6.3)_*$ is an isomorphism, finishing the proof. \square

Next we turn our attention to the equivariant case. This combines the X_γ -relative module defined above, with techniques from [CK23] which were also used in Section 5. To conclude the proof, we use a filtration argument to reduce to Proposition 6.4.

We define an $S \times X_\gamma$ -relative S^1 -equivariant framed module \mathcal{N}^{eq} over $F^{p+\delta}\mathcal{M}_{\tilde{f}, H_\epsilon^a, J}$. Let $\epsilon : S \rightarrow \mathbb{R}$ be a bounded S^1 -equivariant function such that $0 \leq h \leq \epsilon(s)$ and $\epsilon(b_i) = \epsilon_i$. Choose Floer data parametrized by $S \times \mathbb{R} \times S^1$ that is

- equivariant with respect to the diagonal S^1 -action on $S \times S^1$
- the Hamiltonian term is non-increasing in the \mathbb{R} direction
- the Hamiltonian term is equal to $H_\epsilon^a = H + h : S \times S^1 \times M \rightarrow \mathbb{R}$ on the input (positive) end
- the Hamiltonian term is equal to $H^a + \epsilon(s) : S \times M \rightarrow \mathbb{R}$ on the output (negative) end

To define \mathcal{N}^{eq} , we associate to a pair $(q, x) \in ob(F^{p+\delta}\mathcal{M}_{\tilde{f}, H_\epsilon^a, J})$ the moduli of broken pairs (β, u) where

- $\beta : (-\infty, 0] \rightarrow \mathbb{R}$ is a negative, half gradient trajectory of \tilde{f} from q
- $u : \mathbb{R} \times S^1 \rightarrow M$ is a solution from x to some $t \mapsto \gamma(t + \theta)$ to the equation

$$(6.6) \quad \partial_s u + J_{(\beta(-e^s), s, t)}(\partial_t u - X_{H_{(\beta(-e^s), s, t)}})) = 0$$

In other words, u is a continuation trajectory from x to γ “over $\beta(-e^s) : \mathbb{R} \rightarrow S$ ”.

Lemma 6.5. \mathcal{N}^{eq} is a module over $F^{p+\delta}\mathcal{M}_{\tilde{f}, H_\epsilon^a, J}$.

Proof. First of all, observe that for $s \gg 1$, $\beta(s)$ is contained in a slice of the S^1 -action and $(H_{(\beta(s), s, t)}, J_{(\beta(s), s, t)})$ are independent of s on this slice for $s \gg 1$. This follows immediately from the construction of H_ϵ^a, J and of the S^1 -equivariant metric on S , all of which is explained in Section 3.4.2 and Section 4.3.2. Hence (6.6) reduces to the standard Cauchy–Riemann equation for $s \gg 1$.

For $s \ll -1$, the Hamiltonian term of pair $(H_{(\beta(-e^s), s, t)}, J_{(\beta(-e^s), s, t)})$ converges exponentially in s to $H_{(\beta(0), \epsilon(\beta(0)), t)}^a = \overline{H}^a + \epsilon(\beta(0))$, and similarly with its J -term. Similarly to the proof of Lemma 6.2, we observe that there can be no breaking as $s \rightarrow -\infty$ (output). Indeed, any such breaking would cause the appearance of a trajectory from an orbit of $H_\epsilon^a(b, \cdot, \cdot)$ of action less than $p + \delta$ to an orbit of $\overline{H}^a + \epsilon(\beta(0))$ of higher action. This is impossible due to our assumption that the Hamiltonian term has non-increasing derivative in the \mathbb{R} -direction.

We do not spell out the details of transversality and gluing, which follow from similar considerations as for \mathcal{N}^{PSS} in Section 5. \square

We define the evaluation map by

$$(6.7) \quad \begin{aligned} \mathcal{N}^{eq}(q, x) &\rightarrow S \times X_\gamma \\ (\beta, u) &\mapsto (\beta(0), \gamma(\theta)) \end{aligned}$$

where $\gamma(t + \theta)$ is the output asymptotic of u . It carries an equivariant framing relative to the pull-back of the virtual equivariant bundle ν to $S \times X_\gamma$, which we still denote by ν . As a result, we have an S^1 -equivariant map

$$(6.8) \quad F^{p+\delta}HF_S(H_\epsilon^a, \mathbb{S}) := |F^{p+\delta}\mathcal{M}_{\tilde{f}, H_\epsilon^a, J}| \rightarrow \Sigma^\infty S_+ \wedge (X_\gamma)^\nu \simeq (X_\gamma)^\nu.$$

As before, \mathcal{N}^{eq} is empty over $ob(F^{p-\delta}\mathcal{M}_{\tilde{f}, \overline{H}_\epsilon^a, J})$; therefore, (6.8) factors through

$$(6.9) \quad F^{(p-\delta, p+\delta]}HF_S(\overline{H}_\epsilon^a, \mathbb{S}) \rightarrow \Sigma^\infty S_+ \wedge (X_\gamma)^\nu \simeq (X_\gamma)^\nu.$$

Proposition 6.6. (6.9) is an equivalence.

To prove this, we reduce to Proposition 6.4 by using a filtration argument.

Proof. $F^{(p-\delta, p+\delta]} HF_S(\overline{H}_\epsilon^a, \mathbb{S})$ is given by the geometric realization of $\mathcal{M}_{loc} := F^{(p-\delta, p+\delta]} \mathcal{M}_{\tilde{f}, H_\epsilon^a, J}$, which carries another filtration by the value of \tilde{f} . The category $F^r \mathcal{M}_{loc} = F^r F^{(p-\delta, p+\delta]} \mathcal{M}_{\tilde{f}, H_\epsilon^a, J}$ jumps when $r_i = \tilde{f}(b_i)$ for some i , i.e. at the critical values of \tilde{f} . Let X_i denote the S^1 -orbit of b_i , i.e. the critical set of \tilde{f} of index $2i$. Observe that the quotient category $F^{r_i} \mathcal{M}_{loc} / F^{r_i-1} \mathcal{M}_{loc}$ can be identified with the Morse–Bott flow category given by taking the product of X_i with the ordinary flow category $F^{(p-\delta, p+\delta]} \mathcal{M}_{\overline{H}_{\epsilon_i}^a, J}$. Here $F^{(p-\delta, p+\delta]} \mathcal{M}_{\overline{H}_{\epsilon_i}^a, J}$ is the flow category corresponding to restriction of the S -parametric Floer data to b_i . In other words, we take products of every object and the morphism space with X_i . The framing bundles on $X_i \times F^{(p-\delta, p+\delta]} \mathcal{M}_{\overline{H}_{\epsilon_i}^a, J}$ are the sum of these for $F^{(p-\delta, p+\delta]} \mathcal{M}_{\overline{H}_{\epsilon_i}^a, J}$ plus the descending bundle V_i over X_i . Its realization is the same as $X_i^{V_i} \wedge F^{(p-\delta, p+\delta]} HF(\overline{H}_\epsilon^a, \mathbb{S})$.

Similarly, the target $S \times X_\gamma$ is filtered by the value of \tilde{f} , and jumps in the homotopy type only occur at the critical points r_i . Indeed, this filtration depends only on the S -component, and $F^{r_i} \Sigma^\infty S_+ / F^{r_i-1} \Sigma^\infty S_+$ is equivalent to $X_i^{V_i}$ (see [CK23, Sec. 4] for more details). Therefore we have a map

$$(6.10) \quad \begin{aligned} X_i^{V_i} \wedge F^{(p-\delta, p+\delta]} HF(\overline{H}_\epsilon^a, \mathbb{S}) &\simeq |F^{r_i} \mathcal{M}_{loc}| / |F^{r_i-1} \mathcal{M}_{loc}| \rightarrow \\ F^{r_i}(\Sigma^\infty S_+ \wedge (X_\gamma)^\nu) / F^{r_i-1}(\Sigma^\infty S_+ \wedge (X_\gamma)^\nu) &\simeq X_i^{V_i} \wedge X_\gamma^\nu \end{aligned}$$

and it is not difficult to see that this map matches (6.3) smashed with $X_i^{V_i}$. Hence, by Proposition 6.4, the map (6.10) is an equivalence for every i . As a result, by induction, the map

$$(6.11) \quad F^{r_i} HF_S(H_\epsilon^a) \simeq |F^{r_i} \mathcal{M}_{loc}| \rightarrow F^{r_i}(\Sigma^\infty S_+ \wedge (X_\gamma)^\nu)$$

induced by \mathcal{N}^{eq} is an isomorphism for every r_i . This implies the same for their colimit as $i \rightarrow \infty$, which concludes the proof. \square

REFERENCES

- [Abo15] Mohammed Abouzaid. Symplectic cohomology and Viterbo’s theorem. In *Free loop spaces in geometry and topology*, volume 24 of *IRMA Lect. Math. Theor. Phys.*, pages 271–485. Eur. Math. Soc., Zürich, 2015.
- [ACF16] Peter Albers, Kai Cieliebak, and Urs Frauenfelder. Symplectic tate homology. *Proceedings of the London Mathematical Society*, 112(1):169–205, 2016.
- [AD14] Michèle Audin and Mihai Damian. *Morse theory and Floer homology*. Universitext. Springer, London; EDP Sciences, Les Ulis, 2014. Translated from the 2010 French original by Reinie Ern e.
- [AMS21] Mohammed Abouzaid, Mark McLean, and Ivan Smith. Complex cobordism, Hamiltonian loops and global Kuranishi charts. *arXiv e-prints*, page arXiv:2110.14320, October 2021.
- [BB20] Tobias Barthel and Agn es Beaudry. Chromatic structures in stable homotopy theory. In *Handbook of homotopy theory*, CRC Press/Chapman Hall Handb. Math. Ser., pages 163–220. CRC Press, Boca Raton, FL, [2020]  2020.
- [BO09] Fr ed eric Bourgeois and Alexandru Oancea. Symplectic homology, autonomous hamiltonians, and morse-bott moduli spaces. *Duke mathematical journal*, 146(1):71–174, 2009.
- [BO17] Fr ed eric Bourgeois and Alexandru Oancea. S^1 -equivariant symplectic homology and linearized contact homology. *Int. Math. Res. Not. IMRN*, (13):3849–3937, 2017.
- [BP21] Tobias Barthel and Piotr Pstragowski. Morava K-theory and Filtrations by Powers. *arXiv e-prints*, page arXiv:2111.06379, November 2021.
- [CFHW96] K. Cieliebak, A. Floer, H. Hofer, and K. Wysocki. Applications of symplectic homology. II. Stability of the action spectrum. *Math. Z.*, 223(1):27–45, 1996.
- [CHHL23] Dan Cristofaro-Gardiner, Umberto Hryniewicz, Michael Hutchings, and Hui Liu. Proof of Hofer-Wysocki-Zehnder’s two or infinity conjecture. *arXiv e-prints*, page arXiv:2310.07636, October 2023.
- [Cie02] Kai Cieliebak. Handle attaching in symplectic homology and the chord conjecture. *Journal of the European Mathematical Society*, 4:115–142, 2002.
- [CK23] Laurent C ot e and Yusuf Bari s Kartal. Equivariant Floer homotopy via Morse-Bott theory. *arXiv e-prints*, page arXiv:2309.15089, 2023.

- [Gan12] Sheel Ganatra. *Symplectic Cohomology and Duality for the Wrapped Fukaya Category*. ProQuest LLC, Ann Arbor, MI, 2012. Thesis (Ph.D.)—Massachusetts Institute of Technology.
- [Gan19] Sheel Ganatra. Cyclic homology, S^1 -equivariant Floer cohomology, and Calabi-Yau structures. *arXiv e-prints*, page arXiv:1912.13510, December 2019.
- [HHLN23] Rune Haugseng, Fabian Hebestreit, Sil Linskens, and Joost Nuiten. Lax monoidal adjunctions, two-variable fibrations and the calculus of mates. *Proceedings of the London Mathematical Society*, 127(4):889–957, 2023.
- [Lar21] Tim Large. Spectral fukaya categories for liouville manifolds. 2021.
- [PS24] Noah Porcelli and Ivan Smith. Bordism of flow modules and exact lagrangians. *arXiv:2401.11766*, 2024.
- [Seg68] Graeme Segal. Equivariant K -theory. *Inst. Hautes Études Sci. Publ. Math.*, (34):129–151, 1968.
- [Sei08] Paul Seidel. A biased view of symplectic cohomology. In *Current developments in mathematics, 2006*, volume 2006, pages 211–254. International Press of Boston, 2008.
- [Tor23] Takeshi Torii. A perfect pairing for monoidal adjunctions. *Proceedings of the American Mathematical Society*, 151(12):5069–5080, 2023.
- [Tre19] David Treumann. Complex k -theory of mirror pairs. *arXiv preprint arXiv:1909.03018*, 2019.
- [Wur91] Urs Würgler. Morava K -theories: a survey. In *Algebraic topology Poznań 1989*, volume 1474 of *Lecture Notes in Math.*, pages 111–138. Springer, Berlin, 1991.
- [Zha19] Jingyu Zhao. Periodic symplectic cohomologies. *Journal of Symplectic Geometry*, 17(5):1513–1578, 2019.

UNIVERSITÄT BONN, MATHEMATICAL INSTITUTE, BONN, GERMANY
Email address: lcote@math.uni-bonn.de

SCHOOL OF MATHEMATICS, UNIVERSITY OF EDINBURGH, EDINBURGH, UK
Email address: ykartal@ed.ac.uk