

Robust inference for linear regression models with possibly skewed error distribution

Amarnath Nandy, Ayanendranath Basu and Abhik Ghosh*

Interdisciplinary Statistical Research Unit

Indian Statistical Institute, Kolkata, India

amarnath_r@isical.ac.in, ayanbasu@isical.ac.in, abhik.ghosh@isical.ac.in

Abstract

Traditional methods for linear regression generally assume that the underlying error distribution, equivalently the distribution of the responses, is normal. Yet, sometimes real life response data may exhibit a skewed pattern, and assuming normality would not give reliable results in such cases. This is often observed in cases of some biomedical, behavioral, socio-economic and other variables. In this paper, we propose to use the class of skew normal (SN) distributions, which also includes the ordinary normal distribution as its special case, as the model for the errors in a linear regression setup and perform subsequent statistical inference using the popular and robust minimum density power divergence approach to get stable insights in the presence of possible data contamination (e.g., outliers). We provide the asymptotic distribution of the proposed estimator of the regression parameters and also propose robust Wald-type tests of significance for these parameters. We provide an influence function analysis of these estimators and test statistics, and also provide level and power influence functions. Numerical verification including simulation studies and real data analysis is provided to substantiate the theory developed.

Key words: Linear regression; Skew normal (SN) distribution; Density power divergence; Wald-type tests; Influence function; Robustness

1 Introduction

In real life problems, we are often interested in investigating how the different sources of variation (covariates or explanatory variables) affect the desired output (response variable). The linear regression model (LRM) is one of the most popular tools for doing so, where we assume that the covariates (or transformed covariates) have linear relationships with the response. In traditional linear regression,

*Corresponding author

our canonical assumption is that the errors are normally distributed. But in real life, for example in case of real data related to health and behavioral sciences and many related variables (e.g., medical expenditure), skewed patterns may be observed [13]. In such cases, it may be necessary to assume a skewed model distribution for the random error component in the LRM; a few previous attempts have used gamma or generalized gamma distributions for this purpose (see [3], [31], [33]) which are always skewed. However, observed skewness patterns in response distribution may arise due to data contamination (where the true underlying distribution is symmetric) or they can be the inherent characteristics of the response variables. So, we propose to use the class of skew normal distributions – which also includes the ordinary normal distribution as a special case – as the model for the errors in a linear regression model. Recently, several authors have used this skew-normal distribution in regression models where response variables are skewed in nature but all these attempts have used the classical non-robust likelihood based inference procedures (see [2], [17], [27], [28], [30]).

Yet, it is well known that likelihood based inference, for all its efficiency properties, may perform very poorly in terms of robustness against data contamination and other types of anomalies which are not uncommon in modern complex datasets. There can be additional (model-specific) outliers even when the underlying distribution is skewed so that the classical likelihood based inference would again fail to provide correct inference. It is generally difficult, sometimes impossible, to identify and remove such anomalies in an objective manner. So, a robust procedure that automatically takes care of the effects of such data contamination to produce stable inference would be an important and useful alternative in case of practical applications [16, 35, 36, 39]. Some recent attempts to model the outliers in such situations have been made using mixture error distributions; see, [12, 15, 29, 40].

In this paper, we have presented a robust method of fitting the linear regression model under skew normal errors based on the minimization of the density power divergence (Basu et al. (1998) [9]). The corresponding minimum density power divergence estimator (MDPDE for short) is indexed by a single tuning parameter α , which provides a trade-off between the efficiency and the robustness of the method. Recently, Nandy et al. [32] have implemented this method for the skew normal distribution in case of homogeneous data. In our paper, we have extended this idea for the regression set up, where the data are independent but not homogeneous. We have found robust estimators of the regression coefficients as well as the parameters of the error distribution. In addition, we have provided the asymptotic distribution of the estimators and constructed a robust Wald-type test for testing the significance of the regression coefficients. We have also presented an influence function analysis of this method and demonstrated the boundedness of the same for positive α .

The rest of the paper is organized as follows: in Section 2, we have defined our MDPDE for the skew regression model and provided the formula for its asymptotic variance as also an influence function analysis. In Section 3, we have proposed a robust Wald-type test based on the MDPDE and described its usefulness. We have provided the results obtained from different simulation studies

in Section 4. We have also implemented our method on two different real life datasets and we have summarized the results in Section 5. These examples are instructive and help to understand how some of the MDPDEs provide stable inference under the presence of anomalous observations. We have also discussed a method for obtaining the optimal value of the tuning parameter in Section 6. Finally, Section 7 provides some concluding remarks.

2 The model and the proposed estimator

2.1 The linear regression model with skew-normal errors

We first give a brief description of the linear regression model with skew-normal (SN) errors. The skew normal (SN) distribution (see [5], [6], [7]) is defined in terms of three parameters, namely the location parameter $\mu \in \mathbb{R}$, the scale parameter $\sigma \in \mathbb{R}^+$ and the skewness parameter $\gamma \in \mathbb{R}$, and is denoted by $SN(\mu, \sigma, \gamma)$. In particular, if $\mu = 0$ and $\sigma = 1$, it is referred to as the standard SN distribution and is denoted by $SN(\gamma)$. The probability density function (pdf) and the cumulative distribution function (cdf) of the $SN(\mu, \sigma, \gamma)$ distribution are given, respectively, as

$$f(x, \boldsymbol{\eta}) = \frac{2}{\sigma} \phi\left(\frac{x - \mu}{\sigma}\right) \Phi\left(\gamma \frac{x - \mu}{\sigma}\right), \quad x \in \mathbb{R}, \quad (1)$$

$$F(x, \boldsymbol{\eta}) = \Phi\left(\frac{x - \mu}{\sigma}\right) - 2T\left(\frac{x - \mu}{\sigma}, \gamma\right), \quad x \in \mathbb{R}, \quad (2)$$

where $\boldsymbol{\eta} = (\mu, \sigma, \gamma)^T$ is the vector of unknown parameters, ϕ and Φ are the pdf and cdf of the standard normal distribution, respectively, and $T(h, a)$ is the Owen's function defined as

$$T(h, a) = \frac{1}{2\pi} \int_0^a \frac{e^{-\frac{h^2}{2}(1+x^2)}}{1+x^2} dx, \quad h, a \in \mathbb{R}.$$

The mean, variance and skewness (γ_1) of a random variable X having the $SN(\mu, \sigma, \gamma)$ distribution are given by $E(X) = \mu + \sigma\delta\sqrt{\frac{2}{\pi}}$, $Var(X) = \sigma^2\left(1 - \frac{2\delta^2}{\pi}\right)$ and $\gamma_1 = \frac{1}{2}(4 - \pi)\gamma^3\left(\frac{\pi}{2} + (\frac{\pi}{2} - 1)\gamma^2\right)^{-3/2}$, with $\delta = \gamma(1 + \gamma^2)^{-1/2}$. Clearly, the SN distribution is positively and negatively skewed according to the sign of the parameter γ . For the particular case $\gamma = 0$, the SN distribution $SN(\mu, \sigma, 0)$ has skewness zero and coincides with the ordinary (symmetric) normal distribution, $N(\mu, \sigma^2)$, having mean μ and variance σ^2 .

Now, let us consider the skew normal linear regression model

$$y_i = \mathbf{x}_i^T \boldsymbol{\beta} + \epsilon_i, \quad i = 1, 2, \dots, n \quad (3)$$

where $\epsilon_i \sim SN(0, \sigma, \gamma)$, independently. Note that here $\mathbf{x}_i^T \boldsymbol{\beta}$ is not the conditional mean of y_i , but

is the location parameter of the distribution of y_i given \mathbf{x}_i . Here, $\mathbf{x}_i = (x_{i1}, x_{i2}, \dots, x_{ip})$ is the p -dimensional covariate vector corresponding to i -th observation and $\boldsymbol{\beta} = (\beta_1, \beta_2, \dots, \beta_p)$ is the vector of regression coefficients; σ and γ are the scale and skewness parameters of the error distribution. In our case, $\boldsymbol{\theta} = (\boldsymbol{\beta}, \sigma, \gamma)$ and $f_i(y, \boldsymbol{\theta})$ is the density of $SN(\mu_i = \mathbf{x}_i^T \boldsymbol{\beta}, \sigma, \gamma)$ given by

$$f_i(y, \boldsymbol{\theta}) = \frac{2}{\sigma} \phi\left(\frac{y - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right) \Phi\left(\gamma \frac{y - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right), \quad y \in \mathbb{R}. \quad (4)$$

2.2 The minimum density power divergence estimator (MDPDE)

The density power divergence and the corresponding MDPDE for independent and identically distributed (IID) data was first proposed by Basu et al. [9], which was later extended for non-homogeneous data (independent but not identically distributed) by Ghosh and Basu [18], where they considered the standard linear regression model. Later, this technique has been applied to GLM and survival data models [20, 21, 23, 24]. The DPD family [9] is indexed by a single tuning parameter $\alpha \geq 0$, controlling the trade-off between robustness and efficiency of the procedure. For two densities g and f , both being absolutely continuous with respect to some common dominating measure μ , the DPD measure between g and f is defined as

$$d_\alpha(g, f) = \int \left\{ f^{1+\alpha} - \left(1 + \frac{1}{\alpha}\right) g f^\alpha + \frac{1}{\alpha} g^{1+\alpha} \right\}, \quad \text{if } \alpha > 0, \quad (5)$$

$$d_0(g, f) = \lim_{\alpha \downarrow 0} d_\alpha(g, f) = \int g \log \frac{g}{f}. \quad (6)$$

Note that, the DPD at $\alpha = 0$ is nothing but the well-known Kullback-Leiber divergence (KLD) associated with the likelihood approach. Let Y_1, Y_2, \dots, Y_n represent independent and identically distributed observations drawn from the true distribution G (having pdf g). Let $\mathcal{F} = \{F(\cdot, \boldsymbol{\theta}) : \boldsymbol{\theta} \in \Theta \subset \mathbb{R}^p\}$ be the family of distributions modeling it, with $f(\cdot, \boldsymbol{\theta})$ being the pdf of $F(\cdot, \boldsymbol{\theta})$. Minimization of the divergence $d_\alpha(g, f(\cdot, \boldsymbol{\theta}))$ over the parameter space Θ generates the ‘‘best fitting’’ target parameter. As g will always be unknown, we need an empirical estimate of this divergence which may be minimized to obtain the parameter estimate. Note that from (5) one gets

$$d_\alpha(g, f(\cdot, \boldsymbol{\theta})) = \int f^{1+\alpha}(y, \boldsymbol{\theta}) dy - \left(1 + \frac{1}{\alpha}\right) \int f^\alpha(y, \boldsymbol{\theta}) dG(y) + K,$$

where the last term $K = \frac{1}{\alpha} \int g^{1+\alpha}(y) d(y)$ is independent of the parameter $\boldsymbol{\theta}$ and has no role in the optimization. Now, the second term in the above equation can be estimated just by plugging in the empirical estimate of G , namely the empirical cdf G_n obtained from the sample Y_1, \dots, Y_n , and then the MDPDE can be obtained by minimizing the empirical objective function

$$H_{n,\alpha}(\boldsymbol{\theta}) = \int f^{1+\alpha}(y, \boldsymbol{\theta}) dy - \left(1 + \frac{1}{\alpha}\right) \frac{1}{n} \sum_{i=1}^n f^\alpha(Y_i, \boldsymbol{\theta}).$$

In the regression case, Y_1, Y_2, \dots, Y_n are n observations (realizations of the response variable), where $Y_i \sim g_i$ independently. Here, we want to model g_i by $f_i(\cdot, \boldsymbol{\theta})$, where $f_i(\cdot, \boldsymbol{\theta})$ is as defined in (4). Although the distribution of each Y_i is distinct, they share the common parameter $\boldsymbol{\theta}$. However, here the model density is different for each Y_i , so we need to calculate the divergence between data and model separately for each point. Subsequently, we take the average of these individual divergences and we minimize it to estimate $\boldsymbol{\theta}$. So, if $d_\alpha(g_i, f_i(\cdot, \boldsymbol{\theta}))$ be the density power divergence measure between the data and model for the i -th case, then we minimize the objective function

$$\frac{1}{n} \sum_{i=1}^n d_\alpha(g_i, f_i(\cdot, \boldsymbol{\theta}))$$

with respect to $\boldsymbol{\theta} \in \Theta$. In the presence of only one data point Y_i from g_i , the best possible density estimate of g_i is the degenerate density which puts the whole mass on Y_i . Hence, the divergence between g_i and $f_i(\cdot, \boldsymbol{\theta})$ is estimated as

$$\int f_i^{1+\alpha}(y, \boldsymbol{\theta}) dy - \left(1 + \frac{1}{\alpha}\right) f_i^\alpha(Y_i, \boldsymbol{\theta}) + K \quad (7)$$

where K is independent of $\boldsymbol{\theta}$. Dropping the constant, our entire objective function then becomes

$$H_{n,\alpha}(\boldsymbol{\theta}) = \frac{1}{n} \sum_{i=1}^n \left[\int f_i^{1+\alpha}(y, \boldsymbol{\theta}) dy - \left(1 + \frac{1}{\alpha}\right) f_i^\alpha(Y_i, \boldsymbol{\theta}) \right] = \frac{1}{n} \sum_{i=1}^n V_i(Y_i, \boldsymbol{\theta}). \quad (8)$$

Note that, as $\alpha \rightarrow 0$, the corresponding objective function, which is to be minimized, becomes

$$\frac{1}{n} \sum_{i=1}^n [-\ln(f_i(Y_i, \boldsymbol{\theta}))]. \quad (9)$$

Thus the MDPDE with $\alpha = 0$ coincides with the maximum likelihood estimator (MLE) which is obtained by solving the score equation $\sum_{i=1}^n u_i(Y_i, \boldsymbol{\theta}) = \mathbf{0}$, where $u_i(Y_i, \boldsymbol{\theta}) = \frac{\partial \ln(f_i(Y_i, \boldsymbol{\theta}))}{\partial \boldsymbol{\theta}}$ is the score function. In general, the estimating equation for obtaining the MDPDE becomes

$$\sum_{i=1}^n \left[f_i^\alpha(Y_i, \boldsymbol{\theta}) u_i(Y_i, \boldsymbol{\theta}) - \int f_i^{1+\alpha}(y, \boldsymbol{\theta}) u_i(y, \boldsymbol{\theta}) dy \right] = \mathbf{0}. \quad (10)$$

Note that this defines an M-estimator with a model dependent ψ function. In our case, the objective function becomes

$$H_{n,\alpha}(\boldsymbol{\theta}) = \frac{1}{n} \sum_{i=1}^n \left[\int_{-\infty}^{\infty} \left(\frac{2}{\sigma}\right)^{1+\alpha} \phi^{1+\alpha}\left(\frac{y - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right) \Phi^{1+\alpha}\left(\gamma \frac{y - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right) dy - \left(1 + \frac{1}{\alpha}\right) \left(\frac{2}{\sigma}\right)^{\alpha} \phi^{\alpha}\left(\frac{Y_i - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right) \Phi^{\alpha}\left(\gamma \frac{Y_i - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right) \right]. \quad (11)$$

By standard differentiation, we get the estimating equations of our MDPDE to be

$$\sum_{i=1}^n \mathbf{u}_i(Y_i, \boldsymbol{\theta}) \left(\frac{2}{\sigma}\right)^{\alpha} \phi^{\alpha}\left(\frac{Y_i - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right) \Phi^{\alpha}\left(\gamma \frac{Y_i - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right) = \sum_{i=1}^n \boldsymbol{\xi}_{i,\alpha}(\boldsymbol{\theta}), \quad (12)$$

where $\mathbf{u}_i(y, \boldsymbol{\theta})$ is the score function of the SN distribution for the i -th case and has the form

$$\mathbf{u}_i(y, \boldsymbol{\theta}) = \left(\left(\frac{y - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma^2} - \frac{\gamma \phi\left(\frac{\gamma y - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right)}{\sigma \Phi\left(\frac{\gamma y - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right)} \right) \mathbf{x}_i, \frac{(y - \mathbf{x}_i^T \boldsymbol{\beta})^2}{\sigma^3} - \gamma \frac{(y - \mathbf{x}_i^T \boldsymbol{\beta})}{\sigma^2} \frac{\phi\left(\frac{\gamma y - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right)}{\Phi\left(\frac{\gamma y - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right)} - \frac{1}{\sigma}, \frac{(y - \mathbf{x}_i^T \boldsymbol{\beta})}{\sigma} \frac{\phi\left(\frac{\gamma y - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right)}{\Phi\left(\frac{\gamma y - \mathbf{x}_i^T \boldsymbol{\beta}}{\sigma}\right)} \right)^T,$$

and

$$\boldsymbol{\xi}_{i,\alpha}(\boldsymbol{\theta}) = \int \mathbf{u}_i(y, \boldsymbol{\theta}) f_i^{1+\alpha}(y, \boldsymbol{\theta}) dy = \left(\xi_{i,\alpha}^{(1)}(\boldsymbol{\theta}) \quad \xi_{i,\alpha}^{(2)}(\boldsymbol{\theta}) \quad \xi_{i,\alpha}^{(3)}(\boldsymbol{\theta}) \right)^T. \quad (13)$$

Clearly, the estimating equation presented in (12) is very complicated and the involved integral to be computed numerically. Here, we have used the R function `optim` to minimize $H_{n,\alpha}(\boldsymbol{\theta})$.

2.3 The asymptotic distribution and the standard error of the MDPDE

The asymptotic distribution of the MDPDE for the non-homogeneous case was developed by Ghosh and Basu (2013) [18]. They demonstrated that, under some regularity conditions, the minimum DPD estimators are \sqrt{n} -consistent and asymptotically normal. We have used the same theory for our present case. At a given $\alpha \geq 0$, if the corresponding MDPDE based on a random sample of size n is denoted by $\widehat{\boldsymbol{\theta}}_{\alpha,n}$, and the true parameter value is $\boldsymbol{\theta}_0$, we have

$$\sqrt{n} \boldsymbol{\Omega}_n^{-\frac{1}{2}} \boldsymbol{\Psi}_n \left(\widehat{\boldsymbol{\theta}}_{\alpha,n} - \boldsymbol{\theta}_0 \right) \xrightarrow{\mathcal{D}} N_{p+2}(\mathbf{0}_{p+2}, \mathbf{I}_{p+2}),$$

where \mathbf{I}_b is the identity matrix of dimension b , $\mathbf{0}_b$ is a b -dimensional vector with all entries zero, $\boldsymbol{\Psi}_n = \frac{1}{n} \sum_{i=1}^n \mathbf{J}_{\alpha}^{(i)}(\boldsymbol{\theta})$, $\boldsymbol{\Omega}_n = \frac{1}{n} \sum_{i=1}^n \text{Var}[\nabla V_i(Y_i, \boldsymbol{\theta})] = \frac{1}{n} \sum_{i=1}^n \mathbf{K}_{\alpha}^{(i)}(\boldsymbol{\theta})$ (say), and ∇ represents derivative with respect to $\boldsymbol{\theta}$. Here, for SN distributions with $\boldsymbol{\theta} = (\boldsymbol{\beta}, \sigma, \gamma)^T$, the $(p+2) \times (p+2)$ matrices $\mathbf{K}_{\alpha}^{(i)}(\boldsymbol{\theta})$ and $\mathbf{J}_{\alpha}^{(i)}(\boldsymbol{\theta})$ are given by

$$\mathbf{J}_\alpha^{(i)}(\boldsymbol{\theta}) = \int \mathbf{u}_i(y, \boldsymbol{\theta}) \mathbf{u}_i^T(y, \boldsymbol{\theta}) f_i^{1+\alpha}(y, \boldsymbol{\theta}) dy = \begin{pmatrix} N_{i,\alpha}^{(11)}(\boldsymbol{\theta}) & N_{i,\alpha}^{(12)}(\boldsymbol{\theta}) & N_{i,\alpha}^{(13)}(\boldsymbol{\theta}) \\ N_{i,\alpha}^{(12)}(\boldsymbol{\theta}) & N_{i,\alpha}^{(22)}(\boldsymbol{\theta}) & N_{i,\alpha}^{(23)}(\boldsymbol{\theta}) \\ N_{i,\alpha}^{(13)}(\boldsymbol{\theta}) & N_{i,\alpha}^{(23)}(\boldsymbol{\theta}) & N_{i,\alpha}^{(33)}(\boldsymbol{\theta}) \end{pmatrix}, \quad (14)$$

$$\begin{aligned} \mathbf{K}_\alpha^{(i)}(\boldsymbol{\theta}) &= \int \mathbf{u}_i(y, \boldsymbol{\theta}) \mathbf{u}_i^T(y, \boldsymbol{\theta}) f_i^{1+2\alpha}(y, \boldsymbol{\theta}) dy - \boldsymbol{\xi}_{i,\alpha}(\boldsymbol{\theta}) \boldsymbol{\xi}_{i,\alpha}^T(\boldsymbol{\theta}) \\ &= \begin{pmatrix} N_{i,2\alpha}^{(11)}(\boldsymbol{\theta}) - \xi_{i,\alpha}^{(1)}(\boldsymbol{\theta}) \xi_{i,\alpha}^{(1)T}(\boldsymbol{\theta}) & N_{i,2\alpha}^{(12)}(\boldsymbol{\theta}) - \xi_{i,\alpha}^{(1)}(\boldsymbol{\theta}) \xi_{i,\alpha}^{(2)}(\boldsymbol{\theta}) & N_{i,2\alpha}^{(13)}(\boldsymbol{\theta}) - \xi_{i,\alpha}^{(1)}(\boldsymbol{\theta}) \xi_{i,\alpha}^{(3)}(\boldsymbol{\theta}) \\ N_{i,2\alpha}^{(12)}(\boldsymbol{\theta}) - \xi_{i,\alpha}^{(1)}(\boldsymbol{\theta}) \xi_{i,\alpha}^{(2)}(\boldsymbol{\theta}) & N_{i,2\alpha}^{(22)}(\boldsymbol{\theta}) - \xi_{i,\alpha}^{(2)}(\boldsymbol{\theta})^2 & N_{i,2\alpha}^{(23)}(\boldsymbol{\theta}) - \xi_{i,\alpha}^{(2)}(\boldsymbol{\theta}) \xi_{i,\alpha}^{(3)}(\boldsymbol{\theta}) \\ N_{i,2\alpha}^{(13)}(\boldsymbol{\theta}) - \xi_{i,\alpha}^{(1)}(\boldsymbol{\theta}) \xi_{i,\alpha}^{(3)}(\boldsymbol{\theta}) & N_{i,2\alpha}^{(23)}(\boldsymbol{\theta}) - \xi_{i,\alpha}^{(2)}(\boldsymbol{\theta}) \xi_{i,\alpha}^{(3)}(\boldsymbol{\theta}) & N_{i,2\alpha}^{(33)}(\boldsymbol{\theta}) - \xi_{i,\alpha}^{(3)}(\boldsymbol{\theta})^2 \end{pmatrix}, \quad (15) \end{aligned}$$

with $\boldsymbol{\xi}_{i,\alpha}(\boldsymbol{\theta})$ and $f_i(y, \boldsymbol{\theta})$ are as defined in (13) and (4), respectively, and

$$\begin{aligned} N_{i,\alpha}^{(11)}(\boldsymbol{\theta}) &= \int_{-\infty}^{\infty} \left(\left(\frac{y - \mu_i}{\sigma^2} \right) - \frac{\gamma \phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)}{\sigma \Phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)} \right)^2 \mathbf{x}_i \mathbf{x}_i^T f_i^{1+\alpha}(y, \boldsymbol{\theta}) dy \\ N_{i,\alpha}^{(22)}(\boldsymbol{\theta}) &= \int_{-\infty}^{\infty} \left(\frac{(y - \mu_i)^2}{\sigma^3} - \frac{1}{\sigma} - \frac{\gamma(y - \mu_i)}{\sigma^2} \frac{\phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)}{\Phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)} \right)^2 f_i^{1+\alpha}(y, \boldsymbol{\theta}) dy \\ N_{i,\alpha}^{(33)}(\boldsymbol{\theta}) &= \int_{-\infty}^{\infty} \left(\frac{y - \mu_i}{\sigma} \frac{\phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)}{\Phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)} \right)^2 f_i^{1+\alpha}(y, \boldsymbol{\theta}) dy \\ N_{i,\alpha}^{(12)}(\boldsymbol{\theta}) &= \int_{-\infty}^{\infty} \left(\left(\frac{y - \mu_i}{\sigma^2} \right) - \frac{\gamma \phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)}{\sigma \Phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)} \right) \left(\frac{(y - \mu_i)^2}{\sigma^3} - \frac{1}{\sigma} - \frac{\gamma(y - \mu_i)}{\sigma^2} \frac{\phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)}{\Phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)} \right) \mathbf{x}_i f_i^{1+\alpha}(y, \boldsymbol{\theta}) dy \\ N_{i,\alpha}^{(13)}(\boldsymbol{\theta}) &= \int_{-\infty}^{\infty} \left(\left(\frac{y - \mu_i}{\sigma^2} \right) - \frac{\gamma \phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)}{\sigma \Phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)} \right) \left(\frac{y - \mu_i}{\sigma} \frac{\phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)}{\Phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)} \right) \mathbf{x}_i f_i^{1+\alpha}(y, \boldsymbol{\theta}) dy \\ N_{i,\alpha}^{(23)}(\boldsymbol{\theta}) &= \int_{-\infty}^{\infty} \left(\frac{(y - \mu_i)^2}{\sigma^3} - \frac{1}{\sigma} - \frac{\gamma(y - \mu_i)}{\sigma^2} \frac{\phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)}{\Phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)} \right) \left(\frac{y - \mu_i}{\sigma} \frac{\phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)}{\Phi\left(\gamma \frac{y - \mu_i}{\sigma}\right)} \right) f_i^{1+\alpha}(y, \boldsymbol{\theta}) dy \end{aligned}$$

where $\mu_i = \mathbf{x}_i^T \boldsymbol{\beta}$. Here also we can see that the above integrals do not have closed forms, so we must numerically compute them to obtain the asymptotic covariance matrix at different given values of α and $\boldsymbol{\theta}$.

The above asymptotic variance formula can also help us to obtain the standard errors of the MDPDEs in any practical application. For the MDPDE $\hat{\boldsymbol{\theta}}_{\alpha,n} = (\hat{\boldsymbol{\beta}}_{\alpha,n}, \hat{\sigma}_{\alpha,n}, \hat{\gamma}_{\alpha,n})^T$, obtained on the basis of a sample of size n , its standard errors are obtained as $\sqrt{\text{diag}(\boldsymbol{\Sigma}_{\alpha,n}(\hat{\boldsymbol{\theta}}_{\alpha,n}))/n}$, where $\boldsymbol{\Sigma}_{\alpha,n}(\hat{\boldsymbol{\theta}}_{\alpha,n})$ is consistent for $\boldsymbol{\Sigma}_\alpha(\boldsymbol{\theta}_0) = \boldsymbol{\Psi}_n^{-1} \boldsymbol{\Omega}_n \boldsymbol{\Psi}_n^{-1}$.

2.4 Influence function analysis of the MDPDE

The robustness of an estimator can be theoretically examined through the classical influence function (IF) analysis [25]. The IF measures the asymptotic (standardized) bias of the estimator caused by an infinitesimal contamination at a distant contamination point (say y). Therefore, the boundedness of the IF over the contamination point y restricts the extent of possible bias finitely for the corresponding estimator indicating its robust nature. On the other hand, an unbounded IF indicates possibly unbounded bias and non-robustness of the estimator. Further, with similar intuition, the supremum of the absolute IF taken over all possible contamination points naturally indicates the extent of (bias) robustness of the corresponding estimator.

From the theory of the M-estimators [25] or the general MDPDE [9, 10], one can obtain the influence function of the MDPDE functional under the IID set up. Ghosh and Basu [18] have studied the influence functions for the non-homogeneous set up [18]. We will present an influence function analysis along these lines for the estimators of our skew normal linear regression model.

Let, G_i denotes the true distribution of Y_i and $\mathbf{T}_\alpha(G_1, G_2, \dots, G_n)$ be the MDPDE functional defined as the minimizer of (8). Now, we consider the contaminated density $g_{i,\epsilon} = (1 - \epsilon)g_i + \epsilon\delta_{y_i}$, where δ_{y_i} is the degenerate distribution at the point of contamination y_i and let $G_{i,\epsilon}$ be the corresponding distribution function, $i = 1, 2, \dots, n$. Let $\boldsymbol{\theta} = \mathbf{T}_\alpha(G_1, G_2, \dots, G_n)$ and let $\boldsymbol{\theta}_\epsilon^{i_0} = \mathbf{T}_\alpha(G_1, \dots, G_{i_0-1}, G_{i_0,\epsilon}, \dots, G_n)$ be the minimum density power divergence functional with contamination only in the i_0 -th direction, where $G_{i_0,\epsilon} = (1 - \epsilon)G_{i_0} + \epsilon\Delta_{y_{i_0}}$ with $\Delta_{y_{i_0}}$ being the distribution of $\delta_{y_{i_0}}$. Then, along the lines of Ghosh and Basu [18], we can get the influence function of the functional with contamination only in i_0 -th direction as follows:

$$IF_{i_0}(y_{i_0}, T_\alpha, \boldsymbol{\theta}) = \boldsymbol{\Psi}_n^{-1} \frac{1}{n} [\mathbf{u}_{i_0}(y_{i_0}, \boldsymbol{\theta}) f_{i_0}^\alpha(y_{i_0}, \boldsymbol{\theta}) - \boldsymbol{\xi}_{i_0,\alpha}(\boldsymbol{\theta})], \quad (16)$$

where $\boldsymbol{\xi}_{i,\alpha}(\boldsymbol{\theta})$ and $\boldsymbol{\Psi}_n$ are as defined in Section 2.3. On the other hand, if we consider contamination in all directions, the form of the influence function turns out to be

$$IF(y_1, \dots, y_n, T_\alpha, \boldsymbol{\theta}) = \boldsymbol{\Psi}_n^{-1} \frac{1}{n} \sum_{i=1}^n [\mathbf{u}_i(y_i, \boldsymbol{\theta}) f_i^\alpha(y_i, \boldsymbol{\theta}) - \boldsymbol{\xi}_{i,\alpha}(\boldsymbol{\theta})]. \quad (17)$$

2.5 Illustration for a simple set-up

Here we have considered a linear model with a single covariate. Our model is $y_i = x_i\beta + \epsilon_i$, $i = 1, 2, \dots, n$, where $\epsilon_i \sim SN(0, \sigma, \gamma)$ and x is the covariate. Thus, y_i 's are distributed as $SN(\mu_i = x_i\beta, \sigma, \gamma)$, independently of one another. Using the formulas discussed in Section 2.3, we can find the asymptotic relative efficiency (ARE) of the proposed MDPDE which are presented in Table 1 for different values of $\boldsymbol{\theta} = (\beta, \sigma, \gamma)^T$ for the SN linear regression model. We have considered three different

SN distributions for the random error component. From Table 1, we observe that ARE of β and γ are same when we take the error distribution to be $SN(0, 1, 0)$ and $SN(0, 4, 0)$, i.e., we take the estimate of $\gamma = 0$. This is not an accident and we will show later that the relevant estimated quantities are functionally identical in terms of the estimates of σ and β when $\gamma = 0$.

Table 1: Asymptotic Relative Efficiency of the MDPDEs of $(\beta, \sigma, \gamma)^T$ for different SN error distributions. The $\alpha = 0$ case corresponds to the MLE

Error Distribution		α					
		0(MLE)	0.1	0.3	0.5	0.7	1
$SN(0, 1, -2)$	β	100	98.66	91.30	82.25	73.68	62.75
	σ	100	97.52	85.11	71.84	60.94	48.72
	γ	100	98.18	86.76	71.81	58.08	42.35
$SN(0, 1, 0)$	β	100	98.75	92.03	83.57	75.30	64.45
	σ	100	97.49	85.38	72.79	63.01	53.41
	γ	100	98.75	92.03	83.57	75.30	64.45
$SN(0, 4, 0)$	β	100	98.75	92.03	83.57	75.30	64.45
	σ	100	97.49	85.38	72.79	63.01	53.41
	γ	100	98.75	92.03	83.57	75.30	64.45
$SN(0, 1, 2)$	β	100	98.63	91.11	81.85	73.12	62.04
	σ	100	97.98	86.90	74.08	63.13	50.68
	γ	100	97.90	85.13	69.21	55.36	40.15

In our example, the score function becomes

$$\mathbf{u}_i(y, \boldsymbol{\theta}) = \left(\left(\frac{y - \mu_i}{\sigma^2} - \frac{\gamma \phi \left(\gamma \frac{y - \mu_i}{\sigma} \right)}{\sigma \Phi \left(\gamma \frac{y - \mu_i}{\sigma} \right)} \right) x_i, \frac{(y - \mu_i)^2}{\sigma^3} - \gamma \frac{(y - \mu_i)}{\sigma^2} \frac{\phi \left(\gamma \frac{y - \mu_i}{\sigma} \right)}{\Phi \left(\gamma \frac{y - \mu_i}{\sigma} \right)} - \frac{1}{\sigma}, \frac{(y - \mu_i)}{\sigma} \frac{\phi \left(\gamma \frac{y - \mu_i}{\sigma} \right)}{\Phi \left(\gamma \frac{y - \mu_i}{\sigma} \right)} \right)^T, \quad (18)$$

where $\mu_i = x_i \beta$, $\boldsymbol{\theta} = (\beta, \sigma, \gamma)$. When the true value of γ is 0, the integrals within the matrices $\boldsymbol{\Omega}_n$ and $\boldsymbol{\Psi}_n$ become simpler and can be calculated easily. In this case, we obtain, after some algebra, the final form of $\boldsymbol{\Psi}_n$ and $\boldsymbol{\Omega}_n$ as:

$$\boldsymbol{\Psi}_n = \sigma^{-\alpha} \begin{pmatrix} \psi_{11,\alpha} & 0 & \psi_{13,\alpha} \\ 0 & \psi_{22,\alpha} & 0 \\ \psi_{31,\alpha} & 0 & \psi_{33,\alpha} \end{pmatrix}, \quad (19)$$

$$\boldsymbol{\Omega}_n = \sigma^{-2\alpha} \begin{pmatrix} \omega_{11,\alpha} & 0 & \omega_{13,\alpha} \\ 0 & \omega_{22,\alpha} & 0 \\ \omega_{31,\alpha} & 0 & \omega_{33,\alpha} \end{pmatrix}, \quad (20)$$

where

$$\begin{aligned}
\psi_{11,\alpha} &= \phi(\alpha) \frac{\sum_{i=1}^n x_i^2}{n}, & \psi_{13,\alpha} &= \psi_{31,\alpha} = \phi(\alpha) \sqrt{\frac{2}{\pi}} \bar{x}, \\
\psi_{22,\alpha} &= \phi(\alpha) \frac{\alpha^2 + 2}{\alpha + 1}, & \psi_{33,\alpha} &= \phi(\alpha) \frac{2}{\pi}, \quad \phi(\alpha) = (2\pi)^{-\frac{\alpha}{2}} (1 + \alpha)^{-\frac{3}{2}}, \\
\omega_{11,\alpha} &= (2\pi)^{-\alpha} (1 + 2\alpha)^{-\frac{3}{2}} \frac{\sum_{i=1}^n x_i^2}{n}, \\
\omega_{13,\alpha} &= \omega_{31,\alpha} = (2\pi)^{-2\alpha} (1 + 2\alpha)^{-\frac{3}{2}} \sqrt{\frac{2}{\pi}} \bar{x}, \\
\omega_{22,\alpha} &= (2\pi)^{-\alpha} \left[(1 + 2\alpha)^{-\frac{5}{2}} (4\alpha^2 + 2) - \alpha^2 (1 + \alpha)^{-3} \right], \\
\omega_{33,\alpha} &= (2\pi)^{-\alpha} \frac{2}{\pi} (1 + 2\alpha)^{-\frac{3}{2}}.
\end{aligned}$$

Now, after some calculations, the asymptotic variance matrix becomes


$$\boldsymbol{\Sigma}_\alpha(\boldsymbol{\theta}) = \begin{pmatrix} \sigma_{11,\alpha} & 0 & \sigma_{13,\alpha} \\ 0 & \sigma_{22,\alpha} & 0 \\ \sigma_{31,\alpha} & 0 & \sigma_{33,\alpha} \end{pmatrix}, \quad (21)$$

where


$$\begin{aligned}
\sigma_{11,\alpha} &= \sigma(\alpha, \mathbf{x}) \frac{2}{\pi} (1 + 2\alpha)^{-\frac{3}{2}} \left(\frac{\alpha^2 + 2}{\alpha + 1} \right)^2, \\
\sigma_{13,\alpha} &= \sigma_{31,\alpha} = -\sigma(\alpha, \mathbf{x}) \sqrt{\frac{2}{\pi}} \bar{x} (1 + 2\alpha)^{-\frac{3}{2}} \left(\frac{\alpha^2 + 2}{1 + \alpha} \right)^2, \\
\sigma_{22,\alpha} &= \sigma(\alpha, \mathbf{x}) \frac{2}{\pi} s_{xx} \left[(1 + 2\alpha)^{-\frac{5}{2}} (4\alpha^2 + 2) - \alpha^2 (1 + \alpha)^{-3} \right], \\
\sigma_{33,\alpha} &= \sigma(\alpha, \mathbf{x}) \frac{2}{\pi} (1 + 2\alpha)^{-\frac{3}{2}} \left(\frac{\alpha^2 + 2}{\alpha + 1} \right)^2, \\
\sigma(\alpha, \mathbf{x}) &= \frac{(2\pi)^{-\alpha}}{s_{xx} \phi^2(\alpha) \frac{2}{\pi} \left(\frac{\alpha^2 + 2}{1 + \alpha} \right)^2}, \\
\phi(\alpha) &= (2\pi)^{-\frac{\alpha}{2}} (1 + \alpha)^{-\frac{3}{2}}, \quad s_{xx} = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2, \quad \bar{x} = \frac{1}{n} \sum_{i=1}^n x_i.
\end{aligned}$$

From the expressions above, it may be seen that the forms of the asymptotic variances at the estimated β and γ are functionally the same whenever $\gamma = 0$. This is clearly an interesting new observation in the context of the asymptotic distributions of the MDPDEs.


Further, the IFs of the three parameters α , β and γ are plotted in Figures 1a – 1f. It may be clearly observed that these influence functions are bounded for $\alpha > 0$. For $\alpha = 0$, on the other hand, the function is unbounded for all the three parameters, indicating the robustness deficiencies of the maximum likelihood estimator.




(a) IF for β in $i_0 = 10$ th direction




(b) IF for β in all directions




(c) IF for σ in $i_0 = 10$ th direction



(d) IF for σ in all directions



(e) IF for γ in $i_0 = 10$ th direction



(f) IF for γ in all directions

Figure 1: Influence functions (IFs) of the MDPDEs of the parameters in $i_0 = 10$ th direction (left panel) and in all directions (right panel), with $\theta = (3, 2, 2)$ for different α .

3 Robust Wald-type tests for the linear regression model with skew-normal errors

3.1 General theory for composite hypotheses

We now consider the problem of testing the general composite null hypothesis given by

$$H_0 : \boldsymbol{\theta} \in \Theta_0 \text{ against } H_1 : \boldsymbol{\theta} \notin \Theta_0, \quad (22)$$

where Θ_0 is a given subset of the parameter space $\Theta \in \mathbb{R}^{p+2}$. This null parameter space Θ_0 is often defined by a set of r restrictions of the form

$$\mathbf{m}(\boldsymbol{\theta}) = \mathbf{0}_r, \quad (23)$$

where $\mathbf{m} : \mathbb{R}^{p+2} \rightarrow \mathbb{R}^r$ is a known function with $r \leq p+2$. So $\Theta_0 = \{\boldsymbol{\theta} \in \Theta : \mathbf{m}(\boldsymbol{\theta}) = \mathbf{0}_r\}$. Assume that the $(p+2) \times r$ matrix

$$\mathbf{M}(\boldsymbol{\theta}) = \frac{\partial \mathbf{m}^T(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} \quad (24)$$

exists, is continuous in $\boldsymbol{\theta}$, and $\text{rank}(\mathbf{M}(\boldsymbol{\theta})) = r$. An important example is the test of significance for the regression model with the null hypothesis $H_0 : \boldsymbol{\beta} = \mathbf{0}_p$ indicating no covariate effect; this is a special case of (22) with $\mathbf{m}(\boldsymbol{\theta}) = \boldsymbol{\beta}$ so that $\Theta_0 = \{\boldsymbol{\theta} \in \Theta : \mathbf{m}(\boldsymbol{\theta}) = \boldsymbol{\beta} = \mathbf{0}_p\}$ and $\mathbf{M} = [\mathbf{I}_p; O]^T$ with $r = p$, where O is the null matrix of order $2 \times p$.

Now, the Wald-type test statistic from the MDPDE that utilizes the consistent estimator of their asymptotic variance matrix is given by

$$W_n(\widehat{\boldsymbol{\theta}}_{\alpha,n}) = n \mathbf{m}^T(\widehat{\boldsymbol{\theta}}_{\alpha,n}) \left(\mathbf{M}^T(\widehat{\boldsymbol{\theta}}_{\alpha,n}) \boldsymbol{\Sigma}_{\alpha,n}(\widehat{\boldsymbol{\theta}}_{\alpha,n}) \mathbf{M}(\widehat{\boldsymbol{\theta}}_{\alpha,n}) \right)^{-1} \mathbf{m}(\widehat{\boldsymbol{\theta}}_{\alpha,n}), \quad (25)$$

where $\widehat{\boldsymbol{\theta}}_{\alpha,n}$ is the MDPDE of $\boldsymbol{\theta}$ obtained based on a random sample of size n and tuning parameter $\alpha \geq 0$ and utilizes the matrix

$$\boldsymbol{\Sigma}_{\alpha,n}(\widehat{\boldsymbol{\theta}}_{\alpha,n}) = \frac{1}{n} \boldsymbol{\Psi}_n^{-1}(\widehat{\boldsymbol{\theta}}_{\alpha,n}) \boldsymbol{\Omega}_n(\widehat{\boldsymbol{\theta}}_{\alpha,n}) \boldsymbol{\Psi}_n^{-1}(\widehat{\boldsymbol{\theta}}_{\alpha,n}).$$

From the asymptotic distribution of the MDPDE $\widehat{\boldsymbol{\theta}}_{\alpha,n}$ in Section 2.3, it may be observed that W_n asymptotically follows a (central) chi-squared distribution, χ_r^2 , with r degrees of freedom under the null hypothesis in (22). Now, the null hypothesis in (22) is rejected if

$$W_n(\widehat{\boldsymbol{\theta}}_{\alpha,n}) > \chi_{r,\tau}^2, \quad (26)$$

where $\chi_{r,\tau}^2$ is the $100(1 - \tau)$ percentile of a chi-square distribution with r degrees of freedom.

From the general theory of Basu et al. (2018) [11], it may be observed that the MDPDE based Wald-type test is consistent at any fixed alternative. We will check its behavior in the case of contiguous alternative and try to find an approximate power function of the test. Under the contiguous alternative of the form $H_{1,n}$: $\boldsymbol{\theta} = \boldsymbol{\theta}_n$ where $\boldsymbol{\theta}_n = \boldsymbol{\theta}_0 + n^{-1/2}\mathbf{d}$ with $\boldsymbol{\theta}_0 \in \Theta_0$ and $\mathbf{d} \in \mathbb{R}^{p+2} \setminus \{\mathbf{0}_{p+2}\}$, the statistic $W_n(\widehat{\boldsymbol{\theta}}_{\alpha,n})$ asymptotically follows a non-central chi-squared distribution, denoted as $\chi_r^2(\delta)$, having r degrees of freedom and the non-centrality parameter is $\delta = \mathbf{d}^T \mathbf{Q}_\alpha(\boldsymbol{\theta}_0) \mathbf{d}$, with $\mathbf{Q}_\alpha(\boldsymbol{\theta}_0) = \mathbf{M}(\boldsymbol{\theta}_0) [\mathbf{M}^T(\boldsymbol{\theta}_0) \boldsymbol{\Sigma}_\alpha(\boldsymbol{\theta}_0) \mathbf{M}(\boldsymbol{\theta}_0)]^{-1} \mathbf{M}^T(\boldsymbol{\theta}_0)$. Based on this result, an approximate expression of the contiguous power function of the test based on $W_n(\widehat{\boldsymbol{\theta}}_{\alpha,n})$ can be calculated as

$$\phi_\alpha(\boldsymbol{\theta}_n) = 1 - G_{\chi_r^2(\delta)}(\chi_{r,\tau}^2),$$

where $G_{\chi_r^2(\delta)}$ is the cdf of the $\chi_r^2(\delta)$ distribution.

3.2 Influence analysis of the robust Wald-type test

The robustness properties of the MDPDE based Wald-type tests were discussed for non homogeneous cases by Basu et al. (2018) [11] and Ghosh and Basu (2018) [22], which also hold true for our cases. For completeness, we restate the main results briefly.

At the null distribution with $\boldsymbol{\theta}_0 \in \Theta_0$, the first order IF of the Wald-type test statistic becomes identically zero but the second order IFs (either in a single direction or along all directions) have the form

$$IF_{2,i_0}(y_{i_0}, W_n, \boldsymbol{\theta}_0) = 2IF_{i_0}(y_{i_0}, T_\alpha, \boldsymbol{\theta}_0)^T \mathbf{Q}_\alpha(\boldsymbol{\theta}_0) IF_{i_0}(y_{i_0}, T_\alpha, \boldsymbol{\theta}_0), \quad (27)$$

$$IF_2(y_1, \dots, y_n, W_n, \boldsymbol{\theta}_0) = 2IF(y_1, \dots, y_n, T_\alpha, \boldsymbol{\theta}_0)^T \mathbf{Q}_\alpha(\boldsymbol{\theta}_0) IF(y_1, \dots, y_n, T_\alpha, \boldsymbol{\theta}_0), \quad (28)$$

where T_α is the MDPD functional. Note that, this (second order) IF of the MDPDE based Wald-type test statistic directly depends on the IF of corresponding MDPD functional. Based on our earlier exploration in Section 2.4, the influence function of the Wald type test statistic in (28) will then be bounded in the contamination point y for any $\alpha > 0$ which implies the robustness of the proposed test statistic in (25) based on the MDPDE.

Again from the general theory of [11], one can observe the robustness of the level and power of the MDPDE based Wald-type tests for any $\alpha > 0$ through their bounded level influence function (LIF) and the power influence function (PIF). In particular, the LIF of any order is identically zero and the PIF for testing at the significance level τ has the form

$$PIF(y_1, \dots, y_n, W_n, \boldsymbol{\theta}_0) = C_r^* (\mathbf{d}^T \mathbf{Q}_\alpha(\boldsymbol{\theta}_0) \mathbf{d}) \mathbf{d}^T \mathbf{Q}_\alpha(\boldsymbol{\theta}_0) IF(y_1, \dots, y_n, T_\alpha, \boldsymbol{\theta}_0), \quad (29)$$

where $C_r^*(s) = e^{-\frac{s}{2}} \sum_{v=0}^{\infty} s^{v-1} 2^{-v} (2v-s) P(\chi_{r+2v}^2 > \chi_{r,\tau}^2) / v!$. Again the PIF is a linear function of the IF of the MDPDE and hence bounded for all $\alpha > 0$ indicating power robustness of the Wald-type test based on the statistic in (25).

3.3 An illustration: testing for the regression coefficient with one covariate

We now discuss, in detail, a particular testing problem regarding the regression coefficient. Let us consider the linear model with a single covariate as $y = x\beta + \epsilon$, with $\epsilon \sim SN(0, \sigma, \gamma)$. Here, we consider the sample size to be 100 and we generate x from $N(1, 1)$. Very often we want to test the significance of the regression coefficients, i.e. in our context we want to test $H_0 : \beta = 0$. Let us consider a slightly more general problem of testing

$$H_0 : \beta = \beta_0 \text{ against } H_1 : \beta \neq \beta_0, \quad (30)$$

for a pre-fixed real β_0 . The choice $\beta_0 = 0$ gives the test of significance of the covariate. Here σ and γ are unknown nuisance parameters.

In the notation of Section 3.1, we have $\Theta_0 = \{\boldsymbol{\theta} = (\beta_0, \sigma, \gamma)^T : \beta_0 \in \mathbb{R}, \sigma \in \mathbb{R}^+, \gamma \in \mathbb{R}\}$, $r = 1$, $\mathbf{m}(\boldsymbol{\theta}) = \beta - \beta_0$ and $M(\boldsymbol{\theta}) = (1, 0, 0)^T$.

Denoting the MDPDE as $\widehat{\boldsymbol{\theta}}_{\alpha,n} = (\widehat{\beta}_{\alpha,n}, \widehat{\sigma}_{\alpha,n}, \widehat{\gamma}_{\alpha,n})$, our MDPDE based Wald-type test statistics (25) has a simplified form for testing (30) which is given by

$$W_n(\widehat{\boldsymbol{\theta}}_{\alpha,n}) = \frac{n(\widehat{\beta}_{\alpha,n} - \beta_0)^2}{\Sigma_{\alpha,n}^{(11)}(\widehat{\boldsymbol{\theta}}_{\alpha,n})}, \quad (31)$$

where $\Sigma_{\alpha,n}^{(11)}(\boldsymbol{\theta})$ is the (1,1)-th element of $\boldsymbol{\Sigma}_{\alpha,n}(\boldsymbol{\theta})$. Then, under the null hypothesis in (30), $W_n(\widehat{\boldsymbol{\theta}}_{\alpha,n})$ asymptotically follows χ_1^2 distribution and the test can be performed by comparing $W_n(\widehat{\boldsymbol{\theta}}_{\alpha,n})$ with the corresponding critical values. Further, the approximate expression of power function at the contiguous hypothesis of the form $H_{1,n}: \beta = \beta_n$, where $\beta_n = \beta_0 + n^{-1/2}d$, with $d \in \mathbb{R} \setminus \{0\}$, is given by

$$\phi_\alpha(\boldsymbol{\theta}_n) = 1 - G_{\chi_1^2(\delta)}(\chi_{1,\tau}^2), \quad \text{with } \delta = \frac{d^2}{\Sigma_\alpha^{(11)}(\boldsymbol{\theta}_0)}, \quad \boldsymbol{\theta}_0 \in \Theta_0.$$

We have numerically calculated this asymptotic contiguous power for testing $\beta = 3$ at 5% level by using the MDPDE based Wald-type test with different values of α , which is graphically presented in Figure 2 for $\boldsymbol{\theta}_0 = (3, 1, 2)^T$. It is clear that, just like the ARE of the MDPDE, the contiguous power of the MDPDE based test also decreases as α increases but this loss is not quite significant at small $\alpha > 0$.




Figure 2: Asymptotic Contiguous Power of the MDPDE based Wald-type test for testing regression coefficient ($\beta_0 = 3$) at 5% level of significance with $\theta_0 = (3, 1, 2)^T$ and different values of α .

The robustness of the Wald-type test based on the statistic (31) for testing (30) can be studied through the second order influence function of the test statistic and the PIF. From the general formulas presented in Section 3.1, we can easily calculate these measures in the present case of testing (30) as given by

$$IF_{2,i_0}(y_{i_0}, W_n, \theta_0) = 2IF_{i_0}(y_{i_0}, T_\alpha^{(\beta)}, \theta_0)^2 / \Sigma_\alpha^{(11)}(\theta_0). \quad (32)$$

$$IF_2(\mathbf{y}, W_n, \theta_0) = 2IF(\mathbf{y}, T_\alpha^{(\beta)}, \theta_0)^2 / \Sigma_\alpha^{(11)}(\theta_0). \quad (33)$$

$$PIF(\mathbf{y}, W_n, \theta_0) = C_1^* \left(d^2 / \Sigma_\alpha^{(11)}(\theta_0) \right) IF(\mathbf{y}, T_\alpha^{(\beta)}, \theta_0) d / \Sigma_\alpha^{(11)}(\theta_0), \quad (34)$$

where $T_\alpha^{(\beta)}$ is the MDPDE functional corresponding to β and $\mathbf{y} = y\mathbf{1}$. For illustration, we have considered the model $y = x\beta + \epsilon$ with $\epsilon \sim SN(0, \sigma, \gamma)$ and presented the plots of these IF_2 and PIF for 6 different values of α at $\theta_0 = (3, 1, 2)^T$ and $d = 0.01$ in Figures 3 and 4. It is clear from these figures that the MDPDE based Wald-type test statistic (31) has bounded second order IF and PIF for all $\alpha > 0$.

4 Simulation studies

In this section, we present some interesting findings from extraneous simulation studies in order to examine the claimed robustness of the skewed regression model as well as the robust Wald-type test. First, we calculate the empirical bias and MSE (mean squared error) of the estimates of the model parameters based on simulated datasets. Next, we conduct robust Wald-type tests related to regression




Figure 3: Second order influence functions ($IF2$) of the robust Wald-type test statistic in $i_0 = 10$ th and all directions respectively with $\theta = (3, 2, 2)$ for different α .




Figure 4: Power influence function (PIF) of the robust Wald-type test statistic with $\theta = (3, 2, 2)$ for different α .

coefficients and calculate the power and level of the test. We use the R function `optim` to find the MDPDE of the model parameters.

4.1 Performance of the MDPDE

We consider the linear regression model

$$Y_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \epsilon_i \tag{35}$$

where $\epsilon_i \sim SN(0, 1, 2)$. We have taken $(\beta_0, \beta_1, \beta_2) = (1, 2, 3)$. Besides, we simulate x_1 from $N(1, 1)$ and x_2 from $N(-1, 1)$. Under these specifications, we simulate a random sample of size $n = 100$ observations. Based on 500 replications, we have tabulated the empirical biases and MSEs for 6 different α in Table 2. We have used the R package `sn` to simulate random samples from the skew normal distribution.

Next, we consider 5% contamination in the data. In this case, we simulate 5% of the errors from

$SN(-10, 1, 2)$ and rest from $SN(0, 1, 2)$. Again, based on 500 replications, we calculate the bias and MSE which are tabulated in Table 2. The process is then repeated for sample size $n = 200$ and the bias and MSEs are summarized in the same table.

Table 2: Empirical bias and MSE of the estimates of regression parameters under pure data as well as 5% and 10% contaminated data of size n [ϵ denotes the contamination proportion]

ϵ	Parametric MDPDE at different α						Parametric MDPDE at different α						
	0(MLE)	0.1	0.3	0.5	0.7	1	0(MLE)	0.1	0.3	0.5	0.7	1	
	Bias						MSE						
$n = 100$													
0%	β_0	0.0554	0.0604	0.0687	0.0872	0.0904	0.0954	0.0668	0.0807	0.0843	0.0879	0.0893	0.0902
	β_1	0.0001	-0.0003	-0.0004	-0.0004	-0.0019	-0.0025	0.0047	0.0052	0.0055	0.0056	0.0072	0.0085
	β_2	0.0037	0.0038	-0.0054	-0.0055	-0.0055	-0.0061	0.0043	0.0047	0.0055	0.0061	0.0065	0.0074
	σ	-0.0222	-0.0225	-0.0243	-0.0467	-0.0551	-0.0657	0.0157	0.0162	0.0173	0.0178	0.0187	0.0225
	γ	0.0098	0.0545	0.1370	0.2802	0.3501	0.5870	0.1667	0.3844	0.6104	0.8000	1.0094	1.2017
	5%	β_0	-0.1389	-0.1189	0.1090	0.1003	0.0973	0.0981	0.1564	0.1429	0.1368	0.1249	0.1001
β_1		0.0075	0.0064	0.0051	-0.0039	-0.0030	-0.0034	0.0117	0.0105	0.0099	0.0093	0.0086	0.0090
β_2		-0.0089	-0.0075	-0.0064	-0.0059	-0.0057	-0.0063	0.0108	0.0100	0.0096	0.0088	0.0076	0.0080
σ		-0.0782	-0.0725	-0.0658	-0.0598	-0.0575	-0.0659	0.0400	0.0348	0.0313	0.0287	0.0232	0.0243
γ		0.9028	0.8635	0.7318	0.6042	0.4828	0.5385	1.7183	1.6048	1.4652	1.3378	1.1494	1.2246
10%		β_0	-0.2013	-0.1648	-0.1313	0.1261	0.0980	0.0996	0.1824	0.1638	0.1448	0.1370	0.1124
	β_1	0.0086	0.0077	0.0066	-0.0048	-0.0037	-0.0042	0.0219	0.0155	0.0129	0.0108	0.0094	0.0101
	β_2	-0.0103	-0.0095	-0.0086	-0.0071	-0.0063	-0.0072	0.0166	0.0148	0.0115	0.0100	0.0088	0.0095
	σ	-0.0938	-0.0873	-0.0795	-0.0721	-0.0653	-0.0718	0.0528	0.0447	0.0385	0.0305	0.0281	0.0296
	γ	0.9383	0.9031	0.7982	0.6418	0.5435	0.5828	1.8214	1.6925	1.5597	1.4043	1.2295	1.2792
	$n = 200$												
0%	β_0	0.0414	0.0479	0.0544	0.0658	0.0770	0.0824	0.0548	0.0677	0.0753	0.0810	0.0855	0.0882
	β_1	0.0001	-0.0002	-0.0003	-0.0004	-0.0016	-0.0022	0.0036	0.0044	0.0050	0.0054	0.0065	0.0077
	β_2	0.0029	0.0033	-0.0040	-0.0046	-0.0050	-0.0057	0.0038	0.0042	0.0049	0.0056	0.0062	0.0069
	σ	-0.0199	-0.0205	-0.0216	-0.0377	-0.0451	-0.0547	0.0101	0.0108	0.0114	0.0120	0.0124	0.0130
	γ	0.0090	0.0425	0.1060	0.2285	0.3011	0.4861	0.1067	0.2484	0.4404	0.6199	0.7987	0.9171
	5%	β_0	-0.1162	-0.1028	-0.0931	0.0813	0.0732	0.0843	0.1503	0.1392	0.1234	0.1093	0.0902
β_1		0.0058	0.0052	0.0043	-0.0031	-0.0018	-0.0024	0.0101	0.0096	0.0084	0.0073	0.0066	0.0071
β_2		-0.0074	-0.0065	-0.0058	-0.0047	-0.0039	-0.0046	0.0095	0.0087	0.0080	0.0071	0.0053	0.0067
σ		-0.0681	-0.0591	-0.0508	-0.0417	-0.0341	-0.0425	0.0261	0.0193	0.0154	0.0113	0.0098	0.0125
γ		0.7821	0.7032	0.6090	0.5199	0.4181	0.4612	1.2762	1.1549	0.9962	0.8574	0.7877	0.8482
10%		β_0	-0.1423	-0.1283	-0.1002	0.0922	0.0842	0.0913	0.1663	0.1519	0.1392	0.1204	0.1022
	β_1	0.0069	-0.0063	-0.0052	-0.0039	-0.0025	-0.0031	0.0184	0.0143	0.0115	0.0096	0.0082	0.0094
	β_2	-0.0094	-0.0085	-0.0073	-0.0059	-0.0052	-0.0064	0.0119	0.0103	0.0094	0.0083	0.0072	0.0088
	σ	-0.0861	-0.0754	-0.0683	-0.0554	-0.0431	-0.0529	0.0374	0.0295	0.0214	0.0171	0.0159	0.0187
	γ	0.8412	0.8062	0.6950	0.5736	0.4610	0.5013	1.3166	1.2239	1.0952	0.9484	0.8294	0.8618

From Table 2 we observe that, for pure data, the empirical biases and MSEs have increasing trends with the increment of the tuning parameter α . The biases and MSEs also decrease with the sample size n . Under contamination these quantities are high for small values of α ; however, they steadily decline with increasing α (in absolute value for the bias) and eventually stabilize at values close to those of the pure data case (although there appears to be a slight upturn somewhere between $\alpha = 0.7$ and $\alpha = 1$).

4.2 Performance of the robust Wald-type test

Here we take the same set up of the regression model as in Section 4.1. Here we want to test $H_0: \beta_1 = 2$ against $H_{1,n}: \beta_1 = \beta_{1,n}$, where $\beta_{1,n} = 2 + n^{-1/2}d$, for which the Wald-type test statistic W_n has a form similar to that in (31). We first simulate random samples of sizes $n = 100, 200$ and perform the MDPDE based Wald-type test for different α , including the classical Wald test at $\alpha = 0$. Based on 500 replications, we then compute the empirical levels of the tests, measured as the proportion of test statistics exceeding the chi-square critical value among the 500 replications. Subsequently, to compute the empirical power of the tests, we repeat the above exercise but now we calculate the proportion of times $H_{1,n}: \beta_1 = \beta_{1,n}$ is rejected, where $\beta_{1,n} = 2 + n^{-\frac{1}{2}}1.5$. For calculating the empirical power, we have considered the same linear regression model as used in case of level calculation except that we have taken the value of β_1 as $\beta_{1,n} = 2 + n^{-\frac{1}{2}}1.5$. Finally, to illustrate the claimed robustness, we recalculate the level and power of the Wald-type tests after contamination 100 $\epsilon\%$ of each sample in the previous simulation exercises with $\epsilon = 0.05, 0.1$. The contaminated observations generated using the error distribution as $SN(-5, 1, 2)$ and $SN(-3, 1, 2)$ respectively, for the level and power calculations. In Table 3, we report all the resulting empirical levels and powers obtained from different simulation scenarios.

Table 3: Empirical levels and powers of robust Wald-type test statistic under pure data as well as 5% and 10% contaminated data [ϵ denotes the contamination proportion]

Sample size	ϵ	Parametric MDPDE at different α					
		0(MLE)	0.1	0.3	0.5	0.7	1
		Level					
$n = 100$	0%	0.072	0.076	0.084	0.095	0.106	0.119
	5%	0.429	0.325	0.219	0.163	0.112	0.091
	10%	0.467	0.359	0.261	0.185	0.134	0.098
$n = 200$	0%	0.054	0.057	0.062	0.071	0.08	0.094
	5%	0.454	0.331	0.193	0.129	0.094	0.076
	10%	0.482	0.368	0.247	0.161	0.11	0.089
		Power					
$n = 100$	0%	1	1	1	1	0.996	0.963
	5%	0.403	0.498	0.615	0.707	0.829	0.952
	10%	0.395	0.476	0.575	0.696	0.756	0.934
$n = 200$	0%	1	1	1	1	1	1
	5%	0.416	0.597	0.713	0.886	1	1
	10%	0.409	0.567	0.694	0.842	0.92	1

From Table 3, we observe that, for pure data, the empirical levels increase with the increment in α . It can also be observed that, by the time the sample size rises to 200, the observed levels are close to the nominal level of 0.05 for small values of α ; however it gets inflated for larger α . For

contaminated data the levels are high for large α , but steadily drop with increasing α . In case of power the contamination leads to a significant drop in power for smaller α , but at larger values of α the power is much better protected.

5 Real data examples

In this section, we apply our MDPDE based skew normal regression methods to analyze two real life datasets. For each dataset, we calculate the estimates along with the standard errors (SEs) of the regression coefficients (β) as well as the scale parameter (σ) and shape parameter (γ). Apart from these, we also conduct robust Wald-type tests for testing the significance of the covariates (involving the β parameters) and the symmetry of the model (involving the γ parameter).

5.1 Australian Institute of Sports (AIS) data

These data consist of the health measurements of 706 Australian athletes from 12 different sports which were collected at the Australian Institute of Sports (AIS) in 1990. The data were first analyzed by Telford and Cunningham (1991) [37]. A subset of these data (consisting of 207 observations) are now available in the R package `sn` which we use in the following analysis.

There are several biomedical variables related to the health conditions of the athletes included in this dataset. For our analysis, we model the plasma ferritin concentration in blood (denoted as “Fe”) with the body mass index (BMI) and the lean body mass (LBM) as the covariates using the following regression equation:

$$\text{Fe}_i = \beta_0 + \beta_1 \text{BMI}_i + \beta_2 \text{LBM}_i + \epsilon_i, \quad i = 1, \dots, 207,$$

where $\epsilon_i \sim SN(0, \sigma, \gamma)$ independently for each i . We summarize the results in Table 4, where we also provide the p-values of robust Wald-type tests for testing individual parameters being zero.

To explain how the analysis based on the procedures corresponding to larger values of the tuning parameter α provide more stable inference, we offer the following description. Together with the results of the full data (containing 207 observations) analysis, we also provide the analysis based on the outlier deleted data (containing 202 observations) where five cases are deleted based on their outlyingness in terms of the adjusted box plot of the BMI covariate. The results of the full data and outlier deleted data analysis are presented in Tables 4 and 5 respectively.

Contrasting the results of these two tables provides instructive insights. For the full data analysis of the BMI variable, the p-values indicate that for smaller α the covariate BMI is highly significant, although the significance disappears (or at best becomes an extremely tentative one) for larger α (Table 4). Thus different values of α may lead to very different inference on the BMI variable in this case. However, for the outlier deleted data case, the p-values for the same test are remarkably

Table 4: Estimates along with standard error (SE) for AIS data at different α

Parameter	MDPDE at different α							
	0(MLE)	0.05	0.1	0.3	0.5	0.7	0.8	1
Intercept	-46.3478	-46.3477	-46.3477	-46.0384	-45.8145	-45.8965	-46.2416	-46.3238
SE	(13.2494)	(13.1964)	(13.1176)	(13.6866)	(14.2514)	(14.5397)	(14.1700)	(14.6015)
p-value	(0.0005)	(0.0005)	(0.0004)	(0.0008)	(0.0013)	(0.0016)	(0.0011)	(0.0015)
BMI	2.3485	2.3484	2.3482	2.2346	2.1372	1.9937	1.8058	1.6145
SE	(0.8049)	(0.8002)	(0.7969)	(0.8317)	(0.8667)	(0.8854)	(0.8637)	(0.8910)
p-value	(0.0035)	(0.0034)	(0.0032)	(0.0072)	(0.0137)	(0.0243)	(0.0365)	(0.0603)
LBM	0.2280	0.2286	0.2290	0.2745	0.3139	0.3735	0.4536	0.5331
SE	(0.1764)	(0.1753)	(0.1746)	(0.1822)	(0.1899)	(0.1940)	(0.1892)	(0.1952)
p-value	(0.1960)	(0.1918)	(0.1897)	(0.1321)	(0.0983)	(0.0542)	(0.0165)	(0.0093)
σ	70.8381	70.1732	69.6698	66.4086	63.7539	62.4061	62.0975	61.2081
SE	(3.8689)	(3.8601)	(3.8562)	(4.0110)	(4.2578)	(4.5166)	(4.5966)	(4.8078)
γ	10.9054	10.9052	10.9048	9.7672	9.2024	9.5091	10.5502	10.7931
SE	(3.5252)	(3.5419)	(3.5635)	(3.2956)	(3.3856)	(3.9831)	(4.8686)	(5.6060)
p-value	(0.0020)	(0.0022)	(0.0022)	(0.0030)	(0.0066)	(0.0199)	(0.0382)	(0.0642)

Table 5: Estimates along with standard error (SE) for AIS data for different α after excluding the outlier

Parameter	MDPDE at different α							
	0(MLE)	0.05	0.1	0.3	0.5	0.7	0.8	1
Intercept	-45.9629	-45.9619	-45.9584	-45.5282	-46.1271	-46.5730	-46.6960	-46.8758
SE	(12.8445)	(12.9465)	(13.2931)	(13.6445)	(13.9400)	(14.0808)	(14.2659)	(14.4313)
p-value	(3.2E-04)	(3.9E-04)	(4.1E-04)	(6.94E-04)	(9.65E-04)	(1.03E-03)	(1.19E-03)	(1.46E-03)
BMI	1.1241	1.1238	1.0899	1.2867	1.3529	1.5783	1.5875	1.6424
SE	(0.9044)	(0.9086)	(0.9157)	(0.9375)	(0.9523)	(0.9683)	(0.9752)	(0.9928)
p-value	(5.01E-02)	(5.05E-02)	(5.10E-02)	(5.30E-02)	(5.56E-02)	(5.86E-02)	(5.93E-02)	(6.14E-02)
LBM	0.2356	0.2362	0.2486	0.2874	0.3367	0.3986	0.4523	0.5251
SE	(0.1731)	(0.1749)	(0.1773)	(0.1807)	(0.1886)	(0.1929)	(0.1952)	(0.2008)
p-value	(1.98E-03)	(2.43E-03)	(2.70E-03)	(3.65E-03)	(4.76E-03)	(6.52E-03)	(7.05E-03)	(8.91E-03)
σ	61.5903	61.0054	60.3713	60.9935	61.0749	61.2209	61.3342	61.5622
SE	(3.7811)	(3.7964)	(3.7667)	(3.9118)	(4.1457)	(4.4299)	(4.5869)	(4.6985)
γ	11.1081	11.1079	11.0874	10.0807	9.5198	9.6695	10.6575	10.9846
SE	(3.4891)	(3.4994)	(3.5200)	(3.6118)	(3.8193)	(4.0805)	(4.5378)	(4.8609)
p-value	(0.0496)	(0.0511)	(0.0517)	(0.0526)	(0.0540)	(0.0576)	(0.0598)	(0.0633)

uniform over α , all indicating weak significance to lack of significance (Table 5). For the case $\alpha = 0$, the p-values of this test are widely disparate between the full data and outlier deleted cases (0.0035 versus 0.0501). On the other hand, for $\alpha = 1$, the p-value is practically unaffected by the presence or absence of outliers (0.0603 versus 0.0614). Clearly the significance of the BMI variable in the $\alpha = 0$ case is driven by the five outlying observations, which is also the case for other low values of α ; but the effect of the outliers is minimal for large α , demonstrating the stability of the procedures for such

α . On the whole, we repose our faith on the analysis based on large values of α in this case, as they are stable (minimally affected by the presence or absence of outliers) and closely match the consistent results of the analysis for the outlier deleted case, where the conclusions are consistent over different values of α (small and large).

The results of the significance test of the γ parameter show exactly similar trends. A strong significance at low α is pushed to a lack of significance or weak significance through the removal of the outliers. At larger α the p-value remains stable, demonstrating a lack of significance/weak significance with or without the outliers.

For the LBM variable, on the other hand, insignificant p-values are pushed to significance through the removal of outliers for lower values of α , whereas for large values of α the p-values remain stable, exhibiting significance with or without the outliers. All these results, taken together, indicate that the analysis based on larger values of α provide protection against both kinds of errors – false declaration of significance when there is none, and failure to detect true significance – and should therefore be the trusted version of our analysis.

The p-values of the three tests (for the significance of the BMI, the LBM and the test of symmetry) are graphically presented in Figure 5 to give a better visual representation.




Figure 5: p-values for the tests of significance of covariates and test of symmetry in case of AIS data. [Solid black line: Full data, Solid blue line: Outlier deleted data, Dashed red line: level of significance.]

5.2 Crime data

In this example we will once again demonstrate how the presence or absence of a handful of outlying observations can drive the decision for small α , but for large α the outliers have little or no effect. We consider the Crime data (consisting of 51 observations from each of 51 states of the United States of America, measured in 1993) presented in Agresti and Finlay (1997) [4]. The dataset is loaded in R studio from the link: <https://stats.idre.ucla.edu/stat/data/crime.dta>.

Among the available variables within this dataset, we use the number of murders per 1,000,000 people in each state (denoted as “murder”) as our response variable and the percentage of people living under the poverty line (denoted as “poverty”) and percentage of populations who are single parents (denoted as “single”) as our explanatory variables. Thus, we fit the model

$$\text{murder}_i = \beta_0 + \beta_1 \text{poverty}_i + \beta_2 \text{single}_i + \epsilon_i, \quad i = 1, \dots, 51,$$

where the subscript i represents the observation from i th state and ϵ_i is the error component having SN distribution as in (3).

The results obtained from the full data are presented in Table 6. The last (51st) observation is a clear outlier by any standards as observed from the residuals of the regression for any α . The results of the outlier deleted regression (based on the remaining 50 observations) are presented in Table 7. There are many similarities between the comparative analysis of these two tables and the comparative analysis of Tables 4 and 5. An additional observation is that stable results are obtained in Table 6 even for very small (positive) values of α . In case of the “poverty” variable, for example, the insignificance at $\alpha = 0$ is turned into strong significance by the removal of the single outlier; however, for full data


Table 6: Estimates along with standard error (SE) for Crime data for different α

Parameter	MDPDE at different α							
	0(MLE)	0.05	0.1	0.3	0.5	0.7	0.8	1
Intercept	-23.5159	-17.0934	-14.8622	-14.7993	-14.8492	-14.5321	-17.4511	-19.0914
SE	(2.4342)	(1.9853)	(1.6189)	(1.6701)	(1.7170)	(1.6926)	(1.4126)	(0.9116)
p-value	(4.43E-22)	(8.45E-21)	(4.28E-20)	(7.91E-19)	(5.23E-18)	(9.02E-18)	(1.24E-21)	(1.10E-23)
Poverty	0.2805	0.3495	0.3708	0.3730	0.3725	0.3681	0.4604	0.4743
SE	(0.1148)	(0.1083)	(0.1052)	(0.0779)	(0.0804)	(0.0798)	(0.067)	(0.0436)
p-value	(0.1459)	(0.0007)	(8.20E-04)	(6.20E-04)	(1.68E-04)	(3.64E-05)	(3.73E-05)	(3.93E-05)
Single	2.0139	1.4853	1.2701	1.2565	1.2607	1.2427	1.4403	1.5882
SE	(0.2482)	(0.1973)	(0.1625)	(0.1683)	(0.1738)	(0.1724)	(0.1447)	(0.0943)
p-value	(4.85E-04)	(3.95E-04)	(3.73E-04)	(5.48E-04)	(8.35E-04)	(4.08E-04)	(4.72E-04)	(5.60E-04)
σ	8.6706	5.4882	3.7718	3.9108	3.9956	4.0052	3.7316	3.6592
SE	(0.9717)	(0.6713)	(0.5321)	(0.5739)	(0.6249)	(0.6558)	(0.5924)	(0.5411)
γ	8.6820	5.7169	3.3957	3.7598	4.1798	4.9330	6.7388	8.2260
SE	(5.1359)	(3.0184)	(1.6423)	(1.9707)	(2.4778)	(3.3800)	(4.2761)	(5.5665)
p-value	(0.0909)	(0.0413)	(0.0387)	(0.0364)	(0.0316)	(0.0344)	(0.0382)	(0.0406)


even a small positive value of α such as 0.05 exhibits a large drop in p-value compared to $\alpha = 0$, and the removal of the outlier has no effect on the decision in this case (or at larger values of α). Please see also the plots of p-values in Figure 6. This phenomenon is consistent with the observations of other authors (e.g., Basak et al. (2021) [8]) where it has been observed that if the outliers are widely discrepant with the main data cloud, even small (positive) values of α can discount them.

Table 7: Estimates along with standard error (SE) for Crime data for different α after excluding the outlier

Parameter	MDPDE at different α							
	0(MLE)	0.05	0.1	0.3	0.5	0.7	0.8	1
Intercept	-14.9633	-14.9713	-14.9776	-15.0102	-15.0459	-16.4709	-17.5197	-18.7928
SE	(1.2255)	(1.2392)	(1.2483)	(1.3284)	(1.4002)	(1.5119)	(1.6736)	(1.6919)
p-value	(4.15E-25)	(7.23E-25)	(9.94E-25)	(2.31E-24)	(5.42E-24)	(7.11E-24)	(8.54E-24)	(1.03E-23)
Poverty	0.3680	0.3701	0.3707	0.3734	0.3733	0.3971	0.4061	0.4389
SE	(0.0343)	(0.0351)	(0.0355)	(0.0370)	(0.0395)	(0.0409)	(0.0427)	(0.0458)
p-value	(1.19E-05)	(1.39E-05)	(1.47E-05)	(2.19E-05)	(2.39E-05)	(2.86E-05)	(3.23E-05)	(3.71E-05)
Single	1.2859	1.2815	1.2805	1.2760	1.2790	1.2808	1.3416	1.4582
SE	(0.1061)	(0.1079)	(0.1085)	(0.1166)	(0.1341)	(0.1464)	(0.1679)	(0.1940)
p-value	(1.81E-05)	(2.35E-05)	(2.68E-05)	(4.26E-05)	(4.63E-05)	(5.72E-05)	(6.49E-05)	(7.84E-05)
σ	3.6931	3.7411	3.7710	3.8875	3.9528	3.9590	3.7868	3.6008
SE	(0.5163)	(0.5208)	(0.5272)	(0.5369)	(0.5792)	(0.6298)	(0.5768)	(0.6497)
γ	3.2595	3.3314	3.4009	3.6961	4.0558	5.0839	5.9720	7.8440
SE	(1.5748)	(1.6074)	(1.6611)	(1.9551)	(2.4217)	(3.5364)	(3.7488)	(4.3472)
p-value	(0.0292)	(0.0307)	(0.0312)	(0.0325)	(0.0338)	(0.0353)	(0.0376)	(0.0397)



(a) p-value for Poverty



(b) p-value for γ

Figure 6: p-values for the tests of significance of “Poverty” and test of symmetry in case of Crime data. [Solid black line: Full data, Solid blue line: Outlier deleted data, Horizontal dashed red line: level of significance.]

6 Choice of the robustness tuning parameter

The tuning parameter α used in the proposed MDPDE based skewed regression model controls the trade-off between the asymptotic efficiency and the robustness of the procedure. At $\alpha = 0$, the MDPDE coincides with the MLE, which is the most efficient estimator for pure data but may have degraded performance under model misspecification and data contamination. As α increases, so does the stability of the estimator, but at the cost of an efficiency loss under pure data. It is therefore very important to choose the appropriate tuning parameter depending on the situation to get optimal performance. However, the experimenter would not know apriori how much anomaly is present in the data in relation to the assumed model. So a fixed choice of α cannot provide optimal performance in all situations. Thus it will be very useful to have an automatic data-specific method of selecting the tuning parameter that will provide the most appropriate treatment for the data at hand.

Several authors have considered data-specific optimal choices of the tuning parameter α . Warwick and Jones (2005) [38] explored this problem under the IID set up, where they suggested the minimization of an empirical estimate of the asymptotic summed mean square error (AMSE) over α to choose the optimal tuning parameter. This approach was extended to the non-homogeneous setup by Ghosh and Basu (2015) [19]. While this approach generally provides a reasonable solution, it is dependent on the choice of an initial pilot estimator. Hong and Kim (2001) [26] had earlier proposed a method for the optimal tuning parameter selection, where they used the variance component only (rather than the entire mean squared error). As this method does not consider the bias component, it avoids the use of a pilot estimator; however this method suffers from the problem of an occasional non-robust solution. Recently, Basak et al. [8] have proposed an iterative algorithm for finding the optimal value of the tuning parameter in case of both IID and non-homogeneous data which has been called the ‘Iterated Warwick-Jones’ (IWJ) algorithm. This method appears to be able to remove the dependence on the pilot estimator which was inherent in the original Warwick-Jones approach.

In this paper, we will follow the IWJ method for finding the optimal value of α . For the sake of completeness, we state the algorithm briefly, as adapted to our case. Suppose that the observed dataset is obtained from a contamination model $g_i = (1 - \epsilon)f_i(\cdot, \boldsymbol{\theta}^*) + \epsilon\delta_{y_i}$ for each i , where g_i and $f_i(\cdot, \boldsymbol{\theta})$ are defined in Section 2, δ_{y_i} s are some contaminating densities and $\boldsymbol{\theta}^*$ is the target parameter value. Let $\hat{\boldsymbol{\theta}}_{\alpha,n}$ and $\boldsymbol{\theta}^g$ be the MDPDE and the corresponding best fitting parameters of $\boldsymbol{\theta}$ as defined in Section 2. Then the asymptotic summed MSE (AMSE) of the MDPDE $\hat{\boldsymbol{\theta}}_{\alpha,n}$ with respect to the target $\boldsymbol{\theta}^*$ is given by

$$AMSE\left(\hat{\boldsymbol{\theta}}_{\alpha,n}\right) = (\boldsymbol{\theta}^g - \boldsymbol{\theta}^*)^T (\boldsymbol{\theta}^g - \boldsymbol{\theta}^*) + \frac{1}{n} \text{trace} \left[\boldsymbol{\Psi}_{\alpha}^{-1}(\boldsymbol{\theta}^g) \boldsymbol{\Omega}_{\alpha}(\boldsymbol{\theta}^g) \boldsymbol{\Psi}_{\alpha}^{-1}(\boldsymbol{\theta}^g) \right]. \quad (36)$$

The expression on the right hand side of the above equation is a function of two unknown quantities, $\boldsymbol{\theta}^g$ and $\boldsymbol{\theta}^*$. We replace $\boldsymbol{\theta}^g$ by its consistent estimator $\hat{\boldsymbol{\theta}}_{\alpha,n}$. However we do not have an obvious

surrogate for θ^* and therefore we have to use an appropriate robust pilot estimator $\widehat{\theta}_P$ for it; the MDPDEs for $\alpha = 0.5$ and 1 are popular choices for this purpose. Thus, an empirical estimate of the summed AMSE at tuning parameter α is given by

$$\widehat{AMSE}(\widehat{\theta}_{\alpha,n}) = (\widehat{\theta}_{\alpha,n} - \widehat{\theta}_P)^T (\widehat{\theta}_{\alpha,n} - \widehat{\theta}_P) + \frac{1}{n} \text{trace} \left[\Psi_n^{-1}(\widehat{\theta}_{\alpha,n}) \Omega_n(\widehat{\theta}_{\alpha,n}) \Psi_n^{-1}(\widehat{\theta}_{\alpha,n}) \right]. \quad (37)$$

As the estimated summed $\widehat{AMSE}(\widehat{\theta}_{\alpha,n})$ is a function of α , we can minimize this measure over $\alpha \in [0, 1]$ to get the optimal tuning parameter for a given dataset. The algorithm proceeds as follows:

- (1) Start with a suitable robust pilot estimator $\widehat{\theta}_P$ of θ .
- (2) Calculate $\widehat{AMSE}(\widehat{\theta}_{\alpha,n})$ for a large number of equispaced $\alpha \in [0, 1]$.
- (3) Find the value of α that minimizes estimated $\widehat{AMSE}(\widehat{\theta}_{\alpha,n})$ in (37) over the grid of α values considered in the previous step. Denote this by α_1 and let the corresponding estimate of θ be denoted as $\widehat{\theta}_{1,\alpha_1}$.
- (4) Choose $\widehat{\theta}_{1,\alpha_1}$ as the updated pilot estimate of θ for the next step.
- (5) Repeat steps (3) and (4) until there is no further change in the estimate of θ (or, equivalently, the estimate of the tuning parameter α).

The main advantage of this method is that the final choice of optimal α does not depend on the original pilot estimator $\widehat{\theta}_P$. However, we still have to start the process at a suitable robust starting value, and here we use the MDPDE with $\alpha = 0.5$ as recommended by Ghosh and Basu (2015) [19].

Notice that the above derivation of the optimal tuning parameter α uses the minimization of the empirical summed AMSE. This may be viewed as providing an optimal solution to the problem of estimating the parameter vector. While we will follow the same approach for estimating the tuning parameter in the hypothesis testing problem, we will make suitable changes which may enhance the performance. In particular, since the tests of significance relate to individual parameters, we will use the estimated asymptotic mean square error for the variable in question rather than the summed AMSE over all the variables. This will also allow us to remove the interference of other variables which may have very large standard errors (SE) dominating the SE of the variable in question. This may be observed, for example, in case of the BMI and LBM variables in the AIS data example, where the very large value of the SE for the estimate of γ will dominate the summed mean square error and can perhaps lead to incorrect insights for the BMI and LBM variables.

For both the real data examples, the full set of parameters is $\theta = (\beta_0, \beta_1, \beta_2, \sigma, \gamma)$. Our primary interest is in the regression parameters β_1 and β_2 and the skewness parameter γ . When, we are testing

the significance of a particular covariate, say BMI for AIS data, our null hypothesis of interest is given by $H_0 : m(\boldsymbol{\theta}) = \boldsymbol{\theta}[2] = 0$. Now if we have to find the optimal tuning parameter for $m(\boldsymbol{\theta}) = \boldsymbol{\theta}[2]$, the estimated AMSE becomes $\widehat{AMSE} \left(m \left(\hat{\boldsymbol{\theta}}_{\alpha,n} \right) \right) = \left(\hat{\boldsymbol{\theta}}_{\alpha,n}[2] - \hat{\boldsymbol{\theta}}_P[2] \right)^2 + \boldsymbol{\Sigma}_{\alpha,n}^{(22)} \left(\hat{\boldsymbol{\theta}}_{\alpha,n} \right)$. Minimizing this over α , we get the optimal value of the tuning parameter α to be 0.99. Similarly, if we consider the covariate LBM in the AIS data example, the value of the corresponding optimal α becomes 0.93. A similar exercise with the two covariates in case of Crime data leads to common optimum α values of 0.82. The optimal estimates of α for performing the test of hypotheses about the skewness parameter γ are 0.66 and 0.06, respectively, for the AIS data and Crime data examples. Note that the estimated optimal parameter for estimating the skewness parameter in the second real data example is only 0.06, a rather low value; however as mentioned previously, this dataset contained a huge outlier and only a small positive value of α is sufficient to lead to a reversal of the results that are obtained for the MLE ($\alpha = 0$) for the full data case.

7 Concluding remarks

In this paper we have used the minimum density power divergence estimation method to fit linear regression models under skew errors. We have demonstrated, through influence function analysis, simulation studies and real data examples how the MDPDE can provide good fits for the regression model with skew normal errors. In particular this exploration reveals how procedures with large α can withstand the effect of model specific outliers unlike the likelihood based solutions (or other procedures with small values of α) where the decision may be driven by a handful of outliers even in large data sets. So, in future, this method may be extended for the linear regression set up under any other positively skewed distributions, such as extreme valued, gamma, Gompertz etc. Further, the estimation procedure based on MDPDE may be applied to the Cox proportional hazards regression model (specially the frailty model) based on right censored survival data with the frailty distribution being skew normal (SN).

References

- [1] Aeberhard, W. H., Cantoni, E. and Heritier, S. (2014). Robust inference in the Negative Binomial regression model with an application to Falls data. *Biometrics*; 70:920-931.
- [2] Alhamide, A. A., Ibrahim, K. and Alodat, M. T. (2015). Multiple linear regression estimators with skew normal errors. *AIP Conference Proceedings*, 1678, 060013(2015). [doi: 10.1063/1.4931340].

- [3] Agostinelli, C., Bianco, A. M. and Boente, G. (2020). Robust estimation in single-index models when the errors have a unimodal density with unknown nuisance parameter. *Annals of the Institute of Statistical Mathematics*; 72:855-893.
- [4] Agresti, A. and Finlay, B. (1997). *Statistical Methods for the Social Sciences(3rd edition)*. Pearson Education Limited.
- [5] Azzalini, A.(1985). A Class of Distributions which Includes the Normal Ones. *Scandinavian Journal of Statistics*; 12:171-178.
- [6] Azzalini, A.(1986). Further Results on a Class of Distributions Which Includes the Normal Ones. *Statistica*; 46:199-208.
- [7] Azzalini, A.(2005). The Skew-normal Distribution and Related Multivariate Families. *Scandinavian Journal of Statistics*; 32:159-188.
- [8] Basak, S., Basu, A. and Jones, M. C. (2020). On the ‘optimal’ density power divergence tuning parameter. *Journal of Applied Statistics*[doi:10.1080/02664763.2020.1736524].
- [9] Basu, A., Harris, I. R., Hjort, N. L., and Jones, M. C. (1998). Robust and efficient estimation by minimising a density power divergence. *Biometrika*; 85:549-559.
- [10] Basu, A., Shioya, H. and Park, C. (2011). *Statistical Inference: The Minimum Distance Approach*. Chapman & Hall/CRC. Boca Raton, Florida.
- [11] Basu, A., Ghosh, A., Martin, N. and Pardo, L. (2018). Robust Wald-type tests for non-homogeneous observations based on the minimum density power divergence estimator. *Metrika*; 81:493–522.
- [12] Beath, K. J. (2018). A mixture-based approach to robust analysis of generalised linear models. *Journal of Applied Statistics*; 45(12):2256-2268.
- [13] Boos, D. D. (1987). Detecting skewed error from regression residuals. *Technometrics*; 29(1):83-90.
- [14] Cao, X., Wang, G. and Wu, L. (2020). Skew-normal median regression model with applications. *Journal of Physics Conference Series*; 1616 (2020) 012079 [doi:10.1088/1742-6596/1616/1/012079].
- [15] Dogru, F. Z. and Arslan, O. (2017). Robust mixture regression based on the skew t distribution. *Colombian Journal of Statistics*; 40(1):45-64.
- [16] Deb, S. (2017). A Novel Robust R-Squared Measure and Its Applications in Linear Regression. In: Phon-Amnuaisuk, S., Au, T.W., Omar, S. (eds) *Computational Intelligence in Information Systems*.

- CIIS 2016. *Advances in Intelligent Systems and Computing*; 532. Springer, Cham. [doi:10.1007/978-3-319-48517-1_12].
- [17] Guedes, T. A., Rossi, R. M., Martins, A. B. T., Janeiro, V. and Carneiro, J. W. P. (2014). Applying regression models with skew-normal errors to the height of bedding plants of *Stevia rebaudiana* (Bert) Bertoni. *Acta Scientiarum: Technology, Maringa*; 36(3):463-468.
- [18] Ghosh, A. and Basu, A. (2013). Robust estimation for independent non-homogeneous observations using density power divergence with applications to linear regression. *Electronic Journal of Statistics*; 7(2013):2420-2456.
- [19] Ghosh, A. and Basu, A. (2015). Robust estimation for non-homogeneous data and the selection of the optimal tuning parameter: the density power divergence approach. *Journal of Applied Statistics*; 42(9):2056-2072.
- [20] Ghosh, A. and Basu, A. (2016). Robust estimation in generalized linear models: the density power divergence approach. *Test*, 25(2):269–290.
- [21] Ghosh, A., Mandal, A., Martin, N. and Pardo, L. (2016). Influence analysis of robust Wald-type tests. *Journal of Multivariate Analysis*; 147:102–126.
- [22] Ghosh, A. and Basu, A. (2018). Robust bounded influence tests for independent but non-homogeneous observations. *Statistica Sinica*; 28(3):1133-1155.
- [23] Ghosh, A. and Basu, A. (2019). Robust and efficient estimation in the parametric Cox regression model under random censoring. *Statistics in Medicine*; 38(27):1-17.
- [24] Ghosh, A., Basu, A. and Pardo, L. (2019). Robust Wald-type tests under random censoring with applications to clinical trial analyses. *arXiv:1708.09695v2 [stat.ME]*.
- [25] Hampel, F. R., Ronchetti, E., Rousseeuw, P. J. and Stahel W.(1986). *Robust Statistics: The Approach Based on Influence Functions*. New York, USA: John Wiley & Sons.
- [26] Hong, C. and Kim, Y. (2001). Automatic selection of the tuning parameter in the minimum density power divergence estimation. *Journal of the Korean Statistical Society*; 30:453–465.
- [27] Hui-qiong, L. I. and Liu-cang, W. U. (2014). Joint modelling of location and scale parameters of the skew-normal distribution. *Applied Mathematics-A Journal of Chinese Universities*; 29(3):265-272.
- [28] Kheramandi, A. and Rasekh, A. (2015). Estimation in skew-normal linear mixed measurement error models. *Journal of Multivariate Analysis*; 136:1-11.

- [29] Liu, M. and Lin, T-I. (2014). A skew-normal mixture regression model. *Educational and Psychological Measurement*; 74(1):139-162.
- [30] Lyu, W. and Feng, Z. (2018). Estimation and application of skew-normal data for generalized linear regression. *Advances in Intelligent Systems Research*; 151:208-211.
- [31] Malehi, A. S., Pourmotahari, F. and Angali, K. A. (2015). Statistical models for the analysis of skewed healthcare cost data: a simulation study (2015). *Health Economics Review*; 5(11):1-16.
- [32] Nandy, A., Basu, A. and Ghosh, A. (2021). Robust inference for skewed data in health sciences. *Journal of Applied Statistics*; 1-31 [doi.org/10.1080/02664763.2021.1891527].
- [33] Petrinco, M., Barbati, G., Meylan, D., Pagano, E., Gregori, D., Merletti, F. and Marazzi, A. (2012). Robust Gamma regression models for the analysis of health care cost data. *Model Assisted Statistics and Applications*; 7:115-124.
- [34] Punzo, A., Mazza, A. and Maruotti, A. (2018). Fitting insurance and economic data with outliers: a flexible approach based on finite mixtures of contaminated gamma distributions. *Journal of Applied Statistics*; 45(14):2563-2584.
- [35] Rousseeuw, P. J. and Leroy, A. M. (1987). *Robust regression and outlier detection*. USA: John Wiley & Sons.
- [36] Swallow, W. H. and Kianifard, F. (1996). Using robust scale estimates in detecting multiple outliers in linear regression. *Biometrics*; 52(2):545-556.
- [37] Telford, R. D. and Cunningham, R. B. (1991). Sex, Sport, and Body-size Dependency of Hematology in Highly Trained Athletes. *Medicine and Science in Sports and Exercise*; 23(7):788-794.
- [38] Warwick, J. and Jones, M. C. (2005). Choosing a robustness tuning parameter. *Journal of Statistical Computation and Simulation*; 75(7):581-588.
- [39] Zaman, A., Rousseeuw, P. J. and Orhan, M. (2001). Econometric applications of high-breakdown robust regression techniques. *Economic Letters*; 71:1-8.
- [40] Zeller, C. B., Cabral, C. R. B. and Lachos, V. H. (2016). Robust mixture regression modeling based on scale mixtures of skew-normal distributions. *Test*; 25:375-396.