

Refutability as Recursive as Provability

Paola Cattabriga

University of Bologna

[0000-0001-5260-2677](mailto:paola.cattabriga@unibo.it)

Abstract. Gödel numbering is an *arithmetization* of syntax which defines provability by coding a primitive recursive predicate, $\text{Pf}(x, v)$ [19, 190-198]. A multiplicity of researches and results all around this well-known recursive predicate are today widespread in many areas of logic and AI. Not equally investigated is the refutability predicate defined by Gödel numbering within the same primitive recursive status. $\text{Rf}(x, v)$ can be defined as a recursive predicate meaning that x is the Gödel number of a refutation in PA of the formula with Gödel number v . This article proposes a logical investigation of the interactive links between provability and refutability predicates when defined within the same recursive status. The resulting Lemmas are clarifying and open new perspectives for the incompleteness argument and the codings of its underlying notions.

Keywords: decision problem, provability predicate, Gödel numbering.

Introduction

Once the axioms and rules of inference in the calculus of first-order predicates have been established, thanks to Gödel's 1930 completeness, all the theorems can be derived. There are several formal ways to further describe deductive completeness, for example we can list, enumerate or generate all first-order theorems. Predicate calculus is considered semidecidable, since in first-order axiomatic theories we have a computable procedure for listing all predicate formulas that are theorems. We could also think of a machine, like the Turing machine, which writes all the first order theorems step by step on the tape. Despite all this, there are arguments, such as the so-called Church and Turing theorems, which affirm the undecidability of first-order calculus. We have truth tables as a decision method for propositional calculus, but we have no equivalent decision method for predicate calculus. In simple terms, taking any first order formula we cannot always decide whether it is true or false. At the basis of these arguments there is an upstream idea that sees the existence of formulas that are neither true nor false as an impasse. See for example the well-known words

... there are propositions which are neither true nor false but *indeterminate* [17, 126].

A conception which in Lukasiewicz then seems to move from semantic validity to syntactic demonstrability and stated in another form: "... there exist significant expressions, which can neither be proved by our axioms and rules of assertion nor disproved by our axioms and rules of rejection, I call such expressions undecidable with respect to our basis" [18, 100]. This *indeterminacy* is intrinsically linked to the non-denumerability and the infinite generation of formulas.

Is the number of undecidable expressions finite or not? If it is finite, the problem of decision is easily solved: we may accept true expressions as new asserted axioms, and reject false expressions axiomatically. The method, however, is not practicable if the number of undecidable expressions is not finite. We cannot assert or reject an infinity of axioms [18, 100].

Something quite similar seems to echo from Gödel. The completeness theorem establishes an equivalence between validity and provability, but this equivalence

entails, for the decision problem, a reduction of the nondenumerable to the denumerable, since valid refers to the nondenumerable totality of functions, while provable presupposes only the denumerable totality of the formal proofs [13].

From a historical point of view, we can trace a conceptual path that leads to *indeterminacy* in the face of infinity, going back to the Lowenheim-Skolem theorem, the Skolem paradox and the so-called Cantor theorem about the non-denumerability of the set of all the subsets of the natural numbers [20,4,5]. As is known, diagonalization is the basis of Cantor's reasoning and consequently also of Skolem's paradox. Moreover, Gödel's 1931 undecidable formula itself contains the codification of a form of self-referentiality, usually called the Diagonalization Lemma [14][15, 362-363][19, 203]. In this regard we have already pointed out the neglect and omission of the complement of the set or predicate made object of diagonalization [9,8]. Gödel derives the undecidable formula encoding the notion of provability by means of the famous primitive recursive predicate $\text{Pf}(x, v)$ [19, 190-198]. This derivation is achieved by completely neglecting the underlying definability of an equally recursive encoding of refutability [6,7,10]. As noted by Lukasiewicz:

Of two intellectual acts, to assert a proposition and to reject it, only the first has been taken into account in modern formal logic [18, 95].

Indeterminacy leads Lukasiewicz not only to reject the principle of bivalence and to introduce a third truth value, but also to the need to formalize the rejection. This aligns with our exploration of the definition of the complementary predicate of $\text{Pf}(x, v)$, namely $\text{Rf}(x, v)$ [6,10]. Below we reproduce and extend the investigation of the relations between these two predicates, since it seems to provide another way out of *indeterminacy*.

Basic Setup

We shall assume a first order theory which adequately formalizes Peano Arithmetic (see for example the system S , with all the necessary assumptions, in [19] 116-175). Let us call it PA . Numerals, as usual, are defined recursively, $\bar{0}$ is 0 and for any natural number n , $\overline{n+1}$ is $(\bar{n})'$ (where $'$ is the Successor function). For any expression X we use $\ulcorner X \urcorner$ to denote the corresponding Gödel number of X . Let us define the Gödel numbering as follows:

1. First assign different odd numbers to the primitive symbols of the language of PA .
2. Let X be a formal expression X_0, X_1, \dots, X_n , where each X_i , $0 \leq i \leq n$, is a primitive symbol of the language of PA . Then

$$\ulcorner X \urcorner = p_0^{\ulcorner X_0 \urcorner} \cdot p_1^{\ulcorner X_1 \urcorner} \cdot \dots \cdot p_n^{\ulcorner X_n \urcorner}$$

where p_n is the n -th prime number and $p_0 = 2$.

3. Let X be composed by the formal expressions X_0, X_1, \dots, X_n , then

$$\ulcorner X \urcorner = p_0^{\ulcorner X_0 \urcorner} \cdot \dots \cdot p_n^{\ulcorner X_n \urcorner}.$$

For any given formula $\phi(v)$ of PA we then have its Gödel number $n = \ulcorner \phi(v) \urcorner$. This number n has a name in the language of PA , namely \bar{n} , and this name can be substituted back into $\phi(v)$. This self-reference procedure is usually admitted by the so-called Diagonalization Lemma [6,9,8].

1 Refutation Predicate

We shall construct a new predicate by Gödel numbering. The reader can refer to the arithmetization as defined by Mendelson; the new predicate must be considered as a last relations added to the functions and relations (1-26) presented in [19, 190-198]. We shall not reproduce entirely this long list of definitions which is already well-known (see also [14, 162-176]). Let us start recalling some of the definitions involved, precisely only those we need.

$\text{MP}(x, y, z)$: The expression with Gödel number z is a direct consequence of the expressions with Gödel numbers x and y by modus ponens,

$$y = 2^3 * x * 2^{11} * z * 2^5 \wedge \text{Gd}(x) \wedge \text{Gd}(z).$$

$\text{Gen}(x, y)$: The expression with Gödel number y comes from the expression with Gödel number x by the Generalization Rule,

$$(\exists v)_{v < y} (\text{EVbl}(v) \wedge y = 2^3 * 2^3 * 2^{13} * v * 2^5 * x * 2^5 \wedge \text{Gd}(x)).$$

$\text{Ax}(y)$: y is the Gödel number of an axiom of PA :

$$\text{LAX}(y) \vee \text{PRAX}(y).$$

$\text{Neg}(v)$: the Gödel number of $(\neg\alpha)$ if v is the Gödel number of α :

$$\text{Neg}(v) = 2^3 * 2^9 * v * 2^5.$$

$\mathbf{Prf}(x)$: x is the Gödel number of a proof in PA :

$$\begin{aligned} & \exists u_{u < x} \exists v_{v < x} \exists z_{z < x} \exists w_{w < x} ([x = 2^w \wedge \mathbf{Ax}(w)] \vee \\ & [\mathbf{Prf}(u) \wedge \mathbf{Fml}((u)_w) \wedge x = u * 2^v \wedge \mathbf{Gen}((u)_w, v)] \vee \\ & [\mathbf{Prf}(u) \wedge \mathbf{Fml}((u)_z) \wedge \mathbf{Fml}((u)_w) \wedge x = u * 2^v \wedge \mathbf{MP}((u)_z, (u)_w, v)] \vee \\ & [\mathbf{Prf}(u) \wedge x = u * 2^v \wedge \mathbf{Ax}(v)]). \end{aligned}$$

$\mathbf{Pf}(x, v)$: x is the Gödel number of a proof in PA of the formula with Gödel number v :

$$\mathbf{Prf}(x) \wedge v = (x)_{lh(x) \cdot 1}.$$

By means of such definitions, we shall define a new predicate, \mathbf{Rf} .

$\mathbf{Rf}(x, v)$: x is the Gödel number of a proof in PA of the negation of the formula with Gödel number v :

$$\mathbf{Pf}(x, z) \wedge z = \mathbf{Neg}(v).$$

In other terms $\mathbf{Rf}(x, v)$ states x is the Gödel number of a refutation in PA of the formula with Gödel number v . \mathbf{Rf} is *primitive recursive*, as the relations obtained from primitive recursive relations by means of propositional connectives are also primitive recursive [16, 69][19, 180]. For their recursiveness $\mathbf{Pf}(x, v)$ and $\mathbf{Rf}(x, v)$ are *expressible* in PA by the formulas $Pf(x, v)$ and $Rf(x, v)$. Hereafter we will call $Pf(x, v)$ and $Rf(x, v)$ respectively the *proof* and the *refutation* predicate, while “provability” and “refutability” will refer to their existential assertions (section 3).

2 Proof and Refutation connections

From Luckasiewicz’s ‘assertion - rejection’ binomial to our current ‘proof - refutation’ we can see a historical progression, which goes from those systems of demonstration traditionally on the level of the thought and pure symbolic representation up to the systems of effective computation and automated deduction in computer science and artificial intelligence. Automatic theorem proving branch from two traditions of logical inquiry, the first via *resolution* from Herbrand’s theorem and Robinson’s resolution rule, and the second via *tableaux* from Gentzen’s natural deduction. The mechanical generation of first order theorems can be considered as an investigation that starts precisely from the deductive completeness of predicate calculus. The recursive definition of the proof predicate is historically the result of investigations both on the search of an algorithm to solve any mathematical problem and on problems of proof theory. The classical provability problem is that to decide, for a given a first order formula, if it is provable in the formal system [2]. For its recursiveness the proof predicate $\mathbf{Pf}(x, y)$ must be understood as an algorithm, i.e. an effective computation procedure, which outputs those first order arithmetical formulas that are the theorems of PA . The sequence of 45 numbered formulas defining the proof predicate in Gödel’s 1931 looks very much like a computer program, so that Kurt Gödel’s appears today to be like a pioneer of logic programming [12, 97-98][3, 14-19].

To visualize the algorithmic structure of this predicates let us display two simple examples. The first one, a propositional logic case, where x is the Gödel number of a proof of a formula with Gödel number y .

Example 1. Let us suppose we have the proof of $\vdash \neg\neg B \rightarrow B$ as follows.

- (1) $(\neg B \rightarrow \neg\neg B) \rightarrow ((\neg B \rightarrow \neg B) \rightarrow B),$
- (2) $(\neg B \rightarrow \neg B),$
- (3) $(\neg B \rightarrow \neg\neg B) \rightarrow B,$
- (4) $\neg\neg B \rightarrow (\neg B \rightarrow \neg\neg B),$
- (5) $\neg\neg B \rightarrow B.$

The corresponding Gödel numbers for each (1) - (5) would be:

$$\begin{aligned}
g(1) & 2^{33}3^{95}5^{17}11^{91}13^{91}17^{51}19^5 23^{11}29^3 31^3 37^9 41^{51}43^{11}47^9 53^{51}59^5 61^{11}67^{51}71^5 \\
g(2) & 2^{33}3^{95}5^{17}11^{91}13^{51}17^5 \\
g(3) & 2^{33}3^{95}5^{17}11^{91}13^9 17^{51}19^5 23^{11}29^5 \\
g(4) & 2^9 3^9 5^{17}11^{13}13^9 17^{51}19^{11}23^9 29^9 31^{51}37^5 \\
g(5) & 2^{33}3^{95}7^{51}11^{11}13^{51}17^5
\end{aligned}$$

So that in $\text{Pf}(x, y)$ x would be an expression like $2^{g(1)}u 3^{g(2)}u 5^{g(3)}u 7^{g(4)}u 11^{g(5)}$ and $y = g(5)$, (i.e. $\overline{\neg\neg B \rightarrow B}$).

Example 2. As a further simple example, the derivation in eleven steps by $\text{Rf}(x, y)$ of $\vdash \neg(t < t)$ in formal number theory:

- (1) $t < t,$
- (2) $\exists w(w \neq 0 \wedge w + t = t),$
- (3) $b \neq 0 \wedge b + t = t,$
- (4) $b + t = t,$
- (5) $t = 0 + t,$
- (6) $b + t = 0 + t,$
- (7) $b = 0,$
- (8) $b \neq 0,$
- (9) $b = 0 \wedge b \neq 0,$
- (10) $0 = 0 \wedge 0 \neq 0,$
- (11) $\neg(t < t)$

[19, 164]. It would yield for x an expression like $u 2^{g(1)}u 3^{g(2)} \dots 23^{g(10)}u 29^{g(11)}$ and for y $2^{33} 3^{865} 5^{33}$ (i.e. $\overline{\neg t < t}$)¹.

The link from each formula to its Gödel number is a one to one correspondence ensured by the uniqueness of the factorization of the integers into primes [19, 191]. To every primitive recursive predicate there corresponds a primitive recursive characteristic function, and since every primitive recursive function is computable then every recursive predicate is decidable [16, 66-67][11, 51-55]. Accordingly, both the proof predicate $\text{Pf}(x, y)$ and the refutation predicate $\text{Rf}(x, y)$ are also decidable. The first says us whether x is the Gödel number of a proof of the formula with Gödel number y or not, while the second says us whether x is the Gödel number of a refutation of the formula with Gödel number y or not. It therefore seems legitimate to ask, given such effective procedures, are there actually undecidable formulas? Or, in other words, where exactly is *indeterminacy*

¹ We regarded t as to be f_1^0 and $<$ as f_3^2 .

encoded? The connections between the two recursive predicates produces the followings lemmas, that will be useful for answering this questions (see section 3).

Lemma 1. *For any natural number n and for any formula α not both $\mathbf{Rf}(n, \ulcorner \alpha \urcorner)$ and $\mathbf{Pf}(n, \ulcorner \alpha \urcorner)$.*

Proof. Let us suppose to have both $\mathbf{Rf}(n, \ulcorner \alpha \urcorner)$ and $\mathbf{Pf}(n, \ulcorner \alpha \urcorner)$. We should have then $\mathbf{Prf}(n) \wedge \ulcorner \alpha \urcorner = (n)_{lh(n)-1}$ and $\mathbf{Pf}(n, z) \wedge z = \mathbf{Neg}(\ulcorner \alpha \urcorner)$, i.e. $\mathbf{Prf}(n) \wedge \ulcorner \alpha \urcorner = (n)_{lh(n)-1}$ and $\mathbf{Prf}(n) \wedge \mathbf{Neg}(\ulcorner \alpha \urcorner) = (n)_{lh(n)-1}$.

By the definition of $\mathbf{Prf}(x)$ this would mean to have
 $\exists u_{u < n} \exists v_{v < n} \exists z_{z < n} \exists w_{w < n} ([n = 2^w \wedge \mathbf{Ax}(w)] \vee$
 $[\mathbf{Prf}(u) \wedge \mathbf{Fml}((u)_w) \wedge n = u * 2^v \wedge \mathbf{Gen}((u)_w, v)] \vee$
 $[\mathbf{Prf}(u) \wedge \mathbf{Fml}((u)_z) \wedge \mathbf{Fml}((u)_w) \wedge n = u * 2^v \wedge \mathbf{MP}((u)_z, (u)_w, v)] \vee$
 $[\mathbf{Prf}(u) \wedge n = u * 2^v \wedge \mathbf{Ax}(v)])$ and both

$$\ulcorner \alpha \urcorner = (n)_{lh(n)-1} \quad \text{and} \quad \mathbf{Neg}(\ulcorner \alpha \urcorner) = (n)_{lh(n)-1}$$

and hence the four cases

1. $[n = 2^{\ulcorner \alpha \urcorner} \wedge \mathbf{Ax}(\ulcorner \alpha \urcorner)]$ and $[n = 2^{\mathbf{Neg}(\ulcorner \alpha \urcorner)} \wedge \mathbf{Ax}(\mathbf{Neg}(\ulcorner \alpha \urcorner))]$
2. $[\mathbf{Prf}(u) \wedge \mathbf{Fml}((u)_w) \wedge n = u * 2^{\ulcorner \alpha \urcorner} \wedge \mathbf{Gen}((u)_w, \ulcorner \alpha \urcorner)]$ and $[\mathbf{Prf}(u) \wedge \mathbf{Fml}((u)_w) \wedge n = u * 2^{\mathbf{Neg}(\ulcorner \alpha \urcorner)} \wedge \mathbf{Gen}((u)_w, \mathbf{Neg}(\ulcorner \alpha \urcorner))]$
3. $[\mathbf{Prf}(u) \wedge \mathbf{Fml}((u)_z) \wedge \mathbf{Fml}((u)_w) \wedge n = u * 2^{\ulcorner \alpha \urcorner} \wedge \mathbf{MP}((u)_z, (u)_w, \ulcorner \alpha \urcorner)]$ and $[\mathbf{Prf}(u) \wedge \mathbf{Fml}((u)_z) \wedge \mathbf{Fml}((u)_w) \wedge n = u * 2^{\mathbf{Neg}(\ulcorner \alpha \urcorner)} \wedge \mathbf{MP}((u)_z, (u)_w, \mathbf{Neg}(\ulcorner \alpha \urcorner))]$
4. $[\mathbf{Prf}(u) \wedge n = u * 2^{\ulcorner \alpha \urcorner} \wedge \mathbf{Ax}(\ulcorner \alpha \urcorner)]$ and $[\mathbf{Prf}(u) \wedge n = u * 2^{\mathbf{Neg}(\ulcorner \alpha \urcorner)} \wedge \mathbf{Ax}(\mathbf{Neg}(\ulcorner \alpha \urcorner))]$

which are all immediately impossible by the definitions of $\mathbf{Ax}(y)$, $\mathbf{Gen}(x, y)$ and $\mathbf{MP}(x, y, z)$ and thence by the definitions of the axioms of PA , the Generalization Rule and Modus Ponens, because no axiom belongs to PA together with its negation and the two inference rules preserve logical validity.

We now recall the definition of characteristic function. If R is a relation of n arguments, then the characteristic function C_R is defined as follows

$$C_R(x_1, \dots, x_n) = \begin{cases} 0 & \text{if } R(x_1, \dots, x_n) \text{ is true,} \\ 1 & \text{if } R(x_1, \dots, x_n) \text{ is false.} \end{cases}$$

Let us call the characteristic functions of $\mathbf{Pf}(x, v)$ and $\mathbf{Rf}(x, v)$ respectively $C_{\mathbf{Pf}}$ and $C_{\mathbf{Rf}}$. A relation $R(x_1, \dots, x_n)$ is said to be primitive recursive (recursive) if and only if its characteristic function $C_R(x_1, \dots, x_n)$ is primitive recursive (recursive) [19, 179-180]. As $\mathbf{Pf}(x, v)$ and $\mathbf{Rf}(x, v)$ are primitive recursive then also $C_{\mathbf{Pf}}$ and $C_{\mathbf{Rf}}$ are primitive recursive. Every recursive function is representable in PA , thence $C_{\mathbf{Pf}}$ and $C_{\mathbf{Rf}}$ are representable in PA . We shall assume C_{Pf} and C_{Rf} to represent respectively $C_{\mathbf{Pf}}$ and $C_{\mathbf{Rf}}$ in PA .

Lemma 2. For any formula α , and n as the Gödel number of a proof in PA of α

$$\vdash_{PA} C_{Pf}(\overline{n}, \overline{\ulcorner \alpha \urcorner}) = \overline{0} \wedge C_{Rf}(\overline{n}, \overline{\ulcorner \alpha \urcorner}) = \overline{1}$$

Proof. One can easily see that the two conjuncts are true: as n is the Gödel number of a proof in PA of α $C_{Pf}(\overline{n}, \overline{\ulcorner \alpha \urcorner}) = \overline{0}$ is true. By Lemma (1) $Rf(n, \ulcorner \alpha \urcorner)$ does not hold, therefore it is true that n is not the Gödel number of a refutation in PA of α .

Lemma 3. For any formula α , and n as the Gödel number of a refutation in PA of α

$$\vdash_{PA} C_{Rf}(\overline{n}, \overline{\ulcorner \alpha \urcorner}) = \overline{0} \wedge C_{Pf}(\overline{n}, \overline{\ulcorner \alpha \urcorner}) = \overline{1}$$

Proof. One can easily see that the two conjuncts are true: as n is the Gödel number of a refutation in PA of α $C_{Rf}(\overline{n}, \overline{\ulcorner \alpha \urcorner}) = \overline{0}$ is true. By Lemma (1) $Pf(n, \ulcorner \alpha \urcorner)$ does not hold, therefore it is true that n is not the Gödel number of a proof in PA of α .

Lemma 4. For any formula α

(I) not both

$$\vdash_{PA} Pf(\overline{n}, \overline{\ulcorner \alpha \urcorner}) \wedge \vdash_{PA} Rf(\overline{n}, \overline{\ulcorner \alpha \urcorner}),$$

(II) for n as the Gödel number of a refutation in PA of α

$$\vdash_{PA} Rf(\overline{n}, \overline{\ulcorner \alpha \urcorner}) \iff \neg Pf(\overline{n}, \overline{\ulcorner \alpha \urcorner}),$$

(III) for n as the Gödel number of a proof in PA of α

$$\vdash_{PA} Pf(\overline{n}, \overline{\ulcorner \alpha \urcorner}) \iff \neg Rf(\overline{n}, \overline{\ulcorner \alpha \urcorner}).$$

Proof. (I) Immediately by Lemma (1) and the definition of being expressible which holds for both $Pf(x, v)$ and $Rf(x, v)$ ([19] 130).

(II) Let us assume $\vdash_{PA} Rf(\overline{n}, \overline{\ulcorner \alpha \urcorner})$, then Lemma (3) yields $\vdash_{PA} C_{Pf}(\overline{n}, \overline{\ulcorner \alpha \urcorner}) = \overline{1}$. Hence by definition $Pf(\overline{n}, \overline{\ulcorner \alpha \urcorner})$ is false, consequently $\vdash_{PA} \neg Pf(\overline{n}, \overline{\ulcorner \alpha \urcorner})$. Conversely let us assume $\vdash_{PA} \neg Pf(\overline{n}, \overline{\ulcorner \alpha \urcorner})$ then $Pf(\overline{n}, \overline{\ulcorner \alpha \urcorner})$ is false and by Lemma (3) we attain $\vdash_{PA} Rf(\overline{n}, \overline{\ulcorner \alpha \urcorner})$.

(III) Let us assume $\vdash_{PA} Pf(\overline{n}, \overline{\ulcorner \alpha \urcorner})$, then Lemma (2) yields $\vdash_{PA} C_{Rf}(\overline{n}, \overline{\ulcorner \alpha \urcorner}) = \overline{1}$. Hence by definition $Rf(\overline{n}, \overline{\ulcorner \alpha \urcorner})$ is false, consequently $\vdash_{PA} \neg Rf(\overline{n}, \overline{\ulcorner \alpha \urcorner})$. Conversely let us assume $\vdash_{PA} \neg Rf(\overline{n}, \overline{\ulcorner \alpha \urcorner})$ then $Rf(\overline{n}, \overline{\ulcorner \alpha \urcorner})$ is false and by Lemma (2) we attain $\vdash_{PA} Pf(\overline{n}, \overline{\ulcorner \alpha \urcorner})$.

3 Provability and Refutability

In general, an n -ary predicate P is recursively enumerable if it is empty or if there exist n singularly recursive functions $f_1 \dots f_n$ such that for all $x_1 \dots x_n$

$$P(x_1 \dots x_n) \iff \exists y (f_1(y) = x_1 \wedge \dots \wedge f_n(y) = x_n).$$

In addition, every predicate $\exists yP(x_1 \dots x_n y)$ with a recursive kernel P is a recursively enumerable predicate, r.e. predicate, and every r.e. predicate can be represented in this way [16, 188-189] [1, 103]. In other terms, applying an unlimited existential quantifier to a predicate whose kernel is recursive yields a r. e. predicate. This is the case of the famous “Bew(x)”, the first one of the list of notions 1–46 in Gödel’s 1931 of which we cannot say that it is recursive, being defined as an assertion of existence. Let us quote its definition [14, 171]:

45. $xBy \equiv Bw(x) \ \& \ [l(x)] \ Gl \ x = y,$
 x is a PROOF of the FORMULA y .

46. $Bew(x) \equiv (Ey)yBx,$
 x is a PROVABLE FORMULA. .

$(Ey)yBx$ states it does exist a proof y of the formula x , and it is accordingly assumed throughout Gödel’s 1931 argumentation as an assertion of existence, to derive the famous undecidable formula. Let us consider $Bew(x)$, i.e. in our modern notation $\exists yPf(y, x)$, which assumes the existence of a proof y of x , in the light of our lemmas.

There is no doubt that from “ n as the Gödel number of a proof in PA of α ” and $\exists yPf(y, x)$ we obtain $Pf(\bar{n}, \overline{\alpha})$. Accordingly by Lemma (4)(III) $\vdash_{PA} Pf(\bar{n}, \overline{\alpha}) \iff \neg Rf(\bar{n}, \overline{\alpha})$. And, on the other hand, from “ n as the Gödel number of a refutation in PA of α ” and $\exists yRf(y, x)$, we have $Rf(\bar{n}, \overline{\alpha})$, and by Lemma (4)(II) $\vdash_{PA} Rf(\bar{n}, \overline{\alpha}) \iff \neg Pf(\bar{n}, \overline{\alpha})$.

Let us firstly resume the usual way to reason about as follows.

- a) Being $Pf(x, v)$ decidable, for $Pf(\bar{n}, \overline{\alpha})$ we can decide whether n is the Gödel number of a proof in PA of α , namely whether $\vdash_{PA} \alpha$ or $\not\vdash_{PA} \alpha$. But when we assert $\exists xPf(x, \overline{\alpha})$ then we can only generate all the formulas α such that $\vdash_{PA} \alpha$, and we could not be able to generate all the formulas such that $\not\vdash_{PA} \alpha$.
- b) By $Rf(x, v)$, for $Rf(\bar{n}, \overline{\alpha})$ we can decide whether n is the Gödel number of a refutation in PA of α , so we can decide whether $\vdash_{PA} \neg\alpha$ or $\not\vdash_{PA} \neg\alpha$. Likewise, to state $\exists xRf(x, \overline{\alpha})$ surely generate all the formulas α such that $\vdash_{PA} \neg\alpha$, but we could not be able to generate all the formulas such that $\not\vdash_{PA} \neg\alpha$.

But now thanks to Lemma (4) when in PA we assume that $\exists xPf(x, \overline{\alpha})$ (or $\exists xRf(x, \overline{\alpha})$) we know something more i.e. (4)(III) (or (4)(II)).

- a’) If in PA we state that $\exists xPf(x, \overline{\alpha})$ we can add (4)(III), then for n as the Gödel number of a proof in PA of α , we have not only $\vdash_{PA} Pf(\bar{n}, \overline{\alpha})$ and hence $\vdash_{PA} \alpha$, but also $\vdash_{PA} \neg Rf(\bar{n}, \overline{\alpha})$, namely $\not\vdash_{PA} \neg\alpha$.
- b’) When in PA we state $\exists xRf(x, \overline{\alpha})$ we can add (4)(II). That is for n as the Gödel number of a refutation in PA of α , not only $\vdash_{PA} Rf(\bar{n}, \overline{\alpha})$, i.e. $\vdash_{PA} \neg\alpha$, but consequently also $\vdash_{PA} \neg Pf(\bar{n}, \overline{\alpha})$, namely $\not\vdash_{PA} \alpha$.

This also explains our previous results in [6,7,9,10].

Remark 1. We are then able to notice that, being the predicate $\mathbf{Rf}(x, v)$ recursively defined, as much as $\mathbf{Pf}(x, v)$ is, it comes to our aid with respect to Gödel's 1931 "Bew(x)". As noticed in the introduction this is something at all in twilights within Gödel's incompleteness argument. Moreover, the recursive definition of $\mathbf{Rf}(x, v)$ highlights how, apart from the self-referring nested in the definition $Q(x, y)$ [14, 175], the assumption of the existence of a proof y when stating $(\exists y)yBx$ is precisely a codification, or formal expression, of the *indeterminacy*. Indeterminacy, that in the light of the Lemma (4) has no real reason to exist.

As a concluding paragraph, let us explain further how all the four possible cases $\vdash \alpha$, $\not\vdash \alpha$ and $\vdash \neg \alpha$, $\not\vdash \neg \alpha$ are now ruled in PA .

As already mentioned, C_{Pf} is computable and $Pf(x, y)$ decidable. For n a Gödel number of a proof in PA of α , $\langle \bar{n}, \overline{\alpha} \rangle \in Pf \leftrightarrow C_{Pf}(\bar{n}, \overline{\alpha}) = \bar{0}$, and we can decide whether $\vdash_{PA} \alpha$ or $\not\vdash_{PA} \alpha$. See example 1, $\langle 2^{g(1)}u3^{g(2)}u5^{g(3)}u7^{g(4)}u11^{g(5)}, \overline{\neg \neg B \rightarrow B} \rangle \in Pf$, where we can decide whether $\neg \neg B \rightarrow B$ is a theorem or not, in PA . But when we state $\exists x Pf(x, \overline{\alpha})$ we make an assumption of existence, i.e. of a r. e. predicate, that can be represented as follows

$$i) \quad Pf(0, \overline{\alpha}) \vee Pf(1, \overline{\alpha}) \vee Pf(2, \overline{\alpha}) \vee Pf(3, \overline{\alpha}) \vee \dots$$

[11, xxiv-xxv]. In other terms, we have a computable function f which lists all the Gödel numbers of proofs in PA yielding the set of all the ordered couples $\langle f(0), \overline{\alpha} \rangle, \langle f(1), \overline{\alpha} \rangle, \langle f(2), \overline{\alpha} \rangle \dots$ till to reach that \bar{n} such that effectively $\langle \bar{n}, \overline{\alpha} \rangle \in Pf$. See the example 1, then we would be restricted by $i)$ to have a computable function enumerating all the proof in PA till to reach the first couple such that $\langle 2^{g(1)}u3^{g(2)}u5^{g(3)}u7^{g(4)}u11^{g(5)}, \overline{\neg \neg B \rightarrow B} \rangle \in Pf$. Being $\exists x Pf(x, \overline{\alpha})$ only r.e. we can generate the Gödel number of a proof in PA of α , i.e. $\vdash_{PA} \alpha$, but we could not be able to obtain $\not\vdash_{PA} \alpha$.

Respectively, $C_{Rf} \langle \bar{n}, \overline{\alpha} \rangle \in Rf \leftrightarrow C_{Rf}(\bar{n}, \overline{\alpha}) = 0$, and since $Rf(x, y)$ is decidable we can decide whether n is the Gödel number of a refutation in PA of α , i.e. whether $\vdash_{PA} \neg \alpha$ or $\not\vdash_{PA} \neg \alpha$. In example 2, we can decide whether $t < t$ is a refutation or not in PA . But with the assertion $\exists x Rf(x, \overline{\alpha})$ we have a r.e. predicate, such that

$$ii) \quad Rf(0, \overline{\alpha}) \vee Rf(1, \overline{\alpha}) \vee Rf(2, \overline{\alpha}) \vee Rf(3, \overline{\alpha}) \vee \dots,$$

and we can reach $\vdash_{PA} \neg \alpha$, but we could not be able to generate a derivation such that $\not\vdash_{PA} \neg \alpha$. In example 2 we could be able to reach $\langle u2^{g(1)}u3^{g(2)} \dots u23^{g(10)}u29^{g(11)}, \overline{t < t} \rangle \in Rf$ but we could never obtain a derivation such that $\not\vdash_{PA} \neg(t < t)$.

Let us therefore extend our investigation for the assertion of existence. For "n the Gödel number of a proof in PA of α " from $\exists x Pf(x, y)$ we obtain $Pf(\bar{n}, \overline{\alpha})$.

By Lemma (4)(III) $\vdash_{PA} Pf(\bar{n}, \overline{\alpha}) \iff \neg Rf(\bar{n}, \overline{\alpha})$, and from the viewpoint of the effective computability this is equivalent to generating from *i*)

$$iii) \quad \neg Rf(0, \overline{\alpha}) \vee \neg Rf(1, \overline{\alpha}) \vee \neg Rf(2, \overline{\alpha}) \vee \neg Rf(3, \overline{\alpha}) \vee \dots$$

In accordance, from *i*) and *iii*) we can attain both $\vdash_{PA} \alpha$ and $\not\vdash_{PA} \neg\alpha$. On the other side, for “*n* is the Gödel number of a refutation in PA of α ”, $\exists x Rf(x, y)$, we can obtain $Rf(\bar{n}, \overline{\alpha})$. Then by Lemma (4)(II) $\vdash_{PA} Rf(\bar{n}, \overline{\alpha}) \iff \neg Pf(\bar{n}, \overline{\alpha})$, from *ii*)

$$iv) \quad \neg Pf(0, \overline{\alpha}) \vee \neg Pf(1, \overline{\alpha}) \vee \neg Pf(2, \overline{\alpha}) \vee \neg Pf(3, \overline{\alpha}) \vee \dots,$$

and from *ii*) and *iv*) we can obtain both $\not\vdash_{PA} \alpha$ and $\vdash_{PA} \neg\alpha$. Let us notice also that an assertion of existence like $\exists x \neg Pf(x, y)$ would on its turn generate the lists *iv*) and *ii*), while $\exists x \neg Rf(x, y)$ would yield *iii*) and *i*).

By reason of all that, about provability and refutability, *indeterminacy* has no reason to be. And this is also an explanation of the reason behind our previous results in [6,7,9,10]. For objections about theories with more than denumerable models see [9,8].

References

1. Börger, E. Egon. *Computability, complexity, logic*. Elsevier Science Pub. Co., N.Y., 1989.
2. Börger, E., Grädel E., Gurevich Y. *The Classical Decision Problem*. Springer-Verlag, Berlin, 2001.
3. Chaitin, J. Gregory. *The Unknowable*. Springer-Verlag, Singapore, 1999.
4. Cantor, Georg. Über eine Eigenschaft des Inbegriffes aller reellen algebraischen Zahlen. In *Journal für die reine und angewandte Mathematik*, Vol. 77, 1874, pp. 258-262.
5. Cantor, Georg. Über eine elementare Frage der Mannigfaltigkeitslehre. In *Jahres-bericht der Deutschen Mathematiker-Vereinigung*, Vol. 1, 1891, pp. 75-78.
6. Cattabriga, Paola. Beyond Undecidable. Proceedings IC-AI'2000, Vol. III, CSREA pp. 1475-1481, [arXiv:math/0606713](https://arxiv.org/abs/math/0606713).
7. Cattabriga, Paola. Observations concerning Gödel's 1931. [arXiv:math/0306038](https://arxiv.org/abs/math/0306038); Abs. in *Bulletin of Symbolic Logic*, Vol. 13, N. 3, p. 382 (2007).
8. Cattabriga, Paola. Diagonalization as a Definition, Proceedings InterSymp 2019, IAS.
9. Cattabriga, Paola. Uniqueness Violations. Presented at *SLAI-2022*. In *Logic & Artificial Intelligence*, V. Andrunachievici Institute of Mathematics and Computer Science, Edited by Cojocaru Drugus Nikitchenco Muravitsky, Chisinau, 2023, pp. 134-144.
10. Cattabriga, Paola. On Gödel's treatment of the undecidable in 1931. [arXiv:2403.19665](https://arxiv.org/abs/2403.19665). Round Table Revisiting Gödel's Incompleteness Theorem WCP Rome 2024.

11. Davis, Martin. *Computability and Unsolvability*. McGraw-Hill, N.Y., 1958.
12. Davis, Martin. *The Universal Computer*. CRC Press, 2018.
13. Gödel, Kurt. The completeness of the axioms of the functional calculus of logic. In *Collected Works*, Vol. I Publications 1929-1936. Oxford University Press, New York, 1986, pp. 102-123.
14. Gödel, Kurt. On formally undecidable proposition of Principia mathematica and related systems I. 1931. In *Collected Works*, Vol. I Publications 1929-1936. Oxford University Press, New York, 1986, pp. 145-195.
15. Gödel, Kurt. On undecidable proposition of formal mathematical systems. 1934. In *Collected Works*, Vol. I Publications 1929-1936. Oxford University Press, New York, 1986, pp. 346-371.
16. Hermes, Hans. *Enumerability Decidability Computability*. Springer-Verlag, New York, 1969.
17. Lukasiewicz, Jan. 1922 On Determinism. In Jan Lukasiewicz *Selected Works*, North-Holland Publishing, Amsterdam, 1970.
18. Lukasiewicz, Jan. 1951 *Aristotele's syllogistic from the standpoint of modern formal logic*. Clarendon Press, Oxford, 1957.
19. Mendelson, Elliott. *Introduction to Mathematical Logic* Third Edition. The Wadsworth & Brooks, Pacific Grove, California, 1987. See also Fourth Edition, Chapman & Hall, 1997.
20. Skolem, Thoralf. 1922 Some remarks on axiomatized set theory. In *From Frege to Gödel: A Source Book in Mathematical Logic, 1879-1931.*, Harvard University Press, Cambridge, Massachusetts, 1967.