ON THE VORTICITY THRESHOLD FOR STEADY WATER WAVES

EVGENIY LOKHARU

Abstract. This paper presents a comprehensive analysis of two-dimensional water waves characterized by a significant adverse constant vorticity over flows without stagnation points. Surprisingly, we discover qualitative distinctions between this scenario and those featuring favorable vorticity. Our findings reveal that in the presence of large adverse vorticity, neither Stokes waves nor solitary waves can approach extreme waves exhibiting surface stagnation points. Conversely, we demonstrate that global bifurcation curves always lead to unidirectional Stokes and solitary waves with precisely one stagnation point located at the bottom right beneath the crest. In contrast, it is known that any favorable or even small adverse vorticity eliminates the possibility of bottom stagnation, leading to extreme waves exhibiting surface singularities. Additionally, we establish several novel facts, including new bounds for both amplitude and the Froude number.

1. INTRODUCTION

Waves over rotational flows are common in nature and one of the most known mathematical models is Euler equations with a constant vorticity. The assumption of a linear current is not only physically relevant but also simplifies mathematical aspects of the theory. In the present paper we will consider the case of adverse constant vorticity in the unidirectional setting, when no stagnation points are allowed. The corresponding stream function formulation is

$$
\Delta \psi + \omega = 0 \qquad \text{in } D_{\eta}, \tag{1.1a}
$$

$$
\frac{1}{2}|\nabla\psi|^2 + y = r \qquad \text{on } y = \eta,\tag{1.1b}
$$

$$
\psi = 1 \qquad \text{on } y = \eta,\tag{1.1c}
$$

$$
\psi = 0 \qquad \text{on } y = 0. \tag{1.1d}
$$

Note that we use non-dimensional variables, where the mass flux and the gravitational constant are scaled to the unity. Throughout the paper we assume $\omega = -\gamma$ with $\gamma > 0$ to be a large constant.

In what follows we will consider the case when the flow is unidirectional, that is,

$$
\psi_y > 0 \tag{1.1e}
$$

everywhere in D_n . We will prove below that this condition is very restrictive, provided the vorticity ω is a large negative constant, that is adverse. In this case all solutions subject to [\(1.1e\)](#page-0-0) are close to laminar flows. In particular, they are of small-amplitude. Our main results are the following.

Theorem 1.1. For any unidirectional Stokes or solitary wave solution (ψ, η) with $r \in (R(s_c), R(0))$ the following is true.

(i) Everywhere at the surface we have

$$
\psi_y \ge \gamma \check{\eta} - O(\gamma^{-2}) = \sqrt{2\gamma} - O(\gamma^{-1}).
$$

(ii) At every crest we have

$$
\psi_y \ge \sqrt{2\gamma} - O(\gamma^{-\frac{5}{2}}), \quad \hat{\eta} \le \sqrt{\frac{2}{\gamma}} + O(\gamma^{-\frac{7}{2}}).
$$

(iii) The amplitude satisfies

$$
\hat{\eta} - \check{\eta} = O(\gamma^{-2}).
$$

Date: April 9, 2024.

Here $\hat{\eta} = \max \eta$ and $\check{\eta} = \min \eta$ are the water depths at the crest and trough respectively. The first statement (i) forbids surface stagnation points for unidirectional waves. In (ii) we have a bound for $\hat{\eta}$ which is sharp in the following sense. Note that the value $d(0) = \sqrt{2/\gamma}$ coincides with the depth of the unidirectional flow enjoying stagnation at the bottom. There is also an analogue of that $d(s_0)$ for favourable vorticity and it is known that $\hat{\eta} < d(s_0)$ for all unidirectional Stokes waves and favourable vorticity. In the case of adverse and large vorticity, on the other hand, $\hat{\eta}$ may exceed $d(0)$ but nor much as required in (ii). For instance, every unidirectional wave with a bottom stagnation, that are proved to exist in the theorems below, must have $\hat{\eta} > \sqrt{\frac{2}{\gamma}}$.

The last claim (iii) shows that all unidirectional waves with a strong adverse vorticity are of small amplitude. The estimate (iii) is similar to the ones in [\[CSV21,](#page-10-0) [LWW23\]](#page-10-1) for a favourable constant vorticity.

The assumption $r < \gamma + \sqrt{2/\gamma}$ is somewhat natural, since the Bernoulli constant of every laminar flow subject to [\(1.1e\)](#page-0-0) and giving rise to a local bifurcation of Stokes waves satisfies this inequality. The best known upper bound for r was obtained in [\[Lok21\]](#page-10-2) but we prove a sharp bound $r < \gamma + \sqrt{2/\gamma}$ for large γ in Theorem [4.4.](#page-8-0)

The claim of Theorem [1.1](#page-0-1) partly justifies the numerical results of [\[KS08b,](#page-10-3) [KS08a\]](#page-10-4). It was observed there a certain threshold value for the vorticity γ_{\star} such that no unidirectional extreme waves appear through global bifurcations with $\gamma > \gamma_{\star}$, while values $\gamma < \gamma_{\star}$ always lead to extreme waves. When $\gamma > \gamma_{\star}$ one finds instead solutions with stagnation points located at the bottom right below crests. The corresponding bifurcation curves can then be extended further giving rise to solutions with critical layers attached to the bottom; see [\[KKL20\]](#page-10-5), where solitary waves of this type were found.

As a corollary we prove the following existence results.

Theorem 1.2 (The existence of unidirectional solitary waves with bottom stagnation). For any large enough $\gamma > 0$ let C be the global bifurcation curve of unidirectional solitary waves of elevation constructed in [\[Whe13\]](#page-10-6). Then there is a sequence $(\psi^{(j)}, \eta^{(j)}, r^{(j)}) \in \mathcal{C}$ converging to a solitary wave (ψ^*, η^*, r^*) such that $\eta^* \in C^{\infty}(\mathbb{R})$ and $\psi^*_{y} > 0$ everywhere in the fluid expect one point at the bottom right below the crest, where $\psi_y^* = 0$.

A similar statement holds true for bifurcations of Stokes waves.

Theorem 1.3 (The existence of unidirectional solitary waves with bottom stagnation). For any large enough $\gamma > 0$ let C be the analytic global bifurcation curve of unidirectional Stokes waves with a fixed wavelength constructed in [\[Var20,](#page-10-7) Theorem 18]. We assume that this curve starts at any unidirectional laminar flow $(U(y; s); d(s))$ with $0 < s < s_c$. Then there is a sequence $(\psi^{(j)}, \eta^{(j)}, r^{(j)}) \in \mathcal{C}$ converging to a Stokes wave (ψ^*, η^*, r^*) such that $\eta^* \in C^\infty(\mathbb{R})$ and $\psi_y^* > 0$ everywhere in the fluid expect one point at the bottom right below the crest, where $\psi_y^* = 0$.

By a Stokes wave we mean a smooth periodic solution to the problem [\(1.1\)](#page-0-2) whose surface profile has exactly one trough and one crest in each minimal period and is monotone in between. Stokes waves are typical solutions arising as small-amplitude perturbations of laminar flows. There are also large-amplitude Stokes waves and even extreme Stokes waves enjoying surface singularities at every crest, where $\psi_y = 0$. However, as follows from our Theorem [1.1](#page-0-1) such extreme waves (or more precisely connected families of Stokes waves leading to an extreme wave) are not possible when ω is negative and large.

2. Preliminaries

Before proving Theorem [1.1](#page-0-1) we will verify the following more technical result.

Theorem 2.1. There exist absolute constants $\gamma_{\star}, \delta_{\star} > 0$ such that there is no Stokes or solitary wave solution (ψ, η) with $\gamma > \gamma_{\star}$, $r < \gamma + \sqrt{2/\gamma}$ and such that

$$
\min \psi_y(x, \eta(x)) = \sqrt{2\gamma} - \frac{\delta_\star}{\gamma},\tag{2.1}
$$

provided [\(1.1e\)](#page-0-0) holds true everywhere in the fluid.

An important part of the analysis is related to laminar flows subject to [\(1.1e\)](#page-0-0), whose stream functions can be obtained as

$$
U(y; s) = \frac{1}{2}\gamma y^2 + sy, \ \ s > 0.
$$

The corresponding formulas for the depth and the Bernoulli constant are given by

$$
d(s) = \frac{-s + \sqrt{s^2 + 2\gamma}}{\gamma}, \quad R(s) = \frac{1}{2}s^2 + \gamma + d(s).
$$

Thus, for each $s > 0$ the pair $\psi = U(y; s)$ and $\eta = d(s)$ solve [\(1.1\)](#page-0-2) with $r = R(s)$ and satisfy [\(1.1e\)](#page-0-0). For more on unidirectional flows we refer to [\[KKL15\]](#page-10-8).

The formula for $d(s)$ implies that the depth is a monotonically decreasing function of $s > 0$, while $R(s)$ decreases for $s \in [0, s_c]$ and increases for $s \geq s_c$ for some critical value s_c defined by the relation $R'(s) = 0$. This equation has a unique solution $s = s_c$, which satisfies

$$
s_c - \frac{1}{\gamma} + \frac{s_c}{\gamma \sqrt{s_c^2 + 2\gamma}} = 0.
$$

Assuming γ is large, we have

$$
s_c = \frac{1}{\gamma} + O(\gamma^{-3/2}).
$$

For the corresponding depth, we find

$$
d(s_c) = \frac{-s_c + \sqrt{s_c^2 + 2\gamma}}{\gamma} = \sqrt{\frac{2}{\gamma}} - \frac{1}{\gamma^2} + O(\gamma^{-5/2}).
$$

Note also that for any $r \in (R(s_c), R(0)]$ there are exactly two roots $s_-(r) < s_+(r)$ to the equation $R(s) = r$.

The largest possible values for $d = d(s)$ is attained at $s = 0$ and is given by

$$
d(0) = \sqrt{\frac{2}{\gamma}},
$$

while the corresponding Bernoulli constant is

$$
R(0) = \gamma + \sqrt{\frac{2}{\gamma}}.
$$

It is known from [\[KKL15\]](#page-10-8) that an arbitrary non-laminar and unidirectional solution corresponds the Bernoulli constant that is strictly greater than $R(s_c)$. Furthermore, unidirectional laminar flows have Bernoulli constants strictly less than $R(0)$. This explains one of the assumptions we make below regarding the Bernoulli constant, that is

$$
r \in (R(s_c), R(0)).
$$

Using the formula for $d(s_c)$ it is clear that any such r is subject to

$$
r = \gamma + \sqrt{\frac{2}{\gamma}} + O(\gamma^{-2}).
$$
\n(2.2)

Asymptotics of this type are crucial for the proofs. Below we will obtain expansions for $\check{\eta}$ = $\min(\eta)$ and $\psi_y|_{y=0}$ at the bottom.

Proposition 2.2. Every smooth solution (ψ, η) with $r \in (R(s_c), R(0))$ is subject to the bounds

$$
d(s_+(R(0))) < \check{\eta} < d(0). \tag{2.3}
$$

In particular, we get

$$
\check{\eta} = \sqrt{\frac{2}{\gamma}} + O(\gamma^{-2}).\tag{2.4}
$$

Proof. The inequalities

$$
d_{-}(r) < \check{\eta} < d(0)
$$

were proved in [\[KKL15\]](#page-10-8) (note that $d_{-}(r)$ there is exactly $d(s_{+}(r))$). In order to verify [\(2.3\)](#page-2-0) it is left note that $d(s_{+}(R(0))) < d_{-}(r)$ since $s_{+}(R(0)) > s_{+}(r)$ and d is a decreasing function of s. For the remaining part [\(2.4\)](#page-2-1) we need to compute $\bar{s} := s_+(R(0))$, which solves

$$
\frac{1}{2}\bar{s}^2 + \gamma + d(\bar{s}) = R(0).
$$

Using the corresponding formulas for $d(\bar{s})$ and $R(0)$ we can rewrite the last equality as

$$
\frac{1}{2}\bar{s}^2 + \frac{-\bar{s} + \sqrt{\bar{s}^2 + 2\gamma}}{\gamma} = \sqrt{\frac{2}{\gamma}}.
$$

From here one computes $\bar{s} = s_+(R(0)) = O(\gamma^{-1})$ and so

$$
d(s_{+}(R(0))) = \sqrt{\frac{2}{\gamma}} + O(\gamma^{-2}).
$$

This and [\(2.3\)](#page-2-0) together give the desired asymptotics, since $d(0) = \sqrt{2/\gamma}$.

The next statement will be proved only for Stokes waves, which are periodic solutions to the problem with the surface profile being symmetric around each crest and trough and monotone in between. We will assume that $x = -\Lambda$ and $x = 0$ are the coordinates for the trough and crest respectively; here Λ is the half-period. Under given assumptions we have $\psi_x(x, y) < 0$ for all $x \in (-\Lambda, 0)$ and $y \in (0, \eta(x))$. This fact will be used in the proof below. More details on properties of Stokes waves can be found in [\[CS04\]](#page-10-9).

Proposition 2.3. There exist absolute constants $\gamma_0 > 0$ and $C_0 > 0$ such that for any smooth Stokes or solitary wave solution (ψ, η) with $r \in (R(s_c), R(0))$ we have

$$
\psi_y(x,0) \le \frac{C_0}{\gamma} \tag{2.5}
$$

uniformly in $x \in \mathbb{R}$, provided $\gamma > \gamma_0$.

Proof. When proving this and later statements we will only consider the case of a Stokes wave. A similar argument will always work also for solitary waves with no or trivial modifications. In order to obtain an upper bound for ψ_y at the bottom we need to estimate $\psi_y(-\Lambda, 0)$ right below the trough. This is because $\psi_{xy} < 0$ on $(-\Lambda, 0)$ and so $\psi_y(x, 0)$ is decreasing there. To prove this one can apply the Hopf lemma to the harmonic function ψ_x being positive for all $x \in (-\Lambda, 0)$ and $y \in (0, \eta(x))$ and zero for $y = 0$. Furthermore, by the same Hopf lemma one finds $\psi_{xx} \leq 0$ everywhere below the trough and so

 $\psi_{yy} \geq \gamma$

there for all $y \in [0, \eta(-\Lambda)]$ as follows from [\(1.1a\)](#page-0-3). In particular, we see that

$$
\psi_y \ge \psi_y|_{y=0} + \gamma y
$$

everywhere below the trough. Now we integrate this inequality over $[0, \tilde{\eta}]$ and obtain

$$
1 \geq \check{\eta}\psi_y(-\Lambda,0) + \tfrac{1}{2}\gamma\check{\eta}^2.
$$

Next we use [\(2.4\)](#page-2-1) to find

$$
1 \ge \left(\sqrt{\frac{2}{\gamma}} + O(\gamma^{-2})\right) \psi_y(-\Lambda, 0) + \frac{1}{2}\gamma \left(\frac{2}{\gamma} + O(\gamma^{-5/2})\right).
$$

From here one gets $\psi_y(-\Lambda, 0) = O(\gamma^{-1})$ as desired.

Proposition 2.4. There are absolute constants $C_1, C_2 > 0$ such that for any $\delta > 0$ there exists a constant $\gamma_1(\delta)$ depending only on δ with the following property. For any Stokes or solitary wave solution (ψ, η) with $\gamma > \gamma_1(\delta)$, $r \in (R(s_c), R(0))$ and satisfying the inequality

$$
\psi_y \geq \sqrt{2\gamma} - \frac{\delta}{\gamma}
$$

at every crest we have

$$
\psi_y \ge C_1 \gamma \delta^{-1} y \tag{2.6}
$$

everywhere in the fluid. Furthermore, we have

$$
\hat{\eta} \le C_2 \delta \sqrt{\frac{2}{\gamma}} \tag{2.7}
$$

for some another absolute constant C_2 .

Proof. We consider a harmonic function

$$
h=\psi_y-\lambda y
$$

for some $\lambda > 0$ to be chosen. Note that $h \geq 0$ on $y = 0$, while

$$
h\geq \sqrt{2\gamma}-\frac{\delta}{\gamma}-\lambda \hat{\eta}
$$

on $y = \eta$. Note that from the Bernoulli equation we get

$$
\frac{1}{2}\left(\sqrt{2\gamma}-\frac{\delta}{\gamma}\right)^2+\hat{\eta}\leq \gamma+\sqrt{\frac{2}{\gamma}}+O(\gamma^{-2}),
$$

which gives

$$
\hat{\eta} \leq C \delta \sqrt{\frac{2}{\gamma}} + O(\gamma^{-2})
$$

with some absolute C . Thus, we can choose

$$
\lambda=\frac{\gamma}{2C\delta}
$$

and so

$$
h\geq \sqrt{\frac{\gamma}{2}}-\frac{\delta}{\gamma}\geq 0
$$

on the surface if $\gamma \geq \gamma_1(\delta)$ for some γ_1 depending on δ . Now the maximum principle gives the desired inequality. desired inequality.

Proposition 2.5. Let (ψ, η) be an arbitrary smooth solution with $\hat{\eta} := \max \eta < \gamma/4$. Then the inequality

$$
\psi(x,y) < \frac{\psi_x^2(x,y) + \psi_y^2(x,y)}{\gamma}
$$

holds true for all $(x, y) \in D_n$.

Proof. Let us consider the function

$$
f = \frac{1}{2}\psi_x^2 + \frac{1}{2}\psi_y^2 - \frac{\gamma}{2}\psi.
$$

A direct computation shows that

$$
\Delta f = 2\frac{f_x^2 + f_y^2}{\psi_x^2 + \psi_y^2}
$$

and then the minimum of f has to be at the boundary. Note that f is strictly positive everywhere at the bottom, while on the surface $y = \eta$ we have

$$
f = r - \eta - \frac{\gamma}{2} \ge 0.
$$

Here we used the main assumption on $\hat{\eta}$ and the inequality $r \geq \gamma$, which holds for any smooth solution with $r \geq R(s_c)$ (and hence for arbitrary smooth solutions; see [\[KKL15\]](#page-10-8)). Thus, we conclude $f \geq 0$ everywhere in the fluid by the maximum principle, which immediately leads the desired inequality. desired inequality.

Proposition 2.6. There is an absolute constant $C_3 > 0$ such that for any $\delta > 0$ there exists a constant $\gamma_2(\delta)$ depending only on δ with the following property. For any Stokes or solitary wave solution (ψ, η) with $\gamma > \gamma_2(\delta)$, $r \in (R(s_c), R(0))$ and satisfying the inequality

$$
\psi_y \geq \sqrt{2\gamma} - \frac{\delta}{\gamma}
$$

everywhere at the surface we have

$$
y_{\star} < C_3 \frac{\sqrt{\delta}}{\gamma^2}
$$

for any y_{\star} enjoying $\psi_{y}(0, y_{\star}) = 1/\gamma$.

Proof. Given $\delta > 0$ we choose $\gamma_2(\delta)$ to be larger than $\gamma_1(\delta)$ and such that

$$
C_2 \delta \sqrt{\frac{2}{\gamma}} \le \frac{\gamma}{4}
$$

whenever $\gamma > \gamma_2(\delta)$. Here C_2 is the constant from [\(2.7\)](#page-4-0). This allows to apply both Proposition [2.4](#page-3-0) and Proposition [2.5](#page-4-1) to every solution under given assumptions. Let $y_{\star} \in (0, \eta(0))$ be such that $\psi_y(0, y_\star) = 1/\gamma$. Note that by Proposition [2.4](#page-3-0) we have

$$
y \le \frac{C\delta}{\gamma} \psi_y(x, y)
$$

everywhere in the fluid with $C = C_1^{-1}$. We integrate this inequality for $x = 0$ and $y \in [0, y_\star]$, which gives

$$
y_\star^2 \leq \frac{2C\delta}{\gamma}\psi(0,y_\star) < \frac{2C\delta}{\gamma^4}
$$

Here for the last inequality we used Proposition [2.5.](#page-4-1) Now it is left to take the square root. \Box

3. Proof of Theorem [2.1](#page-1-0)

In order to prove the theorem we need to provide δ_{\star} and γ_{\star} . First we will choose δ_{\star} and then γ_{\star} depending on δ_{\star} . Let $\delta_1 > 0$ and γ_3 be some absolute constants such that

$$
r - \gamma - \frac{\sqrt{2\gamma} - \frac{\delta}{\gamma}}{\gamma} > \frac{\delta}{2\gamma^2}
$$
\n(3.1)

.

for all $\delta > \delta_1$, $\gamma > \gamma_3$ and $r \in (R(s_c), R(0))$. This is always possible since such r is subject to the identity

$$
r - \gamma = \sqrt{\frac{2}{\gamma}} + O(\gamma^{-2}).
$$

Next we choose δ_2 such that

$$
C_3\sqrt{\delta} < \frac{\delta}{2} \tag{3.2}
$$

for all $\delta > \delta_2$. Here C_3 is the constant from Proposition [2.6.](#page-5-0) Next, we choose any δ_{\star} such that $\delta_{\star} > \max(\delta_1, \delta_2, C_0^2)$, where C_0 is the constant in [\(2.5\)](#page-3-1). Now let $\gamma_4(\delta_{\star})$ be defined so that

$$
1 - \gamma \left(\sqrt{2\gamma} - \frac{\delta_\star}{\gamma}\right) < 0\tag{3.3}
$$

for all $\gamma > \gamma_4(\delta_\star)$. Finally, we choose any

$$
\gamma_{\star} > \max(\gamma_0, \gamma_1(\delta_{\star}), \gamma_2(\delta_{\star}), \gamma_3, \gamma_4(\delta_{\star})).
$$

We prove the claim of the theorem for chosen above δ_{\star} and γ_{\star} by a contradiction. Let (ψ, η) be a solution satisfying the assumptions and such that

$$
\min \psi_y(x, \eta(x)) = \psi_y(x_0, \eta(x_0)) = \sqrt{2\gamma} - \frac{\delta_\star}{\gamma}
$$

for some x_0 . Next we consider the auxiliary function

$$
g = \frac{1}{2}\psi_x^2 + \frac{1}{2}\psi_y^2 + y - \gamma\psi - \frac{1}{\gamma}\psi_y.
$$

A direct computation shows that

$$
\Delta g + \frac{a}{(\psi_y - \frac{1}{\gamma})^2 + \psi_x^2} g_x + \frac{b}{(\psi_y - \frac{1}{\gamma})^2 + \psi_x^2} g_y = 0,
$$

where a, b are some smooth functions. Thus, the maximum of g has to be achieved either at the upper, lower boundaries or at the points, where $\psi_x = 0$ and $\psi_y = 1/\gamma$. At the upper boundary we get

$$
\max g(x, \eta(x)) = g(x_0, \eta(x_0)) = r - \gamma - \frac{\sqrt{2\gamma} - \frac{\delta_\star}{\gamma}}{\gamma} > \frac{\delta_\star}{2\gamma^2}
$$
(3.4)

by [\(3.1\)](#page-5-1). Note that the maximum of q can not be attained at the lower boundary $y = 0$ since there

$$
g < \frac{1}{2}\psi_y^2 \le \frac{C_0^2}{2\gamma^2}
$$

and $\delta_{\star} > C_0^2$ by the choice. We aim to show that the maximum of g can only be attained at the upper boundary and so we need to consider the points where $\psi_x = 0$ and $\psi_y = 1/\gamma$. Note that $\psi_x(x, y) \neq 0$ for $x \in (-\Lambda, 0)$ and so there are two options $x = 0$ or $x = -\Lambda$ if we take into account the symmetry around $x = 0$. Assume that

$$
\max g = g(-\Lambda, y_{\dagger})
$$

for some $y_{\dagger} \in (0, \tilde{\eta})$. A direct computation gives

$$
g_{yy}(-\Lambda, y_{\dagger}) = -\psi_{xx}(-\Lambda, y_{\dagger})\psi_{yy}(-\Lambda, y_{\dagger}).
$$

Note however that $\psi_{xx}(-\Lambda, y_t) < 0$ and $\psi_{yy}(-\Lambda, y_t) > 0$ so that $g_{yy}(-\Lambda, y_t) > 0$ and no maximum can be there. The other alternative is

$$
\max g = g(0, y_\star).
$$

Here we have

$$
g(0, y_{\star}) < y_{\star} < C_3 \frac{\sqrt{\delta_{\star}}}{\gamma^2}
$$

by Proposition [2.6,](#page-5-0) which is applicable by the choice of δ_{\star} and γ_{\star} . But [\(3.2\)](#page-5-2) holds true for $\delta = \delta_{\star}$ and so we conclude

$$
g(0, y_{\star}) < C_3 \frac{\sqrt{\delta_{\star}}}{\gamma^2} < \frac{\delta_{\star}}{2\gamma^2} < g(x_0, \eta(x_0))
$$

by [\(3.4\)](#page-6-0). Hence the maximum of g is to be attained at the upper boundary. At $x = -\Lambda$ the function $\psi_y(x, \eta(x))$ has a local minimum, while for $x = 0$ we have

$$
g_y(0, \eta(0)) = -\eta'' \psi_y \frac{1 - \gamma \psi_y}{\gamma} < 0.
$$

Here we used [\(3.3\)](#page-5-3) which guarantees that $1 - \gamma \psi_y < 0$ along the upper boundary. Thus, the maximum of g is at some $(x_0, \eta(x_0))$ with $\eta'(x_0) \neq 0$. A direct computation shows that

$$
g_y(x_0, \eta(x_0)) = \frac{1 - \gamma \psi_y}{\gamma (1 + \eta'^2) \psi_y} < 0.
$$

This contradicts the Hopf lemma and completes the proof.

4. Improved bounds

Our aim is to obtain better bounds for the amplitude of a solution as well to improve our central inequality [\(2.1\)](#page-1-1). We will use a similar idea as in the proof of Proposition [2.4](#page-3-0) combined with the following lemma.

Lemma 4.1. For any $\lambda \geq \frac{\gamma}{2}$ $\frac{\gamma}{2}$ the function $\zeta(x) = \psi_y(x, \eta(x)) - \lambda \eta(x)$ attains its minimum at the crest, where $x = 0$.

Proof. A proof of this result is given in [\[KL23,](#page-10-10) Proposition 3.3] (see the second formula after (3.7) there), one just needs to choose $k = -\lambda$ and note that $\mathcal{L}f \geq 0$ when $k \leq \frac{\omega}{2}$ $\frac{\omega}{2}$ or, equivalently, $\lambda \geq \frac{\gamma}{2}$ 2 . The contract of \Box

Lemma 4.2. Assume that $\psi_y \ge \sqrt{2\gamma} - \frac{\delta}{\gamma} > 0$ everywhere at the surface and $\hat{\eta} \le A_0 \sqrt{\frac{2}{\gamma}}$, then

$$
\hat{\eta} \le \sqrt{\frac{2}{\gamma}} \cdot \frac{1}{1 - \frac{\delta}{\sqrt{2\gamma^3}}}.
$$

In particular this bound is independent of $A_0 > 0$.

Proof. We shall consider the harmonic function

$$
h = \psi_y - \lambda y, \quad \lambda = \frac{\sqrt{2\gamma} - \frac{\delta}{\gamma}}{A_0 \sqrt{\frac{2}{\gamma}}}.
$$

The choice of λ ensures that $h \geq 0$ everywhere at the surface, while $h = \psi_y \geq 0$ at the bottom. Thus, h is non-negative everywhere by the maximum principle. This implies

$$
y \le \frac{1}{\lambda} \psi_y
$$

everywhere below the crest. Integrating this inequality over $[0, \hat{\eta}]$, we find

$$
\frac{\hat{\eta}^2}{2} \le \frac{1}{\lambda},
$$

which is equivalent to

$$
\hat{\eta} \le \left(\frac{A_0}{1 - \frac{\delta}{\sqrt{2\gamma^3}}}\right)^{\frac{1}{2}} \sqrt{\frac{2}{\gamma}} = A_1 \sqrt{\frac{2}{\gamma}}.
$$

Repeating the same argument with A_0 replaced by A_1 we find

$$
\hat{\eta} \le A_2 \sqrt{\frac{2}{\gamma}}, \quad A_2 = \left(\frac{A_1}{1 - \frac{\delta}{\sqrt{2\gamma^3}}}\right)^{\frac{1}{2}} \sqrt{\frac{2}{\gamma}}.
$$

This way we obtain a sequence

$$
A_n = \left(\frac{A_{n-1}}{1 - \frac{\delta}{\sqrt{2\gamma^3}}}\right)^{\frac{1}{2}} \sqrt{\frac{2}{\gamma}} \to \frac{1}{1 - \frac{\delta}{\sqrt{2\gamma^3}}},
$$

which gives the desired inequality after passing to the limit. \Box

Lemma 4.3. Assume that $\psi_y \ge \sqrt{2\gamma}(1-\epsilon) > 0$ just at the crest, where $\hat{\eta} \le B_0 \sqrt{\frac{2}{\gamma}}$, then

$$
\hat{\eta} \le \sqrt{\frac{2}{\gamma}} \frac{1}{1 - \epsilon},
$$

provided $|\epsilon|, |1 - B_0| < \epsilon^*$ for some absolute constant $\epsilon^* > 0$.

Proof. The proof is similar to the argument in Lemma [4.2,](#page-7-0) but here we do not require the inequality for ψ_y to hold everywhere at the surface. To overcome this difficulty we will use Lemma [4.1](#page-6-1) to ensure that the function h has the right sign at the surface. More precisely, we define

$$
h = \psi_y - \lambda y, \quad \lambda = \frac{(1 - \epsilon)}{B_0} \gamma.
$$

If ϵ and $|1 - B_0|$ are small, then we have Lemma [4.1](#page-6-1) that forces h to have minimum along the surface exactly at the crest, where $h \geq 0$ by the choice of λ . As before, this leads to the new bound

$$
\hat{\eta} \le B_1 \sqrt{\frac{2}{\gamma}}, \quad B_1 = \left(\frac{B_0}{1-\epsilon}\right)^{1/2}.
$$

Now we consider the convergent sequence

$$
B_n = \left(\frac{B_{n-1}}{1-\epsilon}\right)^{1/2} \to \frac{1}{1-\epsilon}.
$$

To apply this argument for all n we need to ensure that

$$
\frac{(1-\epsilon)}{B_n} \ge \frac{1}{2}.
$$

This is true for the limiting value of B_n , provided ϵ is small enough. It remains to check a finite number of inequalities, which can done by choosing B_0 close enough to 1. This finishes the proof. \Box

Now we can improve the result of Theorem [2.1.](#page-1-0)

Proof of Theorem [1.1.](#page-0-1) Our proof is based on the previous lemmas. We can apply Lemma [4.2](#page-7-0) in view of Theorem [2.1.](#page-1-0) This gives the bound

$$
\hat{\eta} \le \sqrt{\frac{2}{\gamma}} + O(\gamma^{-2}).
$$

Using this inequality inside of the Bernoulli equation at the crest, we get

$$
\frac{1}{2}\psi_y^2 \ge r - \hat{\eta} \ge \gamma + \sqrt{\frac{2}{\gamma}} - O(\gamma^{-2}) - \sqrt{\frac{2}{\gamma}} - O(\gamma^{-2}) = \gamma - O(\gamma^{-2}).
$$

Thus, we conclude

$$
\psi_y \ge \sqrt{2\gamma} (1 - O(\gamma^{-3}))
$$

at the crest as stated in (ii). In order to obtain the bound for $\hat{\eta}$ we just need to apply Lemma [4.3](#page-7-1) for large enough γ . This proves the claim (ii) and also claim (iii), since

$$
\check{\eta} = \sqrt{\frac{2}{\gamma}} - O(\gamma^{-2}).
$$

For the first claim we need to use the monotonicity of the function

$$
\psi_y(x,\eta(x))-\gamma\eta(x)
$$

at the surface, which decays from the trough to the crest. In particular, we get

$$
\psi_y \ge \gamma \eta + \psi_y(0, \hat{\eta}) - \gamma \hat{\eta} \ge \gamma \check{\eta} - O(\gamma^{-2})
$$

as desired.

In a similar way one can also prove the following important result.

Theorem 4.4. There exist no unidirectional Stokes or solitary waves with $r = R(0)$, provided γ is large enough.

The claim of is also true for unidirectional Stokes and solitary waves with a bottom stagnation.

Proof. Our argument is the same for Stokes and solitary waves. Assume the statement is false and such solution exists. Our argument here is essentially based on Lemma [4.3](#page-7-1) from above. We aim to show that $\hat{\eta} \leq \sqrt{\frac{2}{\gamma}}$ and $\psi_y \geq \sqrt{2\gamma}$ at the crest. We start with the bound

$$
\hat{\eta} \le \sqrt{\frac{2}{\gamma}} (1 + \epsilon_0), \ \ \epsilon_0 = A_0 \gamma^{-3}
$$

 \Box

provided by (ii) in Theorem [1.1.](#page-0-1) Now we use the Bernoulli equation and get

$$
\frac{1}{2}\psi_y^2 = R(0) - \hat{\eta} = \gamma + \sqrt{\frac{2}{\gamma}} - \sqrt{\frac{2}{\gamma}}(1 + \epsilon_0) = \gamma - \epsilon_0 \sqrt{\frac{2}{\gamma}}.
$$

Hence, we find

$$
\psi_y \ge \sqrt{2\gamma} \sqrt{1 - \epsilon_0 \sqrt{\frac{2}{\gamma^3}}} \ge \sqrt{2\gamma} (1 - \epsilon'_1),
$$

provided

$$
\epsilon_1' = \epsilon_0 \sqrt{\frac{2}{\gamma^3}} = A_0 \sqrt{2} \gamma^{-\frac{9}{2}} \le 1.
$$
\n(4.1)

Which is true if γ large enough. Now we apply Lemma [4.3](#page-7-1) and obtain

$$
\hat{\eta} \le \sqrt{\frac{2}{\gamma}} \frac{1}{1 - \epsilon_1'} \le \sqrt{\frac{2}{\gamma}} (1 + \epsilon_1), \quad \epsilon_1 = 2\epsilon_1' = \epsilon_0 2\sqrt{\frac{2}{\gamma^3}}
$$

wherever $\epsilon'_1 < \frac{1}{2}$ $\frac{1}{2}$. Thus, we obtained a better bound for $\hat{\eta}$ using the Bernoulli equation, which does not produce an extra error, since $r = R(0)$. We can repeat this argument, which will lead to the sequence

$$
\epsilon_n = \epsilon_{n-1} 2 \sqrt{\frac{2}{\gamma^3}} = \epsilon_0 \left(\frac{2\sqrt{2}}{\gamma^3} \right)^n.
$$

Thus, if γ is large enough, we can satisfy all inequalities such as [\(4.1\)](#page-9-0) and the assumption of Lemma [4.3](#page-7-1) at each step. And after passing to the limit we conclude

$$
\hat{\eta} \le \sqrt{\frac{2}{\gamma}}, \quad \psi_y \ge \sqrt{2\gamma}
$$

to be true at the crest. Note that we always have $\hat{\eta} \geq d_+(r) = d(0) = \sqrt{\frac{2}{\gamma}}$ and so $\hat{\eta} = d(0)$. Now we apply the maximum principle to the function

$$
\psi(x, y) - U(y), \quad U(y) = \frac{\gamma}{2}y^2
$$

defined on the fluid domain $\{(x, y) : 0 \le y \le \eta(x)\}\$. This function is non-negative and is zero at the crest, where the Hopf lemma gives

$$
\psi_y - U_y < 0.
$$
\nsince $\psi_x > \sqrt{2\gamma}$ and $U_y = \sqrt{2}$

But this leads to a contradiction, since $\psi_y \geq \sqrt{2\gamma}$ and $U_y =$ $\sqrt{2\gamma}$ at the crest.

 \Box

5. Global bifurcation of solitary waves

Theorem 5.1 (Nonexistence of solitary waves with large Bernoulli constants). For each large enough γ there exists a constant $R^* \in (R(s_c), R(0))$ with the following property. For every smooth unidirectional solitary wave the corresponding Bernoulli constant is subject to the bound $R(s_c) < r < R^*$.

The claim of is also true for unidirectional Stokes and solitary waves with a bottom stagnation.

Proof. Let γ be large enough so that assumptions of Theorem [2.1](#page-1-0) and Theorem [4.4](#page-8-0) are fulfilled. Then we assume the claim is wrong and there exists a sequence of unidirectional solitary waves $(\psi^{(j)}, \eta^{(j)}, r^{(j)})$ with $r^{(j)} \to R(0)$ as $j \to \infty$. In view of Theorem [2.1](#page-1-0) we have ψ_y being uniformly separated from zero and the slopes are uniformly bounded. This guarantees that the norms of solutions are uniformly bounded by a constant depending only on γ ; see Proposition 3.2 from [\[KL23\]](#page-10-10) providing uniform regularity near the surface. The uniform regularity allows to find a convergent subsequence and we can assume that $\eta^{(j)} \to \eta^{(*)}$ in $C^2(I)$ over any compact interval I. The limiting solution $\eta^{(*)}$ is smooth and satisfies $\eta^{(*)}(crest) \geq d(0)$ and $r^{(*)} = R(0)$. Furthermore, it has to be monotone one each side of the crest. Note that it can not coincide with the subcritical stream solution $(U(y, 0), d(0))$ since it has a different flow force constant, corresponding to the supercritical stream (as a limit of supercritical solitary waves). It also

can not coincide with the supercritical stream since $\eta^{(*)}(crest) \geq d(0)$. Thus, it is a non-trivial solitary wave satisfying $\psi_y^{(*)} > 0$ everywhere in the fluid except, possibly, one point at the bottom right below the crest, where $\psi_y^{(*)} = 0$. However, this is in a contradiction with Theorem $4.4.$

Proof of Theorem [1.2.](#page-1-2) The claim of the theorem follows from [\[Whe13,](#page-10-6) Theorem 1.1] and the inequality $r < R^*$ provided by Theorem [5.1.](#page-9-1) This bound for the Bernoulli constant ensures that the result of our Theorem [2.1](#page-1-0) is applicable to solutions in $\mathcal C$ and so all solitary waves along the bifurcation curve are uniformly bounded in norm, that is $\sup_{\mathcal{C}} ||\eta||_{C^2} < +\infty$. Therefore, using the claim of [\[Whe13,](#page-10-6) Theorem 1.1] we can find a sequence of solutions $(\psi^{(j)}, \eta^{(j)}, r^{(j)}) \in C$ convergent to a smooth solitary wave (ψ^*, η^*, r^*) such that inf $\psi_y^* = 0$, where the infimum is taken over the whole fluid domain. But on the surface $\psi_y^* > 0$ by our Theorem [2.1,](#page-1-0) which only leaves a possibility of the bottom stagnation. However, the bottom stagnation can only happen right below the crest, which finishes the proof. Note that this solitary wave is a non-trivial in view of the Theorem [5.1.](#page-9-1)

6. Global bifurcations of Stokes waves

Proof of Theorem [1.3.](#page-1-3) For a large γ and given a subcritical stream solution $(U(y; s); d(s))$ with $0 < s < s_c$ we consider the global bifurcation curve C of fixed period Stokes waves starting from $(U(y; s); d(s))$. Such curves were constructed in [\[Var20,](#page-10-7) Theorem 18] under a more general setting allowing for critical layers and stagnation points. Then, as directly follows from [\[Var20,](#page-10-7) Theorem 18], there are several alternatives as one move along the bifurcation curve: (i) the norms of solutions become unbounded, (ii) the surface profiles have troughs approaching the bottom, or (iii) a stagnation point is being formed somewhere in the fluid. Note that whenever solutions are unidirectional the corresponding Bernoulli constant is bounded from above by $R(0)$ in view of Theorem [4.4,](#page-8-0) provided γ is large enough. Thus, the second option is not possible since $\check{\eta} \geq d(s_{+}(R(0)))$ (as we explained in the proof of Proposition [2.2\)](#page-2-2). The first option is naturally eliminated by Theorem [2.1,](#page-1-0) which ensures the norms are uniformly bounded. This leaves the third possibility, which, again by Theorem [2.1,](#page-1-0) implies the bottom stagnation. Thus, there exists a sequence $(\psi^{(j)}, \eta^{(j)}, r^{(j)})$ of Stokes waves with $R(s_c) < r^{(j)} < R(0)$ such that $\psi^{(j)}(0,0) \to 0$ as $j \to +\infty$. Now since the norms are uniformly bounded we can always find a converging subsequence. This finishes the proof.

REFERENCES

- [CS04] Adrian Constantin and Walter Strauss. Exact steady periodic water waves with vorticity. Comm. Pure Appl. Math., 57(4):481–527, 2004.
- [CSV21] Adrian Constantin, Walter Strauss, and Eugen Vărvărucă. Large-amplitude steady downstream water waves. Communications in Mathematical Physics, 387(1):237–266, August 2021.
- [KKL15] V. Kozlov, N. Kuznetsov, and E. Lokharu. On bounds and non-existence in the problem of steady waves with vorticity. Journal of Fluid Mechanics, 765, jan 2015.
- [KKL20] V. Kozlov, N. Kuznetsov, and E. Lokharu. Solitary waves on constant vorticity flows with an interior stagnation point. Journal of Fluid Mechanics, 904, October 2020.
- [KL23] Vladimir Kozlov and Evgeniy Lokharu. Global bifurcation and highest waves on water of finite depth. Archive for Rational Mechanics and Analysis, 247(5), September 2023.
- [KS08a] JOY KO and WALTER STRAUSS. Effect of vorticity on steady water waves. Journal of Fluid Mechanics, 608:197–215, July 2008.
- [KS08b] Joy Ko and Walter Strauss. Large-amplitude steady rotational water waves. European Journal of Mechanics - B/Fluids, 27(2):96–109, March 2008.
- [Lok21] Evgeniy Lokharu. On bounds for steady waves with negative vorticity. Journal of Mathematical Fluid Mechanics, 23(2), March 2021.
- [LWW23] Evgeniy Lokharu, Erik Wahlén, and Jörg Weber. On the amplitude of steady water waves with favorable constant vorticity. Journal of Mathematical Fluid Mechanics, 25(3), June 2023.
- [Var20] Kristoffer Varholm. Global bifurcation of waves with multiple critical layers. SIAM Journal on Mathematical Analysis, 52(5):5066–5089, January 2020.
- [Whe13] Miles H. Wheeler. Large-amplitude solitary water waves with vorticity. SIAM J. Math. Anal., 45(5):2937–2994, 2013.

Centre for Mathematical Sciences, Lund University, SE-223 62 Lund, Sweden