Stratifying on Treatment Status

Jinyong Hahn[∗] John Ham† Geert Ridder‡ Shuyang Sheng§

April 9, 2024

Abstract

We investigate the estimation of treatment effects from a sample that is stratified on the binary treatment status. In the case of unconfounded assignment where the potential outcomes are independent of the treatment given covariates, we show that standard estimators of the average treatment effect are inconsistent. In the case of an endogenous treatment and a binary instrument, we show that the IV estimator is inconsistent for the local average treatment effect. In both cases, we propose simple alternative estimators that are consistent in stratified samples, assuming that the fraction treated in the population is known or can be estimated.

Keywoods: treatment effects, stratification on treatment status, choice-based sampling, consistency, asymptotic efficiency.

JEL codes: C21, C83, C14.

1 Introduction

Sample surveys are usually not simple random samples where each member of the population has the same probability of being included in the sample. Often the sample design partitions the population in non-overlapping subpopulations or strata. In each stratum the subpopulation is sampled with equal probability, but the inclusion probability can differ between the strata.

The over- or under-representation of strata is often intentional. Stratified samples can result in more precise estimates of population features. It may also be desirable to overrepresent particular groups in the sample survey. In some surveys the cost of sampling is lower

[∗]Department of Economics, UCLA, Email: hahn@econ.ucla.edu.

[†]Department of Economics, NYU Abu Dhabi, Email: jch18@nyu.edu.

[‡]Department of Economics, USC, Email: ridder@usc.edu.

[§]Department of Economics, UCLA, Email: ssheng@econ.ucla.edu.

if the sample is stratified. An example is the choice of transportation mode where sampling of travelers by transportation mode is less costly than sampling of the general population [\(Domencich and McFadden](#page-16-0) [\(1975\)](#page-16-0)). Finally, the sample may consist of independent samples that are combined in an overall sample. In this case the fraction of the population in each subpopulation may be unknown [\(Ridder and Moffitt](#page-17-0) [\(2007\)](#page-17-0)).

Econometric models specify conditional relations in which endogenous variables are determined by exogenous variables. Stratification on endogenous variables biases parameter estimates in parametric econometric models, while stratification on exogenous variables usually does not result in biased estimates. Stratification on endogenous variables was first discussed in discrete choice models under the name choice-based sampling [\(Manski and Lerman](#page-17-1) [\(1977\)](#page-17-1), [Hausman and Wise](#page-17-2) [\(1981](#page-17-2))). In Section [2](#page-2-0) we show that stratification on a conditionally exogenous treatment status surprisingly results in a biased estimate of the Average Treatment Effect $(ATE).$

We study the estimation of treatment effects from a sample that is stratified on the binary treatment status. We consider the case of unconfounded assignment where the potential outcomes are independent of the treatment assignment given covariates X , and the case that the treatment assignment is endogenous, i.e., is correlated with the potential outcomes, but there is a binary instrument Z that is independent of the potential outcomes, but correlated with the treatment indicator D. We show that in both cases standard estimators of the treatment effect are inconsistent. We also propose simple alternative estimators that are consistent in stratified samples.

In the applied literature, there are many examples where the strata are chosen on the basis of treatment status. For example, [Ham and Khan](#page-16-1) [\(2023\)](#page-16-1) want to measure the effectiveness of a new teaching approach, JAAGO, in urban Bangladesh as compared to government and non-governmental schools. The new approach is implemented in only two schools in Dhaka, so they use the population of JAAGO students as the treatment group. Alternatively, [Azoulay, Zivin, and Wang](#page-16-2) [\(2010\)](#page-16-2) want to assess the impact of collaborator quality on researchers' output. To do this, they compare the output of researchers whose superstar collaborators died versus researchers whose superstar collaborators did not die. They work with non-experimental methods since there can be selection in who works with a superstar. Their data set consists of all researchers who had a superstar co-author die and a random sample of researchers with a superstar co-author who did not die; the size of the complete set of comparison researchers is much larger than their treatment group. Finally, there are a large number of papers that compared trainees earnings with suitable comparison individuals, either to evaluate

non-experimental estimators or to obtain an estimate of the impact of training. In such studies, the group of potential comparisons is much larger than the number of treatments. All of the above papers, and many other studies, face the issue of having a group of treatments but no control group. They create a comparison group from a different data set, and the number of potential comparison individuals is much larger than the number of treatments.

In the unconfounded case independent random samples of the outcome Y and the conditioning variables X are drawn for the treated and the controls. It should not be a surprise that from these samples we can estimate the conditional distributions of Y given X for the treated and the controls. From these conditional distributions we can recover the Conditional Average Treatment Effect (CATE) given X. As shown by [Heckman and Todd](#page-17-3) [\(2009\)](#page-17-3) we can even recover the CATE given the propensity score, because as they show there is a 1-1 relation between the population propensity score and the log-odds ratio estimated from the stratified sample. However when we average the CATE to obtain the ATE, averaging over the distribution of X or that of the propensity score in the stratified sample results in inconsistent estimates of the ATE. We show that there is a simple fix for this problem. For the ATE this fix requires that the fraction of the treated in the population is known or can be estimated. Knowledge of this fraction is not required for the average treatment effects on the treated (ATT).

In the case of an endogenous treatment and a binary instrument Z we show that the Wald ratio estimator is not a consistent estimator of the Local Average Treatment Effect (LATE) if the sample is stratified on the treatment status. We again propose a simple fix that again requires that the fraction treated in the population is known. If the stratification is on Z , then the Wald ratio for the stratified sample recovers the LATE.

In Appendix [A.5](#page-31-0) we also consider the efficiency of stratified sampling in the case of abundant control observations. We show that even if we have a small number of treated, we should use all available controls to estimate the ATT efficiently.

2 Conditionally Exogenous Case

We first discuss the (conditionally) exogenous case where the treatment assignment is unconfounded, i.e., the treatment D is independent of the potential outcomes (Y_0, Y_1) given X. We consider the situation where the sampling distribution is different from the population distribution due to stratification on the treatment status. We note that conditionally on X the treatment indicator is exogenous so that the conditional average treatment effect (CATE) $E[Y_1 - Y_0 | X = x]$ given $X = x$ is correctly identified by the difference of the conditional on

D and X sample averages for the treated and controls even if we stratify on D. Despite this we argue that conventional methods applied to the sample do not identify the average treatment effects (ATE). Curiously, we also note that conventional methods do identify the average treatment effects on the treated (ATT).

Let $Y \equiv DY_1 + (1 - D)Y_0$ denote the observed outcome. Let π^* denote the unconditional probability of sampling $D = 1$.¹ The stratified sample is such that $\pi^* \neq \pi$, where π denotes the population proportion of the treated. Let $q(x)$ denote the population density of X. The density of (Y, D, X) in the population is

$$
[h_1(y|x) P (D = 1|x)]^d [h_0(y|x) P (D = 0|x)]^{1-d} g(x), \qquad (1)
$$

where the h's denote the conditional densities of Y given X and D. Note that we can alternatively write [\(1\)](#page-3-0) as

$$
[h_1(y|x) g(x|D=1)\pi]^d [h_0(y|x) g(x|D=0) (1-\pi)]^{1-d}, \qquad (2)
$$

where we use

$$
g(x|D = 1) = g(x) \frac{P(D = 1|x)}{\pi},
$$

\n
$$
g(x|D = 0) = g(x) \frac{P(D = 0|x)}{1 - \pi},
$$
\n(3)

which follows from Bayes' theorem.

2.1 Relationship Between Sampling Objects and Population Counterparts

The joint density of (Y, D, X) in the stratified sample is

$$
[h_1(y|x) g (x|D=1) \pi^*]^d [h_0(y|x) g (x|D=0) (1-\pi^*)]^{1-d}.
$$

¹We can either draw each unit with probability π^* from the treated and with probability $1 - \pi^*$ from the controls or draw independent samples from the treated and controls with sample sizes such that the fraction treated in the combined sample is π^* . We adopt the former sampling scheme, but all results are the same for the second method.

We may rewrite the joint density in the stratified sample in a form similar to (1) (see [Heckman and Todd,](#page-17-3) [2009](#page-17-3), p.S232):

$$
[h_1(y|x) P^*(D=1|x)]^d [h_0(y|x) P^*(D=0|x)]^{1-d} g^*(x), \qquad (4)
$$

where

$$
g^*(x) \equiv g(x | D = 1) \pi^* + g(x | D = 0) (1 - \pi^*), \tag{5}
$$

$$
P^*(D = 1|x) \equiv \frac{g(x|D = 1)\pi^*}{g^*(x)}
$$

=
$$
\frac{g(x|D = 1)\pi^*}{g(x|D = 1)\pi^* + g(x|D = 0)(1 - \pi^*)}
$$

=
$$
\frac{P(D = 1|x)g(x)\frac{\pi^*}{\pi}}{P(D = 1|x)g(x)\frac{\pi^*}{\pi} + P(D = 0|x)g(x)\frac{1 - \pi^*}{1 - \pi}}
$$

=
$$
\frac{P(D = 1|x)\frac{\pi^*}{\pi}}{P(D = 1|x)\frac{\pi^*}{\pi} + P(D = 0|x)\frac{1 - \pi^*}{1 - \pi}},
$$
(6)

and

$$
P^*(D = 0 | x) \equiv 1 - P^*(D = 1 | x).
$$

Comparing (1) and (4) , it is straightforward to see that the conditional densities of Y given X and D are identical in the population and the stratified on D sample. This implies that

$$
E^*[Y|D=1,X] - E^*[Y|D=0,X] = E[Y|D=1,X] - E[Y|D=0,X],
$$
\n(7)

where E and E^* denote the expectations taken with respect to the population distribution (1) and the stratified on D sampling distribution (4) , respectively. This is the sense in which conditioning or matching on X "works". Conditional means of outcomes for the treated and controls identify $E[Y|D=1, X] - E[Y|D=0, X]$ in the population, and identify $E^*[Y|D=1, X] - E[X|D=0, X]$ $E^{\ast}[Y|D=0,X]$ in the stratified sample. Because of [\(7\)](#page-4-1), the differences of conditional means are equal in the population and the stratified sample. Despite the success of conditioning/matching, we will argue in the next section that conventional methods designed to identify the ATE in the population do not identify the ATE if they are applied in samples that are stratified on the treatment status. For this purpose, it is useful to understand the relationship between the population and sampling objects, which we present below.

Let $\pi(x) \equiv P(D=1|x)$ denote the propensity score from the true population, and let

 $\pi^*(x) \equiv P^*(D=1|x)$ denote the propensity score from the stratified sample. We will try to understand the relationship between some sampling objects and their population counterparts.

We first note that by [\(6\)](#page-4-2)

$$
\pi^*(x) = \frac{\pi(x) \frac{\pi^*}{\pi}}{\pi(x) \frac{\pi^*}{\pi} + (1 - \pi(x)) \frac{1 - \pi^*}{1 - \pi}},
$$
\n(8)

from which we obtain

$$
\pi(x) = \frac{\pi^*(x) \frac{\pi}{\pi^*}}{\pi^*(x) \frac{\pi}{\pi^*} + (1 - \pi^*(x)) \frac{1 - \pi}{1 - \pi^*}}.
$$
\n(9)

Because $\pi^*(x)$ and π^* are identified from the stratified/sampled distribution of (Y, D, X) , we can see that the true propensity score $\pi(x)$ can be identified *if and only* π is known to the researcher.

We can also see from (3) and (5) as well as (9) that

$$
g^*(x) = g(x) \frac{\pi^*}{\pi} \pi(x) + g(x) \frac{1 - \pi^*}{1 - \pi} (1 - \pi(x))
$$

= $g(x) \left[\frac{\pi^*}{\pi} \frac{\pi^*(x) \frac{\pi}{\pi^*}}{\pi^*(x) \frac{\pi}{\pi^*} + (1 - \pi^*(x)) \frac{1 - \pi}{1 - \pi^*}} + \frac{1 - \pi^*}{1 - \pi} \left(1 - \frac{\pi^*(x) \frac{\pi}{\pi^*}}{\pi^*(x) \frac{\pi}{\pi^*} + (1 - \pi^*(x)) \frac{1 - \pi}{1 - \pi^*}} \right) \right]$
= $g(x) \frac{1}{\pi^*(x) \frac{\pi}{\pi^*} + (1 - \pi^*(x)) \frac{1 - \pi}{1 - \pi^*}},$

so

$$
g(x) = \left(\pi^*(x)\frac{\pi}{\pi^*} + (1 - \pi^*(x))\frac{1 - \pi}{1 - \pi^*}\right)g^*(x).
$$
 (10)

Because $\pi^*(x)$, $g^*(x)$, and π^* are identified from the distribution of (Y, D, X) in the stratified sample, we can see that the population density $g(x)$ of X can be identified *if and only if* π is known to the researcher either exactly or by an estimate.

2.2 Matching/Conditioning on X

We now ask whether the success of matching/conditioning as discussed in the previous section translates into the success of conventional methods in identifying the Average Treatment Effect (ATE) in samples that are stratified on the treatment status. For this purpose, we ask whether identification of common treatment effect parameters is possible without knowledge of the population π or $\pi(x)$. Let's first note that if we have a random sample from the population distribution, we can identify the ATE by iterated expectations

$$
E[Y_1 - Y_0] = E\{E[Y|D = 1, X] - E[Y|D = 0, X]\}.
$$
\n(11)

We examine whether the stratified sampling counterpart of (11)

$$
E^*[Y_1 - Y_0] \equiv E^* \{ E^* [Y | D = 1, X] - E^* [Y | D = 0, X] \}
$$

identifies the ATE.

It is convenient to define

$$
\beta(x) \equiv E[Y|D = 1, X = x] - E[Y|D = 0, X = x],
$$

$$
\beta^*(x) \equiv E^*[Y|D = 1, X = x] - E^*[Y|D = 0, X = x].
$$

These are the CATE for the population and stratified sample, respectively. By [\(7\)](#page-4-1) the CATE are equal $\beta(\cdot) = \beta^*(\cdot)$, so that

$$
E[Y_1 - Y_0] = E[\beta(X)] = \int \beta(x) g(x) dx,
$$

$$
E^*[Y_1 - Y_0] = E^*[\beta^*(X)] = \int \beta(x) g^*(x) dx.
$$

Because by [\(10\)](#page-5-1) $g^*(x) \neq g(x)$ in general, we conclude that

$$
E^*[Y_1 - Y_0] = \int \beta(x) g^*(x) dx \neq \int \beta(x) g(x) dx = E[Y_1 - Y_0].
$$
 (12)

The conclusion that the naive estimator does not identify the ATE is intuitive and unsurprising. On the other hand, it is of importance to recognize that the source of the problem is not the failure of matching/conditioning, but is due to averaging the CATE over the wrong distribution of X, i.e., over the stratified sampling distribution g^* instead of g. Note that matching/conditioning "works" even without knowledge of π . If we know π , we can use [\(10\)](#page-5-1)

to identify the ATE using

$$
E[Y_1 - Y_0] = \int \beta(x) g(x) dx = \int \beta^*(x) g(x) dx
$$

=
$$
\int \beta^*(x) \left(\pi^*(x) \frac{\pi}{\pi^*} + (1 - \pi^*(x)) \frac{1 - \pi}{1 - \pi^*} \right) g^*(x) dx
$$

=
$$
E^* \left[\beta^*(X) \left(\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1 - \pi}{1 - \pi^*} \right) \right]
$$

=
$$
E^* \left[\left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) \beta^*(X) \right].
$$

We have a two-step semiparametric estimator characterization of the ATE, in terms of the sampling distribution:

$$
E^* [Y - \beta_1^*(x) | D = 1, X = x] = 0,
$$

\n
$$
E^* [Y - \beta_0^*(x) | D = 0, X = x] = 0,
$$

\n
$$
E^* \left[\left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) (\beta_1^*(X) - \beta_0^*(X)) - \beta \right] = 0,
$$
\n(13)

where β denotes the ATE, and β_d^* $d_d^*(x) = E^*(Y|X = x, D = d), d = 0, 1.2$

2.3 Matching/Conditioning on the Propensity Score

In this section, we argue that the ATE cannot be identified by the naive matching/conditioning on the propensity score. In order to simplify the notation, we will start our analysis with the identification of $E[Y_1]$. Because independence of the potential outcomes and D given X implies independence given the propensity score, it follows by iterated expectations that

$$
E[Y_1] = E[E[Y | D = 1, \pi(X)]]
$$

so a naive approach is to use $E^* [E^* [Y | D = 1, \pi^* (X)]]$. [Heckman and Todd](#page-17-3) [\(2009\)](#page-17-3) noted³ that

$$
\frac{\pi^*(x)}{1-\pi^*(x)} = \frac{\frac{\pi^*}{\pi}}{\frac{1-\pi^*}{1-\pi}} \frac{\pi(x)}{1-\pi(x)}
$$

²This representation assumes that π^* is known. If the π^* is unknown, we can add one more moment $E^*[D - \pi^*] = 0$ to the above system of equations. See [A](#page-18-0)ppendix A for the asymptotic distribution of the implied estimator.

³This can be derived from (6) .

i.e., the sample odds ratio is a monotonically increasing function of the population odds ratio, and argued that conditioning on the sample propensity score should be equivalent to conditioning on the population propensity score. This can be explained in a simpler way. From [\(8\)](#page-5-2) and [\(9\)](#page-5-0), we see that there is a 1-1 relation between $\pi(x)$ and $\pi^*(x)$. It follows that the sigmaalgebras generated by these two random variables are identical, and therefore, conditioning on $\pi(X)$ and conditioning on $\pi^*(X)$ are equivalent, so that

$$
\beta_1^* (\pi^* (X)) \equiv E^* [Y | D = 1, \pi^* (X)] = E^* [Y | D = 1, \pi (X)]. \tag{14}
$$

The same argument that led to [\(7\)](#page-4-1) implies that

$$
E^*[Y|D=1, \pi(X)] = E[Y|D=1, \pi(X)] \equiv \beta_1(\pi(X))
$$
\n(15)

so we can see that the sample conditional mean $E^*[Y|D=1,\pi^*(X)]$ is equal to the corresponding population conditional mean.

Unfortunately, the averaging problem remains. We see that

$$
E^*[F^*[Y|D = 1, \pi^*(X)]] = E^*[\beta_1(\pi(X))]
$$

= $\int \beta_1(\pi(x)) g^*(x) dx \neq \int \beta_1(\pi(x)) g(x) dx = E[Y_1]$

in general because by [\(10\)](#page-5-1) $g^*(x) \neq g(x)$. As before, this problem can be overcome if we know π . It is because

$$
E[Y_1] = E[E^*[Y|D = 1, \pi^*(X)]]
$$

= $\int \beta_1(x) \left(\pi^*(x) \frac{\pi}{\pi^*} + (1 - \pi^*(x)) \frac{1 - \pi}{1 - \pi^*} \right) g^*(x) dx$
= $E^* \left[\beta_1^*(X) \left(\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1 - \pi}{1 - \pi^*} \right) \right]$
= $E^* \left[\beta_1^*(\pi^*(X)) \left(\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1 - \pi}{1 - \pi^*} \right) \right].$

To conclude, we can see that matching/conditioning works only partially, because it does not solve the problem of averaging when estimating the ATE. This implies that we should be careful interpreting [Heckman and Todd](#page-17-3) [\(2009](#page-17-3), p.S231)'s observation that "matching and selection procedures can identify population treatment effects using misspecified estimates of propensity scores fit on choice-based samples" when we estimate the ATE. As we shall see their

statement is fully correct when the target is the ATT.

Similarly to [\(13\)](#page-7-0), we have a two-step semiparametric estimator for the ATE

$$
E^* [D - \pi^* (x) | X = x] = 0,
$$

\n
$$
E^* [Y - \beta_1^* (\pi^* (x)) | D = 1, \pi^* (X) = \pi^* (x)] = 0,
$$

\n
$$
E^* [Y - \beta_0^* (\pi^* (x)) | D = 0, \pi^* (X) = \pi^* (x)] = 0,
$$

\n
$$
E^* \left[\left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) \left(\beta_1^* (\pi^* (X)) - \beta_0^* (\pi^* (X)) \right) - \beta \right] = 0.
$$
 (16)

We discussed two methods of estimating the ATE if the sample is stratified on the treatment status, and proposed fixes to naive estimators. In Appendix [A.1](#page-18-1) and [A.2,](#page-22-0) we use the results in [Newey](#page-17-4) [\(1994](#page-17-4)), and show that the corrected estimators have the same asymptotic distribution. We also present a formula for the asymptotic variance.

2.4 Inverse Propensity Score Weighting

In this section, we consider yet another conventional approach to identify the ATE. We consider the strategy of weighting by the inverse propensity score (IPW), and argue that it does not identify the ATE if it were applied to a sample that is stratified by the treatment status. IPW identifies the ATE by

$$
E[Y_1 - Y_0] = E\left[\frac{DY}{\pi(X)} - \frac{(1 - D)Y}{1 - \pi(X)}\right],
$$

and the naive sample counterpart is

$$
E^* \left[\frac{DY}{\pi^* (X)} - \frac{(1 - D) Y}{1 - \pi^* (X)} \right].
$$

Noting that

$$
E^* \left[\frac{DY}{\pi^*(X)} - \frac{(1-D)Y}{1-\pi^*(X)} \right] = E^* \left[E^* \left[\frac{DY}{\pi^*(X)} - \frac{(1-D)Y}{1-\pi^*(X)} \right] X \right]
$$

= $E^* [E^* [Y | D = 1, X] - E^* [Y | D = 0, X]]$
= $E^* [\beta^*(X)],$

we can conclude using [\(7\)](#page-4-1) that

$$
E^* \left[\frac{DY}{\pi^* (X)} - \frac{(1 - D) Y}{1 - \pi^* (X)} \right] = E^* \left[\beta \left(X \right) \right].
$$

Therefore, the weighting is also an incomplete solution because it does not take care of the averaging. In the end, we want to use a weighted average version

$$
E^* \left[\left(\frac{DY}{\pi^* (X)} - \frac{(1 - D)Y}{1 - \pi^* (X)} \right) \left(\pi^* (X) \frac{\pi}{\pi^*} + (1 - \pi^* (X)) \frac{1 - \pi}{1 - \pi^*} \right) \right],
$$

which again requires knowledge of π .

2.5 Average Treatment Effects on the Treated

In the previous sections, we argued that matching/conditioning identifies the CATE $E[Y | D =$ $1, X]-E[Y|D=0, X]$, although no conventional method based on matching/conditioning succeeds in identifying the ATE. In this section, we consider the ATT, and show that conventional methods work. Under unconfoundedness we have, because $g(x|D = 1) = \pi(x)g(x)/\pi$,

$$
ATT = E[Y_1 - Y_0 | D = 1] = E[E[Y_1 - Y_0 | D = 1, X] | D = 1]
$$

=
$$
E[\beta(X) | D = 1] = \frac{1}{\pi} E[\beta(X) \pi(X)],
$$
 (17)

and its sample counterpart is

$$
ATT^* = \frac{1}{\pi^*} E^* [\beta^* (X) \pi^* (X)].
$$

Note that $\beta^* (\cdot) = \beta (\cdot)$ by [\(7\)](#page-4-1). Moreover, comparing equations [\(3\)](#page-3-1) and [\(6\)](#page-4-2) we have $\pi^* (x) g^* (x) / \pi^* =$ $\pi(x) g(x)/\pi$. Hence,

$$
ATT^* = \frac{1}{\pi^*} E^* [\beta(X) \pi^*(X)] = \frac{1}{\pi^*} \int \beta(X) \pi^*(x) g^*(x) dx
$$

=
$$
\frac{1}{\pi} \int \beta(x) \pi(x) g(x) dx = \frac{1}{\pi} E [\beta(X) \pi(X)] = ATT.
$$
 (18)

This implies that for the ATT, stratification on the treatment status does not make a difference.

We consider several identification strategies that can serve as basis for estimation of the ATT. First, note that because $E^*[Y - \beta_0^*]$ $\mathcal{O}_0^*(X)|D=1, X] = \beta^*(X),$

$$
E^* [\beta^* (X) \pi^* (X)] = E^* [E^* [Y - \beta_0^* (X)] D = 1, X] E^* [D|X]] = E^* [D(Y - \beta_0^* (X))],
$$

which implies that

$$
ATT^* = \frac{1}{\pi^*} E^* [D(Y - \beta_0^*(X))].
$$

This suggests an estimator of the ATT of the form

$$
\frac{1}{n^{-1}\sum_{i=1}^{n}D_{i}} \cdot \frac{1}{n}\sum_{i=1}^{n}D_{i}\left(Y_{i}-\widehat{\beta}_{0}\left(X_{i}\right)\right) = \frac{1}{n_{1}}\sum_{i=1}^{n}D_{i}\left(Y_{i}-\widehat{\beta}_{0}\left(X_{i}\right)\right),\tag{19}
$$

where $\widehat{\beta}_0(X_i)$ denotes a nonparametric estimator of $E^* [Y | X, D = 0]$, and we note that the estimator does not depend on the fraction treated in the population π . This estimator subtracts from the outcome for a treated unit a predicted outcome for the counterfactual. Although intuitive, we are not aware of any reference which establishes the asymptotic distribution of such an estimator. Note that the estimator would be consistent if the data are a random sample from the population. Therefore we can consider its asymptotic distribution as if the sampling distribution is the same as the population distribution. In Appendix [A.3,](#page-26-0) we use the results in [Newey](#page-17-4) [\(1994](#page-17-4)), and show that the estimator has the same asymptotic distribution as the semiparametrically efficient estimator developed in [Hahn](#page-16-3) [\(1998](#page-16-3)).

We investigate whether the ATT can be identified by propensity score conditioning/matching. Because $E[Y_1|D=1] = E[Y|D=1] = E^*[Y|D=1]$, we focus on $E[Y_0|D=1]$, which can be identified in the population by

$$
E[Y_0|D=1] = E[E[Y_0|D=1, \pi(X)]|D=1]
$$

=
$$
E[E[Y_0|D=0, \pi(X)]|D=1] = E[\beta_0(\pi(X))|D=1].
$$

The sample counterpart is

$$
E^* [Y_0 | D = 1] = E^* [E^* [Y_0 | D = 1, \pi^* (X)] | D = 1]
$$

=
$$
E^* [E^* [Y_0 | D = 0, \pi^* (X)] | D = 1] = E^* [\beta_0^* (\pi^* (X)) | D = 1].
$$

Because β_0^* $\chi_{0}^{*}(\pi^{*}(X)) = \beta_{0}(\pi(X))$ by [\(14\)](#page-8-0) and [\(15\)](#page-8-1) and $\pi^{*}(x) g^{*}(x) / \pi^{*} = \pi(x) g(x) / \pi$, we can see that

$$
E^* [\beta_0^* (\pi^* (X)) | D = 1] = \frac{1}{\pi^*} \int \beta_0^* (\pi^* (x)) \pi^* (x) g^* (x) dx
$$

=
$$
\frac{1}{\pi} \int \beta_0 (\pi (x)) \pi (x) g (x) dx
$$

=
$$
E [\beta_0 (\pi (X)) | D = 1].
$$

As before, we can identify the ATT by the propensity score conditioning/matching. The stratifi-

cation has no impact on either the matching or the averaging, the latter because the distribution of $X|D = 1$ is the same in the population and the sample. In Appendix [A.4,](#page-29-0) we consider the implied estimator of the above strategy, and show that the estimator has the same asymptotic distribution as the one discussed above.

Lastly, we consider the estimator based on weighting by the inverse propensity score. Because

$$
E[Y_0|D = 1] = E[E[Y_0|D = 1, X]|D = 1] = E[E[Y_0|X]|D = 1]
$$

$$
= E\left[E\left[\frac{(1 - D)Y}{1 - \pi(X)}\middle|X\right]\middle|D = 1\right] = \frac{1}{\pi}E\left[\frac{(1 - D)Y}{1 - \pi(X)}\pi(X)\right],
$$

the ATT in the population is

$$
E[Y|D = 1] - \frac{1}{\pi}E\left[\frac{(1 - D)Y}{1 - \pi(X)}\pi(X)\right].
$$

In the sample, we have

$$
\frac{1}{\pi^*} E^* \left[\frac{(1 - D)Y}{1 - \pi^*(X)} \pi^*(X) \right] = \frac{1}{\pi^*} E^* \left[E^*[Y_0 | X] \pi^*(X) \right] = \frac{1}{\pi^*} E^* \left[E^*[Y | D = 0, X] \pi^*(X) \right]
$$

$$
= \frac{1}{\pi^*} E^* \left[E[Y | D = 0, X] \pi^*(X) \right] = \frac{1}{\pi} E \left[E[Y | D = 0, X] \pi(X) \right]
$$

$$
= \frac{1}{\pi} E \left[E[Y_0 | X] \pi(X) \right] = E[Y_0 | D = 1].
$$

Again, the ATT can be estimated by the standard estimator based on weighting by the inverse propensity score. The stratification has no impact.

3 Endogenous Case

If the treatment assignment is unconfounded and hence conditionally exogenous, then the conditional distribution of Y given X and $D = d$ is identical in the population and the sample, where the latter is stratified on the treatment status. As a consequence, conditional treatment effects are the same in population and sample. All the complications in the previous section were due to the averaging of conditional treatment effects. In this section, we consider the case of an endogenous treatment and stratification on that endogenous treatment. We have an instrumental variable Z . The instrumental variable is binary and to focus on essentials there are no additional covariates.

The estimand is the Local Average Treatment Effect (LATE). [Angrist, Imbens, and Rubin](#page-16-4) [\(1996\)](#page-16-4) show that the LATE is identified in the population by the Wald ratio

$$
\frac{E[Y|Z=1] - E[Y|Z=0]}{E[D|Z=1] - E[D|Z=0]}.
$$

Alternatively the Wald ratio is the solution for β of the moment condition

$$
E^* \left\{ \begin{bmatrix} 1 \\ Z \end{bmatrix} (Y - \alpha - \beta D) \right\} = 0.
$$

We show that if the sample is stratified on the treatment status, then the sample Wald ratio is not equal to the population Wald ratio.

3.1 Relationship Between Sampling Objects and Population Counterparts

As in the exogenous case, we will first derive the relationship between sampling objects and population counterparts. Note that the conditional distribution of (Y, Z) given $D = d$ is the same in the population and in the sample that is stratified on D . This implies that

$$
E^*[YZ | D = d] = E [YZ | D = d]
$$

$$
E^*[Y (1 - Z) | D = d] = E [Y (1 - Z) | D = d],
$$

and

$$
P^* (Z = 1 | D = d) = P (Z = 1 | D = d)
$$

$$
P^* (Z = 0 | D = d) = P (Z = 0 | D = d)
$$

for $d = 0, 1$.

3.2 The Wald Ratio in the Population and Stratified Sample

The sample Wald ratio is

$$
E^* [Y | Z = 1] - E^* [Y | Z = 0]
$$

$$
E^* [D | Z = 1] - E^* [D | Z = 0].
$$

We can see that

$$
E^*[Y|Z=1] - E^*[Y|Z=0] = \frac{E^*[YZ]}{P^*(Z=1)} - \frac{E^*[Y(1-Z)]}{P^*(Z=0)}
$$

=
$$
\frac{E[YZ|D=1]\pi^* + E[YZ|D=0](1-\pi^*)}{P(Z=1|D=1)\pi^* + P(Z=1|D=0)(1-\pi^*)}
$$

=
$$
\frac{E[Y(1-Z)|D=1]\pi^* + E[Y(1-Z)|D=0](1-\pi^*)}{P(Z=0|D=1)\pi^* + P(Z=0|D=0)(1-\pi^*)}
$$

and

$$
E^*[D|Z=1] - E^*[D|Z=0] = P^*(D=1|Z=1) - P^*(D=1|Z=0)
$$

=
$$
\frac{P(Z=1|D=1)\pi^*}{P(Z=1|D=1)\pi^* + P(Z=1|D=0)(1-\pi^*)}
$$

$$
-\frac{P(Z=0|D=1)\pi^*}{P(Z=0|D=1)\pi^* + P(Z=0|D=0)(1-\pi^*)}.
$$

Therefore, we have

$$
\frac{E^* [Y | Z = 1] - E^* [Y | Z = 0]}{E^* [D | Z = 1] - E^* [D | Z = 0]} = \frac{A}{B},
$$

with

$$
A = \frac{E[YZ|D=1]\pi^* + E[YZ|D=0](1-\pi^*)}{P(Z=1|D=1)\pi^* + P(Z=1|D=0)(1-\pi^*)} - \frac{E[Y(1-Z)|D=1]\pi^* + E[Y(1-Z)|D=0](1-\pi^*)}{P(Z=0|D=1)\pi^* + P(Z=0|D=0)(1-\pi^*)}
$$

and

$$
B = \frac{P(Z=1|D=1)\pi^*}{P(Z=1|D=1)\pi^* + P(Z=1|D=0) (1-\pi^*)} - \frac{P(Z=0|D=1)\pi^*}{P(Z=0|D=1)\pi^* + P(Z=0|D=0) (1-\pi^*)}.
$$

This is different from the population Wald ratio, which is given by the same formula with π in place of π^* .

3.3 Correcting the Sample Wald Ratio

We now present a fix for the sample Wald ratio. The fix is based on the following lemma:

Lemma 1 *For any function* φ (*Y*, *Z*, *D*)*, we have*

$$
E\left[\varphi\left(Y,Z,D\right)\right] = E^* \left[\left(D \frac{\pi}{\pi^*} + (1-D) \frac{1-\pi}{1-\pi^*} \right) \varphi\left(Y,Z,D\right) \right].
$$

Proof. Because the conditional distribution of Y, Z, D is the same in the population and stratified sample we have for any function $h(Y, Z, D)$

$$
E[h(Y, Z, D)|D] = E^* [h(Y, Z, D)|D]
$$

By iterated expectations

$$
E^* \left[\left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) \varphi(Y, Z, D) \right]
$$

\n
$$
= E^* \left[\left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) \varphi(Y, Z, D) \middle| D = 1 \right] \pi^*
$$

\n
$$
+ E^* \left[\left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) \varphi(Y, Z, D) \middle| D = 0 \right] (1 - \pi^*)
$$

\n
$$
= E \left[\varphi(Y, Z, D) \middle| D = 1 \right] \frac{\pi}{\pi^*} \pi^* + E \left[\varphi(Y, Z, D) \middle| D = 0 \right] \frac{1 - \pi}{1 - \pi^*} (1 - \pi^*)
$$

\n
$$
= E \left[\varphi(Y, Z, D) \middle| D = 1 \right] \pi + E \left[\varphi(Y, Z, D) \middle| D = 0 \right] (1 - \pi)
$$

\n
$$
= E \left[\varphi(Y, Z, D) \right].
$$

П

Lemma [1](#page-15-0) implies that the population Wald ratio is the solution for β of the sample moment condition

$$
E^* \left\{ \left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) \begin{bmatrix} 1 \\ Z \end{bmatrix} (Y - \alpha - \beta D) \right\} = 0.
$$

This fix requires knowledge of π for implementation.⁴

Remark 1 *In the exogenous case without* X*, that is, under random assignment, stratification on* D *does not create any problem, because without* X *there is no need for averaging that creates the inconsistency in the conditionally exogenous case. In the endogenous case, the bias is present even without* X*. The intuition can be found in the literature on stratification on endogenous variables, see for instance [Hausman and Wise](#page-17-2) [\(1981\)](#page-17-2).*

 $4F₁$ and 2007 considers a nonparametric estimator of LATE that allows for covariates.

4 Summary

If the treatment assignment is unconfounded stratification on the treatment status does not bias conditional treatment effects. It does bias the average treatment effect that is identified by averaging the conditional treatment effect. We propose an easy fix that however requires that the fraction treated in the population is known or can be estimated. Stratification on the treatment status does not affect the estimation of the average treatment effect on the treated. If the treatment assignment is endogenous, but a valid instrument is available, then stratification on the treatment status biases the IV estimator of the local average treatment effect. If the fraction treated in the population is known or can be estimated, we propose a modified IV estimator that is a consistent for the population LATE.

References

- Angrist, J. D., G. W. Imbens, and D. B. Rubin (1996): "Identification of Causal Effects Using Instrumental Variables," *Journal of the American Statistical Association*, 91(434), 444–455.
- Azoulay, P., J. S. G. Zivin, and J. Wang (2010): "Superstar Extinction," *Quarterly Journal of Economics*, 125(2), 549–589.
- DOMENCICH, T., AND D. L. MCFADDEN (1975): *Urban Travel Demand: A Behavioral Analysis*. North-Holland Publishing.
- FRÖLICH, M. (2007): "Nonparametric IV estimation of local average treatment effects with covariates," *Journal of Econometrics*, 139(1), 35–75.
- Hahn, J. (1998): "On the Role of the Propensity Score in Efficient Semiparametric Estimation of Average Treatment Effects," *Econometrica*, 66(2), 315.
- HAHN, J., AND G. RIDDER (2013): "Asymptotic Variance of Semiparametric Estimators With Generated Regressors," *Econometrica*, 81(1), 315–340.
- Ham, J. C., and S. Khan (2023): "Can JAAGO, A New Education Organization, Address Bangladesh's Gender Differences in Achievement," Manuscript.
- Hausman, J. A., and D. A. Wise (1981): "Stratification on Endogenous Variables and Estimation: The Gary Income Maintenance Experiment," in *Structural Analysis of Discrete Data with Econometric Applications*, ed. by C. Manski, and D. McFadden. MIT Press.
- HECKMAN, J. J., AND P. E. TODD (2009) : "A note on adapting propensity score matching and selection models to choice based samples," *Econometrics Journal*, 12, S230–S234.
- Manski, C. F., and S. R. Lerman (1977): "The Estimation of Choice Probabilities from Choice Based Samples," *Econometrica*, 45(8), 1977.
- Newey, W. K. (1994): "The Asymptotic Variance of Semiparametric Estimators," *Econometrica*, 62(6), 1349–1382.
- RIDDER, G., AND R. MOFFITT (2007): "The Econometrics of Data Combination," in *Handbook of Econometrics*, vol. 6B, pp. 5469–5547. Elsevier.

Appendix

A Asymptotic Variance

A.1 ATE Estimator with Covariate Matching/Conditioning

We derive the asymptotic distribution of the ATE estimator that uses the moments [\(13\)](#page-7-0). The estimator for the ATE is

$$
\widehat{\beta} = \frac{1}{n} \sum_{i=1}^{n} \left(D_i \frac{\pi}{\pi^*} + (1 - D_i) \frac{1 - \pi}{1 - \pi^*} \right) \left(\widehat{\beta}_1 \left(X_i \right) - \widehat{\beta}_0 \left(X_i \right) \right).
$$

Standard arguments (e.g., [Newey](#page-17-4) [\(1994\)](#page-17-4)) can be used to show that the influence function of $\widehat{\beta}$ is equal to a main term

$$
\left(D\frac{\pi}{\pi^*} + (1-D)\frac{1-\pi}{1-\pi^*}\right)(\beta_1(X) - \beta_0(X)) - \beta
$$

plus the adjustment term for the estimation of $(\beta_1(X), \beta_0(X))$. Lemma [2](#page-19-0) below shows that this adjustment term is given by (21) , so the influence function is equal to

$$
\left(D\frac{\pi}{\pi^*} + (1 - D)\frac{1 - \pi}{1 - \pi^*}\right)(\beta_1(X) - \beta_0(X)) - \beta + \left(\pi^*(X)\frac{\pi}{\pi^*} + (1 - \pi^*(X))\frac{1 - \pi}{1 - \pi^*}\right)\left(\frac{D}{\pi^*(X)}(Y - \beta_1(X)) - \frac{1 - D}{1 - \pi^*(X)}(Y - \beta_0(X))\right).
$$
\n(20)

If π^* is estimated, we need to add

$$
E^* \left[\frac{\partial}{\partial \pi^*} \left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) (\beta_1(X) - \beta_0(X)) \right] (D - \pi^*)
$$

=
$$
E^* \left[\left(-D \frac{\pi}{(\pi^*)^2} + (1 - D) \frac{1 - \pi}{(1 - \pi^*)^2} \right) (\beta_1(X) - \beta_0(X)) \right] (D - \pi^*)
$$

=
$$
E^* \left[\left(-\pi^*(X) \frac{\pi}{(\pi^*)^2} + (1 - \pi^*(X)) \frac{1 - \pi}{(1 - \pi^*)^2} \right) (\beta_1(X) - \beta_0(X)) \right] (D - \pi^*)
$$

to (20) above.

Lemma 2 *The adjustment for the estimation of* $(\beta_1(X), \beta_0(X))$ *is equal to*

$$
\left(\pi^*(X)\frac{\pi}{\pi^*} + (1 - \pi^*(X))\frac{1 - \pi}{1 - \pi^*}\right)\left(\frac{D}{\pi^*(X)}(Y - \beta_1(X)) - \frac{1 - D}{1 - \pi^*(X)}(Y - \beta_0(X))\right). \tag{21}
$$

Proof. We derive [\(21\)](#page-19-1) following [Newey](#page-17-4) [\(1994](#page-17-4))'s argument. By [\(13\)](#page-7-0) we can write the ATE as a linear functional in the conditional means of Y given X for the treated and the controls

$$
\beta = \mathbb{E}^* \left[\left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) (h_1(X) - h_2(X)) \right]
$$

= $\mathbb{E}^* \left[\left(\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1 - \pi}{1 - \pi^*} \right) (h_1(X) - h_2(X)) \right]$
= $\mathbb{E}^* [w(X) (h_1(X) - h_2(X))].$

In this expression

$$
h_1(X) \equiv \mathbb{E}^*[Y | D = 1, X], \quad h_2(X) \equiv \mathbb{E}^*[Y | D = 0, X]
$$

and

$$
w(X) \equiv \pi^* (X) \frac{\pi}{\pi^*} + (1 - \pi^* (X)) \frac{1 - \pi}{1 - \pi^*}.
$$
 (22)

Therefore the ATE satisfies the moment equation

$$
\mathbb{E}^* [m(X, \beta, h_1, h_2)] = 0,
$$

where

$$
m(X, \beta, h_1, h_2) = w(X) (h_1(X) - h_2(X)) - \beta = D(X)' h(X) - \beta,
$$

with

$$
D(X) \equiv w(X) (1, -1)', \quad h(X) \equiv (h_1(X), h_2(X))'.
$$
 (23)

We use [Newey](#page-17-4) [\(1994,](#page-17-4) equation 4.1)'s notation $D(X)$ for the functional derivative which is not the treatment dummy D.

Following [Newey](#page-17-4) [\(1994\)](#page-17-4), define a path indexed by the scalar parameter θ for the distribution of (Y, D, X) with density $f^*(\cdot, \theta)$, where $f^*(\cdot, 0) = f^*(\cdot)$ is the density of the distribution of (Y, D, X) in the stratified sample. If \mathbb{E}_{θ}^* denotes an expectation with respect to the distribution with density $f^*(\cdot, \theta)$, then we define the corresponding paths for the projections $h_1(X, \theta) \equiv$ E ∗ $\mathcal{E}_{\theta}[Y|D = 1, X]$ and $h_2(X, \theta) \equiv \mathbb{E}_{\theta}^*$ $\partial_{\theta}[Y|D=0,X].$ The paths $h_1(X,\theta)$ and $h_2(X,\theta)$ are the minimizers of the objective functions

$$
\mathbb{E}_{\theta}^{*}\left[D\left(Y-\widetilde{h}_{1}\left(X,\theta\right)\right)^{2}\right] \quad \text{and} \quad \mathbb{E}_{\theta}^{*}\left[\left(1-D\right)\left(Y-\widetilde{h}_{2}\left(X,\theta\right)\right)^{2}\right],
$$

respectively, so the following orthogonality conditions hold for all functions $\widetilde{h}_1(X, \theta)$ and $\widetilde{h}_2(X,\theta)$:

$$
\mathbb{E}_{\theta}^{*}\left[D\left(Y-h_{1}\left(X,\theta\right)\right)\right)\widetilde{h}_{1}\left(X,\theta\right)\right]=0 \text{ and } \mathbb{E}_{\theta}^{*}\left[\left(1-D\right)\left(Y-h_{2}\left(X,\theta\right)\right)\widetilde{h}_{2}\left(X,\theta\right)\right]=0.
$$

We sum up the orthogonality conditions

$$
\mathbb{E}_{\theta}^{*}\left[D\left(Y-h_{1}\left(X,\theta\right)\right)\widetilde{h}_{1}\left(X,\theta\right)+\left(1-D\right)\left(Y-h_{2}\left(X,\theta\right)\right)\widetilde{h}_{2}\left(X,\theta\right)\right]=0
$$

for all functions $\left(\widetilde{h}_1(X,\theta),\widetilde{h}_2(X,\theta)\right)$. Choose $\left(\widetilde{h}_1(X,\theta),\widetilde{h}_2(X,\theta)\right) = \left(\frac{w(X)}{\pi^*(X,\theta)},-\frac{w(X)}{1-\pi^*(X,\theta)}\right)$, where $\pi^*(X, \theta) \equiv \mathbb{E}_\theta^*$ $_{\theta}^*[D|X]$. We obtain

$$
\mathbb{E}_{\theta}^* \left[w(X) \left(\frac{D}{\pi^*(X, \theta)} \left(Y - h_1(X, \theta) \right) - \frac{(1 - D)}{1 - \pi^*(X, \theta)} \left(Y - h_2(X, \theta) \right) \right) \right] = 0, \quad (24)
$$

which is equivalent to

$$
\mathbb{E}_{\theta}^{*}\left[w(X)\left(\frac{D}{\pi^{*}(X,\theta)}Y-\frac{1-D}{1-\pi^{*}(X,\theta)}Y\right)\right]
$$
\n
$$
=\mathbb{E}_{\theta}^{*}\left[w(X)\left(\frac{D}{\pi^{*}(X,\theta)}h_{1}(X,\theta)-\frac{1-D}{1-\pi^{*}(X,\theta)}h_{2}(X,\theta)\right)\right]
$$
\n
$$
=\mathbb{E}_{\theta}^{*}\left[w(X)\left(h_{1}(x,\theta)-h_{2}(x,\theta)\right)\right].
$$
\n(25)

As in [Newey](#page-17-4) [\(1994,](#page-17-4) equation 4.5) we compute the total derivative of [\(25\)](#page-20-0) (evaluating the derivatives at $\theta = 0$)

$$
\frac{\partial \mathbb{E}_{\theta}^{*}[w(X) (h_{1}(X,\theta) - h_{2}(X,\theta))] }{\partial \theta} = \frac{\partial \mathbb{E}_{\theta}^{*}[w(X) (h_{1}(X) - h_{2}(X))]}{\partial \theta} + \frac{\partial \mathbb{E}^{*}[w(X) (h_{1}(X,\theta) - h_{2}(X,\theta))]}{\partial \theta}, \qquad (26)
$$

where we used the fact that the projections at $\theta = 0$ are equal to $h_1(X)$ and $h_2(X)$. Therefore

combining (26) with (25) , we obtain

$$
\frac{\partial \mathbb{E}^* \left[D(X)' h(X, \theta) \right]}{\partial \theta} = \frac{\partial \mathbb{E}^* \left[w(X) \left(h_1(\theta) - h_2(X, \theta) \right) \right]}{\partial \theta} \n= \frac{\partial \mathbb{E}_{\theta}^* \left[w(X) \left(h_1(X, \theta) - h_2(X, \theta) \right) \right]}{\partial \theta} - \frac{\partial \mathbb{E}_{\theta}^* \left[w(X) \left(h_1(X) - h_2(X) \right) \right]}{\partial \theta} \n= \frac{\partial}{\partial \theta} \left(\mathbb{E}_{\theta}^* \left[w(X) \left(\frac{D}{\pi^*(X, \theta)} Y - \frac{1 - D}{1 - \pi^*(X, \theta)} Y \right) \right] \right) \n- \frac{\partial}{\partial \theta} \left(\mathbb{E}_{\theta}^* \left[w(X) \left(\frac{D}{\pi^*(X, \theta)} h_1(X) - \frac{1 - D}{1 - \pi^*(X, \theta)} h_2(X) \right) \right] \right) \n= \frac{\partial}{\partial \theta} \left(\mathbb{E}_{\theta}^* \left[w(X) \left(\frac{D}{\pi^*(X, \theta)} (Y - h_1(X)) - \frac{1 - D}{1 - \pi^*(X, \theta)} (Y - h_2(X)) \right) \right] \right).
$$

By taking the total derivative of the right hand side at $\theta = 0$ and noting that the second term after the first equality is 0, we derive

$$
\frac{\partial \mathbb{E}^* \left[D \left(X \right)' h(X, \theta) \right]}{\partial \theta} = \mathbb{E}^* \left[w(X) \left(\frac{D}{\pi^* \left(X \right)} \left(Y - h_1 \left(X \right) \right) - \frac{1 - D}{1 - \pi^* \left(X \right)} \left(Y - h_2 \left(X \right) \right) \right) S \left(Y, D, X \right) \right] + \mathbb{E}^* \left[w(X) \left(- \frac{\pi^* \left(X \right) D}{\pi^* \left(X \right)^2} \left(Y - h_1 \left(X \right) \right) - \frac{\pi^* \left(X \right) \left(1 - D \right)}{\left(1 - \pi^* \left(X \right) \right)^2} \left(Y - h_2 \left(X \right) \right) \right) \right] - \mathbb{E}^* \left[w(X) \left(\frac{D}{\pi^* \left(X \right)} \left(Y - h_1 \left(X \right) \right) - \frac{1 - D}{1 - \pi^* \left(X \right)} \left(Y - h_2 \left(X \right) \right) \right) S \left(Y, D, X \right) \right].
$$

In this expression $S(\cdot) \equiv \partial \ln f^*(\cdot, \theta) / \partial \theta |_{\theta=0}$ is the score of the density, and $\pi^*(X) \equiv \partial \pi^*(X, \theta) / \partial \theta |_{\theta=0}$. Therefore by [Newey](#page-17-4) [\(1994,](#page-17-4) Proposition 4) the adjustment to the influence function is

$$
w(X)\left(\frac{D}{\pi^*(X)}(Y - h_1(X)) - \frac{1 - D}{1 - \pi^*(X)}(Y - h_2(X))\right).
$$

The statement of the lemma follows by (22) .

When π^* is known, we can derive the asymptotic variance of the ATE estimator based on the influence function in [\(20\)](#page-18-2). Recall that $\beta(X) = \beta_1(X) - \beta_0(X)$. Rewrite (20) as

$$
\left(\left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) \beta(X) - \beta \right) + \frac{\pi^* (X) \frac{\pi}{\pi^*} + (1 - \pi^* (X)) \frac{1 - \pi}{1 - \pi^*}}{\pi^* (X)} - \frac{\pi^* (X) \frac{\pi}{\pi^*} + (1 - \pi^* (X)) \frac{1 - \pi}{1 - \pi^*}}{1 - \pi^* (X)} (1 - D) \varepsilon_0,
$$
\n(27)

where $\varepsilon_d \equiv Y - \beta_d(X)$. Under unconfoundedness, the asymptotic variance of $\sqrt{n}(\hat{\beta} - \beta)$ is given by (note that the three terms of [\(27\)](#page-21-0) are uncorrelated)

$$
E^* \left[\left(\left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) \beta(X) - \beta \right)^2 \right] + E^* \left[\frac{\left(\pi^* (X) \frac{\pi}{\pi^*} + (1 - \pi^* (X)) \frac{1 - \pi}{1 - \pi^*} \right)^2 \sigma_1^2(X)}{\pi^* (X)} \right] + E^* \left[\frac{\left(\pi^* (X) \frac{\pi}{\pi^*} + (1 - \pi^* (X)) \frac{1 - \pi}{1 - \pi^*} \right)^2 \sigma_0^2(X)}{1 - \pi^* (X)} \right],
$$

where $\sigma_1^2(X) \equiv E^*[\varepsilon_1^2|X]$ and $\sigma_0^2(X) \equiv E^*[\varepsilon_0^2|X]$.

A.2 ATE Estimator with Propensity Score Matching/Conditioning

We can estimate the ATE alternatively based on the moments (16) that use propensity score matching/conditioning. The estimator takes the form

$$
\frac{1}{n}\sum_{i=1}^n \left(D_i \frac{\pi}{\pi^*} + (1-D_i) \frac{1-\pi}{1-\pi^*}\right)\left(\widehat{\beta}_1\left(\widehat{\pi}^*(X_i)\right) - \widehat{\beta}_0\left(\widehat{\pi}^*(X_i)\right)\right).
$$

The influence function of this estimator has as its main term

$$
\left(D\frac{\pi}{\pi^*} + (1-D)\frac{1-\pi}{1-\pi^*}\right)\left(\beta_1(\pi^*(X)) - \beta_0(\pi^*(X))\right) - \beta,\tag{28}
$$

to which we add the adjustment for the estimation of $(\beta_1(\cdot), \beta_0(\cdot))$ and the adjustment for the estimation of $\pi^*(X)$. The first adjustment can be derived following Lemma [2](#page-19-0) as

$$
\left(\pi^*(X)\frac{\pi}{\pi^*} + (1 - \pi^*(X))\frac{1 - \pi}{1 - \pi^*}\right)\left(\frac{D}{\pi^*(X)}(Y - \beta_1(\pi^*(X))) - \frac{1 - D}{1 - \pi^*(X)}(Y - \beta_0(\pi^*(X)))\right).
$$
\n(29)

Lemma [3](#page-23-0) below shows that the second adjustment is given by [\(30\)](#page-23-1).

Lemma 3 *The adjustment for the estimation of* $\pi^*(X)$ *is equal to*

$$
\left(D\frac{\pi}{\pi^*} + (1-D)\frac{1-\pi}{1-\pi^*}\right)(\beta(X) - \beta(\pi^*(X)))
$$
\n
$$
-\frac{\pi^*(X)\frac{\pi}{\pi^*} + (1-\pi^*(X))\frac{1-\pi}{1-\pi^*}}{\pi^*(X)}D(\beta_1(X) - \beta_1(\pi^*(X)))
$$
\n
$$
+\frac{\pi^*(X)\frac{\pi}{\pi^*} + (1-\pi^*(X))\frac{1-\pi}{1-\pi^*}}{1-\pi^*(X)}(1-D)(\beta_0(X) - \beta_0(\pi^*(X))).
$$
\n(30)

Proof. We derive [\(30\)](#page-23-1) following [Hahn and Ridder](#page-16-6) [\(2013,](#page-16-6) Theorem 7). Adopting [Hahn and Ridder](#page-16-6) [\(2013\)](#page-16-6)'s notation, we let

$$
h(D, \mu_1, \mu_2) \equiv \left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) \left(\mu_1 \left(\pi^* (X) \right) - \mu_2 \left(\pi^* (X) \right) \right),
$$

where

$$
\mu_1(v) \equiv E^*[Y|D=1, \pi^*(X)=v] = \beta_1(v), \quad \mu_2(v) \equiv E^*[Y|D=0, \pi^*(X)=v] = \beta_0(v).
$$

Also $E^*[D|\pi^*(X)] = \pi^*(X),$

$$
\frac{\partial h(D, \mu_1, \mu_2)}{\partial \mu_1} = D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*}, \n\frac{\partial h(D, \mu_1, \mu_2)}{\partial \mu_2} = -\left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right), \n\kappa_1(v) = E^* \left[\frac{\partial h(D, \mu_1, \mu_2)}{\partial \mu_1} \middle| \pi^*(X) = v \right] = v \frac{\pi}{\pi^*} + (1 - v) \frac{1 - \pi}{1 - \pi^*}, \n\kappa_2(v) \equiv E^* \left[\frac{\partial h(D, \mu_1, \mu_2)}{\partial \mu_2} \middle| \pi^*(X) = v \right] = -v \frac{\pi}{\pi^*} - (1 - v) \frac{1 - \pi}{1 - \pi^*}, \n\frac{\partial \kappa_1(v)}{\partial v} = \frac{\pi}{\pi^*} - \frac{1 - \pi}{1 - \pi^*}, \n\frac{\partial \kappa_2(v)}{\partial v} = -\frac{\pi}{\pi^*} + \frac{1 - \pi}{1 - \pi^*},
$$

and finally

$$
\pi_1(v) \equiv v,
$$

$$
\pi_2(v) \equiv 1 - v.
$$

Further, we denote

$$
\mu_1(X) \equiv E^*[Y|D = 1, X] = \beta_1(X), \quad \mu_2(X) \equiv E^*[Y|D = 0, X] = \beta_0(X).
$$

Using [Hahn and Ridder](#page-16-6) [\(2013,](#page-16-6) Theorem 7), we can see that the contribution of estimating $\pi^*(X)$ is the sum of the three terms in [\(31\)](#page-24-0), [\(32\)](#page-24-1), and [\(33\)](#page-24-2) times $D - \pi^*(X)$:

$$
E^* \left[\left(\frac{\partial h(D, \mu_1, \mu_2)}{\partial \mu_1} - \kappa_1 (\pi^* (X)) \right) \frac{\partial \mu_1 (\pi^* (X))}{\partial v} \middle| X \right] + E^* \left[\left(\frac{\partial h(D, \mu_1, \mu_2)}{\partial \mu_2} - \kappa_2 (\pi^* (X)) \right) \frac{\partial \mu_2 (\pi^* (X))}{\partial v} \middle| X \right] = 0, \tag{31}
$$

$$
E^* \left[\left(\mu_1 \left(X \right) - \mu_1 \left(\pi^* \left(X \right) \right) \right) \frac{\partial \kappa_1 \left(\pi^* \left(X \right) \right)}{\partial v} \middle| X \right] + E^* \left[\left(\mu_2 \left(X \right) - \mu_2 \left(\pi^* \left(X \right) \right) \right) \frac{\partial \kappa_2 \left(\pi^* \left(X \right) \right)}{\partial v} \middle| X \right]
$$
\n
$$
= \left(\beta_1 \left(X \right) - \beta_1 \left(\pi^* \left(X \right) \right) \right) \left(\frac{\pi}{\pi^*} - \frac{1 - \pi}{1 - \pi^*} \right) - \left(\beta_0 \left(X \right) - \beta_0 \left(\pi^* \left(X \right) \right) \right) \left(\frac{\pi}{\pi^*} - \frac{1 - \pi}{1 - \pi^*} \right)
$$
\n
$$
= \left(\beta \left(X \right) - \beta \left(\pi^* \left(X \right) \right) \right) \left(\frac{\pi}{\pi^*} - \frac{1 - \pi}{1 - \pi^*} \right), \tag{32}
$$

and

$$
-E^* \left[\frac{1}{\pi_1(\pi^*(X))} \left(\mu_1(X) - \mu_1(\pi^*(X)) \right) \kappa_1(\pi^*(X)) \frac{\partial \pi_1(\pi^*(X))}{\partial v} \right] X \right]
$$

\n
$$
-E^* \left[\frac{1}{\pi_2(\pi^*(X))} \left(\mu_2(X) - \mu_2(\pi^*(X)) \right) \kappa_2(\pi^*(X)) \frac{\partial \pi_2(\pi^*(X))}{\partial v} \right] X \right]
$$

\n
$$
= -\frac{1}{\pi^*(X)} \left(\beta_1(X) - \beta_1(\pi^*(X)) \right) \left(\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1 - \pi}{1 - \pi^*} \right)
$$

\n
$$
- \frac{1}{1 - \pi^*(X)} \left(\beta_0(X) - \beta_0(\pi^*(X)) \right) \left(-\pi^*(X) \frac{\pi}{\pi^*} - (1 - \pi^*(X)) \frac{1 - \pi}{1 - \pi^*} \right) (-1)
$$

\n
$$
= -\frac{1}{\pi^*(X)} \left(\beta_1(X) - \beta_1(\pi^*(X)) \right) \left(\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1 - \pi}{1 - \pi^*} \right)
$$

\n
$$
- \frac{1}{1 - \pi^*(X)} \left(\beta_0(X) - \beta_0(\pi^*(X)) \right) \left(\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1 - \pi}{1 - \pi^*} \right).
$$
 (33)

Combining them, we obtain

$$
(\beta(X) - \beta(\pi^*(X))) \left(\frac{\pi}{\pi^*} - \frac{1-\pi}{1-\pi^*}\right) (D - \pi^*(X))
$$

$$
- \frac{1}{\pi^*(X)} (\beta_1(X) - \beta_1(\pi^*(X))) \left(\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1-\pi}{1-\pi^*}\right) (D - \pi^*(X))
$$

$$
- \frac{1}{1-\pi^*(X)} (\beta_0(X) - \beta_0(\pi^*(X))) \left(\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1-\pi}{1-\pi^*}\right) (D - \pi^*(X))
$$
 (34)

as the adjustment for the estimation of $\pi^*(X)$. We can rewrite [\(34\)](#page-25-0) as

$$
(\beta(X) - \beta(\pi^*(X))) \left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} - \left(\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1 - \pi}{1 - \pi^*} \right) \right)
$$

\n
$$
- (\beta_1(X) - \beta_1(\pi^*(X))) \left(\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1 - \pi}{1 - \pi^*} \right) \left(\frac{D}{\pi^*(X)} - 1 \right)
$$

\n
$$
- (\beta_0(X) - \beta_0(\pi^*(X))) \left(\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1 - \pi}{1 - \pi^*} \right) \left(1 - \frac{1 - D}{1 - \pi^*(X)} \right)
$$

\n
$$
= \left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) (\beta(X) - \beta(\pi^*(X)))
$$

\n
$$
- \frac{\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1 - \pi}{1 - \pi^*}}{\pi^*(X)} D (\beta_1(X) - \beta_1(\pi^*(X)))
$$

\n
$$
+ \frac{\pi^*(X) \frac{\pi}{\pi^*} + (1 - \pi^*(X)) \frac{1 - \pi}{1 - \pi^*}}{1 - \pi^*(X)} (1 - D) (\beta_0(X) - \beta_0(\pi^*(X))),
$$

from which we obtain the conclusion. \blacksquare

Adding up [\(28\)](#page-22-1) and [\(29\)](#page-22-2), we can see that the influence function of the infeasible estimator based on known $\pi^*(X)$ is equal to

$$
\left(\left(D \frac{\pi}{\pi^*} + (1 - D) \frac{1 - \pi}{1 - \pi^*} \right) \beta \left(\pi^* (X) \right) - \beta \right) + \frac{\pi^* (X) \frac{\pi}{\pi^*} + (1 - \pi^* (X)) \frac{1 - \pi}{1 - \pi^*}}{\pi^* (X)} - \frac{\pi^* (X) \frac{\pi}{\pi^*} + (1 - \pi^* (X)) \frac{1 - \pi}{1 - \pi^*}}{1 - \pi^* (X)} (1 - D) \tilde{\varepsilon}_0,
$$
\n(35)

where $\tilde{\varepsilon}_d \equiv Y - \beta_d (\pi^*(X))$. Because $\tilde{\varepsilon}_d = \varepsilon_d + \beta_d (X) - \beta_d (\pi^*(X))$, the sum of [\(35\)](#page-25-1) and [\(30\)](#page-23-1)

is equal to

$$
\left(\left(D\frac{\pi}{\pi^*} + (1-D)\frac{1-\pi}{1-\pi^*}\right)\beta(X) - \beta\right) + \frac{\pi^*(X)\frac{\pi}{\pi^*} + (1-\pi^*(X))\frac{1-\pi}{1-\pi^*}}{\pi^*(X)}D\varepsilon_1 - \frac{\pi^*(X)\frac{\pi}{\pi^*} + (1-\pi^*(X))\frac{1-\pi}{1-\pi^*}}{1-\pi^*(X)}(1-D)\varepsilon_0.
$$

This is identical to the influence function [\(27\)](#page-21-0) of the ATE estimator for the case that we match/condition on X.

A.3 ATT Estimator with Covariate Matching/Conditioning

Now we derive the asymptotic variance of the intuitive estimator of the ATT in [\(19\)](#page-11-0)

$$
\widehat{\gamma} = \frac{1}{n_1} \sum_{i=1}^{n} D_i \left(Y_i - \widehat{\beta}_0 \left(X_i \right) \right) = \frac{\frac{1}{n} \sum_{i=1}^{n} D_i \left(Y_i - \widehat{\beta}_0 \left(X_i \right) \right)}{\frac{1}{n} \sum_{i=1}^{n} D_i},\tag{36}
$$

where $\widehat{\beta}_0(X_i)$ is a nonparametric regression of Y on X in the untreated subsample (i.e., $D = 0$). Lemma [4](#page-26-1) derives the adjustment for the estimation of $\beta_0(X)$ in the numerator of [\(36\)](#page-26-2).

Lemma 4 *The adjustment for the estimation of* $\beta_0(X)$ *in the numerator* $\frac{1}{n} \sum_{i=1}^n D_i(X_i - \widehat{\beta}_0(X_i))$ *is equal to*

$$
-\frac{\pi^{*}(X)}{1-\pi^{*}(X)}(1-D)(Y-\beta_{0}(X)).
$$

Proof. We follow the proof of Lemma [2,](#page-19-0) using similar notation. We write the numerator as

$$
\frac{1}{n}\sum_{i=1}^{n}D_{i}\left(Y_{i}-\widehat{\beta}_{0}\left(X_{i}\right)\right)=\frac{1}{n}\sum_{i=1}^{n}D_{i}Y_{i}-\frac{1}{n}\sum_{i=1}^{n}D_{i}\widehat{\beta}_{0}\left(X_{i}\right),\tag{37}
$$

and focus on $\frac{1}{n} \sum_{i=1}^{n} D_i \widehat{\beta}_0 (X_i)$, which estimates

$$
\beta^* \equiv \mathbb{E}^* \left[\pi^* \left(X \right) h_2 \left(X \right) \right],
$$

where

$$
h_2(X) \equiv \mathbb{E}^* [Y | D = 0, X] = \beta_0(X).
$$

So we consider the moment equation

$$
\mathbb{E}^*[m(X,\beta^*,h_2)] = 0,
$$

where

$$
m(X, \beta^*, h_2) = \pi^*(X) h_2(X) - \beta^*.
$$

In [Newey](#page-17-4) [\(1994](#page-17-4), equation 4.1)'s notation, we therefore want to consider \mathbb{E}^* [D (X) h_2 (X)] with $D(X) \equiv \pi^*(X)$, and $D(X) h_2(X)$ is obviously linear in h_2 . Recall that the following orthogonality condition holds for all functions $\widetilde{h}_2(X, \theta)$:

$$
\mathbb{E}_{\theta}^* \left[\left(1 - D \right) \left(Y - h_2 \left(X, \theta \right) \right) \widetilde{h}_2 \left(X, \theta \right) \right] = 0.
$$

Choose $\widetilde{h}_2(X, \theta) = \frac{\pi^*(X)}{1 - \pi^*(X, \theta)}$, where $\pi^*(X, \theta) \equiv \mathbb{E}_{\theta}^*$ $_{\theta}^*[D|X]$, so that

$$
0 = \mathbb{E}_{\theta}^{*} \left[\frac{\pi^{*}(X)(1-D)}{1-\pi^{*}(X,\theta)} (Y - h_{2}(X,\theta)) \right]
$$

or

$$
\mathbb{E}_{\theta}^* \left[\frac{\pi^*(X)(1-D)}{1-\pi^*(X,\theta)} Y \right] = \mathbb{E}_{\theta}^* \left[\frac{\pi^*(X)(1-D)}{1-\pi^*(X,\theta)} h_2(X,\theta) \right] = \mathbb{E}_{\theta}^* \left[\pi^*(X) h_2(X,\theta) \right].
$$

The last expression is useful to compute the derivative as in [Newey](#page-17-4) [\(1994](#page-17-4), equation 4.5). Taking the total derivative (evaluating the derivatives at $\theta = 0$)

$$
\frac{\partial \mathbb{E}_{\theta}^{*} \left[\pi^{*}\left(X\right) h_{2}\left(X,\theta\right) \right]}{\partial \theta} = \frac{\partial \mathbb{E}_{\theta}^{*} \left[\pi^{*}\left(X\right) h_{2}\left(X\right) \right]}{\partial \theta} + \frac{\partial \mathbb{E}^{*} \left[\pi^{*}\left(X\right) h_{2}\left(X,\theta\right) \right]}{\partial \theta},
$$

we obtain

$$
\frac{\partial \mathbb{E}^* \left[\pi^* (X) h_2(X, \theta) \right]}{\partial \theta} = \frac{\partial \mathbb{E}_{\theta}^* \left[\pi^* (X) h_2(X, \theta) \right]}{\partial \theta} - \frac{\partial \mathbb{E}_{\theta}^* \left[\pi^*(X) h_2(X) \right]}{\partial \theta} \n= \frac{\partial}{\partial \theta} \mathbb{E}_{\theta}^* \left[\frac{\pi^*(X) (1 - D)}{1 - \pi^*(X, \theta)} Y \right] - \frac{\partial}{\partial \theta} \mathbb{E}_{\theta}^* \left[\frac{\pi^*(X) (1 - D)}{1 - \pi^*(X, \theta)} h_2(X) \right] \n= \frac{\partial}{\partial \theta} \mathbb{E}_{\theta}^* \left[\frac{\pi^*(X) (1 - D)}{1 - \pi^*(X, \theta)} (Y - h_2(X)) \right] \n= \mathbb{E}^* \left[\frac{\pi^*(X) (1 - D)}{1 - \pi^*(X)} (Y - h_2(X)) S(Y, D, X) \right] \n+ \mathbb{E}^* \left[\pi^*(X) (1 - D) \left(-\frac{\pi^*(X)}{(1 - \pi^*(X))^2} (Y - h_2(X)) \right) \right] \n= \mathbb{E}^* \left[\frac{\pi^*(X) (1 - D)}{1 - \pi^*(X)} (Y - h_2(X)) S(Y, D, X) \right],
$$

so the adjustment for $\frac{1}{n} \sum_{i=1}^{n} D_i \widehat{\beta}_0 (X_i)$ is

$$
\frac{\pi^*(X)(1-D)}{1-\pi^*(X)}\left(Y-h_2(X)\right)=\frac{\pi^*(X)(1-D)}{1-\pi^*(X)}\left(Y-\beta_0(X)\right),
$$

from which we obtain the conclusion. $\quad \blacksquare$

By Lemma [4,](#page-26-1) the numerator $\frac{1}{n} \sum_{i=1}^{n} D_i \left(Y_i - \widehat{\beta}_0 (X_i) \right)$ has the influence function

$$
D(Y - \beta_0(X)) - E^* [\pi^*(X) \beta(X)] - \frac{\pi^*(X)}{1 - \pi^*(X)} (1 - D) (Y - \beta_0(X))
$$

=
$$
D(\beta_1(X) + \varepsilon_1 - \beta_0(X)) - E^* [\pi^*(X) \beta(X)] - \frac{\pi^*(X)}{1 - \pi^*(X)} (1 - D) \varepsilon_0
$$

=
$$
D\beta(X) - E^* [\pi^*(X) \beta(X)] + D\varepsilon_1 - \frac{\pi^*(X)}{1 - \pi^*(X)} (1 - D) \varepsilon_0,
$$
 (38)

where $\varepsilon_d = Y - \beta_d(X)$. In other words,

$$
\frac{1}{n}\sum_{i=1}^{n} \left(D_i \left(Y_i - \widehat{\beta}_0 \left(X_i \right) \right) - E^* \left[\pi^* \left(X \right) \beta \left(X \right) \right] \right)
$$
\n
$$
= \frac{1}{n}\sum_{i=1}^{n} \left(D_i \beta \left(X_i \right) - E^* \left[\pi^* \left(X \right) \beta \left(X \right) \right] + D_i \varepsilon_{1i} - \frac{\pi \left(X_i \right)}{1 - \pi \left(X_i \right)} \left(1 - D_i \right) \varepsilon_{0i} \right) + o_p \left(\frac{1}{\sqrt{n}} \right).
$$

By the delta method, we obtain

$$
\sqrt{n}\left(\frac{1}{n_{1}}\sum_{i=1}^{n}D_{i}\left(Y_{i}-\widehat{\beta}_{0}\left(X_{i}\right)\right)-\frac{E^{*}\left[\pi^{*}\left(X\right)\beta\left(X\right)\right]}{\pi^{*}}\right) \n=\sqrt{n}\left(\frac{\frac{1}{n}\sum_{i=1}^{n}D_{i}\left(Y_{i}-\widehat{\beta}_{0}\left(X_{i}\right)\right)}{\frac{1}{n}\sum_{i=1}^{n}D_{i}}-\frac{E^{*}\left[\pi^{*}\left(X\right)\beta\left(X\right)\right]}{\pi^{*}}\right) \n=\sqrt{n}\left(\frac{\frac{1}{n}\sum_{i=1}^{n}\left(D_{i}\beta\left(X_{i}\right)-E^{*}\left[\pi^{*}\left(X\right)\beta\left(X\right)\right]+D_{i}\varepsilon_{1i}-\frac{\pi^{*}\left(X_{i}\right)}{1-\pi^{*}\left(X_{i}\right)}\left(1-D_{i}\right)\varepsilon_{0i}\right)}{\pi^{*}}\right) \n-\frac{E^{*}\left[\pi^{*}\left(X\right)\beta\left(X\right)\right]}{\left(\pi^{*}\right)^{2}}\frac{1}{\sqrt{n}}\sum_{i=1}^{n}\left(D_{i}-\pi^{*}\right)+o_{p}\left(1\right) \n=\frac{1}{\sqrt{n}}\sum_{i=1}^{n}\left(\frac{D_{i}\beta\left(X_{i}\right)}{\pi^{*}}-\gamma-\frac{\gamma}{\pi^{*}}\left(D_{i}-\pi^{*}\right)+\frac{D_{i}\varepsilon_{1i}-\frac{\pi^{*}\left(X_{i}\right)}{1-\pi^{*}\left(X_{i}\right)}\left(1-D_{i}\right)\varepsilon_{0i}}{\pi^{*}}\right)+o_{p}\left(1\right) \n=\frac{1}{\sqrt{n}}\sum_{i=1}^{n}\left(\frac{D_{i}\left(\beta\left(X_{i}\right)-\gamma\right)}{\pi^{*}}+\frac{D_{i}\varepsilon_{1i}-\frac{\pi^{*}\left(X_{i}\right)}{1-\pi^{*}\left(X_{i}\right)}\left(1-D_{i}\right)\varepsilon_{0i}}{\pi^{*}}\right)+o_{p}\left(1\right),
$$

where

$$
\gamma \equiv E\left[\frac{\pi(X)}{\pi} \beta(X)\right] = E^* \left[\frac{\pi^*(X)}{\pi^*} \beta(X)\right]
$$

is the ATT. This implies that the asymptotic variance of $\sqrt{n}(\hat{\gamma} - \gamma)$ is equal to

$$
E^* \left[\frac{\pi^* \left(X\right) \left(\beta \left(X\right) - \gamma\right)^2}{(\pi^*)^2} + \frac{\pi^* \left(X\right) \sigma_1^2 \left(X\right)}{(\pi^*)^2} + \frac{\pi^* \left(X\right)^2 \sigma_0^2 \left(X\right)}{(\pi^*)^2 \left(1 - \pi^* \left(X\right)\right)} \right],\tag{39}
$$

which is equal to the asymptotic variance bound based on the stratified sample. See [Hahn](#page-16-3) [\(1998\)](#page-16-3). So, the intuitive estimator is efficient. Note that [\(39\)](#page-29-1) is different from the asymptotic variance based on the unstratified population. While the ATT γ can be consistently estimated by [\(36\)](#page-26-2), its asymptotic variance depends on the stratification on the treatment status.

A.4 ATT Estimator with Propensity Score Matching/Conditioning

We now consider the intuitive estimator of the ATT that uses propensity score matching/conditioning

$$
\frac{1}{n_1} \sum_{i=1}^n D_i \left(Y_i - \widehat{\beta}_0 \left(\widehat{\pi} \left(X_i \right) \right) \right) = \frac{\frac{1}{n} \sum_{i=1}^n D_i \left(Y_i - \widehat{\beta}_0 \left(\widehat{\pi} \left(X_i \right) \right) \right)}{\frac{1}{n} \sum_{i=1}^n D_i}.
$$
\n(40)

Lemma 5 *The adjustment for the estimation of* $\pi^*(X)$ *in the numerator* $\frac{1}{n} \sum_{i=1}^n D_i (Y_i - \widehat{\beta}_0 (\widehat{\pi}(X_i)))$ *is equal to*

$$
-\frac{1}{1-\pi^*(X)}\left(\beta_0(X) - \beta_0(\pi^*(X))\right)(D - \pi^*(X)).\tag{41}
$$

Proof. We derive [\(41\)](#page-30-0) following [Hahn and Ridder](#page-16-6) [\(2013,](#page-16-6) Theorem 7).⁵ We adopt Hahn and Ridder [\(2013\)](#page-16-6)'s notation similarly as in Lemma [3,](#page-23-0) except that we let

$$
h(D, Y, \mu_2) \equiv D(Y - \mu_2(\pi^*(X)))
$$

and

$$
\frac{\partial h(D, Y, \mu_2)}{\partial \mu_2} = -D,
$$

\n
$$
\kappa_2(v) \equiv E^* \left[\frac{\partial h(D, Y, \mu_2)}{\partial \mu_2} \middle| \pi^*(X) = v \right] = -v,
$$

\n
$$
\frac{\partial \kappa_2(v)}{\partial v} = -1.
$$

Applying [Hahn and Ridder](#page-16-6) [\(2013,](#page-16-6) Theorem 7), we can see that the adjustment for the estimation of $\pi^*(X)$ is the sum of the three terms in [\(42\)](#page-30-1), [\(43\)](#page-30-2), and [\(44\)](#page-30-3) times $D - \pi^*(X)$:

$$
E^* \left[\left(\frac{\partial h(D, Y, \mu_2)}{\partial \mu_2} - \kappa_2 \left(\pi^*(X) \right) \right) \frac{\partial \mu_2(\pi^*(X))}{\partial v} \middle| X \right] = 0, \tag{42}
$$

$$
E^* \left[\left(\mu_2(X) - \mu_2(\pi^*(X)) \right) \frac{\partial \kappa_2(\pi^*(X))}{\partial v} \middle| X \right] = -(\beta_0(X) - \beta_0(\pi^*(X))), \tag{43}
$$

and

$$
- E^* \left[\frac{1}{1 - \pi^*(X)} \left(\mu_2(X) - \mu_2(\pi^*(X)) \right) \kappa_2(\pi^*(X)) \frac{\partial \pi_2(\pi^*(X))}{\partial v} \right| X \right]
$$

=
$$
- \frac{1}{1 - \pi^*(X)} (\beta_0(X) - \beta_0(\pi^*(X))) \pi^*(X).
$$
 (44)

Combining them, we obtain the adjustment in (41) .

Similarly to [\(38\)](#page-28-0), we can show that the infeasible estimator $\frac{1}{n} \sum_{i=1}^{n} D_i \left(Y_i - \widehat{\beta}_0 \left(\pi^*(X_i) \right) \right)$

⁵[Hahn and Ridder](#page-16-6) [\(2013](#page-16-6), p.333) considered an imputation version, not this version of the estimated propensity score matching for the ATT.

with known $\pi^*(X)$ has the influence function

$$
D(Y - \beta_0(\pi^*(X))) - E^*[\pi^*(X)\beta(\pi^*(X))] - \frac{\pi^*(X)}{1 - \pi^*(X)}(1 - D)(Y - \beta_0(\pi^*(X)))
$$

= $D\beta(\pi^*(X)) - E^*[\pi^*(X)\beta(\pi^*(X))] + D\tilde{\varepsilon}_1 - \frac{\pi^*(X)}{1 - \pi^*(X)}(1 - D)\tilde{\varepsilon}_0,$ (45)

where $\tilde{\varepsilon}_d = Y - \beta_d (\pi^*(X))$. Rewrite the adjustment in [\(41\)](#page-30-0) as

$$
- \left(D - \frac{\pi^*(X)(1 - D)}{1 - \pi^*(X)} \right) (\beta_0(X) - \beta_0(\pi^*(X)))
$$

=
$$
D (\beta(X) - \beta(\pi^*(X))) - D (\beta_1(X) - \beta_1(\pi^*(X)))
$$

+
$$
\frac{\pi^*(X)}{1 - \pi^*(X)} (1 - D) (\beta_0(X) - \beta_0(\pi^*(X))).
$$
 (46)

Because $\tilde{\varepsilon}_d = \varepsilon_d + \beta_d(X) - \beta_d(\pi^*(X))$, summing up [\(45\)](#page-31-1) and [\(46\)](#page-31-2) we obtain the overall influence function of the numerator in (40) as

$$
D\beta(X) - E^*[\pi^*(X)\beta(\pi^*(X))] + D\varepsilon_1 - \frac{\pi^*(X)}{1 - \pi^*(X)}(1 - D)\varepsilon_0,
$$

which is identical to the influence function (38) of the "numerator" in the covariate matching case. Therefore, the ATT estimators that condition on the propensity score or on the covariates have the same asymptotic variance given by (39) . So there is no efficiency gain from using the propensity score to estimate the ATE or ATT in stratified samples. In the random sampling case, [Hahn and Ridder](#page-16-6) [\(2013\)](#page-16-6) come to the same conclusion.

A.5 Choosing the Number of Controls

In applications observations of treated units are often costlier than observations on control units. If the number of treated units is fixed at n_1 which is relatively small but large enough to use an asymptotic approximation, how many controls should we use? In particular should we use all available controls or only a subset? If the fraction treated in the stratified sample is π^* , then the number of controls is $n_0 = \frac{1-\pi^*}{\pi^*} n_1$. The following proposition shows that for the estimator of the ATT in the stratified sample the variance is monotonically increasing in π^* .

Proposition 1 *Under unconfoundedness, the asymptotic variance of* $\sqrt{n_1}(\hat{\gamma} - \gamma)$ *is a monotonically increasing function of* π^* .

Proof. Recall that the asymptotic variance of our ATT estimator $\sqrt{n}(\hat{\gamma} - \gamma)$ is equal to

$$
E^* \left[\frac{\pi^* (X) \left(\beta (X) - \gamma \right)^2}{(\pi^*)^2} + \frac{\pi^* (X) \sigma_1^2 (X)}{(\pi^*)^2} + \frac{\pi^* (X)^2 \sigma_0^2 (X)}{(\pi^*)^2 (1 - \pi^* (X))} \right].
$$

Recalling that

$$
\pi^*(x) = \frac{\pi(x) \frac{\pi^*}{\pi}}{\pi(x) \frac{\pi^*}{\pi} + (1 - \pi(x)) \frac{1 - \pi^*}{1 - \pi}},
$$

$$
g^*(x) = g(x) \left(\pi(x) \frac{\pi^*}{\pi} + (1 - \pi(x)) \frac{1 - \pi^*}{1 - \pi}\right),
$$

we obtain that

$$
E^* \left[\frac{\pi^* (X) \left(\beta (X) - \gamma \right)^2}{(\pi^*)^2} \right]
$$

=
$$
E \left[\frac{\pi (X) \frac{\pi^*}{\pi}}{\pi (X) \frac{\pi^*}{\pi} + (1 - \pi (X)) \frac{1 - \pi^*}{1 - \pi}} \frac{\left(\beta (X) - \gamma \right)^2}{(\pi^*)^2} \left(\pi (X) \frac{\pi^*}{\pi} + (1 - \pi (X)) \frac{1 - \pi^*}{1 - \pi} \right) \right]
$$

=
$$
E \left[\frac{\pi (X)}{\pi \pi^*} \left(\beta (X) - \gamma \right)^2 \right],
$$

$$
E^* \left[\frac{\pi^* (X) \sigma_1^2 (X)}{(\pi^*)^2} \right]
$$

=
$$
E \left[\frac{\pi (X) \frac{\pi^*}{\pi}}{\pi (X) \frac{\pi^*}{\pi} + (1 - \pi (X)) \frac{1 - \pi^*}{1 - \pi}} \frac{\sigma_1^2 (X)}{(\pi^*)^2} \left(\pi (X) \frac{\pi^*}{\pi} + (1 - \pi (X)) \frac{1 - \pi^*}{1 - \pi} \right) \right]
$$

=
$$
E \left[\frac{\pi (X)}{\pi \pi^*} \sigma_1^2 (X) \right],
$$

and

$$
E^* \left[\frac{\pi^* (X)^2 \sigma_0^2(X)}{(\pi^*)^2 (1 - \pi^* (X))} \right]
$$

=
$$
E \left[\frac{\sigma_0^2 (X)}{(\pi^*)^2} \frac{\left(\frac{\pi (X) \frac{\pi^*}{\pi}}{(\pi (X) \frac{\pi^*}{\pi} + (1 - \pi (X))) \frac{1 - \pi^*}{1 - \pi}}\right)^2}{1 - \frac{\pi (X) \frac{\pi^*}{\pi}}{(\pi (X) \frac{\pi^*}{\pi} + (1 - \pi (X))) \frac{1 - \pi^*}{1 - \pi}} \left(\pi (X) \frac{\pi^*}{\pi} + (1 - \pi (X)) \frac{1 - \pi^*}{1 - \pi}\right) \right]
$$

=
$$
E \left[\frac{\pi (X)^2 \sigma_0^2(X)}{\pi^2 (1 - \pi (X))} \frac{1 - \pi^*}{1 - \pi^*} \right].
$$

It follows that the asymptotic variance of $\sqrt{n_1} (\hat{\gamma} - \gamma)$, which should be equal to the asymptotic variance of $\sqrt{n} (\hat{\gamma} - \gamma)$ times π^* , is equal to

$$
E\left[\frac{\pi(X)}{\pi}(\beta(X) - \gamma)^2 + \frac{\pi(X)}{\pi}\sigma_1^2(X) + \frac{\pi(X)^2(1-\pi)\sigma_0^2(X)}{\pi^2(1-\pi(X))}\frac{\pi^*}{1-\pi^*}\right].
$$

 \blacksquare

We conclude that even if we have a small number of treated we should use all available controls to estimate the ATT. There are two caveats: we were unable to prove the proposition for the ATE, and if π^* is close to 0 the asymptotic approximation may not be appropriate.