

# High-dimensional bootstrap and asymptotic expansion

Yuta Koike <sup>\*†</sup>

July 30, 2024

## Abstract

The recent seminal work of Chernozhukov, Chetverikov and Kato has shown that bootstrap approximation for the maximum of a sum of independent random vectors is justified even when the dimension is much larger than the sample size. In this context, numerical experiments suggest that third-moment match bootstrap approximations would outperform normal approximation even without studentization, but the existing theoretical results cannot explain this phenomenon. In this paper, we first show that Edgeworth expansion, if justified, can give an explanation for this phenomenon. Second, we obtain valid Edgeworth expansions in the high-dimensional setting when the random vectors have Stein kernels. Finally, we prove the second-order accuracy of a double wild bootstrap method in this setting. As a byproduct, we find an interesting blessing of dimensionality phenomenon: The single third-moment match wild bootstrap is already second-order accurate in high-dimensions if the covariance matrix has identical diagonal entries and bounded eigenvalues.

*Keywords:* Cornish–Fisher expansion; coverage probability; double bootstrap; Edgeworth expansion; second-order accuracy; Stein kernel.

## 1 Introduction

Let  $X_1, \dots, X_n$  be independent centered random vectors in  $\mathbb{R}^d$  with finite variance. Set

$$S_n := \frac{1}{\sqrt{n}} \sum_{i=1}^n X_i.$$

The aim of this paper is to investigate the accuracy of bootstrap approximation for the maximum type statistics

$$T_n := \max_{1 \leq j \leq d} S_{n,j} \quad \text{and} \quad \|S_n\|_\infty := \max_{1 \leq j \leq d} |S_{n,j}|,$$

when both  $n$  and  $d$  tend to infinity. The seminal work of Chernozhukov, Chetverikov & Kato [17] has established Gaussian type approximations for these statistics under very mild assumptions when the dimension  $d$  is possibly much larger than the sample size  $n$ . To be precise, let  $Z$  be a centered Gaussian vector in  $\mathbb{R}^d$  with the same covariance matrix as  $S_n$ , say  $\Sigma$ . Gaussian analogs of  $T_n$  and  $\|S_n\|_\infty$  are respectively given by

$$Z^\vee := \max_{1 \leq j \leq d} Z_j \quad \text{and} \quad \|Z\|_\infty := \max_{1 \leq j \leq d} |Z_j|.$$

---

<sup>\*</sup>Graduate School of Mathematical Sciences, University of Tokyo

<sup>†</sup>CREST, Japan Science and Technology Agency

Under mild moment assumptions, Chernozhukov, Chetverikov & Kato [17] have shown that

$$\sup_{t \in \mathbb{R}} |P(T_n \leq t) - P(Z^\vee \leq t)| = O\left(\left(\frac{\log^a(dn)}{n}\right)^b\right) \quad (1.1)$$

holds with  $a = 7$  and  $b = 1/8$ . An analogous result also holds for  $\|S_n\|_\infty$ . This result implies that given a significance level  $\alpha \in (0, 1)$ , the probability  $P(T_n \geq c_{1-\alpha}^G)$  is approximately equal to  $\alpha$  as long as  $\log d = o(n^{1/7})$ , where  $c_{1-\alpha}^G$  is the  $(1 - \alpha)$ -quantile of  $Z^\vee$ . Therefore, we can use  $c_{1-\alpha}^G$  as a critical value to construct asymptotically  $(1 - \alpha)$ -level simultaneous confidence intervals or  $\alpha$ -level tests for a high-dimensional vector of parameters; see [4, 22] for details. In practice,  $c_{1-\alpha}^G$  is not computable because  $\Sigma$  is generally unknown, so we need to replace it by an estimate. In [17], this is implemented by the Gaussian wild (or multiplier) bootstrap: Let  $w_1, \dots, w_n$  be i.i.d. standard normal variables independent of the data  $X_1, \dots, X_n$ . Define the Gaussian wild bootstrap version of  $S_n$  as follows:

$$S_n^* := \frac{1}{\sqrt{n}} \sum_{i=1}^n w_i (X_i - \bar{X}), \quad \text{where } \bar{X} = \frac{1}{n} \sum_{i=1}^n X_i. \quad (1.2)$$

We may naturally expect that  $c_{1-\alpha}^G$  would be well-approximated by the  $(1 - \alpha)$ -quantile of the conditional law of  $T_n^* := \max_{1 \leq j \leq d} S_{n,j}^*$  given the data, say  $\hat{c}_{1-\alpha}$ . This is formally justified by [17]: They essentially prove

$$P(T_n \geq \hat{c}_{1-\alpha}) = \alpha + O\left(\left(\frac{\log^a(dn)}{n}\right)^b\right) \quad (1.3)$$

with  $a = 7$  and  $b = 1/8$ . The successive work [19] have improved the convergence rates of (1.1) and (1.3) to  $b = 1/6$ . They also proved the left hand side of (1.1) can be replaced by  $\sup_{A \in \mathcal{R}} |P(S_n \in A) - P(Z \in A)|$ , where  $\mathcal{R} := \{\prod_{j=1}^d [a_j, b_j] : a_j \leq b_j, j = 1, \dots, d\}$  is the class of rectangles in  $\mathbb{R}^d$ .

It is easy to see that the conditional law of  $S_n^*$  given the data is  $N(0, \hat{\Sigma}_n)$ , where  $\hat{\Sigma}_n$  is the sample covariance matrix:  $\hat{\Sigma}_n := n^{-1} \sum_{i=1}^n (X_i - \bar{X})(X_i - \bar{X})^\top$ . Hence, the Gaussian wild bootstrap is essentially a feasible version of normal approximation for  $T_n$ . Then, it is natural to ask whether the approximation accuracy can be improved by more sophisticated bootstrap methods such as the empirical and non-Gaussian wild bootstraps. In the fixed-dimensional setting, it is well-known that standard bootstrap methods improve the approximation accuracy in the coverage probabilities upon normal approximation only when the statistic of interest is asymptotically pivotal (cf. [35, Chapter 3] and [43, Section 3]). However, despite that  $T_n$  and  $\|S_n\|_\infty$  are not asymptotically pivotal in general, numerical experiments suggest that third-moment match bootstrap methods would outperform normal approximation (cf. [21, 24]). To appreciate this, we depict in Fig. 1 the P-P plot for the rejection rate  $P(T_n \geq \hat{c}_{1-\alpha})$  against the nominal significance level  $\alpha$  when  $n = 200$  and  $d = 400$ , where  $\hat{c}_{1-\alpha}$  is computed either the Gaussian wild bootstrap or a wild bootstrap with third-moment match. We can clearly see that the latter performance is much better than the former.

Deng & Zhang [24] tried to explain this phenomenon by showing that convergence rates of third-moment match bootstrap approximations have a better dimension dependence, i.e. they achieve  $a = 5$  and  $b = 1/6$  in (1.3). Later, however, it was shown in [36] that the same convergence rate is achieved by normal approximation, i.e. (1.1) holds with  $a = 5$  and  $b = 1/6$ . Chernozhukov *et al.* [21] have further improved the




Figure 1: PP-plots for the rejection rate  $P(T_n \geq \hat{c}_{1-\alpha})$  against the nominal significance level  $\alpha$  when  $n = 200$  and  $d = 400$ . The rejection rate is evaluated based on 20,000 Monte Carlo iterations. The critical value  $\hat{c}_{1-\alpha}$  is computed by the Gaussian wild bootstrap for the left panel and the wild bootstrap with  $w_1$  generated from the standardized beta distribution with parameters  $\alpha, \beta$  given by (2.12) with  $\nu = 0.1$  for the right panel, respectively. The number of bootstrap replications is 499.  $X_1, \dots, X_n$  are generated from a Gaussian copula model with gamma marginals as in the simulation study of Section 4. The parameter matrix is  $R = (0.2^{|j-k|})_{1 \leq j, k \leq d}$ .

convergence rate to  $a = 5$  and  $b = 1/4$  for both normal and bootstrap approximations. Meanwhile, if we require  $\Sigma$  to be invertible, it is possible to achieve the Berry–Esseen rate  $n^{-1/2}$  up to a log factor even in the high-dimensional setting. Results in this direction first appeared in Fang & Koike [26], where the following result is obtained when  $X_1, \dots, X_n$  are log-concave:

$$\sup_{A \in \mathcal{R}} |P(S_n \in A) - P(Z \in A)| = O\left(\sqrt{\frac{\log^3 d}{n}} \log n\right). \quad (1.4)$$

This rate is known to be optimal up to the  $\log n$  factor in terms of both  $n$  and  $d$ ; see Proposition 1.1 in [26]. This type of results has been further investigated in [23, 39, 44]. In particular, Chernozhukov *et al.* [23] have obtained the above nearly optimal rate when  $\max_{i,j} |X_{ij}|$  is bounded. Here, the boundedness condition can be replaced with the sub-exponential condition by a simple truncation argument; see Appendix A. Further, in some situations, the rate  $n^{-1/2}$  is (nearly) attainable even when  $\Sigma$  is (asymptotically) degenerate; see [27, 29, 45]. Nevertheless, all of these improvements are valid for normal approximation and thus do not explain the superior performances of third-moment match bootstrap approximations.

In this paper, we aim to explain the superior performance of bootstrap approximation in high-dimensions using Edgeworth expansion and related techniques. Our first main result shows that if valid Edgeworth expansions for  $T_n$  and  $T_n^*$  are available, then we have

$$P(T_n \geq \hat{c}_{1-\alpha}) = \alpha + O\left(\sqrt{\frac{\log^2 d}{n}} \log n + \frac{\log^3(dn)}{n} \log n\right), \quad (1.5)$$

provided that  $\hat{c}_{1-\alpha}$  is computed by a third-moment match bootstrap method. Thus the coverage error has a better dimension dependence than the optimal normal approximation rate in (1.4). An analogous result

holds for  $\|S_n\|_\infty$  but we do not need the third-moment matching condition in this case. The next question is when we have valid Edgeworth expansions in the high-dimensional setting. We answer this question by proving the validity of Edgeworth expansion for  $S_n$  when  $X_i$  have Stein kernels (cf. Definition 2.1). This also allows us to derive a valid Edgeworth expansion for the wild bootstrap statistic  $S_n^*$  when the weights  $w_i$  have Stein kernels. In particular, our results cover the simulation setting for Fig. 1 (cf. Example 2.2). Finally, we construct a second-order accurate critical value  $\tilde{c}_{1-\alpha}$  in the sense that

$$P(T_n \geq \tilde{c}_{1-\alpha}) = \alpha + O\left(\frac{\log^a(dn)}{n}\right) \quad (1.6)$$

for some constant  $a > 0$ . A classical solution to this problem is bootstrapping the studentized version of  $S_n$ , but this is impossible in high-dimensions since the sample covariance matrix  $\hat{\Sigma}_n$  is degenerate whenever  $d \geq n$ . Instead, we achieve this by Beran [6]’s double bootstrap method, another classical technique to improve the approximation accuracy for non-pivotal statistics. To prove the second-order accuracy of the double bootstrap, we develop an asymptotic expansion formula of  $P(T_n \geq \hat{c}_{1-\alpha})$  in Theorem 3.3. As a byproduct, we find that the wild bootstrap with third moment match is *already* second-order accurate when  $d \geq n$  and  $\Sigma$  has identical diagonal entries and bounded eigenvalues, revealing the blessing of dimensionality in this context; see Corollary 3.1.

Despite that Edgeworth expansion is a standard tool to analyze the performance of bootstrap in the classical setting (cf. [35]), this approach has not been investigated for the above problem so far. A main reason would be the lack of valid Edgeworth expansion for  $T_n$  and  $\|S_n\|_\infty$  in the high-dimensional setting. While asymptotic expansion for statistics of high-dimensional data has been actively studied in multivariate statistics (see [33] for an overview), results developed there seem inapplicable to our problem. One main reason is that  $T_n$  and  $\|S_n\|_\infty$  may not have any limit distributions as  $n, d \rightarrow \infty$  even after properly scaled. In fact, this is one of the motivations for the development of Chernozhukov–Chetverikov–Kato’s theory. In view of (1.4), we are concerned with Edgeworth expansion of  $P(S_n \in A)$  over  $A \in \mathcal{R}$ . In the fixed-dimensional setting, a valid Edgeworth expansion of  $P(S_n \in A)$  is conventionally derived from an asymptotic expansion of the characteristic function of  $S_n$  via Fourier analysis (see e.g. [7]). Such an argument makes the dimension dependence of the error bound extremely complicated, so it is rarely given explicitly. One exceptional work is Anderson *et al.* [1], but their proof technique seems to inherently require the condition  $d \ll n$  and thus inapplicable to our setting. In fact, in the high-dimensional setting, the geometry of the set  $A$  plays a key role to get an improved dimension dependence of error bounds, and it is unclear how to incorporate such information into Fourier analytic arguments. We also mention the recent work by Zhilova [57] who establishes explicit, computable error bounds for  $\sup_{A \in \mathcal{A}} |P(S_n \in A) - P(S'_n \in A)|$  where  $S'_n$  is another sum of independent random vectors and  $\mathcal{A}$  is either the class of balls or half-spaces. However, apart from other technical issues, these error bounds contain  $1/\sqrt{n}$  terms and cannot be used for second-order analysis.

To circumvent the above issue, we develop valid asymptotic expansions using Stein’s method. The use of Stein’s method for asymptotic expansion was initiated by Barbour [2] who derived an asymptotic expansion of  $E[h(S_n)]$  when  $d = 1$  and  $h$  is a smooth function. To drop the smoothness of the test function  $h$ , the so-called Cramér’s condition is usually assumed in the Fourier analytic approach, but it is unknown how

to (directly) incorporate Cramér's condition into Stein's method based arguments. Instead, we assume that the underlying random vectors have Stein kernels, motivated by the recent development of this approach by Fang & Liu [30] in the univariate case (see Lemma 2.1 *ibidem*). Apart from the technical difficulty, Cramér's condition is violated whenever the underlying statistic has a singular covariance matrix. This is unsuitable for application to bootstrap statistics in high-dimensions, so Stein kernels will be a more appropriate tool for our problem (see Remark 2.6).

The remainder of the paper is organized as follows. In Section 2.1, we give the precise form of claim (1.5). In Section 2.2, we develop valid Edgeworth expansions for  $S_n$  and  $S_n^*$  in high-dimensions. Then, we study the second-order accuracy of bootstrap approximations for  $T_n$  in Section 3: After developing Cornish–Fisher type expansions for  $T_n$  and  $T_n^*$  in Section 3.1, we develop an asymptotic expansion formula for  $P(T_n \geq \hat{c}_{1-\alpha})$  in Section 3.2. Based on this result, we show in Section 3.3 that a double wild bootstrap method is second-order accurate. Section 4 contains a small simulation study. Most proofs are collected in Sections 5 and 6. Exceptions are proofs for properties of Stein kernels, which are given in Appendix C. The appendix also contains other additional proofs and auxiliary results.

**Notation** Throughout the paper, we assume that  $S_n$  has an invertible covariance matrix  $\Sigma$  and denote by  $\sigma_*$  the square root of the minimum eigenvalue of  $\Sigma$ . We also set  $\bar{\sigma} = \max_{j=1,\dots,d} \sqrt{\Sigma_{jj}}$  and  $\underline{\sigma} = \min_{j=1,\dots,d} \sqrt{\Sigma_{jj}}$ . Further,  $w_1, \dots, w_n$  denote i.i.d. random variables independent of  $X_1, \dots, X_n$ . They are used to define the wild bootstrap statistic  $S_n^*$  in (1.2). We always assume  $E[w_1] = 0$  and  $E[w_1^2] = 1$ . Also,  $P^*$  and  $E^*$  denote the conditional probability and expectation given the data  $X_1, \dots, X_n$ , respectively. For  $p \in (0, 1)$ ,  $\hat{c}_p$  denotes the conditional  $p$ -quantile of  $T_n^*$  given the data, i.e.  $\hat{c}_p := \inf\{t \in \mathbb{R} : P^*(T_n^* \leq t) \geq p\}$ .

For a vector  $x \in \mathbb{R}^d$ , we set  $|x| := \sqrt{\sum_{j=1}^d x_j^2}$  and  $x^\vee := \max_{1 \leq j \leq d} x_j$ . We denote by  $\mathbf{1}_d = (1, \dots, 1)^\top \in \mathbb{R}^d$  the all-ones vector in  $\mathbb{R}^d$ . For  $r \in \mathbb{N}$ ,  $(\mathbb{R}^d)^{\otimes r}$  denotes the set of real-valued  $d$ -dimensional  $r$ -arrays  $V = (V_{j_1, \dots, j_r})_{1 \leq j_1, \dots, j_r \leq d}$ . In particular,  $(\mathbb{R}^d)^{\otimes 1} = \mathbb{R}^d$  and  $(\mathbb{R}^d)^{\otimes 2}$  is the set of  $d \times d$  matrices. For  $U \in (\mathbb{R}^d)^{\otimes q}$  and  $V \in (\mathbb{R}^d)^{\otimes r}$ , we set  $U \otimes V := (U_{i_1, \dots, i_q} V_{j_1, \dots, j_r})_{1 \leq i_1, \dots, i_q, j_1, \dots, j_r \leq d} \in (\mathbb{R}^d)^{\otimes (q+r)}$ . We write  $U^{\otimes 2} = U \otimes U$  for short. When  $q = r$ , we also set  $\langle U, V \rangle := \sum_{j_1, \dots, j_r=1}^d U_{j_1, \dots, j_r} V_{j_1, \dots, j_r}$ . In particular, when  $q = r = 1$ ,  $\langle U, V \rangle$  is the Euclidean inner product of  $U$  and  $V$  which we also write  $U \cdot V$ . In addition, we set  $\|V\|_1 := \sum_{j_1, \dots, j_r=1}^d |V_{j_1, \dots, j_r}|$  and  $\|V\|_\infty := \max_{1 \leq j_1, \dots, j_r \leq d} |V_{j_1, \dots, j_r}|$ . Further, for  $x \in \mathbb{R}^d$ , we define  $x^{\otimes r} := (x_{j_1} \cdots x_{j_r})_{1 \leq j_1, \dots, j_r \leq d} \in (\mathbb{R}^d)^{\otimes r}$ . Finally, we set

$$\bar{X}^r := \frac{1}{n} \sum_{i=1}^n X_i^{\otimes r}.$$

Given an  $r$ -times differentiable function  $h : \mathbb{R}^d \rightarrow \mathbb{R}$ , we set  $\nabla^r h(x) := (\partial_{j_1, \dots, j_r} h(x))_{1 \leq j_1, \dots, j_r \leq d} \in (\mathbb{R}^d)^{\otimes r}$  for  $x \in \mathbb{R}^d$ , where  $\partial_{j_1, \dots, j_r} = \frac{\partial^r}{\partial x_{j_1} \cdots \partial x_{j_r}}$ . For  $m \in \mathbb{N} \cup \{\infty\}$ ,  $C_b^m(\mathbb{R}^d)$  denotes the set of bounded  $C^m$  functions with bounded derivatives.

For an invertible matrix  $V$ ,  $\phi_V$  denotes the density of  $N(0, V)$ . We write  $\phi_d = \phi_{I_d}$  for short, where  $I_d$  is the  $d \times d$  identity matrix. Further, we write  $\phi = \phi_1$  for short.  $\Phi$  denotes the standard normal distribution function. Also, for a distribution function  $F : \mathbb{R} \rightarrow [0, 1]$ , its (generalized) inverse is defined as  $F^{-1}(p) =$

$\inf\{t \in \mathbb{R} : F(t) \geq p\}$ ,  $p \in (0, 1)$ . We refer to Appendix A.1 in [10] for useful properties of inverse distribution functions.

For a random vector  $\xi$  and  $p \in (1, \infty)$ , we set  $\|\xi\|_p := (\mathbb{E}[|\xi|^p])^{1/p}$  (recall that  $|\cdot|$  is the Euclidean norm). Further, for  $\alpha > 0$ , we set  $\|\xi\|_{\psi_\alpha} := \inf\{t > 0 : \mathbb{E}[\exp\{(|\xi|/t)^\alpha\}] \leq 2\}$ . For two random vectors  $\xi$  and  $\eta$ , we write  $\xi \stackrel{d}{=} \eta$  if  $\xi$  has the same law as  $\eta$ .

We assume  $d \geq 3$  whenever we consider an expression containing  $\log d$ . A similar convention is applied to  $n$ .

## 2 High-dimensional bootstrap and Edgeworth expansion

### 2.1 Coverage error bounds via Edgeworth expansion

We begin by introducing appropriate (second-order) Edgeworth expansions for  $S_n$  and  $S_n^*$ . The former is standard. That is, our Edgeworth expansion for  $S_n$  is defined as

$$p_n(z) = \phi_\Sigma(z) - \frac{1}{6\sqrt{n}} \langle \mathbb{E}[\overline{X^3}], \nabla^3 \phi_\Sigma(z) \rangle, \quad z \in \mathbb{R}^d.$$

The situation is different for the latter. In the low-dimensional setting, a natural bootstrap version of  $p_n(z)$  would be obtained by replacing  $\Sigma$  and  $\mathbb{E}[\overline{X^3}]$  with their sample counterparts  $\widehat{\Sigma}_n$  and  $\overline{X^3}$ , respectively. However, when  $d \geq n$ ,  $\widehat{\Sigma}_n$  is always degenerate, so  $\phi_{\widehat{\Sigma}_n}$  is not well-defined. For this reason, we consider an Edgeworth expansion ‘‘around  $\phi_\Sigma$ ’’. Formally, our Edgeworth expansion for  $S_n^*$  is defined as

$$\hat{p}_{n,\gamma}(z) = \phi_\Sigma(z) + \frac{1}{2} \langle \overline{X^2} - \Sigma, \nabla^2 \phi_\Sigma(z) \rangle - \frac{\gamma}{6\sqrt{n}} \langle \overline{X^3}, \nabla^3 \phi_\Sigma(z) \rangle, \quad z \in \mathbb{R}^d,$$

where  $\gamma \in \mathbb{R}$  is a constant determined by the construction of  $S_n^*$ . We expect  $\gamma = 1$  for third moment match bootstrap methods.

**Theorem 2.1.** *Suppose that there exist constants  $\gamma \in \mathbb{R}$ ,  $\Delta_n > 0$  and  $\delta_n \in (0, 1)$  such that*

$$\sup_{A \in \mathcal{R}} \left| P(S_n \in A) - \int_A p_n(z) dz \right| \leq \Delta_n \tag{2.1}$$

and

$$\sup_{A \in \mathcal{R}} \left| P^*(S_n^* \in A) - \int_A \hat{p}_{n,\gamma}(z) dz \right| \leq \Delta_n \tag{2.2}$$

with probability at least  $1 - \delta_n$ . Suppose also that there exists a constant  $b > 0$  such that

$$\max_{1 \leq i \leq n} \max_{1 \leq j \leq d} \|X_{ij}\|_{\psi_1} \leq b. \tag{2.3}$$

Then, there exists a universal constant  $C > 0$  such that

$$|P(T_n \geq \hat{c}_{1-\alpha}) - \alpha| \leq C(1 + |\gamma|) \frac{b^2 \log d}{\sigma_*^2 \sqrt{n}} \sqrt{\log n} + 3\Delta_n + \delta_n \tag{2.4}$$

for any  $\alpha \in (0, 1)$ , provided that  $\gamma = 1$  or  $\mathbb{E}[\overline{X^3}] = 0$ . Further, with  $|\hat{c}_{1-\alpha}$  denoting the  $(1 - \alpha)$ -quantile of  $\|S_n\|_\infty$ , we have

$$|P(\|S_n\|_\infty \geq |\hat{c}_{1-\alpha}) - \alpha| \leq C \frac{b^2 \log d}{\sigma_*^2 \sqrt{n}} \sqrt{\log n} + 3\Delta_n + \delta_n \tag{2.5}$$

regardless of the values of  $\gamma$  and  $\mathbb{E}[\overline{X^3}]$ .

**Remark 2.1.** (a) The sub-exponential assumption (2.3) is imposed just for clarity. It is necessary only for deriving concentration inequalities for terms of  $\hat{p}_{n,\gamma}$  (cf. Lemma E.10) and can be replaced by another assumption as soon as such bounds are available.

(b) While Theorem 2.1 is stated for the wild bootstrap, the conclusion remains true for the empirical bootstrap as long as (2.2) is satisfied. However, so far we have no result to ensure (2.2) with a reasonable  $\Delta_n$  for the empirical bootstrap in the high-dimensional setting.

In the next subsection we will see that (2.1) and (2.2) hold with  $\Delta_n \asymp \frac{\log^3(dn)}{n} \log n$  and  $\delta_n = 1/n$  under regularity conditions. Hence we have (1.5) for the third-moment match bootstrap, showing that it could give a better approximation in the coverage probability than the normal approximation. An intuition behind this improvement is as follows. (2.1) and (2.2) imply that  $P^*(S_n^* \in A) - P(S_n \in A)$  is approximately equal to

$$\int_A \{\hat{p}_{n,\gamma}(z) - p_n(z)\} dz = \frac{1}{2} \langle \overline{X^2} - \Sigma, \int_A \nabla^2 \phi_\Sigma(z) dz \rangle - \frac{1}{6\sqrt{n}} \langle \gamma \overline{X^3} - \mathbb{E}[\overline{X^3}], \int_A \nabla^3 \phi_\Sigma(z) dz \rangle \quad (2.6)$$

for every  $A \in \mathcal{R}$ . When  $\gamma = 1$  or  $\mathbb{E}[\overline{X^3}] = 0$ , the right hand side can be written as a sum of centered independent random variables, so a standard argument shows that it is of order  $O_p(n^{-1/2} \log d + n^{-1} \log^{3/2} d)$  for a fixed sequence of  $A \in \mathcal{R}$  (cf. Lemmas E.4 and E.10). It turns out that such estimates give an error bound for  $P(T_n \geq \hat{c}_{1-\alpha})$  of essentially the same order. For  $P(\|S_n\|_\infty \geq |\hat{c}_{1-\alpha}|)$ , it suffices to consider rectangles of the form  $A = [-c, c]^d$  for some  $c > 0$ . In this case, the second term on the right hand side of (2.6) always vanishes since  $A$  is symmetric. Hence we need neither  $\gamma = 1$  nor  $\mathbb{E}[\overline{X^3}] = 0$ .

**Remark 2.2** (Estimation of distribution functions). Deng & Zhang [24] actually focus on the bootstrap estimation error for the distribution function of  $T_n$  in the Kolmogorov distance, i.e.  $\sup_{t \in \mathbb{R}} |P(T_n \leq t) - P^*(T_n^* \leq t)|$ . For this problem, there is a theoretical explanation for why bootstrap approximation outperforms normal approximation in the fixed dimensional setting; see [5, Section 2.1] for details and also [43] for a related discussion. In view of the superior performance of bootstrap approximation reported in the simulation study of [24], we may naturally expect that results in [5] could be extended to the high-dimensional setting. The formal development is left to future research.

**Remark 2.3.** Theorem 2.1 does *not* mean that third-moment match bootstraps work with the weaker requirement  $\log^2 d = o(n/\log n)$  compared to the normal approximation. This is because we usually need at least  $\log^3 d = o(n)$  to have  $\Delta_n$  vanish. It is known that for some high-dimensional linear models, bootstrap for linear contrasts works with a weaker requirement on the model dimension than the normal approximation (see [46]), so it will be interesting to study whether a similar phenomenon occurs for maximum type statistics.

## 2.2 Valid Edgeworth expansion in high-dimensions

Let us formally define the notion of Stein kernel.

**Definition 2.1** (Stein kernel). Let  $\xi$  be a random vector in  $\mathbb{R}^d$  with  $\mathbb{E}[\|\xi\|_\infty] < \infty$ . A measurable function  $\tau : \mathbb{R}^d \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d$  is called a *Stein kernel* for (the law of)  $\xi$  if  $\mathbb{E}[\|\tau(\xi)\|_\infty] < \infty$  and

$$\mathbb{E}[(\xi - \mathbb{E}[\xi]) \cdot \nabla h(\xi)] = \mathbb{E}[\langle \tau(\xi), \nabla^2 h(\xi) \rangle] \quad (2.7)$$

for any  $h \in C_b^2(\mathbb{R}^d)$ .

The concept of Stein kernel was originally introduced in Stein [55, Lecture VI] for the univariate case. Although its partial multivariate extension dates back to [13], general treatments have started in more recent studies of [41, 51], stemming from the discovery of connection to Malliavin calculus due to Nourdin & Peccati [49] (the so-called *Malliavin–Stein method*). We refer to [47] for the recent development.

**Remark 2.4** (Alternative definition). Our definition of Stein kernel is taken from [41]. In the literature, the definition of Stein kernel often requires (2.7) to hold with  $\nabla h$  on the both sides replaced by any bounded  $C^1$  function  $h : \mathbb{R}^d \rightarrow \mathbb{R}^d$  with bounded derivatives. Except for the case  $d = 1$ , this requirement is slightly stronger than ours. Nevertheless, as far as the author knows, this stronger requirement has so far been met by all known constructions of Stein kernels, including all the examples of this paper.

The validity of Edgeworth expansion for  $S_n$  is ensured if the summands have Stein kernels:

**Theorem 2.2** (Edgeworth expansion for  $S_n$ ). *Suppose that  $X_i$  has a Stein kernel  $\tau_i^X$  for every  $i = 1, \dots, n$ . Suppose also that there exists a constant  $b > 0$  such that*

$$\|X_{ij}\|_{\psi_1} \leq b, \quad \|\tau_{i,jk}^X(X_i)\|_{\psi_{1/2}} \leq b^2 \quad (2.8)$$

for all  $i = 1, \dots, n$  and  $j, k = 1, \dots, d$ . Further, assume  $\log^3 d \leq n$ . Then there exists a universal constant  $C > 0$  such that

$$\sup_{A \in \mathcal{R}} \left| P(S_n \in A) - \int_A p_n(z) dz \right| \leq C \frac{b^5 \log^3 d}{\sigma_*^5 n} \log n. \quad (2.9)$$

**Remark 2.5.** Here and below, we do not intend to optimize the dependence of bounds on  $b$  and  $\sigma_*$ .

Below we give a few examples satisfying (2.8).

**Example 2.1** (Log-concave distribution). When  $X_i$  has a log-concave density,  $X_i$  has a Stein kernel  $\tau^X$  and (2.8) is satisfied with  $b = C \max_{1 \leq j \leq d} \sqrt{\text{Var}[X_{ij}]}$  for some universal constant  $C > 0$  by [31, Theorem 2.3 and Proposition 3.2] and Lemma E.5.

**Example 2.2** (Gaussian copula model). Let  $R$  be a  $d \times d$  positive semidefinite symmetric matrix with unit diagonals. Also, for every  $j = 1, \dots, d$ , let  $\mu_j$  be a non-degenerate probability distribution on  $\mathbb{R}$  (i.e.  $\mu_j$  is not the unit mass at a point), and denote by  $F_j$  its distribution function. The Gaussian copula model  $U = (U_1, \dots, U_d)^\top$  with parameter matrix  $R$  and marginal distributions  $\mu_1, \dots, \mu_d$  is defined as  $U_j = F_j^{-1}(\Phi(Z_j))$  for  $j = 1, \dots, d$ , where  $Z \sim N(0, R)$ .

**Proposition 2.1** (Stein kernel of Gaussian copula model). *Suppose that there exists a constant  $\kappa > 0$  such that for every  $j = 1, \dots, d$  and any Borel set  $B \subset \mathbb{R}$ ,*

$$\liminf_{h \downarrow 0} \frac{\mu_j(B^h) - \mu_j(B)}{h} \geq \kappa \min\{\mu_j(B), 1 - \mu_j(B)\}, \quad (2.10)$$

where  $B^h := \{t \in \mathbb{R} : |t - s| < h \text{ for some } s \in B\}$ . Then  $X := U - \mathbb{E}[U]$  has a Stein kernel  $\tau$  and

$$\max_{1 \leq j \leq d} \|X_j\|_{\psi_1} \leq C\kappa^{-1}, \quad \max_{1 \leq j, k \leq d} \|\tau_{jk}(X)\|_{\psi_1} \leq C\kappa^{-2}$$

for some universal constant  $C > 0$ .

The maximal constant  $\kappa$  satisfying (2.10) is called the *Cheeger (isoperimetric) constant* of  $\mu_j$ . We refer to [9, Theorem 1.3] for a useful equivalent formulation in the univariate case. When  $\mu_j$  is log-concave, then (2.10) is satisfied with  $\kappa = 1/\sqrt{3 \text{Var}[X_j]}$  by Proposition 4.1 in [8]. Since the gamma distribution with shape parameter  $\geq 1$  is log-concave, Proposition 2.1 shows that the simulated model in the introduction satisfies the assumptions of Theorem 2.2. We can actually show that any gamma distribution has a positive Cheeger constant; see Proposition C.2.

**Example 2.3** (Multiplicative perturbation). Let  $X$  be a random vector in  $\mathbb{R}^d$  and  $\epsilon$  a centered random variable independent of  $X$  and having a Stein kernel  $\tau$ . Then  $\epsilon X$  has a Stein kernel  $x \mapsto \mathbb{E}[\tau(\epsilon)X^{\otimes 2} \mid \epsilon X = x]$ , provided that  $\mathbb{E}[\|\epsilon X\|_\infty] + \mathbb{E}[\|\tau(\epsilon)X^{\otimes 2}\|_\infty] < \infty$ . This easily follows by applying Lemma C.1 conditional on  $X$ . This type of random vector arises in high-dimensional regression; see [17, Section 4].

Other constructions of multivariate Stein kernels are found in [47, Section 4], although it does not seem straightforward to verify the second condition of (2.8) for them.

**Remark 2.6** (Relation to classical conditions). (a) In the univariate case, if a non-degenerate distribution has a Stein kernel, then it has a non-zero absolutely continuous part; see Proposition C.1. In particular, it must satisfy Cramér's condition. It is worth mentioning that while univariate Stein kernels are often investigated in the existence of density in the literature, a non-degenerate distribution without density can have a Stein kernel. A simple example is the law of  $I\zeta$ , where  $I$  is a Bernoulli variable with success probability  $p \in (0, 1)$  and  $\zeta$  is a standard normal variable independent of  $I$ . In this case, we can easily check that  $I\zeta$  has a Stein kernel  $1_{\mathbb{R} \setminus \{0\}}$ . More interesting examples are given by Example 2.2 since any univariate distribution can be realized as a Gaussian copula model and (2.10) can hold without density; see discussions after [9, Theorem 1.3].

(b) In the multivariate case, a non-degenerate distribution may not satisfy Cramér's condition even when it has a Stein kernel: A simple example is a multivariate normal distribution with singular covariance matrix. This example is indeed important in the high-dimensional setting when analyzing the Gaussian wild bootstrap.

We turn to Edgeworth expansion for  $S_n^*$ . Its validity is ensured if the weight variables have Stein kernels:

**Theorem 2.3** (Edgeworth expansion for  $S_n^*$ ). *Suppose that (2.3) is satisfied. Suppose also that  $w_1$  satisfies either of the following conditions:*

- (i)  $w_1$  has a Stein kernel  $\tau^*$  and there exists a constant  $b_w \geq 1$  such that  $|w_1| \leq b_w$  and  $|\tau^*(w_1)| \leq b_w^2$ .
- (ii)  $w_1 \sim N(0, 1)$ . We set  $b_w = 1$  in this case.

Further, assume  $\log^3 d \leq n$ . Set  $\gamma := \mathbb{E}[w_1^3]$ . Then there exists a universal constant  $C > 0$  such that

$$\sup_{A \in \mathcal{R}} \left| P^*(S_n^* \in A) - \int_A \hat{p}_{n,\gamma}(z) dz \right| \leq C \frac{b_w^5 b^5 \log^3(dn)}{\sigma_*^5 n} \log n \quad (2.11)$$

with probability at least  $1 - 1/n$ .

We can construct a random variable  $w_1$  satisfying Condition (i) and  $E[w_1^3] = 1$  as follows: Let  $\eta$  be a random variable following the beta distribution with parameters  $\alpha, \beta > 0$ . Then  $w := (\eta - E[\eta])/\sqrt{\text{Var}[\eta]}$  satisfies (i) by [42, Example 4.9] and Lemma C.1. Also, we have

$$E[w_1^3] = \frac{2(\beta - \alpha)\sqrt{\alpha + \beta + 1}}{(\alpha + \beta + 2)\sqrt{\alpha\beta}} = \frac{2(1 - 2\mu)\sqrt{1 + \nu}}{(2 + \nu)\sqrt{\mu(1 - \mu)}},$$

where  $\mu = \alpha/(\alpha + \beta)$  and  $\nu = \alpha + \beta$ . From this expression, given a positive constant  $\nu > 0$ , we have  $E[w_1^3] = 1$  if we set

$$\alpha = \nu \frac{c - (2 + \nu)\sqrt{c}}{2c}, \quad \beta = \nu \frac{c + (2 + \nu)\sqrt{c}}{2c} \quad \text{with } c = \nu^2 + 20\nu + 20. \quad (2.12)$$

A drawback of Theorem 2.3 is that two-point distributions do not admit Stein kernels (cf. Proposition C.1). In particular, it does not cover Mammen's wild bootstrap (cf. Eq.(4.1)) examined in the simulation study of [24]. However, the above standardized beta distribution becomes closer to Mammen's two-point distribution as  $\nu$  is closer to 0, and their numerical difference virtually vanishes. Our simulation study shows that the beta wild bootstrap with  $\nu = 0.1$  performs very similarly to Mammen's one.

### 3 Second-order accurate approximation

Our next aim is to construct a second-order accurate critical value  $\hat{c}$  in the sense that (1.6) holds. To accomplish this, we will develop an asymptotic expansion of the bootstrap coverage probability. Such an expansion is conventionally derived with the help of Cornish–Fisher expansion (cf. Section 3.5.2 in [35]), so we first develop such expansions for  $T_n$  and  $T_n^*$  in our setting.

Before starting discussions, we introduce some notation used throughout this section. For  $t \in \mathbb{R}$ , we set  $A(t) := (-\infty, t]^d$ . We denote by  $f_\Sigma$  the density of  $Z^\vee$ , where  $Z \sim N(0, \Sigma)$ . Note that  $f_\Sigma$  is a  $C^\infty$  function since  $\Sigma$  is invertible. Finally, we set  $\varsigma_d := \sqrt{\text{Var}[Z^\vee] \log d}$ . By Lemma E.3,  $\varsigma_d$  is bounded from below by a positive constant depending only on  $\bar{\sigma}$  and  $\underline{\sigma}$ . By Lemma E.1,  $\varsigma_d$  is generally bounded by  $\bar{\sigma}\sqrt{\log d}$ , but we often have  $\varsigma_d = O(1)$  as  $d \rightarrow \infty$ , known as a *superconcentration* phenomenon (cf. [15]). For example, this is the case when  $\Sigma_{jj} = 1$  for all  $j$  and there exists a constant  $C > 0$  such that  $\Sigma_{jk} \leq C/\log(2 + |j - k|)$  for all  $j, k$ . This follows from [15, Theorem 9.12].

#### 3.1 Cornish–Fisher expansion

This section develops Cornish–Fisher type expansions for  $T_n$  and  $T_n^*$ .

**Theorem 3.1** (Cornish–Fisher expansion for  $T_n$ ). *Under the assumptions of Theorem 2.2, let  $\lambda > 0$  be a constant such that  $b/\sigma_* \leq \lambda$ . Then, for any  $\varepsilon \in (0, 1/2)$ , there exist positive constants  $c$  and  $C$  depending only on  $\lambda$  and  $\varepsilon$  such that if*

$$\frac{\varsigma_d^3 \log^3 d}{\sigma_*^3 n} \log n \leq c, \quad (3.1)$$

then

$$\sup_{\varepsilon < p < 1 - \varepsilon} \left| c_p - \left( c_p^G - \frac{Q_n(c_p^G)}{f_\Sigma(c_p^G)} \right) \right| \leq \frac{C}{\sqrt{\log d}} \frac{\varsigma_d^3 \log^3 d}{\sigma_*^2 n} \log n, \quad (3.2)$$

where  $c_p$  is the  $p$ -quantile of  $T_n$  and

$$Q_n(t) := \int_{A(t)} \{p_n(z) - \phi_\Sigma(z)\} dz = -\frac{1}{6\sqrt{n}} \langle \mathbb{E}[\overline{X^3}], \int_{A(t)} \nabla^3 \phi_\Sigma(z) dz \rangle, \quad t \in \mathbb{R}.$$

**Theorem 3.2** (Cornish–Fisher expansion for  $T_n^*$ ). *Under the assumptions of Theorem 2.3, let  $\lambda > 0$  be a constant such that  $b/\sigma_* \leq \lambda$ . Then, for any  $\varepsilon \in (0, 1/2)$ , there exist positive constants  $c$  and  $C$  depending only on  $\lambda, \varepsilon$  and  $b_w$  such that if*

$$\frac{\varsigma_d^3 \log^3(dn)}{\sigma_*^3 n} \log n \leq c, \quad (3.3)$$

then

$$\sup_{\varepsilon < p < 1-\varepsilon} \left| \hat{c}_p - \left( c_p^G - \frac{\hat{Q}_{n,\gamma}(c_p^G)}{f_\Sigma(c_p^G)} \right) \right| \leq \frac{C}{\sqrt{\log d}} \frac{\varsigma_d^3 \log^3(dn)}{\sigma_*^2 n} \log n \quad (3.4)$$

with probability at least  $1 - 1/n$ , where

$$\hat{Q}_{n,\gamma}(t) := \int_{A(t)} \{\hat{p}_{n,\gamma}(z) - \phi_\Sigma(z)\} dz.$$

**Remark 3.1.** We are not getting valid Cornish–Fisher type expansions for  $\|S_n\|_\infty$  and  $\|S_n^*\|_\infty$  because it is not straightforward to derive an adequate bound for the second derivative of the quantile function of  $\|Z\|_\infty$ . Since there is another technical issue to develop asymptotic expansion of  $P(\|S_n\|_\infty \geq |\hat{c}|_{1-\alpha})$  (see Remark 3.3), we do not pursue them in this paper.

### 3.2 Asymptotic expansion of coverage probability

For a  $d \times d$  matrix  $V$ ,  $\text{vec}(V)$  denotes the  $d^2$ -dimensional vector obtained by stacking the columns of  $V$ . For two random vectors  $\xi$  and  $\eta$ , the random vector  $(\xi^\top, \eta^\top)^\top$  will be denoted by  $(\xi, \eta)$  for simplicity.

**Theorem 3.3** (Asymptotic expansion of bootstrap coverage probability). *Suppose that the assumptions of Theorem 3.2 are satisfied. For every  $i = 1, \dots, n$ , set  $Y_i := \text{vec}(X_i^{\otimes 2} - \mathbb{E}[X_i^{\otimes 2}])$  and suppose that the  $(d + d^2)$ -dimensional random vector  $(X_i, Y_i)$  has a Stein kernel  $\bar{\tau}_i$  of the form*

$$\bar{\tau}_i = \begin{pmatrix} \tau_i^X & \tau_i^{XY} \\ \tau_i^{YX} & \tau_i^Y \end{pmatrix} \quad (3.5)$$

with  $\tau_i^X$  an  $(\mathbb{R}^d)^{\otimes 2}$ -valued function and such that

$$\begin{aligned} \max_{1 \leq j, k \leq d} \|\tau_{i,jk}^X(X_i, Y_i)\|_{\psi_{1/2}} &\leq b^2, & \max_{1 \leq j, k \leq d^2} \|\tau_{i,jk}^Y(X_i, Y_i)\|_{\psi_{1/4}} &\leq b^4, \\ \max_{1 \leq j \leq d, 1 \leq k \leq d^2} \left( \|\tau_{i,jk}^{XY}(X_i, Y_i)\|_{\psi_{1/3}} + \|\tau_{i,kj}^{YX}(X_i, Y_i)\|_{\psi_{1/3}} \right) &\leq b^3. \end{aligned} \quad (3.6)$$

Then, for any  $\varepsilon \in (0, 1/2)$ , there exist positive constants  $c$  and  $C$  depending only on  $\lambda, \varepsilon$  and  $b_w$  such that if (3.3) holds, then

$$\sup_{\varepsilon < \alpha < 1-\varepsilon} \left| P(T_n \geq \hat{c}_{1-\alpha}) - (\alpha - (1-\gamma)Q_n(c_{1-\alpha}^G) - \mathbb{E}[R_n(\alpha)]) \right| \leq C \frac{\varsigma_d^3 \log^3(dn)}{\sigma_*^3 n} \log n,$$

where

$$R_n(\alpha) := \frac{1}{\sqrt{n}} \frac{\langle \overline{X^3} \otimes \mathbf{1}_d, \Psi_\alpha^{\otimes 2} \rangle}{2f_\Sigma(c_{1-\alpha}^G)}, \quad \Psi_\alpha := \int_{A(c_{1-\alpha}^G)} \nabla^2 \phi_\Sigma(z) dz.$$

**Remark 3.2** (Univariate case). When  $d = 1$  and  $\Sigma = 1$ , the above asymptotic expansion formula reduces to

$$\begin{cases} \alpha - \frac{\mathbb{E}[X^3]}{6\sqrt{n}} \{2(c_{1-\alpha}^G)^2 + 1\} \phi(c_{1-\alpha}^G) & \text{if } \gamma = 0, \\ \alpha - \frac{\mathbb{E}[X^3]}{2\sqrt{n}} (c_{1-\alpha}^G)^2 \phi(c_{1-\alpha}^G) & \text{if } \gamma = 1. \end{cases}$$

These recover the asymptotic expansion formulae for normal and empirical bootstrap coverage probabilities, respectively; see e.g. [43, Eqs.(2)–(3)] (note that  $c_{1-\alpha}^G = \Phi^{-1}(1 - \alpha) = -\Phi^{-1}(\alpha)$  when  $d = 1$ ).

The new assumption in Theorem 3.3 is the existence of a (nice) Stein kernel for  $(X_i, Y_i)$ . This assumption can be viewed as a counterpart of joint Cramér’s condition for  $X_i$  and  $Y_i$  that is typically imposed to derive a univariate counterpart of Theorem 3.3; see e.g. Eq.(2.54) in [35]. It is natural in this sense, but the verification is not easy in practice. Here, we give one sufficient condition following Mikulincer [48]’s idea of using the Malliavin–Stein method.

**Lemma 3.1.** *Let  $G$  be a standard Gaussian vector in  $\mathbb{R}^{d'}$ . Let  $\psi : \mathbb{R}^{d'} \rightarrow \mathbb{R}^d$  be a locally Lipschitz function such that  $\mathbb{E}[|\psi(G)|^2] < \infty$  and  $\max_{1 \leq j \leq d} \mathbb{E}[|\nabla \psi_j(G)|^2] < \infty$ . Then  $X := \psi(G) - \mathbb{E}[\psi(G)]$  has a Stein kernel  $\tau$  such that*

$$\|\tau_{jk}(X)\|_p \leq \|\nabla \psi_j(G)\|_{2p} \|\nabla \psi_k(G)\|_{2p} \quad (3.7)$$

for all  $p \geq 1$  and  $j, k = 1, \dots, d$ . In addition,

$$\|X_j\|_p \leq \sqrt{p-1} \|\nabla \psi_j(G)\|_p \quad (3.8)$$

for any even integer  $p \geq 2$  and  $j = 1, \dots, d$ .

Moreover, if we further assume  $\mathbb{E}[|\psi(G)|^4] < \infty$  and  $\max_{1 \leq j \leq d} \mathbb{E}[|\psi(G)|^2 |\nabla \psi_j(G)|^2] < \infty$ , then for  $Y = \text{vec}(X^{\otimes 2} - \mathbb{E}[X^{\otimes 2}])$ ,  $(X, Y)$  has a Stein kernel of the form (3.5) and satisfies

$$\begin{aligned} \max_{1 \leq j, k \leq d} \|\tau_{i,jk}^X(X, Y)\|_p &\leq \max_{1 \leq j \leq d} \|\nabla \psi_j(G)\|_{2p}^2, \\ \max_{1 \leq j \leq d, 1 \leq l \leq d^2} (\|\tau_{i,jl}^{XY}(X, Y)\|_p \vee \|\tau_{i,lj}^{YX}(X, Y)\|_p) &\leq 2 \max_{1 \leq j, k, l \leq d} \|\nabla \psi_j(G)\|_{2p} \|X_l \nabla \psi_k(G)\|_{2p}, \\ \max_{1 \leq l, m \leq d^2} \|\tau_{i,lm}^Y(X, Y)\|_p &\leq 4 \max_{1 \leq j, k \leq d} \|X_j \nabla \psi_k(G)\|_{2p}^2 \end{aligned}$$

for all  $p \geq 1$ .

Using this lemma, we give a few examples satisfying (3.6).

**Example 3.1** (Uniformly log-concave distribution). Let  $\varepsilon > 0$ . A probability density function  $f : \mathbb{R}^d \rightarrow [0, \infty)$  is said to be  $\varepsilon$ -uniformly log-concave if there exists a log-concave function  $g : \mathbb{R}^d \rightarrow [0, \infty)$  such that  $q(x) = g(x)e^{-\varepsilon|x|^2/2}$  for all  $x \in \mathbb{R}^d$ . If  $X_i$  has an  $\varepsilon$ -uniformly log-concave density, there exists a 1-Lipschitz function  $\psi_0 : \mathbb{R}^d \rightarrow \mathbb{R}^d$  such that  $X_i$  has the same law as  $\psi_0(\varepsilon^{-1/2}G)$  with  $G \sim N(0, I_d)$  by Caffarelli’s log-concave perturbation theorem (cf. [14, Theorem 11]). Hence  $(X_i, Y_i)$  has a Stein kernel of the form (3.5) and satisfies (3.6) with  $b = C\varepsilon^{-1/2}$  for some universal constant  $C > 0$ . We remark that results with similar natures to Caffarelli’s theorem are available for other distributions. We refer to [32] and references therein.

**Example 3.2** (Gaussian copula model). Consider the same setting as Example 2.2. Proposition 2.1 can be extended as follows.

**Proposition 3.1.** *Set  $Y = \text{vec}(X^{\otimes 2} - \mathbb{E}[X^{\otimes 2}])$ . Under the assumptions of Proposition 2.1,  $(X, Y)$  has a Stein kernel of the form (3.5) and satisfies (3.6) with  $b = C\kappa^{-1}$  for some universal constant  $C > 0$ .*

Now we discuss implications of Theorem 3.3 to the second-order accuracy of standard bootstrap approximations. An easy consequence is that any wild bootstrap approximation is second-order accurate when  $\mathbb{E}[\overline{X^3}] = 0$  as long as  $w_1$  satisfies the assumptions in Theorem 2.3. However, simulation results suggest that the choice of  $w_1$  would affect the performance even when  $\mathbb{E}[\overline{X^3}] = 0$ , so there is still room to investigate.

The following corollary gives a more interesting implication:

**Corollary 3.1.** *Under the assumptions of Theorem 3.3, suppose additionally that  $\mathbb{E}[w_1^3] = 1$ ,  $\varepsilon \geq 2e^{-d/2}$ ,  $\underline{\sigma} = \overline{\sigma} =: \sigma$  and the maximum eigenvalue of  $\Sigma$  is bounded by  $K\sigma^2$  with some constant  $K > 0$ . Then there exist a constant  $C > 0$  depending only on  $\lambda, \varepsilon, K$  and  $b_w$  such that*

$$\sup_{\varepsilon < \alpha < 1 - \varepsilon} |P(T_n \geq \hat{c}_{1-\alpha}) - \alpha| \leq C \left( \frac{\varsigma_d^3 \log^3(dn)}{\sigma_*^3 n} \log n + \frac{\varsigma_d}{\sigma} \sqrt{\frac{\log^3 d}{dn}} \right). \quad (3.9)$$

Observe that the second term on the right hand side of (3.9) is divided by  $\sqrt{d}$ . Hence, Corollary 3.1 implies that the third-moment match wild bootstrap is second-order accurate if  $d \geq n$  and  $\Sigma$  has identical diagonal entries and bounded eigenvalues with respect to  $d$ . This seems to be a new result on the blessing of dimensionality, although too high-dimensionality is harmful due to the first term of the bound.

**Remark 3.3.** The proof of Theorem 3.3 relies crucially on the identity  $\max_{1 \leq j \leq d} x_j + a = \max_{1 \leq j \leq d} (x_j + a)$  for  $x \in \mathbb{R}^d$  and  $a \in \mathbb{R}$ . We will use this identity to get an Edgeworth expansion for  $\tilde{T}_n = T_n + \eta$  with  $\eta$  a sum of independent random variables (see (6.15)), i.e.  $\tilde{T}_n$  is again represented as the maximum of a sum of independent random vectors in this case. We note that this argument is inapplicable to  $\|S_n\|_\infty$ .

### 3.3 Double wild bootstrap

As mentioned in the introduction, the lack of second-order accuracy in standard bootstrap methods is due to the fact that  $T_n$  is not asymptotically pivotal. If we knew the distribution function of  $T_n$ , say  $F_n$ , then  $F_n(T_n)$  would give an (exactly) pivotal statistic. Beran [6] suggested estimating  $F_n$  by the bootstrap distribution function  $\hat{F}_n(t) = P^*(T_n^* \leq t)$  and use  $\hat{F}_n(T_n)$  to construct critical values. This method is called *bootstrap prepivoting*. Note that  $\hat{F}_n$  can be computed by simulating the conditional law of  $T_n^*$  given the data. To estimate the law of  $\hat{F}_n(T_n)$ , we use the following nested double wild bootstrap procedure following [6]: Let  $v_1, \dots, v_n$  be i.i.d. variables independent of everything else and such that  $\mathbb{E}[v_1] = 0$  and  $\mathbb{E}[v_1^2] = 1$ . We define the wild bootstrap statistic of  $S_n^*$  as

$$S_n^{**} = \frac{1}{\sqrt{n}} \sum_{i=1}^n v_i (X_i^* - \bar{X}^*), \quad \text{where } X_i^* = w_i (X_i - \bar{X}), \quad \bar{X}^* = \frac{1}{n} \sum_{i=1}^n X_i^*.$$

Then define  $\hat{F}_n^{**}(t) = P^{**}(T_n^{**} \leq t)$  for  $t \in \mathbb{R}$ , where  $T_n^{**} := \max_{1 \leq j \leq d} S_{n,j}^{**}$  and  $P^{**}$  is the conditional probability given  $X_1, \dots, X_n, w_1, \dots, w_n$ . We regard  $\hat{F}_n^{**}(T_n^*)$  as a bootstrap version of  $\hat{F}_n(T_n)$  and estimate

the law of  $\hat{F}_n(T_n)$  by the conditional law of  $\hat{F}_n^*(T_n^*)$ . Formally, given a significance level  $\alpha \in (0, 1)$ , let  $\hat{\beta}_\alpha$  be the conditional  $(1 - \alpha)$ -quantile of  $\hat{F}_n^*(T_n^*)$  given the data. We expect that  $P(\hat{F}_n(T_n) \geq \hat{\beta}_\alpha) = P(T_n \geq \hat{c}_{\hat{\beta}_\alpha})$  would be close to  $\alpha$ . This is formally justified by the following theorem:

**Theorem 3.4** (Second-order accuracy of double bootstrap coverage probability). *Under the assumptions of Theorem 3.3, assume further that  $(w_1, w_1^2)$  has a Stein kernel  $\bar{\tau}^*$  such that  $\|\bar{\tau}^*(w_1, w_1^2)\|_\infty \leq b_w^4$ . Suppose also that  $v_1$  has a Stein kernel  $\tau^{**}$  and there exists a constant  $b_v \geq 1$  such that  $|v_1| \leq b_v$  and  $|\tau^{**}(v_1)| \leq b_v^2$ . Further, assume  $E[w_1^3] = E[v_1^3] = 1$ . Then, for any  $\varepsilon \in (0, 1/4)$ , there exists a constant  $C > 0$  depending only on  $\lambda, \varepsilon, b_w$  and  $b_v$  such that*

$$\sup_{2\varepsilon < \alpha < 1-2\varepsilon} \left| P\left(T_n \geq \hat{c}_{\hat{\beta}_\alpha}\right) - \alpha \right| \leq C \frac{\varsigma_d^4 \log^3(dn)}{\sigma_*^4 n} \log n. \quad (3.10)$$

The new assumption here is the existence of a bounded Stein kernel for  $(w_1, w_1^2)$ . This assumption is not problematic in practice because beta random variables still work:

**Proposition 3.2.** *Let  $\eta$  be a beta random variable and set  $w := (\eta - E[\eta])/\sqrt{\text{Var}[\eta]}$ . Then  $(w, w^2)$  has a bounded Stein kernel.*

**Remark 3.4** ( $p$ -value). One can easily check that  $T_n \geq \hat{c}_{\hat{\beta}_\alpha}$  is equivalent to  $P^*(\hat{p}_n^* \leq \hat{p}_n) \leq \alpha$ , where  $\hat{p}_n := 1 - \hat{F}_n(T_n)$  and  $\hat{p}_n^* := 1 - \hat{F}_n^*(T_n^*)$  are the  $p$ -values of the first and second level bootstraps, respectively. Hence the  $p$ -value of the double bootstrap method is  $P^*(\hat{p}_n^* \leq \hat{p}_n)$ .

#### 4 Simulation study

This section conducts a small Monte Carlo study to supplement our theoretical findings. We adopt the same simulation design as [24]: We set  $n = 200, d = 400$  and generate the data from a Gaussian copula model, i.e.  $X_1, \dots, X_n$  are i.i.d. with the same law as  $U - E[U]$ , where  $U$  is defined as in Example 2.2. The marginal distributions  $\mu_j$  are the gamma distribution with shape parameter 1 and unit scale. As the parameter matrix  $R$ , we consider two designs: (I)  $R = \rho \mathbf{1}_d^{\otimes 2} + (1 - \rho)I_d$  and (II)  $R = (\rho^{|j-k|})_{1 \leq j, k \leq d}$ . Here, the parameter  $\rho$  is varied as  $\rho \in \{0.2, 0.8\}$ . We compute the rejection rates  $P(T_n \geq \hat{c})$  and  $P(\|S_n\|_\infty \geq \hat{c})$  at the 10% significance level based on 20,000 Monte Carlo iterations, where  $\hat{c}$  is an estimated 90% quantile of the corresponding statistic using various bootstrap methods. In addition, to assess the performance when the skewness of the data is zero, we also consider the case that  $X_i \stackrel{d}{=} U - U'$ , where  $U'$  is an independent copy of  $U$ . To keep the marginal kurtosis at the same level, we change the shape parameter of the gamma distribution to 0.5 in this case.

For the bootstrap methods, we consider the empirical bootstrap (EB), wild bootstrap and double wild bootstrap (DB) methods. For the wild bootstrap, we consider the following 4 types of weight variables:

**GB**  $w_1$  is a standard normal variable.

**MB**  $w_1$  follows Mammen's two point distribution [46]:

$$P\left(w_1 = \frac{\sqrt{5} + 1}{2}\right) = 1 - P\left(w_1 = -\frac{\sqrt{5} - 1}{2}\right) = \frac{\sqrt{5} - 1}{2\sqrt{5}}. \quad (4.1)$$

**RB**  $w_1$  is a Rademacher variable:  $P(w_1 = \pm 1) = 1/2$ .

**BB**  $w_1$  follows the standardized beta distribution with parameters given by (2.12) with  $\nu = 0.1$ .

The double wild bootstrap is implemented with both  $w_1$  and  $v_1$  generated from the standardized beta distribution with parameters given by (2.12) with  $\nu = 0.1$ . Note that our theoretical results are applicable to only GB, BB and DB. We include EB, MB and RB in our assessment because they are commonly used in the literature. The number of bootstrap replications is set to 499 for the first-level bootstrap and 99 for the second-level bootstrap in DB.

We summarize the simulation results in Tables 1 and 2. First, Table 1 reports empirical rejection rates at the 10% level when the laws of  $X_i$  are asymmetric. We find that the difference of performances between GB and BB is largely in line with our Theorem 2.1 except for  $T_n$  in Design (I) with  $\rho = 0.8$ : BB performs better than GB for  $T_n$ , while they perform similarly for  $\|S_n\|_\infty$ . For  $T_n$  in Design (I) with  $\rho = 0.8$ , GB outperforms BB. This phenomenon might be explained as follows: In Design (I), for  $G \sim N(0, R)$ ,  $G^\vee$  has the same law as  $\rho\zeta + (1-\rho)\tilde{G}^\vee$ , where  $\zeta \sim N(0, 1)$  and  $\tilde{G} \sim N(0, I_d)$  are independent. Since  $\text{Var}[\tilde{G}^\vee] = O(1/\sqrt{\log d})$ ,  $G^\vee$  is asymptotically normal as  $d \rightarrow \infty$  in this case. This perhaps imply that  $T_n$  behaves as in the classical setting when  $d$  and  $\rho$  are large. Then, it is known that normal approximation typically outperforms bootstrap approximation without studentization in terms of coverage errors; see [43, Section 3] for details. Turning to the performance of DB, it tends to over-reject but outperforms GB and BB in Design (I). The latter is expected since our theory reveals that DB are second-order accurate while GB and BB are generally not. In Design (II), the performances of DB and BB are comparable. This would be because  $\Sigma$  has no large eigenvalues; see Corollary 3.1.

Next, Table 2 reports empirical rejection rates at the 10% level when the laws of  $X_i$  are symmetric. Recall that our Theorem 3.3 implies that both GB and BB are second-order accurate (at least) for  $T_n$  in this case. Reflecting this fact, GB clearly performs better than the asymmetric case for  $T_n$ . The performance of BB is improved in Design (I) but not in Design (II). The latter would be due to the same reasoning as above, i.e. BB is second-order accurate in Design (II) even when the skewness is not zero by Corollary 3.1. By contrast, the performance of DB is not improved. This is not surprising because DB is already second-order accurate in the asymmetric case and the zero skewness condition would not contribute to its performance. When comparing GB and BB, BB still outperforms GB. This may be due to an effect of kurtosis, but we will need higher-order asymptotic expansions for the formal discussion and leave it to future work.

Finally, we briefly discuss the performances of EB, MB and RB. First, EB tends to under-reject and its performance is not pronounced compared to other methods. In fact, we can observe similar phenomena in the simulation results of [21, 24]. Formally, this does not contradict our theory because we have no valid Edgeworth expansion for EB in high-dimensions, although it is unclear whether this is an artifact of our proof strategy. Next, although MB is not covered by our theory, its performance is similar to BB. This is perhaps explained by the fact that their weights are very close numerically. Third, RB performs remarkably well in the symmetric case. This is already observed in the simulation study of [21] who explain this phenomenon by their Theorem 2.3. Another possible explanation is the match of higher moments, but we have no formal theoretical result for this so far.

Table 1: Rejection rate at the 10% level (Asymmetric case)

$\rho$		EB	GB	MB	RB	BB	DB
(I) $R_{jk} = \rho + (1 - \rho)1_{\{j=k\}}$							
0.2	$T_n$	0.061	0.124	0.080	0.155	0.078	0.114
	$\ S_n\ _\infty$	0.060	0.073	0.082	0.107	0.079	0.114
0.8	$T_n$	0.071	0.090	0.072	0.093	0.071	0.101
	$\ S_n\ _\infty$	0.091	0.091	0.097	0.099	0.097	0.100
(II) $R_{jk} = \rho^{ j-k }$							
0.2	$T_n$	0.065	0.146	0.092	0.195	0.091	0.117
	$\ S_n\ _\infty$	0.061	0.083	0.086	0.123	0.085	0.117
0.8	$T_n$	0.069	0.139	0.089	0.177	0.088	0.113
	$\ S_n\ _\infty$	0.062	0.079	0.084	0.113	0.083	0.112

Table 2: Rejection rate at the 10% level (Symmetric case)

$\rho$		EB	GB	MB	RB	BB	DB
(I) $R_{jk} = \rho + (1 - \rho)1_{\{j=k\}}$							
0.2	$T_n$	0.065	0.076	0.083	0.100	0.082	0.114
	$\ S_n\ _\infty$	0.058	0.067	0.082	0.099	0.082	0.113
0.8	$T_n$	0.089	0.092	0.091	0.096	0.091	0.105
	$\ S_n\ _\infty$	0.082	0.084	0.088	0.093	0.088	0.093
(II) $R_{jk} = \rho^{ j-k }$							
0.2	$T_n$	0.062	0.071	0.085	0.101	0.084	0.114
	$\ S_n\ _\infty$	0.056	0.068	0.084	0.104	0.083	0.119
0.8	$T_n$	0.067	0.076	0.088	0.100	0.086	0.109
	$\ S_n\ _\infty$	0.060	0.070	0.087	0.105	0.086	0.118

## 5 Proofs for Section 2

We use the following notation in the remainder of the paper: For two random variables  $\xi$  and  $\eta$ , we write  $\xi \lesssim \eta$  or  $\eta \gtrsim \xi$  if there exists a *universal* constant  $C > 0$  such that  $\xi \leq C\eta$ . Also, given real numbers  $\theta_1, \dots, \theta_m$ , we use  $C_{\theta_1, \dots, \theta_m}$  to denote positive constants, which depend only on  $\theta_1, \dots, \theta_m$  and may be different in different expressions.

### 5.1 Proof of Theorem 2.1

Without loss of generality, we may assume

$$\frac{b^2 \log d}{\sigma_*^2 \sqrt{n}} \leq \frac{1}{2}. \quad (5.1)$$

Since  $\sigma_*^2 \leq n^{-1} \sum_{i=1}^n \mathbb{E}[X_{i1}^2] \leq 2b^2$ , this particularly yields  $\log^2 d \leq n$ .

Let us prove (2.4). Let  $\mathcal{E}_n$  be the event on which (2.2) holds. We have  $P(\mathcal{E}_n) \geq 1 - \delta_n$  by assumption.

Also, by (2.1),

$$\sup_{t \in \mathbb{R}} P(T_n = t) \leq \Delta_n. \quad (5.2)$$

Next, by Lemma E.4, there exists a universal constant  $C_1 > 0$  such that

$$\sup_{A \in \mathcal{R}} \left\| \int_A \nabla^2 \phi_\Sigma(z) dz \right\|_1 \leq C_1 \frac{\log d}{\sigma_*^2}, \quad \sup_{A \in \mathcal{R}} \left\| \int_A \nabla^3 \phi_\Sigma(z) dz \right\|_1 \leq C_1 \frac{\log^{3/2} d}{\sigma_*^3}.$$

Also, by Lemma E.10, there exists a universal constant  $C_2 > 0$  such that

$$P \left( |\langle \overline{X^2} - \Sigma, V_1 \rangle| > C_2 \|V_1\|_1 b^2 \sqrt{\frac{\log n}{n}} \right) \leq \frac{1}{n}$$

for any  $V_1 \in (\mathbb{R}^d)^{\otimes 2}$  and

$$P \left( \left| \frac{1}{\sqrt{n}} \langle \overline{X^3}, V_2 \rangle - \mathbb{E}[\langle \overline{X^3}, V_2 \rangle] \right| > C_2 \|V_2\|_1 b^3 \frac{\sqrt{\log(dn)}}{n} \right) \leq \frac{1}{n}$$

for any  $V_2 \in (\mathbb{R}^d)^{\otimes 3}$ . Now we set

$$\Delta'_n := C_1 C_2 \left( b^2 \frac{\log d}{2\sigma_*^2} \sqrt{\frac{\log n}{n}} + |\gamma| b^3 \frac{\log^{3/2} d \sqrt{\log(dn)}}{6n\sigma_*^3} \right).$$

For every  $A \in \mathcal{R}$ , recall that we have the decomposition (2.6). Therefore, setting

$$\mathcal{E}_n(A) := \left\{ \left| \int_A \{\hat{p}_{n,\gamma}(z) - p_n(z)\} dz \right| \leq \Delta'_n \right\},$$

we have  $P(\mathcal{E}_n(A)) \geq 1 - 2/n$ , provided that  $\gamma = 1$  or  $\mathbb{E}[\overline{X^3}] = 0$ .

Set  $\bar{\Delta}_n := \Delta_n + \Delta'_n$ . Also, recall that  $c_p$  is the  $p$ -quantile of  $T_n$  for  $p \in (0, 1)$  and thus  $P(T_n < c_p) \leq p \leq P(T_n \leq c_p)$ . Then, if  $2\bar{\Delta}_n < \alpha$ , we have on  $\mathcal{E}_n$

$$1 - \alpha + 2\bar{\Delta}_n \leq P(T_n \leq c_{1-\alpha+2\bar{\Delta}_n}) \leq P^*(T_n^* \leq c_{1-\alpha+2\bar{\Delta}_n}) + 2\Delta_n + \left| \int_{A_1} \{\hat{p}_{n,\gamma}(z) - p_n(z)\} dz \right|,$$

where  $A_1 := \{z \in \mathbb{R}^d : z^\vee \leq c_{1-\alpha+2\bar{\Delta}_n}\}$ . Thus, on  $\mathcal{E}_n \cap \mathcal{E}_n(A_1)$ ,

$$1 - \alpha \leq -2\bar{\Delta}_n + P^*(T_n^* \leq c_{1-\alpha+2\bar{\Delta}_n}) + 2\Delta_n + \Delta'_n \leq P^*(T_n^* \leq c_{1-\alpha+2\bar{\Delta}_n}).$$

This implies  $\hat{c}_{1-\alpha} \leq c_{1-\alpha+2\bar{\Delta}_n}$  on  $\mathcal{E}_n \cap \mathcal{E}_n(A_1)$ . Hence

$$P(T_n \geq \hat{c}_{1-\alpha}) \geq P(T_n \geq c_{1-\alpha+2\bar{\Delta}_n}) - \delta_n - \frac{2}{n} \geq \alpha - 2\bar{\Delta}_n - \delta_n - \frac{2}{n}.$$

Thus  $\alpha - P(T_n \geq \hat{c}_{1-\alpha}) \leq 2\bar{\Delta}_n + \delta_n + 2/n$ . Also, this bound trivially holds if  $2\bar{\Delta}_n \geq \alpha$ . Meanwhile, if  $3\bar{\Delta}_n < 1 - \alpha$ , we have on  $\mathcal{E}_n$

$$P^*(T_n^* \leq c_{1-\alpha-3\bar{\Delta}_n}) \leq P(T_n \leq c_{1-\alpha-3\bar{\Delta}_n}) + 2\Delta_n + \left| \int_{A_2} \{\hat{p}_{n,\gamma}(z) - p_n(z)\} dz \right|,$$

where  $A_2 := \{z \in \mathbb{R}^d : z^\vee \leq c_{1-\alpha-3\bar{\Delta}_n}\}$ . Observe that

$$P(T_n \leq c_{1-\alpha-3\bar{\Delta}_n}) \leq P(T_n < c_{1-\alpha-3\bar{\Delta}_n}) + \Delta_n \leq 1 - \alpha - 2\bar{\Delta}_n,$$

where we used (5.2) for the first inequality. Hence, on  $\mathcal{E}_n \cap \mathcal{E}_n(A_2)$ ,

$$P^*(T_n^* \leq c_{1-\alpha-3\bar{\Delta}_n}) \leq 1 - \alpha - \Delta'_n < 1 - \alpha.$$

Thus  $\hat{c}_{1-\alpha} > c_{1-\alpha-3\bar{\Delta}_n}$  on  $\mathcal{E}_n \cap \mathcal{E}_n(A_2)$ . Hence

$$P(T_n \geq \hat{c}_{1-\alpha}) \leq P(T_n > c_{1-\alpha-3\bar{\Delta}_n}) + \delta_n + \frac{2}{n} \leq \alpha + 3\bar{\Delta}_n + \delta_n + \frac{2}{n}$$

Thus  $P(T_n \geq \hat{c}_{1-\alpha}) - \alpha \leq 3\bar{\Delta}_n + \delta_n + 2/n$  and this trivially holds if  $3\bar{\Delta}_n \geq 1 - \alpha$ . Finally, using (5.1) and  $\sigma_*^2 \leq 2b^2$ , we can easily check  $\Delta'_n \lesssim (1 + |\gamma|) \frac{b^2 \log d}{\sigma_*^2 \sqrt{n}} \sqrt{\log n}$ . All together, we obtain (2.4).

To prove (2.5), observe that  $\int_A \nabla^3 \phi_\Sigma(z) dz = 0$  when  $A$  is of the form  $A = [-c, c]^d$  for some  $c > 0$  because  $\nabla^3 \phi_\Sigma$  is an odd function. Hence, in this case we have  $P(\mathcal{E}_n(A)) \geq 1 - 1/n$  with the second term of  $\Delta'_n$  being 0, regardless of the values of  $\gamma$  and  $E[\bar{X}^3]$ . Thus, (2.5) follows by a similar argument to the above regardless of the values of  $\gamma$  and  $E[\bar{X}^3]$ .  $\square$

## 5.2 Proofs of Theorems 2.2 and 2.3

The proofs are based on the following two abstract error bounds for high-dimensional Edgeworth expansion (the latter is used for the Gaussian wild bootstrap):

**Theorem 5.1** (Error bound for high-dimensional Edgeworth expansion via Stein kernel). *Let  $\xi_1, \dots, \xi_n$  be independent random vectors in  $\mathbb{R}^d$  with mean 0 and finite variance. Set  $W := \sum_{i=1}^n \xi_i$  and  $\Sigma_W := \text{Cov}[W]$ . Suppose that  $\xi_i$  has a Stein kernel  $\tau_i$  and satisfies  $E[\|\xi_i\|_\infty^5] + E[\|\tau_i(\xi_i)\|_\infty^{5/2}] < \infty$  for all  $i = 1, \dots, n$ . Set  $T = \sum_{i=1}^n \tau_i(\xi_i)$ ,  $\bar{T} = T - \Sigma$  and*

$$p_W(z) = \phi_\Sigma(z) + \frac{1}{2} \langle \Sigma_W - \Sigma, \nabla^2 \phi_\Sigma(z) \rangle - \frac{1}{6} \langle E[W^{\otimes 3}], \nabla^3 \phi_\Sigma(z) \rangle, \quad z \in \mathbb{R}^d. \quad (5.3)$$

Then, there exists a universal constant  $C > 0$  such that for any  $t \in (0, 1/2]$ ,

$$\begin{aligned} & \sup_{A \in \mathcal{R}} \left| P(W \in A) - \int_A p_W(z) dz \right| \\ & \leq C |\log t| \frac{\log^2 d}{\sigma_*^4} \left( E \|\bar{T}\|_\infty^2 + E \left\| \sum_{i=1}^n \tau_i(\xi_i)^{\otimes 2} \right\|_\infty + E \left\| \sum_{i=1}^n \xi_i^{\otimes 4} \right\|_\infty \right) \\ & \quad + C \frac{\log^{5/2} d}{\sigma_*^5} \left( E \left\| \bar{T} \otimes \sum_{i=1}^n \xi_i^{\otimes 3} \right\|_\infty + E \left\| \sum_{i=1}^n \xi_i^{\otimes 3} \otimes \tau_i(\xi_i) \right\|_\infty \right) \\ & \quad + C \bar{\sigma} \sqrt{t} \left( \frac{\log d}{\underline{\sigma}} + \frac{\log^2 d}{\sigma_*^3} \|\Sigma_W - \Sigma\|_\infty + \frac{\log^{5/2} d}{\sigma_*^4} \left\| \sum_{i=1}^n E[\xi_i^{\otimes 3}] \right\|_\infty \right). \end{aligned} \quad (5.4)$$

**Theorem 5.2** (Refined Gaussian comparison inequality). *Let  $W$  be a centered Gaussian vector in  $\mathbb{R}^d$  with covariance matrix  $\Sigma_W$ . Then, there exists a universal constant  $C > 0$  such that for any  $t \in (0, 1/2]$ ,*

$$\sup_{A \in \mathcal{R}} \left| P(W \in A) - \int_A p_W(z) dz \right|$$

$$\leq C |\log t| \frac{\log^2 d}{\sigma_*^4} \|\Sigma_W - \Sigma\|_\infty^2 + C \bar{\sigma} \sqrt{t} \left( \frac{\log d}{\underline{\sigma}} + \frac{\log^2 d}{\sigma_*^3} \|\Sigma_W - \Sigma\|_\infty \right),$$

where  $p_W$  is defined by (5.3); note that  $\mathbb{E}[W^{\otimes 3}] = 0$  in the present case.

First we prove Theorems 2.2 and 2.3 using the above results. Below we will frequently use the following identity without reference: For any  $x_1, \dots, x_n \in \mathbb{R}^d$  and  $r \in \mathbb{N}$ ,

$$\left\| \sum_{i=1}^n x_i^{\otimes (2r)} \right\|_\infty = \max_{1 \leq j \leq d} \sum_{i=1}^n x_{ij}^{2r}.$$

This follows from the AM-GM inequality.

**Proof of Theorem 2.2.** Observe that for every  $i = 1, \dots, n$ ,  $(\tau_i^X + (\tau_i^X)^\top)/2$  is a Stein kernel for  $X_i$  and  $\max_{j,k} \|(\tau_{i,jk}^X(X_i) + \tau_{i,kj}^X(X_i))/2\|_{\psi_{1/2}} \lesssim b^2$ . Therefore, we may assume  $\tau_{i,jk}^X = \tau_{i,kj}^X$  for all  $j, k \in \{1, \dots, d\}$  without loss of generality. This particularly implies  $\mathbb{E}[\tau_i(X_i)] = \mathbb{E}[X_i^{\otimes 2}]$ .

We apply Theorem 5.1 with  $\xi_i = X_i/\sqrt{n}$ . Observe that  $\Sigma_W = \Sigma$  and that  $\xi_i$  has a Stein kernel  $\tau_i$  satisfying  $\tau_i(\xi_i) = \tau_i^X(X_i)/n$ .

Let us bound the quantities appearing in the right hand side of (5.4). First, noting that  $\mathbb{E}[\tau_i(\xi_i)] = \mathbb{E}[\xi_i^{\otimes 2}]$ , we have

$$\bar{T} = \sum_{i=1}^n (\tau_i(\xi_i) - \mathbb{E}[\tau_i(\xi_i)]).$$

Thus, by Lemma E.9 with  $K = b^2/n$ ,  $\alpha = 1/2$  and  $r = 2$ , we obtain

$$\mathbb{E}[\|\bar{T}\|_\infty^2] \lesssim \frac{b^4}{n^2} \left( \sqrt{n \log d} + (\log d)^2 \right)^2 \lesssim \frac{b^4 \log d}{n}, \quad (5.5)$$

where the second inequality follows from  $\log^3 d \leq n$ . Next, by Lemma E.7 and Lemma E.9 with  $K = b^2/n$ ,  $\alpha = 1/4$  and  $r = 1$ ,

$$\mathbb{E} \left\| \sum_{i=1}^n \{ \tau_i(\xi_i)^{\otimes 2} - \mathbb{E}[\tau_i(\xi_i)^{\otimes 2}] \} \right\|_\infty \lesssim \frac{b^4}{n^2} \left( \sqrt{n \log d} + (\log d)^4 \right).$$

Therefore,

$$\begin{aligned} \mathbb{E} \left\| \sum_{i=1}^n \tau_i(\xi_i)^{\otimes 2} \right\|_\infty &\leq \left\| \sum_{i=1}^n \mathbb{E}[\tau_i(\xi_i)^{\otimes 2}] \right\|_\infty + \mathbb{E} \left\| \sum_{i=1}^n \{ \tau_i(\xi_i)^{\otimes 2} - \mathbb{E}[\tau_i(\xi_i)^{\otimes 2}] \} \right\|_\infty \\ &\lesssim \frac{b^4}{n} + \frac{b^4 \sqrt{\log d}}{n^{3/2}} + \frac{b^4 (\log d)^4}{n^2} \lesssim \frac{b^4}{n} + \frac{b^4 (\log d)^4}{n^2}, \end{aligned} \quad (5.6)$$

where the last inequality follows from  $\sqrt{\log d} \leq n^{1/6} \leq \sqrt{n}$ . Similarly, we can show that

$$\mathbb{E} \left\| \sum_{i=1}^n \xi_i^{\otimes 4} \right\|_\infty \lesssim \frac{b^4}{n} + \frac{b^4 (\log d)^4}{n^2}, \quad (5.7)$$

$$\mathbb{E} \left\| \sum_{i=1}^n \xi_i^{\otimes 3} \otimes \tau_i(\xi_i) \right\|_\infty \lesssim \frac{b^5}{n^{3/2}} + \frac{b^5 (\log d)^5}{n^{5/2}}. \quad (5.8)$$

In addition, by the Schwarz inequality,

$$\mathbb{E} \left\| \bar{T} \otimes \sum_{i=1}^n \xi_i^{\otimes 3} \right\|_{\infty} \leq \sqrt{\mathbb{E} \|\bar{T}\|_{\infty}^2} \sqrt{\mathbb{E} \left\| \sum_{i=1}^n \xi_i^{\otimes 3} \right\|_{\infty}^2}.$$

Similarly to the proof of (5.6), we can prove

$$\mathbb{E} \left\| \sum_{i=1}^n \xi_i^{\otimes 3} \right\|_{\infty}^2 \lesssim \left( \frac{b^3}{\sqrt{n}} + \frac{b^3(\log d)^3}{n^{3/2}} \right)^2 \lesssim \frac{b^6}{n},$$

where the second inequality follows by the assumption  $\log^3 d \leq n$ . Combining this with (5.5) gives

$$\mathbb{E} \left\| \bar{T} \otimes \sum_{i=1}^n \xi_i^{\otimes 3} \right\|_{\infty} \lesssim \frac{b^5 \sqrt{\log d}}{n}. \quad (5.9)$$

Now, by (5.5)–(5.7),

$$\begin{aligned} & \frac{\log^2 d}{\sigma_*^4} \left( \mathbb{E} \|\bar{T}\|_{\infty}^2 + \mathbb{E} \left\| \sum_{i=1}^n \tau_i(\xi_i)^{\otimes 2} \right\|_{\infty}^2 + \mathbb{E} \left\| \sum_{i=1}^n \xi_i^{\otimes 4} \right\|_{\infty} \right) \\ & \lesssim \frac{\log^2 d}{\sigma_*^4} \left( \frac{b^4 \log d}{n} + \frac{b^4(\log d)^4}{n^2} \right) \lesssim \frac{b^4 \log^3 d}{\sigma_*^4 n}, \end{aligned}$$

where the second inequality follows by the assumption  $\log^3 d \leq n$ . Also, by (5.8) and (5.9),

$$\begin{aligned} & \frac{\log^{5/2} d}{\sigma_*^5} \left( \mathbb{E} \left\| \bar{T} \otimes \sum_{i=1}^n \xi_i^{\otimes 3} \right\|_{\infty} + \mathbb{E} \left\| \sum_{i=1}^n \xi_i^{\otimes 3} \otimes \tau_i(\xi_i) \right\|_{\infty} \right) \\ & \lesssim \frac{\log^{5/2} d}{\sigma_*^5} \left( \frac{b^5 \sqrt{\log d}}{n} + \frac{b^5(\log d)^5}{n^{5/2}} \right) \lesssim \frac{b^5 \log^3 d}{\sigma_*^5 n}, \end{aligned}$$

where we used the assumption  $\log^3 d \leq n$  in the second inequality. All together, we obtain by Theorem 5.1

$$\begin{aligned} & \sup_{A \in \mathcal{R}} \left| P(S_n \in A) - \int_A p_n(z) dz \right| \\ & \lesssim |\log t| \frac{b^4 \log^3 d}{\sigma_*^4 n} + \frac{b^5 \log^3 d}{\sigma_*^5 n} + \bar{\sigma} \sqrt{t} \left( \frac{\log d}{\underline{\sigma}} + \frac{\log^{5/2} d}{\sigma_*^4} \left\| \sum_{i=1}^n \mathbb{E}[\xi_i^{\otimes 3}] \right\|_{\infty} \right) \end{aligned}$$

for any  $t \in (0, 1/2]$ . With  $t = 1/n^2$ , we obtain

$$\bar{\sigma} \sqrt{t} \left( \frac{\log d}{\underline{\sigma}} + \frac{\log^{5/2} d}{\sigma_*^4} \left\| \sum_{i=1}^n \mathbb{E}[\xi_i^{\otimes 3}] \right\|_{\infty} \right) \lesssim \frac{b \log d}{\sigma_* n} + \frac{b^4 \log^{5/2} d}{\sigma_*^4 n^{3/2}} \lesssim \frac{b^5 \log^3 d}{\sigma_*^5 n}.$$

Consequently, we obtain the desired result.  $\square$

**Proof of Theorem 2.3.** First consider Case (i). Set  $\tilde{X}_i := X_i - \bar{X}$  for  $i = 1, \dots, n$ . We apply Theorem 5.1 with  $\xi_i = w_i \tilde{X}_i / \sqrt{n}$  and  $t = 1/n^3$  conditional on the data. Note that conditional on the data,  $\xi_i$  has a Stein kernel  $\tau_i$  satisfying  $\tau_i(\xi_i) = \tau_i^*(w_i) \tilde{X}_i^{\otimes 2} / n$  by Lemma C.1. Therefore, we have

$$\sup_{A \in \mathcal{R}} \left| P^*(S_n^* \in A) - \int_A \tilde{p}_{n,\gamma}(z) dz \right|$$

$$\begin{aligned}
&\lesssim (\log n) \frac{\log^2 d}{\sigma_*^4} \left( \mathbb{E}^* \left[ \|\bar{T}^*\|_\infty^2 \right] + \mathbb{E}^* \left[ \left\| \frac{1}{n^2} \sum_{i=1}^n \tau_i^*(w_i)^2 \tilde{X}_i^{\otimes 4} \right\|_\infty \right] + \mathbb{E}^* \left[ \left\| \frac{1}{n^2} \sum_{i=1}^n w_i^4 \tilde{X}_i^{\otimes 4} \right\|_\infty \right] \right) \\
&+ \frac{\log^{5/2} d}{\sigma_*^5} \left( \mathbb{E}^* \left[ \left\| \bar{T}^* \otimes \frac{1}{n^{3/2}} \sum_{i=1}^n w_i^3 \tilde{X}_i^{\otimes 3} \right\|_\infty \right] + \mathbb{E}^* \left[ \left\| \frac{1}{n^{5/2}} \sum_{i=1}^n w_i^3 \tau_i^*(w_i) \tilde{X}_i^{\otimes 5} \right\|_\infty \right] \right) \\
&+ \frac{\bar{\sigma}}{n^{3/2}} \left( \frac{\log d}{\underline{\sigma}} + \frac{\log^2 d}{\sigma_*^3} \|\widehat{\Sigma}_n - \Sigma\|_\infty + \frac{\log^{5/2} d}{\sigma_*^4} \left\| \frac{1}{n^{3/2}} \sum_{i=1}^n \mathbb{E}[w_i^3] \tilde{X}_i^{\otimes 3} \right\|_\infty \right), \tag{5.10}
\end{aligned}$$

where

$$\tilde{p}_{n,\gamma}(z) = \phi_\Sigma(z) + \frac{1}{2} \langle \widehat{\Sigma}_n - \Sigma, \nabla^2 \phi_\Sigma(z) \rangle - \frac{\gamma}{6n^{3/2}} \sum_{i=1}^n \langle (X_i - \bar{X})^{\otimes 3}, \nabla^3 \phi_\Sigma(z) \rangle$$

and

$$\bar{T}^* := \frac{1}{n} \sum_{i=1}^n \tau_i^*(w_i) \tilde{X}_i^{\otimes 2} - \Sigma.$$

Next, by Lemmas E.9 and E.11, there exists a universal constant  $c$  such that the event

$$\begin{aligned}
\mathcal{E}_n &:= \bigcap_{r=1}^2 \left\{ \left\| \frac{1}{n} \sum_{i=1}^n (X_i^{\otimes r} - \mathbb{E}[X_i^{\otimes r}]) \right\|_\infty \leq cb^r \left( \sqrt{\frac{\log(dn)}{n}} + \frac{\log^r(dn)}{n} \right) \right\} \\
&\cap \bigcap_{r=3}^5 \left\{ \max_{1 \leq j \leq d} \left| \frac{1}{n} \sum_{i=1}^n (|X_{ij}|^r - \mathbb{E}[|X_{ij}|^r]) \right| \leq cb^r \left( \sqrt{\frac{\log(dn)}{n}} + \frac{\log^r(dn)}{n} \right) \right\} \\
&\cap \left\{ \max_{1 \leq j, k \leq d} \frac{1}{n} \sum_{i=1}^n |X_{ij}| |X_{ik}| \mathbb{1}_{\{|X_{ij}| \vee |X_{ik}| > 2b \log n\}} \leq cb^2 \left( \sqrt{\frac{\log(dn)}{n}} + \frac{\log^2(dn)}{n} \right) \right\} \\
&\cap \left\{ \max_{1 \leq j, k \leq d} \frac{1}{n} \sum_{i=1}^n X_{ij}^2 X_{ik}^2 \mathbb{1}_{\{|X_{ij}| \vee |X_{ik}| \leq 2b \log n\}} \leq cb^4 \left( 1 + \frac{\log(dn) \log^4 n}{n} \right) \right\} \tag{5.11}
\end{aligned}$$

occurs with probability at least  $1 - 1/n$ . Recall that  $\log^3 d \leq n$  by assumption. Hence, on  $\mathcal{E}_n$ , we have

$$\|\bar{X}\|_\infty \lesssim b \sqrt{\frac{\log(dn)}{n}} \tag{5.12}$$

and

$$\left\| \frac{1}{n} \sum_{i=1}^n X_i^{\otimes 2} - \Sigma \right\|_\infty \lesssim b^2 \sqrt{\frac{\log(dn)}{n}}. \tag{5.13}$$

Thus, on  $\mathcal{E}_n$ ,

$$\|\widehat{\Sigma}_n - \Sigma\|_\infty \leq \left\| \frac{1}{n} \sum_{i=1}^n X_i^{\otimes 2} - \Sigma \right\|_\infty + \|\bar{X}\|_\infty^2 \lesssim b^2 \sqrt{\frac{\log(dn)}{n}}. \tag{5.14}$$

Meanwhile, for every  $r \in \{2, 3, 4, 5\}$ , we have on  $\mathcal{E}_n$

$$\left\| \frac{1}{n} \sum_{i=1}^n X_i^{\otimes r} \right\|_\infty \leq \max_{1 \leq j \leq d} \frac{1}{n} \sum_{i=1}^n |X_{ij}|^r \lesssim b^r \left( 1 + \frac{\log^r(dn)}{n} \right). \tag{5.15}$$

Combining this bound with (5.12), we have on  $\mathcal{E}_n$

$$\max_{1 \leq j \leq d} \frac{1}{n} \sum_{i=1}^n |\tilde{X}_{ij}|^r \lesssim b^r \left( 1 + \frac{\log^r(dn)}{n} \right). \tag{5.16}$$

Further, by (5.12), (5.15), the construction of  $\mathcal{E}_n$  and  $\log^3 d \leq n$ , we have on  $\mathcal{E}_n$

$$\max_{1 \leq j, k \leq d} \frac{1}{n} \sum_{i=1}^n |\tilde{X}_{ij} \tilde{X}_{ik}| 1_{\{|X_{ij}| \vee |X_{ik}| > 2b \log n\}} \lesssim b^2 \left( \sqrt{\frac{\log d}{n}} + \frac{\log^2(dn)}{n} \right), \quad (5.17)$$

and

$$\max_{1 \leq j, k \leq d} \frac{1}{n} \sum_{i=1}^n \tilde{X}_{ij}^2 \tilde{X}_{ik}^2 1_{\{|X_{ij}| \vee |X_{ik}| \leq 2b \log n\}} \lesssim b^4. \quad (5.18)$$

Now we bound the right hand side of (5.10) on the event  $\mathcal{E}_n$ . First, we have

$$\mathbb{E}^* [\|\bar{T}^*\|_\infty^2] \leq 2 \mathbb{E}^* \left[ \max_{1 \leq j, k \leq d} R_{jk}^2 \right] + 2 \|\hat{\Sigma}_n - \Sigma\|_\infty^2. \quad (5.19)$$

where  $R_{jk} := n^{-1} \sum_{i=1}^n \{\tau_i^*(w_i) - 1\} \tilde{X}_{ij} \tilde{X}_{ik}$ . We decompose  $R_{jk}$  as

$$\begin{aligned} R_{jk} &= \frac{1}{n} \sum_{i=1}^n \{\tau_i^*(w_i) - 1\} \tilde{X}_{ij} \tilde{X}_{ik} \left( 1_{\{|X_{ij}| \vee |X_{ik}| > 2b \log n\}} + 1_{\{|X_{ij}| \vee |X_{ik}| \leq 2b \log n\}} \right) \\ &=: R_{1,jk} + R_{2,jk}. \end{aligned}$$

Since

$$\max_{1 \leq j, k \leq d} |R_{1,jk}| \leq (b_w^2 + 1) \max_{1 \leq j, k \leq d} \frac{1}{n} \sum_{i=1}^n |\tilde{X}_{ij} \tilde{X}_{ik}| 1_{\{|X_{ij}| \vee |X_{ik}| > 2b \log n\}},$$

we have on  $\mathcal{E}_n$

$$\max_{1 \leq j, k \leq d} |R_{1,jk}| \lesssim b_w^2 b^2 \left( \sqrt{\frac{\log d}{n}} + \frac{\log^2(dn)}{n} \right) \quad (5.20)$$

by (5.17). Meanwhile, by Nemirovski's inequality (cf. Lemma 14.24 in [12]),

$$\mathbb{E}^* \left[ \max_{1 \leq j, k \leq d} R_{2,jk}^2 \right] \lesssim \frac{b_w^4 \log d}{n} \max_{1 \leq j, k \leq d} \frac{1}{n} \sum_{i=1}^n \tilde{X}_{ij}^2 \tilde{X}_{ik}^2 1_{\{|X_{ij}| \vee |X_{ik}| \leq 2b \log n\}}.$$

Hence we have on  $\mathcal{E}_n$

$$\mathbb{E}^* \left[ \max_{1 \leq j, k \leq d} R_{2,jk}^2 \right] \lesssim \frac{b_w^4 b^4 \log d}{n} \quad (5.21)$$

by (5.18). Combining (5.19)–(5.21) with (5.14), we obtain

$$\mathbb{E}^* [\|\bar{T}^*\|_\infty^2] \lesssim b_w^4 b^4 \left( \frac{\log(dn)}{n} + \frac{\log^4(dn)}{n^2} \right). \quad (5.22)$$

Consequently, we have

$$\frac{\log^2 d}{\sigma_*^4} \mathbb{E}^* [\|\bar{T}^*\|_\infty^2] \lesssim \frac{b_w^4 b^4}{\sigma_*^4} \left( \frac{\log^3(dn)}{n} + \frac{\log^6(dn)}{n^2} \right) \lesssim \frac{b_w^4 b^4 \log^3(dn)}{\sigma_*^4 n}, \quad (5.23)$$

where we used the assumption  $\log^3 d \leq n$  for the last inequality. Next, we have

$$\left\| \frac{1}{n^2} \sum_{i=1}^n \tau_i^*(w_i)^2 \tilde{X}_i^{\otimes 4} \right\|_\infty \leq b_w^4 \max_{j,k,l,m} \frac{1}{n^2} \sum_{i=1}^n |\tilde{X}_{ij} \tilde{X}_{ik} \tilde{X}_{il} \tilde{X}_{im}| \lesssim \frac{b_w^4 b^4}{n} \left( 1 + \frac{\log^4(dn)}{n} \right), \quad (5.24)$$

where the second inequality follows by the AM-GM inequality and (5.16). Similarly, we can prove

$$\left\| \frac{1}{n^2} \sum_{i=1}^n w_i^4 \tilde{X}_i^{\otimes 4} \right\|_{\infty} \lesssim \frac{b_w^4 b^4}{n} \left( 1 + \frac{\log^4(dn)}{n} \right), \quad (5.25)$$

$$\left\| \frac{1}{n^{3/2}} \sum_{i=1}^n w_i^3 \tilde{X}_i^{\otimes 3} \right\|_{\infty} \lesssim \frac{b_w^3 b^3}{\sqrt{n}} \left( 1 + \frac{\log^3(dn)}{n} \right) \lesssim \frac{b_w^3 b^3}{\sqrt{n}}, \quad (5.26)$$

$$\left\| \frac{1}{n^{5/2}} \sum_{i=1}^n w_i^3 \tau_i^*(w_i) \tilde{X}_i^{\otimes 5} \right\|_{\infty} \lesssim \frac{b_w^5 b^5}{n^{3/2}} \left( 1 + \frac{\log^5(dn)}{n} \right). \quad (5.27)$$

By (5.24)–(5.25),

$$\begin{aligned} & \frac{\log^2 d}{\sigma_*^4} \left( \mathbb{E}^* \left[ \left\| \frac{1}{n^2} \sum_{i=1}^n \tau_i^*(w_i)^2 \tilde{X}_i^{\otimes 4} \right\|_{\infty} \right] + \mathbb{E}^* \left[ \left\| \frac{1}{n^2} \sum_{i=1}^n w_i^4 \tilde{X}_i^{\otimes 4} \right\|_{\infty} \right] \right) \\ & \lesssim \frac{b_w^4 b^4}{\sigma_*^4} \left( \frac{\log^2 d}{n} + \frac{\log^6(dn)}{n^2} \right) \lesssim \frac{b_w^4 b^4 \log^3(dn)}{\sigma_*^4 n}. \end{aligned} \quad (5.28)$$

Also, by (5.27),

$$\begin{aligned} & \frac{\log^{5/2} d}{\sigma_*^5} \mathbb{E}^* \left[ \left\| \frac{1}{n^{5/2}} \sum_{i=1}^n w_i^3 \tau_i^*(w_i) \tilde{X}_i^{\otimes 5} \right\|_{\infty} \right] \lesssim \frac{b_w^5 b^5}{\sigma_*^5} \left( \frac{\log^{5/2} d}{n^{3/2}} + \frac{\log^{15/2}(dn)}{n^{5/2}} \right) \\ & \lesssim \frac{b_w^5 b^5 \log^3(dn)}{\sigma_*^5 n}. \end{aligned} \quad (5.29)$$

Further, by the Schwarz inequality, (5.22) and (5.26),

$$\begin{aligned} & \frac{\log^{5/2} d}{\sigma_*^5} \mathbb{E}^* \left[ \left\| \bar{T}^* \otimes \frac{1}{n^{3/2}} \sum_{i=1}^n w_i^3 \tilde{X}_i^{\otimes 3} \right\|_{\infty} \right] \lesssim \frac{\log^{5/2} d}{\sigma_*^5} \sqrt{b_w^4 b^4 \left( \frac{\log(dn)}{n} + \frac{\log^4(dn)}{n^2} \right)} \cdot \frac{b_w^3 b^3}{\sqrt{n}} \\ & \leq \frac{b_w^5 b^5}{\sigma_*^5} \left( \frac{\log^3(dn)}{n} + \frac{\log^{9/2}(dn)}{n^{3/2}} \right) \lesssim \frac{b_w^5 b^5 \log^3(dn)}{\sigma_*^5 n}. \end{aligned} \quad (5.30)$$

Combining (5.10), (5.14), (5.23), (5.26), (5.28)–(5.30) and  $b_w \geq 1, b/\sigma_* \geq 1$ , we have, on  $\mathcal{E}_n$ ,

$$\begin{aligned} & \sup_{A \in \mathcal{R}} \left| P^*(S_n^* \in A) - \int_A \tilde{p}_{n,\gamma}(z) dz \right| \\ & \lesssim \frac{b_w^5 b^5 \log^3(dn)}{\sigma_*^5 n} \log n + \frac{\bar{\sigma}}{n^{3/2}} \left( \frac{\log d}{\underline{\sigma}} + b^2 \frac{\log^{5/2} d}{\sigma_*^3 \sqrt{n}} + b^3 b_w^3 \frac{\log^{5/2} d}{\sigma_*^4 \sqrt{n}} \right) \lesssim \frac{b_w^5 b^5 \log^3(dn)}{\sigma_*^5 n} \log n. \end{aligned}$$

It remains to prove

$$\sup_{A \in \mathcal{R}} \left| \int_A \tilde{p}_{n,\gamma}(z) dz - \int_A \hat{p}_{n,\gamma}(z) dz \right| \lesssim \frac{b_w^5 b^5 \log^3(dn)}{\sigma_*^5 n} \log n \quad \text{on } \mathcal{E}_n. \quad (5.31)$$

Observe that

$$\sup_{A \in \mathcal{R}} \left| \langle \hat{\Sigma}_n - \bar{X}^2, \int_A \nabla^2 \phi_{\Sigma}(z) dz \rangle \right| \lesssim \|\bar{X}\|_{\infty}^2 \frac{\log d}{\sigma_*^2} \quad (5.32)$$

and

$$\begin{aligned} & \sup_{A \in \mathcal{R}} \left| \frac{1}{n^{3/2}} \sum_{i=1}^n \langle (X_i - \bar{X})^{\otimes 3} - X_i^{\otimes 3}, \int_A \nabla^3 \phi_\Sigma(z) dz \rangle \right| \\ & \lesssim \frac{1}{\sqrt{n}} \left( \left\| \frac{1}{n} \sum_{i=1}^n X_i^{\otimes 2} \right\|_\infty \|\bar{X}\|_\infty + \|\bar{X}\|_\infty^3 \right) \frac{\log^{3/2} d}{\sigma_*^3}. \end{aligned}$$

Also, note that  $|\gamma| \leq b_w \mathbb{E}[w_1^2] = b_w$ . Hence (5.31) follows from (5.12) and (5.15).

Next consider Case (ii). In this case,  $S_n^* \sim N(0, \widehat{\Sigma}_n)$  conditional on the data. Hence, applying Theorem 5.2 and using the bounds (5.12), (5.14) and (5.32), we obtain the desired bound with a simplified argument of the proof for Case (i).  $\square$

Now we turn to the proof of Theorems 5.1 and 5.2. As usual, the proof starts with a smoothing inequality. We will use the following version.

**Lemma 5.1.** *Let  $\mu$  be a finite measure,  $\nu$  a finite signed measure, and  $K$  a probability measure on  $\mathbb{R}^d$ . Let  $\varepsilon > 0$  be a constant such that  $\alpha := K([- \varepsilon, \varepsilon]^d) > 1/2$ . Let  $h : \mathbb{R}^d \rightarrow \mathbb{R}$  be a bounded measurable function. Then we have*

$$\left| \int h d(\mu - \nu) \right| \leq (2\alpha - 1)^{-1} [\gamma^*(h; \varepsilon) + \tau^*(h; \varepsilon) + \alpha \tilde{\tau}^*(h; \varepsilon)],$$

where

$$\gamma^*(h; \varepsilon) = \sup_{y \in \mathbb{R}^d} \gamma(h_y; \varepsilon), \quad \tau^*(h; \varepsilon) = \sup_{y \in \mathbb{R}^d} \tau(h_y; \varepsilon), \quad \tilde{\tau}^*(h; \varepsilon) = \sup_{y \in \mathbb{R}^d} \tilde{\tau}(h_y; \varepsilon),$$

with  $h_y(x) = h(x + y)$ ,

$$\begin{aligned} \gamma(h; \varepsilon) &= \max \left\{ \int M_h(x; \varepsilon) (\mu - \nu) * K(dx), - \int m_h(x; \varepsilon) (\mu - \nu) * K(dx) \right\}, \\ \tau(h; \varepsilon) &= \max \left\{ \int [M_h(x; \varepsilon) - h(x)] \nu(dx), \int [h(x) - m_h(x; \varepsilon)] \nu(dx) \right\}, \\ \tilde{\tau}(h; \varepsilon) &= \sup_{y \in [-\varepsilon, \varepsilon]^d} \left| \int [h(x + y) - h(x)] \nu(dx) \right|, \\ M_h(x; \varepsilon) &= \sup_{y: \|y-x\|_\infty \leq \varepsilon} h(y), \quad m_h(x; \varepsilon) = \inf_{y: \|y-x\|_\infty \leq \varepsilon} h(y), \end{aligned}$$

and  $*$  denotes the convolution of two finite signed measures.

The proof of this lemma is a straightforward modification of [7, Lemma 11.4] and given in Appendix B.1, but its statement contains an important difference from the original one: The bound does not contain the positive part of the signed measure  $\nu$ . This is important for bounding  $\tau^*(h; \varepsilon)$  and  $\tilde{\tau}^*(h; \varepsilon)$  in our setting. To bound these quantities, we will use the following anti-concentration inequality. For  $A = \prod_{j=1}^d [a_j, b_j] \in \mathcal{R}$  and  $u, v \in \mathbb{R}_+^d := [0, \infty)^d$ , define  $A^{u,v} = \prod_{j=1}^d [a_j - u_j, b_j + v_j]$ .

**Lemma 5.2.** *Let  $r \in \mathbb{N}$ . Then*

$$\sup_{A \in \mathcal{R}} \sup_{\varepsilon > 0} \sup_{u, v \in [0, \varepsilon]^d} \frac{1}{\varepsilon} \left\| \int_{A^{u,v} \setminus A} \nabla^r \phi_\Sigma(z) dz \right\|_1 \leq C_r \frac{(\log d)^{(r+1)/2}}{\sigma_*^{r+1}}, \quad (5.33)$$

where  $C_r > 0$  is a constant depending only on  $r$ .

**Proof.** See Appendix B.2. □

We will apply Lemma 5.1 with  $K = N(0, t\Sigma)$ . To bound the quantity  $\gamma^*(h; \varepsilon)$ , we introduce some notation and lemmas. Given a bounded measurable function  $h : \mathbb{R}^d \rightarrow \mathbb{R}$  and  $s \in [0, 1]$ , we define a function  $h_s : \mathbb{R}^d \rightarrow \mathbb{R}$  as

$$h_s(x) = \mathbb{E}[h(\sqrt{1-s}x + \sqrt{s}Z)], \quad x \in \mathbb{R}^d,$$

where  $Z \sim N(0, \Sigma)$ . When  $s > 0$ ,  $h_s(x)$  can be rewritten as

$$h_s(x) = s^{-d/2} \int_{\mathbb{R}^d} h(z) \phi_\Sigma \left( \frac{z - \sqrt{1-s}x}{\sqrt{s}} \right) dz.$$

By this expression,  $h_s$  is infinitely differentiable and

$$\nabla^r h_s(x) = \left( -\sqrt{\frac{1-s}{s}} \right)^r \int_{\mathbb{R}^d} h(\sqrt{1-s}x + \sqrt{s}z) \nabla^r \phi_\Sigma(z) dz \quad (5.34)$$

for any  $r \in \mathbb{N}$ . In particular,  $h_s \in C_b^\infty(\mathbb{R}^d)$ . We will use the following lemmas to bound  $\gamma^*(h; \varepsilon)$ .

**Lemma 5.3.** *Let  $h : \mathbb{R}^d \rightarrow \mathbb{R}$  be a bounded measurable function and  $t \in (0, 1]$ . Then, under the assumptions of Theorem 5.1,*

$$\begin{aligned} & \mathbb{E}[h_t(W)] - \int_{\mathbb{R}^d} h_t(z) p_W(z) dz \\ &= \frac{1}{4} \int_t^1 \left( \int_s^1 \frac{1}{(1-u)^2} \left( \mathbb{E}[\langle \bar{T}^{\otimes 2}, \nabla^4 h_u(W) \rangle] - \sum_{i=1}^n \mathbb{E}[\langle \tau_i(\xi_i)^{\otimes 2}, \nabla^4 h_u(W) \rangle] \right) du \right) ds \\ & \quad - \frac{1}{8} \int_t^1 \left( \int_s^1 \sum_{i=1}^n \frac{\mathbb{E}[\langle \tau_i(\xi_i) \otimes \xi_i^{\otimes 2}, \nabla^4 h_u(W) \rangle]}{(1-u)^2} du \right) ds \\ & \quad + \frac{1}{16} \int_t^1 \left( \int_s^1 \frac{1}{\sqrt{1-u}} \left( \int_u^1 \sum_{i=1}^n \frac{\mathbb{E}[\langle \xi_i^{\otimes 4}, \nabla^4 h_v(W) \rangle]}{(1-v)^{5/2}} dv \right) du \right) ds \\ & \quad + \frac{1}{16} \int_t^1 \left( \int_s^1 \frac{1}{\sqrt{1-u}} \left( \int_u^1 \sum_{i=1}^n \frac{\mathbb{E}[\langle \xi_i^{\otimes 3} \otimes (\bar{T} - \tau_i(\xi_i)), \nabla^5 h_v(W) \rangle]}{(1-v)^{5/2}} dv \right) du \right) ds. \end{aligned} \quad (5.35)$$

Also, under the assumptions of Theorem 5.2,

$$\mathbb{E}[h_t(W)] - \int_{\mathbb{R}^d} h_t(z) p_W(z) dz = \frac{1}{4} \int_t^1 \left( \int_s^1 \frac{\mathbb{E}[\langle (\Sigma_W - \Sigma)^{\otimes 2}, \nabla^4 h_u(W) \rangle]}{(1-u)^2} du \right) ds. \quad (5.36)$$

**Proof.** See Appendix B.3. □

**Lemma 5.4.** *Let  $h = 1_A$  with  $A \in \mathcal{R}$ . Then, for any  $s \in (0, 1)$  and  $r \in \mathbb{N}$ ,*

$$\sup_{x \in \mathbb{R}^d} \|\nabla^r h_s(x)\|_1 \leq C_r \left( \frac{1-s}{\sigma_*^2 s} \log d \right)^{r/2}, \quad (5.37)$$

where  $C_r > 0$  is a constant depending only on  $r$ .

**Proof.** Observe that  $\{z \in \mathbb{R}^d : \sqrt{1-sx} + \sqrt{sz} \in A\} \in \mathcal{R}$  for any  $x \in \mathbb{R}^d$ . Thus, the claim follows from (5.34) and Lemma E.4.  $\square$

**Proof of Theorem 5.1.** We apply Lemma 5.1 to  $\mu, \nu$  and  $K$  defined as

$$\mu(A) = P(\sqrt{1-t}W \in A), \quad \nu(A) = \int_{\mathbb{R}^d} 1_A(\sqrt{1-tz})p_W(z)dz, \quad K(A) = P(\sqrt{t}Z \in A).$$

Since

$$P(\|Z\|_\infty > \bar{\sigma}\sqrt{2\log(2d)}) \leq \sum_{j=1}^d P(|Z_j| > \sigma_j\sqrt{2\log(2d)}) \leq \frac{d}{2}e^{-\log(2d)} = \frac{1}{4},$$

we have  $\alpha := K([- \varepsilon, \varepsilon]^d) \geq 3/4 > 1/2$  with  $\varepsilon = \bar{\sigma}\sqrt{2t\log(2d)}$ . Let  $h = 1_A$  with  $A = \prod_{j=1}^d [a_j, b_j] \in \mathcal{R}$ . Then we have  $M_h(x; \varepsilon) = 1_{A^\varepsilon}(x)$  and  $m_h(x; \varepsilon) = 1_{A^{-\varepsilon}}(x)$ , where we set  $A^r := \prod_{j=1}^d [a_j - r, b_j + r]$  for any  $r \in \mathbb{R}$  with interpreting  $[a, b] = \emptyset$  if  $a > b$ . Hence

$$\gamma(h; \varepsilon) \leq \sup_{h=1_A, A \in \mathcal{R}} \left| \mathbb{E}[h_t(W)] - \int h_t(z)p_W(z)dz \right|$$

and

$$\int [M_h(x; \varepsilon) - h(x)]\nu(dx) \leq \sup_{A \in \mathcal{R}; u, v \in [0, \varepsilon]^d} \left| \int 1_{A^{u, v} \setminus A}(\sqrt{1-tz})p_W(z)dz \right|.$$

Further, for each  $j = 1, \dots, d$ , set

$$\begin{cases} I_j = [a_j + \varepsilon, b_j - \varepsilon], u_j = v_j = \varepsilon & \text{if } a_j + \varepsilon < b_j - \varepsilon, \\ I_j = \{(a_j + b_j)/2\}, u_j = v_j = (b_j - a_j)/2 & \text{if } b_j \leq a_j + 2\varepsilon. \end{cases}$$

Then we have  $\tilde{A} := \prod_{j=1}^d I_j \in \mathcal{R}$ ,  $u := (u_1, \dots, u_d)^\top \in [0, \varepsilon]^d$ ,  $v := (v_1, \dots, v_d)^\top \in [0, \varepsilon]^d$  and

$$\int [h(x) - m_h(x; \varepsilon)]\nu(dx) = \int 1_{\tilde{A}^{u, v} \setminus \tilde{A}}(\sqrt{1-tz})p_W(z)dz.$$

Consequently,

$$\tau(h; \varepsilon) \leq \sup_{A \in \mathcal{R}; u, v \in [0, \varepsilon]^d} \left| \int 1_{A^{u, v} \setminus A}(\sqrt{1-tz})p_W(z)dz \right|.$$

Besides, for any  $x \in \mathbb{R}^d$  and  $y \in [-\varepsilon, \varepsilon]^d$ , we have  $h(x+y) - h(x) = 1_{(A+y) \setminus A}(x) - 1_{A \setminus (A+y)}(x)$ . For each  $j = 1, \dots, d$ , set

$$\begin{cases} I_j = [a_j + y_j, b_j], u_j = 0, v_j = y_j & \text{if } y_j \geq 0, a_j + y_j < b_j, \\ I_j = [a_j, b_j + y_j], u_j = -y_j, v_j = 0 & \text{if } y_j < 0, a_j < b_j + y_j, \\ I_j = \{a_j + y_j\}, u_j = 0, v_j = b_j - a_j & \text{otherwise.} \end{cases}$$

Then we have  $\tilde{A} := \prod_{j=1}^d I_j \in \mathcal{R}$ ,  $u := (u_1, \dots, u_d)^\top \in [0, \varepsilon]^d$ ,  $v := (v_1, \dots, v_d)^\top \in [0, \varepsilon]^d$  and

$$\int 1_{(A+y) \setminus A}(x)\nu(dx) = \int 1_{\tilde{A}^{u, v} \setminus \tilde{A}}(\sqrt{1-tz})p_W(z)dz.$$

Also, observe that  $A \setminus (A + y) = [(A + y) - y] \setminus (A + y)$  and  $A + y \in \mathcal{R}$ . Hence we conclude

$$\tilde{\tau}(h; \varepsilon) \leq 2 \sup_{A \in \mathcal{R}; u, v \in [0, \varepsilon]^d} \left| \int 1_{A^u, v \setminus A}(\sqrt{1-tz}) p_W(z) dz \right|.$$

In addition, observe that  $h_y = 1_{A-y}$  for any  $y \in \mathbb{R}^d$ . As a result, Lemma 5.1 gives

$$\begin{aligned} & \sup_{A \in \mathcal{R}} \left| P(W \in A) - \int_A p_W(z) dz \right| \\ & \leq 2 \sup_{h=1_A, A \in \mathcal{R}} \left| \mathbb{E}[h_t(W)] - \int h_t(z) p_W(z) dz \right| + 6 \sup_{A \in \mathcal{R}; u, v \in [0, \varepsilon]^d} \left| \int 1_{A^u, v \setminus A}(\sqrt{1-tz}) p_W(z) dz \right|. \end{aligned}$$

Note that  $A/\sqrt{1-t} \in \mathcal{R}$  for any  $A \in \mathcal{R}$ . Thus, we have by Lemmas E.2 and 5.2

$$\begin{aligned} & \sup_{A \in \mathcal{R}; u, v \in [0, \varepsilon]^d} \left| \int 1_{A^u, v \setminus A}(\sqrt{1-tz}) p_W(z) dz \right| \\ & \lesssim \frac{\varepsilon}{\sqrt{1-t}} \left( \frac{\sqrt{\log d}}{\underline{\sigma}} + \frac{\log^{3/2} d}{\sigma_*^3} \|\Sigma_W - \Sigma\|_\infty + \frac{\log^2 d}{\sigma_*^4} \left\| \sum_{i=1}^n \mathbb{E}[\xi_i^{\otimes 3}] \right\|_\infty \right) \\ & \lesssim \bar{\sigma} \sqrt{t} \left( \frac{\log d}{\underline{\sigma}} + \frac{\log^2 d}{\sigma_*^3} \|\Sigma_W - \Sigma\|_\infty + \frac{\log^{5/2} d}{\sigma_*^4} \left\| \sum_{i=1}^n \mathbb{E}[\xi_i^{\otimes 3}] \right\|_\infty \right). \end{aligned}$$

Further, by Lemma 5.4, we have for any  $u, v \in (0, 1)$

$$\begin{aligned} & |\mathbb{E}[\langle \bar{T}^{\otimes 2}, \nabla^4 h_u(W) \rangle]| \lesssim \mathbb{E} \|\bar{T}\|_\infty^2 \frac{(1-u)^2 \log^2 d}{u^2 \sigma_*^4}, \\ & \left| \sum_{i=1}^n \mathbb{E}[\langle \tau_i(\xi_i)^{\otimes 2}, \nabla^4 h_u(W) \rangle] \right| \lesssim \mathbb{E} \left\| \sum_{i=1}^n \tau_i(\xi_i)^{\otimes 2} \right\|_\infty \frac{(1-u)^2 \log^2 d}{u^2 \sigma_*^4}, \\ & \left| \sum_{i=1}^n \mathbb{E}[\langle \tau_i(\xi_i) \otimes \xi_i^{\otimes 2}, \nabla^4 h_u(W) \rangle] \right| \lesssim \mathbb{E} \left\| \sum_{i=1}^n \tau_i(\xi_i) \otimes \xi_i^{\otimes 2} \right\|_\infty \frac{(1-u)^2 \log^2 d}{u^2 \sigma_*^4}, \\ & \left| \sum_{i=1}^n \mathbb{E}[\langle \xi_i^{\otimes 4}, \nabla^4 h_v(W) \rangle] \right| \lesssim \mathbb{E} \left\| \sum_{i=1}^n \xi_i^{\otimes 4} \right\|_\infty \frac{(1-v)^2 \log^2 d}{v^2 \sigma_*^4}, \\ & \left| \sum_{i=1}^n \mathbb{E}[\langle \xi_i^{\otimes 3} \otimes \bar{T}, \nabla^5 h_v(W) \rangle] \right| \lesssim \mathbb{E} \left\| \bar{T} \otimes \sum_{i=1}^n \xi_i^{\otimes 3} \right\|_\infty \frac{(1-v)^{5/2} \log^{5/2} d}{v^{5/2} \sigma_*^5}, \\ & \left| \sum_{i=1}^n \mathbb{E}[\langle \xi_i^{\otimes 3} \otimes \tau_i(\xi_i), \nabla^5 h_v(W) \rangle] \right| \lesssim \mathbb{E} \left\| \sum_{i=1}^n \xi_i^{\otimes 3} \otimes \tau_i(\xi_i) \right\|_\infty \frac{(1-v)^{5/2} \log^{5/2} d}{v^{5/2} \sigma_*^5}. \end{aligned}$$

Also, by the AM-GM inequality,

$$\mathbb{E} \left\| \sum_{i=1}^n \tau_i(\xi_i) \otimes \xi_i^{\otimes 2} \right\|_\infty \leq \frac{1}{2} \left( \mathbb{E} \left\| \sum_{i=1}^n \tau_i(\xi_i)^{\otimes 2} \right\|_\infty + \mathbb{E} \left\| \sum_{i=1}^n \xi_i^{\otimes 4} \right\|_\infty \right).$$

Since

$$\int_t^1 \left( \int_s^1 \frac{1}{u^2} du \right) ds \leq |\log t|, \quad \int_t^1 \left( \int_s^1 \frac{1}{\sqrt{1-u}} \left( \int_u^1 \frac{1}{v^{5/2} \sqrt{1-v}} dv \right) du \right) ds \lesssim 1,$$

we obtain the desired result by (5.35).  $\square$

**Proof of Theorem 5.2.** The claim follows by replacing (5.35) with (5.36) in the proof of Theorem 5.1.  $\square$

## 6 Proofs for Section 3

Given a random vector  $W$ , we denote by  $F_W$  the distribution function of  $W^\vee$ .

### 6.1 Proofs of Theorems 3.1 and 3.2

The proofs are based on the following abstract result.

**Proposition 6.1** (Abstract Cornish–Fisher type expansion for maximum statistics). *Let  $\varepsilon \in (0, 1/2)$ . Also, let  $W$  be a random vector in  $\mathbb{R}^d$ . Suppose that there exist arrays  $U \in (\mathbb{R}^d)^{\otimes 2}$ ,  $V \in (\mathbb{R}^d)^{\otimes 3}$  and a constant  $\Delta > 0$  such that*

$$\sup_{t \in \mathbb{R}} \left| P(W^\vee \leq t) - \int_{A(t)} p_{U,V}(z) dz \right| \leq \Delta, \quad (6.1)$$

where  $p_{U,V}(z) = \phi_\Sigma(z) + \langle U, \nabla^2 \phi_\Sigma(z) \rangle + \langle V, \nabla^3 \phi_\Sigma(z) \rangle$ . Set

$$\delta := \|U\|_\infty \frac{\log d}{\sigma_*^2} + \|V\|_\infty \frac{\log^{3/2} d}{\sigma_*^3}, \quad \bar{\Delta} := \Delta + \frac{\varsigma_d}{\sigma_*} (\Delta + \delta) \delta.$$

Then, there exist positive constants  $c$  and  $C$  depending only on  $\varepsilon$  such that if  $\delta + \bar{\Delta} \leq c$ , then

$$\sup_{\varepsilon < p < 1-\varepsilon} \left| F_W^{-1}(p) - \left( F_Z^{-1}(p) - \frac{Q_{U,V}(F_Z^{-1}(p))}{f_\Sigma(F_Z^{-1}(p))} \right) \right| \leq \frac{C}{\sqrt{\log d}} \left( \varsigma_d \bar{\Delta} + \frac{\varsigma_d^3}{\sigma_*^2} (\delta + \bar{\Delta})^2 \right),$$

where

$$Q_{U,V}(t) = \int_{A(t)} \{p_{U,V}(z) - \phi_\Sigma(z)\} dz.$$

First we prove Theorems 3.1 and 3.2 using Proposition 6.1.

**Proof of Theorem 3.1.** First, observe that Lemma E.3 yields

$$\varsigma_d / \sigma_* \gtrsim \underline{\sigma} / \bar{\sigma} \geq \lambda^{-1}. \quad (6.2)$$

Hence, due to (3.1), we may assume

$$\left( 1 + \frac{\varsigma_d^3}{\sigma_*^3} \right) \frac{\log^3 d}{n} \log n \leq 1. \quad (6.3)$$

Then, we have (2.9) by Theorem 2.2. Also, observe that  $E[\bar{X}^3] \lesssim b^3 / \sqrt{n}$ . Hence, in this setting,  $\delta$  and  $\bar{\Delta}$  in Proposition 6.1 are bounded as

$$\delta \leq C_\lambda \sqrt{\frac{\log^3 d}{n}}, \quad \bar{\Delta} \leq C_\lambda \left( 1 + \frac{\varsigma_d}{\sigma_*} \right) \frac{\log^3 d}{n} \log n,$$

where we used (6.3) for the second inequality. Combining these bounds with (6.2) and (6.3) gives

$$\delta + \bar{\Delta} \leq C_\lambda \sqrt{\frac{\log^3 d}{n} \log n}, \quad \bar{\Delta} \leq C_\lambda \frac{\varsigma_d \log^3 d}{\sigma_* n} \log n.$$

Consequently, the desired result follows from Proposition 6.1.  $\square$

**Proof of Theorem 3.2.** By the same reasoning as in the proof of Theorem 3.1, we may assume

$$\left(1 + \frac{\zeta_d^3}{\sigma_*^3}\right) \frac{\log^3(dn)}{n} \log n \leq 1. \quad (6.4)$$

Let  $\mathcal{E}_n$  be the event defined by (5.11). Recall that  $P(\mathcal{E}_n) \geq 1 - 1/n$ . Also, by the proof of Theorem 2.3, we have (2.11) on  $\mathcal{E}_n$ . Further, recall that we have (5.13) and (5.15) on  $\mathcal{E}_n$ . Hence, on  $\mathcal{E}_n$

$$\frac{1}{2} \|\overline{X^2} - \Sigma\|_\infty \frac{\log d}{\sigma_*^2} + \frac{|\gamma|}{6} \|\overline{X^3}\|_\infty \frac{\log^{3/2} d}{\sigma_*^3} \lesssim b^2 \frac{\log^{3/2}(dn)}{\sigma_*^2 \sqrt{n}} + \frac{|\gamma| b^3 \log^{3/2} d}{\sigma_*^3 \sqrt{n}} \leq C_{\lambda, b_w} \sqrt{\frac{\log^3(dn)}{n}}.$$

Consequently, a similar argument to the proof of Theorem 3.1 gives the desired result.  $\square$

Now we turn to the proof of Proposition 6.1. The proof relies on the following lemma.

**Lemma 6.1.** *Let  $Z$  be a centered Gaussian vector in  $\mathbb{R}^d$ . If  $Z^\vee$  has a continuous density  $f$ , then*

$$f(F_Z^{-1}(p)) \geq \frac{1}{4\sqrt{\text{Var}[Z^\vee]}} \min\left\{\frac{p}{\sqrt{2}}, (1-p)^{3/2}\right\} \quad (6.5)$$

for all  $p \in (0, 1)$ . Moreover, if  $\text{Cov}[Z] = \Sigma$ , there exists a universal constant  $C > 0$  such that

$$|(F_Z^{-1})''(p)| \leq C \left(\frac{\text{Var}[Z^\vee]}{\min\{p^2, (1-p)^3\}}\right)^{3/2} \frac{\log d}{\sigma_*^2} \quad (6.6)$$

for all  $p \in (0, 1)$ .

**Proof.** See Appendix D.  $\square$

**Remark 6.1.** Under the first assumption of Lemma 6.1, we can also derive the following Gaussian type isoperimetric inequality for  $Z^\vee$ : For all  $p \in (0, 1)$ ,

$$f(F_Z^{-1}(p)) \geq \frac{1}{\sigma} \phi(\Phi^{-1}(p)), \quad (6.7)$$

where  $\sigma := \max_{1 \leq j \leq d} \sqrt{\text{Var}[Z_j]}$ . In fact, by [3, Proposition 5], (6.7) follows once we prove

$$\phi(\Phi^{-1}(\mathbb{E}[g(Z^\vee)])) \leq \mathbb{E} \left[ \sqrt{\phi(\Phi^{-1}(g(Z^\vee)))^2 + \sigma^2 g'(Z^\vee)^2} \right]$$

for any locally Lipschitz function  $g : \mathbb{R} \rightarrow [0, 1]$ . The latter follows by applying Bobkov's functional Gaussian isoperimetric inequality to the function  $x \mapsto g(\max_{1 \leq j \leq d} (\text{Cov}[Z]^{1/2} x)_j)$  (cf. Eq.(2) of [3]). While (6.7) has a better dependence on  $p$  than (6.5), it is often the case that  $\text{Var}[Z^\vee] = O(1/\log d)$  as already mentioned at the beginning of Section 3, so (6.5) is preferable to (6.7) in terms of the dimension dependence.

**Proof of Proposition 6.1.** Observe that

$$Q_{U,V}(t) = \langle U, \int_{A(t)} \nabla^2 \phi_\Sigma(z) dz \rangle + \langle V, \int_{A(t)} \nabla^3 \phi_\Sigma(z) dz \rangle.$$

Hence, by Lemmas E.4 and 5.2, there exists a universal constant  $C_1 \geq 1$  such that

$$|Q_{U,V}(t)| \leq C_1 \delta \quad (6.8)$$

and

$$|Q_{U,V}(t) - Q_{U,V}(s)| \leq C_1 \delta \frac{\sqrt{\log d}}{\sigma_*} |t - s| \quad (6.9)$$

for all  $t, s \in \mathbb{R}$ . Also, for any  $p \in (\varepsilon, 1 - \varepsilon)$ , we have by (6.1)

$$p \leq F_W(F_W^{-1}(p)) \leq F_Z(F_W^{-1}(p)) + Q_{U,V}(F_W^{-1}(p)) + \Delta \quad (6.10)$$

and

$$p \geq F_W(F_W^{-1}(p)-) \geq F_Z(F_W^{-1}(p)) + Q_{U,V}(F_W^{-1}(p)) - \Delta. \quad (6.11)$$

Combining these bounds with (6.8) gives  $p - \Delta - C_1 \delta \leq F_Z(F_W^{-1}(p)) \leq p + \Delta + C_1 \delta$ . Therefore, provide that  $\Delta + C_1 \delta < \varepsilon/2$ , we have by the mean value theorem and (6.5)

$$|F_Z^{-1}(p \pm (\Delta + C_1 \delta)) - F_Z^{-1}(p)| \leq C_2 \sqrt{\text{Var}[Z^\vee]} (\Delta + C_1 \delta)$$

for some constant  $C_2 \geq 1$  depending only on  $\varepsilon$ . Thus we obtain

$$|F_W^{-1}(p) - F_Z^{-1}(p)| \leq C_2 \sqrt{\text{Var}[Z^\vee]} (\Delta + C_1 \delta).$$

This and (6.9) give

$$|Q_{U,V}(F_W^{-1}(p)) - Q_{U,V}(F_Z^{-1}(p))| \leq C_1 C_2 \delta \frac{\varsigma_d}{\sigma_*} (\Delta + C_1 \delta) =: \Delta'.$$

Combining this with (6.10) and (6.11), we obtain

$$p - Q_{U,V}(F_Z^{-1}(p)) - \Delta - \Delta' \leq F_Z(F_W^{-1}(p)) \leq p - Q_{U,V}(F_Z^{-1}(p)) + \Delta + \Delta'. \quad (6.12)$$

Thus, provided that  $C_1 \delta + \Delta + \Delta' < \varepsilon/2$ , we have by Taylor's theorem and (6.6)

$$\begin{aligned} & \left| F_Z^{-1}(p - Q_{U,V}(F_Z^{-1}(p)) \pm (\Delta + \Delta')) - \left( F_Z^{-1}(p) - \frac{Q_{U,V}(F_Z^{-1}(p)) \mp (\Delta + \Delta')}{f_\Sigma(F_Z^{-1}(p))} \right) \right| \\ & \leq C_3 \frac{\varsigma_d^3}{\sigma_*^2 \sqrt{\log d}} |Q_W(F_Z^{-1}(p)) \mp (\Delta + \Delta')|^2 \end{aligned}$$

for some constant  $C_3 \geq 1$  depending only on  $\varepsilon$ . Combining this with (6.5), (6.8) and (6.12) gives

$$\begin{aligned} & \left| F_W^{-1}(p) - \left( F_Z^{-1}(p) - \frac{Q_{U,V}(F_Z^{-1}(p))}{f_\Sigma(F_Z^{-1}(p))} \right) \right| \\ & \leq \frac{4\sqrt{\text{Var}[Z^\vee]}}{\varepsilon^{3/2}} (\Delta + \Delta') + C_3 \frac{\varsigma_d^3}{\sigma_*^2 \sqrt{\log d}} (C_1 \delta + \Delta + \Delta')^2. \end{aligned}$$

Since  $\Delta + \Delta' \leq C_1^2 C_2 \bar{\Delta}$ , this completes the proof.  $\square$

## 6.2 Proof of Theorem 3.3

**Lemma 6.2.** For any  $r \in \mathbb{N}$  and  $t \in \mathbb{R}$ ,

$$\int_{A(t)^c} \nabla^r \phi_\Sigma(z) dz = - \int_{A(t)} \nabla^r \phi_\Sigma(z) dz.$$

**Proof.** Let  $Z \sim N(0, \Sigma)$ . Then, for any  $x \in \mathbb{R}^d$ ,

$$\int_{A(t)^c} \phi_\Sigma(z+x) dz = P(Z-x \in A(t)^c) = 1 - P(Z-x \in A(t)) = 1 - \int_{A(t)} \phi_\Sigma(z+x) dz.$$

Differentiating the both sides  $r$  times with respect to  $x$  and setting  $x = 0$ , we obtain the desired result.  $\square$

**Lemma 6.3** (Anti-concentration inequality for  $T_n$ ). *Under the assumptions of Theorem 2.2, there exists a universal constant  $C > 0$  such that*

$$P(t \leq T_n \leq t + \varepsilon) \leq C \left( \frac{b^5 \log^3 d}{\sigma_*^5} \frac{1}{n} \log n + \varepsilon \left( \frac{\sqrt{\log d}}{\sigma} + \frac{b^3 \log^2 d}{\sigma_*^4 \sqrt{n}} \right) \right)$$

for all  $t \in \mathbb{R}$  and  $\varepsilon > 0$ .

**Proof.** The claim immediately follows by combining Theorem 2.2 with Lemmas E.2 and 5.2.  $\square$

**Proof of Theorem 3.3.** By Theorems 3.1 and 3.2, there exist positive constants  $c$  and  $C$  depending only on  $\lambda, \varepsilon$  and  $b_w$  such that if (3.3) holds, then we have (3.2) and (3.4) with probability at least  $1 - 1/n$ . In the sequel we assume (3.3) is satisfied with this  $c$  and fix  $\alpha \in (\varepsilon, 1 - \varepsilon)$  arbitrarily. By (6.5) and Lemmas E.4 and E.10

$$\frac{1}{f_\Sigma(c_{1-\alpha}^G)} \left| \hat{Q}_{n,\gamma}(c_{1-\alpha}^G) - \gamma Q_n(c_{1-\alpha}^G) - \frac{1}{2} \langle \bar{X}^2 - \Sigma, \Psi_\alpha \rangle \right| \lesssim \frac{|\gamma| \zeta_d}{\varepsilon^{3/2} \sqrt{\log d}} \frac{b^3 \log^{3/2} d}{\sigma_*^3} \frac{1}{n} \sqrt{\log n}$$

with probability at least  $1 - 1/n$ . Combining this with (3.2) and (3.4), we have

$$\left| \hat{c}_{1-\alpha} - \tilde{c}_{1-\alpha} + \frac{\langle \bar{X}^2 - \Sigma, \Psi_\alpha \rangle}{2f_\Sigma(c_{1-\alpha}^G)} \right| \leq \frac{C_{\lambda,\varepsilon,b_w} \zeta_d^3 \log^3(dn)}{\sqrt{\log d} \sigma_*^2} \frac{1}{n} \log n \quad (6.13)$$

with probability at least  $1 - 2/n$ , where  $\tilde{c}_{1-\alpha} := c_{1-\alpha} + (1 - \gamma)Q_n(c_{1-\alpha}^G)/f_\Sigma(c_{1-\alpha}^G)$ . This and Lemma 6.3 give

$$\begin{aligned} & \left| P(T_n \geq \hat{c}_{1-\alpha}) - P\left(T_n \geq \tilde{c}_{1-\alpha} - \frac{\langle \bar{X}^2 - \Sigma, \Psi_\alpha \rangle}{2f_\Sigma(c_{1-\alpha}^G)}\right) \right| \\ & \leq C_{\lambda,\varepsilon,b_w} \left( \frac{\log^3 d}{n} \log n + \frac{\zeta_d^3 \log^3(dn)}{\sigma_*^3} \frac{1}{n} (\log n) \left( 1 + \sqrt{\frac{\log^3 d}{n}} \right) \right) + \frac{1}{n} \\ & \leq C_{\lambda,\varepsilon,b_w} \frac{\zeta_d^3 \log^3(dn)}{\sigma_*^3} \frac{1}{n} \log n, \end{aligned} \quad (6.14)$$

where the second inequality follows by (3.3). Now, observe that

$$\tilde{T}_n := T_n + \frac{\langle \bar{X}^2 - \Sigma, \Psi_\alpha \rangle}{2f_\Sigma(c_{1-\alpha}^G)} = \max_{1 \leq j \leq d} \frac{1}{\sqrt{n}} \sum_{i=1}^n (X_{ij} + U_i), \quad (6.15)$$

where

$$U_i := \frac{\langle X_i^{\otimes 2} - \mathbb{E}[X_i^{\otimes 2}], \tilde{\Psi}_\alpha \rangle}{\sqrt{n}}, \quad \tilde{\Psi}_\alpha := \frac{\Psi_\alpha}{2f_\Sigma(c_{1-\alpha}^G)}. \quad (6.16)$$

Hence we can derive an Edgeworth expansion for  $\tilde{T}_n$  by applying Theorem 5.1 with  $\xi_i = (X_i + U_i \mathbf{1}_d)/\sqrt{n}$ . By Lemma C.1,  $\xi_i$  has a Stein kernel  $\tau_i$  such that  $\tau_i(\xi_i) = (\tau_i^X(X_i, Y_i) + V_i + V'_i)/n$ , where

$$V_i := \frac{\tau_i^{XY}(X_i, Y_i) \text{vec}(\tilde{\Psi}_\alpha) \mathbf{1}_d^\top + \mathbf{1}_d \text{vec}(\tilde{\Psi}_\alpha)^\top \tau_i^{YX}(X_i, Y_i)}{\sqrt{n}},$$

$$V'_i := \frac{\mathbf{1}_d \text{vec}(\tilde{\Psi}_\alpha)^\top \tau_i^Y(X_i, Y_i) \text{vec}(\tilde{\Psi}_\alpha) \mathbf{1}_d^\top}{n}.$$

We are going to bound the quantities appearing in the right hand side of (5.4). First, by (6.5) and Lemma E.4

$$\|\tilde{\Psi}_\alpha\|_1 \leq C_\varepsilon \frac{\varsigma_d \sqrt{\log d}}{\sigma_*^2}. \quad (6.17)$$

Hence, by Lemma E.6 and (3.3),

$$\|U_i\|_{\psi_{1/2}} \leq C_\varepsilon \frac{b^2 \varsigma_d \sqrt{\log d}}{\sqrt{n} \sigma_*^2} \leq C_{\lambda, \varepsilon, b_w} \frac{b}{\log d}, \quad (6.18)$$

and

$$\max_{j,k} \|V_{i,jk}\|_{\psi_{1/3}} \leq C_\varepsilon \frac{b^3 \varsigma_d \sqrt{\log d}}{\sigma_*^2 \sqrt{n}} \leq C_{\lambda, \varepsilon, b_w} \frac{b^2}{\log d},$$

$$\max_{j,k} \|V'_{i,jk}\|_{\psi_{1/4}} \leq C_\varepsilon \frac{b^4 \varsigma_d^2 \log d}{\sigma_*^4 n} \leq C_{\lambda, \varepsilon, b_w} \frac{b^2}{\log^2 d}. \quad (6.19)$$

These estimates allow us to prove (5.6)–(5.8) with  $b$  replaced by  $C_{\lambda, \varepsilon, b_w} b$  in a similar manner to the proof of Theorem 2.2. Further, observe that

$$\max_j |\mathbb{E}[X_{ij} U_i]| \leq C_{\lambda, \varepsilon} \frac{b^2 \varsigma_d \sqrt{\log d}}{\sqrt{n} \sigma_*}, \quad |\mathbb{E}[U_i^2]| \leq C_{\lambda, \varepsilon, b_w} \frac{b^2}{\sqrt{n} \sigma_* \sqrt{\log d}}.$$

Combining these estimates with (6.19), we can also prove (5.5) with  $b$  replaced by  $C_{\lambda, \varepsilon} b \sqrt{\varsigma_d / \sigma_*}$  similarly to the proof of Theorem 2.2. All together, we can proceed as in the proof of Theorem 2.2 and then obtain

$$\sup_{t \in \mathbb{R}} \left| P(\tilde{T}_n \leq t) - \int_{A(t)} (p_n(z) + q_n(z)) dz \right| \leq C_{\lambda, \varepsilon, b_w} \frac{\varsigma_d^2 \log^3 d}{\sigma_*^2 n} \log n,$$

where

$$q_n(z) = \frac{1}{2n} \sum_{i=1}^n \langle 2\mathbb{E}[X_i U_i] \mathbf{1}_d^\top + \mathbb{E}[U_i^2] \mathbf{1}_d^{\otimes 2}, \nabla^2 \phi_\Sigma(z) \rangle - \frac{1}{6n^{3/2}} \sum_{i=1}^n \langle \mathbb{E}[(X_i + U_i \mathbf{1}_d)^{\otimes 3} - X_i^{\otimes 3}], \nabla^3 \phi_\Sigma(z) \rangle.$$

Therefore, in view of Theorem 2.2 and (5.2), it remains to prove

$$\left| \int_{A(\tilde{c}_{1-\alpha}^c)} p_n(z) dz - \int_{A(c_{1-\alpha}^c)} p_n(z) dz + (1-\gamma) Q_n(c_{1-\alpha}^G) \right| \leq C_{\lambda, \varepsilon, b_w} \frac{\varsigma_d^2 \log^3(dn)}{\sigma_*^2 n}, \quad (6.20)$$

$$\left| \int_{A(\tilde{c}_{1-\alpha})^c} q_n(z) dz + \mathbb{E}[R_n(\alpha)] \right| \leq C_{\lambda, \varepsilon, b_w} \frac{\varsigma_d^3 \log^3(dn)}{\sigma_*^3 n} \log n. \quad (6.21)$$

Let us prove (6.20). By Lemma 5.2,

$$\left| \int_{A(\tilde{c}_{1-\alpha})^c} p_n(z) dz - \int_{A(c_{1-\alpha})^c} p_n(z) dz + \{F_Z(\tilde{c}_{1-\alpha}) - F_Z(c_{1-\alpha})\} \right| \lesssim \frac{|1 - \gamma| b^3 \log^2 d}{\sigma_*^4 \sqrt{n}} \left| \frac{Q_n(c_{1-\alpha}^G)}{f_\Sigma(c_{1-\alpha}^G)} \right|.$$

Also, by Taylor's theorem and Lemma D.1,

$$|F_Z(\tilde{c}_{1-\alpha}) - F_Z(c_{1-\alpha}) - (1 - \gamma)Q_n(c_{1-\alpha}^G)| \lesssim \frac{|1 - \gamma| \log d}{\sigma_*^2} \left| \frac{Q_n(c_{1-\alpha}^G)}{f_\Sigma(c_{1-\alpha}^G)} \right|^2.$$

Further, Lemma E.4 and (6.5) yield

$$\left| \frac{Q_n(c_{1-\alpha}^G)}{f_\Sigma(c_{1-\alpha}^G)} \right| \lesssim \frac{\varsigma_d}{\sqrt{\varepsilon^3 \log d}} \cdot \frac{b^3}{\sqrt{n}} \cdot \frac{\log^{3/2} d}{\sigma_*^3} \leq C_{\lambda, \varepsilon} \frac{\varsigma_d \log d}{\sqrt{n}}. \quad (6.22)$$

Combining these three estimates gives (6.20). Next, to prove (6.21), consider the following decomposition:

$$\begin{aligned} & \int_{A(\tilde{c}_{1-\alpha})^c} q_n(z) dz \\ &= \frac{1}{n} \sum_{i=1}^n \langle \mathbb{E}[X_i U_i] \mathbf{1}_d^\top, \int_{A(\tilde{c}_{1-\alpha})^c} \nabla^2 \phi_\Sigma(z) dz \rangle + \frac{1}{2n} \sum_{i=1}^n \langle \mathbb{E}[U_i^2] \mathbf{1}_d^{\otimes 2}, \int_{A(\tilde{c}_{1-\alpha})^c} \nabla^2 \phi_\Sigma(z) dz \rangle \\ & \quad - \frac{1}{6n^{3/2}} \sum_{i=1}^n \langle \mathbb{E}[(X_i + U_i \mathbf{1}_d)^{\otimes 3} - X_i^{\otimes 3}], \int_{A(\tilde{c}_{1-\alpha})^c} \nabla^3 \phi_\Sigma(z) dz \rangle \\ &=: I + II + III. \end{aligned}$$

We can rewrite  $I$  as

$$\begin{aligned} I &= \frac{1}{n} \sum_{i=1}^n \sum_{j,k=1}^d \mathbb{E}[X_{ij} U_i] \int_{A(\tilde{c}_{1-\alpha})^c} \partial_{jk} \phi_\Sigma(z) dz \\ &= \frac{1}{n^{3/2}} \sum_{i=1}^n \sum_{j,k,l,m=1}^d \mathbb{E}[X_{ij} X_{il} X_{im}] \tilde{\Psi}_{\alpha, lm} \int_{A(\tilde{c}_{1-\alpha})^c} \partial_{jk} \phi_\Sigma(z) dz \\ &= \frac{1}{\sqrt{n}} \langle \mathbb{E}[\overline{X^3}] \otimes \mathbf{1}_d, \tilde{\Psi}_\alpha \otimes \int_{A(\tilde{c}_{1-\alpha})^c} \nabla^2 \phi_\Sigma(z) dz \rangle = -\frac{1}{\sqrt{n}} \langle \mathbb{E}[\overline{X^3}] \otimes \mathbf{1}_d, \tilde{\Psi}_\alpha \otimes \int_{A(\tilde{c}_{1-\alpha})} \nabla^2 \phi_\Sigma(z) dz \rangle, \end{aligned}$$

where we used Lemma 6.2 for the last equality. We are going to prove  $\tilde{c}_{1-\alpha}$  in the last expression can be replaced by  $c_{1-\alpha}^G$ . By Lemma 5.2 and (6.17),

$$\left\| \tilde{\Psi}_\alpha \otimes \left( \int_{A(\tilde{c}_{1-\alpha})} \nabla^2 \phi_\Sigma(z) dz - \int_{A(c_{1-\alpha}^G)} \nabla^2 \phi_\Sigma(z) dz \right) \right\|_1 \lesssim \frac{\varsigma_d \log^2 d}{\sigma_*^5} |\tilde{c}_{1-\alpha} - c_{1-\alpha}^G|.$$

Also, by (3.2) and (6.22),

$$|\tilde{c}_{1-\alpha} - c_{1-\alpha}^G| \leq C_{\lambda, \varepsilon, b_w} \left( \frac{\varsigma_d \log d}{\sqrt{n}} + \frac{\varsigma_d^3 \log^{5/2}(dn)}{\sigma_*^2 n} \log n \right).$$

Consequently, we deduce

$$|I + \mathbb{E}[R_n(\alpha)]| \leq C_{\lambda, \varepsilon, b_w} \frac{b^3 \varsigma_d \log^2 d}{\sigma_*^5 \sqrt{n}} |\tilde{c}_{1-\alpha} - c_{1-\alpha}^G| \leq C_{\lambda, \varepsilon, b_w} \frac{\varsigma_d^3 \log^3(dn)}{\sigma_*^3 n} \log n,$$

where we also used (3.3) and (6.2) for the last inequality. Meanwhile, by Lemma E.4 and (6.18),

$$|II| \leq C_\varepsilon \frac{b^4}{n} \cdot \frac{\varsigma_d^2 \log d \log d}{\sigma_*^4 \sigma_*^2} \leq C_{\lambda, \varepsilon} \frac{\varsigma_d^2 \log^2 d}{\sigma_*^2 n}$$

and

$$\begin{aligned} |III| &\lesssim \frac{1}{\sqrt{n}} \max_{1 \leq i \leq n} (\| \mathbb{E}[X_i^{\otimes 2} U_i] \|_\infty + \| \mathbb{E}[X_i U_i^2] \|_\infty + | \mathbb{E}[U_i^3] |) \frac{\log^{3/2} d}{\sigma_*^3} \\ &\leq C_{\lambda, \varepsilon, b_w} \frac{b^4 \varsigma_d \sqrt{\log d} \log^{3/2} d}{n \sigma_*^2 \sigma_*^3} \leq C_{\lambda, \varepsilon, b_w} \frac{\varsigma_d \log^2 d}{\sigma_* n}. \end{aligned}$$

All together, we complete the proof.  $\square$

### 6.3 Proof of Corollary 3.1

First, replacing  $X_i$  by  $X_i/\sigma$ , we may assume  $\sigma = 1$  without loss of generality. Note that we have  $\sigma_*^{-1} \lesssim \lambda$  under this assumption. Next, set  $\rho := \max_{1 \leq j < k \leq d} |\Sigma_{jk}| < 1$ . Note that we have  $1 - \rho^2 \geq \sigma_*^2$  because  $1 - \rho^2$  coincides with the minimum principal minor of  $\Sigma$  of size 2.

We begin by proving the following inequalities for every  $\alpha \in (\varepsilon, 1 - \varepsilon)$ :

$$|c_{1-\alpha}^G| \leq C_\varepsilon \sqrt{\log d}, \quad \exp(-(c_{1-\alpha}^G)^2/2) \leq C_{\lambda, \varepsilon, K} d^{-1} \sqrt{\log d}. \quad (6.23)$$

The first one is an immediate consequence of Lemma A.6 in [4]. Meanwhile, by Eq.(4.2.9) in [40] (see also Eq.(4.2.1) in [40]), we have for any  $u > 0$

$$|P(Z^\vee \leq u) - \Phi(u)^d| \leq C_\lambda \sum_{1 \leq j < k \leq d} |\Sigma_{jk}| \exp\left(-\frac{u^2}{1 + |\Sigma_{jk}|}\right).$$

Set  $\mathcal{I} := \{(j, k) \in \{1, \dots, d\}^2 : |\Sigma_{jk}| > u^{-2}\}$ . Observe that  $dK^2 \geq \sum_{j, k=1}^d \Sigma_{jk}^2 \geq u^{-2} \sum_{(j, k) \in \mathcal{I}} |\Sigma_{jk}|$ . Hence we obtain

$$\begin{aligned} |P(Z^\vee \leq u) - \Phi(u)^d| &\leq C_\lambda \sum_{(j, k) \in \mathcal{I}} |\Sigma_{jk}| e^{-u^2/(1+\rho)} + C_\lambda \sum_{(j, k) \notin \mathcal{I}} |\Sigma_{jk}| e^{-u^2/(1+u^{-2})} \\ &\leq C_\lambda d K^2 u^2 e^{-u^2/(1+\rho)} + C_\lambda d^{3/2} K e^{-u^2}. \end{aligned}$$

Further, observe that  $1 - (\varepsilon/2)^{1/d} < -d^{-1} \log(\varepsilon/2) \leq 1/2$ . Thus, with  $u = \Phi^{-1}((\varepsilon/2)^{1/d}) = -\Phi^{-1}(1 - (\varepsilon/2)^{1/d})$ , we have by Lemma 10.3 in [11]

$$e^{-u^2/2} = \sqrt{2\pi} \phi(u) \leq -\sqrt{2\pi} d^{-1} \log(\varepsilon/2) \sqrt{2 \log(-d/\log(\varepsilon/2))} \leq C_\varepsilon d^{-1} \sqrt{\log d}. \quad (6.24)$$

In addition, by the well-known inequality  $\Phi(-s) = 1 - \Phi(s) \leq e^{-s^2/2}$  for all  $s \geq 0$ , we deduce  $u \leq \sqrt{-2 \log(1 - (\varepsilon/2)^{1/d})} \leq C_\varepsilon \sqrt{\log d}$ . Hence we obtain

$$P(Z^\vee \leq u) \leq \varepsilon/2 + C_{\lambda, \varepsilon} \left( d^{-(1-\rho)/(1+\rho)} K^2 (\log d)^{1+1/(1+\rho)} + K d^{-1/2} \log d \right).$$

Therefore, if the second term on the right hand side is less than  $\varepsilon/2$ , then  $P(Z^\vee \leq u) \leq \varepsilon < 1 - \alpha$ , so  $c_{1-\alpha}^G > u$ . Hence the second bound in (6.23) follows by (6.24). Otherwise, we have  $d \leq d_0$  for some constant  $d_0$  depending only on  $\lambda, \varepsilon$  and  $K$ , so the second bound in (6.23) trivially holds with sufficiently large  $C_{\lambda, \varepsilon, K}$ .

Now we turn to the main body of the proof. Since the left hand side of (3.9) is bounded by 1, we may assume (3.3) holds with the constant  $c$  in Theorem 3.3. Also, we have  $f_\Sigma(c_{1-\alpha}^G) \geq C_\varepsilon \varsigma d / \sqrt{\log d}$  by (6.5). Thus, the proof completes once we show that

$$|\mathbb{E}[\langle \overline{X^3} \otimes \mathbf{1}_d, \Psi_\alpha^{\otimes 2} \rangle]| \leq C_{\lambda, \varepsilon, K} d^{-1/2} \log^2 d \quad (6.25)$$

for any  $\alpha \in (\varepsilon, 1 - \varepsilon)$ . Observe that

$$\langle \overline{X^3} \otimes \mathbf{1}_d, \Psi_\alpha^{\otimes 2} \rangle = \frac{1}{n} \sum_{i=1}^n \langle X_i^{\otimes 3} \otimes \mathbf{1}_d, \Psi_\alpha^{\otimes 2} \rangle = \frac{1}{n} \sum_{i=1}^n \langle X_i^{\otimes 2}, \Psi_\alpha \rangle \langle X_i \otimes \mathbf{1}_d, \Psi_\alpha \rangle.$$

Thus, by the Schwarz inequality

$$|\mathbb{E}[\langle \overline{X^3} \otimes \mathbf{1}_d, \Psi_\alpha^{\otimes 2} \rangle]| \leq \sqrt{\frac{1}{n} \sum_{i=1}^n \mathbb{E}[\langle X_i^{\otimes 2}, \Psi_\alpha \rangle^2] \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\langle X_i \otimes \mathbf{1}_d, \Psi_\alpha \rangle^2]}.$$

We have

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}[\langle X_i^{\otimes 2}, \Psi_\alpha \rangle^2] = \frac{1}{n} \sum_{i=1}^n \langle \mathbb{E}[X_i^{\otimes 4}], \Psi_\alpha^{\otimes 2} \rangle \leq b^4 \|\Psi_\alpha\|_1^2 \lesssim \lambda^4 \log^2 d,$$

where the last inequality follows from Lemma E.4. Also,

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\langle X_i \otimes \mathbf{1}_d, \Psi_\alpha \rangle^2] &= \frac{1}{n} \sum_{i=1}^n \sum_{j, k, l, m=1}^d \mathbb{E}[X_{ij} X_{ik}] \Psi_{\alpha, jl} \Psi_{\alpha, km} \\ &= \sum_{j, k, l, m=1}^d \Sigma_{jk} \Psi_{\alpha, jl} \Psi_{\alpha, km} \leq K \sum_{j=1}^d \left( \sum_{l=1}^d \Psi_{\alpha, jl} \right)^2. \end{aligned}$$

Therefore, (6.25) follows once we show

$$\sum_{j=1}^d \left( \sum_{l=1}^d \Psi_{\alpha, jl} \right)^2 \leq C_{\lambda, \varepsilon, K} d^{-1} \log^2 d. \quad (6.26)$$

Below we write  $t = c_{1-\alpha}^G$  for short. Fix  $j \in \{1, \dots, d\}$  arbitrarily. A straightforward computation shows

$$\Psi_{\alpha, jj} = -\mathbb{E}[(\Sigma^{-1} Z)_j 1_{\{\max_{k: k \neq j} Z_k \leq t\}} \mid Z_j = t] \phi(t),$$

where  $Z \sim N(0, \Sigma)$ . Hence

$$|\Psi_{\alpha, jj}| \leq \mathbb{E}[|(\Sigma^{-1} Z)_j| \mid Z_j = t] \phi(t) \leq (\Sigma^{-1})_{jj} |t| \phi(t) + \mathbb{E}[|(\Sigma^{-1})_{j, -j} \cdot Z_{-j}| \mid Z_j = t] \phi(t), \quad (6.27)$$

where  $(\Sigma^{-1})_{j,-j} := ((\Sigma^{-1})_{jk})_{1 \leq k \leq d: k \neq j}$  and  $Z_{-j} := (Z_k)_{1 \leq k \leq d: k \neq j}$ . It is well-known that the conditional distribution of  $Z_{-j}$  given  $Z_j = t$  is  $N(t\Sigma_{j,-j}, \Sigma_{-j,-j} - \Sigma_{j,-j}^{\otimes 2})$  with  $\Sigma_{j,-j} := (\Sigma_{jk})_{1 \leq k \leq d: k \neq j}$  and  $\Sigma_{-j,-j} := (\Sigma_{kl})_{1 \leq k, l \leq d: k, l \neq j}$  (see e.g. Theorem 1.2.5 in [34]). Therefore,

$$\mathbb{E}[(\Sigma^{-1})_{j,-j} \cdot Z_{-j} \mid Z_j = t] = (\Sigma^{-1})_{j,-j} \cdot t\Sigma_{j,-j} = t \sum_{k:k \neq j} (\Sigma^{-1})_{jk} \Sigma_{jk} = t(1 - (\Sigma^{-1})_{jj})$$

and

$$\begin{aligned} \text{Var}[(\Sigma^{-1})_{j,-j} \cdot Z_{-j} \mid Z_j = t] &= (\Sigma^{-1})_{j,-j}^\top (\Sigma_{-j,-j} - \Sigma_{j,-j}^{\otimes 2}) (\Sigma^{-1})_{j,-j} \\ &= \sum_{k,l:k,l \neq j} (\Sigma^{-1})_{jk} (\Sigma^{-1})_{jl} (\Sigma_{kl} - \Sigma_{jk} \Sigma_{jl}) \\ &= \sum_{k:k \neq j} (\Sigma^{-1})_{jk} (-(\Sigma^{-1})_{jj} \Sigma_{kj} - \Sigma_{jk} + \Sigma_{jk} (\Sigma^{-1})_{jj}) \\ &= (\Sigma^{-1})_{jj} - 1. \end{aligned}$$

Consequently,

$$|\mathbb{E}[(\Sigma^{-1})_{j,-j} \cdot Z_{-j} \mid Z_j = t]| \leq |t(1 - (\Sigma^{-1})_{jj})| + \sqrt{(\Sigma^{-1})_{jj} - 1} \leq C_\lambda(1 + |t|). \quad (6.28)$$

Combining (6.27) and (6.28) with (6.23) gives

$$|\Psi_{\alpha,jj}| \leq C_\lambda(1 + |t|)\phi(t) \leq C_{\lambda,\varepsilon,K} d^{-1} \log d. \quad (6.29)$$

Next, fix  $l \in \{1, \dots, d\} \setminus \{j\}$  arbitrarily. Then we have

$$|\Psi_{\alpha,jl}| \leq P\left(\max_{k:k \neq j,l} Z_k \leq t \mid Z_j = t, Z_l = t\right) \phi_{\Sigma[j,l]}(t, t) \leq \phi_{\Sigma[j,l]}(t, t),$$

where  $\Sigma[j, l] = (\Sigma_{pq})_{p,q \in \{j,l\}}$ . Hence

$$|\Psi_{\alpha,jl}| \leq \frac{1}{2\pi\sqrt{1-\rho^2}} \exp\left(-\frac{t^2}{1+\Sigma_{jl}}\right).$$

Let  $\mathcal{I}_j := \{k \in \{1, \dots, d\} : |\Sigma_{jk}| > t^{-2}\}$ . Then we have

$$K^2 \geq \sum_{k=1}^d \Sigma_{jk}^2 \geq \sum_{k \in \mathcal{I}_j} \Sigma_{jk}^2 > t^{-4} \#\mathcal{I}_j,$$

where  $\#\mathcal{I}_j$  is the number of elements in  $\mathcal{I}_j$ . Hence we obtain

$$\begin{aligned} \sum_{l:l \neq j} |\Psi_{\alpha,jl}| &\leq C_\lambda \sum_{l \in \mathcal{I}_j} e^{-t^2/(1+\rho)} + C_\lambda \sum_{l \notin \mathcal{I}_j} e^{-t^2/(1+t^{-2})} \leq C_\lambda (K^2 t^4 e^{-t^2/(1+\rho)} + d e^{-t^2}) \\ &\leq C_{\lambda,\varepsilon,K} \left( d^{-2/(1+\rho)} (\log d)^{2+1/(1+\rho)} + d^{-1} \log d \right) \leq C_{\lambda,\varepsilon,K} d^{-1} \log d, \end{aligned}$$

where the third inequality follows from (6.23). Combining this with (6.29) gives (6.26).  $\square$

#### 6.4 Proof of Theorem 3.4

For  $\alpha \in (0, 1)$ , we denote by  $\hat{c}_{1-\alpha}^*$  the  $(1 - \alpha)$ -quantile of  $T_n^{**}$  under  $P^{**}$ .

**Lemma 6.4.** *Under the assumptions of Theorem 3.4, there exist positive constants  $c$  and  $C$  depending only on  $\lambda, \varepsilon, b_w$  and  $b_v$  such that if (3.3) holds, then*

$$\sup_{\varepsilon < \alpha < 1 - \varepsilon} P \left( \left| P^*(T_n^* \geq \hat{c}_{1-\alpha}^*) - (\alpha - R_n(\alpha)) \right| > C \frac{\zeta_d^3 \log^3(dn)}{\sigma_*^3} \frac{1}{n} \log n \right) \leq \frac{5}{n}.$$

**Proof.** The proof is basically a straightforward modification of that of Theorem 3.3. We only give a sketch of the proof with emphasis on relatively major changes.

Fix  $\alpha \in (\varepsilon, 1 - \varepsilon)$  arbitrarily. First, it is not difficult to see that an analogous result to Theorem 3.2 holds for  $T_n^{**}$ . Thus, by a similar argument to the proof of (6.13), we can find a constant  $c$  depending only on  $\lambda, \varepsilon, b_w$  and  $b_v$  and an event  $\mathcal{E}_n^*(\alpha)$  satisfying the following conditions:

(i) If (3.3) holds, then we have on  $\mathcal{E}_n^*(\alpha)$

$$\left| \hat{c}_{1-\alpha}^* - \hat{c}_{1-\alpha} + \frac{1}{2f_\Sigma(e_{1-\alpha}^G)} \left\langle \frac{1}{n} \sum_{i=1}^n (w_i^2 - 1) X_i^{\otimes 2}, \Psi_\alpha \right\rangle \right| \leq \frac{C_{\lambda, \varepsilon, b_w, b_v} \zeta_d^3 \log^3(dn)}{\sqrt{\log d} \sigma_*^2} \frac{1}{n} \log n.$$

(ii) (2.11) holds on  $\mathcal{E}_n^*(\alpha)$ .

(iii)  $P(\mathcal{E}_n^*(\alpha)) \geq 1 - 1/n^2$ .

In the sequel we assume (3.3) is satisfied with the above  $c$ . Then, by a similar argument to the proof of (6.14), we obtain

$$\left| P^*(T_n^* \geq \hat{c}_{1-\alpha}^*) - P^*(T_n^* \geq \hat{c}_{1-\alpha} - J_n^*(\alpha)) \right| \leq C_{\lambda, \varepsilon, b_w, b_v} \frac{\zeta_d^3 \log^3(dn)}{\sigma_*^3} \frac{1}{n} \log n + P^*(\mathcal{E}_n^*(\alpha)^c),$$

where  $J_n^*(\alpha) := \langle n^{-1} \sum_{i=1}^n (w_i^2 - 1) X_i^{\otimes 2}, \tilde{\Psi}_\alpha \rangle$  and  $\tilde{\Psi}_\alpha$  is defined as in (6.16). Since  $P(P^*(\mathcal{E}_n^*(\alpha)^c) \geq 1/n) \leq n \mathbb{E}[P^*(\mathcal{E}_n^*(\alpha)^c)] \leq 1/n$  by Markov's inequality and (iii), we conclude

$$\left| P^*(T_n^* \geq \hat{c}_{1-\alpha}^*) - P^*(T_n^* \geq \hat{c}_{1-\alpha} - J_n^*(\alpha)) \right| \leq C_{\lambda, \varepsilon, b_w, b_v} \frac{\zeta_d^3 \log^3(dn)}{\sigma_*^3} \frac{1}{n} \log n$$

with probability at least  $1 - 1/n$ . As in the proof of Theorem 3.3, we derive an Edgeworth expansion for  $T_n^* + J_n^*(\alpha)$  by applying Theorem 5.1 with  $\xi_i = (w_i \tilde{X}_i + (w_i^2 - 1) U_i \mathbf{1}_d) / \sqrt{n}$  conditional on the data, where  $\tilde{X}_i := X_i - \bar{X}$  and  $U_i := \langle X_i^{\otimes 2}, \tilde{\Psi}_\alpha \rangle / \sqrt{n}$ . Conditional on the data,  $\xi_i$  has a Stein kernel  $\tau_i$  such that

$$\tau_i(\xi_i) = \frac{\bar{\tau}_{i,11}(w_i, w_i^2) \tilde{X}_i^{\otimes 2} + \bar{\tau}_{i,12}(w_i, w_i^2) V_i + \bar{\tau}_{i,21}(w_i, w_i^2) V_i^\top + \bar{\tau}_{i,22}(w_i, w_i^2) V_i'}{n},$$

where  $V_i := \tilde{X}_i U_i \mathbf{1}_d^\top$  and  $V_i' := U_i^2 \mathbf{1}_d^{\otimes 2}$ . It is not difficult to check that we have the estimates corresponding to (5.24)–(5.27) in the present setting with probability at least  $1 - 1/n$  by a similar argument to the proof of Theorem 2.3. Meanwhile, by the Schwarz inequality,

$$\left\| \frac{1}{n} \sum_{i=1}^n (\bar{\tau}_{i,12}(w_i, w_i^2) V_i + \bar{\tau}_{i,21}(w_i, w_i^2) V_i^\top) \right\|_\infty \leq \frac{2b_w^3}{\sqrt{n}} \sqrt{\frac{1}{n} \sum_{i=1}^n \langle X_i^{\otimes 2}, \tilde{\Psi}_\alpha \rangle^2} \sqrt{\max_{1 \leq j \leq d} \frac{1}{n} \sum_{i=1}^n \tilde{X}_{ij}^2}$$

and

$$\left\| \frac{1}{n} \sum_{i=1}^n \bar{\tau}_{i,22}(w_i, w_i^2) V_i' \right\|_{\infty} \leq \frac{b_w^4}{n} \frac{1}{n} \sum_{i=1}^n \langle X_i^{\otimes 2}, \tilde{\Psi}_{\alpha} \rangle^2.$$

Observe that  $\langle X_i^{\otimes 2}, \tilde{\Psi}_{\alpha} \rangle^2 = \langle X_i^{\otimes 4}, \tilde{\Psi}_{\alpha}^{\otimes 2} \rangle$ . Hence, by (6.17) and Lemma E.10,

$$\frac{1}{n} \sum_{i=1}^n \langle X_i^{\otimes 2}, \tilde{\Psi}_{\alpha} \rangle^2 \leq C_{\varepsilon} \frac{b_w^4 \zeta_d^2 \log d}{\sigma_*^4} \leq C_{\lambda, \varepsilon} \frac{b^2 \zeta_d^2 \log d}{\sigma_*^2}$$

with probability at least  $1 - 1/n$ . Combining these estimates with (5.16) and the argument to prove (5.22), we obtain the estimate corresponding to (5.22) with  $b_w b$  replaced by  $C_{\lambda, \varepsilon, b_w, b_v} b \sqrt{\zeta_d / \sigma_*}$  with probability at least  $1 - 2/n$ . All together, we can proceed as in the proof of Theorem 2.3 and then obtain

$$\sup_{t \in \mathbb{R}} \left| P^*(T_n^* + J_n^*(\alpha) \leq t) - \int_{A(t)} (\hat{p}_{n,1}(z) + \hat{q}_n(z)) dz \right| \leq C_{\lambda, \varepsilon, b_w, b_v} \frac{\zeta_d^3 \log^3(dn)}{\sigma_*^3 n} \log n$$

with probability at least  $1 - 3/n$ , where

$$\hat{q}_n(z) = \frac{1}{2n} \sum_{i=1}^n \langle 2\tilde{X}_i U_i \mathbf{1}_d^{\top} + U_i^2 \mathbf{1}_d^{\otimes 2}, \nabla^2 \phi_{\Sigma}(z) \rangle - \frac{1}{6n^{3/2}} \sum_{i=1}^n \langle \mathbb{E}^*[(w_i \tilde{X}_i + U_i \mathbf{1}_d)^{\otimes 3}] - \tilde{X}_i^{\otimes 3}, \nabla^3 \phi_{\Sigma}(z) \rangle.$$

The remaining proof is a minor modification of the proof of (6.21), so we omit the details.  $\square$

**Lemma 6.5.** *Under the assumptions of Theorem 3.4, there exists a constant  $C > 0$  depending only on  $\lambda$  and  $\varepsilon$  such that*

$$|\mathbb{E}[R_n(\alpha + \delta)] - \mathbb{E}[R_n(\alpha)]| \leq C \delta \frac{\zeta_d^3}{\sigma_*^3} \sqrt{\frac{\log^3 d}{n}}. \quad (6.30)$$

for any  $\alpha \in (\varepsilon, 1 - 2\varepsilon)$  and  $\delta \in (0, \varepsilon]$ .

**Proof.** By Lemma E.4 and (6.5),

$$|\mathbb{E}[R_n(\alpha + \delta)] - \mathbb{E}[R_n(\alpha)]| \leq C_{\lambda, \varepsilon} \frac{b^3}{\sqrt{n}} \left( \frac{\log^2 d}{\sigma_*^4} \left| \frac{1}{f_{\Sigma}(c_{1-\alpha-\delta}^G)} - \frac{1}{f_{\Sigma}(c_{1-\alpha}^G)} \right| + \frac{\log d}{\sigma_*^2} \frac{\zeta_d}{\sqrt{\log d}} |\Psi_{\alpha+\delta} - \Psi_{\alpha}| \right).$$

Noting that  $(F_Z^{-1})'(p) = 1/f_{\Sigma}(c_p^G)$  for all  $p \in (0, 1)$ , we obtain by the mean value theorem and (6.6)

$$\left| \frac{1}{f_{\Sigma}(c_{1-\alpha-\delta}^G)} - \frac{1}{f_{\Sigma}(c_{1-\alpha}^G)} \right| \leq C_{\varepsilon} \delta \frac{\zeta_d^3}{\sigma_*^2 \sqrt{\log d}}.$$

Also, by Lemma 5.2, the mean value theorem and (6.5),

$$|\Psi_{\alpha+\delta} - \Psi_{\alpha}| \lesssim \frac{\log^{3/2} d}{\sigma_*^3} |c_{1-\alpha-\delta}^G - c_{1-\alpha}^G| \leq C_{\varepsilon} \delta \frac{\zeta_d \log d}{\sigma_*^3}.$$

Combining these bounds gives (6.30).  $\square$

**Proof of Theorem 3.4.** Denote by  $c_1$  and  $C_1$  the constants  $c$  and  $C$  in Theorem 3.3, respectively. Also, denote by  $c_2$  and  $C_2$  the constants  $c$  and  $C$  in Lemma 6.4, respectively. Since the left hand side of (3.10) is bounded by 1, we may assume (3.3) holds with  $c = c_1 \wedge c_2$  without loss of generality. Then, for each  $\alpha \in (\varepsilon, 1 - \varepsilon)$ , the event

$$\mathcal{E}_n(\alpha) := \left\{ \left| P^*(T_n^* \geq \hat{c}_{1-\alpha}^*) - (\alpha - R_n(\alpha)) \right| \leq C_2 \frac{\varsigma_d^3 \log^3(dn)}{\sigma_*^3 n} \log n \right\}$$

occurs with probability at least  $1 - 5/n$ . Meanwhile, by (3.3), (6.5) and Lemmas E.4 and E.9, there exists a constant  $C_3 > 0$  depending only on  $\lambda, \varepsilon$  such that the event

$$\mathcal{E}_n := \left\{ \sup_{\varepsilon < \alpha < 1-\varepsilon} |R_n(\alpha) - \mathbb{E}[R_n(\alpha)]| \leq C_3 \frac{\varsigma_d \log^3(dn)}{\sigma_* n} \right\}$$

occurs with probability at least  $1 - 1/n$  and

$$\sup_{\varepsilon < \alpha < 1-\varepsilon} |\mathbb{E}[R_n(\alpha)]| \leq C_3 \frac{\varsigma_d}{\sigma_*} \sqrt{\frac{\log^3 d}{n}}. \quad (6.31)$$

Further, let  $C_4$  be the constant  $C$  in Lemma 6.5. Set

$$\Delta_n := C_2 \frac{\varsigma_d^3 \log^3(dn)}{\sigma_*^3 n} \log n + C_3 \frac{\varsigma_d \log^3(dn)}{\sigma_* n} + C_3 C_4 \frac{\varsigma_d^4 \log^3 d}{\sigma_*^4 n}.$$

Since the left hand side of (3.10) is bounded by 1, we may assume without loss of generality

$$C_3 \frac{\varsigma_d}{\sigma_*} \sqrt{\frac{\log^3 d}{n}} + 3\Delta_n \leq \varepsilon. \quad (6.32)$$

Now fix  $\alpha \in (2\varepsilon, 1 - 2\varepsilon)$  arbitrarily. Set  $\alpha_{n,1} := \alpha - \Delta_n$  and  $\alpha'_{n,1} := \alpha_{n,1} + \mathbb{E}[R_n(\alpha_{n,1})]$ . By (6.31) and (6.32),  $\alpha_{n,1}, \alpha'_{n,1} \in (\varepsilon, 1 - \varepsilon)$ . Hence, on  $\mathcal{E}_n(\alpha'_{n,1}) \cap \mathcal{E}_n$ ,

$$\begin{aligned} P^*(\hat{F}_n^*(T_n^*) > 1 - \alpha'_{n,1}) &\leq P^*(T_n^* \geq \hat{c}_{1-\alpha'_{n,1}}^*) \leq \alpha'_{n,1} - R_n(\alpha'_{n,1}) + C_2 \frac{\varsigma_d^3 \log^3(dn)}{\sigma_*^3 n} \log n \\ &\leq \alpha_{n,1} + \mathbb{E}[R_n(\alpha_{n,1})] - \mathbb{E}[R_n(\alpha'_{n,1})] + C_3 \frac{\varsigma_d \log^3(dn)}{\sigma_* n} + C_2 \frac{\varsigma_d^3 \log^3(dn)}{\sigma_*^3 n} \log n \leq \alpha, \end{aligned}$$

where the last inequality follows from (6.30) and (6.31). This yields  $\hat{\beta}_\alpha \leq 1 - \alpha'_{n,1}$  on  $\mathcal{E}_n(\alpha'_{n,1}) \cap \mathcal{E}_n$ . Hence

$$\begin{aligned} P(T_n \geq \hat{\beta}_\alpha) &\geq P(T_n \geq \hat{c}_{1-\alpha'_{n,1}}) - \frac{6}{n} \geq \alpha'_{n,1} - \mathbb{E}[R_n(\alpha'_{n,1})] - C_{\lambda,\varepsilon,b_w} \frac{\varsigma_d^3 \log^3(dn)}{\sigma_*^3 n} \log n \\ &\geq \alpha + \mathbb{E}[R_n(\alpha_{n,1})] - \mathbb{E}[R_n(\alpha'_{n,1})] - C_{\lambda,\varepsilon,b_w,b_v} \frac{\varsigma_d^4 \log^3(dn)}{\sigma_*^4 n} \log n \\ &\geq \alpha - C_{\lambda,\varepsilon,b_w,b_v} \frac{\varsigma_d^4 \log^3(dn)}{\sigma_*^4 n} \log n, \end{aligned} \quad (6.33)$$

where the second inequality is by Theorem 3.3, the third by  $\Delta_n \leq C_{\lambda,\varepsilon,b_w,b_v} \frac{\varsigma_d^4 \log^3 d}{\sigma_*^4 n} \log n$  and the fourth by (6.30) and (6.31). Similarly, with  $\alpha_{n,2} := \alpha + 2\Delta_n$  and  $\alpha'_{n,2} := \alpha_{n,2} + \mathbb{E}[R_n(\alpha_{n,2})]$ , we have on  $\mathcal{E}_n(\alpha'_{n,2}) \cap \mathcal{E}_n$

$$P^*(\hat{F}_n^*(T_n^*) > 1 - \alpha'_{n,2} - \Delta_n) \geq P^*(\hat{F}_n^*(T_n^*) \geq 1 - \alpha'_{n,2}) = P^*(T_n^* \geq \hat{c}_{1-\alpha'_{n,2}}^*)$$

$$\geq \alpha'_{n,2} - R_n(\alpha'_{n,2}) - C_2 \frac{\varsigma_d^3 \log^3(dn)}{\sigma_*^3 n} \log n > \alpha.$$

Hence  $\hat{\beta}_\alpha > 1 - \alpha'_{n,2} - \Delta_n$  on  $\mathcal{E}_n(\alpha'_{n,2}) \cap \mathcal{E}_n$ . Therefore, a similar argument to (6.33) yields

$$P(T_n \geq \hat{c}_{\hat{\beta}_\alpha}) \leq \alpha + C_{\lambda, \varepsilon, b_w, b_v} \frac{\varsigma_d^4 \log^3(dn)}{\sigma_*^4 n} \log n.$$

Combining this and (6.33) gives the desired result.  $\square$

## Appendix

### A Nearly optimal high-dimensional CLT under the sub-exponential condition

**Theorem A.1.** *Set  $\sigma_j := \sqrt{\text{Var}[S_{n,j}]}$  for  $j = 1, \dots, d$ . Suppose that there exists a constant  $B \geq 1$  such that  $\max_{i,j} \|X_{ij}/\sigma_j\|_{\psi_1} \leq B$  and  $\max_j n^{-1} \sum_{i=1}^n \mathbb{E}[(X_{ij}/\sigma_j)^2] \leq B^2$ . Then there exists a universal constant  $C > 0$  such that*

$$\sup_{A \in \mathcal{R}} |P(S_n \in A) - P(Z \in A)| \leq \frac{C}{\rho_*^2} \sqrt{\frac{B^2 \log^3(dn)}{n}} \log n, \quad (\text{A.1})$$

where  $Z \sim N(0, \text{Cov}[S_n])$  and  $\rho_*^2$  is the minimum eigenvalue of the correlation matrix of  $S_n$ .

**Proof.** As announced, the proof is a combination of [23, Theorem 2.1] and a simple truncation argument used in [36, Section 5.2] and [29, Section 4.3]. Denote by  $\delta$  the left hand side of (A.1). Considering  $X_{ij}/\sigma_j$  instead of  $X_{ij}$ , we may assume  $\sigma_j = 1$  for all  $j$  without loss of generality. Further, since  $\delta \leq 1$ , we may also assume

$$\frac{1}{\rho_*^2} \sqrt{\frac{B^2 \log^3(dn)}{n}} \log n \leq 1. \quad (\text{A.2})$$

Next, let  $\kappa_n := 2B \log n$ . For  $i = 1, \dots, n$  and  $j = 1, \dots, d$ , define  $\hat{X}_{ij} := X_{ij} 1_{\{|X_{ij}| \leq \kappa_n\}} - \mathbb{E}[X_{ij} 1_{\{|X_{ij}| \leq \kappa_n\}}]$  and set  $\hat{X}_i = (\hat{X}_{i1}, \dots, \hat{X}_{id})^\top$  and  $\hat{S}_n := n^{-1/2} \sum_{i=1}^n \hat{X}_i$ . Note that  $\max_{1 \leq i \leq n} \|\hat{X}_i\|_\infty \leq 2\kappa_n$ . Then, by a similar argument to the proof of Eq.(4.19) in [29], we obtain

$$\delta \lesssim \frac{1}{n} + \frac{B \log(dn) \sqrt{\log d}}{\sqrt{n}} \log n + \hat{\delta},$$

where  $\hat{\delta} := \sup_{A \in \mathcal{R}} |P(\hat{S}_n \in A) - P(Z \in A)|$ . Since  $\rho_*^2 \leq \sigma_1^2 = 1$ , it remains to prove

$$\hat{\delta} \lesssim \frac{1}{\rho_*^2} \sqrt{\frac{B^2 \log^3(dn)}{n}} \log n. \quad (\text{A.3})$$

We prove this bound by applying Theorem 2.1 in [23] with  $\psi = 2\kappa_n$ . This gives

$$\hat{\delta} \lesssim (\log n) \left( \Delta_0 + \sqrt{\frac{B^2 \log^3 d}{n \rho_*^4} + \frac{\kappa_n^2 \log^2 d}{n \rho_*^2}} \right) + \frac{\kappa_n \log^{3/2} d}{\rho_* \sqrt{n}}, \quad (\text{A.4})$$

where  $\Delta_0 := \frac{\log d}{\rho_*^2} \|\text{Cov}(\hat{S}_n) - \text{Cov}(S_n)\|_\infty$ . By (A.2) and  $\rho_* \leq 1$ ,

$$\frac{\kappa_n^2 \log^2 d}{n \rho_*^2} = \frac{4}{\rho_*^2} \sqrt{\frac{B^2 \log^3 d}{n}} \sqrt{\frac{B^2 (\log d) (\log^2 n)}{n}} \log n \leq \frac{4}{\rho_*^2} \sqrt{\frac{B^2 \log^3(dn)}{n}}$$

and

$$\frac{\kappa_n \log^{3/2} d}{\rho_* \sqrt{n}} \leq \frac{2}{\rho_*^2} \sqrt{\frac{B^2 \log^3(dn)}{n}} \log n.$$

Further, by Eq.(26) in [36] and (A.2),

$$\Delta_0 \lesssim \frac{\log d}{\rho_*^2} e^{-\kappa_n/(2B)} B^2 \log n = \frac{1}{\rho_*^2} \frac{B^2 \log d}{n} \log n \leq \frac{1}{\rho_*^2} \sqrt{\frac{B^2 \log^3(dn)}{n}}.$$

Consequently, we obtain (A.3) from (A.4). □

## B Proofs of the auxiliary results in Section 5.2

### B.1 Proof of Lemma 5.1

As already mentioned, the proof is a straightforward modification of [7, Lemma 11.4]. Let

$$\delta := \sup \left\{ \left| \int h_y d(\mu - \nu) \right| : y \in \mathbb{R}^d \right\}.$$

Assume first that

$$\delta = \sup \left\{ \int h_y d(\mu - \nu) : y \in \mathbb{R}^d \right\}. \quad (\text{B.1})$$

Then, given any  $\eta > 0$ , there exists a vector  $z \in \mathbb{R}^d$  such that  $\int h_z d(\mu - \nu) \geq \delta - \eta$ . In this case, we have

$$\begin{aligned} & \int_{[-\varepsilon, \varepsilon]^d} \left[ \int M_{h_z}(y+x; \varepsilon)(\mu - \nu)(dy) \right] K(dx) \\ & \geq \int_{[-\varepsilon, \varepsilon]^d} \left[ \int h_z(y) \mu(dy) - \int M_{h_z}(y+x; \varepsilon) \nu(dy) \right] K(dx) \\ & = \int_{[-\varepsilon, \varepsilon]^d} \left[ \int h_z(y) (\mu - \nu)(dy) - \int \{M_{h_z}(y+x; \varepsilon) - h_z(y)\} \nu(dy) \right] K(dx) \\ & \geq \int_{[-\varepsilon, \varepsilon]^d} \left[ \delta - \eta - \int \{M_{h_z}(y+x; \varepsilon) - h_z(y+x)\} \nu(dy) - \int \{h_z(y+x) - h_z(y)\} \nu(dy) \right] K(dx) \\ & \geq \int_{[-\varepsilon, \varepsilon]^d} [\delta - \eta - \tau^*(h; \varepsilon) - \tilde{\tau}^*(h; \varepsilon)] K(dx) = \alpha [\delta - \eta - \tau^*(h; \varepsilon) - \tilde{\tau}^*(h; \varepsilon)] \end{aligned}$$

and

$$\begin{aligned} & \int_{\mathbb{R}^d \setminus [-\varepsilon, \varepsilon]^d} \left[ \int M_{h_z}(y+x; \varepsilon)(\mu - \nu)(dy) \right] K(dx) \\ & \geq \int_{\mathbb{R}^d \setminus [-\varepsilon, \varepsilon]^d} \left[ \int h_z(y+x) \mu(dy) - \int M_{h_z}(y+x; \varepsilon) \nu(dy) \right] K(dx) \\ & = \int_{\mathbb{R}^d \setminus [-\varepsilon, \varepsilon]^d} \left[ \int h_z(y+x) (\mu - \nu)(dy) - \int \{M_{h_z}(y+x; \varepsilon) - h_z(y+x)\} \nu(dy) \right] K(dx) \\ & \geq \int_{\mathbb{R}^d \setminus [-\varepsilon, \varepsilon]^d} [-\delta - \tau^*(h; \varepsilon)] K(dx) = (1 - \alpha) [-\delta - \tau^*(h; \varepsilon)]. \end{aligned}$$

Consequently, we obtain

$$\gamma^*(h; \varepsilon) \geq \int M_{h_z}(x; \varepsilon)(\mu - \nu) * K(dx) \geq (2\alpha - 1)\delta - \tau^*(h; \varepsilon) - \alpha \tilde{\tau}^*(h; \varepsilon) - \alpha \eta.$$

Letting  $\eta \downarrow 0$ , we obtain the desired result. If instead of (B.1) we have

$$\delta = \sup \left\{ - \int h_y d(\mu - \nu) : y \in \mathbb{R}^d \right\},$$

then, given any  $\eta > 0$ , we can find  $z \in \mathbb{R}^d$  such that  $-\int h_z d(\mu - \nu) \geq \delta - \eta$ . Now look at  $-h_z$  (instead of  $h_z$ ) and note that  $M_{-h_y}(\cdot; \varepsilon) = -m_{h_y}(\cdot; \varepsilon)$  and

$$\int \{h_y - m_{h_y}(x; \varepsilon)\} \nu(dx) = \int \{M_{-h_y}(x; \varepsilon) - (-h_y)\} \nu(dx)$$

for every  $y \in \mathbb{R}^d$ . Proceeding exactly as above, we obtain

$$\gamma^*(h; \varepsilon) \geq - \int m_{h_z}(x; \varepsilon) (\mu - \nu) * K(dx) \geq (2\alpha - 1)\delta - \tau^*(h; \varepsilon) - \alpha \tilde{\tau}^*(h; \varepsilon) - \alpha \eta.$$

Thus we complete the proof.  $\square$

## B.2 Proof of Lemma 5.2

We divide the proof into four steps.

**Step 1.** First we reduce the proof to the case  $\Sigma = I_d$ . Let  $Z \sim N(0, \Sigma)$  and  $Z' \sim N(0, \Sigma - \sigma_*^2 I_d)$ . Then, for any  $A \in \mathcal{R}$ ,  $u, v \in \mathbb{R}_+^d$  and  $x \in \mathbb{R}^d$ , we have

$$\int_{A^{u,v} \setminus A} \phi_\Sigma(x+z) dz = \mathbb{E}[1_{A^{u,v} \setminus A}(Z-x)] = \mathbb{E} \left[ \int_{\mathbb{R}^d} 1_{A^{u,v} \setminus A}(\sigma_* z + Z') \phi_d(z+x/\sigma_*) dz \right].$$

Differentiating the both sides  $r$  times with respect to  $x$  and then setting  $x = 0$ , we obtain

$$\int_{A^{u,v} \setminus A} \nabla^r \phi_\Sigma(z) dz = \frac{1}{\sigma_*^r} \mathbb{E} \left[ \int_{\mathbb{R}^d} 1_{A^{u,v} \setminus A}(\sigma_* z + Z') \phi_d(z) dz \right].$$

Observe that  $1_{A^{u,v} \setminus A}(\sigma_* z + Z') = 1_{(\sigma_*^{-1}(A-Z'))^{u/\sigma_*, v/\sigma_*} \setminus (\sigma_*^{-1}(A-Z'))}(z)$  and  $\sigma_*^{-1}(A-Z') \in \mathcal{R}$ . Hence

$$\left\| \int_{A^{u,v} \setminus A} \nabla^r \phi_\Sigma(z) dz \right\|_1 \leq \frac{1}{\sigma_*^r} \sup_{A \in \mathcal{R}} \left\| \int_{A^{u/\sigma_*, v/\sigma_*} \setminus A} \nabla^r \phi_d(z) dz \right\|_1.$$

Therefore, the claim for general  $\Sigma$  follows from that for  $\Sigma = I_d$ .

**Step 2.** In this and the next steps, we show that the quantity inside  $\sup_{A \in \mathcal{A}}$  on the left hand side of (5.33) can be replaced by a weighted surface integral of  $\nabla^r \phi_\Sigma$  over the boundary of  $A$ . Note that an analogous result for the case  $r = 0$  is standard in the literature; see e.g. Proposition 1.1 in [53]. For  $A \in \mathcal{R}$ ,  $u, v \in \mathbb{R}_+^d$  and  $\varepsilon > 0$ , set

$$I_A(u, v) = \left\| \int_{A^{u,v} \setminus A} \nabla^r \phi_d(z) dz \right\|_1, \quad K(\varepsilon) = \sup_{A \in \mathcal{R}; u, v \in [0, \varepsilon]^d} \frac{I_A(u, v)}{\varepsilon}.$$

In this step, we prove

$$\sup_{\varepsilon > 0} K(\varepsilon) = \limsup_{\varepsilon \downarrow 0} K(\varepsilon).$$

Take  $\varepsilon > 0$  arbitrarily. For any  $A \in \mathcal{R}$  and  $u, v \in \mathbb{R}_+^d$ , observe that  $A^{u/2, v/2} \in \mathcal{R}$  and  $A^{u, v} \setminus A$  is the disjoint union of  $(A^{u/2, v/2})^{u/2, v/2} \setminus (A^{u/2, v/2})$  and  $A^{u/2, v/2} \setminus A$ . This implies that

$$\sup_{A \in \mathcal{R}; u, v \in [0, \varepsilon]^d} I_A(u, v) \leq 2 \sup_{A \in \mathcal{R}; u, v \in [0, \varepsilon/2]^d} I_A(u, v),$$

and thus  $K(\varepsilon) \leq K(\varepsilon/2)$ . Repeating this procedure gives  $K(\varepsilon) \leq K(\varepsilon/2^n)$  for  $n = 1, 2, \dots$ . Hence

$$K(\varepsilon) \leq \limsup_{n \rightarrow \infty} K(\varepsilon/2^n) \leq \limsup_{\eta \downarrow 0} K(\eta).$$

Since  $\varepsilon$  is arbitrary, we obtain the desired result.

**Step 3.** For any Borel set  $A \subset \mathbb{R}^{d-1}$ ,  $j \in \{1, \dots, d\}$  and  $s \in \mathbb{R}$ , define

$$J_{A,j}(s) = \int_A \nabla^r \phi_d(z|_{z_j=s}) dz_1 \cdots \widehat{dz_j} \cdots dz_d,$$

where  $z|_{z_j=s} = (z_1, \dots, z_{j-1}, s, z_{j+1}, \dots, z_d)^\top$  and  $\widehat{dz_j}$  means that  $dz_j$  is omitted. Then, for  $u, v \in \mathbb{R}^d$  and  $A = \prod_{j=1}^d [a_j, b_j] \in \mathcal{R}$ , set

$$L_A(u, v) = \sum_{j=1}^d \{u_j J_{A^j, j}(a_j) + v_j J_{A^j, j}(b_j)\},$$

where  $A^j = \prod_{k:k \neq j} [a_k, b_k]$ . In this step, we prove

$$\limsup_{\varepsilon \downarrow 0} K(\varepsilon) = \limsup_{\varepsilon \downarrow 0} \sup_{A \in \mathcal{R}; u, v \in [0, \varepsilon]^d} \frac{\|L_A(u, v)\|_1}{\varepsilon}.$$

Fix  $A = \prod_{j=1}^d [a_j, b_j] \in \mathcal{R}$ ,  $\varepsilon > 0$  and  $u, v \in [0, \varepsilon]^d$ . For  $j_1, \dots, j_r \in \{1, \dots, d\}$ , we set

$$A_{j_1, \dots, j_r}^{u, v} = \{x \in A^{u, v} : x_j \notin [a_j, b_j] \text{ for } j \in \{j_1, \dots, j_r\} \text{ and } x_j \in [a_j, b_j] \text{ for } j \notin \{j_1, \dots, j_r\}\}.$$

Then,  $A^{u, v} \setminus A = \bigcup_{r=1}^d \bigcup_{1 \leq j_1 < \dots < j_r \leq d} A_{j_1, \dots, j_r}^{u, v}$  and this is a disjoint union. Hence we have

$$I_A(u, v) = \left\| \sum_{r=1}^d \sum_{1 \leq j_1 < \dots < j_r \leq d} \int_{A_{j_1, \dots, j_r}^{u, v}} \nabla^r \phi_d(z) dz \right\|_1.$$

One can easily check that

$$\sup_{A \in \mathcal{R}; u, v \in [0, \varepsilon]^d} \sum_{r=2}^d \sum_{1 \leq j_1 < \dots < j_r \leq d} \int_{A_{j_1, \dots, j_r}^{u, v}} \|\nabla^r \phi_d(z)\|_1 dz = O(\varepsilon^2) \quad \text{as } \varepsilon \downarrow 0.$$

Hence we obtain

$$\limsup_{\varepsilon \downarrow 0} K(\varepsilon) = \limsup_{\varepsilon \downarrow 0} \sup_{A \in \mathcal{R}; u, v \in [0, \varepsilon]^d} \left\| \frac{1}{\varepsilon} \sum_{j=1}^d \int_{A_j^{u, v}} \nabla^r \phi_d(z) dz \right\|_1.$$

For any  $j \in \{1, \dots, d\}$ , observe that  $A_j^{u,v}$  is the disjoint union of

$$\underline{A}_j^u := [a_1, b_1] \times \cdots \times [a_{j-1}, b_{j-1}] \times [a_j - u_j, a_j] \times [a_{j+1}, b_{j+1}] \times \cdots \times [a_d, b_d]$$

and

$$\overline{A}_j^v := [a_1, b_1] \times \cdots \times [a_{j-1}, b_{j-1}] \times (b_j, b_j + v_j] \times [a_{j+1}, b_{j+1}] \times \cdots \times [a_d, b_d].$$

Therefore, we have

$$\begin{aligned} & \left\| \sum_{j=1}^d \int_{A_j^{u,v}} \nabla^r \phi_d(z) dz - L_A(u, v) \right\|_1 \\ &= \left\| \sum_{j=1}^d \left( \int_{\underline{A}_j^u} \{\nabla^r \phi_d(z) - \nabla^r \phi_d(z|_{z_j=a_j})\} dz + \int_{\overline{A}_j^v} \{\nabla^r \phi_d(z) - \nabla^r \phi_d(z|_{z_j=b_j})\} dz \right) \right\|_1 \\ &\leq 2\varepsilon \sum_{j=1}^d \int_{\mathbb{R}^{d-1}} \sup_{s, t \in \mathbb{R}: |s-t| \leq \varepsilon} \|\nabla^r \phi_d(z|_{z_j=s}) - \nabla^r \phi_d(z|_{z_j=t})\|_1 dz_1 \cdots \widehat{dz}_j \cdots dz_d. \end{aligned}$$

Thus,

$$\limsup_{\varepsilon \downarrow 0} \sup_{A \in \mathcal{R}; u, v \in [0, \varepsilon]^d} \frac{1}{\varepsilon} \left\| \sum_{j=1}^d \int_{A_j^{u,v}} \nabla^r \phi_d(z) dz - L_A(u, v) \right\|_1 = 0.$$

This gives the desired result.

**Step 4.** It remains to prove

$$\limsup_{\varepsilon \downarrow 0} \sup_{A \in \mathcal{R}; u, v \in R_+(\varepsilon)} \frac{\|L_A(u, v)\|_1}{\varepsilon} \leq C_r (\log d)^{(r+1)/2}. \quad (\text{B.2})$$

We first note that if  $A$  is an orthant, i.e.  $a_j = -\infty$  for all  $j$ , then (B.2) immediately follows from the fundamental theorem of calculus and Lemma E.4. In fact, we have in this case

$$\frac{\|L_A(u, v)\|_1}{\varepsilon} \leq \left\| \int_A \nabla^{r+1} \phi_d(z) dz \right\|_1$$

for any  $\varepsilon > 0$  and  $u, v \in [0, \varepsilon]^d$ . In the following we show that the proof is essentially reduced to this case by a similar argument to the proof of [26, Lemma 2.2]. For every  $q \in \{1, \dots, r\}$ , set

$$\mathcal{N}_q(r) = \{(\nu_1, \dots, \nu_q) \in \mathbb{Z}^q : \nu_1, \dots, \nu_q \geq 0, \nu_1 + \cdots + \nu_q = r\}.$$

Also, for any  $m \in \mathbb{N}$ , let

$$\mathcal{J}_m(d) = \{(j_1, \dots, j_m) \in \{1, \dots, d\}^m : j_1, \dots, j_m \text{ are distinct}\}.$$

Then, for any  $A \in \mathcal{R}$ ,  $j \in \{1, \dots, d\}$  and  $s \in \mathbb{R}$ , we have

$$\|J_{A^j, j}(s)\|_1 = \sum_{j_1, \dots, j_r=1}^d \left| \int_{A^j} \partial_{j_1, \dots, j_r} \phi_d(z|_{z_j=s}) dz_1 \cdots \widehat{dz}_j \cdots dz_d \right|$$

$$\begin{aligned}
&\leq C_r \sum_{q=1}^r \sum_{(\nu_1, \dots, \nu_q) \in \mathcal{N}_q(r)} \sum_{(j_1, \dots, j_q) \in \mathcal{J}_q(d)} \left| \int_{A^j} \partial_{j_1}^{\nu_1} \cdots \partial_{j_q}^{\nu_q} \phi_d(z|_{z_j=s}) dz_1 \cdots \widehat{dz_j} \cdots dz_d \right| \\
&=: C_r \sum_{q=1}^r \sum_{\nu=(\nu_1, \dots, \nu_q) \in \mathcal{N}_q(r)} \sum_{\mathbf{j}=(j_1, \dots, j_q) \in \mathcal{J}_q(d)} \Lambda_j(\nu, \mathbf{j}).
\end{aligned}$$

For each  $r = 1, \dots, q$ , the cardinality of the set  $\mathcal{N}_q(r)$  is bounded by a constant depending only on  $r$ . Therefore, to prove (B.2), it suffices to show that

$$\sum_{j=1}^d \sum_{\mathbf{j}=(j_1, \dots, j_r) \in \mathcal{J}_q(d)} \Lambda_j(\nu, \mathbf{j}) \leq C_r (\log d)^{(r+1)/2} \quad (\text{B.3})$$

for any (fixed)  $q \in \{1, \dots, r\}$ ,  $\nu = (\nu_1, \dots, \nu_q) \in \mathcal{N}_q(r)$ ,  $A = \prod_{j=1}^d [a_j, b_j] \in \mathcal{R}$  and  $s \in \{a_j, b_j\}$ ,  $j = 1, \dots, d$ .

To prove (B.3), we introduce additional notation. For a non-negative integer  $m$ ,  $H_m$  denotes the  $m$ -th Hermite polynomial, i.e.  $H_m(t) = (-1)^m \phi(t)^{-1} \phi^{(m)}(t)$ . When  $m \geq 1$ , we set  $h_m(t) = H_{m-1}(t) \phi(t)$ . Also, we denote by  $t_m$  the maximum root of  $H_m$ . For example,  $t_1 = 0, t_2 = 1, t_3 = \sqrt{3}$ . Finally, set  $M_m := \max_{0 \leq t \leq t_m} |H_{m-1}(t)| < \infty$  and define

$$\tilde{h}_m(t) = M_m \phi(t) 1_{[0, t_m]}(t) + h_m(t) 1_{(t_m, \infty)}(t).$$

The function  $\tilde{h}_m$  satisfies the following properties by Lemma A.1 in [26]:

$$\tilde{h}_m \text{ is decreasing on } [0, \infty). \quad (\text{B.4})$$

$$|h_m(t)| \leq \tilde{h}_m(|t|) \text{ for all } t \in \mathbb{R}. \quad (\text{B.5})$$

Now, we fix  $\mathbf{j} = (j_1, \dots, j_q) \in \mathcal{J}_q(d)$  and  $j \in \{1, \dots, d\}$  for a while. Set

$$\nu = \begin{cases} \nu_p & \text{if } j = j_p \text{ for some } p \in \{1, \dots, q\}, \\ 0 & \text{otherwise.} \end{cases}$$

Then we have

$$\begin{aligned}
\Lambda_j(\nu, \mathbf{j}) &= |h_{\nu+1}(s)| \left( \prod_{p: j_p \neq j} |h_{\nu_p}(b_{j_p}) - h_{\nu_p}(a_{j_p})| \right) \prod_{k: k \neq j_1, \dots, j_q, j} \{\Phi(b_k) - \Phi(a_k)\} \\
&\leq \tilde{h}_{\nu+1}(|s|) \left( \prod_{p: j_p \neq j} (\tilde{h}_{\nu_p}(|b_{j_p}|) + \tilde{h}_{\nu_p}(|a_{j_p}|)) \right) \prod_{k: k \neq j_1, \dots, j_q, j} \{\Phi(b_k) + \Phi(-a_k) - 1\},
\end{aligned}$$

where the last inequality follows from (B.5) and the identity  $1 - \Phi(t) = \Phi(-t)$ . Set  $c_k = |a_k| \wedge |b_k|$  for  $k = 1, \dots, d$ . Then we have  $\Phi(b_k) + \Phi(-a_k) - 1 \leq \min\{\Phi(b_k), \Phi(-a_k)\} \leq \Phi(c_k)$ . Combining this with (B.4) gives

$$\Lambda_j(\nu, \mathbf{j}) \leq 2^q \tilde{h}_{\nu+1}(c_j) \left( \prod_{p: j_p \neq j} \tilde{h}_{\nu_p}(c_{j_p}) \right) \prod_{k: k \neq j_1, \dots, j_q, j} \Phi(c_k).$$

Now, observe that  $\tilde{h}_m(t) \leq C_m(1 + t^{m-1})\phi(t)$  for any  $m \in \mathbb{N}$  and  $t \geq 0$  by construction. Hence, if  $\max_{p=1, \dots, q} c_{j_p} \leq \sqrt{4(r+1) \log d}$ ,

$$\Lambda_j(\nu, \mathbf{j}) \leq \begin{cases} C_r (\log d)^{(r-q+1)/2} \left( \prod_{p=1}^q \phi(c_{j_p}) \right) \prod_{k: k \neq j_1, \dots, j_q} \Phi(c_k) & \text{if } j \in \{j_1, \dots, j_q\}, \\ C_r (\log d)^{(r-q)/2} \phi(c_j) \left( \prod_{p=1}^q \phi(c_{j_p}) \right) \prod_{k: k \neq j_1, \dots, j_q, j} \Phi(c_k) & \text{otherwise,} \end{cases}$$

where we used the identity  $\sum_{p=1}^q \nu_p = r$ . On the other hand, if  $\max_{p=1, \dots, q} c_{j_p} > \sqrt{4(r+1) \log d}$ ,

$$\Lambda_j(\nu, \mathbf{j}) \leq C_r \prod_{p=1}^q e^{-c_{j_p}^2/4} \leq C_r d^{-r-1}.$$

Consequently,

$$\begin{aligned} & \sum_{j=1}^d \sum_{\mathbf{j}=(j_1, \dots, j_q) \in \mathcal{J}_q(d)} \Lambda_j(\nu, \mathbf{j}) \\ & \leq C_r + C_r \sum_{(j_1, \dots, j_q) \in \mathcal{J}_q(d)} (\log d)^{(r-q+1)/2} \left( \prod_{p=1}^q \phi(c_{j_p}) \right) \prod_{k: k \neq j_1, \dots, j_q} \Phi(c_k) \\ & \quad + C_r \sum_{(j_1, \dots, j_q) \in \mathcal{J}_q(d)} \sum_{j: j \neq j_1, \dots, j_q} (\log d)^{(r-q)/2} \phi(c_j) \left( \prod_{p=1}^q \phi(c_{j_p}) \right) \prod_{k: k \neq j_1, \dots, j_q, j} \Phi(c_k). \end{aligned}$$

With  $A' = \prod_{j=1}^d (-\infty, c_j]$ , we can rewrite the right hand side of the above inequality as

$$\begin{aligned} & C_r + C_r (\log d)^{(r-q+1)/2} \sum_{(j_1, \dots, j_q) \in \mathcal{J}_q(d)} \int_{A'} \partial_{j_1, \dots, j_q} \phi_d(z) dz \\ & \quad + C_r (\log d)^{(r-q)/2} \sum_{(j_1, \dots, j_{q+1}) \in \mathcal{J}_{q+1}(d)} \int_{A'} \partial_{j_1, \dots, j_{q+1}} \phi_d(z) dz. \end{aligned}$$

This quantity is bounded by

$$C_r \left( 1 + (\log d)^{(r-q+1)/2} \left\| \int_{A'} \nabla^q \phi_d(z) dz \right\|_1 + (\log d)^{(r-q)/2} \left\| \int_{A'} \nabla^{q+1} \phi_d(z) dz \right\|_1 \right). \quad (\text{B.6})$$

For any  $m \in \mathbb{N}$ , observe that

$$\left\| \int_{A'} \nabla^m \phi_d(z) dz \right\|_1 = \lim_{a \rightarrow -\infty} \left\| \int_{\prod_{j=1}^d [a, c_j]} \nabla^m \phi_d(z) dz \right\|_1 \leq \sup_{A \in \mathcal{R}} \left\| \int_A \nabla^m \phi_d(z) dz \right\|_1.$$

Therefore, by Lemma E.4, the quantity in (B.6) is bounded by  $C_r (\log d)^{(r+1)/2}$ . This gives (B.3).  $\square$

### B.3 Proof of Lemma 5.3

The proof of (5.35) is an almost straightforward multi-dimensional extension of that of [30, Lemma 2.1], and the proof of (5.36) is its simplification. The following lemma will play a key role in our argument.

**Lemma B.1.** *Let  $\xi$  be a centered random vector in  $\mathbb{R}^d$ . Suppose that  $\xi$  has a Stein kernel  $\tau$  such that  $\mathbb{E}[\|\xi\|_\infty^3] + \mathbb{E}[\|\tau(\xi) \otimes \xi^{\otimes 2}\|_\infty] < \infty$ . Then,  $\mathbb{E}[\|\tau(\xi) \otimes \xi\|_\infty] < \infty$  and for any  $f \in C_b^4(\mathbb{R}^d)$ ,*

$$\mathbb{E}[\langle \tau(\xi) \otimes \xi, \nabla^3 f(\xi) \rangle] = \frac{1}{2} (\mathbb{E}[\langle \xi^{\otimes 3}, \nabla^3 f(\xi) \rangle] - \mathbb{E}[\langle \tau(\xi) \otimes \xi^{\otimes 2}, \nabla^4 f(\xi) \rangle]). \quad (\text{B.7})$$

**Proof.** Take  $j, k, l \in \{1, \dots, d\}$  arbitrarily. Then we have

$$\begin{aligned} \mathbb{E}[\tau_{jk}(\xi)\xi_l] &= \mathbb{E}[\tau_{jk}(\xi)\xi_l; |\xi_l| \leq 1] + \mathbb{E}[\tau_{jk}(\xi)\xi_l; |\xi_l| > 1] \\ &\leq \mathbb{E}[\tau_{jk}(\xi)] + \mathbb{E}[\tau_{jk}(\xi)|\xi_l^2] \leq \mathbb{E}[\|\tau(\xi)\|_\infty] + \mathbb{E}[\|\tau(\xi) \otimes \xi^{\otimes 2}\|_\infty] < \infty. \end{aligned}$$

Hence  $\mathbb{E}[\|\tau(\xi) \otimes \xi\|_\infty] < \infty$ .

Next we prove (B.7). By a standard approximation argument, it suffices to prove (B.7) when  $f$  is compactly supported. For every  $j = 1, \dots, d$ , define a function  $g_j : \mathbb{R}^d \rightarrow \mathbb{R}$  as

$$g_j(x) = \langle x^{\otimes 2}, \nabla^2 \partial_j f(x) \rangle = \sum_{u,v=1}^d x_u x_v \partial_{juv} f(x), \quad x \in \mathbb{R}^d.$$

Observe that  $g_j$  is compactly supported because  $f$  is compactly supported. For  $j, k \in \{1, \dots, d\}$  and  $x \in \mathbb{R}^d$ , we have

$$\partial_k g_j(x) = 2 \sum_{v=1}^d x_v \partial_{jkv} f(x) + \sum_{u,v=1}^d x_u x_v \partial_{jkuv} f(x).$$

Hence we obtain

$$\begin{aligned} \mathbb{E}[\langle \tau(\xi) \otimes \xi, \nabla^3 f(\xi) \rangle] &= \sum_{j,k,v=1}^d \mathbb{E}[\tau_{jk}(\xi)\xi_v \partial_{jkv} f(\xi)] \\ &= \frac{1}{2} \sum_{j,k=1}^d \mathbb{E}[\tau_{jk}(\xi) \partial_k g_j(\xi)] - \frac{1}{2} \sum_{j,k,u,v=1}^d \mathbb{E}[\tau_{jk}(\xi)\xi_u \xi_v \partial_{jkuv} f(\xi)]. \end{aligned}$$

The second term on the last line is equal to  $\frac{1}{2} \mathbb{E}[\langle \tau(\xi) \otimes \xi^{\otimes 2}, \nabla^4 f(\xi) \rangle]$ . To evaluate the first term, define a function  $G : \mathbb{R}^d \rightarrow \mathbb{R}$  as  $G(x) = \sum_{k=1}^d x_k \int_0^1 g_k(\theta x) d\theta$ ,  $x \in \mathbb{R}^d$ . Then, using the relation  $\partial_k g_j = \partial_j g_k$ , one can easily verify  $\partial_j G = g_j$  for all  $j = 1, \dots, d$ . Moreover,  $G$  is bounded. In fact, since  $g_1, \dots, g_d$  are compactly supported, there exists a constant  $K > 0$  such that  $g_j(x) = 0$  for all  $j$  whenever  $\|x\|_\infty > K$ . Then, for any  $x \in \mathbb{R}^d$  with  $\|x\|_\infty > K$ ,

$$|G(x)| \leq \sum_{k=1}^d |x_k| \int_0^{K/\|x\|_\infty} |g_k(\theta x)| d\theta \leq dK \max_{1 \leq k \leq d} \sup_{x \in \mathbb{R}^d} |g_k(x)|.$$

It is straightforward to check that this estimate is still true when  $\|x\|_\infty \leq K$ . Hence  $G$  is bounded. As a result,  $G \in C_b^2(\mathbb{R}^d)$  and thus

$$\sum_{j,k=1}^d \mathbb{E}[\tau_{jk}(\xi) \partial_k g_j(\xi)] = \sum_{j,k=1}^d \mathbb{E}[\tau_{jk}(\xi) \partial_{kj} G(\xi)] = \sum_{j=1}^d \mathbb{E}[\xi_j \partial_j G(\xi)] = \sum_{j=1}^d \mathbb{E}[\xi_j g_j(\xi)]$$

$$= \mathbb{E}[\langle \xi^{\otimes 3}, \nabla^3 f(\xi) \rangle],$$

where the second equality follows from the definition of Stein kernel. Combining these identities gives the desired result.  $\square$

**Proof of Lemma 5.3.** First we prove (5.35). Without loss of generality, we may assume  $(\xi_i)_{i=1}^n$  and  $Z \sim N(0, \Sigma)$  are independent. Also, it suffices to prove the claim when  $h \in C_b^\infty(\mathbb{R}^d)$ . To see this, let  $t_1 = t/2$  and  $t_2 = t/(2-t)$ . One can easily check that  $t_1, t_2 \in (0, 1]$  and  $h_t = (h_{t_1})_{t_2}$ . Since  $h_{t_1} \in C_b^\infty(\mathbb{R}^d)$ , the general case follows by applying the claim to  $h = h_{t_1}$  and  $t = t_2$ .

Set  $W(s) = \sqrt{1-s}W + \sqrt{s}Z$  for every  $s \in [0, 1]$ . Then we have  $\mathbb{E}[h_t(W)] = \mathbb{E}[h(W(t))]$  and  $\mathbb{E}[h(Z)] = \mathbb{E}[h(W(1))]$ . Therefore, by the fundamental theorem of calculus, we obtain

$$\begin{aligned} \mathbb{E}[h_t(W)] - \mathbb{E}[h(Z)] &= - \int_t^1 \frac{\partial}{\partial s} \mathbb{E}[h(W(s))] ds \\ &= \frac{1}{2} \int_t^1 \left( \frac{\mathbb{E}[W \cdot \nabla h(W(s))]}{\sqrt{1-s}} - \frac{\mathbb{E}[Z \cdot \nabla h(W(s))]}{\sqrt{s}} \right) ds. \end{aligned}$$

By the multivariate Stein identity,

$$\frac{\mathbb{E}[Z \cdot \nabla h(W(s))]}{\sqrt{s}} = \mathbb{E}[\langle \Sigma, \nabla^2 h(W(s)) \rangle].$$

Also, since  $x \mapsto \mathbb{E}[T | W = x]$  is a Stein kernel for  $W$ , we have

$$\frac{\mathbb{E}[W \cdot \nabla h(W(s))]}{\sqrt{1-s}} = \mathbb{E}[\langle T, \nabla^2 h(W(s)) \rangle].$$

Consequently,

$$\mathbb{E}[h_t(W)] - \mathbb{E}[h(Z)] = \frac{1}{2} \int_t^1 \mathbb{E}[\langle \bar{T}, \nabla^2 h(W(s)) \rangle] ds. \quad (\text{B.8})$$

Next, for  $t < s < 1$ , by the fundamental theorem of calculus again, we have

$$\begin{aligned} &\mathbb{E}[\langle \bar{T}, \nabla^2 h(W(s)) \rangle] - \mathbb{E}[\langle \bar{T}, \nabla^2 h(Z) \rangle] \\ &= \frac{1}{2} \int_s^1 \left( \frac{\mathbb{E}[\langle \bar{T} \otimes W, \nabla^3 h(W(u)) \rangle]}{\sqrt{1-u}} - \frac{\mathbb{E}[\langle \bar{T} \otimes Z, \nabla^3 h(W(u)) \rangle]}{\sqrt{u}} \right) du. \end{aligned}$$

Since  $Z$  is independent of  $(W, T)$ , we have by the multivariate Stein identity

$$\frac{\mathbb{E}[\langle \bar{T} \otimes Z, \nabla^3 h(W(u)) \rangle]}{\sqrt{u}} = \mathbb{E}[\langle \bar{T} \otimes \Sigma, \nabla^4 h(W(u)) \rangle].$$

Meanwhile, we rewrite  $\mathbb{E}[\langle \bar{T} \otimes W, \nabla^3 h(W(u)) \rangle]$  as

$$\begin{aligned} \mathbb{E}[\langle \bar{T} \otimes W, \nabla^3 h(W(u)) \rangle] &= \sum_{i=1}^n \mathbb{E}[\langle \bar{T} \otimes \xi_i, \nabla^3 h(W(u)) \rangle] \\ &= \sum_{i=1}^n \mathbb{E}[\langle \bar{T}^{(i)} \otimes \xi_i, \nabla^3 h(W(u)) \rangle] + \sum_{i=1}^n \mathbb{E}[\langle \tau_i(\xi_i) \otimes \xi_i, \nabla^3 h(W(u)) \rangle], \end{aligned}$$

where  $\bar{T}^{(i)} = \bar{T} - \tau_i(\xi_i)$ . By Lemma B.1, the second term on the last line can be rewritten as

$$\begin{aligned} & \sum_{i=1}^n \mathbb{E}[\langle \tau_i(\xi_i) \otimes \xi_i, \nabla^3 h(W(u)) \rangle] \\ &= \frac{1}{2} \sum_{i=1}^n \left( \mathbb{E}[\langle \xi_i^{\otimes 3}, \nabla^3 h(W(u)) \rangle] - \sqrt{1-u} \mathbb{E}[\langle \tau_i(\xi_i) \otimes \xi_i^{\otimes 2}, \nabla^4 h(W(u)) \rangle] \right). \end{aligned}$$

Further, for  $i = 1, \dots, n$ , since  $\tau_i$  is a Stein kernel for  $\xi_i$  and  $\xi_i$  is independent of  $\bar{T}^{(i)}$  and  $W - \xi_i$ , we have

$$\mathbb{E}[\langle \bar{T}^{(i)} \otimes \xi_i, \nabla^3 h(W(u)) \rangle] = \sqrt{1-u} \mathbb{E}[\langle \bar{T}^{(i)} \otimes \tau_i(\xi_i), \nabla^4 h(W(u)) \rangle].$$

Hence

$$\begin{aligned} & \sum_{i=1}^n \mathbb{E}[\langle \bar{T}^{(i)} \otimes \xi_i, \nabla^3 h(W(u)) \rangle] \\ &= \sqrt{1-u} \left( \sum_{i=1}^n \mathbb{E}[\langle \bar{T} \otimes \tau_i(\xi_i), \nabla^4 h(W(u)) \rangle] - \sum_{i=1}^n \mathbb{E}[\langle \tau_i(\xi_i)^{\otimes 2}, \nabla^4 h(W(u)) \rangle] \right) \\ &= \sqrt{1-u} \left( \mathbb{E}[\langle \bar{T} \otimes T, \nabla^4 h(W(u)) \rangle] - \sum_{i=1}^n \mathbb{E}[\langle \tau_i(\xi_i)^{\otimes 2}, \nabla^4 h(W(u)) \rangle] \right). \end{aligned}$$

Consequently, we conclude

$$\begin{aligned} & \mathbb{E}[\langle \bar{T}, \nabla^2 h(W(s)) \rangle] - \mathbb{E}[\langle \bar{T}, \nabla^2 h(Z) \rangle] \\ &= \frac{1}{2} \int_s^1 \left( \mathbb{E}[\langle \bar{T}^{\otimes 2}, \nabla^4 h(W(u)) \rangle] - \sum_{i=1}^n \mathbb{E}[\langle \tau_i(\xi_i)^{\otimes 2}, \nabla^4 h(W(u)) \rangle] \right) du \\ &+ \frac{1}{4} \int_s^1 \sum_{i=1}^n \left( \frac{\mathbb{E}[\langle \xi_i^{\otimes 3}, \nabla^3 h(W(u)) \rangle]}{\sqrt{1-u}} - \mathbb{E}[\langle \tau_i(\xi_i) \otimes \xi_i^{\otimes 2}, \nabla^4 h(W(u)) \rangle] \right) du. \end{aligned} \tag{B.9}$$

Fix  $i \in \{1, \dots, n\}$  and  $0 < u < 1$ . By the fundamental theorem of calculus and multivariate Stein identity again, we have

$$\begin{aligned} & \mathbb{E}[\langle \xi_i^{\otimes 3}, \nabla^3 h(W(u)) \rangle] - \mathbb{E}[\langle \xi_i^{\otimes 3}, \nabla^3 h(Z) \rangle] \\ &= \frac{1}{2} \int_u^1 \left( \frac{\mathbb{E}[\langle \xi_i^{\otimes 3} \otimes W, \nabla^4 h(W(v)) \rangle]}{\sqrt{1-v}} - \mathbb{E}[\langle \xi_i^{\otimes 3} \otimes \Sigma, \nabla^5 h(W(v)) \rangle] \right) dv. \end{aligned}$$

We rewrite  $\mathbb{E}[\langle \xi_i^{\otimes 3} \otimes W, \nabla^4 h(W(v)) \rangle]$  as

$$\begin{aligned} \mathbb{E}[\langle \xi_i^{\otimes 3} \otimes W, \nabla^4 h(W(v)) \rangle] &= \sum_{j=1}^n \mathbb{E}[\langle \xi_i^{\otimes 3} \otimes \xi_j, \nabla^4 h(W(v)) \rangle] \\ &= \mathbb{E}[\langle \xi_i^{\otimes 4}, \nabla^4 h(W(v)) \rangle] + \sum_{j:j \neq i} \mathbb{E}[\langle \xi_i^{\otimes 3} \otimes \xi_j, \nabla^4 h(W(v)) \rangle]. \end{aligned}$$

For  $j \neq i$ ,  $\xi_j$  is independent of  $\xi_i^{\otimes 3}$  and  $W - \xi_j$ , so we obtain by the definition of Stein kernel

$$\mathbb{E}[\langle \xi_i^{\otimes 3} \otimes \xi_j, \nabla^4 h(W(v)) \rangle] = \sqrt{1-v} \mathbb{E}[\langle \xi_i^{\otimes 3} \otimes \tau_j(\xi_j), \nabla^5 h(W(v)) \rangle].$$

Hence,

$$\begin{aligned} & \mathbb{E}[\langle \xi_i^{\otimes 3} \otimes W, \nabla^4 h(W(v)) \rangle] \\ &= \mathbb{E}[\langle \xi_i^{\otimes 4}, \nabla^4 h(W(v)) \rangle] + \sqrt{1-v} \mathbb{E}[\langle \xi_i^{\otimes 3} \otimes (T - \tau_i(\xi_i)), \nabla^5 h(W(v)) \rangle]. \end{aligned}$$

Overall, we conclude

$$\begin{aligned} & \mathbb{E}[\langle \xi_i^{\otimes 3}, \nabla^3 h(W(u)) \rangle] - \mathbb{E}[\langle \xi_i^{\otimes 3}, \nabla^3 h(Z) \rangle] \\ &= \frac{1}{2} \int_u^1 \frac{\mathbb{E}[\langle \xi_i^{\otimes 4}, \nabla^4 h(W(v)) \rangle]}{\sqrt{1-v}} dv + \frac{1}{2} \int_u^1 \mathbb{E}[\langle \xi_i^{\otimes 3} \otimes (\bar{T} - \tau_i(\xi_i)), \nabla^5 h(W(v)) \rangle] dv. \end{aligned} \quad (\text{B.10})$$

Further, note that we have by integration by parts

$$\begin{aligned} \int_{\mathbb{R}^d} h_t(z) \nabla^r \phi_\Sigma(z) dz &= (-\sqrt{1-t})^r \int_{\mathbb{R}^d} \mathbb{E}[\nabla^r h(\sqrt{1-t}z + \sqrt{t}Z)] \phi_\Sigma(z) dz \\ &= (-\sqrt{1-t})^r \mathbb{E}[\nabla^r h(Z)] \end{aligned}$$

for any  $r \in \mathbb{N}$ . Hence

$$\frac{1-t}{2} \mathbb{E}[\langle \bar{T}, \nabla^2 h(Z) \rangle] = \frac{1-t}{2} \langle \Sigma_W - \Sigma, \mathbb{E}[\nabla^2 h(Z)] \rangle = \frac{1}{2} \int_{\mathbb{R}^d} h_t(z) \langle \Sigma_W - \Sigma, \nabla^2 \phi_\Sigma(z) \rangle dz$$

and

$$\frac{\mathbb{E}[\langle \xi_i^{\otimes 3}, \nabla^3 h(Z) \rangle]}{8} \int_t^1 \left( \int_s^1 \frac{1}{\sqrt{1-u}} du \right) ds = -\frac{1}{6} \int_{\mathbb{R}^d} h_t(z) \langle \mathbb{E}[\xi_i^{\otimes 3}], \nabla^3 \phi_\Sigma(z) \rangle dz.$$

Consequently,

$$\begin{aligned} & \mathbb{E}[h(Z)] + \frac{1-t}{2} \mathbb{E}[\langle \bar{T}, \nabla^2 h(Z) \rangle] + \sum_{i=1}^n \frac{\mathbb{E}[\langle \xi_i^{\otimes 3}, \nabla^3 h(Z) \rangle]}{8} \int_t^1 \left( \int_s^1 \frac{1}{\sqrt{1-u}} du \right) ds \\ &= \int_{\mathbb{R}^d} h_t(z) p_W(z) dz. \end{aligned} \quad (\text{B.11})$$

Finally, observe that

$$\nabla^r h_s(W) = (1-s)^{r/2} \mathbb{E}[\nabla^r h(W(s)) \mid W] \quad (\text{B.12})$$

for any  $r \in \mathbb{N}$  and  $s \in [0, 1]$ . Combining (B.8)–(B.12) gives (5.35).

Next we prove (5.36). As above, we may assume that  $W$  and  $Z$  are independent and  $h \in C_b^\infty(\mathbb{R}^d)$ . Also, we define  $W(s)$  as above. Then, by the proof of (B.8),

$$\mathbb{E}[h_t(W)] - \mathbb{E}[h(Z)] = \frac{1}{2} \int_t^1 \mathbb{E}[\langle \Sigma_W - \Sigma, \nabla^2 h(W(s)) \rangle] ds.$$

Also, applying the proof of (B.8) to the function  $x \mapsto \langle \Sigma_W - \Sigma, \nabla^2 h(x) \rangle$  instead of  $h$  gives

$$\mathbb{E}[\langle \Sigma_W - \Sigma, \nabla^2 h(W(s)) \rangle] - \mathbb{E}[\langle \Sigma_W - \Sigma, \nabla^2 h(Z) \rangle] = \frac{1}{2} \int_s^1 \mathbb{E}[\langle (\Sigma_W - \Sigma)^{\otimes 2}, \nabla^4 h(W(u)) \rangle] du.$$

Combining these two identities with (B.12) gives (5.36).  $\square$

## C Properties of Stein kernel

### C.1 Basic properties

**Lemma C.1.** *Let  $\xi$  be a random vector in  $\mathbb{R}^d$  with a Stein kernel  $\tau$ . Then, for any  $a \in \mathbb{R}^p$  and  $p \times d$  matrix  $V$ ,  $V\xi + a$  has a Stein kernel given by  $x \mapsto \mathbb{E}[V\tau(\xi)V^\top \mid V\xi + a = x]$ .*

**Proof.** Straightforward from the definition of Stein kernel.  $\square$

**Proposition C.1.** *Let  $\xi$  be a centered random variable having a Stein kernel  $\tau$ . Then  $\tau(\xi) \geq 0$  a.s. Moreover, if  $P(\xi \neq 0) > 0$ , then the event  $E := \{\tau(\xi) > 0\}$  occurs with a positive probability and the conditional law of  $\xi$  given  $E$  is absolutely continuous. In particular, the law of  $\xi$  has a non-zero absolutely continuous part.*

**Proof.** The asserted claims are shown by essentially the same arguments as those in the proofs of [50, Proposition 2.9.4] and [50, Theorem 10.1.1]. We give the details for the sake of completeness.

Let  $B$  be a bounded Borel subset of  $\mathbb{R}$ . First we prove

$$\mathbb{E} \left[ \xi \int_0^\xi 1_B(y) dy \right] = \mathbb{E}[1_B(\xi)\tau(\xi)]. \quad (\text{C.1})$$

Consider a Borel measure  $\mu$  on  $\mathbb{R}$  given by  $\mu = \mathcal{L}_\xi + \mathcal{L}$ , where  $\mathcal{L}_\xi$  is the law of  $\xi$  and  $\mathcal{L}$  is the Lebesgue measure on  $\mathbb{R}$ . Then we have  $\mu(K) < \infty$  for any compact set  $K \subset \mathbb{R}$ . Hence  $\mu$  is regular by Theorem 2.18 in [54]. Also, note that  $\mu(B) < \infty$  by the boundedness of  $B$ . Therefore, by Lusin's theorem (see Theorem 2.24 in [54]), for every  $m \in \mathbb{N}$ , there exists a compactly supported continuous function  $g_m : \mathbb{R} \rightarrow [-1, 1]$  such that  $\mu(\{y \in \mathbb{R} : 1_B(y) \neq g_m(y)\}) < 1/m$ . Now define a function  $G_m : \mathbb{R} \rightarrow \mathbb{R}$  as  $G_m(x) = \int_0^x g_m(y) dy$ ,  $x \in \mathbb{R}$ . Then  $G_m$  is a bounded  $C^1$  function with  $G'_m = g_m$ , so  $\mathbb{E}[\xi G_m(\xi)] = \mathbb{E}[g_m(\xi)\tau(\xi)]$  by the definition of Stein kernel. Since  $|G_m(\xi) - \int_0^\xi 1_B(y) dy| \leq 1/m$  by construction,  $\mathbb{E}[\xi G_m(\xi)]$  converges to the quantity on the left hand side of (C.1) as  $m \rightarrow \infty$ . Meanwhile, since  $g_m(\xi) \rightarrow 1_B(\xi)$  a.s. as  $m \rightarrow \infty$  by construction, the dominated convergence theorem gives  $\mathbb{E}[g_m(\xi)\tau(\xi)] \rightarrow \mathbb{E}[1_B(\xi)\tau(\xi)]$  as  $m \rightarrow \infty$ . So we obtain (C.1).

Let us prove the first claim. Observe that  $x \int_0^x 1_B(y) dy \geq 0$  for all  $x \in \mathbb{R}$ . Hence  $\mathbb{E}[1_B(\xi)\tau(\xi)] \geq 0$  by (C.1). By the dominated convergence theorem, this is still true even if  $B$  is unbounded. Hence  $\tau(\xi) \geq 0$  a.s.

It remains to prove the second claim. First, if  $\tau(\xi) = 0$  a.s., then (C.1) implies  $\xi \int_0^\xi 1_B(y) dy = 0$  a.s. (recall the above argument). Taking  $B = [-K, K]$  for some  $K > 0$  gives  $\xi(|\xi| \wedge K) = 0$  a.s. Letting  $K \rightarrow \infty$ , we obtain  $\xi = 0$  a.s. By contraposition,  $P(\xi \neq 0) > 0$  means  $P(E) > 0$ . Next, if the Lebesgue measure of  $B$  is zero, then  $\mathbb{E}[1_B(\xi)\tau(\xi)] = 0$  by (C.1). Since  $\mathbb{E}[1_B(\xi)\tau(\xi)] = \mathbb{E}[1_B(\xi)\tau(\xi)1_E]$ , we obtain  $1_B(\xi)\tau(\xi)1_E = 0$  a.s. Since  $\tau(\xi) > 0$  on  $E$ , this implies  $1_B(\xi)1_E = 0$  a.s. Hence  $P(\xi \in B \mid E) = \mathbb{E}[1_B(\xi)1_E]/P(E) = 0$ . This implies that the conditional law of  $\xi$  given  $E$  is absolutely continuous.  $\square$

### C.2 Existence results

#### C.2.1 Proof of Lemma 3.1

The proof uses the Malliavin–Stein method. We refer to [50, Chapter 2] for undefined notation and concepts used below.

Let  $H = \mathbb{R}^d$  be the Hilbert space equipped with the canonical inner product. Consider an isonormal Gaussian process over  $H$  given by  $W(h) = h \cdot G$ ,  $h \in H$ . We consider Malliavin calculus with respect to

$W$ . First, approximating  $\psi$  by a Lipschitz function and applying Proposition 2.3.8 in [50] and Lemma 1.2.3 in [52], we obtain  $X_j \in \mathbb{D}^{1,2}$  and  $DX_j = \nabla\psi_j(G)$  for every  $j = 1, \dots, d$ . Therefore, by Proposition 3.7 in [51], the map  $\tau : \mathbb{R}^d \rightarrow (\mathbb{R}^d)^{\otimes 2}$  defined by  $\tau_{jk}(x) = \mathbb{E}[-DL^{-1}X_j \cdot DX_k \mid X = x]$  for  $x \in \mathbb{R}^d$  and  $j, k = 1, \dots, d$  gives a Stein kernel for  $X$ . For any  $p \geq 1$  and  $j, k = 1, \dots, d$ , we have

$$\mathbb{E}[|\tau_{jk}(X)|^p] \leq \sqrt{\mathbb{E}[|DL^{-1}X_j|^{2p}] \mathbb{E}[|DX_k|^{2p}]} \leq \sqrt{\mathbb{E}[|DX_j|^{2p}] \mathbb{E}[|DX_k|^{2p}]},$$

where the first inequality is by the Jensen and Schwarz inequalities and the second by Lemma 5.3.7 in [50]. If  $p$  is an even integer, we also have  $\mathbb{E}[X_j^p] \leq (p-1)^{p/2} \mathbb{E}[|DX_j|^p]$  by Lemma 5.3.7 in [50]. Since  $DX_j = \nabla\psi_j(G)$  for each  $j$ , this completes the first part of the proof. The second part can be shown in a similar way and thus we omit it.  $\square$

### C.2.2 Proof of Propositions 2.1 and 3.1

Denote by  $r_j$  the  $j$ -th row vector of  $R^{1/2}$ . Then  $Z$  has the same law as  $(r_1 \cdot G, \dots, r_d \cdot G)^\top$  with  $G \sim N(0, I_d)$ . Hence we may assume  $X$  is of the form  $X = \psi(G)$  with  $\psi : \mathbb{R}^d \rightarrow \mathbb{R}^d$  defined as  $\psi_j(x) = F_j^{-1}(\Phi(r_j \cdot x))$  for  $j = 1, \dots, d$  and  $x \in \mathbb{R}^d$ . To apply Lemma 3.1, we need to prove  $\psi_j$  are locally Lipschitz and compute its gradient. To see this, we note that  $F_j^{-1}$  is absolutely continuous and satisfies  $|(F_j^{-1})'(t)| \leq (\kappa \min\{t, 1-t\})^{-1}$  a.e. by (2.10). This follows from arguments in Section 5.3 of [10] (see also Propositions A.17 and A.19 in [10]). Consequently,  $F_j^{-1}$  is locally Lipschitz. This implies  $\psi_j$  is locally Lipschitz since  $\Phi$  is Lipschitz, and its gradient is given by  $\nabla\psi_j(x) = \phi(r_j \cdot x) f_j(F_j^{-1}(\Phi(r_j \cdot x)))^{-1} r_j$  a.e. Since  $|r_j|^2 = R_{jj} = 1$ , we obtain

$$|\nabla\psi_j(G)| = \phi(Z_j) f_j(F_j^{-1}(\Phi(Z_j)))^{-1} \leq \frac{\phi(Z_j)}{\kappa \min\{\Phi(Z_j), 1 - \Phi(Z_j)\}} \leq \frac{1 + |Z_j|}{\kappa},$$

where the last inequality follows by Birnbaum's inequality. Hence  $\|\nabla\psi_j(G)\|_p \lesssim \sqrt{p}\kappa^{-1}$ . Combining this with Lemmas 3.1 and E.5 gives the desired results.  $\square$

### C.2.3 Cheeger constant of the gamma distribution

**Proposition C.2.** *Any gamma distribution has a positive Cheeger constant.*

**Proof.** Let  $\mu$  be the gamma distribution with shape  $\nu$  and rate  $\alpha$ . If  $\nu \geq 1$ , then  $\mu$  is log-concave, so the claim follows by Proposition 4.1 in [8]. When  $\nu < 1$ , the density  $f$  of  $\mu$  satisfies  $\inf_{0 < t < M} f(t) > 0$  for every  $M > 0$ . Hence, in view of Theorem 1.3 in [9], it suffices to prove  $\liminf_{p \uparrow 1} f(F^{-1}(p))/(1-p) > 0$ , where  $F$  is the distribution function of  $\mu$ . Since

$$\frac{d}{dp} f(F^{-1}(p)) = \frac{\nu - 1}{F^{-1}(p)} - \alpha \rightarrow -\alpha \quad (p \uparrow 1),$$

we have by L'Hôpital's rule  $\liminf_{p \uparrow 1} f(F^{-1}(p))/(1-p) = \alpha > 0$ . This completes the proof.  $\square$

### C.2.4 Proof of Proposition 3.2

**Lemma C.2** (Gaussian type isoperimetric inequality for the beta distribution). *Let  $f$  and  $F$  be the density and distribution function of the beta distribution with parameters  $\alpha, \beta$ , respectively. Then there exists a constant  $c > 0$  such that  $f(F^{-1}(p)) \geq c\phi(\Phi^{-1}(p))$  for all  $p \in (0, 1)$ .*

**Proof.** Since  $\inf_{\varepsilon < p < 1-\varepsilon} f(F^{-1}(p))/\phi(\Phi^{-1}(p)) > 0$  for any  $\varepsilon \in (0, 1/2)$ , it suffices to prove

$$\lim_{p \downarrow 0} \frac{f(F^{-1}(p))}{\phi(\Phi^{-1}(p))} = \infty \quad \text{and} \quad \lim_{p \uparrow 1} \frac{f(F^{-1}(p))}{\phi(\Phi^{-1}(p))} = \infty.$$

By symmetry it is enough to prove the former. When  $\alpha \leq 1$ ,  $\lim_{p \downarrow 0} f(F^{-1}(p)) > 0$  while  $\lim_{p \downarrow 0} \phi(\Phi^{-1}(p)) = 0$ , so the claim is obvious. When  $\alpha > 1$ , by L'Hôpital's rule, it suffices to prove

$$\lim_{p \downarrow 0} \frac{\alpha - 1}{-F^{-1}(p)\Phi^{-1}(p)} = \infty.$$

First, observe that  $F(t) \geq C_{\alpha, \beta} t^\alpha$  for any  $0 < t < 1/2$ . Hence  $F^{-1}(p) \leq (C_{\alpha, \beta}^{-1} p)^{1/\alpha}$  for any  $0 < p < F^{-1}(1/2)$ . Next, by the well-known inequality  $\Phi(-t) = 1 - \Phi(t) \leq e^{-t^2/2}$  for all  $t \geq 0$ , we deduce  $\sqrt{-2 \log p} \geq -\Phi^{-1}(p)$  for all  $0 < p < 1/2$ . Since  $\lim_{p \downarrow 0} p^{1/\alpha} \sqrt{-2 \log p} = 0$  and  $-F^{-1}(p)\Phi^{-1}(p) \geq 0$  for all  $0 < p < 1/2$ , we conclude  $\lim_{p \downarrow 0} \{-F^{-1}(p)\Phi^{-1}(p)\} = 0$ . This gives the desired result.  $\square$

**Proof of Proposition 3.2.** Let  $F$  be the distribution function of  $\eta$ . Then  $\eta \stackrel{d}{=} F^{-1}(\Phi(G))$  with  $G \sim N(0, 1)$ . Hence, in view of Lemmas 3.1 and C.1, it suffices to show that  $(F^{-1} \circ \Phi)'$  is bounded. This immediately follows from Lemma C.2.  $\square$

#### D Proof of Lemma 6.1

**Lemma D.1.** *There exists a universal constant  $C > 0$  such that  $\sup_{t \in \mathbb{R}} |f'_\Sigma(t)| \leq C \sigma_*^{-2} \log d$ .*

**Proof.** Observe that

$$F_Z(t) = \int_{(-\infty, t]^d} \phi_\Sigma(z) dz = \int_{(-\infty, 0]^d} \phi_\Sigma(z + t \mathbf{1}_d) dz$$

for all  $t \in \mathbb{R}$ . Differentiating this equation twice with respect to  $t$  gives

$$f'_\Sigma(t) = \int_{(-\infty, 0]^d} \langle \mathbf{1}_d^{\otimes 2}, \nabla^2 \phi_\Sigma(z + t \mathbf{1}_d) \rangle dz = \int_{(-\infty, t]^d} \langle \mathbf{1}_d^{\otimes 2}, \nabla^2 \phi_\Sigma(z) \rangle dz.$$

Hence the desired result follows from Lemma E.4.  $\square$

**Proof of Lemma 6.1.** Let us prove (6.5). We write  $F = F_Z$  and  $x = F^{-1}(p)$  for short. First we consider the case  $p \geq 1/2$ . Since the function  $\mathbb{R}^d \ni z \mapsto z^\vee \in \mathbb{R}$  is convex,  $\Phi^{-1} \circ F$  is concave by Corollary A.2.9 in [56]. Hence, for any  $y > x$ ,

$$\frac{(\Phi^{-1} \circ F)(y) - (\Phi^{-1} \circ F)(x)}{y - x} \leq (\Phi^{-1} \circ F)'(x) = \frac{f(x)}{\phi(\Phi^{-1}(F(x)))}.$$

Let  $y = F^{-1}(p) + c$  with  $c$  a positive constant specified later. Then,

$$\frac{f(F^{-1}(p))}{\phi(\Phi^{-1}(p))} \geq \frac{\Phi^{-1}(F(F^{-1}(p) + c)) - \Phi^{-1}(p)}{c}.$$

Thus, we need to choose  $c$  so that  $F(F^{-1}(p) + c)$  has an appropriate lower bound. Noting  $p \geq 1/2$ , we have

$$\begin{aligned} 1 - F(F^{-1}(p) + c) &= P(Z^\vee > F^{-1}(p) + c) \leq P(Z^\vee > F^{-1}(1/2) + c) \\ &\leq \frac{\mathbb{E}[(Z^\vee - F^{-1}(1/2))^2]}{c^2} \leq \frac{2 \text{Var}[Z^\vee]}{c^2}, \end{aligned}$$

where we used the following inequality for the last bound: For any random variable  $Y$  and its median  $m$ ,  $\mathbb{E}[(Y - m)^2] = \text{Var}[Y] + (\mathbb{E}[Y] - m)^2 \leq 2 \text{Var}[Y]$ . Thus, letting  $c = \sqrt{4 \text{Var}[Z^\vee]/(1-p)}$ , we obtain  $F(F^{-1}(p) + c) \geq 1 - (1-p)/2 = (1+p)/2$ . Consequently,

$$f(F^{-1}(p)) \geq \phi(\Phi^{-1}(p)) \frac{\Phi^{-1}((1+p)/2) - \Phi^{-1}(p)}{2\sqrt{\text{Var}[Z^\vee]}} \sqrt{1-p}. \quad (\text{D.1})$$

Also, by the fundamental theorem of calculus,

$$\Phi^{-1}((1+p)/2) - \Phi^{-1}(p) = \int_p^{(1+p)/2} \frac{1}{\phi(\Phi^{-1}(u))} du \geq \frac{1-p}{2\phi(\Phi^{-1}(p))}, \quad (\text{D.2})$$

where the last inequality holds because  $\phi \circ \Phi^{-1}$  is decreasing on  $[1/2, 1)$ . Combining (D.1) with (D.2) gives (6.5).

Next consider the case  $p < 1/2$ . Since the function  $\log \circ \Phi$  is increasing and concave,  $\log \circ F = (\log \circ \Phi) \circ (\Phi^{-1} \circ F)$  is concave. Hence  $(\log \circ F)' = f/F$  is non-increasing. Thus we obtain

$$\frac{f(F^{-1}(p))}{p} \geq \frac{f(F^{-1}(1/2))}{F(F^{-1}(1/2))} \geq \frac{1}{2^{5/2} \sqrt{\text{Var}[Z^\vee]}},$$

where the last inequality follows from (6.5) for  $p = 1/2$  which was already proved in the above.

It remains to prove (6.6) when  $\text{Cov}[Z] = \Sigma$ . An elementary computation shows

$$(F_Z^{-1})'(u) = \frac{1}{f_\Sigma(F_Z^{-1}(u))}, \quad (F_Z^{-1})''(u) = -\frac{f'_\Sigma(F_Z^{-1}(u))}{f_\Sigma(F_Z^{-1}(u))^3}$$

for all  $u \in (0, 1)$ . Thus, the desired result follows from (6.5) and Lemma D.1.  $\square$

## E Technical tools

### E.1 Inequalities related to multivariate normal distributions

Let  $Z$  be a centered Gaussian vector in  $\mathbb{R}^d$ . Set  $\bar{\sigma} := \max_{1 \leq j \leq d} \sqrt{\text{Var}[Z_j]}$  and  $\underline{\sigma} := \min_{1 \leq j \leq d} \sqrt{\text{Var}[Z_j]}$ .

**Lemma E.1.**  $\mathbb{E}[Z^\vee] \leq \bar{\sigma} \sqrt{2 \log d}$  and  $\text{Var}[Z^\vee] \leq \bar{\sigma}^2$ .

**Proof.** The first bound follows from [11, Theorem 2.5]. The second one follows from [11, Theorem 5.8].  $\square$

**Lemma E.2** (Nazarov's inequality). *If  $\underline{\sigma} > 0$ , then for any  $x \in \mathbb{R}^d$  and  $\varepsilon > 0$ ,*

$$P\left(0 < \max_{1 \leq j \leq d} (Z_j - x_j) \leq \varepsilon\right) \leq \frac{\varepsilon}{\underline{\sigma}} (\sqrt{2 \log d} + 2).$$

**Proof.** See [20, Theorem 1].  $\square$

**Lemma E.3.** *There exists a universal constant  $c > 0$  such that  $\sqrt{\text{Var}[Z^\vee]} \log d \geq c \underline{\sigma}^2 / \bar{\sigma}$ .*

**Proof.** By a straightforward modification of the proof of Theorem 1.8 in [25], we can prove

$$\sqrt{\text{Var}[Z^\vee]} (\underline{\sigma} + \mathbb{E}[Z^\vee]) \geq c' \underline{\sigma}^2$$

for some universal constant  $c' > 0$ . In fact, this follows by applying the arguments in the proof of Theorem 1.8 in [25] to  $\mathbf{X} = Z/\bar{\sigma}$  with  $t = \sqrt{1 - \underline{\sigma}^2/(4m^2)}$  when  $m := \mathbb{E}[Z^\vee] > \underline{\sigma}/2$  (Lemma 2.2 in [25] is applied with  $\lambda = \underline{\sigma}^2/(4m\bar{\sigma})$ ). Then, since  $\mathbb{E}[Z^\vee] \leq \bar{\sigma}\sqrt{2\log d}$  by Lemma E.1, we obtain the desired result.  $\square$

**Lemma E.4** (Anderson–Hall–Titterton’s bound). *For any  $r \in \mathbb{N}$ ,*

$$\sup_{A \in \mathcal{R}} \left\| \int_A \nabla^r \phi_\Sigma(z) dz \right\|_1 \leq C_r \frac{\log^{r/2} d}{\sigma_*^r},$$

where  $C_r > 0$  is a constant depending only on  $r$ .

**Proof.** Let  $Z \sim N(0, \Sigma)$  and  $Z' \sim N(0, \Sigma - \sigma_*^2 I_d)$ . Then, for any  $A \in \mathcal{R}$  and  $x \in \mathbb{R}^d$ , we have

$$\int_A \phi_\Sigma(x + z) dz = \mathbb{E}[1_A(Z - x)] = \mathbb{E} \left[ \int_{\mathbb{R}^d} 1_A(\sigma_* z + Z') \phi_d(z + x/\sigma_*) dz \right].$$

Differentiating the both sides  $r$  times with respect to  $x$  and then setting  $x = 0$ , we obtain

$$\int_A \nabla^r \phi_\Sigma(z) dz = \frac{1}{\sigma_*^r} \mathbb{E} \left[ \int_{\mathbb{R}^d} 1_A(\sigma_* z + Z') \nabla^r \phi_d(z) dz \right].$$

Hence

$$\left\| \int_A \nabla^r \phi_\Sigma(z) dz \right\|_1 \leq \frac{1}{\sigma_*^r} \mathbb{E} \left[ \left\| \int_{\mathbb{R}^d} 1_A(\sigma_* z + Z') \nabla^r \phi_d(z) dz \right\|_1 \right] \leq C_r \frac{\log^{r/2} d}{\sigma_*^r},$$

where the last inequality follows by Lemma 2.2 in [26] because  $\{z \in \mathbb{R}^d : \sigma_* z + Z' \in A\} \in \mathcal{R}$ .  $\square$

## E.2 Inequalities related to sub-Weibull norms

**Lemma E.5.** *Let  $\xi$  be a random variable. Suppose that there is a constant  $A > 0$  such that  $\|\xi\|_p \leq Ap^{1/\alpha}$  for all  $p \geq 1$ . Then  $\|\xi\|_{\psi_\alpha} \leq C_\alpha A$ .*

**Proof.** See Lemma A.5 in [37].  $\square$

**Lemma E.6.** *For any  $\alpha \in (0, 1)$ , there exists a constant  $C_\alpha > 0$  depending only on  $\alpha$  such that*

$$\left\| \sum_{i=1}^n \xi_i \right\|_{\psi_\alpha} \leq C_\alpha \sum_{i=1}^n \|\xi_i\|_{\psi_\alpha}$$

for any random variables  $\xi_1, \dots, \xi_n$ .

**Proof.** This follows from Lemma C.2 in [16] and the triangle inequality for the Orlicz norm associated with a convex function.  $\square$

**Lemma E.7.** *Let  $\xi_1, \xi_2$  be two random variables such that  $\|\xi_1\|_{\psi_{\alpha_1}} + \|\xi_2\|_{\psi_{\alpha_2}} < \infty$  for some  $\alpha_1, \alpha_2 > 0$ . Then we have  $\|\xi_1 \xi_2\|_{\psi_\alpha} \leq \|\xi_1\|_{\psi_{\alpha_1}} \|\xi_2\|_{\psi_{\alpha_2}}$ , where  $\alpha > 0$  is defined by the equation  $1/\alpha = 1/\alpha_1 + 1/\alpha_2$ .*

**Proof.** See [38, Proposition D.2].  $\square$

**Lemma E.8.** Let  $\xi_1, \dots, \xi_n$  be independent random variables such that  $\max_{1 \leq i \leq n} \|\xi_i\|_{\psi_\alpha} \leq K$  for some  $K > 0$  and  $\alpha \in (0, 1]$ . Then, there is a constant  $C_\alpha > 0$  depending only on  $\alpha$  such that for any  $p \geq 1$ ,

$$\left\| \sum_{i=1}^n (\xi_i - \mathbb{E}[\xi_i]) \right\|_p \leq C_\alpha K \left( \sqrt{pn} + p^{1/\alpha} \right).$$

**Proof.** See Lemma 2.1 in [28]. □

**Lemma E.9.** Let  $Y_1, \dots, Y_n$  be independent random vectors in  $\mathbb{R}^d$ . Suppose that there exist constants  $K > 0$  and  $\alpha \in (0, 1]$  such that  $\max_{1 \leq i \leq n} \max_{1 \leq j \leq d} \|Y_{ij}\|_{\psi_\alpha} \leq K$ . Then, there exists a constant  $C_\alpha > 0$  depending only on  $\alpha$  such that

$$\mathbb{E} \left\| \sum_{i=1}^n (Y_i - \mathbb{E}[Y_i]) \right\|_\infty^r \leq C_\alpha^r K^r \left( \sqrt{nr \log d} + (r \log d)^{1/\alpha} \right)^r \quad (\text{E.1})$$

for any  $r \geq 1$  and

$$P \left( \left\| \sum_{i=1}^n (Y_i - \mathbb{E}[Y_i]) \right\|_\infty > C_\alpha K \left( \sqrt{an \log(dn)} + a^{1/\alpha} \log^{1/\alpha}(dn) \right) \right) \leq \frac{1}{n^a} \quad (\text{E.2})$$

for any  $a \geq 1$ .

**Proof.** By Lemma E.8, we have for any  $p \geq 2$

$$\max_{1 \leq j \leq d} \left\| \sum_{i=1}^n (Y_{ij} - \mathbb{E}[Y_{ij}]) \right\|_p \leq C'_\alpha K \left( \sqrt{pn} + p^{1/\alpha} \right),$$

where  $C'_\alpha > 0$  depends only on  $\alpha$ . Therefore, with  $p = r \log d$ , we have

$$\begin{aligned} \mathbb{E} \left\| \sum_{i=1}^n (Y_i - \mathbb{E}[Y_i]) \right\|_\infty^r &\leq \left( \mathbb{E} \left\| \sum_{i=1}^n (Y_i - \mathbb{E}[Y_i]) \right\|_\infty^p \right)^{r/p} \leq d^{r/p} \max_{1 \leq j \leq d} \left\| \sum_{i=1}^n (Y_{ij} - \mathbb{E}[Y_{ij}]) \right\|_p^r \\ &\leq d^{r/p} (C'_\alpha)^r K^r \left( \sqrt{nr \log d} + (r \log d)^{1/\alpha} \right)^r. \end{aligned}$$

Since  $d^{r/p} = e^{r \log d/p} = e \leq e^r$ , we obtain (E.1) with  $C_\alpha = e C'_\alpha$ . Also, by the union bound, Markov's inequality and (E.2), we have for any  $t > 0$  and  $p \geq 2$

$$P \left( \left\| \sum_{i=1}^n (Y_i - \mathbb{E}[Y_i]) \right\|_\infty > t \right) \leq d \left( t^{-1} C'_\alpha K \left( \sqrt{pn} + p^{1/\alpha} \right) \right)^p.$$

Applying this estimate with  $p = a \log(dn)$  and  $t = e C'_\alpha K \left( \sqrt{pn} + p^{1/\alpha} \right)$ , we obtain (E.2) □

**Lemma E.10.** Let  $r \in \mathbb{N}$ . If (2.3) is satisfied, there exists a constant  $C_r > 0$  depending only on  $r$  such that

$$P \left( \left| \frac{1}{n} \sum_{i=1}^n \langle X_i^{\otimes r}, V \rangle - \mathbb{E}[\langle X_i^{\otimes r}, V \rangle] \right| > a^r C_r \|V\|_1 b^r \sqrt{\frac{\log n}{n}} \right) \leq \frac{1}{n^a}$$

for any  $a \geq 1$  and  $V \in (\mathbb{R}^d)^{\otimes r}$ .

**Proof.** By Lemmas E.6 and E.7, for every  $i = 1, \dots, n$ ,

$$\|\langle X_i^{\otimes r}, V \rangle\|_{\psi_{1/r}} \leq C_r \sum_{j_1, \dots, j_k=1}^d \|X_{ij_1} \cdots X_{ij_r}\|_{\psi_{1/r}} |V_{j_1, \dots, j_k}| \leq C_r \|V\|_1 \max_j \|X_{ij}\|_{\psi_1}^r.$$

Therefore, by Lemma E.8, there exists a constant  $C'_r > 0$  depending only on  $r$  such that

$$\left\| \frac{1}{n} \sum_{i=1}^n (\langle X_i^{\otimes r}, V \rangle - \mathbb{E}[\langle X_i^{\otimes r}, V \rangle]) \right\|_p \leq C'_r \|V\|_1 b^r \left( \sqrt{\frac{p}{n}} + \frac{p^r}{n} \right)$$

for any  $p \geq 1$ . Hence, with  $p = a \log n$ , we have by Markov's inequality

$$P \left( \left| \frac{1}{n} \sum_{i=1}^n (\langle X_i^{\otimes r}, V \rangle - \mathbb{E}[\langle X_i^{\otimes r}, V \rangle]) \right| > e C'_r \|V\|_1 b^r \left( \sqrt{\frac{p}{n}} + \frac{p^r}{n} \right) \right) \leq e^{-p} = \frac{1}{n^a}.$$

Noting  $n^{-1} \log^r n = \sqrt{n^{-1} \log n} \cdot n^{-1/2} \log^{r-1/2} n \leq C_r \sqrt{n^{-1} \log n}$ , we complete the proof.  $\square$

**Lemma E.11.** *If (2.3) is satisfied, there exists a universal constant  $C > 0$  such that for any  $a \geq 1$ ,*

$$P \left( \max_{1 \leq j, k \leq d} \frac{1}{n} \sum_{i=1}^n |X_{ij}| |X_{ik}| 1_{\{|X_{ij}| \vee |X_{ik}| > 2b \log n\}} > C b^2 \left( \sqrt{\frac{a \log(dn)}{n}} + \frac{a^2 \log^2(dn)}{n} \right) \right) \leq \frac{1}{n^a} \quad (\text{E.3})$$

and

$$P \left( \max_{1 \leq j, k \leq d} \frac{1}{n} \sum_{i=1}^n X_{ij}^2 X_{ik}^2 1_{\{|X_{ij}| \vee |X_{ik}| \leq 2b \log n\}} > C b^4 \left( 1 + a \frac{\log(dn) \log^4 n}{n} \right) \right) \leq \frac{1}{n^a}. \quad (\text{E.4})$$

**Proof.** Let us prove (E.3). Write  $Y_{i,jk} := |X_{ij}| |X_{ik}| 1_{\{|X_{ij}| \vee |X_{ik}| > 2b \log n\}}$  for short. Since  $\|Y_{i,jk}\|_{\psi_{1/2}} \leq b^2$ , we have by Lemma E.9

$$\max_{1 \leq j, k \leq d} \frac{1}{n} \left| \sum_{i=1}^n (Y_{i,jk} - \mathbb{E}[Y_{i,jk}]) \right| \lesssim b^2 \left( \sqrt{\frac{a \log(dn)}{n}} + \frac{a^2 \log^2(dn)}{n} \right)$$

with probability at least  $1 - 1/n^a$ . Moreover, observe that

$$|\mathbb{E}[Y_{i,jk}]| \leq \max_{1 \leq j, k \leq d} \sqrt{\mathbb{E}[X_{ij}^2 X_{ik}^2] P(|X_{ij}| > 2b \log n)} \lesssim \frac{b^2}{n}.$$

Hence we conclude

$$\max_{1 \leq j, k \leq d} \frac{1}{n} \left| \sum_{i=1}^n Y_{i,jk} \right| \lesssim b^2 \left( \sqrt{\frac{a \log(dn)}{n}} + \frac{a \log^2(dn)}{n} \right)$$

with probability at least  $1 - 1/n^a$ . This proves (E.3).

Next we prove (E.4). Set

$$\zeta := \max_{1 \leq j, k \leq d} \frac{1}{n} \sum_{i=1}^n X_{ij}^2 X_{ik}^2 1_{\{|X_{ij}| \vee |X_{ik}| \leq 2b \log n\}}.$$

By Lemma E.5 in [19] with  $\eta = 3$  and  $B = 16b^4 \log^4 n/n$ , we have for any  $t > 0$

$$P\left(\zeta \geq 4\mathbb{E}[\zeta] + 16\frac{b^4 \log^4 n}{n}t\right) \leq e^{-t}.$$

Further,  $\mathbb{E}[\zeta] \lesssim b^4 + \frac{b^4 \log^4 n}{n} \log d$  by Lemma 9 in [18]. Hence there exists a universal constant  $C_2 > 0$  such that

$$P\left(\zeta \geq C_2 b^4 \left(1 + \frac{\log^4 n}{n}(\log d + t)\right)\right) \leq e^{-t}$$

for any  $t > 0$ . Applying this with  $t = a \log n$  gives (E.4).  $\square$

**Acknowledgments** The author thanks Xiao Fang and Ryo Imai for valuable discussions about the subject of this paper. This work was partly supported by JST CREST Grant Number JPMJCR2115 and JSPS KAKENHI Grant Numbers JP22H00834, JP22H00889, JP22H01139, JP24K14848.

## References

- [1] Anderson, N. H., Hall, P. & Titterton, D. (1998). Edgeworth expansions in very-high-dimensional problems. *J. Statist. Plann. Inference* **70**, 1–18.
- [2] Barbour, A. (1986). Asymptotic expansions based on smooth functions in the central limit theorem. *Probab. Theory Relat. Fields* **72**, 289–303.
- [3] Barthe, F. & Maurey, B. (2000). Some remarks on isoperimetry of Gaussian type. *Ann. Inst. Henri Poincaré Probab. Stat.* **36**, 419–434.
- [4] Belloni, A., Chernozhukov, V., Chetverikov, D., Hansen, C. & Kato, K. (2018). High-dimensional econometrics and regularized GMM. Working paper. arXiv: 1806.01888.
- [5] Beran, R. (1982). Estimated sampling distributions: The bootstrap and competitors. *Ann. Statist.* **10**, 212–225.
- [6] Beran, R. (1988). Prepivoting test statistics: A bootstrap view of asymptotic refinements. *J. Amer. Statist. Assoc.* **83**, 687–697.
- [7] Bhattacharya, R. N. & Rao, R. R. (2010). *Normal approximation and asymptotic expansions*. SIAM.
- [8] Bobkov, S. G. (1999). Isoperimetric and analytic inequalities for log-concave probability measures. *Ann. Probab.* **27**, 1903–1921.
- [9] Bobkov, S. G. & Houdré, C. (1997). Isoperimetric constants for product probability measures. *Ann. Probab.* **25**, 184–205.
- [10] Bobkov, S. G. & Ledoux, M. (2019). *One-dimensional empirical measures, order statistics, and Kantorovich transport distances*, vol. 261 of *Memoirs of the American Mathematical Society*. American Mathematical Society.
- [11] Boucheron, S., Lugosi, G. & Massart, P. (2013). *Concentration inequalities: A nonasymptotic theory of independence*. Oxford University Press.
- [12] Bühlmann, P. & van de Geer, S. (2011). *Statistics for high-dimensional data*. Springer.

- [13] Cacoullos, T. & Papathanasiou, V. (1992). Lower variance bounds and a new proof of the central limit theorem. *J. Multivariate Anal.* **43**, 173–184.
- [14] Caffarelli, L. A. (2000). Monotonicity properties of optimal transportation and the FKG and related inequalities. *Comm. Math. Phys.* **214**, 547–563.
- [15] Chatterjee, S. (2014). *Superconcentration and related topics*. Springer.
- [16] Chen, X. & Kato, K. (2019). Randomized incomplete  $U$ -statistics in high dimensions. *Ann. Statist.* **47**, 3127–3156.
- [17] Chernozhukov, V., Chetverikov, D. & Kato, K. (2013). Gaussian approximations and multiplier bootstrap for maxima of sums of high-dimensional random vectors. *Ann. Statist.* **41**, 2786–2819.
- [18] Chernozhukov, V., Chetverikov, D. & Kato, K. (2015). Comparison and anti-concentration bounds for maxima of Gaussian random vectors. *Probab. Theory Related Fields* **162**, 47–70.
- [19] Chernozhukov, V., Chetverikov, D. & Kato, K. (2017). Central limit theorems and bootstrap in high dimensions. *Ann. Probab.* **45**, 2309–2353.
- [20] Chernozhukov, V., Chetverikov, D. & Kato, K. (2017). Detailed proof of Nazarov’s inequality. Unpublished paper. arXiv: 1711.10696.
- [21] Chernozhukov, V., Chetverikov, D., Kato, K. & Koike, Y. (2022). Improved central limit theorem and bootstrap approximation in high dimensions. *Ann. Statist.* **50**, 2562–2586.
- [22] Chernozhukov, V., Chetverikov, D., Kato, K. & Koike, Y. (2023). High-dimensional data bootstrap. *Annu. Rev. Stat. Appl.* **10**, 427–449.
- [23] Chernozhukov, V., Chetverikov, D. & Koike, Y. (2023). Nearly optimal central limit theorem and bootstrap approximations in high dimensions. *Ann. Appl. Probab.* **33**, 2374–2425.
- [24] Deng, H. & Zhang, C.-H. (2020). Beyond Gaussian approximation: Bootstrap for maxima of sums of independent random vectors. *Ann. Statist.* **48**, 3643–3671.
- [25] Ding, J., Eldan, R. & Zhai, A. (2015). On multiple peaks and moderate deviations for the supremum of a Gaussian field. *Ann. Probab.* **43**, 3468–3493.
- [26] Fang, X. & Koike, Y. (2021). High-dimensional central limit theorems by Stein’s method. *Ann. Appl. Probab.* **31**, 1660–1686.
- [27] Fang, X. & Koike, Y. (2022). Sharp high-dimensional central limit theorems for log-concave distributions. *Ann. Inst. Henri Poincaré Probab. Stat. (forthcoming)*. Working paper is available at arXiv: 2207.14536.
- [28] Fang, X. & Koike, Y. (2023). From  $p$ -Wasserstein bounds to moderate deviations. *Electron. J. Probab.* **28**, 1–52.
- [29] Fang, X., Koike, Y., Liu, S.-H. & Zhao, Y.-K. (2023). High-dimensional central limit theorems by Stein’s method in the degenerate case. Working paper. arXiv: 2305.17365v1.
- [30] Fang, X. & Liu, S.-H. (2022). Edgeworth expansion by Stein’s method. Working paper. arXiv: 2211.04174.

- [31] Fathi, M. (2019). Stein kernels and moment maps. *Ann. Probab.* **47**, 2172–2185.
- [32] Fathi, M., Mikulincer, D. & Shenfeld, Y. (2024). Transportation onto log-Lipschitz perturbations. *Calc. Var. Partial Differential Equations* **63**, 61.
- [33] Fujikoshi, Y. & Ulyanov, V. V. (2020). *Non-asymptotic analysis of approximations for multivariate statistics*. Springer.
- [34] Fujikoshi, Y., Ulyanov, V. V. & Shimizu, R. (2010). *Multivariate statistics*. Wiley.
- [35] Hall, P. (1992). *The bootstrap and Edgeworth expansion*. Springer.
- [36] Koike, Y. (2021). Notes on the dimension dependence in high-dimensional central limit theorems for hyperrectangles. *Jpn. J. Stat. Data Sci.* **4**, 643–696.
- [37] Koike, Y. (2023). High-dimensional central limit theorems for homogeneous sums. *J. Theoret. Probab.* **36**, 1–45.
- [38] Kuchibhotla, A. K. & Chakraborty, A. (2022). Moving beyond sub-Gaussianity in high-dimensional statistics: Applications in covariance estimation and linear regression. *Inf. Inference* **11**, 1389–1456.
- [39] Kuchibhotla, A. K. & Rinaldo, A. (2020). High-dimensional CLT for sums of non-degenerate random vectors:  $n^{-1/2}$ -rate. Working paper. arXiv: 2009.13673.
- [40] Leadbetter, M. R., Lindgren, G. & Rootzén, H. (1983). *Extremes and related properties of random sequences and processes*. Springer.
- [41] Ledoux, M., Nourdin, I. & Peccati, G. (2015). Stein’s method, logarithmic Sobolev and transport inequalities. *Geom. Funct. Anal.* **25**, 256–306.
- [42] Ley, C., Reinert, G. & Swan, Y. (2017). Stein’s method for comparison of univariate distributions. *Probab. Surv.* **14**, 1–52.
- [43] Liu, R. Y. & Singh, K. (1987). On a partial correction by the bootstrap. *Ann. Statist.* **15**, 1713–1718.
- [44] Lopes, M. E. (2022). Central limit theorem and bootstrap approximation in high dimensions with near  $1/\sqrt{n}$  rates. *Ann. Statist.* **50**, 2492–2513.
- [45] Lopes, M. E., Lin, Z. & Müller, H.-G. (2020). Bootstrapping max statistics in high dimensions: Near-parametric rates under weak variance decay and application to functional and multinomial data. *Ann. Statist.* **48**, 1214–1229.
- [46] Mammen, E. (1993). Bootstrap and wild bootstrap for high dimensional linear models. *Ann. Statist.* **21**, 255–285.
- [47] Mijoule, G., Raič, M., Reinert, G. & Swan, Y. (2023). Stein’s density method for multivariate continuous distributions. *Electron. J. Probab.* **28**, 1–40.
- [48] Mikulincer, D. (2022). A CLT in Stein’s distance for generalized Wishart matrices and higher-order tensors. *Int. Math. Res. Not. IMRN* **2022**, 7839–7872.
- [49] Nourdin, I. & Peccati, G. (2009). Stein’s method on Wiener chaos. *Probab. Theory Related Fields* **145**, 75–118.

- [50] Nourdin, I. & Peccati, G. (2012). *Normal approximations with Malliavin calculus: From Stein's method to universality*. Cambridge University Press.
- [51] Nourdin, I., Peccati, G. & Swan, Y. (2014). Entropy and the fourth moment phenomenon. *J. Funct. Anal.* **266**, 3170–3207.
- [52] Nualart, D. (2006). *The Malliavin calculus and related topics*. Springer, 2nd edn.
- [53] Raič, M. (2019). A multivariate Berry–Esseen theorem with explicit constants. *Bernoulli* **25**, 2824–2853.
- [54] Rudin, W. (1987). *Real and complex analysis*. McGraw-Hill, 3rd edn.
- [55] Stein, C. (1986). *Approximate computation of expectations*. Institute of Mathematical Statistics.
- [56] van der Vaart, A. W. & Wellner, J. A. (2023). *Weak convergence and empirical processes*. Springer, 2nd edn.
- [57] Zhilova, M. (2022). New Edgeworth-type expansions with finite sample guarantees. *Ann. Statist.* **50**, 2545–2561.