

12th Workshop on the CKM Unitarity Triangle
Santiago de Compostela, 18-22 September 2023

Measurement of the CP violating phase ϕ_s and $\phi_s^{sq\bar{q}}$ at LHCb

Melissa Maria Cruz Torres on behalf of the LHCb collaboration
email: melissa.cruz@unah.edu.hn¹

¹Universidad Nacional Autónoma de Honduras, Honduras

April 9, 2024

Abstract

Precise measurements of the $B_s^0 - \bar{B}_s^0$ mixing parameters provide a powerful test of the Standard Model, offering potential hints to new physics. The LHCb collaboration has performed updated measurements of the CP -violating phases $\phi_s^{c\bar{c}s}$ and $\phi_s^{s\bar{s}s}$, which supersede previous results. Also, an alternative approach to determine $\Delta\Gamma_s$ is presented, bringing a new tool that may help to resolve the tension observed between measurements made in $B_s^0 \rightarrow J/\psi\phi$ by LHC experiments.

1 Introduction

One of the key goals of the LHCb experiment is the measurement of the CP -violating phase, ϕ_s , that originates from the interference of the mixing and direct decay of B_s mesons to CP eigenstates. Within the Standard Model ϕ_s is predicted to be equal to $-2\beta_s$, where $\beta_s \equiv \arg[-(V_{ts}V_{tb}^*)/(V_{cs}V_{cb}^*)]$, ignoring subleading penguin contributions, and where V_{ij} represents the Cabibbo-Kobayashi-Maskawa matrix elements [1]. In this sense, precise measurements of this phase enhance the potentiality to probe physics beyond the Standard Model [2]

2 Measurement of ϕ_s in $B_s^0 \rightarrow J/\psi K^+ K^-$ decays

The study of the time-dependent CP asymmetry in decays modes with transitions of the form $b \rightarrow c\bar{c}s$ has been performed by several experiments. It provides

insights into understanding the CP violation phenomena. The world average value of the CP -violating phase, $\phi_s^{c\bar{c}s}$, is found to be -0.049 ± 0.019 rad [3], which is dominated by the LHCb results in the decay channel $B_s^0 \rightarrow J/\psi h^+ h^-$ ($h = K$ or π) [4]. An update of the phase, $\phi_s^{c\bar{c}s}$, as well of the physics parameters $|\lambda|$, $\Delta\Gamma_s$, $\Delta\Gamma_s - \Gamma_d$ and the B_s^0 mass difference Δm_s , have been performed using the golden channel $B_s^0 \rightarrow J/\psi K^+ K^-$ in the vicinity of $\phi(1020)$, which is reported in this talk. These results supersede the previous ones. The dataset used includes the full data sample from 2015 to 2018, collected by the LHCb detector corresponding to an integrated luminosity of 6 fb^{-1} at $\sqrt{s} = 13 \text{ TeV}$.

The analysis strategy consists in performing a flavour-tagger time-dependent angular analysis, where four polarization states are identified, namely $A_0, A_{||}, A_{\perp}$ for the P-wave and A_s for the S-wave, regarding to the polarization states of the $K^+ K^-$ system. Candidates with $K^+ K^-$ invariant masses in the range $[900, 1050] \text{ MeV}/c^2$ from $B_s^0 \rightarrow J/\psi(\rightarrow \mu^+ \mu^-) K^+ K^-$ decays are selected following the strategy described in reference [4]. Experimental improvements benefit the updated measurements, as to be the flavour-tagging algorithms, the B_s^0 decay-time resolution model as well as in the determination of the particle identification. A gradient-boosted decision tree (BDT) classifier is trained separately for each year from 2016 to 2018, improving the signal to background ratio by a factor of 50. The Cross-feed contamination due to pion and proton misidentification is handled by appropriate particle identification requirements and mass constraints.

The data sample is divided into 48 independent sub samples, corresponding to six bins in the $\phi(1020)$ region, two trigger configurations and four year of data taking. By performing a maximum likelihood fit, a total yield of about 349 000 signal decays are extracted. The fit to the invariant mass is shown in Fig. 1.

To determine the phase ϕ_s in $B_s^0 \rightarrow J/\psi K^+ K^-$, a weighted simultaneous fit to the distributions of decay time and decay angles ($\cos\theta_K, \cos\theta_{\mu}, \phi_h$) in the helicity basis is performed in the 48 sub samples. Along with the determination of ϕ_s , the physics parameters and the polarization amplitudes $A_k = |A_k| e^{-i\delta_k}$ are also determined. The sub-index, k , in the latter, stands for polarization states of the $K^+ K^-$ system. The probability density function for the signal takes into consideration the decay-time and angular efficiencies, decay-time resolution and flavour tagging. The results from the fit are shown in Table 1 [6]. The background subtracted data distribution with fit projections overlaid can be seen in Fig 2.

The results are in good agreement with LHCb Run 1 [5] and 2015+2016 [4] measurements. The obtained values for ϕ_s , $\Delta\Gamma_s$ and $\Gamma_s - \Gamma_d$ represent the most precise measurements to date and are in good agreement with SM expectations. No evidence for CP violation is found. Results also show no evidence for polarization dependence of ϕ_s . Combination of all LHCb ϕ_s measurements of B_s^0 decays via $b \rightarrow c\bar{c}s$ is shown in Figure 3. The combined value is $\phi_s = -0.031 \pm 0.018$ rad.




Figure 1: Invariant mass distribution of $B_s^0 \rightarrow J/\psi K^+ K^-$ for the full data sample. The projection of the maximum likelihood fit is overlaid [6].

3 CP violation measurements in the penguin-mediated decay $B_s^0 \rightarrow \phi\phi$

B mesons decays with Flavour-Changing Neutral Current (FCNC) transitions provides a very sensitive ground to search for new physics. One of the benchmark channels to study at LHCb, in FCNC decays, is the penguin-dominated process $B_s^0 \rightarrow \phi\phi$ where $\phi \rightarrow K^+ K^-$ as loop contributions could reveal new sources of CP violation. A time-dependent angular analysis, using the full Run 2 dataset, is performed on this channels to determine the CP -violating parameters $\phi_s^{s\bar{s}s}$ and $|\lambda|$. Any deviation of their expected values, 0 or 1 respectively, could be an indication of new physics entering in the penguin decay or the B_s^0 mixing. The analysis is performed using the full Run 2 data sample.

The $B_s^0 \rightarrow \phi\phi$ candidates are selected in the [5150,5600] MeV/c^2 mass range, yielding about 16,000 of signal events. The three polarization states B_s^0 are considered, namely A_0 , A_{\parallel} and A_{\perp} and both, polarization-dependent and polarization-independent, scenarios are evaluated. The parameters $\phi_{s,i}$ and $|\lambda_i|$ are defined by the equation

$$\frac{q}{p} \cdot \frac{\bar{A}_i}{A_i} = \eta_i |\lambda_i| e^{-i\phi_{s,i}}, \quad (3.1)$$

where η_i is the CP eigenvalue of the polarization state i , q and p are complex

Table 1: Physics parameters results. The first uncertainty is statistical and the second systematic [6]

Parameter	Result
ϕ_s [rad]	$-0.039 \pm 0.022 \pm 0.006$
$ \lambda $	$1.001 \pm 0.011 \pm 0.005$
$\Gamma_s - \Gamma_d$ [ps $^{-1}$]	$-0.0056 \begin{smallmatrix} +0.0013 \\ -0.0015 \end{smallmatrix} \pm 0.0014$
$\Delta\Gamma_s$ [ps $^{-1}$]	$0.0845 \pm 0.0044 \pm 0.0024$
Δm_s [ps $^{-1}$]	$17.743 \pm 0.033 \pm 0.009$
$ A_\perp ^2$	$0.2463 \pm 0.0023 \pm 0.0024$
$ A_0 ^2$	$0.5179 \pm 0.0017 \pm 0.0032$
$\delta_\perp - \delta_0$ [rad]	$2.903 \begin{smallmatrix} +0.075 \\ -0.074 \end{smallmatrix} \pm 0.048$
$\delta_\parallel - \delta_0$ [rad]	$3.146 \pm 0.061 \pm 0.052$




Figure 2: Background-subtracted data distribution with fit projections for decay-time and decay-angles [6].

numbers relating the B_s^0 mass eigenstates to the flavour eigenstates. The differential decay rate is written as

$$\frac{d^4\Gamma(t, \vec{\Omega})}{dt d\vec{\Omega}} \propto \sum_{k=1}^6 h_k(t) f_s(\vec{\Omega}), \quad (3.2)$$

where t is the decay time and $\vec{\Omega} = (\theta_1, \theta_2, \chi)$ refer to the helicity angles of the K^+ mesons in the ϕ rest frame. χ is the angle between the $\phi \rightarrow KK$ decay




Figure 3: ϕ_s combination for all LHCb results [4].

planes. The angular function $f_k(\vec{\Omega})$ and the time-dependent function $h_k(t)$ are defined as given in references [7] and [8], respectively. After applying a maximum-likelihood fit to the distribution of t , $\vec{\Omega}$ and the initial B_s^0 state, the polarization-independent measurements of the CP -violation parameters are reported to be $\phi_s^{s\bar{s}s} = -0.074 \pm 0.069$ rad and $|\lambda| = 1.009 \pm 0.030$ in combination with Run 1 results [4]. This is the most precise measurement of CP violation in $B_s^0 \rightarrow \phi\phi$ decays to date [9] as can be seen in Fig 4. The results solely due the Run 2 data are reported to be $\phi_s^{s\bar{s}s} = -0.042 \pm 0.075 \pm 0.009$ rad and $|\lambda| = 1.004 \pm 0.030 \pm 0.009$, where the first uncertainty is statistical and the second systematic [9]. The total measurable observables can be seen in Table 2. For the polarization-dependent scenario, no dependence of the CP -violation parameters in the polarization states is found.

Table 2: Measured observables in the polarization-independent fit for the full Run 2 dataset [9].

Parameter	Result
$\phi_s^{s\bar{s}s}$ [rad]	$-0.042 \pm 0.075 \pm 0.009$
$ \lambda $	$1.004 \pm 0.030 \pm 0.009$
$ A_0 ^2$	$0.384 \pm 0.007 \pm 0.003$
$ A_\perp ^2$	$0.310 \pm 0.006 \pm 0.003$
$\delta_\parallel - \delta_0$ [rad]	$2.463 \pm 0.029 \pm 0.009$
$\delta_\perp - \delta_0$ [rad]	$2.769 \pm 0.105 \pm 0.011$




Figure 4: Comparison of $\phi_s^{s\bar{s}s}$ measurements by LHCb experiment. Results presented in this work are denoted as Run 1 + Run 2, 9 fb^{-1} and Run 2, 6 fb^{-1} [9]. The blue vertical line represents the SM prediction.

4 A measurement of $\Delta\Gamma_s$ from $B_s^0 \rightarrow J/\psi\eta'$ and $B_s^0 \rightarrow J/\psi\pi^+\pi^-$ decays

As explained before, the measurement of the $B_s^0 - \bar{B}_s^0$ mixing parameters offer a powerful test of the Standard Model. Particularly, $\Delta\Gamma_s$ have been determined experimentally using the golden channel $B_s^0 \rightarrow J/\psi\phi$ by ATLAS [10], CMS [11] and LHCb [6] experiments. The results are precise but in tension with each other. In this analysis, an alternative approach to determine $\Delta\Gamma_s$ that follows closely the formalism proposed in [12], is presented.

Given that CP -odd modes measure the heavy mass eigenstate lifetime ($\tau_H = 1/\Gamma_H$) and CP -even modes measure the light mass eigenstate lifetime ($\tau_L = 1/\Gamma_L$), $\Delta\Gamma_s$ can be determined from the decay-width difference between a CP -odd and a CP -even B_s^0 state. In this study it is used the CP -even decay $B_s^0 \rightarrow J/\psi\eta'$ and the CP -odd decay $B_s^0 \rightarrow J/\psi\pi^+\pi^-$, where $J/\psi \rightarrow \mu^+\mu^-$ and $\eta' \rightarrow \rho^0\gamma$.

If CP violation is considered negligible the time dependent rate can be expressed as

$$\Gamma(B_s^0(t) \rightarrow f) \propto e^{-\Gamma_s t} \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) + \eta_{CP} \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) \right], \quad (4.1)$$

where η_{CP} is 1 or -1 for CP -odd or CP -even states, respectively. By integrating Eq. 4.1 over a time range $[t_1, t_2]$, and performing their ratio, it is obtained the following equation

$$R_i = \frac{N_L}{N_H} \propto \frac{[e^{-\Gamma_s t(1+y)}]_{t_1}^{t_2}}{[e^{-\Gamma_s t(1-y)}]_{t_1}^{t_2}} \cdot \frac{(1-y)}{(1+y)}, \quad (4.2)$$

the parameter y is defined to be $\Delta\Gamma_s/2\Gamma_s$ and experimental corrections are taken into account into; R_i is corrected by the relative efficiency in each decay time bin as $R_i = A_i \frac{N_L^{RAW}}{N_H^{RAW}}$. A total of 8 time bins are chosen, whose ranges are determined using simulation samples. Similar yields are expected in each bin and it follows the purpose of optimizing the sensitivity of $\Delta\Gamma_s$. Finally, $\Delta\Gamma_s$ is determined from χ^2 minimization of the expression above (Eq.4.2), where $\Delta\Gamma_s$ and a normalization factor are free parameters.

The results are presented in Table 3. The weighted average is reported to be $\Delta\Gamma_s = 0.087 \pm 0.012 \pm 0.0009 \text{ ps}^{-1}$, where the first uncertainty is statistical and the second systematic [13]. The study was performed using the full data sample of proton-proton collisions collected by the LHCb between 2011 and 2018 corresponding to a integrated luminosity of 9 fb^{-1} in the center-of-mass energy $\sqrt{s} = 7, 8$ and 13 TeV . The comparison of the results by years of datasets are shown in Fig. 5.




Figure 5: Measurements of $\Delta\Gamma_s$ for the sets of data taking and the weighted average. The orange band represents 1σ error [13]

In summary, the value obtained for $\Delta\Gamma_s$ is in agreement with the HFLAV average value, $\Delta\Gamma_s = 0.074 \pm 0.006 \text{ ps}^{-1}$ [14], obtained from time-dependent angular

analyses of $B_s^0 \rightarrow J/\psi\phi$, where the initial flavour of the state is tagged. The value reported also agrees with the HFLAV average, $\Delta\Gamma_s = 0.083 \pm 0.005 \text{ ps}^{-1}$ [14], that includes constraints from other untagged effective lifetime measurements. $\Delta\Gamma_s$ is measured for the first time using the decay mode $B_s^0 \rightarrow J/\psi\eta'$.

Table 3: $\Delta\Gamma_s$ results and χ^2 probability for the four datasets [13].

Dataset	$\Delta\Gamma_s$ [ps^{-1}]	$P(\chi^2)$
2011&12	0.039 ± 0.026	0.83
2015&16	0.081 ± 0.022	0.77
2017	0.117 ± 0.024	0.57
2018	0.102 ± 0.021	0.78

References

- [1] Kobayashi, M. and Maskawa, T. CP-violation in the renormalizable theory of weak interaction. Prog. Theor. Phys 1973, 49(1973), 652. Cabibbo, N. Unitary symmetry and leptonic decays. Phys. Rev. Lett. 1963, 10(1963), 531. <https://doi.org/10.1143/PTP.49.652> <https://doi.org/10.1103/PhysRevLett.10.531>
- [2] Buras, A. J. Flavour theory. PoS EPS-HEP2009, 024 (2009); <https://doi.org/10.48550/arXiv.0910.1032>
- [3] Workman, R. L. et al. Particle Data Group. Review of particle physics. Prog. Theor. Exp. Phys. 2022, 083C01 (2022); <https://pdg.lbl.gov/> <https://doi.org/10.1093/ptep/ptac097>
- [4] The LHCb collaboration., Aaij, R., et al. Updated measurement of time-dependent CP-violating observables in $B_s^0 \rightarrow J/\psi K^+ K^-$ decays. Eur. Phys. J. 2019, C79 (2019), 706. Erratum ibid. C80 (2020), 601.; <https://doi.org/10.1140/epjc/s10052-019-7159-8> <https://doi.org/10.1140/epjc/s10052-020-7875-0>
- [5] The LHCb collaboration., Aaij, R., et al. Precision measurement of CP violation in $B_s^0 \rightarrow J/\psi K^+ K^-$ decays. Phys. Rev. Lett 2015, 114 (2015), 041801; <https://doi.org/10.1103/PhysRevLett.114.041801>
- [6] The LHCb collaboration., Aaij, R., et al. Improved measurement of CP violation parameters in $B_s^0 \rightarrow J/\psi K^+ K^-$ decays in the vicinity of the $\phi(1020)$ resonance. Phys. Rev. Lett 2024, 132 (2024), 051802; <https://doi.org/10.1103/PhysRevLett.132.051802>
- [7] The LHCb collaboration., Aaij, R., et al. Measurement of CP violation in the $B_s^0 \rightarrow \phi\phi$ decay and search for the $B_s^0 \rightarrow \phi\phi$ decay. JHEP 2019, 12 (2019), 155; [https://doi.org/10.1007/JHEP12\(2019\)155](https://doi.org/10.1007/JHEP12(2019)155)

- [8] Pree du, T. Search for a Strange Phase in Beautiful Oscillations, PhD thesis. Vrije U., Amsterdam, 2010.
- [9] The LHCb collaboration., Aaij, R., et al. Precision measurement of CP violation in the penguin-mediated decay $B_s^0 \rightarrow \phi\phi$. Phys. Rev. Lett 2023, 131 (2019), 171802; <https://doi.org/10.1103/PhysRevLett.131.171802>
- [10] The ATLAS collaboration., Aad, G., et al. Measurement of the CP -violating phase ϕ_s in $B_s^0 \rightarrow J/\psi\phi$ decays in ATLAS at 13 TeV. Eur. Phys. J. 2021, C81 (2021), 342; <https://doi.org/10.1140/epjc/s10052-021-09011-0>
- [11] The CMS collaboration., Sirunyan, A. M., et al. Measurement of the CP -violating phase ϕ_s in $B_s^0 \rightarrow J/\psi\phi \rightarrow \mu^+\mu^-K^+K^-$ channel in proton-proton collisions at $\sqrt{s} = 13$ TeV. Phys. Lett. 2021, B816 (2021), 136188; <https://doi.org/10.1016/j.physletb.2021.136188>
- [12] Fleischer, R. and Knegjens, R. Effective lifetime of B_s decays and $B_s^0 - \bar{B}_s^0$ mixing parameter. Eur. Phys. J 2011, C71 (2022), 1789; <https://doi.org/10.1140/epjc/s10052-011-1832-x>
- [13] The LHCb collaboration., Aaij, R., et al. A measurement of $\Delta\Gamma_s$. Phys. arXiv:2310.12649; <https://doi.org/10.48550/arXiv.2310.12649>
- [14] Amhis, Y. et al. Averages of b-hadron, c-hadron, and τ -lepton properties as of 2021. Phys. Rev. 2023, D107 (2023), 052008; <https://doi.org/10.1103/PhysRevD.107.052008>