

General position sets, colinear sets, and Sierpiński product graphs

Jing Tian ^{a,b}
 Sandi Klavžar ^{b,c,d}

^a School of Science, Zhejiang University of Science and Technology,
 Hangzhou, Zhejiang 310023, PR China

`jingtian526@126.com`

^b Institute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia

^c Faculty of Mathematics and Physics, University of Ljubljana, Slovenia

^d Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia

`sandi.klavzar@mf.uni-lj.si`

Abstract

Let $G \otimes_f H$ denote the Sierpiński product of G and H with respect to the function f . The Sierpiński general position number $\text{gps}(G, H)$ is introduced as the cardinality of a largest general position set in $G \otimes_f H$ over all possible functions f . Similarly, the lower Sierpiński general position number $\underline{\text{gps}}(G, H)$ is the corresponding smallest cardinality. The concept of vertex-colinear sets is introduced. Bounds for the general position number in terms of extremal vertex-colinear sets, and bounds for the (lower) Sierpiński general position number are proved. The extremal graphs are investigated. Formulas for the (lower) Sierpiński general position number of the Sierpiński products with K_2 as the first factor are deduced. It is proved that if $m, n \geq 2$, then $\text{gps}(K_m, K_n) = m(n - 1)$ and that if $n \geq 2m - 2$, then $\underline{\text{gps}}(K_m, K_n) = m(n - m + 1)$.

Keywords: general position set; colinear set; Sierpiński product of graphs; Sierpiński general position number

AMS Subj. Class. (2020): 05C12, 05C76

1 Introduction

General position sets were introduced to graph theory in [16], but Chandran S.V. and Parthasarathy in [3] earlier studied the concept of geodetic irredundant sets which turned out to be an equivalent concept to general position sets. Moreover, yet much earlier Körner [14] studied the general position sets of hypercubes (in a completely different context thought).

The problem of finding a largest general position set in a graph, that is, the general position number of the graph, is NP-hard [16]. In [1], general position sets of a graph were characterized. Afterwards, the problem of determining the general position number of (classes of) graphs received a wide attention, see [7, 19, 24, 26]. General position sets were generalized to d -position sets [10] and to Steiner general position sets [8]. The edge version of general position sets were investigated in [12, 17, 22], while for the monophonic version see [21].

A lot of attention has been given to the general position sets in Cartesian product graphs, see [11, 13, 23]. In this paper we are interested in general position sets in the recently introduced, attractive graph operation called the Sierpiński product. The operation was introduced by Kovič, Pisanski, Zemljič, and Žitnik in [15] with the idea to generalize the Sierpiński graphs [9]. The investigation of the latter graphs up to 2017 is summarized in the survey [6], some of the recent research of Sierpiński graphs can be found, for instance, in [18, 20, 25], see also the references therein. The Sierpiński product was further investigated in [4, 5], where the (upper) Sierpiński domination number and the (upper) Sierpiński metric dimension were respectively introduced. Following this line of investigation, in this paper we introduce the Sierpiński general position number $\text{gps}(G, H)$, and the lower Sierpiński general position number $\underline{\text{gps}}(G, H)$, of graphs G and H .

In the rest of the paper we proceed as follows. In the next section we give definitions and recall known results needed. In Section 3 we introduce a new concept into the theory of graph general position sets, the vertex-colinear sets. This concept is useful while studying the general position sets in Sierpiński product graphs, but we believe to be also of independent interest. In particular, we prove bounds for the general position number of a graph in terms of extremal vertex-colinear sets and show their usefulness for graphs with bridges. In Section 4, we present bounds for the (lower) Sierpiński general position number for general graphs and give formulas for these numbers for the Sierpiński product graphs with K_2 as the first factor. In Section 5, we consider the Sierpiński product of complete graphs $K_m \otimes_f K_n$. We determine their Sierpiński general position number, more precisely, if $m, n \geq 2$, then $\text{gps}(K_m, K_n) = m(n - 1)$. For the lower Sierpiński general position number we prove that $\underline{\text{gps}}(K_m, K_n) = m(n - m + 1)$ provided that $n \geq 2m - 2$. The

assumption $n \geq 2m - 2$ is required as demonstrated by our last result which asserts that $\underline{\text{gps}}(K_6, K_9) = 25$.

2 Preliminaries

In this section we define concepts, notation, and results needed, and introduce the (lower) Sierpiński general position number. We begin by basic definitions, and follow by introducing the general position sets and the Sierpiński product graphs.

For a positive integer k we set $[k] = \{1, \dots, k\}$. Unless stated otherwise, all graphs $G = (V(G), E(G))$ in the paper are connected. The *order* $n(G)$ of G is equal $|V(G)|$. The *degree* of a vertex u , $\deg_G(u)$, is the number of adjacent vertices of u in G . Vertices of degree one are called *leaves*. The number of leaves of G will be denoted by $\ell(G)$. For a graph G , let $v_k(G)$ denote the number of vertices of G of degree k , cf. [2]. An edge e of G is a *bridge* if $G - e$ is disconnected. A vertex of G is *simplicial* if its neighbourhood induces a complete subgraph. If $S \subseteq V(G)$, then the subgraph of G induced by S is denoted by $G[S]$. The *distance* $d_G(u, v)$ between vertices u and v of G is the number of edges on a shortest u, v -path. The *interval* between vertices u and v is

$$I_G[u, v] = \{w : d_G(u, v) = d_G(u, w) + d_G(w, v)\}.$$

A subgraph H of G is *isometric* if for each pair of vertices $u, v \in V(H)$ we have $d_H(u, v) = d_G(u, v)$ and is *convex* if whenever $u, v \in V(H)$ and P is a shortest u, v -path in G , then P lies completely in H .

Let $X \subseteq V(G)$. Then X is a *general position set* of G if as soon as P is a shortest u, v -path, we have $V(P) \cap X = \{u, v\}$. The cardinality of a largest general position set of G is the *general position number*, $\text{gp}(G)$, of G . A general position set X of cardinality $\text{gp}(G)$ is referred to as a *gp-set* of G . Clearly, if G has at least two vertices, then $\text{gp}(G) \geq 2$.

Let G and H be graphs and let $f: V(G) \rightarrow V(H)$ be a function. The *Sierpiński product* of G and H (with respect to f) is the graph $G \otimes_f H$ with vertices

- $V(G \otimes_f H) = V(G) \times V(H)$,

and with edges

- $(g, h)(g, h')$, where $g \in V(G)$ and $hh' \in E(H)$, and
- $(g, f(g'))(g', f(g))$, where $gg' \in E(G)$.

The second type of the just defined edges will be called the *connecting edges* of $G \otimes_f H$. Note that for each vertex $g \in V(G)$, the subgraph of $G \otimes_f H$ induced by the set of vertices $\{(g, h) : h \in V(H)\}$, is isomorphic to H ; it will be denoted by gH .

Let G and H be connected graphs and denote by H^G the set of functions from $V(G)$ to $V(H)$. We introduce the *Sierpiński general position number*, $\text{gps}(G, H)$, as the cardinality of a largest general position set in $G \otimes_f H$ over all possible functions $f \in H^G$ and the *lower Sierpiński general position number*, $\underline{\text{gps}}(G, H)$, as the corresponding smallest cardinality. That is,

$$\text{gps}(G, H) = \max_{f \in H^G} \{\text{gp}(G \otimes_f H)\} \quad \text{and} \quad \underline{\text{gps}}(G, H) = \min_{f \in H^G} \{\text{gp}(G \otimes_f H)\}.$$

At the end of the section, we will call up several known results needed later on. For the first one, known as *Isometric Cover Lemma*, we need the following definition. A set of subgraphs $\{H_1, \dots, H_k\}$ is an *isometric cover* of G if each H_i , $i \in [k]$, is isometric in G and $\bigcup_{i=1}^k V(H_i) = V(G)$.

Theorem 2.1 [16, Theorem 3.1] *If $\{H_1, \dots, H_k\}$ is an isometric cover of G , then*

$$\text{gp}(G) \leq \sum_{i=1}^k \text{gp}(H_i).$$

Theorem 2.2 [16, Theorem 3.6] *If S is the set of simplicial vertices of a block graph G , then $\text{gp}(G) = |S|$.*

If G is a connected graph, $S \subseteq V(G)$, and $\mathcal{P} = \{S_1, \dots, S_t\}$ a partition of S , then \mathcal{P} is *distance-constant* if for any $i, j \in [t]$, $i \neq j$, the distance $d_G(x, y)$, where $x \in S_i$ and $y \in S_j$, is independent of the selection of x and y . A distance-constant partition \mathcal{P} is *in-transitive* if $d_G(S_i, S_k) \neq d_G(S_i, S_j) + d_G(S_j, S_k)$ holds for $i, j, k \in [p]$. The following characterization of general position sets will be used either implicitly or explicitly in the rest of the paper.

Theorem 2.3 [1, Theorem 3.1] *Let G be a connected graph. Then $S \subseteq V(G)$ is a general position set if and only if the components of $G[S]$ are complete subgraphs, the vertices of which form an in-transitive, distance-constant partition of S .*

The last retrieved result is about the metric structure of Sierpiński product graphs.

Theorem 2.4 [5, Theorem 4.1] *If G and H are connected graphs, $f \in H^G$, and $g \in V(G)$, then gH is a convex subgraph of $G \otimes_f H$.*

3 Colinear sets

In this section we introduce a new concept into the theory of graph general position sets, the vertex-colinear sets. We believe that the concept is interesting in its own right. Moreover, from our aspect it turns out to be extremely useful for studying the general position sets in Sierpiński product graphs.

Let G be a connected graph and $u \in V(G)$. A set $S \subseteq V(G)$ is a u -colinear set if S is a general position set such that $u \notin S$ and $y \notin I_G[x, u]$ for any $x, y \in S$. We are interested in u -colinear sets of largest cardinality and hence set

$$\xi_G(u) = \max\{|S| : S \text{ is a } u\text{-colinear set}\}.$$

Further, let

$$\begin{aligned}\xi_m(G) &= \min\{\xi_G(u) : u \in V(G)\}, \\ \xi_M(G) &= \max\{\xi_G(u) : u \in V(G)\}.\end{aligned}$$

To illustrate these concepts, we state the following easy result.

Proposition 3.1 *If T is a tree with $n(T) \geq 3$ and $u \in V(T)$, then $\xi_T(u) = \ell(T) - 1$ if u is a leaf, and $\xi_T(u) = \ell(T)$, otherwise. In particular, $\xi_m(T) = \ell(T) - 1$ and $\xi_M(T) = \ell(T)$.*

Proof. Assume first that u is a leaf and let S be a u -colinear set. Let u' be the support vertex adjacent to u . In the following arguments we use the consequence of Theorem 2.2 asserting that the general position number of a tree is the number of its leaves. If $\deg_T(u') > 2$, then $\text{gp}(T - u) = \ell(T) - 1$, and then the leaves of $T - u$ form a largest u -colinear set. Assume next that $\deg_T(u') = 2$. Then $T - u$ contains $\ell(T)$ leaves (one of them being u'). However, if u' lies in a u -colinear set, then no other vertex lies in such a set. Hence we can again conclude that $\xi_T(u) = \ell(T) - 1$. Since $n(T) \geq 3$, there exists a vertex u which is not a leaf. Then the leaves of T form a u -colinear set and hence $\xi_T(u) = \ell(T)$ and so also $\xi_M(T) = \ell(T)$. \square

The above invariants fulfil the following chain of inequalities involving the general position number. This in part demonstrate that our definitions are meaningful, further reasons for their introduction will be given later on.

Theorem 3.2 *If G is a connected graph of order at least 2 and $u \in V(G)$, then*

$$\xi_m(G) \leq \xi_G(u) \leq \xi_M(G) \leq \text{gp}(G) \leq 2\xi_m(G).$$

Proof. The left three inequalities follow directly from definitions. To prove the last inequality, let u be a vertex of G such that $\xi_G(u) = \xi_m(G)$ and let X be an arbitrary gp-set of G . We distinguish two cases.

Assume first that $u \in X$. Since X is a general position set of G , we infer that $X \setminus \{u\}$ is a u -colinear set. Then it follows that $\xi_G(u) \geq |X| - 1$, hence $|X| \leq \xi_m(G) + 1$. Since the order of G is at least two, $\xi_m(G) \geq 1$ and thus we have $|X| \leq \xi_m(G) + \xi_m(G) \leq 2\xi_m(G)$.

Assume second that $u \notin X$. In this case we partition X into two sets X_1 and X_2 as follows. For every $x \in X$, we put each vertex from $(I_G[x, u] \setminus \{x\}) \cap X$ into X_2 . This defines X_2 , and then $X_1 = X \setminus X_2$. We claim that each of X_1 and X_2 is a u -colinear set. By the definition, this clearly holds for X_1 . Consider now two vertices $x, x' \in X_2$ and suppose by way of contradiction that there exists a shortest x, u -path $P_{x,u}$ that contains x' . Then $d_G(x, u) = d_G(x, x') + d_G(x', u)$. Since $x \in X_2$, there exists a vertex $x'' \in X$ and a shortest x'', u -path $P_{x'',u}$. We are going to show that x'', x , and x' lie on a common shortest path. If not, then $d_G(x'', x') < d_G(x'', x) + d_G(x, x')$, but then

$$\begin{aligned} d_G(x'', u) &= d_G(x'', x) + d_G(x, u) \\ &= d_G(x'', x) + d_G(x, x') + d_G(x', u) \\ &> d_G(x'', x') + d_G(x', u) \\ &= d_G(x'', u), \end{aligned}$$

which is not possible. We have thus seen that also X_2 is a u -colinear set. Therefore,

$$\xi_G(u) \geq \max\{|X_1|, |X_2|\} \geq \frac{1}{2}|X| = \frac{1}{2}\text{gp}(G),$$

hence $\text{gp}(G) \leq 2\xi_G(u) = 2\xi_m(G)$. \square

We now give some examples which demonstrate that in the inequality chain of Theorem 3.2 the inequalities can be sharp or not. If $n \geq 2$, then $\xi_m(K_{n,n}) = \xi_M(K_{n,n}) = \text{gp}(K_{n,n})$, hence for these graphs the left three equalities hold. On the other hand, $\xi_M(G)$ can be smaller than $\text{gp}(G)$. For instance, $\xi_M(K_n) = n - 1 = \text{gp}(K_n) - 1$ for $n \geq 2$. In addition, if P is the Petersen graph, then $\xi_m(P) = \xi_M(P) = 5$ while $\text{gp}(P) = 6$.

Consider next the cycle chain graphs C_{2k}^ℓ , $k, \ell \geq 2$, where C_{2k}^ℓ consists of ℓ cycles C_{2k} sharing a vertex, such that a middle cycle shares its diametral vertices with its two neighboring cycles. See Fig. 1 where C_6^5 is shown. Consider the vertex u of C_{2k}^ℓ as shown in Fig. 1. Then $\xi_{C_{2k}^\ell}(u) = 2$ and hence $\xi_m(C_{2k}^\ell) = 2$. On the other hand, for the vertex v we have $\xi_{C_{2k}^\ell}(v) = 4$ which in turn implies that $\xi_M(C_{2k}^\ell) = 4$.




Figure 1: The graph C_6^5 .

Thus $\xi_m(G)$ can be strictly smaller than $\xi_M(G)$. Moreover, the vertex v of C_{2k}^ℓ also demonstrates that in Theorem 3.2, $\xi_G(u)$ can be equal to $2\xi_m(G)$.

We conclude this section by demonstrating how colinear sets can be used while determining the general position number of graphs containing bridges.

Proposition 3.3 *Let $e = u_1u_2$ be a bridge of a connected graph G , and G_1, G_2 the two components of $G - e$, where $u_i \in V(G_i)$, $i \in [2]$. Then $\text{gp}(G) \geq \xi_{G_1}(u_1) + \xi_{G_2}(u_2)$. Moreover, the equality holds in block graphs containing at least one bridge.*

Proof. Let X_1 be a u_1 -colinear set of G_1 with cardinality $\xi_{G_1}(u_1)$, and let X_2 be an analogous u_2 -colinear set of G_2 with cardinality $\xi_{G_2}(u_2)$. Then it is straightforward to check that $X_1 \cup X_2$ is a general position set of G , hence $\text{gp}(G) \geq \xi_{G_1}(u_1) + \xi_{G_2}(u_2)$.

Consider the block graph G with bridges, and let $e = u_1u_2$ be an arbitrary bridge of G . Let G_1 and G_2 be the two components of $G - e$, where $u_1 \in V(G_1)$ and $u_2 \in V(G_2)$. From the above, we have $\text{gp}(G) \geq \xi_{G_1}(u_1) + \xi_{G_2}(u_2)$. Let S be the set of simplicial vertices of G and set $S_1 = S \cap V(G_1)$ and $S_2 = S \cap V(G_2)$. Since the vertices from S_i , $i \in [2]$, are simplicial vertices in G_i , we infer that S_i is a u_i -colinear set of G_i . Clearly, $|S_1| + |S_2| = |S|$ and by Theorem 2.2 we know that $\text{gp}(G) = |S|$. Putting all these fact together we have

$$\text{gp}(G) \geq \xi_{G_1}(u_1) + \xi_{G_2}(u_2) \geq |S_1| + |S_2| = |S| = \text{gp}(G),$$

which proves that $\text{gp}(G) = \xi_{G_1}(u_1) + \xi_{G_2}(u_2)$. \square

Next, we give two examples of non-block graphs showing that the inequality of Proposition 3.3 can be sharp or not. Consider the graph H as shown in Fig. 2. It is straightforward to see that $\text{gp}(H) = 6$ and $\xi_{H_1}(u_1) + \xi_{H_2}(u_2) = 3 + 2 = 5$, hence $\text{gp}(G)$ can be larger than $\xi_{G_1}(u_1) + \xi_{G_2}(u_2)$. On the other hand, to see that $\text{gp}(G)$ can be equal to $\xi_{G_1}(u_1) + \xi_{G_2}(u_2)$ on non-block graphs with bridges, consider the graph H' as shown in Fig. 2. We have $\text{gp}(H') = 6$ and $\xi_{H'_1}(u_1) + \xi_{H'_2}(u_2) = 3 + 3 = 6$.




Figure 2: The graphs H and H' .

4 Sierpiński products of arbitrary graphs

In this section we bound the (lower) Sierpiński general position number for general graphs. Then we give formulas for the (lower) Sierpiński general position number for the Sierpiński product of graphs with K_2 as the first factor. We begin with the following simple, but useful lemma.

Lemma 4.1 *Let X be a general position set of $K_2 \otimes_f H$, where $V(K_2) = [2]$ and $f(2) = u \in V(H)$. If $|X \cap V(1H)| \geq 2$ and $(1, u) \in X$, then $X \cap 2H = \emptyset$.*

Proof. Let $(1, v)$ be a vertex of X different from $(1, u)$. Then every vertex from $2H$, say $(2, x)$, is an endvertex of a shortest $(1, v)$, $(2, x)$ -path passing through $(1, u)$, hence $(2, x) \notin X$. \square

Theorem 4.2 *If G and H are two connected graphs of order at least 2, then*

$$\text{gp}(H) \leq \underline{\text{gps}}(G, H) \leq \text{gps}(G, H) \leq n(G)\text{gp}(H).$$

Moreover, $\text{gps}(G, H) = n(G)\text{gp}(H)$ if and only if $\text{gp}(H) = \xi_M(H)$.

Proof. Let $G \otimes_f H$ be an arbitrary Sierpiński product of graphs G and H . By Theorem 2.4, the product $G \otimes_f H$ contains $n(G)$ convex subgraphs gH , $g \in V(G)$. Since a convex subgraph is an isometric subgraph, these subgraphs form an isometric cover. Therefore, by Theorem 2.1, we have $\text{gps}(G, H) \leq n(G)\text{gp}(H)$, hence the right most inequality holds. The convexity of the subgraphs gH also implies the left inequality, while the middle inequality is obvious.

Assume now that $\text{gps}(G, H) = n(G)\text{gp}(H)$. Consider an arbitrary function $f \in H^G$. Using once more the convexity of the subgraphs gH , there exists a gp-set X of $G \otimes_f H$, such that $|V(gH) \cap X| = \text{gp}(H)$ for each $g \in V(G)$. Consider a fixed subgraph gH and let $gg' \in E(G)$. Since $|V(gH) \cap X| = \text{gp}(H)$ and $\text{gps}(G, H) = n(G)\text{gp}(H)$, Lemma 4.1 implies that $(g, f(g')) \notin X$. Setting $u = (g, f(g'))$ we obtain,

having in mind Lemma 4.1 again, that $\xi_{gH}(u) = \text{gp}(H)$ which in turn implies that $\xi_M(H) = \text{gp}(H)$.

Assume next that $\xi_M(H) = \text{gp}(H)$. Let u be a vertex of H such that $\xi_H(u) = \xi_M(H)$ and let $f \in H^G$ be the identity function $f(g) = u$ for any $g \in V(G)$. Each gH has at least $\xi_M(H)$ vertices from some gp-set of the graph $G \otimes_f H$, hence $\text{gp}(G \otimes_f H) \geq n(G)\xi_M(H) = n(G)\text{gp}(H)$. Thus we have $\text{gps}(G, H) \geq n(G)\text{gp}(H)$. Since gH is convex in $G \otimes_f H$, Theorem 2.1 gives $\text{gp}(G \otimes_f H) \leq n(G)\text{gp}(H)$ and then $\text{gps}(G, H) \leq n(G)\text{gp}(H)$. From the above, we conclude that $\text{gps}(G, H) = n(G)\text{gp}(H)$. \square

Corollary 4.3 *If G is a connected graph with $n(G) \geq 2$ and T is a tree with $n(T) \geq 3$, then $\text{gps}(G, T) = n(G)\ell(T)$.*

Proof. Since $n(T) \geq 3$, Proposition 3.1 gives $\text{gp}(T) = \ell(T) = \xi_M(T)$. Hence Theorem 4.2 yields the conclusion. \square

Theorem 4.4 *If H is a connected graph with $n(H) \geq 2$, then the following assertions hold.*

- (i) $\text{gps}(K_2, H) = 2\xi_m(H)$.
- (ii) $\text{gps}(K_2, H) = 2\xi_M(H)$.

Proof. Set $V(K_2) = [2]$ for the rest of the proof.

(i) Let $f \in H^{K_2}$ be an arbitrary function, say $f(1) = u$ and $f(2) = v$. By Proposition 3.3 we have $\text{gp}(K_2 \otimes_f H) \geq \xi_H(v) + \xi_H(u) \geq \xi_m(H) + \xi_m(H)$. It follows that $\text{gps}(K_2, H) \geq 2\xi_m(H)$.

To prove the reverse inequality, let u be a vertex of H such that $\xi_H(u) = \xi_m(H)$. Define $f \in H^{K_2}$ by $f(1) = f(2) = u$ and let X be a gp-set of $K_2 \otimes_f H$. If $V(1H) \cap X = \emptyset$ or $V(2H) \cap X = \emptyset$, then $\text{gp}(K_2 \otimes_f H) \leq \text{gp}(H)$ holds and then Theorem 3.2 gives $\text{gp}(K_2 \otimes_f H) \leq 2\xi_m(H)$. Assume second that $V(1H) \cap X \neq \emptyset$ and $V(2H) \cap X \neq \emptyset$. Then we claim that $|V(1H) \cap X| \leq \xi_H(u)$ and $|V(2H) \cap X| \leq \xi_H(u)$. By establishing this claim, the $\text{gp}(K_2 \otimes_f H) \leq 2\xi_m(H)$ follows. Suppose on the contrary that this is not the case and assume without loss of generality that $|V(1H) \cap X| \geq \xi_H(u) + 1$. Then there exist two vertices $(1, h), (1, h') \in V(1H) \cap X$, where $h \neq h'$, such that the vertices $(1, h)$, $(1, h')$, and $(2, u)$ lie on a common shortest path. If $(2, h'')$ is an arbitrary vertex from $V(2H) \cap X$, then the vertices $(1, h)$, $(1, h')$, and $(2, h'')$ lie on a common shortest path. It is impossible because we have assumed that $V(2H) \cap X \neq \emptyset$. This contradiction proves the claim.

(ii) Let $f \in H^{K_2}$ be an arbitrary function, say $f(1) = u$ and $f(2) = v$ and let X be an arbitrary gp-set of $K_2 \otimes_f H$. We claim that $\text{gp}(K_2 \otimes_f H) \leq 2\xi_M(H)$. Suppose

on the contrary that $\text{gp}(K_2 \otimes_f H) \geq 2\xi_M(H) + 1$. Since the order of H at least 2, Theorem 3.2 implies that $2\xi_M(H) \geq \text{gp}(H) \geq 2$. Then $\text{gp}(K_2 \otimes_f H) \geq \text{gp}(H) + 1 \geq 3$ and thus $|X \cap V(1H)| \geq 2$ or $|X \cap V(2H)| \geq 2$.

Assume, without loss of generality, that $|X \cap V(1H)| \geq 2$. If $(1, v) \in X$, then Lemma 4.1 implies that $X \cap V(2H) = \emptyset$ and then, using Theorem 3.2 once more, $\text{gp}(K_2 \otimes_f H) \leq \text{gp}(H) \leq 2\xi_m(H) \leq 2\xi_M(H)$, a contradiction. Hence assume in the rest that $(1, v) \notin X$. Since $\text{gp}(K_2 \otimes_f H) \geq \text{gp}(H) + 1$, we get $|X \cap V(2H)| \geq 1$. But as argued in (i), this is not possible. This contradiction implies that $|X \cap V(2H)| \leq \xi_H(u) \leq \xi_M(H)$. By symmetry we also have that if $|X \cap V(2H)| \geq 2$, then $|X \cap V(1H)| \leq \xi_H(v) \leq \xi_M(H)$ holds. We can thus conclude that $\text{gps}(K_2, H) \leq 2\xi_M(H)$.

To complete the argument we need to demonstrate that $\text{gps}(K_2, H) \geq 2\xi_M(H)$. Let x be a vertex of H such that $\xi_H(x) = \xi_M(H)$ and let the function $f \in H^{K_2}$ be defined by $f(1) = f(2) = x$. By Proposition 3.3, it follows that $\text{gp}(K_2 \otimes_f H) \geq 2\xi_H(x) = 2\xi_M(H)$. Then we have $\text{gps}(K_2, H) \geq 2\xi_M(H)$, and we are done. \square

5 Sierpiński products of complete graphs

In this section we consider the Sierpiński product of complete graphs $K_m \otimes_f K_n$. In the first main result we determine their Sierpiński general position number. For the lower Sierpiński general position number we prove that $\text{gps}(K_m, K_n) = m(n - m + 1)$ provided that $n \geq 2m - 2$. The latter assumption is required as demonstrated by our last result which asserts that $\text{gps}(K_6, K_9) = 25$.

Throughout this section we will assume that $V(K_m) = [m]$ and $V(K_n) = \{x_i : i \in [n]\}$.

Theorem 5.1 *If $m, n \geq 2$, then*

$$\text{gps}(K_m, K_n) = m(n - 1).$$

Proof. Let $f \in K_n^{K_m}$ be defined by $f(i) = x_1$ for $i \in [m]$. Let $X = \{(i, x_j) : i \in [m], 2 \leq j \leq n\}$. Since f is an identify function, for any two vertices $(i, x_j), (i', x'_j) \in X$, where $i \neq i'$, we have $d_{K_m \otimes_f K_n}((i, x_j), (i', x'_j)) = 3$. By Theorem 2.3, X is a general position set, hence $\text{gp}(K_m \otimes_f K_n) \geq m(n - 1)$. It follows that $\text{gps}(K_m, K_n) \geq \text{gp}(K_m \otimes_f K_n) \geq m(n - 1)$.

To prove equality, suppose on the contrary that $\text{gps}(K_m, K_n) \geq m(n - 1) + 1$. It means that there exists a function $f' \in K_n^{K_m}$ and an index $i \in [m]$, such that $V(iK_n)$ completely lies in some general position set S of $K_m \otimes_{f'} K_n$. Since $m, n \geq 3$, there exists a vertex $(i', x_j) \in S$, where $i \neq i'$. Assume that $f'(i') = x_p$ and let $q \in [n]$ be such that $q \neq p$. Then the vertices $(i, x_q) \in S$, $(i, x_p) \in S$, and $(i', x_j) \in S$ lie on a

common shortest path (of length 2) of $K_m \otimes_{f'} K_n$, a contradiction. This contradiction implies that $\text{gp}(K_m \otimes_{f'} K_n) \leq m(n-1)$, hence we have $\text{gps}(K_m, K_n) = m(n-1)$. \square

In the rest we focus on the lower Sierpiński general position number.

Lemma 5.2 *Let $m, n \geq 2$ and let $f \in K_n^{K_m}$. Then the following hold.*

(i) *If $m \leq n$, then $\text{gp}(K_m \otimes_f K_n) \geq v_{n-1}(K_m \otimes_f K_n) \geq (n-m+1)m$. In particular,*

$$\underline{\text{gps}}(K_m, K_n) \geq (n-m+1)m.$$

(ii) *If $m \geq n$, then $\text{gp}(K_m \otimes_f K_n) \geq \max\{2(n-1), m\}$. In particular,*

$$\underline{\text{gps}}(K_m, K_n) \geq \max\{2(n-1), m\}.$$

Proof. Let $f \in K_n^{K_m}$, and set $Z = K_m \otimes_f K_n$ for the rest of the proof.

(i) Note that a vertex of Z is of degree $n-1$ if and only if it is not incident to a connecting edge. For each $i \in [m]$, let X_i be the set of vertices from $V(Z) \cap V(iK_n)$ of degree $n-1$. (Note that it is possible that $X_i = \emptyset$.) Then X_i clearly induces a complete subgraph. Moreover, if $i \neq j$, and if $(i, x_k) \in X_i$ and $(j, x_\ell) \in X_j$, then $d_Z((i, x_k), (j, x_\ell)) = 3$. Theorem 2.3 implies that $\bigcup_{i=1}^m X_i$ is a general position set. This in turn yields $\text{gp}(K_m \otimes_f K_n) \geq v_{n-1}(K_m \otimes_f K_n)$ because each vertex not in $\bigcup_{i=1}^m X_i$ is of degree at least n . Moreover, since each iK_n contains at most $m-1$ connecting edges, we have $|X_i| \geq n-m+1$, so we also have $v_{n-1}(K_m \otimes_f K_n) \geq (n-m+1)m$.

The above argument holds true for any function $f \in K_n^{K_m}$, hence we have $\underline{\text{gps}}(K_m, K_n) \geq (n-m+1)m$.

(ii) Assume first that f is an identity function, without loss of generality let $f(i) = x_1$ for $i \in [m]$. By Theorem 2.3 we then get that $V(Z) \setminus \{(1, x_1), \dots, (m, x_1)\}$ is a general position set, hence in this case $\text{gp}(Z) \geq 2(n-1)$. Assume second that f is not an identity function and assume without loss of generality that $f(1) = x_1$ and $f(2) = x_2$. Now we claim that $S = V(1K_n) \cup V(2K_n) \setminus \{(1, x_2), (2, x_1)\}$ is a general position set. To prove this claim it suffices to show (again in view of Theorem 2.3) that if $(1, x_i), (2, x_j) \in S$, then $d_Z((1, x_i), (2, x_j)) = 3$. Suppose on the contrary that $d_Z((1, x_i), (2, x_j)) = 2$. Then there exists a vertex (p, x_k) , where $k \neq 1, 2$, such that $(1, x_i)(p, x_k) \in E(Z)$ and $(2, x_j)(p, x_k) \in E(Z)$. But this means that $f(1) = x_k$ and $f(2) = x_k$, a contradiction since we have assumed that $f(1) \neq f(2)$. This contradiction proves the claim. We conclude that S is a general position set of Z and as $|S| = 2(n-1)$ we have proved that $\text{gp}(Z) \geq 2(n-1)$.

To prove that $\text{gp}(Z) \geq m$, we may assume without loss of generality that the function f is non-decreasing with respect to the indices of x_i s. That is, let $k_1 \geq \dots \geq k_n \geq 0$, where $k_1 + \dots + k_n = m$, and set $k_0 = 0$. Then for $i \in [n]$,

$$f(j) = x_i \quad \text{for } j \in \{k_1 + \dots + k_{i-1} + 1, \dots, k_1 + \dots + k_i\}.$$

For $i \in [n]$, set

$$X_i = \{(j, x_i) : j \in \{k_1 + \dots + k_{i-1} + 1, \dots, k_1 + \dots + k_i\}\}.$$

Note that for some i we can have $X_i = \emptyset$, but in any case $\sum_{i=1}^n |X_i| = m$.

We claim that $X = \bigcup_{i=1}^n X_i$ is a general position set of Z . Note first that X_i , $i \in [n]$, induces a complete graph K_{k_i} of Z . Therefore, in view of Theorem 2.3, to prove that X is a general position set, it suffices to demonstrate that if $(j, x_i) \in X_i$ and $(j', x_{i'}) \in X_{i'}$, where $i \neq i'$, then $d_Z((j, x_i), (j', x_{i'})) = 3$. Since (j, x_i) and $(j', x_{i'})$ are not adjacent, suppose on the contrary that $d_Z((j, x_i), (j', x_{i'})) = 2$. Then there exists a vertex (p, x_q) , where $p \neq j, j'$, such that $(j, x_i)(p, x_q) \in E(Z)$ and $(j', x_{i'})(p, x_q) \in E(Z)$. Then it follows that $f(j) = x_q$ and $f(j') = x_q$, contradicting our assumption. This contradiction proves the claim, hence $\text{gp}(Z) \geq m$.

Again, the above arguments hold true for any function $f \in K_n^{K_m}$, hence we have $\underline{\text{gps}}(K_m, K_n) \geq \max\{2(n-1), m\}$. \square

For the lower Sierpiński general position number of two complete graphs we have the following result, where we need to assume that the second factor is relatively large with respect to the first factor.

Theorem 5.3 *If $m \geq 2$ and $n \geq 2m - 2$, then*

$$\underline{\text{gps}}(K_m, K_n) = m(n - m + 1).$$

Proof. Using Lemma 5.2(i) once more we have $\underline{\text{gps}}(K_m, K_n) \geq m(n - m + 1)$.

To prove that $\underline{\text{gps}}(K_m, K_n) = m(n - m + 1)$, consider the function $f \in K_n^{K_m}$ defined by $f(i) = x_i$ for $i \in [m]$. Setting $G = K_m \otimes_f K_n$ we are going to show that $\text{gp}(G) \leq m(n - m + 1)$.

Suppose on the contrary that $\text{gp}(G) \geq m(n - m + 1) + 1$ and let R be a gp-set of G . Then it follows that there exists an index $i \in [m]$ such that $|V(iK_n) \cap R| \geq n - m + 2 \geq 2$. Assume, without loss of generality, that $|V(1K_n) \cap R|$ is as large as possible.

We claim that if $(1, x_j) \in V(1K_n) \cap R$, then $V(jK_n) \cap R = \emptyset$, $j \in [m]$. Indeed, since $|V(1K_n) \cap R| \geq 2$, there exists a vertex $(1, x_{j'}) \in R$, where $j' \neq j$ and $j' \in [m]$. Then the vertices $(1, x_{j'})$, $(1, x_j)$, (j, x_1) , and $(j, x_p) \in V(jK_n)$ form a shortest path, hence we conclude that $(j, x_p) \notin R$ for $p \in [n]$.

As we have assumed that $\text{gp}(G) \geq m(n-m+1)+1$, we can write $|V(1K_n) \cap R| = n - m + 1 + k$, where $k \geq 1$ (and $k \leq m - 1$). Then $|V(sK_n) \cap R| \leq n - m + 1 + k$, where $2 \leq s \leq m$. By the above, G contains at least k copies iK_n such that $V(iK_n) \cap R = \emptyset$. Then we have

$$\begin{aligned} |R| &\leq (n - m + 1 + k) + (m - k - 1)(n - m + 1 + k) \\ &= m(n - m + 1) + k(2m - n - 1 - k) \\ &< m(n - m + 1) + 1. \end{aligned}$$

Here the last inequality holds because $n \geq 2m - 2$ and $k \geq 1$ which implies that $2m - n - 1 - k \leq 2 - 1 - k = 1 - k \leq 0$. This contradiction implies that $\text{gp}(G) \leq m(n - m + 1)$. We conclude that $\underline{\text{gps}}(K_m, K_n) = m(n - m + 1)$. \square

In Theorem 5.3 we have assumed that $n \geq 2m - 2$. The following result explains why this assumption cannot be avoided in general.

Proposition 5.4 $\underline{\text{gps}}(K_6, K_9) = 25$.

Proof. By Lemma 5.2(i), we have $\underline{\text{gps}}(K_6, K_9) \geq 24$. To prove that $\underline{\text{gps}}(K_6, K_9) = 25$, we consider two cases depending on the function $f \in K_9^{K_6}$.

Case 1. f is injective.

In this case we may assume, without loss of generality, that $f(i) = x_i$ for $i \in [6]$. The obtained graph $K_6 \otimes_f K_9$ is shown in Fig. 3, where the edges of complete subgraphs iK_9 , $i \in [6]$, and the edges of the factors K_6 and K_9 are not drawn for a clearer picture.

We can check that the set consisting of black vertices from Fig. 3 is a general position set of cardinality 25 of $K_6 \otimes_f K_9$. Hence $\text{gp}(K_6 \otimes_f K_9) \geq 25$. To prove the reverse inequality, suppose there exists a general position set X of $K_6 \otimes_f K_9$ with $|X| \geq 26$. Let $X_i = X \cap V(iK_9)$, $i \in [6]$. We may assume without loss of generality that $|X_1| \geq |X_i|$ for $i \in \{2, \dots, 6\}$. Clearly, $|X_1| \geq 5$. If $|X_1| = 5$, then for some $i \in \{2, \dots, 6\}$ we have $X_i = \emptyset$. This implies that $|X| \leq 25$. Hence we must have $|X_1| = 4 + k$, where $2 \leq k \leq 5$. Then there are k copies iK_9 with no vertex from X . It follows that $|X| \leq (4 + k) + (6 - 1 - k)(4 + k) \leq 25$. We can conclude that $\text{gp}(K_6 \otimes_f K_9) \leq 25$ and thus $\text{gp}(K_6 \otimes_f K_9) = 25$.

Case 2. f is not injective.

In this case we may assume, without loss of generality, that $f(1) = f(2)$. Then in each of the copies iK_9 , $i \in \{3, 4, 5, 6\}$, the graph $K_6 \otimes_f K_9$ contains at least five vertices of degree 8. Since in each of $1K_9$ and $2K_9$ there are at least four such vertices, we can see (using the argument from the proof of Lemma 5.2(i)) that $\text{gp}(K_6 \otimes_f K_9) \geq 8 + 20 = 28$. \square




Figure 3: The Sierpiński product graph $K_6 \otimes_f K_9$, where $f(i) = x_i$ for $i \in [6]$

6 Concluding remarks

In Proposition 3.3 we have bounded from below the general position number of graphs with bridges. It would be interesting to characterize the graphs that attain the equality in the proved bound.

In view of Theorem 4.2 it would be interesting to characterize the graphs G with $\text{gp}(G) = \xi_M(G)$. Such graphs are, for instance, grid graphs $P_n \square P_m$, $n, m \geq 3$, for which we have $\text{gp}(P_n \square P_m) = 4 = \xi_M(P_n \square P_m)$. Similarly, in view of Theorem 4.4 it would be interesting to study the graphs G with $\xi_m(G) = \xi_M(G)$.

In Theorem 5.3 we have proved that $\underline{\text{gps}}(K_m, K_n) = m(n - m + 1)$ if $m \geq 2$ and $n \geq 2m - 2$. It remains as an open problem to determine $\underline{\text{gps}}(K_m, K_n)$ for the cases when $n < 2m - 2$. In Proposition 5.4 the particular case of $\underline{\text{gps}}(K_6, K_9)$ has been solved.

Acknowledgements

This work was supported by the Slovenian Research Agency ARIS (research core funding P1-0297 and projects N1-0285, N1-0218).

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Our manuscript has no associated data.

References

- [1] B.S. Anand, U. Chandran S.V., M. Changat, S. Klavžar, E.J. Thomas, Characterization of general position sets and its applications to cographs and bipartite graphs, *Appl. Math. Comput.* 359 (2019) 84–89.
- [2] M.C. Cai, A remark on the number of vertices of degree k in a minimally k -edge-connected graph, *Discrete Math.* 104 (1992) 221–226.
- [3] U. Chandran S.V., G.J. Parthasarathy, The geodesic irredundant sets in graphs, *Int. J. Math. Combin.* 4 (2016) 135–143.
- [4] M.A. Henning, S. Klavžar, E. Kleszcz, M. Pilśniak, The Sierpiński domination number, *Ars Math. Contemp.* (2023) doi.org/10.26493/1855-3974.2892.f07.
- [5] M.A. Henning, S. Klavžar, I.G. Yero, Resolvability and convexity properties in the Sierpiński product of graphs, *Mediterr. J. Math.* 21(1) (2024) Paper No. 3.
- [6] A.M. Hinz, S. Klavžar, S.S. Zemljich, A survey and classification of Sierpiński-type graphs, *Discrete Appl. Math.* 217 (2017) 565–600.
- [7] V. Iršič, S. Klavžar, G. Rus, J. Tuite, General position polynomials, *Results Math.* 79 (2024) Paper No. 110.
- [8] S. Klavžar, D. Kuziak, I. Peterin, I.G. Yero, A Steiner general position problem in graph theory, *Comput. App. Math.* 40 (2021) Paper No. 223.
- [9] S. Klavžar, U. Milutinović, Graphs $S(n, k)$ and a variant of the Tower of Hanoi problem, *Czechoslovak Math. J.* 47(122) (1997) 95–104.

- [10] S. Klavžar, D.F. Rall, I.G. Yero, General d -position sets, *Ars. Math. Contemp.* 21 (2021) Paper #P1.03.
- [11] S. Klavžar, G. Rus, The general position number of integer lattices, *Appl. Math. Comput.* 390 (2021) Paper No. 125664.
- [12] S. Klavžar, E. Tan, Edge general position sets in Fibonacci and Lucas cubes, *Bull. Malays. Math. Sci. Soc.* 46 (2023) Paper No. 120.
- [13] D. Korže, A. Vesel, General position sets in two families of Cartesian product graphs, *Mediterr. J. Math.* 20 (2023) Paper No. 203.
- [14] J. Körner, On the extremal combinatorics of the Hamming space, *J. Comb. Theory Ser. A* 71 (1995) 112–126.
- [15] J. Kovič, T. Pisanski, S.S. Zemljič, A. Žitnik, The Sierpiński product of graphs, *Ars Math. Contemp.* 23(1) (2023) Paper No. 1.
- [16] P. Manuel, S. Klavžar, A general position problem in graph theory, *Bull. Aust. Math. Soc.* 98 (2018) 177–187.
- [17] P. Manuel, R. Prabha, S. Klavžar, The edge general position problem, *Bull. Malays. Math. Sci. Soc.* 45 (2022) 2997–3009.
- [18] M.K. Menon, M.R. Chithra, K.S. Savitha, Security in Sierpiński graphs, *Discrete Appl. Math.* 328 (2023) 10–15.
- [19] B. Patkós, On the general position problem on Kneser graphs, *Ars Math. Contemp.* 18 (2020) 273–280.
- [20] M.A.D.R. Palma, A.J. León, S. Dantas, On the AVDTC of Sierpiński-type graphs, *Discrete Appl. Math.* 346 (2024) 10–29.
- [21] E.J. Thomas, U. Chandran S.V., J. Tuite, G. Di Stefano, On monophonic position sets in graphs, *Discrete Appl. Math.* (2023) doi.org/10.1016/j.dam.2023.02.021.
- [22] J. Tian, S. Klavžar, E. Tan, Extremal edge general position sets in some graphs, *Graph Combin.* 40 (2024) Paper No. 40.
- [23] J. Tian, K. Xu, The general position number of Cartesian products involving a factor with small diameter, *Appl. Math. Comput.* 403 (2021) Paper No. 126206.

- [24] J. Tian, K. Xu, D. Chao, On the general position numbers of maximal outer-plane graphs, *Bull. Malays. Math. Sci. Soc.* 46 (2023) Paper No. 198.
- [25] J. Varghese, V. Anu, A.S. Lakshmanan, Domination parameters of generalized Sierpiński graphs, *AKCE Int. J. Graphs Comb.* 21 (2024) 4–10.
- [26] Y. Yao, M. He, S. Ji, On the general position number of two classes of graphs, *Open Math.* 20 (2022) 1021–1029.