# A GEOMETRIC STUDY OF BZ OPERATOR ON REPRESENTATIONS OF $GL_n$ OVER NON-ARCHIMEDEAN FIELD

## TAIWANG DENG

ABSTRACT. In this article, we geometrically study the partial Bernstein-Zelevinsky operator introduced in the author's thesis, which generalizes the original Bernstein-Zelevinsky operator. We relate the partial Bernstein-Zelevinsky operator to the geometric induction of Lusztig and then perform explicit computations in special cases. Finally, we develop a symmetric reduction to the previously mentioned special cases

# CONTENTS

1.	Introduction	1
2.	New Poset Structure on Multisegments	3
3.	Canonical Basis and Quantum Algebras	7
4.	Partial BZ operator and Poincaré's series	13
5.	A formula for Lusztig's product	16
6.	Multisegments of Grassmanian Type	28
7.	Grassmanian case	34
8.	Parabolic Case	36
9.	Calculation of Partial BZ operator	38
Ap	pendix A. Minimal Degree Terms in Partial BZ operator	44
Ref	ferences	46

## 1. INTRODUCTION

Zelevinsky [17] classifies the admissible irreducible representations of  $\operatorname{GL}_n(F)$  in terms of multi-segments. More precisely, given a multisegment **a**, one can attach to it an irreducible representation  $L_{\mathbf{a}}$ , described as the unique irreducible sub-representation in some standard representation  $\pi(\mathbf{a})$  constructed by parabolic induction. In other words, in the Grothendieck group  $\mathcal{R}$  of the category of admissible representations, we have

$$\pi(\mathbf{a}) = L_{\mathbf{a}} + \sum_{\mathbf{b} < \mathbf{a}} m(\mathbf{b}, \mathbf{a}) L_{\mathbf{b}}, \quad m(\mathbf{b}, \mathbf{a}) \in \mathbb{Z}_{\geq 0},$$

*Key words and phrases.* Parabolic induction, Graded nilpotent classes, Kazhdan-Lusztig polynomials, BZ operator, Symmetric reduction, Schubert varieties.

where "<" is suitable partial order imposed on the set of multi-segments. In [2, 4.5], Bernstein and Zelevinsky define an operator  $\mathscr{D}$  (We call it BZ operator) to be an algebra homomorphism

$$\mathscr{D}: \mathcal{R} \to \mathcal{R},$$

which plays a crucial role in Zelevinsky's classification theorem. We introduced a partial analogue of the BZ operator in [6, Definition 2.19] (this is part of the author's thesis [14]).

In this article, we study the problem of computing the partial BZ operator  $\mathscr{D}^k(L_{\mathbf{a}})$  of the irreducible representation  $L_{\mathbf{a}}$  attached to a multisegment  $\mathbf{a}$ . There are two motivations for this. The first is to use these computations to calculate the multiplicities in certain induced representation  $L_{\mathbf{a}} \times L_{\mathbf{b}}$  in a forthcoming work. The second is also related to forthcoming work, in which we study the relations between the category of equivariant perverse sheaves on quiver varieties (in the sense of [6, §4]) and the category of perverse sheaves on flag varieties base on ideas of [6, §4]. We understand that this is a dual version of [5]. We further want to relate the partial BZ operator to the translation functor in representation of real Lie groups based on work of the present paper.

Let us discuss in more details about the content of the paper. From Zelevinsky's classification theorem, we can write

$$L_{\mathbf{a}} = \sum_{\mathbf{b}} \tilde{m}_{\mathbf{b},\mathbf{a}} \pi(\mathbf{a}).$$

Thus, the task boils down to calculating:

$$\mathscr{D}^k(\pi(\mathbf{a})) = \sum_{\mathbf{b}} n_{\mathbf{b},\mathbf{a}} L_{\mathbf{b}}, \qquad n_{\mathbf{b},\mathbf{a}} \ge 0.$$

We first introduce a new poset structure  $\leq_k$  on the set of multisegments so that we show the equivalence between  $n_{\mathbf{b},\mathbf{a}} > 0$  and  $\mathbf{b} \leq_k \mathbf{a}$ , cf. Proposition 2.6.

The primary result of this paper is the interpretation of the coefficient  $n_{\mathbf{b},\mathbf{a}}$  as the value at q = 1 of some Poincaré series of the Lusztig product of two explicit perverse sheaves on the space of graded nilpotent classes, cf. Theorem 4.8. This relies crucially on the work of Lusztig, as discussed in section 3. Note that the category of equivariant perverse sheaves should be understood as the dual of the category of representations of  $\operatorname{GL}_n(F)$ . Given that the BZ operator is related to the Jacquet functor, on the dual side, it should relate to geometric induction.

In section 5, we compute these Lusztig products as the push forward by a projection  $\beta''$ , cf. Theorem 5.19, of some concrete perverse sheaf on some space of graded nilpotent classes. This is our second main results.

Section 6-7 are dedicated to a concrete study of the coefficient  $n(\mathbf{b}, \mathbf{a})$  in the Grassmanian case, from which we deduce some combinatorial formulae for it, cf. Proposition 7.6.

In section 8, we continue to study the case where the multisegments are of Grassmanian type. In this case the projection  $\beta''$  is simple (cf. Proposition 8.6), being the natural projection

$$GL_n/P \to GL_n/P'$$

with  $P \subseteq P'$  two parabolic subgroups. The constructions and proofs in this case are very close to the case of Grassmanian type.

Finally in the last section 9, we obtain a complete formula for  $\mathscr{D}^k(L_{\mathbf{a}})$  in the general case, cf. Corollary 9.16. It's worth noting that to achieve this, we require a reduction process to the parabolic cases (cf. Proposition 9.6) in the style of [6, §6.2]. This elucidates the extensive discussion thereof.

Acknowledgements This paper is part of my thesis at University Paris 13, which is funded by the program DIM of the region Ile de France. I would like to thank my advisor Pascal Boyer for his keen interest in this work and his continuing support and countless advice. It was rewritten during my stay as a postdoc at Max Planck Institute for Mathematics and Yau Mathematical Sciences Center, I thank their hospitality. In addition, I would like to thank Alberto Mínguez ,Vincent Sécherre, Yichao Tian and Bin Xu for their helpful discussions on the subject.

## 2. New Poset Structure on Multisegments

We keep the same notations as in [6, §2]. Recall that in [6, Definition 2.12] we have introduced a partial order  $\leq$  on the set of multisegments and the finite set

$$S(\mathbf{a}) = \{\mathbf{b} : \mathbf{b} \le \mathbf{a}\}$$

following [17, 7.1]. We also recall that in [6, Notation 2.15] the standard representation  $\pi(\mathbf{a})$  was attached to a multisegment  $\mathbf{a}$  and its decomposition in the Grothendieck group into irreducible ones was considered:

$$\pi(\mathbf{a}) = \sum_{\mathbf{b} \in S(\mathbf{a})} m(\mathbf{b}, \mathbf{a}) L_{\mathbf{b}}, \quad m(\mathbf{b}, \mathbf{a}) \ge 0$$
(1)

where  $L_{\mathbf{a}}$  denotes the irreducible representation attached to  $\mathbf{a}$ . Here we have taken the cuspidal representation  $\rho = 1$  to be the trivial representation of  $GL_1$ , an assumption that we will make throughout this paper. In this section we define a new poset structure  $\leq_k$  depending on an integer k on the set of multisegments and show that the term  $L_{\mathbf{b}}$ appears in  $\mathscr{D}^k(\pi(\mathbf{a}))$  (cf. [6, Definition 2.19]) if and only if  $\mathbf{b} \leq_k \mathbf{a}$ . Let  $b(\Delta)$  (resp.  $e(\Delta)$ ) denote the beginning (resp. the end) of a segment  $\Delta$ . We also use the notation

$$b(\mathbf{a})(\text{ resp. } e(\mathbf{a}))$$
 (2)

to denote the multi-set of beginnings (resp. ends) in a multisegment **a**. If  $\Delta = [i, j]$  we set  $\Delta^- = [i, j - 1]$ .

**Definition 2.1.** Let  $k \in \mathbb{Z}$  and  $\Delta$  be a segment, we define

$$\Delta^{(k)} = \begin{cases} \Delta^-, & \text{if } e(\Delta) = k; \\ \Delta, & \text{otherwise} \end{cases}$$

For a multisegment  $\mathbf{a} = \{\Delta_1, \cdots, \Delta_r\}$ , we define

$$\mathbf{a}^{(k)} = \{\Delta_1^{(k)}, \cdots, \Delta_r^{(k)}\}.$$

**Definition 2.2.** For a well ordered multisegment  $\mathbf{a} = \{\Delta_1, \dots, \Delta_s\}$ with  $\Delta_1 \leq \dots \leq \Delta_s$ , let

$$\mathbf{a}(k) := \{\Delta \in \mathbf{a} : e(\Delta) = k\} = \{\Delta_{i_0}, \Delta_{i_0+1}, \cdots, \Delta_{i_1}\}.$$

Now let  $\Gamma \subseteq \mathbf{a}(k)$ , let

$$\mathbf{a}(k)_{\Gamma} := (\mathbf{a}(k) \setminus \Gamma) \cup \{\Delta^{(k)} : \Delta \in \Gamma\},\$$

and

$$\mathbf{a}_{\Gamma} := (\mathbf{a} \setminus \mathbf{a}(k)) \cup \mathbf{a}(k)_{\Gamma}$$

We say  $\mathbf{b} \preceq_k \mathbf{a}$  if there exist a multisegment  $\mathbf{c} \in S(\mathbf{a})$  such that

 $\mathbf{b} \leq \mathbf{a}_{\Gamma}(cf. [6, \text{Definition } 2.12])$ 

for some  $\Gamma$ .

Lemma 2.3. We have

$$\mathscr{D}^{k}(\pi(\mathbf{a})) = \pi(\mathbf{a}) + \sum_{\Gamma \subseteq \mathbf{a}(k), \Gamma \neq \emptyset} \pi(\mathbf{a}_{\Gamma}).$$
(3)

*Proof.* Let

$$\mathbf{a} = \{\Delta_1, \cdots, \Delta_r, \Delta_{r+1}, \cdots, \}.$$

Then

$$\pi(\mathbf{a}) = \prod_{i=1}^{r} L_{\Delta_i} \times \prod_{i>r} L_{\Delta_i}$$

and

$$\mathscr{D}^{k}(\pi(\mathbf{a})) = \prod_{i=1}^{r} (L_{\Delta_{i}} + L_{\Delta_{i}^{(k)}}) \times \prod_{i>r} L_{\Delta_{i}}$$
$$= \pi(\mathbf{a}) + \sum_{\Gamma \subseteq \mathbf{a}(k), \Gamma \neq \emptyset} \pi(\mathbf{a}_{\Gamma}).$$

	_

Recall that in [6, Notation 2.16], we use the notation  $\mathcal{R}$  to denote the Grothendieck group of all finite length unipotent representations of  $GL_n(F)$  with n = 0, 1, 2, ... for a non-archimedean local field F. Note that  $\mathcal{R}$  is a polynomial algebra endowed with a natural Hopf algebra structure ([6, Proposition 2.17 and Corollary 2.18]).

**Lemma 2.5.** Let  $\mathbf{b} \in S(\mathbf{a})$ , then  $\pi(\mathbf{a}) - \pi(\mathbf{b}) \ge 0$  in  $\mathcal{R}$ .

*Proof.* By choosing a maximal chain of multisegments between  $\mathbf{a}$  and  $\mathbf{b}$ , we can assume that

$$\mathbf{a} = \{\Delta_1, \cdots, \Delta_r\},\$$
$$\mathbf{b} = (\mathbf{a} \setminus \{\Delta_j, \Delta_k\}) \cup \{\Delta_j \cap \Delta_k, \Delta_j \cup \Delta_k\}.$$

Then by [15, section 4.6],

$$\pi(\mathbf{a}) = \pi(\mathbf{b}) + L_{\Delta_1} \times \cdots \times \widehat{L}_{\Delta_j} \times \cdots \times \widehat{L}_{\Delta_k} \times \cdots \times L_{\Delta_r} \times L_{\{\Delta_j, \Delta_k\}}.$$

# Proposition 2.6. Let

$$\mathscr{D}^{k}(\pi(\mathbf{a})) = \sum_{\mathbf{b}} n_{\mathbf{b},\mathbf{a}} L_{\mathbf{b}}.$$
(4)

Then  $n_{\mathbf{b},\mathbf{a}} > 0$  if and only if  $\mathbf{b} \leq_k \mathbf{a}$ .

Proof. By [6, Theorem 2.22] the coefficients  $n_{\mathbf{b},\mathbf{a}}$  in (4) are all nonnegative (but notice that the  $n_{\mathbf{b},\mathbf{a}}$  are not the  $n_{\mathbf{b},\mathbf{a}}$  in loc.cit.). Let  $\mathbf{b} \leq_k \mathbf{a}$ , then by definition we have  $\mathbf{b} \leq \mathbf{a}_{\Gamma}$  for some  $\Gamma$ . Therefore  $m_{\mathbf{b},\mathbf{a}_{\Gamma}} > 0$  (cf. (1)), now we have  $n_{\mathbf{b},\mathbf{a}} > 0$  by (3). Conversely, if  $n_{\mathbf{b},\mathbf{a}} > 0$ , then by (3),  $\mathbf{b} \leq \mathbf{a}_{\Gamma}$  for some  $\Gamma$ .

**Corollary 2.8.** We have  $\mathbf{b} \preceq_k \mathbf{a}$  if and only if  $\mathscr{D}^k(\pi(\mathbf{a})) - \pi(\mathbf{b}) \ge 0$ in  $\mathcal{R}$ .

Proof. By Proposition 2.6  $\mathbf{b} \leq_k \mathbf{a}$  implies  $\mathbf{b} \leq \mathbf{a}_{\Gamma}$  for some  $\Gamma \subseteq \mathbf{a}(k)$ . By Lemma 2.5  $\mathbf{b} \leq \mathbf{a}_{\Gamma}$  implies that  $\pi(\mathbf{a}_{\Gamma}) - \pi(\mathbf{b}) \geq 0$  in  $\mathcal{R}$ . Since  $\mathscr{D}^k(\pi(\mathbf{a})) - \pi(\mathbf{a}_{\Gamma}) \geq 0$  by (3), we have  $\mathscr{D}^k(\pi(\mathbf{a})) - \pi(\mathbf{b}) \geq 0$ . Conversely, if  $\mathscr{D}^k(\pi(\mathbf{a})) - \pi(\mathbf{b}) \geq 0$ , we have  $n(\mathbf{b}, \mathbf{a}) > 0$ , hence  $\mathbf{b} \leq_k \mathbf{a}$  by Proposition 2.6.

**Definition 2.9.** We define for  $\mathbf{b} \leq \mathbf{a}$ ,

$$\ell(\mathbf{b}, \mathbf{a}) = \max_{n} \{ n : \mathbf{a} = \mathbf{b}_0 \ge \mathbf{b}_1 \dots \ge \mathbf{b}_n = \mathbf{b} \},\$$

and  $\ell(a) = \ell(\mathbf{a}_{\min}, \mathbf{a}).$ 

**Proposition 2.10.** For any  $\mathbf{b} \leq_k \mathbf{a}$ , there exists  $\mathbf{c} \in S(\mathbf{a})$ , and some subset  $\Gamma \subseteq \mathbf{c}(k)$ , such that

$$\mathbf{b} = \mathbf{c}_{\Gamma}.$$

Conversely, if  $\mathbf{b} = \mathbf{c}_{\Gamma}$  for some  $\mathbf{c} \in S(\mathbf{a})$ , then  $\mathbf{b} \preceq_k \mathbf{a}$ .

*Proof.* For the converse part, suppose  $\mathbf{c} \neq \mathbf{a}$ , by (3), we have  $\mathscr{D}^k(\pi(\mathbf{c})) - \pi(\mathbf{b}) \geq 0$  in  $\mathcal{R}$ . By Lemme 2.5, we know that  $\pi(\mathbf{a}) - \pi(\mathbf{c}) \geq 0$  in  $\mathcal{R}$ , hence  $\mathscr{D}^k(\pi(\mathbf{a})) - \mathscr{D}^k(\pi(\mathbf{c})) \geq 0$  by [6, Theorem 2.22]. Therefore  $n_{\mathbf{b},\mathbf{a}} > 0$ . Hence we have  $\mathbf{b} \leq_k \mathbf{a}$ .

For the direct part, suppose that  $\mathbf{b} \leq_k \mathbf{a}$ , hence  $\mathbf{b} < \mathbf{a}_{\Gamma_1}$  for some  $\Gamma_1$ . We prove by induction on  $\ell(\mathbf{b}, \mathbf{a}_T)$ . If  $\ell(\mathbf{b}, \mathbf{a}_{\Gamma_1}) = 0$ , then  $\mathbf{b} = \mathbf{a}_{\Gamma_1}$ , we are done. Now let  $\mathbf{b} < \mathbf{d} \leq \mathbf{a}_{\Gamma_1}$  such that  $\ell(\mathbf{b}, \mathbf{d}) = 1$ , by induction,

$$\mathbf{d} = \mathbf{c}'_{\Gamma_0},$$

for some  $\mathbf{c}' \in S(\mathbf{a})$ . Note that by replacing  $\mathbf{c}'$  by  $\mathbf{a}$ , we can assume that  $\mathbf{d} = \mathbf{a}_{\Gamma_1}$  and  $\ell(\mathbf{b}, \mathbf{a}_{\Gamma_1}) = 1$ .

By definition **b** is obtained by applying the elementary operation to a pair of segments  $\{\Delta \leq \Delta'\}$  in  $\mathbf{a}_T$ . Now we set out to construct **c**.

• If  $\{\Delta, \Delta'\} \subseteq \mathbf{a} \setminus \{\Delta^{(k)} : \Delta \in \Gamma_1\} \subseteq \mathbf{a}$ , let **c** be the multisegment obtained by applying the elementary operations to  $\{\Delta, \Delta'\}$ . And we have

$$\mathbf{b} = \mathbf{c}_{\Gamma_1}$$

If {Δ, Δ'} ∩ {Δ<sup>(k)</sup> : Δ ∈ Γ<sub>1</sub>} = {Δ'}, then {Δ, Δ'<sup>+</sup>} ∈ a let
c be the multisegment obtained by applying the elementary operations to {Δ, Δ'<sup>+</sup>}. Then let

$$\Gamma = (\Gamma_1 \setminus \{\Delta'^+\}) \cup \{\Delta \cup \Delta'^+\}$$

and we have

$$\mathbf{b} = \mathbf{c}_{\Gamma}.$$

• If  $\{\Delta, \Delta'\} \cap \{\Delta^{(k)} : \Delta \in \Gamma_1\} = \{\Delta\}$ , then  $\{\Delta^+, \Delta'\} \in \mathbf{a}$  let **c** be the multisegment obtained by applying the elementary operations to  $\{\Delta^+, \Delta'\}$ . Then let

$$\Gamma = (\Gamma_1 \setminus \{\Delta^+\}) \cup \{\Delta \cap \Delta'\}$$

and we have

$$\mathbf{b} = \mathbf{c}_{\Gamma}$$
.

Hence we are done.

We use the notation  $\mathcal{O}$  to denote the set of multisegments with support in  $\mathbb{Z}(\text{cf. }[6, \text{ Definition } 2.14]).$ 

# **Proposition 2.11.** The relation $\leq_k$ defines a poset structure on $\mathcal{O}$ .

*Proof.* By definition we have  $\mathbf{a} \leq_k \mathbf{a}$  for any  $\mathbf{a} \in \mathcal{O}$ . Suppose  $\mathbf{a}_1 \leq_k \mathbf{a}_2, \mathbf{a}_2 \leq_k \mathbf{a}_3$ , we want to show that  $\mathbf{a}_1 \leq_k \mathbf{a}_3$ . By proposition 2.10, there exists  $\mathbf{c} \in S(\mathbf{a}_2)$  and  $\Gamma_1 \subseteq \mathbf{c}(k)$ , such that

$$\mathbf{a}_1 = \mathbf{c}_{\Gamma_1}.$$

Note that by Corollary 2.8, the fact  $\mathbf{a}_2 \leq_k \mathbf{a}_3$  implies  $\mathscr{D}^k(\pi(\mathbf{a}_3)) - \pi(\mathbf{a}_2) \geq 0$ . Hence we have  $n(\mathbf{a}_3, \mathbf{c}) > 0$ , therefore  $\mathbf{c} \leq_k \mathbf{a}_3$  by Proposition 2.6. In turn there exists a multisegment  $\mathbf{c}' \in S(\mathbf{a}_3)$  and  $\Gamma_2 \subseteq \mathbf{c}'(k)$ , such that

$$\mathbf{c} = \mathbf{c}'_{\Gamma_2}$$

Since we have  $\mathbf{c}(k) \subseteq \mathbf{c}'(k)$ , we take

$$\Gamma_3 := \Gamma_1 \cup \Gamma_2 \subseteq \mathbf{c}'(k).$$

Now we get

$$\mathbf{a}_1 = \mathbf{c}'_{\Gamma_3}$$

which implies  $\mathbf{a}_1 \leq_k \mathbf{a}_3$  by Proposition 2.10. Finally, if  $\mathbf{a} \leq_k \mathbf{b}$  and  $\mathbf{b} \leq_k \mathbf{a}$ , then by definition we have  $\mathbf{a} = \mathbf{b}$ .

# **Definition 2.12.** We let

$$\Gamma(\mathbf{a},k) = \{\mathbf{b} : \mathbf{b} \preceq_k \mathbf{a}\}.$$

# 3. CANONICAL BASIS AND QUANTUM ALGEBRAS

In this section, we recall the results of Lusztig [12] on canonical basis while keeping the notations of [6] and draw the connection to the algebra  $\mathcal{R}$  base on [10]. No new results are contained in this section.

**Definition 3.1.** Let  $\mathbb{N}^{(\mathbb{Z})}$  be the semi-group of sequences  $(d_j)_{j \in \mathbb{Z}}$  of non negative integers which are zero for all but finitely many j. Let  $\alpha_i$  be the element whose *i*-th term is 1 and other terms are zero.

**Definition 3.2.** We define a symmetric bilinear form on  $\mathbb{N}^{(\mathbb{Z})}$  given by

$$(\alpha_i, \alpha_j) = \begin{cases} 2, & \text{for } i = j; \\ -1, & \text{for } |i - j| = 1; \\ 0, & \text{otherwise} \end{cases}$$

**Definition 3.3.** Let q be an indeterminate and  $\mathbb{Q}(q^{1/2})$  be the fractional field of  $\mathbb{Z}[q^{1/2}]$ . Let  $U_q^{\geq 0}$  be the  $Q(q^{1/2})$ -algebra generated by the elements  $E_i$  and  $K_i^{\pm 1}$  for  $i \in \mathbb{Z}$  with the following relations:

$$K_i K_j = K_j K_i, \ K_i K_i^{-1} = 1;$$
  

$$K_i E_i = q^{1/2(\alpha_i, \alpha_j)} E_i K_i;$$
  

$$E_i E_j = E_j E_i, \ if \ |i - j| > 1;$$
  

$$E_i^2 E_j - (q^{1/2} + q^{-1/2}) E_i E_j E_i + E_j E_i^2 = 0, \ if \ |i - j| = 1.$$

and let  $U^+$  be the subalgebra generated by the  $E_i$ 's.

*Remark:* This is the + part of the quantized enveloping algebra U associated by Drinfeld and Jimbo to the root system  $A_{\infty}$  of  $SL_{\infty}$ . And for q = 1, this specializes to the classical enveloping algebra of the nilpotent radical of a Borel subalgebra.

**Definition 3.4.** We define a new order on the set of segments  $\Sigma$ 

$$\begin{bmatrix} [j,k] \triangleleft [m,n], & \text{if } k < n, \\ [j,k] \triangleright [m,n], & \text{if } j < m, n = k. \end{bmatrix}$$

We also denote  $[j,k] \triangleleft [m,n]$  or [j,k] = [m,n] by  $\trianglelefteq [m,n]$ .

*Remark:* This should be compared with [6, Definition 2.5].

**Lemma 3.5.** The algebra  $U_q^+$  is  $\mathbb{N}^{(\mathbb{Z})}$ -graded via the weight function  $\operatorname{wt}(E_i) = \alpha_i$ . Moreover, for a given weight  $\alpha$ , the homogeneous component of  $U_q^+$  with weight  $\alpha$  is of finite dimension, and its basis are naturally parametrized by the multisegments of the same weight.

*Proof.* Let  $\mathbf{a} = \sum_{s=1}^{r} m_{i_s, j_s}[i_s, j_s]$  be a multisegment of weight  $\alpha$  (cf. [6, Definition 3.1]), note that here we identify the weight  $\varphi_{[i]}$  with  $\alpha_i$ , and that

 $[i_1, j_1] \leq \cdots \leq [i_r, j_r]$  (cf. Definition 3.4)

Then we associate to  $\mathbf{a}$  the element

$$(E_{j_1}\cdots E_{i_1})\cdots (E_{j_r}\cdots E_{i_r}).$$

In this way we obtain the desired parametrization.

Notation 3.6. For  $x \in U^+$  be an element of degree  $\alpha$ , we will denote  $\operatorname{wt}(x) = \alpha$ .

**Example 3.7.** For  $i \leq j$ , let  $\alpha_{ij} = \alpha_i + \cdots + \alpha_j$ . Consider the homogeneous components of  $U^+$  with weight  $\alpha = 2\alpha_{12}$ , whose basis is given by

$$E_1 E_2 E_1 E_2, E_1 E_1 E_2 E_2.$$

The element  $E_1E_2E_1E_2$  is parametrized by the multisegment [1]+[1,2]+[2], while  $E_1E_1E_2E_2$  is parametrized by the multisegment 2[1]+2[2].

In [12], Lusztig has defined certain bases for  $U_q^+$  associated to the orientations of a Dynkin diagram, called PBW( Poincaré-Birkhoff-Witt) basis, which specializes to the classical PBW type bases. Following [10], we describe the PBW-basis as follows.

# Definition 3.8. We define

 $E([i]) = E_i, \ E([i, j]) = [E_j[\cdots [E_{i+1}, E_i]_{q^{1/2}} \cdots ]_{q^{1/2}}]_{q^{1/2}},$ where  $[x, y]_{q^{1/2}} = xy - q^{-1/2(\text{wt}(x), \text{wt}(y))}yx.$  More generally, let  $\mathbf{a} = \sum_s a_{i_s, j_s}[i_s, j_s]$  be a multisegment, such that  $[i_1, j_1] \lhd \cdots \lhd [i_r, j_r](\text{ cf. Definition 3.4}),$ 

we define

$$E(\mathbf{a}) = \frac{1}{\prod_{s} [a_{i_s, j_s}]_{q^{1/2}}!} E([i_1, j_1])^{a_{i_1, j_1}} \cdots E([i_r, j_r])^{a_{i_r, j_r}},$$

here  $[m]_{q^{1/2}} = \frac{q^{1/2m} - q^{-1/2m}}{q^{1/2} - q^{-1/2}}$  for  $m \in \mathbb{Z}$  and  $[m]_{q^{1/2}}! = [m]_{q^{1/2}}[m - 1]_{q^{1/2}} \cdots [2]_{q^{1/2}}.$ 

**Definition 3.9.** Let  $x \mapsto \overline{x}$  be the involution defined as the unique ring automorphism of  $U_q^+$  defined by

$$\overline{q^{1/2}} = q^{-1/2}, \ \overline{E_i} = E_i.$$

**Proposition 3.10.** (cf. [12, Theorem 8.10]) Let  $\mathcal{L} := \bigoplus_{\mathbf{a} \in \mathcal{O}} \mathbb{Z}[q^{1/2}] E(\mathbf{a}) \subseteq$ 

 $U_q^+$ . Then there exists a unique  $\mathbb{Q}(q^{1/2})$ -basis  $\{G(\mathbf{a}) : \mathbf{a} \in \mathcal{O}\}$  of  $U_q^+$  such that

$$G(\mathbf{a}) = G(\mathbf{a}), \ G(\mathbf{a}) = E(\mathbf{a}) \ modulo \ q^{1/2} \mathcal{L}.$$

This is called Lusztig's canonical basis.

Lusztig also gave a geometric description of his canonical basis in terms of the orbital varieties  $\overline{O}_{\mathbf{a}}$  [6, Proposition 3.3].

**Definition 3.11.** Let  $\mathbb{A}$  be the group ring of  $\overline{\mathbb{Q}}_{\ell}^*$  over  $\mathbb{Z}$ . Let  $\mathbf{K}_{\varphi}$  be the Grothendieck group over  $\mathbb{A}$  of the category of constructible  $G_{\varphi}$ equivariant  $\mathbb{Q}_{\ell}$  sheaves over  $E_{\varphi}$  (cf. [6, Definition 3.2]), considered as a variety over a finite field  $\mathbb{F}_q$ .

Given a weight function  $\varphi$  [6, Definition 3.2], we use  $S(\varphi)$  to denote the finite set of multisegments with weight  $\varphi$ . We also use S to denote the space of weight functions.

**Lemma 3.12.** (cf. [12, §9.4]) The A-module  $\mathbf{K}_{\varphi}$  admits a basis { $\gamma_{\mathbf{a}}$  :  $\mathbf{a} \in S(\varphi)$ } indexed by the  $G_{\varphi}$  orbits of  $E_{\varphi}$ , where  $\gamma_{\mathbf{a}}$  corresponds to the constant sheaf  $\overline{\mathbb{Q}}_{\ell}$  on the orbit  $O_{\mathbf{a}}$ , extending by 0 to the complement.

**Definition 3.13.** Let  $\varphi = \varphi_1 + \varphi_2 \in S$ . We define a diagram of varieties

$$E_{\varphi_1} \times E_{\varphi_2} \xleftarrow{\beta} E' \xrightarrow{\beta'} E'' \xrightarrow{\beta''} E_{\varphi},$$
 (5)

where

 $E'' := \{ (T, W) : W = \bigoplus W_i, \ W_i \subseteq V_{\varphi,i}, \ T(W_i) \subseteq W_{i+1}, \ \dim(W_i) = \varphi_2(i) \}, \\ E' := \{ (T, W, \mu, \mu') : (T, W) \in E'', \ \mu : W \simeq V_{\varphi_2}, \ \mu' : V_{\varphi}/W \simeq V_{\varphi_1} \}, \\ and$ 

 $\beta''((T,W)) = W, \ \beta'((T,W,\mu,\mu')) = (T,W), \ \beta((T,W,\mu,\mu')) = (T_1,T_2),$ such that

$$T_1 = \mu' \circ T \circ \mu'^{-1}, \ T_2 = \mu \circ T \circ \mu^{-1}$$

**Proposition 3.15.** (cf. [12, §6.1]) The group  $G_{\varphi} \times G_{\varphi_1} \times G_{\varphi_2}$  acts naturally on the varieties in the diagram (5) with  $G_{\varphi}$  acting trivially on  $E_{\varphi_1} \times E_{\varphi_2}$  and  $G_{\varphi_1} \times G_{\varphi_2}$  acting trivially on  $E_{\varphi}$ . And all the maps there are compatible with such actions. Moreover, we have

## TAIWANG DENG

- (1): The morphism  $\beta'$  is a principle  $G_{\varphi_1} \times G_{\varphi_2}$ -fibration.
- (2): The morphism  $\beta$  is a locally trivial trivial fibration with smooth connected fibers.
- (3): The morphism  $\beta''$  is proper.

**Example 3.16.** Let  $\varphi_1 = \chi_1$  (the singleton supported at the integer  $1 \in \mathcal{O}$ ) and  $\varphi_2 = \chi_2$ . Then  $\varphi = \chi_1 + \chi_2$  and

$$E_{\varphi_1} = E_{\varphi_2} = 0, \ E_{\varphi} = \overline{\mathbb{F}}_q.$$

Moreover, we have

$$E'' = \{(T, W) : W = V_{\varphi_2}, T \in \overline{\mathbb{F}}_q\} \simeq \overline{\mathbb{F}}_q,$$

and

$$E' = \{ (T, W, \mu, \mu') : (T, W) \in E'', \mu, \mu' \in \overline{\mathbb{F}}_q^{\times} \} \simeq \overline{\mathbb{F}}_q \times (\overline{\mathbb{F}}_q^{\times})^2.$$

**Corollary 3.17.** (cf. [12, §9.5]) Let  $\mathbf{a} \in S(\varphi_1), \mathbf{a}' \in S(\varphi_2)$  and  $\varphi = \varphi_1 + \varphi_2$  be a weight function. There exists a simple perverse sheaf(up to shift)  $\mathcal{P}$  on  $E_{\varphi}$  such that

$$\beta^*(IC(\overline{O}_{\mathbf{a}}) \otimes IC(\overline{O}_{\mathbf{a}'})) = \beta'^*(\mathcal{P}).$$

**Example 3.18.** As in Example 3.16, let  $\mathbf{a} = \{[1]\}, \mathbf{a}' = \{[2]\}, then$  $IC(\overline{O}_{\mathbf{a}}) = \overline{\mathbb{Q}}_{\ell}, IC(\overline{O}_{\mathbf{a}'}) = \overline{\mathbb{Q}}_{\ell}.$ 

Hence if we let

$$\mathcal{P} = \overline{\mathbb{Q}}_{\ell},$$

then

$$\beta^*(IC(\overline{O}_{\mathbf{a}}) \otimes IC(\overline{O}_{\mathbf{a}'})) = \beta''^*(\mathcal{P}).$$

**Definition 3.19.** We define a multiplication

$$IC(\overline{O}_{\mathbf{a}}) \star IC(\overline{O}_{\mathbf{a}'}) = \beta_*''(\mathcal{P}).$$

**Example 3.20.** As in the Example 3.16, we have

$$IC(O_{\mathbf{a}}) \star IC(O_{\mathbf{a}'}) = \beta''_{*}(\mathcal{P}) = IC(E_{\varphi}),$$

note that here  $\beta''$  is an isomorphism.

**Proposition 3.21.** (cf. [12, §9.5]) Let  $\mathbf{a} \in S(\varphi_1), \mathbf{a}' \in S(\varphi_2) \varphi = \varphi_1 + \varphi_2$  be a weight function. We associate to the intersection cohomology complex  $IC(\overline{O}_{\mathbf{a}})$ 

$$\tilde{\gamma}_{\mathbf{a}} = \sum_{\mathbf{b} \ge \mathbf{a}} p_{\mathbf{b}, \mathbf{a}}(q) \gamma_{\mathbf{b}},$$

where  $p_{\mathbf{b},\mathbf{a}}(q)$  is the formal alternative sum of eigenvalues of the Frobenius map on the stalks of the cohomology sheaves of  $IC(\overline{O}_{\mathbf{a}})$  at any  $\mathbb{F}_q$ rational point of  $O_{\mathbf{b}}$ . Moreover, the multiplication  $\star$  gives a  $\mathbb{A}$ -bilinear map

$$\mathbf{K}_{\varphi_1} imes \mathbf{K}_{\varphi_2} o \mathbf{K}_{\varphi},$$

which defines an associative algebra structure over  $\mathbf{K} = \bigoplus_{\varphi} \mathbf{K}_{\varphi}$ .

**Proposition 3.22.** (12, Proposition 9.8 and Theorem 9.13)

- The elements  $\gamma_i := \gamma_{[i]}$  for all  $i \in \mathbb{Z}$  generate the algebra  $\mathbf{K}$  over  $\mathbb{A}$ .
- Let  $U_{\mathbb{A}}^{\geq 0} = U_q^{\geq 0} \otimes_{\mathbb{Z}} \mathbb{A}$ . Then we have a unique  $\mathbb{A}$ -algebra morphism  $\Gamma : \mathbf{K} \to U_{\mathbb{A}}^{\geq 0}$  such that

$$\Gamma(\gamma_j) = K_j^{-j} E_j;$$

for all  $j \in \mathbb{Z}$ . Moreover, for  $\varphi \in S$ , let

$$d(\varphi) = \sum_{i \in \mathbb{Z}} (\varphi(i) - 1)\varphi(i)/2 - \sum_{i \in \mathbb{Z}} \varphi(i)\varphi(i+1).$$
(6)

Then there is an  $\mathbb{A}$ -linear map  $\Theta: K_{\varphi} \to U^+_{\mathbb{A}}$ , such that

$$\Gamma(\xi) = q^{1/2d(\varphi)} K(\varphi) \Theta(\xi),$$
  
-  $\prod K^{-i\varphi(i)}$ 

where  $K(\varphi) = \prod_{i \in \mathbb{Z}} K_i^{-i\varphi(i)}$ . • We have

$$\Gamma(\gamma_{\mathbf{c}}) = q^{1/2(r-\delta_{\mathbf{c}})} K(\varphi_{\mathbf{c}}) E(\mathbf{c}),$$

where

$$r = \sum_{i} \varphi_{\mathbf{c}}(i)(\varphi_{\mathbf{c}}(i) - 1)(2i - 1)/2 - \sum_{i} i\varphi_{\mathbf{c}}(i - 1)\varphi_{\mathbf{c}}(i),$$

and  $\delta_{\mathbf{c}}$  is the co-dimension of the orbit  $O_{\mathbf{c}}$  in  $E_{\varphi_{\mathbf{c}}}$ .

• We have

$$\Theta(\gamma_{\mathbf{a}}) = q^{1/2 \dim(O_{\mathbf{a}})} E(\mathbf{a}), \ \Theta(\tilde{\gamma}_{\mathbf{a}}) = q^{1/2 \dim(O_{\mathbf{a}})} G(\mathbf{a}).$$

Hence

$$G(\mathbf{a}) = \sum_{\mathbf{b} \ge \mathbf{a}} P_{\mathbf{b}, \mathbf{a}}(q) E(\mathbf{b}).$$

**Proposition 3.23.** [11, §4.3] The canonical basis of  $U_q^+$  are almost orthogonal with respect to a scalar product introduced by Kashiwara [8], which are given by

$$(E(\mathbf{a}), E(\mathbf{b})) = \frac{(1-q)^{\deg(\mathbf{a})}}{\prod_{i \le j} h_{a_{ij}}(q)} \delta_{\mathbf{a}, \mathbf{b}}$$

where  $\mathbf{a} = \sum_{i \leq j} a_{ij}[i, j], h_k(z) = (1-z) \cdots (1-z^k)$  and  $\delta$  is the Kronecker symbol. And we have

$$(G(\mathbf{a}), G(\mathbf{b})) = \delta_{\mathbf{a}, \mathbf{b}} \mod q^{1/2} \mathbb{A}.$$

**Notation 3.24.** We denote by  $\{E^*(\mathbf{a})\}$  and  $\{G^*(\mathbf{a})\}$  the dual basis of  $\{E(\mathbf{a})\}$  and  $\{G(\mathbf{a})\}$  with respect to the Kashiwara scalar product.

*Remark:* Note that  $\{G^*(\mathbf{a})\}$  is referred as the dual canonical basis.

**Proposition 3.25.** (cf. [10, §3.4]) Let  $\mathbf{a} = \sum_{i \leq j} a_{ij}[i, j]$ . Then

• We have

$$E^{*}(\mathbf{a}) = \frac{\prod_{i \le j} h_{a_{ij}}(q)}{(1-q)^{\deg(\mathbf{a})}} E(\mathbf{a}) = \prod_{ij}^{\longrightarrow} q^{1/2\binom{a_{ij}}{2}} E^{*}([i,j])^{a_{ij}},$$

here the product is taken with respect to the order  $\leq$ .

• *And* 

$$E^*(\mathbf{a}) = \sum_{\mathbf{b} \le \mathbf{a}} P_{\mathbf{a}, \mathbf{b}}(q) G^*(\mathbf{b}).$$

**Example 3.26.** Let  $\mathbf{a} = [1] + [2]$ . Then

$$E^*(\mathbf{a}) = E([1])E([2]) = E(\mathbf{a}),$$

and

$$\begin{split} G^*([1,2]) &= E^*([1,2]),\\ G^*(\mathbf{a}) &= E^*(\mathbf{a}) - q^{1/2} E^*([1,2]). \end{split}$$

Finally, we establish the relation of between the algebras  $\mathcal{R}$  and  $U^+$ .

**Definition 3.27.** Let B be the polynomial algebra generated by the set of coordinate functions  $\{t_{ij} : i < j\}$ . Following [10, §2.6], we write  $t_{ii} = 1, t_{ij} = 0$  if i > j, and index the non-trivial  $t_{i,j}$ 's by segments, namely,  $t_{[ij]} = t_{i,j-1}$  for i < j.

**Proposition 3.28.** [6, Corollary 2.18], We have an algebra isomorphism  $\phi : B \simeq \mathcal{R}$  by identifying  $t_{[ij]}$  with  $L_{[ij]}$  for all i < j.

**Definition 3.29.** ([1], [10, §3.5]) Let  $B_q$  be the quantum analogue of *B* generate by  $\{T_{ij} : i < j\}$ , where  $T_{ij}$  is considered as the q-analogue of  $t_{ij}$ . Also, we write  $T_{ii} = 1$  and  $T_{ij} = 0$  if i > j. And we will indexed the non-trivial  $T_{ij}$  by  $T_{[i,j-1]}$ . The generators  $T_s$ 's satisfies the following relations. Let s > s' be two segments. Then

$$T_{s'}T_s = \left\{ \begin{array}{ll} q^{-1/2(\text{wt}(s'),\text{wt}(s))}T_sT_{s'} + (q^{-1/2} - q^{1/2})T_{s\cap s'}T_{s\cup s'}, & \text{if $s$ and $s'$ are linked,} \\ q^{-1/2(\text{wt}(s'),\text{wt}(s))}T_sT_{s'}, & \text{otherwise} \ . \end{array} \right.$$

**Proposition 3.30.** [10, §3.5] There exist an algebra isomorphic morphism

$$\iota: U_q^+ \to B_q,$$

given by  $\iota(E^*([i,j])) = T_{[i,j]}$ . Moreover, for  $\mathbf{a} = \sum_{i \leq j} a_{ij}[i,j]$ , we have

$$\iota(E^*(\mathbf{a})) = \prod_{i \le j} q^{1/2\binom{a_{ij}}{2}} T^{a_{ij}}_{[i,j]},$$

here the multiplication is taken with respect to the order <.

**Example 3.31.** Let  $\mathbf{a} = [1] + [2]$ , then

$$u(E^*(\mathbf{a})) = T_{[1]}T_{[2]}.$$

**Proposition 3.32.** [10, Theorem 12] By specializing at q = 1, the dual canonical basis  $\{G^*(\mathbf{a}) : \mathbf{a} \in \mathcal{O}\}$  gives rise to a well defined basis for B, denoted by  $\{g^*(\mathbf{a}) : \mathbf{a} \in \mathcal{O}\}$ . Moreover, the morphism  $\phi$  (cf. Proposition 3.28) sends  $g^*(\mathbf{a})$  to  $L_{\mathbf{a}}$  for all  $\mathbf{a} \in \mathcal{O}$ .

# 4. PARTIAL BZ OPERATOR AND POINCARÉ'S SERIES

In this section we will deduce a geometric description for the partial BZ operator, using results of last section.

**Definition 4.1.** [9, §2.2] Kashiwara [8] introduced some q-derivations  $E'_i$  in  $\operatorname{End}(U^+_q)$  for all  $i \in \mathbb{Z}$  satisfying

$$E'_i(E_j) = \delta_{ij}, \ E'_i(uv) = E'_i(u)v + q^{-1/2(\alpha_i, \operatorname{wt}(u))}uE'_i(v).$$

**Example 4.2.** Simple calculation shows that

$$E'_{i}(E([j,k])) = \delta_{i,k}(1-q)E([j,k-1]),$$

by taking dual, we get

$$E'_{i}(E^{*}([j,k])) = \delta_{i,k}E^{*}([j,k-1]),$$

Proposition 4.3. We have

$$(E'_i(u), v) = (u, E_i v)$$

where (,) is the scalar product introduced in Proposition 3.28.

Note that by identifying the algebra  $U_q^+$  and  $B_q$  via  $\iota$ , we get a version of q-derivations in End $(B_q)$ .

**Definition 4.4.** By specializing at q = 1, the q derivation  $E'_i$  gives a derivation  $e'_i$  of the algebra B by

$$e'_i(t_{[jk]}) = \delta_{ik} t_{[j,k-1]}, \ e'_i(uv) = e'_i(u)v + ue'_i(v).$$

**Proposition 4.5.** Let

$$D^i := \sum_{n=0}^{\infty} \frac{1}{n!} e_i^{\prime n}.$$

Then the morphism  $D^i: B \to B$  is an algebraic morphism. Moreover, if we identify the algebras  $\mathcal{R}$  and B via  $\phi$  (cf. Proposition 3.28), then the morphism  $D^i$  coincides with the partial BZ operator  $\mathcal{D}^i$ .

*Proof.* For  $n \in \mathbb{N}$ , we have

$$e'^{n}(uv) = \sum_{r+s=n} \binom{n}{r} e'^{r}_{i}(u) e'^{s}_{i}(v),$$

therefore

$$D^{i}(uv) = \sum_{n=0}^{\infty} \frac{1}{n!} \sum_{r+s=n} \binom{n}{r} e_{i}^{\prime r}(u) e_{i}^{\prime s}(v) = D^{i}(u) D^{i}(v).$$

Finally, to show that  $D^i$  and  $\mathscr{D}^i$  coincides, it suffices to prove that

 $\phi \circ D^{i}(t_{[j,k]}) = \mathscr{D}^{i} \circ \phi(t_{[j,k]}),$ 

but we have

$$D^{i}(t_{[j,k]} = t_{[j,k]} + \delta_{i,k}t_{[j,k-1]},$$

and

$$\mathscr{D}^i(L_{[j,k]}) = L_{j,k} + \delta_{i,k} L_{[j,k-1]}.$$

Therefore, we have

$$\phi \circ D^{i}(t_{[j,k]}) = \mathscr{D}^{i} \circ \phi(t_{[j,k]}).$$

*Remark:* Without specializing at q = 1, the operator  $D^i$  is not an algebraic morphism. To get an algebraic morphism at the level of  $U_q^+$ , one should consider not only the summation of the iteration of  $e'_i$ 's but all the derivations, which gives rise to an embedding into the quantum shuffle algebras, cf. [9].

Next we show how to determine  $\mathscr{D}^i(L_{\mathbf{a}})$  by the algebra **K** of Lusztig.

# **Lemma 4.6.** Let $n \in \mathbb{N}$ , and $\mathbf{d} \in \mathcal{O}$ . Then we have

$$E_i^n G(\mathbf{b}) = \sum_{\mathbf{d}} (E_i^n G(\mathbf{b}), E^*(\mathbf{d})) E(\mathbf{d}) = \sum_{\mathbf{d}} (G(\mathbf{b}), E_i'^n E^*(\mathbf{d})) E(\mathbf{d}),$$

where (,) is the Kashiwara scalar product. Moreover, for each **b** such that  $(G(\mathbf{b}), E'_i E^*(\mathbf{d})) \neq 0$ , we have

$$\operatorname{wt}(\mathbf{d}) = \operatorname{wt}(\mathbf{b}) + n\alpha_i.$$

*Proof.* This is by definition.

**Corollary 4.7.** Let  $\mathbf{b} \preceq_k \mathbf{d}$  such that  $wt(\mathbf{d}) = wt(\mathbf{b}) + n\alpha_i$ . Then  $L_{\mathbf{b}}$  appears as a factor of  $\frac{1}{r!}e_i'^r(\pi(\mathbf{d}))$  if and only if r = n.

*Proof.* For each  $\mathbf{b} \leq_k \mathbf{d}$ , the representation  $L_{\mathbf{b}}$  is a factor of  $\mathscr{D}^i(\pi(\mathbf{d}))$ . Now by Proposition 4.5,  $\mathscr{D}^i = \sum_r \frac{1}{r!} e_i'^r$ , moreover, by Lemma 4.6,

factors of  $\frac{1}{r!}e_i'^r(\pi(\mathbf{d}))$  always have weight  $\operatorname{wt}(\mathbf{d}) - r\alpha_i$ . Therefore we are done.

**Theorem 4.8.** Let  $\mathbf{b} \preceq_k \mathbf{a}$ , then there exists  $\mathbf{c} \in S(\mathbf{a})$  such that  $\mathbf{c} = \mathbf{b} + \ell[k]$ . Then

$$n_{\mathbf{b},\mathbf{a}} = \sum_{i} \dim \mathcal{H}^{2i}(IC(\overline{O}_{\ell[k]}) \star IC(\overline{O}_{\mathbf{b}}))_{\mathbf{a}}.$$

*Proof.* Note that by Proposition 3.22, we have

$$\Gamma(\tilde{\gamma}_{\ell[k]} \star \tilde{\gamma}_{\mathbf{b}}) = \Gamma(\tilde{\gamma}_{\ell[k]}) \Gamma(\tilde{\gamma}_{\mathbf{b}})$$
  
=  $q^{1/2(d(\varphi_{\ell[k]}) + d(\varphi_{\mathbf{b}}))} K(\varphi_{\ell[k]}) K(\varphi_{\mathbf{b}}) \Theta(\tilde{\gamma}_{\ell[k]}) \Theta(\tilde{\gamma}_{\mathbf{b}})$ 

$$= q^{1/2(d(\varphi_{\ell[k]}) + d(\varphi_{\mathbf{b}}))} K(\varphi_{\ell[k]} + \varphi_{\mathbf{b}}) q^{1/2(\dim(O_{\ell[k]}) + \dim(O_{\mathbf{b}}))} G(\ell[k]) G(\mathbf{b}).$$

Since we have

$$d(\varphi_{\ell[k]}) = \dim(O_{\ell[k]}) = 0, \ G(\ell[k]) = E(\ell[k]) = \frac{1}{[\ell]_{q^{1/2}}!} E_k^{\ell}, \ \varphi_{\ell[k]} + \varphi_{\mathbf{b}} = \varphi_{\mathbf{a}},$$
so

$$\Gamma(\tilde{\gamma}_{\ell[k]} \star \tilde{\gamma}_{\mathbf{b}}) = \frac{1}{[\ell]_{q^{1/2}}!} q^{1/2(d(\varphi_{\mathbf{b}}) + \dim(O_{\mathbf{b}}))} K(\varphi_{\mathbf{a}}) E_k^{\ell} G(\mathbf{b}).$$

And

$$\Gamma(\gamma_{\mathbf{d}}) = q^{1/2d(\varphi_{\mathbf{d}})} K(\varphi_{\mathbf{d}}) \Theta(\gamma_{\mathbf{d}})$$
$$= q^{1/2(d(\varphi_{\mathbf{d}}) + \dim(O_{\mathbf{d}}))} K(\varphi_{\mathbf{d}}) E(\mathbf{d}).$$

Now write

$$\tilde{\gamma}_{\ell[k]} \star \tilde{\gamma}_{\mathbf{b}} = \sum_{\mathbf{b} \preceq_k \mathbf{d}, \varphi_{\mathbf{d}} = \varphi_{\mathbf{a}}} p_{\mathbf{d}, \mathbf{b}}(q) \gamma_{\mathbf{d}},$$

with

$$p_{\mathbf{d},\mathbf{b}}(q) = \sum_{i} q^{i} \mathcal{H}^{2i}(IC(\overline{O}_{\ell[k]}) \star IC(\overline{O}_{\mathbf{b}}))_{\mathbf{d}}.$$

Applying  $\Gamma$  gives

$$\frac{1}{[\ell]_{q^{1/2}!}}q^{1/2(d(\varphi_{\mathbf{b}})+\dim(O_{\mathbf{b}}))}K(\varphi_{\mathbf{a}})E_{k}^{\ell}G(\mathbf{b}) = \sum_{\mathbf{b} \leq_{k}\mathbf{d},\varphi_{\mathbf{d}}=\varphi_{\mathbf{a}}} p_{\mathbf{d},\mathbf{b}}(q)q^{1/2(d(\varphi_{\mathbf{d}})+\dim(O_{\mathbf{d}}))}K(\varphi_{\mathbf{d}})E(\mathbf{d}).$$

Hence

$$E_k^{\ell}G(\mathbf{b}) = [\ell]_{q^{1/2}}! \sum_{\mathbf{b} \leq_k \mathbf{d}, \varphi_{\mathbf{d}} = \varphi_{\mathbf{a}}} p_{\mathbf{d}, \mathbf{b}}(q) q^{1/2(d(\varphi_{\mathbf{d}}) + \dim(O_{\mathbf{d}}) - d(\varphi_{\mathbf{b}}) - \dim(O_{\mathbf{b}}))} E(\mathbf{d}),$$

now compare with Lemma 4.6, we get

$$(G(\mathbf{b}), E_i^{\prime n} E^*(\mathbf{d})) = [\ell]_{q^{1/2}}! p_{\mathbf{d}, \mathbf{b}}(q) q^{1/2(d(\varphi_{\mathbf{d}}) + \dim(O_{\mathbf{d}}) - d(\varphi_{\mathbf{b}}) - \dim(O_{\mathbf{b}}))}.$$

Finally, we write

$$\frac{1}{[\ell]_{q^{1/2}}!}E_i'^n E^*(\mathbf{d}) = \sum_{\mathbf{b}} n_{\mathbf{b},\mathbf{d}}(q)G^*(\mathbf{b}),$$

by applying the scalar product, we get

$$n_{\mathbf{b},\mathbf{d}}(q) = (G(\mathbf{b}), \frac{1}{[\ell]_{q^{1/2}}!} E_i^{\prime n} E^*(\mathbf{d})) = p_{\mathbf{d},\mathbf{b}}(q) q^{1/2(d(\varphi_{\mathbf{d}}) + \dim(O_{\mathbf{d}}) - d(\varphi_{\mathbf{b}}) - \dim(O_{\mathbf{b}}))}.$$

Hence, by specializing at q = 1, we have

$$n_{\mathbf{b},\mathbf{d}} = p_{\mathbf{d},\mathbf{b}}(1).$$

Now take  $\mathbf{d} = \mathbf{a}$ , we get the formula in our theorem.

#### TAIWANG DENG

## 5. A FORMULA FOR LUSZTIG'S PRODUCT

In this section we find a geometric way to calculate Lusztig's product in special case, which allows us to determine the partial BZ operator in the following sections.

**Definition 5.1.** Let  $k \in \mathbb{Z}$ . We say that **a** satisfies the assumption  $(\mathbb{A}_k)$  if it satisfies the following conditions <sup>1</sup>

(1): We have

 $\max\{b(\Delta) : \Delta \in \mathbf{a}\} + 1 < \min\{e(\Delta) : \Delta \in \mathbf{a}\}.$ 

(2): Moreover, we have  $\varphi_{e(\mathbf{a})}(k) \neq 0$  and  $\varphi_{e(\mathbf{a})}(k+1) = 0$ .

**Lemma 5.2.** Let **a** be a multisegment satisfying the assumption  $(\mathbb{A}_k)$ . Then **a** is of parabolic type. Moreover, The set  $S(\varphi_{\mathbf{a}})$  contains a unique maximal element satisfying the assumption  $(\mathbb{A}_k)$ , denoted by  $\mathbf{a}_{Id}$ .

Proof. Let  $b(\mathbf{a}) = \{k_1 \leq \cdots \leq k_r\}, e(\mathbf{a}) = \{\ell_1 \leq \cdots \leq \ell_r\}.$ 

Then by [7, Proposition 3.32], there exists an element  $w \in S_r^{J_1(\mathbf{a}), J_2(\mathbf{a})}$ , such that

$$\mathbf{a} = \sum_{j} [k_j, \ell_{w(j)}].$$

Let

$$\mathbf{a}_{\mathrm{Id}} = \sum_{j} [k_j, \ell_j].$$

By [7, Proposition 3.28] (The proposition is stated without proof but a proof can be given exactly as Proposition 3.18 in loc.cit..),  $\mathbf{a} \leq \mathbf{a}_{\text{Id}}$ . Finally,  $\mathbf{a}_{\text{Id}}$  depends only on  $b(\mathbf{a})$  and  $e(\mathbf{a})$ , not on  $\mathbf{a}$ , which shows that  $\mathbf{a}_{\text{Id}}$  is the maximal element in  $S(\varphi_{\mathbf{a}})$  satisfying the assumption  $(\mathbb{A}_k)$ .  $\Box$ 

Recall that in [6, Definition 5.4] we introduced a subset  $\hat{S}(\mathbf{a})_k$  of  $S(\mathbf{a})$ and further introduced a subset  $S(\mathbf{a})_k$  of  $\tilde{S}(\mathbf{a})_k$  in [6, Definition 5.9].

**Lemma 5.3.** Suppose that **a** is a multisegment satisfying the hypothesis  $(\mathbb{A}_k)$ , then

(1):  $\tilde{S}(\mathbf{a})_k = S(\mathbf{a}),$ (2): we have

$$X_{\mathbf{a}}^{k} = Y_{\mathbf{a}} = \coprod_{\mathbf{c} \in S(\mathbf{a})} O_{\mathbf{c}},$$

where the variety  $X_{\mathbf{a}}^k$  is defined in [6, Definition 5.20], which is a suitable sub-variety of  $E_{\varphi}$  and  $Y_{\mathbf{a}}$  is defined to be the union of all orbits indexed by elements of  $S(\mathbf{a})$ .

<sup>&</sup>lt;sup>1</sup>Since here we only work with the partial BZ operator  $\mathscr{D}^k$  with  $k \in \mathbb{Z}$ , for every multisegment, we can always use the reduction method to increase the length of segments from the left, so that at some point we arrive at the situation of our assumption  $(\mathbb{A}_k)$ , therefore we do not lose the generality. For a more precise description of such a process, we refer to [6, §6.2]

*Proof.* Note that by assumption

$$\max\{b(\Delta): \Delta \in \mathbf{a}\} < \min\{e(\Delta): \Delta \in \mathbf{a}\}.$$

This ensures that for any  $\mathbf{c} \in S(\mathbf{a})$ , we have  $\varphi_{e(\mathbf{c})}(k) = \varphi_{e(\mathbf{a})}(k)$ , hence by definition  $\mathbf{c} \in \tilde{S}(\mathbf{a})_k$ . This proves (1), and (2) follows from (1).  $\Box$ 

**Lemma 5.4.** Let **a** be a multisegment satisfying the assumption  $(\mathbb{A}_k)$ and  $\mathbf{a} = \mathbf{a}_{\text{Id}}$ . Let  $\ell \in \mathbb{N}$  such that  $\ell \leq \varphi_{e(\mathbf{a})}(k)$  and  $\varphi \in S$  such that

$$\varphi + \ell \chi_{[k]} = \varphi_{\mathbf{a}}.$$

Then for  $\mathbf{b} \in S(\varphi)$ , we have  $\mathbf{b} \preceq_k \mathbf{a}$  if and only if  $\mathbf{b}^{(k)} \leq \mathbf{a}^{(k)}$  (cf. Definition 2.1) and  $\varphi_{e(\mathbf{b})}(k-1) = \ell + \varphi_{e(\mathbf{a})}(k-1)$ .

*Proof.* Let  $\mathbf{b} \in S(\varphi)$  such that  $\mathbf{b} \preceq_k \mathbf{a}$ , then by Proposition 2.10  $\mathbf{b} = \mathbf{c}_{\Gamma}$  for some  $\mathbf{c} \in S(\mathbf{a})$  and  $\Gamma \subseteq \mathbf{c}(k)$ . Therefore

$$\mathbf{b}^{(k)} = \mathbf{c}^{(k)} \le \mathbf{a}^{(k)}$$

by Lemma 5.3. By definition of  $\mathbf{c}_{\Gamma}$ ,

$$\varphi_{e(\mathbf{b})}(k-1) = \ell + \varphi_{e(\mathbf{c})}(k-1).$$

Now applying the fact that **a** satisfies the assumption  $(\mathbb{A}_k)$ , we deduce that

$$\varphi_{e(\mathbf{c})}(k-1) = \varphi_{e(\mathbf{a})}(k-1).$$

Conversely, let  $\mathbf{b} \in S(\varphi)$  be a multisegment such that  $\mathbf{b}^{(k)} \leq \mathbf{a}^{(k)}$  and  $\varphi_{e(\mathbf{b})}(k-1) = \ell + \varphi_{e(\mathbf{a})}(k-1)$ .

We deduce from  $\mathbf{b}^{(k)} \leq \mathbf{a}^{(k)}$  that

$$\mathbf{b} \le \mathbf{a}^{(k)} + \varphi_{e(\mathbf{b})}(k)[k],$$

from which we obtain

$$\varphi_{\mathbf{b}} = \varphi_{\mathbf{a}^{(k)}} + \varphi_{e(\mathbf{b})}(k)\chi_{[k]}.$$

By assumption

$$\varphi_{\mathbf{b}} + \ell \chi_{[k]} = \varphi_{\mathbf{a}}.$$

Combining with the formula

$$\varphi_{\mathbf{a}} = \varphi_{\mathbf{a}^{(k)}} + \varphi_{e(\mathbf{a})}(k)\chi_{[k]},$$

we have

$$\varphi_{e(\mathbf{a})}(k) = \varphi_{e(b)}(k) + \ell.$$

Now that for any  $\Delta \in \mathbf{a}$ , if  $e(\Delta) = k$ , then  $b(\mathbf{a}) \leq k - 1$ . Therefore we have

$$\varphi_{e(\mathbf{a}^{(k)})}(k-1) = \varphi_{e(\mathbf{a})}(k-1) + \varphi_{e(\mathbf{a})}(k).$$

Applying the formula  $\varphi_{e(\mathbf{b})}(k-1) = \ell + \varphi_{e(\mathbf{a})}(k-1), \ \varphi_{e(\mathbf{b}^{(k)})}(k-1) = \varphi_{e(\mathbf{a}^{(k)})}(k-1)$ , we get

$$\varphi_{e(\mathbf{b}^{(k)})}(k-1) = \varphi_{e(\mathbf{b})}(k-1) + \varphi_{e(\mathbf{b})}(k).$$

Such a formula implies that for  $\Delta \in \mathbf{b}$ , if  $e(\Delta) = k$ , then  $b(\Delta) \leq k - 1$ .

Let  $b(\mathbf{a}) = \{k_1 \leq \cdots \leq k_r\}, e(\mathbf{a}) = \{\ell_1 \leq \cdots \leq \ell_r\}$ . The assumption that  $\mathbf{a} = \mathbf{a}_{\text{Id}}$  implies that

$$\mathbf{a} = \sum_{i} [k_i, \ell_i]$$

Suppose that

$$\mathbf{a}(k) = \{ [k_i, \ell_i] : i_0 \le i \le i_1 \}.$$

Take  $\Gamma = \{ [k_i, \ell_i] : i_0 + \ell \le i \le i_1 \}$  and  $\mathbf{a}' = \mathbf{a}_{\Gamma}$ 

$$\mathbf{a} = \mathbf{a}_{\Gamma}$$

Then  $\mathbf{a}' \leq_k \mathbf{a}$ . Note that  $\mathbf{a}'$  is a multisegment of parabolic type which corresponds to the identify in some symmetric group, cf. [7, Notation 3.25]. Finally, [7, Proposition 3.28] implies that  $\mathbf{b} \in S(\mathbf{a}')$ .

Recall that the variety  $X_{\mathbf{a}}^k$  in Lemma 5.3 admits a fibration  $\alpha$  over the Grassmanian  $Gr(\ell_{\mathbf{a},k}, V_{\varphi})$  (cf. [6, Proposition 5.28]) with

$$\ell_{\mathbf{a},k} = \sharp \{ \Delta \in \mathbf{a} : e(\Delta) = k \}, ( \text{ cf. } [6, \text{ Notation 5.12}] ).$$

For fixed  $W \in Gr(\ell_{\mathbf{a},k}, V_{\varphi})$ , let  $(X_{\mathbf{a}}^k)_W$  be the fiber of  $\alpha$  over W (cf. [6, Notation 5.30]). By [6, Proposition 5.35], we have an open immersion

$$\tau_W : (X^k_{\mathbf{a}})_W \to (Z^{k,\mathbf{a}})_W \times \operatorname{Hom}(V_{\varphi,k-1},W).$$

Here  $Z^{k,\mathbf{a}}$  is a sub-variety of a certain fiber bundles  $\tilde{Z}^k$  over the Grassmanian  $Gr(\ell_{\mathbf{a},k}, V_{\varphi})$  (cf. [6, Proposition 5.24]) and  $(Z^{k,\mathbf{a}})_W$  is the fiber over W.

**Lemma 5.5.** Assume that **a** is a multisegment satisfying  $(\mathbb{A}_k)$ . Let  $r \leq \varphi_{e(\mathbf{a})}(k)$  and  $\mathbf{d} = \mathbf{a} + r[k+1]$ . Then we have  $X_{\mathbf{d}}^{k+1} = Y_{\mathbf{d}}$  and for a fixed subspace W of  $V_{\varphi_{\mathbf{d}},k+1}$  of dimension r, the open immersion

$$\tau_W : (X^{k+1}_{\mathbf{d}})_W \to (Z^{k+1,\mathbf{d}})_W \times \operatorname{Hom}(V_{\varphi_{\mathbf{d}},k},W)$$

is an isomorphism.

*Proof.* Note that our assumption on **a** ensures that  $X_{\mathbf{d}}^{k+1} = Y_{\mathbf{d}}$  since we have  $\mathbf{d}_{\min} \in \tilde{S}(\mathbf{d})_{k+1}$  (cf. [6, Notation 5.11]). It suffices to show that  $\tau_W$  is surjective. Let  $(T^{(k)}, T_0) \in (Z^{k+1,\mathbf{d}})_W \times \operatorname{Hom}(V_{\varphi_{\mathbf{d}},k}, W)$ , by fixing a splitting  $V_{\varphi_{\mathbf{d}},k+1} = W \oplus V_{\varphi_{\mathbf{d}},k+1}/W$ , we define

$$T'|_{V_{\varphi_{\mathbf{d}},k}} = T_0 \oplus T^{(k+1)}|_{V_{\varphi_{\mathbf{d}},k}},$$
$$T'|_{V_{\varphi_{\mathbf{d}},k+1}} = T^{(k+1)}|_{V_{\varphi_{\mathbf{d}},k+1}/W} \circ p_W,$$
$$T'|_{V_{\varphi_{\mathbf{d}},i}} = T^{(k+1)}, \text{ for } i \neq k, k+1,$$

where  $p_w: V_{\varphi_{\mathbf{d}}} \to V_{\varphi_{\mathbf{d}},k}/W$  is the canonical projection. Then we have  $T' \in Y_{\mathbf{d}}$  hence  $T' \in (X_{\mathbf{d}}^{k+1})_W$ . Now since by construction we have  $\tau_W(T') = (T^{(k)}, T_0)$ , we are done.

**Definition 5.6.** Assume that **a** is a multisegment satisfying  $(\mathbb{A}_k)$  and  $\mathbf{d} = \mathbf{a} + r[k+1]$  for some  $r \leq \varphi_{e(\mathbf{a})}(k)$ . Let  $\mathfrak{X}_{\mathbf{d}}$  be the open subvariety of  $X_{\mathbf{d}}^{k+1}$  consisting of those orbits  $O_{\mathbf{c}}$  with  $\mathbf{c} \in S(\mathbf{d})$ , such that  $\varphi_{e(\mathbf{c})}(k) + r = \varphi_{e(\mathbf{a})}(k)$ .

**Definition 5.7.** Let V be a vector space and  $\ell_1 < \ell_2 < \dim(V)$  be two integers. We define

$$Gr(\ell_1, \ell_2, V) = \{ (U_1, U_2) : U_1 \subseteq U_2 \subseteq V, \dim(U_1) = \ell_1, \dim(U_2) = \ell_2 \}.$$

**Definition 5.8.** Let  $\ell$  be an integer and **a** be a multisegment. We let

$$E''_{\mathbf{a}} = \{ (T', W') : T' \in Y_{\mathbf{a}}, W' \in Gr(\ell, \ker(T'|_{V_{(2\mathbf{a},k)}})) \}.$$

Note that we have a canonical morphism

$$\alpha': E''_{\mathbf{a}} \to Gr(\ell, \varphi_{e(\mathbf{a})}(k), V_{\varphi_{\mathbf{a}}, k})$$

sending (T', W') to  $(W', \ker(T'|_{V_{\varphi_{\mathbf{a}},k}}))$ .

**Proposition 5.9.** The morphism  $\alpha'$  is a fibration.

*Proof.* The morphism  $\alpha'$  is equivariant under the action of  $GL(V_{\varphi_{\mathbf{a}},k})$ . The same proof as in [6, Proposition 5.28] shows that the morphism  $\alpha'$  is actually a  $P_{(U_1,U_2)}$  bundle, where  $P_{(U_1,U_2)}$  is a subgroup of  $GL(V_{\varphi_{\mathbf{a}},k})$  which fixes the given element  $(U_1, U_2)$ . Now we take a Zariski neighborhood  $\mathfrak{U}$  of  $(U_1, U_2)$  over which we have the trivialization

$$\gamma: \alpha'^{-1}(\mathfrak{U}) \simeq \alpha'^{-1}((U_1, U_2)) \times \mathfrak{U},$$

such an isomorphism comes from a section

$$s: \mathfrak{U} \to GL(V_{\varphi_{\mathbf{a}},k}), \ s((U_1, U_2)) = Id,$$

$$\tag{7}$$

given by  $\gamma((T, W')) = [(g^{-1}T, g^{-1}W'), \alpha'((T, W'))]$ , where  $g = s(\alpha'((T, W')))$ . We remark that the existence of the section s is guaranteed by local triviality of  $GL(V_{\varphi_{\mathbf{a}},k}) \to GL(V_{\varphi_{\mathbf{a}},k})/P_{(U_1,U_2)}$ , cf. [13], § 4.

Note that we use  $(\mathfrak{X}_{\mathbf{d}})_W$  to denote the intersection of  $\mathfrak{X}_{\mathbf{d}}$  with the fiber above W along the morphism  $\alpha$  discussed before Lemma 5.5.

**Proposition 5.10.** Assume that **a** is a multisegment satisfying  $(\mathbb{A}_k)$ and  $\mathbf{d} = \mathbf{a} + r[k+1]$  for some  $r \leq \varphi_{e(\mathbf{a})}(k)$ . Let  $\ell \in \mathbb{N}$  such that  $r + \ell = \varphi_{e(\mathbf{a})}(k)$  and W a subspace of  $V_{\varphi_{\mathbf{d}},k+1}$  such that  $\dim(W) = r$ . We have a canonical projection

$$p: (\mathfrak{X}_{\mathbf{d}})_W \to E''_{\mathbf{a}} \tag{8}$$

where for  $T \in (\mathfrak{X}_{\mathbf{d}})_W$  with  $\tau_W(T) = (T_1, T_0) \in (Z^{k+1,\mathbf{d}})_W \times \operatorname{Hom}(V_{\varphi_{\mathbf{d}},k}, W)$ , we define  $p(T) = (T_1, \ker(T_0|_{W_1}))$ , where  $W_1 = \ker(T_1|_{V_{\varphi_{\mathbf{d}},k}})$  (Note that here we identify  $(Z^{k+1,\mathbf{d}})_W$  with  $Y_{\mathbf{a}}$ , see the remark after [6, Proposition 5.31]). Moreover, let  $U_1 \subseteq U_2 \subseteq V_{\varphi_{\mathbf{d}},k}$  be subspaces such that  $\dim(U_1) = \ell$ ,  $\dim(U_2) = \varphi_{e(\mathbf{a})}(k)$ , then p is a fibration with fiber

$$\{T \in \operatorname{Hom}(V_{\varphi_{\mathbf{d}},k}, W) : \ker(T|_{U_2}) = U_1\}$$

*Proof.* We show that p is well defined. Since by definition of  $(\mathfrak{X}_{\mathbf{d}})_W$ ,

 $\dim(W) + \dim(\ker(T_0|_{\ker(T_1|_{V_{\varphi_{\mathbf{d}},k}})})) = \dim(\ker(T_1|_{V_{\varphi_{\mathbf{d}},k}})),$ 

hence in order to see that

$$\ell = \dim(\ker(T_0|_{\ker(T_1|_{V_{\varphi_{\mathbf{d}},k}})})),$$

it suffices to show that

$$\varphi_{e(\mathbf{a})}(k) = \dim(\ker(T_1|_{V_{\varphi_{\mathbf{a}},k}}))$$

This follows from the fact that  $\mathbf{a} = \mathbf{d}^{(k+1)}$ . Finally, we show that p is a fibration. Note that by definition, the fiber of p is isomorphic to

 $\{T \in \operatorname{Hom}(V_{\varphi_{\mathbf{d}},k}, W) : \ker(T|_{U_2}) = U_1\}.$ 

So it suffices to show that it is locally trivial. To show this, we consider the open subset  $\mathfrak{U}$  in  $E''_{\mathbf{a}}$  as constructed in the proof of Proposition 5.9. Let s be the section in (7). We construct a trivialization for p:

 $\varrho: p^{-1}(\alpha'^{-1}(\mathfrak{U})) \to \alpha'^{-1}(\mathfrak{U}) \times \{T \in \operatorname{Hom}(V_{\varphi_{\mathbf{d}},k}, W) : \ker(T|_{U_2}) = U_1\}$ with  $\varrho(T) = [(T_1, W'), g^{-1}(T_0)]$ , where  $g = s((W', W_1)), W_1 = \ker(T_1|_{V_{\varphi_{\mathbf{d}},k}})$ . Note that given

 $[(T_1, W'), T_0] \in \alpha'^{-1}(\mathfrak{U}) \times \{T \in \operatorname{Hom}(V_{\varphi_{\mathbf{d}}, k}, W) : \ker(T|_{U_2}) = U_1\},$ 

then  $(W', W_1) \in \mathfrak{U}$  with  $W_1 = \ker(T_1|_{V_{\varphi_{\mathbf{d}},k}})$ . It follows that we can take the section along s in (7) to obtain  $g = s((W', W_1))$ . Let  $T'_0 = gT_0$ . Then  $T = \tau_W^{-1}((T_1, T'_0)) \in p^{-1}(\alpha'^{-1}(\mathfrak{U}))$ .

**Definition 5.11.** Let  $\ell + \dim(W) = \varphi_{e(\mathbf{a})}(k)$ . We define  $\mathfrak{Y}_{\mathbf{a}}$  to be the set of pairs (T, U) satisfying

(1):  $U \in Gr(\ell, V_{\varphi_{\mathbf{a}},k}), T \in End(V_{\varphi_{\mathbf{a}}}/U)$  of degree 1;

(2):  $T \in O_{\mathbf{b}}$  for some  $\mathbf{b} \leq_k \mathbf{a}$  (cf. Definition 2.2).

(

And we have a canonical projection

$$\tau: E''_{\mathbf{a}} \to \mathfrak{Y}_{\mathbf{a}}$$

for  $(T, U) \in E''_{\mathbf{a}}$ , we associate

$$\sigma((T,U)) = (T',U)$$

where  $T' \in \operatorname{End}(V_{\varphi_{\mathbf{a}}}/U)$  is the quotient of T. Also, we have a morphism  $\sigma': \mathfrak{Y}_{\mathbf{a}} \to Gr(\ell, \varphi_{e(\mathbf{a})}(k), V_{\varphi_{\mathbf{a}}, k}),$ (9)

by

$$\sigma'((T,U)) = (U, \pi^{-1}(\ker(T|_{V_{(2n,k)}/U}))).$$

where  $\pi: V_{\varphi_{\mathbf{a}},k} \to V_{\varphi_{\mathbf{a}},k}/U$  is the canonical projection.

**Lemma 5.12.** We have for  $(T, U) \in \mathfrak{Y}_{\mathbf{a}}$ ,

- (1):  $\gamma_k(T) \in Z^{k,\mathbf{a}}$  (cf. [6, Definition 5.25]),
- (2):  $T|_{V_{\varphi_{\mathbf{a}},k-1}}$  is surjective,

(3): dim(ker(
$$T|_{V_{\omega_{\mathbf{a}},k}/U}$$
)) = dim( $W$ ),

where the morphism  $\gamma_k : X_k^{\mathbf{a}} \to Z^{k,\mathbf{a}}$  is defined in [6, After Definition 5.26].

*Proof.* (1)To show  $\gamma_k(T) \in Z^{k,\mathbf{a}}$ , it suffices to show that for  $\mathbf{b} \leq \mathbf{a}^k$  a implies that  $\mathbf{b}^{(k)} \leq \mathbf{a}^{(k)}$ . Note that by Proposition 2.10, there exists  $\mathbf{c} \in S(\mathbf{a})$  and  $\Gamma \subseteq \mathbf{c}(k)$ , such that

$$\mathbf{b} = \mathbf{c}_{\Gamma}.$$

Now by Lemma 5.3, we have  $\mathbf{c} \in \tilde{S}(\mathbf{a})_k$ , which implies that

$$\mathbf{b}^{(k)} = \mathbf{c}^{(k)} \le \mathbf{a}^{(k)}.$$

(2)By definition, for any  $(T, U) \in \mathfrak{Y}_{\mathbf{a}}$  we have  $T \in O_{\mathbf{b}}$  for some  $\mathbf{b} \leq_k \mathbf{a}$ . By the fact that  $\mathbf{a}$  satisfies the assumption  $(\mathbb{A}_k)$ , any  $\mathbf{c} \in S(\mathbf{a})$  also satisfies  $(\mathbb{A}_k)$ , hence  $\mathbf{b} = \mathbf{c}_{\Gamma}$  cannot contain a segment which starts at k, therefore  $T|_{V_{\varphi_{\mathbf{a}},k-1}}$  is surjective.

(3) Note that by the definition of  $\mathfrak{Y}$ , for  $(T, U) \in \mathfrak{Y}_{\mathbf{a}}$ , we have  $T \in O_{\mathbf{b}}$  for some  $\mathbf{b} \leq_{\mathbf{k}} \mathbf{a}$ . Now it follows

$$\ker(T|_{V_{\varphi_{\mathbf{a}},k}/U}) = \varphi_{e(\mathbf{a})}(k) - \ell = \dim(W).$$

**Proposition 5.13.** Let **a** be a multisegment satisfying the assumption  $(\mathbb{A}_k)$ . Then the morphism  $\sigma'$  is a fibration. Moreover, if we assume that  $\mathbf{a} = \mathbf{a}_{Id}$  (cf. Lemma 5.2), then the morphism  $\sigma$  is also a fibration.

Proof. We first show that  $\sigma'$  is locally trivial. We observe that the group  $GL(V_{\varphi_{\mathbf{a}},k})$  acts both on the source and target of  $\sigma'$  in such a way that  $\sigma'$  is  $GL(V_{\varphi_{\mathbf{a}},k})$ -equivariant. As in the proof of Proposition 5.9, let  $\mathfrak{U} \subseteq Gr(\ell, \varphi_{e(\mathbf{a})}(k), V_{\varphi_{\mathbf{a}},k})$  be a neighborhood of a given element  $(U_1, U_2)$  such that we have a section

$$s: \mathfrak{U} \to GL(V_{\varphi_{\mathbf{a}},k}), \ s((U_1, U_2)) = Id.$$

Note that in this case we have a natural trivialization of  $\sigma'$  by

$$\sigma':\beta'^{-1}(\mathfrak{U})\simeq\mathfrak{U}\times\beta'^{-1}((U_1,U_2))$$

by  $\sigma'((T,U)) = [(U, \pi^{-1}(\ker(T|_{V_{\varphi_{\mathbf{a}},k}}))), g^{-1}((T,U))]$  with  $g = s((U, \pi^{-1}(\ker(T|_{V_{\varphi_{\mathbf{a}},k}}))))$ . Finally, we show that  $\sigma$  is surjective and locally trivial.

We observe that  $\alpha' = \sigma' \sigma$  and  $\sigma$  preserves fibers. Now we fix a neighborhood  $\mathfrak{U}$  as above and get a commutative diagram

where  $\delta([x,T]) = [x,\sigma(T)]$  for any  $x \in \mathfrak{U}$  and  $T \in \alpha'^{-1}((U_1,U_2))$ . Therefore to show that  $\sigma$  is locally trivial, it suffices to show that it is

locally trivial when restricted to the fiber  $\alpha'^{-1}((U_1, U_2))$ . Note that we have

$$\alpha'^{-1}((U_1, U_2)) \simeq \{T \in Y_{\mathbf{a}} : \ker(T|_{V_{\varphi_{\mathbf{a}},k}}) = U_2\} \simeq (X_{\mathbf{a}}^k)_{U_2} \hookrightarrow Y_{\mathbf{a}^{(k)}} \times \operatorname{Hom}(V_{\varphi_{\mathbf{a}},k-1}, U_2)$$
  
and

$$\begin{split} \sigma'^{-1}((U_1, U_2)) &\simeq \{T : T \in \operatorname{End}(V_{\varphi_{\mathbf{a}}}/U_1) \text{ of degree } 1, \ker(T|_{V_{\varphi_{\mathbf{a}},k}/U_1}) = U_2/U_1; \\ T \in O_{\mathbf{b}}, \text{ for some } \mathbf{b} \preceq_k \mathbf{a} \} \hookrightarrow Y_{\mathbf{a}^{(k)}} \times \operatorname{Hom}(V_{\varphi_{\mathbf{a}},k-1}, U_2/U_1). \end{split}$$

Note that the canonical morphism

$$\operatorname{Hom}(V_{\varphi_{\mathbf{a}},k-1},U_2) \to \operatorname{Hom}(V_{\varphi_{\mathbf{a}},k-1},U_2/U_1)$$

is a fibration. Hence to show that

$$\alpha'^{-1}((U_1, U_2)) \to \sigma'^{-1}((U_1, U_2))$$

is a fibration, it suffices to show that  $\sigma|_{\alpha'^{-1}((U_1,U_2))}$  is surjective with isomorphic fibers everywhere . Let  $(T, U_1) \in \sigma'^{-1}((U_1, U_2))$  with

$$\tau_{U_2/U_1}(T) = (T_0, q_0) \in Y_{\mathbf{a}^{(k)}} \times \operatorname{Hom}(V_{\varphi_{\mathbf{a}}, k-1}, U_2/U_1),$$

where  $\tau_{U_2/U_1}$  is the morphism defined in [6, After Definition 5.26](see also the remark before [6, Definition 5.26]). We fix a splitting  $U_2 \simeq U_2/U_1 \oplus U_1$ . Now to give  $(T', U_1) \in \sigma^{-1}((T, U_1))$  amounts to give  $q_1 \in \operatorname{Hom}(V_{\varphi_{\mathbf{a}},k-1}, U_1)$  such that

$$au_{U_2}(T') = (T_0, q_0 \oplus q_1).$$

Note that by Lemma 5.4, the condition  $\mathbf{a} = \mathbf{a}_{\text{Id}}$  implies that T' lies in  $(X^k_{\mathbf{a}})_{U_2}$  if and only if  $q_1$  satisfies

$$\dim(\ker(q_0 \oplus q_1|_{\ker(T_0|_{V_{\varphi_{\mathbf{a}},k-1}})})) = \varphi_{e(\mathbf{a})}(k-1),$$

which is an open condition. Therefore  $\sigma$  is surjective. By definition of  $\mathfrak{Y}_{\mathbf{a}}$ ,

$$\dim(\ker(q_0|_{\ker(T_0|_{V_{\varphi_{\mathbf{a}},k-1}})})) = \varphi_{e(\mathbf{a})}(k-1) + \ell,$$

therefore if we denote  $W_1 = \ker(q_0|_{\ker(T_0|_{V_{\varphi_{\mathbf{a}},k-1}})})$ , then  $q_1$  satisfies that

$$\dim(\ker(q_1|_{\ker(T_0|_{V_{\varphi_{\mathbf{a}},k-1}})}) \cap W_1) = \varphi_{e(\mathbf{a})}(k-1).$$

Such a condition is independent of the pair  $(T_0, q_0)$  since we always have  $\dim(\ker(T_0|_{V_{\varphi_{\mathbf{a},k-1}}})) = \varphi_{e(\mathbf{a})}(k-1) + \varphi_{e(\mathbf{a})}(k)$  and  $\dim(W_1) = \varphi_{e(\mathbf{a})}(k-1) + \ell$ .

We return to the morphism p and  $\sigma$ .

**Lemma 5.14.** Note that an element of  $G_{\varphi_d}$  stabilizes  $(\mathfrak{X}_d)_W$  if and only if it stabilizes W. Let  $G_{\varphi_d,W}$  be the stabilizer of W, then for  $\mathbf{c} \leq \mathbf{d}$ , and  $T \in O_{\mathbf{c}} \cap (\mathfrak{X}_d)_W$ , we have

$$O_{\mathbf{c}} \cap (\mathfrak{X}_{\mathbf{d}})_W = G_{\varphi_{\mathbf{d}},W}T.$$

*Proof.* Recall that by [7, Proposition 2.9], we have



where  $\ell_{k+1} = \varphi_{e(\mathbf{d})}(k+1)$ . Note that we have

 $G_{\varphi_{\mathbf{d}},W} = \cdots \times G_{\varphi_{\mathbf{d}},k} \times P_W \times G_{\varphi_{\mathbf{d}},k+2} \times \cdots,$ 

where  $G_{\varphi_{\mathbf{d}},i} = GL(V_{\varphi_{\mathbf{d}},i})$ . From this diagram we observe that there is a one to one correspondance between the  $G_{\varphi_{\mathbf{d}}}$  orbits on  $X_{\mathbf{d}}^{k+1}$  and  $G_{\varphi_{\mathbf{d}},W}$ orbits on  $\alpha^{-1}(W)$ . Finally, since  $\mathfrak{X}_{\mathbf{d}}$  is an open subvariety consisting of  $G_{\varphi_{\mathbf{d}}}$  orbits, we are done.

Definition 5.15. The canonical projection

$$\pi: V_{\varphi_{\mathbf{d}}} \to V_{\varphi_{\mathbf{d}}}/W$$

induces a projection

$$F_*: G_{\varphi_{\mathbf{d}}, W} \to G_{\varphi_{\mathbf{a}}},$$

where we identify  $V_{\varphi_{\mathbf{d}}}/W$  with  $V_{\varphi_{\mathbf{a}}}$ .

**Proposition 5.16.** Assume that **a** is a multisegment satisfying  $(\mathbb{A}_k)$ and  $\mathbf{d} = \mathbf{a} + r[k+1]$  for some  $r \leq \varphi_{e(\mathbf{a})}(k)$ . The morphism p is equivariant under the action of  $G_{\varphi_{\mathbf{d}},W}$  and  $G_{\varphi_{\mathbf{a}}}$  via  $\pi_*$ , i.e,

$$p(gx) = \pi_*(g)p(x).$$

Moreover, it induces a one to one correspondance between orbits.

*Proof.* Note that for  $T \in (\mathfrak{X}_{\mathbf{d}})_W$ , such that  $\tau_W(T) = (T_1, T_0) \in (Z^{k+1,\mathbf{d}})_W \times \operatorname{Hom}(V_{\varphi_{\mathbf{d}},k}, W)$ , let

$$U_1 = \ker(T_1|_{V_{\varphi_d,k}}), \ U_0 = \ker(T_0|_{U_1})$$

we have

$$p(T) = (T_1, U_0).$$

Now it follows from the definition that we have

$$p(gT) = \pi_*(g)p(T).$$

Hence p sends orbits to orbits. It remains to show that the pre-image of an orbit is an orbit instead of a union of orbits.

We proved in Proposition 5.10 that

$$p^{-1}p(T) = \{(T_1, q) : q \in \operatorname{Hom}(V_{\varphi_{\mathbf{d}}, k}, W), \ker(q|_{U_1}) = U_0\},\$$

note that here we identify elements of  $(\mathfrak{X}_{\mathbf{d}})_W$  with its image under  $\tau_W$ . Let  $(T_1, q) \in p^{-1}p(T)$ . Then we want to find  $g \in G_{\varphi_{\mathbf{d}},W}$  such that  $g(T_1, T_0) = (T_1, q)$ . Note that by fixing a splitting  $V_{\varphi_{\mathbf{d}},k+1} =$ 

 $W \oplus V_{\varphi_{\mathbf{d}},k+1}/W$ , we can choose  $g \in G_{\varphi_{\mathbf{d}}}$  such that  $g_i = Id \in GL(V_{\varphi_{\mathbf{d}},i})$  for all  $i \neq k+1$ , and

$$g_{k+1} = \begin{pmatrix} g_1 & g_{12} \\ 0 & Id_{V_{\varphi_{\mathbf{d}},k+1}/W} \end{pmatrix} \in P_W,$$

where  $g_1 \in GL(W)$ , and  $g_{12} \in \text{Hom}(V_{\varphi_d,k+1}/W,W)$ . By hypothesis, the restrictions of q and  $T_0$  to  $U_1$  are surjective with kernel  $U_0$ . So we can choose  $g_1 \in GL(W)$ , such that

$$g_1T_0(v) = q(v)$$
, for all  $v \in U_1$ .

Finally for  $v_1 \in V_{\varphi_{\mathbf{d}},k+1}/W$ , our assumption on **a** implies that  $T_1|V_{\varphi_{\mathbf{d}},k}$  is surjective. Hence there exists  $v \in V_{\varphi_{\mathbf{d}},k}$  such that  $T_1(v) = v_1$ . Then we define

$$g_{12}(v_1) = q(v) - g_1 T_0(v).$$

We check that this is well defined, i.e, for another  $v' \in V_{\varphi_{\mathbf{d}},k}$  such that  $T_1(v') = v_1$ , we have

$$q(v) - g_1 T_0(v) = q(v') - g_1 T_0(v'),$$

this is the same as to say that

$$q(v - v') = g_1 T_0(v - v').$$

We observe that  $T_1(v - v') = 0$ , hence  $v - v' \in U_1$ , now  $q(v - v') = g_1 T_0(v - v')$  follows from our definition of  $g_1$ . Under such a choice, we have

$$g((T_1, T_0)) = (T_0, q).$$

Hence we are done.

**Proposition 5.17.** The morphism  $\sigma$  is equivariant under the action of  $G_{\varphi_{\mathbf{a}}}$ . Assume that  $\mathbf{a}$  is a multisegment which satisfies the assumption  $(\mathbb{A}_k)$ . Let  $\varphi \in \mathcal{S}$  such that

$$\varphi + \ell \chi_{[k]} = \varphi_{\mathbf{a}},$$

where  $\chi$  is the characteristic function at k. Then there exists a one to one correspondence between the  $G_{\varphi_{\mathbf{a}}}$ -orbits on  $\mathfrak{Y}_{\mathbf{a}}$  and the set

$$S := \{ \mathbf{b} \in S(\varphi) : \mathbf{b} \preceq_k \mathbf{a} \}.$$

Moreover, for each orbit  $\mathfrak{Y}_{\mathbf{a}}(\mathbf{b})$  indexed by  $\mathbf{b}$  on  $\mathfrak{Y}_{\mathbf{a}}$ ,  $\sigma^{-1}(\mathfrak{Y}_{\mathbf{a}}(\mathbf{b}))$  is irreducible hence contains a unique orbit in  $E''_{\mathbf{a}}$  as (Zariski) open subset.

*Proof.* The fact that  $\sigma$  is equivariant under the action of  $G_{\varphi_{\mathbf{a}}}$  follows directly from the definition. To show that the  $G_{\varphi_{\mathbf{a}}}$ -orbits on  $\mathfrak{Y}_{\mathbf{a}}$  is indexed by S, consider the morphism

$$p': \mathfrak{Y}_{\mathbf{a}} \to Gr(\ell, V_{\varphi_{\mathbf{a}}, k}), \ (T, U) \mapsto U$$

By [7, Proposition 2.9], we have the following diagram



such that p' is a  $GL_{\varphi_{\mathbf{a},k}}$  bundle. Moreover, the same proof as in Lemma 5.14 shows that the orbits on  $\mathfrak{Y}_{\mathbf{a}}$  are in one to one correspondence with that of the fibers

$$p'^{-1}(U) \simeq \{T \in \operatorname{End}(V_{\varphi_{\mathbf{a}}}/U) : T \text{ is of degree } 1, T \in O_{\mathbf{b}} \text{ for some } \mathbf{b} \preceq_{k} \mathbf{a}\},\$$
  
under the action of stabilizer  $G_{\varphi_{\mathbf{a}},U}$  of  $U$ . Let  $\varphi \in \mathcal{S}$  be the such that  $\varphi + \ell \chi_{[k]} = \varphi_{\mathbf{a}}$ . Then by identifying  $V_{\varphi}$  with  $V_{\varphi_{\mathbf{a}}}/U$ , we can view  $p'^{-1}(U)$   
as an open subvariety of  $E_{\varphi}$ . Note that we are identifying orbits with  
orbits by the canonical projection

$$G_{\varphi_{\mathbf{a}},U} \to G_{\varphi}.$$

Now it follows that the fibers are parametrized by the set S. Finally, let  $\mathbf{b} \in S$ . We have to show that  $\sigma^{-1}(\mathfrak{Y}_{\mathbf{a}}(\mathbf{b}))$  is irreducible, which is a consequence of the following lemma.

**Lemma 5.18.** Let  $\mathbf{a}, \mathbf{b}$  be the multisegments as above. Then there exists a bijection between the set

$$Q(\mathbf{a}, \mathbf{b}) = \{ \mathbf{c} \in S(\mathbf{a}) : \mathbf{b} = \mathbf{c}_{\Gamma} \text{ for some } \Gamma \subseteq \mathbf{c}(k) \}$$

and the orbits in  $\sigma^{-1}(\mathfrak{Y}_{\mathbf{a}}(\mathbf{b}))$  which respects the poset structure given by

$$\mathbf{c} \mapsto E_{\mathbf{a}}''(\mathbf{c}^{\sharp}),$$

where for  $\mathbf{b} = \mathbf{c}_{\Gamma}$ ,

$$\mathbf{c}^{\sharp} = (\mathbf{c} \setminus \mathbf{c}(k)) \cup \Gamma \cup \{\Delta^{+} : \Delta \in \mathbf{c}(k) \setminus \Gamma\},\$$

and  $E''_{\mathbf{a}}(\mathbf{c}^{\sharp})$  is the orbit indexed by  $\mathbf{c}^{\sharp}$  on  $E''_{\mathbf{a}}$ . Moreover, the set  $Q(\mathbf{a}, \mathbf{b})$  contains a unique minimal element.

*Proof.* We constructed in (8) of Proposition 5.10 a morphism p. Consider the composition

$$(\mathfrak{X}_{\mathbf{d}})_W \xrightarrow{p} E''_{\mathbf{a}} \xrightarrow{\sigma} \mathfrak{Y}_{\mathbf{a}},$$

which sends  $(O_{\mathbf{c}})_W$  to  $\mathfrak{Y}(\mathbf{b})$ , where  $\mathbf{b} = \mathbf{c}^{(k,k+1)}$  for  $\mathbf{c} \in S(\mathbf{d})$ . Hence we have

$$\mathbf{b} = (\mathbf{c}^{(k+1)})_{\Gamma}$$

for  $\Gamma = \{\Delta \in \mathbf{c} : e(\Delta) = k\}$ . Note that  $\mathbf{c} \in S(\mathbf{d})$  implies that  $\mathbf{c}^{(k+1)} \leq \mathbf{a} = \mathbf{d}^{(k+1)}$ . Conversely, for  $\mathbf{c} \in Q(\mathbf{a}, \mathbf{b})$ , such that

$$\mathbf{b} = \mathbf{c}_{\Gamma}$$

there is a unique element

$$\mathbf{c}' = \mathbf{c}^{\sharp}$$

in  $S(\mathbf{d})$  such that  $O_{\mathbf{c}'} \subseteq \mathfrak{X}_{\mathbf{d}}$  and  $\mathbf{c} = \mathbf{c}'^{(k+1)}$ . Therefore we conclude that there is a bijection between the  $G_{\varphi_{\mathbf{a}}}$ -orbits in  $\sigma^{-1}(\mathfrak{Y}_{\mathbf{a}}(\mathbf{b}))$  and  $Q(\mathbf{a}, \mathbf{b})$ . Finally, for

Finally, for

$$\varphi_{\mathbf{a}} = \varphi_{\mathbf{b}} + \ell \chi_{[k]},$$

we show by induction on  $\ell$  that the set  $Q(\mathbf{a}, \mathbf{b})$  contains a unique minimal element.

For case  $\ell = 1$ , let

$$\mathbf{b}(k) := \{ \Delta \in \mathbf{b} : e(\Delta) = k \} = \{ \Delta_1 \preceq \cdots \preceq \Delta_h \}.$$

and  $\mathbf{c}_i = (\mathbf{b} \setminus \Delta_i) \cup \Delta_i^+$ . Then

$$Q(\mathbf{a},\mathbf{b}) \subseteq {\mathbf{c}_i : i = 1, \cdots, h}$$

and  $\mathbf{c}_h$  is minimal in the latter, which implies that  $\mathbf{c}_h \in Q(\mathbf{a}, \mathbf{b})$  and is minimal. In general, let

$$\varphi = \varphi_{\mathbf{b}} + \chi_{[k]}.$$

Note that there exists  $\mathbf{c}' \in S(\varphi)$  satisfying the assumption  $(\mathbb{A}_k)$  and  $\Gamma' \subseteq \mathbf{c}'(k)$  such that

$$\mathbf{b} = \mathbf{c}'_{\Gamma'}.$$

In fact, by assumption

 $\mathbf{b} = \mathbf{c}_{\Gamma}$ 

for some  $\mathbf{c} \in S(\mathbf{a})$  and  $\Gamma \subseteq \mathbf{c}(k)$ . Let

$$\Gamma \supseteq \Gamma_1$$
,

such that  $\ell = \sharp \Gamma = \sharp \Gamma' + 1$  and

$$\mathbf{c}' = \mathbf{c}_{\Gamma_1},$$

then we have

$$\mathbf{b} = \mathbf{c}'_{\Gamma \setminus \Gamma_1}.$$

Now we apply our induction to the case

 $Q_1 := \{ \mathbf{c} \in S(\varphi) : \mathbf{c} \text{ satisfies the assumption } (\mathbb{A}_k), \mathbf{b} = \mathbf{c}_{\Gamma} \text{ for some } \Gamma \subseteq \mathbf{c}(k) \},\$ 

from which we know that there exists a unique minimal element  $\mathbf{c}_1$  in  $Q_1$ . Now by assumption

$$\mathbf{b}_1 \leq \mathbf{c}' \preceq_k \mathbf{a},$$

and by induction the set  $Q(\mathbf{a}, \mathbf{b}_1)$  contains a unique element  $\mathbf{b}_2$ . We claim that  $\mathbf{b}_2$  is minimal in  $Q(\mathbf{a}, \mathbf{b})$ . In fact, let  $\mathbf{e} \in Q(\mathbf{a}, \mathbf{b})$ , then

$$\mathbf{b} = \mathbf{e}_{\Gamma'}$$

for some  $\Gamma' \subseteq \mathbf{e}(k)$ . Again let

$$\Gamma'_1 \subseteq \Gamma', \mathbf{e}' = \mathbf{e}_{\Gamma'_1}$$

such that  $\ell = \sharp \Gamma' = \sharp \Gamma'_1 + 1$ . Now we obtain

$$\mathbf{e}' \in Q_1, \ \mathbf{b} = \mathbf{e}'_{\Gamma' \setminus \Gamma'_1}.$$

By minimality of  $\mathbf{c}_1$ , we know that

 $\mathbf{c}_1 \leq \mathbf{e}'.$ 

Note that this implies  $\mathbf{c}_1 \preceq \mathbf{e}'$ , and by transitivity of poset relation, we get  $\mathbf{c}_1 \preceq_k \mathbf{e}$ . Now we apply Proposition 2.10 to get

$$\mathbf{c}_1 = \mathbf{f}_{\Gamma''},$$

for some  $\mathbf{f} \in S(\mathbf{e})$  and  $\Gamma'' \subseteq \mathbf{f}(k)$ . Again we deduce from induction that

$$\mathbf{f} \geq \mathbf{c}_2$$
.

Hence  $\mathbf{c}_2 \leq \mathbf{e}$ .

Now we return to the calculation of product of perverse sheaves, cf. Corollary 3.17.

**Theorem 5.19.** Let **a** be a multisegment satisfying the assumption  $(\mathbb{A}_k)$  and  $\mathbf{b} \leq_k \mathbf{a}$  such that

$$\varphi_{\mathbf{a}} = \varphi_{\mathbf{b}} + \ell \chi_{[k]}, \quad \ell \in \mathbb{N}.$$

Let **c** the minimal element in  $Q(\mathbf{a}, \mathbf{b})$  and  $E''_{\mathbf{a}}(\mathbf{c})$  be the  $G_{\varphi_{\mathbf{a}}}$  orbit indexed by **c** in  $E''_{\mathbf{a}}$  (cf. Definition 5.8). Then we have

$$IC(\overline{O}_{\mathbf{b}}) \star IC(\overline{O}_{\ell[k]}) = \beta_*''(IC(\overline{E_{\mathbf{a}}''(\mathbf{c}^{\sharp})})).$$

*Proof.* First of all, by definition

$$E'' = \{ (T, U) : T \in E_{\varphi_{\mathbf{a}}}, \ T(U) = 0, \ \dim(U) = \ell \},\$$

therefore we have

$$E''_{\mathbf{a}} \subseteq E''.$$

Furthermore, the variety  $E_{\mathbf{a}}''$  is open in E''. In fact, consider the canonical morphism

 $\beta'': E'' \to E_{\varphi_{\mathbf{a}}},$ 

then  $E''_{\mathbf{a}} = \beta''^{-1}(Y_{\mathbf{a}})$ . Since  $Y_{\mathbf{a}}$  is open in  $E_{\varphi_{\mathbf{a}}}$ ,  $E''_{\mathbf{a}}$  is open in E''. Now we have two morphisms

$$\sigma\beta' : \beta'^{-1}(E''_{\mathbf{a}}) \to \mathfrak{Y}_{\mathbf{a}}, \beta : E' \to E_{\varphi_{\mathbf{b}}} \times E_{\varphi_{\ell[k]}} \simeq E_{\varphi_{\mathbf{b}}}.$$

We claim that  $\beta^{-1}(O_{\mathbf{b}}) \cap \beta'^{-1}(E''_{\mathbf{a}}) = \beta'^{-1}\sigma^{-1}(\mathfrak{Y}_{\mathbf{a}}(\mathbf{b}))$ , where  $\mathfrak{Y}_{\mathbf{a}}(\mathbf{b})$  is the orbit in  $\mathfrak{Y}_{\mathbf{a}}(\mathbf{b})$  under the action of  $G_{\varphi_{\mathbf{a}}}$ .

By definition of  $\beta$ ,

$$\beta^{-1}(O_{\mathbf{b}}) \cap \beta'^{-1}(E_{\mathbf{a}}'') = \{ (T, W, \mu, \mu') | \mu : W \simeq V_{\varphi_{\ell[k]}}, \ \mu' : V_{\varphi_{\mathbf{a}}}/W \simeq V_{\varphi_{\mathbf{b}}},$$
$$T \in O_{\mathbf{f}} \text{ for some } \mathbf{f} \in S(\mathbf{a}), \mathbf{b} \preceq_k \mathbf{f} \}.$$

Now by definition of  $\sigma$  and  $\beta'$ , we know that  $\beta^{-1}(O_{\mathbf{b}}) \cap \beta'^{-1}(E''_{\mathbf{a}}) = \beta'^{-1}\sigma^{-1}(\mathfrak{Y}_{\mathbf{a}}(\mathbf{b}))$ . Now by Proposition 5.17,  $\sigma^{-1}(\mathfrak{Y}_{\mathbf{a}}(\mathbf{b}))$  contains  $E''_{\mathbf{a}}(\mathbf{c}^{\sharp})$ 

as the unique open orbit, where **c** is the minimal element in  $Q(\mathbf{a}, \mathbf{b})$ . Therefore we conclude that

$$\beta^{\prime*}(IC(\overline{E_{\mathbf{a}}^{\prime\prime}(\mathbf{c}^{\sharp})})) = \beta^{*}(IC(\overline{O_{\mathbf{b}}}) \otimes IC(E_{\varphi_{\ell[k]}})).$$

Now by definition

$$IC(\overline{O}_{\mathbf{b}}) \star IC(\overline{O}_{\ell[k]}) = \beta_*''(IC(\overline{E_{\mathbf{a}}''(\mathbf{c}^{\sharp})})).$$

# 6. Multisegments of Grassmanian Type

In order to precisely describe the previous corollary concerning Lusztig's product in the Grassmanian case in the next section, we generalize the construction in [6, §5.3] to get more general results concerning the the set  $S(\mathbf{a})$  for general multisegment  $\mathbf{a}$ .

Let V a  $\mathbb{C}$  vector space of dimension  $r + \ell$  and  $Gr_r(V)$  be the variety of r-dimensional subspaces of V.

**Definition 6.1.** By a partition of  $\ell$ , we mean a sequence  $\lambda = (\ell_1, \dots, \ell_r)$ for some r, where  $\ell_i \in \mathbb{N}$ ,  $0 \leq \ell_1 \leq \dots \ell_r \leq \ell$ . And for  $\mu = (\mu_1, \dots, \mu_s)$  be another partition, we say  $\mu \leq \lambda$  if and only if  $\mu_i \leq \lambda_i$ for all  $i = 1, \dots, Let \mathcal{P}(\ell, r)$  be the set of partitions of  $\ell$  into r parts.

**Definition 6.2.** We fix a complete flag

$$0 = V^0 \subset V^1 \subset \cdots \subset V^{r+\ell} = V.$$

This flag provides us a stratification of the variety  $Gr_r(V)$  by Schubert varieties, labeling by partitions, denoted by  $\overline{X}_{\lambda}$ ,

$$\overline{X}_{\lambda} = \{ U \in Gr_r(V) : \dim(U \cap V^{\ell_i + i}) \ge i, \text{ for all } i = 1, \cdots, r \}.$$

Lemma 6.3. (cf. [16]) We have

$$\mu \le \lambda \Longleftrightarrow \overline{X}_{\mu} \subseteq \overline{X}_{\lambda}.$$

And the Schubert cell

$$X_{\lambda} = \overline{X}_{\lambda} - \sum_{\mu < \lambda} \overline{X}_{\mu}$$

is open in  $\overline{X}_{\lambda}$ .

**Definition 6.4.** Let  $\Omega^{r,\ell}$  be the set

$$\Omega^{r,\ell} = \{ (a_1, \cdots, a_m; b_0, \cdots, b_{m-1}) : \sum_i a_i = r, \sum_j, b_j = \ell,$$
  
for  $0 < i < m, a_i > 0, b_i > 0 \}.$ 

Lemma 6.5. (cf. [16]) There exists a bijection

$$\Omega^{r,\ell} \to \mathcal{P}(\ell,r),$$

which sends  $(a_1, \dots, a_m; b_0, \dots, b_{m-1})$  to a partition of  $\ell$  given by  $b_0, b_0 + b_1, \dots, b_0 + \dots + b_{m-1}$ , and that the elements  $b_0 + \dots + b_{i-1}$  figures in  $\lambda$  with multiplicity  $a_i$ .

Notation 6.6. From now on, we will also write

$$\lambda = (a_1, \cdots, a_m; b_0, \cdots, b_{m-1}),$$

with notations as in the previous lemma.

We introduce the formula in [16] to calculate the Kazhdan Lusztig polynomials for Grassmannians.

**Definition 6.7.** Let  $\lambda = (a_1, \dots, a_m; b_0, \dots, b_{m-1})$  be a partition. Following [16], we represent a partition as a broken line in the plane (x, y), i.e., the graph of the piecewise-linear function  $y = \lambda(x)$  which equals |x| for large |x|, has everywhere slope  $\pm 1$ , and whose ascending and decreasing segments are precisely  $b_0, \dots, b_{m-1}$  and  $a_1, \dots, a_m$ , respectively. Moreover, we call the local maximum and minimum of the graph  $y = \lambda(x)$  the peaks and depressions of  $\lambda$ .



FIGURE 1.

**Lemma 6.8.** (cf. [16]) For  $\lambda, \mu \in \Omega^{r,\ell}$ , then

$$\lambda \ge \mu \iff \lambda(x) \ge \mu(x), \text{ for all } x.$$

From now on until the end of this section, let

$$J = \{\sigma_i : i = 1, \cdots, r - 1\} \cup \{\sigma_i : i = r + 1 \cdots, r + \ell - 1\},\$$

and fix

$$\mathbf{a} := \mathbf{a}_{\mathrm{Id}}^{J,\emptyset} = \{\Delta_1, \cdots, \Delta_r, \cdots, \Delta_{r+\ell}\}$$

to be a multisegment of parabolic type  $(J, \emptyset)$  (cf. [7, Definition 3.26]) such that

$$e(\Delta_i) = k - 1$$
, for  $i = 1, \cdots, r$ ,

and

$$e(\Delta_i) = k$$
, for  $i = r+1, \cdots, r+\ell$ .

**Definition 6.9.** Then to each partition  $\lambda \in \Omega^{r_1,\ell_1}$  such that  $r_1 \geq r$  and  $r_1 + \ell_1 = r + \ell$ , we associate

$$\mathbf{a}_{\lambda} = \sum_{i=1}^{b_0} [b(\Delta_i), k] + \sum_{i=b_0+1}^{b_0+a_1} [b(\Delta_i), k-1] + \cdots + \sum_{i=b_0+a_1\cdots+b_{j-1}+a_j}^{b_0+a_1\cdots+b_{j-1}+a_j} [b(\Delta_i), k-1] + \sum_{i=b_0+\cdots+b_{j-1}+a_j+1}^{b_0+a_1\cdots+b_j} [b(\Delta_i), k] + \cdots$$

**Definition 6.10.** Let  $r, n \in \mathbb{N}$  such that  $r \leq n$ . Let

 $R_{r}(n) = \{(x_{1}, \dots, x_{r}) : 1 \leq x_{1} < \dots < x_{r} \leq n\}.$ (1): Let  $x = (x_{1}, \dots, x_{r_{1}}) \in R_{r_{1}}(n)$  and  $x' = (x'_{1}, \dots, x'_{r_{2}}) \in R_{r_{2}}(n)$  such that  $r_{1} \geq r_{2}$ . We say  $x \supseteq x'$  if  $\{x_{1}, \dots, x_{r_{1}}\} \supseteq \{x'_{1}, \dots, x'_{r_{2}}\}.$ (2): Let  $x = (x_{1}, \dots, x_{r}) \in R_{r}(n)$  and  $x' = (x'_{1}, \dots, x'_{r}) \in R_{r}(n).$ We say  $x \ge x'$  if  $x_{i} \ge x'_{i}$  for all  $i = 1, \dots, r$ .
(3): We define  $x \succeq y$ , if  $x \ge y' \supseteq y$  for some y'.

*Remark:* The set  $R_r(n)$  is a poset with respect to the relation  $\geq$ , while the set  $\bigcup_{r \leq n} R_r(n)$  is a poset with respect to the relation  $\supseteq$ .

**Proposition 6.11.** For  $J = \{\sigma_i : i = 1, \dots, r-1\} \cup \{\sigma_i : i = r + 1, \dots, r+\ell-1\}$ , we have an isomorphism of posets

$$\varsigma_1: S_{r+\ell}^{J,\emptyset} \to R_r(r+\ell),$$

by associating the element w with  $x_w := (w^{-1}(1), \cdots, w^{-1}(r)).$ 

*Proof.* Note that by definition (cf. [7, Definition3.20]),

 $S_{r+\ell}^{J,\emptyset} = \{ w \in S_{r+\ell} : w^{-1}(1) < \dots < w^{-1}(r) \text{ and } w^{-1}(r+1) < \dots < w^{-1}(r+\ell) \}.$ 

Therefore,  $\varsigma$  is a bijection. This preserves the partial order by [3, Proposition 2.4.8].

**Definition 6.12.** For  $\lambda \in \Omega^{r,\ell}$  and  $\lambda' \in \Omega^{r_1,\ell_1}$  such that  $r + \ell = r_1 + \ell_1$ . We define  $\lambda \supseteq \lambda'$  if and only if  $x_\lambda \supseteq x_{\lambda'}$ , and  $\lambda \succeq \lambda'$  if and only if  $x_\lambda \succeq x_{\lambda'}$ .

**Definition 6.13.** Let  $\lambda = (a_1, \dots, a_m; b_0, \dots, b_{m-1})$ , consider the set  $\{b(\Delta) : \Delta \in \mathbf{a}_{\lambda}, e(\Delta) = k-1\} = \{x_1 < \dots < x_r\},\$ 

here we have r segments ending in k-1 since  $\sum_{i} a_{i} = r$ , we associate  $\lambda$  with the element

$$x_{\lambda} := (x_1, \cdots, x_r).$$

This allows us to get a morphism  $\varsigma_2 : \Omega^{r,\ell} \to R_r(r+\ell)$  sending  $\lambda$  to  $x_{\lambda}$ .

**Lemma 6.14.** The map  $\varsigma_2$  is an isomorphism of posets.

*Proof.* To see that  $\varsigma_2$  is a bijection, we only need to construct an inverse. Given  $x = (x_1, \dots, x_r) \in R_r(r+\ell)$ , we have  $y = (y_1, \dots, y_\ell) \in R_\ell(r+\ell)$ such that  $\{1, \dots, r+\ell\} = \{x_1, \dots, x_r, y_1, \dots, y_\ell\}$ . We can associate a multisegment to x:

$$\mathbf{a}_{x} = \sum_{j=1}^{r} [b(\Delta_{x_{j}}), k-1] + \sum_{j=1}^{\ell} [b(\Delta_{y_{j}}), k].$$

Note that this allows us to construct a partition  $\lambda(x) \in \Omega^{r,\ell}$  by counting the segments ending in k-1 and k respectively.

A simple calculation shows that if we write  $\lambda \in \Omega^{r,\ell}$  is identified via Lemma 6.5 with a partition  $(\ell_1, \dots, \ell_r)$  such that  $0 \leq \ell_1 \leq \dots \leq \ell_r$ , then

$$\varsigma_2(\lambda) = (\ell_1 + 1, \cdots, \ell_r + r).$$

It follows from [4, Page 6, Paragraph 3] that

$$\mu \ge \lambda \Leftrightarrow \varsigma_2(\mu) \ge \varsigma_2(\lambda).$$

**Proposition 6.15.** For  $\lambda \in \Omega^{r,\ell}$ , we have  $\mathbf{a}_{\lambda} \in S(\mathbf{a})$ , moreover, all the elements in  $S(\mathbf{a})$  are of this form. Moreover, we have  $S(\mathbf{a}_{\lambda}) = \{a_{\mu} : \mu \geq \lambda\}$ .

*Proof.* Let  $w \in S^{J,\emptyset}$ , by definition, we have

$$w^{-1}(1) < \dots < w^{-1}(r), \ w^{-1}(r+1) < \dots < w^{-1}(r+\ell).$$

By definition, we have

$$\Phi_{J,\emptyset}(w) = \sum_{j} [b(\Delta_{j}), e(\Delta_{w(j)})]$$
  
=  $\sum_{j} [b(\Delta_{w^{-1}(j)}), e(\Delta_{j})]$   
=  $\sum_{j=1}^{r} [b(\Delta_{w^{-1}(j)}), k-1] + \sum_{j=r+1}^{r+\ell} [b(\Delta_{w^{-1}(j)}), k]$   
=  $\mathbf{a}_{\varsigma_{2}^{-1}(x_{w})}$ 

Now that  $\varsigma_2^{-1} \circ \varsigma_1$  preserves the partial order, we have

$$S(\mathbf{a}_{\lambda}) = \{\mathbf{a}_{\mu} : \mu \ge \lambda\}$$

by [7, Proposition 3.18].

Example 6.16. For example, for  $r = 1, \ell = 3$ , with  $J = \{\sigma_2, \sigma_3\}$  and  $\mathbf{a} = \mathbf{a}_{\text{Id}}^{J,\emptyset} = [1, 4] + [2, 5] + [3, 5] + [4, 5].$ 

Let  $\lambda = (a_1, a_2; b_0, b_1) = (1, 0; 2, 1)$ , then  $\mathbf{a}_{\lambda} = [1, 5] + [2, 5] + [3, 4] + [4, 5]$ . This corresponds to the element  $\varsigma_1^{-1} \circ \varsigma_2(\lambda) = \sigma_1 \sigma_2$  in  $S_4^{J,\emptyset}$ .



FIGURE 2.

**Proposition 6.17.** Let  $\lambda, \mu \in \Omega^{r,\ell}$  such that  $\lambda < \mu$ . We have

$$P_{\mathbf{a}_{\lambda},\mathbf{a}_{\mu}}(q) = P_{\lambda,\mu}(q).$$

*Proof.* We can also prove this proposition in the following way. Let  $w, v \in S_{r+\ell}^{J,\emptyset}$ , such that

$$\lambda = \varsigma_2^{-1} \varsigma_1(w), \ \mu = \varsigma_2^{-1} \varsigma_1(v).$$

Let  $P_J$  be the parabolic subgroup of  $GL_n$ , then by fixing an element in  $V_0 \in Gr_r(\mathbb{C}^{r+\ell})$ , we can identify  $P_J \setminus GL_n$  with  $Gr_r(\mathbb{C}^{r+\ell})$ . Moreover, the *B*-orbits  $P_J \setminus wB$  corresponds to the varieties  $X_\lambda$ , see [4] for a precise description. Hence we have

$$P_{\lambda,\mu}(q) = P_{w,v}^{J,\emptyset}(q) = P_{a_{\lambda},\mathbf{a}_{\mu}}(q).$$

			٦
	_	_	

# **Definition 6.18.** Let $\lambda \in \Omega^{r,\ell}$ .

(1): We define  $\Gamma(\lambda) = \{ \mu \in \Omega^{r_1,\ell_1} : r_1 + \ell_1 = r + \ell, r_1 \ge r, \ \mu \succeq \lambda \}.$ and  $\Gamma^{\mu}(\lambda) = \{ \mu' : \mu \ge \mu', \mu' \succeq \lambda \},$   $\Gamma^{\mu}_1(\lambda) = \{ \mu' : \mu \ge \mu', \mu' \supseteq \lambda \}.$ (2): For  $\mu \in \Gamma(\lambda)$ , we define  $S^{\mu}(\lambda) = \{ \lambda' \in \Omega^{r,\ell} : \lambda' \ge \lambda, \mu \succeq \lambda' \},$ and let  $S^{\mu}(\lambda) = \{ \lambda' \in \Omega^{r,\ell} : \lambda' \ge \lambda, \mu \supseteq \lambda' \}.$ 

**Proposition 6.19.** Let  $\lambda \in \Omega^{r,\ell}$  and  $\mu \in \Omega^{r_1,\ell_1}$  with  $r_1 \geq r$  and  $r_1 + \ell_1 = r + \ell$ . Then  $\pi(\mathbf{a}_{\mu})$  appears as a summand of  $\mathscr{D}^k(\pi(\mathbf{a}_{\lambda}))$  if and only if  $\mu \in \Gamma(\lambda)$ .

*Proof.* Let  $x_{\lambda} = (x_1^{\lambda}, \dots, x_r^{\lambda}) = \varsigma_2(\lambda)$  and  $y_{\lambda} = (y_1^{\lambda}, \dots, y_{\ell}^{\lambda}) \in R_r(r+\ell)$  such that

$$\{1,\cdots,r+\ell\} = \{x_1^{\lambda},\cdots,x_r^{\lambda},y_1^{\lambda},\cdots,y_\ell^{\lambda}\}.$$

By Proposition 6.15, we have

$$\mathbf{a}_{\lambda} = \sum_{j=1}^{r} [b(\Delta_{x_j^{\lambda}}), k-1] + \sum_{j=1}^{\ell} [b(\Delta_{y_j^{\lambda}}), k].$$

Therefore

$$\mathscr{D}^k(\pi(\mathbf{a}_{\lambda})) = \pi(\mathbf{a}_{\lambda}) + \sum_{y \supseteq x_{\lambda}} \pi(\mathbf{a}_{\varsigma_2^{-1}(y)}).$$

By Lemma 2.5,  $\pi(\mu)$  is a summand of  $\mathscr{D}^k(\pi(\mathbf{a}_{\lambda}))$  if and only if  $\mu \geq \varsigma_2^{-1}(y)$  for some  $y \supseteq x_{\lambda}$ , i.e,  $\mu \succeq \lambda$ .

**Corollary 6.20.** We have  $\mu \succeq \lambda$  if and only if  $\mathbf{a}_{\mu} \preceq_{k} \mathbf{a}_{\lambda}$ .

*Proof.* By Corollary 2.8,  $\mathbf{a}_{\mu} \leq_k \mathbf{a}_{\lambda}$  if and only if  $\mathscr{D}^k(\pi(\mathbf{a}_{\lambda})) - \pi(\mathbf{a}_{\mu}) \geq 0$  in  $\mathcal{R}$ , which is equivalent to say that  $\mu \leq \lambda$  by the previous proposition.

**Proposition 6.21.** Let  $\lambda \in \Omega^{r,\ell}$  and  $\mu \in \Omega^{r_1,\ell_1}$ . Then we have  $\mathbf{a}_{\mu} = (\mathbf{a}_{\lambda})_{\Gamma}$  for some  $\Gamma \subseteq \mathbf{a}_{\lambda}(k)$ . if and only if we have  $\mu \supseteq \lambda$ .

*Proof.* Let  $x_{\lambda} = (x_1^{\lambda}, \cdots, x_r^{\lambda}) = \varsigma_2(\lambda)$  and  $y_{\lambda} = (y_1^{\lambda}, \cdots, y_{\ell}^{\lambda}) \in R_r(r+\ell)$  such that

$$1, \cdots, r+\ell\} = \{x_1^{\lambda}, \cdots, x_r^{\lambda}, y_1^{\lambda}, \cdots, y_\ell^{\lambda}\}.$$

By Proposition 6.15, we have

{

$$\mathbf{a}_{\lambda} = \sum_{j=1}^{r} [b(\Delta_{x_{j}^{\lambda}}), k-1] + \sum_{j=1}^{\ell} [b(\Delta_{y_{j}^{\lambda}}), k].$$

Hence

$$\mathbf{a}_{\lambda}(k) = \sum_{j=1}^{\ell} [b(\Delta_{y_j^{\lambda}}), k].$$

Let  $\Gamma = \sum_{m=1}^{t} [b(\Delta_{y_{j_m}^{\lambda}}), k]$ . If  $\mathbf{a}_{\mu} = (\mathbf{a}_{\lambda})_{\Gamma}$ , then

$$\mathbf{a}_{\mu} = \sum_{j=1}^{r} [b(\Delta_{x_{j}^{\lambda}}), k-1] + \sum_{m=1}^{r} [b(\Delta_{y_{j_{m}}^{\lambda}}), k-1] + \sum_{j \notin \{j_{1}, \cdots, j_{t}\}} [b(\Delta_{y_{j}^{\lambda}}), k].$$

Therefore

$$x_{\mu} \supseteq x_{\lambda}$$

as a set. The converse is also true.

#### TAIWANG DENG

## 7. GRASSMANIAN CASE

As before, let

$$J = \{\sigma_i : i = 1, \cdots, r - 1\} \cup \{\sigma_i : i = r + 1 \cdots, r + \ell - 1\},\$$

and

 $\mathbf{a} := \mathbf{a}_{\mathrm{Id}}^{J,\emptyset} = \{\Delta_1, \cdots, \Delta_r, \cdots, \Delta_{r+\ell}\}$ 

be a multisegment of parabolic type  $(J, \emptyset)$  (cf. [7, Definition 3.26]) such that

$$e(\Delta_i) = k - 1$$
, for  $i = 1, \dots, r_i$ 

and

$$e(\Delta_i) = k$$
, for  $i = r + 1, \cdots, r + \ell$ .

Moreover, for  $\lambda \in \mathcal{P}(\ell, r)$ , let  $x_{\lambda} = (x_1^{\lambda}, \cdots, x_r^{\lambda}) = \varsigma_2(\lambda) \in R_r(r+\ell)$ and  $y_{\lambda} = (y_1^{\lambda}, \cdots, y_{\ell}^{\lambda}) \in R_{\ell}(r+\ell)$  such that

$$\{1,\cdots,r+\ell\} = \{x_1^{\lambda},\cdots,x_r^{\lambda},y_1^{\lambda},\cdots,y_\ell^{\lambda}\}.$$

It follows from Proposition 6.15 that we have

$$\mathbf{a}_{\lambda} = \sum_{j=1}^{r} [b(\Delta_{x_{j}^{\lambda}}), k-1] + \sum_{j=1}^{\ell} [b(\Delta_{y_{j}^{\lambda}}), k].$$

Let  $0 < r_0 \le \ell$  and  $r_1 = r + r_0$ ,  $\ell_1 = \ell - r_0$ .

**Proposition 7.1.** Let  $\mu \in \mathcal{P}(\ell_1, r_1)$ . Then there exists  $\mu^{\flat} \in \mathcal{P}(\ell, r)$ , such that

$$\{\mathbf{b} \in S(\mathbf{a}) : \mathbf{a}_{\mu} \preceq_{k} \mathbf{b}\} = \{\mathbf{a}_{\lambda} : \lambda \in \mathcal{P}(\ell, r), \ \lambda \leq \mu^{\flat}\}.$$

More explicitly, if  $x_{\mu} = (x_1^{\mu}, \cdots, x_{r_1}^{\mu}) = \varsigma_2(\mu)$ , then

$$x_{\mu^{\flat}} = \varsigma_2(\mu^{\flat}) = (x_{r_0+1}^{\mu}, \cdots, x_{r_1}^{\mu}).$$

*Proof.* By Lemma 5.18, the set

$$\{\mathbf{b} \in S(\mathbf{a}) : \mathbf{a}_{\mu} \preceq_{k} \mathbf{b}\}$$

contains a unique minimal element  $\mathbf{a}_{\mu^{\flat}} \in S(\mathbf{a})$  for some  $\mu^{\flat} \in \mathcal{P}(\ell, r)$ . Therefore we have

$$\{\mathbf{b} \in S(\mathbf{a}) : \mathbf{a}_{\mu} \preceq_{k} \mathbf{b}\} = \{\mathbf{a}_{\lambda} : \lambda \in \mathcal{P}(\ell, r), \ \lambda \leq \mu^{\flat}\}.$$

Note that if we write

$$\mathbf{a}_{\mu} = \sum_{j=1}^{r_1} [b(\Delta_{x_j^{\mu}}), k-1] + \sum_{j=1}^{\ell_1} [b(\Delta_{y_j^{\mu}}), k],$$

then

$$\mathbf{a}_{\mu^{\flat}} = \sum_{j=1}^{r_0} [b(\Delta_{x_j^{\mu}}), k] + \sum_{j=r_0+1}^{r_1} [b(\Delta_{x_j^{\mu}}), k-1] + \sum_{j=1}^{\ell_1} [b(\Delta_{y_j^{\mu}}), k]$$

is the minimal element in  $S(\mathbf{a})$  satisfying

$$\mathbf{a}_{\mu} = (\mathbf{a}_{\mu^{\flat}})_{\Gamma}, \quad (cf. \text{ Definition } 2.2)$$

for some  $\Gamma \subseteq \mathbf{a}_{\mu^{\flat}}(k)$ .

Definition 7.2. Let

 $J_1 = \{\sigma_i : i = 1, \cdots, r-1\} \cup \{\sigma_i : i = r+1, \cdots, r_1-1\} \cup \{\sigma_i : r_1+1, \cdots, r+\ell-1\},\$ and

 $\mathbf{a}_1 =: \mathbf{a}_{\mathrm{Id}}^{J_1,\emptyset} = \{\Delta_1, \cdots, \Delta_{r_1}, \Delta_{r_1+1}^+, \cdots, \Delta_{r+\ell}^+\},\$ 

where  $\mathbf{a} = \{\Delta_1, \cdots, \Delta_{r+\ell}\}$  with  $\Delta_1 \leq \Delta_2 \leq \cdots \leq \Delta_r$  (cf. Definition 3.4).

**Lemma 7.3.** Let  $d = a + \ell_1[k+1]$ , then

- we have  $\mathbf{a} = \mathbf{a}_1^{(k+1)}$ ;
- and  $\mathfrak{X}_{\mathbf{d}} = \prod_{w \in S_{r+\ell}^{J_1,\emptyset}} O_{\mathbf{a}_w}$ , where  $\mathbf{a}_w = \mathbf{a}_w^{J_1,\emptyset} \in S(\mathbf{a}_1)$  is the element

associated to w by Lemma [7, Lemma 3.27].

*Proof.* Note that by definition we have

$$\mathbf{a} = \mathbf{a}_1^{(k+1)}$$

By definition  $\mathfrak{X}_{\mathbf{d}}$  consists of the orbits  $O_{\mathbf{c}}$  with  $\mathbf{c} \in S(\mathbf{d})$  such that  $\varphi_{e(\mathbf{c})}(k) + \ell_1 = \varphi_{e(\mathbf{a})}(k)$ , and the latter condition implies that there exists  $w \in S_{r+\ell}^{J_1,\emptyset}$  such that  $\mathbf{c} = \mathbf{a}_w^{J_1,\emptyset}$ .

**Proposition 7.4.** Let  $\mathbf{d} = \mathbf{a} + \ell_1[k+1]$  and  $W \subseteq V_{\varphi_{\mathbf{d}},k+1}$  such that  $\dim(W) = \ell_1$  (which implies that  $W = V_{\varphi_d,k+1}$ ). Then the composition of morphisms

$$\mathfrak{X}_{\mathbf{d}} = (\mathfrak{X}_{\mathbf{d}})_W \xrightarrow{p} E''_{\mathbf{a}} \xrightarrow{\beta''} E_{\varphi_{\mathbf{a}}},$$

sends  $O_{\mathbf{a}_w} \cap (\mathfrak{X}_{\mathbf{d}})_W$  to  $O_{\mathbf{a}_w^{(k+1)}}$ .

*Proof.* This is by definition.

**Proposition 7.5.** Let  $\mu \in \mathcal{P}(\ell_1, r_1)$  and  $x_{\mu} = \varsigma_2(\mu) = (x_1^{\mu}, \cdots, x_{r_1}^{\mu})$ ,  $y_{\mu} = (y_1^{\mu}, \cdots, y_{\ell_1}^{\mu})$  such that

$$\{1, \cdots, r_1 + \ell_1\} = \{x_1^{\mu}, \cdots, x_{r_1}^{\mu}, y_1^{\mu}, \cdots, y_{\ell_1}^{\mu}\}.$$

Then

$$(\mathbf{a}_{\mu^{\flat}})^{\sharp} = \sum_{j=1}^{r_0} [b(\Delta_{x_j^{\mu}}), k] + \sum_{j=r_0+1}^{r_1} [b(\Delta_{x_j^{\mu}}), k-1] + \sum_{j=1}^{\ell_1} [b(\Delta_{y_j^{\mu}}), k+1],$$

for definition of  $(\mathbf{a}_{\mu^{\flat}})^{\sharp}$ , cf. Lemma 5.18.

*Proof.* Note that by Proposition 7.1,

$$\mathbf{a}_{\mu^{\flat}} = \sum_{j=1}^{r_0} [b(\Delta_{x_j^{\mu}}), k] + \sum_{j=r_0+1}^{r_1} [b(\Delta_{x_j^{\mu}}), k-1] + \sum_{j=1}^{\ell_1} [b(\Delta_{y_j^{\mu}}), k]$$

and

$$\mathbf{a}_{\mu} = (\mathbf{a}_{\mu^{\flat}})_{\Gamma}$$

for 
$$\Gamma = \sum_{j=1}^{r_0} [b(\Delta_{x_j^{\mu}}), k]$$
. Now by construction in Lemma 5.18,  
 $r_0$   $r_1$   $\ell_1$ 

$$(\mathbf{a}_{\mu^{\flat}})^{\sharp} = \sum_{j=1}^{r_0} [b(\Delta_{x_j^{\mu}}), k] + \sum_{j=r_0+1}^{r_1} [b(\Delta_{x_j^{\mu}}), k-1] + \sum_{j=1}^{r_1} [b(\Delta_{y_j^{\mu}}), k+1].$$

# Proposition 7.6. We have

$$n(\mathbf{a}_{\mu}, \mathbf{a}_{\mu^{\flat}}) = \sharp \{ \mathbf{c} \in S(\mathbf{a}_1) : \mathbf{c}^{(k+1)} = \mathbf{a}_{\mu^{\flat}}, \mathbf{c} \ge (\mathbf{a}_{\mu^{\flat}})^{\sharp} \}.$$

Proof. Consider the composed morphism

$$h: \mathfrak{X}_{\mathbf{d}} = (\mathfrak{X}_{\mathbf{d}})_W \xrightarrow{p} E''_{\mathbf{a}} \xrightarrow{\beta''} E_{\varphi_{\mathbf{a}}},$$

then the orbits contained in  $h^{-1}(O_{\mathbf{a}_{\mu^\flat}})$  is indexed by the set

$$\{\mathbf{c} \in S(\mathbf{a}_1) : \mathbf{c}^{(k+1)} = \mathbf{a}_{\mu^\flat}, \mathbf{c} \ge (\mathbf{a}_{\mu^\flat})^\sharp\}$$

Note that by Theorem 5.19 and Theorem 4.8, the number

$$n(\mathbf{a}_{\mu}, \mathbf{a}_{\mu^{\flat}}) = \sum_{i} \dim \mathcal{H}^{2i}(\beta_{*}^{\prime\prime}(IC(\overline{E_{\mathbf{a}}^{\prime\prime}((\mathbf{a}_{\mu^{\flat}})^{\sharp}))))_{x}$$

for some  $x \in O_{\mathbf{a}_{\mu^{\flat}}}$ . Finally, note that the morphism  $\beta''$  is smooth when restricted to the variety  $\beta''^{-1}(O_{\mathbf{a}_{\mu^{\flat}}})$ . Moreover, the fibers are open in some Schubert variety, therefore, we are reduced to the counting of orbits.

# 8. PARABOLIC CASE

In this section, we deduce a formula for calculating the coefficient  $n(\mathbf{b}, \mathbf{a})$  in the Parabolic cases. Most results in this section are stated without proof since the proof are just direct generalizations of the Grassmanian case.

Let

$$J \subseteq S$$

be a subset of generators and

$$\mathbf{a} = \mathbf{a}_{\mathrm{Id}}^{J,\emptyset}$$

be some multisegment of parabolic type  $(J, \emptyset)$  associated to the identity satisfying that  $\ell_{\mathbf{a},k} \neq 0, \ell_{\mathbf{a},k+1} = 0$  (cf. [6, Notation 5.12]). Recall that

$$\ell_{\mathbf{a},k} = \sharp \{ \Delta \in \mathbf{a} : e(\mathbf{a}) = k \}.$$

In particular, we have

$$\ell_{\mathbf{a},k} = \ell_{e(\mathbf{a}),k}.$$

**Definition 8.1.** Let  $\mathbf{a}(k) = \{\Delta_1, \dots, \Delta_{\ell_k}\}$  with  $\Delta_1 \leq \dots \leq \Delta_{\ell_k}$  and  $r_0 \in \mathbb{N}$  with  $1 \leq r_0 \leq \ell_k$ . Then let

$$\mathbf{a}_{1} = (\mathbf{a} \setminus \mathbf{a}(k)) \cup \{\Delta \in \mathbf{a}(k) : \Delta \trianglelefteq \Delta_{\ell_{k}-r_{0}}\} \cup \{\Delta^{+} \in \mathbf{a}(k) : \Delta \trianglerighteq \Delta_{\ell_{k}-r_{0}+1}\},\\ \mathbf{a}_{2} = (\mathbf{a} \setminus \mathbf{a}(k)) \cup \{\Delta^{-} \in \mathbf{a}(k) : \Delta \trianglelefteq \Delta_{r_{0}}\} \cup \{\Delta \in \mathbf{a}(k) : \Delta \trianglerighteq \Delta_{r_{0}+1}\}$$

and  $J_i(r_0, k)(i = 1, 2)$  be a subset of S such that  $\mathbf{a}_i$  is a multisegment of parabolic type  $(J_i(r_0, k), \emptyset)$ . Moreover, let

$$\mathbf{a}_{\mathrm{Id}}^{J_i(r_0,k),\emptyset} = \mathbf{a}_i, \text{ for } i = 1, 2.$$

**Lemma 8.2.** Let  $\ell_1 = \ell_k - r_0$  and  $\mathbf{d} = \mathbf{a} + \ell_1[k+1]$ , then

• we have  $\mathbf{a} = \mathbf{a}_1^{(k+1)}$ ; • and  $\mathfrak{X}_{\mathbf{d}} = \prod_{w \in S_n^{J_1(r_0,k),\emptyset}} O_{\mathbf{a}_w}$ , where  $\mathbf{a}_w = \mathbf{a}_w^{J_1(r_0,k),\emptyset} \in S(\mathbf{a}_1)$  is the

element associated to w by [7, Lemma 3.27].

**Proposition 8.3.** Let  $w \in S_n^{J_2(r_0,k),\emptyset}$ . Then there exists  $w^{\flat} \in S_n^{J,\emptyset}$ , such that

$$\{\mathbf{b} \in S(\mathbf{a}) : \mathbf{a}_w \preceq_k \mathbf{b}\} = \{\mathbf{a}_v : v \in S_n^{J,\emptyset}, v \leq w^\flat\}.$$

More explicitly, if  $\mathbf{a}_w(k-1) = \{\Delta_1, \cdots, \Delta_{\ell_{k-1}}\}$  with  $\Delta_1 \leq \cdots \leq \Delta_{\ell_{k-1}}$ , then

$$\mathbf{a}_{w^{\flat}} = (\mathbf{a}_w \setminus \mathbf{a}_w(k-1)) \cup \{\Delta^+ \in \mathbf{a}_w(k-1) : \Delta \trianglelefteq \Delta_{r_0}\} \cup \{\Delta \in \mathbf{a}_w(k-1) : \Delta \trianglerighteq \Delta_{r_0+1}\}$$

**Proposition 8.4.** Let  $w \in S_n^{J_2(r_0,k),\emptyset}$ . Then

$$(\mathbf{a}_{w^{\flat}})^{\sharp} = (\mathbf{a}_{w^{\flat}} \setminus \mathbf{a}_{w}(k)) \cup \{\Delta^{+} : \Delta \in \mathbf{a}_{w}(k)\}$$

for definition of  $(\mathbf{a}_{w^{\flat}})^{\sharp}$ , cf. Lemma 5.18.

**Definition 8.5.** Let  $t_w \in S_n^{J_1(\ell_k - r_0, k), \emptyset}$  be the element such that

$$\mathbf{a}_{t_w} = (\mathbf{a}_{w^\flat})^\sharp.$$

**Proposition 8.6.** Let  $P_J$  and  $P_{J_1(\ell_k-r_0,k)}$  be the parabolic subgroups corresponding to J,  $J_1(\ell_k - r_0, k)$  respectively. Consider the natural morphism

$$\pi: P_{J_1(\ell_k - r_0, k)} \backslash GL_n \to P_J \backslash GL_n.$$

Then

$$n(\mathbf{a}_w, \mathbf{a}_v) = \sum_i \dim \mathcal{H}^{2i}(\pi_*(IC(\overline{P_{J_1(\ell_k - r_0, k)} t_w B})))_x$$

for some  $x \in P_J v B$ .

*Proof.* Consider the composed morphism

$$h: \mathfrak{X}_{\mathbf{d}} = (\mathfrak{X}_{\mathbf{d}})_{W} \xrightarrow{p} E_{\mathbf{a}}'' \xrightarrow{\beta''} E_{\varphi_{\mathbf{a}}}.$$

This proposition can be deduced in a similar fashion as in the symmetric cases [6, Corollary 4.15]. We omit the details.  $\Box$ 

## TAIWANG DENG

# 9. CALCULATION OF PARTIAL BZ OPERATOR

In this section, we restrict ourselves to the case of multisegment of parabolic type.

**Definition 9.1.** Let  $J_1 \subseteq J_2 \subseteq S$  be two subsets of generators of  $S_n$ . Let  $v \in S_n^{J_1,\emptyset}, w \in S_n^{J_2,\emptyset}$ , we define  $\theta_{J_2}^{J_1}(w,v)$  to be the multiplicities of  $IC(\overline{P_{J_2}wB})$  in  $\pi_*(IC(\overline{P_{J_1}vB}))$ , where

$$\pi: P_{J_1} \backslash GL_n \to P_{J_2} \backslash GL_n$$

be the canonical projection.

*Remark:* By [7, Proposition 2.20], in case when  $J_1 = \emptyset$ ,  $J_2 = \{s_i\}$  we have  $\theta_{J_2}^{J_1}(w, v) = \mu(s_i w, v)$  if  $\ell(v) \leq \ell(s_i v)$ , where  $\mu(x, y)$  is the coefficient of degree  $(\ell(y) - \ell(x) - 1)/2$  in  $P_{x,y}(q)$ .

**Proposition 9.2.** Let  $J \subseteq S$  be a subset of generators in  $S_n$ . Let  $k \in \mathbb{Z}$  and **a** be a multisegment satisfies all the assumptions we made before Definition 8.1. Then for any  $w \in S_n^{J,\emptyset}$ , we have

$$\mathscr{D}^{k}(L_{\Phi(w)}) = \sum_{r_{0}=0}^{\ell_{\mathbf{a},k}} \sum_{v \in S_{n}^{J_{2}(r_{0},k),\emptyset}} \theta_{J}^{J_{1}(\ell_{\mathbf{a},k}-r_{0},k)}(w,t_{v})L_{\Phi(v)}.$$

*Proof.* Note that by Proposition 2.6,

$$\mathscr{D}^k(\pi(\Phi(w))) = \sum_{\mathbf{b} \preceq_k \Phi(w)} n(\mathbf{b}, \mathbf{a}) L_{\mathbf{b}}.$$

Note that by Proposition 2.10,  $\mathbf{b} \leq_k \Phi(w)$  implies that

$$\mathbf{b} = \Phi(v),$$

for some  $v \in J_2(\ell_{\mathbf{a},k} - r_0, k)$ . Moreover, according to Proposition 8.6

$$n(\Phi(v), \Phi(w)) = \sum_{i} \dim \mathcal{H}^{2i}(\pi_*(IC(\overline{P_{J_1(\ell_{\mathbf{a},k}-r_0,k)}t_vB})))_a$$

for some  $x \in P_J w B$ . In fact, by the decomposition theorem, we have

$$\pi_*(IC(\overline{P_{J_1(\ell_{\mathbf{a},k}-r_0,k)}t_vB}) = \bigoplus_{u \in S_n^J} \oplus_i IC(\overline{P_J uB})^{h_i(u,t_v)}[d_u^i]$$
(10)

therefore

$$\theta_J^{J_1(\ell_{\mathbf{a},k}-r_0,k)}(u,t_v) = \sum_i h_i(u,t_v).$$

Furthermore, we denote

$$\theta_J^{J_1(\ell_{\mathbf{a},k}-r_0,k)}(u,t_v)(q) = \sum_i h_i(u,t_v)q^{-d_u^i/2}$$

By localizing at a point of  $P_J w B$  and applying proper base change, we get

$$\sum_{S_{J_1(\ell_{\mathbf{a},k}-r_0,k)}\setminus S_J\ni\rho} q^{\ell(\rho)} P^{J_1(\ell_{\mathbf{a},k}-r_0,k),\emptyset}_{\rho w,t_v}(q) = \sum_u \theta^{J_1(\ell_{\mathbf{a},k}-r_0,k)}_J(u,t_v)(q) P^{J,\emptyset}_{w,u}(q).$$
(11)

Here  $\ell(\rho)$  denotes the length function on symmetric group. Now we return to the formula

$$\pi(\Phi(w)) = \sum_{u} P_{w,u}^{J,\emptyset}(1) L_{\Phi(u)}.$$
 (12)

By induction, we can assume that for u > w, we have that  $L_{\Phi(v)}$  appears in  $\mathscr{D}^k(L_{\Phi(u)})$  with multiplicity  $\theta_J^{J_1(\ell_{\mathbf{a},k}-r_0,k)}(u,t_v)$ . Then by applying the derivation  $\mathscr{D}^k$  to (12), on the right hand side we get the multiplicity of  $L_{\Phi(v)}$  given by

$$x + \sum_{u > w} \theta_J^{J_1(\ell_{\mathbf{a},k} - r_0,k)}(u, t_v) P_{w,u}^{J,\emptyset}(1),$$

where x denotes the multiplicity of  $L_{\Phi(v)}$  in the operator  $\mathscr{D}^k(L_{\Phi(w)})$ . On the right hand side, applying [6, Corollary 5.38], we get

$$\sum_{S_{J_1(\ell_{\mathbf{a},k}-r_0,k)}\setminus S_J\ni\rho} P^{J_1(\ell_{\mathbf{a},k}-r_0,k),\emptyset}_{\rho w,t_v}(1).$$

Now compare with (11) to get  $x = \theta_J^{J_1(\ell_{\mathbf{a},k}-r_0,k)}(w,t_v)$ .

From now on we consider the operator  $\mathscr{D}^k(L_{\mathbf{c}})$  for a general multisegment  $\mathbf{c}$  such that  $\ell_{\mathbf{c},k} > 0$ .

**Proposition 9.6.** There exists a multisegment  $\mathbf{c}'$  which is of parabolic type  $(J_1(\mathbf{c}), \emptyset)(cf.$  [7, Definition 3.26]). and a sequence of integers  $k_1, \ldots, k_r, k_{r+1}, \ldots, k_{r+\ell}$  such that  $L_{\mathbf{c}}$  is the minimal degree term with multiplicity one in

$$^{k_1}\mathscr{D}\cdots {}^{k_r}\mathscr{D}\mathscr{D}^{k_{r+1}}\cdots \mathscr{D}^{k_{r+\ell}}(L_{\mathbf{c}'}),$$

and

$$\ell_{\mathbf{c}',i} = \ell_{\mathbf{c},i}, \quad \text{if } i \le k,$$
  
$$\ell_{\mathbf{c},k+1} = 0,$$
  
$$k_i > k+1, \quad \text{if } i > r.$$

*Proof.* Let  $i_0 = \min\{i : \ell_{b(\mathbf{c}),i} > 1\}$  and  $\Delta_0 = \max\{\Delta \in \mathbf{c} : b(\Delta) = i_0\}$ . Then replace all segments  $\Delta \in \mathbf{c}$  with  $b(\mathbf{c}) < i_0$  by  $^+\Delta$  and  $\Delta_0$  by  $^+\Delta$  to get a new multisegment  $\mathbf{c}_1$ . Let  $\{i \in b(\mathbf{c}) : i < i_0\} = \{j_1 < \cdots < j_r\}$ , then we have  $L_{\mathbf{c}}$  is the minimal degree terms in

$$^{j_1-1}\mathscr{D}\cdots ^{j_r-1}\mathscr{D}(L_{\mathbf{c}_1}),$$

#### TAIWANG DENG

Repeat this procedure to get  $\mathbf{c}_0$  together with a sequence of integers  $k_1, \dots, k_r$  such that  $L_{\mathbf{c}}$  is the minimal degree term with multiplicity one in

$$^{k_1} \mathscr{D} \cdots {}^{k_r} \mathscr{D}(L_{\mathbf{c}_0}).$$

Suppose that  $\ell_{\mathbf{c}_0,k+1} > 0$ . Now replace all segments  $\Delta$  in  $\mathbf{c}_0$  with  $e(\Delta) > k$  by  $\Delta^+$  to obtain  $\mathbf{c}'$ , we are done.

Definition 9.7. We define

$$\Gamma^{i}(\mathbf{a},k) = \{\mathbf{b} \in \Gamma(\mathbf{a},k) : \deg(\mathbf{b}) + i = \deg(\mathbf{a})\},\$$

where  $\Gamma(\mathbf{a}, k)$  is defined in Definition 2.12.

**Definition 9.8.** Let **a** be a multisegment and  $k, k_1 \in \mathbb{Z}$ . Then we define

$$\Gamma^{i}(\mathbf{a},k)_{k_{1}} = \{ \mathbf{b} \in \Gamma^{i}(\mathbf{a},k) : \mathbf{b} \in S(\mathbf{b})_{k_{1}}, \mathbf{b}^{(k_{1})} \in \Gamma^{i}(\mathbf{a}^{(k_{1})},k) \}, \\ \Gamma(\mathbf{a},k)_{k_{1}} = \bigcup_{i} \Gamma^{i}(\mathbf{a},k)_{k_{1}}.$$

More generally for a sequence of integers  $k_1, \dots, k_r$ , we define

 $\Gamma(\mathbf{a},k)_{k_1,\cdots,k_r} = \{ \mathbf{b} \preceq_k \mathbf{a} : \mathbf{b}^{(k_1,\cdots,k_{i-1})} \in \Gamma(\mathbf{a}^{(k_1,\cdots,k_{i-1})},k)_{k_i} \text{ for } 1 \leq i \leq r \}.$ Similarly, we can define  $_{k_1}\Gamma(\mathbf{a},k)$  and  $_{k_1,\cdots,k_r}\Gamma(\mathbf{a},k).$ 

*Remark:* We can also define  $_{k_{r+1},\cdots,k_{r+\ell}}(\Gamma(\mathbf{a},k)_{k_1,\cdots,k_r})$ , cf. [6, Definition 6.6].

**Lemma 9.9.** Let  $k_1 \neq k - 1$ , then the map

$$\psi_{k_1} : \Gamma(\mathbf{a}, k)_{k_1} \to \Gamma(\mathbf{a}^{(k_1)}, k)$$
$$\mathbf{b} \mapsto \mathbf{b}^{(k_1)}$$

is bijective.

*Proof.* In fact we have  $\Gamma^i(\mathbf{a}, k) = S(\mathbf{a}_i)$  where  $\mathbf{a}_i$  is constructed in the following way: let  $\mathbf{a}(k) = \{\Delta_1 \succeq \cdots \succeq \Delta_r\}$ , then

$$\mathbf{a}_i = (\mathbf{a} \setminus \mathbf{a}(k)) \cup \{\Delta_j^- : j \le i\} \cup \{\Delta_j : j > i\}.$$

Note that  $\Gamma^{i}(\mathbf{a}, k) = S(\mathbf{a}_{i})$ , which implies that we have

$$\Gamma^i(\mathbf{a},k)_{k_1} = S(\mathbf{a}_i)_{k_1}.$$

Finally, note that by [6, Proposition 5.39] we have a bijection

$$\psi_{k_1}: S(\mathbf{a}_i)_{k_1} \to S(\mathbf{a}_i^{(k_1)})$$

Note that  $k_1 \neq k - 1, k$  implies that  $\mathbf{a}_i^{(k_1)} \in \Gamma^i(\mathbf{a}^{(k_1)}, k)$  and  $\Gamma(\mathbf{a}^{(k_1)}, k) = \bigcup_i S(\mathbf{a}_i^{(k_1)}).$ 

If  $k_1 = k$ , then

$$\Gamma(\mathbf{a},k)_k = S(\mathbf{a})_k, \quad \Gamma(\mathbf{a}^{(k_1)},k) = S(\mathbf{a}^{(k)}).$$

**Lemma 9.10.** Let  $k_1, k \in \mathbb{Z}$  then the map

$$_{k_1}\psi:_{k_1}\Gamma(\mathbf{a},k)\to\Gamma(^{(k_1)}\mathbf{a},k)$$
  
 $\mathbf{b}\mapsto ^{(k_1)}\mathbf{b}$ 

is bijective.

*Proof.* If  $k_1 \neq k$ , the proof is the same as that of the previous lemma. Consider the case where  $k_1 = k$ . Let  $\mathbf{a}(k) = \{\Delta_1 \succeq \cdots \succeq \Delta_{r_0} \succ [k] = \cdots = [k]\}$ . Then for  $i \leq \ell_k$ , we have

$$\mathbf{a}_i = (\mathbf{a} \setminus \mathbf{a}(k)) \cup \{\Delta_j^- : j \le i\} \cup \{\Delta_j : j > i\}$$

where  $\Delta_j = [k]$  if  $j > r_0$ . And we have  $\Gamma^i(\mathbf{a}, k) = S(\mathbf{a}_i)$ . By definition, we have  $\mathbf{b} \in {}_k \Gamma^i(\mathbf{a}, k)$  if and only if

$$\mathbf{b} \in {}_k S(\mathbf{b}), \quad {}^{(k)}\mathbf{b} \in \Gamma^i({}^{(k)}\mathbf{a}, k).$$

Since  ${}^{(k)}\mathbf{a}(k) = \{\Delta_1, \cdots, \Delta_{r_0}\}$ , we know that for  $\mathbf{b} \in {}_k\Gamma^i(\mathbf{a}, k)$ , we must have  $i \leq r_0$ . Let

$$({}^{(k)}\mathbf{a})_i = ({}^{(k)}\mathbf{a} \setminus {}^{(k)}\mathbf{a}(k)) \cup \{\Delta_j^- : j \le i\} \cup \{\Delta_j : r_0 \ge j > i\}.$$

Then we have  $\Gamma^i({}^{(k)}\mathbf{a}, k) = S(({}^{(k)}\mathbf{a})_i)$  and

$$^{(k)}\mathbf{a}_i = (^{(k)}\mathbf{a})_i$$

Finally, we conclude that  $\mathbf{b} \in {}_{k}\Gamma^{i}(\mathbf{a}, k)$  if and only if  $\mathbf{b} \in {}_{k}S(\mathbf{a}_{i})$ . Since the map

$$_kS(\mathbf{a}_i) \to S(^{(k)}\mathbf{a}_i)$$

is bijective, we are done.

**Proposition 9.11.** Let  $\mathbf{b}, \mathbf{c}$  be two multisegments and  $k_1 \in \mathbb{Z}$  such that

$$\mathbf{b} = {}^{(k_1)} \mathbf{c}, \quad \mathbf{c} \in {}_{k_1}S(\mathbf{c}).$$

If we write

$$\mathscr{D}^{k}(L_{\mathbf{c}}) = L_{\mathbf{c}} + \sum_{\mathbf{d} \in \Gamma(\mathbf{c}, k) \setminus \{\mathbf{c}\}} \tilde{n}(\mathbf{d}, \mathbf{c}) L_{\mathbf{d}},$$
(13)

then

$$\mathscr{D}^{k}(L_{\mathbf{b}}) = L_{\mathbf{b}} + \sum_{\mathbf{d} \in_{k_{1}} \Gamma(\mathbf{c}, k) \setminus \{\mathbf{c}\}} \tilde{n}(\mathbf{d}, \mathbf{c}) L_{(k_{1})}_{\mathbf{d}}.$$

*Proof.* -Suppose that  $\deg(\mathbf{c}) = \deg(\mathbf{b}) + 1$ . In fact, by Corollary A.4, we have

$$^{k_1}\mathscr{D}(L_{\mathbf{c}}) = L_{\mathbf{c}} + L_{\mathbf{b}}$$

By applying the derivation  $\mathscr{D}^k$  and using the fact  $\mathscr{D}^k({}^{k_1}\mathscr{D}) = {}^{k_1}\mathscr{D}\mathscr{D}^k$ , we have

$$\mathscr{D}^{k}(L_{\mathbf{c}}) + \mathscr{D}^{k}(L_{\mathbf{b}}) = L_{\mathbf{c}} + L_{\mathbf{b}} + \sum_{\mathbf{d} \in \Gamma(\mathbf{c},k) \setminus \{\mathbf{c}\}} \tilde{n}(\mathbf{d},\mathbf{c})^{k_{1}} \mathscr{D}(L_{\mathbf{d}})$$

## TAIWANG DENG

By assumption that  $deg(\mathbf{b}) + 1 = deg(\mathbf{c})$ , we have

$${}^{k_1} \mathscr{D}(L_{\mathbf{d}}) = L_{\mathbf{d}} + L_{^{(k_1)}\mathbf{d}} \text{ or } L_{\mathbf{d}}$$

where  ${}^{k_1}\mathscr{D}(L_{\mathbf{d}}) = L_{\mathbf{d}} + L_{(k_1)_{\mathbf{d}}}$  if and only if  $\mathbf{d} \in {}_{k_1}S(\mathbf{d})$  and  $\deg({}^{(k_1)}\mathbf{d}) = \deg(\mathbf{d}) - 1$ . This is equivalent to say that  $\mathbf{d} \in {}_{k_1}\Gamma(\mathbf{a}, k)$ .

-For general case, consider

$$\{\Delta \in \mathbf{c} : b(\Delta) = k_1\} = \{\Delta_1 \succeq \cdots \succeq \Delta_r\}.$$

Now by Proposition A.1 and Proposition 2.6,

$$^{k_1}\mathscr{D}(L_{\mathbf{c}}) = L_{\mathbf{b}} + \sum_{\ell_{\mathbf{d},k_1} > \ell_{\mathbf{b},k_1}} \tilde{n}(\mathbf{d},\mathbf{c})L_{\mathbf{d}},$$

for some  $\tilde{n}(\mathbf{d}, \mathbf{c}) \in \mathbb{N}$ .

If  $k_1 \neq k$ , then We observe that for any **d** such that  $\ell_{\mathbf{d},k_1} > \ell_{\mathbf{b},k_1}$  and  $\mathbf{d}' \leq_k \mathbf{d}$ , we have

$$\ell_{\mathbf{d}',k_1} > \ell_{\mathbf{b},k_1}$$

which implies that  $L_{\mathbf{d}'}$  can not be a summand of  $\mathscr{D}^k(L_{\mathbf{b}})$ . Therefore

$$\mathscr{D}^k(L_{\mathbf{b}})$$

is the sum of all irreducible representations  $L_{\mathbf{d}''}$  contained in  $\mathscr{D}^k({}^{k_1}\mathscr{D})(L_{\mathbf{c}})$  satisfying

$$\ell_{\mathbf{d}'',k_1} = \ell_{\mathbf{b},k_1}$$

Applying the derivation  ${}^{k_1}\mathcal{D}$  to (13), we get

$$^{k_1}\mathscr{D}\mathscr{D}^k(L_{\mathbf{c}}) = {}^{k_1}\mathscr{D}(L_{\mathbf{c}}) + \sum_{\mathbf{d} \preceq_k \mathbf{c}} \tilde{n}(\mathbf{d}, \mathbf{c})({}^{k_1}\mathscr{D})(L_{\mathbf{d}}).$$

Note that in this case the sub-quotient of  ${}^{k_1}\mathscr{D}\mathscr{D}^k(L_c)$  consisting of irreducible representations  $L_{\mathbf{d}''}$  satisfying

$$\ell_{\mathbf{d}'',k_1} = \ell_{\mathbf{b},k_1}$$

is given by

$$L_{\mathbf{b}} + \sum_{\mathbf{d} \in_{k_1} \Gamma(\mathbf{c}, k) \setminus \{\mathbf{c}\}} \tilde{n}(\mathbf{d}, \mathbf{c}) L_{(k_1)_{\mathbf{d}}}$$

Compare the equation  ${}^{k_1}\mathscr{D}\mathscr{D}^k(L_{\mathbf{c}}) = \mathscr{D}^k({}^{k_1}\mathscr{D})(L_{\mathbf{c}})$  gives the results. If  $k_1 = k$ , consider

$$\{\Delta \in \mathbf{c} : b(\Delta) = k_1\} = \{\Delta_1 \succeq \cdots \succeq \Delta_r\}.$$

Let  $\mathbf{c}'$  be the multisegment obtained by replacing all segments  $\Delta$  in  $\mathbf{c}$  such that  $b(\Delta) < k_1$  by  $^+\Delta$ , and  $\Delta_1$  by  $^+\Delta_1$ . Then there exists

$$k_2 = k_1 - 1 > k_3 > \dots > k_r$$

such that

$$\mathbf{c} = {}^{(k_r, \cdots, k_2)} \mathbf{c}'.$$

and

$$\mathbf{b} = {}^{(k_r, \cdots, k_3, k_1, k_2, k_1)} \mathbf{c}'.$$

Let  $\mathbf{b}' = {}^{(k_1)} \mathbf{c}'$ , then by induction on  $f_{b(\mathbf{c})}(k)$ , we can assume that

$$\mathscr{D}^{k}(L_{\mathbf{b}'}) = L_{\mathbf{b}'} + \sum_{\mathbf{d} \in_{k} \Gamma(\mathbf{c}',k) \setminus \mathbf{c}'} \tilde{n}(\mathbf{d},\mathbf{c}') L_{(k)\mathbf{d}}.$$

Applying what we have proved before, we get

$$\mathscr{D}^{k}(L_{\mathbf{b}}) = L_{\mathbf{b}} + \sum_{\mathbf{d} \in_{k_{r}, \cdots, k_{3}, k, k_{2}, k} \Gamma(\mathbf{c}', k) \setminus \{\mathbf{c}'\}} \tilde{n}(\mathbf{d}, \mathbf{c}') L_{(k_{r}, \cdots, k_{3}, k, k_{2}, k)} \mathbf{d}.$$

Also, we have

$$\mathscr{D}^{k}(L_{\mathbf{c}}) = L_{\mathbf{c}} + \sum_{\mathbf{d} \in_{k_{r}, \cdots, k_{3}, k_{2}} \Gamma(\mathbf{c}', k) \setminus \{\mathbf{c}'\}} \tilde{n}(\mathbf{d}, \mathbf{c}') L_{(k_{r}, \cdots, k_{3}, k_{2})} \mathbf{d}.$$

Since for any multisegment  $\mathbf{d}$ , we have

 $^{(k,k_r,\cdots,k_3,k_2)}\mathbf{d} = ^{(k_r,\cdots,k_3,k,k_2,k)} \mathbf{d},$ 

it remains to show that

$$_{k_r,\cdots,k_3,k,k_2,k}\Gamma(\mathbf{c}',k) =_{k,k_r,\cdots,k_3,k_2} \Gamma(\mathbf{c}',k).$$

By definition and the following lemma, we can assume that r = 2. In this case we argue by contradiction. Suppose that  $\mathbf{d} \in_{k,k-1,k} \Gamma^i(\mathbf{c}',k)$ and  $\mathbf{d} \notin_{k,k-1} \Gamma(\mathbf{c}',k)$ , which is equivalent to say that  $\mathbf{d} \notin_{k,k-1} S(\mathbf{d})$ . Note that  $\mathbf{d} \notin_{k,k-1} S(\mathbf{d})$  implies that there exists two linked segments  $\{\Delta, \Delta'\}$ , such that

$$b(\Delta) = k, \quad b(\Delta') = k - 1.$$

Then  ${}^{(k-1,k)}\mathbf{d}$  contains the pair of segments  $\{-\Delta, -\Delta'\}$ . The fact that  ${}^{(k-1,k)}\mathbf{d} \in {}_kS({}^{(k-1,k)}\mathbf{d})$  implies that  ${}^{-}\Delta' = \emptyset$ , i.e.  $\Delta' = [k-1]$ . However, this implies that  ${}^{(k,k-1,k)}\mathbf{d} \notin \Gamma^i({}^{(k,k-1,k)}\mathbf{c}',k)$  since  $\deg({}^{(k,k-1,k)}\mathbf{d}) + i = \deg({}^{(k,k-1,k)}\mathbf{a}) + 1$ , which is a contradiction.

Conversely, assume that  $\mathbf{d} \in {}_{k,k-1}\Gamma(\mathbf{c}',k)$  and  $\mathbf{d} \notin {}_{k,k-1,k}\Gamma^{i}(\mathbf{c}',k)$ , which by definition is equivalent to  $\mathbf{d} \notin {}_{k,k-1,k}S(\mathbf{d})$ . Note that  $\mathbf{d} \notin {}_{k,k-1,k}S(\mathbf{d})$  implies that  $\mathbf{d} \notin {}_{k}S(\mathbf{d})$ , which contradicts to  $\mathbf{d} \in {}_{k,k-1}S(\mathbf{d})$ .

**Lemma 9.13.** Let k > k - 1 > k' be two integers. Then for any multisegment  $\mathbf{c}$ , we have

$$_{k,k'}\Gamma(\mathbf{c},k) = _{k',k}\Gamma(\mathbf{c},k).$$

*Proof.* Note that since for any multisegment **d** 

$$^{(k',k)}\mathbf{d} = {}^{k,k'}\mathbf{d}$$

the fact

$$_{k,k'}\Gamma(\mathbf{c},k) = {}_{k',k}\Gamma(\mathbf{c},k)$$

is equivalent to

$$\mathbf{d} \in {}_{k,k'}S(\mathbf{d}) \Leftrightarrow \mathbf{d} \in {}_{k',k}S(\mathbf{d})$$

for all  $\mathbf{d} \in {}_{k,k'}\Gamma(\mathbf{c},k)$ . But for any multisegment  $\mathbf{d}$  and k > k-1 > k', we have

$$\mathbf{d} \in {}_{k,k'}S(\mathbf{d}) \Leftrightarrow \mathbf{d} \in {}_{k',k}S(\mathbf{d})$$

Hence we are done.

**Proposition 9.14.** Let  $k_1 \neq k - 1, k, k + 1$ . Let **b**, **c** be two multisegments such that

$$\mathbf{b} = \mathbf{c}^{(k_1)}, \quad \mathbf{c} \in S(\mathbf{c})_{k_1}.$$

If we write

$$\mathscr{D}^{k}(L_{\mathbf{c}}) = L_{\mathbf{c}} + \sum_{\mathbf{d} \in \Gamma(\mathbf{c}, k) \setminus \{\mathbf{c}\}} \tilde{n}(\mathbf{d}, \mathbf{c}) L_{\mathbf{d}},$$
(14)

then

$$\mathscr{D}^{k}(L_{\mathbf{b}}) = L_{\mathbf{b}} + \sum_{\mathbf{d} \in \Gamma(\mathbf{c},k)_{k_{1}} \setminus \{\mathbf{c}\}} \tilde{n}(\mathbf{d},\mathbf{c}) L_{d^{(k_{1})}}.$$

*Proof.* The proof is the same as Proposition 9.11.

Now let  $\mathbf{c}' = \Phi(w)$  for some  $w \in S_n^{J,\emptyset}$ .

Corollary 9.16. We have

$$\mathscr{D}^{k}(L_{\mathbf{a}}) = \sum_{r_{0}=0}^{\ell_{\mathbf{a},k}} \sum_{v \in S_{n}^{J_{2}(r_{0},k),\emptyset}, \Phi(v) \in k_{1}, \dots, k_{r}(\Gamma(\Phi(w),k)_{k_{r+1},\dots,k_{r+\ell})} \\ \theta_{J}^{J_{1}(\ell_{k}-r_{0},k)}(w,t_{v}) L_{(k_{1},\dots,k_{r})\Phi(v)}^{(k_{r+1},\dots,k_{r+\ell})}$$

Notation 9.17. For  $\mathbf{b} \leq_k \mathbf{a}$ , we denote

$$\theta_k(\mathbf{b}, \mathbf{a}) = \theta_J^{J_1(\ell_k - r_0, k)}(w, t_v)$$

if  $\mathbf{b} = {}^{(k_1, \cdots, k_r)} \Phi(v)^{(k_{r+1}, \cdots, k_{r+\ell})}$ . Otherwise, put  $\theta_k(\mathbf{b}, \mathbf{a}) = 0$ .

*Remark:* The same way we define  $_k\theta(\mathbf{b},\mathbf{a})$  by the formula

$$({}^{k}\mathscr{D})(L_{\mathbf{a}}) = \sum_{\mathbf{b}} {}_{k}\theta(\mathbf{b}, \mathbf{a})L_{\mathbf{b}}$$

And let

 $\Gamma(k, \mathbf{a}) = \{ \mathbf{b} : {}_k\theta(\mathbf{b}, \mathbf{c}) \neq 0 \text{ for some } \mathbf{c} \in S(\mathbf{a}) \},\$ 

it shares similar properties with  $\Gamma(\mathbf{a}, k)$ .

Appendix A. Minimal Degree Terms in Partial BZ operator

**Proposition A.1.** (i): Suppose that a satisfies the hypothesis  $H_k(\mathbf{a})$  (cf. [6, Definition 5.3]).

Then  $\mathscr{D}^k(L_{\mathbf{a}})$  contains in  $\mathcal{R}$  a unique irreducible representation of minimal degree, which is  $L_{\mathbf{a}^{(k)}}$ , and it appears with multiplicity one.

44

(ii): If **a** fails to satisfy the hypothesis  $H_k(\mathbf{a})$ , then  $L_{\mathbf{a}^{(k)}}$  will not appear in  $\mathscr{D}^k(L_{\mathbf{a}})$ , and the irreducible representations appearing are all of degree  $> \deg(\mathbf{a}^{(k)})$ .

*Proof.* Let  $\mathbf{a} = \{\Delta_1 \preceq \cdots \preceq \Delta_r\}$ , such that

$$e(\Delta_1) \leq \cdots < e(\Delta_i) = \cdots = e(\Delta_j) < \cdots \leq e(\Delta_r),$$

with  $k = e(\Delta_i)$ .

We prove the proposition by induction on  $\ell(\mathbf{a})$  (cf. Definition 2.9). For,  $\ell(\mathbf{a}) = 0$ , which means that  $\mathbf{a} = \mathbf{a}_{\min}$ , in this case  $\mathbf{a}$  satisfies the  $H_k(\mathbf{a})$ , and

$$\mathscr{D}^{k}(L_{\mathbf{a}}) = \mathscr{D}^{k}(\pi(\mathbf{a})) = \Delta_{1} \times \cdots \times (\Delta_{i} + \Delta_{i}^{-}) \times \cdots \times (\Delta_{j} + \Delta_{j}^{-}) \times \cdots$$

which contains

$$L_{\mathbf{a}^{(k)}} = \pi(\mathbf{a}^{(k)}) = \Delta_1 \times \cdots \times \Delta_i^- \times \cdots \Delta_j^- \times \cdots$$

Hence we are done in this case.

Next we consider general **a**. We write

$$\pi(\mathbf{a}) = L_{\mathbf{a}} + \sum_{\mathbf{b} < \mathbf{a}} m(\mathbf{b}, \mathbf{a}) L_{\mathbf{b}}.$$
 (15)

Now applying  $\mathscr{D}^k$  to both sides and consider only the lowest degree terms, on the left hand side, we get

$$\pi(\mathbf{a}^{(k)}) = \Delta_1 \times \cdots \times \Delta_{i-1} \times \Delta_i^- \times \cdots \times \Delta_j^- \times \cdots \Delta_r.$$
(16)

By [6, Theorem 2.22], both sides are a nonnegative sum of irreducible representations, then

• If a satisfies the hypothesis  $H_k(\mathbf{a})$ , on the right hand side by combining [6, Lemma 5.5] and induction, we know that for all  $\mathbf{b} < \mathbf{a}, \mathscr{D}^k(L_{\mathbf{b}})$  does not contain  $L_{\mathbf{a}^{(k)}}$  as subquotient. Hence  $\mathscr{D}^k(L_{\mathbf{a}})$  must contain  $L_{\mathbf{a}^{(k)}}$  with multiplicity one. We have to show that it does not contain other subquotients of  $\pi(\mathbf{a}^{(k)})$ . Note that by induction, we have the following formula

$$\pi(\mathbf{a}^{(k)}) = X + \sum_{\mathbf{c} \in S(\mathbf{a})_k \setminus \mathbf{a}} m(\mathbf{c}, \mathbf{a}) L_{\mathbf{c}^{(k)}},$$

where X denotes the minimal degree terms in  $\mathscr{D}^k(L_{\mathbf{a}})$ . Now apply [6, Corollary 5.40], we conclude that  $X = L_{\mathbf{a}^{(k)}}$ .

• Now if **a** fails to satisfy the hypothesis  $H_k(\mathbf{a})$ ,  $\mathbf{a} \notin S(\mathbf{a})_k$ , combining [6, Proposition 5.39] and induction, we know that there exists  $\mathbf{b} \in S(\mathbf{a})_k$ , such that  $\mathbf{a}^{(k)} = \mathbf{b}^{(k)}$  and  $\mathscr{D}^k(L_{\mathbf{b}})$  contains  $L_{\mathbf{a}^{(k)}}$  as a subquotient with multiplicity one.

Now by the Lemma 2.5,  $\pi(\mathbf{a}) - \pi(\mathbf{b})$  is a nonnegative sum of irreducible representations which contain  $L_{\mathbf{a}}$ : by the positivity of partial BZ operator, we obtain a nonnegative sum of

#### TAIWANG DENG

irreducible representations after applying  $\mathscr{D}^k$ . Now

$$\mathscr{D}^k(\pi(\mathbf{a}) - \pi(\mathbf{b})) = \pi(\mathbf{a}^{(k)}) - \pi(\mathbf{b}^{(k)}) + \text{ higher degree terms}$$

contains only terms of degree > deg( $\mathbf{a}^{(k)}$ ), so does  $\mathscr{D}^k(L_{\mathbf{a}})$ . This finishes our arguments.

**Corollary A.4.** Let **a** be a multisegment such that  $\varphi_{e(\mathbf{a})}(k) = 1$ . Then

- If a ∈ S(a)<sub>k</sub>, then D<sup>k</sup>(L<sub>a</sub>) = L<sub>a</sub> + L<sub>a<sup>(k)</sup></sub>.
   If a ∉ S(a)<sub>k</sub>, then D<sup>k</sup>(L<sub>a</sub>) = L<sub>a</sub>.

*Proof.* First of all, we observe that the highest degree term in  $\mathscr{D}^k(L_{\mathbf{a}})$ is given by  $L_{\mathbf{a}}$ . In fact, we have

$$\mathscr{D}^{k}(\pi(\mathbf{a})) = \mathscr{D}^{k}(L_{\mathbf{a}}) + \sum_{\mathbf{b} < \mathbf{a}} m(\mathbf{b}, \mathbf{a}) \mathscr{D}^{k}(L_{\mathbf{b}}),$$

meanwhile we have

$$\mathscr{D}^k(\pi(\mathbf{a})) = \pi(\mathbf{a}) + \text{ lower terms.}$$

By induction on  $\ell(\mathbf{a})$  we conclude that the highest degree terms in  $\mathscr{D}^k(L_{\mathbf{a}})$  is  $L_{\mathbf{a}}$ .

If  $\mathbf{a} \in S(\mathbf{a})_k$ , then Proposition A.1 implies that the minimal degree term of  $\mathscr{D}^k(L_{\mathbf{a}})$ , but since  $\deg(\mathbf{a}^{(k)}) = \deg(\mathbf{a}) - 1$ , therefore we must have

$$\mathscr{D}^{k}(L_{\mathbf{a}}) = L_{\mathbf{a}} + L_{\mathbf{a}^{(k)}}$$

On the contrary, if  $\mathbf{a} \notin S(\mathbf{a})_k$ , then by (ii) of the Proposition A.1, we know that all irreducible representations appearing in  $\mathscr{D}^k(L_{\mathbf{a}})$  are of degree > deg( $\mathbf{a}^{(k)}$ ) = deg( $\mathbf{a}$ ) - 1, which implies

$$\mathscr{D}^k(L_{\mathbf{a}}) = L_{\mathbf{a}}.$$

## References

- [1] Arkady Berenstein and Andrei Zelevinsky. String bases for quantum groups of type  $A_r$ . In I. M. Gelfand Seminar, volume 16, Part 1 of Adv. Soviet Math., pages 51-89. Amer. Math. Soc., Providence, RI, 1993.
- [2] I. N. Bernstein and A. V. Zelevinsky. Induced representations of reductive p-adic groups. I. Ann. Sci. École Norm. Sup. (4), 10(4):441-472, 1977.
- [3] Anders Björner and Francesco Brenti. Combinatorics of Coxeter groups, volume 231 of Graduate Texts in Mathematics. Springer, New York, 2005.
- [4] Michel Brion. Lectures on the geometry of flag varieties. In Topics in cohomological studies of algebraic varieties, Trends Math., pages 33-85. Birkhäuser, Basel, 2005.
- [5] Kei Yuen Chan and Kayue Daniel Wong. On the Lefschetz principle for  $GL(n, \mathbb{C})$  and  $GL(m, \mathbb{Q}_p)$ . arXiv:2305.15766, 2023.
- [6] Taiwang Deng. Study of multiplicities in induced representations of  $GL_n$ through a symmetric reduction. Manuscripta Math., 171(1-2):23–72, 2023.

- [7] Taiwang Deng. Some geometric consequences of the symmetric reduction. arXiv preprint arXiv:2404.00759, 2024.
- [8] M. Kashiwara. On crystal bases of the Q-analogue of universal enveloping algebras. Duke Math. J., 63(2):465-516, 1991.
- [9] Bernard Leclerc. Dual canonical bases, quantum shuffles and q-characters. Math. Z., 246(4):691-732, 2004.
- [10] Bernard Leclerc, Maxim Nazarov, and Jean-Yves Thibon. Induced representations of affine Hecke algebras and canonical bases of quantum groups. In *Studies in memory of Issai Schur (Chevaleret/Rehovot, 2000)*, volume 210 of *Progr. Math.*, pages 115–153. Birkhäuser Boston, Boston, MA, 2003.
- [11] Bernard Leclerc, Jean-Yves Thibon, and Eric Vasserot. Zelevinsky's involution at roots of unity. J. Reine Angew. Math., 513:33–51, 1999.
- [12] G. Lusztig. Canonical bases arising from quantized enveloping algebras. J. Amer. Math. Soc., 3(2):447–498, 1990.
- [13] Jean-Pierre Serre. Espaces fibrés algébriques (d'après André Weil). In Séminaire Bourbaki, Vol. 2, pages Exp. No. 82, 305–311. Soc. Math. France, Paris, 1995.
- [14] Deng Taiwang. Parabolic induction and geometry of orbital varieties for GL(n). arXiv preprint arXiv:1603.06387, 2016.
- [15] A. V. Zelevinskii. The p-adic analogue of the Kazhdan-Lusztig conjecture. Funktsional. Anal. i Prilozhen., 15(2):9–21, 96, 1981.
- [16] A. V. Zelevinskii. Small resolutions of singularities of Schubert varieties. Funktsional. Anal. i Prilozhen., 17(2):75–77, 1983.
- [17] A. V. Zelevinsky. Induced representations of reductive p-adic groups. II. On irreducible representations of GL(n). Ann. Sci. École Norm. Sup. (4), 13(2):165–210, 1980.

BEIJING INSTITUTE OF MATHEMATICAL SCIENCES AND APPLICATIONS (BIMSA), TRAFFIC LIGHT GATE, NO. 544 HEFANGKOU VILLAGE, HUAIROU DISTRICT, BEIJING

Email address: dengtaiw@bimsa.cn