

Extremal minimal bipartite matching covered graphs

Amit Kumar Mallik *

Indian Institute of Technology Bombay
amit.km287@gmail.com

Ajit A. Diwan

Indian Institute of Technology Bombay
aad@cse.iitb.ac.in

Nishad Kothari[†]

Indian Institute of Technology Madras
nishadkothari@gmail.com

April 12, 2024

Dedicated to Matching Theory by László Lovász and Michael D. Plummer

Abstract

A connected graph, on four or more vertices, is *matching covered* if every edge is present in some perfect matching. An ear decomposition theorem (similar to the one for 2-connected graphs) exists for bipartite matching covered graphs due to Hetyei. From the results and proofs of Lovász and Plummer, that rely on Hetyei's theorem, one may deduce that any minimal bipartite matching covered graph has at least $2(m - n + 2)$ vertices of degree two (where *minimal* means that deleting any edge results in a graph that is not matching covered); such a graph is said to be *extremal* if it attains the stated lower bound.

In this paper, we provide a complete characterization of the class of extremal minimal bipartite matching covered graphs. In particular, we prove that every such graph G is obtained from two copies of a tree devoid of degree two vertices, say T and T' , by adding edges — each of which joins a leaf of T with the corresponding leaf of T' .

Apart from the aforementioned bound, there are four other bounds that appear in, or may be deduced from, the work of Lovász and Plummer. Each of these bounds leads to a notion of extremality. In this paper, we obtain a complete characterization of all of these extremal classes and also establish relationships between them. Two of our characterizations are in the same spirit as the one stated above. For the remaining two extremal classes, we reduce each of them to one of the already characterized extremal classes using standard matching theoretic operations.

*The results herein shall appear in the Master's Thesis of the author.

[†]Supported by IC&SR IIT Madras.

Contents

1	Introduction and summary	2
1.1	Minimality, bounds and corresponding notions of extremality	3
1.2	Characterizations of $\mathcal{H}_2, \mathcal{H}_3$ and \mathcal{H}_4 using halin trees	4
1.3	Proofs of easier implications	6
1.4	Characterizations of \mathcal{H}_0 and \mathcal{H}_1	8
1.5	Containment poset for the extremal classes	10
1.6	Organization of the rest of this paper	11
2	Ear decomposition theory	11
3	Lower bounds established by Lovász and Plummer	12
4	Properties of extremal graphs	13
4.1	Balanced 2-cut property	15
5	An induction tool using balanced 2-cuts	17
6	Main Theorem: Characterization of \mathcal{H}_2	20
7	Other notions of extremality	21
7.1	Characterization of \mathcal{H}_3	22
7.2	Characterization of \mathcal{H}_4	22
7.3	Characterization of \mathcal{H}_0	23
7.4	Characterization of \mathcal{H}_1	26

1 Introduction and summary

All graphs considered here are loopless; however, we allow parallel/multiple edges. For notation and terminology, we largely follow Bondy and Murty [1]. For a graph $G := (V, E)$, its *order* is the number of vertices denoted by n , and its *size* is the number of edges denoted by m . We use the notation $G[A, B]$ to denote a bipartite graph with specified color classes A and B .

A graph is *matchable* if it has a perfect matching and an edge is *matchable* if it belongs to some perfect matching. A subgraph H of a graph G is *conformal* if $G - H$ is matchable. A connected graph of order four or more is *matching covered* if every edge is matchable. There is an elegant ear decomposition theory for the class of bipartite matching covered graphs, due to Hetyei [5], that may be viewed as a refinement of the more well-known ear decomposition theory for the larger class of 2-connected graphs due to Whitney [11]. We describe this below.

An *ear of H in G* is an odd path of G whose ends are in H but is otherwise disjoint from H . For a bipartite graph G , a sequence of subgraphs (G_0, G_1, \dots, G_r) is an *ear decomposition of G* if: (i) $C := G_0$ is an (even) cycle, (ii) $G_{i+1} = G_i \cup P_{i+1}$ where P_{i+1} is an ear of G_i for

each $i \in \{0, 1, \dots, r - 1\}$, and (iii) $G_r = G$. It is easy to observe that each subgraph G_i is a conformal matching covered subgraph of G , and that $r = m - n$. We refer to (C, P_1, \dots, P_r) as the *associated ear sequence*, or simply an *ear sequence* of G . Figure 1 shows a bipartite matching covered graph and an ear sequence for the same.



Figure 1: a bipartite matching covered graph and its ear decomposition

The following is the aforementioned result by Heteyi that we will find useful in order to establish that certain bipartite graphs are indeed matching covered.

Theorem 1.1. [EAR DECOMPOSITION THEOREM]

A bipartite graph is matching covered if and only if it admits an ear decomposition.

There is also a generalization of the above theorem for nonbipartite matching covered graphs due to Lovász and Plummer [9]. However, we do not describe this here since our work focuses on bipartite matching covered graphs.

1.1 Minimality, bounds and corresponding notions of extremality

A matching covered graph is *minimal* if deletion of any edge results in a graph that is not matching covered. We use \mathcal{H} to denote the class of minimal bipartite matching covered graphs. Using the ear decomposition theory, Lovász and Plummer [9] proved that each member of \mathcal{H} has at least $m - n + 2$ pairwise nonadjacent 2-edges — where a *2-edge* is an edge whose each end has degree two. Thus, such a graph has at least $2(m - n + 2)$ degree two vertices. Furthermore, for each of these invariants, one may also deduce lower bounds solely in terms of n ; in particular, that each member of \mathcal{H} has at least $\frac{n+10}{6}$ 2-edges and at least $\frac{n}{2} + 2$ vertices of degree two; see Corollaries 7.9 and 7.1. They also proved an upper bound — namely, that each member of \mathcal{H} , distinct from C_4 , has at most $\frac{3n-6}{2}$ edges.

Each of the bounds stated in the above paragraph leads to a notion of extremality for minimal bipartite matching covered graphs. As we have five different bounds, we have five notions of extremality as defined in Table 1. For instance, \mathcal{H}_2 denotes the class of *2-vertex extremal* minimal bipartite matching covered graphs — that is, those members of \mathcal{H} that satisfy $|V_2| = 2(m - n + 2)$ where V_2 is the set of degree two vertices. Likewise, we use E_2 to denote the set of 2-edges.

In this paper, we completely characterize all of the five classes of “extremal” minimal bipartite matching covered graphs in terms of special trees. Our characterizations are reminiscent of similar characterization(s) of “extremal” minimal 2-connected graphs — where minimality is defined with respect to edge deletion; we briefly describe some of these results below before stating our characterizations.

Class	Property	Notation
<i>2-edge extremal</i>	$ E_2 = m - n + 2$	\mathcal{H}_0
<i>2-edge n-extremal</i>	$ E_2 = \frac{n+10}{6}$	\mathcal{H}_1
<i>2-vertex extremal</i>	$ V_2 = 2(m - n + 2)$	\mathcal{H}_2
<i>2-vertex n-extremal</i>	$ V_2 = \frac{n}{2} + 2$	\mathcal{H}_3
<i>edge extremal</i>	$ E = \frac{3n-6}{2}$	\mathcal{H}_4

Table 1: Definitions of different notions of extremality within \mathcal{H}

Dirac [3] proved that a minimal 2-connected graph has at least $\frac{n+4}{3}$ vertices of degree two. A minimal 2-connected graph is *extremal* if it satisfies this lower bound with equality. Oxley [10] gave a generation theorem for extremal minimal 2-connected graphs. Karpov [7] then characterized this class of graphs in terms of special trees. In particular, they proved that a graph is extremal minimal 2-connected if and only if it can be obtained from two copies of a tree, each of whose non-leaves is of degree three, by identifying the corresponding leaves as per some fixed isomorphism; see Figure 2 for an example.




Figure 2: illustration of Karpov’s characterization of extremal minimal 2-connected graphs

A tree is said to be a *halin tree* if all of its non-leaves have degree at least three, or equivalently, if it has no vertices of degree two; see Figure 3 for an example. We call them halin trees because a *Halin graph* is a (planar) graph that is obtained from a planar embedding of such a tree by adding a cycle each of whose edges joins two leaves that appear consecutively in the cyclic order (as per the planar embedding). Halin graphs, first introduced by Halin [4], as examples of minimal 3-connected graphs, have been studied extensively in the literature. Halin trees are precisely the “homeomorphically irreducible” trees; finding such spanning trees in a cubic graph is a well-studied problem; see [6]. A halin tree is *cubic* if the corresponding Halin graph is cubic. To put it differently, a tree is a cubic halin tree if all of its non-leaves have degree exactly three; see Figure 2 for an example. Thus, the aforementioned Karpov’s characterization of extremal minimal 2-connected graphs draws a bijection between this graph class and cubic halin trees.

1.2 Characterizations of \mathcal{H}_2 , \mathcal{H}_3 and \mathcal{H}_4 using halin trees

We are now ready to state our characterizations of “extremal” minimal bipartite matching covered graphs — that are similar to Karpov’s characterization. However, in our case,

identifying the corresponding leaves is not the correct operation.

For two disjoint copies of a halin tree, say T and T' , let H be the (bipartite) graph obtained from $T \cup T'$ by adding a matching each of whose edges joins a leaf of T with the corresponding leaf of T' as per some fixed isomorphism between T and T' . We say that H is obtained from T and T' by *isomorphic leaf matching*, or simply, that H is obtained from T by isomorphic leaf matching.

Note that, since K_2 is the smallest halin tree, C_4 is the smallest graph that may be obtained using the operation defined above. The following observation is easily proved and it implies that, for every other graph obtained using this operation, one may recover the two copies of the halin tree by simply deleting the set of 2-edges.

Proposition 1.2. *If a graph H , distinct from C_4 , is obtained from a halin tree T by isomorphic leaf matching, then $H - E_2$ has precisely two components, each of which is isomorphic to T .* \square

We first focus on the 2-vertex extremal class \mathcal{H}_2 (see Table 1) and obtain the following characterization establishing a bijection between 2-vertex extremal bipartite matching covered graphs and halin trees.

Theorem 1.3. [MAIN THEOREM: CHARACTERIZATION OF \mathcal{H}_2]

A graph is a 2-vertex extremal minimal bipartite matching covered graph if and only if it is obtained from a halin tree by isomorphic leaf matching.




Figure 3: a member of \mathcal{H}_2 — deleting the 2-edges results in two copies of a halin tree

Figure 3 shows an illustration of the above theorem. In this figure, and relevant figures henceforth, we adopt the following color conventions: the red vertices have degree three or more, the green vertices are of degree two, and the cyan edges are precisely the 2-edges.

Thereafter, we will use our Main Theorem (1.3) to derive characterizations of the extremal classes \mathcal{H}_3 and \mathcal{H}_4 as stated below; see Figure 4 for an example of each.

Theorem 1.4. [CHARACTERIZATION OF \mathcal{H}_3]

A graph is a 2-vertex n -extremal minimal bipartite matching covered graph if and only if it is obtained from a cubic halin tree by isomorphic leaf matching.

By a *star*, we mean $K_{1,p}$ where $p \geq 2$. Observe that these, except $K_{1,2}$, comprise a restricted subclass of halin trees; they play a crucial role in our characterization of \mathcal{H}_4 stated below.

Theorem 1.5. [CHARACTERIZATION OF \mathcal{H}_4]

A graph is an edge extremal minimal bipartite matching covered graph if and only if it is obtained from a star by isomorphic leaf matching.




Figure 4: (a) a member of \mathcal{H}_3 : deleting the 2-edges results in two copies of a cubic halin tree
(b) a member of \mathcal{H}_4 : deleting the 2-edges results in two copies of a star

In the next subsection, we prove the easier implications of the characterizations stated above (that is, Theorems 1.3, 1.4 and 1.5).

1.3 Proofs of easier implications

We begin by proving that any graph obtained from a nontrivial tree by isomorphic leaf matching is a minimal bipartite matching covered graph. We shall find ear decompositions useful to establish the matching covered property; however, for minimality, we need an easy observation pertaining to cuts.

Given a graph $G := (V, E)$, for any $W \subseteq V$, the *cut* comprising the edges joining W and its complement $\overline{W} := V - W$ is denoted by $\partial(W)$. A k -*cut* refers to a cut of cardinality k . We use abbreviated notations $\partial(v) := \partial(\{v\})$ for a vertex v , and $\partial(H) := \partial(V(H))$ for a subgraph H .

An edge e of a matching covered graph G is *removable* if $G - e$ is also matching covered. Thus, a matching covered graph is minimal if and only if it has no removable edges. Since matching covered graphs are 2-connected, we immediately observe the following.

Lemma 1.6. *In a matching covered graph, any edge that participates in a 2-cut is not removable.* \square

The above lemma is the aforementioned observation that helps us in establishing minimality in the proof of the following result; we shall find it useful later as well.

Proposition 1.7. *Any graph obtained from a nontrivial tree by isomorphic leaf matching is a minimal bipartite matching covered graph.*

Proof. Let G be a graph obtained from two copies of a nontrivial tree, say T and T' , by isomorphic leaf matching. We will first prove, by induction on the order of T , that G is a bipartite matching covered graph. If T is a path then G is an even cycle, and we are done.

Now, let T be a tree that is not a path and let v denote a leaf. Let P be the maximal path starting at v , each of whose internal vertices has degree two in T . Let u be the end of P distinct from v . Observe that $d_T(u) \geq 3$. Let $v'P'u'$ be the path corresponding to vPu in T' . Let $H := G - V(P - u) - V(P' - u')$. Observe that H is obtained from the tree $T - V(P - u)$ by isomorphic leaf matching. By the induction hypothesis, H is a bipartite matching covered graph. Note that G may be obtained from H by adding the ear $uPvv'P'u'$. By the ear decomposition theorem (1.1), G is a bipartite matching covered graph.

Now, we prove minimality. Observe that any edge e of G , whose each end has degree at least three, either belongs to T or belongs to T' . Adjust notation so that $e \in T$. Let e' be the copy of e in T' . Note that $\{e, e'\}$ is a 2-cut of G . Thus, by Lemma 1.6, e is not removable. We thus infer that G is minimal. \square




Figure 5: a counterexample to the converse of Proposition 1.7

We note that the converse of the above proposition does not hold. The graph G , shown in Figure 5, is a minimal bipartite matching covered graph that can not be obtained from a tree by isomorphic leaf matching. To see this, suppose to the contrary that G is obtained from two copies of a tree, say T and T' , by isomorphic leaf matching. Then, as per the proof of Proposition 1.7, the labeled edge e belongs either to T or to T' as both ends of e have degree greater than two; furthermore, e belongs to a 2-cut of G . However, $G - e$ is 2-connected. This is a contradiction. We shall revisit this example in Section 4.1.

In light of Proposition 1.7, in order to prove the easier implications of Theorems 1.3, 1.4 and 1.5, we simply need to argue that if we choose a tree T from the corresponding class (as in that theorem's statement), then the resulting minimal bipartite matching covered graph G (obtained by isomorphic leaf matching) satisfies the desired extremality notion. We first note that if T is the star $K_{1,2}$ then $G = C_6$ is an edge extremal minimal bipartite matching covered graph. All of the remaining cases are handled by the following theorem — which we prove using straightforward counting arguments.

Theorem 1.8. *Any graph G obtained from a halin tree T by isomorphic leaf matching is a minimal bipartite matching covered graph that is 2-vertex extremal. Furthermore:*

- (i) if T is a cubic halin tree then G is also 2-vertex n -extremal, whereas
- (ii) if T is a star then G is also edge extremal.

Proof. Let $G := (V, E)$ be a graph obtained from two copies of a halin tree, say T and T' , by isomorphic leaf matching. By Proposition 1.7, G is a minimal bipartite matching covered graph. It remains to prove “extremality”. We use V_2 and E_2 to denote $V_2(G)$ and $E_2(G)$, respectively.

Since T is a halin tree, the non-leaves of T and T' comprise those vertices of G that have degree at least three, whereas the leaves of T and T' comprise the remaining vertices of G (that is, its vertices of degree two). In particular, there are no 2-edges in $T \cup T'$, and $T \cup T' = G - E_2$.

Now, $|E(T)| = |V(T)| - 1$. So, $|E(T \cup T')| = n - 2$. On the other hand, $|E(T \cup T')| = |E| - |E_2| = m - |E_2|$. Thus, $|E_2| = m - n + 2$. Since E_2 is a matching of G , and the corresponding matched vertices comprise the set V_2 , we get $|V_2| = 2|E_2| = 2(m - n + 2)$. Hence, G is 2-vertex extremal. It remains to prove statements (i) and (ii).

First, suppose that T is a cubic halin tree. So, $T \cup T'$ has $|V_2|$ leaves and $n - |V_2|$ vertices of degree three. By handshaking lemma, $3(n - |V_2|) + |V_2| = 2|E(T \cup T')| = 2(n - 2)$. By simplifying, we get $|V_2| = \frac{n}{2} + 2$; whence G is 2-vertex n -extremal.

Now, suppose that T is a star $K_{1,p}$, where $p \geq 3$. Then, $n = 2(p + 1)$. Observe that $m = 3p = \frac{3n-6}{2}$. Ergo, G is edge extremal. \square

We now switch our attention to the remaining “extremal” classes \mathcal{H}_0 and \mathcal{H}_1 .

1.4 Characterizations of \mathcal{H}_0 and \mathcal{H}_1

In general, members of \mathcal{H}_0 and \mathcal{H}_1 may not be obtained from a tree by isomorphic leaf matching, unlike the members of $\mathcal{H}_2, \mathcal{H}_3$ and \mathcal{H}_4 . To see this, we discuss a couple of examples that are shown in Figure 6; using the ear decomposition theorem (1.1) and Lemma 1.6, the reader may easily verify that both graphs belong to \mathcal{H} . By counting, one may infer these graphs belong to \mathcal{H}_0 and \mathcal{H}_1 , respectively. For the graph shown in Figure 6a, deleting all of the 2-edges (displayed in blue) results in precisely two components that are trees but nonisomorphic. On the other hand, for the graph shown in Figure 6b, deleting all of the 2-edges results in two isomorphic trees; however, the resultant graph is not obtained by isomorphic leaf matching operation. Interestingly, we are able to reduce members of \mathcal{H}_0 and \mathcal{H}_1 to members of \mathcal{H}_2 and \mathcal{H}_4 , respectively, using the notion of “partial retract” that we describe next.

Let G be a graph and v be a vertex of degree two that has two distinct neighbours, say v_1 and v_2 . Let G' denote the graph obtained from G by contracting the two edges vv_1 and vv_2 . We say that G' is obtained from G by *bicontraction* (of v), and we denote it by $G' := G/v$; see Figure 15. Furthermore, if each of v_1 and v_2 has degree at least three in G , we say that G' is obtained from G by *restricted bicontraction* (of v); see Figure 16. Note




Figure 6: extremal graphs that may not be obtained from a tree by isomorphic leaf matching

that restricted bicontraction may be performed if and only if the subgraph of G induced by its degree two vertices has an isolated vertex.

We define the *partial retract* of G as the graph \widehat{G} obtained by repeatedly applying the restricted bicontraction operation until the resulting graph has the property that the subgraph induced by its degree two vertices has no isolated vertex. Observe that given a graph G , its partial retract \widehat{G} is unique; Figure 7 shows an example.

Using the notion of partial retract, we are able to relate the extremal class \mathcal{H}_0 with the extremal class \mathcal{H}_2 , as stated below.

Theorem 1.9. [CHARACTERIZATION OF \mathcal{H}_0]

A graph G , distinct from C_4 , is a 2-edge extremal minimal bipartite matching covered graph if and only if its partial retract \widehat{G} is a 2-vertex extremal minimal bipartite matching covered graph.




Figure 7: an illustration of Theorem 1.9 that relates \mathcal{H}_0 and \mathcal{H}_2

To see an example of the above theorem, consider the graph G shown in Figure 7a and its partial retract \widehat{G} shown in Figure 7b. The reader may verify using Theorem 1.1 and Lemma 1.6 that G and \widehat{G} belong to \mathcal{H} . Note that $G \in \mathcal{H}_0$ as $|E_2(G)| = m - n + 2$. On the other hand, $\widehat{G} \in \mathcal{H}_2$ as $|V_2(\widehat{G})| = 2(m - n + 2)$.

Likewise, using the notion of partial retract, we are able to relate the extremal class \mathcal{H}_1 with the extremal class \mathcal{H}_4 , as stated below.

Theorem 1.10. [CHARACTERIZATION OF \mathcal{H}_1]

A graph G is a 2-edge n -extremal minimal bipartite matching covered graph if and only if its

partial retract \widehat{G} is an edge extremal minimal bipartite matching covered graph and $\Delta(G) = 3$.



Figure 8: an illustration of Theorem 1.10 that relates \mathcal{H}_1 and \mathcal{H}_4

To see an example of the above theorem, consider the graph G shown in Figure 8a and its partial retract \widehat{G} shown in Figure 8b. The reader may verify using Theorem 1.1 and Lemma 1.6 that G and \widehat{G} belong to \mathcal{H} . Note that $G \in \mathcal{H}_1$ as $|E_2(G)| = \frac{n+10}{6}$. On the other hand, $\widehat{G} \in \mathcal{H}_4$ as $|E(\widehat{G})| = \frac{3n-6}{2}$.

We have thus stated our characterizations for all of the extremal classes shown in Table 1. We now proceed to discuss further relations between them.

1.5 Containment poset for the extremal classes

In order to discuss the containment relationships between different extremal classes, we shall find it convenient to exclude the cycle graphs. It follows from the definitions of the extremal classes (see Table 1) that neither of \mathcal{H}_0 and \mathcal{H}_1 contains any cycle graph, that C_4 is the only cycle in \mathcal{H}_2 as well as in \mathcal{H}_3 , whereas C_6 is the only cycle in \mathcal{H}_4 . In light of this, we let $\mathcal{H}_2^* := \mathcal{H}_2 - C_4$, $\mathcal{H}_3^* := \mathcal{H}_3 - C_4$ and $\mathcal{H}_4^* := \mathcal{H}_4 - C_6$.




Figure 9: The Θ graph




Figure 10: Containment poset

Figure 10 shows the containment poset for all of the extremal classes. For instance, the arrow from \mathcal{H}_1 to \mathcal{H}_0 , labelled as 7.11, is to be read as follows: Corollary 7.11 states that $\mathcal{H}_1 \subset \mathcal{H}_0$. Additionally, our characterizations imply that $\mathcal{H}_1 \cap \mathcal{H}_2 = \mathcal{H}_3 \cap \mathcal{H}_4 = \{\Theta\}$ where Θ is the graph shown in Figure 9; see Corollaries 7.5 and 7.12.

1.6 Organization of the rest of this paper

In the rest of the paper, we will prove the difficult directions of Theorems 1.3, 1.4 and 1.5, and we will also prove Theorems 1.9 and 1.10.

We first focus on the class \mathcal{H}_2 . In order to prove the Main Theorem (1.3), we need some known properties of bipartite matching covered graphs and minimal bipartite matching covered graphs; these are discussed in Sections 2 and 3. In particular, in Section 3, we state and prove the lower bound on the number of 2-edges due to Lovász and Plummer [9], and we deduce an easy lower bound on the number of degree two vertices. In Section 4, we enforce equality in its proof to infer some useful properties of members of \mathcal{H}_2 — one of which is the existence of “balanced” 2-cuts. We then develop an induction tool using balanced 2-cuts in Section 5. In Section 6, we prove the Main Theorem (1.3).

Finally, in Section 7, we address the other notions of extremality. In Subsections 7.1 and 7.2, we first state and prove the bounds corresponding to the classes \mathcal{H}_3 and \mathcal{H}_4 , respectively; thereafter, we enforce equality in their proofs to deduce their characterizations (Theorems 1.4 and 1.5) using the Main Theorem (1.3). In Subsection 7.3, we prove our characterization of \mathcal{H}_0 (Theorem 1.9). Lastly, in Subsection 7.4, we prove our characterization of \mathcal{H}_1 (Theorem 1.10) using the characterization of \mathcal{H}_0 .

2 Ear decomposition theory

Recall that a subgraph H of a graph G is conformal if $G - H$ is matchable. It is easy to see that, in a matching covered graph, there is a conformal cycle containing any two adjacent edges. As an immediate consequence, we get the following.

Proposition 2.1. *Let G be a matching covered graph, let u be a vertex of degree at least three and let $e \in \partial(u)$. Then there is a conformal cycle C such that $u \in C$ but $e \notin C$. \square*

In fact, Little [8] proved the stronger statement that any two edges (not necessarily adjacent) lie in a conformal cycle; we will not require this.

Recall that an ear of a subgraph H in a graph G is an odd path of G whose ends are in H but is otherwise disjoint from H . We will now discuss the proof of the Ear Decomposition Theorem (1.1) as it appears in Lovász and Plummer [9, Theorem 4.1.6]. In particular, we extract a couple of lemma statements that are implicit in their proof. The first of these shows that “adding an ear” preserves the matching covered property. For a subgraph H and an ear P of H , the subgraph $H \cup P$ is said to be obtained by adding the ear P to H .

Lemma 2.2. *Let G be a bipartite graph, H be a matching covered subgraph and P be an ear of H in G . Then $H \cup P$ is matching covered.*

The second lemma establishes the existence of an ear of any conformal matching covered proper subgraph H with the additional constraint that any specified edge “sticking out of H ” is included in the ear.

Lemma 2.3. *Let G be a bipartite matching covered graph, H be any conformal matching covered subgraph and $e \notin H$ denote an edge that has at least one end in H . Then there exists an ear of H , say P_e , containing e such that $H \cup P_e$ is a conformal (matching covered) subgraph of G .*

The reader may verify that the above two lemmas imply the the following stronger version of the Ear Decomposition Theorem (1.1) that is also stated in Lovász and Plummer [9, pg 126].

Theorem 2.4. *For any conformal matching covered subgraph H of a bipartite matching covered graph G , there exists an ear decomposition of G in which H appears as one of the subgraphs. \square*

In the next section, we discuss the proofs of the lower bounds (on the number of 2-edges) due to Lovász and Plummer [9, Theorem 4.2.7]. These proofs rely heavily on the ear decomposition theory, and we will need some of their details to prove our results.

3 Lower bounds established by Lovász and Plummer

Recall that, for a matching covered graph G , an edge e is removable if $G - e$ is matching covered. Thus, G is minimal if and only if it has no removable edges. It follows from Theorem 1.1 that, given any ear decomposition of a bipartite matching covered graph G , every trivial ear (that is, an ear comprising a single edge) is a removable edge in G . This observation, coupled with Theorem 2.4, yields the following; see Lovász and Plummer [9, Theorem 4.2.1].

Lemma 3.1. *Every conformal matching covered subgraph of a minimal bipartite matching covered graph is an induced subgraph. \square*

In order to establish their lower bound on the number of 2-edges in a minimal bipartite matching covered graph, Lovász and Plummer proved a lower bound for each element (the starting conformal cycle and each ear) of an ear sequence. The following is their lemma for ears.

Lemma 3.2. *Let G be a minimal bipartite matching covered graph and let $(C, P_1, P_2, \dots, P_r)$ be any ear sequence. Then each ear P_i contains a 2-edge of G .*

Note that the above lemma (see Lovász and Plummer [9, Lemma 4.2.4]) refers to degrees in the original graph G . Adding a nontrivial ear P_i clearly creates a 2-edge in the resultant subgraph. However, one may add ears later thereby increasing the degrees of internal vertices of P_i and possibly destroying all 2-edges of P_i in the final graph G . The above lemma states that at least one such 2-edge survives in G . We now switch our attention to conformal cycles.

Let G be a minimal bipartite matching covered graph that is not a cycle. Using Lemma 3.2

and some standard ear decomposition tricks, one may infer that each conformal cycle C of G contains a pair of 2-edges that are separated in C by vertices that have degree at least three (in G). Interestingly, the same holds even for cycles that are not conformal; see Lovász and Plummer [9, Corollary 4.2.5].

A matching M is *induced* if the subgraph induced by $V(M)$ contains precisely the edges of M . The facts stated in the above paragraph, along with Lemma 3.1, imply the following (that is easily verified in the case in which the graph itself is a cycle of order six or more).

Corollary 3.3. *In a minimal bipartite matching covered graph, distinct from C_4 , each conformal cycle contains a pair of 2-edges that comprise an induced matching.*

We now invoke Lemma 3.2 and Corollary 3.3, and the fact that the length of any ear sequence is $m - n + 1$, to deduce the following lower bound result that is a mild strengthening of what is stated in Lovász and Plummer; see [9, Theorem 4.2.7].

Theorem 3.4. *A minimal bipartite matching covered graph, distinct from C_4 , has an induced matching of size at least $m - n + 2$, each of whose members is a 2-edge.*

Proof. Let G be a minimal bipartite matching covered graph and (C, P_1, \dots, P_r) be any ear sequence. By Corollary 3.3, the conformal cycle C has a pair of 2-edges, say e and e' , that comprise an induced matching of G . By Lemma 3.2, each ear P_i has a 2-edge, say e_i . Observe that each end of e and e' has both its neighbours in C . Likewise, each end of e_i has both its neighbours in P_i . Furthermore, the ends of each ear P_i have degree at least three in G . All of these observations imply that the set $F := \{e, e', e_1, e_2, \dots, e_r\}$ is an induced matching of G and each of its members is a 2-edge. Also, $r = m - n$. Thus, $|F| = m - n + 2$. \square

The following lower bound on the number of vertices of degree two comes for free.

Corollary 3.5. *In a minimal bipartite matching covered graph, $|V_2| \geq 2(m - n + 2)$.*

Proof. The statement clearly holds for C_4 . Now, let G be a minimal bipartite matching covered graph, distinct from C_4 , and let F denote a maximum induced matching, each of whose members is a 2-edge. Clearly, $|V_2| \geq 2|F|$. By Theorem 3.4, $|F| \geq m - n + 2$. Thus, $|V_2| \geq 2(m - n + 2)$. \square

Throughout the next three sections, for the sake of brevity, we shall use the term *extremal* to refer to 2-vertex extremal — that is, those minimal bipartite matching covered graphs that satisfy the bound in Corollary 3.5 with equality.

4 Properties of extremal graphs

In this section, we establish some useful properties of this class including the “balanced 2-cut” property that is crucial for our inductive proof of Theorem 1.3 that characterizes this class. To this end, we shall enforce equality in the proofs of Corollary 3.5 and Theorem 3.4.

Lemma 4.1. *Every extremal minimal bipartite matching covered graph, distinct from C_4 , satisfies the following:*

- (i) E_2 is a perfect matching of $G[V_2]$ and $|E_2| = m - n + 2$,
- (ii) each ear in any ear decomposition has exactly one 2-edge, and
- (iii) each conformal cycle has precisely two 2-edges.

Proof. Let G be an extremal minimal bipartite matching covered graph, distinct from C_4 , and let F denote an induced matching of maximum cardinality, each of whose members is a 2-edge, as in the proof of Corollary 3.5. Since, by definition, $|V_2| = 2(m - n + 2)$, equality holds everywhere in the proof of Corollary 3.5. In particular, $|F| = m - n + 2$ and $|V_2| = 2|F|$; thus $V_2 = V(F)$. Observe that, since F is an induced matching, $E_2 = F$. Consequently, E_2 is a perfect matching of $G[V_2]$ and $|E_2| = m - n + 2$. This proves (i).

Now, let (C, P_1, \dots, P_r) be any ear sequence of G . Since $|F| = m - n + 2$, equality holds everywhere in the proof of Theorem 3.4. Furthermore, using the fact that $|E_2| = m - n + 2$, we infer that the conformal cycle C contributes exactly two 2-edges, and each ear P_i contributes precisely one 2-edge, to the set E_2 . This proves (ii) and (iii). \square

Using Lemma 4.1 (i), we immediately deduce the following containment.

Corollary 4.2. $\mathcal{H}_2 - C_4 \subset \mathcal{H}_0$. \square

We now derive, using the above lemma and simple counting arguments, a few additional properties of this extremal class. In order to do so, we need some notation and terminology that we define next.

Following Lovász and Plummer [9], we call an edge (of a graph G) a *3-edge* if each of its ends has degree at least three, and we use $E_3(G)$ to denote the set comprising these edges. In the same spirit, we use $V_3(G)$ to denote the set of vertices of degree at least three. Technically, it should be denoted by $V_{\geq 3}(G)$. However, throughout this paper, we distinguish between vertices of degree two and the rest — that is, vertices of degree three or more. So, for the sake of brevity, we drop “ \geq ” from the notation. Finally, we use $E_{3,2}(G)$ to denote the set of those edges that join a vertex of degree two with a vertex of degree at least three. As usual, if the graph G is clear from the context, we shall drop G from these notations. Note that $E_{3,2} = \partial(V_3) = \partial(V_2)$.

Now, let G be an extremal minimal bipartite matching covered graph distinct from C_4 . By definition, $|V_3| = n - |V_2| = 3n - 2m - 4$. By Lemma 4.1, each vertex of degree two has exactly one neighbour of degree at least three; consequently, $|E_{3,2}| = |V_2| = 2(m - n + 2)$. Lastly, $|E_3| = m - |E_2| - |E_{3,2}| = 3n - 2m - 6$. Also, since $|V_3| = 3n - 2m - 4$, it follows that $|E_3| = |V_3| - 2$. This proves the following.

Corollary 4.3. *Every extremal minimal bipartite matching covered graph, distinct from C_4 , satisfies the following:*

- (i) $|V_3| = 3n - 2m - 4$,

(ii) $|E_{3,2}| = 2m - 2n + 4$, and

(iii) $|E_3| = 3n - 2m - 6 = |V_3| - 2$. □

We shall now use Lemma 4.1 to prove the “balanced 2-cut” property of extremal minimal bipartite matching covered graphs that was alluded to at the beginning of this section.

4.1 Balanced 2-cut property

For a cut $\partial(W)$ of a bipartite graph $G[A, B]$, we classify the edges of $\partial(W)$ into two types, depending upon the color classes of their ends in W . We use $\partial^A(W)$ to denote those edges of $\partial(W)$ whose end in W belongs to A . The set $\partial^B(W)$ is defined analogously. The cut $\partial(W)$ is *balanced* if $|\partial^A(W)| = |\partial^B(W)|$. In our work, balanced 2-cuts play a crucial role. Note that a 2-cut $\partial(W)$ is balanced if $|\partial^A(W)| = |\partial^B(W)| = 1$. We are now ready to prove the balanced 2-cut property stated below.

Theorem 4.4. [BALANCED 2-CUT PROPERTY]

In an extremal minimal bipartite matching covered graph, each 3-edge participates in a balanced 2-cut with some other 3-edge.

Proof. Let $G[A, B]$ denote an extremal minimal bipartite matching covered graph and let $e := ab$ denote any 3-edge so that $a \in A$. We intend to show the existence of another 3-edge, say f , such that $\{e, f\}$ is a balanced 2-cut.

By Proposition 2.1, there is a conformal cycle C such that $a \in C$ and $e \notin C$. Now, let H be a maximal conformal matching covered subgraph of G such that $a \in H$ and $e \notin H$. By Lemma 3.1, H is induced; thus $b \notin H$. In particular, $e \in \partial^A(H)$. We shall demonstrate that $\partial(H)$ is the desired balanced 2-cut. In order to do so, we state and prove specific claims; within their proofs, we shall invoke ear decomposition results (namely, Lemma 2.3 and Theorem 2.4) several times without mentioning it explicitly.

4.4.1. $\partial^A(H) = \{e\}$.

Proof. Suppose not; then there is an edge $f \neq e$ in $\partial^A(H)$. Now, there is an ear P_f of H containing f . Observe that $e \notin P_f$ since P_f is odd. Thus, $H' := H \cup P_f$ is a larger conformal matching covered subgraph of G such that $a \in H'$ and $e \notin H'$; this contradicts the maximality of H . This proves that $\partial^A(H) = \{e\}$. □

In the proofs of the next two claims, every instance of 2-edge refers to the condition of being a 2-edge in the graph G similar to our earlier discussion (after the proof of Lemma 3.2).

4.4.2. $|\partial^B(H)| = 1$.

Proof. Since G is 2-connected, $|\partial^B(H)| \geq 1$. Now suppose that f_1 and f_2 are distinct edges in $\partial^B(H)$ and let b_1 and b_2 denote their (not necessarily distinct) ends in H , respectively. Now, there is an ear P' of H containing f_1 (which must also contain e). Furthermore, there is an ear P_2 of $H \cup P'$ containing f_2 . Let $x \in A$ be the end of P_2 distinct from b_2 (see

Figure 11a). Observe that $x \notin H$ since $\partial^A(H) = \{e\}$ and $e \in P'$. Thus, x is an internal vertex of P' . Consequently, x splits P' into two paths, say $P := aP'x$ and $P_1 := xP'b_1$. Note that P_2 is an ear of $H \cup (P + P_1) = H \cup P'$; likewise, P_1 is an ear of $H \cup (P + P_2)$. By Lemma 4.1 (ii), each of P' , P_1 and P_2 has exactly one 2-edge. Since P_1 is a subpath of P' , we conclude that P is free of 2-edges. Furthermore, by Lemma 4.1 (i), each vertex of P has degree at least three in G .




Figure 11: illustrations for the proof of statement 4.4.2

Now, let y be the neighbour of x in P . As discussed above, $d_G(y) \geq 3$. So, there is an ear P_y of $H \cup (P + P_1) \cup P_2$ starting from y . Let z be the other end of P_y . Observe that $z \notin H$ as $\partial^A(H) = \{e\}$.

Now, suppose that $z \in P$ (see Figure 11b). Note that the conformal matching covered subgraph $H \cup (P + P_1) \cup P_y$ can also be obtained from H by adding the ears $aPzP_yyxP_1b_1$ and yPz in that order. However, yPz has no 2-edges since, as noted earlier, each vertex of P has degree at least three in G . This contradicts Lemma 4.1 (ii). Thus, $z \notin P$.

Thus, $z \in P_1 \cup P_2$. Adjust notation so that $z \in P_1$ (see Figure 11c). Let $Q := aP_yP_yzP_1b_1$ and $Q' := zP_1xP_2b_2$ (see Figure 11d). Observe that the conformal matching covered subgraph $H \cup (P + P_1) \cup P_2 \cup P_y$ can also be obtained from H by adding the ears Q, Q' and xy in that order. In particular, $H \cup Q \cup Q'$ is a conformal matching covered subgraph of G that is not induced. This contradicts Lemma 3.1. Thus, $|\partial^B(H)| = 1$. \square

Let f denote the unique edge of $\partial^B(H)$. Note that $\partial(H) = \{e, f\}$ is indeed a balanced 2-cut. It remains to prove the following claim.

4.4.3. f is a 3-edge.

Proof. We let $f := uv$ where $v \in V(H) \cap B$. As H is matching covered, $d_H(v) \geq 2$; thus $d_G(v) \geq 3$. Suppose to the contrary that $d_G(u) = 2$. By Lemma 4.1 (i), the neighbour of u distinct from v , say w , has degree two in G . Let P be an ear of H containing f . Observe that $uw, e \in P$. By Lemma 4.1 (ii), uw is the only 2-edge of P . Furthermore, by Lemma 4.1 (i), each internal vertex of P , distinct from u and w , has degree at least three in G .




Figure 12: illustrations for the proof of statement 4.4.3

Since $d_G(b) \geq 3$, there is an ear P' of $H \cup P$ starting from b (see Figure 12a). Let $x \in A$ be the other end of P' . Observe that x is an internal vertex of P that is distinct from u . Let $Q := abP'xPv$ and $Q' := bP'x$ (see Figure 12b). Observe that the conformal matching covered subgraph $H \cup P \cup P'$ can also be obtained from H by adding the ears Q and Q' in that order. It follows from the preceding paragraph that each vertex of Q' has degree at least three in G . Consequently, the ear Q' has no 2-edges. This contradicts Lemma 4.1 (ii). Thus, $d_G(u) \geq 3$ and f is indeed a 3-edge. \square

This completes the proof of Theorem 4.4. \square

Note that Theorem 4.4 does not hold for all minimal bipartite matching covered graphs — that is, it requires the extremality hypothesis. As we noted in Section 1, the graph G shown in Figure 5 is a minimal bipartite matching covered graph. Observe that e is a 3-edge that is not contained in any 2-cut since $G - e$ is 2-connected.

In the next section, we establish an induction tool based on balanced 2-cuts. This tool, coupled with Theorem 4.4, will be used in Section 6 to prove the Main Theorem (1.3).

5 An induction tool using balanced 2-cuts

We begin this section by defining an operation that “breaks” a given bipartite graph, that has a balanced 2-cut, into “smaller” bipartite graphs, and its converse operation.

Let $G[A, B]$ denote a bipartite graph that has a balanced 2-cut, say $F := \partial(X)$. Let $H_1 := G[X]$ and $H_2 := G[\overline{X}]$. As F is balanced, for each $i \in \{1, 2\}$, let $a_i \in A$ and $b_i \in B$ denote the ends of the edges of F in H_i , as shown in Figure 13a. Let G_1 be the graph obtained from H_1 by adding the ear $a_1v_1u_1b_1$ (of length three), as shown in Figure 13b. The

graph G_2 is obtained from H_2 analogously. We say that the (bipartite) graphs G_1 and G_2 are obtained from G by a *balanced 2-cut decomposition*, or simply by a *2-cut decomposition*, across the 2-cut F . Note that u_1v_1 is a 2-edge in G_1 ; likewise, u_2v_2 is a 2-edge in G_2 .

Conversely, let G_1 and G_2 be disjoint bipartite graphs, each of which has a 2-edge, say u_1v_1 and u_2v_2 , respectively. Let a_1 denote the neighbour of v_1 that is distinct from u_1 , and let b_1 denote the neighbour of u_1 that is distinct from v_1 , as shown in Figure 13b. The vertices a_2 and b_2 are defined analogously. Let G be the graph obtained from $G_1 \cup G_2$ by deleting the vertices u_1, v_1, u_2 and v_2 , and adding the edges a_1b_2 and a_2b_1 , as shown in Figure 13a. We say that the (bipartite) graph G is obtained from G_1 and G_2 by *2-edge splicing* across the pair of 2-edges $\{u_1v_1, u_2v_2\}$. Note that $\{a_1b_2, a_2b_1\}$ is a balanced 2-cut of G .

Observe that the 2-cut decomposition operation is the “inverse” of the 2-edge splicing operation — that is, G_1 and G_2 are obtained from G by 2-cut decomposition across a balanced 2-cut if and only if G is obtained from G_1 and G_2 by 2-edge splicing across a pair of 2-edges. We now show that these operations preserve the matching covered, minimality and extremality properties.




Figure 13: illustration of balanced 2-cut decomposition and 2-edge splicing

Theorem 5.1. [BALANCED 2-CUT INDUCTION TOOL]

Let G be a bipartite graph that has a balanced 2-cut F , and let G_1 and G_2 be the (bipartite) graphs obtained from G by a 2-cut decomposition across F . Then the following statements hold:

- (i) G is matching covered if and only if G_1 and G_2 are both matching covered;
- (ii) furthermore, G is minimal if and only if G_1 and G_2 are both minimal; and
- (iii) finally, G is extremal if and only if G_1 and G_2 are both extremal.

Proof. We adopt all of the notations from the above definition of 2-cut decomposition, as shown in Figure 13. We will prove the three statements one by one.

5.1.1. G is matching covered if and only if G_1 and G_2 are both matching covered.

Proof. First suppose that G is matching covered. Let C denote a conformal cycle containing a_1b_2 . Note that $a_2b_1 \in C$ as well. Let (C, P_1, \dots, P_r) be an ear sequence of G ; its existence

is guaranteed by Theorem 2.4. Observe that each ear P_i is a subgraph of exactly one of H_1 and H_2 . Let $P_{11}, P_{12}, \dots, P_{1r_1}$ denote those ears that are subgraphs of H_1 and appear in that order in the ear sequence (C, P_1, \dots, P_r) . Let $C_1 := (C \cap H_1) + a_1v_1u_1b_1$. Observe that $(C_1, P_{11}, P_{12}, \dots, P_{1r_1})$ is an ear sequence of G_1 . Thus, by the Ear Decomposition Theorem (1.1), the bipartite graph G_1 is matching covered. An analogous argument proves that G_2 is matching covered.

Conversely, suppose that G_1 and G_2 are matching covered. Let C_i be a conformal cycle of G_i containing the 2-edge u_iv_i for $i \in \{1, 2\}$. Let $(C_1, P_{11}, P_{12}, \dots, P_{1r_1})$ and $(C_2, P_{21}, P_{22}, \dots, P_{2r_2})$ be ear sequences of G_1 and G_2 , respectively; these exist due to Theorem 2.4. Let $C := (C_1 - u_1 - v_1) + (C_2 - u_2 - v_2) + a_1b_2 + a_2b_1$. Observe that $(C, P_{11}, P_{12}, \dots, P_{1r_1}, P_{21}, P_{22}, \dots, P_{2r_2})$ is an ear sequence of G . Thus, by the Ear Decomposition Theorem (1.1), G is matching covered. \square

Henceforth, we assume that G is matching covered, or equivalently that G_1 and G_2 are both matching covered.

5.1.2. *G is minimal if and only if G_1 and G_2 are both minimal.*

Proof. First suppose that G is minimal. We claim that G_1 is also minimal. Suppose not, and let e be a removable edge of G_1 . Note that $e \in H_1$ since each of the remaining three edges has an end of degree two. Observe that $G_1 - e$ and G_2 are obtained by a 2-cut decomposition of $G - e$. Consequently, by 5.1.1, the graph $G - e$ is matching covered; this contradicts the minimality of G . Thus, G_1 is minimal. Likewise, G_2 is minimal.

Conversely, suppose that G_1 and G_2 are both minimal. We claim that G is minimal. Suppose not, and let e be a removable edge of G . Note that, since F is 2-cut, e belongs to exactly one of H_1 and H_2 . Adjust notation so that $e \in H_1$. As in the preceding paragraph, $G_1 - e$ and G_2 are obtained by a 2-cut decomposition of $G - e$. Thus, by 5.1.1, the graph $G_1 - e$ is matching covered; this contradicts the minimality of G_1 . Hence, G is minimal. \square

Henceforth, we assume that G is a minimal bipartite matching covered graph, or equivalently that G_1 and G_2 are both minimal bipartite matching covered graphs.

5.1.3. *G is extremal if and only if G_1 and G_2 are both extremal.*

Proof. In order to argue extremality, we find it convenient to use $\lambda := |V_2(G)| - 2(m - n + 2)$ to denote the ‘‘surplus’’ vertices of degree two in G . Analogously, we define λ_1 and λ_2 for the graphs G_1 and G_2 , respectively. By Corollary 3.5, since all three graphs are minimal bipartite matching covered graphs, the quantities λ , λ_1 and λ_2 are non-negative. Note that G is extremal if and only if $\lambda = 0$. Likewise, for $i \in \{1, 2\}$, the graph G_i is extremal if and only if $\lambda_i = 0$. Thus, it suffices to prove that $\lambda = 0$ if and only if $\lambda_1 = 0$ and $\lambda_2 = 0$.

For $i \in \{1, 2\}$, we let $n_i := |V(G_i)|$ and $m_i := |E(G_i)|$. The reader may find it useful to look at Figure 13. Note that $n_1 + n_2 = n + 4$ and $m_1 + m_2 = m + 4$. Observe that each vertex w of G corresponds to a vertex of G_1 or of G_2 (that is distinct from u_1, v_1, u_2 and v_2)

whose degree is the same as that of w . This implies that $|V_2(G_1)| + |V_2(G_2)| = |V_2(G)| + 4$. Using these three equalities, and the definitions of λ, λ_1 and λ_2 :

$$\begin{aligned}\lambda_1 + \lambda_2 &= |V_2(G_1)| + |V_2(G_2)| - 2(m_1 + m_2 - n_1 - n_2 + 4) \\ &= |V_2(G)| + 4 - 2(m - n + 4) \\ &= |V_2(G)| - 2(m - n + 2) = \lambda\end{aligned}$$

Thus, since $\lambda, \lambda_1, \lambda_2$ are non-negative, $\lambda = 0$ if and only if $\lambda_1 = 0$ as well as $\lambda_2 = 0$. \square

This completes the proof of Theorem 5.1. \square

6 Main Theorem: Characterization of \mathcal{H}_2

In this section, we prove our Main Theorem (1.3) that provides a characterization of extremal minimal bipartite matching covered graphs. Before doing so, we prove a lemma that will be used in the base case of our inductive proof.

Lemma 6.1. *Every extremal minimal bipartite matching covered graph, that has precisely two vertices of degree at least three, is obtained from a star by isomorphic leaf matching.*

Proof. Let G be an extremal minimal bipartite matching covered graph with precisely two vertices of degree at least three, say a and b . Let (C, P_1, \dots, P_r) be an ear sequence. Clearly, G is not a cycle; thus $r \geq 1$. Also, each end of each ear P_i has degree at least three in G . Thus, every ear is an ab -path. Consequently, $a, b \in C$. By Lemma 4.1 (ii), each P_i has length three. Furthermore, by Lemma 4.1 (iii), the vertices a and b split C into two paths — each of length three. In particular, C is a 6-cycle. Observe that G is indeed obtained from the star $K_{1,r+2}$ by isomorphic leaf matching. \square

We will now use the balanced 2-cut property (Theorem 4.4) and the balanced 2-cut induction tool (Theorem 5.1), along with the above lemma, to prove our characterization of the class of extremal minimal bipartite matching covered graphs.

Theorem 1.3. [MAIN THEOREM: CHARACTERIZATION OF \mathcal{H}_2]

A graph G belongs to \mathcal{H}_2 if and only if it is obtained from a halin tree by isomorphic leaf matching.

Proof. The reverse implication is simply the main statement of Theorem 1.8 that we have already proved.

For the forward direction, we let $G[A, B]$ denote a member of \mathcal{H}_2 , and we proceed by induction on the order of G . First suppose that $|V_3| \leq 2$. Since $|A| = |B|$, it follows that $|V_3| \in \{0, 2\}$. If $|V_3| = 0$, then G is a cycle and $|V| = |V_2| = 2(m - n + 2) = 4$; thus G is C_4 . On the other hand, if $|V_3| = 2$, we invoke Lemma 6.1. In either case, we conclude that G is obtained from a halin tree by isomorphic leaf matching.

Now, suppose that $|V_3| \geq 3$. By Corollary 4.3 (iii), we note that $|E_3| = |V_3| - 2 \geq 1$, and we let a_1b_2 denote a 3-edge. By Theorem 4.4, there exists another 3-edge b_1a_2 such that

$\{a_1b_2, b_1a_2\}$ is a balanced 2-cut, say $\partial(X)$, where $\{a_1, b_1\} \subseteq X \subset V(G)$. Now, let G_1 and G_2 be the graphs obtained by 2-cut decomposition across $\partial(X) = \{a_1b_2, b_1a_2\}$. In particular, G_1 is obtained from $G[X]$ by adding the ear $a_1v_1u_1b_1$; likewise, G_2 is obtained from $G[\overline{X}]$ by adding the ear $a_2v_2u_2b_2$. See Figure 13 for an illustration.

By Theorem 5.1, each of G_1 and G_2 belongs to \mathcal{H}_2 . Since each of a_1, b_1, a_2 and b_2 has degree at least three in G , it follows that each of X and \overline{X} has at least four vertices. This implies that $6 \leq |V(G_i)| < |V(G)|$ for each $i \in \{1, 2\}$. Thus, by the induction hypothesis, each of G_1 and G_2 is obtained from a halin tree by isomorphic leaf matching; also, neither of them is isomorphic to C_4 . By Proposition 1.2, for each $i \in \{1, 2\}$, the graph $G_i - E_2(G_i)$ has precisely two components that are isomorphic halin trees, say T_i and T'_i . We let $\phi_i : V(T_i) \mapsto V(T'_i)$ denote the corresponding isomorphism as per the isomorphic leaf matching used to obtain G_i .




Figure 14: illustration for the proof of the Main Theorem

Since u_1v_1 is a 2-edge of G_1 , we may adjust notation so that $v_1, a_1 \in T_1$ and $u_1, b_1 \in T'_1$. See Figure 14. Likewise, we adjust notation so that $u_2, b_2 \in T_2$ and $v_2, a_2 \in T'_2$. Note that $\phi_1(v_1) = u_1$. Consequently, $\phi_1(a_1) = b_1$. Likewise, $\phi_2(b_2) = a_2$. We let $T := (T_1 - v_1) + (T_2 - u_2) + a_1b_2$ and $T' := (T'_1 - u_1) + (T'_2 - v_2) + a_2b_1$. Note that T and T' are halin trees. Observe that $\phi : V(T) \mapsto V(T')$, as defined below, is an isomorphism between T and T' .

$$\phi(v) := \begin{cases} \phi_1(v) & \text{if } v \in V(T_1) \\ \phi_2(v) & \text{if } v \in V(T_2) \end{cases}$$

Finally, note that G is obtained from T and T' by isomorphic leaf matching as per the above defined isomorphism. This completes the proof of the Main Theorem. \square

7 Other notions of extremality

We now shift our focus to the remaining notions of extremality — that is, the extremal classes $\mathcal{H}_3, \mathcal{H}_4, \mathcal{H}_0$ and \mathcal{H}_1 — in that order.

7.1 Characterization of \mathcal{H}_3

We begin by deducing the lower bound on the number of degree two vertices in a minimal bipartite matching covered graph solely in terms of n , that was stated in Subsection 1.1, from the lower bound of Lovász and Plummer (Corollary 3.5).

Corollary 7.1. *In a minimal bipartite matching covered graph, $|V_2| \geq \frac{n}{2} + 2$.*

Proof. Using the fact that $n = |V_2| + |V_3|$, and the handshaking lemma:

$$m - n = \left(\sum_{v \in V_3} \frac{d(v)}{2} + |V_2| \right) - n \geq \left(\frac{3}{2}|V_3| + |V_2| \right) - n = \frac{1}{2}|V_3|$$

We invoke Corollary 3.5 to infer that $|V_2| \geq 2(m - n + 2) \geq |V_3| + 4 = n - |V_2| + 4$. Consequently, $|V_2| \geq \frac{n}{2} + 2$. \square

We are now ready to prove our characterization for \mathcal{H}_3 . To this end, we shall enforce equality in the proof of the above corollary.

Theorem 1.4. [CHARACTERIZATION OF \mathcal{H}_3]

A graph G belongs to \mathcal{H}_3 if and only if it is obtained from a cubic halin tree by isomorphic leaf matching.

Proof. The reverse implication is simply statement (i) of Theorem 1.8 that we have already proved. It remains to prove the forward implication.

Let G be a member of \mathcal{H}_3 . That is, G is a minimal bipartite matching covered graph such that $|V_2| = \frac{n}{2} + 2$. Thus, G satisfies each inequality in the proof of Corollary 7.1 with equality. Firstly, $|V_2| = 2(m - n + 2)$; thus $G \in \mathcal{H}_2$. By the Main Theorem (1.3), G is obtained from a halin tree T by isomorphic leaf matching. Secondly, each vertex in V_3 has degree precisely three in G . This implies that each non-leaf of T is cubic. Thus, T is a cubic halin tree. \square

We end this section by noting the following proper containment that is established within the above proof.

Corollary 7.2. $\mathcal{H}_3 \subset \mathcal{H}_2$. \square

7.2 Characterization of \mathcal{H}_4

We will follow the same approach as in the previous subsection. We begin by deducing the upper bound $|E| \leq \frac{3n-6}{2}$ (see Lovász and Plummer [9, Theorem 4.2.3]), that was stated in Subsection 1.1, from the lower bound of Lovász and Plummer (Corollary 3.5).

Corollary 7.3. *A minimal bipartite matching covered graph, distinct from C_4 , has at most $\frac{3n-6}{2}$ edges.*

Proof. Let $G[A, B]$ denote a minimal bipartite matching covered graph distinct from C_4 . Note that $n \geq 6$ if and only if $n \leq \frac{3n-6}{2}$. Consequently, if G is a cycle, $m = n \leq \frac{3n-6}{2}$.

Now suppose that G is not a cycle; in other words, $|V_3| \geq 1$. Since $|A| = |B|$, we infer $|V_3| \geq 2$. Using Corollary 3.5 and the fact that $n = |V_2| + |V_3|$, we have $2(m - n + 2) \leq |V_2| = n - |V_3|$. Since $|V_3| \geq 2$, we get $2(m - n + 2) \leq n - 2$. On simplifying, we conclude $m \leq \frac{3n-6}{2}$. \square

We shall now enforce equality in the proof of the above corollary to prove our characterization of \mathcal{H}_4 .

Theorem 1.5. [CHARACTERIZATION OF \mathcal{H}_4]

A graph G belongs to \mathcal{H}_4 if and only if it is obtained from a star by isomorphic leaf matching.

Proof. The reverse implication is settled by statement (ii) of Theorem 1.8 for stars $K_{1,p}$ where $p \geq 3$, and the paragraph above it for $K_{1,2}$. It remains to prove the forward implication.

Let G be a member of \mathcal{H}_4 . That is, G is a minimal bipartite matching covered graph such that $|E| = \frac{3n-6}{2}$. If G is a cycle, then $n = 6$ and G is C_6 which is obtained from the star $K_{1,2}$ by isomorphic leaf matching.

Now, suppose that G is not a cycle. Consequently, G satisfies all of the inequalities in the last paragraph of the proof of Corollary 7.3 with equality. In particular, $|V_2| = 2(m - n + 2)$ and $|V_3| = 2$. In other words, G is a member of \mathcal{H}_2 that has precisely two vertices of degree at least three. Thus, by Lemma 6.1, we conclude that G is obtained from a star by isomorphic leaf matching. \square

We end this section with a couple of easy consequences. The first of them is the following proper containment that is already established within the above proof.

Corollary 7.4. $\mathcal{H}_4 - C_6 \subset \mathcal{H}_2$. \square

Note that Θ , shown in Figure 9, belongs to \mathcal{H}_3 as well as \mathcal{H}_4 . Conversely, let G be any graph in $\mathcal{H}_3 \cap \mathcal{H}_4$. Since $G \in \mathcal{H}_4$, by Theorem 1.5, G is obtained from a star $T := K_{1,p}$ by isomorphic leaf matching. On the other hand, since $G \in \mathcal{H}_3$, by Theorem 1.4, T is a cubic halin tree. Thus $p = 3$; in other words, G is Θ . This proves the following.

Corollary 7.5. $\mathcal{H}_4 \cap \mathcal{H}_3 = \{\Theta\}$. \square

7.3 Characterization of \mathcal{H}_0

Recall the bicontraction and the restricted bicontraction operations as defined in Subsection 1.4. In this subsection, we will first show that restricted bicontraction preserves 2-edge extremality. Then, we will use this to deduce our characterization of \mathcal{H}_0 .

In order to achieve the above, we first define the inverse of restricted bicontraction as well as of bicontraction. Observe that the bicontraction operation (when applicable) results in a vertex of degree at least two, whereas the restricted bicontraction operation (when applicable) results in a vertex of degree at least four.

Let G' be a graph with a vertex v' of degree two or more. We partition the edges of $\partial(v')$ into two sets F_1 and F_2 each of size at least one. Let G be obtained from $G' - v'$ by (i) introducing three new vertices v, v_1 and v_2 , (ii) adding two edges vv_1 and vv_2 , and (iii) for each $i \in \{1, 2\}$, adding an edge joining v and v_i for each $uv \in F_i$. See Figure 15 for an illustration. We say that G is obtained from G' by *bisplitting* the vertex v' . Observe that, in G , the vertex v has degree two and each of its neighbours v_1 and v_2 has degree at least two.

Furthermore, if v' has degree at least four and each of the sets F_1 and F_2 has size at least two, we say that G is obtained from G' by *restricted bisplitting* of the vertex v' . In this case, note that the vertex v has degree two and each of its neighbours v_1 and v_2 has degree at least three. See Figure 16 for an illustration.




Figure 15: an illustration of bicontraction and bisplitting

Observe that G' is obtained from G by bicontraction if and only if G may be obtained from G' by bisplitting. Likewise, G' is obtained from G by restricted bicontraction if and only if G may be obtained from G' by restricted bisplitting. The following lemma pertaining to bicontraction/bisplitting is easily proved, and may also be deduced from [2, Propositions 2.1 and 2.3].

Lemma 7.6. [BICONTRACTION PRESERVES MATCHING COVERED PROPERTY]

Let G be a graph that has a vertex of degree two, say v , that has two distinct neighbours each of which has degree two or more, and let $G' := G/v$. Then G is matching covered if and only if G' is matching covered. \square




Figure 16: an illustration of restricted bicontraction and restricted bisplitting

However, bicontraction may not preserve minimality. For instance, the graph shown in Figure 3 belongs to \mathcal{H} , but any application of the bicontraction operation results in a

graph that has a removable edge. On the other hand, the restricted bicontraction operation preserves minimality as well as 2-edge extremality, as proved below.

Theorem 7.7. [RESTRICTED BICONTRACTION PRESERVES 2-EDGE EXTREMALITY]

Let G be a bipartite graph that has a vertex of degree two, say v , each of whose neighbours has degree three or more, and let $G' := G/v$. Then the following statements hold:

- (i) G is matching covered if and only if G' is matching covered;
- (ii) furthermore, G is minimal if and only if G' is minimal; and
- (iii) finally, G belongs to \mathcal{H}_0 if and only if G' belongs to \mathcal{H}_0 .

Proof. We adopt notation from the definition of restricted bisplitting, as shown in Figure 16. Observe that (i) follows immediately from Lemma 7.6. Henceforth, we assume that G is matching covered, or equivalently, that G' is matching covered. We will prove the other two statements one-by-one.

7.7.1. G is minimal if and only if G' is minimal.

Proof. First suppose that G' is not minimal, and let e be a removable edge. Note that $d_{G-e}(v_i) \geq d_G(v_i) - 1 \geq 2$ since the operation is a restricted bicontraction. Consequently, $G' - e$ is obtained from $G - e$ by bicontracting v . Thus, by Lemma 7.6, $G - e$ is matching covered. In other words, G is not minimal.

Conversely, suppose that G is not minimal, and let e be a removable edge. Note that $e \notin \partial(v)$ since $d(v) = 2$. Now, observe that $G' - e$ is obtained from $G - e$ by bicontracting v . Thus, by Lemma 7.6, $G' - e$ is matching covered. Consequently, G' is not minimal. \square

Henceforth, we assume that G is a minimal bipartite matching covered graph, or equivalently, that G' is a minimal bipartite matching covered graph. Recall that G belongs to \mathcal{H}_0 if $|E_2(G)| = m - n + 2$.

7.7.2. G belongs to \mathcal{H}_0 if and only if G' belongs to \mathcal{H}_0 .

Proof. Let $n' := |V(G')|$ and $m' := |E(G')|$. Note that $m' = m - 2$ and $n' = n - 2$. Thus, $m' - n' = m - n$. By definition of restricted bicontraction operation, $d_G(v_1) \geq 3$ and $d_G(v_2) \geq 3$; subsequently $d_{G'}(v') \geq 3$. These observations imply that $|E_2(G')| = |E_2(G)|$. \square

This completes the proof of Theorem 7.7 \square

Recall the definition of partial retract from Subsection 1.4. The following is an immediate consequence of the above theorem.

Corollary 7.8. A graph G belongs to \mathcal{H}_0 if and only if its partial retract \widehat{G} belongs to \mathcal{H}_0 . \square

We are now ready to prove our characterization of the extremal class \mathcal{H}_0 in terms of the extremal class \mathcal{H}_2 , as restated below.

Theorem 1.9. [CHARACTERIZATION OF \mathcal{H}_0]

A graph G , distinct from C_4 , belongs to \mathcal{H}_0 if and only if its partial retract \widehat{G} belongs to \mathcal{H}_2 .

Proof. First suppose that \widehat{G} belongs to \mathcal{H}_2 . Observe that either $\widehat{G} = G$ or \widehat{G} has a vertex of degree at least four. In either case, $\widehat{G} \neq C_4$. By the containment established in Corollary 4.2, we infer that $\widehat{G} \in \mathcal{H}_0$. Now, by Corollary 7.8, we conclude that $G \in \mathcal{H}_0$.

Now suppose that G belongs to \mathcal{H}_0 . We invoke Corollary 7.8 to infer that $\widehat{G} \in \mathcal{H}_0$. Thus, by definition, $|E_2(\widehat{G})| = \widehat{m} - \widehat{n} + 2$, where $\widehat{n} := |V(\widehat{G})|$ and $\widehat{m} := |E(\widehat{G})|$. By definition of partial retract, the subgraph of \widehat{G} induced by its degree two vertices has no isolated vertex. Applying the handshaking lemma to this subgraph gives us: $|V_2(\widehat{G})| \leq 2|E_2(\widehat{G})| = 2(\widehat{m} - \widehat{n} + 2)$. However, by the lower bound established in Corollary 3.5, $|V_2(\widehat{G})| \geq 2(\widehat{m} - \widehat{n} + 2)$. Thus, $|V_2(\widehat{G})| = 2(\widehat{m} - \widehat{n} + 2)$; by definition, $\widehat{G} \in \mathcal{H}_2$. \square

We now switch our attention to the only remaining extremal class.

7.4 Characterization of \mathcal{H}_1

We begin by deducing a lower bound on the number of 2-edges in a minimal bipartite matching covered graph solely in terms of n (see Lovász and Plummer [9, Lemma 4.2.4 (b)]), that was stated in Subsection 1.1, from the lower bound of Lovász and Plummer (Theorem 3.4).

Corollary 7.9. *In a minimal bipartite matching covered graph, $|E_2| \geq \frac{n+10}{6}$.*

Proof. Let G be a minimal bipartite matching covered graph. By Theorem 3.4:

$$|E_2| \geq m - n + 2 \tag{1}$$

Now, using the handshaking lemma, and the fact that $d(v) \geq 3$ for each $v \in V_3$, we have: $2m \geq 3|V_3| + 2|V_2| = 3n - |V_2|$. In particular:

$$2m \geq 3n - |V_2| \tag{2}$$

By counting the edges incident with vertices of degree two in two different ways, we have: $2|V_2| = 2|E_2| + |E_{23}| = |E_2| + m - |E_3|$. Now, using the fact that $|E_3| \geq 0$, we have:

$$|E_2| + m \geq 2|V_2| \tag{3}$$

Now, by scaling and adding the above inequalities as (1) \times 5 + (2) \times 2 + (3) \times 1, we arrive at the desired inequality: $6|E_2| \geq n + 10$. \square

Next, we invite the reader to make a simple observation pertaining to the restricted bicontraction operation.

Lemma 7.10. *If a graph G has a vertex of degree two, say v , each of whose neighbours has degree at least three, then $|E_3(G)| = |E_3(G/v)|$. Consequently, $|E_3(G)| = |E_3(\widehat{G})|$. \square*

We are now ready to prove our characterization of the extremal class \mathcal{H}_1 in terms of the extremal class \mathcal{H}_4 , as restated below.

Theorem 1.10. [CHARACTERIZATION OF \mathcal{H}_1]

A graph G belongs to \mathcal{H}_1 if and only if its partial retract \widehat{G} belongs to \mathcal{H}_4 and $\Delta(G) = 3$.

Proof. For the reverse direction, let G be a graph such that $\widehat{G} \in \mathcal{H}_4$ and $\Delta(G) = 3$. In what follows, we shall demonstrate that all of the inequalities that appear in the proof of Corollary 7.3 hold with equality. Firstly, by the characterization of \mathcal{H}_4 obtained in Theorem 1.5, we infer that $|E_3(\widehat{G})| = 0$. Thus, by Lemma 7.10, $|E_3(G)| = |E_3(\widehat{G})| = 0$. Secondly, by the containment established in Corollary 7.4, $\widehat{G} \in \mathcal{H}_2$. By Theorem 1.9, $G \in \mathcal{H}_0$. In particular, $|E_2(G)| = m - n + 2$. Lastly, as $\Delta(G) = 3$, clearly $d_G(v) = 3$ for each $v \in V_3$. The reader may thus verify that equality holds in all of the inequalities discussed in the proof of Corollary 7.9. Hence, $|E_2(G)| = \frac{n+10}{6}$ and $G \in \mathcal{H}_1$.

For the forward direction, let G be a graph in \mathcal{H}_1 . Consequently, G satisfies all of the inequalities that appear in the proof of Corollary 7.9 with equality. Firstly, $|E_2(G)| = m - n + 2$; thus $G \in \mathcal{H}_0$. Consequently, by Theorem 1.9, $\widehat{G} \in \mathcal{H}_2$. Secondly, $|E_3(G)| = 0$. Thus, by Lemma 7.10, $|E_3(\widehat{G})| = 0$. By Corollary 4.3 (iii), we deduce that $|V_3(\widehat{G})| = 2$. Hence, by Lemma 6.1 and Theorem 1.8 (ii), we conclude that $\widehat{G} \in \mathcal{H}_4$. Finally, $d_G(v) = 3$ for each $v \in V_3(G)$. Thus, $\Delta(G) \leq 3$. Note that if G is a cycle then $n = m = E_2 = \frac{n+10}{6}$ which implies that $n = 2$; a contradiction. Thus, G is not a cycle; whence $\Delta(G) = 3$. \square

We end this section, and this paper, with a couple of easy consequences. The first of them is the following proper containment that is already established within the above proof.

Corollary 7.11. $\mathcal{H}_1 \subset \mathcal{H}_0$. \square

Note that Θ , shown in Figure 9, belongs to \mathcal{H}_1 as well as \mathcal{H}_2 . Conversely, let $G \in \mathcal{H}_1 \cap \mathcal{H}_2$. Since $G \in \mathcal{H}_2$, by Lemma 4.1 (i), E_2 is a perfect matching of $G[V_2]$ which implies that there is no vertex of degree two each of whose neighbours has degree three or more. Thus, $\widehat{G} = G$. On the other hand, since $G \in \mathcal{H}_1$, by Theorem 1.10, $G \in \mathcal{H}_4$. Furthermore, $\Delta(G) = 3$. Observe that Θ is the only graph in \mathcal{H}_4 with maximum degree three. Thus, $G = \Theta$; this proves the following.

Corollary 7.12. $\mathcal{H}_1 \cap \mathcal{H}_2 = \{\Theta\}$. \square

References

- [1] J. A. Bondy and U. S. R. Murty. *Graph Theory*. Springer, 2008.
- [2] M. H. Carvalho, C. L. Lucchesi, and U. S. R. Murty. Graphs with independent perfect matchings. *J. Graph Theory*, 48:19–50, 2005.
- [3] Gabriel A Dirac. Minimally 2-connected graphs. 1967.

- [4] Rudolf Halin. Studies on minimally n -connected graphs. In *Combinatorial Mathematics and its Applications (Proc. Conf., Oxford, 1969)*, pages 129–136, 1971.
- [5] Gábor Hetyei. Rectangular configurations which can be covered by 2×1 rectangles. *Pécsi Tan. Foisk. Közl.*, 8:351–367, 1964.
- [6] Arthur Hoffmann-Ostenhof, Kenta Noguchi, and Kenta Ozeki. On homeomorphically irreducible spanning trees in cubic graphs. *Journal of Graph Theory*, 89(2):93–100, 2018.
- [7] DV Karpov. Minimal k -connected graphs with minimal number of vertices of degree k . *Journal of Mathematical Sciences*, 212:666–682, 2016.
- [8] Charles H. C. Little. A theorem on connected graphs in which every edge belongs to a 1-factor. *Journal of the Australian Mathematical Society*, 18(4):450–452, 1974.
- [9] L. Lovász and M. D. Plummer. *Matching Theory*. Number 29 in Annals of Discrete Mathematics. Elsevier Science, 1986.
- [10] James G Oxley. On some extremal connectivity results for graphs and matroids. *Discrete Mathematics*, 41(2):181–198, 1982.
- [11] H. Whitney. 2-isomorphic graphs. *American Journal of Mathematics*, 57:245–254, 1933.