

CORRESPONDENCES BETWEEN GRAPH C^* -ALGEBRAS

RASMUS BENTMANN AND RALF MEYER

ABSTRACT. We describe proper correspondences from graph C^* -algebras to arbitrary C^* -algebras by K-theoretic data. If the target C^* -algebra is a graph C^* -algebra as well, we may lift an isomorphism on a certain invariant to correspondences back and forth that are inverse to each other up to ideal-preserving homotopy. This implies a classification theorem for purely infinite graph C^* -algebras up to stable isomorphism.

1. INTRODUCTION

Graph C^* -algebras form a class of C^* -algebras that are well-behaved in various ways. For instance, they are all nuclear and satisfy the Universal Coefficient Theorem. Moreover, their ideal structure and K-theory is well-understood in terms of the underlying graph, and the K-theory has some special properties: the K_1 -group of a graph C^* -algebra is always free, and the exponential map associated to a (gauge-invariant) ideal in a graph C^* -algebra always vanishes. In addition, every simple graph C^* -algebra is either approximately finite-dimensional or purely infinite. (See [32] for (references for) these well known facts). Hence it is natural to explore what the general classification theory of C^* -algebras has to say about C^* -algebras with these features. Among other places, this has been done in [15–19, 21] for the mixed cases and in [4, 8, 9, 28] for the (strongly) purely infinite case. While many results have been proved, a classification of nonunital graph C^* -algebras along these lines is not yet in reach.

The approach to classification using the symbolic dynamics associated to the underlying graph has been very successful recently, culminating in the complete classification of unital graph C^* -algebras (see [20] and the references therein, and [34, 35] for earlier results). Unitality is, however, a strong restriction for graph C^* -algebras because the unitisation of a graph C^* -algebra need not be a graph C^* -algebra. A graph C^* -algebra is unital if and only if the underlying graph has finitely many vertices. It seems difficult to generalise the ideas of [20] to the nonunital case.

The present article introduces a new approach to classify graph C^* -algebras. The goal is to understand (proper) C^* -correspondences out of graph C^* -algebras well enough to be able to establish existence and uniqueness results for proper C^* -correspondences lifting isomorphisms on suitable invariants. We achieve this with uniqueness up to (ideal-preserving) homotopy. This is the main technical achievement of the article. Uniqueness up to homotopy is, however, too weak for Elliott’s Approximate Intertwining Argument. Hence our results do not suffice for a general classification of graph C^* -algebras up to stable isomorphism. For purely infinite graph C^* -algebras, however, it implies an equivalence in Kirchberg’s KK-theory, which is enough to apply Kirchberg’s Classification Theorem [24]. So we get a classification result for all purely infinite (possibly nonunital) graph C^* -algebras

2010 *Mathematics Subject Classification.* Primary 46L35; Secondary 18E30, 19K35.

Supported by the DFG grant “Classification of non-simple purely infinite C^* -algebras.” This work is part of the project Graph Algebras partially supported by EU grant HORIZON-MSCA-SE-2021 Project 101086394.

up to stable isomorphism. In general, our method only gives a classification theorem for graph C^* -algebras up to ideal-preserving homotopy equivalence.

We now describe the sections of this article. In Section 2, we collect some basic results about proper C^* -correspondences, their induced maps on ideal lattices and their relation to $*$ -homomorphisms into stabilised C^* -algebras. Section 3 studies proper C^* -correspondences out of graph C^* -algebras. The main result says that any homomorphism between the monoids of equivalence classes of projections lifts to a proper C^* -correspondence if the domain is a graph C^* -algebra. Section 4 relates the monoid of equivalence classes of projections to the ideal lattice and the ordered K_0 -groups of ideals, especially in the case of graph C^* -algebras. Section 6 adds information related to K_1 to the picture. We show that a pair of maps on equivalence classes of projections and on K_1 is realised by a proper C^* -correspondence if and only if a certain variant of the obstruction class of [9] vanishes. In Section 7, we combine our new results with Kirchberg's Classification Theorem to classify purely infinite graph C^* -algebras. In Sections 8 and 9, we allow arbitrary graph C^* -algebras again and prove an ideal-preserving homotopy equivalence instead of an isomorphism under the same assumptions as in Section 7. More precisely, Section 8 only proves a homotopy equivalence preserving gauge-invariant ideals. Section 9 contains some extra arguments that allow to prove an ideal-preserving homotopy equivalence for all ideals, gauge-invariant or not.

2. CORRESPONDENCES VERSUS HOMOMORPHISMS INTO STABILISATIONS

This section contains results for general σ -unital C^* -algebras, which we are going to specialise to graph C^* -algebras in later sections. Let A and B be two C^* -algebras. We are going to show that proper C^* -correspondences between A and B are "equivalent" to $*$ -homomorphisms from A to the C^* -stabilisation $B \otimes \mathbb{K}$ when A is σ -unital. Here $\mathbb{K} := \mathbb{K}(\ell^2\mathbb{N})$ is the C^* -algebra of compact operators on a separable Hilbert space. If A is C^* -stable and both A and B are σ -unital, then full, proper correspondences are equivalent to nondegenerate $*$ -homomorphisms $A \rightarrow B \otimes \mathbb{K}$. Let $\mathbb{I}(A)$ be the lattice of closed ideals of A . A $*$ -homomorphism $\varphi: A \rightarrow B \otimes \mathbb{K}$ induces a map $\mathbb{I}(\varphi): \mathbb{I}(A) \rightarrow \mathbb{I}(B)$. We describe it in terms of the proper C^* -correspondence corresponding to φ . All this is either easy or follows from the proofs by Mingo and Phillips [29] of the Kasparov Stabilisation Theorem and the Brown–Green–Rieffel Theorem.

First we recall some definitions. An A, B -correspondence is a (right) Hilbert B -module \mathcal{H} with a nondegenerate $*$ -homomorphism φ from A to the C^* -algebra $\mathbb{B}(\mathcal{H})$ of adjointable operators on \mathcal{H} . It is *proper* if $\varphi(A)$ is contained in the C^* -algebra $\mathbb{K}(\mathcal{H})$ of compact operators on \mathcal{H} . Let $\langle \mathcal{H} | \mathcal{H} \rangle \triangleleft B$ be the ideal generated by the inner products $\langle x | y \rangle$ for $x, y \in \mathcal{H}$. Call \mathcal{H} *full* if $\langle \mathcal{H} | \mathcal{H} \rangle = B$. An *isomorphism* between two A, B -correspondences \mathcal{H}_1 and \mathcal{H}_2 is a unitary operator $u: \mathcal{H}_1 \xrightarrow{\sim} \mathcal{H}_2$ of right Hilbert B -modules that intertwines the left actions of A , that is, $u(a \cdot \xi) = a \cdot (u\xi)$ for all $a \in A$, $\xi \in \mathcal{H}_1$. The *direct sum* $\mathcal{H}_1 \oplus \mathcal{H}_2$ of two correspondences is defined in the obvious way. It is again proper if \mathcal{H}_1 and \mathcal{H}_2 are proper. And

$$(2.1) \quad \langle \mathcal{H}_1 \oplus \mathcal{H}_2 | \mathcal{H}_1 \oplus \mathcal{H}_2 \rangle = \langle \mathcal{H}_1 | \mathcal{H}_1 \rangle + \langle \mathcal{H}_2 | \mathcal{H}_2 \rangle,$$

so that $\mathcal{H}_1 \oplus \mathcal{H}_2$ is full if \mathcal{H}_1 or \mathcal{H}_2 is full. The direct sum operation descends to isomorphism classes, is associative, and has the zero element $\{0\}$. Thus isomorphism classes of proper A, B -correspondences with the operation \oplus induced by the direct sum of correspondences form a monoid. We denote it by $\mathcal{P}(A, B)$.

An A, B -correspondence \mathcal{H} induces a map $\mathbb{I}(\mathcal{H}): \mathbb{I}(A) \rightarrow \mathbb{I}(B)$ by mapping $I \triangleleft A$ to the ideal $\langle I\mathcal{H} | I\mathcal{H} \rangle \triangleleft B$ generated by the inner products $\langle ax | by \rangle$ with $a, b \in I$,

$x, y \in \mathcal{H}$. Since $I^* \cdot I = I$, we have

$$\langle I\mathcal{H} | I\mathcal{H} \rangle = \langle \mathcal{H} | I\mathcal{H} \rangle = \langle I\mathcal{H} | \mathcal{H} \rangle.$$

Therefore, the map $\mathbb{I}(\mathcal{H})$ commutes with arbitrary suprema, that is,

$$\mathbb{I}(\mathcal{H})\left(\bigvee_{I \in S} I\right) = \left(\bigvee_{I \in S} \mathbb{I}(\mathcal{H})(I)\right)$$

for any set S of ideals in A , where $\bigvee_{I \in S} I = \overline{\sum_{I \in S} I}$ is the supremum of S in $\mathbb{I}(A)$. As a consequence, the map $\mathbb{I}(\mathcal{H})$ is monotone and preserves the zero ideals. It need not commute with intersections, however. And $\mathbb{I}(\mathcal{H})(A) = \langle \mathcal{H} | \mathcal{H} \rangle$ is only equal to B if \mathcal{H} is full.

Let $\mathcal{H}^\infty := \mathcal{H} \otimes \ell^2(\mathbb{N})$ or, equivalently, the direct sum of countably many copies of \mathcal{H} . There is an isomorphism $\mathbb{K}(\mathcal{H}) \otimes \mathbb{K} \xrightarrow{\sim} \mathbb{K}(\mathcal{H}^\infty)$, which maps $S \otimes T \in \mathbb{K}(\mathcal{H}) \otimes \mathbb{K}$ to the unique operator $S \otimes T$ on $\mathcal{H} \otimes \ell^2\mathbb{N}$ that satisfies $(S \otimes T)(x \otimes y) := S(x) \otimes T(y)$ for all $x \in \mathcal{H}$, $y \in \ell^2\mathbb{N}$. In particular, $B \otimes \mathbb{K} \cong \mathbb{K}(B^\infty)$ because $B \cong \mathbb{K}(B)$ when we view B as a Hilbert B -module in the obvious way.

A *-homomorphism $\varphi: A \rightarrow B \otimes \mathbb{K}$ induces a proper A, B -correspondence \mathcal{H}_φ as follows. The underlying Hilbert B -module is $\mathcal{H} := \varphi(A) \cdot B^\infty \subseteq B^\infty$, and A acts nondegenerately on \mathcal{H} through φ . If φ is nondegenerate, then $\mathcal{H} = B^\infty$ and thus \mathcal{H}_φ is full. The next proposition reverses this construction if A is σ -unital, that is, if A has a sequential approximate unit.

Proposition 2.2. *If A is σ -unital, then any proper A, B -correspondence \mathcal{H} is isomorphic to \mathcal{H}_φ for a *-homomorphism $\varphi: A \rightarrow B \otimes \mathbb{K}$. If A and B are σ -unital and A is C^* -stable, then any proper, full A, B -correspondence \mathcal{H} is isomorphic to \mathcal{H}_φ for a nondegenerate *-homomorphism $\varphi: A \rightarrow B \otimes \mathbb{K}$.*

Proof. First assume only that A is σ -unital. Since the map $A \rightarrow \mathbb{K}(\mathcal{H})$ is nondegenerate, it maps a sequential approximate unit in A to one in $\mathbb{K}(\mathcal{H})$. The Hilbert module \mathcal{H} is countably generated if and only if $\mathbb{K}(\mathcal{H})$ is σ -unital. So \mathcal{H} is countably generated. Then the Kasparov Stabilisation Theorem gives a unitary operator $\mathcal{H} \oplus B^\infty \cong B^\infty$ and, in particular, an adjointable isometry $S: \mathcal{H} \rightarrow B^\infty$. Then $\varphi(a) := SaS^*$ for $a \in A$ defines a *-homomorphism $\varphi: A \rightarrow B \otimes \mathbb{K}$. By construction, $\varphi(A) \cdot B^\infty = S(\mathcal{H})$ and S is an isomorphism of A, B -correspondences from \mathcal{H} to \mathcal{H}_φ .

Now assume, in addition, that $A = A_0 \otimes \mathbb{K}$ is C^* -stable and that B is σ -unital. The nondegenerate left action of A on \mathcal{H} induces an action of $\mathbb{K} \subseteq \mathcal{M}(A \otimes \mathbb{K})$. Let $p \in \mathbb{K}$ be a rank-1 projection and $\mathcal{H}_0 := (1 \otimes p)\mathcal{H}$. The matrix units in \mathbb{K} allow to identify $\mathcal{H} \cong \mathcal{H}_0^\infty$. Since \mathcal{H} is full, so is \mathcal{H}_0 . Since \mathcal{H} is countably generated, so is \mathcal{H}_0 . Since $B = \mathbb{K}(B)$ is σ -unital, B is a countably generated Hilbert B -module. Hence the proof of the Brown–Green–Rieffel Theorem in [29] gives a unitary operator $S: \mathcal{H} \xrightarrow{\sim} \mathcal{H}_0^\infty \xrightarrow{\sim} B^\infty$. Now the *-homomorphism $\varphi: A \rightarrow B \otimes \mathbb{K}$, $a \mapsto SaS^*$, is nondegenerate, and S is an isomorphism from \mathcal{H} to \mathcal{H}_φ . \square

If $A = \mathbb{C}$, then a proper \mathbb{C}, B -correspondence is just a Hilbert B -module \mathcal{H} with $\text{id}_\mathcal{H} \in \mathbb{K}(\mathcal{H})$. We call such Hilbert B -modules *proper*. Let $\mathbb{M}_\infty(B) = \bigcup_{n=0}^\infty \mathbb{M}_n(B)$ denote the set of infinite matrices over B with finitely many nonzero entries.

Theorem 2.3. *Let B be a C^* -algebra. A Hilbert B -module \mathcal{H} is proper if and only if there is a projection $p \in \mathbb{M}_\infty(B)$ with $\mathcal{H} \cong p \cdot B^\infty$. Let $p_1, p_2 \in \mathbb{M}_\infty(B)$ be projections. Unitary operators $p_1 B^\infty \rightarrow p_2 B^\infty$ are in bijection with Murray–von Neumann equivalences between p_1 and p_2 , that is, $u \in \mathbb{M}_\infty(B)$ with $uu^* = p_2$ and $u^*u = p_1$.*

Proof. This follows from well known results if B is unital (see [38, Theorem 15.4.2]). The statements that we claim still hold if B is nonunital (see also [5, Proposition 4.2]). We include a proof here for the convenience of the reader.

First assume that $\mathcal{H} \cong pB^m$ for a projection $p \in \mathbb{M}_m(B)$. We let $V: \mathcal{H} \rightarrow B^m$ be the resulting isometric embedding. It is adjointable with the adjoint $B^m \rightarrow \mathcal{H}$, $x \mapsto p \cdot x$. The compositions V^*V and VV^* are the identity map on \mathcal{H} and left multiplication by p , respectively. The latter is a compact operator on B^m because $\mathbb{K}(B^m) = \mathbb{M}_m(B)$. Since VV^* is compact, so is $V = VV^*V$ and hence also $V^*V = \text{id}_{\mathcal{H}}$.

Conversely, assume that $\text{id}_{\mathcal{H}}$ is compact. Then there is a finite-rank approximation $\sum_{j=1}^m |x_j\rangle\langle y_j|$ with, say,

$$\left\| \sum_{j=1}^m |x_j\rangle\langle y_j| - \text{id}_{\mathcal{H}} \right\| < 1$$

Then $\sum_{j=1}^m |x_j\rangle\langle y_j|$ is invertible in $\mathbb{K}(\mathcal{H})$; let $s \in \mathbb{K}(\mathcal{H})$ be its inverse. Let

$$y := (|y_j|)_{j=1, \dots, m}: \mathcal{H} \rightarrow B^m, \quad x := (|x_j\rangle)_{j=1, \dots, m}: B^m \rightarrow \mathcal{H};$$

these are adjointable maps with $(s \circ x) \circ y = \text{id}_{\mathcal{H}}$. Then $e = y \circ (s \circ x): B^m \rightarrow B^m$ is idempotent, and adjointable as a composition of adjointable operators. Hence its range is complementable. The operators e and y have the same range and y^* is surjective because $(s \circ x) \circ y = \text{id}_{\mathcal{H}}$. As a consequence, the operator y has a polar decomposition. This gives an isometry $V: \mathcal{H} \rightarrow B^m$. Since $\text{id}_{\mathcal{H}}$ is compact, so is $V = V \circ \text{id}_{\mathcal{H}}$ and hence VV^* . Thus VV^* is left multiplication by some $p \in \mathbb{M}_m(B)$. This is a projection whose image is isomorphic to \mathcal{H} via V . \square

Thus $\mathcal{P}(B) := \mathcal{P}(\mathbb{C}, B)$ is canonically isomorphic to the monoid of Murray–von Neumann equivalence classes of projections in $\mathbb{M}_{\infty}(B)$ with the usual operation \oplus , even if B is not unital.

Proper correspondences may be composed by tensor product. This descends to a category structure on isomorphism classes of proper correspondences. In particular, there is a canonical composition $\mathcal{P}(A) \times \mathcal{P}(A, B) \rightarrow \mathcal{P}(B)$. It is additive in both variables. Equivalently, a proper A, B -correspondence \mathcal{H} induces a monoid homomorphism $\mathcal{H}_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$, $[\mathcal{M}] \mapsto [\mathcal{M} \otimes_A \mathcal{H}]$, and these homomorphisms satisfy $(\mathcal{H}_1)_* + (\mathcal{H}_2)_* = (\mathcal{H}_1 \oplus \mathcal{H}_2)_*$.

Next, we define “unitary equivalence” of $*$ -homomorphisms $A \rightarrow B \otimes \mathbb{K}$ in such a way that it corresponds exactly to unitary equivalence of the corresponding proper A, B -correspondences. Let $R_{\varphi} := \varphi(A) \cdot B \otimes \mathbb{K}$. This is a right ideal. We also view it as a Hilbert module over $B \otimes \mathbb{K}$.

Definition 2.4. Let $\varphi_0, \varphi_1: A \rightarrow B \otimes \mathbb{K}$ be two $*$ -homomorphisms. A *unitary equivalence* from φ_0 to φ_1 is a unitary operator $u: R_{\varphi_0} \xrightarrow{\sim} R_{\varphi_1}$ with $u\varphi_0(a)u^* = \varphi_1(a)$ for all $a \in A$, as operators on the Hilbert B -module R_{φ_1} .

A unitary multiplier \bar{u} of $B \otimes \mathbb{K}$ with $u\varphi_0(a)u^* = \varphi_1(a)$ in $B \otimes \mathbb{K}$ for all $a \in A$ restricts to a unitary operator $u: R_{\varphi_0} \xrightarrow{\sim} R_{\varphi_1}$ that is a unitary equivalence from φ_0 to φ_1 . Thus we allow more unitary equivalences than usual.

The argument that identifies the multiplier algebra of $\mathbb{K}(\mathcal{H})$ for a Hilbert B -module \mathcal{H} with $\mathbb{B}(\mathcal{H})$ gives a bijection between unitary equivalences $\varphi_1 \rightarrow \varphi_2$ and isomorphisms of correspondences $\mathcal{H}_{\varphi_1} \rightarrow \mathcal{H}_{\varphi_2}$. Thus isomorphism classes of proper A, B -correspondences are in bijection with unitary equivalence classes of $*$ -homomorphisms $A \rightarrow B \otimes \mathbb{K}$. And isomorphism classes of proper, full $A \otimes \mathbb{K}, B$ -correspondences are in bijection with unitary equivalence classes of nondegenerate $*$ -homomorphisms $A \otimes \mathbb{K} \rightarrow B \otimes \mathbb{K}$, for the standard notion of unitary equivalence.

A $*$ -homomorphism $\varphi: A \rightarrow B \otimes \mathbb{K}$ induces a map $\mathbb{I}(\varphi): \mathbb{I}(A) \rightarrow \mathbb{I}(B \otimes \mathbb{K}) \cong \mathbb{I}(B)$ by mapping $I \triangleleft A$ to the ideal in $B \otimes \mathbb{K}$ generated by $\varphi(I) \subseteq B \otimes \mathbb{K}$. A routine computation identifies this map with the map $\mathbb{I}(\mathcal{H}_{\varphi})$ defined above.

For C*-algebra classification, the equivalence relation of unitary equivalence for *-homomorphisms is too fine, even with our mild modification. We should use asymptotic or approximate unitary equivalence or homotopy. A *homotopy* from φ_0 to φ_1 is a *-homomorphism $\Phi: A \rightarrow C([0, 1], B \otimes \mathbb{K})$ with $\text{ev}_t \circ \Phi = \varphi_t$ for $t = 0, 1$. An *asymptotic unitary equivalence* from φ_0 to φ_1 is a family of unitary operators $u_t: R_{\varphi_0} \xrightarrow{\sim} R_{\varphi_1}$ for $t \in [0, \infty)$ that is *-strongly continuous in the sense that $t \mapsto u_t(x) \in R_{\varphi_1}$ and $t \mapsto u_t^*(y) \in R_{\varphi_0}$ are norm-continuous for $x \in R_{\varphi_0}$, $y \in R_{\varphi_1}$ and that satisfies

$$\lim_{t \rightarrow \infty} u_t \varphi_0(a) u_t^* = \varphi_1(a)$$

for all $a \in A$, as operators on the Hilbert B -module R_{φ_1} . An *approximate unitary equivalence* from φ_0 to φ_1 is a family of unitary operators $u_n: R_{\varphi_0} \xrightarrow{\sim} R_{\varphi_1}$ for $n \in \mathbb{N}$ that satisfies

$$\lim_{n \rightarrow \infty} u_n \varphi_0(a) u_n^* = \varphi_1(a)$$

for all $a \in A$. For proper A, B -correspondences, these three notions correspond to the following:

Definition 2.5. A *homotopy* from \mathcal{H}_0 to \mathcal{H}_1 is an $A, C([0, 1], B)$ -correspondence \mathcal{H} together with isomorphisms $\mathcal{H} \otimes_{\text{ev}_t} B \cong \mathcal{H}_t$ for $t = 0, 1$. An *asymptotic isomorphism* from \mathcal{H}_0 to \mathcal{H}_1 is a *-strongly continuous family of unitary operators $(u_t)_{t \in [0, \infty)}$ between the underlying Hilbert B -modules that satisfies

$$\lim_{t \rightarrow \infty} u_t(a \cdot_0 u_t^* x) = a \cdot_1 x$$

for all $a \in A$, $x \in \mathcal{H}_1$; here \cdot_i denotes the left action of A on \mathcal{H}_i . An *approximate isomorphism* is defined similarly, replacing the indexing set $[0, \infty)$ by \mathbb{N} .

3. CORRESPONDENCES OUT OF A GRAPH C*-ALGEBRA

Let $G = (r, s: E \rightrightarrows V)$ be a directed graph with sets of vertices and edges V and E , respectively, and range and source maps r and s . The *graph C*-algebra of G* is the universal C*-algebra $C^*(G)$ generated by mutually orthogonal projections p_v for $v \in V$ and partial isometries t_e for $e \in E$ subject to the relations

$$(CK1) \quad t_e^* t_e = p_{s(e)} \text{ for all } e \in E;$$

$$(CK2) \quad \sum_{r(e)=v} t_e t_e^* = p_v \text{ for all } v \in V \text{ with } 0 < |r^{-1}(v)| < \infty;$$

$$(CK3) \quad t_e t_e^* \leq p_{r(e)} \text{ for all } e \in E \text{ and } t_e t_f^* = 0 \text{ for all } e, f \in E \text{ with } e \neq f.$$

We call $v \in V$ *regular* if $0 < |r^{-1}(v)| < \infty$; let $V_{\text{reg}} \subseteq V$ be the subset of regular vertices. The graph G is *regular* if all vertices are regular. Then the relation (CK2) is imposed for all vertices. More generally, the graph G is *row-finite* if $|r^{-1}(v)| < \infty$ for all $v \in V$. Then the relation (CK3) is redundant. Any graph C*-algebra $C^*(G)$ is stably isomorphic to $C^*(G')$ for a regular graph G' , which is constructed in [30] by “adding tails” to G . Since we only care about graph C*-algebras up to stable isomorphism, we may as well assume G to be regular. Actually, our methods work equally well for row-finite graphs. We assume all graphs in the following to be row-finite, sometimes tacitly.

Theorem 3.1 ([3, Theorems 3.5 and 7.1]). *Let G be a row-finite graph. Each projection $q \in \mathbb{M}_\infty(C^*(G))$ is equivalent to a finite sum $\sum_{v \in V} c_v p_v$ of vertex projections with $(c_v)_{v \in V} \in \mathbb{N}[V]$, that is, $c_v \in \mathbb{N}$ for all $v \in V$ and $c_v \neq 0$ only for finitely many $v \in V$.*

Theorem 3.2. *Let G be a row-finite graph and B a C*-algebra. Proper $C^*(G), B$ -correspondences are equivalent to families $((\mathcal{H}_v)_{v \in V}, (U_v)_{v \in V_{\text{reg}}})$, where each \mathcal{H}_v is a proper Hilbert B -module and U_v is a unitary operator $\bigoplus_{e \in E^v} \mathcal{H}_{s(e)} \xrightarrow{\sim} \mathcal{H}_v$. Here “equivalent” means an equivalence of categories, where arrows from $((\mathcal{H}_{1,v}), (U_{1,v}))$*

to $((\mathcal{H}_{2,v}), (U_{2,v}))$ are families of unitaries $W_v: \mathcal{H}_{1,v} \xrightarrow{\sim} \mathcal{H}_{2,v}$ for $v \in V$ that make the following diagrams commute for all $v \in V_{\text{reg}}$:

$$(3.3) \quad \begin{array}{ccc} \bigoplus_{e \in E^v} \mathcal{H}_{1,s(e)} & \xrightarrow{U_{1,v}} & \mathcal{H}_{1,v} \\ \bigoplus_{e \in E^v} W_{s(e)} \downarrow & & \downarrow W_v \\ \bigoplus_{e \in E^v} \mathcal{H}_{2,s(e)} & \xrightarrow{U_{2,v}} & \mathcal{H}_{2,v} \end{array}$$

For any monoid homomorphism $\varphi: \mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B)$, there is a proper $C^*(G)$, B -correspondence \mathcal{H} with $\mathcal{H}_* = \varphi$.

Proof. Fix $v \in V_{\text{reg}}$. Let $E^v := \{e \in E \mid r(e) = v\}$ and let $m_v = |E^v|$ be the number of edges with range v . Totally order these edges, and let $u_v \in \mathbb{M}_{1,m_v}(C^*(G))$ be the row matrix with entries t_e for $e \in E^v$. The relations (CK1) and (CK2) say that $u_v u_v^* = \sum_{e \in E^v} t_e t_e^* = p_v$ and $u_v^* u_v = (t_e^* t_f)_{e,f \in E^v} = (\delta_{e,f} p_{s(e)})_{e,f \in E^v}$, the orthogonal direct sum $\bigoplus_{e \in E^v} p_{s(e)}$ of the projections $p_{s(e)}$ for $e \in E^v$. That is, u_v is a partial isometry with range and source projections p_v and $\bigoplus_{e \in E^v} p_{s(e)}$. Since any vertex in $r(E)$ is regular, the unitaries u_v for $v \in V_{\text{reg}}$ contain the partial isometries t_e for all edges.

Now we prove that a proper $C^*(G)$, B -correspondence \mathcal{H} is equivalent to a family $((\mathcal{H}_v)_{v \in V}, (U_v)_{v \in V_{\text{reg}}})$. Let $\mathcal{H}_v := p_v \cdot \mathcal{H} \subseteq \mathcal{H}$. These are proper Hilbert B -modules because $p_v \in \mathbb{K}(\mathcal{H})$ for a proper $C^*(G)$, B -correspondence. The partial isometry u_v for $v \in V_{\text{reg}}$ acts by a unitary operator $U_v: \bigoplus_{e \in E^v} \mathcal{H}_{s(e)} \xrightarrow{\sim} \mathcal{H}_v$. This maps \mathcal{H} to a family $((\mathcal{H}_v), (U_v))$. Let \mathcal{H}_1 and \mathcal{H}_2 be proper $C^*(G)$, B -correspondences and let $W: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ be an isomorphism. This restricts to isomorphisms of Hilbert B -modules $W_v: \mathcal{H}_{1,v} \rightarrow \mathcal{H}_{2,v}$ because W intertwines the left action of the projections p_v . The diagrams (3.3) commute because W intertwines the left action of the generators t_e for $e \in E$ and hence of the matrices u_v for $v \in V_{\text{reg}}$.

Conversely, let \mathcal{H}_v for $v \in V$ be proper Hilbert B -modules and let

$$U_v: \bigoplus_{e \in E^v} \mathcal{H}_{s(e)} \xrightarrow{\sim} \mathcal{H}_v$$

be unitaries for $v \in V_{\text{reg}}$. Let $\mathcal{H} := \bigoplus_{v \in V} \mathcal{H}_v$. Let $P_v \in \mathbb{K}(\mathcal{H})$ be the projection onto the summand \mathcal{H}_v . Let T_e for $e \in E^v$ be the restriction of U_v to the summand $\mathcal{H}_{s(e)}$. The operators P_v and T_e on \mathcal{H} satisfy the Cuntz–Krieger relations (CK1) and (CK2). So there is a unique $*$ -homomorphism $C^*(G) \rightarrow \mathbb{K}(\mathcal{H})$ mapping p_v to P_v and t_e to T_e . This is nondegenerate because it maps $(\bigoplus_{v \in F} p_v)_F$ to an approximate unit in $\mathbb{K}(\mathcal{H})$. Thus the family $((\mathcal{H}_v), (U_v))$ comes from a proper $C^*(G)$, B -correspondence. Let $((\mathcal{H}_{1,v}), (U_{1,v}))$ and $((\mathcal{H}_{2,v}), (U_{2,v}))$ be two such families and let $(W_v)_{v \in V}$ be a family of unitaries $W_v: \mathcal{H}_{1,v} \xrightarrow{\sim} \mathcal{H}_{2,v}$ making the diagrams (3.3) commute for all $v \in V_{\text{reg}}$. Then $W := \bigoplus_{v \in V} W_v$ is a unitary $\bigoplus_{v \in V} \mathcal{H}_{1,v} \rightarrow \bigoplus_{v \in V} \mathcal{H}_{2,v}$, which intertwines the left actions of the generators p_v for $v \in V$ and t_e for $e \in E$.

The functors from the category of proper $C^*(G)$, B -correspondences to the category of families $((\mathcal{H}_v), (U_v))$ and back are easily seen to be inverse to each other up to natural isomorphism. That is, they form an equivalence of categories.

Now let $\varphi: \mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B)$ be a homomorphism. We construct a family $((\mathcal{H}_v), (U_v))$ from it. For $v \in V$, let \mathcal{H}_v be a proper Hilbert B -module that represents $\varphi[p_v] \in \mathcal{P}(B)$. Let $v \in V_{\text{reg}}$. The equivalence u_v in $\mathbb{M}_\infty(C^*(G))$ implies that $\sum_{e \in E^v} [p_{s(e)}] = [p_v]$ holds in $\mathcal{P}(C^*(G))$. Since φ is a homomorphism, this implies $\sum_{e \in E^v} \varphi[p_{s(e)}] = \varphi[p_v]$. Equivalently, there is a unitary U_v between the corresponding proper Hilbert B -modules $\bigoplus_{e \in E^v} \mathcal{H}_{s(e)}$ and \mathcal{H}_v . The Hilbert modules \mathcal{H}_v for $v \in V$ and the unitaries U_v for $v \in V_{\text{reg}}$ describe a proper $C^*(G)$, B -correspondence \mathcal{H} ,

which induces a map $\mathcal{H}_* : \mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B)$. We claim that $\mathcal{H}_* = \varphi$. By construction, $\mathcal{H}_*[p_v] = [\mathcal{H}_v] = \varphi[p_v]$ for all $v \in V$. The subset of $\mathcal{P}(C^*(G))$ where \mathcal{H}_* and φ coincide is a submonoid. Hence it contains all finite sums of the vertex projections. Any class in $\mathcal{P}(C^*(G))$ is such a finite sum by Theorem 3.1. Hence $\mathcal{H}_* = \varphi$ holds on all of $\mathcal{P}(C^*(G))$. \square

Let $\varphi : \mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B)$ be a homomorphism. There may be several proper $C^*(G), B$ -correspondences \mathcal{H} with $\mathcal{H}_* = \varphi$. By Theorem 3.2, \mathcal{H} is given by proper Hilbert B -modules \mathcal{H}_v for $v \in V$ and unitaries $U_v : \bigoplus_{e \in E^v} \mathcal{H}_{s(e)} \rightarrow \mathcal{H}_v$ for $v \in V_{\text{reg}}$. We must have $[\mathcal{H}_v] = \varphi[p_v]$ in $\mathcal{P}(B)$, which determines \mathcal{H}_v up to isomorphism. Given any family of Hilbert B -module isomorphisms $W_v : \mathcal{H}_v \rightarrow \mathcal{H}'_v$ for $v \in V$, there are unique unitaries U'_v for $v \in V_{\text{reg}}$ that make (3.3) commute, so that (W_v) induces an isomorphism $(\mathcal{H}_v, U_v) \cong (\mathcal{H}'_v, U'_v)$. Thus it does not matter how we choose the Hilbert B -modules \mathcal{H}_v in their isomorphism classes. But the choice of the operators U_v does matter. They contain information about the action of the C^* -correspondence on K_1 . We shall make this more precise later. Here we prepare this with some simple observations.

Remark 3.4. By Theorem 2.3, any proper Hilbert B -module is isomorphic to $p \cdot B^\infty$ for some projection $p \in \mathbb{M}_\infty(B)$, and unitary operators between proper Hilbert B -modules are equivalent to equivalences of projections. Hence we may replace the families $((\mathcal{H}_v)_{v \in V}, (U_v)_{v \in V_{\text{reg}}})$ by families $((P_v)_{v \in V}, (U_v)_{v \in V_{\text{reg}}})$ consisting of projections in $\mathbb{M}_\infty(B)$ and equivalences between them.

Lemma 3.5. *Let \mathcal{H}_v for $v \in V$ be Hilbert B -modules. Let $U_v : \bigoplus_{e \in E^v} \mathcal{H}_{s(e)} \xrightarrow{\sim} \mathcal{H}_v$ for $v \in V_{\text{reg}}$ be unitaries. Families of unitaries $U'_v : \bigoplus_{e \in E^v} \mathcal{H}_{s(e)} \xrightarrow{\sim} \mathcal{H}_v$ for $v \in V_{\text{reg}}$ are in bijection with families of unitaries $\Upsilon_v \in \mathcal{U}(\mathcal{H}_v)$ for $v \in V_{\text{reg}}$, by mapping U'_v to $\Upsilon_v := U'_v U_v^* \in \mathcal{U}(\mathcal{H}_v)$ and $\Upsilon_v \in \mathcal{U}(\mathcal{H}_v)$ to $U'_v := \Upsilon_v U_v$. An isomorphism between the $C^*(G), B$ -correspondences described by $((\mathcal{H}_v), (U_v))$ and $((\mathcal{H}_v), (\Upsilon_v U_v))$ is a family of unitaries $W_v \in \mathcal{U}(\mathcal{H}_v)$ for $v \in V$ with*

$$\Upsilon_v = W_v \circ U_v \circ \left(\bigoplus_{e \in E^v} W_{s(e)}^* \right) \circ U_v^*$$

for all $v \in V_{\text{reg}}$.

Proof. The bijective correspondence between U'_v and Υ_v is trivial. The condition $\Upsilon_v = W_v \circ U_v \circ \left(\bigoplus_{e \in E^v} W_{s(e)}^* \right) \circ U_v^*$ is equivalent to the commutativity of the diagram (3.3) with $U_{1,v} = U_v$ and $U_{2,v} = \Upsilon_v U_v$. \square

Lemma 3.5 allows us to describe homotopies and asymptotic and approximate isomorphisms between $C^*(G), B$ -correspondences. Since homotopic projections are equivalent, two correspondences that are homotopic must have isomorphic Hilbert modules \mathcal{H}_v . The same holds for asymptotically or approximately isomorphic correspondences. Assume for simplicity that they even have the same Hilbert modules \mathcal{H}_v . By Lemma 3.5, the two correspondences are described by unitaries $U_v, \Upsilon_v U_v : \bigoplus_{e \in E^v} \mathcal{H}_{s(e)} \rightarrow \mathcal{H}_v$ for some $\Upsilon_v \in \mathcal{U}(\mathcal{H}_v)$ for $v \in V_{\text{reg}}$, respectively. A homotopy between them consists of norm-continuous homotopies $(\Upsilon_{v,t})_{t \in [0,1]}$ in $\mathcal{U}(\mathcal{H}_v)$ for all $v \in V_{\text{reg}}$ and unitaries $W_{v,j} \in \mathcal{U}(\mathcal{H}_v)$ for $j = 0, 1$ and $v \in V$, such that

$$\begin{aligned} \Upsilon_{v,0} &= W_{v,0} \circ U_v \circ \left(\bigoplus_{e \in E^v} W_{s(e),0}^* \right) \circ U_v^*, \\ \Upsilon_{v,1} &= W_{v,1} \circ \Upsilon_v U_v \circ \left(\bigoplus_{e \in E^v} W_{s(e),1}^* \right) \circ U_v^*. \end{aligned}$$

In other words, $(\Upsilon_{v,t})_{v \in V_{\text{reg}}}$ for $t \in [0, 1]$ yields a homotopy between two correspondences that are isomorphic to those corresponding to $(U_v)_{v \in V_{\text{reg}}}$ and $(\Upsilon_v U_v)_{v \in V_{\text{reg}}}$.

An asymptotic isomorphism between the two correspondences is given by a family of norm-continuous paths of $W_{v,t} \in \mathcal{U}(\mathcal{H}_v)$ for $t \in [0, \infty[$, $v \in V$, such that

$$\lim_{t \rightarrow \infty} W_{v,t} \circ U_v \circ \left(\bigoplus_{e \in E^v} W_{s(e),t}^* \right) \circ U_v^* = \Upsilon_v$$

for all $v \in V_{\text{reg}}$; there is no constraint on $W_{v,0}$. An approximate isomorphism between them is similar, but instead $W_{v,t} \in \mathcal{U}(\mathcal{H}_v)$ is needed only for $t \in \mathbb{N}$, $v \in V$. We can make this even more explicit:

Lemma 3.6. *Let V be countable and let $((\mathcal{H}_v), (U_v))$ and $((\mathcal{H}_v), (\Upsilon_v U_v))$ describe two $C^*(G), B$ -correspondences. They are approximately isomorphic if and only if for each finite subset $F \subseteq V$ and each $\varepsilon > 0$ there is a family of unitaries $W_v \in \mathcal{U}(\mathcal{H}_v)$ for $v \in V$ with*

$$\left\| \Upsilon_v - W_v \circ U_v \circ \left(\bigoplus_{e \in E^v} W_{s(e)}^* \right) \circ U_v^* \right\| < \varepsilon$$

for all $v \in F \cap V_{\text{reg}}$.

Proof. Write $V = \bigcup_{n \in \mathbb{N}} F_n$ as an increasing union of finite subsets. Let the unitaries $W_{v,n}$ be as in the statement of the lemma for $F_n \subseteq V$ and $\varepsilon = 2^{-n}$. These unitaries give the desired approximate isomorphism. \square

3.1. The AF-case. In this short subsection, we show that the techniques above already suffice to classify C^* -correspondences from $C^*(G)$ to B if our graph C^* -algebra is AF. This should be expected because the following sections are mainly about incorporating $K_1(C^*(G))$ into the picture, which is trivial in the AF-case.

Definition 3.7. A *cycle* in a directed graph G is a closed path e_1, \dots, e_n , that is, $s(e_1) = r(e_n)$, $s(e_2) = r(e_1)$, \dots , $s(e_n) = r(e_{n-1})$. A cycle is *simple* if the vertices $s(e_1), \dots, s(e_n)$ are all distinct.

Assume for some time that G has no cycles. Equivalently, $C^*(G)$ is an AF-algebra. Then the invariant $\mathcal{P}(C^*(G))$ determines $C^*(G)$ up to isomorphism. We are going to prove that proper $C^*(G), B$ -correspondences are classified by the induced map $\mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B)$ up to approximate isomorphism:

Proposition 3.8. *Let G be a directed graph without cycles. Let B be a C^* -algebra. Identify proper $C^*(G), B$ -correspondences with families $((\mathcal{H}_v)_{v \in V}, (U_v)_{v \in V_{\text{reg}}})$. Assume that $((\mathcal{H}_v), (U_v))$ and $((\mathcal{H}'_v), (U'_v))$ induce the same map $\mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B)$.*

- (1) *Let $F \subseteq V_{\text{reg}}$ be a finite subset. Then $((\mathcal{H}'_v), (U'_v))$ is isomorphic to a proper $C^*(G), B$ -correspondence $((\mathcal{H}''_v), (U''_v))$ with $\mathcal{H}''_v = \mathcal{H}_v$ for all $v \in V$ and $U''_v = U_v$ for $v \in F$.*
- (2) *If G is countable, then $((\mathcal{H}_v), (U_v))$ and $((\mathcal{H}'_v), (U'_v))$ are approximately isomorphic.*

Proof. We have already explained how to arrange $\mathcal{H}'_v = \mathcal{H}_v$ for all $v \in V$. Define $x \prec y$ for $x, y \in V$ if there is a path from x to y in G . This is a partial order on V because G has no cycles. Let $F \subseteq V_{\text{reg}}$ be finite. We may enumerate $F \cong \{1, \dots, n\}$ so that $x \prec y$ implies $x \leq y$ in $\{1, \dots, n\}$. We construct the isomorphism from $((\mathcal{H}'_v), (U'_v))$ to $((\mathcal{H}''_v), (U''_v))$ by induction on n , the number of points in F . We assume that we have such an isomorphism for $\{1, \dots, n-1\}$. Since isomorphisms of correspondences may be composed, we may assume that already $U'_i = U_i$ for $i = 1, \dots, n-1$. Now we define an isomorphism $(W_v)_{v \in V}$ by $W_n := U_n(U'_n)^*$ and

$W_v := 1$ for all $v \neq n$ and, in particular, for all $v \in V \setminus F$. By Lemma 3.6, this is an isomorphism from $((\mathcal{H}'_v), (U'_v))$ to $((\mathcal{H}'_v), (U''_v))$ with

$$U''_v := W_v \circ U'_v \circ \left(\bigoplus_{r(e)=v} W_{s(e)}^* \right).$$

If $v < n$, then $s(e) < n$ for all $e \in E$ with $r(e) = v$, so that $U''_v = U'_v = U_v$. If $v = n$, then $s(e) < n$ for all $e \in E$ with $r(e) = n$, so that $U''_n = W_n U'_n = U_n$. This proves the first statement. If G is countable, we may enumerate $V_{\text{reg}} \cong \mathbb{N}$. The isomorphisms $((\mathcal{H}'_v), (U'_v)) \cong ((\mathcal{H}'_v), (U''_v))$ for the finite subsets $\{1, \dots, n\} \subseteq V$ form an approximate isomorphism between the $C^*(G), B$ -correspondences $((\mathcal{H}'_v), (U'_v))$ and $((\mathcal{H}'_v), (U_v))$ because $*$ -polynomials in the generators p_v, s_e are dense in $C^*(G)$. So the first statement implies the second one. \square

4. PROJECTIONS VERSUS K-THEORY DIAGRAMS

For a C^* -algebra B that satisfies “stable weak cancellation,” we are going to describe $\mathcal{P}(B)$ using the ideal structure of B and the K_0^+ -monoids for certain ideals of B . Graph C^* -algebras and purely infinite C^* -algebras have stable weak cancellation. This clarifies the relationship between $\mathcal{P}(B)$ and invariants that have already been used in the classification of nonsimple C^* -algebras.

Let $p \in \mathbb{M}_\infty(B)$ be a projection. The ideal $\text{supp } p$ generated by p is the closed ideal generated by products $b_1^* p b_2$ with $b_1, b_2 \in B^\infty$. This is equal to the ideal generated by inner products of vectors in the Hilbert module $p \cdot B^\infty$ and to the ideal generated by the matrix coefficients of p . Any projection $p \in \mathcal{P}(B)$ belongs to $\mathbb{M}_\infty(\text{supp } p)$ and is a full projection there by construction of $\text{supp } p$. Equivalent projections generate the same ideal. Hence we get a map

$$\text{supp}: \mathcal{P}(B) \rightarrow \mathbb{I}(B).$$

It is *additive*, that is, $\text{supp } 0 = 0$ and

$$(4.1) \quad \text{supp}(p \oplus q) = \text{supp } p + \text{supp } q$$

for all $p, q \in \mathcal{P}(B)$. Hence it is also *monotone*: if $p, q \in \mathcal{P}(B)$, then

$$(4.2) \quad p \leq q \implies \text{supp } p \subseteq \text{supp } q;$$

here $p \leq q$ by definition if there is $p' \in \mathcal{P}(B)$ with $p \oplus p' = q$.

Definition 4.3. Let $\mathbb{I}_p(B) \subseteq \mathbb{I}(B)$ be the range of the map supp . We also say that an ideal *is generated by a projection* if it is equal to $\text{supp } p$ for some projection $p \in \mathbb{M}_\infty(B)$. Let $\mathbb{I}_p^\infty(B) \subseteq \mathbb{I}(B)$ be the larger subset of all ideals in B that are generated by some set of projections in $\mathbb{M}_\infty(B)$.

The lattice $\mathbb{I}(B)$ is isomorphic to the lattice of open subsets of the topological space \widehat{B} of irreducible representations of B (see [14, §3.2]). Let $\widehat{I} \subseteq \widehat{B}$ denote the open subset of \widehat{B} corresponding to $I \in \mathbb{I}(B)$. This space is indeed homeomorphic to the space of irreducible representations of I .

Lemma 4.4. *Let B be a C^* -algebra and $I \in \mathbb{I}(B)$. Then $I \in \mathbb{I}_p(B)$ if and only if $I \in \mathbb{I}_p^\infty(B)$ and \widehat{I} is quasi-compact.*

Proof. Let $p \in \mathbb{M}_n(B)$ for some $n \in \mathbb{N}$ be idempotent. Let I be the ideal generated by p . An irreducible representation ϱ of B remains nondegenerate on I if and only if $\varrho_n(p) \neq 0$, if and only if $\|\varrho_n(p)\| \geq 1$, where ϱ_n is the corresponding irreducible representation of $\mathbb{M}_n(B)$. So $\widehat{I} \subseteq \widehat{B}$ is the set of all $[\varrho]$ with $\|\varrho_n(p)\| \geq 1$. This subset is quasi-compact by [14, Proposition 3.3.7]. Conversely, let I be generated by some set of projections $(p_\alpha)_{\alpha \in S}$ and let $\widehat{I} \subseteq \widehat{B}$ be quasi-compact. For each finite

subset $F \subseteq S$, let I_F be the ideal generated by the projections p_α for $\alpha \in F$. Then $\bigcup I_F = I$. Since \widehat{I} is quasi-compact, there is some finite subset $F \subseteq S$ with $\widehat{I}_F = \widehat{I}$ and hence $I_F = I$. Thus I is generated by the projection $\bigoplus_{\alpha \in F} p_\alpha$. \square

Remark 4.5. A C^* -algebra has the *ideal property* if $\mathbb{I}(B) = \mathbb{I}_p^\infty(B)$, that is, any ideal is generated by the projections that it contains. Then an ideal $I \triangleleft B$ belongs to $\mathbb{I}_p(B)$ if and only if \widehat{I} is quasi-compact by Lemma 4.4. Any C^* -algebra of real rank 0 has the ideal property by [10, Corollary 2.8 and Proposition 2.9].

Remark 4.6. If B is separable and purely infinite, then $I \triangleleft B$ belongs to $\mathbb{I}_p(B)$ if and only if \widehat{I} is quasi-compact (see [31, Proposition 2.7]).

Definition 4.7 ([3, 11]). A C^* -algebra B has *stable weak cancellation* if projections p, q in $\mathbb{M}_\infty(B)$ with $\text{supp } p = \text{supp } q$ and $[p] = [q]$ in $K_0(\text{supp } p)$ are equivalent.

The equality $[p] = [q]$ in $K_0(\text{supp } p)$ means that there is a projection $r \in \mathbb{M}_\infty(\text{supp } p)$ so that $p \oplus r$ and $q \oplus r$ are equivalent, briefly, $p \oplus r \sim q \oplus r$. So stable weak cancellation holds if and only if $p \sim q$ follows if $p \oplus r \sim q \oplus r$ for a projection $r \in \mathbb{M}_\infty(\text{supp } p \cap \text{supp } q)$: this can only happen if $\text{supp } p = \text{supp } q$.

If B is simple, then either $p = 0$ or $\text{supp } p = B$. Thus stable weak cancellation for simple B means cancellation for all nonzero projections. It allows that there is a full (equivalently, nonzero) projection $0^f \in \mathcal{P}(B)$ representing 0 in $K_0(B)$. Then $\mathcal{P}(B)$ does not have cancellation because $0^f + 0^f = 0 + 0^f$. Thus stable weak cancellation is strictly weaker than cancellation in the monoid $\mathcal{P}(B)$.

Graph C^* -algebras have stable weak cancellation by [3, Corollary 7.2].

Theorem 4.8. *Purely infinite C^* -algebras have stable weak cancellation.*

Proof. This follows from [36, Theorem 4.1.4]. \square

Our next goal is to prove the following: if B has stable weak cancellation, then the monoid $\mathcal{P}(B)$ contains the same information as the diagram of monoids $K_0^+(I)$ for $I \in \mathbb{I}_p(B)$ with the canonical maps $K_0^+(I) \rightarrow K_0^+(J)$ for $I, J \in \mathbb{I}_p(B)$ with $I \subseteq J$. We first make the statement more precise.

Definition 4.9. Let (L, \leq) be a partially ordered set. A *diagram of (semi)groups* over it consists of (semi)groups X_W for $W \in L$ and homomorphisms $\iota_{W_2, W_1} : X_{W_1} \rightarrow X_{W_2}$ for $W_1 \leq W_2$ in L , such that $\iota_{W, W} = \text{id}_{X_W}$ for all $W \in L$ and $\iota_{W_3, W_2} \circ \iota_{W_2, W_1} = \iota_{W_3, W_1}$ if $W_1 \leq W_2 \leq W_3$ in L .

Given diagrams $X = (X_W, \varphi_{W_2, W_1})$ and $Y = (Y_W, \psi_{W_2, W_1})$ over L , we let $\text{Hom}_L(X, Y)$ be the (semi)group of natural transformations of diagrams; its elements are families of homomorphisms $\varphi_W : X_W \rightarrow Y_W$ for $W \in L$ that intertwine the maps φ_{W_2, W_1} and ψ_{W_2, W_1} for $W_1 \subseteq W_2$ in L .

Let (L_1, \leq) and (L_2, \leq) be partially ordered sets and $\psi : L_1 \rightarrow L_2$ an order-preserving map. Let $(X_W, \varphi_{W_2, W_1})_{W \in L_2}$ be a diagram of (semi)groups over L_2 . Then $\psi^*((X_W, \varphi_{W_2, W_1})_{W \in L_2})$ is the diagram of (semi)groups over L_1 consisting of the (semi)groups $X_{\psi(W)}$ for $W \in L_1$ and the homomorphisms $\varphi_{\psi(W_2), \psi(W_1)}$ for $W_1, W_2 \in L_1$ with $W_1 \leq W_2$.

Definition 4.10. Let B be a C^* -algebra. Let $K_0^+(B, \mathbb{I}_p)$ be the diagram over $\mathbb{I}_p(B)$ consisting of the Abelian monoids $K_0^+(I)$ for $I \in \mathbb{I}_p(B)$ and the homomorphisms $K_0^+(I) \rightarrow K_0^+(J)$ induced by the inclusion maps $I \rightarrow J$ for $I, J \in \mathbb{I}_p(B)$ with $I \subseteq J$. We define $K_0(B, \mathbb{I}_p)$ and $K_1(B, \mathbb{I}_p)$ similarly, replacing K_0^+ by K_0 and K_1 .

We recall the definition of an order unit:

Definition 4.11. An *order unit* of a commutative semigroup M (with its natural preorder \leq) is an element u of M such that for any element x of M , there is a

positive integer n with $x \leq n \cdot u$. A homomorphism $\varphi: M \rightarrow M'$ preserves order units if $\varphi(u)$ for an order unit u in M is an order unit in M' . (If $\varphi(u)$ is an order unit for some order unit u in M , then this happens for all order units u in M .)

Let A and B be C*-algebras and let $\psi: \mathbb{I}_p(A) \rightarrow \mathbb{I}_p(B)$ be monotone. Then the pull-back diagram $\psi^*(\mathbb{K}_0^+(B, \mathbb{I}_p))$ consists of the monoids $\mathbb{K}_0^+(\psi(I))$ for $I \in \mathbb{I}_p(A)$ with the canonical maps $\mathbb{K}_0^+(\psi(I_1)) \rightarrow \mathbb{K}_0^+(\psi(I_2))$ for $I_1 \subseteq I_2$.

Theorem 4.12. *Let A and B be C*-algebras. Let B have stable weak cancellation. Then there is a natural bijection between monoid homomorphisms $f: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$ and pairs $(\psi, (\varphi_I)_{I \in \mathbb{I}_p(A)})$, where $\psi: \mathbb{I}_p(A) \rightarrow \mathbb{I}_p(B)$ is monotone and*

$$\varphi = (\varphi_I)_{I \in \mathbb{I}_p(A)}: \mathbb{K}_0^+(A, \mathbb{I}_p) \rightarrow \psi^* \mathbb{K}_0^+(B, \mathbb{I}_p)$$

is a morphism of monoid diagrams such that each φ_I preserves order units. The map ψ in such a pair is automatically additive, that is, $\psi(I_1 + I_2) = \psi(I_1) + \psi(I_2)$ for all $I_1, I_2 \in \mathbb{I}_p(A)$.

A monoid isomorphism clearly preserves order units. Hence Theorem 4.12 implies that $\mathcal{P}(A) \cong \mathcal{P}(B)$ if and only if there are an isomorphism of partially ordered sets $\psi: \mathbb{I}_p(A) \xrightarrow{\sim} \mathbb{I}_p(B)$ and an isomorphism $\mathbb{K}_0^+(A, \mathbb{I}_p) \cong \psi^*(\mathbb{K}_0^+(B, \mathbb{I}_p))$ as diagrams of monoids over $\mathbb{I}_p(A)$. This is what was meant by saying that $\mathcal{P}(A)$ and the diagram $\mathbb{K}_0^+(I)_{I \in \mathbb{I}_p(A)}$ contain equivalent information.

We prepare the proof of Theorem 4.12 by two lemmas.

Lemma 4.13. *Let B be a C*-algebra, $I \in \mathbb{I}_p(B)$, and $p \in \mathcal{P}(I)$. Then $\text{supp } p = I$ if and only if p is an order unit in $\mathcal{P}(I)$. When B has stable weak cancellation, this happens if and only if the image of p in $\mathbb{K}_0^+(I)$ is an order unit.*

Proof. Since $I \in \mathbb{I}_p(B)$, there is $q \in \mathcal{P}(B)$ with $I = \text{supp } q$. We fix such a q throughout the proof.

Assume that p is an order unit in $\mathcal{P}(I)$. Then $q \leq n \cdot p$ for some $n \in \mathbb{N}_{>0}$ and hence $I = \text{supp } q \subseteq n \cdot \text{supp } p = \text{supp } p$. Then $I = \text{supp } p$. Conversely, let $\text{supp } p = I$. Let $x \in \mathcal{P}(I)$. We must show that $x \leq n \cdot p$ for some $n \in \mathbb{N}_{>0}$. Equivalently, $x \cdot I^\infty$ is a direct summand in $(p \cdot I^\infty)^n$ as a Hilbert I -module.

Let $\mathcal{H} := p \cdot I^\infty$ as a right Hilbert I -module. Then $\mathbb{K}(\mathcal{H}) \cong p(I \otimes \mathbb{K})p$. Let \mathcal{H}^* be the dual Hilbert $I, \mathbb{K}(\mathcal{H})$ -bimodule. Then $\mathcal{H}^* \otimes_{\mathbb{K}(\mathcal{H})} \mathcal{H} \cong I$ and $\mathcal{H} \otimes_I \mathcal{H}^* \cong \mathbb{K}(\mathcal{H})$. So $x \cdot I^\infty$ is a direct summand in $(p \cdot I^\infty)^n$ if and only if $x \cdot I^\infty \otimes_I \mathcal{H}^*$ is a direct summand in $(p \cdot I^\infty)^n \otimes_I \mathcal{H}^* = \mathbb{K}(\mathcal{H})^n$ as a Hilbert $\mathbb{K}(\mathcal{H})$ -module. The C*-algebra $\mathbb{K}(x \cdot I^\infty \otimes_I \mathcal{H}^*)$ is isomorphic to $\mathbb{K}(x \cdot I^\infty)$ and hence still unital. So $x \cdot I^\infty \otimes_I \mathcal{H}^*$ is a proper Hilbert $\mathbb{K}(\mathcal{H})$ -module. By Theorem 2.3, $x \cdot I^\infty \otimes_I \mathcal{H}^*$ is a direct summand in $(p \cdot I^\infty)^n$ for some $n \in \mathbb{N}_{>0}$. This finishes the proof that $\text{supp } p = I$ if and only if p is an order unit in $\mathcal{P}(I)$.

Let p be an order unit in $\mathcal{P}(I)$. If $x \in \mathbb{K}_0^+(I)$, then $x = [\bar{x}]$ for some $x \in \mathcal{P}(I)$. By assumption, $\bar{x} \leq n \cdot p$ for some $n \in \mathbb{N}_{>0}$. Hence $x \leq n \cdot [p]$. Thus $[p] \in \mathbb{K}_0^+(I)$ is an order unit in $\mathbb{K}_0^+(I)$. Conversely, let $[p] \in \mathbb{K}_0^+(I)$ be an order unit. Let $x \in \mathcal{P}(I)$. Then $x \oplus q$ is full in I . There is $n \in \mathbb{N}_{>0}$ with $[x \oplus q] \leq n \cdot [p]$ in $\mathbb{K}_0^+(I)$, that is, $[x \oplus q] \oplus [y] = n \cdot [p]$ for some $y \in \mathcal{P}(I)$. Since B has stable weak cancellation and both sides are full in I , this implies $x \oplus q \oplus y \sim n \cdot p$ in $\mathcal{P}(I)$. So $x \leq n \cdot p$. Thus p is also an order unit in $\mathcal{P}(I)$. \square

Definition 4.14. For $p, q \in \mathcal{P}(B)$, we write $p \prec q$ if and only if $\text{supp } p \subseteq \text{supp } q$.

Lemma 4.15. *Let $p, q \in \mathcal{P}(B)$. Then $p \prec q$ if and only if $p \leq n \cdot q$ for some $n \in \mathbb{N}$.*

Proof. If $p \leq n \cdot q$, then $\text{supp } p \subseteq \text{supp } n \cdot q = \text{supp } q$. Conversely, let $\text{supp } p \subseteq \text{supp } q$. By Lemma 4.13, q is an order unit for $\mathcal{P}(\text{supp } p)$. Hence $p \leq n \cdot q$ for some $n \in \mathbb{N}$. \square

Proof of Theorem 4.12. Let $f: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$ be a homomorphism. The map $\text{supp}: \mathcal{P}(A) \rightarrow (\mathbb{I}_p(A), +)$ is a surjective monoid homomorphism. Hence $\mathbb{I}_p(A)$ is isomorphic to the quotient of $\mathcal{P}(A)$ by the congruence relation defined by $\text{supp } p = \text{supp } q$. Lemma 4.15 describes this congruence relation and also the partial order defined by $\text{supp } p \subseteq \text{supp } q$ in a purely algebraic way. So $\text{supp } p \subseteq \text{supp } q$ implies $\text{supp } f(p) \subseteq \text{supp } f(q)$ and similarly with equality instead of inclusion. As a result, f induces a well defined map $\psi: \mathbb{I}_p(A) \rightarrow \mathbb{I}_p(B)$ that is both additive and monotone.

Let $I \in \mathbb{I}_p(A)$. Then there is $q \in \mathcal{P}(A)$ with $I = \text{supp } q$. Then $\psi(I) = \text{supp } f(q)$. Lemma 4.15 shows that $p \in \mathcal{P}(A)$ satisfies $p \in \mathcal{P}(I)$ if and only if there is $n \in \mathbb{N}$ with $p \leq n \cdot q$. Hence f maps $\mathcal{P}(I)$ to $\mathcal{P}(\psi(I))$. This induces a group homomorphism $K_0(I) \rightarrow K_0(\psi(I))$, which restricts to a monoid homomorphism $\varphi_I: K_0^+(I) \rightarrow K_0^+(\psi(I))$. It maps the order unit q in $\mathcal{P}(I)$ to $f(q)$, which is an order unit in $\mathcal{P}(\psi(I))$ by Lemma 4.13 because $\psi(I) = \text{supp } f(q)$. These homomorphisms for $I \in \mathbb{I}_p(A)$ form a morphism of diagrams over $\mathbb{I}_p(A)$ that preserves order units.

Conversely, let $\psi: \mathbb{I}_p(A) \rightarrow \mathbb{I}_p(B)$ be a monotone map and let $\varphi_I: K_0^+(I) \rightarrow K_0^+(\psi(I))$ for $I \in \mathbb{I}_p(A)$ be monoid homomorphisms that preserve order units and form a morphism of diagrams. Assume that B has stable weak cancellation. Let $p \in \mathcal{P}(A)$. Let $I := \text{supp } p$. Choose $q \in \mathcal{P}(\psi(I))$ representing $\varphi_I[p]$ in $K_0^+(\psi(I))$. Since φ_I preserves order units, $\text{supp } q = \psi(I)$. Since B has stable weak cancellation, q is unique as an element of $\mathcal{P}(B)$. So $f(p) := q$ well-defines a map $f: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$.

Let $p_1, p_2 \in \mathcal{P}(A)$. We claim that $f(p_1 + p_2) = f(p_1) + f(p_2)$. Let $p_3 := p_1 + p_2$. For $j = 1, 2, 3$, define $I_j := \text{supp } p_j$. By Lemma 4.13, $[p_j] \in K_0^+(I_j)$ is an order unit. Therefore, $\varphi_{I_j}[p_j] \in K_0^+(\psi(I_j))$ is an order unit. Since B has stable weak cancellation, $q_j := f(p_j) \in \mathcal{P}(B)$ is the unique element with support ideal $\psi(I_j)$ and $\varphi_{I_j}[p_j] = [q_j]$ in $K_0^+(\psi(I_j))$. The relation $p_1 + p_2 = p_3$ holds in $K_0^+(I_3)$ when we tacitly view p_1, p_2, p_3 as elements of $\mathcal{P}(I_3)$. Since φ_{I_3} is additive, this implies the relation $[q_1 + q_2] = [q_1] + [q_2] = [q_3]$ in $K_0^+(\psi(I_3))$. Since the maps φ_I preserve order units, $[q_1 + q_2] = [q_3]$ is an order unit for $K_0^+(\psi(I_3))$. Then Lemma 4.13 implies that $\psi(I_3)$ is the support ideal of q_3 and of $q_1 + q_2$. The support of $q_1 + q_2$ is contained in $\psi(I_1) + \psi(I_2)$. So $\psi(I_3) \subseteq \psi(I_1) + \psi(I_2)$. Since ψ is monotone by assumption, it follows that ψ must be additive. And stable weak cancellation for B shows that $q_1 + q_2 = q_3$ holds in $\mathcal{P}(B)$. That is, $f: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$ is a homomorphism. It is easy to see that the two constructions above are inverse to each other, giving the asserted bijection. \square

Corollary 4.16. *Let A and B be C^* -algebras. Assume that B is purely infinite. There is a natural bijection between homomorphisms $f: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$ and pairs $(\psi, (\varphi_I)_{I \in \mathbb{I}_p(A)})$ with a monotone map $\psi: \mathbb{I}_p(A) \rightarrow \mathbb{I}_p(B)$ and a morphism of diagrams of Abelian groups $\varphi_I: K_0(A, \mathbb{I}_p) \rightarrow \psi^* K_0(B, \mathbb{I}_p)$.*

Proof. Since any $I \in \mathbb{I}_p(B)$ is also purely infinite, any class in $K_0(I)$ is represented by a projection in I (see [36, Theorem 4.1.4]). Thus $K_0^+(B, \mathbb{I}_p) = K_0(B, \mathbb{I}_p)$, which is a diagram of Abelian groups. Any element of an Abelian group is an order unit. Theorem 4.8 says that B has stable weak cancellation. So Theorem 4.12 applies here and gives the remaining statements with K_0^+ instead of K_0 . If $I \in \mathbb{I}_p(A)$, then $K_0(I)$ is isomorphic to the Grothendieck group of $K_0^+(I)$. Hence there is a bijection between homomorphisms $K_0^+(I) \rightarrow K_0^+(J)$ and $K_0(I) \rightarrow K_0(J)$ for $I \in \mathbb{I}_p(A)$ and $J \in \mathbb{I}_p(B)$. \square

Remark 4.17. If $I \in \mathbb{I}_p^\infty(B)$, then the set of $J \in \mathbb{I}_p(B)$ with $J \subseteq I$ is directed, and the closure of its union is equal to I . If $I \in \mathbb{I}_p(B)$, then this directed set has I as a maximal element. Therefore, any monoid homomorphism $\mathbb{I}_p(A) \rightarrow \mathbb{I}_p(B)$ extends uniquely to a monoid homomorphism $\mathbb{I}_p^\infty(A) \rightarrow \mathbb{I}_p^\infty(B)$ that commutes also

with infinite suprema. We also have $K_0^+(I) = \varinjlim K_0^+(J)$ and $K_i(I) = \varinjlim K_i(J)$ for $i = 1, 2$, where J runs through the directed set of all $J \in \mathbb{I}_p(B)$ with $J \subseteq I$. Thus a diagram morphism $K_0^+(A, \mathbb{I}_p) \rightarrow \psi^* K_0^+(B, \mathbb{I}_p)$ or $K_i(A, \mathbb{I}_p) \rightarrow \psi^* K_i(B, \mathbb{I}_p)$, $i = 0, 1$, extends uniquely to a diagram morphism $K_0^+(A, \mathbb{I}_p^\infty) \rightarrow \psi^* K_0^+(B, \mathbb{I}_p^\infty)$ or $K_i(A, \mathbb{I}_p^\infty) \rightarrow \psi^* K_i(B, \mathbb{I}_p^\infty)$. Hence we may replace the indexing set $\mathbb{I}_p(A)$ by $\mathbb{I}_p^\infty(A)$ in Theorem 4.12 and Corollary 4.16 provided we add the condition that the map on \mathbb{I}_p^∞ preserves infinite suprema. The condition in Theorem 4.12 that the map $\varphi_I: K_0^+(I) \rightarrow K_0^+(\psi(I))$ preserves order units is empty for $I \in \mathbb{I}_p^\infty(A) \setminus \mathbb{I}_p(A)$ because $K_0^+(I)$ has no order unit in this case.

5. SOME HOMOLOGICAL ALGEBRA WITH K-THEORY DIAGRAMS

We first recall well known results on the ideal structure of graph C*-algebras and about their K-theory diagrams $K_0(C^*(G), \mathbb{I}_p)$, $K_1(C^*(G), \mathbb{I}_p)$ and $K_0^+(C^*(G), \mathbb{I}_p)$.

Definition 5.1. A subset $H \subseteq V$ is *hereditary* if for any edge $e \in E$, $r(e) \in H$ implies $s(e) \in H$. It is *saturated* if any $v \in V_{\text{reg}}$ with $s(e) \in H$ for all edges e with $r(e) = v$ satisfies $v \in H$.

The intersection of any set of hereditary and saturated subsets is again hereditary and saturated. The union may fail to be hereditary and saturated, but then there is a smallest hereditary and saturated subset containing it. Thus the hereditary and saturated subsets of V form a complete lattice, which we denote by $\mathbb{I}(G)$.

Definition 5.2. For a subset $V_2 \subseteq V$, let $\langle V_2 \rangle \subseteq V$ be the smallest hereditary and saturated subset containing V_2 . If $V_2 = \{v\}$, we briefly denote this by $\langle v \rangle$.

There is an order-isomorphism between $\mathbb{I}(G)$ and the lattice $\mathbb{I}^\Gamma(C^*(G))$ of gauge-invariant ideals in $C^*(G)$ because G is row-finite (see [7, Theorem 4.1] or [6, Theorem 3.6]). It maps a hereditary and saturated subset $H \subseteq V$ to the ideal generated by the projections p_v for all $v \in H$. Its inverse maps a gauge-invariant ideal I to the set of $v \in V$ with $p_v \in I$. We write $C^*(G)_H$ for the ideal corresponding to $H \in \mathbb{I}(G)$. It is Morita–Rieffel equivalent to the graph C*-algebra of the restriction $G|_H$ of G to H .

Write $q \in \mathcal{P}(C^*(G))$ as $\sum_{v \in V} c_v p_v$ as in Theorem 3.1. Then $\text{supp } q$ is equal to the ideal generated by the projections p_v with $c_v \neq 0$. An ideal of this form is automatically gauge-invariant. The hereditary, saturated subset of V that corresponds to $\text{supp } q$ is the smallest one that contains all v with $c_v \neq 0$. It is called the (hereditary, saturated) *support* of c . The following theorem summarises the findings:

Theorem 5.3. *Let G be a graph. An ideal of $C^*(G)$ belongs to $\mathbb{I}_p^\infty(C^*(G))$ if and only if it is gauge-invariant, and $\mathbb{I}_p^\infty(C^*(G)) \cong \mathbb{I}(G)$ as complete lattices. This isomorphism maps the subset $\mathbb{I}_p(C^*(G))$ onto the partially ordered set of finitely generated hereditary, saturated subsets of V ; we denote this by $\mathbb{I}_p(G)$. The support map $\text{supp}: \mathcal{P}(C^*(G)) \rightarrow \mathbb{I}_p(C^*(G)) \cong \mathbb{I}_p(G)$ maps the class of the projection $\sum c_v p_v$ for $c \in \mathbb{N}[V]$ to the hereditary, saturated support of c .*

The free Abelian groups $\mathbb{Z}[W]$ for $W \in \mathbb{I}_p(G)$ form a diagram of groups over $\mathbb{I}_p(G)$, which we denote by $\mathbb{Z}[V, \mathbb{I}_p]$. Similarly, let $W_{\text{reg}} := W \cap V_{\text{reg}}$ for $W \in \mathbb{I}_p(G)$. The free Abelian groups $\mathbb{Z}[W_{\text{reg}}]$ for $W \in \mathbb{I}_p(G)$ form a diagram of groups over $\mathbb{I}_p(G)$ denoted by $\mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p]$. We define a morphism of diagrams $M_G^{\mathbb{I}_p}: \mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p] \rightarrow \mathbb{Z}[V, \mathbb{I}_p]$ to consist of the maps

$$M_G^W: \mathbb{Z}[W_{\text{reg}}] \rightarrow \mathbb{Z}[W], \quad M_G^W((c_w)_{w \in W})(v) := \sum_{\{e \in E: s(e)=w\}} c_r(e).$$

These maps form a diagram morphism because if $W \subseteq V$ is hereditary, then $M_G^W = M_G^V|_{\mathbb{Z}[W]}$. Roughly speaking, $M_G^{\mathbb{I}_p}$ multiplies with the *adjacency matrix* of G , that is, with the matrix whose v, w -entry is the number of edges $v \leftarrow w$ in G .

The results on the K-theory of graph C^* -algebras provide natural exact sequences

$$(5.4) \quad 0 \rightarrow K_1(C^*(G|_W)) \xrightarrow{\kappa|_W} \mathbb{Z}[W_{\text{reg}}] \xrightarrow{\text{id}-M_G^W} \mathbb{Z}[W] \xrightarrow{\pi|_W} K_0(C^*(G|_W)) \rightarrow 0$$

for all $W \in \mathbb{I}_p^\infty(G)$ (see [33, Theorem 3.2]). These are compatible with the diagram structure over $\mathbb{I}_p(G)$, so that (5.4) gives an exact sequence of diagrams

$$(5.5) \quad 0 \rightarrow K_1(C^*(G), \mathbb{I}_p) \xrightarrow{\kappa} \mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p] \xrightarrow{\text{id}-M_G^{\mathbb{I}_p}} \mathbb{Z}[V, \mathbb{I}_p] \xrightarrow{\pi} K_0(C^*(G), \mathbb{I}_p) \rightarrow 0.$$

In addition, $K_0^+(C^*(G|_W)) = \pi|_W(\mathbb{N}[W]) \subseteq K_0(C^*(G|_W))$ by Theorem 3.1.

Diagrams of Abelian groups over $\mathbb{I}_p(G)$ form an Abelian category. This allows us to do homological algebra with them. We shall need the extension groups in this category, which we denote by $\text{Ext}_{\mathbb{I}_p(G)}^n$ for $n \in \mathbb{N}$. The case $n = 0$ gives $\text{Ext}^0 = \text{Hom}$.

Proposition 5.6. *The diagrams $\mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p]$ and $\mathbb{Z}[V, \mathbb{I}_p]$ are projective. So (5.5) is the beginning of a projective resolution of $K_0(C^*(G), \mathbb{I}_p)$.*

Proof. For $v \in V$, let $\langle v \rangle \subseteq V$ be the hereditary and saturated subset generated by v . So $\langle v \rangle \in \mathbb{I}_p(G)$. Let $(X_W, \varphi_{W_2, W_1})_{W \in \mathbb{I}_p(G)}$ be a diagram of groups over $\mathbb{I}_p(G)$. A family $x_v \in X_{\langle v \rangle}$ for $v \in V$ defines a diagram morphism $\xi: \mathbb{Z}[V, \mathbb{I}_p] \rightarrow (X_W)_{W \in \mathbb{I}_p(G)}$, which consists of the maps

$$\xi_W: \mathbb{Z}[W] \rightarrow X_W, \quad \sum_{v \in W} c_v \delta_v \mapsto \sum_{v \in W} c_v \varphi_{W, \langle v \rangle}(x_v).$$

Any diagram morphism $\xi: \mathbb{Z}[V, \mathbb{I}_p] \rightarrow (X_W)_{W \in \mathbb{I}_p(G)}$ is of this form for a unique family of $x_v \in X_{\langle v \rangle}$ for $v \in V$, namely, $x_v := \xi_{\langle v \rangle}(\delta_v)$. So

$$\text{Hom}_{\mathbb{I}_p(G)}(\mathbb{Z}[V, \mathbb{I}_p], (X_W)) \cong \prod_{v \in W} X_{\langle v \rangle}.$$

This functor is exact. That is, $\mathbb{Z}[V, \mathbb{I}_p]$ is projective. A similar argument works for $\mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p]$ or, more generally, for $\mathbb{Z}[W, \mathbb{I}_p]$ for any subset W of V . \square

The resolution in (5.5) allows us to compute $\text{Ext}_{\mathbb{I}_p(G)}^n(K_0(C^*(G), \mathbb{I}_p), Y)$ for any diagram of groups Y over $\mathbb{I}_p(G)$ and $n = 0, 1, 2$ as the cohomology groups of the cochain complex

$$(5.7) \quad 0 \rightarrow \text{Hom}_{\mathbb{I}_p(G)}(\mathbb{Z}[V, \mathbb{I}_p], Y) \xrightarrow{(\text{id}-M_G^{\mathbb{I}_p})^*} \text{Hom}_{\mathbb{I}_p(G)}(\mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p], Y) \xrightarrow{\kappa^*} \text{Hom}_{\mathbb{I}_p(G)}(K_1(C^*(G), \mathbb{I}_p), Y) \rightarrow 0.$$

This works for $n = 0, 1, 2$, regardless whether or not $K_1(C^*(G), \mathbb{I}_p)$ is projective.

We now compare the resolution above to the constructions in [9]. That article only considers graph C^* -algebras with finitely many ideals and, without loss of generality, assumes all graphs to be regular. While each ideal in $C^*(G)$ is generated by a projection in that case, only some ideals are used in [9]. Call an ideal $I \triangleleft B$ *irreducible* if $I = \overline{\sum_{\alpha \in S} J_\alpha}$ for a family of ideals J_α implies $J_\alpha = I$ for some $\alpha \in S$. The irreducible ideals form a partially ordered set. The invariant that is used in [9] is the diagram consisting of the $\mathbb{Z}/2$ -graded Abelian groups $K_*(I)$ for all *irreducible* ideals $I \triangleleft B$. If B has only finitely many ideals, then for each primitive ideal $\mathfrak{p} \triangleleft B$ there is a minimal open subset of the primitive ideal space that contains \mathfrak{p} . The ideals corresponding to these minimal open subsets are exactly the irreducible ideals.

Remark 5.8. The Cantor set is the primitive ideal space of a commutative AF-algebra and hence of a graph C*-algebra. All open subsets of the Cantor set are reducible. So restricting to irreducible ideals is not reasonable for C*-algebras with infinitely many ideals, not even if they are graph C*-algebras.

For a graph C*-algebra with finitely many ideals, let $\mathbb{I}_s(G) \subseteq \mathbb{I}_p(G) = \mathbb{I}(G)$ be the subset corresponding to the irreducible ideals in $C^*(G)$. We may restrict the exact sequence of diagrams (5.5) to the subset $\mathbb{I}_s(G) \subseteq \mathbb{I}_p(G)$. The restricted diagrams $\mathbb{Z}[V, \mathbb{I}_s]$ and $K_1(C^*(G), \mathbb{I}_s)$ are shown in [9] to be projective objects in the Abelian category of diagrams of Abelian groups over $\mathbb{I}_s(G)$. Hence the groups $\text{Ext}_{\mathbb{I}_s(G)}^n(K_0(C^*(G), \mathbb{I}_s), \psi^*K_1(B, \mathbb{I}_s))$ in this Abelian category of diagrams vanish for $n \geq 3$ and are computed exactly as in (5.7) with \mathbb{I}_s instead of \mathbb{I}_p if $n = 0, 1, 2$.

The invariant over \mathbb{I}_s clearly contains less information than the invariant over \mathbb{I}_p . This does not mean, however, that lifting a map on the invariant over \mathbb{I}_p is easier than lifting a map on the invariant over \mathbb{I}_s : a given extension to \mathbb{I}_p may put constraints on the desired C*-correspondence.

Remark 5.9. Let G be a regular graph with only finitely many hereditary saturated subsets. Then any irreducible hereditary saturated subset is generated by a single vertex $v \in V$. The converse does not hold, however, because the union of two hereditary and saturated subsets of V need not remain hereditary. For instance, let G be the graph $1 \rightarrow 3 \leftarrow 2$. Then G is generated by the single vertex 3. But the corresponding ideal $C^*(G) \triangleleft C^*(G)$ is not irreducible: it is the sum of the ideals corresponding to $\{1\}$ and $\{2\}$. This sum contains 3 because any hereditary and saturated subset that contains 1 and 2 also contains 3.

Remark 5.10. The machinery of homological algebra in triangulated categories does not apply to the invariant $K_*(B, \mathbb{I}_p)$, not even if the ideal lattice of B is finite. This remark explains why. First we make our invariant more precise. Let X be a locally quasi-compact topological space and let (L, \leq) be the partially ordered set of quasi-compact open subsets of X . Let B be a C*-algebra over X . If $\text{Prim}(B) \cong X$ and B has the ideal property, then $L = \mathbb{I}_p(B)$. Diagrams of $\mathbb{Z}/2$ -graded countable Abelian groups over (L, \leq) form a stable Abelian category, and mapping B to the diagram $K_*(B, L) := (K_*(I))_{I \in L}$ is a stable homological functor into this category. This invariant is, however, not “universal,” so that the homological algebra machinery developed in [25–28] does not apply directly to it. Let $I_1, I_2 \in L$ and assume $I_1 \cap I_2 \in L$. We have $I_1 + I_2 \in L$ anyway. The boundary map ∂ in the Mayer–Vietoris sequence

$$\cdots \rightarrow K_*(I_1 \cap I_2) \rightarrow K_*(I_1) \oplus K_*(I_2) \rightarrow K_*(I_1 + I_2) \xrightarrow{\partial} K_{*-1}(I_1 \cap I_2) \rightarrow \cdots$$

is a natural transformation from $K_*(I_1 + I_2)$ to $K_{*-1}(I_1 \cap I_2)$, which is not yet encoded in our Abelian category of diagrams. Hence our invariant cannot be universal.

Despite Remark 5.10, we shall work with the invariant $K_*(B, \mathbb{I}_p)$ and use homological algebra in the category of diagrams over $\mathbb{I}_p(G)$. This works quite well to describe correspondences from graph C*-algebras to B . In fact, we shall use $\mathcal{P}(B)$ instead of $K_0^+(B, \mathbb{I}_p)$ most of the time. These invariants are equivalent by Theorem 4.12 if B has stable weak cancellation. We do not need to assume stable weak cancellation when we work directly with $\mathcal{P}(B)$.

6. ADDING K_1

Theorem 3.2 only uses the invariant $\mathcal{P}(B)$, which is not fine enough for classification. Now we add information about K_1 . We develop a criterion when a pair of

maps $\mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B)$ and $K_1(C^*(G), \mathbb{I}_p) \rightarrow \psi^*K_1(B, \mathbb{I}_p)$ is induced by a proper $C^*(G), B$ -correspondence. It uses a variant of the obstruction class in [9].

Let \mathcal{H} be a proper $C^*(G), B$ -correspondence. Describe \mathcal{H} through proper Hilbert B -modules \mathcal{H}_v for $v \in V$ and unitaries $U_v: \bigoplus_{e \in E^v} \mathcal{H}_{s(e)} \xrightarrow{\sim} \mathcal{H}_v$ for $v \in V_{\text{reg}}$ as in Theorem 3.2. We are going to describe the homomorphisms

$$\mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B), \quad \mathbb{I}(C^*(G)) \rightarrow \mathbb{I}_p(B), \quad K_1(C^*(G), \mathbb{I}_p) \rightarrow \psi^*K_1(B, \mathbb{I}_p)$$

induced by \mathcal{H} . By Theorem 3.1, the homomorphism $\mathcal{H}_*: \mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B)$, $[\mathcal{M}] \mapsto [\mathcal{M} \otimes_{C^*(G)} \mathcal{H}]$, is determined by its values

$$\mathcal{H}_*[p_v] = \mathcal{H}_*[p_v \cdot C^*(G)] = [p_v \cdot \mathcal{H}] = [\mathcal{H}_v].$$

The homomorphism $\mathcal{H}_*: \mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B)$ induces a map

$$\psi: \mathbb{I}_p(G) \cong \mathbb{I}_p(C^*(G)) \rightarrow \mathbb{I}_p(B)$$

as in the proof of Theorem 4.12; this part of the proof works without stable weak cancellation. Namely, let $W \in \mathbb{I}_p(G)$ be a finitely generated, hereditary, saturated subset of V . Let $F \subseteq W$ be any finite generating set for W . Then the projection $\bigoplus_{v \in F} p_v$ generates the ideal $C^*(G)_W$ corresponding to W . Hence

$$\psi(W) := \text{supp } \varphi_0 \left(\sum_{v \in F} [p_v] \right) = \bigvee_{v \in F} \text{supp } \varphi_0 [p_v]$$

This ideal is equal to $\bigvee_{v \in W} \text{supp } \varphi_0 [p_v]$ and does not depend on the choice of F .

By definition, $\psi(W)$ is the ideal in B that is generated by the inner product on the proper Hilbert B -module $\bigoplus_{v \in F} p_v \cdot \mathcal{H} = \bigoplus_{v \in F} \mathcal{H}_v$. Therefore, $C^*(G)_W \cdot \mathcal{H}$ is a full Hilbert $\psi(W)$ -module. As such, it is nondegenerate as a right $\psi(W)$ -module, so that

$$(6.1) \quad C^*(G)_W \cdot \mathcal{H} \subseteq \mathcal{H} \cdot \psi(W).$$

We briefly call a $C^*(G), B$ -correspondence \mathcal{H} satisfying (6.1) $\mathbb{I}_p(G)$ -covariant.

Equation (6.1) implies that \mathcal{H} restricts to a proper $C^*(G)_W, \psi(W)$ -correspondence and thus induces a grading-preserving group homomorphism

$$\mathcal{H}_*|_W: K_*(C^*(G)_W) \rightarrow K_*(\psi(W)).$$

These homomorphisms for $W \in \mathbb{I}_p(G)$ form a morphism of diagrams

$$\mathcal{H}_*: K_*(C^*(G), \mathbb{I}_p) \rightarrow \psi^*K_*(B, \mathbb{I}_p).$$

We shall not use this on K_0 in the following because it contains less information than the homomorphism $\mathcal{H}_*: \mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B)$ (compare Theorem 4.12). We now describe the map $\mathcal{H}_*|_W: K_1(C^*(G), \mathbb{I}_p) \rightarrow \psi^*K_1(B, \mathbb{I}_p)$. Since the map $\kappa|_W$ in (5.4) is injective, we may identify $K_1(C^*(G)_W) \cong K_1(C^*(G|_W))$ with the kernel of the map $\text{id} - M_G^W: \mathbb{Z}[W_{\text{reg}}] \rightarrow \mathbb{Z}[W]$. Let $x = \sum c_v \delta_v \in \mathbb{Z}[W_{\text{reg}}]$ belong to this kernel. Write $x = \sum c_v^+ \delta_v - \sum c_v^- \delta_v$ with $c_v^\pm \geq 0$, $c_v = c_v^+ - c_v^-$ and $c_v^+ = 0$ or $c_v^- = 0$ for all $v \in W_{\text{reg}}$. Set $c_v = 0$ for $v \in V_{\text{reg}} \setminus W$. The equation $(\text{id} - M_G^W)(x) = 0$ says that

$$(6.2) \quad \sum_{v \in V_{\text{reg}}} c_v^+ \delta_v + \sum_{e \in E} c_{r(e)}^- \delta_{s(e)} = \sum_{e \in E} c_{r(e)}^+ \delta_{s(e)} + \sum_{v \in V_{\text{reg}}} c_v^- \delta_v.$$

Equivalently, the domains and codomains of the partial isometry

$$u_x := \bigoplus_{v \in W_{\text{reg}}} u_v^{\oplus c_v^+} \oplus \bigoplus_{v \in W_{\text{reg}}} (u_v^*)^{\oplus c_v^-}$$

are equal up to a permutation of the summands. Let u'_x be the product of u_x with the permutation matrix that makes it a unitary element in the corner of $\mathbb{M}_\infty(C^*(G)_W)$

associated to the projection

$$p_x := \bigoplus_{v \in V_{\text{reg}}} p_v^{\oplus c_v^+} + \bigoplus_{e \in E} p_{s(e)}^{\oplus c_{r(e)}^-}.$$

Then u'_x has a class in $K_1(C^*(G)_W)$. The exact sequence (5.5) is proved by showing that any class in $K_1(C^*(G)_W)$ is of this form for a unique $x \in \ker(\text{id} - M_G^W)$. Let $P_v \in \mathbb{K}(\mathcal{H})$ be the projection onto \mathcal{H}_v and let U_v be as in Theorem 3.2. Define

$$P_x := \bigoplus_{v \in V_{\text{reg}}} P_v^{\oplus c_v^+} \oplus \bigoplus_{e \in E} P_{s(e)}^{\oplus c_{r(e)}^-},$$

and define a unitary U'_x in $P_x \mathbb{K}(\mathcal{H}^\infty) P_x$ in the same way as u'_x . This is how $u'_x \in p_x \mathbb{M}_\infty(C^*(G)) p_x$ acts on $P_x \mathcal{H}^\infty$, and the latter is a Hilbert $\psi(W)$ -module. The map $\mathcal{H}_*|_W: K_1(C^*(G)_W) \rightarrow K_1(\psi(W))$ maps x to $[U'_x]$.

Now let $\varphi_0: \mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B)$ be a monoid homomorphism. Let

$$\psi: \mathbb{I}_p(G) \cong \mathbb{I}_p(C^*(G)) \rightarrow \mathbb{I}_p(B)$$

be induced by φ_0 . And let $\varphi_1: K_1(C^*(G), \mathbb{I}_p) \rightarrow \psi^* K_1(B, \mathbb{I}_p)$ be a morphism of diagrams over $\mathbb{I}_p(G)$. When is there a proper $C^*(G), B$ -correspondence \mathcal{H} that induces the maps φ_0 and φ_1 ? Our criterion needs a variant of the obstruction class in [9]. Our variant lives in the Ext^2 -group

$$\mathfrak{Ert}^2(G, \psi^* K_1(B, \mathbb{I}_p)) := \text{Ext}_{\mathbb{I}_p(G)}^2(K_0(C^*(G), \mathbb{I}_p), \psi^* K_1(B, \mathbb{I}_p)).$$

The extension group in question is described in (5.7) as the cokernel of the map

$$\kappa^*: \text{Hom}_{\mathbb{I}_p(G)}(\mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p], \psi^* K_1(B, \mathbb{I}_p)) \rightarrow \text{Hom}_{\mathbb{I}_p(G)}(K_1(C^*(G), \mathbb{I}_p), \psi^* K_1(B, \mathbb{I}_p))$$

induced by the canonical injective diagram morphism $\kappa: K_1(C^*(G), \mathbb{I}_p) \rightarrow \mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p]$ in (5.5). So elements in $\mathfrak{Ert}^2(G, \psi^* K_1(B, \mathbb{I}_p))$ are represented by diagram morphisms from $K_1(C^*(G), \mathbb{I}_p)$ to $\psi^* K_1(B, \mathbb{I}_p)$, and such a diagram morphism represents 0 if and only if it factors through κ .

Definition 6.3. Let $\varphi_0: \mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B)$ be a homomorphism. It induces a map $\psi: \mathbb{I}_p(G) \cong \mathbb{I}_p(C^*(G)) \rightarrow \mathbb{I}_p(B)$. Let

$$\varphi_1: K_1(C^*(G), \mathbb{I}_p) \rightarrow \psi^* K_1(B, \mathbb{I}_p)$$

be a morphism of diagrams. By Theorem 3.2, there is a correspondence \mathcal{H} that lifts φ_0 . The *obstruction class* $\Delta(\varphi_0, \varphi_1) \in \mathfrak{Ert}_{\mathbb{I}_p(G)}^2(G, \psi^* K_1(B, \mathbb{I}_p))$ of (φ_0, φ_1) is the image of the diagram morphism $\varphi_1 - \mathcal{H}_*: K_1(C^*(G), \mathbb{I}_p) \rightarrow \psi^* K_1(B, \mathbb{I}_p)$.

As the name indicates, the obstruction class obstructs the existence of a proper correspondence that lifts both maps φ_0 and φ_1 :

Proposition 6.4. *The obstruction class does not depend on the choice of the proper correspondence \mathcal{H} lifting φ_0 . It vanishes if there is a proper correspondence \mathcal{H} that induces the given maps φ_0 and φ_1 on \mathcal{P} and K_1 .*

Proof. By definition, $\Delta(\varphi_0, \varphi_1) = 0$ if \mathcal{H} induces φ_0 and φ_1 on \mathcal{P} and K_1 . So the second statement follows if the obstruction class does not depend on the choice of \mathcal{H} . This is what we are going to prove.

Let \mathcal{H}' be another proper correspondence lifting φ_0 . This involves isomorphic Hilbert B -modules \mathcal{H}_v because $[\mathcal{H}_v] = \varphi_0[p_v]$. So \mathcal{H}' is isomorphic to the proper correspondence for a family $((\mathcal{H}_v), (U'_v))$ with the same proper Hilbert B -modules \mathcal{H}_v . By Lemma 3.5, the unitaries U'_v are of the form $U'_v = \Upsilon_v U_v$ with $\Upsilon_v \in \mathcal{U}(\mathbb{K}(\mathcal{H}_v))$ for $v \in V_{\text{reg}}$. Each Υ_v has a class in $K_1(\mathbb{K}(\mathcal{H}_v))$. The C^* -algebra $\mathbb{K}(\mathcal{H}_v)$ is Morita-Rieffel equivalent through \mathcal{H}_v to the ideal in B generated by the inner products of

elements in \mathcal{H}_v . This ideal is equal to $\psi(\langle v \rangle) \triangleleft B$, where $\langle v \rangle \subseteq V$ is the hereditary and saturated subset generated by v . Thus \mathcal{H}' yields a diagram morphism

$$(6.5) \quad [\Upsilon]: \mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p] \rightarrow \psi^*(\mathbf{K}_1(B, \mathbb{I}_p)), \quad \delta_v \mapsto [\Upsilon_v],$$

compare the proof of Proposition 5.6. We claim that

$$(6.6) \quad \mathcal{H}_* - \mathcal{H}'_* = [\Upsilon] \circ \kappa \quad \text{as maps } \mathbf{K}_1(\mathbf{C}^*(G), \mathbb{I}_p) \rightarrow \psi^* \mathbf{K}_1(B, \mathbb{I}_p)$$

with the map κ in (5.5). So \mathcal{H} and \mathcal{H}' have the same image in $\mathfrak{Ert}^2(G, \psi^* \mathbf{K}_1(B, \mathbb{I}_p))$, as desired. To prove (6.6), choose a hereditary and saturated subset $W \subseteq V$ and $x \in \mathbf{K}_1(\mathbf{C}^*(G)_W)$. Write $\kappa(x) = \sum_{v \in W_{\text{reg}}} (c_v^+ - c_v^-) \delta_v$ with $c_v^\pm \geq 0$. Let $v \in W_{\text{reg}}$.

The Hilbert B -modules $\mathcal{F}_v := \bigoplus_{v \in V} \mathcal{H}_v^{\oplus c_v^+} \oplus \bigoplus_{e \in E^v} \mathcal{H}_{s(e)}^{\oplus c_v^-}$ and $\bigoplus_{v \in V} \mathcal{H}_v^{\oplus c_v^-} \oplus \bigoplus_{e \in E^v} \mathcal{H}_{s(e)}^{\oplus c_v^+}$ are isomorphic by a permutation Σ of the summands because $(\text{id} - M_G^W) \kappa(x) = 0$. Since their inner products take values in $\psi(W) \triangleleft B$, they are Hilbert $\psi(W)$ -modules. The unitaries

$$U_x := \bigoplus_{v \in V_{\text{reg}}} U_v^{\oplus c_v^+} \oplus \bigoplus_{v \in V_{\text{reg}}} (U_v^*)^{\oplus c_v^-},$$

$$U'_x := \bigoplus_{v \in V_{\text{reg}}} (\Upsilon_v U_v)^{\oplus c_v^+} \oplus \bigoplus_{v \in V_{\text{reg}}} (U_v^* \Upsilon_v^*)^{\oplus c_v^-}$$

are isomorphisms between these Hilbert $\psi(W)$ -modules as well. So $\Sigma U_x, \Sigma U'_x \in \mathcal{U}(\mathcal{F}_v)$. By definition, $\mathcal{H}_*(x)$ and $\mathcal{H}'_*(x)$ are the classes in $\mathbf{K}_1(\psi(W))$ defined by the unitaries ΣU_x and $\Sigma U'_x$. They differ by the class of $\bigoplus_{v \in V_{\text{reg}}} \Upsilon_v^{\oplus c_v^+} \oplus \bigoplus_{v \in V_{\text{reg}}} (\Upsilon_v^*)^{\oplus c_v^-}$. This is $[\Upsilon](\kappa(x))$ because $[\Upsilon_v^*] = -[\Upsilon_v]$ in $\mathbf{K}_1(\psi(W))$. \square

The vanishing of the obstruction class is necessary for the existence of a proper correspondence \mathcal{H} that lifts (φ_0, φ_1) . We are going to show that it is also sufficient under an extra assumption on B .

Definition 6.7. A unital \mathbf{C}^* -algebra B is \mathbf{K}_1 -surjective or \mathbf{K}_1 -injective, respectively, if the map $\pi_0(\mathcal{U}(B)) \rightarrow \mathbf{K}_1(B)$ is surjective or injective, respectively.

Proposition 6.8. *Let G be a graph, B a \mathbf{C}^* -algebra, and let $\varphi_0: \mathcal{P}(\mathbf{C}^*(G)) \rightarrow \mathcal{P}(B)$ and $\varphi_1: \mathbf{K}_1(\mathbf{C}^*(G), \mathbb{I}_p) \rightarrow \psi^* \mathbf{K}_1(B, \mathbb{I}_p)$ be given, where $\psi: \mathbb{I}_p(G) \rightarrow \mathbb{I}_p(B)$ is induced by φ_0 . For $v \in V$, let \mathcal{H}_v be proper Hilbert B -modules with $[\mathcal{H}_v] = \varphi_0[p_v]$. Assume that the unital \mathbf{C}^* -algebras $\mathbb{K}(\mathcal{H}_v)$ for $v \in V$ are \mathbf{K}_1 -surjective. There is a proper $\mathbf{C}^*(G)$, B -correspondence \mathcal{H} that induces the given maps φ_0 and φ_1 if and only if $\Delta(\varphi_0, \varphi_1) = 0$.*

Proof. One direction is proven already in Proposition 6.4, without assumption on B . Here we prove the converse. So assume that the obstruction class vanishes. We shall modify a proper $\mathbf{C}^*(G)$, B -correspondence \mathcal{H} that lifts φ_0 so that it also lifts φ_1 . The vanishing of the obstruction class means that

$$\varphi_1 - \mathcal{H}_* = \alpha \circ \kappa: \mathbf{K}_1(\mathbf{C}^*(G), \mathbb{I}_p) \rightarrow \psi^* \mathbf{K}_1(B, \mathbb{I}_p)$$

for a diagram morphism $\alpha: \mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p] \rightarrow \psi^* \mathbf{K}_1(B, \mathbb{I}_p)$. Describe \mathcal{H} by a family $((\mathcal{H}_v), (U_v))$ as above. Let $v \in V_{\text{reg}}$ and let $\langle v \rangle \subseteq V$ be the hereditary and saturated subset generated by v . Since $\psi(\langle v \rangle) \triangleleft B$ is Morita–Rieffel equivalent to $\mathbb{K}(\mathcal{H}_v)$ and $\mathbb{K}(\mathcal{H}_v)$ is \mathbf{K}_1 -surjective, $\alpha(\delta_v) \in \mathbf{K}_1(\psi(\langle v \rangle)) \cong \mathbf{K}_1(\mathbb{K}(\mathcal{H}_v))$ is represented by some $\Upsilon_v \in \mathcal{U}(\mathbb{K}(\mathcal{H}_v))$.

Define $U'_v := \Upsilon_v \circ U_v$. The family $((\mathcal{H}_v), (U'_v))$ gives another proper $\mathbf{C}^*(G)$, B -correspondence \mathcal{H}' lifting the same map φ_0 . Let $[\Upsilon_v]$ denote the class of Υ_v in $\mathbf{K}_1(\psi(\langle v \rangle)) \cong \mathbf{K}_1(\mathbb{K}(\mathcal{H}_v))$. As above, this defines a morphism of diagrams $\mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p] \rightarrow \psi^* \mathbf{K}_1(B, \mathbb{I}_p)$. The proof of Proposition 6.4 shows that $\mathcal{H}'_* - \mathcal{H}_*$ is equal to $[\Upsilon_v] \circ \kappa = \alpha \circ \kappa = \varphi_1 - \mathcal{H}_*$. As a result, $\mathcal{H}'_* = \varphi_1$. \square

Now assume that the target B is a graph C*-algebra as well. The following theorem allows us to apply all of the previous results to study A, B -correspondences:

Theorem 6.9. *Let $G_2 = (r, s: E_2 \rightrightarrows V_2)$ be another regular directed graph and let B be its graph C*-algebra. Let $p \in \mathcal{P}(B)$. Then B has stable weak cancellation and the unital C*-algebra $p(B \otimes \mathbb{K})p$ is K_1 -bijective.*

Proof. Graph C*-algebras have stable weak cancellation by [3, Corollary 7.2]. By Theorem 3.1, any $p \in \mathcal{P}(B)$ is equivalent to $\sum_{v \in V_2} c_v [p_v]$ for a finitely supported function $(c_v)_{v \in V_2} \in \mathbb{N}[V]$.

For a finite subset $V'_2 \subseteq V_2$, let B' be the C*-subalgebra of B generated by $\{t_e : r(e) \in V'_2\} \cup \{p_v : v \in V'_2\}$. These form a directed set of C*-subalgebras of B , and their union is dense in B . Therefore, $p(B \otimes \mathbb{K})p$ is the inductive limit of the C*-algebras $p(B' \otimes \mathbb{K})p$. Both K_1 and the set of connected components of the unitary group are continuous for inductive limits. Therefore, it is enough to prove that $p(B' \otimes \mathbb{K})p$ is K_1 -bijective for all finite subsets $V'_2 \subseteq V_2$. We may assume without loss of generality that V'_2 contains $\text{supp}(c_v)$, so that $p \in B' \otimes \mathbb{K} \subseteq B \otimes \mathbb{K}$. The C*-algebra B' may be identified with the C*-algebra of a finite graph (compare [33, Lemma 1.2]). Here we may take the graph with vertex set $V'_2 \cup s(r^{-1}(V'_2))$ and edge set $r^{-1}(V'_2)$. Therefore, it suffices to prove that $p(B \otimes \mathbb{K})p$ is K_1 -bijective when B is the graph C*-algebra of a finite graph. We assume this from now on.

The advantage in the finite case is that there are only finitely many gauge-invariant ideals. Therefore, there is an increasing chain of gauge-invariant ideals $0 = I_0 \subset I_1 \subset I_2 \subset \dots \subset I_\ell = B$ so that each subquotient I_{k+1}/I_k for $k = 0, \dots, \ell-1$ has no gauge-invariant ideals. It is known that these subquotients I_{k+1}/I_k are again Morita equivalent to graph C*-algebras, so that they have stable weak cancellation. In addition, they are either AF or of the form $C(\mathbb{T}, \mathbb{M}_n)$ for some $n \in \mathbb{N}_{\geq 1}$ or purely infinite simple. In the first two cases, $p \cdot (I_{k+1}/I_k \otimes \mathbb{K})p$ has stable rank 1 (see [33, Proposition 5.4]). In all three cases, $p \cdot (I_{k+1}/I_k \otimes \mathbb{K}) \cdot p$ is extremally rich and has weak cancellation (see [11]). Brown and Pedersen show in [11] that extremally rich C*-algebras with weak cancellation have the following properties:

- K_1 -surjectivity by [11, Theorem 4.4];
- good index theory by [11, Theorem 5.1];
- K_1 -injectivity and weak K_0 -surjectivity by [11, Theorem 6.7].

It is shown in [11] that the class of C*-algebras with these four properties – weak K_0 -surjectivity, K_1 -surjectivity, K_1 -injectivity, and good index theory – is closed under extensions (see [11, Propositions 6.6 and 7.2.4]). Therefore, $p(B \otimes \mathbb{K})p$ has these four properties. In particular, it is K_1 -bijective. \square

7. THE PURELY INFINITE CASE

At this point, our results combined with Kirchberg's Classification Theorem imply a criterion when there is a C*-algebra isomorphism $C^*(G) \cong B$, even without assuming B to be a graph C*-algebra:

Theorem 7.1. *Let G be a countable graph for which $C^*(G)$ is purely infinite. Let B be a separable, nuclear, strongly purely infinite C*-algebra such that all ideals of B are in the bootstrap class and such that $\text{Prim}(B)$ has a basis consisting of quasi-compact open subsets. Assume that there are an isomorphism $\psi: \mathbb{I}_p(G) \rightarrow \mathbb{I}_p(B)$ of partially ordered sets and isomorphisms of diagrams $\varphi_j: K_j(C^*(G), \mathbb{I}_p) \xrightarrow{\sim} \psi^* K_j(B, \mathbb{I}_p)$ for $j = 0, 1$ for which the obstruction class vanishes. Then $C^*(G)$ and B are stably isomorphic, and vice versa.*

The proof of the converse direction (“vice versa”) is easy. The proof of the interesting direction requires some preparatory results.

Lemma 7.2. *Let X be a topological space. Let $\mathbb{C}\mathbb{O}(X) \subseteq \mathbb{O}(X)$ be the set of compact-open subsets. Assume that $\mathbb{C}\mathbb{O}(X)$ is a basis. Let $\mathcal{F}(X)$ denote the set of subsets of $\mathbb{C}\mathbb{O}(X)$ which are closed under finite unions and subsets. Then the map $U \mapsto F_U := \{V \in \mathbb{C}\mathbb{O}(X) \mid V \subseteq U\}$ is an isomorphism of partially ordered sets $\mathbb{O}(X) \cong \mathcal{F}(X)$ with inverse $S \mapsto \bigcup S$.*

Proof. The maps $F: \mathbb{O}(X) \rightarrow \mathcal{F}(X)$ and $\bigcup: \mathcal{F}(X) \rightarrow \mathbb{O}(X)$ are well defined monotone maps. One composite is the identity as $U = \bigcup \{V \in \mathbb{C}\mathbb{O}(X) : V \subseteq U\}$ for all $U \in \mathbb{O}(X)$ by assumption on X ; the other is the identity because

$$S \subseteq \{V \in \mathbb{C}\mathbb{O}(X) : V \subseteq \bigcup S\} \subseteq S$$

for all $S \in \mathcal{F}(X)$, where the first inclusion is obvious and the second is seen as follows: if $V \subseteq \bigcup S$ is compact, then $V \subseteq \bigcup T$ for some finite subset $T \subseteq S$ and then $V \in S$ by assumption on S . \square

Proposition 7.3. *Let B be a separable purely infinite C^* -algebra such that the topology on $\text{Prim}(B)$ has a basis consisting of compact-open subsets. Let $\mathcal{F}(\mathbb{I}_p(B))$ denote the set of subsets of $\mathbb{I}_p(B)$ which are closed under finite unions and subsets. Then $I \mapsto F_I := \{J \in \mathbb{I}_p(B) \mid J \subseteq I\}$ defines an isomorphism of partially ordered sets $\mathbb{I}(B) \cong \mathcal{F}(\mathbb{I}_p(B))$ with inverse $S \mapsto \overline{\sum S}$.*

Proof. [31, Proposition 2.7] says that $\mathbb{I}_p(B)$ is the set of all ideals $I \triangleleft B$ for which \widehat{I} is quasi-compact. Now we can deduce the result from Lemma 7.2, using the isomorphism $\mathbb{O}(\text{Prim}(A)) \cong \mathbb{I}(A)$, $U \mapsto \bigcap \text{Prim}(A) \setminus U$ (see [14, §3.2]). \square

Proof of Theorem 7.1. Since $C^*(G)$ and B are purely infinite, Corollary 4.16 shows that an isomorphism $\varphi_0: K_0(C^*(G), \mathbb{I}_p) \xrightarrow{\sim} \psi^* K_0(B, \mathbb{I}_p)$ is equivalent to an isomorphism $\mathcal{P}(C^*(G)) \cong \mathcal{P}(B)$. Then the obstruction class in the statement is well defined. If it vanishes, Proposition 6.8 gives a proper $C^*(G)$, B -correspondence \mathcal{H} that induces the given isomorphisms φ_0 and φ_1 . The proper correspondence \mathcal{H} has a class in the equivariant KK-group $\text{KK}(\text{Prim } C^*(G); C^*(G), B)$. It induces an isomorphism on the K-theory of all ideals in $\mathbb{I}_p(B) \cong \mathbb{I}_p(C^*(G))$. By the Universal Coefficient Theorem, it induces E-theory equivalences on these ideals.

Every ideal in $C^*(G)$ or in B is generated by its projections. This follows for $C^*(G)$ because every ideal in a purely infinite graph C^* -algebra is gauge-invariant and for B by [31, Proposition 2.7]. It also follows from [31, Proposition 2.7] that $\text{Prim}(C^*(G))$ has a basis consisting of quasi-compact open subsets. Hence Proposition 7.3 shows that ψ extends canonically to a lattice isomorphism $\mathbb{I}(C^*(G)) \xrightarrow{\sim} \mathbb{I}(B)$. Thus $\text{Prim}(C^*(G)) \cong \text{Prim}(B)$. The quasi-compact open subsets form a basis for the topology on these primitive ideal spaces, and the correspondence \mathcal{H} induces E-theory equivalences on those ideals that correspond to quasi-compact open subsets. Thus [12, Theorems 3.2 and 3.10] imply that \mathcal{H} induces a $\text{Prim } C^*(G)$ -equivariant E-theory equivalence. Then the corresponding equivariant KK-class is also invertible by [22, Theorem 6.2]. Hence Kirchberg's Classification Theorem allows to lift it to a stable isomorphism between $C^*(G)$ and B . \square

Remark 7.4. The constructed stable isomorphism $C^*(G) \otimes \mathbb{K} \cong B \otimes \mathbb{K}$ also gives a proper $C^*(G)$, B -correspondence. So it corresponds to a particular choice of the proper Hilbert B -modules \mathcal{H}_v and the unitaries U_v in our description of such correspondences. Of course, it is very hard to find these unitaries directly.

If the target C^* -algebra B is known to be a graph C^* -algebra as well, then the proof of Theorem 7.1 simplifies considerably. Then Theorem 8.7 and Proposition 7.3 show that the C^* -correspondence above is an ideal-preserving stable homotopy equivalence. This forces it to be an ideal-preserving KK-equivalence.

8. HOMOTOPY CLASSES OF CORRESPONDENCES

In this section, we first study when two proper $C^*(G), B$ -correspondences \mathcal{H} and \mathcal{H}' are homotopic. Secondly, we use this to prove a criterion for two graph C^* -algebras to be homotopy equivalent.

Lemma 8.1. *Let \mathcal{H} and \mathcal{H}' be homotopic. Then they induce the same maps*

$$\begin{aligned} \varphi_0: \mathcal{P}(C^*(G)) &\rightarrow \mathcal{P}(B), & \psi: \mathbb{I}_p^\infty(C^*(G)) &\rightarrow \mathbb{I}_p^\infty(B), \\ \varphi_1: K_1(C^*(G), \mathbb{I}_p) &\rightarrow \psi^*K_1(B, \mathbb{I}_p). \end{aligned}$$

The entire homotopy $(\mathcal{H}_t)_{t \in [0,1]}$ preserves all ideals in $\mathbb{I}_p^\infty(C^*(G))$ in the sense that each \mathcal{H}_t induces the same map $\mathbb{I}_p^\infty(C^*(G)) \rightarrow \mathbb{I}_p^\infty(B)$.

Proof. The maps $\mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B)$ induced by \mathcal{H} and \mathcal{H}' are the same because homotopic projections are Murray–von Neumann equivalent. Therefore, \mathcal{H} and \mathcal{H}' – and also \mathcal{H}_t for $t \in [0, 1]$ – induce the same map $\mathbb{I}_p^\infty(C^*(G)) \rightarrow \mathbb{I}_p^\infty(B)$. Since K_1 is defined using homotopy of unitaries, it follows that the diagram morphisms $K_1(C^*(G), \mathbb{I}_p) \rightarrow \psi^*K_1(B, \mathbb{I}_p)$ induced by \mathcal{H} and \mathcal{H}' are equal as well. Even more, they are constant along the homotopy (\mathcal{H}_t) . \square

Let \mathcal{H} and \mathcal{H}' be $C^*(G), B$ -correspondences that induce the same maps

$$\varphi_0: \mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B), \quad \varphi_1: K_1(C^*(G), \mathbb{I}_p) \rightarrow \psi^*K_1(B, \mathbb{I}_p).$$

The first condition implies that \mathcal{H} and \mathcal{H}' also induce the same map $\psi: \mathbb{I}_p(C^*(G)) \rightarrow \mathbb{I}_p(B)$, which is needed to formulate the second condition. Describe \mathcal{H} and \mathcal{H}' by families $((\mathcal{H}_v), (U_v))$ and $((\mathcal{H}'_v), (U'_v))$, respectively, as in Theorem 3.2. Since $[\mathcal{H}_v] = [\mathcal{H}'_v]$ in $\mathcal{P}(B)$, the Hilbert B -modules \mathcal{H}_v and \mathcal{H}'_v are isomorphic. For simplicity, we replace \mathcal{H}' by an isomorphic correspondence so that these Hilbert modules become equal. Then Lemma 3.5 gives unitaries $\Upsilon_v \in \mathbb{K}(\mathcal{H}_v)$ for $v \in V_{\text{reg}}$ with $U'_v = \Upsilon_v U_v$.

Next, we describe an obstruction for \mathcal{H} and \mathcal{H}' to be homotopic. This obstruction will turn out to be sharp under an extra assumption on B . Let $v \in V_{\text{reg}}$. Let $\langle v \rangle \in \mathbb{I}_p(G)$ be the hereditary and saturated subset generated by v . Since this corresponds to the ideal in $C^*(G)$ generated by p_v , it follows that $\psi(\langle v \rangle) \triangleleft B$ is the ideal generated by the matrix coefficients of vectors in \mathcal{H}_v . Thus \mathcal{H}_v is a Morita–Rieffel equivalence between $\mathbb{K}(\mathcal{H}_v)$ and $\psi(\langle v \rangle)$. As in the proof of Proposition 6.8, we let $[\Upsilon_v]$ denote the class of Υ_v in $K_1(\psi(\langle v \rangle)) \cong K_1(\mathbb{K}(\mathcal{H}_v))$. We combine these classes for $v \in V_{\text{reg}}$ into a morphism of diagrams $[\Upsilon]: \mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p] \rightarrow \psi^*K_1(B, \mathbb{I}_p)$ as in (6.5). Since \mathcal{H} and \mathcal{H}' induce the same map φ_1 , (6.6) implies $[\Upsilon] \circ \kappa = 0$.

We have described the group

$$\mathfrak{Ert}^1(G, \psi^*K_1(B, \mathbb{I}_p)) := \text{Ext}_{\mathbb{I}_p(G)}^1(K_0(C^*(G), \mathbb{I}_p), \psi^*K_1(B, \mathbb{I}_p))$$

as the first cohomology group of the cochain complex in (5.7). The equation $[\Upsilon] \circ \kappa = 0$ says that $[\Upsilon]$ is a cocycle and thus defines a class in $\mathfrak{Ert}^1(G, \psi^*K_1(B, \mathbb{I}_p))$.

Proposition 8.2. *If \mathcal{H} and \mathcal{H}' are homotopic $C^*(G), B$ -correspondences, then $[\Upsilon]$ becomes 0 in $\mathfrak{Ert}^1(G, \psi^*K_1(B, \mathbb{I}_p))$. In particular, $[\Upsilon]$ does not depend on the choices made in its construction.*

Proof. By the remarks after Lemma 3.5, \mathcal{H} and \mathcal{H}' are homotopic if and only if we may find unitaries $W_{v,0}, W_{v,1} \in \mathcal{U}(\mathcal{H}_v)$ for all $v \in V_{\text{reg}}$ so that

$$\begin{aligned} \Upsilon_{v,0} &:= W_{v,0} \circ U_v \circ \left(\bigoplus_{e \in E^v} W_{s(e),0}^* \right) \circ U_v^*, \\ \Upsilon_{v,1} &:= W_{v,1} \circ \Upsilon_v U_v \circ \left(\bigoplus_{e \in E^v} W_{s(e),1}^* \right) \circ U_v^* \end{aligned}$$

are homotopic in $\mathcal{U}(\mathcal{H}_v)$. Equivalently, Υ_v is homotopic to

$$(8.3) \quad W_v \cdot U_v \left(\bigoplus_{e \in E^v} W_{s(e)}^* \right) U_v^*,$$

where $W_v := W_{v,1}^* W_{v,0}$. Assume first that this is the case. In the same way that the unitaries $\Upsilon_v \in \mathcal{U}(\mathcal{H}_v)$ for $v \in V_{\text{reg}}$ give a map $[\Upsilon]: \mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p] \rightarrow \psi^* \mathbb{K}_1(B, \mathbb{I}_p)$, the unitaries W_v give a map $[W]: \mathbb{Z}[V, \mathbb{I}_p] \rightarrow \psi^* \mathbb{K}_1(B, \mathbb{I}_p)$. The class of the unitary in (8.3) in $\mathbb{K}_1(\mathbb{K}(\mathcal{H}_v)) \cong \mathbb{K}_1(\psi(\langle v \rangle))$ is $[W_v] - \sum_{e \in E^v} [W_{s(e)}]$. Therefore, $[W] \circ (\text{id} - M_G^{\mathbb{I}_p}) = [\Upsilon]$ as diagram morphisms $\mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p] \rightarrow \psi^* \mathbb{K}_1(B, \mathbb{I}_p)$. This witnesses that $[\Upsilon]$ represents zero in $\mathfrak{Krt}^1(G, \psi^* \mathbb{K}_1(B, \mathbb{I}_p))$. \square

Theorem 8.4. *Describe a $C^*(G)$, B -correspondences \mathcal{H} by a family $((\mathcal{H}_v), (U_v))$ as in Theorem 3.2. Let*

$$\varphi_0: \mathcal{P}(C^*(G)) \rightarrow \mathcal{P}(B), \quad \varphi_1: \mathbb{K}_1(C^*(G), \mathbb{I}_p) \rightarrow \psi^* \mathbb{K}_1(B, \mathbb{I}_p)$$

be the maps induced by \mathcal{H} . Assume that the unital C^* -algebras $\mathbb{K}(\mathcal{H}_v)$ for $v \in V_{\text{reg}}$ are \mathbb{K}_1 -bijective. Then the construction in Proposition 8.2 gives a bijection between the set of homotopy classes of $C^*(G)$, B -correspondences that induce the same maps φ_0 and φ_1 and the group $\mathfrak{Krt}^1(G, \psi^* \mathbb{K}_1(B, \mathbb{I}_p))$.

Proof. First let \mathcal{H}' and \mathcal{H}'' be $C^*(G)$, B -correspondences that induce the same maps φ_0 and φ_1 as \mathcal{H} . We are going to prove that they are homotopic. Describe \mathcal{H}' and \mathcal{H}'' by families $((\mathcal{H}'_v), (U'_v))$ and $((\mathcal{H}''_v), (U''_v))$ and replace them by isomorphic correspondences with $\mathcal{H}'_v = \mathcal{H}''_v = \mathcal{H}_v$ for all $v \in V$. Then there are unitaries Υ'_v and Υ''_v for $v \in V_{\text{reg}}$ with $U'_v = \Upsilon'_v U_v$ and $U''_v = \Upsilon''_v U_v$. Assume that $[\Upsilon'] = [\Upsilon'']$ holds in $\mathfrak{Krt}^1(G, \psi^* \mathbb{K}_1(B, \mathbb{I}_p))$. Then there is a map $\beta: \mathbb{Z}[V, \mathbb{I}_p] \rightarrow \psi^* \mathbb{K}_1(B, \mathbb{I}_p)$ with $\beta \circ (\text{id} - M_G^{\mathbb{I}_p}) = [\Upsilon''] - [\Upsilon'] = [(\Upsilon')^* \Upsilon'']$. The proof of Proposition 6.8 shows that β is specified by elements $\beta_v \in \mathbb{K}_1(\psi(\langle v \rangle)) \cong \mathbb{K}_1(\mathbb{K}(\mathcal{H}_v))$ for $v \in V$. Since $\mathbb{K}(\mathcal{H}_v)$ is \mathbb{K}_1 -surjective, these \mathbb{K}_1 -classes are already represented by unitaries $W_v \in \mathcal{U}(\mathcal{H}_v)$. The equation $\beta \circ (\text{id} - M_G^{\mathbb{I}_p}) = [(\Upsilon')^* \Upsilon'']$ says that the unitaries in (8.3) and $(\Upsilon'_v)^* \Upsilon''_v$ represent the same class in $\mathbb{K}_1(\psi(\langle v \rangle)) \cong \mathbb{K}_1(\mathbb{K}(\mathcal{H}_v))$. Since $\mathbb{K}(\mathcal{H}_v)$ is \mathbb{K}_1 -injective, $(\Upsilon'_v)^* \Upsilon''_v$ and $W_v \cdot U_v \left(\bigoplus_{e \in E^v} W_{s(e)}^* \right) U_v^*$ are homotopic in $\mathbb{K}(\mathcal{H}_v)$. Similarly, $\Upsilon'_v W_v$ is

homotopic to $\Upsilon'_v W_v$. So Υ''_v is homotopic in $\mathbb{K}(\mathcal{H}_v)$ to $W_v \cdot \Upsilon'_v U_v \left(\bigoplus_{e \in E^v} W_{s(e)}^* \right) U_v^*$.

The latter family together with \mathcal{H}_v for $v \in V$ describes a $C^*(G)$, B -correspondence that is isomorphic to \mathcal{H}' , and the homotopy of unitaries gives a homotopy between the latter and \mathcal{H}'' . So \mathcal{H}' and \mathcal{H}'' are homotopic.

Next, take any class in $\mathfrak{Krt}^1(G, \psi^* \mathbb{K}_1(B, \mathbb{I}_p))$. This is represented by some diagram morphism $v: \mathbb{Z}[V_{\text{reg}}, \mathbb{I}_p] \rightarrow \psi^* \mathbb{K}_1(B, \mathbb{I}_p)$ with $v \circ \kappa = 0$. For $v \in V_{\text{reg}}$, the value of this map on $\delta_v \in \mathbb{Z}[\langle v \rangle]$ is a class in $\mathbb{K}_1(\psi(\langle v \rangle)) \cong \mathbb{K}_1(\mathbb{K}(\mathcal{H}_v))$. The latter is represented by a unitary $\Upsilon_v \in \mathbb{K}(\mathcal{H}_v)$ because $\mathbb{K}(\mathcal{H}_v)$ is \mathbb{K}_1 -bijective. The family $((\mathcal{H}_v), (\Upsilon_v U_v))$ describes a $C^*(G)$, B -correspondence \mathcal{H}' that induces the same map φ_0 . Since $v \circ \kappa = 0$, it also induces the same diagram morphism $\mathbb{K}_1(C^*(G), \mathbb{I}_p) \rightarrow \psi^* \mathbb{K}_1(B, \mathbb{I}_p)$. By construction, \mathcal{H}' reproduces the class in $\mathfrak{Krt}^1(G, \psi^* \mathbb{K}_1(B, \mathbb{I}_p))$ that we started with. \square

Now let G and G' be two graphs. We denote the sets of vertices and edges of G, G' by V, V' and E, E' , respectively. Let $A := C^*(G)$ and $B := C^*(G')$. Assume that there are an order isomorphism $\psi: \mathbb{I}_p(G) \cong \mathbb{I}_p(A) \xrightarrow{\sim} \mathbb{I}_p(B) \cong \mathbb{I}_p(G')$ and

diagram isomorphisms

$$\begin{aligned}\varphi_0^+ : K_0^+(A, \mathbb{I}_p) &\xrightarrow{\sim} \psi^* K_0^+(B, \mathbb{I}_p), \\ \varphi_1 : K_1(A, \mathbb{I}_p) &\xrightarrow{\sim} \psi^* K_1(B, \mathbb{I}_p).\end{aligned}$$

Since graph C*-algebras have stable weak cancellation by Theorem 6.9 and monoid isomorphisms preserve order units, ψ and φ_0^+ come from a monoid isomorphism

$$\varphi_0 : \mathcal{P}(A) \xrightarrow{\sim} \mathcal{P}(B)$$

by Theorem 4.12. When is there a stable homotopy equivalence between A and B that induces the pair of maps φ_0 and φ_1 ? In order to lift this pair of maps to proper correspondences, we need the obstruction classes of (φ_0, φ_1) and $(\varphi_0^{-1}, \varphi_1^{-1})$ to vanish. We are going to prove that this condition is also sufficient for a homotopy equivalence. First, we prove that one of these two obstruction classes vanishes if and only if the other does so.

Lemma 8.5. *The obstruction class of (φ_0, φ_1) vanishes if and only if the obstruction class of $(\varphi_0^{-1}, \varphi_1^{-1})$ vanishes.*

Proof. By symmetry, it suffices to prove one implication. So assume that the obstruction class of (φ_0, φ_1) vanishes. Then Proposition 6.8 gives an A, B -correspondence \mathcal{H}_{AB} that lifts (φ_0, φ_1) because graph C*-algebras are K_1 -surjective by Theorem 6.9. Theorem 3.2 gives a B, A -correspondence \mathcal{H}_{BA} that lifts φ_0^{-1} , but we do not yet control the map on K_1 induced by \mathcal{H}_{BA} . The composite B, B -correspondence $\mathcal{H}_{AB} \otimes_A \mathcal{H}_{BA}$ induces the map $\varphi_0 \circ \varphi_0^{-1} = \text{id}_{\mathcal{P}(B)}$ on $\mathcal{P}(B)$. So we may use it to compute the obstruction class for the maps $\text{id}_{\mathcal{P}(B)}$ and $\text{id}_{K_1(B, \mathbb{I}_p)}$. We may also use the identity correspondence on B to compute this obstruction class. Since the obstruction class is well defined by Proposition 6.4, both computations give the same result. The first one gives $\text{id}_{K_1(B, \mathbb{I}_p)} - \varphi_1 \circ \mathcal{H}_{BA, *}$ because $\mathcal{H}_{AB, *} = \varphi_1$ on K_1 . The second one gives just 0. Thus $\text{id}_{K_1(B, \mathbb{I}_p)} - \varphi_1 \circ \mathcal{H}_{BA, *}$ represents 0 in $\mathfrak{K}t^2(G', K_1(B, \mathbb{I}_p))$. This remains so when we compose with the isomorphism φ_1^{-1} . That is, $\varphi_1^{-1} - \mathcal{H}_{BA, *}: K_1(B, \mathbb{I}_p) \rightarrow (\psi^{-1})^* K_1(A, \mathbb{I}_p)$ represents 0 in $\mathfrak{K}t^2(G', (\psi^{-1})^* K_1(A, \mathbb{I}_p))$. At the same time, $\varphi_1^{-1} - \mathcal{H}_{BA, *}$ is exactly the obstruction class of $(\varphi_0^{-1}, \varphi_1^{-1})$. \square

Lemma 8.6. *A self-correspondence \mathcal{H}_A of a graph C*-algebra $A = C^*(G)$ that induces the identity maps on $\mathcal{P}(A)$ and $K_1(A, \mathbb{I}_p)$ is a stable homotopy equivalence. In addition, all maps in the homotopies involved here induce the identity map on the lattices of gauge-invariant ideals.*

Proof. Since \mathcal{H}_A induces the identity map on $\mathcal{P}(A)$, we may assume without loss of generality that the underlying Hilbert modules \mathcal{H}_v are $p_v A$ for all vertices v . Then \mathcal{H}_A is described through unitaries $\Upsilon_v \in p_v A p_v = \mathbb{K}(\mathcal{H}_v)$ for $v \in V_{\text{reg}}$. Namely, its representation in Theorem 3.2 involves the family of Hilbert modules $(p_v A)_{v \in V}$ and unitaries $(\Upsilon_v u_v)_{v \in V_{\text{reg}}}$ with the canonical unitaries u_v . Since $\mathcal{H}_v = \mathcal{H}'_v = p_v A$, the underlying right Hilbert module of \mathcal{H}_A is isomorphic to $\bigoplus_v p_v A = A$. Thus \mathcal{H}_A is isomorphic to the correspondence associated to the *-homomorphism $\sigma: A \rightarrow A$ defined by $\sigma(p_v) = p_v$ for $v \in V$ and $\sigma(t_e) = \Upsilon_{r(e)} t_e$ for $e \in E$.

Now let $\sigma': A \rightarrow A$ be defined by the conditions $\sigma'(p_v) = p_v$ for all $v \in V$ and $\sigma'(t_e) = \Upsilon_{r(e)}^* t_e$ for all $e \in E$. Then $\sigma' \circ \sigma(p_v) = p_v$ and $\sigma' \circ \sigma(t_e) = \Upsilon_{r(e)}^* \sigma(\Upsilon_{r(e)}) t_e$. Since σ induces the identity map on $K_1(A, \mathbb{I}_p)$, the unitaries $\Upsilon_{r(e)}^* \sigma(\Upsilon_{r(e)}) \in p_v A p_v$ represent zero in $K_1(p_v A p_v)$. Since $p_v C^*(A) p_v$ is K_1 -injective for any vertex v of the graph by Theorem 6.9, the unitary $\Upsilon_{r(e)}^* \sigma(\Upsilon_{r(e)})$ in $p_v A p_v$ is homotopic to the unit element p_v . This gives us a homotopy between $\sigma' \circ \sigma$ and the identity map (compare Theorem 8.4). Since σ and $\sigma' \circ \sigma$ induce the identity map on the invariants,

it follows that σ' does so as well. Then we may switch the roles of σ and σ' to show that $\sigma \circ \sigma'$ is homotopic to the identity map as well. Then σ is a homotopy equivalence. Since all gauge-invariant ideals are generated by the projections that they contain, and since all maps in the homotopies induce the same maps on $\mathcal{P}(A)$, it follows that all maps in the homotopies induce the identity map on the lattices of gauge-invariant ideals. \square

Theorem 8.7. *Let G and G' be two countable graphs, such that there are an order isomorphism $\psi: \mathbb{I}_p(G) \cong \mathbb{I}_p(A) \xrightarrow{\sim} \mathbb{I}_p(B) \cong \mathbb{I}_p(G')$ and diagram isomorphisms*

$$\begin{aligned} \varphi_0^+ : K_0^+(A, \mathbb{I}_p) &\xrightarrow{\sim} \psi^* K_0^+(B, \mathbb{I}_p), \\ \varphi_1 : K_1(A, \mathbb{I}_p) &\xrightarrow{\sim} \psi^* K_1(B, \mathbb{I}_p). \end{aligned}$$

These isomorphisms induce a monoid isomorphism $\varphi_0: \mathcal{P}(A) \xrightarrow{\sim} \mathcal{P}(B)$ as above. Assume that the obstruction class of (φ_0, φ_1) vanishes. Then there is a stable homotopy equivalence between $C^(G)$ and $C^*(G')$ that induces the maps φ_0^+ and φ_1 . The homotopies in this equivalence automatically preserve gauge-invariant ideals, that is, the induced maps on $\mathbb{I}^\mathbb{T}(C^*(G))$ and $\mathbb{I}^\mathbb{T}(C^*(G'))$ are constant along the homotopies.*

Proof. Since the obstruction class of (φ_0, φ_1) vanishes, the same holds for the inverse maps $(\varphi_0^{-1}, \varphi_1^{-1})$ by Lemma 8.5. Since graph C^* -algebras are K_1 -surjective by Theorem 6.9, Proposition 6.8 gives an A, B -correspondence \mathcal{H}_{AB} and a B, A -correspondence \mathcal{H}_{BA} that induce the maps (φ_0, φ_1) and $(\varphi_0^{-1}, \varphi_1^{-1})$ on the invariants. Let $\mathcal{H}_A := \mathcal{H}_{AB} \otimes_B \mathcal{H}_{BA}$ and $\mathcal{H}_B := \mathcal{H}_{BA} \otimes_A \mathcal{H}_{AB}$ be the composite A, A - and B, B -correspondences. Both induce the identity maps on \mathcal{P} and on the K_1 -diagram. Thus they are stable homotopy equivalences by Lemma 8.6. Thus \mathcal{H}_{AB} and \mathcal{H}_{BA} are both left and right invertible up to homotopy. This forces them to be stable homotopy equivalences. In addition, the lemma shows that the homotopies implicit in this all induce the identity map on the lattice of gauge-invariant ideals. \square

9. THE CIRCLE COMPONENTS

The homotopy equivalence in Theorem 8.7 has the extra property that it preserves the gauge-invariant ideals. A graph C^* -algebra may have gauge-invariant subquotients that are Morita equivalent to $C(\mathbb{T})$. In this case, there are many ideals that are not gauge-invariant. In this section, we show that we may arrange the homotopy equivalence to also preserve these ideals. As before, we let G be a row-finite graph. We are going to use some results of Hong and Szymański [23]. Therefore, we assume G to be countable from now on. This ensures that ideals in $C^*(G)$ are prime if and only if they are primitive, and this may go wrong in general (see [1, 2]). The following notation is introduced in [23] to describe the primitive ideal space of $C^*(G)$. Beware that [23] uses the opposite convention regarding the source and range maps on the graph.

Definition 9.1. Write $w \leq v$ for $w, v \in V$ if there is a path in G from w to v , and $w \sim v$ if $w \leq v$ and $v \leq w$.

The relation \leq is a preorder. So \sim is an equivalence relation.

Definition 9.2. A subset $M \subseteq V$ is called a *maximal tail* if

- (MT1) if $e \in E$ and $s(e) \in M$, then $r(e) \in M$;
- (MT2) if $v \in M \cap V_{\text{reg}}$, then there is $e \in E$ with $r(e) = v$ and $s(e) \in M$;
- (MT3) for all $v, w \in M$, there is $y \in M$ with $v \geq y$ and $w \geq y$.

If M is a maximal tail, let $\Omega(M) = V \setminus M$. Let $\mathcal{M}(G)$ be the set of all maximal tails in G . Let $\mathcal{M}_\gamma(G) \subseteq \mathcal{M}(G)$ be the subset of all $M \in \mathcal{M}(G)$ with the extra property that each cycle in $G|_M$ has an entry in $G|_M$. Let $\mathcal{M}_\tau(G) := \mathcal{M}(G) \setminus \mathcal{M}_\gamma(G)$.

Condition (MT1) holds if and only if $v \geq w$ and $w \in M$ implies $v \in M$. Conditions (MT1) and (MT2) say exactly that $\Omega(M)$ is hereditary and saturated. It is shown in [23] that the gauge-invariant ideal corresponding to a hereditary saturated subset $W \in \mathbb{I}(G)$ is prime in the complete lattice $\mathbb{I}^\mathbb{T}(\mathbb{C}^*(G))$ of gauge-invariant ideals if and only if $V \setminus W$ satisfies (MT3) as well as (MT1) and (MT2). Equivalently, the map $M \mapsto \Omega(M)$ is a bijection between $\mathcal{M}(G)$ and the set of prime ideals in the lattice $\mathbb{I}(G)$.

Proposition 9.3 ([23]). *Let $M \in \mathcal{M}_\tau(G)$. Then M contains a unique simple cycle, and this cycle has no entry in M . Let $v \in M$ be a vertex on this cycle. Then $\mathbb{C}^*(G)_{\Omega(M)} \triangleleft \mathbb{C}^*(G)$ is a prime ideal in $\mathbb{C}^*(G)$ and the subquotient*

$$\mathbb{C}^*(G)|_M := \mathbb{C}^*(G)_{\langle \Omega(M) \cup \{v\} \rangle} / \mathbb{C}^*(G)_{\Omega(M)}$$

is gauge-simple and Morita–Rieffel equivalent to $\mathbb{C}(\mathbb{T})$. More precisely, the image of the corner $p_v \mathbb{C}^(G) p_v$ in $\mathbb{C}^*(G) / \mathbb{C}^*(G)_{\Omega(M)}$ is isomorphic to $\mathbb{C}(\mathbb{T})$.*

The corner $p_v(\mathbb{C}^*(G) / \mathbb{C}^*(G)_{\Omega(M)}) p_v$ is Morita–Rieffel equivalent to the ideal in $\mathbb{C}^*(G) / \mathbb{C}^*(G)_{\Omega(M)}$ generated by the image of p_v . This ideal is gauge-invariant and corresponds to the smallest hereditary and saturated subset of V containing $\Omega(M)$ and v . Then it must be $\mathbb{C}^*(G)|_M$. We call $\mathbb{C}^*(G)|_M$ *the gauge-simple subquotient of $\mathbb{C}^*(G)$ associated to $M \in \mathcal{M}_\tau(G)$* .

Now we return to the situation of Theorem 8.7. That is, G and G' are countable graphs, $A = \mathbb{C}^*(G)$, $B = \mathbb{C}^*(G')$, and there are an order isomorphism $\psi: \mathbb{I}_p(G) \cong \mathbb{I}_p(A) \xrightarrow{\sim} \mathbb{I}_p(B) \cong \mathbb{I}_p(G')$ and diagram isomorphisms

$$\begin{aligned} \varphi_0^\dagger: \mathbb{K}_0^+(A, \mathbb{I}_p) &\xrightarrow{\sim} \psi^* \mathbb{K}_0^+(B, \mathbb{I}_p), \\ \varphi_1: \mathbb{K}_1(A, \mathbb{I}_p) &\xrightarrow{\sim} \psi^* \mathbb{K}_1(B, \mathbb{I}_p). \end{aligned}$$

These isomorphisms induce a monoid isomorphism $\varphi_0: \mathcal{P}(A) \xrightarrow{\sim} \mathcal{P}(B)$. We assume that the obstruction class of (φ_0, φ_1) vanishes.

Lemma 9.4. *The order isomorphism $\psi: \mathbb{I}_p(G) \xrightarrow{\sim} \mathbb{I}_p(G')$ extends uniquely to an order isomorphism $\bar{\psi}: \mathbb{I}(G) = \mathbb{I}_p^\infty(G) \xrightarrow{\sim} \mathbb{I}_p^\infty(G') = \mathbb{I}(G')$. The extension maps the subsets $\mathcal{M}_\gamma(G)$ and $\mathcal{M}_\tau(G)$ bijectively onto $\mathcal{M}_\gamma(G')$ and $\mathcal{M}_\tau(G')$, respectively.*

Proof. As in the proof of Proposition 7.3, the extension $\bar{\psi}$ exists by Lemma 7.2. The order isomorphism $\bar{\psi}$ restricts to a bijection between the prime ideals. Thus it induces a bijection $\psi: \mathcal{M}(G) \xrightarrow{\sim} \mathcal{M}(G')$. We must show that $H \in \mathcal{M}(G)$ belongs to $\mathcal{M}_\tau(G)$ if and only if $\psi(H) \in \mathcal{M}(G')$ belongs to $\mathcal{M}_\tau(G')$. To do this, we characterise through the diagrams $\mathbb{K}_0^+(\mathbb{C}^*(G), \mathbb{I}_p)$ and $\mathbb{K}_1(\mathbb{C}^*(G), \mathbb{I}_p)$ when $H \in \mathcal{M}(G)$ belongs to $\mathcal{M}_\tau(G)$. First, Remark 4.17 shows that φ_0^\dagger and φ_1 extend to isomorphisms of diagrams $\mathbb{K}_0^+(\mathbb{C}^*(G), \mathbb{I}_p^\infty) \xrightarrow{\sim} \mathbb{K}_0^+(\mathbb{C}^*(G'), \mathbb{I}_p^\infty)$ and $\mathbb{K}_1(\mathbb{C}^*(G), \mathbb{I}_p^\infty) \xrightarrow{\sim} \mathbb{K}_1(\mathbb{C}^*(G'), \mathbb{I}_p^\infty)$. Theorem 5.3 identifies $\mathbb{I}_p^\infty(\mathbb{C}^*(G))$ with $\mathbb{I}(G)$, the lattice of hereditary saturated subsets.

Let $M \in \mathcal{M}(G)$. Then $H := \Omega(M) = V \setminus M \in \mathbb{I}(G)$. If $M \in \mathcal{M}_\tau(G)$, then we define $H_2 := \langle \Omega(M) \cup \{v\} \rangle \in \mathbb{I}(G)$ as in Proposition 9.3. Then $H \subsetneq H_2$. If $K \in \mathbb{I}(G)$ also satisfies $H \subsetneq K$, then $K \cap H_2 = H$ or $K \cap H_2 = H_2$ because $K \cap H_2 \in \mathbb{I}(G)$ and the subquotient $\mathbb{C}^*(G)|_M := \mathbb{C}^*(G)_{H_2} / \mathbb{C}^*(G)_H$ has no gauge-invariant ideals. The case $K \cap H_2 = H$ is impossible because H is prime. It follows that $H_2 \in \mathbb{I}(G)$ is the unique minimal element of the set $\{K \in \mathbb{I}(G) : H \subsetneq K\}$. Therefore, the existence of such a unique minimal element H_2 is necessary for $M \in \mathcal{M}_\tau(G)$. This property is

preserved by order isomorphisms. We assume from now on that such an H_2 exists and, in this generality, we define

$$C^*(G)|_M := C^*(G)_{H_2}/C^*(G)_H.$$

A quotient of a graph C^* -algebra by a gauge-invariant ideal is again a graph C^* -algebra (see [7, Theorem 4.1]). Any gauge-invariant ideal in a graph C^* -algebra is again isomorphic to a graph C^* -algebra. This is first claimed in [13, Lemma 1.6] and correctly proven in [37]. The construction in [13] is correct for row-finite graphs, which is the case we need. It is easy to see that the isomorphism in [13] is also \mathbb{T} -equivariant. Therefore, $C^*(G)|_M$ is \mathbb{T} -equivariantly isomorphic to the graph C^* -algebra of some graph G_M . This subquotient of $C^*(G)$ has no gauge-invariant ideals because H_2 is the unique minimal element of $\{K \in \mathbb{I}(G) : H \subsetneq K\}$. Equivalently, the graph G_M is cofinal. Now there are three possible cases:

- (1) G_M has no cycles; then $C^*(G_M)$ is a simple AF-algebra;
- (2) G_M has more than one simple cycle; then $C^*(G_M)$ is simple purely infinite;
- (3) G_M contains exactly one simple cycle.

In the third case, the unique simple cycle cannot have an entry because then $C^*(G_M)$ would be simple by [6, Proposition 5.1], and then we must be in one of the first two cases, by the well known dichotomy for simple graph C^* -algebras. So $M \in \mathcal{M}_\tau(G)$ in the third case, that is, $C^*(G_M)$ is Morita–Rieffel equivalent to $C(\mathbb{T})$.

The description of the graph for a quotient in [7, Theorem 4.1] shows that any vertex in G_M is also a vertex in H_2 . Therefore, Theorem 3.1 implies that any projection in $C^*(G_M)$ lifts to a projection in $C^*(G)_{H_2}$. That is, the map $K_0^+(C^*(G)_{H_2}) \rightarrow K_0^+(C^*(G_M))$ is surjective. This remains so on the Grothendieck groups. Therefore, the K-theory long exact sequence for the C^* -algebra extension $C^*(G)_H \twoheadrightarrow C^*(G)_{H_2} \twoheadrightarrow C^*(G)|_M$ looks as follows:

$$(9.5) \quad \begin{array}{ccccc} K_0(C^*(G)_H) & \xrightarrow{i_0} & K_0(C^*(G)_{H_2}) & \twoheadrightarrow & K_0(C^*(G)|_M) \\ \uparrow & & & & \downarrow 0 \\ K_1(C^*(G)|_M) & \xleftarrow{i_1} & K_1(C^*(G)_{H_2}) & \xleftarrow{\quad} & K_1(C^*(G)_H). \end{array}$$

If $M \in \mathcal{M}_\tau(G)$, then $C^*(G)|_M$ is Morita–Rieffel equivalent to $C(\mathbb{T})$ and therefore $K_0^+(C^*(G)|_M) \cong \mathbb{N}$ and $K_1(C^*(G)|_M) \cong \mathbb{Z}$. In contrast, if $M \notin \mathcal{M}_\tau(G)$, then either $C^*(G)|_M$ is purely infinite and $K_0^+(C^*(G)|_M)$ is a group, or $C^*(G)|_M$ is AF and $K_1(C^*(G)|_M) \cong 0$. Now we check that these three cases can be distinguished looking only at the maps

$$K_0^+(C^*(G)_H) \rightarrow K_0^+(C^*(G)_{H_2}), \quad K_i(C^*(G)_H) \rightarrow K_i(C^*(G)_{H_2})$$

for $i = 0, 1$ induced by the inclusion $H \subsetneq H_2$. Indeed, if $M \in \mathcal{M}_\tau(G)$, then the quotient of $K_0^+(C^*(G)_{H_2})$ by the image of $K_0^+(C^*(G)_H)$ cannot be a group because it maps to the monoid $K_0^+(C^*(G)|_M) \cong \mathbb{N}$, which is not a group, and it cannot happen that the map $K_0(C^*(G)_H) \rightarrow K_0(C^*(G)_{H_2})$ is injective and $K_1(C^*(G)_H) \rightarrow K_1(C^*(G)_{H_2})$ is surjective, because that would force $K_1(C^*(G)|_M)$ to vanish. If $M \notin \mathcal{M}_\tau(G)$ and $C^*(G)|_M$ is purely infinite simple, then the quotient of $K_0^+(C^*(G)_{H_2})$ by the image of $K_0^+(C^*(G)_H)$ is a group. And if $M \notin \mathcal{M}_\tau(G)$ and $C^*(G)|_M$ is AF, then the map $K_0(C^*(G)_H) \rightarrow K_0(C^*(G)_{H_2})$ is injective and $K_1(C^*(G)_H) \rightarrow K_1(C^*(G)_{H_2})$ is surjective. Thus we can read off from $\mathbb{I}(G)$ and the diagrams $K_0^+(C^*(G), \mathbb{I})$ and $K_1(C^*(G), \mathbb{I})$ whether or not $M \notin \mathcal{M}_\tau(G)$. Since these invariants do not distinguish our two graphs, it follows that the bijection $\psi: \mathcal{M}(G) \xrightarrow{\sim} \mathcal{M}(G')$ maps $\mathcal{M}_\tau(G)$ onto $\mathcal{M}_\tau(G')$. \square

Theorem 9.6. *Let G and G' be two countable graphs. Let*

$$\psi: \mathbb{I}_p(G) \cong \mathbb{I}_p(A) \xrightarrow{\sim} \mathbb{I}_p(B) \cong \mathbb{I}_p(G')$$

be an order isomorphism and let

$$\begin{aligned} \varphi_0^+ &: K_0^+(A, \mathbb{I}_p) \xrightarrow{\sim} \psi^* K_0^+(B, \mathbb{I}_p), \\ \varphi_1 &: K_1(A, \mathbb{I}_p) \xrightarrow{\sim} \psi^* K_1(B, \mathbb{I}_p) \end{aligned}$$

be diagram isomorphisms. These isomorphisms induce a monoid isomorphism $\varphi_0: \mathcal{P}(A) \xrightarrow{\sim} \mathcal{P}(B)$. Assume that the obstruction class of (φ_0, φ_1) vanishes. Then there is an A - B -correspondence \mathcal{H}_{AB} that induces the isomorphisms φ_0^+ and φ_1 and that induces a Morita–Rieffel equivalence between the gauge-simple subquotients of A and B associated to M and $\psi(M)$ for all $M \in \mathcal{M}_\tau(G)$. In addition, there is a B - A -correspondence \mathcal{H}_{BA} that is inverse to \mathcal{H}_{AB} up to homotopy, and such that all the correspondences in the homotopies between the composite correspondences $\mathcal{H}_{AB} \otimes_B \mathcal{H}_{BA}$ and the identity induce the identity correspondences on the gauge-simple subquotients associated to all $M \in \mathcal{M}_\tau(G)$, and similarly for $\mathcal{H}_{BA} \otimes_A \mathcal{H}_{AB}$.

Proof. First, we fix $M \in \mathcal{M}_\tau(G)$. Theorem 8.7 shows that there is a $C^*(G)$ - $C^*(G')$ -correspondence \mathcal{H}_{AB} that induces the maps ψ , φ_0^+ and φ_1 on our invariants. In particular, this correspondence induces correspondences between the gauge-invariant ideals $C^*(G)_H$ and $C^*(G')_{\psi(H)}$ for all $H \in \mathbb{I}_p(G)$; this extends automatically to $H \in \mathbb{I}(G)$. And then it also induces a correspondence $\mathcal{H}_{AB,M}$ between the subquotients $C^*(G)|_M$ and $C^*(G')|_{\psi(M)}$. Since our correspondence is a homotopy equivalence that preserves the gauge-invariant ideals, it induces an isomorphism on $K_*(C^*(G)_H)$ and $K_*(C^*(G')_{\psi(H)})$ in (9.5). By the Five Lemma, it induces an isomorphism $K_*(C^*(G)|_M) \xrightarrow{\sim} K_*(C^*(G')|_{\psi(M)})$ as well. This isomorphism on $K_0(C^*(G)|_M) \cong \mathbb{Z}$ maps the positive cone \mathbb{N} into itself. Thus it is the identity map.

Now we again describe $C^*(G)|_M$ as the graph C^* -algebra of another graph G_M , which has the vertex set $H_2 \setminus H$. The $C^*(G)|_M$ - $C^*(G')|_{\psi(M)}$ -correspondence induced by \mathcal{H}_{AB} is described through proper Hilbert $C^*(G')|_{\psi(M)}$ -modules $\mathcal{H}_{AB,v}$ for $v \in H_2 \setminus H$ and unitaries U_v for $v \in H_2 \setminus H$ that are regular in the graph G_M . Our correspondence induces an isomorphism $\mathcal{P}(C^*(G)|_M) \xrightarrow{\sim} \mathcal{P}(C^*(G')|_{\psi(M)})$ and an isomorphism $K_1(C^*(G)|_M) \xrightarrow{\sim} K_1(C^*(G')|_{\psi(M)})$. Since $C^*(G)|_M$ and $C^*(G')|_{\psi(M)}$ are Morita–Rieffel equivalent to $C(\mathbb{T})$, we know that $\mathcal{P}(C^*(G)|_M) \cong \mathbb{N}$ and $\mathcal{P}(C^*(G')|_{\psi(M)}) \cong \mathbb{N}$. Let v and v' be vertices that belong to the unique simple cycles in M and $\psi(M)$, respectively. By Proposition 9.3, the projections p_v and $p_{v'}$ generate $\mathcal{P}(C^*(G)|_M)$ and $\mathcal{P}(C^*(G')|_{\psi(M)})$, respectively, and the corners $p_v C^*(G)|_M p_v$ and $p_{v'} C^*(G')|_{\psi(M)} p_{v'}$ are both isomorphic to $C(\mathbb{T})$. The image $(\varphi_0^+)_M[p_v]$ in $\mathcal{P}(C^*(G')|_{\psi(M)})$ is the class of the proper Hilbert $C^*(G')|_{\psi(M)}$ -module $\mathcal{H}_{AB,v}$. Since this class is another generator of $\mathcal{P}(C^*(G')|_{\psi(M)}) \cong \mathbb{N}$, it follows that $\mathcal{H}_{AB,v}$ is isomorphic to $p_{v'} C^*(G')|_{\psi(M)}$. We fix such an isomorphism to identify the C^* -algebra $\mathbb{K}(\mathcal{H}_{AB,v})$ with $p_{v'} C^*(G')|_{\psi(M)} p_{v'}$ and thus with $C(\mathbb{T})$.

Our correspondence contains a unital $*$ -homomorphism

$$p_v C^*(G)|_M p_v \rightarrow p_{v'} C^*(G')|_{\psi(M)} p_{v'}.$$

We claim that we may change the $C^*(G)$ - $C^*(G')$ -correspondence \mathcal{H}_{AB} so that this $*$ -homomorphism becomes an isomorphism, while keeping the other desirable properties of \mathcal{H}_{AB} already achieved. We already know that our $*$ -homomorphism induces an isomorphism

$$K_1(p_v C^*(G)|_M p_v) \xrightarrow{\sim} K_1(p_{v'} C^*(G')|_{\psi(M)} p_{v'}).$$

In other words, the generator $z \in C(\mathbb{T})$ of $K_1(p_v C^*(G)|_M p_v) \cong K_1(C(\mathbb{T})) \cong \mathbb{Z}$ is mapped to a generator of $K_1(p_{v'} C^*(G')|_{\psi(M)} p_{v'}) \cong C(\mathbb{T}) \cong \mathbb{Z}$. This means that we

get a unitary in $C(\mathbb{T})$ of winding number ± 1 . This is a product of $z^{\pm 1}$ by a unitary $\zeta \in C(\mathbb{T})$ that is homotopic to 1. Then ζ lifts to a unitary $\hat{\zeta} \in p_{v'}C^*(G')p_{v'}$ that is still homotopic to the identity. We may multiply the unitary U_v in our description of \mathcal{H}_{AB} by $\hat{\zeta}^*$ without changing its homotopy class or the induced maps on K_0^+ and K_1 because $\hat{\zeta}$ is homotopic to the identity. The element $z \in C(\mathbb{T}) \subseteq C^*(G|_M)$ is the product of the edges in the cycle in M based at z . So it contains the unitary U_v exactly once. Therefore, changing U_v in this way ensures that the new $C^*(G)|_M$ - $C^*(G')|_{\psi(M)}$ -correspondence now maps the generator $z \in C(\mathbb{T})$ to $z^{\pm 1}$. Thus it induces an isomorphism

$$(9.7) \quad C(\mathbb{T}) \cong p_v C^*(G)|_M p_v \xrightarrow{\sim} p_{v'} C^*(G')|_{\psi(M)} p_{v'} \cong C(\mathbb{T}).$$

The change that was needed to achieve this for a fixed $M \in \mathcal{M}_\tau(G)$ only occurred at a single vertex v , which is supported in a cycle in the difference set $H_2 \setminus H$ corresponding to M . Different circle components in the primitive ideal space involve disjoint cycles and therefore do not interfere with each other. Therefore, we may arrange the induced maps $p_v C^*(G)|_M p_v \xrightarrow{\sim} p_{v'} C^*(G')|_{\psi(M)} p_{v'}$ to be isomorphisms for all $M \in \mathcal{M}_\tau(G)$. Equivalently, the induced $C^*(G)|_M$ - $C^*(G')|_{\psi(M)}$ -correspondences become Morita–Rieffel equivalences simultaneously for $M \in \mathcal{M}_\tau(G)$.

Lemma 8.6 gives a B - A -correspondence \mathcal{H}_{BA} that is inverse to \mathcal{H}_{AB} up to homotopy. Even more, the homotopies that are implicit here automatically preserve all gauge-invariant ideals because equivalent projections generate the same ideal. Therefore, for each $M \in \mathcal{M}_\tau(G)$, the induced correspondences $\mathcal{H}_{AB,M}$ and $\mathcal{H}_{BA,M}$ between $C^*(G)|_M$ and $C^*(G')|_{\psi(M)}$ are also inverse to each other up to homotopy. Now we change the correspondence \mathcal{H}_{BA} in the same way as we changed \mathcal{H}_{AB} , to a homotopic correspondence so that the induced map $C(\mathbb{T}) \cong p_{v'} C^*(G')|_{\psi(M)} p_{v'} \xrightarrow{\sim} p_v C^*(G)|_M p_v \cong C(\mathbb{T})$ maps the unitary generator z to $z^{\pm 1}$. Then the composite of this *-isomorphism with the one in (9.7) also sends z to $z^{\pm 1}$. Since the composite induces the identity map on $K_1(C(\mathbb{T}))$, it must be the identity map on $C(\mathbb{T})$. We have arranged the composite correspondence to be homotopic to the identity correspondence on $C^*(G)$. This homotopy induces a homotopy of *-endomorphisms of $C(\mathbb{T})$ that connects the identity with itself. It maps the generator z to $\zeta_t \cdot z$ for a continuous path of unitaries $\zeta_t \in C(\mathbb{T})$ with $\zeta_0 = \zeta_1 = 1$. Now we may lift these unitaries and correct the $C^*(G)$ - $C^*(G)$ -correspondences $(\mathcal{H}_t)_{t \in [0,1]}$ in the homotopy so that \mathcal{H}_0 and \mathcal{H}_1 remain the same and each \mathcal{H}_t induces the identity map on $p_v C^*(G)|_M p_v$ for all $M \in \mathcal{M}_\tau(G)$ and $v \in M$ in the unique cycle in M . Thus the composite correspondence $\mathcal{H}_{AB} \otimes_{C^*(G')} \mathcal{H}_{BA}$ is homotopic to the identity correspondence through an ideal-preserving homotopy. The same may be done for the homotopy that joins the other composite $\mathcal{H}_{BA} \otimes_{C^*(G)} \mathcal{H}_{AB}$ to the identity correspondence on $C^*(G')$. \square

REFERENCES

- [1] Gene Abrams and Mark Tomforde, *A class of C^* -algebras that are prime but not primitive*, Münster J. Math. **7** (2014), no. 2, 489–514, doi: 10.17879/58269761901. MR **3426227**
- [2] Gene D. Abrams and Mark Tomforde, *The prime spectrum and primitive ideal space of a graph C^* -algebra*, Internat. J. Math. **25** (2014), no. 7, 1450070, 22, doi: 10.1142/S0129167X14500700. MR **3238092**
- [3] Pere Ara, M. A. Moreno, and Enrique Pardo, *Nonstable K -theory for graph algebras*, Algebr. Represent. Theory **10** (2007), no. 2, 157–178, doi: 10.1007/s10468-006-9044-z. MR **2310414**
- [4] Sara E. Arklint, Rasmus Bentmann, and Takeshi Katsura, *Reduction of filtered K -theory and a characterization of Cuntz–Krieger algebras*, J. K-Theory **14** (2014), no. 3, 570–613, doi: 10.1017/is014009013jkt281. MR **3349327**
- [5] Tathagata Banerjee and Ralf Meyer, *Noncommutative coarse geometry*, J. Noncommut. Geom. **13** (2019), no. 3, 1117–1149, doi: 10.4171/JNCG/346. MR **4030362**

- [6] Teresa Bates, Jeong H. Hong, Iain Raeburn, and Wojciech Szymański, *The ideal structure of the C^* -algebras of infinite graphs*, Illinois J. Math. **46** (2002), no. 4, 1159–1176, available at <http://projecteuclid.org/euclid.ijm/1258138472>. MR **1988256**
- [7] Teresa Bates, David Pask, Iain Raeburn, and Wojciech Szymański, *The C^* -algebras of row-finite graphs*, New York J. Math. **6** (2000), 307–324, available at <http://nyjm.albany.edu/j/2000/6-14.html>. MR **1777234**
- [8] Rasmus Bentmann, *Kirchberg X -algebras with real rank zero and intermediate cancellation*, J. Noncommut. Geom. **8** (2014), no. 4, 1061–1081, doi: 10.4171/JNCG/178. MR **3310939**
- [9] Rasmus Bentmann and Ralf Meyer, *A more general method to classify up to equivariant KK -equivalence*, Doc. Math. **22** (2017), 423–454, available at <https://www.math.uni-bielefeld.de/documenta/vol-22/14.html>. MR **3628788**
- [10] Lawrence G. Brown and Gert K. Pedersen, *C^* -algebras of real rank zero*, J. Funct. Anal. **99** (1991), no. 1, 131–149, doi: 10.1016/0022-1236(91)90056-B. MR **1120918**
- [11] ———, *Non-stable K -theory and extremally rich C^* -algebras*, J. Funct. Anal. **267** (2014), no. 1, 262–298, doi: 10.1016/j.jfa.2014.04.003. MR **3206515**
- [12] Marius Dădărlat and Ralf Meyer, *E -Theory for C^* -algebras over topological spaces*, J. Funct. Anal. **263** (2012), no. 1, 216–247, doi: 10.1016/j.jfa.2012.03.022. MR **2920847**
- [13] Klaus Deicke, Jeong Hee Hong, and Wojciech Szymański, *Stable rank of graph algebras. Type I graph algebras and their limits*, Indiana Univ. Math. J. **52** (2003), no. 4, 963–979, doi: 10.1512/iumj.2003.52.2350. MR **2001940**
- [14] Jacques Dixmier, *Les C^* -algèbres et leurs représentations*, Deuxième édition. Cahiers Scientifiques, Fasc. XXIX, Gauthier-Villars Éditeur, Paris, 1969. MR **0246136**
- [15] Søren Eilers, Gunnar Restorff, and Efren Ruiz, *Classification of extensions of classifiable C^* -algebras*, Adv. Math. **222** (2009), no. 6, 2153–2172, doi: 10.1016/j.aim.2009.07.014. MR **2562779**
- [16] ———, *On graph C^* -algebras with a linear ideal lattice*, Bull. Malays. Math. Sci. Soc. (2) **33** (2010), no. 2, 233–241, available at http://math.usm.my/bulletin/html/vol133_2_7.html. MR **2666426**
- [17] ———, *Classifying C^* -algebras with both finite and infinite subquotients*, J. Funct. Anal. **265** (2013), no. 3, 449–468, doi: 10.1016/j.jfa.2013.05.006. MR **3056712**
- [18] ———, *The ordered K -theory of a full extension*, Canad. J. Math. **66** (2014), no. 3, 596–625, doi: 10.4153/CJM-2013-015-7. MR **3194162**
- [19] ———, *Strong classification of extensions of classifiable C^* -algebras*, Bull. Korean Math. Soc. **59** (2022), no. 3, 567–608, doi: 10.4134/BKMS.b210047. MR **4432429**
- [20] Søren Eilers, Gunnar Restorff, Efren Ruiz, and Adam P. W. Sørensen, *The complete classification of unital graph C^* -algebras: geometric and strong*, Duke Math. J. **170** (2021), no. 11, 2421–2517, doi: 10.1215/00127094-2021-0060. MR **4302548**
- [21] Søren Eilers and Mark Tomforde, *On the classification of nonsimple graph C^* -algebras*, Math. Ann. **346** (2010), no. 2, 393–418, doi: 10.1007/s00208-009-0403-z. MR **2563693**
- [22] James Gabe, *Lifting theorems for completely positive maps*, J. Noncommut. Geom. **16** (2022), no. 2, 391–421, doi: 10.4171/jncg/479. MR **4478259**
- [23] Jeong Hee Hong and Wojciech Szymański, *The primitive ideal space of the C^* -algebras of infinite graphs*, J. Math. Soc. Japan **56** (2004), no. 1, 45–64, doi: 10.2969/jmsj/1191418695. MR **2023453**
- [24] Eberhard Kirchberg, *Das nicht-kommutative Michael-Auswahlprinzip und die Klassifikation nicht-einfacher Algebren*, C^* -Algebras (Münster, 1999), Springer, Berlin, 2000, pp. 92–141, doi: 10.1007/978-3-642-57288-3. MR **1796912**
- [25] Ralf Meyer, *Homological algebra in bivariant K -theory and other triangulated categories. II*, Tbil. Math. J. **1** (2008), 165–210, available at http://www.tcms.org.ge/Journals/TMJ/Volume1/Xpapers/tmj1_9.pdf. MR **2563811**
- [26] Ralf Meyer and Ryszard Nest, *C^* -Algebras over topological spaces: the bootstrap class*, Münster J. Math. **2** (2009), 215–252, available at <http://nbn-resolving.de/urn:nbn:de:hbz:6-10569452982>. MR **2545613**
- [27] ———, *Homological algebra in bivariant K -theory and other triangulated categories. I*, Triangulated categories (Thorsten Holm, Peter Jørgensen, and Raphaël Rouquier, eds.), London Math. Soc. Lecture Note Ser., vol. 375, Cambridge Univ. Press, Cambridge, 2010, pp. 236–289, doi: 10.1017/CBO9781139107075.006. MR **2681710**
- [28] ———, *C^* -Algebras over topological spaces: filtrated K -theory*, Canad. J. Math. **64** (2012), 368–408, doi: 10.4153/CJM-2011-061-x. MR **2953205**
- [29] James A. Mingo and William J. Phillips, *Equivariant triviality theorems for Hilbert C^* -modules*, Proc. Amer. Math. Soc. **91** (1984), no. 2, 225–230, available at <http://www.jstor.org/stable/2044632>. MR **740176**

- [30] Paul S. Muhly and Mark Tomforde, *Adding tails to C^* -correspondences*, Doc. Math. **9** (2004), 79–106, available at <http://www.math.uni-bielefeld.de/documenta/vol1-09/05.html>. MR **2054981**
- [31] Cornel Pasnicu and Mikael Rørdam, *Purely infinite C^* -algebras of real rank zero*, J. Reine Angew. Math. **613** (2007), 51–73, doi: 10.1515/CRELLE.2007.091. MR **2377129**
- [32] Iain Raeburn, *Graph algebras*, CBMS Regional Conference Series in Mathematics, vol. 103, Amer. Math. Soc., Providence, RI, 2005. MR **2135030**
- [33] Iain Raeburn and Wojciech Szymański, *Cuntz–Krieger algebras of infinite graphs and matrices*, Trans. Amer. Math. Soc. **356** (2004), 39–59, doi: 10.1090/S0002-9947-03-03341-5MR **2020023**
- [34] Gunnar Restorff, *Classification of Cuntz–Krieger algebras up to stable isomorphism*, J. Reine Angew. Math. **598** (2006), 185–210, doi: 10.1515/CRELLE.2006.074. MR **2270572**
- [35] Mikael Rørdam, *Classification of Cuntz–Krieger algebras*, *K-Theory* **9** (1995), no. 1, 31–58, doi: 10.1007/BF00965458. MR **1340839**
- [36] ———, *Classification of nuclear, simple C^* -algebras*, Classification of nuclear C^* -algebras. Entropy in operator algebras, Encyclopaedia of Mathematical Sciences, vol. 126, Springer, Berlin, 2002, pp. 1–145, doi: 10.1007/978-3-662-04825-2_1. MR **1878881**
- [37] Efren Ruiz and Mark Tomforde, *Ideals in graph algebras*, *Algebr. Represent. Theory* **17** (2014), no. 3, 849–861, doi: 10.1007/s10468-013-9421-3. MR **3254772**
- [38] Niels Erik Wegge-Olsen, *K-Theory and C^* -algebras*, Oxford Science Publications, The Clarendon Press Oxford University Press, New York, 1993. MR **1222415**

Email address: rasmusbentmann@gmail.com

Email address: rmeyer2@uni-goettingen.de

MATHEMATISCHES INSTITUT, GEORG-AUGUST UNIVERSITÄT GÖTTINGEN, BUNSENSTRASSE 3–5,
37073 GÖTTINGEN, GERMANY