

GLOBAL WELL-POSEDNESS OF THE NONLINEAR HARTREE EQUATION FOR INFINITELY MANY PARTICLES WITH SINGULAR INTERACTION

SONAE HADAMA AND YOUNGHUN HONG

ABSTRACT. The nonlinear Hartree equation (NLH) in the Heisenberg picture admits steady states of the form $\gamma_f = f(-\Delta)$ representing quantum states of infinitely many particles. In this article, we consider the time evolution of perturbations from a large class of such steady states via the three-dimensional NLH. We prove that if the interaction potential w has finite measure and initial states have finite relative entropy, then solutions preserve the relative free energy, and they exist globally in time. This result extends the important work of Lewin and Sabin [31] to singular interaction cases.

1. INTRODUCTION

1.1. Nonlinear Hartree equation for infinitely many quantum particles. The nonlinear Hartree equation (NLH) is a fundamental equation describing the mean-field dynamics of a large number of quantum particles. In the Heisenberg picture, by a non-negative self-adjoint operator γ on $L^2(\mathbb{R}^3)$, it is given by

$$i\partial_t\gamma = [-\Delta + w * \rho_\gamma, \gamma], \quad (1.1)$$

where $[A, B] = AB - BA$ is the Lie bracket. Here, w is a pair interaction potential on \mathbb{R}^3 , and $\rho_\gamma(x) = \gamma(x, x)$ is the total density of a quantum observable γ , where $\gamma(x, x')$ denotes the integral kernel of the operator γ , i.e.,

$$(\gamma\phi)(x) = \int_{\mathbb{R}^d} \gamma(x, x')\phi(x')dx'.$$

The convolution $w * \rho_\gamma$ represents the self-generated potential which acts as a multiplication operator in (1.1). By the spectral representation $\gamma = \sum_{j=1}^{\infty} \lambda_j |\phi_j\rangle\langle\phi_j|$ with an orthonormal set $\{\phi_j\}_{j=1}^{\infty}$ in $L^2(\mathbb{R}^3)$, the NLH (1.1) is equivalent to the system of (possibly) infinitely many coupled equations for orthonormal functions,

$$i\partial_t\phi_j = -\Delta\phi_j + \left(w * \sum_{k=1}^{\infty} \lambda_k |\phi_k|^2 \right) \phi_j, \quad \forall j \in \mathbb{N}. \quad (1.2)$$

The NLH (1.1) can be used for both boson and fermion particles. When a bosonic system is cooled down to nearly zero temperature, almost all particles occupy the same quantum

Date: April 11, 2024.

state, forming a Bose-Einstein condensate. In this case, they can be described by the single wave equation

$$i\partial_t\phi = -\Delta\phi + (w * |\phi|^2)\phi.$$

For rigorous derivation including the delta interaction, we refer to [19–22]. However, when the temperature of the total system is not sufficiently low, the multi-particle model must be taken into account [1]. On the other hand, for fermions, the Hartree-Fock equation is first derived by the mean-field approximation with the additional exchange term and with the additional condition $0 \leq \gamma \leq 1$ (or $0 \leq \lambda_j \leq 1$), which comes from Pauli's exclusion principle [3, 4]. However, the exchange term is sometimes dropped because it is treated as a smaller order term in the semi-classical approximation. In this context, the NLH (1.1) is called the *reduced* Hartree-Fock equation. In the finite-trace case, or when the particle number is finite, the global well-posedness of the Hartree(-Fock) equation has established [9–11, 36]. Recently, the global well-posedness and the small data scattering of the more general Kohn-Sham equation is proved [33].

One of the advantages of considering the operator-valued equation (1.1) is that contrary to the multi-particle system (1.2), it admits steady states of infinitely many particles. Precisely, any Fourier multiplier operator $\gamma_f = f(-\Delta)$, with symbol $f(|\xi|^2) \in L^1(\mathbb{R}^3)$ such that $f \geq 0$ and $f \not\equiv 0$, solves the NLH (1.1), because it has constant density. Moreover, it has infinite particle number, i.e., $\text{Tr}(\gamma_f) = \int_{\mathbb{R}^d} \rho_{\gamma_f} dx = \infty$. Such steady states include the following physically important examples.

- $\mathbb{1}(-\Delta \leq \mu)$ with chemical potential $\mu > 0$ (Fermi gas at zero temperature).
- $\frac{1}{e^{(-\Delta - \mu)/T} + 1}$ with $\mu \in \mathbb{R}$ and $T > 0$ (Fermi gas at positive temperature).
- $\frac{1}{e^{(-\Delta - \mu)/T} - 1}$ with $\mu < 0$ and $T > 0$ (Bose gas at positive temperature).
- $\frac{1}{e^{(-\Delta - \mu)/T}}$ with $\mu \in \mathbb{R}$ and $T > 0$ (Boltzmann gas at positive temperature).

Having steady states of infinitely many particles, it is natural to investigate the dynamics of their (possibly large) perturbations, governed by the NLH (1.1). In the celebrated work of Lewin and Sabin [30, 31], the rigorous mathematical formulation was *first* introduced. Moreover, global well-posedness of large perturbations is proved in the zero/positive temperature [31], and the 2D small data scattering in the positive temperature case was established [30]. Since then, the global well-posedness is extended to the singular interaction case in the zero temperature case [12]. In addition, the small scattering is shown in higher dimensions with singular interactions [13, 26]. We also remark that an alternative formulation of (1.1) introducing the random field is proposed in de Suzzoni [35]. In this direction, small data scattering is given in [14–16, 32]. See also the recent work [18, 28] in different settings.

1.2. Statement of the main result. The main goal of this article is to extend the global well-posedness of the NLH (1.1) for infinite particles in the positive temperature case to a larger class of singular interaction potentials as well as to a larger class of reference states.

For the setup of the problem, following Lewin and Sabin [30, 31], we fix a reference state of the form

$$\gamma_f = f(-\Delta) \quad \text{with } f = (S')^{-1},$$

where S is admissible in the following sense.

Definition 1.1 (\mathcal{H} -admissible). *A function $S : [0, 1] \rightarrow \mathbb{R}^+$ is called \mathcal{H} -admissible if*

- S is continuous on $[0, 1]$, and is differentiable on $(0, 1)$;
- $-S'$ is operator monotone on $(0, 1)$, and not constant;
- $\lim_{x \rightarrow 0^+} S'(x) = +\infty$ and $\lim_{x \rightarrow 1^-} S'(x) \leq 0$;
- $\|(S')_+\|_{L^{\frac{3}{2}}(0,1)} < \infty$.

For such a reference state, the *relative entropy* is defined by

$$\mathcal{H}(\gamma|\gamma_f) = \mathcal{H}_S(\gamma|\gamma_f) := -\text{Tr}(S(\gamma) - S(\gamma_f) - S'(\gamma_f)(\gamma - \gamma_f)). \quad (1.3)$$

Note that by construction, $\mathcal{H}(\gamma|\gamma_f)$ is non-negative for any $0 \leq \gamma \leq 1$. For more details and its key properties of the relative entropy, we refer to [29, 30] and Section 3 of this paper. Let

$$\mathcal{K}_f := \left\{ \gamma : \text{self-adjoint on } L^2(\mathbb{R}^3), 0 \leq \gamma \leq 1 \text{ and } \mathcal{H}(\gamma|\gamma_f) < \infty \right\}. \quad (1.4)$$

denote the collection of operators having finite relative entropy, called the *free energy space*. Then, given a finite signed measure w on \mathbb{R}^3 , we define the *relative free energy* by

$$\mathcal{F}_f(\gamma|\gamma_f) := \mathcal{H}(\gamma|\gamma_f) + \frac{1}{2} \iint_{\mathbb{R}^3 \times \mathbb{R}^3} w(x-y) \rho_{\gamma-\gamma_f}(x) \rho_{\gamma-\gamma_f}(y) dx dy. \quad (1.5)$$

This quantity is well-defined on the free energy space \mathcal{K}_f (see Proposition 3.6).

Our main result asserts that under weaker assumptions on w and f , the NLH (1.1) preserves the relative free energy and is globally well-posed in the free energy space.

Theorem 1.2. *Suppose that $w \in \mathcal{M}(\mathbb{R}^3)$ is a finite measure on \mathbb{R}^d such that $\widehat{w} \geq 0$, $S : [0, 1] \rightarrow \mathbb{R}^+$ is \mathcal{H} -admissible, and that $f := (S')^{-1}$ satisfies*

$$\|\langle \xi \rangle^{\frac{3}{2}+\epsilon} f(|\xi|^2)\|_{L^2_\xi(\mathbb{R}^3)} < \infty$$

for small $\epsilon > 0$. Then, given initial data $\gamma_0 \in \mathcal{K}_f$, there exists a unique global solution $\gamma(t)$ to the NLH (1.1) with initial data γ_0 such that $\gamma(t) \in \mathcal{K}_f$ and $\mathcal{F}_f(\gamma(t)|\gamma_f) = \mathcal{F}_f(\gamma_0|\gamma_f)$ for all $t \in \mathbb{R}$.

Remark 1.3. (i) In 1D, the analogous global well-posedness (see Theorem A.1) can be proved by minor modifications of the proof in Lewin and Sabin [31]. One of the reasons is that the $L^2(\mathbb{R})$ -norm of the density $\rho_{\gamma-\gamma_f}$ of a perturbation can be controlled by some operator norm $\|\langle \nabla \rangle^{\frac{1+\epsilon}{4}} (\gamma - \gamma_f) \langle \nabla \rangle^{\frac{1+\epsilon}{4}}\|_{\mathfrak{S}^2}$, where $\|\cdot\|_{\mathfrak{S}^2}$ is the Schatten-2 norm, and we can establish a proper local well-posedness using the corresponding operator class. Then, combining with Klein's inequality (3.2), one can deduce the continuity of the initial data-to-solution map for the perturbation equation. It allows us to justify the conservation of the relative free

energy (see (1.5)) taking the limit of those for regular solutions.

(ii) On the other hand, in 2D and 3D cases, the density $\rho_{\gamma-\gamma_f}$ is bounded by a higher regularity norm $\|\langle \nabla \rangle^{\frac{d+\epsilon}{4}}(\gamma-\gamma_f)\langle \nabla \rangle^{\frac{d+\epsilon}{4}}\|_{\mathfrak{S}^2}$, while the relative entropy controls $\|(\gamma-\gamma_f)\langle \nabla \rangle\|_{\mathfrak{S}^2}$ by Klein's inequality. For this reason, in multi-dimensions, we are restricted to using the operator norm to develop a proper local well-posedness. Moreover, in the 3D case, one of our key estimates, that is, the Strichartz estimates for density functions (Lemma 2.2), also require higher regularity.

(iii) Intuitively, global well-posedness seems easier to show in 2D. However, we are not able to prove it, because endpoint failures in some inequalities arise in several places. Precisely, the Schatten-2 norm of $|\nabla|^{\frac{1}{2}}Q_0|\nabla|^{\frac{1}{2}}$ is used in (4.8), but then with these half-derivatives, Lemma 2.2 fails in 2D. We speculate that this is a technical issue, but we postpone work on the 2D problem in future work.

1.3. Ideas of Proof. The general outline of the proof follows from Lewin and Sabin [30]. Analyzing the equation

$$i\partial_t Q = [-\Delta + w * \rho_Q, Q + \gamma_f] \quad (1.6)$$

for the perturbation $Q(t) = \gamma(t) - \gamma_f$, we will show that solutions to the NLH (1.1), evolving from the free energy space, preserve the relative free energy, and that the conserved quantity prevents solutions from blowing up in finite time. However, several new ingredients are introduced to include singular pair interactions.

First of all, our proof is based on the key observation that the relative entropy has a better control in high frequencies. Indeed, it is already proved in Lewin and Sabin [30] that the relative entropy has a lower bound

$$\mathcal{H}(\gamma|\gamma_f) \geq \int_0^1 \text{Tr}(-\Delta - \mu)(\mathbb{1}(\gamma \geq \lambda) - \Pi_\mu^-)d\lambda, \quad (1.7)$$

where $\mu = S'(\lambda)$ and $\Pi_\mu^- = \mathbb{1}(-\Delta \leq \mu)$ (see [30, (86)]). In the right hand side lower bound, since $(-\Delta - \mu)$ is degenerate on the frequency sphere $|\xi| = \sqrt{\mu}$, it is natural to decompose the integral with respect to μ , and write

$$\mathcal{H}(\gamma, \gamma_f) \geq \sum_{\pm} \int_0^1 \text{Tr}(-\Delta - \mu)\Pi_{2\mu}^{\pm}(\mathbb{1}(\gamma \geq \lambda) - \Pi_\mu^-)\Pi_{2\mu}^{\pm}d\lambda, \quad (1.8)$$

where $\Pi_{2\mu}^- = \mathbb{1}(-\Delta \leq 2\mu)$ and $\Pi_{2\mu}^+ = \mathbb{1}(-\Delta > 2\mu)$. On the other hand, from the expression

$$\gamma - \gamma_f = \int_0^1 (\mathbb{1}(\gamma \geq \lambda) - \Pi_\mu^-)d\lambda \quad (1.9)$$

for the perturbation, obtained by functional calculus, the density also can be decomposed into

$$\rho_{\gamma-\gamma_f} = \sum_{\pm} \int_0^1 \rho_{\Pi_{2\mu}^{\pm}(\mathbb{1}(\gamma \geq \lambda) - \Pi_\mu^-)\Pi_{2\mu}^{\pm}}d\lambda + \sum_{\pm} \int_0^1 \rho_{\Pi_{2\mu}^{\pm}(\mathbb{1}(\gamma \geq \lambda) - \Pi_\mu^-)\Pi_{2\mu}^{\mp}}d\lambda.$$

We observe that the high-high frequency contribution of the density can be controlled by that of the lower bound in (1.8), because it includes a trace norm $\text{Tr}(-\Delta - \mu)\Pi_{2\mu}^+(\mathbb{1}(\gamma \geq$

$\lambda) - \Pi_{\mu}^{-})\Pi_{2\mu}^{+}$ but also $(-\Delta - \mu)$ is comparable with $-\Delta$. For the remaining part of the density, one can take an advantage of smoothness from the low frequency cut-off.

Secondly, we employ the equivalent density function formulation for perturbations, motivated by Lewin and Sabin [30] but modified adopting the idea in the first author's work [26]. In general, for global well-posedness, proper local well-posedness is required to construct solutions satisfying the conservation of the relative free energy. However, in our setting, an issue is that it seems that there is no Banach space of operators equipped with the norm controlled by the relative entropy. Indeed, by Klein's inequality $\mathcal{H}(\gamma, \gamma_f) \geq C_{\text{Klein}} \text{Tr}(1 - \Delta)(\gamma - \gamma_f)^2$ (see [30]), one may speculate that the operator space equipped with the norm $(\text{Tr}(1 - \Delta)(\gamma - \gamma_f)^2)^{1/2}$ would be a possible candidate, but it is not good enough when w is too singular. To overcome this problem, we reformulate the equation as follows. Let $S_V(t, t_0)$ denote the linear propagator for the linear Schrödinger equation from time t_0 with a time-dependent potential

$$i\partial_t u = -\Delta u + Vu, \quad (1.10)$$

where $V = V(t, x) : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$. In other words, $\phi(t) = S_V(t, t_0)\phi$ is the solution to the equation (1.10) with $\phi(t_0) = \phi$. Then, it is deduced from the NLH (1.6) for perturbations that

$$Q(t) = S_{w*\rho_Q}(t, 0)_* Q_0 - i \int_0^t S_{w*\rho_Q}(t, t_1)_* [w * \rho_Q(t_1), \gamma_f] dt_1, \quad (1.11)$$

where $\gamma_0 = Q_0 + \gamma_f$ and \mathcal{L}_* is the abbreviated notation for the operator

$$\mathcal{L}_* A = \mathcal{L} A \mathcal{L}^*.$$

Consequently, taking density of (1.11), we obtain the equation

$$\rho(t) = \rho_{S_{w*\rho}(t, 0)_* Q_0} - i \int_0^t \rho_{S_{w*\rho}(t, t_1)_* [w*\rho(t_1), \gamma_f]} dt_1. \quad (1.12)$$

Now, unlike (1.6), the equation (1.12) has an unknown function ρ only. Therefore, for local well-posedness, it suffices to choose a standard mixed norm function space where solutions live. The idea of using the density function formulation was introduced earlier via the wave operator in [29, 30], but (1.12) becomes relatively easier to deal with.

Finally, for the well-posedness of the density formulation (1.12), we prove and make use of Strichartz estimates for density functions of the linear perturbative Schrödinger flow from the free energy space \mathcal{K}_f . Indeed, by the perturbative nature of the contraction mapping argument, one may hope to show local well-posedness taking a function space, with possibly the strongest norm, in which the free evolution $\rho_{e^{it\Delta} Q_0 e^{-it\Delta}}$, with $\gamma_0 = Q_0 + \gamma_f \in \mathcal{K}_f$, lies. For this, we combine Strichartz estimates in [12] and the Lieb-Thirring inequality [30, Theorem 7], and also employ the relative entropy bound (1.8).

1.4. Outline of the paper. The rest of the paper is organized as follows. In Section 2, we provide preliminary definitions and basic Strichartz estimates for operators and density functions associated with the free Schrödinger flow. In Section 3, we review the relative entropy and the relative free energy from Lewin and Sabin [29, 30], and extend the Lieb-Thirring inequality (see Proposition 3.6). In Section 4, we develop the Strichartz estimates for density functions associated with the *perturbed* linear flow (Proposition 4.5), which is the key analysis tool in our paper. Then, in Section 5, using the estimates in the previous section, we establish the local well-posedness of the density formulation (1.12) for the NLH. Finally, in Section 6, making the conservation of the relative free energy rigorous, we establish global well-posedness. In addition, in Appendix A, we give a simpler proof of global well-posedness in the 1D case, and illustrate why the Strichartz estimates (Proposition 4.5) are introduced for the 3D problem.

1.5. Acknowledgement. Y. Hong was supported by National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT) (No. RS-2023-00208824 and No. RS-2023-00219980). S. Hadama was supported by JST, the establishment of university fellowships towards the creation of science technology innovation, Grant Number JPMJFS2123.

2. PRELIMINARIES

2.1. Operator spaces. Throughout this paper, for $1 \leq p \leq \infty$, the p -th Schatten class \mathfrak{S}^p is defined to be the completion of compact operators on $L^2(\mathbb{R}^3)$ equipped with the norm

$$\|\gamma\|_{\mathfrak{S}^p} := \begin{cases} (\mathrm{Tr}|\gamma|^p)^{\frac{1}{p}} & \text{if } 1 \leq p < \infty, \\ \|\gamma\|_{\mathcal{L}(L^2_x(\mathbb{R}^3))} & \text{if } p = \infty, \end{cases}$$

where $|\gamma| = \sqrt{\gamma\gamma^*}$, γ^* is the adjoint of γ and $\mathcal{L}(H)$ is the operator norm on the Hilbert space H . In particular, \mathfrak{S}^1 is the trace class, \mathfrak{S}^2 is the Hilbert-Schmidt class, and \mathfrak{S}^∞ is the Banach space of all compact operators.

In our analysis, operator norms based on the Schatten-2 norm, or the Hilbert-Schmidt norm, are particularly useful. Indeed, the Schatten-2 norm is friendlier to Fourier analysis, because it can be identified with the L^2 -norm for operator kernels as functions,

$$\|\gamma\|_{\mathfrak{S}^2}^2 = \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |\gamma(x, x')|^2 dx dx' = \|\gamma(x, x')\|_{L^2_{x, x'}}^2.$$

In this regard, for $\alpha \geq 0$, let \mathcal{H}^α be the Hilbert-Schmidt Sobolev space equipped with the norm

$$\begin{aligned} \|\gamma\|_{\mathcal{H}^\alpha} &:= \|\langle \nabla \rangle^\alpha \gamma \langle \nabla \rangle^\alpha\|_{\mathfrak{S}^2} = \left\{ \iint_{\mathbb{R}^3 \times \mathbb{R}^3} |\langle \nabla_x \rangle^\alpha \langle \nabla_{x'} \rangle^\alpha \gamma(x, x')|^2 dx dx' \right\}^{1/2} \\ &= \left\{ \frac{1}{(2\pi)^6} \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \langle \xi \rangle^{2\alpha} \langle \xi' \rangle^{2\alpha} |\hat{\gamma}(\xi, \xi')|^2 d\xi d\xi' \right\}^{1/2}. \end{aligned}$$

2.2. Strichartz estimates for the free flow. We recall the Strichartz estimates for the free flow [27],

$$\|e^{it\Delta}u_0\|_{L_t^q(\mathbb{R};L_x^r)} \lesssim \|u_0\|_{L_x^2} \quad (2.1)$$

provided that (q, r) is Strichartz-admissible, that is, (q, r) satisfying

$$\frac{2}{q} + \frac{3}{r} = \frac{3}{2} \quad \text{and} \quad 2 \leq q, r \leq \infty.$$

To capture dispersive properties at the operator level, we introduce the Strichartz norm for the operator kernel defined by

$$\|\gamma(t)\|_{\mathcal{S}(I)} := \sup_{(q,r): \text{admissible}} \|\gamma(t, x, x')\|_{L_t^q(I; L_x^r L_{x'}^2)} + \|\gamma(t, x, x')\|_{L_t^q(I; L_x^r L_{x'}^2)},$$

where $I \subset \mathbb{R}$. For $\alpha \geq 0$, we define the Sobolev-Strichartz norm by

$$\|\gamma(t)\|_{\mathcal{S}^\alpha(I)} := \|\langle \nabla \rangle^\alpha \gamma(t) \langle \nabla \rangle^\alpha\|_{\mathcal{S}(I)},$$

and let $\mathcal{S}^\alpha(I)$ denote the subspace of $L_t^\infty(I; \mathcal{H}^\alpha)$ satisfying $\|\gamma(t)\|_{\mathcal{S}^\alpha(I)} < \infty$. Then, the following Strichartz estimates hold (see [12] for instance).

Lemma 2.1 (Strichartz estimates for operator kernels). *There exists $c_0 > 0$ such that*

$$\begin{aligned} \|e^{it\Delta}\gamma_0 e^{-it\Delta}\|_{\mathcal{S}^\alpha(\mathbb{R})} &\leq c_0 \|\gamma_0\|_{\mathcal{H}^\alpha}, \\ \left\| \int_0^t e^{i(t-t_1)\Delta} \gamma(t_1) e^{-i(t-t_1)\Delta} dt_1 \right\|_{\mathcal{S}^\alpha(\mathbb{R})} &\leq c_0 \|\gamma\|_{L_t^{\tilde{q}}(\mathbb{R}; L_x^{\tilde{r}} L_{x'}^2)}, \end{aligned}$$

provided that (\tilde{q}, \tilde{r}) is Strichartz-admissible.

For densities, we will make use of the following Strichartz estimates.

Lemma 2.2 (Strichartz estimates for density functions). *For sufficiently small $\epsilon > 0$, there exists $c_0 = c_0(\epsilon) > 0$ such that*

$$\|\rho e^{it\Delta} \gamma_0 e^{-it\Delta}\|_{L_t^\infty(\mathbb{R}; L_x^2) \cap L_t^2(\mathbb{R}; \dot{H}_x^{1+\epsilon})} \leq c_0 \left\| |\nabla|^{\frac{1}{2}} \gamma_0 |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}}.$$

Remark 2.3. For density functions, much wider ranges of Strichartz estimates for general Schatten class initial data have been established in various settings since the important work of Frank-Lewin-Lieb-Seiringer [24] and Frank-Sabin [25] (see also [?, 5, 6]). However, in this article, we only employ the above Hilbert-Schmidt type version, because it is simpler and easier to apply together with Lemma 2.2.

Proof of Lemma 2.2. Note that the Fourier transform of ρ_γ is given by

$$\int_{\mathbb{R}^3} \left\{ \frac{1}{(2\pi)^6} \iint_{\mathbb{R}^3 \times \mathbb{R}^3} \widehat{\gamma}(\xi_1, \xi_2) e^{ix \cdot (\xi_1 + \xi_2)} d\xi_1 d\xi_2 \right\} e^{ix \cdot \xi} dx = \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} \widehat{\gamma}(\xi_1, \xi - \xi_1) d\xi_1.$$

Hence, by the Plancherel theorem and the Hölder inequality, we obtain that

$$\begin{aligned} \|\rho_\gamma\|_{L_x^2} &\lesssim \left\{ \sup_{\xi \in \mathbb{R}^3} \int_{\mathbb{R}^3} \frac{d\xi_1}{|\xi - \xi_1| \langle \xi - \xi_1 \rangle^{\frac{1+2\epsilon}{2}} |\xi_1| \langle \xi_1 \rangle^{\frac{1+2\epsilon}{2}}} \right\}^{\frac{1}{2}} \left\| |\nabla|^{\frac{1}{2}} \gamma |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}} \\ &\lesssim \left\| |\nabla|^{\frac{1}{2}} \gamma |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}}. \end{aligned}$$

Thus, by unitarity of $e^{it\Delta}$, it follows that

$$\|\rho_{e^{it\Delta}\gamma_0 e^{-it\Delta}}\|_{L_t^\infty(\mathbb{R}; L_x^2)} \lesssim \left\| |\nabla|^{\frac{1}{2}} \gamma_0 |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}}.$$

Similarly, we write the space-time Fourier transform of the density $\rho_{e^{it\Delta}\gamma_0 e^{-it\Delta}}$ as

$$\begin{aligned} &\int_{\mathbb{R}} \left\{ \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} e^{-it(|\xi_1|^2 - |\xi - \xi_1|^2)} \widehat{\gamma}_0(\xi_1, \xi - \xi_1) d\xi_1 \right\} e^{-it\tau} dt \\ &= \frac{1}{(2\pi)^2} \int_{\mathbb{R}^3} \widehat{\gamma}_0(\xi_1, \xi - \xi_1) \delta(\tau + |\xi_1|^2 - |\xi - \xi_1|^2) d\xi_1. \end{aligned}$$

Then, it follows that

$$\begin{aligned} \|\rho_{e^{it\Delta}\gamma_0 e^{-it\Delta}}\|_{L_t^2 \dot{H}_x^{1+\epsilon}} &\sim \left\| |\xi|^{1+\epsilon} \int_{\mathbb{R}^3} \widehat{\gamma}_0(\xi - \xi_1, \xi_1) \delta(\tau + |\xi_1|^2 - |\xi - \xi_1|^2) d\xi_1 \right\|_{L_\tau^2 L_\xi^2} \\ &\leq \left\{ \sup_{\xi \in \mathbb{R}^3} |\xi|^{2+2\epsilon} \int_{\mathbb{R}^3} \frac{\delta(\tau + |\xi_1|^2 - |\xi - \xi_1|^2)}{|\xi - \xi_1| \langle \xi - \xi_1 \rangle^{\frac{1+2\epsilon}{2}} |\xi_1| \langle \xi_1 \rangle^{\frac{1+2\epsilon}{2}}} d\xi_1 \right\}^{\frac{1}{2}} \\ &\quad \cdot \left\| |\nabla|^{\frac{1}{2}} \gamma_0 |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}}. \end{aligned}$$

By changing the variables,

$$\begin{aligned} &|\xi|^{2+2\epsilon} \int_{\mathbb{R}^d} \frac{\delta(\tau + |\xi_1|^2 - |\xi - \xi_1|^2)}{|\xi - \xi_1| \langle \xi - \xi_1 \rangle^{\frac{1+2\epsilon}{2}} |\xi_1| \langle \xi_1 \rangle^{\frac{1+2\epsilon}{2}}} d\xi_1 \\ &= \frac{|\xi|^{1+2\epsilon}}{2} \int_{\tau - |\xi|^2 + 2\xi \cdot \xi_1 = 0} \frac{d\sigma_{\xi_1}}{|\xi - \xi_1| \langle \xi - \xi_1 \rangle^{\frac{1+2\epsilon}{2}} |\xi_1| \langle \xi_1 \rangle^{\frac{1+2\epsilon}{2}}} \lesssim 1, \end{aligned}$$

where $d\sigma_{\xi_1}$ is the standard measure on the plane $\tau - |\xi|^2 + 2\xi \cdot \xi_1 = 0$. This proves the lemma. \square

3. GENERALIZED RELATIVE ENTROPY

Following Lewin-Sabin [29, 31], we review the rigorous construction of the quantum relative entropy and its basic properties, and provide a refined Lieb-Thirring inequality (Proposition 3.6).

3.1. A heuristic argument. We begin with some formal calculations illustrating a motivation to introduce the relative entropy and the relative free energy. First, we note that solutions to the NLH (1.1) formally preserve the average energy

$$\mathcal{E}(\gamma) = \text{Tr}((-\Delta)\gamma) + \frac{1}{2} \int_{\mathbb{R}^3} w(x-y)\rho_\gamma(x)\rho_\gamma(y)dx dy$$

and a generalized entropy¹

$$\mathcal{T}_S(\gamma) := -\text{Tr}(S(\gamma))$$

for “any” function S . For $\gamma = \gamma_f + Q$, we formally expand the conserved quantity $\mathcal{T}_S(\gamma) - \mathcal{T}_S(\gamma_f) + \mathcal{E}(\gamma) - \mathcal{E}(\gamma_f)$ as

$$\begin{aligned} & -\text{Tr}(S(\gamma_f + Q) - S(\gamma_f) - (-\Delta)Q) + \frac{1}{2} \int_{\mathbb{R}^3} w(x-y)\rho_Q(x)\rho_Q(y)dx dy \\ & + \left\{ \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} f(|\xi|^2)d\xi \right\} \text{Tr}(Q). \end{aligned}$$

Since $\text{Tr}(Q) = \text{Tr}(\gamma) - \text{Tr}(\gamma_f)$ is formally conserved, so is

$$\mathcal{F}_f(\gamma_f + Q|\gamma_f) := \underbrace{-\text{Tr}(S(\gamma_f + Q) - S(\gamma_f) - (-\Delta)Q)}_{= \mathcal{H}_S(\gamma_f + Q|\gamma_f) = \text{relative entropy}} + \frac{1}{2} \int_{\mathbb{R}^d} w(x-y)\rho_Q(x)\rho_Q(y)dx dy.$$

However, in order to employ this formal conservation law for global well-posedness, it must satisfy some coercivity property. Therefore, we assume that $\hat{w} \geq 0$ and take $-\Delta = S'(\gamma_f)$ with $S'' \leq 0$. Then, by the formal Taylor series expansion, we have

$$\mathcal{H}_S(\gamma_f + Q|\gamma_f) = -\text{Tr}(S(\gamma_f + Q) - S(\gamma_f) - S'(\gamma_f)Q) \approx -\frac{1}{2} \text{Tr}(QS''(\gamma_f)Q) \geq 0. \quad (3.1)$$

3.2. General theory. Suppose that $S : [0, 1] \rightarrow \mathbb{R}^+$ is \mathcal{H} -admissible (see Definition 1.1), and define the *relative entropy* by

$$\mathcal{H}_S(A|B) := -\text{Tr}(S(A) - S(B) - S'(B)(A - B)) \in [0, \infty]$$

for two finite-dimensional Hermitian matrices $0 \leq A, B \leq 1$. By the assumptions, $\mathcal{H}_S(A|B)$ is monotone in the sense that $\mathcal{H}_S(XAX^*|XBX^*) \leq \mathcal{H}_S(A|B)$ for any linear map $X : H \rightarrow \tilde{H}$ such that $X^*X \leq 1$, where H and \tilde{H} are finite-dimensional vector spaces [29, Theorem 1]. Thus, the relative entropy can be extended to infinite-dimensional spaces as

$$\mathcal{H}_S(A|B) := \lim_{k \rightarrow \infty} \mathcal{H}_S(P_k A P_k | P_k B P_k) \in [0, \infty]$$

for self-adjoint operators $0 \leq A, B \leq 1$ on a separable Hilbert space H , where P_k is an increasing sequence of finite-rank projections on H such that $P_k \uparrow 1$ strongly. An important fact is that $\mathcal{H}_S(A|B)$ is well-defined, since it is independent of the choice of the sequence $P_k \uparrow 1$ [29, Theorem 2 (1)]. Moreover, the relative entropy satisfies the following properties.

Theorem 3.1 (Properties of the relative entropy [31, Theorem 2]). *Suppose that $S : [0, 1] \rightarrow \mathbb{R}^+$ is \mathcal{H} -admissible and that $0 \leq A, B \leq 1$ are self-adjoint on a Hilbert space H .*

¹because a solution can be written formally as $\gamma(t) = U(t)_* \gamma(0)$ for some unitary operator $U(t)$.

(i) (Monotonicity) If $X : H \rightarrow \tilde{H}$ is linear and $X^*X \leq 1$, then

$$\mathcal{H}_S(A|B) \geq \mathcal{H}_S(XAX^*|XBX^*).$$

(ii) (Approximation) If $X_k : H \rightarrow \tilde{H}_k$ is linear, $X_k^*X_k \leq 1$ and $X_k^*X_k \rightarrow 1$ strongly in H , then

$$\mathcal{H}_S(A|B) = \lim_{k \rightarrow \infty} \mathcal{H}_S(X_kAX_k^*|X_kBX_k^*).$$

(iii) (Weak lower semi-continuity) If $A_k \rightarrow A$ and $B_k \rightarrow B$ weakly-* in $\mathcal{B}(H)$, then

$$\mathcal{H}_S(A|B) \leq \liminf_{k \rightarrow \infty} \mathcal{H}_S(A_k|B_k).$$

By the assumptions, the relative entropy is non-negative (see also (3.1)). Indeed, even more than that, it enjoys a better coercivity property to measure the difference between two operators.

Theorem 3.2 (Klein's inequality [31, Theorem 3]). *Under the assumptions in Theorem 3.1, there exists $C_S > 0$, depending only on S , such that*

$$\mathcal{H}_S(A|B) \geq C_S \text{Tr}(1 + |S'(B)|)(A - B)^2.$$

3.3. Perturbation from quantum steady states. Now, coming to our setting, we fix a reference state of the form $\gamma_f = f(-\Delta)$ with $f = (S')^{-1}$ such that $0 \leq \gamma_f \leq 1$. By the conditions in Definition 1.1, for a self-adjoint operator $0 \leq \gamma \leq 1$ on $L^2(\mathbb{R}^3)$, we define its relative entropy from the reference state γ_f by

$$\mathcal{H}(\gamma|\gamma_f) = \mathcal{H}_S(\gamma|\gamma_f) := \lim_{k \rightarrow \infty} \mathcal{H}(P_k\gamma P_k|P_k\gamma_f P_k),$$

where P_k is a finite-rank projector such that $P_k \uparrow 1$ strongly in $L^2(\mathbb{R}^3)$. Then, collecting self-adjoint operators having finite relative entropy, we define the *relative free energy space* by

$$\mathcal{K}_f := \left\{ \gamma : L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3) \mid 0 \leq \gamma = \gamma^* \leq 1 \text{ and } \mathcal{H}(\gamma|\gamma_f) < \infty \right\}.$$

We note that Klein's inequality (Theorem 3.2) is stated as

$$\mathcal{H}(\gamma|\gamma_f) \gtrsim \|(\nabla)(\gamma - \gamma_f)\|_{\mathfrak{S}^2}^2 \geq \|\gamma - \gamma_f\|_{\mathcal{H}^{\frac{1}{2}}}^2. \quad (3.2)$$

Hence, $\gamma \in \mathcal{K}_f$ is a compact perturbation of γ_f .

The following lemma asserts that the relative entropy can be approximated by that of a finite-rank smooth perturbation.

Lemma 3.3 (Partial finite-dimensional approximation). *Suppose that $S : [0, 1] \rightarrow \mathbb{R}^+$ is \mathcal{H} -admissible, and $\gamma_f = f(-\Delta)$ with $f = (S')^{-1}$. Then, for any $\gamma = \gamma_f + Q \in \mathcal{K}_f$, there exist finite-rank smooth operators Q_k such that $0 \leq \gamma_f + Q_k \leq 1$ and*

$$\lim_{k \rightarrow \infty} |\mathcal{H}(\gamma_f + Q_k|\gamma_f) - \mathcal{H}(\gamma_f + Q|\gamma_f)| + \lim_{k \rightarrow \infty} \|Q_k - Q\|_{\mathfrak{S}^2} = 0.$$

Proof. First, we claim that the relative entropy $\mathcal{H}(\gamma_f + Q|\gamma_f)$ can be approximated by that for a regular perturbation. To see this, for small $\delta > 0$, let Π'_δ be the Fourier multiplier with symbol $\mathbb{1}_{\{\delta \leq f(|\xi|^2) \leq 1-\delta\}}$. Then, it is easy to check that $0 \leq \gamma_f + \Pi'_\delta Q \Pi'_\delta \leq 1$. Moreover, by the definition of the relative entropy and the monotonicity (Theorem 3.1 (ii)), we have

$$\mathcal{H}(\gamma_f + \Pi'_\delta Q \Pi'_\delta|\gamma_f) = \mathcal{H}(\Pi'_\delta(\gamma_f + Q)\Pi'_\delta|\Pi'_\delta\gamma_f\Pi'_\delta) \rightarrow \mathcal{H}(Q + \gamma_f|\gamma_f) \text{ as } \delta \rightarrow 0.$$

Thus, for any $\epsilon > 0$, we can take $\delta > 0$ such that neither $-(1-\delta)$ nor δ are not eigenvalues of Q and $|\mathcal{H}(\gamma_f + Q|\gamma_f) - \mathcal{H}(\gamma_f + \Pi'_\delta Q \Pi'_\delta|\gamma_f)| \leq \epsilon$.

Next, using the spectral decomposition $Q = \sum_{n=1}^{\infty} a_n^+ |u_n^+\rangle\langle u_n^+| - \sum_{n=1}^{\infty} a_n^- |u_n^-\rangle\langle u_n^-|$, where $(u_n^\pm)_{n=1}^{\infty}$ are orthonormal sets in L_x^2 and $a_n^\pm \geq 0$, we introduce the finite-rank cut-off

$$Q_k := P_k Q P_k = Q_k^+ - Q_k^- := \sum_{n=1}^k a_n^+ |u_n^+\rangle\langle u_n^+| - \sum_{n=1}^k a_n^- |u_n^-\rangle\langle u_n^-|.$$

We claim that $0 \leq Q_k + \gamma_f \leq 1$. Indeed, we have $\gamma_f + Q_k^+ \leq 1$ on $\text{Ran} Q_k^+$, where $\text{Ran} A$ denotes the image of A , because otherwise, there exist $u \in \text{Ran} Q_k^+$ such that $\|u\|_{L_x^2}^2 < \langle u | (\gamma_f + Q_k^+) u \rangle = \langle u | (\gamma_f + Q) u \rangle$. Thus, $\gamma_f + Q_k^+ \leq 1$ on L_x^2 . Similarly, one can show that $\gamma_f - Q_k^- \geq 0$. Therefore, it follows that $0 \leq \gamma_f - Q_k^- \leq \gamma_f + Q_k \leq \gamma_f + Q_k^+ \leq 1$. By the claim, we also have $0 \leq \gamma_f + \Pi'_\delta Q_k \Pi'_\delta \leq 1$, so the relative entropy is well-defined for $\gamma_f + \Pi'_\delta Q_k \Pi'_\delta$.

It remains to show that

$$\lim_{k \rightarrow \infty} \mathcal{H}(\gamma_f + \Pi'_\delta Q_k \Pi'_\delta|\gamma_f) = \mathcal{H}(\gamma_f + \Pi'_\delta Q \Pi'_\delta|\gamma_f).$$

Note that expanding by the definition,

$$\begin{aligned} & \mathcal{H}(\gamma_f + \Pi'_\delta Q \Pi'_\delta|\gamma_f) - \mathcal{H}(\gamma_f + \Pi'_\delta Q_k \Pi'_\delta|\gamma_f) \\ &= \mathcal{H}(\Pi'_\delta(\gamma_f + Q)\Pi'_\delta|\Pi'_\delta\gamma_f\Pi'_\delta) - \mathcal{H}(\Pi'_\delta(\gamma_f + Q_k)\Pi'_\delta|\Pi'_\delta\gamma_f\Pi'_\delta). \end{aligned} \quad (3.3)$$

Thus, we may work on the Hilbert space $\Pi'_\delta L_x^2$, not on L_x^2 . For the proof, we will employ [29, Theorem 3], which requires strict upper and lower bounds. Indeed, it is obvious that $\delta \leq \Pi'_\delta \gamma_f \Pi'_\delta \leq 1 - \delta$ on $\Pi'_\delta L_x^2$. We claim that there exists small $\delta' > 0$ such that

$$\delta' \leq \Pi'_\delta(\gamma_f + Q)\Pi'_\delta \leq 1 - \delta' \quad (3.4)$$

on $\Pi'_\delta L_x^2$. Indeed, if the lower bound in (3.4) does not hold, there exists $u_n \in \Pi'_\delta L_x^2$ such that $\|u_n\|_{L_x^2} = 1$ and $\langle u_n | (\gamma_f + Q) u_n \rangle \leq \frac{1}{n}$. Then, $\langle u_n | Q u_n \rangle = \langle u_n | (\gamma_f + Q) u_n \rangle - \langle u_n | \gamma_f u_n \rangle \leq \frac{1}{n} - (1 - \delta) \rightarrow -(1 - \delta)$ as $n \rightarrow \infty$. On the other hand, since $Q \geq -\gamma_f \geq -(1 - \delta)$ on $\Pi'_\delta L_x^2$, it implies that $-(1 - \delta)$ is an eigenvalue of Q , which deduces a contradiction. Similarly, one can show the upper bound in (3.4). Therefore, applying [29, Theorem 3] to (3.3) with (3.4), we obtain

$$\begin{aligned} & |\mathcal{H}(\gamma_f + \Pi'_\delta Q \Pi'_\delta|\gamma_f) - \mathcal{H}(\gamma_f + \Pi'_\delta Q_k \Pi'_\delta|\gamma_f)| \\ & \lesssim_{\delta'} \|\Pi'_\delta(Q - Q_k)\Pi'_\delta\|_{\mathfrak{S}^2}^2 + \|\Pi'_\delta Q \Pi'_\delta\|_{\mathfrak{S}^2} \|\Pi'_\delta(Q - Q_k)\Pi'_\delta\|_{\mathfrak{S}^2} \rightarrow 0 \text{ as } k \rightarrow \infty, \end{aligned}$$

which proves the lemma. \square

A very important remark is that in this setting, the relative entropy has a useful lower bound (see Lemma 3.4 below). To see this, we recall the notion of the relative kinetic energy from a Fermi sea; we refer to [23] for more detailed rigorous construction. Let $\Pi_\mu^- = \mathbb{1}_{(-\Delta \leq \mu)}$ be the Fermi sea with a chemical potential $\mu > 0$. For a quantum state $0 \leq \gamma = \Pi_\mu^- + Q \leq 1$ with a finite-rank smooth perturbation Q , its relative kinetic energy is defined by

$$\mathrm{Tr}(-\Delta - \mu)(\gamma - \Pi_\mu^-) = \mathrm{Tr}(-\Delta - \mu)Q.$$

Note that $\mathrm{Tr}(-\Delta - \mu)Q$ is non-negative, because $\mathrm{Tr}(-\Delta - \mu)Q = \mathrm{Tr}(-\Delta - \mu)\Pi_\mu^+ \gamma \Pi_\mu^+ + \mathrm{Tr}(\Delta + \mu)\Pi_\mu^-(1 - \gamma)\Pi_\mu^- \geq 0$, where $\Pi_\mu^+ = 1 - \Pi_\mu^- = \mathbb{1}_{(-\Delta > \mu)}$. Moreover, the inequality

$$\|Q|\Delta + \mu|^{\frac{1}{2}}\|_{\mathfrak{S}^2}^2 \leq \mathrm{Tr}(-\Delta - \mu)Q \quad (3.5)$$

holds. We define the operator class \mathcal{K}_μ having finite relative energy from Π_μ^- by

$$\mathcal{K}_\mu := \Pi_\mu^- + \mathcal{X}_\mu,$$

where \mathcal{X}_μ is the completion of the smooth finite-rank self-adjoint operators on $L^2(\mathbb{R}^d)$ such that $-\Pi_\mu^- \leq Q \leq \Pi_\mu^+$ (so, $0 \leq \Pi_\mu^- + Q \leq 1$) with respect to the norm

$$\|Q\|_{\mathcal{X}_\mu} := \|Q\| + \|Q|\Delta + \mu|^{\frac{1}{2}}\|_{\mathfrak{S}^2} + \sum_{\pm} \| |\Delta + \mu|^{\frac{1}{2}} \Pi_\mu^\pm Q \Pi_\mu^\pm |\Delta + \mu|^{\frac{1}{2}} \|_{\mathfrak{S}^1}.$$

Then, by density, $\mathrm{Tr}(-\Delta - \mu)Q \geq 0$ and (3.5) holds for all $\gamma \in \mathcal{K}_\mu$.

Lemma 3.4 (Lower bound for the relative entropy [31, (86)]). *If $S : [0, 1] \rightarrow \mathbb{R}^+$ is \mathcal{H} -admissible and $\gamma_f = f(-\Delta)$ with $f = (S')^{-1}$, then*

$$\mathcal{H}(\gamma|\gamma_f) \geq \int_0^1 \mathrm{Tr}(-\Delta - \mu)Q_\mu d\lambda,$$

where $\mu = S'(\lambda)$ and $Q_\mu = \mathbb{1}(\gamma \geq \lambda) - \Pi_\mu^-$.

3.4. Lieb-Thirring inequalities. The fundamental Lieb-Thirring inequality states that if $d = 3$ and $\frac{5}{3} \leq r \leq 3$, then

$$\|\rho_\gamma\|_{L^r} \lesssim \| |\nabla| \gamma |\nabla| \|_{\mathfrak{S}^1}^{\frac{3(r-1)}{2r}} \quad (3.6)$$

for any self-adjoint operator γ such that $0 \leq \gamma \leq 1$. In Frank-Lewin-Lieb-Seiringer [23], extending to quantum states perturbed from Fermi seas, it is shown that for $\Pi_\mu^- + Q \in \mathcal{K}_\mu$,

$$\mathrm{Tr}(-\Delta - \mu)Q \gtrsim \int_{\mathbb{R}^3} \left\{ (\rho_{\Pi_\mu^-} + \rho_Q(x))^{\frac{5}{3}} - (\rho_{\Pi_\mu^-})^{\frac{5}{3}} - \frac{5}{3}(\rho_{\Pi_\mu^-})^{\frac{2}{3}}\rho_Q(x) \right\} dx. \quad (3.7)$$

From this, we derive a form that is convenient to use in our setting.

Lemma 3.5 (Lieb-Thirring inequality for perturbations from a Fermi sea at zero temperature). *For any $\mu > 0$ and $\Pi_\mu^- + Q \in \mathcal{K}_\mu$, we have*

$$\|\rho_Q\|_{H^{\frac{1}{2}}} \lesssim \mathrm{Tr}((-\Delta - \mu)Q) + \left\{ \mathrm{Tr}((-\Delta - \mu)Q) \right\}^{\frac{3}{4}} + (\mu^{\frac{1}{4}} + \mu^{\frac{1}{2}}) \left\{ \mathrm{Tr}((-\Delta - \mu)Q) \right\}^{\frac{1}{2}}.$$

Proof. For $\Pi_\mu^- + Q \in \mathcal{K}_\mu$, we split $Q = \Pi_{2\mu}^+ Q \Pi_{2\mu}^+ + \Pi_{2\mu}^+ Q \Pi_{2\mu}^- + \Pi_{2\mu}^- Q \Pi_{2\mu}^+ + \Pi_{2\mu}^- Q \Pi_{2\mu}^-$, and estimate them separately.

(Step 1. High-high term) We claim that

$$\|\rho_{\Pi_{2\mu}^+ Q \Pi_{2\mu}^+}\|_{H^{\frac{1}{2}}} \lesssim \text{Tr}((-\Delta - \mu)Q) + \left\{ \text{Tr}((-\Delta - \mu)Q) \right\}^{\frac{3}{4}}.$$

Indeed, since $0 \leq \Pi_{2\mu}^+ Q \Pi_{2\mu}^+ = \Pi_{2\mu}^+ \gamma \Pi_{2\mu}^+ \leq 1$ is self-adjoint and non-negative, it follows from (3.6) with $r = 2$ that

$$\|\rho_{\Pi_{2\mu}^+ Q \Pi_{2\mu}^+}\|_{L^2} \lesssim \| |\nabla| \Pi_{2\mu}^+ Q \Pi_{2\mu}^+ |\nabla| \|_{\mathfrak{S}^1}^{\frac{3}{4}} \sim (\text{Tr}(-\Delta - \mu)Q)^{\frac{3}{4}}.$$

On the other hand, by the spectral decomposition $\Pi_{2\mu}^+ Q \Pi_{2\mu}^+ = \sum_{j=1}^{\infty} c_{\mu;j} |\phi_{\mu;j}\rangle \langle \phi_{\mu;j}|$ with $\{c_{\mu;j}\}_{j=1}^{\infty} \subset [0, 1]$ and an orthonormal set $\{\phi_{\mu;j}\}_{j=1}^{\infty}$, the simple triangle inequality, the fractional Leibniz rule and the Sobolev inequality, we obtain that

$$\begin{aligned} \||\nabla|^{\frac{1}{2}} \rho_{\Pi_{2\mu}^+ Q \Pi_{2\mu}^+}\|_{L^2} &\leq \sum_{j=1}^{\infty} c_{\mu;j} \||\nabla|^{\frac{1}{2}} (|\phi_{\mu;j}|^2)\|_{L^2} \lesssim \sum_{j=1}^{\infty} c_{\mu;j} \|\phi_{\mu;j}\|_{L^6} \||\nabla|^{\frac{1}{2}} \phi_{\mu;j}\|_{L^3} \\ &\lesssim \sum_{j=1}^{\infty} c_{\mu;j} \|\nabla \phi_{\mu;j}\|_{L^2}^2 = \text{Tr}(|\nabla| \Pi_{2\mu}^+ Q \Pi_{2\mu}^+ |\nabla|) \lesssim \text{Tr}((-\Delta - \mu)Q). \end{aligned}$$

(Step 2. High-low and low high terms) By symmetry, it suffices to consider the high-low term $\Pi_{2\mu}^+ Q \Pi_{2\mu}^-$. For the proof, we apply the Plancherel theorem with

$$\hat{\rho}_{\Pi_{2\mu}^+ Q \Pi_{2\mu}^-}(\xi) = \frac{1}{(2\pi)^3} \int_{|\xi - \xi_1| \leq \sqrt{2\mu} \leq |\xi_1|} \hat{Q}(\xi_1, \xi - \xi_1) d\xi_1$$

and the Hölder inequality to obtain

$$\|\rho_{\Pi_{2\mu}^+ Q \Pi_{2\mu}^-}\|_{H^{\frac{1}{2}}}^2 \lesssim \left\{ \sup_{\xi \in \mathbb{R}^3} \langle \xi \rangle \int_{|\xi - \xi_1| \leq \sqrt{2\mu} \leq |\xi_1|} \frac{d\xi_1}{|\xi_1|^2} \right\} \|(|\nabla| \Pi_{2\mu}^+ Q)(x, x')\|_{L_x^2 L_{x'}^2}^2.$$

We observe that

$$\langle \xi \rangle \int_{|\xi - \xi_1| \leq \sqrt{2\mu} \leq |\xi_1|} \frac{d\xi_1}{|\xi_1|^2} \lesssim \int_{|\xi - \xi_1| \leq \sqrt{2\mu} \leq |\xi_1|} \frac{1}{|\xi_1|^2} + \frac{1}{|\xi_1|} + \frac{|\xi - \xi_1|}{|\xi_1|^2} d\xi_1 \sim \mu^{\frac{1}{2}} + \mu.$$

On the other hand, we have by (3.5) that

$$\|(|\nabla| \Pi_{2\mu}^+ Q)(x, x')\|_{L_x^2 L_{x'}^2}^2 = \| |\nabla| \Pi_{2\mu}^+ Q \|_{\mathfrak{S}^2}^2 \sim \| |-\Delta - \mu|^{\frac{1}{2}} Q \|_{\mathfrak{S}^2}^2 \lesssim \text{Tr}((-\Delta - \mu)Q).$$

Thus, it follows that

$$\|\rho_{\Pi_{2\mu}^+ Q \Pi_{2\mu}^-}\|_{H^{\frac{1}{2}}} \lesssim (\mu^{\frac{1}{4}} + \mu^{\frac{1}{2}}) \left\{ \text{Tr}((-\Delta - \mu)Q) \right\}^{\frac{1}{2}}.$$

(Step 3. Low-low term) Note that $0 \leq \Pi_\mu^- + (\Pi_{2\mu}^- Q \Pi_{2\mu}^-) = \Pi_{2\mu}^- \gamma \Pi_{2\mu}^- \leq 1$. Hence, it follows from the Lieb-Thirring inequality (3.7) that

$$\begin{aligned} \mathrm{Tr}(-\Delta - \mu)Q &\geq \mathrm{Tr}(-\Delta - \mu)\Pi_{2\mu}^- Q \Pi_{2\mu}^- \\ &\gtrsim \int_{\mathbb{R}^3} \left\{ (\rho_{\Pi_\mu^-} + \rho_{\Pi_{2\mu}^- Q \Pi_{2\mu}^-}(x))^{\frac{5}{3}} - (\rho_{\Pi_\mu^-})^{\frac{5}{3}} - \frac{5}{3}(\rho_{\Pi_\mu^-})^{\frac{2}{3}}\rho_{\Pi_{2\mu}^- Q \Pi_{2\mu}^-}(x) \right\} dx \\ &= (\rho_{\Pi_\mu^-})^{\frac{5}{3}} \int_{\mathbb{R}^3} \left\{ \left(1 + \frac{\rho_{\Pi_{2\mu}^- Q \Pi_{2\mu}^-}(x)}{\rho_{\Pi_\mu^-}} \right)^{\frac{5}{3}} - 1 - \frac{5}{3} \frac{\rho_{\Pi_{2\mu}^- Q \Pi_{2\mu}^-}(x)}{\rho_{\Pi_\mu^-}} \right\} dx. \end{aligned}$$

Note that $0 \leq \rho_{\Pi_{2\mu}^- Q \Pi_{2\mu}^-}(x) \lesssim \rho_{\Pi_{2\mu}^-} \sim \mu^{\frac{3}{2}}$ for all $x \in \mathbb{R}^3$, because $0 \leq \Pi_{2\mu}^- Q \Pi_{2\mu}^- \leq \Pi_{2\mu}^-$. Thus, Taylor's theorem yields

$$\begin{aligned} \mathrm{Tr}(-\Delta - \mu)Q &\gtrsim (\rho_{\Pi_\mu^-})^{\frac{5}{3}} \int_{\mathbb{R}^3} \left(\frac{\rho_{\Pi_{2\mu}^- Q \Pi_{2\mu}^-}(x)}{\rho_{\Pi_\mu^-}} \right)^2 dx \\ &\sim (\rho_{\Pi_\mu^-})^{-\frac{1}{3}} \|\rho_{\Pi_{2\mu}^- Q \Pi_{2\mu}^-}\|_{L^2}^2 \sim \mu^{-\frac{1}{2}} \|\rho_{\Pi_{2\mu}^- Q \Pi_{2\mu}^-}\|_{L^2}^2. \end{aligned} \tag{3.8}$$

Consequently, we conclude that

$$\| |\nabla|^{\frac{1}{2}} \rho_{\Pi_{2\mu}^- Q \Pi_{2\mu}^-} \|_{L_x^2} \lesssim \mu^{\frac{1}{4}} \|\rho_{\Pi_{2\mu}^- Q \Pi_{2\mu}^-}\|_{L_x^2} \lesssim \mu^{\frac{1}{2}} \left\{ \mathrm{Tr}((-\Delta - \mu)Q) \right\}^{\frac{1}{2}},$$

since the Fourier transform of $\rho_{\Pi_{2\mu}^- Q \Pi_{2\mu}^-}$ is supported in $|\xi| \leq 2\sqrt{2\mu}$. \square

The Lieb-Thirring inequality [30, Theorem 7] asserts that $\rho_{\gamma - \gamma_f} \in L^2 + L^{\frac{5}{3}}$ when $\gamma \in \mathcal{K}_f$. As an application of Lemma 3.5, it is extended as:

Proposition 3.6. *Assume that $S : [0, 1] \rightarrow \mathbb{R}^+$ is \mathcal{H} -admissible and $f = (S')^{-1}$. Then, for any $0 \leq \gamma \leq 1$, we have*

$$\|\rho_{\gamma - \gamma_f}\|_{H^{\frac{1}{2}}} \lesssim \mathcal{H}(\gamma|\gamma_f) + \mathcal{H}(\gamma|\gamma_f)^{\frac{1}{2}}.$$

Proof. By the integral representation (1.9), we have

$$\|\rho_Q\|_{H^{\frac{1}{2}}} \leq \int_0^1 \|\rho_{Q_\mu}\|_{H^{\frac{1}{2}}} d\lambda,$$

where $Q_\mu := \mathbf{1}(\gamma \geq \lambda) - \Pi_\mu^-$. Then, applying Lemma 3.5 to $\|\rho_{Q_\mu}\|_{H^{\frac{1}{2}}}$ and Hölder's inequality in λ and (1.7), we conclude that $\|\rho_Q\|_{H^{\frac{1}{2}}} \lesssim \mathcal{H}(\gamma|\gamma_f) + \mathcal{H}(\gamma|\gamma_f)^{\frac{3}{4}} + \mathcal{H}(\gamma|\gamma_f)^{\frac{1}{2}}$. \square

4. DENSITY FUNCTION ESTIMATES FOR LINEAR SCHRÖDINGER FLOWS

In this section, we develop various density function estimates for linear Schrödinger flows perturbed by time-dependent real-valued potentials in

$$\mathcal{B}_R^{1+\epsilon}(I) := \{g = g(t, x) : \|g\|_{\mathcal{D}^{1+\epsilon}(I)} \leq R\},$$

where $\epsilon > 0$ is a sufficiently small number, $R > 0$, $I \subset \mathbb{R}$, and

$$\|g\|_{\mathcal{D}^{1+\epsilon}(I)} := \|g\|_{L_t^\infty(I; L_x^2) \cap L_t^2(I; \dot{H}_x^{1+\epsilon})}.$$

In later sections, they will be employed for the local well-posedness of the nonlinear problem.

4.1. Hilbert-Schmidt Sobolev initial data. As a first step, we extend the density function estimates for the free flow (see Lemma 2.1 and 2.2), and show continuous dependence on potential perturbations.

Lemma 4.1 (Density function estimates for linear flows I). *Let $R \geq 1$. Suppose that $t_0 \in I$ and $0 < |I| \ll R^{-\frac{4}{1+2\epsilon}}$ for sufficiently small $\epsilon > 0$. Then, for any $V, V' \in \mathcal{B}_R^{1+\epsilon}(I)$, we have*

$$\|\rho_{S_V(t,t_0)\gamma_0 S_{V'}(t,t_0)^*}\|_{\mathcal{D}^{1+\epsilon}(I)} \leq c_1 \|\ |\nabla|^{\frac{1}{2}} \gamma_0 |\nabla|^{\frac{1}{2}}\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}}, \quad (4.1)$$

$$\|\rho_{S_V(t,t_0)\star\gamma_0} - \rho_{S_{V'}(t,t_0)\star\gamma_0}\|_{\mathcal{D}^{1+\epsilon}(I)} \leq c_1 |I|^{\frac{1+2\epsilon}{4}} \|\ |\nabla|^{\frac{1}{2}} \gamma_0 |\nabla|^{\frac{1}{2}}\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}} \|V - V'\|_{L_t^2(I; H_x^{1+\epsilon})}. \quad (4.2)$$

The following lemma is elementary but useful in our analysis.

Lemma 4.2. *For small $\epsilon > 0$, we have*

$$\|\ |\nabla|^{\frac{1}{2}} V \gamma |\nabla|^{\frac{1}{2}}\|_{L_t^1(I; \mathcal{H}^{\frac{1+2\epsilon}{4}})} \lesssim |I|^{\frac{1+2\epsilon}{4}} \|V\|_{L_t^2(I; H_x^{1+\epsilon})} \|\ |\nabla|^{\frac{1}{2}} \gamma |\nabla|^{\frac{1}{2}}\|_{S^{\frac{1+2\epsilon}{4}}(I)}.$$

Proof. By the fractional Leibniz rule and the Sobolev inequality, we obtain

$$\begin{aligned} \|\ |\nabla|^{\frac{1}{2}} (Vu)\|_{W_x^{\frac{1+2\epsilon}{4}}} &\lesssim \|V\|_{W_x^{\frac{3+2\epsilon}{4}, \frac{12}{5-2\epsilon}}} \|u\|_{L_x^{\frac{12}{1+2\epsilon}}} + \|V\|_{L_x^{\frac{6}{1-2\epsilon}}} \|\ |\nabla|^{\frac{1}{2}} u\|_{W_x^{\frac{1+2\epsilon}{4}, \frac{3}{1+\epsilon}}} \\ &\lesssim \|V\|_{H_x^{1+\epsilon}} \|\ |\nabla|^{\frac{1}{2}} u\|_{W_x^{\frac{1+2\epsilon}{4}, \frac{3}{1+\epsilon}}}, \end{aligned}$$

and consequently,

$$\|\ |\nabla|^{\frac{1}{2}} V \gamma |\nabla|^{\frac{1}{2}}\|_{L_t^1(I; \mathcal{H}^{\frac{1+2\epsilon}{4}})} \lesssim |I|^{\frac{1+2\epsilon}{4}} \|V\|_{L_t^2(I; H_x^{1+\epsilon})} \|\ |\nabla|^{\frac{1}{2}} \gamma |\nabla|^{\frac{1}{2}}\|_{L_t^{\frac{4}{1-2\epsilon}}(I; W_x^{\frac{1+2\epsilon}{4}, \frac{3}{1+\epsilon}} H_x^{\frac{1+2\epsilon}{4}})}.$$

Since $(\frac{4}{1-2\epsilon}, \frac{3}{1+\epsilon})$ is admissible, the lemma follows. \square

Proof of Lemma 4.1. For (4.1), let $\gamma(t) := S_V(t, t_0)\gamma_0 S_{V'}(t, t_0)^*$ solving the integral equation

$$\gamma(t) = e^{i(t-t_0)\Delta} \gamma_0 e^{-i(t-t_0)\Delta} - i \int_{t_0}^t e^{i(t-t_1)\Delta} (V\gamma - \gamma V')(t_1) e^{-i(t-t_1)\Delta} dt_1.$$

Then, it follows from Strichartz estimates for the free flow (see Lemma 2.1 and 2.2) and Lemma 4.2 that

$$\begin{aligned} &\|\rho_\gamma\|_{\mathcal{D}^{1+\epsilon}(I)} + \|\ |\nabla|^{\frac{1}{2}} \gamma |\nabla|^{\frac{1}{2}}\|_{S^{\frac{1+2\epsilon}{4}}(I)} \\ &\leq c_0 \|\ |\nabla|^{\frac{1}{2}} \gamma_0 |\nabla|^{\frac{1}{2}}\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}} + c_0 \|\ |\nabla|^{\frac{1}{2}} V \gamma |\nabla|^{\frac{1}{2}}\|_{L_t^1(I; \mathcal{H}^{\frac{1+2\epsilon}{4}})} + c_0 \|\ |\nabla|^{\frac{1}{2}} V' \gamma |\nabla|^{\frac{1}{2}}\|_{L_t^1(I; \mathcal{H}^{\frac{1+2\epsilon}{4}})} \\ &\leq c_0 \|\ |\nabla|^{\frac{1}{2}} \gamma_0 |\nabla|^{\frac{1}{2}}\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}} + 2\tilde{c}_0 |I|^{\frac{1+2\epsilon}{4}} R \|\ |\nabla|^{\frac{1}{2}} \gamma |\nabla|^{\frac{1}{2}}\|_{S^{\frac{1+2\epsilon}{4}}(I)}. \end{aligned}$$

Hence, taking an interval I such that $|I| = (4\tilde{c}_0 R)^{-\frac{4}{1+2\epsilon}}$, we prove that

$$\|\rho_\gamma\|_{\mathcal{D}^{1+\epsilon}(I)} + \|\ |\nabla|^{\frac{1}{2}} \gamma |\nabla|^{\frac{1}{2}}\|_{S^{\frac{1+2\epsilon}{4}}(I)} \leq 2c_0 \|\ |\nabla|^{\frac{1}{2}} \gamma_0 |\nabla|^{\frac{1}{2}}\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}}. \quad (4.3)$$

For (4.2), we write the difference between $\gamma(t) := S_V(t, t_0)_* \gamma_0$ and $\gamma'(t) := S_{V'}(t, t_0)_* \gamma_0$ as

$$\gamma(t) - \gamma'(t) = -i \int_{t_0}^t e^{i(t-t_1)\Delta} ([V - V', \gamma] + [V', \gamma - \gamma'])(t_1) e^{i(t-t_1)\Delta} dt_1.$$

Indeed, repeating the argument to show (4.3) and by the choice of T , one can show that

$$\begin{aligned} & \|\rho_\gamma - \rho_{\gamma'}\|_{\mathcal{D}^{1+\epsilon}(I)} + \left\| |\nabla|^{\frac{1}{2}} (\gamma - \gamma') |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{S}^{\frac{1+2\epsilon}{4}}(I)} \\ & \leq \tilde{c}_0 |I|^{\frac{1+2\epsilon}{4}} \|V - V'\|_{L_t^2(I; H_x^{1+\epsilon})} \left\| |\nabla|^{\frac{1}{2}} \gamma |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{S}^{\frac{1+2\epsilon}{4}}(I)} \\ & \quad + \tilde{c}_0 |I|^{\frac{1+2\epsilon}{4}} \|V'\|_{L_t^2(I; H_x^{1+\epsilon})} \left\| |\nabla|^{\frac{1}{2}} (\gamma - \gamma') |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{S}^{\frac{1+2\epsilon}{4}}(I)} \\ & \leq 2c_0 \tilde{c}_0 |I|^{\frac{1+2\epsilon}{4}} \left\| |\nabla|^{\frac{1}{2}} \gamma_0 |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}} \|V - V'\|_{L_t^2(I; H_x^{1+\epsilon})} + \frac{1}{2} \left\| |\nabla|^{\frac{1}{2}} (\gamma - \gamma') |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{S}^{\frac{1+2\epsilon}{4}}(I)}. \end{aligned}$$

Thus, (4.2) follows. \square

4.2. Perturbation from a Fermi sea at zero temperature. Next, we consider the linear Schrödinger flows from initial data having finite relative kinetic energy, and prove analogous density function estimates.

Lemma 4.3 (Density function estimates for linear flows II). *Let $R \geq 1$. Suppose that $t_0 \in I$ and $0 < |I| \ll R^{-\frac{4}{1+2\epsilon}}$ for sufficiently small $\epsilon > 0$. Then, for any $\gamma_0 = \Pi_\mu^- + Q_0 \in \mathcal{K}_\mu$ with $\mu > 0$, $V, V' \in \mathcal{B}_R^{1+\epsilon}(I)$ and $t_0 \in I$, we have*

$$\|\rho_{S_V(t, t_0)_* Q_0}\|_{\mathcal{D}^{1+\epsilon}(I)} \leq c_2 \{ \text{Tr}(-\Delta - \mu) Q_0 + 1 + \mu^{\frac{3}{2}+\epsilon} \} \quad (4.4)$$

and

$$\begin{aligned} & \|\rho_{S_V(t, t_0)_* Q_0} - \rho_{S_{V'}(t, t_0)_* Q_0}\|_{\mathcal{D}^{1+\epsilon}(I)} \\ & \leq c_2 |I|^{\frac{1+2\epsilon}{4}} \{ \text{Tr}(-\Delta - \mu) Q_0 + 1 + \mu^{\frac{3}{2}+\epsilon} \} \|V - V'\|_{L_t^2(I; H_x^{1+\epsilon})}. \end{aligned} \quad (4.5)$$

For the proof, we need the following lemma.

Lemma 4.4 (Boundedness of the linear propagator in Sobolev norms). *Let $\alpha \in (1, \frac{3}{2})$, $R \geq 1$ and $0 < T \ll R^{-\frac{4}{2\alpha-1}}$. Then, for any $V, V' \in \mathcal{B}_R^\alpha(I)$ and $t_0 \in I$, we have*

$$\sup_{t \in I} \|S_V(t, t_0)\|_{\mathcal{L}(\dot{H}_x^\alpha)} \leq 2, \quad (4.6)$$

$$\sup_{t \in I} \|S_V(t, t_0) - S_{V'}(t, t_0)\|_{\mathcal{L}(\dot{H}_x^\alpha)} \leq \tilde{c} |I|^{\frac{2\alpha-1}{4}} \|V - V'\|_{L_t^2(I; \dot{H}_x^\alpha)}, \quad (4.7)$$

where $\|\cdot\|_{\mathcal{L}(\dot{H}_x^\alpha)}$ is the operator norm on \dot{H}_x^α .

Proof. Let $u(t) = S_V(t, t_0)u_0$ and $u'(t) = S_{V'}(t, t_0)u_0$. Then, applying Strichartz estimates (2.1) to

$$u(t) = e^{i(t-t_0)\Delta} u_0 - i \int_{t_0}^t e^{i(t-t_1)\Delta} (Vu)(t_1) dt_1,$$

we obtain that

$$\|u\|_{C_t(I; \dot{H}_x^\alpha)} \leq \|u_0\|_{\dot{H}_x^\alpha} + c \left\| |\nabla|^\alpha (Vu) \right\|_{L_t^{\frac{4}{1+2\alpha}}(I; L_x^{\frac{3}{3-\alpha})}}.$$

Note that by the fractional Leibniz rule,

$$\left\| |\nabla|^\alpha (Vu) \right\|_{L_t^{\frac{4}{1+2\alpha}}(I; L_x^{\frac{3}{3-\alpha}})} \lesssim |I|^{\frac{2\alpha-1}{4}} \|V\|_{L_t^2(I; \dot{H}_x^\alpha)} \|u\|_{C_t(I; \dot{H}_x^\alpha)}.$$

Hence, it follows that

$$\|u\|_{C_t(I; \dot{H}_x^\alpha)} \leq \|u_0\|_{\dot{H}_x^\alpha} + \tilde{c}|I|^{\frac{2\alpha-1}{4}} R \|u\|_{C_t(I; \dot{H}_x^\alpha)}.$$

Then, taking an interval I such that $|I| \leq (2\tilde{c}R)^{-\frac{4}{2\alpha-1}}$, we prove (4.6).

For (4.7), we consider the difference

$$(u - u')(t) = -i \int_{t_0}^t e^{i(t-t_1)\Delta} |\nabla|^s (Vu - V'u')(t_1) dt_1.$$

Indeed, by Strichartz estimates and the fractional Leibniz rule as before, we have

$$\begin{aligned} \|u - u'\|_{C_t(I; \dot{H}_x^\alpha)} &\leq c|I|^{\frac{2\alpha-1}{4}} \|V - V'\|_{L_t^2(I; \dot{H}_x^\alpha)} \|u\|_{C_t(I; \dot{H}_x^\alpha)} \\ &\quad + \tilde{c}|I|^{\frac{2\alpha-1}{4}} \|V'\|_{L_t^2(I; \dot{H}_x^\alpha)} \|u - u'\|_{C_t(I; \dot{H}_x^\alpha)}. \end{aligned}$$

Then, by the choice of I , (4.7) follows. \square

Proof of Lemma 4.3. We decompose the initial perturbation into the high and the low frequency parts as

$$Q_0 = (Q_0 - \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^-) + \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^-.$$

For the high frequency contribution, we apply Lemma 4.1 to obtain

$$\|\rho_{S_V(t, t_0)_* (Q_0 - \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^-)}\|_{\mathcal{D}^{1+\epsilon}(I)} \lesssim \left\| |\nabla|^{\frac{1}{2}} (Q_0 - \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^-) |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}}$$

and

$$\begin{aligned} &\left\| \rho_{S_V(t, t_0)_* (Q_0 - \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^-)} - \rho_{S_{V'}(t, t_0)_* (Q_0 - \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^-)} \right\|_{\mathcal{D}^{1+\epsilon}(I)} \\ &\lesssim |I|^{\frac{1+2\epsilon}{4}} \|V - V'\|_{L_t^2(I; H_x^{1+\epsilon})} \left\| |\nabla|^{\frac{1}{2}} (Q_0 - \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^-) |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}}. \end{aligned}$$

Hence, it suffices to show that

$$\begin{aligned} &\left\| |\nabla|^{\frac{1}{2}} \Pi_{2\mu}^+ Q_0 \Pi_{2\mu}^+ |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}} + \left\| |\nabla|^{\frac{1}{2}} \Pi_{2\mu}^+ Q_0 \Pi_{2\mu}^- |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}} \\ &\lesssim \text{Tr}(-\Delta - \mu) Q_0 + 1 + \mu^{\frac{3}{2}+\epsilon}. \end{aligned} \tag{4.8}$$

For the high-high frequency component, we have

$$\left\| |\nabla| \Pi_{2\mu}^+ Q_0 \Pi_{2\mu}^+ |\nabla| \right\|_{\mathfrak{S}^2} \lesssim \left\| |\Delta + \mu|^{\frac{1}{2}} \Pi_{2\mu}^+ Q_0 \Pi_{2\mu}^+ |\Delta + \mu|^{\frac{1}{2}} \right\|_{\mathfrak{S}^1} \leq \text{Tr}(-\Delta - \mu) Q_0$$

while

$$\begin{aligned} \left\| |\nabla|^{\frac{1}{2}} \Pi_{2\mu}^+ Q_0 \Pi_{2\mu}^+ |\nabla|^{\frac{1}{2}} \right\|_{\mathfrak{S}^2} &= \left\{ \text{Tr}(Q_0 |\nabla| \Pi_{2\mu}^+ Q_0 \Pi_{2\mu}^+ |\nabla|) \right\}^{\frac{1}{2}} \lesssim \|Q_0\|_{\mathfrak{S}^2}^{\frac{1}{2}} \left\| |\nabla| \Pi_{2\mu}^+ Q_0 \Pi_{2\mu}^+ |\nabla| \right\|_{\mathfrak{S}^1}^{\frac{1}{2}} \\ &\lesssim \left\{ \text{Tr}(-\Delta - \mu) Q_0 \right\}^{\frac{1}{2}} \lesssim \text{Tr}(-\Delta - \mu) Q_0 + 1. \end{aligned}$$

Hence, complex interpolation yields $\left\| |\nabla|^{\frac{1}{2}} \Pi_{2\mu}^+ Q_0 \Pi_{2\mu}^+ |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}} \lesssim \text{Tr}(-\Delta - \mu) Q_0 + 1$. On the other hand, for the high-low component, by (3.5), we have

$$\left\| |\nabla|^{\frac{1}{2}} \Pi_{2\mu}^+ Q_0 \Pi_{2\mu}^- |\nabla|^{\frac{1}{2}} \right\|_{\mathcal{H}^{\frac{1+2\epsilon}{4}}} \lesssim \mu^{\frac{1}{4}} \langle \mu \rangle^{\frac{1+2\epsilon}{8}} \left\| |\Delta + \mu|^{\frac{1}{2}} Q_0 \right\|_{\mathfrak{S}^2} \lesssim \text{Tr}(-\Delta - \mu) Q_0 + 1 + \mu^{\frac{3+2\epsilon}{4}}.$$

Therefore, (4.8) follows.

For the low frequency contribution, using the formula

$$S_V(t, t_0) = e^{i(t-t_0)\Delta} - i \int_{t_0}^t S_V(t, t_1) V(t_1) e^{i(t_1-t_0)\Delta} dt_1,$$

we write $S_V(t, t_0)_* \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^-$ as

$$\begin{aligned} & e^{i(t-t_0)\Delta} \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- \left(e^{-i(t-t_0)\Delta} + i \int_{t_0}^t e^{-i(t_1-t_0)\Delta} V(t_1) S_V(t, t_1)^* dt_1 \right) \\ & + \left(-i \int_{t_0}^t S_V(t, t_1) V(t_1) e^{i(t_1-t_0)\Delta} dt_1 \right) \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- S_V(t, t_0)^*. \end{aligned} \quad (4.9)$$

Thus, we have

$$\begin{aligned} \|\rho_{S_V(t, t_0)_* \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^-}\|_{\mathcal{D}^{1+\epsilon}(I)} &\leq \|\rho_{e^{i(t-t_0)\Delta} \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- e^{-i(t-t_0)\Delta}}\|_{\mathcal{D}^{1+\epsilon}(I)} \\ &\quad + \int_I \|\rho_{e^{i(t-t_0)\Delta} \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- e^{-i(t_1-t_0)\Delta} V(t_1) S_V(t, t_1)^*}\|_{\mathcal{D}^{1+\epsilon}(I)} dt_1 \\ &\quad + \int_I \|\rho_{S_V(t, t_1) V(t_1) e^{i(t_1-t_0)\Delta} \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- S_V(t, t_0)^*}\|_{\mathcal{D}^{1+\epsilon}(I)} dt_1 \\ &=: A_\mu + B_\mu + C_\mu. \end{aligned}$$

For A_μ , we note that the Fourier transform of $\rho_{e^{i(t-t_0)\Delta} \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- e^{-i(t-t_0)\Delta}}$ is supported in frequency $|\xi| \leq 2\sqrt{2\mu}$ and that

$$0 \leq \Pi_\mu^- + e^{i(t-t_0)\Delta} \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- e^{-i(t-t_0)\Delta} = e^{i(t-t_0)\Delta} \Pi_{2\mu}^- \gamma_0 \Pi_{2\mu}^- e^{-i(t-t_0)\Delta} \leq 1.$$

Hence, it follows from the inequality (3.8) with $0 \leq T \leq 1$ that

$$\begin{aligned} A_\mu &\lesssim \langle \mu \rangle^{\frac{1+\epsilon}{2}} \|\rho_{e^{i(t-t_0)\Delta} \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- e^{-i(t-t_0)\Delta}}\|_{L_t^\infty(I; L_x^2)} \\ &\lesssim \langle \mu \rangle^{\frac{1+\epsilon}{2}} \mu^{\frac{1}{4}} \{\text{Tr}(-\Delta - \mu) Q_0\}^{\frac{1}{2}} \lesssim \text{Tr}(-\Delta - \mu) Q_0 + 1 + \mu^{\frac{3}{2}+\epsilon}. \end{aligned}$$

For B_μ and C_μ , by Lemma 4.1 and Lemma 4.4, we obtain that

$$\begin{aligned} B_\mu + C_\mu &\lesssim \|e^{i(t_1-t_0)\Delta} \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- e^{-i(t_1-t_0)\Delta} V(t_1)\|_{L_{t_1}^1(I; \mathcal{H}^{\frac{3+2\epsilon}{4}})} \\ &\quad + \|V(t_1) e^{i(t_1-t_0)\Delta} \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- S_V(t_1, t_0)^*\|_{L_{t_1}^1(I; \mathcal{H}^{\frac{3+2\epsilon}{4}})} \\ &\lesssim \|\langle \nabla \rangle^{\frac{3+2\epsilon}{4}} V(t) e^{i(t-t_0)\Delta} \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- \langle \nabla \rangle^{\frac{3+2\epsilon}{4}}\|_{L_t^1(I; \mathfrak{S}^2)} \\ &\lesssim \|\langle \nabla \rangle^{\frac{3+2\epsilon}{4}} V(t) \Pi_{2\mu}^- \|_{L_t^1(I; \mathfrak{S}^2)} \|Q_0\|_{\mathcal{L}(L^2)} \|\Pi_{2\mu}^- \langle \nabla \rangle^{\frac{3+2\epsilon}{4}}\|_{\mathcal{L}(L^2)} \\ &\lesssim (1 + \mu)^{\frac{3+2\epsilon}{8}} \|\langle \nabla \rangle^{\frac{3+2\epsilon}{4}} V(t) \Pi_{2\mu}^- \|_{L_t^1(I; \mathfrak{S}^2)}. \end{aligned}$$

Note that by direct calculation, the Fourier transform of the kernel of $V \Pi_{2\mu}^-$ is given by $\hat{V}(\xi + \xi') \mathbb{1}_{(|\xi'|^2 \leq 2\mu)}$. Hence, by the Plancherel theorem, it follows that

$$\|\langle \nabla_x \rangle^{\frac{3+2\epsilon}{4}} (V \Pi_{2\mu}^-)(x, x')\|_{L_{x, x'}^2} \sim \|\langle \xi \rangle^{\frac{3+2\epsilon}{4}} \hat{V}(\xi + \xi') \mathbb{1}_{(|\xi'|^2 \leq 2\mu)}\|_{L_{\xi, \xi'}^2} \lesssim (1 + \mu)^{\frac{9+2\epsilon}{8}} \|V\|_{H_x^{1+\epsilon}},$$

and so

$$B_\mu + C_\mu \lesssim (1 + \mu)^{\frac{3+\epsilon}{2}} |I|^{\frac{1}{2}} \|V\|_{L_t^2(I; H_x^\epsilon)} \leq (1 + \mu)^{\frac{3+\epsilon}{2}} |I|^{\frac{1}{2}} R \lesssim (1 + \mu)^{\frac{3+\epsilon}{2}} |I|^{\frac{1-2\epsilon}{4}}.$$

Collecting all, we prove the first inequality (4.4).

For the difference, by (4.9), we write

$$\begin{aligned} & S_V(t, t_0)_* \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- - S_{V'}(t, t_0)_* \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- \\ &= i \int_{t_0}^t e^{i(t-t_0)\Delta} \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- e^{-i(t_1-t_0)\Delta} (V(t_1) S_V(t, t_1)^* - V'(t_1) S_{V'}(t, t_1)^*) dt_1 \\ &\quad - i \int_{t_0}^t (S_V(t, t_1) V(t_1) - S_{V'}(t, t_1) V'(t_1)) e^{i(t_1-t_0)\Delta} \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- S_V(t, t_0)^* dt_1 \\ &\quad - i \int_{t_0}^t S_{V'}(t, t_1) V'(t_1) e^{i(t_1-t_0)\Delta} \Pi_{2\mu}^- Q_0 \Pi_{2\mu}^- (S_V(t, t_0)^* - S_{V'}(t, t_0)^*) dt_1, \end{aligned}$$

and expand further using

$$S_V(t, t_1) V(t_1) - S_{V'}(t, t_1) V'(t_1) = (S_V(t, t_1) - S_{V'}(t, t_1)) V(t_1) + S_{V'}(t, t_1) (V - V')(t_1)$$

in the integral. Then, repeating the above estimates for B_μ and C_μ , one can prove the second inequality (4.5). \square

4.3. Finite relative entropy initial data. Finally, we establish the density estimates when the initial state has finite relative entropy.

Proposition 4.5 (Density function estimates for linear flows III). *Let $R \geq 1$. Suppose that $t_0 \in I$ and $0 < |I| \ll R^{-\frac{4}{1+2\epsilon}}$ for sufficiently small $\epsilon > 0$. Then, for any $\gamma_0 = \gamma_f + Q_0 \in \mathcal{K}_f$, $V, V' \in \mathcal{B}_R^{1+\epsilon}(I)$, we have*

$$\|\rho_{S_V(t, t_0)_* Q_0}\|_{\mathcal{D}^{1+\epsilon}(I)} \leq c_2 \{ \mathcal{H}(\gamma_0 | \gamma_f) + 1 + \|\mu^{\frac{3}{2}+\epsilon}\|_{L_\lambda^1(0,1)} \}$$

and

$$\begin{aligned} & \|\rho_{S_V(t, t_0)_* Q_0} - \rho_{S_{V'}(t, t_0)_* Q_0}\|_{\mathcal{D}^{1+\epsilon}(I)} \\ & \leq c_2 |I|^{\frac{1+2\epsilon}{4}} \{ \mathcal{H}(\gamma_0 | \gamma_f) + 1 + \|\mu^{\frac{3}{2}+\epsilon}\|_{L_\lambda^1(0,1)} \} \|V - V'\|_{L_t^2(I; H_x^{1+\epsilon})}. \end{aligned}$$

Proof. Applying Lemma 4.3 to the integral representation of an initial perturbation

$$S_V(t, t_0)_* Q_0 = \int_0^1 S_V(t, t_0)_* Q_0^{(\mu)} d\lambda, \quad (4.10)$$

where $\mu = S'(\lambda)$, $Q_0^{(\mu)} = \mathbf{1}_{(\gamma_0 \geq \lambda)} - \Pi_\mu^-$ and $\Pi_\mu^- = \mathbf{1}_{(\gamma_f \geq \lambda)}$, we obtain

$$\begin{aligned} \|\rho_{S_V(t, t_0)_* Q_0}\|_{\mathcal{D}^{1+\epsilon}(I)} & \leq \int_0^1 \|\rho_{S_V(t, t_0)_* Q_0^{(\mu)}}\|_{\mathcal{D}^{1+\epsilon}(I)} d\lambda \\ & \leq c_2 \int_0^1 \{ \text{Tr}(-\Delta - \mu) Q_0^{(\mu)} + 1 + \mu^{\frac{3}{2}+\epsilon} \} d\lambda \end{aligned}$$

and

$$\begin{aligned}
& \|\rho_{S_V(t,t_0)\star Q_0} - \rho_{S_{V'}(t,t_0)\star Q_0}\|_{\mathcal{D}^{1+\epsilon}(I)} \\
& \leq \int_0^1 \|\rho_{S_V(t,t_0)\star Q_0^{(\mu)}} - \rho_{S_{V'}(t,t_0)\star Q_0^{(\mu)}}\|_{\mathcal{D}^{1+\epsilon}(I)} d\lambda \\
& \leq c_2 |I|^{\frac{1+2\epsilon}{4}} \|V - V'\|_{L_t^2(I; H_x^{1+\epsilon})} \int_0^1 \{\text{Tr}(-\Delta - \mu)Q_0^{(\mu)} + 1 + \mu^{\frac{3}{2}+\epsilon}\} d\lambda.
\end{aligned}$$

Therefore, the proposition follows from Lemma 3.4. \square

5. LOCAL WELL-POSEDNESS OF THE NLH WITH FINITE RELATIVE ENTROPY INITIAL DATA

In this section, we establish the local well-posedness of the nonlinear Hartree equation for perturbations around a reference state of infinite particles. For mathematical formulation, we employ the linear propagator $S_{w*\rho_Q}(t, t_1)$ to write the perturbation equation (1.6) with initial data Q_0 as

$$Q(t) = S_{w*\rho_Q}(t, 0)\star Q_0 - i \int_0^t S_{w*\rho_Q}(t, t_1)\star [w * \rho_Q(t_1), \gamma_f] dt_1. \quad (5.1)$$

Then, taking density, we derive an equation

$$g(t) = \rho_{S_{w*g}(t, 0)\star Q_0} - i \rho \left(\int_0^t S_{w*g}(t, t_1)\star [w * g(t_1), \gamma_f] dt_1 \right) \quad (5.2)$$

with only one type of unknown as a density function. Using this equation, we establish the following local well-posedness.

Proposition 5.1 (Local well-posedness of the Hartree equation). *Let $\epsilon > 0$ be sufficiently small. Suppose that $S : [0, 1] \rightarrow \mathbb{R}^+$ is \mathcal{H} -admissible, and set $f := (S')^{-1}$. Moreover, we assume that*

$$\|\langle \xi \rangle^{\frac{3}{2}+\epsilon} f(|\xi|^2)\|_{L_\xi^2} < \infty. \quad (5.3)$$

Let w be a finite measure on \mathbb{R}^3 . Take large

$$R \geq 2c_2(1 + \|w\|_{\mathcal{M}}) \left(\mathcal{H}(\gamma_0|\gamma_f) + 1 + \|\mu^{\frac{3}{2}+\epsilon}\|_{L^1(0,1)} \right) + 1, \quad (5.4)$$

where $\mu = S'(\lambda)$ and c_2 is a constant given in Proposition 4.5. Then, for any $\gamma_0 = Q_0 + \gamma_f \in \mathcal{K}_f$, we have the following.

- (i) (Local well-posedness) There exist $0 < T \ll R^{-\frac{4}{1+2\epsilon}}$, depending only on R and $\|\langle \xi \rangle^{\frac{3+\epsilon}{2}} f(|\xi|^2)\|_{L^2}$ (see (5.8) below), and a unique solution $g \in L_t^2(I; H_x^{1+\epsilon})$ to the density function equation (5.2) such that $\|g\|_{L_t^2(I; H_x^{1+\epsilon})} \leq R$, where $I = [-T, T]$.
- (ii) (Construction of solution to NLH) Let g be the solution to (5.2) in (i). Then,

$$Q(t) := S_{w*g}(t, 0)\star Q_0 - i \int_0^t S_{w*g}(t, t_1)\star [w * g(t_1), \gamma_f] dt_1 \in C_t(I; \mathcal{H}^{\frac{1}{2}})$$

is a unique solution to the perturbation equation (5.1) such that $\|\rho_Q\|_{L_t^2(I; H_x^{1+\epsilon})} \leq R$.

Remark 5.2. The assumption (5.3) is sufficient for Strichartz estimates in Proposition 4.5. Indeed, by Hölder's inequality, we have

$$\|\langle \xi \rangle^{\frac{\epsilon}{2}} f(|\xi|^2)\|_{L^1_\xi(\mathbb{R}^3)} \leq \|\langle \xi \rangle^{-\frac{3+\epsilon}{2}}\|_{L^2_\xi(\mathbb{R}^3)} \|\langle \xi \rangle^{\frac{3+\epsilon}{2}} f(|\xi|^2)\|_{L^2_\xi(\mathbb{R}^3)} \lesssim \|\langle \xi \rangle^{\frac{3+\epsilon}{2}} f(|\xi|^2)\|_{L^2_\xi(\mathbb{R}^3)},$$

but one can also check that by changing variables, $\|\langle \xi \rangle^{\frac{\epsilon}{2}} f(|\xi|^2)\|_{L^1_\xi} < \infty$ if and only if $\|\mu^{\frac{3}{2}+\frac{\epsilon}{4}}\|_{L^1_\lambda(0,1)} < \infty$. Therefore, from the assumption (5.3), replacing ϵ by $\frac{\epsilon}{4}$, we obtain $\|\mu^{\frac{3}{2}+\epsilon}\|_{L^1_\lambda(0,1)} < \infty$ and $\|\langle \xi \rangle^{\frac{3}{2}+4\epsilon} f(|\xi|^2)\|_{L^2_\xi} < \infty$.

Proof. For (i), we aim to show that

$$\Phi(g)(t) := \rho_{S_{w*g}(t,0)_*Q_0} - i\rho \left(\int_0^t S_{w*g}(t,t_1)_*[w * g(t_1), \gamma_f] dt_1 \right)$$

is contractive on the ball $\mathcal{B}_R^{1+\epsilon}(I) := \{g : \|g\|_{\mathcal{D}^{1+\epsilon}(I)} \leq R\}$, where $I = [-T, T]$ and $T > 0$ is a small number to be chosen later. Indeed, for $g \in \mathcal{B}_R^{1+\epsilon}(I)$, applying Proposition 4.5 for the first term and Lemma 4.1 for the integral term in $\Phi(g)$, we have

$$\|\Phi(g)\|_{\mathcal{D}^{1+\epsilon}(I)} \leq \frac{R}{2} + 2c_1 \|(w * g)\gamma_f\|_{L^1_t(I; \mathcal{H}^{\frac{3+2\epsilon}{4}})}.$$

On the right hand side, by the fractional Leibniz rule and the Plancherel theorem,

$$\begin{aligned} \|(w * g)\gamma_f\|_{\mathcal{H}^{\frac{3+2\epsilon}{4}}} &= \|\langle \nabla_x \rangle^{\frac{3+2\epsilon}{4}} ((w * g)\gamma_f \langle \nabla_x \rangle^{-\frac{3+2\epsilon}{4}})(x, x')\|_{L^2_{x'} L^2_x} \\ &\lesssim \|g\|_{H_x^{1+\epsilon}} \|\langle \xi \rangle^{\frac{3}{2}+\epsilon} f(|\xi|^2)\|_{L^2_\xi}, \end{aligned} \quad (5.5)$$

since $\|w\|_{\mathcal{M}} < \infty$. Hence, it follows that

$$\|\Phi(g)\|_{\mathcal{D}^{1+\epsilon}(I)} \leq \frac{R}{2} + \tilde{c}_1 T^{\frac{1}{2}} R \|\langle \xi \rangle^{\frac{3}{2}+\epsilon} f(|\xi|^2)\|_{L^2_\xi}. \quad (5.6)$$

Similarly, by Proposition 4.5 and Lemma 4.1, we estimate the difference

$$\begin{aligned} \Phi(g)(t) - \Phi(g')(t) &= \rho_{S_{w*g}(t,0)_*Q_0} - \rho_{S_{w*g'}(t,0)_*Q_0} \\ &\quad - i\rho \left(\int_0^t \{S_{w*g}(t,t_1)_*[w * g(t_1), \gamma_f] - S_{w*g'}(t,t_1)_*[w * g(t_1), \gamma_f]\} dt_1 \right) \\ &\quad - i\rho \left(\int_0^t S_{w*g'}(t,t_1)_*[w * (g - g')(t_1), \gamma_f] dt_1 \right) \end{aligned}$$

with $g, g' \in \mathcal{B}_R(I)$. Then, one can show that

$$\begin{aligned} &\|\Phi(g) - \Phi(g')\|_{\mathcal{D}^{1+\epsilon}(I)} \\ &\leq \frac{1}{2} \|g - g'\|_{L^2_t(I; H_x^{1+\epsilon})} + c_1 T^{\frac{1+2\epsilon}{4}} \|(w * g)\gamma_f\|_{L^1_t(I; \mathcal{H}^{\frac{3+2\epsilon}{4}})} \|g - g'\|_{L^2_t(I; H_x^{1+\epsilon})} \\ &\quad + c_1 T^{\frac{1+2\epsilon}{4}} \|(w * (g - g'))\gamma_f\|_{L^1_t(I; \mathcal{H}^{\frac{3+2\epsilon}{4}})} \\ &\leq \left\{ \frac{1}{2} + \tilde{c}_1 T^{\frac{1+2\epsilon}{4}} (R + 1) T^{\frac{1}{2}} \|\langle \xi \rangle^{\frac{3}{2}+\epsilon} f(|\xi|^2)\|_{L^2_\xi} \right\} \|g - g'\|_{L^2_t(I; H_x^{1+\epsilon})}, \end{aligned} \quad (5.7)$$

where (5.5) is used in the last step. Therefore, taking

$$T = \min \left\{ \frac{1}{16\tilde{c}_1^2 \| \langle \xi \rangle^{\frac{3}{2} + \epsilon} f(|\xi|^2) \|_{L_\xi^2}^2}, (R+1)^{-\frac{4}{1+2\epsilon}} \right\} \quad (5.8)$$

in (5.6) and (5.7), we conclude that $\Phi(g)$ is contractive on $\mathcal{B}_R^{1+\epsilon}(I)$.

For (ii), using the fixed point $g \in \mathcal{B}_R^{1+\epsilon}(I)$ such that $g = \Phi(g)$, we construct a quantum state

$$Q_g(t) := S_{w*g}(t, 0)_* Q_0 - i \int_0^t S_{w*g}(t, t_1)_* [w * g(t_1), \gamma_f] dt_1. \quad (5.9)$$

By the completely same argument as the proof of [26, Lemma 2.1], we have $Q(t) \in C(I; \mathcal{H}^{\frac{1}{2}})$. Next, taking the density of the equation (5.9) and repeating the estimates in the proof of (i), one can show that ρ_{Q_g} is a fixed point for Φ in $\mathcal{B}_R^{1+\epsilon}(I)$. Then, by uniqueness for the density equation (5.2), it implies that $g = \rho_{Q_g}$ and Q_g is the solution to the operator equation (5.1). For uniqueness, we assume that $\tilde{Q}(t)$ is a solution to (5.1) with the same initial data Q_0 such that $\|\rho_{\tilde{Q}}\|_{\mathcal{D}^{1+\epsilon}(I)} < \infty$. Then, by the standard argument, it follows that $\rho_{\tilde{Q}}$ is also a fixed point for (5.2) in $\mathcal{B}_R^{1+\epsilon}(I)$. Therefore, we conclude that $\rho_{\tilde{Q}} = \rho_Q$ and $\tilde{Q} = Q$ in $C_t(I; \mathcal{H}^{\frac{1}{2}})$. \square

6. GLOBAL WELL-POSEDNESS OF THE NLH: PROOF OF THE MAIN THEOREM

In this last section, we show that the local-in-time solution $\gamma(t)$ to the NLH (1.1) with initial data $\gamma_0 = \gamma_f + Q_0 \in \mathcal{K}_f$, obtained in the previous section, preserves the relative free energy $\mathcal{F}_f(\gamma(t)|\gamma_f) = \mathcal{F}_f(\gamma_0|\gamma_f)$, and then we employ the conservation law to establish its global existence. For the proof, we follow the strategy in Lewin-Sabin [31]; we construct a sequence $\{Q_n(t)\}_{n=1}^\infty$ of solutions to the regularized equation

$$\begin{cases} i\partial_t Q_n = [-\Delta + w_n * \rho_{Q_n}, Q_n + \gamma_{f_n}], \\ Q_n(0) = Q_{0;n} \end{cases} \quad (6.1)$$

obeying the relative free energy conservation law, and we prove the conservation law for $\gamma(t)$ sending $n \rightarrow \infty$.

6.1. Choice of the regularized equation. First, we take a sequence of $\{w_n\}_{n=1}^\infty$ of regular interaction potentials such that

$$C_c^\infty(\mathbb{R}^3) \ni w_n \rightarrow w \quad \text{in } \mathcal{M}(\mathbb{R}^3). \quad (6.2)$$

For \mathcal{H} -admissible $S : [0, 1] \rightarrow \mathbb{R}^+$, it is shown in [31, Lemma 10] that there exists a sequence $\{S_n\}_{n=1}^\infty$ of \mathcal{H} -admissible functions on $[0, 1]$ such that

$$S'_n(x) \leq S'(x) \quad \text{on } [0, 1], \quad (6.3)$$

$$S_n^{(j)}(x) \rightarrow S^{(j)}(x) \quad \text{uniformly on any compact subset of } (0, 1) \text{ for } j = 1, 2, \quad (6.4)$$

and

$$\lim_{n \rightarrow \infty} \|f_n(|\xi|^2) - f(|\xi|^2)\|_{L^1(\mathbb{R}^3)} = \lim_{n \rightarrow \infty} \|\langle \xi \rangle^{\frac{3}{2}+\epsilon} (f_n(|\xi|^2) - f(|\xi|^2))\|_{L^2(\mathbb{R}^3)} = 0, \quad (6.5)$$

where $f = (S')^{-1}$ and $f_n = (S'_n)^{-1}$. Note that (6.3) implies $f(\lambda) \leq f_n(\lambda)$ for all λ .

A sequence $\{Q_{0;n}\}_{n=1}^\infty$ of smooth finite-rank initial perturbations is also constructed in [31, Lemma 10], but we include an improved convergence for density functions (see (6.6) below).

Lemma 6.1 (Construction of initial data sequence). *Suppose that $S : [0, 1] \rightarrow \mathbb{R}^+$ is \mathcal{H} -admissible, $\{S_n\}_{n=1}^\infty$ satisfies (6.3) and (6.5), and $\gamma_0 = \gamma_f + Q_0 \in \mathcal{K}_f$. For $R \geq 1$, we assume that $\|V\|_{L_t^2(I; H_x^{1+\epsilon})} \leq R$, $I \ni t_0$ and $0 < |I| \ll R^{-\frac{4}{1+2\epsilon}}$. Then, there exist a sequence $\{Q_{0;n}\}_{n=1}^\infty$ of finite-rank smooth operators such that the followings hold:*

$$\lim_{n \rightarrow \infty} \|\rho_{S_V(t, t_0)_* Q_0} - \rho_{S_V(t, t_0)_* Q_{0;n}}\|_{\mathcal{D}^{1+\epsilon}(I)} + \|Q_0 - Q_{0;n}\|_{\mathfrak{S}^2} = 0, \quad (6.6)$$

$$\lim_{n \rightarrow \infty} \mathcal{H}_{S_n}(\gamma_{f_n} + Q_{0;n} | \gamma_{f_n}) = \mathcal{H}_S(\gamma_f + Q_0 | \gamma_f). \quad (6.7)$$

Proof. (Step 1: Proof of (6.7)) By Lemma 3.3, there exists a sequence $\{\tilde{Q}_{0;n}\}_{n=1}^\infty$ of finite-rank smooth operators such that $\mathcal{H}_S(\gamma_f + \tilde{Q}_{0;n} | \gamma_f) \rightarrow \mathcal{H}_S(\gamma_f + Q_0 | \gamma_f)$ and $0 \leq \gamma_f + \tilde{Q}_{0;n} \leq 1$. Then, for $Q_{0;n} := X_n \tilde{Q}_{0;n} X_n$ with $X_n := (\gamma_{f_n} / \gamma_f)^{\frac{1}{2}}$ by (6.3), we have

$$0 \leq \gamma_{f_n} + Q_{0;n} = X_n(\gamma_f + \tilde{Q}_{0;n})X_n \leq X_n^2 \leq 1. \quad (6.8)$$

By the monotonicity of the relative entropy (Theorem 3.1 (i)) and the construction of S_n (see the proof of [30, Lemma 10]), it follows that

$$\begin{aligned} \mathcal{H}_S(\gamma_f + \tilde{Q}_{0;n} | \gamma_f) &\geq \mathcal{H}_S(\gamma_{f_n} + Q_{0;n} | \gamma_{f_n}) = \mathcal{H}_{S_n}(\gamma_{f_n} + Q_{0;n} | \gamma_{f_n}) + \mathcal{H}_{S-S_n}(\gamma_{f_n} + Q_{0;n} | \gamma_{f_n}) \\ &\geq \mathcal{H}_{S_n}(\gamma_{f_n} + Q_{0;n} | \gamma_{f_n}), \end{aligned}$$

and consequently,

$$\limsup_{n \rightarrow \infty} \mathcal{H}_{S_n}(\gamma_{f_n} + Q_{0;n} | \gamma_{f_n}) \leq \mathcal{H}_S(\gamma_f + Q_0 | \gamma_f).$$

On the other hand, by the weak lower semi-continuity (see Theorem 3.1), we have

$$\begin{aligned} \mathcal{H}_S(\gamma_f + Q | \gamma_f) &\leq \liminf_{n \rightarrow \infty} \mathcal{H}_S(\gamma_{f_n} + Q_{0;n} | \gamma_{f_n}) \\ &= \liminf_{n \rightarrow \infty} \{\mathcal{H}_{S_n}(\gamma_{f_n} + Q_{0;n} | \gamma_{f_n}) + \mathcal{H}_{\varphi_n}(\gamma_{f_n} + Q_{0;n} | \gamma_{f_n})\}, \end{aligned}$$

where $\varphi_n := S - S_n$. For the second term, we note that by (6.8), the monotonicity and Lemma 3.3,

$$0 \leq \mathcal{H}_{\varphi_n}(\gamma_{f_n} + Q_{0;n} | \gamma_{f_n}) \leq \mathcal{H}_{\varphi_n}(\gamma_f + \tilde{Q}_{0;n} | \gamma_f).$$

Hence, it suffices to show that $\mathcal{H}_{\varphi_n}(\gamma_f + \tilde{Q}_{0;n} | \gamma_f) \rightarrow 0$. Recalling the definitions of Π'_j and P_k and the construction of approximating operators from the proof of Lemma 3.3, we write $\tilde{Q}_{0;n} = \Pi'_{1/j_n} P_{j_n} Q_0 P_{j_n} \Pi'_{1/j_n}$ with $j_n \rightarrow \infty$. Hence, we have

$$\mathcal{H}_{\varphi_n}(\gamma_f + \tilde{Q}_{0;n} | \gamma_f) = \mathcal{H}_{\varphi_n}(\Pi'_{1/j_n} (\gamma_f + P_{j_n} Q_0 P_{j_n}) \Pi'_{1/j_n} | \Pi'_{1/j_n} \gamma_f \Pi'_{1/j_n}).$$

Note that [17, Theorem 4] enables us to write

$$\begin{aligned} & \mathcal{H}_{\varphi_n}(\Pi'_{1/j_n} \gamma \Pi'_{1/j_n} | \Pi'_{1/j_n} \gamma_f \Pi'_{1/j_n}) \\ &= \text{Tr} \left(\varphi_n(\Pi'_{1/j_n} \gamma \Pi'_{1/j_n}) - \varphi_n(\Pi'_{1/j_n} \gamma_f \Pi'_{1/j_n}) - \varphi'_n(\Pi'_{1/j_n} \gamma_f \Pi'_{1/j_n}) \Pi'_{1/j_n} P_{j_n} Q_0 P_{j_n} \Pi'_{1/j_n} \right), \end{aligned}$$

where $\gamma = \gamma_f + P_{j_n} Q_0 P_{j_n}$. By replacing $(\varphi_n)_{n=1}^\infty$ by its subsequence if necessary, we may assume that

$$\sup_{t \in [S'(1 - \frac{1}{j_n}), S'(\frac{1}{j_n})]} |\varphi_n(t)| + |\varphi'_n(t)| + |\varphi''_n(t)| \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

sufficiently rapidly. There exist $\tilde{\varphi}_n \in C_c^\infty((0, 1))$ satisfying $\varphi_n = \tilde{\varphi}_n$ on $[S'(1 - \frac{1}{j_n}), S'(\frac{1}{j_n})]$.

Then, $\tilde{\varphi}_n$ satisfies $\tilde{\varphi}_n^{(j)} \rightarrow 0$ uniformly on \mathbb{R} for $j = 0, 1, 2$ sufficiently rapidly. On the one hand, by (6.4), we get

$$\left| \text{Tr} \left(\tilde{\varphi}'_n(\Pi'_{1/j_n} \gamma_f \Pi'_{1/j_n}) \Pi'_{1/j_n} P_{j_n} Q_0 P_{j_n} \Pi'_{1/j_n} \right) \right| \leq \|\tilde{\varphi}'_n\|_{L^\infty(\mathbb{R})} \|P_{j_n} Q_0 P_{j_n}\|_{\mathfrak{S}^1} \rightarrow 0$$

as $n \rightarrow \infty$. On the other hand, by [2, Theorem 1.6.1], we obtain

$$\left| \text{Tr} \left(\tilde{\varphi}_n(\Pi'_{1/j_n} \gamma \Pi'_{1/j_n}) - \tilde{\varphi}_n(\Pi'_{1/j_n} \gamma_f \Pi'_{1/j_n}) \right) \right| \lesssim \|\tilde{\varphi}_n\|_{B_{\infty,1}^1(\mathbb{R})} \|P_{j_n} Q_0 P_{j_n}\|_{\mathfrak{S}^1} \rightarrow 0$$

as $n \rightarrow \infty$, since $\|\tilde{\varphi}_n\|_{B_{\infty,1}^1(\mathbb{R})} \lesssim \|\tilde{\varphi}_n\|_{L^\infty(\mathbb{R})} + \|\tilde{\varphi}'_n\|_{L^\infty(\mathbb{R})} \rightarrow 0$.

(Step 2: Proof of (6.6)) First, we write

$$\begin{aligned} \|\rho_{S_V(t, t_0)_*}(Q_0 - Q_{0;n})\|_{\mathcal{D}^{1+\epsilon}(I)} &\leq \|\rho_{S_V(t, t_0)_*}(Q_0 - \Pi_{k_n}^- Q_0 \Pi_{k_n}^-)\|_{\mathcal{D}^{1+\epsilon}(I)} \\ &\quad + \|\rho_{S_V(t, t_0)_*}(\Pi_{k_n}^- (Q_0 - Q_{0;n}) \Pi_{k_n}^-)\|_{\mathcal{D}^{1+\epsilon}(I)} \\ &\quad + \|\rho_{S_V(t, t_0)_*}(\Pi_{k_n}^- Q_{0;n} \Pi_{k_n}^- - Q_{0;n})\|_{\mathcal{D}^{1+\epsilon}(I)} \\ &=: (\text{I})_n + (\text{II})_n + (\text{III})_n, \end{aligned}$$

where $\{k_n\}_{n=1}^\infty$, with $k_n \rightarrow \infty$, is a sequence to be chosen later. For the first term, by the representation (4.10), we have

$$(\text{I})_n \leq \int_{n \geq 2\mu} + \int_{n < 2\mu} \|\rho_{S_V(t, t_0)_*}(Q_0^{(\mu)} - \Pi_n^- Q_0^{(\mu)} \Pi_n^-)\|_{\mathcal{D}^{1+\epsilon}(I)} d\lambda =: A_n + B_n,$$

where $Q_0^{(\mu)} = \mathbb{1}_{(\gamma_0 \geq \lambda)} - \mathbb{1}_{(\gamma_f \geq \lambda)}$, $\Pi_\mu^- = \mathbb{1}_{(\gamma_f \geq \lambda)}$ and $\mu = S'(\lambda)$. For A_n , applying Lemma 4.1 to the decomposition

$$\begin{aligned} Q_0^{(\mu)} - \Pi_n^- Q_0^{(\mu)} \Pi_n^- &= \Pi_n^+ Q_0^{(\mu)} \Pi_n^+ + \Pi_n^- \Pi_{2\mu}^+ Q_0^{(\mu)} \Pi_n^+ + \Pi_{2\mu}^- Q_0^{(\mu)} \Pi_n^+ \\ &\quad + \Pi_n^+ Q_0^{(\mu)} \Pi_{2\mu}^- + \Pi_n^+ Q_0^{(\mu)} \Pi_{2\mu}^+ \Pi_n^-. \end{aligned}$$

and by symmetry, we obtain that

$$A_n \lesssim \int_{n \geq 2\mu} \|\Pi_n^+ Q_0^{(\mu)} \Pi_n^+\|_{\mathcal{H}^{\frac{3+2\epsilon}{4}}} + \|\Pi_n^- \Pi_{2\mu}^+ Q_0^{(\mu)} \Pi_n^+\|_{\mathcal{H}^{\frac{3+2\epsilon}{4}}} + \|\Pi_{2\mu}^- Q_0^{(\mu)} \Pi_n^+\|_{\mathcal{H}^{\frac{3+2\epsilon}{4}}} d\lambda.$$

We observe that if $n \geq 2\mu$, then

$$\|\Pi_n^+ Q_0^{(\mu)} \Pi_n^+\|_{\mathcal{H}^{\frac{3+2\epsilon}{4}}} \lesssim n^{-\frac{1-2\epsilon}{4}} \text{Tr}(-\Delta - \mu) Q_0^{(\mu)}.$$

Moreover, by cyclicity of the trace and (3.5), we prove that

$$\begin{aligned}
\|\Pi_n^- \Pi_{2\mu}^+ Q_0^{(\mu)} \Pi_n^+\|_{\mathcal{H}^{\frac{1}{2}}} &\lesssim n^{-\frac{1}{4}} \|\Delta + \mu|^{\frac{1}{4}} \Pi_\mu^+ Q_0^{(\mu)} \Pi_\mu^+ |\Delta + \mu|^{\frac{1}{2}}\|_{\mathfrak{S}^2} \\
&= n^{-\frac{1}{4}} \left\{ \text{Tr}(|\Delta + \mu|^{\frac{1}{2}} \Pi_\mu^+ Q_0^{(\mu)} \Pi_\mu^+ |\Delta + \mu|^{\frac{1}{2}} \Pi_\mu^+ Q_0^{(\mu)} \Pi_\mu^+ |\Delta + \mu|^{\frac{1}{2}}) \right\}^{\frac{1}{2}} \\
&\lesssim n^{-\frac{1}{4}} \|\Delta + \mu|^{\frac{1}{2}} \Pi_\mu^+ Q_0^{(\mu)} \Pi_\mu^+ |\Delta + \mu|^{\frac{1}{2}}\|_{\mathfrak{S}^2}^{\frac{1}{2}} \|Q_0^{(\mu)} |\Delta + \mu|^{\frac{1}{2}}\|_{\mathfrak{S}^2}^{\frac{1}{2}} \\
&\lesssim n^{-\frac{1}{4}} \left\{ \text{Tr}(-\Delta - \mu) Q_0^{(\mu)} \right\}^{\frac{3}{4}} \lesssim n^{-\frac{1}{4}} \left\{ \text{Tr}(-\Delta - \mu) Q_0^{(\mu)} + 1 \right\}.
\end{aligned}$$

Thus, by complex interpolating with $\|\Pi_n^- \Pi_{2\mu}^+ Q_0^{(\mu)} \Pi_n^+\|_{\mathcal{H}^1} \lesssim \text{Tr}(-\Delta - \mu) Q_0^{(\mu)}$, we obtain

$$\begin{aligned}
\|\Pi_n^- \Pi_{2\mu}^+ Q_0^{(\mu)} \Pi_n^+\|_{\mathcal{H}^{\frac{3+2\epsilon}{4}}} &\leq \|\Pi_n^- \Pi_{2\mu}^+ Q_0^{(\mu)} \Pi_n^+\|_{\mathcal{H}^{\frac{1}{2}}}^{\frac{1-2\epsilon}{2}} \|\Pi_n^- \Pi_{2\mu}^+ Q_0^{(\mu)} \Pi_n^+\|_{\mathcal{H}^1}^{\frac{1+2\epsilon}{2}} \\
&\lesssim n^{-\frac{1-2\epsilon}{8}} \left\{ \text{Tr}(-\Delta - \mu) Q_0^{(\mu)} + 1 \right\},
\end{aligned}$$

and similarly,

$$\begin{aligned}
\|\Pi_{2\mu}^- Q_0^{(\mu)} \Pi_n^+\|_{\mathcal{H}^{\frac{3+2\epsilon}{4}}} &\lesssim n^{-\frac{1-2\epsilon}{8}} (1 + \mu)^{\frac{3+2\epsilon}{8}} \|Q_0^{(\mu)} |\Delta + \mu|^{\frac{1}{2}}\|_{\mathfrak{S}^2} \\
&\lesssim n^{-\frac{1-2\epsilon}{8}} \left\{ \text{Tr}(-\Delta - \mu) Q_0^{(\mu)} + 1 + \mu^{\frac{3+2\epsilon}{4}} \right\}.
\end{aligned}$$

Therefore, it follows from the lower bound for the relative entropy (Lemma 3.4) and Hölder's inequality that $A_n \rightarrow 0$. For B_n , applying Strichartz estimates (Lemma 4.3), we prove that

$$B_n \lesssim \int_{n < 2\mu} \left(\text{Tr}(-\Delta - \mu) Q_0^{(\mu)} + 1 + \mu^{\frac{3}{2} + \epsilon} \right) d\lambda.$$

Note that $n < 2\mu = 2S'(\lambda)$ if and only if $0 \leq \lambda < (S')^{-1}(\frac{n}{2}) = f(\frac{n}{2}) \rightarrow 0$ as $n \rightarrow \infty$. Therefore, it follows that $B_n \rightarrow 0$.

For (II)_n, by $Q_{0;n} := X_n \tilde{Q}_{0;n} X_n$, the construction of $\tilde{Q}_{0;n}$ in the proof of Lemma 3.3 and the Lemma 4.1, we obtain

$$(II)_n \lesssim \|\Pi_{k_n}^+ Q_{0;n}\|_{\mathcal{H}^{\frac{3+2\epsilon}{4}}} \lesssim |S'(\frac{1}{n})|^{\frac{3}{2} + \epsilon} \|\Pi_{k_n}^+ Q_0\|_{\mathfrak{S}^2} \rightarrow 0$$

taking a sufficiently rapidly increasing sequence $k_n \rightarrow \infty$.

For (III)_n, by choosing j_n and ℓ_n sufficiently large (recall that X_n depends on the choice of ℓ_n), we can assume that

$$(III)_n \lesssim \|\Pi_{k_n}^- (Q_0 - Q_{0;n}) \Pi_{k_n}^-\|_{\mathcal{H}^{\frac{3+2\epsilon}{4}}} \lesssim k_n^{\frac{3}{2} + \epsilon} \|Q_0 - Q_{0;n}\|_{\mathfrak{S}^2} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Therefore, collecting all, we complete the proof. \square

6.2. Proof of global well-posedness. Now, we are ready to prove our main result (Theorem 1.2).

6.2.1. *Construction of approximate solutions.* Suppose that $\gamma_0 = \gamma_f + Q_0 \in \mathcal{K}_f$, and take $\{w_n\}_{n=1}^\infty$, $\{S_n\}_{n=1}^\infty$ and $\{Q_{0;n}\}_{n=1}^\infty$ satisfying (6.2)-(6.5) and the properties in Lemma 6.1. Then, by Proposition 5.1, one can construct the solution $Q(t) \in C_t(I; \mathcal{H}^{\frac{1}{2}})$ to the perturbation equation (1.6) with initial data Q_0 such that $\|\rho_Q\|_{\mathcal{D}^{1+\epsilon}(I)} \leq R$, where $R > 0$ is given by (5.4), $I = [-T, T]$ and $0 < T \ll R^{-\frac{4}{1+2\epsilon}}$. Moreover, one can employ Proposition 5.1 again to show that replacing $T > 0$ by a smaller number independent of $n \gg 1$ if necessary, for large $n \gg 1$, the regularized equation (6.1) has a unique solution $Q_n(t) \in C_t(I; \mathcal{H}^{\frac{1}{2}})$ such that $\|\rho_{Q_n}\|_{\mathcal{D}^{1+\epsilon}(I)} \leq R$.

We claim that for sufficiently large $n \gg 1$, $Q_n(t)$ stays regular on the interval $I = [-T, T]$, where $Q(t)$ exists in $C_t(I; \mathfrak{S}^4)$ and $\|Q\|_{\mathfrak{S}^4} = \|\langle \nabla \rangle^4 Q\|_{\mathfrak{S}^1}$. Indeed, it is obvious that

$$\begin{aligned} \|Q_n(t)\|_{\mathfrak{S}^1} &\leq \|S_{w_n * \rho_{Q_n}}(t) * Q_{0;n}\|_{\mathfrak{S}^1} + \int_0^t \|S_{w_n * \rho_{Q_n}}(t, \tau) * [w_n * \rho_{Q_n}(t_1), \gamma_{f_n}]\|_{\mathfrak{S}^1} dt_1 \\ &\leq \|Q_{0;n}\|_{\mathfrak{S}^1} + 2 \int_0^t \|(w_n * \rho_{Q_n})(t_1) \gamma_f\|_{\mathfrak{S}^1} dt_1, \end{aligned}$$

On the right hand side, by Birman-Solomjak's inequality (see [7], [8]; see also [34, Theorem 4.5]), we have

$$\|(w_n * \rho_{Q_n}) \gamma_{f_n}\|_{\mathfrak{S}^4} \lesssim \|w_n * \rho_{Q_n}\|_{\ell^1 L^2} \|f_n\|_{\ell^1 L^2} \lesssim \|w_n\|_{L^1} \|f_n\|_{\ell^1 L^2} \|Q_n\|_{\mathfrak{S}^1}.$$

Thus, it follows that

$$\|Q_n(t)\|_{\mathfrak{S}^1} \leq \|Q_{0;n}\|_{\mathfrak{S}^1} + \|w_n\|_{L^1} \|f_n\|_{\ell^1 L^2} \int_0^t \|Q_n(t_1)\|_{\mathfrak{S}^1} dt_1,$$

and Gronwall's inequality yields

$$\|Q_n\|_{C_t(I; \mathfrak{S}^1)} \leq C_n. \quad (6.9)$$

Moreover, estimating

$$Q_n(t) = e^{it\Delta} Q_{0;n} e^{-it\Delta} - i \int_0^t e^{i(t-t_1)\Delta} [w_n * \rho_{Q_n}, Q_n + \gamma_{f_n}](t_1) e^{-i(t-t_1)\Delta} dt_1$$

by trivial estimates, we obtain

$$\begin{aligned} \|Q_n(t)\|_{\mathfrak{S}^4} &\leq \|Q_{0;n}\|_{\mathfrak{S}^4} + \int_0^t \|(w_n * \rho_{Q_n}) Q_n(t_1)\|_{\mathfrak{S}^4} + \|Q_n(w_n * \rho_{Q_n})(t_1)\|_{\mathfrak{S}^4} dt_1 \\ &\quad + \int_0^t \|(w_n * \rho_{Q_n})(t_1) \gamma_{f_n} - \gamma_{f_n}(w_n * \rho_{Q_n})(t_1)\|_{\mathfrak{S}^4} dt_1. \end{aligned}$$

Note that for $t_1 \in I$, by (6.9),

$$\begin{aligned} &\|(w_n * \rho_{Q_n}) Q_n(t_1)\|_{\mathfrak{S}^4} + \|Q_n(w_n * \rho_{Q_n})(t_1)\|_{\mathfrak{S}^4} \\ &\lesssim \|\langle \nabla \rangle^4 w_n * \rho_{Q_n}(t_1)\|_{L^\infty} \|Q_n(t_1)\|_{\mathfrak{S}^4} \lesssim \|\langle \nabla \rangle^4 w_n\|_{L^\infty} \|Q_n(t_1)\|_{\mathfrak{S}^1} \|Q_n(t_1)\|_{\mathfrak{S}^4} \\ &\leq C_n \|\langle \nabla \rangle^4 w_n\|_{L^\infty} \|Q_n(t_1)\|_{\mathfrak{S}^4} \end{aligned}$$

and by Birman-Solomjak's inequality as above,

$$\begin{aligned} \|(w_n * \rho_{Q_n})(t_1)\gamma_{f_n} - \gamma_{f_n}(w_n * \rho_{Q_n})(t_1)\|_{\mathfrak{S}^4} &\lesssim \|\langle \nabla \rangle^4 w_n\|_{L^1} \|\langle \nabla \rangle^4 f_n\|_{\ell^1 L^2} \|Q_n(t_1)\|_{\mathfrak{S}^1} \\ &\leq C_n \|\langle \nabla \rangle^4 w_n\|_{L^1} \|\langle \nabla \rangle^4 f_n\|_{\ell^1 L^2}. \end{aligned}$$

Therefore, by Gronwall's inequality, it follows that $\|Q_n\|_{C_t(I; \mathfrak{S}^4)} < \infty$.

6.2.2. *Approximation by Q_n .* Next, we claim that

$$\lim_{n \rightarrow \infty} \|\rho_{Q_n} - \rho_Q\|_{\mathcal{D}^{1+\epsilon}(I)} + \lim_{n \rightarrow \infty} \|Q_n - Q\|_{C_t(I; \mathfrak{S}^2)} = 0, \quad (6.10)$$

$$\mathcal{H}_S(\gamma_f + Q(t)|\gamma_f) \leq \liminf_{n \rightarrow \infty} \mathcal{H}_{S_n}(\gamma_{f_n} + Q_n(t)|\gamma_{f_n}). \quad (6.11)$$

Indeed, by the same argument as Step 2 in the proof of Lemma 6.1, we obtain (6.11). For the difference

$$\begin{aligned} (Q - Q_n)(t) &= (S_{w * \rho_Q}(t, 0) \star Q_0 - S_{w_n * \rho_{Q_n}}(t, 0) \star Q_0) \\ &\quad + S_{w_n * \rho_{Q_n}}(t, 0) \star (Q_0 - Q_{0;n}) \\ &\quad - i \int_0^t \{S_{w * \rho_Q}(t, t_1) \star [w * \rho_Q(t_1), \gamma_f] - S_{w_n * \rho_{Q_n}}(t, t_1) \star [w * \rho_Q(t_1), \gamma_f]\} dt_1 \\ &\quad - i \int_0^t S_{w * \rho_{Q_n}}(t, t_1) \star ([w * \rho_Q(t_1), \gamma_f] - [w_n * \rho_{Q_n}(t_1), \gamma_{f_n}]) dt_1, \end{aligned} \quad (6.12)$$

we apply Proposition 4.5 and Lemma 4.1 again. Then, repeating the estimates in (5.5), one can show that

$$\begin{aligned} \|\rho_{Q_n} - \rho_Q\|_{\mathcal{D}^{1+\epsilon}(I)} &\lesssim o_n(1) + \frac{1}{2} \|\rho_{Q_n} - \rho_Q\|_{L_t^2(I; H_x^\alpha)} \\ &\quad + \|\langle \xi \rangle^{\frac{3}{2}+\epsilon} (f_n - f)(|\xi|^2)\|_{L^2} + \|w_n - w\|_{\mathcal{M}}, \end{aligned}$$

where the implicit constant in the above inequality is independent of $n \gg 1$ but may depend on the quantities R , $\|\rho_Q\|_{\mathcal{D}^{1+\epsilon}(I)}$, $\|w\|_{\mathcal{M}}$, $\|\langle \xi \rangle^{\frac{3}{2}+\epsilon} f(|\xi|^2)\|_{L^2}$ and $\|\mu^{\frac{3}{2}+\epsilon}\|_{L^1(0,1)}$. Therefore, we obtain $\|\rho_{Q_n} - \rho_Q\|_{\mathcal{D}^{1+\epsilon}(I)} \rightarrow 0$. Then, having this density convergence, coming back to (6.12), one can show that $Q_n \rightarrow Q$ in $C_t(I; \mathcal{H}^{\frac{1}{2}})$ by the boundedness of the linear propagator (Lemma 4.4) and the convergence of the initial data $Q_{0;n} \rightarrow Q_0$ in $\mathcal{H}^{\frac{1}{2}}$ (see (6.6)).

6.2.3. *Proof of the conservation law.* For each $t' \in [-T, T]$, the density convergence (6.10) implies the potential energy convergence

$$\iint_{\mathbb{R}^3 \times \mathbb{R}^3} w(x-y) \rho_{Q_n}(t', x) \rho_{Q_n}(t', y) dx dy \rightarrow \iint_{\mathbb{R}^3 \times \mathbb{R}^3} w(x-y) \rho_Q(t', x) \rho_Q(t', y) dx dy.$$

Therefore, by the lower semi-continuity of the relative entropy (6.11) and the conservation of the relative free energy for regular solutions [30, Proposition 7], it follows that

$$\begin{aligned} \mathcal{F}_f(\gamma_f + Q(t')|\gamma_f) &\leq \liminf_{n \rightarrow \infty} \mathcal{F}_{f_n}(\gamma_{f_n} + Q_n(t')|\gamma_{f_n}) \\ &= \lim_{n \rightarrow \infty} \mathcal{F}(\gamma_f + Q_{0;n}|\gamma_f) = \mathcal{F}(\gamma_f + Q_0|\gamma_f). \end{aligned}$$

Next, repeating the same argument to the backward evolution from $t = t'$ to $t = 0$, one can show the reverse inequality $\mathcal{F}(\gamma_f + Q_0|\gamma_f) \leq \mathcal{F}(\gamma_f + Q(t')|\gamma_f) + o_n(1)$. Since t' is arbitrary, these prove the conservation of the relative free energy.

6.2.4. *Proof of global well-posedness.* By the conservation law, we have $\mathcal{F}(\gamma_f + Q_0|\gamma_f) \gtrsim \mathcal{H}(\gamma_f + Q(T)|\gamma_f)$. Then, making the solution evolve forward/backward-in-time from $t = \pm T$, one can show that the solution exists until $t = \pm 2T$ with the relative free energy conservation law. Repeating this, one can extend the existence time arbitrarily long.

APPENDIX A. 1D CASE

The same problem can be formulated in lower dimensional cases ($d = 1, 2$), where the equation (1.1), the perturbation equation (1.6), the relative entropy $\mathcal{H}_S(\gamma|\gamma_f)$ (see (1.3)), the relative free energy $\mathcal{F}_f(\gamma|\gamma_f)$ (see (1.5)) and the operator space \mathcal{K}_f (see (1.4)) are defined by the same way. However, the last condition in \mathcal{H} -admissibility (see Definition 1.1) is replaced by $\|(S')_+\|_{L^{\frac{d}{2}}(0,1)} < \infty$. We refer to Lewin-Sabin [31] for more details.

In this appendix, we provide the statement of the analogous global well-posedness in 1D and give a sketch of the proof.

Theorem A.1. *Let $d = 1$. Suppose that $S : [0,1] \rightarrow \mathbb{R}^+$ is \mathcal{H} -admissible, and that $f := (S')^{-1}$ satisfies $\|\langle \xi \rangle^{\frac{1+\epsilon}{2}} f(|\xi|^2)\|_{L^2_\xi} < \infty$ for small $\epsilon > 0$, and $w \in \mathcal{M}(\mathbb{R})$ is a finite measure on \mathbb{R}^d such that $\|\hat{w}_-\|_{L^\infty} < \frac{K_{\text{LT},S}}{\sqrt{2\pi}}$, where $K_{\text{LT},S}$ is the best constant for the Lieb-Thirring inequality $\mathcal{H}_S(\gamma|\gamma_f) \geq K_{\text{LT},S} \|\rho_{\gamma-\gamma_f}\|_{L^2}^2$ (see [31, Lemma 9]). Then, given initial data $\gamma_0 \in \mathcal{K}_f$, there exists a unique global solution $\gamma(t)$ to the NLH $i\partial_t \gamma = [-\Delta + w * \rho_\gamma, \gamma]$ with initial data γ_0 such that $\gamma(t) \in \mathcal{K}_f$ and $\mathcal{F}_f(\gamma(t)|\gamma_f) = \mathcal{F}_f(\gamma_0|\gamma_f)$ for all $t \in \mathbb{R}$.*

Sketch of Proof. (Step 0. Preliminaries) Repeating the proof of Lemma 2.2, one can show its 1D analogue,

$$\|\rho_{e^{it\Delta}\gamma_0 e^{-it\Delta}}\|_{\mathcal{D}^{\frac{3+\epsilon}{4}}(I)} \lesssim \|\langle \nabla \rangle^{\frac{1+\epsilon}{4}} \gamma_0 \langle \nabla \rangle^{\frac{1+\epsilon}{4}}\|_{\mathfrak{S}^2}, \quad (\text{A.1})$$

where $\|g\|_{\mathcal{D}^s(I)} := \|g\|_{L_t^\infty(I; L_x^2) \cap L_t^2(I; \dot{H}_x^s)}$ and $\|\gamma\|_{\mathcal{H}^\alpha} := \|\langle \nabla \rangle^\alpha \gamma \langle \nabla \rangle^\alpha\|_{\mathfrak{S}^2}$.

(Step 1. Local well-posedness) Let $T > 0$ be a small number to be chosen later. For sufficiently small $\epsilon > 0$, we set

$$\mathcal{B}_R^{\frac{1+\epsilon}{4}}(I) := \left\{ (Q, g) : \|Q\|_{C_t(I; \mathcal{H}^{\frac{1+\epsilon}{4}})} + \|g\|_{\mathcal{D}^{\frac{3+\epsilon}{4}}(I)} \leq R \right\},$$

where $R \gg \|Q_0\|_{\mathcal{H}^{\frac{1+\epsilon}{4}}} + \|w\|_{\mathcal{M}_x} + \|\langle \xi \rangle^{\frac{1+\epsilon}{2}} f(|\xi|^2)\|_{L^2_\xi}$ and $I = [-T, T]$, and define $\Phi = (\Phi_1, \Phi_2)$ by

$$\begin{cases} \Phi_1(Q, g)(t) := e^{it\Delta} Q_0 e^{-it\Delta} - i \int_0^t e^{i(t-t_1)\Delta} [w * g, Q + \gamma_f](t_1) e^{-i(t-t_1)\Delta} dt_1, \\ \Phi_2(Q, g)(t) := \rho_{\Phi_1(Q, g)(t)}. \end{cases}$$

Then, by the unitarity of the free flow, the density estimate in Step 0, and by symmetry for the inhomogeneous terms, it follows that

$$\begin{aligned} & \|\Phi_1(Q, g)\|_{C_t(I; \mathcal{H}^{\frac{1+\epsilon}{4}})} + \|\Phi_2(Q, g)\|_{\mathcal{D}^{\frac{3+\epsilon}{4}}(I)} \\ & \lesssim \|Q_0\|_{\mathcal{H}^{\frac{1+\epsilon}{4}}} + \|(w * g)Q\|_{L_t^1(I; \mathcal{H}^{\frac{1+\epsilon}{4}})} + \|(w * g)\gamma_f\|_{L_t^1(I; \mathcal{H}^{\frac{1+\epsilon}{4}})}. \end{aligned}$$

Note that by the fractional Leibniz rule and the Sobolev inequality,

$$\begin{aligned} \|(w * g)Q\|_{L_t^1(I; \mathcal{H}^{\frac{1+\epsilon}{4}})} &= \|\langle \nabla_x \rangle^{\frac{1+\epsilon}{4}} \langle \nabla_{x'} \rangle^{\frac{1+\epsilon}{4}} ((w * g)(t, x)Q(t, x, x'))\|_{L_t^1(I; L_x^2 L_{x'}^2)} \\ &\lesssim T^{\frac{1}{2}} \|w * g\|_{L_t^2 W_x^{4, \frac{1+\epsilon}{4}}} \|\langle \nabla_{x'} \rangle^{\frac{1+\epsilon}{4}} Q\|_{C_t(I; L_x^4 L_{x'}^2)} \\ &\quad + T^{\frac{1}{2}} \|w * g\|_{L_t^2 L_x^\infty} \|\langle \nabla_x \rangle^{\frac{1+\epsilon}{4}} \langle \nabla_{x'} \rangle^{\frac{1+\epsilon}{4}} Q\|_{C_t(I; L_x^2 L_{x'}^2)} \\ &\lesssim T^{\frac{1}{2}} \|w\|_{\mathcal{M}_x} \|g\|_{\mathcal{D}^{\frac{3+\epsilon}{4}}(I)} \|Q\|_{C_t(I; \mathcal{H}^{\frac{1+\epsilon}{4}})} \end{aligned}$$

and

$$\begin{aligned} \|(w * g)\gamma_f\|_{L_t^1(I; \mathcal{H}^{\frac{1+\epsilon}{4}})} &= \|\langle \nabla_x \rangle^{\frac{1+\epsilon}{4}} \langle \nabla_{x'} \rangle^{\frac{1+\epsilon}{4}} ((w * g)(t, x)f(|\xi|^2)^\vee(x - x'))\|_{L_t^1(I; L_x^2 L_{x'}^2)} \\ &\lesssim T^{\frac{1}{2}} \|w\|_{\mathcal{M}_x} \|g\|_{\mathcal{D}^{\frac{3+\epsilon}{4}}(I)} \|\langle \xi \rangle^{\frac{1+\epsilon}{2}} f(|\xi|^2)\|_{L_\xi^2}. \end{aligned}$$

Therefore, taking $0 < T \ll \frac{1}{R^2}$, we prove that Φ maps from $\mathcal{B}_R^{\frac{1+\epsilon}{4}}(I)$ to itself. Similarly estimating the difference, one can show that Φ is contractive in $\mathcal{B}_R^{\frac{1+\epsilon}{4}}(I)$. Then, we conclude that the perturbation equation $i\partial_t Q = [-\Delta + w * \rho_Q, Q + \gamma_f]$ has a unique strong solution in $C_t(I; \mathcal{H}^{\frac{1+\epsilon}{4}})$. In addition, by repeating the same analysis, one can prove the continuity of map $Q_0 \mapsto (Q, \rho_Q)$ from $\mathcal{H}^{\frac{1+\epsilon}{4}}$ to $\mathcal{B}_R^{\frac{1+\epsilon}{4}}(I)$.

(Step 3. Proof of conservation law and global well-posedness) Once we have the local well-posedness, we can prove the conservation of the relative free energy

$$\mathcal{F}_f(\gamma|\gamma_f) = \mathcal{H}(\gamma|\gamma_f) + \frac{1}{2} \iint_{\mathbb{R} \times \mathbb{R}} w(x-y) \rho_{\gamma-\gamma_f}(x) \rho_{\gamma-\gamma_f}(y) dx dy$$

and global well-posedness following the proof of Lewin and Sabin [31]. Indeed, it is already shown that initial data having finite relative entropy can be approximated by a smooth finite-rank operator with the convergence $\|Q_{n,0} - Q_0\|_{\mathcal{H}^{\frac{1}{2}}} \rightarrow 0$ (see [31, Lemma 10]). In 1D, by (A.1), this convergence is sufficient to obtain the convergence in density,

$$\lim_{n \rightarrow \infty} \|\rho_{e^{it\Delta} Q_{n,0} e^{-it\Delta}} - \rho_{e^{it\Delta} Q_0 e^{-it\Delta}}\|_{\mathcal{D}^{\frac{3+\epsilon}{4}}(I)} = 0.$$

For more details, we refer to [31, Section 7]. \square

REFERENCES

- [1] G.L. Aki, J. Dolbeault and C. Sparber, *Thermal effects in gravitational Hartree systems*, Ann. Henri Poincaré **12** (2011), no. 6, 1055–1079.
- [2] A.B. Aleksandrov and V.V. Peller, *Operator Lipschitz functions*. Uspekhi Mat. Nauk **71** (2016), no.4, 3–106; translation in Russian Math. Surveys **71** (2016), no.4, 605–702.

- [3] N. Benedikter, V. Jakšić, M. Porta and B. Schlein, *Mean-field evolution of fermionic mixed states*, Comm. Pure Appl. Math. **69** (2016), no. 12, 2250–2303.
- [4] N. Benedikter, M. Porta and B. Schlein, *Mean-field evolution of fermionic systems*, Comm. Math. Phys. **331** (2014), no. 3, 1087–1131.
- [5] N. Bez, Y. Hong, S. Lee, S. Nakamura and Y. Sawano, *On the Strichartz estimates for orthonormal systems of initial data with regularity*, Adv. Math. **354** (2019), 106736, 37 pp.
- [6] N. Bez, S. Lee and S. Nakamura, *Strichartz estimates for orthonormal families of initial data and weighted oscillatory integral estimates*, Forum Math. Sigma **9** (2021), Paper No. e1, 52 pp.
- [7] M.Š. Birman and M.Z. Solomjak, *Estimates of singular numbers of integral operators. III. Operators in unbounded domains*, Vestnik Leningrad. Univ. **24** (1969), no.1, 35–48.
- [8] M.Š. Birman and M.Z. Solomjak, *Quantitative analysis in Sobolev imbedding theorems and applications to spectral theory*, American Mathematical Society Translations, Series 2, **114**, American Mathematical Society, Providence, R.I., 1980.
- [9] A. Bove, G. Da Prato, G. Fano, *An existence proof for the Hartree-Fock time-dependent problem with bounded two-body interaction*, Comm. Math. Phys. **37** (1974), 183–191.
- [10] A. Bove, G. Da Prato, G. Fano, *On the Hartree-Fock time-dependent problem*, Comm. Math. Phys. **49** (1976), 25–33.
- [11] J.M. Chadam, *The time-dependent Hartree-Fock equations with Coulomb two-body interaction*, Comm. Math. Phys. **46** (1976), 99–104.
- [12] T. Chen, Y. Hong and N. Pavlović, *Global well-posedness of the NLS system for infinitely many fermions*, Arch. Ration. Mech. Anal. **224** (2017), no. 1, 91–123.
- [13] T. Chen, Y. Hong, N. Pavlović, *On the scattering problem for infinitely many fermions in dimensions $d \geq 3$ positive temperature*, Ann. Inst. H. Poincaré Anal. Non Linéaire **35** (2018), no. 2, 393–416.
- [14] C. Collot, A.-S. de Suzzoni, *Stability of equilibria for a Hartree equation for random fields*, J. Math. Pures Appl. (9) **137** (2020), 70–100.
- [15] C. Collot, E. Danesi, A.-S. de Suzzoni, C. Malézé, *Stability of homogeneous equilibria of the Hartree-Fock equation, for its equivalent formulation for random fields*, (2023), arXiv:2310.03442.
- [16] C. Collot, A.-S. de Suzzoni, *Stability of steady states for Hartree and Schrödinger equations for infinitely many particles*, Ann. H. Lebesgue **5** (2022), 429–490.
- [17] A. Deuchert, C. Hainzl and R. Seiringer, *Note on a family of monotone quantum relative entropies*, Lett. Math. Phys. **105** (2015), no.10, 1449–1466.
- [18] X. Dong, *The Hartree equation with a constant magnetic field: well-posedness theory*, Lett. Math. Phys. **111** (2021), no.4, Paper No. 101.
- [19] L. Erdős, B. Schlein and H.-T. Yau, *Derivation of the Gross-Pitaevskii hierarchy for the dynamics of Bose-Einstein condensate*, Comm. Pure Appl. Math. **59** (2006), no. 12, 1659–1741.
- [20] L. Erdős, B. Schlein and H.-T. Yau, *Derivation of the cubic non-linear Schrödinger equation from quantum dynamics of many-body systems*, Invent. Math. **167** (2007), no. 3, 515–614.
- [21] L. Erdős, B. Schlein and H.-T. Yau, *Rigorous derivation of the Gross-Pitaevskii equation with a large interaction potential*, J. Amer. Math. Soc. **22** (2009), no. 4, 1099–1156.
- [22] L. Erdős, B. Schlein and H.-T. Yau, *Derivation of the Gross-Pitaevskii equation for the dynamics of Bose-Einstein condensate*, Ann. of Math. (2) **172** (2010), no. 1, 291–370.
- [23] R.L. Frank, M. Lewin, E.H. Lieb and R. Seiringer, *A positive density analogue of the Lieb-Thirring inequality*, Duke Math. J. **162** (2013), no.3, 435–495.
- [24] R.L. Frank, M. Lewin, E.H. Lieb and R. Seiringer, *Strichartz inequality for orthonormal functions*, J. Eur. Math. Soc. (JEMS) **16** (2014), no.7, 1507–1526.

- [25] R.L. Frank and J. Sabin, *Restriction theorems for orthonormal functions, Strichartz inequalities, and uniform Sobolev estimates*, Amer. J. Math. **139** (2017), no. 6, 1649–1691.
- [26] S. Hadama, *Asymptotic stability of a wide class of stationary solutions for the Hartree and Schrödinger equations for infinitely many particles*, (2023), arXiv:2308.15929.
- [27] M. Keel and T. Tao, *Endpoint Strichartz estimates*, Amer. J. Math. **120** (1998), no. 5, 955–980.
- [28] S. Hadama and T. Yamamoto, *Probabilistic Strichartz estimates in Schatten classes and their applications to the Hartree equation*, arXiv:2311.02713.
- [29] M. Lewin and J. Sabin, *A family of monotone quantum relative entropies* Lett. Math. Phys. **104** (2014), no.6, 691–705.
- [30] M. Lewin and J. Sabin, *The Hartree equation for infinitely many particles, II: Dispersion and scattering in 2D*, Anal. PDE **7** (2014), no.6, 1339–1363.
- [31] M. Lewin and J. Sabin, *The Hartree equation for infinitely many particles I. Well-posedness theory*, Comm. Math. Phys. **334** (2015), no.1, 117–170.
- [32] C. Malézé, *A Scattering Result Around a Non-Localised Equilibria for the Quintic Hartree Equation for Random Fields*, (2023), arXiv:2309.10500.
- [33] F. Pusateri, I.M. Sigal, *Long-time behaviour of time-dependent density functional theory*, Arch. Ration. Mech. Anal. **241** (2021), no. 1, 447–473.
- [34] B. Simon, *Trace ideals and their applications, Second edition*. Mathematical Surveys and Monographs. **120**. American Mathematical Society. (2005)
- [35] A.-S. de Suzzoni, *An equation on random variables and systems of fermions*, (2015) arXiv:1507.06180.
- [36] S. Zagatti, *The Cauchy problem for Hartree-Fock time-dependent equations*, Ann. Inst. H. Poincaré Phys. Théor. **56** (1992), 357–374.

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KITA-SHIRAKAWA, SAKYO-KU, KYOTO, JAPAN 606-8502.

Email address: hadama@kurims.kyoto-u.ac.jp

DEPARTMENT OF MATHEMATICS, CHUNG-ANG UNIVERSITY, SEOUL 06974, KOREA

Email address: yhhong@cau.ac.kr