

On degree power sum in P_k -free graphs

Jiangdong Ai*, Fankang He†, Yihang Liu‡, Bo Ning§

Abstract

Let G be a graph on n vertices with degree sequence (d_1, d_2, \dots, d_n) . For a real $p \geq 1$, let $D_p(G) = \sum_{i=1}^n d_i^p$. A Turán-type problem of degree power sum was initiated by Caro and Yuster [5]: determining the function $D_p(n, H) := \max\{D_p(G) : G \text{ is an } n\text{-vertex } H\text{-free graph}\}$. They obtained some exact values for certain graphs H . For a path P_k , they mentioned that “a close examination of the proof of Theorem 1.2 shows that the value of $n_0(k)$ in the statement of the theorem is $O(k^2)$ ”, namely, they could show the n -vertex P_k -free graph with maximum degree power sum is $W_{n, k-1, \lfloor \frac{k}{2} \rfloor - 1} = K_{\lfloor \frac{k}{2} \rfloor - 1} \vee \left((n - \lceil \frac{k}{2} \rceil) K_1 \cup K_{1+k-2\lfloor \frac{k}{2} \rfloor} \right)$ when $n \geq ck^2$ for some constant c . In this note, we improve their result to a linear size of k by a different approach. The bound is tight up to a constant factor.

1 Introduction

Let G be a graph on n vertices whose degree sequence is (d_1, d_2, \dots, d_n) . For a real $p \geq 1$, we denote the *degree power sum* of G by $D_p(G) := \sum_{i=1}^n d_i^p$. This parameter, which is also known as the *general zeroth-order Randić index*, is well-studied in chemical graph theory, see the comprehensive survey [2]. Caro and Yuster [5] initiated a Turán-type problem of degree power sum: determining the function $D_p(n, H) := \max\{D_p(G) : G \text{ is an } n\text{-vertex } H\text{-free graph}\}$. Since then, a lot of research on this Turán-type problem has appeared, see [3–10].

In this note, we focus on this Turán-type problem when the forbidden graph is a path. Caro and Yuster [5] determined $D_p(n, P_k)$ for any integers $k \geq 2$, $p \geq 2$ and sufficiently large n , and characterized the corresponding extremal graphs. Define $W_{n, k, s}$ to be a graph on n vertices, in which its vertex set can be partitioned into three subsets X, Y, Z such that $|X| = s$, $|Y| = k - 2s$, $|Z| = n - (k - s)$, and the edge set consists of all possible edges between X and Z and all edges in $X \cup Y$.

Theorem 1 ([5]). *For two integers $k \geq 4$ and $p \geq 2$, there exists a positive integer $n_0 = n_0(k)$ such that for all $n \geq n_0$, the following holds. If G is an n -vertex P_k -free graph with maximum degree power sum $D_p(G)$, then G is $W_{n, k-1, t}$ where $t = \lfloor \frac{k}{2} \rfloor - 1$.*

*Corresponding author. School of Mathematical Sciences and LPMC, Nankai University, Tianjin 300071, P.R. China. Email: jd@nankai.edu.cn.

†School of Mathematical Sciences and LPMC, Nankai University, Tianjin 300071, P.R. China. Email: hefankang@mail.nankai.edu.cn.

‡School of Mathematical Sciences and LPMC, Nankai University, Tianjin 300071, P.R. China. Email: liuyihang@mail.nankai.edu.cn.

§College of Computer Science, Nankai University, Tianjin 300350, P.R. China. Email: bo.ning@nankai.edu.cn. Partially supported by the NSFC grants (Nos. 12371350 and 11971346).

Note that Caro and Yuster obtained the above result for integers $p \geq 2$. A natural question arising from Theorem 1 is to determine, for fixed real p , how large $n_0(k)$ should be as a function of k . Caro and Yuster mentioned that “a close examination of the proof of Theorem 1.2 shows that the value of $n_0(k)$ in the statement of the theorem is $O(k^2)$ ” (quoted from [5]). In this note, by a different method, we prove that $n_0(k)$ can be as small as $10k$. And our result extends Theorem 1 to any real $p \geq 2$. We remark that the bound is tight up to a constant factor. Note that K_{k-1} , rather than $W_{k-1,k-1,t}$ where $t = \lfloor \frac{k}{2} \rfloor - 1$, attains the maximum degree power sum among all $(k-1)$ -vertex P_k -free graphs. This yields $n_0(k) \geq k$.

Theorem 2. *For an integer $k \geq 4$ and real $p \geq 2$, let G be an n -vertex P_k -free graph with the maximum degree power sum $D_p(G)$. If $n \geq 10k$, then G is $W_{n,k-1,t}$ where $t = \lfloor \frac{k}{2} \rfloor - 1$.*

Let us outline the main ideas behind the proof of Theorem 2. Given an n -vertex P_k -free graph G with the largest degree power sum, we will first show that there are a few components. This will allow us to use structural information to guarantee the existence of a large component G_i . Next, we will show that if there is any other component G_j , then we can use G_i to “absorb” G_j to get a larger degree power sum which contradicts the maximality of the degree power sum of G .

2 Proofs

Recently, Ai, Lei, Ning, and Shi [1] proved two connected versions of Theorem 1. Based on these results, we characterize some structures in the proof of Theorem 2.

Lemma 1 ([1]). *Let $k \geq 4$ and $t = \lfloor \frac{k}{2} \rfloor - 1$ and let $p \geq 2$. The following holds for all $n \geq k$. If G is a connected n -vertex P_k -free graph with maximum degree power sum $D_p(G)$, then G is $W_{n,k-1,1}$ or $W_{n,k-1,t}$.*

Lemma 2 ([1]). *Let $k \geq 4$ and $t = \lfloor \frac{k}{2} \rfloor - 1$ and let $p \geq 2$. The following holds for all $n \geq 2k$. If G is a connected n -vertex P_k -free graph with maximum degree power sum $D_p(G)$, then G is $W_{n,k-1,t}$.*

The following lemma is useful in the proof of the main theorem.

Lemma 3. *Let $a \geq b > 0$ and $p \geq 2$. Then $(a+b)^p \geq a^p + pa^{p-1}b + b^p$. In particular, $(a+b)^p \geq a^p + (p+1)b^p$.*

Proof. We first show that $(x+1)^p \geq x^p + px^{p-1} + 1$ when $x \geq 1$. Let $f(x) = (x+1)^p - x^p - px^{p-1} - 1$ and so $f(1) = 2^p - p - 2 \geq 0$. Taking derivative of $f(x)$, we get $f'(x) = p((x+1)^{p-1} - x^{p-1} - (p-1)x^{p-2})$. Note that $f'(x) \geq 0$ due to convexity of x^{p-1} . Hence, $f(x) \geq 0$ when $x \geq 1$ as desired. Then

$$(a+b)^p = b^p \left(\frac{a}{b} + 1 \right)^p \geq b^p \left(\left(\frac{a}{b} \right)^p + p \left(\frac{a}{b} \right)^{p-1} + 1 \right) = a^p + pa^{p-1}b + b^p \geq a^p + (p+1)b^p.$$

This completes the proof. ■

Proof of Theorem 2. Let G be an n -vertex P_k -free graph which attains the maximum degree power sum $D_p(G)$ where $p \geq 2$. Let $t = \lfloor \frac{k}{2} \rfloor - 1$ and let G_1, G_2, \dots, G_ℓ be all components of G . Let $n_i = |G_i|$ for $i \in [\ell]$. By Lemma 1, we can immediately get the following claim.

Claim 1. For any component G_i , where $1 \leq i \leq \ell$, if $n_i \leq k - 1$, then G_i is a clique; if $n_i \geq k$, then $G_i \in \{W_{n_i, k-1, 1}, W_{n_i, k-1, t}\}$.

Proof. When $n_i \leq k - 1$, then any n_i -vertex graph is P_k -free. Hence, K_{n_i} would attain the largest degree power sum among all n_i -vertex graphs. When $n_i \geq k$, by Lemma 1, we have $G_i \in \{W_{n_i, k-1, 1}, W_{n_i, k-1, t}\}$. ■

In the rest of this proof, we say that a component G_i is a *Type-A* component if $G_i = W_{n_i, k-1, 1}$, and is a *Type-B* component if $G_i = W_{n_i, k-1, t}$ where $t = \lfloor \frac{k}{2} \rfloor - 1$.

Claim 2. For any two components G_1 and G_2 of order $n_1 \leq n_2 \leq k - 1$, we have $n_1 + n_2 \geq k$.

Proof. Suppose that $n_1 + n_2 \leq k - 1$. Then G_i must be a clique for $i = 1, 2$. Note that $D_p(G_1 \cup G_2) = n_1(n_1 - 1)^p + n_2(n_2 - 1)^p$. If we replace $G_1 \cup G_2$ with $K_{n_1+n_2}$ and denote the new graph by G' . Then G' is still P_k -free since $n_1 + n_2 \leq k - 1$. And $D_p(K_{n_1+n_2}) = (n_1 + n_2)(n_1 + n_2 - 1)^p$. Note that $D_p(G_1 \cup G_2) = n_1(n_1 - 1)^p + n_2(n_2 - 1)^p \leq (n_1 + n_2)((n_1 - 1)^p + (n_2 - 1)^p) < D_p(K_{n_1+n_2})$, as there holds the inequality $x^p + y^p < (x + y + 1)^p$ where $x \geq 0, y \geq 0$ and $p \geq 2$. Now, we proved G' has a larger degree power sum than G , a contradiction. This proves the claim. ■

Claim 3. There exists at most one component of order at most $k - 2$.

Proof. Suppose to the contrary that there exist two components, say G_1 and G_2 , of order $n_1 \leq k - 2$ and $n_2 \leq k - 2$, respectively. By Claim 2, $n_1 + n_2 \geq k$. Let $n'_1 = k - 1$ and $n'_2 = n_1 + n_2 - k + 1$. Note that $n'_2 \leq k - 3$, then $K_{n'_1} \cup K_{n'_2}$ contains no P_k . Recall that $D_p(K_r) = r(r - 1)^p$. Then $D_p(K_{n'_1} \cup K_{n'_2}) > D_p(G_1 \cup G_2)$ by convexity argument. This means that if we replace $G_1 \cup G_2$ with $K_{n'_1} \cup K_{n'_2}$, then we get an n -vertex P_k -free graph with a larger degree power sum, a contradiction. ■

Claim 4. There exist at most five components of order $k - 1$.

Proof. Suppose to the contrary that there exist six components, say $G_i, i = [6]$ of order $n_i = k - 1, i \in [6]$. Then

$$\begin{aligned} D_p(W_{6k-6, k-1, t}) - D_p\left(\bigcup_{i=1}^6 G_i\right) &= \left(\lfloor \frac{k}{2} \rfloor - 1\right) (6k - 7)^p + (6k - \lceil \frac{k}{2} \rceil - 6) \left(\lfloor \frac{k}{2} \rfloor - 1\right)^p \\ &\quad + \left(1 + k - 2\lfloor \frac{k}{2} \rfloor\right) \left(\lceil \frac{k}{2} \rceil - 1\right)^p - 6(k - 1)(k - 2)^p \\ &> 6^p \left(\lfloor \frac{k}{2} \rfloor - 1\right) \left(k - \frac{7}{6}\right)^p - 6(k - 1)(k - 2)^p \\ &\geq 6^2 \left(\frac{k}{2} - \frac{3}{2}\right) (k - 2)^p - 6(k - 1)(k - 2)^p \\ &= (12k - 48)(k - 2)^p \geq 0. \end{aligned}$$

This means that we can replace $\bigcup_{i=1}^6 G_i$ with $W_{6k-6, k-1, t}$ to get an n -vertex P_k -free graph with a larger degree power sum, a contradiction. ■

Claim 5. There exists at most one Type-A component.

Proof. Suppose to the contrary that there exist two Type-A components, say G_1 and G_2 , of order n_1 and n_2 , respectively. Then by Claim 1 and Lemma 2, $k \leq n_1 \leq 2k - 1$ and $k \leq n_2 \leq 2k - 1$. Let $n'_1 = k - 1$ and $n'_2 = n_1 + n_2 - k + 1$, let $G'_1 = W_{n'_1, k-1, 1}$ and $G'_2 = W_{n'_2, k-1, 1}$. Recall that $D_p(W_{n, k-1, t}) = (n - k + 2) + (k - 3)^p + (n - 1)^p$ holds for $n \geq k - 2$. Then $D_p(G'_1 \cup G'_2) > D_p(G_1 \cup G_2)$ by convexity argument. Now we can replace $G_1 \cup G_2$ with $G'_1 \cup G'_2$ to get an n -vertex P_k -free graph with a larger degree power sum, a contradiction. \blacksquare

Claim 6. There exists at most one Type-B component.

Proof. Suppose to the contrary that there exist two Type-B components, say G_1 and G_2 , of order n_1 and n_2 , respectively. W.O.L.G., suppose $n_1 \geq n_2$. By Claim 1, $n_1 \geq k$ and $n_2 \geq k$. Then $D_p(G_i) = (\lfloor \frac{k}{2} \rfloor - 1)(n_i - 1)^p + (n_i - \lceil \frac{k}{2} \rceil)(\lfloor \frac{k}{2} \rfloor - 1)^p + (1 + k - 2\lfloor \frac{k}{2} \rfloor)(\lceil \frac{k}{2} \rceil - 1)^p$, $i = 1, 2$. Let $G_0 = W_{n_1+n_2, k-1, t}$. Note that

$$\begin{aligned}
D_p(G_0) - D_p(G_1 \cup G_2) &= \left(\lfloor \frac{k}{2} \rfloor - 1 \right) \left((n_1 + n_2 - 1)^p - (n_1 - 1)^p - (n_2 - 1)^p \right) \\
&\quad + \lceil \frac{k}{2} \rceil \left(\lfloor \frac{k}{2} \rfloor - 1 \right)^p - \left(1 + k - 2\lfloor \frac{k}{2} \rfloor \right) \left(\lceil \frac{k}{2} \rceil - 1 \right)^p \\
&> \left(\lfloor \frac{k}{2} \rfloor - 1 \right) \left((n_1 + n_2 - 2)^p - (n_1 - 1)^p - (n_2 - 1)^p \right) \\
&\quad + \lceil \frac{k}{2} \rceil \left(\lfloor \frac{k}{2} \rfloor - 1 \right)^p - \left(1 + k - 2\lfloor \frac{k}{2} \rfloor \right) \left(\lceil \frac{k}{2} \rceil - 1 \right)^p \\
&\geq \left(\lfloor \frac{k}{2} \rfloor - 1 \right) p(n_2 - 1)^p + \lceil \frac{k}{2} \rceil \left(\lfloor \frac{k}{2} \rfloor - 1 \right)^p \\
&\quad - \left(1 + k - 2\lfloor \frac{k}{2} \rfloor \right) \left(\lceil \frac{k}{2} \rceil - 1 \right)^p \tag{1} \\
&\geq 2(k - 1)^p - 2 \left(\lceil \frac{k}{2} \rceil - 1 \right)^p + \lceil \frac{k}{2} \rceil \left(\lfloor \frac{k}{2} \rfloor - 1 \right)^p \\
&> 0.
\end{aligned}$$

The inequality (1) holds due to Lemma 3 by letting $a = n_1 - 1$ and $b = n_2 - 1$. Then we can replace $G_1 \cup G_2$ with G_0 to get an n -vertex P_k -free graph with a larger degree power sum, a contradiction. \blacksquare

Claim 7. If $n_i \geq 2k$, $n_j \leq k - 1$ and G_i is of Type-B, then we can replace $G_i \cup G_j$ with $W_{n_i+n_j, k-1, t}$ to get a larger degree power sum.

Proof. Let $G_0 = W_{n_i+n_j, k-1, t}$. Then

$$\begin{aligned}
D_p(G_0) - D_p(G_i \cup G_j) &= \left(\lfloor \frac{k}{2} \rfloor - 1 \right) \left((n_i + n_j - 1)^p - (n_i - 1)^p \right) \\
&\quad + n_j \left(\lfloor \frac{k}{2} \rfloor - 1 \right)^p - n_j(n_j - 1)^p \\
&\geq \left(\lfloor \frac{k}{2} \rfloor - 1 \right) p(n_i - 1)^{p-1} n_j + n_j \left(\lfloor \frac{k}{2} \rfloor - 1 \right)^p - n_j(n_j - 1)^p \tag{2} \\
&\geq p \left(\frac{k}{2} - \frac{3}{2} \right) n_j(2k - 1)^{p-1} - n_j(n_j - 1)^p + n_j \left(\lfloor \frac{k}{2} \rfloor - 1 \right)^p
\end{aligned}$$

$$\begin{aligned}
&\geq n_j(2k-6) \left(k - \frac{1}{2}\right)^{p-1} - n_j(k-2)^p + n_j \left(\left\lfloor \frac{k}{2} \right\rfloor - 1\right)^p \\
&> 0.
\end{aligned}$$

The inequality (2) holds due to Lemma 3 by letting $a = n_i - 1$ and $b = n_j$. This completes the proof. \blacksquare

Claim 8. If $n_i \geq 2k$, G_i is of Type-B, and G_j is of Type-A, then we can replace $G_i \cup G_j$ with $W_{n_i+n_j, k-1, t}$ to get a larger degree power sum.

Proof. Let $G_0 = W_{n_i+n_j, k-1, t}$. By Lemma 2 and Claim 1, we have $k \leq n_j \leq 2k-1$. Then

$$\begin{aligned}
D_p(G_0) - D_p(G_1 \cup G_2) &= \left(\left\lfloor \frac{k}{2} \right\rfloor - 1\right) \left((n_i + n_j - 1)^p - (n_i - 1)^p\right) + n_j \left(\left\lfloor \frac{k}{2} \right\rfloor - 1\right)^p \\
&\quad - (n_j - k + 2) - (k-3)^{p+1} - (n_j - 1)^p \\
&> \left(\left\lfloor \frac{k}{2} \right\rfloor - 1\right) p(n_i - 1)^{p-1} n_j + n_j \left(\left\lfloor \frac{k}{2} \right\rfloor - 1\right)^p \\
&\quad - (n_j - k + 2) - (k-3)^{p+1} - (n_j - 1)^p \tag{3} \\
&\geq \left(\left\lfloor \frac{k}{2} \right\rfloor - 1\right) n_j (n_i - 1)^{p-1} - (k-3)^{p+1} \\
&\quad + \left(\left\lfloor \frac{k}{2} \right\rfloor - 1\right) n_j (n_i - 1)^{p-1} - (n_j - 1)^p - (n_j - k + 2) \\
&\geq \left(\frac{k}{2} - \frac{3}{2}\right) k(2k-1)^{p-1} - (k-3)^{p+1} \\
&\quad + n_j (n_i - 1)^{p-1} - (n_j - 1)^p - (n_j - k + 2) \\
&\geq (k-3) k \left(k - \frac{1}{2}\right)^{p-1} - (k-3)^{p+1} \\
&\quad + n_j (n_j - 1)^{p-1} - (n_j - 1)^p - (n_j - k + 2) \\
&> 0.
\end{aligned}$$

The inequality (3) holds due to Lemma 3 by letting $a = n_i - 1$ and $b = n_j$. This completes the proof. \blacksquare

Claim 9. There is no component of order at most $k-1$. And there is no Type-A component.

Proof. We first show that there is a component of order at least $2k$. By Claim 3 and Claim 4, there are at most six components of order at most $k-1$. By Claim 5, there is at most one Type-A component. Lemma 2 tells us that the order of a Type-A component is at most $2k-1$. Combined with the assumption that $n \geq 10k$, there is a component of order at least $10k - 6(k-1) - 2k + 1 = 2k + 7$, which is of Type-B.

If there is a component of order at most $k-1$, then by Claim 7, we can get an n -vertex P_k -free graph with a larger degree power sum. If there is a Type-A component, then by Claim 8, we can get an n -vertex P_k -free graph with a larger degree power sum. Now, we are done. \blacksquare

As a result, G has only one component which is of Type-B, i.e., G is $W_{n, k-1, t}$. This proves the theorem. \blacksquare

References

- [1] J. Ai, H. Lei, B. Ning, and Y. Shi. Graph operations and a unified method for kinds of Turán-type problems on paths, cycles and matchings. *arXiv preprint arXiv:2312.08226*, 2023.
- [2] A. Ali, I. Gutman, E. Milovanović, and I. Milovanović. Sum of powers of the degrees of graphs: extremal results and bounds. *MATCH Commun. Math. Comput. Chem.*, 80(1):5–84, 2018.
- [3] B. Bollobás and V. Nikiforov. Degree powers in graphs with forbidden subgraphs. *Electron. J. Combin.*, 11(1):Research Paper 42, 8, 2004.
- [4] B. Bollobás and V. Nikiforov. Degree powers in graphs: the Erdős-Stone theorem. *Combin. Probab. Comput.*, 21(1-2):89–105, 2012.
- [5] Y. Caro and R. Yuster. A Turán type problem concerning the powers of the degrees of a graph. *Electron. J. Combin.*, 7:Research Paper 47, 14, 2000.
- [6] Y. Chang, X. Chen, and S. Zhang. Extremal results on degree powers in some classes of graphs. *arXiv preprint arXiv:2312.07005*, 2023.
- [7] R. Gu, X. Li, and Y. Shi. Degree powers in C_5 -free graphs. *Bull. Malays. Math. Sci. Soc.*, 38(4):1627–1635, 2015.
- [8] Y. Lan, H. Liu, Z. Qin, and Y. Shi. Degree powers in graphs with a forbidden forest. *Discrete Math.*, 342(3):821–835, 2019.
- [9] O. Pikhurko and A. Taraz. Degree sequences of F -free graphs. *Electron. J. Combin.*, 12:Research Paper 69, 12, 2005.
- [10] L. Zhang. Degree powers in $K_{s,t}$ -minor free graphs. *Discrete Math.*, 345(4):Paper No. 112783, 9, 2022.