

Elliptic Virtual Structure Constants and Generalizations of BCOV-Zinger Formula to Projective Fano Hypersurfaces

Masao Jinzenji⁽¹⁾, Ken Kuwata⁽²⁾

⁽¹⁾*Department of Mathematics, Okayama University
Okayama, 700-8530, Japan*

⁽²⁾*Department of General Education
National Institute of Technology, Kagawa College
Chokushi, Takamatsu, 761-8058, Japan*

e-mail address: ⁽¹⁾ pcj70e4e@okayama-u.ac.jp
⁽²⁾ kuwata-k@t.kagawa-nct.ac.jp

April 16, 2024

Abstract

In this paper, we propose a recipe for B-model computation of genus 1 Gromov-Witten invariants of Calabi-Yau and Fano Projective Hypersurfaces. Our formalism can be applied equally to both Calabi-Yau and Fano cases. In Calabi-Yau case, drastic cancellation of terms used in our formalism occurs and it results in another representation of BCOV-Zinger formula for projective Calabi-Yau hypersurfaces.

1 Introduction

Evaluation of higher genus Gromov-Witten invariants has been one of the most exciting and challenging topics in the study of topological strings and mirror symmetry.

In 1993, Bershadsky, Cecotti, Ooguri and Vafa evaluated genus 1 Gromov-Witten invariants of quintic Calabi-Yau 3-fold and several other Calabi-Yau 3-folds by combining the idea of holomorphic anomaly equation and mirror symmetry [1]. Soon after the paper appeared, they generalized the method presented in [1] to evaluate higher genus ($g \geq 2$) Gromov-Witten invariants of Calabi-Yau 3-folds [2]. Attempts to generalize their results to higher dimensional Calabi-Yau manifolds can be seen in [19].

On the other hand, study of evaluation of higher genus Gromov-Witten invariants of Fano manifolds began after the celebrated paper [17], which derived associativity equation of quantum cohomology ring by using analysis of Chow ring of moduli space of genus 0 stable curves with four marked points, appeared. It enabled us to evaluate genus 0 Gromov-Witten invariants of Fano manifolds. In 1997, Getzler derived Getzler's equation, a kind of generalization of associativity equation to genus 1 Gromov-Witten invariants, from analysis of Chow ring of moduli space of genus 1 stable curves with four marked points [6]. In the same year, Eguchi, Hori and Xiong proposed Virasoro Conjecture, which has potential ability to evaluate all genus Gromov-Witten invariants of Fano manifolds by combining with topological recursion relation [8]. These are results which use differential equations of generating function of Gromov-Witten invariants.

There exists another trend of evaluation of Gromov-Witten invariants. In 1994, Kontsevich directly evaluated genus 0 Gromov-Witten invariants of Projective hypersurfaces by combining the idea of moduli space of stable

maps with Bott residue formula [16]. Actual computational process of this method is extremely hard, but it shedded light on geometrical study on Gromov-Witten invariants. In 1997, Graber and Pandharipande generalized the method in [16] to the case of higher genus Gromov-Witten invariants of projective space [7]. In [7], we can see that crucial point of generalization is consideration on Hodge bundle on moduli space of higher genus stable curves. As far as we know, the results in [7] is also applicable to evaluate higher genus local Gromov-Witten invariants of projective space [5], but not applicable to the case of compact projective hypersurface (at least compact Calabi-Yau hypersurface). Let us summarize this trend by the word “localization”.

We summarize here situations of study of Gromov-Witten invariants around 90’s. At the genus 0 level, mirror symmetry conjecture became tractable by invention of localization method. But in order to attack BCOV conjecture in mathematics sense, localization method can not be applied, at least to the case of compact Calabi-Yau manifold. On the other hand, associativity equation is not effective for evaluation of genus 0 Gromov-Witten invariants of Calabi-Yau manifolds. Getzler’s equation and Virasoro conjecture are also not effective for evaluation of higher genus Gromov-Witten invariants of Calabi-Yau manifolds.

In 2009, Zinger introduced the idea of reduced Gromov-Witten invariants and by combining this idea with a kind of localization, he derived a closed formula for generating function of genus 1 Gromov-Witten invariants of projective Calabi-Yau hypersurfaces [20]. Since then, there appeared many literatures in mathematics on higher genus Gromov-Witten invariants of Calabi-Yau 3-folds. We call the formula proved in [20] BCOV-Zinger formula, because Zinger’s formula in the case of quintic 3-fold is nothing but the formula proposed in [1]. From a lot of experimental computations, we believe his formula is correct, but it is hard for us to understand his logic used in the proof of his formula. This may be so because we originally have physics origin.

The motivation of this paper is “to construct our original approach to genus 1 Gromov-Witten invariants of projective hypersurfaces from the point of view of the theory of moduli space of quasimaps we have constructed so far”.

Let us briefly review our study of classical mirror symmetry of projective hypersurface. Let M_N^k be degree k non-singular hypersurface in CP^{N-1} . We denote by h hyperplane class in $H^{1,1}(CP^{N-1}, \mathbf{C})$. Quantum cohomology ring of M_N^k is defined by the following multiplication rule:

$$\mathcal{O}_h \cdot \mathcal{O}_{h^{N-2-m}} = \sum_{d=0}^{\infty} \frac{1}{k} \langle \mathcal{O}_h \mathcal{O}_{h^{N-2-m}} \mathcal{O}_{h^{m-1+(N-k)d}} \rangle_{0,d} e^{dt} \mathcal{O}_{h^{N-1-m-(N-k)d}}, \quad (1.1)$$

where $\langle \mathcal{O}_h \mathcal{O}_{h^{N-2-m}} \mathcal{O}_{h^{m-1+(N-k)d}} \rangle_{0,d}$ is genus 0 and degree d Gromov-Witten invariant of M_N^k .

In [3], we defined $L_m^{N,k,d}$ by $\frac{1}{k} \langle \mathcal{O}_h \mathcal{O}_{h^{N-2-m}} \mathcal{O}_{h^{m-1+(N-k)d}} \rangle_{0,d}$, which is structure constant of quantum cohomology ring of M_N^k and proved recursive formulas that describe $L_m^{N,k,d}$ in terms of weighted homogeneous polynomial of $L_{m'}^{N+1,k,d'}$ ’s ($d' \leq d$) up to $d = 3$. These recursive formulas take the same form if $N - k \geq 2$. But in the case of $N - k = 1, 0$, we have to modify these recursive formulas. Then we applied the recursive formula for the $N - k \geq 2$ cases formally to the $N - k = 1, 0$ case and found that resulting “virtual structure constant” becomes expansion coefficient of the generalized hypergeometric series used in mirror computation of Gromov Witten invariants of M_N^k and of derivative of the mirror map. We then denote by $\tilde{L}_m^{k,k,d}$ the resulting number and call it “virtual structure constant.” This is the beginning of the notion: “virtual structure constant” and it plays the role of B-model analogue of the structure constant (Gromov-Witten invariant) of quantum cohomology ring of M_N^k .

In [10, 11], we showed that the virtual structure constant $\tilde{L}_m^{k,k,d}$ is given as intersection number $\frac{d}{k} \cdot w(\mathcal{O}_{h^{k-2-m}} \mathcal{O}_{m-1})_{0,d}$ of moduli space of quasimaps from CP^1 with two marked points to CP^{k-1} , which we denote by $\widetilde{Mp}_{0,2}(k, d)^1$. Then we began to call $w(\mathcal{O}_{h^a} \mathcal{O}_{h^b})_{0,d}$, which is defined as intersection number of $\widetilde{Mp}_{0,2}(k, d)$ (it was generalized to general hypersurface M_N^k), “virtual structure constant”, by which we mean B-model analogue of Gromov-Witten invariants.

In [14], we constructed moduli space of quasimaps from CP^1 with $2 + n$ marked points to CP^{N-1} , which we denote by $\widetilde{Mp}_{0,2|n}(N, d)$, and defined multi-point virtual structure constant $w(\mathcal{O}_{h^a} \mathcal{O}_{h^b} | \prod_{p=0}^{N-2} (\mathcal{O}_{h^p})^{m_p})_{0,d}$ for M_N^k as intersection number of $\widetilde{Mp}_{0,2|n}(N, d)$. It is expected to play the role of B-model analogue of genus 0 Gromov-Witten invariant $\langle \mathcal{O}_{h^a} \mathcal{O}_{h^b} | \prod_{p=0}^{N-2} (\mathcal{O}_{h^p})^{m_p} \rangle_{0,d}$ of M_N^k but it generically does not coincide with the corresponding Gromov-Witten invariant. Then we defined “mirror map”, which is defined as generating function

¹For another approach of moduli space of quasimaps, please refer to [4]

of $w(\mathcal{O}_{h^a} \mathcal{O}_1 | \prod_{p=0}^{N-2} (\mathcal{O}_{h^p})^{m_p})_{0,d}$ (see the next section for details), and conjectured that generating function of multi-point virtual structure constants is translated into generating function of Gromov-Witten invariants via coordinate change of deformation parameters given by the mirror map (see also the next section for details). This conjecture enables us to do non-trivial mirror computation of Gromov-Witten invariants for Fano projective hypersurface, even though mirror map used in mirror computation of structure constants of small quantum cohomology ring of Fano projective hypersurface is trivial. This construction is the starting point of this paper.

At this stage, we discuss geometrical merit of introducing virtual structure constant, or moduli space of quasimaps. First of all, geometrical structure of moduli space of quasimaps is far simpler than the one of moduli space of stable maps. This enables us to write down short proof of mirror theorem of projective hypersurface M_N^k [12]. Let us illustrate advantage of moduli space of quasimaps by taking localization computation as example. In localization computation of genus 0 and degree d Gromov-Witten invariant, it is given as sum of contributions from tree graphs with d edges. We can easily see that number of tree graphs with d edges increases very fast as d rises. Therefore, computation of Gromov-Witten invariant via localization gets harder and harder as d rises. On the contrary, in localization computation of virtual structure constant of degree d , it is represented by a multi-variable residue integral given as contribution from line graph with d edges. Line graph with d edges is unique. Therefore, number of graphs needed for evaluation of genus 0 virtual structure constant of degree d is always one. Moreover, it is represented by multivariable residue integral, which is more handy for use of computers than the usual localization computation, which includes summation of characters of torus action. As can be seen in discussion in [14], mirror transformation from Gromov-Witten invariants to virtual structure constants can be interpreted as process of “cutting useless edges of tree graph into line graph “. In other words, mirror transformation from A-model to B-model is interpreted as process of “linearization of moduli space” in our context.

In this paper, we present final results of our trial to find multi-variable residue integral formula that corresponds to this “linearization of moduli space “ at the level of genus 1 Gromov-Witten invariants of projective hypersurface.

We don’t remember exactly when we began this trial, but it was around 2010. First, we tried to rewrite results of genus 1 Gromov-Witten invariants of CP^2 obtained from method of Graber and Pandharipande into residue integrals, and translated them into B-model form by using our mirror map presented in [14] (we had already obtained the formula of multi-point virtual structure constants). As was expected, this process had the effect of “cutting useless edges of tree graph with elliptic vertex and one loop graph “, and we observed that expected residue integral representation would be written as sum of contributions from graphs of type (i) and type (ii) presented in Section 3 of this paper. But we also found that we also have to introduce type (iii) and (iv) graphs presented in Section 3 in order to cancel error terms that appear in rewriting localization formulas of moduli space of stable maps into residue integrals. We called informally these error terms “diagonal anomaly”. It was very difficult to control this diagonal anomaly, and we temporarily stopped our trial. In spring of 2020, we managed to rewrite BCOV-Zinger formula as multivariable residue integral by using type (i) graph and colored one loop graphs. But residue integrals that correspond to colored one loop graphs contained geometrically unnatural factors from the point of view of localization of moduli space of quasimaps, and we didn’t publish these formulas as an official article. Then around winter of 2020, we found how to control “diagonal anomaly” in the CP^2 case, and wrote down final form of residue integral representation of elliptic virtual structure constants of CP^2 . But unfortunately, it had little connection with the residue integral representation of BCOV-Zinger formula we had at that time. In spring of 2023, adding new member Kuwata, we began to generalize the residue integral representation of elliptic virtual structure constants of CP^2 to projective Fano hypersurfaces. Generalization to higher dimensional projective spaces was easy. But it was hard to find satisfactory formulas for the case of Fano hypersurface M_N^k ($N > k$) with k greater than 1. In March of 2024, Kuwata found satisfactory residue integral representation of elliptic virtual structure constants of degree 1, which is applicable to all M_N^k ’s ($N > k$). Within two weeks from this discovery, we reached the residue integral representation of elliptic virtual structure constants presented in this paper. Moreover, we found that these formulas are also applicable to Calabi-Yau hypersurface M_k^k and result in geometrically natural residue integral representation of BCOV-Zinger formula. Then we decided to write down this paper as soon as possible.

As was explained in the previous paragraph, the process of construction of our residue integral formulas is purely heuristic. Therefore, we only write down set-up of our mirror computation, integrands of residue integrals associated with graphs and results of numerical tests of our conjectural formulas.

This paper is organised as follows. In Section 2, we explain set-up of our mirror computation of elliptic Gromov-Witten invariants of projective Fano hypersurfaces based on the results presented in [14]. In Section

3, we present our residue integral representation of elliptic virtual structure constants. In Subsection 3.1, we introduce four types of graphs used in our construction. In Subsection 3.2, we present integrands of residue integral representation associated with each type of graphs. In Section 4, we test our residue integral representation by numerical computation using computers. In Subsection 4.1, we present results in the case of Fano projective hypersurfaces and observe agreement with known results. In Subsection 4.2, we present results in the case of Calabi-Yau hypersurface M_k^k and obtain another representation of BCOV-Zinger formula. In Appendix A, we collect our numerical results for Fano 3-folds M_5^k ($k = 1, 2, 3, 4$) in Tables.

This paper is written from stand point of physicists, and the results presented are all conjectures from the point of view of mathematicians. In the sequel of future works, we try to give proofs of these conjectures. As can be seen from the formulas in this paper, we have to use mathematical processing software like Maple or Mathematica in order to obtain numerical results from our residue integrals. We plan to publish pdf copy of worksheet of maple program in Masao Jinzenji's home page of ReserchGate [15].

Acknowledgment We would like to thank Prof. Y. Kanda, Prof. S. Kobayashi, Prof. H. Iritani and Prof. K. Ono for valuable discussions.

2 Set-Up of Our Computation

Let M_N^k be a non-singular degree k hypersurface in CP^{N-1} . Apparently, its complex dimension is $N - 2$. In this paper, we compute genus 1 (elliptic) Gromov-Witten invariants of M_N^k ($N \geq k$) by using the mirror map constructed from multi-point virtual structure constants of M_N^k [14]. We denote by $h \in H^{1,1}(M_N^k, \mathbf{C})$ the hyperplane class of M_N^k .

Let us briefly review conjectures proposed in [14]. We first introduce the following polynomials:

$$\begin{aligned} e_k(x, y) &:= \prod_{j=0}^k (jx + (k-j)y), \\ w_a(x, y) &:= \frac{x^a - y^a}{x - y} = \sum_{j=0}^{a-1} x^j y^{a-1-j}, \end{aligned} \tag{2.2}$$

which play the role of building blocks of residue integral representation of genus 0 multi-point virtual structure constant $w(\mathcal{O}_{h^a} \mathcal{O}_{h^b} | \prod_{p=0}^{N-2} (\mathcal{O}_{h^p})^{m_p})_{0,d}$ of degree d . If $d \geq 1$, it is explicitly given as follows.

$$\begin{aligned} &w(\mathcal{O}_{h^a} \mathcal{O}_{h^b} | \prod_{p=0}^{N-2} (\mathcal{O}_{h^p})^{m_p})_{0,d} \\ &= \frac{1}{(2\pi\sqrt{-1})^{d+1}} \oint_{C_{z_0}} dz_0 \oint_{C_{z_1}} dz_1 \cdots \oint_{C_{z_d}} dz_d \\ &\quad \times (z_0)^a \left(\frac{\prod_{j=1}^d e_k(z_{j-1}, z_j)}{\prod_{i=1}^{d-1} k z_i (2z_i - z_{i-1} - z_{i+1})} \right) (z_d)^b \left(\prod_{p=0}^{N-2} \left(\sum_{n=1}^d w_p(z_{n-1}, z_n) \right)^{m_p} \right) \\ &\quad \times \prod_{q=0}^d \frac{1}{(z_q)^N}. \end{aligned} \tag{2.3}$$

In the above residue integral, $\oint_{C_{z_0}} dz_0$ and $\oint_{C_{z_d}} dz_d$ represent operation of taking residues at $z_0 = 0$ and $z_d = 0$ respectively. On the other hand, $\oint_{C_{z_j}} dz_j$ ($j = 1, 2, \dots, d-1$) means taking residues at $z_j = 0$ and $z_j = \frac{z_{j-1} + z_{j+1}}{2}$. In this case, the result of integration does not depend on order of integration.

If $d = 0$, it is 0 except for the following cases.

$$w(\mathcal{O}_{h^a}\mathcal{O}_{h^b}|\mathcal{O}_{h^c})_{0,0} = k\delta_{a+b+c,N-2}. \quad (2.4)$$

If $d \geq 1$, we can see the following condition holds.

$$w(\mathcal{O}_{h^a}\mathcal{O}_{h^b}|\prod_{p=0}^{N-2}(\mathcal{O}_{h^p})^{m_p})_{0,d} \neq 0 \implies a + b + \sum_{p=0}^{N-2}(p-1)m_p = N - 3 + (N - k)d. \quad (2.5)$$

Line Graph with d Edges and $(d+1)$ Normal Vertices



Figure 1: The Graph used in Computing Genus 0 Degree d Virtual Structure Constants

We can easily see the following characteristics of $w(\mathcal{O}_{h^a}\mathcal{O}_{h^b}|\prod_{p=0}^{N-2}(\mathcal{O}_{h^p})^{m_p})_{0,d}$:

$$w(\mathcal{O}_{h^a}\mathcal{O}_{h^b}|\prod_{p=0}^{N-2}(\mathcal{O}_{h^p})^{m_p})_{0,d} = \delta_{m_0,0} \cdot d^{m_1} w(\mathcal{O}_{h^a}\mathcal{O}_{h^b}|\prod_{p=2}^{N-2}(\mathcal{O}_{h^p})^{m_p})_{0,d} \quad (d \geq 1), \quad (2.6)$$

since we have $w_0(x, y) = 0$ and $w_1(x, y) = 1$. Next, we introduce “perturbed two-point function:

$$w(\mathcal{O}_{h^a}\mathcal{O}_{h^b})_0(x^0, x^1, \dots, x^{N-2}) := \sum_{m_0=0}^{\infty} \cdots \sum_{m_{N-2}=0}^{\infty} \sum_{d=0}^{\infty} w(\mathcal{O}_{h^a}\mathcal{O}_{h^b}|\prod_{p=0}^{N-2}(\mathcal{O}_{h^p})^{m_p})_{0,d} \prod_{q=0}^{N-2} \frac{(x^q)^{m_q}}{m_q!}. \quad (2.7)$$

(2.4) and (2.6) lead us to the following simplification:

$$\begin{aligned} & w(\mathcal{O}_{h^a}\mathcal{O}_{h^b})_0(x^0, x^1, \dots, x^{N-2}) \\ &= kx^{N-2-a-b} + \sum_{m_2=0}^{\infty} \cdots \sum_{m_{N-2}=0}^{\infty} \sum_{d=1}^{\infty} e^{dx^1} w(\mathcal{O}_{h^a}\mathcal{O}_{h^b}|\prod_{p=2}^{N-2}(\mathcal{O}_{h^p})^{m_p})_{0,d} \prod_{q=2}^{N-2} \frac{(x^q)^{m_q}}{m_q!}. \end{aligned} \quad (2.8)$$

Therefore, we only consider $w(\mathcal{O}_{h^a}\mathcal{O}_{h^b}|\prod_{p=2}^{N-2}(\mathcal{O}_{h^p})^{m_p})_{0,d}$ from now on. In [14], we defined “mirror map”, or coordinate change of deformation variables as follows.

$$t^p(x^1, x^2, \dots, x^{N-2}) := \frac{1}{k} w(\mathcal{O}_{h^{N-2-p}}\mathcal{O}_1)_0 \quad (p = 0, 1, \dots, N-2). \quad (2.9)$$

(2.8) tells us that the above mirror map has the following structure:

$$t^p(x^1, x^2, \dots, x^{N-2}) = x^p + \frac{1}{k} \sum_{m_2=0}^{\infty} \cdots \sum_{m_{N-2}=0}^{\infty} \sum_{d=1}^{\infty} e^{dx^1} w(\mathcal{O}_{h^{N-2-p}}\mathcal{O}_1|\prod_{p=2}^{N-2}(\mathcal{O}_{h^p})^{m_p})_{0,d} \prod_{q=2}^{N-2} \frac{(x^q)^{m_q}}{m_q!}. \quad (2.10)$$

This structure allows us to invert the mirror map:

$$x^p = x^p(t^0, t^1, \dots, t^{N-2}) \quad (p = 0, 1, \dots, N-2). \quad (2.11)$$

At this stage, we introduce Gromov-Witten invariant $\langle \prod_{a=0}^{N-2}(\mathcal{O}_{h^a})^{m_a} \rangle_{g,d}$ of genus g and degree d . In this paper, genus g is limited to take the value 0 or 1. We omit here rigorous definition of this invariant, but we

mention some of its characteristics. In general, it is non-zero only if the following condition is satisfied.

$$\begin{aligned} \langle \prod_{a=0}^{N-2} (\mathcal{O}_{h^a})^{m_a} \rangle_{0,d} \neq 0 &\implies \sum_{a=0}^{N-2} m_a(a-1) = N-5 + d(N-k), \\ \langle \prod_{a=0}^{N-2} (\mathcal{O}_{h^a})^{m_a} \rangle_{1,d} \neq 0 &\implies \sum_{a=0}^{N-2} m_a(a-1) = d(N-k). \end{aligned} \quad (2.12)$$

If $d = 0$ and $g = 0$, it is zero except for the following case:

$$\langle \mathcal{O}_{h^a} \mathcal{O}_{h^b} \mathcal{O}_{h^c} \rangle_{0,d} = k \cdot \delta_{N-2, a+b+c}. \quad (2.13)$$

If $d = 0$ and $g = 1$, it is zero except for the following case:

$$\langle \mathcal{O}_h \rangle_{1,0} = -\frac{1}{24} \int_{M_N^k} h \wedge c_{N-3}(T' M_N^k), \quad (2.14)$$

where $c_{N-3}(T' M_N^k)$ is the second-top Chern class of $T' M_N^k$, the homomorphic tangent bundle of M_N^k . If $d \geq 1$, it satisfies the following equality:

$$\langle \prod_{a=0}^{N-2} (\mathcal{O}_{h^a})^{m_a} \rangle_{g,d} = \delta_{m_0,0} d^{m_1} \langle \prod_{a=2}^{N-2} (\mathcal{O}_{h^a})^{m_a} \rangle_{g,d}. \quad (2.15)$$

Then we restate the conjecture on genus 0 Gromov-Witten invariants of M_N^k proposed in [14]. As was done in (2.7), we introduce perturbed two-point genus 0 Gromov-Witten invariant:

$$\begin{aligned} &\langle \mathcal{O}_{h^a} \mathcal{O}_{h^b} \rangle_0(t^0, t^1, \dots, t^{N-2}) \\ := &\sum_{m_0=0}^{\infty} \dots \sum_{m_{N-2}=0}^{\infty} \sum_{d=0}^{\infty} \langle \mathcal{O}_{h^a} \mathcal{O}_{h^b} \prod_{p=0}^{N-2} (\mathcal{O}_{h^p})^{m_p} \rangle_{0,d} \prod_{q=0}^{N-2} \frac{(t^q)^{m_q}}{m_q!} \\ = &kt^{N-2-a-b} + \sum_{m_2=0}^{\infty} \dots \sum_{m_{N-2}=0}^{\infty} \sum_{d=1}^{\infty} e^{dt^1} \langle \mathcal{O}_{h^a} \mathcal{O}_{h^b} \prod_{p=2}^{N-2} (\mathcal{O}_{h^p})^{m_p} \rangle_{0,d} \prod_{q=2}^{N-2} \frac{(t^q)^{m_q}}{m_q!}. \end{aligned} \quad (2.16)$$

Then the conjecture is given as follows:

Conjecture 1 [14]

$$\langle \mathcal{O}_{h^a} \mathcal{O}_{h^b} \rangle_0(t^0, t^1, \dots, t^{N-2}) = w(\mathcal{O}_{h^a} \mathcal{O}_{h^b})_0(x^0(t^*), x^1(t^*), \dots, x^{N-2}(t^*)), \quad (2.17)$$

where $x^p(t^*)$ is abbreviation of $x^p(t^0, t^1, \dots, t^{N-2})$ given in (2.11).

This conjecture gives us a recipe for computing genus 0 Gromov-Witten invariants of M_N^k and it has been confirmed numerically for low degrees in many examples.

In this paper, we also introduce elliptic multi-point virtual structure constant $w(\prod_{p=0}^{N-2} (\mathcal{O}_{h^p})^{m_p})_{1,d}$ of degree d , whose definition in the $d \geq 1$ case will be given in the next section. In the $d = 0$ case, it is 0 except for the following case.

$$w(\mathcal{O}_h)_{1,0} := \langle \mathcal{O}_h \rangle_{1,0} = -\frac{1}{24} \int_{M_N^k} h \wedge c_{N-3}(T' M_N^k). \quad (2.18)$$

In the $d \geq 1$ case, we can show the following characteristics of $w(\prod_{p=0}^{N-2} (\mathcal{O}_{h^p})^{m_p})_{1,d}$ by using the definition which will be given in the next section.

$$w\left(\prod_{p=0}^{N-2} (\mathcal{O}_{h^p})^{m_p}\right)_{1,d} = \delta_{m_0,0} d^{m_1} w\left(\prod_{p=2}^{N-2} (\mathcal{O}_{h^p})^{m_p}\right)_{1,d}. \quad (2.19)$$

We then introduce generating function of $w(\prod_{p=0}^{N-2} (\mathcal{O}_{h^p})^{m_p})_{1,d}$'s:

$$\begin{aligned}
& F_{1,vir.}^{N,k,B}(x^0, x^1, \dots, x^{N-2}) \\
& := \sum_{d=0}^{\infty} \sum_{m_0=0}^{\infty} \cdots \sum_{m_{N-2}=0}^{\infty} w\left(\prod_{p=0}^{N-2} (\mathcal{O}_{h^p})^{m_p}\right)_{1,d} \prod_{q=0}^{N-2} \frac{(x^q)^{m_q}}{m_q!} \\
& = -\frac{1}{24} \left(\int_{M_N^k} h \wedge c_{N-3}(T' M_N^k) \right) x^1 + \sum_{d=1}^{\infty} \sum_{m_2=0}^{\infty} \cdots \sum_{m_{N-2}=0}^{\infty} e^{dx^1} w\left(\prod_{p=2}^{N-2} (\mathcal{O}_{h^p})^{m_p}\right)_{1,d} \prod_{q=2}^{N-2} \frac{(x^q)^{m_q}}{m_q!}. \quad (2.20)
\end{aligned}$$

In going from the second line to the third line, we used (2.18) and (2.19). On the other hand, we also prepare generating function of genus 1 Gromov-Witten invariants of M_N^k .

$$\begin{aligned}
& F_1^{N,k,A}(t^0, t^1, \dots, t^{N-2}) \\
& := \sum_{d=0}^{\infty} \sum_{m_0=0}^{\infty} \cdots \sum_{m_{N-2}=0}^{\infty} \langle \prod_{p=0}^{N-2} (\mathcal{O}_{h^p})^{m_p} \rangle_{1,d} \prod_{q=0}^{N-2} \frac{(t^q)^{m_q}}{m_q!} \\
& = -\frac{1}{24} \left(\int_{M_N^k} h \wedge c_{N-3}(T' M_N^k) \right) t^1 + \sum_{d=1}^{\infty} \sum_{m_2=0}^{\infty} \cdots \sum_{m_{N-2}=0}^{\infty} e^{dt^1} \langle \prod_{p=2}^{N-2} (\mathcal{O}_{h^p})^{m_p} \rangle_{1,d} \prod_{q=2}^{N-2} \frac{(t^q)^{m_q}}{m_q!}. \quad (2.21)
\end{aligned}$$

With this set-up, we state our main conjecture in this paper.

Conjecture 2 (Main Conjecture)

$$F_1^{N,k,A}(t^0, t^1, \dots, t^{N-2}) = F_{1,vir.}^{N,k,B}(x^0(t^*), x^1(t^*), \dots, x^{N-2}(t^*)), \quad (2.22)$$

where $x^p(t^*)$ is the inversion of the mirror map given in (2.11).

Together with the explicit definition of the elliptic multi-point veirtual structure constants in the next section, this gives us a recipe for computing genus 1 Gromov-Witten invariants of M_N^k . In Section 4, we test this conjecture by comparing our predictions with known results of genus 1 Gromov-Witten invariants of M_N^k .

3 Perturbative Definitions of Elliptic Multi-Point Virtual Structure Constants

3.1 Graphs

In order to explain structure of Graphs used in our computation, we first introduce partition of positive integer d .

$$\sigma = (d_1, d_2, \dots, d_{l(\sigma)}) \quad (d_1 \geq d_2 \geq \dots \geq d_{l(\sigma)} > 0, \sum_{i=1}^{l(\sigma)} d_i = d), \quad (3.23)$$

where we call $l(\sigma)$ length of the partition σ . We denote by P_d set of partitions of positive integer d .

$$P_d := \{ \sigma = (d_1, d_2, \dots, d_{l(\sigma)}) \mid d_1 \geq d_2 \geq \dots \geq d_{l(\sigma)} > 0, \sum_{i=1}^{l(\sigma)} d_i = d \}. \quad (3.24)$$

For later use, we define symmetry factor associated with $\sigma \in P_d$:

$$\text{Sym}(\sigma) := \frac{(l(\sigma) - 1)!}{\prod_{j=1}^d \text{mul}(\sigma; j)!} \quad (\sigma \in P_d), \quad (3.25)$$

where $\text{mul}(\sigma; j)$ is multiplicity of i ($1 \leq i \leq d$) that appears in $\sigma = (d_1, d_2, \dots, d_{l(\sigma)})$. We also define the following rational number:

$$R_{N,k}(d) := \frac{(N-1)kd - 2(Nk-1)}{2kd^2}, \quad (3.26)$$

which plays a role of symmetry factor for specific graphs.

In our computation, we use graphs with normal edges represented as “—” and three types of vertices.

- (i) Normal Vertex
- (ii) Elliptic Vertex
- (iii) Cluster Vertex of degree d

These are graphically represented by the following symbols.




Figure 2: Vertices used in Our Construction

Single edge is assigned with degree 1. Single normal vertex, elliptic vertex and cluster vertex of degree d are assigned with degree 0, 0 and d respectively. Graphs used in computing elliptic virtual constants of degree d are classified by the following four types.

- (i) Star graph associated with $\sigma \in P_d$ having elliptic vertex as its center
- (ii) Loop graph with d edges and d normal vertices ($d \geq 2$)
- (iii) Star graph associated with $\sigma \in P_{d-f}$ ($1 \leq f \leq d-1$) having cluster vertex of degree f as its center
- (iv) Graph which consists of a single cluster vertex of degree d

We write down examples of these four types in the following figures.

Type (i) associated with $\sigma = (3,3,2,1) \in P_9$ of degree 9




Figure 3: An example of Type (i) Graph

Type (ii) Loop graph of degree 5




Figure 4: An example of Type (ii) Graph

Type (iii) associated with $\sigma = (3,2,1) \in P_6$ of degree 8




Figure 5: An example of Type (iii) Graph

Type (iv) of degree 4



Figure 6: An example of Type (iv) Graph

We then write down all the graphs used in computing elliptic virtual constants of $d = 1, 2, 3$.




Figure 7: Graphs used in Computing Elliptic Degree 1, 2 Virtual Structure Constants




Figure 8: Graphs used in Computing Elliptic Degree 3 Virtual Structure Constants

We can easily see that number of graphs used in computing elliptic virtual constants of degree d (which we denote by N_d) is given by,

$$N_d = \begin{cases} 2 & (d = 1) \\ \sum_{j=1}^d (P_j)^\# + 1 & (d \geq 2) \end{cases} \quad (3.27)$$

Therefore, generating function of N_d is given by,

$$\begin{aligned} 1 + \sum_{d=1}^{\infty} N_d q^d &= \frac{q^2}{1-q} + \frac{1}{1-q} \left(\prod_{m=1}^{\infty} \frac{1}{1-q^m} \right) \\ &= 1 + 2q + 5q^2 + 8q^3 + 13q^4 + 20q^5 + 31q^6 + 46q^7 + 68q^8 + 98q^9 + 140q^{10} + \dots \end{aligned} \quad (3.28)$$

We can see that number of graphs increases rather slowly as d rises.

3.2 Integrands of Residue Integrals

Elliptic virtual structure constant $w(\prod_{a=0}^{N-2} (\mathcal{O}_{h^a})^{m_a})_{1,d}$ is computed by summing up contributions from these graphs of degree d . Contribution from each graph is given as residue integral of multi variables determined from the graph. Let us write down integrand of the residue integral associated with the four types of graphs,

We begin with type (i) graph associated with $\sigma = (d_1, d_2, \dots, d_l) \in P_d$. In this case, we prepare d complex variables z_0 and $z_{i,j}$ ($1 \leq i \leq l$, $1 \leq j \leq d_i$). z_0 is associated with center elliptic vertex and $z_{i,j}$ is associated with vertex in a tail of star graph with d_i edges. It is well-known that total Chern class of $T'M_N^k$ is given by,

$$c(T'M_N^k) = \frac{(1+h)^N}{1+kh}, \quad (3.29)$$

and top chern class $c_{N-2}(T'M_N^k)$ is given as h^{N-2} multiplied by coefficient of h^{N-2} in $c(T'M_N^k)$. Then we define $c_T(z)$, which is monomial in z , by the following equality.

$$c_T(h) = c_{N-2}(T'M_N^k). \quad (3.30)$$

Then integrand associated with the graph is given as follows.

$$\begin{aligned} & \text{Sym}(\sigma) \frac{1}{24} \frac{c_T(z_0)}{(kz_0)^{l-1}} \left(\prod_{i=1}^l \frac{e_k(z_0, z_{i,1})}{z_{i,1} - z_0} \right) \left(\prod_{i=1}^l \left(\prod_{j=1}^{d_i-1} \frac{e_k(z_{i,j}, z_{i,j+1})}{(2z_{i,j} - z_{i,j-1} - z_{i,j+1})kz_{i,j}} \right) \right) \\ & \times \left(\frac{1}{(z_0)^N} \right) \left(\prod_{i=1}^l \prod_{j=1}^{d_i} \frac{1}{(z_{i,j})^N} \right) \left(\prod_{a=2}^{N-2} \left(\sum_{i=1}^l \sum_{j=1}^{d_i} w_a(z_{i,j-1}, z_{i,j}) \right)^{m_a} \right), \\ & (z_{i,0} = z_0 \ (i = 1, 2, \dots, l)). \end{aligned} \quad (3.31)$$

Integrand for type (ii) loop graph of degree d is given by,

$$\begin{aligned} & \frac{1}{2d} \left(\prod_{j=1}^d \frac{e_k(z_j, z_{j+1})}{(2z_j - z_{j-1} - z_{j+1})kz_j} \right) \left(\prod_{a=2}^{N-2} \left(\sum_{j=1}^d w_a(z_j, z_{j+1}) \right)^{m_a} \right) \cdot \left(\prod_{j=1}^d \frac{1}{(z_j)^N} \right), \\ & (z_0 = z_d, z_{d+1} = z_1). \end{aligned} \quad (3.32)$$

Integrand for type (iii) graph of degree d with cluster vertex of degree f ($1 \leq f \leq d-1$) associated with $\sigma = (d_1, d_2, \dots, d_l) \in P_{d-f}$ is given as follows.

$$\begin{aligned} & \text{Sym}(\sigma) \frac{1}{24} \left(-\frac{N-1}{N} \frac{1}{w^N} - \frac{N+1}{N} \frac{1}{(z_0)^N} \right) \frac{1}{(w-z_0)^2} \frac{e_k(w, z_0)}{kw} \left(\frac{e(z_0, z_0)}{kz_0} \right)^{f-1} \\ & \times \frac{1}{(kz_0)^{l-1}} \left(\prod_{i=1}^l \frac{e_k(z_0, z_{i,1})}{z_{i,1} - z_0} \right) \left(\prod_{i=1}^l \left(\prod_{j=1}^{d_i-1} \frac{e_k(z_{i,j}, z_{i,j+1})}{(2z_{i,j} - z_{i,j-1} - z_{i,j+1})kz_{i,j}} \right) \right) \left(\frac{1}{(z_0)^{N(f-1)}} \right) \left(\prod_{i=1}^l \prod_{j=1}^{d_i} \frac{1}{(z_{i,j})^N} \right) \\ & \times \left(\prod_{a=2}^{N-2} \left(w_a(w, z_0) + (f-1)w_a(z_0, z_0) + \sum_{i=1}^l \sum_{j=1}^{d_i} w_a(z_{i,j-1}, z_{i,j}) \right)^{m_a} \right), \\ & (z_{i,0} = z_0 \ (i = 1, 2, \dots, l)). \end{aligned} \quad (3.33)$$

Lastly, integrand for type (iv) graph of degree d is given by,

$$\begin{aligned} & \frac{1}{24} R_{N,k}(d) \left(\frac{e_k(z_0, z_0)}{kz_0} \right)^d \left(\prod_{a=2}^{N-2} (dw_a(z_0, z_0))^{m_a} \right) \frac{1}{(z_0)^{Nd+1}} \\ & = \frac{1}{24} R_{N,k}(d) \frac{k^{kd} \prod_{a=2}^{N-2} (da)^{m_a}}{z_0}. \end{aligned} \quad (3.34)$$

Let Γ be the graph of degree d introduced so far, and Graph_d be set of all the graphs of degree d from type (i) to (iv). Then we denote by f_Γ the integrand defined above. In the following, we define a map $\text{Res} : f_\Gamma \rightarrow \mathbf{R}$, which corresponds to operation of taking residue of f_Γ .

Definition 1 $Res : f_\Gamma \rightarrow \mathbf{R}$ is defined for each type of graphs as follows.

- (i) We first take residue of f_Γ at $z_0 = 0$. Next we take residue of the resulting function at $z_{i,j} = 0$ and $z_{i,j} = \frac{z_{i,j-1} + z_{i,j+1}}{2}$ and add them up sequently in ascending order of j ($1 \leq j \leq d_i - 1$). Lastly, we take residue of the resulting function at $z_{i,d_i} = 0$. The order among different i 's does not matter.
- (ii) We take residue of f_Γ at $z_j = 0$ and $z_j = \frac{z_{j-1} + z_{j+1}}{2}$ and add them up sequently in ascending order of j ($1 \leq j \leq d$).
- (iii) We first take residue of f_Γ at $w = z_0$. Then remaining process is the same as the case of type (i).
- (iv) We take residue of f_Γ at $z_0 = 0$.

Then elliptic multi-point veirtual structure constant is defined as follows.

Definition 2

$$w\left(\prod_{a=2}^{N-2} (\mathcal{O}_{h^a})^{m_a}\right)_{1,d} := \sum_{\Gamma \in Graph_d} Res(f_\Gamma) \quad (d \geq 1). \quad (3.35)$$

In the next section, we explicitly compute $w\left(\prod_{a=2}^{N-2} (\mathcal{O}_{h^a})^{m_a}\right)_{1,d}$'s by using this definition for various M_N^k 's and test Conjecture 2 by comparing our predictions with known results.

4 Numerical Tests by Various Examples

4.1 Tests for Fano Hypersurfaces

First, we test Conjecture 2 by using projective surface $CP^2 = M_4^1$, quadric surface M_4^2 and cubic surface M_4^3 as examples. We present here explicit processes of computation in the CP^2 case. In this case, generating function of elliptic virtual structure constants $F_{1,vir}^{4,1,B}$ has the following structure,

$$F_{1,vir}^{4,1,B}(x^0, x^1, x^2) = -\frac{1}{8}x^1 + \sum_{d=1}^{\infty} e^{dx^1} w((\mathcal{O}_{h^2})^{3d})_{1,d} \frac{(x^2)^{3d}}{(3d)!}, \quad (4.36)$$

where we used $c(T'M_4^1) = (1+h)^3 = 1+3h+3h^2$ and the condition:

$$w((\mathcal{O}_{h^2})^a)_{1,d} \neq 0 \implies a = (N-k)d = 3d. \quad (4.37)$$

We abbreviate $w((\mathcal{O}_{h^2})^a)_{1,d}$ to w_a and present the results of computation by using Definition 2 in Table 1. Therefore, $F_{1,vir}^{4,1,B}$ is explicitly given by,

$$\begin{aligned} & F_{1,vir}^{4,1,B}(x^0, x^1, x^2) \\ = & -\frac{1}{8}x^1 - \frac{1}{16}e^{x^1}(x^2)^3 - \frac{7}{80}e^{2x^1}(x^2)^6 - \frac{77789}{362880}e^{3x^1}(x^2)^9 - \frac{21344159}{31933440}e^{4x^1}(x^2)^{12} - \frac{15774542951}{6604416000}e^{5x^1}(x^2)^{15} - \dots \end{aligned} \quad (4.38)$$

On the other hand, the mirror map is evaluated by using (2.3) as follows.

$$\begin{aligned} t^0 &= x^0 + \frac{1}{2}(x^2)^2 e^{x^1} + \frac{8}{15}(x^2)^5 e^{2x^1} + \frac{983}{840}(x^2)^8 e^{3x^1} + \frac{4283071}{1247400}(x^2)^{11} e^{4x^1} + \frac{4019248213}{340540200}(x^2)^{14} e^{5x^1} + \dots, \\ t^1 &= x^1 + \frac{1}{2}(x^2)^3 e^{x^1} + \frac{7}{10}(x^2)^6 e^{2x^1} + \frac{2593}{1512}(x^2)^9 e^{3x^1} + \frac{2668063}{498960}(x^2)^{12} e^{4x^1} + \frac{120501923}{6306300}(x^2)^{15} e^{5x^1} + \dots, \\ t^2 &= x^2 + \frac{1}{4}(x^2)^4 e^{x^1} + \frac{33}{70}(x^2)^7 e^{2x^1} + \frac{16589}{12600}(x^2)^{10} e^{3x^1} + \frac{143698921}{32432400}(x^2)^{13} e^{4x^1} + \frac{75631936691}{4540536000}(x^2)^{16} e^{5x^1} + \dots. \end{aligned} \quad (4.39)$$

In order to obtain $F_1^{4,1,A}(t^0, t^1, t^2)$ from Conjecture 2, we only need $x^1 = x^1(t^1, t^2), x^2 = x^2(t^1, t^2)$ since both $F_{1, \text{vir.}}^{4,1,B}$ and $F_1^{4,1,A}$ do not depend on x^0 and t^0 . After inverting (4.39), they are given as follows:

$$\begin{aligned} x^1 &= t^1 - \frac{1}{2}(t^2)^3 e^{t^1} - \frac{3}{40}(t^2)^6 e^{2t^1} - \frac{3827}{30240}(t^2)^9 e^{3t^1} - \frac{4914517}{19958400}(t^2)^{12} e^{4t^1} - \frac{10460869973}{18162144000}(t^2)^{15} e^{5t^1} - \dots \\ x^2 &= t^2 - \frac{1}{4}(t^2)^4 e^{t^1} - \frac{27}{280}(t^2)^7 e^{2t^1} - \frac{7811}{50400}(t^2)^{10} e^{3t^1} - \frac{82505777}{259459200}(t^2)^{13} e^{4t^1} - \frac{1014012107}{1320883200}(t^2)^{16} e^{5t^1} - \dots, \end{aligned} \quad (4.40)$$

By substituting (4.40) into $F_{1, \text{vir.}}^{4,1,B}(x^1, x^2)$, we obtain from Conjecture 2,

$$\begin{aligned} F_1^{4,1,A}(t^1, t^2) &= F_{1, \text{vir.}}^{4,1,B}(x^1(t^1, t^2), x^2(t^1, t^2)) \\ &= -\frac{1}{8}t^1 + \frac{1}{362880}(t^2)^9 e^{3t^1} + \frac{1}{2128896}(t^2)^{12} e^{4t^1} + \frac{173}{2594592000}(t^2)^{15} e^{5t^1} + \dots. \end{aligned} \quad (4.41)$$

Then we have,

$$\begin{aligned} \langle (\mathcal{O}_{h^2})^3 \rangle_{1,1} &= 0, \langle (\mathcal{O}_{h^2})^6 \rangle_{1,2} = 0, \langle (\mathcal{O}_{h^2})^9 \rangle_{1,3} = \frac{1}{362880} \cdot 9! = 1, \langle (\mathcal{O}_{h^2})^{12} \rangle_{1,4} = \frac{1}{2128896} \cdot 12! = 225, \\ \langle (\mathcal{O}_{h^2})^{15} \rangle_{1,5} &= \frac{173}{2594592000} \cdot 15! = 87192. \end{aligned} \quad (4.42)$$

These results are summarized in Table 1, where $\langle (\mathcal{O}_{h^2})^a \rangle_{1,d}$ is abbreviated to $N_{d,a}^1$.

Table 1: M_4^1

d	a	$N_{d,a}^1$	w_a
1	3	0	$-\frac{3}{8}$
2	6	0	-63
3	9	1	-77789
4	12	225	-320162385
5	15	87192	-3123359504298

Thses results coincide with those presented in the paper [6] by Getzler. In this paper, we computed only up to $d = 5$ because running time of computers to evaluate residue integrals increases rather fast as degree rises.

As for quadric surface and cubic surface, we only present structure of $F_{1, \text{vir.}}^{N,k,B}$'s, mirror maps and Tables of w_a 's and $N_{d,a}^1$'s. Processes of computation are the same as those of M_4^1 .

$F_{1, \text{vir.}}^{4,2,B}$ and $F_{1, \text{vir.}}^{4,3,B}$ have the following structure.

$$\begin{aligned} F_{1, \text{vir.}}^{4,2,B}(x^0, x^1, x^2) &= -\frac{1}{6}x^1 + \sum_{d=1}^{\infty} e^{dx^1} w_{2d} \frac{(x^2)^{2d}}{(2d)!}, \\ F_{1, \text{vir.}}^{4,3,B}(x^0, x^1, x^2) &= -\frac{1}{8}x^1 + \sum_{d=1}^{\infty} e^{dx^1} w_d \frac{(x^2)^d}{d!}. \end{aligned} \quad (4.43)$$

Mirror map of M_4^2 is given by,

$$\begin{aligned} t^0 &= x^0 + 2x^2 e^{x^1} + 10(x^2)^3 e^{2x^1} + \frac{320}{3}(x^2)^5 e^{3x^1} + \frac{53856}{35}(x^2)^7 e^{4x^1} + \frac{74056288}{2835}(x^2)^9 e^{5x^1} + \dots, \\ t^1 &= x^1 + 3(x^2)^2 e^{x^1} + \frac{131}{6}(x^2)^4 e^{2x^1} + \frac{12329}{45}(x^2)^6 e^{3x^1} + \frac{121475}{28}(x^2)^8 e^{4x^1} + \frac{370005883}{4725}(x^2)^{10} e^{5x^1} + \dots, \\ t^2 &= x^2 + 2(x^2)^3 e^{x^1} + \frac{313}{15}(x^2)^5 e^{2x^1} + \frac{10764}{35}(x^2)^7 e^{3x^1} + \frac{15178391}{2835}(x^2)^9 e^{4x^1} + \frac{458817242}{4455}(x^2)^{11} e^{5x^1} + \dots, \end{aligned} \quad (4.44)$$

and that of M_4^3 is given as follows.

$$\begin{aligned}
t^0 &= x^0 + 6e^{x^1} + 144x^2e^{2x^1} + 7398(x^2)^2e^{3x^1} + 520344(x^2)^3e^{4x^1} + 43392510(x^2)^4e^{5x^1} + \dots, \\
t^1 &= x^1 + 21x^2e^{x^1} + \frac{1611}{2}(x^2)^2e^{2x^1} + 52191(x^2)^3e^{3x^1} + \frac{16915311}{4}(x^2)^4e^{4x^1} + 388313757(x^2)^5e^{5x^1} + \dots, \\
t^2 &= x^2 + 21(x^2)^2e^{x^1} + 1305(x^2)^3e^{2x^1} + 106056(x^2)^4e^{3x^1} + \frac{49255533}{5}(x^2)^5e^{4x^1} + \frac{4964539329}{5}(x^2)^6e^{5x^1} + \dots.
\end{aligned} \tag{4.45}$$

Tables of w_a 's and $N_{d,a}^1$'s are given as follows.

Table 2: M_4^2

d	a	$N_{d,a}^1$	$N_{d,a}^1/2^a$	w_a
1	2	0	0	-1
2	4	0	0	$-\frac{262}{3}$
3	6	0	0	$-\frac{98632}{3}$
4	8	256	1	-29153744
5	10	40960	40	-47360066944

Table 3: M_4^3

d	a	$N_{d,a}^1$	$N_{d,a}^1/3^a$	w_a
1	1	0	0	$-\frac{21}{8}$
2	1	0	0	$-\frac{1611}{8}$
3	3	27	1	$-\frac{156465}{4}$
4	4	2187	27	$-\frac{50682753}{4}$
5	5	183708	756	-5815337247

In these tables, $N_{d,a}^1/k^a$'s are presented because Poincare dual of a point in M_N^k is given by $\frac{1}{k}h^2$ instead of h^2 . Therefore, $N_{d,a}^1/k^a$'s turn out to be integers.

Next, we test Conjecture 2 for $N = 5$ cases, i.e., complex Fano 3-folds: $M_5^1, M_5^2, M_5^3, M_5^4$. In these cases, we compute elliptic virtual structure constant $w((\mathcal{O}_{h^2})^a(\mathcal{O}_{h^3})^b)_{1,d}$ and genus 1 Gromov-Witten invariant $\langle (\mathcal{O}_{h^2})^a(\mathcal{O}_{h^3})^b \rangle_{1,d}$. These are non-zero only if the following condition is satisfied:

$$a + 2b = (N - k)d = (5 - k)d. \tag{4.46}$$

We abbreviate these to $w_{a,b}$ and $N_{d,a,b}^1$ and present results of numerical computations by using Conjecture 2 in Tables. These are collected in Appendix A. In 3-fold cases, $N_{d,a,b}^1$ is not always an integer because of degenerate contributions from genus 0 Gromov-Witten invariants $N_{d,a,b}^0 = \langle (\mathcal{O}_{h^2})^a(\mathcal{O}_{h^3})^b \rangle_{0,d}$. It is non zero only if (4.46) is satisfied. According to the paper [18] by Pandharipande, the contribution is given by,

$$-\frac{1}{24}((N - k)d - 2)N_{d,a,b}^0. \tag{4.47}$$

Hence we expect that,

$$\frac{1}{24}((N - k)d - 2)N_{d,a,b}^0 + N_{d,a,b}^1, \tag{4.48}$$

becomes integer because it is expected to count number of genus 1 curves in M_N^k that satisfies ‘‘passing-through’’ conditions imposed by operator insertions. We can observe that this combination is integer-valued in the Tables presented in Appendix A. Results of $M_5^1 = CP^3$ coincide with those presented in the paper [6] by Getzler, and results of M_5^3 coincide with those obtained from Virasoro Conjecture by Eguchi et al. [9]. In comparing our results

with those presented in [9], we have to divide $N_{d,a,b}^1$ and $N_{d,a,b}^0$ by $k^{a+b} = 3^{a+b}$. This is because fundamental integral cohomology classes of $H^{2,2}(M_5^k, \mathbf{C})$ and $H^{3,3}(M_5^k, \mathbf{C})$ are given by $\frac{1}{k}h^2$ and $\frac{1}{k}h^3$ respectively. In [9], $\frac{1}{k}h^2$ and $\frac{1}{k}h^3$ are used as cohomology class that corresponds to operator insertions, but we use h^2 and h^3 instead. Hence we have to divide our results by k^{a+b} . We omit mirror maps used in deriving our results of Fano 3-folds, which we leave to readers as exercises.

4.2 Tests for Calabi-Yau Hypersurfaces and Relation to BCOV-Zinger formula

In this subsection, we consider Calabi-Yau hypersurface M_k^k in CP^{k-1} . Let us introduce the following power series in e^x , which was originally introduced in [3].

$$\tilde{L}_m^{k,k}(e^x) := 1 + \sum_{d=1}^{\infty} \frac{w(\mathcal{O}_{h^{k-2-m}}\mathcal{O}_{h^{m-1}}|\mathcal{O}_h)_{0,d}}{k} e^{dx}. \quad (4.49)$$

In this case, we also introduce two-point virtual structure constant defined by,

$$\begin{aligned} & w(\mathcal{O}_{h^a}\mathcal{O}_{h^b})_{0,d} \\ = & \frac{1}{(2\pi\sqrt{-1})^{d+1}} \oint_{C_{z_0}} dz_0 \oint_{C_{z_1}} dz_1 \cdots \oint_{C_{z_d}} dz_d (z_0)^a \left(\frac{\prod_{j=1}^d e_k(z_{j-1}, z_j)}{\prod_{i=1}^{d-1} k z_i (2z_i - z_{i-1} - z_{i+1})} \right) (z_d)^b \left(\prod_{j=0}^d \frac{1}{(z_j)^N} \right), \end{aligned} \quad (4.50)$$

where integration paths are the same as those used in defining $w(\mathcal{O}_{h^a}\mathcal{O}_{h^b}|\prod_{p=0}^{N-2}(\mathcal{O}_{h^p})^{m_p})_{0,d}$. It satisfies the condition:

$$w(\mathcal{O}_{h^a}\mathcal{O}_{h^b})_{0,d} \neq 0 \implies a + b = k - 3. \quad (4.51)$$

Then we define two-point virtual structure constant perturbed by x :

$$w(\mathcal{O}_{h^{k-2-m}}\mathcal{O}_{h^{m-1}})_0(x) := kx + \sum_{d=1}^{\infty} e^{dx} w(\mathcal{O}_{h^{k-2-m}}\mathcal{O}_{h^{m-1}})_{0,d}. \quad (4.52)$$

As was shown in [11, 13], the mirror map $t = t(x)$ used in mirror computation of Gromov-Witten invariants of M_k^k is given by,

$$t(x) = \frac{1}{k} w(\mathcal{O}_{h^{k-3}}\mathcal{O}_1)_0(x). \quad (4.53)$$

In [12], we proved that Gromov-Witten invariant $\langle \mathcal{O}_{h^{k-2-m}}\mathcal{O}_{h^{m-1}} \rangle_{0,d}$ is computed via the following equality:

$$kt + \sum_{d=1}^{\infty} e^{dt} \langle \mathcal{O}_{h^{k-2-m}}\mathcal{O}_{h^{m-1}} \rangle_{0,d} = w(\mathcal{O}_{h^{k-2-m}}\mathcal{O}_{h^{m-1}})_0(x(t)), \quad (4.54)$$

where $x = x(t)$ is inversion of the mirror map $t = t(x)$. We can easily see the following relation.

$$w(\mathcal{O}_{h^a}\mathcal{O}_{h^b}|\mathcal{O}_h)_{0,d} = d \cdot w(\mathcal{O}_{h^a}\mathcal{O}_{h^b})_{0,d}. \quad (4.55)$$

Hence the generating function $\tilde{L}_m^{k,k}(e^x)$ is related to $w(\mathcal{O}_{h^{k-2-m}}\mathcal{O}_{h^{m-1}})_0(x)$ in the following way.

$$\tilde{L}_m^{k,k}(e^x) := \frac{d}{dx} \frac{w(\mathcal{O}_{h^{k-2-m}}\mathcal{O}_{h^{m-1}})_0(x)}{k}. \quad (4.56)$$

Especially, we have,

$$\tilde{L}_1^{k,k}(e^x) := \frac{dt}{dx}. \quad (4.57)$$

We also mention the following equality that was proved in [13].

$$\tilde{L}_0^{k,k}(e^x) := 1 + \sum_{d=1}^{\infty} \frac{w(\mathcal{O}_{h^{k-2}}\mathcal{O}_{h^{-1}}|\mathcal{O}_h)_{0,d}}{k} e^{dx} = \sum_{d=0}^{\infty} \frac{(kd)!}{(d!)^k} e^{dx}. \quad (4.58)$$

With this set-up, BCOV-Zinger formula[1, 20] is given as follows.

$$\begin{aligned} F_1^{k,k,B}(x) &= -\frac{1}{24} \left(\int_{M_k^k} c_{k-3}(T'M_k^k) \wedge h \right) x + \frac{1}{24} \chi(M_k^k) \log \left(\tilde{L}_0^{k,k}(e^x) \right) \\ &\quad - \frac{k-1}{48} \log(1 - k^k e^x) - \sum_{p=0}^{\frac{k-3}{2}} \frac{(k-1-2p)^2}{8} \log \left(\tilde{L}_p^{k,k}(e^x) \right) \quad (k : \text{odd}), \\ F_1^{k,k,B}(x) &= -\frac{1}{24} \left(\int_{M_k^k} c_{k-3}(T'M_k^k) \wedge h \right) x + \frac{1}{24} \chi(M_k^k) \log \left(\tilde{L}_0^{k,k}(e^x) \right) \\ &\quad - \frac{k-4}{48} \log(1 - k^k e^x) - \sum_{p=0}^{\frac{k-4}{2}} \frac{(k-2p)(k-2p-2)}{8} \log \left(\tilde{L}_p^{k,k}(e^x) \right) \quad (k : \text{even}), \end{aligned} \quad (4.59)$$

where $\chi(M_k^k)$ is Euler number of M_k^k given by,

$$\chi(M_k^k) = \int_{M_k^k} c_{k-2}(T'M_k^k). \quad (4.60)$$

In [20], Zinger proved the following theorem:

Theorem 1 (Zinger [20])

$$\begin{aligned} F_1^{k,k,A}(t) &:= -\frac{1}{24} \left(\int_{M_k^k} c_{k-3}(T'M_k^k) \wedge h \right) t + \sum_{d=1}^{\infty} \langle * \rangle_{1,d} e^{dt} \\ &= F_1^{k,k,B}(x(t)), \end{aligned} \quad (4.61)$$

where $\langle * \rangle_{1,d}$ is Gromov-Witten invariant of M_k^k of genus 1 and of degree d with no operator insertions and $x = x(t)$ is inversion of the mirror map $t = t(x)$ given in (4.53).

At this stage, we look back at Conjecture 2. In the $N = k$ case, $w(\prod_{p=2}^{N-2} (\mathcal{O}_{h^p})^{m_p})_{1,d}$ is non-zero only if $m_2 = m_3 = \dots = m_{N-2} = 0$. Hence $F_{1,vir.}^{k,k,B}$ has the following structure:

$$\begin{aligned} F_{1,vir.}^{k,k,B}(x^1) &= -\frac{1}{24} \left(\int_{M_k^k} h \wedge c_{k-3}(T'M_k^k) \right) x^1 + \sum_{d=1}^{\infty} e^{dx^1} w(*)_{1,d}, \end{aligned} \quad (4.62)$$

where the symbol $*$ represents having no operator insertions. On the other hand, $F_1^{k,k,A}$ also has the same structure:

$$\begin{aligned} F_1^{k,k,A}(t^1) &= -\frac{1}{24} \left(\int_{M_k^k} h \wedge c_{k-3}(T'M_k^k) \right) t^1 + \sum_{d=1}^{\infty} e^{dt^1} \langle * \rangle_{1,d}. \end{aligned} \quad (4.63)$$

Therefore, we only need the mirror map that relates x^1 to t^1 .

$$t^1(x^1) = x^1 + \frac{1}{k} \sum_{d=1}^{\infty} e^{dx^1} w(\mathcal{O}_{h^{k-3}}\mathcal{O}_1|*)_{0,d}. \quad (4.64)$$

Then Conjecture 2 asserts the following equality:

$$F_1^{k,k,A}(t^1) = F_{1,vir.}^{k,k,B}(x^1(t^1)). \quad (4.65)$$

On the other hand, the mirror map which relates x to t in (4.53) has the structure:

$$t(x) = x + \frac{1}{k} \sum_{d=1}^{\infty} e^{dx^1} w(\mathcal{O}_{h^{k-3}\mathcal{O}_1})_{0,d}. \quad (4.66)$$

By comparing the r.h.s.'s of (2.3) and (4.50), we can easily see the equality:

$$w(\mathcal{O}_{h^{k-3}\mathcal{O}_1|*})_{0,d} = w(\mathcal{O}_{h^{k-3}\mathcal{O}_1})_{0,d}. \quad (4.67)$$

holds. In this way, we can identify t^1 and x^1 in Conjecture 2 with t and x in Theorem 1 respectively. This naturally leads us to the following conjecture.

Conjecture 3

$$F_{1,vir.}^{k,k,B}(x) = F_1^{k,k,B}(x). \quad (4.68)$$

In the following, we present results of numerical tests of this conjecture.

By Definition 2, $w(*)_{1,d}$ in (4.62) is given by sum of residues of integrands f_Γ 's associated with $\Gamma \in \text{Graph}_d$,

$$w(*)_{1,d} = \sum_{\Gamma \in \text{Graph}_d} \text{Res}(f_\Gamma). \quad (4.69)$$

We denote by $\text{Graph}_d^{(i)}, \dots, \text{Graph}_d^{(iv)}$ sets of graphs of type (i), \dots , (iv) of degree d respectively. We also denote by Γ_d^{loop} and Γ_d^{point} , the unique graphs that belong to $\text{Graph}_d^{(ii)}$ and $\text{Graph}_d^{(iv)}$ respectively². Then the above equality is rewritten as follows.

$$\begin{aligned} w(*)_{1,d} &= \sum_{\Gamma \in \text{Graph}_d^{(i)}} \text{Res}(f_\Gamma) + \sum_{\Gamma \in \text{Graph}_d^{(ii)}} \text{Res}(f_\Gamma) + \sum_{\Gamma \in \text{Graph}_d^{(iii)}} \text{Res}(f_\Gamma) + \sum_{\Gamma \in \text{Graph}_d^{(iv)}} \text{Res}(f_\Gamma) \\ &= \sum_{\Gamma \in \text{Graph}_d^{(i)}} \text{Res}(f_\Gamma) + \text{Res}(f_{\Gamma_d^{loop}}) + \sum_{\Gamma \in \text{Graph}_d^{(iii)}} \text{Res}(f_\Gamma) + \text{Res}(f_{\Gamma_d^{point}}), \end{aligned} \quad (4.70)$$

By numerical computations for lower k 's and d 's, we obtain the following conjectures.

Conjecture 4 For any positive d and any $\Gamma \in \text{Graph}_d^{(iii)}$, $\text{Res}(f_\Gamma)$ vanishes.

Conjecture 5 The following equality holds.

$$\sum_{d=1}^{\infty} \left(\sum_{\Gamma \in \text{Graph}_d^{(i)}} \text{Res}(f_\Gamma) \right) e^{dx} = \frac{1}{24} \chi(M_k^k) \log \left(\tilde{L}_0^{k,k}(e^x) \right). \quad (4.71)$$

On the other hand, $\text{Res}(f_{\Gamma_d^{point}})$ is immediately computed from (3.34).

$$\begin{aligned} \text{Res}(f_{\Gamma_d^{point}}) &= \frac{1}{24} R_{k,k}(d) k^{kd} \\ &= \frac{1}{24} \left(\frac{(k-1)}{2} \frac{1}{d} - \frac{k^2-1}{k} \frac{1}{d^2} \right) k^{kd}. \end{aligned} \quad (4.72)$$

²Of course, we have to note that $\text{Graph}_1^{(ii)} = \emptyset$.

Hence we obtain,

$$\sum_{d=1}^{\infty} \text{Res}(f_{\Gamma_d^{point}})e^{dx} = -\frac{k-1}{48} \log(1-k^k e^x) + \frac{k^2-1}{24k} \int_{-\infty}^x \log(1-k^k e^s) ds. \quad (4.73)$$

By combining these conjectures, we obtain,

$$\begin{aligned} F_{1, virt.}^{k,k,B}(x) &= -\frac{1}{24} \left(\int_{M_k^k} c_{k-3}(T' M_k^k) \wedge h \right) x + \frac{1}{24} \chi(M_k^k) \log \left(L_0^{k,k}(e^x) \right) \\ &\quad + \sum_{d=2}^{\infty} \text{Res}(f_{\Gamma_d^{loop}})e^{dx} - \frac{k-1}{48} \log(1-k^k e^x) + \frac{k^2-1}{24k} \int_{-\infty}^x \log(1-k^k e^s) ds. \end{aligned} \quad (4.74)$$

This is another representation of BCOV-Zinger formula obtained from Conjecture 3. Therefore, Conjecture 3 is restated as the following identities.

Conjecture 6

$$\begin{aligned} &\sum_{d=2}^{\infty} \text{Res}(f_{\Gamma_d^{loop}})e^{dx} + \frac{k^2-1}{24k} \int_{-\infty}^x \log(1-k^k e^s) ds \\ &= -\sum_{p=0}^{\frac{k-3}{2}} \frac{(k-1-2p)^2}{8} \log \left(\tilde{L}_p^{k,k}(e^x) \right) \quad (k : \text{odd}), \\ &\sum_{d=2}^{\infty} \text{Res}(f_{\Gamma_d^{loop}})e^{dx} - \frac{1}{16} \log(1-k^k e^x) + \frac{k^2-1}{24k} \int_{-\infty}^x \log(1-k^k e^s) ds \\ &= -\sum_{p=0}^{\frac{k-4}{2}} \frac{(k-2p)(k-2p-2)}{8} \log \left(\tilde{L}_p^{k,k}(e^x) \right) \quad (k : \text{even}). \end{aligned} \quad (4.75)$$

We numerically confirmed these identities for $k = 4, 5, 6, 7, 8$ cases up to $d = 5$. We present here numerical data up to $d = 5$ in the $k = 5$ (quintic 3-fold) case.

Generating function of loop amplitudes is given as follows.

$$\begin{aligned} &\sum_{d=2}^{\infty} \text{Res}(f_{\Gamma_d^{loop}})e^{dx} \\ &= -\frac{1174875}{4} e^{2x} - \frac{6913090625}{9} e^{3x} - \frac{31054165371875}{16} e^{4x} - 5008379074144375 e^{5x} - \dots \end{aligned} \quad (4.76)$$

We also have,

$$\begin{aligned} &\frac{1}{5} \int_{-\infty}^x \log(1-5^5 e^s) ds \\ &= -625 e^x - \frac{1953125}{4} e^{2x} - \frac{6103515625}{9} e^{3x} - \frac{19073486328125}{16} e^{4x} - 2384185791015625 e^{5x} - \dots \end{aligned} \quad (4.77)$$

Then, the l.h.s. of (4.75) turns out to be,

$$\begin{aligned} &\sum_{d=2}^{\infty} \text{Res}(f_{\Gamma_d^{loop}})e^{dx} + \frac{1}{5} \int_{-\infty}^x \log(1-5^5 e^s) ds \\ &= -625 e^x - 782000 e^{2x} - \frac{4338868750}{3} e^{3x} - 3132978231250 e^{4x} - 7392564865160000 e^{5x} - \dots \end{aligned} \quad (4.78)$$

On the other hand, $\log\left(\tilde{L}_0^{5.5}(e^x)\right)$ and $\log\left(\tilde{L}_1^{5.5}(e^x)\right)$ are given as follows.

$$\begin{aligned}
& \log\left(\tilde{L}_0^{5.5}(e^x)\right) \\
&= 120e^x + 106200e^{2x} + 155136000e^{3x} + 280511415000e^{4x} + 571399451565120e^{5x} + \dots, \\
& \log\left(\tilde{L}_1^{5.5}(e^x)\right) \\
&= 770e^x + 1139200e^{2x} + \frac{6816105500}{3}e^{3x} + 5143910802500e^{4x} + 12499531924059520e^{5x} + \dots. \quad (4.79)
\end{aligned}$$

And the r.h.s. of (4.75) becomes,

$$\begin{aligned}
& -2\log\left(\tilde{L}_0^{5.5}(e^x)\right) - \frac{1}{2}\log\left(\tilde{L}_1^{5.5}(e^x)\right) \\
&= -625e^x - 782000e^{2x} - \frac{4338868750}{3}e^{3x} - 3132978231250e^{4x} - 7392564865160000e^{5x} - \dots. \quad (4.80)
\end{aligned}$$

In this way, the identity (4.75) in the $k = 5$ case is numerically confirmed up to $d = 5$.

A Tables for N=5 cases

Table 4: M_5^1

d	(a,b)	$N_{d,a,b}^0$	$N_{d,a,b}^1$	$\frac{2d-1}{12}N_{d,a,b}^0 + N_{d,a,b}^1$	$w_{a,b}$
1	(0,2)	1	$-\frac{1}{12}$	0	$-\frac{7}{12}$
1	(2,1)	1	$-\frac{1}{12}$	0	$-\frac{5}{6}$
1	(4,0)	2	$-\frac{1}{6}$	0	$-\frac{7}{6}$
2	(0,4)	0	0	0	$-\frac{76}{3}$
2	(2,3)	1	$-\frac{1}{4}$	0	$-\frac{853}{12}$
2	(4,2)	4	-1	0	-198
2	(6,1)	18	$-\frac{9}{2}$	0	$-\frac{1097}{2}$
2	(8,0)	92	-23	0	$-\frac{4541}{3}$
3	(0,6)	1	$-\frac{5}{12}$	0	$-\frac{19959}{4}$
3	(2,5)	5	$-\frac{25}{12}$	0	$-\frac{62338}{3}$
3	(4,4)	30	$-\frac{25}{2}$	0	$-\frac{516827}{3}$
3	(6,3)	190	$-\frac{469}{6}$	1	$-\frac{1068442}{3}$
3	(8,2)	1312	$-\frac{1598}{3}$	14	$-\frac{4408330}{3}$
3	(10,1)	9864	-3960	150	$-\frac{18159922}{3}$
3	(12,0)	80160	-31900	1500	$-\frac{74719852}{3}$
4	(0,8)	4	$-\frac{4}{3}$	1	$-\frac{7111330}{3}$
4	(2,7)	58	$-\frac{179}{6}$	4	$-\frac{26141813}{2}$
4	(4,6)	480	-248	32	-71830274
4	(6,5)	4000	$-\frac{6070}{3}$	310	$-\frac{1182256279}{3}$
4	(8,4)	35104	$-\frac{51772}{3}$	3220	-2159333004
4	(10,3)	327888	-156594	34674	$-\frac{35458691818}{3}$
4	(12,2)	3259680	-1515824	385656	$-\frac{193936379144}{3}$
4	(14,1)	34382544	-15620216	4436268	-353359995764
4	(16,0)	383306880	-170763640	52832040	-1930689790136
5	(0,10)	105	$-\frac{147}{4}$	42	$-\frac{8363354113}{4}$
5	(2,9)	1265	$-\frac{2379}{4}$	354	$-\frac{28682135389}{2}$
5	(4,8)	13354	$-\frac{13047}{2}$	3492	$-\frac{196198477325}{2}$
5	(6,7)	139098	$-\frac{132549}{2}$	38049	$-\frac{2010681907978}{3}$
5	(8,6)	1492616	-677808	441654	$-\frac{13724961403006}{3}$
5	(10,5)	16744080	-7179606	5378454	$-\frac{93619004917238}{3}$
5	(12,4)	197240400	-79637976	68292324	-212735629674372
5	(14,3)	2440235712	-928521900	901654884	$-\frac{4348697671027760}{3}$
5	(16,2)	31658432256	-11385660384	12358163808	-9873859605646752
5	(18,1)	429750191232	-146713008096	175599635328	$-\frac{201722432909390752}{3}$
5	(20,0)	6089786376960	-1984020394752	2583319387968	$-\frac{1373530281059327936}{3}$

Table 5: M_5^2

d	(a,b)	$N_{d,a,b}^0$	$N_{d,a,b}^1$	$\frac{3d-2}{24}N_{d,a,b}^0 + N_{d,a,b}^1$	$w_{a,b}$
1	(1,1)	4	$-\frac{1}{6}$	0	$-\frac{13}{6}$
1	(3,0)	8	$-\frac{1}{3}$	0	-3
2	(0,3)	8	$-\frac{4}{3}$	0	$-\frac{287}{3}$
2	(2,2)	16	$-\frac{8}{3}$	0	-264
2	(4,1)	64	$-\frac{32}{3}$	0	$-\frac{2174}{3}$
2	(6,0)	320	$-\frac{160}{3}$	0	$-\frac{5956}{3}$
3	(1,4)	64	$-\frac{56}{3}$	0	$-\frac{104500}{3}$
3	(3,3)	320	$-\frac{280}{3}$	0	$-\frac{429196}{3}$
3	(5,2)	2048	$-\frac{1792}{3}$	0	$-\frac{1759552}{3}$
3	(7,1)	15104	$-\frac{13216}{3}$	0	$-\frac{7209584}{3}$
3	(9,0)	123904	$-\frac{108416}{3}$	0	$-\frac{29527616}{3}$
4	(0,6)	384	-160	0	$-\frac{18667312}{3}$
4	(2,5)	2560	$-\frac{3200}{3}$	0	$-\frac{101879272}{3}$
4	(4,4)	18944	$-\frac{22912}{3}$	256	$-\frac{555449168}{3}$
4	(6,3)	163840,	$-\frac{194048}{3}$	3584	$-\frac{3026251616}{3}$
4	(8,2)	1583104	$-\frac{1849856}{3}$	43008	$-\frac{16485590720}{3}$
4	(10,1)	16687104	-6440960	512000	$-\frac{89806327616}{3}$
4	(12,0)	189358080	-72652800	6246400	-163085218816
5	(1,7)	27136	$-\frac{41792}{3}$	768	$-\frac{28726121392}{3}$
5	(3,6)	229376	$-\frac{331264}{3}$	13824	$-\frac{195282001984}{3}$
5	(5,5)	2232320	$-\frac{3049984}{3}$	192512	$-\frac{1326874482304}{3}$
5	(7,4)	24391680	-10660352	2551808	$-\frac{9013280450048}{3}$
5	(9,3)	291545088	-123583488	34336768	$-\frac{61226330115584}{3}$
5	(11,2)	3750199296	-1553444864	477913088	-138652119786496
5	(13,1)	51384877056	-20917362688	6916112384	$-\frac{2826429058966016}{3}$
5	(15,0)	744875950080	-299359264768	104115208192	$-\frac{19209989184830464}{3}$

Table 6: M_5^3

d	(a,b)	$N_{d,a,b}^0$	$N_{d,a,b}^1$	$\frac{d-1}{12}N_{d,a,b}^0 + N_{d,a,b}^1$	$w_{a,b}$
1	(0,1)	18	0	0	$-\frac{21}{2}$
1	(2,0)	45	0	0	$-\frac{27}{2}$
2	(0,2)	54	$-\frac{9}{2}$	0	$-\frac{30501}{4}$
2	(2,1)	378	$-\frac{63}{2}$	0	-2862
2	(4,0)	2187	$-\frac{729}{4}$	0	$-\frac{2187}{2}$
3	(0,3)	648	-81	27	-299943
3	(2,2)	7452	-1161	81	-1188027
3	(4,1)	65610	-10449	486	$-\frac{9537669}{2}$
3	(6,0)	623295	$-\frac{200475}{2}$	3645	-19201644
4	(0,4)	15552	-1701	2187	$-\frac{279086715}{2}$
4	(2,3)	248832	-48357	13851	-740281275
4	(4,2)	2991816	-616734	131220	-3968742582
4	(6,1)	37161504	-7846956	1443420	-21341675475
4	(8,0)	491956902	$-\frac{211336371}{2}$	17321040	$-\frac{229871126583}{2}$
5	(0,5)	583200	972	195372	-92893454856
5	(2,4)	11955600	-2005236	1979964	-617546315223
5	(4,3)	183760488	-35803377	25450119	$-\frac{8281495651131}{2}$
5	(6,2)	2838367332	-584014293	362108151	-27840191130297
5	(8,1)	45746559378	-9717064074	5531789052	-187479083534526
5	(10,0)	776682421065	-169540839261	89353301094	-1263878784214992

Table 7: M_5^4

d	(a,b)	$N_{d,a,b}^0$	$N_{d,a,b}^1$	$\frac{d-2}{24}N_{d,a,b}^0 + N_{d,a,b}^1$	$w_{a,b}$
1	(1,0)	320	$\frac{40}{3}$	0	$-\frac{344}{3}$
2	(0,1)	3888	0	0	$-\frac{84848}{3}$
2	(2,0)	27200	0	0	$-\frac{222080}{3}$
3	(1,1)	672768	-24192	3840	-32895232
3	(3,0)	8388608	$-\frac{971776}{3}$	25600	$-\frac{390036992}{3}$
4	(0,2)	18323712	-861696	665280	-13842672128
4	(2,1)	284802048	-14229504	9504000	$-\frac{221036603392}{3}$
4	(4,0)	5100273664	$-\frac{905199616}{3}$	123289600	$-\frac{1171415220224}{3}$
5	(1,2)	9830744064	-148801536	1080041472	$-\frac{113572126965760}{3}$
5	(3,1)	206561083392	-4948770816	20871364608	-252561662754816
5	(5,0)	4821100789760	-202932748288	399704850432	$-\frac{5041038692581376}{3}$

References

- [1] M. Bershadsky, S. Cecotti, H. Ooguri,; C. Vafa. *Holomorphic anomalies in topological field theories*, Nuclear Phys. B405 (1993), no.2–3, 279–304.
- [2] M. Bershadsky, S. Cecotti, H. Ooguri,; C. Vafa. *Kodaira-Spencer theory of gravity and exact results for quantum string amplitudes*. Comm. Math. Phys.165 (1994), no.2, 311–427.
- [3] A. Collino, M. Jinzenji. *On the structure of the small quantum cohomology rings of projective hypersurfaces*. Comm. Math. Phys. 206 (1999), no. 1, 157–183.
- [4] I. Ciocan-Fontanine, B. Kim. *Wall-crossing in genus zero quasimap theory and mirror maps*. Algebr. Geom. 1 (2014), no. 4, 400–448.
- [5] T.-M. Chiang, A. Klemm, S.-T. Yau, E. Zaslow. *Local mirror symmetry: calculations and interpretations*. Adv. Theor. Math. Phys. 3 (1999), no. 3, 495–565.
- [6] E. Getzler. *Intersection theory on $\overline{\mathcal{M}}_{1,4}$ and elliptic Gromov-Witten invariants*. J. Amer. Math. Soc. 10 (1997), no. 4, 973–998.
- [7] T. Graber; R. Pandharipande. *Localization of virtual classes..* Invent. Math.135(1999), no.2, 487–518.
- [8] T. Eguchi, K.Hori, C.-S. Xiong. *Quantum Cohomology and Virasoro Algebra*. Phys. Lett. B402 (1997), no.1–2, 71–80.
- [9] T. Eguchi, M.Jinzenji, C.-S. Xiong. *Quantum cohomology and free-field representation*. Nuclear Phys. B510 (1998), no.3, 608–622.
- [10] M. Jinzenji. *Virtual structure constants as intersection numbers of moduli space of polynomial maps with two marked points*. Lett. Math. Phys.86(2008), no.2-3, 99–114.
- [11] M. Jinzenji. *Mirror Map as Generating Function of Intersection Numbers: Toric Manifolds with Two Kähler Forms*, arXiv:1006.0607, Comm. Math. Phys. 323(2013), no. 2, 747–811.
- [12] M. Jinzenji. *Geometrical Proof of Generalized Mirror Transformation of Projective Hypersurfaces*. Internat. J. Math. 34 (2023), no. 2, 2350006, 17 pp.
- [13] M. Jinzenji. *Classical Mirror Symmetry*. SpringerBriefs Math. Phys. 29, Springer, Singapore, 2018. viii+140 pp.
- [14] M. Jinzenji, M. Shimizu. *Multi-Point Virtual Structure Constants and Mirror Computation of CP^2 -model*. Commun. Number Theory Phys. 7 (2013), no. 3, 411–468.
- [15] M. Jinzenji. ResearchGate homepage. <https://www.researchgate.net/profile/Masao-Jinzenji>
- [16] M. Kontsevich. *Enumeration of rational curves via torus actions*. The moduli space of curves (Texel Island, 1994), 335–368, Progr. Math., 129, Birkhauser Boston, Boston, MA, 1995.
- [17] M. Kontsevich, Yu. Manin, *Gromov-Witten classes, quantum cohomology, and enumerative geometry*. Comm. Math. Phys.164(1994), no.3, 525–562.
- [18] R. Pandharipande. *Hodge integrals and degenerate contributions*. Comm. Math. Phys.208(1999), no.2, 489–506.
- [19] K. Sugiyama. *The Calabi-Yau moduli and the Toda equations*. Nuclear Phys. B459(1996), no.3, 693–723.
- [20] A. Zinger. *The reduced genus-one Gromov-Witten invariants of Calabi-Yau hypersurfaces*. J. Amer. Math. Soc. 22 (2009), 691-737.